

## Scientific history, sampling approach, and physical characterization of the Camp Century sub-glacial material, a rare archive from beneath the Greenland Ice Sheet

Paul R. Bierman<sup>1,2</sup>, Andrew J. Christ<sup>1\*</sup>, Catherine M. Collins<sup>1,2</sup>, Halley M. Mastro<sup>1,2</sup>, Juliana Souza<sup>1,2</sup>, Pierre-Henri Blard<sup>3,4</sup>, Stefanie Brachfeld<sup>5</sup>, Zoe R. Courville<sup>6</sup>, Tammy M. Rittenour<sup>7</sup>, Elizabeth K. Thomas<sup>8</sup>, Jean-Louis Tison<sup>4</sup>,  
5 and Francois Fripiat<sup>4</sup>

<sup>1</sup> Rubenstein School of the Environment and Natural Resources, University of Vermont, Burlington, VT, 05401, USA

<sup>2</sup> Gund Institute for Environment, University of Vermont, Burlington, VT, 05401, USA

<sup>3</sup> Centre de Recherches Pétrographiques et Géochimiques, CNRS, Université de Lorraine, 54500 Nancy, France

10 <sup>4</sup> Laboratoire de Glaciologie, Université Libre de Bruxelles, Brussels, 1050 Bruxelles, Belgium

<sup>5</sup> Earth and Environmental Studies, Montclair State University, Montclair, NJ 07043, USA

<sup>6</sup> US Army Corps of Engineers Cold Regions Research and Engineering Laboratory, Hanover, NH 03755, USA

<sup>7</sup> Department of Geosciences, Utah State University, Logan, UT 84322, USA

<sup>8</sup> Department of Geology, University at Buffalo, Buffalo, NY 14260, USA

15 \*Current affiliation: US Reinsurance Analytics, Aon plc, Denver, CO 80206, USA

*Correspondence to:* Paul Bierman, pbierman@uvm.edu

**Abstract.** Basal materials in ice cores hold information about paleoclimate conditions, glacial processes, and the  
20 timing of past ice-free intervals, all of which aid understanding of ice-sheet stability and its contribution to sea-  
level rise in a warming climate. Only a few cores have been drilled through ice sheets into the underlying  
sediment and bedrock, producing limited material for analysis. The last of three Camp Century ice cores, which  
the US Army collected in northwest Greenland from 1963-1966 CE, recovered about 3.5 meters of sub-glacial  
25 material including ice and sediment. Here, we document the scientific history of the Camp Century sub-glacial  
material, and present our recent core-cutting, sub-sampling, and processing methodology and results for this  
unique archive.

In 1972 CE, curators at the Buffalo, New York Ice Core Laboratory cut the original core sections into 32  
segments each about 10-cm long. Since then, two segments were lost and are unaccounted for, two were thawed,  
and two were cut as pilot samples in 2019 CE. Except for the two thawed segments, the rest of the extant core  
30 remained frozen since collection. In 2021 CE, we documented, described, and then cut each of the remaining  
frozen archived segments (n=26). We saved an archival half and cut the working half into eight oriented sub-  
samples under controlled temperature and light conditions for physical, geochemical, isotopic, sedimentological,  
magnetic, and biological analyses. Our approach was designed to maximize sample usage for multi-proxy

35 analysis, minimize contamination, and preserve archive material for future analyses of this legacy sub-glacial material.

Grain size, bulk density, sedimentary features, magnetic susceptibility, ice content, as well as pore-ice pH and conductivity, suggest that the basal sediment contains five stratigraphic units. We interpret these stratigraphic units as representing different depositional environments in sub-glacial or ice-free conditions: from bottom to top, a diamicton with sub-horizontal ice lenses (Unit 1); vertically-fractured ice with dispersed fine-grain sediments (< 20% in mass) (Unit 2); a normally graded bed of pebbles to very fine sand in an icy matrix (Unit 3); bedded very fine to fine sand (Unit 4); and stratified medium to coarse sand (Unit 5). Plant macrofossils are present in all samples and most abundant in Units 3 and 4; insect remains are present in some samples (Units 1, 3, and 5).

Our approach provides a working template for future studies of ice-core basal materials because it includes intentional planning of core sub-sampling, processing methodologies, and archiving strategies to optimize the collection of paleoclimate, glacial process, geochemical, geochronological, and sediment properties from archives of limited size. Our work benefited from a carefully curated and preserved archive, allowing for the application of analytical techniques not available in 1966 CE. Preserving uncontaminated core material for future analyses that use currently unavailable tools and techniques is an important consideration for rare archive materials such as these from Camp Century.

## 50 **1. Introduction**


Ice cores are paleoclimate archives that offer valuable insights into past atmospheric composition and temperature via the analysis of trapped air and physical and chemical properties of ice. Ice cores typically recover clean glacial ice ('clear ice'), and only a few ice cores have reached the ice-bed interface to recover basal materials such as sediment-laden ice (sometimes referred to as 'silty ice' or 'basal ice'), sub-glacial sediment, and sub-glacial bedrock (Talalay, 2013).

Isotopic, mineralogic, and geochemical analysis of basal materials from ice cores yields important information about basal ice-sheet processes (Goossens et al., 2016; Souchez et al., 2000; Herron and Langway, 1982; Gow and Meese, 1996; Verbeke et al., 2002; Tison et al., 1994), sub-glacial geology and sediment provenance (Blard et al., 2023; Fountain et al., 1981; Licht and Hemming, 2017; Weis et al., 1997), the timing and duration of ice-free intervals or grounding-line retreat (Schaefer et al., 2016; Venturelli et al., 2020), and the ecosystems that occupied formerly ice-free landscapes now buried beneath the ice (Christ et al., 2021; Tison et al., 1998; Souchez et al., 2006; Willerslev et al., 2007). However, the amount of material retrieved from beneath ice sheets is minimal; sometimes only tens of grams of rock or sediment are available for analysis (Talalay, 2013). Given the specific requirements for sample mass, grain size, and handling needed to enable a wide variety of analyses, strategic sampling and sample distribution protocols are critical to optimize results obtained from such limited quantities of archive material.

The United States Army drilled three ice cores between 1960 to 1966 CE in northwestern Greenland at Camp Century. The last core (1963-1966) was the first to reach the bed of any ice sheet and was also the first to recover a core of sub-glacial material. Until 2023 and the start of Greendrill coring, the Camp Century core contained the longest sub-glacial sedimentary sequence recovered from Greenland (Hansen and Langway, 1966) (Figures 1A, 2A). Because the sub-glacial material was not extensively sampled at the time of collection and has

since been stored frozen, we were able to design a detailed, intentional sub-sampling and analysis plan for multiple geochemical, physical, magnetic, biological, and isotopic proxies.

The purpose of this paper is four-fold. First, we compile and summarize the historical record of the collection, handling, and prior analyses of the Camp Century sub-glacial material to provide clear context for ongoing and future analyses (Section 2). Second, we detail and explain our approach to cutting, sub-sampling, and distributing this unique sediment archive to optimize sample usage for multi-proxy analysis, avoid contamination, and preserve material for future studies (Section 3). Third, we present our documentation of the archive and the data collected during the sub-sampling and initial processing of the sediment, which allow us to describe the stratigraphy of the Camp Century sub-glacial material (Section 4). Fourth, we provide initial interpretations of the depositional history of the material and provide lessons learned for handling, sub-sampling, and storing rare archives of ice-core basal materials (Section 5).




**Figure 1: Location and context of the Camp Century core. (A) Overview map of Greenland showing the locations of Camp Century (red circle) and other deep ice cores (dark circles). (B) Schematic layout of the Camp Century installation with the location of the drill trench marked with a red star (Daugherty, 1963)**

## 2. The scientific history of ice-core drilling at Camp Century and the sub-glacial material core

In the 1950s CE, the U.S. military was concerned that the Arctic was warming (Thomis, 1955) although warming then turned out to be short-lived. Such warming had implications for the ongoing militarization of the Arctic during the Cold War, but little was then known about past changes in Arctic climate (Doel et al., 2017). To address this knowledge gap, Henri Bader and James Bender, two civilian scientists from the Snow, Ice, and Permafrost Research Establishment (SIPRE) in the U.S. Army Corps of Engineers, lobbied for drilling deep cores into and through the Greenland Ice Sheet (Bader, 1962). Bader had overseen drilling of the first U.S. ice core in Alaska from the Taku Glacier in 1950 CE and Bender had suggested that bubbles trapped in glacial ice preserved bits of ancient atmosphere, the analysis of which could resolve questions about fossil fuel combustion and its

relation to climate change (Anon, 1959). Along with Chester Langway, who at the time was a junior researcher at SIPRE, they oversaw drilling of ice cores into the Greenland Ice Sheet at Site 2 (about 100 km northeast and up ice from Camp Century) in 1956 CE and 1957 CE. Although recovery was poor at Site 2 and the cores were only  
100 a few hundred meters long, they made advances in drilling technology.

Analyses on small continuous sections of the 1957 Site 2 ice core demonstrated that the ice preserved records of past climate including annual variations of stable water isotopes (Epstein and Sharp, 1959; Langway, 1958). The full potential of extracting paleoclimatic information from a deep core was revealed when Langway joined forces with Willi Dansgaard in 1967 to perform systematic stratigraphic water isotope analysis of the  
105 Camp Century clear-ice core. Dansgaard, who had been considering the precipitation vs. temperature relationship in stable water isotopes since 1953 (Dansgaard, 1953 and 1954) had the only lab in the world at that time capable of analyzing thousands of samples in a timely fashion. The first 100,000 year-long paleoclimatic record (based on stable water isotopes) was published in 1969 (Dansgaard, 1969). Later, Werner Berner, who was provided Camp Century ice by Langway, extracted gasses from bubbles in the ice and quantified their concentration (Berner,  
110 1978).



115 **Figure 2: Photographs of the Camp Century sub-glacial material core. (A) Historic photograph of the**  
**Camp Century sub-glacial material core segments (by tube number) prior to initial cutting in October**  
**1972 CE (Fountain et al., 1981). (B) Photograph in A with core piece (from cutting in 1972 CE) identifiers**  
**overlain. (C) New core photographs (2021 CE) and 2D rendering of photogrammetric 3D models of the**  
**segments rearranged into their original positions. Shaded cylinders: green – no 3D model available, red –**  
**missing sample, blue – thawed sample. (D) New core photographs (2021 CE) showing interior cut faces of**  
120 **the archived halves of each piece. Note: the photos of interior cut faces (in D) are not all oriented the same**  
**manner as the original core (A and B). Scale bar in upper right of each panel.**

Camp Century was a U.S. Army base carved into the surface of the Greenland Ice Sheet. It was built as  
part of the American militarization of the Arctic (Clark, 1966). The base was ~200 km inland from the ice margin  
in northwest Greenland at 1890 m asl (Figure 1A). Construction of the camp began in June 1959 CE and was  
completed in October 1960 CE. Camp Century was constructed in snow and firn using cut-and-cover methods  
and could house up to 200 people in heated barracks nestled within the snow tunnels. From 1961 until Northern  
Hemisphere summer 1963 CE, the base was occupied year-round – powered and heated by a portable nuclear  
reactor that was removed in 1964 CE (Corliss, 1968). The U.S. military operated Camp Century with a science-  
driven engineering program to assess novel and applied technologies that allowed people to live and work in  
extreme environments. Much of this scientific effort was applied and directed toward better understanding snow  
compaction and firn deformation, both on the surface of the ice sheet and in the tunnels of Camp Century, which  
deformed at rates of 0.1 to 1.0 m/yr (Abele, 1964).

A drilling trench was included in the plans for the base (Figure 1B). Personnel from SIPRE, the  
predecessor of the Cold Regions Research and Engineering Laboratory (CRREL), conducted research on the  
physical properties of ice and snow and led collection of the Camp Century ice core (Figure 3A). A new thermal  
ice drill was designed for Camp Century and deployed from 1960 to 1964 CE, but it was plagued with problems  
related to the removal and collection of meltwater from the borehole (Bader, 1962; Hansen, 1994). The thermal  
drill extracted three ice cores: between 1960 and 1961 CE the first hole recovered 186 m, in 1962 CE the second  
hole recovered 238 m, and in 1963 CE the third hole recovered 264 m. The last hole was then filled with a fluid  
mixture of 12% trichloroethylene and 88% Arctic diesel fuel when Camp Century closed for the winter and  
drilling halted (Ragone and Finelli, 1972). In 1964 CE, the fluid-filled third hole was extended to a depth of 535  
m. Starting in 1965 CE, an electromechanical drill that used ethylene glycol to melt ice chips in addition to the  
drilling fluid (Langway and Hansen, 1970; Ueda and Garfield, 1968) advanced more rapidly and extended the  
third core to 1002 m. In 1966 CE, drilling resumed with the electromechanical drill. Silty ice was encountered  
near the bottom of the hole and coring was extended into the ice-sheet bed.

145 The sub-glacial material recovered in July 1966 is what we describe in this paper. As material came out  
of the hole, it was placed into 1.5 m aluminized tubes. These tube numbers are used as primary sample identifiers.  
The ice-sediment interface is in core tube 1059; the base of the Camp Century core is in tube 1063. The original  
drill log (Table 2 and Figure 4) from July 2<sup>nd</sup>, 1966 CE, describes the sub-glacial sediment as:

150 “[Core tube 1059] 30 cm from top silty ice stops and frozen sand begins. [Core tube  
1060] Appears to be a column of frozen sand with some rocks as large as 5 cm in  
diameter. Permacrete down to 80 cm then a silty ice lens for rest of core. Red, grey,

maroon, and green rocks as large as 5 cm in diameter. [Core tube 1061] Silty ice lens down to 0.80 m. Sand and large rocks after 0.80 m. [Core tube 1062] Large rocks and sand, rocks, and sand. [Core tube 1063] Sand and large rocks (17 cm).”


155 There are discrepancies in the reported drilling depths and thicknesses of clear ice, silty  
ice, and sub-glacial sediment between the original core log, early literature, and later studies of  
the Camp Century basal materials (Table 1). The core tube log appears to have been amended  
with multiple, different entries (in differing pencil weights and ink colors) for depth of core below  
the surface (Table 2 and Figure 4). While the reported “depth below the surface” of the sub-ice  
160 material equates to 4.5 m of penetration (Table 2), the sum of the reported lengths of sediment  
recovered (on the drilling log) in each core tube is 3.55 m. Thus, it is possible that core recovery  
was incomplete. However, the original core logs (both tabular and graphical) have notes that read  
“These depths are in error”. Given the rigidity of the frozen material, core compaction is unlikely.  
Drilling depths were not simple to discern accurately as coring was done by wireline and not  
165 using rods. Additionally, there are several subtraction errors on the hand-written sheet (Figure 4).

We conclude it is not possible with the information available in 2024 CE to figure out  
whether recovery of the sub-ice material was complete or not. Furthermore, the depths and ice  
thicknesses documented in the core log are slightly different than those reported by (Hansen and  
Langway, 1966):

170 “...at a depth of 1,370.5 meters, ice containing silt bands and small pebbles was  
encountered. On July 4, after drilling through 16.9 meters of this material, the interface at  
the bottom of the ice cap was reached at a depth of 1,387.4 meters. The bottom material  
is frozen till; 3.55 meters of this material was recovered. The total depth of the hole is  
1,391 meters.”

175 Early reports about drilling activities at Camp Century (Ueda and Garfield, 1968) and investigations of  
the basal materials (Herron and Langway, 1979; Whalley and Langway, 1980; Fountain et al., 1981) are in closer  
agreement with Hansen and Langway, (1966) and report that the thickness of the silty ice and sub-glacial  
material are 16 m and 3.5 m, respectively (Table 1). Note that while most publications report the sub-glacial core  
was completed on July 4, 1966, the drilling log has the date of July 2, 1966.

180 Later in the Northern Hemisphere summer of 1966 CE, Camp Century was abandoned, and the ice and  
sub-glacial cores were transported frozen (likely with dry ice as was used for previous cores, but with no record  
of temperature monitoring) from Greenland to the United States and stored in the CRREL freezers in and/or near  
Hanover, New Hampshire (Figure 3). At some point between 1966 CE and 1972 CE, the five core tubes  
185 containing the sub-glacial sediment were photographed alongside each other; the only known photograph of the  
intact sub-glacial archive is presented in Fountain et al., (1981) (Figures 2A, 3B).



190

**Figure 3: Timeline of the scientific history of the Camp Century sub-glacial sediment including changes in storage location and core cutting and sub-sampling. The duration of drilling campaigns at other Greenland deep ice core sites (Fig 1A) is shown for comparison. Photo credits: (A) (Langway, 2008; Fountain et al., 1981)(C-G) A. Christ.**

195

TABLE 1: Discrepancies between reported drilling depths and thicknesses

Source	Clean glacial ice			Silty ice			Subglacial sediment		
	Top depth (masl)	Bottom depth (masl)	Thickness (m)	Top depth (masl)	Bottom depth (masl)	Thickness (m)	Top depth (masl)	Bottom depth (masl)	Thickness (m)
Original core log	0	1354.86	1354.86	1354.86	1373.2	17.34	1373.2	1377.7	3.55
Hansen and Langway, 1966	0	1370.5 <sup>a</sup>	1370.5	1370.5 <sup>a</sup>	1387.4 <sup>a</sup>	16.9	1387.4	1391	3.55
Ueda and Garfield, 1968 <sup>b</sup>	0	1356.6	1356.6	NR	NR	NR	1356.6	1360	3.4
Heron and Langway, 1979	0	1359 <sup>c</sup>	1359	1359 <sup>c</sup>	1375	16	NR	NR	NR
Whalley and Langway, 1982	NR	NR	NR	NR	NR	NR	NR	NR	3.5
Fountain et al. 1981	NR	NR	NR	NR	NR	16	NR	NR	3.5
Recent documentation (2019 - 2021 CE)	NR	NR	1368	NR	NR	17	NR	NR	3.44

NR: not reported; masl: meters above sea level

<sup>a</sup> Calculated based on reported thickness

<sup>b</sup> Reported values in feet were converted to meters. This source did not mention silty ice and included only the depths/thickness of ice and sub-glacial material.

<sup>c</sup> Calculated based on reported drilling depth

Table 2: Transcribed core tube log of sub-ice materials

CORE TUBE LOG								
Date	2-Jul-66	Observer	Donovan	Location	Camp Century			
Unexplained notes written on side <sup>a</sup>	Core Tube No.	Length of Core (m)	Depth of Core Below Surface, Top (m)		Condition of Core <sup>b</sup>	Missing or Removed Sections <sup>b</sup>	Reason <sup>b</sup>	Remarks
	897	1055	1.48	1369.26	2 pieces silty			1059 to 1063 is sub-ice
		1056	1.45	1370.74	3 pieces silty			
	898	1057	0.26	1372.19				
			1.19	1373.5	1 piece silty			
		1058	0.22	1374.69	Silty; a considerable portion of sand on end			These depths are inferred, possibly due to the fact that some core was left down the hole
			1.18	1371.92	Silty piece			
	899	1059	0.64	1373	30 cm from top silty ice starts and frozen sand begins			Cut into 10 cm pieces - 10/13/72 <sup>c</sup>
		1060	0.96	1373.64	Appears to be a column of frozen sand with some rocks as large as 5 cm in diameter. Permacrete down to 80 cm then a silty ice lens for rest of core			= c, d
		1061	0.97	1376.73	Red grey maroon and green rocks as large as 5 cm in size. Silty ice lens down to 0.8m			= c, d
					Sand and large rocks after 0.80 m			
	900	1062	0.19	1377.7	Large rocks and sand			= c, d
			0.26	1377.89	Rocks and sand			
	901	1063	0.83	1377.7	Sand and large rocks (10 cm)			= c, d
				0.83 <sup>e</sup>				
				1378.53 <sup>e</sup>				
				1377.70 <sup>f</sup>				
				1373.20 <sup>f</sup>				
				3.50 <sup>g</sup>				

<sup>a</sup> Notes written on the side of the log that do not correspond to a table heading that might refer to a box number in which samples were stored.

<sup>b</sup> Condition of core descriptions were written across the two columns to the right.

<sup>c</sup> This note was written in a different ink color and clearly was added later.

<sup>d</sup> Interpreted to mean 'same as above'.

<sup>e</sup> Purpose of these notes is unclear.

<sup>f</sup> Upper and lower depth range of sub-ice sediment.

<sup>g</sup> Sub-ice sediment thickness.



CORE TUBE LOG

DATE: 2 July 66 OBSERVER: Donovan LOCATION: Camp Century

Core Tube No.	Length of Core (m)	Depth of Core Below Surface, Top (m)	Condition of Core	Missing or Removed Sections	Reason	Remarks
897 { 1055 ✓	1.48	1369.26	2 pieces silty			1059 to 1063 is sub-ice
1056 ✓	1.45	1370.74	3 pieces silty			
898 {	26	1372.19				These depths are incorrect, probably due to the fact that some core pieces were left dry in the
	1057 ✓	1.19	1373.50	1 piece silty		
899 {	22	1374.69	silty a considerable portion of sand on end			Cut in 10 cm pieces 10/13/72
	1058 ✓	1.18	1371.92	silty 1 piece		
1059 ✓	.64	1373.00	30 cm from top silty ice starts and frozen sand begins			
1060 ✓	.96	1373.64	appears to be a column of frozen sand with some rocks as large as 5 cm in diameter Permafrost down to 80 cm. Then a silty ice lens for rest of core red grey maroon + green rocks as large as 5 cm in dia.			
900 {	1061 ✓	.97	silty Ice lens down to .80 m Sand + large rocks after .80 m			
	.19	1371.70	large rocks + sand			
	1062 ✓	.26	1377.81	rocks + sand		
901 {	1063 ✓	0.83m	Sand + large rocks (17 cm)			
		8.733 1373.53				
		1377.70				
		1373.20				
		3.50				

pieces

Figure 4: Scan of original drilling log from Camp Century courtesy of J.P. Steffensen.

200

In October 1972 CE, the sub-glacial material from each core tube was cut into 32 segments, each about 10 cm long (Figure 3C, Figure 4). The cutting of the sub-glacial material preceded Chester Langway's move from CRREL to the University at Buffalo in 1974. The date and sample name were recorded on tags kept with each cylindrically-shaped segment in glass jars. Each segment was given a sample name that included the tube number and a sequential number or letter (e.g., 1059-4, 1061-D1). The reasoning behind the sample naming

205

scheme is not explained in any existing records but appears to be related to sedimentological properties seen on the exterior of the core (e.g., sediment and ice content, grain size). The entirety of the Camp Century basal ice and sub-glacial material was then transferred to the U.S. National Science Foundation (NSF) Ice Core Storage Facility at University at Buffalo where it remained from 1974 to c. 1993 CE.

210

There was renewed interest in the Camp Century ice core basal materials in the late 1970s and early 1980s CE. Susan L. Herron's dissertation research (at University at Buffalo) analyzed the ice fabric, debris content, and gas concentrations of the silty ice (Herron and Langway, 1982, 1979). Basic petrological descriptions and some Rb-Sr dating were performed on exterior pebbles and cobbles from core tubes 1062 and 1063 (Fountain et al., 1981). A scanning electron microscope (SEM) analysis of grain shape was conducted on sand from an unspecified portion of the sub-glacial material (Whalley and Langway, 1980). Along with five samples of the basal silty ice retained with the Camp Century ice core, segment 1060-A2 was cut during the 1980s CE for investigation for microfossils (Figure 3D) (Harwood, 1986). That sample held abundant freshwater diatoms and a few marine diatoms. After Harwood (1986), there is no documented investigation of the Camp Century sub-glacial material.

220

In 1993 CE, the ice cores stored at University at Buffalo, including most of the Camp Century ice core archive, were moved to Denver when the U.S. National Science Foundation contract for ice-core curation and storage was awarded to the University of Colorado for a new, purpose-built ice-core storage facility. At or after that time, the lowermost 131 m of the Camp Century ice core, including the silty ice and sub-glacial material, were transferred from University at Buffalo to the ice core facility at the Centre for Ice and Climate at the Niels Bohr Institute (NBI) at the University of Copenhagen, where they have remained in frozen storage in the original boxes from the University at Buffalo (Figure 3E). Most U.S. researchers and National Science Foundation staff concluded that the lower 131 m of ice, silty ice, and subglacial sediment were lost because they were no longer with the rest of the ice core in Denver.

230


In 2019, the NBI centralized all ice core samples in a new single freezer storage facility, and in the process of doing a full inventory, the curator of ice cores in Copenhagen found the samples of sub-glacial material, and in the NBI archives he also found the original core logbooks and paper log rolls. He then brought this discovery to the attention of the ice-core community. Later in 2019 CE, sedimentological properties on the exterior of core segments were described and two pilot samples were cut using a diamond-wire saw from the upper-most (1059-4) and lower-most (1063-7) segments of the sub-glacial material (Figure 3F) (Christ et al., 2021). Multiple geochemical and isotopic analyses of these two pilot samples showed that the Camp Century sub-glacial material held a multi-million-year-old record of glacial and vegetation history of northwest Greenland (Christ et al., 2021). In 2021 CE, 55 years after the sub-glacial material core was collected and 49 years to the month after it was cut into ~10-cm segments, we documented, cut, and sub-sampled the remaining frozen basal material for multiple analyses (Figure 3G). In 2023, Christ et al. (2023) determined, using luminescence measurements, that the upper-most sample of sub-glacial material was last exposed to sunlight at  $416 \pm 38$  ka, during the Marine Isotope Stage 11 interglacial.

240

### 3. Methods

#### 245 3.1 Sample documentation prior to sub-sample cutting

We documented the archived inventory of core segments of the Camp Century sub-glacial material in the freezer laboratory (-20 °C) at the Ice Core Facility of the Centre for Ice and Climate, NBI, at the University of Copenhagen, Denmark. We recorded the label name, mass, and dimensions for each core segment found in the archive (Table S1). We noted ice content, grain size, and visually observable cryo-sedimentary structures for each sample. We tracked the storage orientation (i.e., top direction) of each segment during the entirety of handling and cutting. We then compared the segment name, dimensions, and storage orientation to the original core-tube logs and core photograph (Fountain et al., 1981) to reconstruct the length of material recovered in each core tube and to identify samples that were previously sampled, thawed, or missing. We estimate that there were 3.44 m of recovered core material based on what is physically present in the inventory and estimating the length of samples that were thawed (n=2) and those now missing (n=2) from the archive.



**Figure 5: Sample cutting in 2021 CE.** A. Scaled schematic of how samples were cut using the band saw (bright purple lines) and diamond wire saw (bright green lines) in the -20 °C climate-controlled laboratory at NBI. Samples were cut under red light conditions (red light symbol) until sub-sample e was cut and packaged in aluminum foil. All numbers refer to dimension in cm, h = height. B. Sub-samples color-coded according to intended analyses. Geochronometry measurements in quadrants a and b include (U-Th)/He and solid-state measurements by LA-ICP-MS.

For the remaining samples not analyzed in the pilot study (Christ et al., 2021), we took approximately 20 digital photographs of each frozen core segment from all directions (excluding the thawed and missing samples). Using these photographs, we constructed three-dimensional photogrammetric models with 3DF ZephyrLite<sup>®</sup> software using default settings to produce a textured mesh for each sample and exported them as 3D objects. We imported the textured meshes into Adobe Dimension<sup>®</sup> and compared them against the original core photographs

270 (Fountain et al., 1981) to reconstruct each core tube into its original orientation, determine if samples were stored  
upside down, and to identify any missing exterior pebbles (images archived at:  
[https://www.morphosource.org/concern/cultural\\_heritage\\_objects/000583438](https://www.morphosource.org/concern/cultural_heritage_objects/000583438)). We note that the core  
reconstruction using 3D photogrammetric models and comparison against historical photographs occurred  
following sediment sub-sampling due to time constraints during the 2021 CE sub-sampling event. Using Adobe  
275 Photoshop® we calculated the color of the frozen sediment by extracting the average RGB value of the exterior  
and interior cut face of each core segment from digital photographs, converting that RGB value to a HEX color  
code, and then matching to the best fit Munsell soil color (Table S2).

In the climate-controlled freezer laboratory at NBI, we measured the mass and dimensions (height,  
radius) of each frozen core segment to calculate the volume and bulk density. Using this data, we derived an  
280 estimated percent ice content by mass, assuming that this material is a two-component mixture of ice and  
sediment with densities of 0.9 g/cm<sup>3</sup> and 2.7 g/cm<sup>3</sup>, respectively (Table S3). We assumed that core segments had  
cylindrical geometries for the purpose of calculating the volume. To calculate the volume of oddly shaped  
segments, we measured the dimensions of specific individual segments, approximating the segment volume to the  
volume of “best-fit” simple 3-D geometries.

### 285 3.2 Sub-sample cutting design, shipping, and distribution


We designed our core-cutting method to ensure sufficient sample material for varied analyses, retain the  
orientation of samples during storage, avoid further chemical and/or biological contamination and exposure to  
light, and save archival material at NBI for future analysis. Using the mass and dimensional measurements, and  
derived bulk density and ice content for each core segment, we trigonometrically calculated the dimensions of  
290 individual frozen sub-samples needed to satisfy sample-size and mass requirements for multiple analyses (Table  
3, Figure 5). We cut samples into eight oriented sub-samples (a-g) (Figure 5A) using two different saws in the -20  
°C cold-room at NBI: (1) a Gryphon C-40 CR Tall Diamond Band Saw (“AquaSaw XL”) with a 1 mm-thick  
stainless steel diamond blade (Gryphon 301SS-42) using ethylene glycol as a cutting fluid and (2) a Well  
Diamond-Wire Saw, Model 6234 with a 0.5 mm diameter diamond wire according to the methods of Tison  
295 (1994). Smaller pieces and those where we needed to avoid glycol contamination were cut with the diamond wire  
saw.

We cut core segments from the entire sub-glacial section of the core (e.g., segments 1059-5 through  
1063-8). After cutting each sample, we removed the bandsaw from the freezer and cleaned it by wiping down and  
washing working surfaces with tap water and a clean sponge, and then allowed to air dry. We drained and  
300 replaced the glycol cutting fluid after every other sample. We saved two aliquots of the ethylene glycol cutting  
fluid for chemical analysis to identify any potential contamination to pore-ice and biological analyses. We  
replaced the bandsaw blade when it became dull, typically every 2 to 3 samples. We replaced the diamond wire  
when necessary, typically after a total cutting length of ~30 cm for the sediment-rich samples, or if the wire  
snapped during cutting. To prevent light exposure of the interior prism for luminescence measurements, we cut  
305 sub-samples (a) through (e) in red-light conditions using photography darkroom lamps and red LED bicycle  
lights. We handled all sub-samples with nitrile gloves and wrapped sub-sample (f2), which was devoted to  
biological analyses, in plastic wrap.

Table 3: Analytical requirements for intended analyses


Count	Medium	Analysis	Purpose	Allocated sub-sample(s)	Thermal state	Minimum mass of untreated sample material required	Grain size (µm)	Special handling	Designated Laboratory
1	Frozen intact sample	Luminescence dating	Chronology	e	Frozen	40 g		Collected and measured in red light conditions	USU
2	Sediment	Cosmogenic 10Be, 26Al quartz	Chronology, surface weathering and erosion	a,b	Dried	60 g	250-850		UVM, Columbia
3	Sediment	Cosmogenic 21Ne quartz	Chronology, surface weathering and erosion	a,b	Dried	3 g	250-850		Columbia, CRPG
4	Sediment	Cosmogenic 36Cl feldspar	Chronology, surface weathering and erosion	a,b	Dried	20 g	>850		Columbia
5	Sediment	Meteorite 10Be	Chronology, surface weathering and erosion	a,b	Dried	2 g	bulk, 250-500, 500-850		UVM, CRPG
6	Sediment	Background radiation and bulk geochemistry	Luminescence dose rates, 36Cl, physical characterization	d	Dried	10 g	bulk		USU
7	Pore ice meltwater	Water stable isotopes	Paleoclimate	a	Liquid	0.25 mL		Stored in plastic, filtered	UW
8	Frozen intact sample	Ancient DNA	Paleoecology	f2	Frozen	what is available	bulk	Wrapped in plastic wrap	NBI
9	Pore ice meltwater	Dissolved organic geochemistry	Paleoecology, paleoclimate	b	Liquid	5 mL		Stored in glass, filtered	UVM
10	Bulk macrofossils	Bulk organic geochemistry (δ13C, δ15N)	Paleoecology, paleoclimate	a,b	Wet	0.05 g	bulk organic material from 63-125		UW
11	Sediment	Diatom identification	Paleoecology, paleoclimate	a,b,g	Wet	10 g	bulk		UC Boulder
12	Sediment	Foraminifera identification	Paleoecology, paleoclimate	a,b,g	Wet	10 g	bulk		UC Boulder
13	Sediment	Lipid biomarkers	Paleoecology, paleoclimate	a,b	Wet	20 g	<63		Univ at Buffalo
14	Sediment	Macrofossil assemblage	Paleoecology, paleoclimate	a,b,g	Wet	50 g	bulk		UVM, Columbia
15	Sediment	Pollen assemblages	Paleoecology, paleoclimate	a,b,g	Wet	20 g			UVM, Columbia
16	Frozen oriented cube	γ, Natural Remanent Magnetization	Physical properties, magnetostratigraphy	f1	Frozen	6.5 cm <sup>3</sup>		Orientation maintained	Montclair State Univ. Columbia
17	Sediment	Ar/Ar dating of hornblende	Sediment provenance, subglacial geology	a,b	Dried	50 g	125-250		
18	Sediment	Automated quantitative mineralogy (AQM)	Sediment provenance, subglacial geology	a,b	Dried	5 g	47-63, 63-125, 125-250		GEUS
19	Sediment	U-Pb dating of zircon, apatite, rutile	Sediment provenance, subglacial geology	a,b	Dried	20 g	47-63, 63-125, 125-250		GEUS
20	Sediment	U-Th/He dating of apatite	Sediment provenance, subglacial geology	a,b	Dried	100 g	125-250		Uconn
21	Sediment	Grain size distribution	Sedimentology, stratigraphy, physical characterization	a,b	Dried	not size dependent	bulk		UVM
22	Sediment	Particle size and shape	Sedimentology, stratigraphy, physical characterization	a,b	Dried	5 g	bulk		UVM, CRPG
23	Frozen intact sample	µCT scanning	Sedimentology, stratigraphy, physical characterization	a,b	Frozen	n/a	n/a	Orientation maintained	CRREL
24	Frozen intact sample	3D photogrammetric models	Sedimentology, stratigraphy, physical characterization	Uncut segment	Frozen	n/a	n/a	Orientation maintained	UVM
25	Frozen intact sample	Ice content	Sedimentology, stratigraphy, physical characterization	a,b	Frozen and dried	not size dependent	n/a		UVM
26	Frozen intact sample	Clay mineralogy SEM, XRD	Surface weathering and erosion	c	Frozen	5 g	<63		UVM
27	Pore ice meltwater	Water cation geochemistry	Surface weathering and erosion	a	Liquid	2 mL		Stored in plastic, filtered	Univ. Arizona, UVM
28	Pore ice meltwater	Water anion chemistry	Surface weathering and erosion	a	Liquid	2 mL		Stored in plastic, filtered	Williams College

USU Utah State University  
 UVM University of Vermont  
 CRPG Centre de RecherchesPétrographiqueset Géochimiques  
 UW University of Washington  
 NBI Neils Bohr Institute  
 UC Boulder University of Colorado, Boulder  
 CRREL Cold Regions Research and Engineering Laboratory



310

Figure 6: Sample cutting, sub-sampling, and processing flow diagram. White boxes show sample materials, orange boxes indicate major processing steps, and green hexagons show analyses.



315 **Figure 7: Each tube of the sub-glacial section of the core. Comparison of the photographs taken between 1966-1972 CE (black and white, Fountain, 1981), the 3D photogrammetry model, and the segments cut for this study in 2021 CE for this study.**

Using the bandsaw, we cut each core segment in half along the z-axis to produce an archival half and a working half (Figure 5A). From the working half, we cut the outer quadrants (sub-samples [a] and [b]) to isolate an interior prism of the core segment. We recorded the dimensions (width, length, height [h]) of sub-samples (a), (b), and the interior rectangular prism. Then, to isolate pristine surfaces, we cut the top and bottom ~1 cm of the interior prism using the diamond-wire saw and saved the pieces as sub-samples (c) and (d). Then, we cut the interior prism using the diamond-wire saw to remove a sub-sample with dimensions of 2 x 2 x (h-2) cm (e) for luminescence measurements. The h-2 cm length is the result of cutting off the top 1 cm and bottom 1 cm from the interior rectangular prism. We wrapped sub-sample (e) in two layers of aluminum foil to secure the sediment from light exposure and placed in a separate plastic sample bag with the sample orientation kept intact. Then, in normal light conditions using the diamond-wire saw, we cut a ~1.8 x 2 x (h-2) cm slice and from that slice cut a 2 x 2 x 1.8 cm cube of sediment (sub-sample [f1]) for paleomagnetic measurements. We placed sub-sample (f1) into a labeled plastic paleomagnetic cube with the core orientations maintained. The remaining outer piece, sub-sample (g), was kept for macro- and microfossil identification and stored in a separate bag. While most samples were ~10 cm-tall cylinders, we applied slightly different procedures for segments that were previously cut or thawed for sampling, had odd or small geometries, or were too delicate to cut according to the cutting design (Table 4). For all samples, we saved any loose material from the storage container as a separate 'residual' sub-sample for method testing and replicate measurements. We recorded the masses of each sub-sample (a-g), residual material, and archive half (Table S1). The cut interior face of the archived half and sub-samples (a) and (b) were photographed and then described including ice content, sedimentological properties, and sedimentary features.

We placed sub-samples (a-d) in the same plastic bag with sample orientations maintained (e.g., up and down direction). All sub-samples (a-g) were kept in the storage freezer at NBI (-30 °C) prior to transfer from NBI to the University of Vermont. The archive halves of all samples remain in the NBI ice core facility in plastic bags inside of glass jars in closed foam boxes shielded from light. We packed sub-samples into insulated shipping containers lined with brine freeze packs that constituted 50% of the interior container volume and allowed to equilibrate at -30 °C for two days before being sealed for transport. We transported all sub-samples from NBI via commercial air travel on a direct flight from Copenhagen, Denmark to Newark, New Jersey to Burlington, Vermont, where samples were promptly transferred into a -20 °C chest freezer with back. Within 1 hour of arrival, we transferred samples into a -20 °C chest freezer at Montclair State University and allowed them to cool for two days. We repacked the samples into insulated shipping containers with brine freeze packs and transported them by car from Montclair, New Jersey to Burlington, Vermont, where samples were promptly transferred into a -20 °C chest freezer with back-up power and high limit alarms for storage. At all times, samples remained frozen.

We distributed sub-samples (e) and (d) for the luminescence and bulk chemistry measurements. We placed sub-sample (e) into labeled, opaque resealable bags and then shipped them frozen to Utah State University for luminescence measurements. We thawed each sub-sample (d) overnight in a 65 °C oven, massed each again. Then, the dried sediment was shipped to Utah State University for bulk geochemistry and background radiation analyses that will be used for luminescence dose-rate calculations, and interpretation of cosmogenic <sup>36</sup>Cl concentrations.



Table 4: Samples with modified cutting or sub-sampling design (2021)		
Sample	Alteration	Reason
1059-7	Thickness of sub-samples (c) and (d) reduced to 0.5 cm	Thin segment; optimized amount of frozen material for (e,f, and g)
1060-A2	Swapped the positions of sub-samples (e) and (f) in the inner prism	Previous cutting exposed inner portion of core to light. Re-arranged to avoid light contamination for (e)
1060-B	All sub-samples collected as grab samples	Sample was thawed and preserved as bulk sediment
1061-C	Sub-samples (e) and (f2) contained two fragments	Sample had fractured during storage
1061-D2	Swapped the positions of sub-samples (e) and (f) in the inner prism	Previous cutting exposed inner portion of core to light. Re-arranged to avoid light contamination for (e)
1061-D4	Not sub-sampled	Sample was missing from archive
1061-D5	Lower ice/rock fragment kept as sub-sample (x)	Sample had odd geometry
1062-4	Sub-samples (e) and (f2) contained two fragments	Sample fractured during wire cutting
1063-1	Sub-samples (a), (b), and (g) taken from fragments on outer part of core	Odd geometry, samples fragmented along ice lens planes
1063-3	Not sub-sampled	Sample was missing from archive
1063-4	Sub-sample (f1) collected from lower slice of inner prism b not kept	Large cobble in sample
1063-5	Swapped the positions of sub-samples (e) and (f) in the inner prism	Previous cutting exposed inner portion of core to light
1063-8	Sub-sample (b) not kept, sub-samples (f1) and (f2) cut from outside of inner prism	small sample

*Note: this table is relevant to deviations from the cutting and sub-sampling protocols during the 2021 sampling event. Details about previous cutting, sampling activity or alterations to the sediment archive prior to 2021 is described in the text, Tables 5 and S9, and Figure 8.*

### 3.3 Analysis of frozen sediment

Using frozen sub-samples, we measured magnetic properties (f1) and collected micro-computed tomography ( $\mu$ CT) scans (a, b). We measured the magnetic susceptibility of the frozen (f1) sub-samples from each core segment at Montclair State University. Susceptibility measurements were collected in an applied field of 200 A/m and at two frequencies (976 Hz and 3904 Hz) using an AGICO MFK2A Kappabridge (Table S4). Here, we report the measurements made at 976 Hz ( $\chi_{976\text{Hz}}$ ) as the low-field susceptibility.

Samples were then measured for paleomagnetic inclination, declination, and intensity of remanence at alternating field demagnetization (AFD) levels of 0 mT (natural remnant magnetization [NRM]), 10 mT, and 20

mT using an AGICO JR-6 Spinner Magnetometer (Table S5) and D-tech alternating field demagnetization unit. Samples were kept frozen between AFD treatments. However, by the 20 mT step many of the frozen samples had partially deformed inside their plastic cubes, and we therefore discontinued AFD treatments. We corrected the orientations of samples that were noted to have been stored upside down at NBI by rotating the NRM vector 180° around the x-axis in the sample coordinate system (Fig. 8; Tables 5, S9).

We collected  $\mu$ CT scans from the (a) and (b) sub-samples to create a digital, three-dimensional record and to non-destructively observe and measure sedimentological properties (e.g., porosity, grain size and grain shape) and stratigraphic features at sub-mm-scale resolution. We scanned sub-samples (a) and (b) from each segment using a Bruker SkyScan1173  $\mu$ CT scanner fit for use in a -10 °C cold room at CRREL in Hanover, New Hampshire. Full-sized, ~10 cm-tall samples were scanned at 71  $\mu$ m/pixel resolution in two overlapping 7.9 cm-tall scans for the bottom and the top of each sub-sample to ensure that the entire sample was scanned.

### 3.4 Processing and analysis of thawed sediment

Following full documentation and  $\mu$ CT scanning of sub-samples (a) and (b), we thawed these sub-samples from each segment to extract pore ice meltwater, save aliquots of bulk material, and isolate specific grain-size fractions used for varied analyses. For each core segment, we placed sub-samples (a) and (b) in separate sealed plastic and glass containers, respectively, to avoid contamination for pore-ice measurements designated for each sub-sample. Plastic was used for sub-sample (a) because pore ice in this sub-sample was analyzed for stable water isotopes, bulk geochemistry, and major ion chemistry. Glass was used for sub-sample (b) because pore ice in this sub-sample was analyzed for dissolved organic geochemistry.

We measured the frozen mass of each sub-sample, thawed the sub-samples overnight in a 4°C refrigerator, and then re-measured the thawed mass to ensure that no water was lost due to evaporation. After thawing, we transferred the sediment into two 250 mL HPDE bottles that we centrifuged for 20 minutes to isolate porewater from the solids. The liquid supernatant from sub-sample (a) was transferred into a 50 mL plastic vial, and the supernatant from sub-sample (b) was transferred into a glass test tube. We recorded the mass of the melted pore ice for both sub-samples. Melted pore ice from sub-sample (a) was then stored frozen at -15 °C immediately after it was recovered. We measured pH and conductivity in melted pore ice from sub-sample (b) using calibrated hand-held Myron L conductivity and pH meters (Table S6). Melted pore ice from sub-sample (b) was transferred to a glass jar and stored frozen at -15 °C.

We recombined, gently mixed, and homogenized the wet thawed sediment from sub-samples (a) and (b) and recorded the recombined mass. We saved two bulk aliquots of the recombined sediment. First, ~10% by mass of the wet sediment was stored in a plastic bag at 4°C as wet bulk archive for other biological analyses (e.g., pollen, microfossils including diatoms, macrofossil characterization). Second, 20% by mass of the wet sediment was oven-dried at 65°C and saved as a dried bulk archive for geological and sedimentological analyses (grain shape, particle size, meteoric <sup>10</sup>Be, rock magnetism, Fe-oxide analysis); the mass was recorded before and after drying to calculate the water lost (Table S7). We calculated the % ice content by mass for each sample by subtracting the total mass of water removed for pore ice meltwater samples and the estimated loss of water from drying from the total thawed sediment mass (using the % water lost from the bulk geological aliquot) (Table S7).

Then we wet sieved (with deionized, 17 Mohm [DI] water) the remaining thawed sediment and stored in plastic bins the >2000, 850-2000, 500-850, 250-500, 125-250, and 63-125  $\mu\text{m}$  grain size fractions. The <63  $\mu\text{m}$  fraction was stored in an acid-washed 5-gallon bucket and allowed to settle for at least 72 hours before decanting water via suction. We transferred the settled wet <63  $\mu\text{m}$  fraction into plastic bags and stored it frozen at -15  $^{\circ}\text{C}$  and then shipped the frozen samples to the University at Buffalo for lipid biomarker analyses. For grain sizes >63  $\mu\text{m}$ , we visually inspected each grain-size fraction for macrofossils and transferred macrofossil specimens floating in water using disposable pipettes into petri dishes for observation and photographic documentation under a dissecting microscope. We transferred macrofossils from each grain size fraction to glass vials filled with DI water for storage at 4 $^{\circ}\text{C}$ . After removing macrofossils, we dried each grain-size fraction in plastic bins at 60 $^{\circ}\text{C}$  overnight, and then measured its dry mass. The dried sediment was stored in plastic bags at room temperature. We calculated the grain-size distribution of each sample using the estimated total dry mass of each sample, the dried masses of each grain size fraction >63  $\mu\text{m}$ , and the estimated the <63  $\mu\text{m}$  mass (Table S7).

For samples composed primarily of ice rather than sediment, we adapted a different processing flow. Using a handsaw, we cut sub-samples from 1060-C4 to separate frozen sediment (1060-C4sediment) from underlying sediment-laden ice (1060-C4ice). For the ice-rich samples (1060-C4ice through 1061-D5) only one quadrant (a or b), whichever was smaller by mass, was processed according to the procedure above. The remaining quadrant remains frozen and in storage for future high-resolution stable water isotope analysis and other analyses.

## 4. Results

This paper reports fundamental observations that we made during core processing and develops a core stratigraphy to provide context for research performed on sub-samples allocated from the core. We report data for physical properties directly measured on the core as it was processed. Other data from geochemical, isotopic, and geochronologic analyses will be reported elsewhere by scientists making those measurements. Below we detail our documentation efforts and results from processing the different sub-samples for an array of analyses (Table 3).

### 4.1 Sample preservation

Based on our measurements of cut core segments and our reconstruction based on a photograph taken before the core was cut, the Camp Century sub-glacial core is 3.44 m long, which is slightly less than what was originally reported (3.55 m). This shortening could be the result of sublimation-induced drying and shrinkage. Comparison of photographs taken before 1972 CE and those taken in 2021 CE shows that there have been changes in the shape of individual core segments including increased surface relief and deeper grooves on the surface. However, our work reveals that most of the core (28 of 32 segments) is almost intact; below we detail modifications or sub-sampling of the archived segments prior to this study (Tables 5, S9). Two segments (1061-D4, 1063-3) are missing without documentation of their current location; no information exists showing that they were used for earlier analysis. Two segments were previously thawed and have been refrozen: 1060-B is loose, dry sediment and 1061-B is loosely compacted ice particles. Table S9 outlines differences in core segment diameter between the top and bottom of the segments and evidence for an icy-flattened side/bottom of the core segments, both indicating melting, ice flow/deformation, and possible sublimation during storage.

**Table 5: Evidence of sub-sampling or modification of archive prior to 2021**

Segment	Modification	Evidence
1059-4	Previously cut and sub-sampled	Sub-sample cut and analyzed as part of the pilot study (Christ et al., 2021).
1060-A2	Cut and sub-sampled for diatoms	A quadrant of the segment was cut; the interior of that segment was sampled; diatom and other microfossil counts from this sample are presented in Harwood (1986).
1060-B	Thawed and re-frozen	No documented explanation.
1060-C3	1 missing exterior pebble; stored upside-down	Cavity observed on outside of segment in recent photographs and 3D photogrammetric model where a pebble is present in the historic photograph of the core tubes (Fountain et al., 1981); however there is no documented analysis of this pebble. Comparison of recent photographs and 3D models with historic photograph indicated this sample was stored upside-down.
1061-B	Thawed and re-frozen	No documented explanation.
1061-D2	Cut and sub-sampled for diatoms (inferred)	A quadrant of the segment was cut; the interior of that segment was sampled in a similar fashion to segment 1060-A2; however there are no published diatom counts from this segment.
1061-D3	Stored upside-down	Comparison of recent photographs and 3D models with historic photograph, as well as color banding observed on the cut faces of this segment indicated it was stored upside-down.
1061-D4	Missing from archive	No documented explanation.
1062-1	2 missing exterior pebbles	2 cavities observed on outside of segment in recent photographs and 3D photogrammetric model where pebbles are present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981.
1062-2	Stored upside-down	Comparison of recent photographs and 3D models with historic photograph indicated this sample was stored upside-down.
1062-3	Stored upside-down	Comparison of recent photographs and 3D models with historic photograph indicated this sample was stored upside-down.
1063-1	1 missing exterior pebble	Cavity observed on outside of segment in recent photographs where a pebble is present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981.
1063-2	1 missing exterior pebble	Cavity observed on outside of segment in recent photographs and 3D photogrammetric model where a pebble is present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981.
1063-3	Missing from archive	No documented explanation.
1063-4	3 missing exterior pebbles; stored upside-down	2 cavities observed on outside of segment in recent photographs and 3D photogrammetric model where pebbles are present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981. Comparison of recent photographs and 3D models with historic photograph indicated this sample was stored upside-down.
1063-5	1 missing exterior pebble; cut and sub-sampled for diatoms (inferred)	A quadrant of the segment was cut; the interior of that segment was sampled in a similar fashion to segment 1060-A2; however there are no published diatom counts from this segment in Harwood (1986). Cavity observed on outside of segment in recent photographs and 3D photogrammetric model where a pebble is present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981.
1063-6	1 missing exterior pebble	Cavity observed on outside of segment where a pebble is present in the historic photograph of the core tubes and analyzed in Fountain et al., 1981.
1063-7	Previously cut and sub-sampled	Sub-sample cut and analyzed as part of the pilot study (Christ et al., 2021).

We confirmed past sampling and investigation of some core segments (Table 5, Fig. 8). A quadrant from segments 1060-A2, 1061-D2, and 1063-5 was previously cut and ~1 cm<sup>3</sup> samples were collected from the interior of the cut quadrant of the core (Figure 3D). These samples were presumably collected for diatom analysis as  
445 Harwood (1986) reported abundant freshwater diatoms in segment 1060-A2; however, there is no documented record of diatom analysis in the other two segments. The outer ~3 cm of two 10-cm long segments (1059-4, 1063-7) were cut, thawed, and analyzed separately in 2019 CE for a pilot study (Christ et al., 2021). We identified cavities on the exterior of several core segments, some of which correspond to pebbles that were analyzed in Fountain et al., (1981): 1062-1 (2 pebbles), 1063-1 (1 pebble), 1063-2 (1 pebble), 1063-4 (3 pebbles), 1063-5 (1  
450 pebble), 1063-6 (1 pebble).

One pebble from the exterior of segment 1060-C3 is missing but there is no documentation of earlier sampling activity for this segment. It is possible that this is the clast Herb Ueda mentions in his oral history as having sent to a colleague in Japan (Shoemaker, 2002). Based on the comparison of the original core photograph


(Fountain et al., 1981) with recent photographs, 3D photogrammetric models, and visual inspection of the core interior, we identified six core segments that were stratigraphically inverted during storage: 1060-C3, 1061-D3, 1062-2, 1062-3, 1063-4, and 1063-7. Additionally, one side of several segments had a flat icy surface, which may have formed due to partial thawing and refreezing of these segments when they were removed for sampling or photography in the past (e.g., Fig. 3C; Table S9). All other segments were cylindrical and therefore had not deformed or melted.

## 460 4.2 Sedimentology and stratigraphy

Based on the grain-size distribution, the pore-ice content, and sedimentary structures seen on the core cut faces, we identified five stratigraphic units in the Camp Century sub-glacial material core (Figure 8). At the base of the core, Unit 1 (215 to 344 cm depth below the ice-sediment interface; segments 1063-8 through 1061-D5) is an olive gray (5Y 4/2) homogeneous, poorly-sorted, clay-rich diamict (all colors are Munsell designations). Pebbles and cobbles are sub-angular to sub-rounded. Pebbles are composed of a variety of lithologies (including granite, gabbro, tonalite, and sandstone); several cobbles in this unit were previously sampled and petrologically analyzed (Fountain et al., 1981). Unit 1 is  $11 \pm 4\%$  ice by mass and has sub-horizontal, mm-scale ice lenses with thicknesses ranging from 1 to 2 mm and spacing ranging from 1 to 5 cm. Ice lenses generally do not intersect pebbles and some ice lenses intersect each other. The contact between Unit 1 and 2 is not readily observable due to the fractured and non-cylindrical geometry of core segment 1061-D5 but is suggested by a shift from clay-rich diamict to sediment-laden ice at 215 cm depth.

Unit 2 (112 to 215 cm depth; segments 1061-D5 through 1060-C4) is composed of olive (5Y 4/4) sediment-laden ice, having sub-vertical inclusions of ice that are longer than they are wide and without sediment. The sediment-laden ice of Unit 2 is clearly distinct from the rest of the sub-glacial material (Herron and Langway, 1979). Ice lenses commonly crosscut each other and show tight folded structures based on visual observations. The lower half of Unit 2 (158 to 215 cm depth) displays horizontal color banding between light olive brown (2.5Y 5/6) ice and brown (10 YR 3/2) ice that likely corresponds to greater sediment content that is mostly silt and clay (72 to 78%) with minor amounts of fine to medium sand and rare pebbles. Segment 1061-B (140 to 158 cm depth) was previously thawed and refrozen so there is no observable stratigraphy or structure in that segment. The upper part of Unit 2 (112 to 140 cm depth) is moderate orange (10Y 3/2) silty ice with brown (7.5YR 2/4) sub-vertical ice-filled lenses. Sediment content is low (4 to 10% by mass) and is dominated by silt and clay (84%) with minor amounts of fine sand and rare pebbles in an ice matrix. Sediment content and grain size decreases in the upper 70 cm of Unit 2. The contact between Unit 2 and Unit 3 (112 cm depth, segment 1060-C4) has pebbly gravel resting over a very irregular ice surface.

Unit 3 (85 to 112 cm depth; segments 1060-C4 sediment through 1060-C1) is dark olivish gray sediment (2.5 GY 4/2) that fines upward from a pebble gravel to sand and mud in a dark amber (7.5Y 3/4) ice-supported matrix (22 to 49% ice by mass); sediment is concentrated in deformed bands separated by ice. This unit has abundant macrofossils. Unit 4 (55 to 85 cm depth; segments 1060-C1 through 1060-B) is dark greenish gray (10Y 4/2) very fine and fine sand with slightly inclined bedding. Due to prior melting of segment 1060-B we cannot evaluate the ice content within much of this unit. There is a gradational contact between Unit 4 and Unit 5 at ~55 cm. Unit 5 (0 to 55 cm depth; segments 1060-A2 through 1059-4) is olive gray (5Y 4/2) bedded medium and coarse sand that is 15% ice by mass.



495 **Figure 8: Results compiled from sample documentation and processing (from left to right). A) core tube**  
**numbers and sample names created when the core was cut in 1972 CE; B) evidence of altered, previously**  
**investigated, or missing samples based on literature, photographs, and recent documentation; C)**  
**stratigraphic units and core log; color corresponds to the average soil color of each sample segment; vf –**  
**very fine sand, f – fine sand, m – medium sand, c – coarse sand, vc – very coarse sand; D) grain-size**  
**distribution (by mass) measured from wet sieving; dashed pattern in 63-125  $\mu\text{m}$  for segments 1059-4 and**  
**1063-7 denote <125  $\mu\text{m}$  fraction; E) percent sediment and ice content (by mass); F) frozen bulk density; G)**  
**magnetic susceptibility ( $\chi$ ) measured at 976Hz; and H) pore ice pH and conductivity. Gaps in datasets due**  
**to thawed or missing segments, insufficient pore water, non-cylindrical geometries, or analyses not made**  
**on pilot samples. Dates in column B from Christ et al., 2021, 2023.**

### 4.3 Physical characteristics

505 Bulk density and  $\chi_{976\text{Hz}}$  measurements (Figure 8F, G) both correspond to the stratigraphy and sediment  
content with generally higher values in the sediment-rich units and lower values in ice-rich units. Bulk density is  
 $2.2 \pm 0.2 \text{ g/cm}^3$  (mean  $\pm 1\sigma$ ) in Unit 1, decreases to  $1.1 \pm 0.2 \text{ g/cm}^3$  in Unit 2, increases to  $1.7 \pm 0.3 \text{ g/cm}^3$  in Unit  
3, and then increases to  $2.3 \pm 0.1 \text{ g/cm}^3$  in Units 4 and 5. At some levels, there is an apparent discrepancy between

510 density and ice content, which is mainly due to our sub-sampling approach used to measure the ice proportion, while the bulk density was done on longer full ice cylinders at lower resolution. Moreover, the contact between Unit 2 and 3 is not horizontal but tilted; several cobbles are also embedded in the top of unit 2.  $\chi_{976\text{Hz}}$  values are greatest in Unit 1 ( $1.93 \pm 0.22 \times 10^{-6} \text{ m}^3/\text{kg}$ ), lowest in Unit 2 ( $0.16 \pm 0.14 \times 10^{-6} \text{ m}^3/\text{kg}$ ) where sediment is substantially diluted by ice, variable in Unit 3 ( $1.27 \pm 0.59 \times 10^{-6} \text{ m}^3/\text{kg}$ ), and similar through Units 4 and 5 ( $1.03 \pm 0.31 \times 10^{-6} \text{ m}^3/\text{kg}$ ).

#### 515 4.4 Paleomagnetic measurements

Paleomagnetic inclinations in Units 1, 3, 4 and 5 are highly variable (Table S5). We do not consider the silty ice of Unit 2, > 80% ice, to be a reliable geomagnetic field recorder. The steepest inclinations were observed at the 0 mT (NRM) AFD level, which may reflect the acquisition of a viscous remanence in the downward direction (+Z in the sample coordinate system) while in storage at NBI (Figure 8G). Inclinations at the 10 mT and 520 20 mT AFD levels are shallower than the expected geocentric axial dipole value of  $84^\circ$  at the site latitude, which could result from compaction of sub-glacial diamict, post-depositional slumping, disturbance from permafrost, or partial thawing of the sample during the AFD treatments and measurements.


Identification of normal or reversed polarity is possible from sub-glacial and grounding line sediment (Wilson et al., 2012; Hodson et al., 2016), though challenging in the Camp Century core due to our limited 525 number of AFD levels. Most of our samples display positive inclinations, consistent with normal polarity, with the possibility of 3 reversed polarity or excursion intervals in Units 1 and 2 recorded in samples 1061-D1, 1062-3, and 1061-D3. Although the AF demagnetization data are limited, these samples display inclinations that become progressively more negative at higher AF levels, consistent with removal of vertically downward overprint, or inclinations that remain moderately negative at all AF demagnetization levels.

#### 530 4.5 Pore ice pH and conductivity

In general, melted pore ice pH and conductivity are inversely related throughout the core (Figure 8H). Conductivity is the lowest in Unit 2, with the highest percentage of ice by mass, and higher in the other units, with maximum values occurring where the percentage of fine grains is the highest. In contrast, pH is the inverse of the bulk density profile, with lower values in sediment-rich units (i.e., Units 1-3, 5) and higher values in ice-rich units (i.e., Unit 2). Unit 1 generally has the lowest pH (mean  $\pm 1\sigma$ :  $6.96 \pm 0.07$ ) and highest conductivity 535 values ( $653 \pm 250 \mu\text{S}$ , maximum:  $1066 \mu\text{S}$ ), while Unit 2 has the highest and most variable pH ( $7.37 \pm 0.35$ ) and lowest conductivity values ( $129 \pm 85 \mu\text{S}$ ; minimum:  $52.1 \mu\text{S}$ ). In Unit 3, conductivity increases up-core from 168 to  $502 \mu\text{S}$  but displays stable pH values ( $7.11 \pm 0.10$ ). Units 4 and 5 have similar pH ( $7.14 \pm 0.11$ ) and low conductivity ( $174 \pm 53 \mu\text{S}$ ).

540 The controls on pore ice pH and conductivity likely differ between the units due to varying sediment and ice content as well as depositional and pore-water history. Regression analysis shows no significant correlation between any combination of conductivity, pH, and grain size, but there is distinct clustering; for example, Figure 9 shows that water isolated from Unit 1 has little variance in pH but large variance in conductivity. The former is likely the result of sediment buffering capacity and the latter is possibly the result of sediment/water interactions 545 during weathering and the competing processes of dilution by meteoric waters and the chemical weathering of the sediment. In contrast, the sediment-laden ice of Unit 2 has consistently low conductivity but a wide variance in

pH, likely the result of low buffering capacity and the paucity of sediment grains that in the presence of liquid water provide ions that would increase conductivity.



550 **Figure 9. Bivariate plot of pH versus conductivity grouped by stratigraphic units.**


#### 4.6 Macrofossils

Macrofossil remains are present in all sub-samples we examined (Figure 10). Initial observation and photographic documentation of isolated macrofossils in sub-samples (a) and (b) indicates that plant macrofossil remains are present in all units and most abundant in Units 3 and 4, and that insect remains are present in Units 1, 3, and 5. Macrofossils are least abundant in Unit 2. The quality of macrofossil preservation is consistently better in the largest size fractions, 850-2000 µm and >2000 µm. In Unit 1, samples hold primarily woody debris, and bryophyte leaves or stem fragments. Near the interface between Unit 1 and 2, sample 1062-1 has insect remains, including chironomid larval head capsules. Unit 2 contains sparse plant remains, mostly fragments of bryophyte leaves. Within Unit 3, a partial chironomid pupa was recovered from sample 1060-C1. Macrofossils in Unit 4, sample 1060-B, are abundant and distinct from those of samples below, including fungal sclerotia. Within Unit 5, samples 1060-A1 and 1059-5 have chironomid larval head capsules.

555

560





565 **Figure 10: Examples of macrofossils recovered during wet sieving: A) *Dryas octopetala* leaf (1060-A2); B) *Cenococcum geophilum* sclerotium [red rectangle] (1060-C2); C) *Cyperaceae* seed (1060-C1); D) *Characeae* oospore [red rectangle] (1060-C1); E) partial chironomid pupa (1060-C1); F) chironomid larval head capsule (1062-1).**

## 5. Implications

### 570 5.1 Initial interpretation of sub-glacial core stratigraphy and paleo-environments

The Camp Century sub-glacial material preserves evidence for different surface and depositional processes in both glaciated and ice-free environments. The absence of stratification, variable grain size, shape, and poor sorting suggest that the diamicton at the base of the core (Unit 1) was most likely sub-glacially

575 deposited as a basal till. The fragmented macrofossil remains in Unit 1 are derived from a non-glaciated  
landscape and were incorporated into the till most likely either by overriding ice or surface mass movement. The  
presence of mm-scale ice lenses throughout Unit 1 are consistent with permafrost activity.

580 The contact between Units 1 and 2 marks a shift from predominantly sediment to ice-dominated sub-  
glacial materials. Unit 2, sediment-laden ice, has near vertical ice-filled fractures. The origin of the ice in Unit 2  
is not clear based on physical observations alone. It could be remnant basal glacial ice, residual firn, or represent  
a thermokarst pond or other small sequentially frozen water body or could be as a lens of permafrost ground ice  
or segregation ice. Further analysis, including chronological and stable water isotope data, will better show how  
585 this unit formed. The unconformable contact between Units 2 and 3, along with upward fining in grain size and  
abundant plant macrofossils, suggests the sediment in Unit 3 might have been deposited by slumping (or by  
another process that had sufficient power to erode Unit 2). The very fine to fine-grained bedded sand with  
abundant macrofossils in Unit 4 suggest subsequent subaerial sediment deposition by low-energy flowing water.  
590 The coarsening in grain size up-core from Unit 4 to 5 suggests sediment deposition by higher-energy water. The  
abundance of well preserved and delicate macrofossil fragments demonstrates the presence of a subaerial  
ecosystem during the deposition of Units 3, 4, and 5. Luminescence dating of the uppermost sample in Unit 5  
indicates it was last exposed to light  $416 \pm 38$  ka, coincident with the Marine Isotope Stage 11 interglacial (Christ  
et al., 2023).

The preservation of delicate frozen macrofossils and bedding structures suggests that the internal  
structures of the Camp Century sub-glacial sediment in Units 3, 4 and 5 have not been internally deformed –  
future fabric and/or micro-morphology analysis could affirm the validity of this observation. Basal borehole  
temperatures at Camp Century ( $-13.0$  °C at time of drilling,  $-11.8$  °C corrected for pressure-melting using local  
595 ice thickness) are far below the pressure melting point (MacGregor et al., 2016), indicating that the northwestern  
Greenland Ice Sheet is presently cold-based and therefore non-erosive. The basal thermal state, and thus erosional  
character, of this sector of the ice sheet has likely varied in the time since the sediment was deposited and over-  
ridden by ice; however, we have no way to determine if or how much overlying sediment was initially present  
and then eroded after deposition.

600

## **5.2 Lessons learned about the acquisition and sampling of basal ice core materials**

The Camp Century sub-glacial core is a unique archive. It was the first sub-glacial archive recovered  
from beneath any ice sheet and remains well preserved. Apart from two missing samples and two previously  
thawed samples, the rest of the archive was still frozen and intact in 2019 CE when we began our work.

605 While the historic documentation of earlier cutting, sampling, and core handling activities is incomplete,  
the excellent preservation of this archive allowed us to reconstruct the original orientation of the sub-glacial  
material and then design and execute a detailed and comprehensive sampling and analysis plan. We archived,  
handled, and processed this sediment to ensure that future studies are not compromised either due to  
contamination or inadvertent loss of grain-size fractions useful for varied analytical techniques that have yet to be  
610 developed.

The framework that we devised in this study draws upon the sub-sampling approaches commonly used for ice cores, lake, and marine sediment cores, which save archival material and allocate samples for a variety of analyses (e.g., Souney et al., 2014; McKay et al., 2019; Priscu and SALSA Science Team, 2021; Hodson et al., 2016). These large community efforts today devise sampling plans that consider analytical techniques prior to  
615 core collection. In contrast, we devised our plan 55 years after core recovery. We share this historical documentation, intentional core cutting and sub-sampling design, and results as a guide for the many on-going and future analyses of the Camp Century sub-glacial archive that will shed light on the paleoclimate conditions, ice-free ecosystems, surface and sub-glacial processes, and chronology of sediment deposition.

This analytical framework could also be useful as a baseline for the study of other sub-glacial materials, since our preliminary results (Christ et al., 2021) demonstrate that small amounts of rare, archival sub-ice  
620 material can be used to answer pressing questions in paleoclimate, such as the long-term history of ice sheets. The approach we outline here is currently being used not only for the US National Science Foundation funded project (Lead investigator, Bierman, University of Vermont) to understand the sub-ice material from Camp Century but also by a European Research Council Project (Lead investigator, Dahl-Jensen, University of  
625 Copenhagen) focused on basal ice and sub-glacial material from several ice cores including historical ones from Camp Century and GRIP.

Our work with the Camp Century core suggests ways in which future sub-glacial materials could be acquired and treated to minimize contamination and maximize scientific impact. Based on our experiences, we make five specific suggestions

- 630 1. Collecting and preserving sediment and the top of bedrock without light exposure allows them to be dated by luminescence and prevents degradation of biochemical records. While not originally collected or stored in red light and light-proof conditions, we were able to sub-sample the Camp Century core segments in light-safe manner. More recent coring campaigns are implementing luminescence-safe sample collection procedures (e.g., EastGRIP, NEEM, GreenDrill).
- 635 2. Extensive documentation of core recovery, including accurate assessment of percent recovery during drilling is also important for understanding the continuity of core sections. Obtaining and preserving information about core orientation is essential for paleomagnetic and structural studies.
- 640 3. Maintaining detailed records of transport and storage conditions as well as sample allocation, treatment, and usage from time of core collection onward is important for future analyses. Considering ways to minimize contamination in sample collection, storage, and transport is key for biologic studies. For example, a modern carpet beetle made its way into one of the Camp Century samples sometime over the last 50+ years.
- 645 4. To facilitate analysis of biomarkers, including lipids and DNA, drilling and sampling in a way that avoids inadvertent contamination is important. Advances could include the use of hydrocarbon-free drilling fluid and the maintenance of sterile conditions for collection of DNA and RNA samples – realizing of course limitations of the polar environment although such work has been pioneered in sub-ice Antarctic lakes (Hodson et al., 2016; Priscu and SALSA Science Team, 2021). Retention of representative samples of drilling fluids is key for determining blank levels for later chemical analyses.
- 650 5. Performing bulk density measurements on cores immediately after they are extracted and again when they are cut will provide information about changes over time as cores are stored in freezers.

Working with the Camp Century sub-glacial archive has taught us the importance documenting all sampling activities and the value of sustained funding and support for ice-core storage facilities that make it possible for future researchers to come back to a core many decades later.

### 5.3 Outstanding Research Questions

655 Although recent (Christ et al., 2021, 2023) and prior research (Whalley and Langway, 1980; Fountain et  
al., 1981; Harwood, 1986) involving the Camp Century sub-glacial core has already revealed much about the  
history of northwestern Greenland, there is much more to do and learn. Christ et al. (2021) and (2023) provide a  
glimpse into the richness and complexity of interpreting sub-glacial sediment archives using modern analytical  
660 techniques and large, international, collaborative teams. We now know that the uppermost sample was exposed at  
the surface during MIS 11 and that the till at the base of the core has been buried no more than 3.2 Myr, but we  
have little knowledge of the age of the units in between. More temporal information will come as cosmogenic  
and luminescence analyses are completed on the rest of the core. Almost the entire core contains microfossils,  
which reveal the composition of interglacial ecosystems. Although many microfossils have been identified using  
reference collections (Figure 10), biochemical analyses in process using biomarkers, including DNA and lipids,  
665 will further characterize biologic molecules preserved in the core materials. Geochemical analysis of both  
sediment and porewater will refine our understanding of water-rock interaction in the past while  
geochronological measurements will enhance our understanding of sediment provenance.

The most important implication of the Camp Century sub-glacial core is not the specific knowledge that  
will be gained about the history of ice, ecosystems, climate, and sediments of northwestern Greenland, but the  
670 realization that sub-glacial archives hold such a rich, albeit complex, record of the past. It is this complexity that  
provides the richness but demands the application of many and varied analytical techniques as we begin to  
demonstrate here. Combined with analysis of the basal and clear ice above as well as the bedrock below, cores  
collected from beneath ice sheets are poised to provide the information needed to understand the behavior of the  
Greenland and Antarctic ice sheets in the past and thus allow better prediction of their future and as a result, ours.  
675 Recovering more cores through the ice and into the sediment and bedrock below should be a high priority of the  
cryosphere science community.

### 5.4 Sample Availability

The archive half of the Camp Century sub-ice and basal ice core remains frozen at the NBI in  
Copenhagen and will be accessible for further analyses upon application to the ice core facility there and a plan  
680 for collaboration with Danish scientists whom now control, via their sample distribution committee, access to the  
core materials. Similarly, material from Camp Century presently in the United States and not used consumed for  
analyses, both at UVM and with investigators at institutions listed in Table 3, will be sent to the National Science  
Foundation Ice Core Facility for preservation and future distribution. It will be available, after current NSF  
support ends, by application and approval of a sample request to the sample allocation committee with the  
685 collaboration and support of American scientists.

### Acknowledgements

Funding for core analysis provided by NSF-OPP-2114629, 2114630, 2114632 to Bierman, Rittenour, and Thomas. Laboratory support at the University of Vermont from NSF-EAR-1735676 and NSF-EAR-1735676 to Bierman and Corbett. Christ's work in part supported by post-doctoral fellowship from University of Vermont  
690 Gund Institute for Environment. François Fripiat and Jean-Louis Tison received funding from the FNRS-FRS for the CDR.J.0080.22 AEROBIC project, and the subventions Jaumotte-Demoulin for the GENIAL project. We thank the staff of the Ice Core Facility in Copenhagen, particularly Dorthe Dahl-Jensen and JP Steffensen, for preserving the sub-ice core since 1994 and for providing support during sampling.

### **Code/Data availability**

695 We provide data in the manuscript and supplemental tables. Photogrammetric models of most core segments are available for viewing and download at this online, public repository:  
[https://www.morphosource.org/concern/cultural\\_heritage\\_objects/000583438](https://www.morphosource.org/concern/cultural_heritage_objects/000583438)

### **Author contributions**

AC and PB designed the sample processing methodology with input from TR and ET. JLT, PHB, FF, AC did core  
700 logging, cut the core and made initial observations and interpretation. AC and SB conducted  $\chi$  and NRM measurements. AC wrote the initial draft of the manuscript and drafted the initial figures with input and data from PB, HM, JS, and CC. All authors read and edited several drafts of the manuscript. PB did several rounds of revision of text and figures with the authors and submitted the manuscript.

### **Competing interests**

705 The contact author declares that at least one of the (co-)authors is a member of the editorial board of The Cryosphere.

### **Disclaimer**

The findings and conclusions in this article are those of the authors and do not necessarily represent the views of their corresponding affiliated organization.

### **References**

- Abele, G.: SR-62 Construction of a Snow Runway at Camp Century for Wheel Landings with Lightweight Aircraft., 1964.
- Anon: 'Ancient Air' held in polar ice cores, Tallahassee Democr., 1959.
- Bader, H.: Special Report 58: Scope, Problems, and Potential Value of deep ice Core Drilling in Ice Sheets.,  
715 Hanover., 1962.
- Berner, W., et al. "Past Atmospheric Composition and Climate, Gas Parameters Measured on Ice Cores." Nature, vol. 276, no. 5683, 1978, pp. 53–55, <https://doi.org/10.1038/276053a0>.
- Blard, P.-H., Protin, M., Tison, J.-L., Fripiat, F., Dahl-Jensen, D., Steffensen, J. P., Mahaney, W. C., Bierman, P. R., Christ, A. J., Corbett, L. B., Debaille, V., Rigaudier, T., Claeys, P. and Team, A.: Basal debris of the NEEM ice  
720 core, Greenland: a window into sub-ice sheet geology, basal ice processes and ice sheet oscillations, J. Glaciol.,

- 69(276), 1011–1029, <https://doi.org/doi: 10.1017/jog.2022.122>, 2023.
- Christ, A. J., Bierman, P. R., Schaefer, J. M., Dahl-Jensen, D., Steffensen, J. P., Corbett, L. B., Peteet, D., Thomas, E. K., Steig, E. J., Rittenour, T. M., Tison, J.-L., Blard, P. H., Perdrial, N., Dethier, D., Lini, A., Hidy, A. J., Caffee, M. W. and Southon, J. R.: A multi-million-year-old record of Greenland vegetation and glacial history preserved in sediment beneath 1.4 km of ice at Camp Century, Proc. Natl. Acad. Sci., 118(13), e2021442118, <https://doi.org/10.1073/pnas.2021442118>, 2021.
- 725 Christ, A. J., Rittenour, T. M., Bierman, P. R., Keisling, B. A., Knutz, P. C., Thomsen, T. B., Keulen, N., Fosdick, J. C., Hemming, S. R., Tison, J., Blard, P., Steffensen, J. P., Caffee, M. W. and Corbett, L. B.: Deglaciation of northwestern Greenland during Marine Isotope Stage 11, Science, 381(6655), 330–335, <https://doi.org/10.1126/science.ade4248>, 2023.
- 730 Clark, E.: Technical Report 174: Camp Century Evolution of Concept and History of Design Construction and performance., 1966.
- Corliss, W. R.: Power Reactors in Small Packages, United States. <https://www.osti.gov/biblio/1158799>, 1968.
- Dansgaard, W.: The Abundance of O18 in Atmospheric Water and Water Vapour, Tellus, 5, 461–469, <https://doi.org/10.1111/j.2153-3490.1953.tb01076.x>, 1953.
- 735 Dansgaard, W.: The O18-abundance in fresh water, Geochimica et Cosmochimica Acta, 6, 241–260, [https://doi.org/10.1016/0016-7037\(54\)90003-4](https://doi.org/10.1016/0016-7037(54)90003-4), 1954.
- Dansgaard, W., Johnsen, S. J., Møller, J., and Langway, C. C.: One Thousand Centuries of Climatic Record from Camp Century on the Greenland Ice Sheet, Science, 166, 377–381, <https://doi.org/10.1126/science.166.3903.377>, 1969.
- 740 Daugherty, C. M.: City Under the Ice, The Macmillan Company, New York., 1963.
- Doel, R. E., Harper, K. C. and Heymann, M.: Exploring Greenland: Cold War Science and Technology on Ice, edited by Ronald E. Doel, K. C. Harper, and M. Heymann, Palgrave Macmillan, New York., 2017.
- Epstein, S. and Sharp, R.: Oxygen Isotope Studies, National Acad. Sci. IGY Bull. #21 (Transactions, Am. Geophys. Union), 40(1), 81–84, 1959.
- 745 Fountain, J., Usselman, T. M., Wooden, J. and Langway, C. C.: Evidence of the bedrock beneath the Greenland ice sheet near Camp Century, Greenland., J. Glaciol., 27(95), 193–197, <https://doi.org/10.1017/S0022143000011370>, 1981.
- Goossens, T., Sapart, C. J., Dahl-Jensen, D., Popp, T., Amri, S. El and Tison, J. L.: A comprehensive interpretation of the NEEM basal ice build-up using a multi-parametric approach, Cryosph., 10(2), 553–567, <https://doi.org/10.5194/tc-10-553-2016>, 2016.
- 750 Gow, A. J. and Meese, D. A.: Nature of basal debris in the GISP2 and Byrd ice cores and its relevance to bed processes, Ann. Glaciol., 22, 134–140, <https://doi.org/10.3189/1996aog22-1-134-140>, 1996.
- Hansen, B. L.: Deep Core Drilling in Ice, Mem. Natl. Inst. Polar Res. Spec. Issue, (49), 5–8, 1994.
- 755 Hansen, B. L. and Langway, C.: Deep core drilling in ice and core analysis at Camp Century, Greenland, 1961–66, Antarct. J. United States, Sept–Oct, 207–208, 1966.
- Harwood, D. M.: Do Diatoms beneath the Greenland Ice Sheet Indicate Interglacials Warmer than Present?,

- Arctic, 39(4), 304–308, 1986.
- 760 Herron, S. and Langway, C. C.: The debris-laden ice at the bottom of the Greenland Ice Sheet, *J. Glaciol.*, 23(89), 193–207, 1979.
- Herron, S. L. and Langway, C. C.: A Comparison of Ice Fabrics and Textures at Camp Century, Greenland and Byrd Station, Antarctica, *Ann. Glaciol.*, 3(Mardia 1972), 118–124, <https://doi.org/10.3189/s0260305500002639>, 1982.
- 765 Hodson, T. O., Powell, R. D., Brachfeld, S. A., Tulaczyk, S. and Scherer, R. P.: Physical processes in Subglacial Lake Whillans, West Antarctica: Inferences from sediment cores, *Earth Planet. Sci. Lett.*, 444, 56–63, <https://doi.org/10.1016/J.EPSL.2016.03.036>, 2016.
- Langway, C.: *The History of Early Polar Ice Cores.*, 2008.
- Langway, C. C.: A 400 Meter Deep Ice Core in Greenland: Preliminary Report, *J. Glaciol.*, 3(23), 217–217, <https://doi.org/10.3189/s0022143000024278>, 1958.
- 770 Langway, C. C. and Hansen, B. L.: Drilling Through the Ice Cap: Probing Climate for a Thousand Centuries, *Bull. At. Sci.*, 26(10), 62–66, <https://doi.org/10.1080/00963402.1970.11457874>, 1970.
- Licht, K. J. and Hemming, S. R.: Analysis of Antarctic glaciogenic sediment provenance through geochemical and petrologic applications, *Quat. Sci. Rev.*, 164, 1–24, <https://doi.org/10.1016/J.QUASCIREV.2017.03.009>, 2017.
- 775 MacGregor, J. ., Fahnestock, M. A., Catania, G., Aschwanden, A., Clow, G. D., Colgan, W. T., Gogineni, S. P., Morlighem, M., Nowicki, S., Paden, J., Price, S. F. and Seroussi, H.: A synthesis of the basal thermal state of the Greenland Ice Sheet, *J. Geophys. Res. Earth Surf.*, 121, 1328–1350, <https://doi.org/10.1002/2013JF002871>. Received, 2016.
- 780 Mckay, R. M., Santis, L. De, Kulhanek, D. K., Ash, J. L., Beny, F., Browne, I. M., Cortese, G., Sousa, I. M. C. de, Dodd, J. P., Esper, O. M., Gales, J. A., Harwood, D. M., Ishino, S., Keisling, B. A., Kim, S., Kim, S., Laberg, J. S., Leckie, R. M., Müller, J., Patterson, M. O., Romans, B. W., Romero, O. E., Sangiorgi, F., Seki, O., Shevenell, A. E., Singh, S. M., Sugisaki, S. T., Flierdt, T. van de, Peer, T. E. van, Xiao, W. and Xiong, Z.: Expedition 374 methods, *Proc. Int. Ocean Discov. Progr.*, 374, 1–54, <https://doi.org/https://doi.org/10.14379/iodp.proc.374.102.2019>, 2019.
- 785 Priscu, J. C. and SALSA Science Team: in review, Scientific Access into Mercer Subglacial Lake: Scientific Objectives, Drilling Operations and Initial Observations, *Ann. Glaciol.*, 62(85–86), 340–352, <https://doi.org/https://doi.org/10.1017/aog.2021.10>, 2021.
- Ragone, S. E. and Finelli, R.: Special Report 167: Procedures for Removing Surface Contaminants From Deep Ice Cores., 1972.
- 790 Rantanen, M., Karpechko, A. Y., Lipponen, A., Ruosteenoja, K., Vihma, T., Laaksonen, A., Nordling, K. and Hyvärinen, O.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, 3(168), 1–10, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- 795 Schaefer, J. M., Finkel, R. C., Balco, G., Alley, R. B., Caffee, M. W., Briner, J. P., Young, N. E., Gow, A. J. and Schwartz, R.: Greenland was nearly ice-free for extended periods during the Pleistocene, *Nature*, 540(7632), 252–255, <https://doi.org/10.1038/nature20146>, 2016.

- Shoemaker, B.: Herb Ueda Interview, Polar Oral Hist. Proj., 51 <http://hdl.handle.net/1811/44677>, 2002.
- 800 Souchez, R., Vandenschrick, G., Lorrain, R. and Tison, J.-L.: Basal ice formation and deformation in central Greenland: a review of existing and new ice core data, in *Deformation of Glacial Materials*, edited by A. J. Maltman, B. Hubbard, and M. J. Hambrey, pp. 13–22, Geological Society, Special Publications, London, <https://doi.org/10.1201/b14059-6>, 2000.
- Souchez, R., Jouzel, J., Landais, A., Chappellaz, J., Lorrain, R. and Tison, J. L.: Gas isotopes in ice reveal a vegetated central Greenland during ice sheet invasion, *Geophys. Res. Lett.*, 33(24), 20–23, <https://doi.org/10.1029/2006GL028424>, 2006.
- 805 Souney, J. M., Twickler, M. S., Hargreaves, G. M., Bencivengo, B. M., Kippenhan, M. J., Johnson, J. A., Cravens, E. D., Neff, P. D., Nunn, R. M., Orsi, A. J., Popp, T. J., Rhoades, J. F., Vaughn, B. H., Voigt, D. E., Wong, G. J. and Taylor, K. C.: Core handling and processing for the WAIS Divide ice-core project, *Ann. Glaciol.*, 55(68), 15–26, <https://doi.org/10.3189/2014AoG68A008>, 2014.
- Talalay, P. G.: Subglacial till and bedrock drilling, *Cold Reg. Sci. Technol.*, 86, 142–166, <https://doi.org/10.1016/j.coldregions.2012.08.009>, 2013.
- 810 Thomis, W.: Arctic Is Warming Up But Word Hasn't Reached Yanks at Outposts, *Chicago Dly. Trib.*, 5, 1955.
- Tison, J.-L., Souchez, R., Wolff, E. W., Moore, J. C., Legrand, M. R. and de Angelis, M.: Is a periglacial biota responsible for enhanced dielectric response in basal ice from the Greenland Ice Core Project ice core? *Jean-Louis, J. Geophys. Res.*, 103(98), 18885–18894, 1998.
- 815 Tison, J. L.: Diamond wire-saw cutting technique for investigating textures and fabrics of debris-laden ice and brittle ice, *J. Glaciol.*, 40(135), 410–414, <https://doi.org/10.1017/S0022143000007498>, 1994.
- Tison, J. L., Thorsteinsson, T., Lorrain, R. D. and Kipfstuhl, J.: Origin and development of textures and fabrics in basal ice at Summit, Central Greenland, *Earth Planet. Sci. Lett.*, 125(1–4), 421–437, [https://doi.org/10.1016/0012-821X\(94\)90230-5](https://doi.org/10.1016/0012-821X(94)90230-5), 1994.
- Ueda, H. T. and Garfield, D. E.: Special Report 126: Drilling through the Greenland Ice Sheet., 1968.
- 820 Venturelli, R. A., Siegfried, M. R., Roush, K. A., Li, W., Burnett, J., Zook, R., Fricker, H. A., Priscu, J. C., Leventer, A. and Rosenheim, B. E.: Mid-Holocene Grounding Line Retreat and Readvance at Whillans Ice Stream, West Antarctica, *Geophys. Res. Lett.*, 47(15), e2020GL088476, <https://doi.org/https://doi.org/10.1029/2020GL088476>, 2020.
- 825 Verbeke, V., Lorrain, R., Johnsen, S. J. and Tison, J. L.: A multiple-step deformation history of basal ice from the dye 3 (Greenland) core: New insights from the CO<sub>2</sub> and CH<sub>4</sub> content, *Ann. Glaciol.*, 35(May 2021), 231–236, <https://doi.org/10.3189/172756402781817248>, 2002.
- Weis, D., Demaiffe, D., Souchez, R., Gow, A. J. and Meese, D. A.: Ice sheet development in Central Greenland: Implications from the Nd, Sr and Pb isotopic compositions of basal material, *Earth Planet. Sci. Lett.*, 150(1–2), 161–169, [https://doi.org/10.1016/s0012-821x\(97\)00073-3](https://doi.org/10.1016/s0012-821x(97)00073-3), 1997.
- 830 Whalley, W. B. and Langway, C. C.: A scanning electron microscope examination of subglacial quartz grains from Camp Century core, Greenland - a preliminary study, *J. Glaciol.*, 25(91), 125–131, 1980.
- Willerslev, E., Cappellini, E., Boomsma, W., Nielsen, R., Hebsgaard, M. B., Brand, T. B., Hofreiter, M., Bunce, M., Poinar, H. N., Dahl-Jensen, D., Johnsen, S., Steffensen, J. P., Bennike, O., Schwenninger, J. L., Nathan, R.,



835 Armitage, S., De Hoog, C. J., Alfimov, V., Christl, M., Beer, J., Muscheler, R., Barker, J., Sharp, M., Penkman, K.  
E. H., Haile, J., Taberlet, P., Gilbert, M. T. P., Casoli, A., Campani, E. and Collins, M. J.: Ancient biomolecules  
from deep ice cores reveal a forested southern Greenland, *Science*, 317(5834), 111–114,  
<https://doi.org/10.1126/science.1141758>, 2007.

840 Wilson, G. S., Levy, R. H., Naish, T. R., Powell, R. D., Florindo, F., Ohneiser, C., Sagnotti, L., Winter, D. M.,  
Cody, R., Henrys, S., Ross, J., Krissek, L., Niessen, F., Pompillio, M., Scherer, R., Alloway, B. V., Barrett, P. J.,  
Brachfeld, S., Browne, G., Carter, L., Cowan, E., Crampton, J., DeConto, R. M., Dunbar, G., Dunbar, N., Dunbar,  
R., von Eynatten, H., Gebhardt, C., Giorgetti, G., Graham, I., Hannah, M., Hansaraj, D., Harwood, D. M.,  
Hinnov, L., Jarrard, R. D., Joseph, L., Kominz, M., Kuhn, G., Kyle, P., Läufer, A., McIntosh, W. C., McKay, R.,  
845 Maffioli, P., Magens, D., Millan, C., Monien, D., Morin, R., Paulsen, T., Persico, D., Pollard, D., Raine, J. I.,  
Riesselman, C., Sandroni, S., Schmitt, D., Sjunneskog, C., Strong, C. P., Talarico, F., Taviani, M., Villa, G.,  
Vogel, S., Wilch, T., Williams, T., Wilson, T. J. and Wise, S.: Neogene tectonic and climatic evolution of the  
Western Ross Sea, Antarctica — Chronology of events from the AND-1B drill hole, *Glob. Planet. Change*, 96–  
97, 189–203, <https://doi.org/10.1016/j.gloplacha.2012.05.019>, 2012.

### **Supplementary materials**

850 Table S1: Sample depths and frozen sub-sample masses

Table S2: Soil color

Table S3: Bulk density

Table S4: Magnetic susceptibility

Table S5: Natural Remanent Magnetization

855 Table S6: Pore ice pH and conductivity

Table S7: Sediment and ice content

Table S8: Grain size distribution

Table S9: Samples altered during storage

Figure S1: Comparison of estimated and measured ice content