



# Magnesium isotope fractionation processes during seafloor serpentization and implications for serpentinite subduction

Sune G. Nielsen<sup>1,2,3</sup>, Frieder Klein<sup>4</sup>, Horst R. Marschall<sup>5</sup>, Philip A. E. Pogge von Strandmann<sup>6</sup>, Maureen Auro<sup>1</sup>

5 <sup>1</sup>NIRVANA Labs, Woods Hole Oceanographic Institution, 02543 Woods Hole, MA, USA

<sup>2</sup>Department of Geology and Geophysics, Woods Hole Oceanographic Institution, 02543 Woods Hole, MA, USA

<sup>3</sup>Centre de Recherches Pétrographiques et Géochimiques, CNRS, Université de Lorraine, 15 rue Notre Dame des Pauvres, 54501 Vandoeuvre lès Nancy, France.

<sup>4</sup>Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, 02543 Woods Hole, MA, USA

10 <sup>5</sup>Institut für Geowissenschaften, Goethe Universität Frankfurt, Frankfurt am Main, Germany

<sup>6</sup>Institute of Geosciences, Johannes Gutenberg University, 55122 Mainz, Germany

*Correspondence to:* Sune G. Nielsen (snielsen@whoi.edu)

**Abstract.** Studies of magnesium (Mg) isotope ratios in subduction zone lavas have revealed small, but significant offsets from  
15 the mantle value with enrichments in the heavy isotopes. However, the very high concentration of Mg in the mantle contrasts with much lower concentrations in the subducted igneous crust and oceanic sediments, making these subduction components unlikely vehicles of the Mg isotope anomalies in arc lavas. Only serpentinites, which in various proportions form part of oceanic plates, have high Mg contents comparable to fresh mantle rocks and have thus been considered a potential source of exotic Mg in the source of arc magmas.

20 In this study we analyzed serpentinite samples from different oceanic settings for their Mg isotopic compositions. The majority of samples are indistinguishable from the depleted mantle ( $\delta^{26}\text{Mg} = -0.24 \pm 0.04 \text{ ‰}$ ) irrespective of their origin. Only a small number of seafloor-weathered serpentinites are slightly enriched in the heavy isotopes (up to  $\delta^{26}\text{Mg} = -0.14 \pm 0.03 \text{ ‰}$ ), implying that bulk serpentinites are unlikely sources of isotopically anomalous Mg in subduction zones.

We also developed a partial-dissolution method in which 5 % acetic acid for 180 minutes was shown to fully dissolve the  
25 minerals brucite and iowaite while leaving the serpentine mineral chrysotile essentially undissolved.

Partial dissolution of 11 bulk serpentinite samples revealed a Mg isotopic composition of brucite ( $\pm$ iowaite) that is systematically  $\sim 0.25 \text{ ‰}$  heavier than that of coexisting serpentine. Thus, preferential breakdown of brucite and/or iowaite in a subducted slab prior to serpentine could preferentially release isotopically heavy Mg, which could subsequently be transported into the source region of arc magmas. Such a scenario would require brucite/iowaite breakdown to occur at pressures in excess  
30 of 3 GPa and produce fluids with very high concentrations of Mg that could be transported to arc magma source regions.

Whether these conditions are met in nature has yet to be experimentally investigated.



## 1 Introduction

The release of volatiles from the subducting slab is an important process driving arc volcanism. However, how substantial amounts of water can be transported to sufficient depths prior to being released into the mantle is highly debated. Some studies argue that sediments and hydrothermally altered oceanic crust (AOC) are the primary sources of H<sub>2</sub>O, as these are directly exposed to seawater at the seafloor and are naturally located at the boundary between the slab and the mantle wedge in the subduction zone (Tatsumi, 1986; Hacker, 2008). It is unclear whether hydrous minerals such as lawsonite and phengite are sufficiently abundant and stable in many slabs to carry the amounts of water required to fuel arc volcanism (e.g. Schmidt and Poli, 1998). Additionally, serpentinite has been proposed as a vehicle to transport water to a sufficient depth to eventually trigger the formation of arc magmas (e.g. Rüpke et al., 2004). Serpentinite forms during the alteration of ultramafic rocks by water at a range of temperatures and pressures (<100–600 °C; (Lazar, 2020)). Serpentinite can form at mid-ocean ridges when mantle rocks are tectonically uplifted to crustal levels where it interacts with heated seawater, such as on slow- and ultraslow-spreading ridges in the Atlantic, Indian, and Arctic oceans (Bach and Früh-Green, 2010; Humphris and Klein, 2018). Serpentinite can also form in tectonic windows and fracture zones of fast-spreading crust and is believed to form along deep-reaching faults in oceanic trenches (Ranero et al., 2003; Mével and Stadoumi, 1996). While direct evidence for serpentinite subduction has been found in the slow spreading crust subducting underneath the Lesser Antilles (Klein et al., 2017), subduction of serpentinite in intermediate- to fast-spreading oceanic plates in the Pacific is chiefly based on geophysical inference (Ranero et al., 2003).

Geochemical evidence for serpentinite subduction remains controversial. One example of a geochemical proxy for serpentinite subduction is magnesium (Mg) isotope compositions. Much of this attention has been spurred by the discovery of  $\delta^{26}\text{Mg}$  ( $\delta^{26}\text{Mg}$  is the deviation of  $^{26}\text{Mg}/^{24}\text{Mg}$  from that of the standard DSM-3 (Galy et al., 2003) in ‰) values slightly higher than the mantle in some arc lavas (Teng et al., 2016; Li et al., 2017). While it has been argued that these high  $\delta^{26}\text{Mg}$ -values may be accounted for by fractional crystallization and crustal assimilation processes (Brewer et al., 2018), others conclude that they must originate from the sub-arc mantle source region (Teng et al., 2016; Li et al., 2017; Chen et al., 2023b). In this case, the high  $\delta^{26}\text{Mg}$  values have been challenging to explain via addition of sediment and AOC into arc lava source regions, because crustal rocks have much lower Mg contents than the mantle wedge (Li and Schoonmaker, 2014; Rudnick and Gao, 2003; McDonough and Sun, 1995). In contrast, Mg contents of peridotite are much higher and Mg is conserved during serpentinitization of peridotite (Klein and Le Roux, 2020), which means that there is potential for even moderate amounts of subducted serpentinite to modify the overall Mg isotopic budget of arc lavas. Two potential processes have been argued to connect serpentinite with the Mg isotope budget of subduction zones: (1) release of fluids from the slab into the sub-arc mantle that are enriched in the heavy Mg isotopes due to the breakdown of serpentine (Teng et al., 2016) or (2) Mg isotope fractionation during serpentinitization at the seafloor that could cause serpentinites to be enriched in the heavy isotopes relative to the otherwise homogenous mantle rocks (Zhao et al., 2023).



65 Both hypotheses provide some challenges. It has been shown in experiments at pressures (P) and temperatures (T) relevant to  
subduction zones that fluids released from crustal slab lithologies (sediments and AOC) carry very little Mg (Kessel et al.,  
2005; Spandler et al., 2007; Carter et al., 2015; Manning, 2004), possibly due to low overall Mg solubility in fluids at these T  
and P. Experimental investigations and theoretical models of Mg solubility in fluids expelled during serpentinite breakdown  
at various subduction zone conditions (1–10 GPa and 300 to 1200 °C) predict that these can be both enriched and depleted in  
70 Mg (Stalder et al., 2001) and potentially cause Mg-metasomatism of adjacent gabbroic lithologies as in the Ligurian Alps of  
Italy (Codillo et al., 2022). Thus, there is no consensus to what extent fluids expelled from serpentinite breakdown affect the  
Mg isotope budget of arc lava source regions.

In some cases, the Mg isotope composition of serpentinite display bulk-rock  $\delta^{26}\text{Mg}$  values slightly higher than the mantle  
(Zhao et al., 2023; Wang et al., 2023; Beinlich et al., 2014; Liu et al., 2017), but there is disagreement as to the processes  
75 responsible for these heavy isotope enrichments. Some argue that seafloor weathering is the main process responsible for these  
values (Wang et al., 2023; Liu et al., 2017; Li et al., 2023), whereas others conclude that the serpentinization reaction itself is  
associated with Mg loss that causes isotope fractionation (Zhao et al., 2023). There is, hence, a need for a more comprehensive  
investigation that distinguishes the Mg isotope effects of primary serpentinization and secondary seafloor weathering.

In addition to serpentine, oceanic serpentinite contains up to 15 wt.% of brucite ((Mg,Fe)(OH)<sub>2</sub>) (Klein et al., 2020) and  
80 theoretical considerations have indicated that the Mg isotope compositions of brucite and serpentine can be substantially  
fractionated relative to each other, as well as relative to the fluids from which they precipitate (Wang et al., 2019; Wimpenny  
et al., 2014). Given that the thermodynamic stability fields of brucite and serpentine are different (Lazar, 2020), the increase  
in pressure and temperature during subduction could provide an additional mechanism for preferential mobilization of  
isotopically fractionated Mg from serpentinites in subduction zones. However, to date no study has investigated whether  
85 brucite and serpentine in serpentinite follow predicted Mg isotope fractionation patterns during serpentinization.

Here we present Mg isotope data for a set of serpentinites from mid-ocean ridge settings and subduction-zone forearc regions  
revealing no bulk Mg isotope change associated with serpentinization. We also perform partial dissolution experiments, where  
only brucite is dissolved, which reveal that brucite is systematically higher in  $\delta^{26}\text{Mg}$  than coexisting serpentine.

## 2 Samples and methods

### 90 2.1 Sample descriptions

Serpentinites included in this study were recovered from mid-ocean ridge, off-axis, and subduction-zone settings. Samples  
recovered during Ocean Drilling Program (ODP) Legs 153 (Site 920B) and 209 (Sites 1268A, 1271B, 1272A, and 1274A) on  
the slow-spreading Mid-Atlantic Ridge include partly to completely serpentinized harzburgite and dunite. Their secondary  
mineral assemblages are dominated by serpentine, brucite, and magnetite in addition to minor chlorite and iowaite (Klein et  
95 al., 2020), and traces of Ni-bearing sulfides and alloys. Brucite, which formed together with serpentine at the expense of olivine  
in mesh texture, is partially and locally completely replaced by iowaite.



Samples from Deep Sea Drilling Program Leg 82 (Site 558)) were recovered off the Mid-Atlantic Ridge in 34.7 Ma crust (37°46.24' N, 37°20.61' W). Harzburgite is completely serpentinized and mainly composed of serpentine, iowaite, and maghemite. In contrast to samples from ODP Legs 153 and 209, iowaite preferentially occurs in late-stage veins that cut across mesh and bastite texture.

Samples from the Puerto Rico Trench were recovered during Cruise 19 of RV Chain (Bowin *et al.*, 1966; Klein *et al.*, 2017). Two dredges from the North Wall of the Puerto Rico Trench, which is currently undergoing subduction beneath the Caribbean Plate, include highly weathered serpentinized peridotite (dredge D10) composed of lizardite, chrysotile, minor antigorite, chlorite, hematite, goethite, clay minerals, and quartz, and Si-metasomatized serpentinite (dredge D2) chiefly composed of antigorite and talc. Dating of zircon in mafic veins that cut sample D10-9 yielded an age of 114.8 Ma, indicating that these serpentinites were originally formed at the Cretaceous Mid-Atlantic Ridge and were subsequently transported by seafloor spreading to their current location (Klein *et al.*, 2017).

Samples from the Mariana forearc were drilled during ODP Leg 125 and 195 from Conical Seamount and South Chamorro Seamount, respectively (D'antonio and Kristensen, 2004; Fryer *et al.*, 1992). These samples are partly to completely serpentinized peridotites chiefly composed of serpentine (lizardite, chrysotile, antigorite), brucite, iowaite, chlorite, and magnetite (Kahl *et al.*, 2015; Klein *et al.*, 2020). In contrast to other samples included in this study, peridotite from South Chamorro and Conical was serpentinized in the shallow mantle wedge by sediment-derived pore fluids evolved during subduction of the Pacific plate (Kahl *et al.*, 2015; Nielsen *et al.*, 2015; Debret *et al.*, 2019).

## 2.2 Partial dissolution experiments

In order to investigate whether brucite, iowaite and serpentine are characterized by different Mg isotope compositions, we developed a partial dissolution protocol designed to preferentially dissolve brucite and iowaite, whereas serpentine was left in the residue. The isotope composition of serpentine could thereby be calculated via mass balance using the bulk Mg isotope composition and concentration.


We performed dissolution experiments of pure single mineral materials of brucite, iowaite and chrysotile using acetic acid to determine the kinetics of their dissolution at different acetic acid strengths. For brucite and iowaite we conducted 4 separate experiments using 5 %, 10 %, 20 % and 40 % acetic acid at ambient room temperature (~20°C), respectively. Sample aliquots were taken out after 1 min, 10 min, 100 min, 360 min (=6 h) and 1440 min (=24 h) and the Mg concentrations were determined (Fig. 1). If less than 100 % of the mineral had dissolved, then a mineral sample aliquot was also taken for Mg isotopic analysis. For chrysotile only one experiment was performed at 5 % acetic acid strength with samples taken at 80 min and 1200 min. The Mg isotope compositions were not determined for this experiment as only a few percent of the serpentine had dissolved during the experiment (Fig. 1).

We selected 11 serpentinite samples for partial dissolution. Samples were selected based on their brucite and iowaite contents, lack of extensive seafloor weathering (Klein *et al.*, 2020), as well as being completely or almost completely serpentinized.



130 Thus, no or only trivial amounts of primary Mg bearing minerals (i.e. olivine, pyroxenes) remained, which allowed us to use  
the bulk Mg isotope compositions to reconstruct the Mg isotope composition of serpentine in these samples via mass balance.  
Concentrations of Na, Mg, Ca, Fe, Al, Cr, Mn, Co, Ni, and Zn in the partial dissolution samples were determined on a Thermo  
iCap quadrupole inductively couple plasma mass spectrometer (ICP-MS) located in the WHOI Plasma Facility. Concentrations  
were determined via reference to serially diluted USGS reference materials AGV-2 and BHVO-2 run at the beginning and end  
135 of each sequence. Instrumental drift was corrected by monitoring the signal of indium that was added to each sample and  
reference material in a known quantity. Following previous studies from our group that use identical methods for concentration  
determination we infer that the external reproducibility of these analyses are ~7 % (1sd) (Shu et al., 2022; Wang et al., 2022).

**Figure 1: Fraction of Mg dissolved in single mineral dissolution experiments. Acetic acid strength is color coded. The fraction of total Mg dissolved for brucite was normalized to the values for the end of the experiment since no solids remained, whereas chrysotile**



140 was calculated by assuming an idealized stoichiometric formula of  $Mg_3(Si_2O_5)(OH)_4$ . Uncertainties for the chrysotile  
are smaller than symbol sizes.

### 2.3 Magnesium separation and isotope measurements

Purification of Mg from sample matrix was performed as previously detailed (Chen et al., 2023a; Pogge Von Strandmann et  
145 al., 2011). Briefly, the dissolved samples were purified using a two-stage cation exchange column procedure, using AG50W  
X-12 ion exchange resin, eluting the Mg using 2M  $HNO_3$ .

Magnesium isotope compositions were measured using a Nu Plasma 3 MC-ICP-MS at the LOGIC laboratories in London,  
using a sample-standard bracketing method relative to DSM-3. Based on the repeated purification and analysis of reference




materials USGS BCR-2 ( $\delta^{26}\text{Mg} = -0.24 \pm 0.08 \text{ ‰}$ , 2sd,  $n = 11$ ), JP-1 ( $-0.23 \pm 0.07 \text{ ‰}$ , 2sd,  $n = 3$ ), and seawater ( $\delta^{26}\text{Mg} = -0.82 \pm 0.04 \text{ ‰}$ , 2sd,  $n = 15$ ), the accuracy of our measurements is indistinguishable from other studies (Foster et al., 2010; Pogge Von Strandmann et al., 2011), and the long-term external precision on  $\delta^{26}\text{Mg}$  is  $\pm 0.08 \text{ ‰}$  (2sd).

### 3 Results and discussion

#### 3.1 Partial dissolution experiments

The single-mineral dissolution experiments reveal that both brucite and iowaite dissolve rapidly in all acetic acid concentrations (Figs. 1 and 2). Brucite dissolved in less than 1 minute in all experiments, whereas iowaite required more than 100 minutes to fully dissolve in 5 % acetic acid. Chrysotile did not dissolve appreciably over the 1200 minutes we conducted the experiment in 5 % acetic acid (Fig. 1). The experiments where iowaite had partially dissolved exhibit a significant Mg isotopic difference as a function of the fraction of Mg dissolved (Fig. 2), suggesting that partial dissolution of iowaite is associated with Mg isotope fractionation, whereby the light isotope is preferentially released into solution first. However, we infer that the net Mg isotope fractionation factor quickly approaches 0 as a larger fraction of iowaite is dissolved. Based on these results we chose to conduct the partial dissolution of the bulk serpentinites for 180 minutes in 5 % acetic acid. This configuration likely ensures full dissolution of brucite and iowaite, whereas dissolution of serpentine would be minimal. The single mineral dissolution experiments imply that it is not possible to chemically separate brucite and iowaite and, therefore, the partial dissolution of bulk serpentinites provides information about the Mg isotope difference between brucite/iowaite and serpentine.



**Figure 2: Results of iowaite single mineral partial dissolution experiments. The fraction of total Mg dissolved was normalized to the values for the end of the experiment since no solids remained. Experiments without full mineral dissolution in 5 % and 10 % acetic acid were analyzed for Mg isotopes (a), whereas the fully dissolved mineral was measured once (black symbol in panel a).**



### 170 3.2 Bulk serpentinite data

Magnesium isotope compositions of 33 bulk serpentinite samples (Table 1) reveal only very minor variation, with the majority of samples exhibiting values within uncertainty of the mantle value (Figure 3). Only a few of the dredged serpentinite samples (those from the Puerto Rico trench) exhibit values that extend up to  $\delta^{26}\text{Mg} = -0.14 \pm 0.03 \text{ ‰}$ , slightly higher than the mantle ( $\delta^{26}\text{Mg} = -0.24 \pm 0.04 \text{ ‰}$ , (Hin et al., 2017; Teng et al., 2010)).

175 As previously shown based on thermogravimetric analysis, mineralogical constraints, and major element data, dredged serpentinites typically record various degrees of seafloor weathering (Klein et al., 2020), which has been argued to result in Mg isotope fractionation towards more positive  $\delta^{26}\text{Mg}$ -values (Wang et al., 2023; Liu et al., 2017). The serpentinite samples that were not affected by weathering, all of which were recovered by scientific ocean drilling, record no Mg isotope deviations from the mantle, which is consistent with the conservation of Mg during serpentinization (Klein et al., 2020; Klein and Le  
180 Roux, 2020). Magnesium loss and Mg isotope fractionation during serpentinization, which has been argued for in the recent literature (Zhao et al., 2023), was possibly caused by (seafloor) weathering prior to metamorphic re-equilibration of these tectonically exhumed rocks or by metasomatism. The fact that these rocks contain negligible brucite (Zhao et al., 2023) is also a strong indication that they were subject to post-serpentinization modifications, e.g., via seafloor weathering or Si-  
metasomatism.

185

Table 1: Bulk Mg isotope compositions of serpentinites


Sample name	Location	Sampling method	$\delta^{25}\text{Mg}$ (‰)	2se	$\delta^{26}\text{Mg}$ (‰)	2se	n
D2-1	Puerto Rico trench	dredge	-0.12	0.02	-0.24	0.03	1
D2-2	Puerto Rico trench	dredge	-0.09	0.02	-0.17	0.02	2
D2-4	Puerto Rico trench	dredge	-0.09	0.03	-0.17	0.04	1
D2-5	Puerto Rico trench	dredge	-0.13	0.02	-0.26	0.06	2
D2-6	Puerto Rico trench	dredge	-0.11	0.04	-0.22	0.03	2
D10-1	Puerto Rico trench	dredge	-0.11	0.01	-0.24	0.04	1
D10-3	Puerto Rico trench	dredge	-0.13	0.03	-0.25	0.04	1
D10-7	Puerto Rico trench	dredge	-0.11	0.02	-0.23	0.03	1
D10-8	Puerto Rico trench	dredge	-0.08	0.04	-0.14	0.03	2
D10-9	Puerto Rico trench	dredge	-0.07	0.03	-0.19	0.03	1
D10-12	Puerto Rico trench	dredge	-0.08	0.01	-0.17	0.03	1
D10-15	Puerto Rico trench	dredge	-0.10	0.03	-0.15	0.04	1
D10-16	Puerto Rico trench	dredge	-0.11	0.02	-0.22	0.02	1
209-1268A-8R1-28-35	15°20FZ	drilling	-0.14	0.01	-0.29	0.02	1
209-1272A-14R1-43-53	15°20FZ	drilling	-0.13	0.04	-0.24	0.04	3



209-1268A-4R1-44-55	15°20FZ	drilling	-0.11	0.01	-0.24	0.03	1
209-1271B-17R1-61-69	15°20FZ	drilling	-0.13	0.04	-0.24	0.05	2
209-1272A-21R1-88-100	15°20FZ	drilling	-0.11	0.01	-0.27	0.03	1
209-1272A-27R2-78-88	15°20FZ	drilling	-0.10	0.02	-0.21	0.03	1
209-1274A-16R2-26-38	15°20FZ	drilling	-0.15	0.02	-0.26	0.02	1
195-1200A-17R2-76-79	South Chamorro Seamount	drilling	-0.13	0.03	-0.28	0.02	1
195-1200A-11R1-47-49	South Chamorro Seamount	drilling	-0.12	0.04	-0.25	0.02	1
195-1200A-13R1-121-124	South Chamorro Seamount	drilling	-0.08	0.03	-0.18	0.03	1
125-779A-10R2-51-53	Conical seamount	drilling	-0.12	0.02	-0.28	0.04	1
125-779A-17R4-32-34	Conical seamount	drilling	-0.17	0.03	-0.26	0.04	1
125-779A-31R2-85-87	Conical seamount	drilling	-0.12	0.03	-0.25	0.02	1
82-558Z-42R1-9-11	Azores off axis	drilling	-0.10	0.01	-0.22	0.03	1
82-558Z-42R1-133-135	Azores off axis	drilling	-0.12	0.02	-0.23	0.04	1
82-558Z-48R1-45-46	Azores off axis	drilling	-0.10	0.03	-0.25	0.03	1
153-920B-2R1-80-82	MARK	drilling	-0.11	0.03	-0.26	0.03	1
153-920B-5R2-35-38	MARK	drilling	-0.06	0.02	-0.18	0.03	1
153-920B-10R1-82-86	MARK	drilling	-0.09	0.04	-0.20	0.04	1
153-920B-12R2-140-143	MARK	drilling	-0.15	0.02	-0.24	0.04	1

190

195





**Figure 3: Bulk-rock Mg isotopic compositions of 33 seafloor serpentinites. Error bars are 2 $\sigma$ . Circle symbols are drill cores, whereas squares are dredged (i.e. Puerto Rico trench). The average mantle value is based on literature data (Teng et al., 2016; Hin et al., 2017).**

200 The Mg isotopic compositions are quasi invariant regardless of the tectonic environment that the serpentinization took place in (i.e. mid-ocean ridge, subduction zone forearc), which is consistent with previous results from similar environments (Wang et al., 2023; Liu et al., 2017). This result implies that the serpentinizing fluid composition, i.e. seawater or fluids sourced from dehydration of sediments (Nielsen et al., 2015), or the olvine-to-orthopyroxene ratio of the protolith, do not affect the Mg isotope budget of serpentinite (Pogge Von Strandmann et al., 2015). Hence, serpentinite subduction does not introduce  
 205 isotopically anomalous Mg into subduction zones and serpentinite is unlikely to form an endmember bulk source of the positive  $\delta^{26}\text{Mg}$ -values observed in some arcs (Teng et al., 2016; Li et al., 2017). Should these positive  $\delta^{26}\text{Mg}$ -values in arcs be generated by processes involving serpentinite, then these signatures must be generated in the subduction zone itself by partial mobilization of Mg from serpentinite associated with Mg isotope fractionation. Two different scenarios could achieve this: (1) either material with low  $\delta^{26}\text{Mg}$ -values is removed from serpentinite at relatively low pressures and temperatures, leaving higher  
 210  $\delta^{26}\text{Mg}$  values in the residue that could then contribute to the arc-lava source regions; or (2) material with positive  $\delta^{26}\text{Mg}$ -values is released from serpentinite and is transported into the source of arc magmas (Teng et al., 2016).

One potentially critical parameter for these scenarios could be, if the dominant Mg-bearing minerals in serpentinite, i.e. serpentine and brucite, were associated with different Mg isotope compositions. This has indeed been suggested based on theoretical calculations and experiments (Wang et al., 2019; Wimpenny et al., 2014; Wimpenny et al., 2010; Gao et al., 2018).  
 215 However, the exact isotope fractionation between these two phases is not agreed upon with evidence pointing towards brucite exhibiting both higher and lower  $\delta^{26}\text{Mg}$ -values than serpentine at equilibrium.

### 3.3. Mg isotope fractionation between major serpentinization minerals

Of the 11 samples for which acetic-acid partial dissolution was performed, all recorded  $\delta^{26}\text{Mg}$  values are within uncertainty of  
 220 or higher than the bulk measurement (Table 2). We confirmed through thermogravimetric analysis (TGA) of 8 samples that complete dissolution of brucite and iowaite occurred, while serpentine was retained (Fig. S1).

**Table 2: Elemental concentrations and Mg isotope compositions of leached serpentinites**

Sample	Mg	Fe	Al	Ca	Cr	Mn	Co	Ni	Zn	$\delta^{26}\text{Mg}$	2 $\sigma$
209-1272A-27R2-78-88	6.30	1.62	0.047	0.009	246	379	30	622	14	-0.22	0.03
209-1272A-21R1-88-100	4.55	0.95	0.064	0.004	332	283	33	826	8	-0.01	0.02
209-1274A-16R2-26-28	6.19	1.70	0.033	0.017	188	413	20	696	17	-0.06	0.05
209-1271B 17R1 61-69	6.53	0.69	0.009	0.004	86	440	21	1365	10	-0.04	0.02
82-558Z-48R1-45-46	4.96	1.77	0.007	0.012	25	233	83	2143	19	-0.26	0.02
82-558Z-42R1-133-135	3.80	1.09	0.042	0.011	292	172	83	1795	11	-0.06	0.03
195-1200A-11R1-47-49	5.35	1.22	0.009	0.177	68	375	54	1068	18	-0.10	0.03



195-1200A-13R1-121-124	6.61	1.65	0.028	0.200	59	456	32	521	20	-0.07	0.01
195-1200A-17G-76-79	4.71	1.26	0.018	0.127	66	316	47	793	12	-0.10	0.03
153-920B-12R2-140-143	6.26	0.63	0.059	0.006	26	468	26	1424	9	-0.13	0.02
125-779A-10R2-51-53	3.76	1.24	0.001	0.050	10	213	29	729	10	-0.16	0.03

**Major elements (Mg, Fe, Al, Ca) in weight %, all others in µg/g**

225

Although the procedure dissolves both brucite and iowaite while leaving serpentine essentially undissolved, we note that iowaite in mesh texture is generally thought to form from replacement of brucite (D'antonio and Kristensen, 2004; Klein et al., 2020). Given that the cation stoichiometry between these two minerals is essentially the same (molar Mg/(Mg+Fe) ≈ 0.8), we infer that all the Mg included in iowaite likely was initially present in brucite. Therefore, the majority of Mg in the leachates reflects the Mg contained in brucite during initial serpentinization. The single-mineral dissolution experiments reveal that minor amounts of Mg from serpentine (~4 %) does dissolve within the first hours of reacting 5 % acetic acid with chrysotile, whereas only little additional Mg was released when extending the experiment to 20 hours (Fig. 1). Assuming that ~4 % of serpentine-bound Mg dissolved during our bulk serpentinite leaching experiments, we calculate that 12–22 % of the Mg released during the experiments originated from serpentine (i.e.  $fMg_{serp,leach}$ , Table 3). Similarly, we use the bulk serpentinite and total leached Mg contents of the samples, while assuming that all Mg in the samples are either located in serpentine, iowaite, or brucite, to calculate the fraction of Mg in the bulk sample accommodated by serpentine (i.e.  $fMg_{serp,bulk}$ ). There is no significant correlation between the fraction of Mg released from serpentine and the Mg isotope composition of the leach (Fig. 4), suggesting that partial dissolution of serpentine did not cause substantial isotope fractionation. In any case, such a process would likely be associated with kinetic isotope fractionation similar to what was observed for partial iowaite dissolution. That would result in lower  $\delta^{26}Mg$ -values in the leachate, opposite to the values higher than the bulk recorded in the leach experiments (Table 2).

Because all samples leached were fully serpentinized without any remaining olivine or pyroxene, we can use mass balance to calculate the Mg isotope compositions of the major Mg bearing serpentinization minerals. We use the following two mass balance equations to determine the Mg isotope compositions of the two phases in the serpentinites (i.e.  $\delta^{26}Mg_{serp}$  and  $\delta^{26}Mg_{b+i}$ ):

245

$$fMg_{serp,bulk} * \delta^{26}Mg_{serp} + fMg_{b+i,bulk} * \delta^{26}Mg_{b+i} = \delta^{26}Mg_{bulk} \quad (\text{eq. 1})$$

$$fMg_{serp,leach} * \delta^{26}Mg_{serp} + fMg_{b+i,leach} * \delta^{26}Mg_{b+i} = \delta^{26}Mg_{leach} \quad (\text{eq. 2})$$

Equation 2 assumes that no Mg isotope fractionation occurred during leaching, which is the case for brucite and iowaite, since they were both fully dissolved. It is unclear if the Mg isotope fractionation observed for iowaite can be translated to serpentine or indeed whether any substantial effect would be present after 3 hours of reaction (as opposed to the 1–10 minutes where we observed small effects). The calculated Mg isotope compositions of the mineral phases (Table 3) allows an assessment of the net Mg isotope fractionation ( $\Delta^{26}Mg$ ) between brucite/iowaite and serpentine (Table 3).



255

**Table 3: Mg isotope mass balance of leached serpentinites**

Sample	Mg <sub>bulk</sub> *	Mg <sub>residue</sub>	Mg <sub>serp.bulk</sub> (4% diss.)	fMg <sub>serp.bulk</sub>	fMg <sub>b+i.bulk</sub>	fMg <sub>serp.leach</sub>	fMg <sub>b+i.leach</sub>	δ <sup>26</sup> Mg <sub>serp</sub>	δ <sup>26</sup> Mg <sub>b+i</sub>	Δ <sup>26</sup> Mg
82-558Z-48R1, 45-46	22.98	18.01	18.76	0.82	0.18	0.151	0.849	-0.24	-0.26	-0.02
82-558Z-42R1, 133-135	23.84	20.04	20.87	0.88	0.12	0.220	0.780	-0.26	0.00	+0.25
125-779A-10R-2, 51-53	24.13	20.37	21.22	0.88	0.12	0.225	0.775	-0.30	-0.13	+0.17
153-920B-12R2, 140-143	24.33	18.07	18.83	0.77	0.23	0.120	0.880	-0.28	-0.11	+0.17
195-1200A-11R1 47-49	24.67	19.32	20.12	0.82	0.18	0.150	0.850	-0.29	-0.06	+0.23
195-1200A-13R1 121-124	25.06	18.45	19.22	0.77	0.23	0.116	0.884	-0.22	-0.05	+0.17
195-1200A-17G 76-79	24.64	19.93	20.76	0.84	0.16	0.177	0.823	-0.32	-0.05	+0.27
209-1271B 17R1 61-69	24.69	18.16	18.92	0.77	0.23	0.116	0.884	-0.31	-0.01	+0.30
209-1272A-21R1 88-100	23.99	19.44	20.25	0.84	0.16	0.178	0.822	-0.33	0.06	+0.38
209-1272A-27R2 78-88	23.85	17.55	18.28	0.77	0.23	0.116	0.884	-0.21	-0.22	-0.01
209-1274A-16R2 26-28	24.77	18.58	19.36	0.78	0.22	0.125	0.875	-0.32	-0.02	+0.31

**Mg concentrations (bulk, residue and serpentine) in weight %. Mg<sub>residue</sub> calculated from leached Mg in Table 2**


\* - bulk Mg concentrations previously published (Paulick et al., 2006; Klein et al., 2017)

260 Although most samples contained iowaite (Supplement), one sample (82-558Z-48R1, 45-46) presented iowaite in monomineralic veins that cut across pseudomorphic textures and likely formed via co-precipitation from mixing of seawater and a more alkaline solution. This sample is one of the two with Δ<sup>26</sup>Mg < 0. We, therefore, conclude that late-stage formation of iowaite veins likely caused precipitation of isotopically light Mg relative to what initially precipitated as brucite during serpentinization. Unfortunately, we do not have a thin section for the other sample with Δ<sup>26</sup>Mg < 0 (209-1272A-27R2 78-88) to corroborate this conclusion, but we infer that this sample also contains late stage iowaite veins.

265 Apart from the samples with late-stage iowaite veins, all samples reveal systematic and similar Mg isotope fractionation between serpentine and brucite of Δ<sup>26</sup>Mg = +0.25 ± 0.15 ‰ (2sd), implying that brucite formed during serpentinization is isotopically about 0.25 ‰ heavier than co-existing serpentine. This result contrasts Δ<sup>26</sup>Mg ~ -0.1 to -0.8 inferred for brucite dissolution during post serpentinization alteration (Li et al., 2023) and is in stark contrast to calculations of Mg isotope equilibrium between brucite and the serpentine mineral lizardite that imply an equilibrium isotope fractionation at 300 °C of  
 270 Δ<sup>26</sup>Mg ~ -1 ‰ (Wang et al., 2019). On the other hand, brucite-precipitation experiments have revealed that brucite preferentially incorporates heavy Mg isotopes relative to the solution from which it precipitates at room temperature, with Δ<sup>26</sup>Mg ~ +0.5 ‰ (Wimpenny et al., 2014). Given that brucite in the examined serpentinites likely precipitated at temperatures of 150–300°C (Klein et al., 2014) and that isotope fractionation factors (equilibrium and/or kinetic) roughly scale somewhere between 1/T and 1/T<sup>2</sup> (Chacko et al., 2001; Bigeleisen and Mayer, 1947), it is expected that fractionation factors are about one  
 275 quarter (for 1/T<sup>2</sup> at 300°C) to three quarters (for 1/T at 150°C) of those found at room temperature. This range would correspond to Δ<sup>26</sup>Mg = +0.13 to +0.38 ‰, identical to what we find here empirically (Table 3). Our data, therefore, imply that brucite formation proceeds through precipitation from a fluid that sourced its Mg from dissolution of olivine (or previously formed serpentine since these are isotopically very similar). Since most Mg during serpentinization forms serpentine, it is expected



that serpentine broadly retains the Mg isotope composition of the bulk starting material. On the contrary, brucite should express  
 280 the Mg isotope fractionation occurring during precipitation, since only a small fraction of the Mg in solution would likely be  
 used to form brucite. These observations imply that brucite and serpentine in serpentinite do not form at thermodynamic Mg  
 isotope equilibrium. Given that brucite and serpentine often form intergrowths suggesting simultaneous formation (Klein et  
 al., 2020; Hostetler et al., 1966), we infer that both minerals precipitate from the same fluid where any Mg isotope fractionation  
 occurring during serpentine precipitation is attenuated due to almost quantitative uptake of Mg from the solution.



285

**Figure 4: Fraction of total Mg leached that originated from serpentine ( $fMg_{\text{serp,leach}}$ ) plotted against measured Mg isotope composition of leach. Samples in dark red are those in which monomineralic veins of iowaite were identified. Arrows indicate mixing between brucite (with an assumed  $\delta^{26}\text{Mg} = 0\text{‰}$ ) and partially dissolved serpentine (with an assumed  $\delta^{26}\text{Mg} = -0.29\text{‰}$ ). The three lines denote congruent serpentine dissolution (full line), dissolution isotope fractionation of  $-0.21\text{‰}$  (short-dashed line), and dissolution isotope fractionation of  $-0.71\text{‰}$  (long-dashed line).**

290

### 3.4. Implications for subduction zone cycling of Mg isotopes

As outlined in the introduction, several volcanic arcs have been investigated that exhibit small, but measurable positive Mg isotope anomalies relative to that of the upper mantle, which has been inferred to be related to subduction of isotopically heavy serpentinite (Teng et al., 2016; Li et al., 2017). However, our data for unweathered serpentinite, together with several other  
 295 data sets (Wang et al., 2023; Liu et al., 2017){Li, 2023 #5724}, reveal that bulk serpentinite is unfractionated relative to the mantle. This demonstrates that the bulk Mg content of serpentinite cannot be responsible for the positive Mg isotope anomalies observed in some volcanic arcs. However, given that the mineral brucite exhibits on average  $\delta^{26}\text{Mg} = -0.04\text{‰}$  (excluding the iowaite vein samples, Table 3), significantly higher than those observed in most arc lavas ( $\delta^{26}\text{Mg} < -0.1\text{‰}$ ), preferential mobilization of brucite-bound Mg from the slab could be responsible for the observed Mg isotope anomalies in some arc  
 300 magmas. If such a process was to operate, then breakdown of brucite would need to form a fluid rich in Mg that could operate in three distinct scenarios of slab-to-mantle transport:

(1) migrate into the mantle wedge and cause melting in the wedge peridotite – in this case the brucite breakdown would have to occur at  $\geq 4$  GPa, such that the fluids can be transported into the sub-arc region.



(2) migrate into the eclogite or metasediment layer and cause melting there, requiring similar pressures of brucite breakdown  
305 as in (1). However, crustal melts forming from sediments and eclogite are tonalitic to rhyolitic (Hermann and Rubatto, 2009;  
Klemme et al., 2002), limiting the efficiency of Mg transport.

(3) metasomatise the slab-mantle interface as part of a melange (forming serpentine, chlorite,  $\pm$ talca, amphiboles) at various  
depths. These metasomatic/mixed materials could then be transported into the arc-magma source region – by diapirism, partial  
melting or dehydration.

310 Studies of the relative stability of brucite and serpentine reveal that assemblages containing both minerals typically are stable  
to temperatures of  $\sim$ 400–550°C at 2–6 GPa (Kempf et al., 2020; Lazar, 2020; Johannes, 1968), whereas serpentine on its own  
can be stable to higher temperatures in the same pressure range (Ferrand, 2019; Ulmer and Trommsdorff, 1995; Wunder and  
Schreyer, 1997). These stability ranges are compatible with brucite being preferentially mobilized at depths of  $\sim$ 100-150 km  
within the lithospheric mantle of the slab in cold and intermediate subduction zones (Van Keken and Wilson, 2023). Similarly,  
315 previously inferred P-T ranges of fluid release from the slab may also support preferential breakdown of brucite as a source of  
fluids to arc magma source regions (Hacker, 2008; Chermia et al., 2015; Konrad-Schmolke et al., 2016). On the other hand,  
even though brucite breakdown may produce a free aqueous fluid, it is currently uncertain if such a fluid would carry much  
Mg (Codillo et al., 2022; Kessel et al., 2005; Spandler et al., 2007; Carter et al., 2015; Manning, 2004; Stalder et al., 2001).

It is also possible that none of the conditions required for brucite to induce Mg isotope anomalies in arc magmas are met. In  
320 this case, serpentinite is unlikely to explain variations in arc lava Mg isotope variations and an alternative process is required.  
For example, the  $\delta^{26}\text{Mg}$  observed in (differentiated) arc lavas may not be a magma source signal reflecting the composition of  
the mantle wedge, but may instead be a result of magmatic fractionation in the crust of the upper plate (Brewer et al., 2018).  
Such a process is testable through detailed Mg isotope measurements of arc magmatic rocks.

325 **Author contribution:** SGN, FK and HRM conceived the study. SGN and FK designed the mineral dissolution experiments  
and MA carried them out. MA and SGN collected concentration data, FK collected TGA data, and PAEPS collected Mg  
isotope data. SGN prepared the manuscript with contributions from all co-authors.

**Competing interests:** The authors declare that they have no conflict of interest.

330

**Acknowledgements:** This study was supported by funds from NSF grant EAR-1829546 to SGN and the Joint Initiative  
Awards Fund from the Andrew W. Mellon Foundation to SGN and FK.

#### 4. References

Bach, W. and Früh-Green, G. L.: Alteration of the oceanic lithosphere and implications for seafloor processes, *Elements*, 6,  
335 173-178, 2010.



- Beinlich, A., Mavromatis, V., Austrheim, H., and Oelkers, E. H.: Inter-mineral Mg isotope fractionation during hydrothermal ultramafic rock alteration—Implications for the global Mg-cycle, *Earth Planet. Sci. Lett.*, 392, 166-176, 2014.
- Bigeleisen, J. and Mayer, M. G.: Calculation of Equilibrium Constants for Isotopic Exchange Reactions, *Journal of Chemical Physics*, 15, 261-267, 1947.
- 340 Brewer, A. W., Teng, F.-Z., and Mullen, E.: Magnesium isotopes as a tracer of crustal materials in volcanic arc magmas in the northern Cascade Arc, *Frontiers in Earth Science*, 6, 21, 2018.
- Carter, L. B., Skora, S., Blundy, J. D., De Hoog, J. C. M., and Elliott, T.: An Experimental Study of Trace Element Fluxes from Subducted Oceanic Crust, *J. Petrol.*, 56, 1585-1605, 10.1093/petrology/egv046, 2015.
- Chacko, T., Cole, D. R., and Horita, J.: Equilibrium oxygen, hydrogen and carbon isotope fractionation factors applicable to  
345 geologic systems, *Reviews in mineralogy and geochemistry*, 43, 1-81, 2001.
- Chemia, Z., Dolejš, D., and Steinle-Neumann, G.: Thermal effects of variable material properties and metamorphic reactions in a three-component subducting slab, *Journal of Geophysical Research: Solid Earth*, 120, 6823-6845, 2015.
- Chen, B.-B., Li, S.-L., von Strandmann, P. A. P., Wilson, D. J., Zhong, J., Ma, T.-T., Sun, J., and Liu, C.-Q.: Behaviour of Sr, Ca, and Mg isotopes under variable hydrological conditions in high-relief large river systems, *Geochim. Cosmochim. Acta.*,  
350 343, 142-160, 2023a.
- Chen, L., Li, D.-Y., Deng, J.-H., Li, S.-Z., Somerville, I., Chen, Y.-X., Zhao, Z.-F., An, W., and Li, X.-H.: Fe–Mg Isotopes Trace the Mechanism of Crustal Recycling and Arc Magmatic Processes in the Neo-Tethys Subduction Zone, *Journal of Geophysical Research: Solid Earth*, 128, e2023JB026778, 2023b.
- Codillo, E., Klein, F., Dragovic, B., Marschall, H., Baxter, E., Scambelluri, M., and Schwarzenbach, E.: Fluid-Mediated Mass  
355 Transfer Between Mafic and Ultramafic Rocks in Subduction Zones, *Geochemistry, Geophysics, Geosystems*, 23, e2021GC010206, 2022.
- D’Antonio, M. and Kristensen, M.: Serpentine and brucite of ultramafic clasts from the South Chamorro Seamount (Ocean Drilling Program Leg 195, Site 1200): inferences for the serpentinization of the Mariana forearc mantle, *Mineralogical Magazine*, 68, 887-904, 2004.
- 360 Debret, B., Albers, E., Walter, B., Price, R., Barnes, J., Beunon, H., Facq, S., Gillikin, D. P., Mattielli, N., and Williams, H.: Shallow forearc mantle dynamics and geochemistry: new insights from IODP Expedition 366, *Lithos*, 326, 230-245, 2019.
- Ferrand, T. P.: Seismicity and mineral destabilizations in the subducting mantle up to 6 GPa, 200 km depth, *Lithos*, 334-335, 205-230, 2019.
- Foster, G. L., Pogge von Strandmann, P. A. E., and Rae, J. W. B.: Boron and magnesium isotopic composition of seawater,  
365 *Geochem. Geophys. Geosyst.*, 11, Artn Q08015, 2010.
- Fryer, P., Pearce, J., and Stokking, L.: 36. A synthesis of Leg 125 drilling of serpentine seamounts on the Mariana and Izu–Bonin forearcs, *Proceedings of the Ocean Drilling Program, Scientific Results*, 593-614,



- Galy, A., Yoffe, O., Janney, P. E., Williams, R. W., Cloquet, C., Alard, O., Halicz, L., Wadhwa, M., Hutcheon, I. D., and Ramon, E.: Magnesium isotope heterogeneity of the isotopic standard SRM980 and new reference materials for magnesium-  
370 isotope-ratio measurements, *J. Anal. At. Spectrom.*, 18, 1352-1356, 2003.
- Gao, C., Cao, X., Liu, Q., Yang, Y., Zhang, S., He, Y., Tang, M., and Liu, Y.: Theoretical calculation of equilibrium Mg isotope fractionations between minerals and aqueous solutions, *Chem. Geol.*, 488, 62-75, 2018.
- Hacker, B. R.: H<sub>2</sub>O subduction beyond arcs, *Geochemistry, Geophysics, Geosystems*, 9, <https://doi.org/10.1029/2007GC001707>, 2008.
- 375 Hermann, J. and Rubatto, D.: Accessory phase control on the trace element signature of sediment melts in subduction zones, *Chem. Geol.*, 265, 512-526, Doi 10.1016/J.Chemgeo.2009.05.018, 2009.
- Hin, R. C., Coath, C. D., Carter, P. J., Nimmo, F., Lai, Y.-J., Pogge von Strandmann, P. A. E., Willbold, M., Leinhardt, Z. M., Walter, M. J., and Elliott, T.: Magnesium isotope evidence that accretional vapour loss shapes planetary compositions, *Nature*, 549, 511, 10.1038/nature23899, 2017.
- 380 Hostetler, P. B., Coleman, R. G., Mumpton, F. A., and Evans, B. W.: Brucite in alpine serpentinites, *American Mineralogist*, 51, 75-98, 1966.
- Humphris, S. E. and Klein, F.: Progress in deciphering the controls on the geochemistry of fluids in seafloor hydrothermal systems, *Annual review of marine science*, 10, 315-343, 2018.
- Johannes, W.: Experimental investigation of the reaction forsterite+ H<sub>2</sub>O $\rightleftharpoons$  serpentine+ brucite, *Contrib. Mineral. Petrol.*, 19, 309-315, 1968.
- 385 Kahl, W.-A., Jöns, N., Bach, W., Klein, F., and Alt, J. C.: Ultramafic clasts from the South Chamorro serpentine mud volcano reveal a polyphase serpentinization history of the Mariana forearc mantle, *Lithos*, 227, 1-20, 2015.
- Kempf, E. D., Hermann, J., Reusser, E., Baumgartner, L. P., and Lanari, P.: The role of the antigorite+ brucite to olivine reaction in subducted serpentinites (Zermatt, Switzerland), *Swiss J Geosci*, 113, 1-36, 2020.
- 390 Kessel, R., Schmidt, M. W., Ulmer, P., and Pettke, T.: Trace element signature of subduction-zone fluids, melts and supercritical liquids at 120-180 km depth, *Nature*, 437, 724-727, 10.1038/nature03971, 2005.
- Klein, F. and Le Roux, V.: Quantifying the volume increase and chemical exchange during serpentinization, *Geology*, 48, 552-556, 10.1130/g47289.1, 2020.
- Klein, F., Humphris, S., and Bach, W.: Brucite formation and dissolution in oceanic serpentinite, *Geochemical Perspectives Letters*, 16, 1-5, 2020.
- 395 Klein, F., Marschall, H. R., Bowring, S. A., Humphris, S. E., and Horning, G.: Mid-ocean ridge serpentinite in the Puerto Rico Trench: From seafloor spreading to subduction, *J. Petrol.*, 58, 1729-1754, 2017.
- Klein, F., Bach, W., Humphris, S. E., Kahl, W.-A., Jöns, N., Moskowicz, B., and Berquó, T.: Magnetite in seafloor serpentinite - Some like it hot, *Geology*, 42, 135-138, 2014.
- 400 Klemme, S., Blundy, J. D., and Wood, B. J.: Experimental constraints on major and trace element partitioning during partial melting of eclogite, *Geochim. Cosmochim. Acta.*, 66, 3109-3123, Doi 10.1016/S0016-7037(02)00859-1, 2002.



- Konrad-Schmolke, M., Halama, R., and Manea, V. C.: Slab mantle dehydrates beneath Kamchatka—yet recycles water into the deep mantle, *Geochemistry, Geophysics, Geosystems*, 17, 2987-3007, 2016.
- Lazar, C.: Using silica activity to model redox-dependent fluid compositions in serpentinites from 100 to 700 C and from 1 to 405 20 kbar, *J. Petrol.*, 61, ega101, 2020.
- Li, S.-G., Yang, W., Ke, S., Meng, X., Tian, H., Xu, L., He, Y., Huang, J., Wang, X.-C., and Xia, Q.: Deep carbon cycles constrained by a large-scale mantle Mg isotope anomaly in eastern China, *National Science Review*, 4, 111-120, 2017.
- Li, X., Li, S., Zhang, Z., Zhong, Y., and Li, D.: Magnesium isotopic fractionation during post-serpentinization alteration: Implications for arc and oceanic Mg cycles, *Chem. Geol.*, 121866, 2023.
- 410 Li, Y. H. and Schoonmaker, J. E.: Chemical Composition and Mineralogy of Marine Sediments, in: *Treatise on geochemistry*, edited by: Turekian, K. K., and Holland, H. D., 1-32, 2014.
- Liu, P.-P., Teng, F.-Z., Dick, H. J., Zhou, M.-F., and Chung, S.-L.: Magnesium isotopic composition of the oceanic mantle and oceanic Mg cycling, *Geochim. Cosmochim. Acta.*, 206, 151-165, 2017.
- Manning, C. E.: The chemistry of subduction-zone fluids, *Earth Planet. Sci. Lett.*, 223, 1-16, 2004.
- 415 McDonough, W. F. and Sun, S.-s.: The composition of the Earth, *Chem. Geol.*, 120, 223-253, 1995.
- Mével, C. and Staudoumi, C.: Hydrothermal alteration of the upper mantle section at Hess Deep, *Proc. ODP Sci. Results*, 147, 293-309, 1996.
- Nielsen, S. G., Klein, F., Kading, T., Blusztajn, J., and Wickham, K.: Thallium as a Tracer of Fluid-Rock Interaction in the Shallow Mariana Forearc, *Earth Planet. Sci. Lett.*, 430, 416-426, 2015.
- 420 Paulick, H., Bach, W., Godard, M., De Hoog, J. C. M., Suhr, G., and Harvey, J.: Geochemistry of abyssal peridotites (Mid-Atlantic Ridge, 15°20'N, ODP Leg 209): Implications for fluid/rock interaction in slow spreading environments, *Chem. Geol.*, 234, 179-210, 2006.
- Pogge von Strandmann, P. A. E., Dohmen, R., Marschall, H. R., Schumacher, J. C., and Elliott, T.: Extreme Magnesium Isotope Fractionation at Outcrop Scale Records the Mechanism and Rate at which Reaction Fronts Advance, *J. Petrol.*, 56, 33-425 58, 10.1093/petrology/egu070, 2015.
- Pogge von Strandmann, P. A. E., Elliott, T., Marschall, H. R., Coath, C., Lai, Y.-J., Jeffcoate, A. B., and Ionov, D. A.: Variations of Li and Mg isotope ratios in bulk chondrites and mantle xenoliths, *Geochim. Cosmochim. Acta.*, 75, 5247-5268, 2011.
- Ranero, C. R., Phipps Morgan, J., McIntosh, K., and Reichert, C.: Bending-related faulting and mantle serpentinization at the 430 Middle America trench, *Nature*, 425, 367-373, 2003.
- Rudnick, R. L. and Gao, S.: Composition of the Continental Crust, in: *Treatise on Geochemistry*, edited by: Holland, H. D., and Turekian, K. K., Pergamon, Oxford, 1-64, 2003.
- Rüpke, L. H., Morgan, J. P., Hort, M., and Connolly, J. A. D.: Serpentine and the subduction zone water cycle, *Earth Planet. Sci. Lett.*, 223, 17-34, 10.1016/j.epsl.2004.04.018, 2004.



- 435 Schmidt, M. W. and Poli, S.: Experimentally based water budgets for dehydrating slabs and consequences for arc magma generation, *Earth Planet. Sci. Lett.*, 163, 361-379, 1998.
- Shu, Y., Nielsen, S. G., Le Roux, V., Wörner, G., Blusztajn, J., and Auro, M.: Sources of dehydration fluids underneath the Kamchatka arc, *Nat Commun*, 13, 4467, 10.1038/s41467-022-32211-5, 2022.
- Spandler, C., Mavrogenes, J., and Hermann, J.: Experimental constraints on element mobility from subducted sediments using  
440 high-P synthetic fluid/melt inclusions, *Chem. Geol.*, 239, 228-249, 2007.
- Stalder, R., Ulmer, P., Thompson, A., and Günther, D.: High pressure fluids in the system MgO–SiO<sub>2</sub>–H<sub>2</sub>O under upper mantle conditions, *Contrib. Mineral. Petrol.*, 140, 607-618, 2001.
- Tatsumi, Y.: Formation of the Volcanic Front in Subduction Zones, *Geophys. Res. Lett.*, 13, 717-720, 1986.
- Teng, F.-Z., Hu, Y., and Chauvel, C.: Magnesium isotope geochemistry in arc volcanism, *Proceedings of the National  
445 Academy of Sciences*, 113, 7082-7087, 2016.
- Teng, F. Z., Li, W. Y., Ke, S., Marty, B., Dauphas, N., Huang, S. C., Wu, F. Y., and Pourmand, A.: Magnesium isotopic composition of the Earth and chondrites, *Geochim. Cosmochim. Acta.*, 74, 4150-4166, 10.1016/j.gca.2010.04.019, 2010.
- Ulmer, P. and Trommsdorff, V.: Serpentine Stability to Mantle Depths and Subduction-Related Magmatism, *Science*, 268, 858-861, 1995.
- 450 van Keken, P. E. and Wilson, C. R.: An introductory review of the thermal structure of subduction zones: III—Comparison between models and observations, *Prog Earth Planet Sc*, 10, 57, 2023.
- Wang, W., Zhou, C., Liu, Y., Wu, Z., and Huang, F.: Equilibrium Mg isotope fractionation among aqueous Mg<sup>2+</sup>, carbonates, brucite and lizardite: Insights from first-principles molecular dynamics simulations, *Geochim. Cosmochim. Acta.*, 250, 117-129, 2019.
- 455 Wang, Y., Lu, W., Costa, K. M., and Nielsen, S. G.: Beyond anoxia: Exploring sedimentary thallium isotopes in paleo-redox reconstructions from a new core top collection, *Geochim. Cosmochim. Acta.*, 333, 347-361, 2022.
- Wang, Y., Deng, J., Liao, R., Chen, L., Li, D., Liu, H., and Sun, W.: Magnesium isotopic composition of the Mariana forearc serpentinite: Implications for Mg isotopic composition of the mantle wedge and Mg isotopic fractionation during mantle wedge serpentinitization, *Chem. Geol.*, 624, 121428, 2023.
- 460 Wimpenny, J., Colla, C. A., Yin, Q.-Z., Rustad, J. R., and Casey, W. H.: Investigating the behaviour of Mg isotopes during the formation of clay minerals, *Geochim. Cosmochim. Acta.*, 128, 178-194, 2014.
- Wimpenny, J., Gíslason, S. R., James, R. H., Gannoun, A., Von Strandmann, P. A. P., and Burton, K. W.: The behaviour of Li and Mg isotopes during primary phase dissolution and secondary mineral formation in basalt, *Geochim. Cosmochim. Acta.*, 74, 5259-5279, 2010.
- 465 Wunder, B. and Schreyer, W.: Antigorite: High-pressure stability in the system MgO–SiO<sub>2</sub>–H<sub>2</sub>O (MSH), *Lithos*, 41, 213-227, 1997.
- Zhao, M.-S., Chen, Y.-X., Xiong, J.-W., Zheng, Y.-F., Zha, X.-P., and Huang, F.: Element mobility and Mg isotope fractionation during peridotite serpentinitization, *Geochim. Cosmochim. Acta.*, 340, 21-37, 2023.