

We thank the reviewer for their comments. We have made the suggested corrections to the text.

Line 131 (now line 136): Fixed reference authorship

Line 311 (now line 328): Made recommended change to text

We thank the reviewer for their comments. We have made many of the suggested corrections to the text, all points are discussed below. When referring to line numbers below “line X” refers to lines in the original document while “new line Y” refers to line numbers in the updated document.

General points:

“The author focus on specific LT and averages, but it would be more meaningful to quantify the change in day-to-day variability between the two cases (The authors mention several times the importance of day-to-day variability). While the text mentions day-to-day variability the study itself focuses on variability in general during the specific time periods. The study would gain if ability to capture the day-to-day variability would be added.”

- We agree that this would be an interesting study, but is beyond the scope of the current paper.

“If the authors want to demonstrate the value of adding middle atmospheric observations to better specify the ionospheric variability, it is surprising that the focus is on 10 LT. The authors should either motivate it or add later local times especially in the afternoon.”

- We focused on two different local times, 10:00 LT and 14:00 LT. These two times were chosen because of the observed changes in the ionosphere. Motivation for these times can be found in the new manuscript on lines XXX and XXX.

“The condition during the time period needs to be described in one short section at the beginning (inc. F10.7, PW evolution & SSW peak, geomagnetic activity). Would the results be different without the change in PW activity. How general is the result?”

Wasn't there some minor geomagnetic activity as well during the time period?"

- This was included in the start of Section 3, and is now in the first paragraph of Section 3. The atmospheric behavior for the time periods used in this study is also discussed in detail in the paper referenced at the end of this section.

"Please make the longitudinal range the same in the plots, so that 0 deg is in the same location e.g. Fig 1 and Fig 3."

- The longitude ranges differ in the figures so that the longitudes of interest can be easily observed. While we appreciate that it is annoying to have figures with different longitude ranges, we felt it was more important to chose ranges that ensured the selected longitudes of interest were not split across the edge of the figure.

Specific comments:

Line 12/new line 12-13: This line was changed to "variability induced by wind-dynamo coupling through electric conductivity and ion-neutral interactions in the upper thermosphere." to clarify the intention of the text.

Line 20/new line 20: Changed to "the short-term variability in the E-region"

Lines 27-28/new lines 27-28: Added "with respect to the long-term mean (calculated using data from 1967-1989)"

Line 35/new line 36: Added "When studied at a fixed local time (LT)," to provide the specific context for this sentence.

Line 52/new line 52: changed "forecasting skills" to "predictions"

as suggested.

Line 54/new line 54: clarified by adding: “global root mean square error in zonal wind”

Line 62/new line 62: made suggested change.

Line 64/new line 64: added “during boreal winter”

Line 66/new line 66: CITS was first defined in the previous paragraph (new line 48).

Line 90-92/new line 92-93: clarified by adding a sentence: “Due to the sparseness of observations, especially in the critical UMLT, analysis fields at a cadence higher than 6 hours are indistinguishable from intermediate forecasts, and we augmented the 6-hour analysis with 3-hourly forecasts; this approach has also been used in (Sassi et al. 2020).”

Line 99/new line 101: This would be a different study that would require the use of WACCMX+DART. While this alternative approach would also likely be interesting, our method is more adaptable to a mechanistic study like the one presented here. Using WACCMX+DART would not be mechanistic (allowing us to easily see the theoretical effect on the ionosphere driven by MA data assimilation), but an interactive study that makes it more difficult to isolate the driving mechanisms.

Line 105/new line 107: This is not possible to do because of the model infrastructure. NAVGEM uses a spectral triangular formulation (hence, T119) and WACCM is a grid-point model. Attempting to use the same formulation of units and grid resolution for both models would be incorrect in their implementations.

Line 114/new lines 116-118: I am not sure what kind of “zonal convection” the reviewer is referring to. The SAMI3 description was updated to add: “...a two-dimensional model of the ionosphere that handles plasma dynamics and chemical evolution along magnetic field lines (varying in latitude and altitude). SAMI3 extends SAMI2 by adding the longitudinal dimension, which includes zonal transport.” If the reviewer is referring to high-latitude plasma convection, then this is handled by driving SAMI3 with Weimer.

Line 122 /new lines 125-128: provided a more thorough explanation of modified apex coordinates.

Line 125: Further details about the standard SAMI3 set up are available in the references.

Line 129/new line 133: The software allows for different way to couple the ionosphere to the thermosphere. For this particular study, we are using a one-way coupled configuration for practical reasons, but the software developed at NRL is also capable of supporting two-way coupling. We changed the wording to say: “The software infrastructure that extends SAMI3 to allow either one-way or two-way coupling with the atmosphere”. We also updated the wording in the subsequent paragraph to make it clear that we are using a one-way coupled set-up, and why we chose to do so.

Line 139/new line 143-149: reworded introduction entirely, taking this into account.

Caption Fig 4: Adjusted wording to say: “along a SAMI3 slice with a longitude of  $\sim 90^\circ$  E at the geographic equator”

Line 156/new line 171: changed wording to “more complex longitudinal structure”

Line 164: It may be possible that there are changes in the phases in the migrating tides that are affecting the ionosphere. However, our approach is to examine the impacts in a local time framework. This makes the phase changes of the migrating tides unimportant, since they can't be observed in this frame of reference.

Figure 1: 10LT and 16LT are used as representative of a complex behavior and exemplify the point of our study: lack of MA observations impacts the ions distribution and transport. The reviewer is correct in pointing out that more advanced studies can be done, but this is a simple exemplification of many other facets. It is also important to examine the impact of MA observations at different local times. While this selection of case studies is limited, we think it is important to focus on more than just one LT region.

Line 169/new lines 186-187: changed to: “Once the vertical ExB drifts lift the plasma to magnetic field lines with higher apex altitudes, the change in the plasma pressure gradient along these longer field lines will cause the ions to diffuse downwards along the magnetic field lines to lower altitudes and higher latitudes”

Line 171: We also look at the ExB drift directly. See Figure 3 and Line 200 (new line 200-210).

Line 180: The reviewer may be correct, but for this study it doesn't matter: the goal of the study is not to investigate the detailed processes operating at this time; it is a sensitivity study to demonstrate that the lack of MA observations is impactful on the ionospheric properties. To ensure this is clearer, we added “at this time and location” to the end of this sentence.

Sect 3.2: This wording refers to the presence or absence of a SSW. This is made more clear now by rewriting the introduction to the case studies in the first paragraph of Section 3.

Line 254 and elsewhere: Replaced “day-to-day” with “intra-day” throughout the text.

Line 255/new line 271: It would be incorrect to remove one parenthesis, since two are needed for closure.

Line 274/new line 290: The SSW is first mentioned on Line 151. The new rewording now brings it up in the first paragraph of Section 3.

Line 276/new line 292: Both an SSW period and a period without SSW were examined to see if the influence of MA observations would be important during extreme times, normal times, or not be important at all. This is now clarified in the first paragraph of Section 3.

Line 281/new line 296: There may be other things, but the MA observations specifically affected the non-migrating solar tides, as is possible due to the nature of this controlled mechanistic experiment. This is why we are talking about their impacts in this paragraph.

Line 284/new line 300: Added reasoning behind discussing these wave patterns in particular to the text.

Figure 9: We agree with the reviewer, and the text states that the NOOBS example shows an instance without the EIA present. The figure caption also says that in the absence of an EIA, the start will mark the peak density in this region. The figure caption

has been further clarified to avoid confusion.

Table 1 and 2: Adjusted the wording in new lines 310-318 more clearly explain that we are examining the low-latitude VTEC distribution by grouping data into three types of EIA configurations: no EIA, an EIA with a higher northern peak, and an EIA with a higher southern peak.

Line 313/new line 330: Added text relating the E-region conductivity to the transport caused by the ExB drift.

Line 320: :)

Discussion: We decided not to do this, because we are specifically interested in the way the MA drives the F-region ionosphere. Because the MA is below the ionosphere, we do not expect two way coupling to change the conclusions we found here.



# On the importance of middle atmosphere observations on ionospheric dynamics using WACCM-X and SAMI3

Fabrizio Sassi<sup>1,2</sup>, Angeline G. Burrell<sup>1</sup>, Sarah E. McDonald<sup>1</sup>, Jennifer Tate<sup>3</sup>, and John P. McCormack<sup>1,4</sup>

<sup>1</sup>Naval Research Laboratory - Space Science Division, 4555 Overlook Ave SW, Washington DC 20375, USA

<sup>2</sup>Heliophysics Division, Goddard Space Flight Center, 8800 Greenbelt Rd., Greenbelt MD 20771, USA

<sup>3</sup>Computational Physics Inc., 8001 Braddock Rd, Suite 210, Springfield VA 22151, USA

<sup>4</sup>Heliophysics Division, Science Mission Directorate, 300 Hidden Figures Way SW, Washington DC 20546, USA

**Correspondence:** Angeline G. Burrell (angeline.burrell@nrl.navy.mil)

**Abstract.** Recent advances in atmospheric observations and modelling have enabled the investigation of thermosphere-ionosphere interactions as a whole atmosphere problem. This study examines how dynamical variability in the middle atmosphere (MA) affects ~~day-to-day~~ intra-day changes in the thermosphere and ionosphere. Specifically, this study investigates ionosphere-thermosphere interactions during different time periods of January 2013 using the Specified Dynamics Whole Atmosphere Community Climate Model, eXtended version (SD-WACCM-X) coupled to the Naval Research Laboratory (NRL) ionosphere of the Sami3 is Another Model of the Ionosphere (SAMI3) model. To represent the weather of the day, the coupled thermosphere-ionosphere system is nudged below 90 km toward the atmospheric specifications provided by the Navy Global Environmental Model for High-Altitude (NAVEM-HA). Hindcast simulations during January 2013 are carried out with the full data set of observations normally assimilated by NAVEM-HA, and with a degraded dataset where observations above 40 km are not assimilated. Ionospheric regions with statistically significant changes are identified using key ionospheric properties, including the electron density, peak electron density, and height of the peak electron density. Ionospheric changes show a spatial structure that illustrates the impact of two different types of coupling between the thermosphere and the ionosphere: variability induced by wind-dynamo coupling through electric conductivity and ion-neutral interactions in the upper thermosphere. The two simulations presented in this study show that changing the state of the MA affects ionosphere-thermosphere coupling through changes in the behavior and amplitude of non-migrating tides, resulting in improved key ionospheric specifications.

## 1 Introduction

The development of whole atmosphere models (ground to exobase) in the last decade along with the availability of high-altitude atmospheric data assimilation has brought the geospace community closer to important understandings of the prediction of short-term variability ( $\leq 10$  days) in the lower E-region. When the Sun is quiescent, a large fraction of ~~this~~ the short-term variability in the E-region is driven by lower atmosphere weather from below. Because of wind-dynamo coupling in this region, neutral variability affects the state of the ionosphere both locally and at higher altitudes.

The impact of atmospheric weather on the behavior of the coupled thermosphere-ionosphere system is a topic that has been actively investigated for nearly a century. From small-scale gravity waves (Hines, 1960) to planetary-scale waves (Forbes and

Zhang, 1997), meteorological weather impacts the ionosphere either through electrodynamics (e.g.,  $\mathbf{E} \times \mathbf{B}$  drifts) or collisional interactions whose impacts greatly affect ion transport along field lines. Forbes et al. (2000) show that the fraction of ionospheric variability (obtained from the critical frequency at the F-peak,  $f_oF2$ ) due to meteorological conditions depends on the period of the meteorological forcing itself, and varies between 20% and 35% with respect to the long-term mean (calculated using data from 1967–1989) during solar minimum conditions; modeling studies tend to estimate a higher fraction for the meteorological contribution, approaching 50% (e.g. Liu et al., 2013). The role of migrating and non-migrating tides in determining the variability and structure of the ionosphere on many time scales, from days to seasons, is prominent and well established (Hagan et al., 2007; Liu, 2016; Sassi et al., 2019, and references therein). Migrating solar tides are associated with a modulation of the daily variability of vertical ion drifts at the geomagnetic equator (Millward et al., 2001; Fang et al., 2013). In particular, the migrating semidiurnal solar tide has been associated with a shift of meridional ion drifts (which are directed vertically at the geomagnetic equator), and consequently of the peak electron density, to earlier local times during days immediately following a Sudden Stratospheric Warming (SSW) (e.g., Goncharenko et al., 2010; McDonald et al., 2015; Fuller-Rowell et al., 2017). ~~Non-migrating~~ When studied at a fixed local time (LT), non-migrating tides have instead been associated with the zonal structure of the ionosphere (Forbes et al., 2008; Immel et al., 2006) and have been found to be very sensitive to the background meteorological conditions of the lower atmosphere (McDonald et al., 2018; Sassi et al., 2020). In addition to solar tides, lunar tides are known to be impactful on the field-aligned neutral winds that help shape the structure of the F-peak ionosphere (Pedatella and Maute, 2015). Furthermore, a statistical analysis that aggregates the response of the Equatorial Ionization Anomaly (EIA) over many SSW (Wu et al., 2021) shows that the timing of the modulation of the EIA following an SSW is affected by the lunar phase.

Such theoretical progress in understanding the coupling between the thermosphere and the ionosphere has, in part, been possible because of momentous advances in the development of whole atmosphere numerical models during the last decades that also include electrodynamic interactions (e.g., Liu et al., 2010, 2018). At the same time, the development of these modeling capabilities has benefited from the availability and utilization of observations well into the upper atmosphere and the extension of forecast/assimilation systems that provide specifications in the upper atmosphere (Eckermann et al., 2009; Wang et al., 2011; McCormack et al., 2017; Eckermann et al., 2018). Thus, investigations of the Coupled Ionosphere Thermosphere System (CITS) have been guided and validated by observations (Jin et al., 2012), both through nudging techniques (e.g., Sassi et al., 2013) and full data assimilation (Wang et al., 2011; Pedatella et al., 2018). In most cases, the need for upper atmospheric observations is not limited to merely validating forecast and hindcast simulations, but rather extends to fundamental advances in theoretical understanding of whole atmosphere interactions (Liu, 2016; Sassi et al., 2019, for reviews). Observations are critical to evaluate and improve ~~forecast-skill~~ predictions in the whole atmosphere: Liu et al. (2009) demonstrated that the troposphere plays a critical role in controlling error growth at higher altitudes. Pedatella et al. (2013) used a data assimilation system to reduce the ~~error-growth~~ global root mean square error in zonal wind up to 40% in the Upper Mesosphere and Lower Thermosphere (UMLT) and in a later study, Pedatella et al. (2019) used a whole atmosphere model with a data assimilation system to demonstrate that the error growth in the lower thermosphere saturates within 5 days.

Observations of the middle atmosphere (MA), which consists of both the stratosphere and mesosphere, are primarily obtained from instruments carried onboard space-borne platforms. Unfortunately, many research platforms have surpassed their expected lifetime, such as the Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) instrument onboard the NASA Thermosphere Ionosphere Mesosphere Energetic Dynamics (TIMED) satellite, while critical operational platforms are not being replaced (e.g., the Defense Meteorological Satellite Program (DMSP)) (e.g., Erwin and Berger, 2021). As noted by Sassi et al. (2020), the potential lack of observations in the MA has significant implications ~~for the day-to-day to capture~~ intra-day variability of neutral winds in the lower thermosphere. In particular, the amplitude of some non-migrating solar tides and of traveling planetary scale waves can change by a factor of two during boreal winter. It stands to reason that such prominent changes of the thermospheric weather will have consequences for the structure and variability of the ionosphere. To the best of our knowledge the fundamental question of how numerical simulations of physical properties of the CITS are impacted by data loss in the middle atmosphere remains elusive and poorly described.

This study seeks to answer the fundamental question of whether the loss of MA observations is potentially consequential for predictions of thermosphere-ionosphere coupling. In Section 2, the different models used in this study are presented, and details about the run configurations are provided. Section 3 presents the results of the model runs, Section 4 presents a discussion of the results and Section 5 the conclusions.

## 2 Models

Three models are used in this study: the Navy Global Environmental Model (NAVGEN-HA), the Whole Atmosphere Community Climate Model, eXtended version (WACCM-X), and the Sami3 is Another Model of the Ionosphere (SAMI3). Details about these models and how they have been integrated to effectively model the CITS are presented below.

### 2.1 NAVGEM-HA

The Navy Global Environmental Model (NAVGEN) is the Navy's operational global numerical weather prediction (NWP) system. NAVGEN couples a semi-implicit semi-Lagrangian atmospheric model with a hybrid four-dimensional variational (4DVAR) data assimilation system that is based on the NRL Atmospheric Variational Data Assimilation System–Accelerated Representer (NAVDAS-AR) (Kuhl et al., 2013). This study uses a version of NAVGEN developed especially for High Altitude research (NAVGEN-HA). The standard version of NAVGEN only covers 0-80 km, but NAVGEN-HA has an extended altitude range that reaches 116 km and includes additional physical parameterizations for high altitude processes such as ozone and water vapor photochemistry, as well as sub-grid scale gravity wave drag. For a detailed description, see McCormack et al. (2017) and Eckermann et al. (2018). In addition to the full suite of operational observations and space-based sensor data, NAVGEN-HA assimilates temperature profiles between 30 and 100 km altitude (provided by the TIMED SABER instrument), temperature, ozone, and water vapor profiles (provided by the Aura Microwave Limb Sounder), and Special Sensor Microwave Imager/Sounder (SSMIS) Upper Atmospheric Sounding (UAS) channel radiances (Hoppel et al., 2013). The standard NAVGEN-HA configuration has T119 triangular spectral resolution (e.g., Laprise, 1992) ( $1^\circ$  latitude/longitude grid

90 spacing), with 74 vertical levels (L74) having an effective spacing of  $\sim 2$  km in the stratosphere,  $\sim 3$  km in the mesosphere and  $\sim 4$  km in the lower thermosphere. The present study uses the NAVGEM-HA global synoptic analyses of key atmospheric state variables (including temperature, winds, geopotential height, ozone, water vapor, vorticity, and divergence) every 6 hours. These atmospheric state variables have been augmented with 3-hour forecasts to provide 3-hourly output. [Due to the sparseness of observations, especially in the critical UMLT, analysis fields at a cadence higher than 6 hours are indistinguishable from](#)  
95 [intermediate forecasts, and we augmented the 6-hour analysis with 3-hourly forecasts; this approach has also been used in \(Sassi et al., 2020\).](#) This greater output frequency is needed to resolve sub-diurnal variability that is crucially important to represent the ~~day-to-day~~ [intra-day](#) variability in the upper atmosphere and ionosphere.

Two sets of NAVGEM-HA atmospheric specifications have been generated for January-February 2013. The first is a reference set with all observations from the ground to 100 km; this hybrid data assimilation with MA observations is referred to  
100 as *hybma* (*hybrid MA*). A second set of NAVGEM-HA simulations have been generated excluding all observations above 40 km; this theory-based set is referred to as *noobs* (*no MA observations*). Temperature, winds and surface pressure generated by *hybma* and *noobs* NAVGEM-HA are used to nudge the atmosphere of SD-WACCM-X (see Sect. 2.2). This is exactly the same dataset of atmospheric specifications that has been used in the Sassi et al. (2020) study.

## 2.2 SD-WACCM-X

105 To determine the neutral atmospheric response to the weather of the day from the lower atmosphere throughout the thermosphere, we use the Whole Atmosphere Community Climate Model, eXtended version (WACCM-X; Liu et al. (2010)). WACCM-X horizontal resolution is  $\sim 2^\circ$  in latitude and longitude. The vertical resolution, while variable in the lower atmosphere, is set to be a quarter of the local pressure scale height in the thermosphere, with 108 vertical levels from the ground to about 500 km. To aid in the investigation of whole atmosphere connections, WACCM-X can be configured to use atmospheric  
110 specifications to constrain its meteorology (winds and temperature) from the ground to any altitude; this model configuration is referred to as Specified Dynamics (SD). The reader is referred to Sassi et al. (2020) for an exhaustive discussion of the day-to-day variability of these simulations.

As in Sassi et al. (2020), the *hybma* and *noobs* atmospheric specifications generated by NAVGEM-HA are used to nudge the SD-WACCM-X meteorology. In the remainder of this article, except when stated otherwise, *hybma* and *noobs* are used as  
115 abbreviations to refer to the SD-WACCM-X simulations with the corresponding NAVGEM-HA nudging meteorology.

## 2.3 SAMI3

SAMI3 is ~~a longitudinal~~ [an](#) extension of SAMI2 (Huba et al., 2000), a two-dimensional model of the ionosphere [that handles plasma dynamics and chemical evolution along magnetic field lines \(varying in latitude and altitude\).](#) SAMI3 [extends SAMI2 by adding the longitudinal dimension, which includes zonal transport.](#) SAMI3 models the plasma and chemical evolution of  
120 seven ion species ( $H^+$ ,  $He^+$ ,  $N^+$ ,  $O^+$ ,  $N_2^+$ ,  $NO^+$  and  $O_2^+$ ) in the altitude range extending from 70 km to  $\sim 8$  Earth radii and magnetic latitudes up to  $\pm 88^\circ$ . Unlike other ionospheric models, SAMI3 solves the ion continuity and momentum equations for all seven of the ion species, and solves the temperature equations for the electrons and three ion species ( $H^+$ ,  $He^+$  and

O<sup>+</sup>). Ion inertia is included in the ion momentum equation for motion along the geomagnetic field. SAMI3 is the only global ionosphere code that models full plasma transport for all of these ion species, including the molecular ions. Another unique feature of SAMI3 is its non-uniform, nonorthogonal fixed grid. The original version of SAMI3 used a tilted dipole model of the geomagnetic field. However, SAMI3 has been upgraded to allow use of Magnetic Apex coordinates (Richmond, 1995), ~~so that an accurate representation of the geomagnetic field~~ (VanZandt et al., 1972; Richmond, 1995; Laundal and Richmond), which take into account both the ellipsoidal shape of the Earth and the secular variation of the Earth's main magnetic field - provided by the International Geomagnetic Reference Field (IGRF-13) is considered; Alken et al. (2021)) - when tracing the magnetic field lines. SAMI3 includes a potential solver to self-consistently solve for the electric fields. The potential equation is derived from current conservation in magnetic coordinates, using Ohm's law and including gravity-driven currents. The perpendicular electric field is used self consistently in SAMI3 to calculate the  $\mathbf{E} \times \mathbf{B}$  drifts driven by the neutral wind in the low- to mid-latitude ionosphere (Huba et al., 2010). At high latitudes SAMI3 uses an empirical model (e.g., Weimer, 1995, 2005), and the solar extreme ultraviolet (EUV) irradiance is determined from the EUVAC model, as described in (Solomon and Qian, 2005). The software infrastructure that extends SAMI3 ~~capability to be fully interactive to allow either one-way or two-way coupling~~ with the atmosphere consists of interpolation software of the Earth System Modeling Framework (ESMF, <https://earthsystemmodeling.org/>) and an extension of neutral field properties to the plasmasphere (McDonald et al., 2018).

In this study we use the SD-WACCM-X winds along with the NRLMSIS 2.0 (Emmert et al., 2021) temperature and composition to define SAMI3 neutral properties. We chose this ~~set-up one-way coupled set-up~~ (similar to the one used in McDonald et al. (2015)), because we want to isolate ~~as much as possible~~ the effects of neutral winds ~~from other factors; a~~. A more realistic and nonlinear system that includes the influences of meteorological winds, temperature and composition is bound to be far more complex and, making it challenging to isolate the effects of neutral winds from the MA on the ionosphere. The output cadence of the SAMI3 fields used for the analysis described in this study is 15 minutes.

### 3 Results

This section presents three ~~local time (LT) cases: two cases case studies: two~~ during a dynamically active week (1–10 January ~~2013~~) and one during a dynamically quiet week (21–30 January 2013). The primary difference between these two weeks is the presence of a SSW during the dynamically active week. We chose these two weeks to determine whether the influence of MA observations would be important only during times of extreme atmospheric forcing, also at times of normal atmospheric forcing, or not at all. The solar and geomagnetic forcing during both periods is similar, with the  $F_{10.7}$  (10.7 cm radio-flux) at a moderate level (typically below 110 solar flux units, with the exception of a large spike to 168 solar flux units on 1 January 2013) and  $K_p < 4$  (see [https://lasp.colorado.edu/lisird/data/noaa\\_radio\\_flux](https://lasp.colorado.edu/lisird/data/noaa_radio_flux) and [https://www-app3.gfz-potsdam.de/kp\\_index/Kp\\_ap\\_since\\_1932.txt](https://www-app3.gfz-potsdam.de/kp_index/Kp_ap_since_1932.txt)). The atmospheric dynamical behavior of January-February 2013 is discussed in (Sassi et al., 2020).

The two case studies during the dynamically active week examine different local time (LT) sectors: one in the morning (10:00 LT) and one another in the afternoon (16:00 LT) and an. The third case from the dynamically quiet week focuses on the afternoon (16:00 LT) case during a dynamically quiet time (21–30 January). We have examined other LTs (not shown).


which do not add more information to the two LTs studied here: 10:00 LT and 16:00 LT fully cover the representation of the ionospheric response paths described below.

Together, they illustrate different paths by which the neutral atmosphere may affect the ionosphere. The different paths will be examined for both *hybma* and *noobs* simulations (see Section 2.2). The first case (Section 3.1) is a morning scenario showing a longitudinally complex behavior with changes of both the lower thermospheric neutral wind and the F-peak winds that prominently affect the ionosphere through wind-dynamo coupling in the E-region causing electrodynamic interactions in the F-region (Heelis, 2004) and modulation of the EIA from inter-hemispheric ion transport at the F-peak, respectively. The second case (Section 3.2) demonstrates that the different specification of the MA state also leads to changes in the thermospheric winds in the F-region, which interact collisionally with the ions and alter the ion transport along the magnetic field lines (e.g., Rishbeth, 1972; Burrell and Heelis, 2012). The third case shows that the MA state affects the ionosphere and thermosphere during dynamically quiet times as well as dynamically active times.

### 3.1 Case 1: 10:00 LT - dynamically active

Figure 1 illustrates the  $N_m F_2$  at 10:00 LT obtained from averaging the 15-minute output for ten days between January 1 and January 10, 2013, i.e., covering the time period immediately prior and following the SSW of January 6, 2013. In each panel, the location of the magnetic equator is marked by a solid black line and the longitude sector of interest is marked by a dashed black line. Both model runs show that the structure of the  $N_m F_2$  is geomagnetically oriented, as the peak values at each longitude lie just north of the magnetic equator. In *hybma* (Figure 1a) the  $N_m F_2$  is broadly uniform north of the magnetic equator at about  $175 \times 10^4 - 200 \times 10^4 \text{ cm}^{-3}$ , with a relative maximum near  $80^\circ \text{E}$  just exceeding  $200 \times 10^4 \text{ cm}^{-3}$ . In contrast, the  $N_m F_2$  in *noobs* (Figure 1b) shows a more complex longitudinal structure north of the magnetic equator. The  $N_m F_2$  displays several peaks in longitude, all of about the same magnitude and just exceeding  $200 \times 10^4 \text{ cm}^{-3}$ . The longitudinally dependent changes are better illustrated by examining the difference between the model results. Figure 1c shows *noobs* minus *hybma*  $N_m F_2$ . The difference in peak density clearly shows that the largest effects introduced by removing MA observations through the *noobs* run at this LT are over the Pacific Ocean and over the Indian Ocean. At the magnetic equator the differences in the two longitudinal regions have opposite signs: a negative anomaly flanked by positive differences in the Pacific Ocean, contrariwise over the Indian Ocean. Overall, the magnitude of the differences are up to  $\pm 50 \times 10^4 \text{ cm}^{-3}$ . Such longitudinal structure at a fixed LT introduced by changes to the lower atmospheric forcing can only be explained by differences in amplitude of non-migrating solar tides (Forbes et al., 2006).

Focusing on the region over the Pacific Ocean near  $225^\circ \text{E}$ , this region shows statistically significant differences between the *noobs* and *hybma* runs: a decrease of  $N_m F_2$  at the magnetic equator is flanked by increases in the  $N_m F_2$  at latitudes corresponding to the peak of the EIA. The symmetry of this structure about the magnetic equator implies a change to the ‘fountain effect’ (e.g., Hanson and Moffett, 1966): the EIA is maintained through  $\mathbf{E} \times \mathbf{B}$  drifts that lift to higher altitudes ions created at or near the magnetic equator at lower altitudes. ~~Opposed by gravity, the~~ Once the vertical  $\mathbf{E} \times \mathbf{B}$  vertical lift runs out near the F-peak where the ions then drifts lift the plasma to magnetic field lines with higher apex altitudes, the change in the plasma pressure gradient along these longer field lines will cause the ions to diffuse downwards along the magnetic field



**Figure 1.**  $N_m F_2$  in units of  $10^4 \text{cm}^{-3}$  for all locations at 10:00 LT averaged for 1-10 January 2013: (a) *hybma*; (b) *noobs*; (c) difference *noobs* minus *hybma*. The oblique dash black lines highlight the position of SAMI3 slices whose longitude at the geographic equator are  $\sim 230^\circ$  and  $\sim 90^\circ$  and are discussed in text. The magnetic equator is illustrated by a solid black line across longitude. Green contours in (c) identify the locations where the anomalies are statistically significant at least at 95% confidence level using a t-test.

190 lines to lower altitudes and higher latitudes. To verify that a stronger fountain effect is taking place at this location, Figure 2 shows the electron density on nearby SAMI3 slice (longitude at the geographic equator  $\sim 230^\circ$ ) that cuts through the region of the largest differences (see oblique dash line in Figure 1). The electron density in *hybma* (Figure 2a) shows the typical arced behavior, with isopleths elevated near the magnetic equator with respect to the adjacent latitudes. The largest electron density is found just below 300 km northward of the geographic equator, with a peak value in excess of  $180 \times 10^4 \text{cm}^{-3}$ . This is consistent with the presence of both an active fountain effect and a northward field-aligned neutral wind. The difference plot (Figure 2b) shows reduced electron density by about  $15\text{--}30 \times 10^4 \text{cm}^{-3}$  at the magnetic equator, and increased values in excess of  $60 \times 10^4 \text{cm}^{-3}$  around  $5\text{--}10^\circ \text{N}$ . This structure is explained by an increased uplift of ions at the magnetic equator followed by transport along geomagnetic field lines due to the effects of neutral winds and gravity. In both simulations,  $\mathbf{E} \times \mathbf{B}$  meridional drifts at the magnetic equator lift ions into the upper F-region; in the case of the *noobs* simulation, a stronger uplift of ions followed by gravity and neutral wind transport results in a removal of charges from the magnetic equator and an accumulation at higher latitudes [at this time and location](#). Note that the accumulation of charges is greater in the northern hemisphere, which

195


200

(at this local time) is likely solely the result of forcing by the neutral wind at the  $h_m F_2$  along these magnetic field lines (Burrell et al., 2012).

If our interpretation of Figure 2 is correct, the *noobs* simulation ought to be showing a more prominent upward drift in the morning hours when compared to *hybma*. Figure 3 shows the 10-day average ((a,b) 1–10 January 2013) of the SAMI3 meridional  $\mathbf{E} \times \mathbf{B}$  drift at the magnetic equator as a function of longitude and LT. At longitudes around  $230^\circ\text{E}$  (which corresponds to the location of SAMI3 slice in Figure 2 ) and at 10:00 LT, Figure 3 shows larger  $\mathbf{E} \times \mathbf{B}$  drifts in *noobs* than in *hybma*. Since the resulting meridional  $\mathbf{E} \times \mathbf{B}$  drift is upward during the day, the electric field must be directed toward the East, resulting in an accumulation of positive (negative) charges at dusk (dawn). Thus, our model results illustrate that the simulation without MA observations (*noobs*) produces a stronger eastward electric field compared to the simulation with full MA observations, accumulating more charges at the dusk/dawn sectors. The stronger upward  $\mathbf{E} \times \mathbf{B}$  drifts lift the F-region plasma to higher altitudes where they then diffuse downwards and polewards along the magnetic field lines. Figure 3 results then confirm that the larger EIA seen in Figure 1c is due to a stronger fountain effect, which was created by unrealistic influences from the MA in the *noobs* experiment.

We now turn to the large differences over the Indian Ocean in Figure 1. The resulting difference field at this longitude (Figure 1c) shows an increase of  $N_m F_2$  just north of the magnetic equator flanked by negative anomalies in both hemispheres. Overall, there is a hemispheric asymmetry, with the largest differences clearly in the Northern Hemisphere. This difference pattern is the opposite of the one shown at the longitude over the Pacific Ocean. There are two mechanisms that may explain these differences: a difference of the meridional  $\mathbf{E} \times \mathbf{B}$  drifts resulting in a decrease of the drifts in *noobs*, and a change to the neutral wind driven ion transport at the F-peak. Referring to Figure 3(a,b) we notice that at this longitude ( $\sim 90^\circ$ ) the meridional  $\mathbf{E} \times \mathbf{B}$  in the *noobs* simulation is smaller than in *hybma*. Figure 4 shows the electron density on the SAMI3 slice over the Indian Ocean in *hybma* (Figure 4a) and the differences *noobs* minus *hybma* (Figure 4b). The electron density in the reference simulation (*hybma*) shows the typical arced structure with elevated isopleth of density at the magnetic equator and lowered isopleths in the subtropics: they result from the competing uplift of ions by the meridional  $\mathbf{E} \times \mathbf{B}$  drifts at the magnetic equator and gravity. The difference of electron density between *noobs* and *hybma* just north of the magnetic equator agrees with the  $N_m F_2$  differences in Figure 1. In this case, a weakening of the fountain effect via  $\mathbf{E} \times \mathbf{B}$  drifts at this longitude (Figure 3a,b) creates the pattern in Figure 1c and Figure 4: both the positive and negative anomalies are necessary to explain the effect of the weakening of the  $\mathbf{E} \times \mathbf{B}$  drift. However, this does not explain the substantial asymmetry of the changes across the magnetic equator in Figure 1c. Thus, we now look at the role of the neutral wind. The field line transport at 300 km (near the  $F_2$  peak) in this longitudinal sector (Figure 5a,b) is northward in both simulations, resulting in ions moved northward across the magnetic equator towards the northern EIA. The difference field *noobs* minus *hybma* in Figure 5c show a large positive (northward) difference in *noobs* just west of the longitude of the SAMI3 slice in Figure 4. This pattern of neutral winds illustrates that field-aligned transport in the *noobs* simulation is stronger than in *hybma*: the neutral winds around 300 km in *noobs* move ions across the magnetic equator and accumulate more ions just north of the magnetic equator setting up the north-south asymmetry in the EIA difference field that is shown in Figure 1c around  $90^\circ\text{E}$ .






**Figure 2.** Electron density (dene) at 10:00 LT in units of  $10^4 \text{ cm}^{-3}$  along SAMI3 slice whose longitude at the equator is  $\sim 230^\circ$  averaged for the same period as in Figure 1. The vertical black line identifies the magnetic equator. (a) electron density in *hybma*; (b) difference *noobs* minus *hybma*.

### 3.2 Case 2: 16:00 LT - dynamically active

We now turn our attention to the afternoon hours, with Figure 6 showing the  $N_m F_2$  at 16:00 LT. This case illustrates another instance where the *noobs* and *hybma* simulations differ. At this LT, the  $N_m F_2$  peaks broadly between  $315^\circ\text{E}$  to  $90^\circ\text{E}$  longitude in the northern hemisphere. The  $N_m F_2$  in excess of  $250 \times 10^4 \text{ cm}^{-3}$  is broader and more prominent in *hybma* (Figure 6a), and the minimum of  $N_m F_2$  at the magnetic equator extends over more longitudes and is deeper in *hybma* than in *noobs* (Figure 6b).

Referring again to Figure 3(a,b) we notice that in the mid-afternoon (16:00 LT) around  $15^\circ\text{E}$  longitude the meridional  $\mathbf{E} \times \mathbf{B}$  drift is stronger in *hybma* when compared to *noobs*. Therefore, in the simulation with MA observations, the electric field is more eastward (i.e., positive) when MA observations are used, resulting in stronger uplift of ions toward the F-peak. Figure 7 shows the magnetic field-aligned component of the neutral wind at 300 km as a function of location at 16:00 LT. Note that the *noobs* run has a stronger northward component along the SAMI3 slice crossing over Africa ( $\sim 16^\circ$ ) north of the magnetic equator, and a weaker northward component to the south of the magnetic equator. Thus, the *noobs* run has a larger neutral wind magnitude




**Figure 3.** 10-day average of  $\mathbf{E} \times \mathbf{B}$  ion drifts in  $\text{m s}^{-1}$  obtained from the *hybma* (a,c) and *noobs* (b,d) simulations. (a,b) are the results of averaging during January 1–10, 2013; (c,d) show the drifts averaged during January 21–30, 2013. The dash black lines identify the approximate longitude (vertical lines) of the morning cases examined for 10:00 LT (horizontal line) and discussed in Figure 1. The dot black lines identify the approximate longitude (vertical lines) of the afternoon cases examined for 16:00 LT (horizontal line) and discussed in Figure 6 and Figure 8.

pushing the plasma northward along the field lines and a weaker fountain effect counteracting the interhemispheric transport. The difference in field parallel transport around the African continent SAMI3 slice, coupled with the different strength of the fountain circulation, explain the hemispheric asymmetry seen in the  $N_m F_2$  at these longitudes between the two simulations.

### 3.3 Case 3: 16:00 LT - dynamically quiet


The case studies discussed in Secs. 3.1 and 3.2 illustrate the ionospheric response when the atmosphere is dynamically disturbed, such as during a SSW. As discussed in Section 1, anomalous ionospheric behavior is to be expected during these times. We demonstrate now that while this behavior is prominent during dynamically active times, the effect of MA observations is also detectable and statistically significant during dynamically quiet times.

Figure 8 is the equivalent of Figure 6 but averaged during the last 10 days of January (21–30 January 2013). This time period is well after the disturbance of the SSW on January 6. We chose the mid-afternoon (16:00 LT) because at this time the electron




**Figure 4.** Electron density (dene) at 10:00 LT in units of  $10^4 \text{ cm}^{-3}$  along a SAMI3 slice with a longitude at the equator of  $\sim 90^\circ$ ;  $E$  at the geographic equator. Electron density is averaged for the same period as in Figure 1. The vertical black line identifies the magnetic equator. (a) electron density in *hybma*; (b) difference *noobs* minus *hybma*.


concentration is higher. During the dynamically quiet times of the end of the month, the largest differences between *hybma* and *noobs* shift over the Indonesian maritime region (dashed line). As in the case of the dynamically disturbed days shown in  
 260 Figure 6, there is a pronounced local minimum of  $N_m F_2$  south of the magnetic equator in the *hybma* simulation, which results in a difference field with a positive  $N_m F_2$  anomaly near the magnetic equator and negative  $N_m F_2$  anomalies at sub-tropical latitudes. We note that these anomalies are statistically significant. Similar to the dynamically disturbed times earlier in the month, the anomalous  $N_m F_2$  later in the month is somewhat asymmetric about the magnetic equator, indicating a continued role of the neutral wind in transporting ions across the magnetic equator (though to a lesser degree). Overall, Figure 8 shows  
 265 that the differences in  $N_m F_2$  between the *hybma* and *noobs* simulations are not due to the characteristics of neutral dynamics during a SSW, but rather due to the lack of MA observations in the *noobs* simulation, which plays a role during all types of background conditions.




**Figure 5.** Neutral wind  $V_{n,q}$  parallel to geomagnetic field lines at 300 km and 10:00 LT. Units are  $\text{m s}^{-1}$ ;  $V_{n,q}$  is averaged during 1–10 January 2013. The solid black line indicates the location of the magnetic equator, and the dashed line is the location of the SAMI3 slice crossing over the Indian Ocean. (a)  $V_{n,q}$  for *hybma* (b)  $V_{n,q}$  for *noobs*; (c)  $V_{n,q}$  difference *noobs* minus *hybma*.



**Figure 6.**  $N_m F_2$  in units of  $10^4 \text{ cm}^{-3}$  for all locations at 16:00 LT averaged for 1-10 January 2013: (a) *hybma*; (b) *noobs*; (c) difference *noobs* minus *hybma*. The oblique dash black line highlights the position of SAMI3 slice which crosses the geographic Equator at longitude  $\sim 15^\circ$ . The magnetic equator is described by a solid black line across longitude. Green contours in panel (c) identify the locations where the differences are statistically significant at least at 95% level using a t-test. Notice that compared to Figure 1, longitudes have been rotated to have  $0^\circ \text{E}$  at the center.



**Figure 7.** Neutral wind  $V_{n,q}$  parallel to geomagnetic field lines at 300 km and 16:00 LT. Units are  $\text{m s}^{-1}$ ;  $V_{n,q}$  is averaged during 1–10 January 2013. The solid black line indicates the location of the magnetic equator, and the dashed line is the location of the SAMI3 slice crossing the geographic equator around longitude  $\sim 16^\circ$ . (a)  $V_{n,q}$  for *hybma* (b)  $V_{n,q}$  for *noobs*; (c)  $V_{n,q}$  difference *noobs* minus *hybma*.



**Figure 8.** As in Figure 6,  $N_m F_2$  in units of  $10^4 \text{ cm}^{-3}$  for all locations at 16:00 LT averaged for 21-30 January 2013: (a) *hybma*; (b) *noobs*; (c) difference *noobs* minus *hybma*. The dash black line highlights the position of SAMI3 slice crossing the geographic equator over the Indian Ocean at longitude  $\sim 117^\circ$ . The magnetic equator is described by a solid black line across longitude. Green contours in panel (c) identify the locations where the differences are statistically significant at least at 95% level using a t-test.

#### 4 Discussion

This study uses two model experiments that couple a thermosphere climate model (WACCM-X), nudged with atmospheric specifications (NAVGEM-HA), to an ionospheric model (SAMI3). The two experiments differ only in the observations that were used to generate the atmospheric specifications: the *hybma* experiment includes all observations up to the UMLT and the *noobs* experiment does not use MA observations above about 40 km. The goal of this study is to demonstrate that a potential future loss of MA observations will have significant negative consequences for our ability to predict the [day-to-day-intra-day](#) variability of the neutral atmosphere and ionosphere, and is not only impactful for research on the neutral dynamics of the lower thermosphere (as demonstrated by Sassi et al. (2020)). The results focused on the  $N_m F_2$  at a fixed LT, as the peak electron density at the F-peak is a well-understood and appropriate diagnostic for ionospheric behavior. Using the  $N_m F_2$  at fixed LT also allows an easier interpretation of the differences in terms of differences of non-migrating solar tides, as discussed by Forbes et al. (2006).

The results show that the difference of  $N_m F_2$  between the two simulations is statistically significant (better than 95% confidence level) at the selected locations and LT. In the morning (10:00 LT, Section 3.1), the largest differences between the two model runs is seen over the Pacific and Indian Oceans: a reduction of electric density in *noobs* near the magnetic equator is accompanied by an increase at adjacent latitudes in the Pacific Ocean, while the opposite signature is seen in the Indian Ocean region. The difference in behavior of  $N_m F_2$  is found to be caused by a variation in the fountain effect when the MA measurements are not incorporated: stronger (weaker) uplift of ions over the Pacific (Indian) Ocean when MA observations are removed. At different solar local times and geographic locations, the neutral winds introduce in both simulations a hemispheric asymmetry to the equatorial plasma distribution.

During the afternoon (16:00 LT, Section 3.2), the changes of  $N_m F_2$  over the African continent are similar to those seen in the morning over the Indian Ocean. Examination of the changes along a magnetic meridian shows in fact that the afternoon anomalies are due to the decrease in the magnitude of the meridional  $\mathbf{E} \times \mathbf{B}$  drift in *noobs* when the MA observations are not considered. The magnetic field-aligned component of the neutral wind at 300 km (an altitude chosen as it is just below the peak  $h_m F_2$ ) shows that the longitudinal structure of the lower thermospheric neutral wind has also been affected by the differences in MA forcing. This, in turn, changes the magnitude of the interhemispheric transport at this local time, affecting the longitudinal structure of the  $N_m F_2$ .


The first two cases illustrated (Sections 3.1 and 3.2) include the occurrence of a SSW in the stratosphere, which is known to produce anomalous behavior in the thermosphere and ionosphere; we therefore examined also a dynamically quiet case towards the end of January 2013 (Section 3.3). Overall, the effect of removing MA observations from the atmospheric specifications during a dynamically quiet period is consistent with the more dynamically disturbed cases, although showing a more muted response. This indicates that a dynamically disturbed atmosphere is not required for MA observations to impact predictions of ionospheric structure, although a disturbed state can mediate the magnitude of the impact.

The zonally articulated response in  $N_m F_2$  in all cases due to the exclusion of MA observations indicates that there are differences in the amplitude of non-migrating solar tides. The zonal structure of the difference field indicates prominent wave-3



or wave-4 modes at fixed LT, which is consistent with past findings. For instance, Sassi et al. (2020) showed that the simulation without MA observations (i.e., *noobs*) produces much larger non-migrating tides DE3 and DE2. These solar tides are expected to be visible in  $N_m F_2$  as wave-4 and wave-3 patterns, [which are known to impact the structure of the F-region ionosphere](#) (Immel et al., 2006), when plotted at fixed LT.

Finally, we compare the results of these runs to TEC observations made at a nearby longitude slice just eastward of the one shown in Figure 6 at  $110^\circ$  apex longitude. This longitude slice lies over portions of Africa and Europe that are sufficiently instrumented to provide coverage of the [lower latitude region that includes the](#) EIA. For each day of the model simulations, the Vertical TEC (VTEC) was selected within  $3^\circ$  (for the model runs) or within  $5^\circ$  (for the observational data) and between 15:45 and 16:15 LT. Then, the mean VTEC was calculated at a  $1^\circ$  latitude resolution. Figure 9(right) shows the geographic longitude of the observations and nearby model data used in our analysis. Figure 9(left) shows the result of this averaging process, with the *noobs* run marked by navy circles, the *hybma* run marked by light blue squares, and the observed VTEC from MIT Haystack (Rideout and Coster, 2006) marked by slate blue triangles.



**Figure 9.** (left) Mean VTEC as a function of magnetic apex latitude at 16:00 LT at  $110^\circ$  magnetic apex longitude on 22 January 2013. The *noobs* results are marked by navy circles, the *hybma* results are marked by light blue squares, and the observation data are marked by slate blue triangles. The EIA peak and trough locations (or only a central peak, [as is the case in the \*noobs\* results](#)) are marked by orange stars. (right) Geographic longitude and magnetic latitudes of all available data (light blue circles), and the selected data (dark blue circles) used to assemble the statistics of Tables 1 and 2; the corresponding model data (orange circle) are also shown. The panel was constructed for 1–10 January 2013, and provided here for reference to show that observational data is broadly located at  $\sim 36^\circ$ E.

Figure 9 shows a day where the two model runs have differing [EIA structures, and low-latitude VTEC distributions, with](#) neither of them fully agree with the observed data. The first level of disagreement is the bias between the model and observations. Depending on the neutral atmospheric composition and density, the SAMI3 model tends to overestimate the ionospheric plasma density. When MSIS is used to provide the background density, as is the case in Figure 9, this difference is on the order of  $\sim 10$  TECU near the equator. These differences are most obvious in the VTEC, as it is vertically integrated and exacerbates the problem. Figure 9 also shows the three potential configurations of the [EIA low-latitude VTEC](#) encountered during this study: no dual EIA peak (No, shown by the *noobs* data), a higher northern EIA peak (North, shown by the observational

data), and a higher southern EIA peak (South, shown by the *hybma* data). We will compare the state of the EIA between the observations and the model runs by determining the times when the modeled and observed EIA had the same configuration.

Tables 1 and 2 show the number of days in the month of January 2013 where each model run had a particular EIA configuration, and how this compares to the observed EIA configuration. Comparing the two tables shows that introducing MA observations in the *hybma* run improves the agreement between the observations and the model. This occurs specifically with an increase of the number of days both the model and observations show an EIA with a dominant northern peak: *noobs* had six days in this EIA configuration, three of which agreed with the observations and *hybma* had 13 days in this EIA configuration, eight of which agreed with the observations. As this is the dominant state in the observations (14 days in total), it greatly improves the overall agreement. This area of change confirms that the improved lower atmospheric boundary does a better job specifying the upper thermospheric wind, which drives the EIA hemispheric asymmetries.

**Table 1.** EIA configuration agreement between *noobs* and observed VTEC at 110° magnetic apex longitude, 16:00 LT.

<i>noobs</i>	Observed			
	South	North	No	Marginal Total
South	3	10	6	19
North	2	3	1	6
No	0	1	5	6
Marginal Total	5	14	12	31

**Table 2.** EIA configuration agreement between *hybma* and observed VTEC at 110° magnetic apex longitude, 16:00 LT.

<i>hybma</i>	Observed			
	South	North	No	Marginal Total
South	2	4	5	11
North	3	8	2	13
No	0	2	5	7
Marginal Total	5	14	12	31

## 5 Conclusions

The results of this study support the thesis presented in the introduction: the absence of MA ([above 40 km](#)) observations has important and statistically significant consequences on our ability to predict the structure and variability of the ionosphere and upper atmosphere. The impact of the MA observations on the E-region conductivity ([which drives the meridional E×B drift](#)) and thermospheric neutral winds demonstrate that this region is an essential piece that must be understood to fully grapple with interactions between the neutral atmosphere and the ionosphere. At lower latitudes, multiple types of interactions

between the thermosphere and ionosphere can lead to ion transport that affects the F-region peak density distribution. Model runs driven with incomplete MA observations may have anomalous neutral wind at E-region altitudes that lead to an aberrant fountain effect through electrodynamic interactions. They may also have atypical F-region neutral winds that cause abnormal hemispheric electron density distributions, such as a higher electron density in one hemisphere or a decrease in the seasonal EIA asymmetries.

Strictly speaking, these results apply only to the modeling system we have examined. Other modeling systems (for example, systems that assimilate ionospheric data) may see different impacts on the ionospheric state. It is possible that assimilating sufficient data from appropriate ionospheric and thermospheric data sets could mitigate the need for MA observations for ionospheric predictions. To the best of our knowledge, this is the first study that attempts to investigate how a future lack of observations in the MA potentially impacts the upper atmosphere and ionosphere. Given the importance for civilian and military applications of high fidelity predictions in the upper atmosphere and ionosphere, it is necessary to continue and expand these studies, including examining results from other modeling systems.

*Code and data availability.* SD-WACCM-X simulation output and NAVGEM-HA atmospheric specifications are archived at the Naval Research Laboratory. The source code and forcing datasets of the SD-WACCM-X is available from the NCAR website <http://www.cesm.ucar.edu/>. GPS TEC used in this study is publicly available from the Madrigal database and was converted into magnetic coordinates using apexpy (van der Meeren et al., 2023).

*Author contributions.* FS conceptualized this study and its methodology, led the writing, and performed the model analysis. AGB led the physical analysis and contributed to both the writing and model analysis. SEM contributed to the physical and model analysis, as well as editing. JLT performed model runs and contributed to the writing. JPM contributed to the writing and model analysis.

*Competing interests.* The authors declare that no competing interests are present.

*Acknowledgements.* FS, AGB, and SEM acknowledge the support of the Office of Naval Research. This work is supported in part by a grant of computer time from the DOD High Performance Computing Modernization Program at the US Navy DOD Supercomputing Resource Center (NAVO).

GPS TEC data products and access through the Madrigal distributed data system are provided to the community by the Massachusetts Institute of Technology under support from U.S. National Science Foundation grant AGS-1242204. Data for the TEC processing is provided by the following organizations: UNAVCO, Scripps Orbit and Permanent Array Center, Institut Geographique National, France, International GNSS Service, The Crustal Dynamics Data Information System (CDDIS), National Geodetic Survey, Instituto Brasileiro de Geografia e Estatística, RAMSAC CORS of Instituto Geográfico Nacional del la República Argentina, Arecibo Observatory, Low-Latitude Ionospheric

365 Sensor Network (LISN), Topcon Positioning Systems, Inc., Canadian High Arctic Ionospheric Network, Institute of Geology and Geophysics, Chinese Academy of Sciences, China Meteorology Administration, Centro di Niveau des Eaux Littorales Recherche Sismogiche, Système d'Observation du (SONEL), RENAG : REseau NAational GPS permanent, and GeoNet—the official source of geological hazard information for New Zealand.

## References

- 370 Alken, P., Thébaud, E., Beggan, C. D., Amit, H., Aubert, J., Baerenzung, J., Bondar, T. N., Brown, W. J., Califf, S., Chambodut, A., Chulliat, A., Cox, G. A., Finlay, C. C., Fournier, A., Gillet, N., Grayver, A., Hammer, M. D., Holschneider, M., Huder, L., Hulot, G., Jager, T., Kloss, C., Korte, M., Kuang, W., Kuvshinov, A., Langlais, B., Léger, J. M., Lesur, V., Livermore, P. W., Lowes, F. J., Macmillan, S., Magnes, W., Manda, M., Marsal, S., Matzka, J., Metman, M. C., Minami, T., Morschhauser, A., Mound, J. E., Nair, M., Nakano, S., Olsen, N., Pavón-Carrasco, F. J., Petrov, V. G., Ropp, G., Rother, M., Sabaka, T. J., Sanchez, S., Saturnino, D., Schnepf, N. R., Shen, X.,
- 375 Stolle, C., Tangborn, A., Tøffner-Clausen, L., Toh, H., Torta, J. M., Varner, J., Vervelidou, F., Vigneron, P., Wardinski, I., Wicht, J., Woods, A., Yang, Y., Zeren, Z., and Zhou, B.: International Geomagnetic Reference Field: the thirteenth generation, *Earth, Planets and Space*, 73, 49, <https://doi.org/10.1186/s40623-020-01288-x>, 2021.
- Burrell, A. and Heelis, R.: The influence of hemispheric asymmetries on field-aligned ion drifts at the geomagnetic equator, *Geophys. Res. Lett.*, 39, L19 101, <https://doi.org/10.1029/2012GL053637>, 2012.
- 380 Burrell, A., Heelis, R., and Stoneback, R.: Equatorial longitude and local time variations of topside magnetic field-aligned ion drifts at solar minimum, *J. Geophys. Res.*, 117, A04 304, <https://doi.org/10.1029/2011JA017264>, 2012.
- Eckermann, S. D., Hoppel, K. W., Coy, L., McCormack, J. P., Siskind, D., Nielsen, K., Kochenash, A., Stevens, M. H., Englert, C. R., Singer, W., and Hervig, M.: High-Altitude data assimilation system experiments for the northern summer mesosphere season of 2007, *J. Atmos. Sol.-Terr. Phys.*, 71, 531–551, <https://doi.org/10.1016/j.jastp.2008.09.036>, 2009.
- 385 Eckermann, S. D., Ma, J., Hoppel, K. W., Kuhl, D., Allen, D. R., Doyle, J., Viner, K. C., Ruston, B. C., Baker, N. L., Swadley, S. D., Whitcomb, T. R., Reynolds, C. A., Xu, L., Kaifler, N., Keifler, B., Reid, I. M., Murphy, D. J., and Love, P. T.: High-Altitude (0-100 km) global atmospheric reanalysis system: Description and application to the 2014 austral winter of the Deep Propagating Gravity Wave Experiment (DEEPWAVE), *Mon. Wea. Rev.*, 146, 2639–2666, <https://doi.org/10.1175/MWR-D-17-0386.1>, 2018.
- Emmert, J. T., Drob, D. P., Picone, J. M., Siskind, D. E., Jones Jr., M., Mlynczak, M. G., Bernath, P. F., Chu, X., Doornbos, E., Funke, B., Goncharenko, L. P., Hervig, M. E., Schwartz, M. J., Sheese, P. E., Vargas, F., Williams, B. P., and Yuan, T.: NRLMSIS 2.0: A
- 390 Whole-Atmosphere Empirical Model of Temperature and Neutral Species Densities, *Earth and Space Science*, 8, e2020EA001 321, <https://doi.org/10.1029/2020EA001321>, 2021.
- Erwin, S. and Berger, B.: A race against time to replace aging military weather satellites, <https://spacenews.com/a-race-against-time-to-replace-aging-military-weather-satellites/>, 2021.
- 395 Fang, T., Akmaev, R., Fuller-Rowell, T., Wu, F., Maruyama, N., and Millward, G.: Longitudinal and day-to-day variability in the ionosphere from lower atmosphere tidal forcing, *Geophys. Res. Lett.*, 40, 2523–2528, <https://doi.org/10.1002/grl.50550>, 2013.
- Forbes, J. and Zhang, X.: Quasi 2-day oscillation of the ionosphere: A statistical study, *J. Atmos. Sol.-Terr. Phys.*, 59, 1025–1034, [https://doi.org/10.1016/S1364-6826\(96\)00175-7](https://doi.org/10.1016/S1364-6826(96)00175-7), 1997.
- Forbes, J., Palo, S., and Zhang, X.: Variability of the ionosphere, *J. Atmos. Sol.-Terr. Phys.*, 62, 685 – 693, [https://doi.org/10.1016/S1364-](https://doi.org/10.1016/S1364-6826(00)00029-8)
- 400 6826(00)00029-8, 2000.
- Forbes, J., Russell, J., Miyahara, S., Zhang, X., Palo, S., Mlynczak, M., Mertens, C., and Hagan, M.: Troposphere-thermosphere tidal coupling as measured by the SABER instrument on TIMED during July–September 2002, *J. Geophys. Res.*, 111, A10S06, <https://doi.org/10.1029/2005JA011492>, 2006.
- Forbes, J., Zhang, X., Palo, S., Russell, J., Mertens, C., and Mlynczak, M.: Tidal variability in the ionospheric dynamo region, *J. Geophys. Res.*, p. A02310, <https://doi.org/10.1029/2007JA012737>, 2008.
- 405

- Fuller-Rowell, T., Fang, T., Wang, H., Matthias, V., Hoffmann, P., Hocke, K., and Studer, S.: Impact of Migrating Tides on Electrodynamics During the January 2009 Sudden Stratospheric Warming, in: *Ionospheric space weather: longitude and hemispheric dependencies and lower atmosphere forcing*, edited by Fuller-Rowell, T., Yizengaw, E., Doherty, P., and Basu, S., vol. 220 of *Geophysical Monograph Series*, pp. 165–174, American Geophysical Union, <https://doi.org/10.1002/9781118929216.ch14>, 2017.
- 410 Goncharenko, L., Coster, A., Chau, L., and Valladares, C.: Impact of sudden stratospheric warmings on equatorial ionization anomaly, *J. Geophys. Res.*, 115, <https://doi.org/10.1029/2010JA015400>, 2010.
- Hagan, M., Maute, A., Roble, R., Richmond, A., Immel, T., and England, S.: Connections between deep tropical clouds and the Earth's ionosphere, *Geophys. Res. Lett.*, 34, L20 109, <https://doi.org/10.1029/2007GL030142>, 2007.
- Hanson, W. and Moffett, R.: Ionization transport effects in the equatorial  $F$  region, *J. Geophys. Res.*, 71, 5559–5572, 415 <https://doi.org/10.1029/JZ071i023p05559>, 1966.
- Heelis, R.: Electrodynamics in the low and middle latitude ionosphere: a tutorial, *J. Atmos. Sol.-Terr. Phys.*, 66, 825—838, <https://doi.org/10.1016/j.jastp.2004.01.034>, 2004.
- Hines, C.: Internal atmospheric gravity waves at ionospheric heights, *Canadian Journal of Physics*, 38, 1441–1481, <https://doi.org/10.1139/p60-150>, 1960.
- 420 Hoppel, K. W., Eckermann, S. D., Coy, L., Nedoluha, G., and Allen, D. R.: Evaluation of SSMIS upper atmosphere sounding channels for high-altitude data assimilation, *Mon. Wea. Rev.*, 141, 3314–3330, <https://doi.org/10.1175/MWR-D-13-00003.1>, 2013.
- Huba, J., Joyce, G., and Fedder, J.: Sami2 is Another Model of the Ionosphere (SAMI2): A new low-latitude ionosphere model, *J. Geophys. Res.*, 105, 23 035–23 053, <https://doi.org/10.1029/2000JA000035>, 2000.
- Huba, J., Joyce, G., Krall, J., Siefring, C., and Bernhardt, P.: Self-consistent modeling of equatorial dawn density depletions, *Geophys. Res. Lett.*, 37, L03 104, <https://doi.org/10.1029/2009GL041492>, 2010.
- 425 Immel, T., E., S., England, S., Henderson, S., Hagan, M., Mende, S., Frey, H., Swenson, C., and Paxton, L.: Control of equatorial ionospheric morphology by atmospheric tides, *Geophys. Res. Lett.*, 33, L15 108, <https://doi.org/10.1029/2006GL026161>, 2006.
- Jin, H., Miyoshi, Y., Pancheva, D., Mukhtarov, P., Fujiwara, H. J., and Shinagawa, H.: Response of migrating tides to the stratospheric sudden warming in 2009 and their effects on the ionosphere studied by a whole atmosphere-ionosphere model GAIA with COSMIC and 430 TIMED/SABER observations, *J. Geophys. Res.*, 117, A10 323, <https://doi.org/10.1029/2012JA017650>, 2012.
- Kuhl, D., Rosmond, T., Bishop, C., McLay, J., and Baker, N.: Comparison of hybrid ensemble/4DVAR and 4DVAR within the NAVDAS-AR data assimilation framework, *Mon. Wea. Rev.*, 141, 2740–2758, <https://doi.org/10.1175/MWR-D-12-00182.1>, 2013.
- Laprise, R.: The resolution of spectral models, *Bull. Amer. Meteor. Soc.*, 73, 1453–1455, 1992.
- Laundal, K. M. and Richmond, A. D.: Magnetic Coordinate Systems, 206, 27–59, <https://doi.org/10.1007/s11214-016-0275-y>.
- 435 Liu, H.-L.: Variability and predictability of the space environment as related to lower atmosphere forcing, *Space Wea.*, 14, 634–658, <https://doi.org/10.1002/2016SW001450>, 2016.
- Liu, H.-L., Sassi, F., and Garcia, R.: Error growth in a whole atmosphere climate model, *J. Atmos. Sci.*, 66, 173–186, <https://doi.org/10.1175/2008JAS2825.1>, 2009.
- Liu, H.-L., Foster, B., Hagan, M., McInerney, J., Maute, A., Qian, L., Richmond, A., Roble, R., Solomon, S., Garcia, R., Kinnison, D., Marsh, 440 D., Smith, A., Richter, J., Sassi, F., and Oberheide, J.: Thermosphere extension of the Whole Atmosphere Community Climate Model, *J. Geophys. Res.*, 115, <https://doi.org/10.1029/2010JA015586>, 2010.
- Liu, H.-L., Yudin, V., and Roble, R.: Day-to-day ionospheric variability due to lower atmosphere perturbations, *Geophys. Res. Lett.*, 40, 665–670, <https://doi.org/10.1002/grl.50125>, 2013.

- Liu, H.-L., Baredn, C., Foster, B., Lauritzen, P., Liu, J., Lu, G., Marsh, D., Maute, A., McInerney, J., Pedatella, N., Qian, L.,  
445 Richmond, A., Roble, R., Solomon, S., Vitt, F., and Wang, W.: Development and validation of the Whole Atmosphere Community Climate Model with thermosphere and ionosphere extension (WACCM-X 2.0), *J. Adv. Model. Earth Syst.*, 10, 381–402, <https://doi.org/10.1002/2017MS001232>, 2018.
- McCormack, J., Hoppel, K., Kuhl, D., de Wit, R., Stober, G., Espy, P., Baker, N., Brown, P., Fritts, D., Jacobi, C., Janches, D., Mitchell, N.,  
450 Ruston, B., Swadley, S., Viner, K., Whitcomb, T., and Hibbins, R.: Comparison of mesospheric winds from high-altitude meteorological analysis system and meteor radar observations during the boreal winters of 2009-2010 and 2012-2013, *J. Atmos. Sol.-Terr. Phys.*, 154, 132–166, <https://doi.org/10.1016/j.jastp.2016.12.007>, 2017.
- McDonald, S., Sassi, F., and Mannucci, A.: SAMI3/SD-WACCM-X simulations of ionospheric variability during northern winter 2009, *Space Wea.*, 13, <https://doi.org/10.1002/2015SW001223>, 2015.
- McDonald, S., Sassi, F., Tate, J., McCormack, J., Kuhl, D., Drob, D., Metzler, C., and Mannucci, A.: Impact of non-migrating tides on  
455 the low latitude ionosphere during a sudden stratospheric warming event in January 2010, *J. Atmos. Sol.-Terr. Phys.*, 171, 188–200, <https://doi.org/10.1016/j.jastp.2017.09.012>, 2018.
- Millward, G. H., Muller-Wodrag, I. C. F., Aylward, A. D., Fuller-Rowell, T. J., Richmond, A. D., and Moffett, R. J.: An investigation into the influence of tidal forcing on F region equatorial vertical ion drift using a global ionosphere-thermosphere model with coupled electrodynamics, *J. Geophys. Res.*, 106, 24 733–24 744, <https://doi.org/10.1029/2000JA000342>, 2001.
- 460 Pedatella, N. and Maute, A.: Impact of semidiurnal tide on the midlatitude thermospheric wind and ionosphere during sudden stratospheric warmings, *J. Geophys. Res.*, 120, 10,740–10,753, <https://doi.org/10.1002/2015JA021986>, 2015.
- Pedatella, N., Reader, K., Anderson, J., and Liu, H.-L.: Application of data assimilation in the whole atmosphere community climate model to the study of day-to-day variability in the middle and upper atmosphere, *Geophys. Res. Lett.*, 40, 4469–4474, <https://doi.org/10.1002/grl.50884>, 2013.
- 465 Pedatella, N., Liu, H.-L., Marsh, D., Reader, K., Anderson, J., Chau, J., Goncharenko, L., and Siddiqui, T.: Analysis and hindcast experiments of the 2009 sudden stratospheric warming in WACCMX+DART, *J. Geophys. Res.*, 123, <https://doi.org/10.1002/2017JA025107>, 2018.
- Pedatella, N., Liu, H.-L., Marsh, D., Reader, K., and Anderson: Error growth in the mesosphere and lower thermosphere based on hindcast experiments in a whole atmosphere model, *Space Wea.*, 17, <https://doi.org/https://doi.org/10.1029/2019SW002221>, 2019.
- Richmond, A.: Ionospheric electrodynamics using magnetix APEX coordinates, *J. of Geomag. and Geoelec.*, 47, 191–212,  
470 <https://doi.org/10.1002/2017JA025107>, 1995.
- Rideout, W. and Coster, A. J.: Automated GPS processing for global total electron content data, *GPS Solutions*, 10, 219–228, <https://doi.org/10.1007/s10291-006-0029-5>, 2006.
- Rishbeth, H.: Thermospheric winds and the F-region: A review, *J. Atmos. Sol.-Terr. Phys.*, 34, 1–47, [https://doi.org/10.1016/0021-9169\(72\)90003-7](https://doi.org/10.1016/0021-9169(72)90003-7), 1972.
- 475 Sassi, F., Liu, H.-L., Ma, J., and Garcia, R.: The lower thermosphere during the northern hemisphere winter of 2009: A modeling study using high-altitude data assimilation products in WACCM-X, *J. Geophys. Res.*, 118, <https://doi.org/10.1002/jgrd.50632>, 2013.
- Sassi, F., McCormack, J., and McDonald, S.: Whole atmosphere coupling on intraseasonal and interseasonal time scales: a potential source of increased predictive capability, *Radio Sci.*, 54, 913–933, <https://doi.org/10.1029/2019RS006847>, 2019.
- Sassi, F., McCormack, J., Tate, J., Kuhl, D., and Baker, N.: Assessing the impact of middle atmosphere observations on day-to-day variability  
480 in lower thermospheric winds using WACCM-X, *J. Atmos. Sol.-Terr. Phys.*, In press, <https://doi.org/10.1016/j.jastp.2020.105486>, 2020.

- Solomon, S. and Qian, L.: Solar extreme-ultraviolet irradiance for general circulation models, *J. Geophys. Res.*, 110, A10306, <https://doi.org/10.1029/2005JA011160>, 2005.
- van der Meeren, C., Laundal, K. M., Burrell, A. G., Lamarche, L. L., Starr, G., Reimer, A. S., and Morschhauser, A.: *aburrell/apexpy: ApexPy Version 2.0.1*, <https://doi.org/10.5281/zenodo.7818719>, 2023.
- 485 VanZandt, T. E., Clark, W. L., and Warnock, J. M.: Magnetic Apex Coordinates: A Magnetic Coordinate System for the Ionospheric F2 Layer, *Journal of Geophysical Research*, 77, 2406–2411, <https://doi.org/10.1029/JA077i013p02406>, 1972.
- Wang, H., Fuller-Rowell, T., Akmaev, R., Hu, M., Kliet, D., and Iredell, M.: First simulations with a whole atmosphere data assimilation and forecast system: The January 2009 major sudden stratospheric warming, *J. Geophys. Res.*, 116, <https://doi.org/10.1029/2011JA017081>, 2011.
- 490 Weimer, D.: Models of high-latitude electric potentials derived with a least error fit of spherical harmonic coefficients, *J. Geophys. Res.*, 100, 19595–19607, <https://doi.org/10.1029/95JA01755>, 1995.
- Weimer, D.: Improved ionospheric electrodynamics models and application to calculating Joule heating rates, *J. Geophys. Res.*, 110, A05306, <https://doi.org/10.1029/2004JA010884>, 2005.
- 495 Wu, T.-Y., Liu, J.-Y., Chang, L., Lin, C.-H., and Chiu, Y.-C.: Equatorial ionization anomaly response to lunar phase and stratospheric sudden warming, *Scientific Reports*, 11, 14695, <https://doi.org/10.1038/s41598-021-94326-x>, 2021.