

Stability requirements of ~~observation systems~~ satellites to detect long-term stratospheric ozone trends based upon Monte Carlo simulations

Mark Weber¹

¹Institut für Umweltphysik, Universität Bremen FB1, Bremen, Germany

Correspondence: Mark Weber (weber@uni-bremen.de)

Abstract. For new ~~observing systems, particularly satellites,~~ satellite instruments, specifications on the stability required for climate variables are provided in order to be useful for certain applications, for instance, deriving long-term trends. The stability is usually stated in units of percent per decade (%/dec) and is often associated with or termed instrument drift. A stability requirement of 3%/dec or better has been recently stated for tropospheric and stratospheric ozone. However, the way this number is derived is not clear. In this study we use Monte Carlo simulations to investigate how a stability requirement translates into uncertainties of long-term trends depending on the lifetime of individual observing systems, which are merged into timeseries, and the period of available observations. Depending on the need to observe a certain trend over a given period, e.g. typically +1%/dec for total ozone and +2%/dec for stratospheric ozone over thirty years, stability for observation systems can be properly specified and justified in order to achieve statistical significance in the observed long-term trend. Assuming a typical mean lifetime of seven years for an individual observing system and a stability of 3%/dec results in a 2%/dec trend uncertainty over a period of 30 years, which is barely sufficient for stratospheric ozone but too high for total ozone. Having two or three observing systems simultaneously ~~time~~-reduces the uncertainty by 30% and 42%, respectively. Such redundancies may be more efficient than developing satellite instruments with higher long-term stability to reduce long-term trend uncertainties. The method presented here is applicable to any variable of interest for which long-term changes are to be detected.

15 1 Introduction

Ninety percent of ozone in the overhead (total) column resides in the stratosphere and protects the earth system from harmful UV radiation (Zerefos et al., 2023). The phase-out of ozone-depleting substances (ODS) according to the Montreal Protocol and its Amendments, lead to recovery of stratospheric ozone in some ~~region~~ regions of the atmosphere after the late 1990s (Hassler et al., 2022). ~~Most-Statistically~~ significant trends are observed in the upper stratosphere by up to about +2%/dec (Godin-Beekmann et al., 2022). In the extratropics, total column ozone has increased by a rate of up to about +1%/dec (Coldewey-Egbers et al., 2022; Weber et al., 2022). The reported statistical trend uncertainty at the 2σ level for ozone profiles are typically of the same order than the trend itself, about 1-2%/dec, similarly for total column ozone at about 0.5-1%/dec (LOTUS, 2019; Godin-Beekmann et al., 2022; Weber et al., 2018, 2022).

Uncertainties in ozone trends reported recently (e.g. Hassler et al., 2022; Godin-Beekmann et al., 2022; Weber et al., 2022) are determined from the statistics of trend regression alone. Bourassa et al. (2014) reported added uncertainties due to drifts of the OSIRIS satellite to the statistical uncertainty, here 3%/dec, which reduced the atmospheric region (altitude and latitude) of the atmosphere showing significant positive trends. Direct comparison between various merged ozone profiles show that the drifts in the difference between zonal mean single-instrument datasets are on the order of 3%/dec (2σ) but not always statistically significant (Rahpoe et al., 2015; Hubert et al., 2016). Similarly, the spread in the recent total ozone trends from the available merged datasets are on the order of $\pm 0.5\%/dec$, which are indicative of drifts between the merged datasets %/dec (2σ). The apparent drifts are not only due to changes in the instrument performance but can also depend on ~~the way~~ how the individual datasets are merged into the long-term dataset (Frith et al., 2017; Weber et al., 2022). Since multiple time series are available, the statistical trend uncertainty can be further reduced by averaging trends or calculating trends from the the median or mean of datasets (e.g. Steinbrecht et al., 2017; LOTUS, 2019; Godin-Beekmann et al., 2022; Weber et al., 2022).

Ozone is expected to recover from the decrease in ozone-depleting substances but long-term changes also depend on the evolution of greenhouse gases (GHG) as well as on the feedback mechanism between ozone and climate (Hassler et al., 2022). The Vienna Convention for the Protection of the Ozone Layer (VIENNA, 1985), which set among others the framework for the Montreal Protocol signed in 1987, also stated the need for continuing observation of ozone, related species, and climate gases. For long-term observations of ozone and other trace species, specific requirements on accuracy and stability for observing systems were defined and updated over the years (e.g. GCOS-01, 2001; IGACO-04, 2004; GCOS-11, 2011; CCI, 2021; CMUG, 2022; GCOS-22, 2022). For trend detection, a stability requirement of 3%/dec ~~were was~~ stated initially for stratospheric ozone and 1%/dec for total ozone (GCOS-01, 2001). Main application ~~area areas~~ defined were trends and operational meteorology for stratospheric ozone (GCOS-01, 2001). The 3%/dec value is also currently defined as the threshold value for stratospheric profiles and total column of ozone, meaning that ~~observing system will be only a satellite instrument will only be~~ useful if this threshold is not exceeded (CCI, 2021; GCOS-22, 2022; CMUG, 2022). In addition to the threshold value, a breakthrough (2%/dec) and target value (1%/dec) is provided (GCOS-22, 2022). No specification in these documents is given if the stability requirement is stated as one (1σ) or two standard deviations (2σ). With regard to the observed long-term ozone trends, these values make only sense if they are defined as 2σ (two standard deviations).

It is not clear how ~~the specification these specifications~~ are derived, but very likely they are ~~leaned~~ derived from the observed ~~ozone statistical~~ statistical ozone trend uncertainties and the expectation that the requirements should be close but lower than these uncertainties. The question we ~~rise~~ raise here in the paper is how such a requirement can be justified from the need to achieve a certain trend uncertainty over a given period of time. From the ozone recovery perspective, the time range since ODS peaked in the stratosphere is close to thirty years. A similar but different question was addressed by Weatherhead et al. (1998). Based upon the noise of data and serial correlation, they determined on statistical grounds how many years we need to observe a significant trend, which is the case when twice the trend uncertainty is lower than the trend magnitude (2σ significance).

We use Monte Carlo ~~simulation~~ simulations to investigate how a stability requirement translates into uncertainties of long-term trends depending on the lifetime of individual ~~observing systems, merged into~~ satellite instruments, which are merged into

long-term timeseries, and the overall period of available observations. In Section 2, the scheme of the Monte Carlo simulation is described. Results are presented in Section 3, and discussed in Section 4. We close with concluding remarks in Section 5.

60 2 Monte Carlo simulations

The scheme of Monte Carlo simulations of time series and derived trend uncertainties due to a given stability requirement is shown in Fig. 1 (e.g. Guimarães Couto et al., 2013). We assume that a simulated timeseries is zero without any drift and bias after subtraction of the start value is zero. Each individual observing system with a given lifetime (segment) of which we compose a long-term merged dataset (timeseries), has varying drifts in %/dec, which follow a Gaussian distribution with a given 1σ stability requirement, e.g. 1.5%/dec as shown in panel a. The variability seen in the sample of one million long-term timeseries (panel b) are therefore solely due to the instrument drifts following the stability distribution. As an example four single timeseries over 30 years are shown in red. Each segment is seven years long corresponding to a constant lifetime. For simplicity, an overlap period is not considered here (Weatherhead et al., 2017).

For each of the timeseries ~~a trend was determined and its distribution of the sample~~, a simple linear trend was fit and the distribution of trends and its width representing the trend uncertainty given by the standard deviation of the distribution is shown in panel c. Here the trends are calculated for a thirty year period. A stability requirement of 1.5%/dec (1σ) and a lifetime of 7 years for each observing system results in a trend uncertainty of 0.77%/dec after 30 years (close to half the value of the stability requirement). With a required limit of stability of 3%/dec (2σ) for ozone, trends lower than about 1.5%/dec (2σ) are not detectable after 30 years (simply doubling the numbers given in the previous sentence).

Some additional modifications in the simulation were introduced. In a merged dataset the individual observing systems have varying lifetimes. An expected lifetime of seven years is typical for current satellite systems, e.g. TROPOMI (Veefkind et al., 2012). Several satellite instruments measured for a decade or more (e.g. SAGE II: 21 years, OSIRIS: more than 20 years, OMI: 19 years, GOME-2A: 12 years, SCIAMACHY, MIPAS, GOMOS: 10 years). In a different setup we therefore vary the lifetimes of individual segments following a Poisson distribution with a mean of seven years (Fig. 2). Later we will show also results for instruments with a longer mean lifetime of 12 years. When merging datasets, usually segments of a single observation system with some extended records are used. In our case we set a lower limit of five years to be included in the timeseries.

Additional uncertainties come from biases between individuals segments. Biases can be corrected when sufficient overlap exists between observing systems as discussed in detail in Weatherhead et al. (2017). In our simulations we do not consider any overlap periods, but we allow for a Gaussian distributed bias from one segment to the other. Figure 3 shows the samples of timeseries assuming 0, 1, and 2% bias (1σ), respectively. In both simulations varying lifetimes with a mean of 7 years and Poisson distributed (Fig. 2) were assumed.

With a stability requirement of 1.5%/dec the trend uncertainty after 30 years increases slightly from 0.77 to 0.83%/dec using varying lifetimes instead of a constant lifetime of seven years (Figs. 2b and 3a). A larger change in uncertainties is seen by adding a 1% bias between segments, resulting into an increase to 1.1%/dec. This means that, for ozone observed with a 3%/dec stability ~~in observing systems~~ and a typical bias of 1% between them, the trend uncertainty is 2.2%/dec over 30 years.




Figure 1. Monte-Carlo simulation of 30-year trend uncertainties assuming a stability requirement of 3%/dec (2σ) for individual observing systems with a lifetime of seven years. Panel a: Gaussian instrumental drift distribution assuming a stability requirement of 3%/dec (2σ). Panel b: Ten thousand timeseries (blue lines) from the one million sample of simulated timeseries, composed of seven-year individual segments (single instrument observations), are shown. Red lines show four examples of timeseries from the sample. Panel c: Distribution of linear trends derived from all timeseries.

3 Results

The dependence of the trend uncertainty as a function of observation years, assuming a stability of 1.5%/dec, is shown in Fig. 4. The different colours in the plot represents lifetimes for the individual observing systems varying from 5 to 20 years. For a given stability requirement and period of the timeseries, the trend uncertainties increases with lifetime. This is on first sight somewhat unexpected as we normally desire to have observing or satellite missions to operate as long as possible. A continuous drift over a longer time period will cause larger deviations of the timeseries from the truth and increases trend uncertainties. It is also evident that the reduction in trend uncertainties with time slows considerably after two decades.




Figure 2. Distribution of lifetimes for single observing systems based upon a Poisson distribution with a mean of seven years. Segments with lifetimes of less than 5 years are not allowed in the timeseries and the distribution is set to zero below five years.

So far we considered the case of one single mission contributing to the time series. If we have several parallel missions with identical geographical coverage for long-term monitoring, the trend uncertainties can be strongly reduced. To investigate this effect, mean timeseries were constructed by averaging n randomly generated timeseries, with n being interpreted as the number of parallel missions. The result is shown in Fig. 5. Trend uncertainties are reduced by a factor of $\frac{1}{\sqrt{n}}$ from the single mission case. With two and three missions in parallel over the entire period, the trend uncertainty is reduced by nearly 30% and 42%, respectively. Redundancy in ~~observing systems~~ satellites is therefore more effective in reducing trend uncertainties than improving the stability of ~~an observing system~~ a satellite instrument. This is an important consideration in a long-term monitoring program (Harris et al., 2015).

~~Trend uncertainty in %/dec as a function of observation years for various stability requirements (panels a to c), different mean lifetimes of single observing systems (solid and dashed lines), and biases between consecutive observation systems (colors).~~

Figure 6 displays the impact of biases between segments on trend uncertainty as a function of the timeseries length. Here we ~~assume again the~~ show in panels a to c, results for various stability requirements of 0.5, 1.0, and 1.5%/dec stability and (all 1σ), which corresponds to the GCOS-22 requirements for target, breakthrough, and threshold, assuming they are given as 2σ values, and results are shown for variable lifetimes with a mean of 7 (solid) and 12 years (dashed line). If there are no biases (black curves in all panels), then the trend uncertainty ~~decrease~~ decreases faster with time for lower lifetimes as already demonstrated in Fig. 4. If the mean bias gets larger, this reverses and trend uncertainty decreases faster for timeseries with segments having longer lifetimes. With a bias between consecutive ~~observation systems~~ satellite instruments, the trend uncertainty initially increases with time and starts decreasing after a couple years, while the trend uncertainties of bias-free timeseries immediately decrease with additional years of data. A too large bias causes the trend uncertainty to be always larger than the stability requirement (red and blue curves in panel a). One of the requirements for long-term monitoring is




Figure 3. Samples of 30-year timeseries composed of segments with varying lifetimes following a Poisson distribution with a mean and standard deviation of 7 years. Ten thousand timeseries (blue lines) from the one million sample of simulated timeseries, composed of seven-year individual segments (single instrument observations), are shown in each panel. Red lines show four examples of timeseries from the sample. Panel a: no bias between segments; panel b and c: biases drawn from a Gaussian distribution with $\sigma = 1\%$ and 2% , respectively.

to keep biases between consecutive ~~observing systems~~ satellite observations as low as possible. Biases can be considerably




Figure 4. Trend uncertainties as a function of observation period assuming a stability requirement of 1.5%/dec for individual segments with a constant lifetime. Different colours show results for lifetimes varying from 5 to 20 years.




Figure 5. Trend uncertainties as a function of observation period and number of parallel observing systems satellites ($n = 1, 2, 3, \text{ and } 5$) assuming a constant lifetime of 7 years and a stability of 1.5%/dec (1σ) for each system satellite.

reduced when the overlap period of two observation systems or consecutive satellite missions is sufficiently long as discussed in Weatherhead et al. (2017). Redundancy in observing systems observations again helps to reduce effect from the biases.

4 Discussion

We present here a simple approach of Monte Carlo simulations to specify stability requirements dependent on the need to observe a certain trend in merged datasets after a given period of time. Stratospheric ozone is expected to recover since the end of the 1990s (somewhat less than thirty years ago) when stratospheric halogens released from ozone-depleting substances reached maximum. ~~A rule of thumb is that after thirty years the trend uncertainty is about 2%/dec assuming a typical~~ While there are many possible combinations of satellite instruments, we show that a reasonable assumption of repeated single instruments with a mean lifetime of 7 years and a stability requirement of 3%/dec of the individual missions or observing systems of which the timeseries is composed of (2σ) would yield a trend uncertainty of about 2%/dec (2σ) after thirty years of observations. If the bias between consecutive observing systems is negligible due to a good bias correction in the case of a sufficient overlap between consecutive missions, the uncertainty improves to about 1.5%/dec.

Our simple approach ~~of course~~ neglects the larger likelihood that instrument drifts with time are not necessarily linear nor ~~drifts are normal~~ are drifts normally distributed. Largest changes in satellite performance ~~is~~ are usually in the beginning or near the end of the mission. When the ambient environment of a satellite instrument changes from the clean room at the ground to space conditions shortly after launch, calibration settings changes and if these changes are not adequately accounted for in the trace gas retrieval, causes data drifts. In the first few years, instruments in space outgas that which change the optical performance of the instrument more rapidly than in later years. In general drifts can be caused by any changes in the optical performance (e.g. Bourassa et al., 2014). For limb based satellite data, pointing errors to the tangent height can increase with time, particularly in the later part of the mission period (Bourassa et al., 2018). Near the end of the mission, instruments are aged and suffer from loss in thermal and/or power stability which may enhance again the instrumental drift in the observational data.

The multiple merged satellite ozone data records are usually not entirely independent from each other, as the same satellite missions are used in different merged timeseries (e.g. Steinbrecht et al., 2017; LOTUS, 2019; Godin-Beekmann et al., 2022; Weber et al., 2022). In such a case the trend uncertainty reduces more slowly than with $1/\sqrt{n}$, which is only valid for n truly independent timeseries.

145 5 Conclusions

Using a simple Monte Carlo timeseries simulation scheme, we were able to estimate the required stability of an observing instrument with a given lifetime that will be needed to reach a certain trend uncertainty in a long-term timeseries consisting of ~~many of those systems or~~ series of satellites. This requirement does not only depend on the length of the timeseries and lifetime, but also on the biases between consecutive observing systems. The time range of the ozone recovery phase is now approaching 30 years in the next few years. With a typical mean lifetime of about 7 years for ~~an observing system, in particular~~ satellites, a 3%/dec (2σ) stability (GCOS-22, 2022) adds about 1.5%/dec (2σ) uncertainty to the statistical long-term trend uncertainty after 30 years. ~~For total ozone the GCOS requirement is insufficient and must be higher and on the order of 1%/dec.~~

~~The time range of the ozone recovery phase is approaching 30 years in the next few years.~~ Assuming a mean residual bias of about 1% between consecutive measurement systems (after overlap corrections), the resulting long-term trend uncertainty increases to ~~2%/dec over this period. For~~ 1%/dec. Combining the statistical (2%/dec, Godin-Beekmann et al., 2022) and drift uncertainty (2%/dec) as a Gaussian sum yields an overall trend uncertainty of about 3%/dec, which is larger than the largest positive trends observed in the upper stratosphere (+2%/dec). For total ozone with statistical trend uncertainties of about 0.5%/dec(2 σ), the total uncertainty will be about 2%/dec after 30 years. For long-term trend assessments the current threshold requirement of 3%/dec (2 σ), as stated in GCOS-22 (2022), is too high and it is recommended to be reduced to 1%/dec (2 σ).

155

160 For the next decades, the ~~next decades the~~ long-term trend uncertainty will only decrease rather slowly, so that redundancy in ~~observing systems~~ satellite instruments is very effective in reducing the impact from instrument drifts. Uncertainties due to instrument drifts can be cut nearly in half with three parallel ~~observing systems~~ satellites. All numbers provided here are lower limits since measurements errors as well as uncertainties due to data merging have not been accounted for.




Figure 6. Trend uncertainty in %/dec as a function of observation years for various stability requirements (panels a to c), different mean lifetimes of single satellite instruments (solid and dashed lines), and biases between consecutive observation systems (colors). Panels a to c show trend uncertainties for the 2σ GCOS-22 stability requirements of 1%/dec (target), 2%/dec (breakthrough), and 3%/dec (threshold), respectively.

Code and data availability. Codes in IDL (Interactive Data Language) and simulated data are available upon request to the author.

165 *Author contributions.* The single author did all the work.

Competing interests. The author is associate editor of this journal.

Acknowledgements. This work was carried out as part of the OREGANO (Ozone Recovery from Merged Observational Data and Model Analysis) project funded by the European Space Agency under the Contract 4000137112/22/I-AG. The author is also grateful for financial support by the State of Bremen via the University of Bremen.

170 References

- Bourassa, A. E., Degenstein, D. A., Randel, W. J., Zawodny, J. M., Kyrölä, E., McLinden, C. A., Sioris, C. E., and Roth, C. Z.: Trends in stratospheric ozone derived from merged SAGE II and Odin-OSIRIS satellite observations, *Atmospheric Chemistry and Physics*, 14, 6983–6994, <https://doi.org/10.5194/acp-14-6983-2014>, 2014.
- Bourassa, A. E., Roth, C. Z., Zawada, D. J., Rieger, L. A., McLinden, C. A., and Degenstein, D. A.: Drift-corrected Odin-OSIRIS ozone product: algorithm and updated stratospheric ozone trends, *Atmospheric Measurement Techniques*, 11, 489–498, <https://doi.org/10.5194/amt-11-489-2018>, 2018.
- CCI: Ozone cci - User Requirement Document (URD), Version 3.1, https://climate.esa.int/media/documents/Ozone_cci_urd_v3.1_version_05032021.pdf, last access: November, 21, 2023, 2021.
- CMUG: Climate Modelling User Group [CMUG] Deliverable 1.1 Meeting the needs of the Climate Community - Requirements, https://admin.climate.esa.int/media/documents/CMUG_Baseline_Requirements_D1.1_v3.0.pdf, last access: November 30, 2023, 2022.
- 180 Coldewey-Egbers, M., Loyola, D. G., Lerot, C., and Van Roozendaal, M.: Global, regional and seasonal analysis of total ozone trends derived from the 1995-2020 GTO-ECV climate data record, *Atmospheric Chemistry and Physics*, 22, 6861–6878, <https://doi.org/10.5194/acp-22-6861-2022>, 2022.
- Frith, S. M., Stolarski, R. S., Kramarova, N. A., and McPeters, R. D.: Estimating uncertainties in the SBUV Version 8.6 merged profile ozone data set, *Atmospheric Chemistry and Physics*, 17, 14 695–14 707, <https://doi.org/10.5194/acp-17-14695-2017>, 2017.
- 185 GCOS-01: WMO/CEOS report on a strategy for integrating satellite and ground-based observations of ozone, WMO GAW-140, <https://library.wmo.int/idurl/4/36995>, last access: November 21, 2023, 2001.
- GCOS-11: Systematic Observation Requirements for Satellite-based Products for Climate Supplemental details to the satellite-based component of the Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC - 2011 update, WMO GCOS-254, <https://library.wmo.int/idurl/4/48411>, last access: December 1, 2023, 2011.
- 190 GCOS-22: The 2022 GCOS ECVs Requirements, WMO GCOS-245, <https://library.wmo.int/idurl/4/58111>, last access: November, 21, 2023, 2022.
- Godin-Beekmann, S., Azouz, N., Sofieva, V. F., Hubert, D., Petropavlovskikh, I., Effertz, P., Ancellet, G., Degenstein, D. A., Zawada, D., Froidevaux, L., Frith, S., Wild, J., Davis, S., Steinbrecht, W., Leblanc, T., Querel, R., Tourpali, K., Damadeo, R., Maillard Barras, E., Stübi, R., Vigouroux, C., Arosio, C., Nedoluha, G., Boyd, I., Van Malderen, R., Mahieu, E., Smale, D., and Sussmann, R.: Updated trends of the stratospheric ozone vertical distribution in the 60°S-60°N latitude range based on the LOTUS regression model, *Atmospheric Chemistry and Physics*, 22, 11 657–11 673, <https://doi.org/10.5194/acp-22-11657-2022>, 2022.
- 195 Guimarães Couto, P. R., Carreteiro Damasceno, J., and Pinheiro de Oliveira, S.: Monte Carlo Simulations Applied to Uncertainty in Measurement, chap. 2, pp. 27–51, IntechOpen Limited, <https://doi.org/10.5772/53014>, 2013.
- 200 Harris, N. R. P., Hassler, B., Tummon, F., Bodeker, G. E., Hubert, D., Petropavlovskikh, I., Steinbrecht, W., Anderson, J., Bhartia, P. K., Boone, C. D., Bourassa, A., Davis, S. M., Degenstein, D., Delcloo, A., Frith, S. M., Froidevaux, L., Godin-Beekmann, S., Jones, N., Kurylo, M. J., Kyrölä, E., Laine, M., Leblanc, S. T., Lambert, J.-C., Liley, B., Mahieu, E., Maycock, A., de Mazière, M., Parrish, A., Querel, R., Rosenlof, K. H., Roth, C., Sioris, C., Staehelin, J., Stolarski, R. S., Stübi, R., Tamminen, J., Vigouroux, C., Walker, K. A., Wang, H. J., Wild, J., and Zawodny, J. M.: Past changes in the vertical distribution of ozone - Part 3: Analysis and interpretation of trends, *Atmospheric Chemistry and Physics*, 15, 9965–9982, <https://doi.org/10.5194/acp-15-9965-2015>, 2015.
- 205

- Hassler, B., Young, P. J., Ball, W. T., Damadeo, R., Keeble, J., Barras, E. M., Sofieva, V., and Zeng, G.: Update on Global Ozone: Past, Present, and Future, in: *Scientific Assessment of Ozone Depletion: 2022*, chap. 3, World Meteorological Organization/UNEP, <https://library.wmo.int/idurl/4/58360>, wMO GAW-58, 2022.
- 210 Hubert, D., Lambert, J.-C., Verhoelst, T., Granville, J., Keppens, A., Baray, J.-L., Bourassa, A. E., Cortesi, U., Degenstein, D. A., Froidevaux, L., Godin-Beekmann, S., Hoppel, K. W., Johnson, B. J., Kyrölä, E., Leblanc, T., Lichtenberg, G., Marchand, M., McElroy, C. T., Murtagh, D., Nakane, H., Portafaix, T., Querel, R., Russell III, J. M., Salvador, J., Smit, H. G. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Stübi, R., Swart, D. P. J., Taha, G., Tarasick, D. W., Thompson, A. M., Urban, J., van Gijssel, J. A. E., Van Malderen, R., von der Gathen, P., Walker, K. A., Wolfram, E., and Zawodny, J. M.: Ground-based assessment of the bias and long-term stability of 14 limb and occultation ozone profile data records, *Atmospheric Measurement Techniques*, 9, 2497–2534, <https://doi.org/10.5194/amt-9-2497-2016>, 2016.
- 215 IGACO-04: The Changing Atmosphere - An Integrated Global Atmospheric Chemistry Observation Theme for the IGOS Partnership, WMO GAW-159, ESA SP-1282, https://igaco-o3.fmi.fi/files/IGACO_Theme_Report_wmo_gaw159.pdf, last access: November 21, 2023, 2004.
- LOTUS: SPARC/IO3C/GAW Report on Long-term Ozone Trends and Uncertainties in the Stratosphere, Petropavlovskikh, I., Godin-Beekmann, S., Hubert, D., Damadeo, R., Hassler, B., and Sofieva, V. (eds.), SPARC Report No. 9, WCRP-17/2018, GAW Report No. 241, <https://doi.org/10.17874/f899e57a20b>, 2019.
- 220 Rahpoe, N., Weber, M., Rozanov, A. V., Weigel, K., Bovensmann, H., Burrows, J. P., Laeng, A., Stiller, G., von Clarmann, T., Kyrölä, E., Sofieva, V. F., Tamminen, J., Walker, K., Degenstein, D., Bourassa, A. E., Hargreaves, R., Bernath, P., Urban, J., and Murtagh, D. P.: Relative drifts and biases between six ozone limb satellite measurements from the last decade, *Atmospheric Measurement Techniques*, 8, 4369–4381, <https://doi.org/10.5194/amt-8-4369-2015>, 2015.
- Steinbrecht, W., Froidevaux, L., Fuller, R., Wang, R., Anderson, J., Roth, C., Bourassa, A., Degenstein, D., Damadeo, R., Zawodny, J., Frith, S., McPeters, R., Bhartia, P., Wild, J., Long, C., Davis, S., Rosenlof, K., Sofieva, V., Walker, K., Rahpoe, N., Rozanov, A., Weber, M., Laeng, A., von Clarmann, T., Stiller, G., Kramarova, N., Godin-Beekmann, S., Leblanc, T., Querel, R., Swart, D., Boyd, I., Hocke, K., Kämpfer, N., Maillard Barras, E., Moreira, L., Nedoluha, G., Vigouroux, C., Blumenstock, T., Schneider, M., García, O., Jones, N., Mahieu, E., Smale, D., Kotkamp, M., Robinson, J., Petropavlovskikh, I., Harris, N., Hassler, B., Hubert, D., and Tummon, F.: An update on ozone profile trends for the period 2000 to 2016, *Atmospheric Chemistry and Physics*, 17, 10 675–10 690, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-17-10675-2017)
- 225 17-10675-2017, 2017.
- 230 Veefkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- 235 VIENNA: The Vienna Convention for the Protection of the Ozone Layer, <https://www.ozone.unep.org/treaties/vienna-convention>, last access: November 30, 2023, 1985.
- Weatherhead, E. C., Reinsel, G. C., Tiao, G. C., Meng, X.-L., Choi, D., Cheang, W.-K., Keller, T., DeLuisi, J., Wuebbles, D. J., Kerr, J. B., Miller, A. J., Oltmans, S. J., and Frederick, J. E.: Factors affecting the detection of trends: Statistical considerations and applications to environmental data, *Journal of Geophysical Research: Atmospheres*, 103, 17 149–17 161, <https://doi.org/10.1029/98JD00995>, 1998.
- 240 Weatherhead, E. C., Harder, J., Araujo-Pradere, E. A., Bodeker, G., English, J. M., Flynn, L. E., Frith, S. M., Lazo, J. K., Pilewskie, P., Weber, M., and Woods, T. N.: How long do satellites need to overlap? Evaluation of climate data stability from overlapping satellite records, *Atmospheric Chemistry and Physics*, 17, 15 069–15 093, <https://doi.org/10.5194/acp-17-15069-2017>, 2017.

- 245 Weber, M., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Burrows, J. P., Long, C. S., and Loyola, D.: Total ozone trends from 1979 to 2016 derived from five merged observational datasets - the emergence into ozone recovery, *Atmospheric Chemistry and Physics*, 18, 2097–2117, <https://doi.org/10.5194/acp-18-2097-2018>, 2018.
- Weber, M., Arosio, C., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Tourpali, K., Burrows, J. P., and Loyola, D.: Global total ozone recovery trends attributed to ozone-depleting substance (ODS) changes derived from five merged ozone datasets, *Atmospheric Chemistry and Physics*, 22, 6843–6859, <https://doi.org/10.5194/acp-22-6843-2022>, 2022.
- 250 Zerefos, C., Fountoulakis, I., Eleftheratos, K., and Kazantzidis, A.: Long-term variability of human health-related solar ultraviolet-B radiation doses from the 1980s to the end of the 21st century, *Physiological Reviews*, 103, 1789–1826, <https://doi.org/10.1152/physrev.00031.2022>, 2023.