

# Emerging[R2C1] extreme Saharan-dust events expand northward over the Atlantic and Europe prompting record-breaking PM<sub>10</sub> and PM<sub>2.5</sub> episodes

Sergio Rodríguez<sup>1,\*</sup>, Jessica López-Darias<sup>1</sup>

5 <sup>1</sup>Consejo Superior de Investigaciones Científicas, IPNA CSIC, Tenerife, Canary Islands, Spain.

Correspondence to: Sergio Rodríguez (sergio.rodriguez@csic.es)

**Abstract.** Unprecedented extreme Saharan-dust (duxt) events have recently expanded northward, from subtropical NW Africa to the NE Atlantic and SW Europe, affecting the with severe impacts on the Canary Islands, mainland Spain and continental Portugal [R1C4]. These six historic duxt episodes occurred [R1C2] on 3-5 February 2020, 22-29 February 2020, 15-21 February 2021, 14-17 January 2022, 29 January - 1 February 2022 and 14-20 March 2022. We analysed data of 341 Governmental Air Quality Monitoring Stations (AQMS) of Spain (330) and Portugal (11) [R1C4], where PM<sub>10</sub> and PM<sub>2.5</sub> are measured with European EN-standards, and found that during duxt events PM<sub>10</sub> concentrations are underestimated due to technical limitations of some PM<sub>10</sub> monitors to properly measure extremely high concentrations. We assessed the consistency of PM<sub>10</sub> and PM<sub>2.5</sub> data and reconstructed 1414 1690 [R1C4] PM<sub>10</sub> (1h average) data of 48 and 7 AQMS of Spain and Portugal, respectively, by using our novel *duxt-r* method. During duxt events, 1-hour average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations were within the range 1000-6000 µg/m<sup>3</sup> and 400-1200 µg/m<sup>3</sup>, respectively. The meteorological and dust modelling reanalyses show that the intense winds leading to massive dust plumes occurred within meteorological dipoles formed by a blocking anticyclone over western Europe and a cut-off low located at the southwest, near the Canary Islands, Cape Verde or into the Sahara. These cyclones reached this region by two main paths: deviated southward from the Atlantic mid-latitude westerly circulation by the blocking anticyclone over western Europe [improvement] or deviated northward from the tropical belt. The analysis of the 2000-2022 PM<sub>10</sub> and PM<sub>2.5</sub> time series shows that these six duxt events have no precedent in this region, in the Canary Islands, mainland Spain and mainland Portugal [improvement]. The 22-29 February 2020 event led to (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> concentrations [improvement] within the range 600-1840 µg/m<sup>3</sup> and 200-404 µg/m<sup>3</sup>, respectively, being the most intense dust episodes ever recorded in the Canary Islands. The 14-20 March 2022 event led to (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> values within the range 500-3070 µg/m<sup>3</sup> and 100-690 µg/m<sup>3</sup> in south-eastern, 200-1000 µg/m<sup>3</sup> and 60-260 µg/m<sup>3</sup> in central and 150-500 µg/m<sup>3</sup> and 75-130 µg/m<sup>3</sup> in northern regions of mainland Spain and within the ranges 200-650 µg/m<sup>3</sup> and 30-70 µg/m<sup>3</sup> in continental Portugal, and to 150-430 µg/m<sup>3</sup> and 30-80 µg/m<sup>3</sup> in the Canary Islands [improvement], respectively, being the most intense dust episode ever recorded in these regions mainland Spain [R1C4]. The analysis of the 2000-2022 PM<sub>10</sub> and PM<sub>2.5</sub> time series shows that these six duxt events have no precedent in Spain [improvement]. All duxt events occurred during northern-hemisphere meteorological anomalies characterised by subtropical anticyclones shifted to higher latitudes,

anomalous low pressures expanding beyond the tropical belt and amplified mid-latitude Rossby-waves. undulation  
[improvement] these overall features resembling the anomalies of the atmospheric circulation linked to the anthropogenic  
global warming [R2C1] New studies have reported on recent record beating PM<sub>10</sub> and PM<sub>2.5</sub> episodes linked to dipoles induced  
extreme dust events from North Africa and Asia, in a paradoxical context of multidecadal decrease of dust emissions, a topic  
5 that requires further investigations. [R2C1 R2C2]

## 1 Introduction

Airborne dust aerosol particles are a key component of the Earth System influencing on climate (Kok et al., 2023), ecosystems  
(Yu et al., 2015), fisheries (Rodríguez et al., 2023) and human health (Domínguez-Rodríguez et al., 2021; Tong et al., 2023).  
[R2C8] Major dust sources are located in North Africa, the Middle East and inner Asia (Prospero et al., 2002), accounting for  
10 ≈ 75% of global emissions; secondary sources are located in northern and southern America, southern Africa, Australia and at  
high latitudes (Kok et al., 2023). Because these sources are located in arid regions they have usually been considered as “natural  
dust-sources” (e.g. the bed of naturally dried ancient lakes (Ginoux et al., 2012; Prospero et al., 2002)); however a growing  
body of evidences is showing that human actions such as the soil disruption by traditional grazing and agriculture (Katra, 2020;  
Mulitza et al., 2010; Vukovic et al., 2021), mining (Rodríguez et al., 2011; Zafra-Pérez et al., 2023), drying of water courses  
15 and lakes (Ginoux et al., 2012; Govarchin-Ghale et al., 2021), the expansion of intensive agriculture (Lambert et al., 2020)  
and wildfires (Yu and Ginoux, 2022) are contributing [improvement] to increase dust emissions.

Dust storms cause huge socio-economic impacts linked to [R2C9] loose of visibility, road traffic disruption and accidents,  
deviation of air travel or closure of maritime and air navigation space, cardiovascular and respiratory diseases, loose of soil  
and a decrease in solar energy production (Cañadillas-Ramallo et al., 2022; Domínguez-Rodríguez et al., 2021; Middleton et  
20 al., 2021; Miri and Middleton, 2022; Pi et al., 2020). For this reason, a set of operational “dust services” are available to  
forecast and monitor dust activity by modelling and satellite observations (Mona et al., 2023). In-situ concentrations of PM<sub>10</sub>  
and PM<sub>2.5</sub> (respirable particulate matter -PM- smaller than 10 and 2.5 microns, respectively) regularly measured in air quality  
monitoring networks are commonly used to assess dust impacts and to validate dust models (Mona et al., 2023). In southern  
Europe, Saharan dust events tend to increase PM<sub>10</sub> concentrations up to typical values within the range 40-90 µg/m<sup>3</sup> (24h  
25 average values), [R2C11] being while dust events with (24h average) PM<sub>10</sub> >100 µg/m<sup>3</sup> unusual (Millán-Martínez et al., 2021;  
Pey et al., 2013).

Understanding how climate change is affecting dust emissions is a challenge as these emissions are also affected by the natural  
atmospheric variability (as traced by ENSO, NAO, AMO and other climatic indexes (Evan et al., 2016)), as well as by the  
changes in atmospheric circulation and soil properties (e.g. humidity and biological crust (Rodríguez-Caballero et al., 2022))  
30 as the atmosphere warms due to the increasing concentrations of greenhouse gases.

Current climate models are unable to reproduce the historical increase in atmospheric dust loads observed in paleorecords (Kutuzov et al., 2019; Preunkert et al., 2019). Based on models constraining dust emissions, it has been estimated that the global dust mass load in the modern climate is  $\approx 56\%$  higher than in pre-industrial times (Kok et al., 2023), with a maximum dust load in the mid-1980s and a subsequent decrease attributed to a decrease in North African and Asian dust emissions linked to a slowdown of the atmospheric circulation interconnected to global warming (Evan et al., 2016; Jiang et al., 2023; Liu et al., 2020; Middleton, 2019; Ridley et al., 2014; Xie et al., 2023) [improvement]. In this scenario of decreasing dust trend in North Africa and Asia, a series of unexpected extreme dust-events have recently occurred.

In March 2018 a “*record-breaking Saharan dust plume*” crossed the Eastern Mediterranean (Kaskaoutis et al., 2019), leading to  $PM_{10}$  values of up to (1h-average)  $6000 \mu\text{g}/\text{m}^3$  (Solomos et al., 2018), a three-fold increase in hospital admissions (Lorentzou et al., 2019; Monteiro et al., 2022) and an accelerated snow melting in the Caucasus (Dumont et al., 2020). In June 2020 the so-called “*Godzilla record-breaking trans-Atlantic African dust plume*” (Bi et al., 2023; Francis et al., 2020, 2022; Pu and Jin, 2021) led to (24h average) values  $PM_{10} = 453 \mu\text{g}/\text{m}^3$  in the Caribbean and  $PM_{10}$  and  $PM_{2.5}$  values = 135 and  $74 \mu\text{g}/\text{m}^3$  in southern United States (Yu et al., 2021), respectively. In March 2021, two “*record-breaking dust events*” in China (Gui et al., 2022; Zhang et al., 2023). In November 2021, an “*extreme dust storm*” in Uzbekistan led to (1h average)  $PM_{10}$  and  $PM_{2.5}$  concentrations of up to  $4575 \mu\text{g}/\text{m}^3$  and  $705 \mu\text{g}/\text{m}^3$  (Nishonov et al., 2023; Xi et al., 2023) .

In this study we present a set of unprecedented extreme dust-events that have recently (2020-2022) expanded northward, from North Africa, to the Atlantic and Europe prompting record-breaking  $PM_{10}$  and  $PM_{2.5}$  episodes in Spain. The observed increase in dust activity in the this region Western Euro-Mediterranean region has recently been studied based on meteorological modelling reanalysis [improvement] and aerosol optical depth - satellite measurements (Cuevas et al., 2023). We also focused on the analysis of the consistency of  $PM_{10}$  and  $PM_{2.5}$  data in the Governmental Air Quality Monitoring Networks during the extreme dust events due to the importance of having suitable data in the public databases used for health effects studies, model validation and constrains etc... (Mona et al., 2023). Understanding these emerging [R2C1] extreme dust events is crucial for this region, since climate projections forecast the expansion of the North African drylands toward the northwest, increasing the risk of desertification of Spain and Portugal as the subtropical anticyclones expand in a warming climate (Cresswell-Clay et al., 2022; Guiot and Cramer, 2016) with an associated increase in the desert dust load (Liu et al., 2024) (Gomez et al., 2023) [improvement].

## 2 Methodology

### 2.1 Data of PM<sub>10</sub> and PM<sub>2.5</sub>

We analysed the 2000-2022 data of PM<sub>10</sub> and PM<sub>2.5</sub> recorded in the Governmental Air Quality Monitoring Network of Spain and Portugal [R1C4]. The data from Spain were recorded in 330 air quality stations distributed across the 17 Autonomous Regions of Spain and the Autonomous Spanish city of Ceuta, whereas the data from Portugal were collected in 11 stations distributed across Norte, Centro, Lisboa, Vale do Tejo, Alentejo, Algarve, Madeira and Azores regions [R1C4].

These stations are integrated in the European Air Quality Monitoring Network, which is the largest European infrastructure for PM<sub>10</sub> and PM<sub>2.5</sub> monitoring following standardized methods for measurements, quality assurance (QA) and quality control (QC) (EN-16450:2017 and EN-12341:2015). In these stations, high temporal (10 to 60 minutes) resolution PM<sub>10</sub> and PM<sub>2.5</sub> data are obtained using automatic monitors based on different principle of measurement, such as beta attenuation, tapered element oscillating microbalance and optical particle sizers (Rodríguez et al., 2012), with technical specifications accomplishing the EN-16450:2017 standard. Data of 24h average PM<sub>10</sub> and PM<sub>2.5</sub> are also obtained with the gravimetric reference method (EN-12341:2015), used for QA/QC assessments and for converting the PM<sub>10</sub> and PM<sub>2.5</sub> data obtained with the automatic devices to gravimetric equivalent data by using the European reference protocols (EN-16450:2017). The PM<sub>10</sub> and PM<sub>2.5</sub> data that we used were provided by the Ministry of Ecological Transition and Demographic Challenge and the air quality departments of the Autonomous Regions of Spain and the Agência Portuguesa do Ambiente [R1C4].

### 2.2 Complementary modelling and satellite data

[R1C5] For each specific study event, we also used data of the National Center for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) meteorological reanalysis (Kalnay et al., 1996), of the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2) model dust reanalysis (Gelaro et al., 2017) and of satellite Visible Infrared Imaging Radiometer Suite (VIIRS) [R1C5] images sensor [R2C12] onboard the Suomi polar satellite.

## 3 Results and discussion

The set of extreme dust events, to which we will refer with the acronyms *duxt* or *dx* episodes, occurred on 3-5 February 2020 (dx-01), 22-29 February 2020 (dx-02), 15-21 February 2021 (dx-03), 14-17 January 2022 (dx-04), 29 January - 1 February 2022 (dx-05), and 14–20 March 2022 (dx-06). These duxt events were characterised by dark and reddish “apocalyptic” skies (Fig.1A-1C). National Spanish and international media (Fig.1), the European Copernicus (Fig.1D) and NASA - Earth Observatory (Fig.1E) platforms reported on these historic events and their impacts on socio economical activities. During the

impact of the duxt-02 event in the Canary Islands, record-breaking temperatures occurred, wildfires were favoured by the windy and dry conditions, solar production energy dropped by a 70%, maritime and air navigation space was closed, and thousands of flights were cancelled with huge economic implications due to the transfer of tourists between the Canary Islands and Europe (Fig.1)(Cuevas et al., 2021).

**ALJAZEERA**  
Thousands evacuated as Canary Isles sandstorm fuels wildfires 24 Feb 2020

**EL PAÍS**  
La calima, el viento y el fuego cierran el espacio aéreo y marítimo en Canarias 24 FEB 2020 -

El episodio de calima de Canarias bate récords de temperatura 24 FEB 2020 -

**The Guardian** 24 Feb 2020

Tourists stranded in Canary Islands after Saharan sandstorm blows in  
Dozens of flights cancelled due to poor visibility, leaving holidaymakers stuck at airports



**NEWS** **B B C**



Spain skies turn orange after Saharan dust cloud sweeps over country Europe - 15 March 2022

**EL PAÍS**



No es Marte, es Murcia: el polvo del Sáhara tiñe de rojo el cielo de España 15 MAR 2022 - 11:32 CET



**INDEPENDENT**

**The Washington Post** February 24, 2020  
Canary Islands sandstorm grounds flights, closes schools as Sahara dust moves into open Atlantic  
Dust from Sahara turns sky in Spain to 'Bladerunner' orange



**A) Almería, 15 March 2022**

Figure 1. News on the extreme dust events impacting mainland Spain and the Canary Islands published in international and Spanish national media. Picture of the Cabo de Gata in south-eastern mainland Spain (Almería province) the 15 of March 2022 (dx-06) (taken and provided by Eva de Mas Castroverde).



**B ) Tenerife, dx-02, 23 feb 2020**




**C ) Tenerife, dx-05, 30 jan 2022**

**D ) "Historic" Saharan dust episode in western Europe – CAMS predictions accurate**

15th March 2022

Southwestern Europe is currently experiencing an exceptional Saharan dust episode, which has been turning skies across the region



Sentinel-3 visible imagery showing Saharan dust over southwest Europe on 15 March 2022.


Source: European Union, Copernicus Sentinel-3 imagery

**E ) An Atmospheric River of Dust**

March 15, 2022



On March 15, 2022, a plume of Saharan dust was blown out of North Africa and across the Mediterranean into Western Europe. The dust turned skies orange, blanketed cities, impaired air quality, and stained ski slopes.



The image above, acquired on March 15 by the Visible Infrared Imaging Radiometer Suite (VIIRS) on the NOAA-20 spacecraft, shows the dust plume moving out of Algeria and over the Iberian Peninsula.

**Figure 1 (continue). Pictures of Tenerife (Canary Islands) the 23 of February 2020 (dx-02) and 30 of January 2022 (dx-05) (taken by the authors). Composite of the websites of Copernicus (<https://atmosphere.copernicus.eu/historical-saharan-dust-episode-western-europe-cams-predictions-accurate>) and Earth Observatory (<https://earthobservatory.nasa.gov/images/149588/an-atmospheric-river-of-dust>) reporting on the historic dust event the 15<sup>th</sup> of March 2022.**

### 3.1 Assessment and reconstruction of PM<sub>x</sub> data during the dust events

We analysed the data of PM<sub>10</sub> and PM<sub>2.5</sub> (PM<sub>x</sub>) of 330 Air Quality Monitoring Stations (AQMS) of Spain and of 11 AQMS of Portugal [R1C4]. We found that during the dust events, the ½ hour and 1 hour resolution data of PM<sub>10</sub> increased up to reach a rather constant ‘saturation’ value, that in most of cases was somewhat lower than 1000 µg/m<sup>3</sup> (in many cases between 900 and 1000 µg/m<sup>3</sup>), no values above this threshold appear in the data records. In these cases, PM<sub>10</sub> concentrations remained constant during the period of extremely high dust concentrations (typically 5 to 30 hours), a behaviour that was not generally observed in PM<sub>2.5</sub>, which exhibited a regular variability (with values < 1000 µg/m<sup>3</sup>) and even increases in the periods when PM<sub>10</sub> remained (un-consistently) constant. This behaviour can be observed in the time series of PM<sub>10</sub> (Fig.2A1-A3) and PM<sub>2.5</sub> (Fig.2B1-B3) linked to the dx-01, dx-02, dx-04, dx-05 and dx-06 events (dx-03 is not included in Fig.2 for the sake of brevity). This saturation threshold close to ≈1000 µg/m<sup>3</sup> is the upper operation limit of some PM<sub>10</sub> monitors and is also the top value of the validation data flag in some data recording commercial software used in many governmental air quality monitoring networks, which assume that PM<sub>10</sub> concentrations above this threshold may suffer underestimation due to the high load of particles (e.g. accumulated in the filter-tape of the beta instrument or particle coincidence problems in the optical particle sizers leading to a [R2C13] loss ~~loose~~ of sensitivity) and consequently do not record values above this threshold. In some AQMS the saturation threshold was found at 500 µg/m<sup>3</sup> and even at 200 µg/m<sup>3</sup>. In fact, the EN-16450, EN-12341 and EN-14907 accreditations certifications [improvement] for PM<sub>10</sub> and/or PM<sub>2.5</sub> monitors available in the European market are obtained for specific ranges, whose most frequent upper limit is 1000 µg/m<sup>3</sup> for many monitors (e.g. Met One™ BAM-1020, Comde Derenda™ APM 2 and Thermo Fisher Scientific™ 5014i, 5030i SHARP, TEOM 1405-F and 1405-DF), although is as low as 200 µg/m<sup>3</sup> for some equipment’s (e.g. [improvement] FAI™ Swam 5a) and is, in contrast [improvement], as high as 10000 µg/m<sup>3</sup> for other devices (e.g. PALAST™ Fidas 200 and 200E) according to the certifications agencies (e.g. TÜV, see <https://www.qal1.de/>). In all these cases of PM<sub>10</sub> data affected by saturation, we reconstructed the PM<sub>10</sub> concentrations with the method described in Fig. 3, which we have called *duxt-r* “PM<sub>x</sub> evaluation and reconstruction method based on ratios during extreme dust events”.


In this *duxt-r* method: 1) we first identified the invalid 1 hour (or ½ hour or 10 minutes time resolution) PM<sub>10</sub> data affected by saturation and the associated invalid PM<sub>2.5</sub>/PM<sub>10</sub> ratios, highlighted with red circles in the example of the dx-04 event shown in Fig. 3A and 3B, 2) then, the PM<sub>2.5</sub>/PM<sub>10</sub> ratio during the PM<sub>10</sub> saturation period was estimated by [R1C6] linear interpolation between the last valid PM<sub>2.5</sub>/PM<sub>10</sub> data before saturation and the first valid PM<sub>2.5</sub>/PM<sub>10</sub> after saturation [R2.5/10(i)], highlighted with green points in Fig. 3B, 3) as results of this interpolation the PM<sub>2.5</sub>/PM<sub>10</sub> ratios we used is not constant, it changes hour by hour, e.g. from 0.184 to 0.162 in the example shown in Fig. 3B. Finally, the 1-hour (or ½ hour or 10 minutes) resolution



PM<sub>10</sub> concentrations were determined with equation-01. [RIC7] The method was validated with comparison with data recorded in a few PM<sub>10</sub> monitors not affected by saturation (described below).

$$PM_{10} = \frac{PM_{2.5}}{R_{2.5/10}(t)} \quad (1)$$

We found that the PM<sub>2.5</sub>/PM<sub>10</sub> ratio during the dust events were within the range 0.16 to 0.22 in most of AQMS, a value lower than that observed during regular dust events (typically ≈0.3) and much lower than that observed in environments affected by secondary particle formation and by vehicle exhaust and other combustion emissions (typically within 0.6-0.9) (Rodríguez and López-Darias, 2021).



10 **Figure 2. Time series of 1h average PM<sub>10</sub> (raw) (A1-A3) and PM<sub>2.5</sub> (B1-B) and evaluated and reconstructed PM<sub>10</sub> data (C1-C3) during February 2020, January 2022 and March 2022, indicating the dust events. The number of AQMS included in each plot is shown. PM<sub>10</sub> (raw) means original raw (non-reconstructed) data (A1-A3). Data of PM<sub>2.5</sub> are also raw (B1-B3). PM<sub>10</sub> data of plots C1-C3 combines measured valid and reconstructed data.**




Figure 3. Evaluation, reconstruction and validation of data with the duxt-r method. Plots (A-C) shows data of El Charco site (Fuerteventura) during dx-04 event (14-17 January 2022): (A) data of  $\text{PM}_{10}$  (raw, i.e. including saturated values) and  $\text{PM}_{2.5}$ , (B) the measured  $\text{PM}_{2.5}/\text{PM}_{10}$  (raw) ratio and the interpolated  $\text{PM}_{2.5}/\text{PM}_{10}$  ratio [ $R_{2.5/10}(i)$ ], and (C)  $\text{PM}_{2.5}$ ,  $\text{PM}_{10}$  (raw) and  $\text{PM}_{10}$  reconstructed data. Invalid data due to  $\text{PM}_{10}$  underestimation are highlighted with red circle (A and B) and red points (C and D). D1-D4 scatterplot of  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  data highlighting valid data (black circle), invalid  $\text{PM}_{10}$  data (red circle) and reconstructed  $\text{PM}_{10}$  data (green circle). E1-E4 reconstructed versus measured  $\text{PM}_{10}$  data. **Green number in plot C indicates some of the values of the  $\text{PM}_{2.5}/\text{PM}_{10}$  interpolation [R1C6].**

In our database, we replaced the PM<sub>10</sub> saturated data by the new PM<sub>10</sub> re-constructed data, i.e. the red (saturated) points shown in Fig. 3C were replaced by the green re-constructed points. At each site (AQMS), the consistency of the reconstructed data was assessed by analysing the scatter plot of the PM<sub>2.5</sub> versus PM<sub>10</sub> data (Fig. 3D1-3D4). As example, the results obtained in 5 AQMS located in Almería province (Mediterraneo AQMS) and Murcia (Mompean AQMS) in south-eastern Spain, Salamanca province (Salamanca-6 AQMS) in central northern Spain, and in Fuerteventura (Canary Islands) (Fig. 3D1-3D4) is shown (Fig.3D1-3D4). With this method, the red PM<sub>10</sub>-saturated data shown in Fig. 3D1-3D4 were replaced by the green (reconstructed) data.

Because of the technical manufacturing specification, the automatic PM<sub>10</sub> and PM<sub>2.5</sub> monitors of two AQMS were able to 10 record valid and consistent PM<sub>10</sub> and PM<sub>2.5</sub> data higher than 1000 µg/m<sup>3</sup>. We used these records to validate this methodology (Fig.3E1, 3E2 and 3E4). At these sites, we re-constructed the PM<sub>10</sub> concentrations above 1000 µg/m<sup>3</sup> as if they had experienced saturation and then we compared the “reconstructed versus the measured PM<sub>10</sub> concentrations”; for this comparison we also included data between 500 and 1000 µg/m<sup>3</sup> for the purpose of having a larger dataset (i.e. all PM<sub>10</sub> data > 500 µg/m<sup>3</sup>). We found that the difference between re-constructed and measured PM<sub>10</sub> concentrations ranged between 2 and 9% (Fig.3E1-3E4). 15 In other few AQMS using beta attenuation devices able to provide PM<sub>10</sub> > 1000 µg/m<sup>3</sup>, we found a low PM<sub>10</sub> variability above this threshold (indicating [R1C8] loose of sensitivity due to mass overload in the filter tape) that was inconsistent with the variability in PM<sub>2.5</sub> and that resulted in PM<sub>2.5</sub>/PM<sub>10</sub> ratios similar to those affected by the saturation as described above (red circles in Fig.3A); at these sites we also reconstructed the PM<sub>10</sub> with the *duxt-r* method (Fig.3). Finally, the PM<sub>10</sub> data measured with the automatic monitors (measured [R2C16] and [R2C17] reconstructed) were converted to gravimetric equivalent data by 20 intercomparison with PM<sub>10</sub> data obtained with the gravimetric reference method (EN-12341:2015; 24h sampling, available during 7 to 25 days/month depending on the AQMS) using the standardised procedure (EN-16450:2017).

The new dataset obtained with this *duxt-r* method (PM<sub>10</sub> measured [R2C16] and [R2C17] reconstructed and then converted to gravimetric equivalent) evidences that 1-hour average PM<sub>10</sub> data that appeared as “saturated” at 1000 µg/m<sup>3</sup> actually reached values close to 6000 µg/m<sup>3</sup> in the dx-02 event, close to 1400 µg/m<sup>3</sup> in the event dx-03 [improvement], close to 2000 µg/m<sup>3</sup> in 25 the events dx-01, dx-04 and dx-05, and between 3000 and 4500 µg/m<sup>3</sup> in the events dx-06 (Fig.2C1-2C3).

By applying this methodology, we reconstructed a total of 1690 [R1C4] hourly PM<sub>10</sub> data: 1537 hourly PM<sub>10</sub> data belonged to (i) 48 AQMS of Spain, distributed between the regions Canary Islands (39), Andalucía (5), Murcia (1) and Castilla y León (2) and Madrid (2) and (ii) 153 hourly PM<sub>10</sub> data belonged to 7 AQMS of Portugal, distributed between the regions Lisboa – Vale do Tejo (6) and Alentejo (1) [R1C4]. The data we reconstructed with this method are already available in public data bases of 30 the Governmental Air Quality Networks of Spain, the Ministry of Ecological Transition and the European Environment Agency. The PM<sub>10</sub> data of other 44 AQMS that also experienced PM<sub>10</sub> saturation could not be reconstructed due to the lack of

simultaneous PM<sub>2.5</sub> measurements, a total of 655 hourly data in these AQMS located in the Canary Islands (14), Andalucía (10), Extremadura (2), Castilla y León (11) and Murcia (7). Just to illustrate the huge importance of re-constructing the data, a brief comparison (for a few AQMS) of the 24h average PM<sub>10</sub> concentrations calculated with saturated PM<sub>10</sub> data vs reconstructed PM<sub>10</sub> data: (1) 948 vs 3070  $\mu\text{g}/\text{m}^3$  15 March 2022 in Almería province (Mediterraneo AQMS), (2) 740 vs 1840  $\mu\text{g}/\text{m}^3$  23 February 2020 in Gran Canaria (Playa del Inglés AQMS), (3) 1238 vs 1684  $\mu\text{g}/\text{m}^3$  23 February 2020 in Tenerife (Tomé Cano AQMS, affected by filter mass overload [improvement]), (4) 577 vs 1421  $\mu\text{g}/\text{m}^3$  23 February 2020 in Tenerife (Piscina Municipal AQMS), and (5) 527 vs 621  $\mu\text{g}/\text{m}^3$  15 March 2022 in Granada province (Palacio de Congresos AQMS). The maximum 1-hour average PM<sub>10</sub> and PM<sub>2.5</sub> recorded during dx-01 to dx-06 are in a selection of AQMS is shown in Fig.S1 and S2 [improvement].

10

### 3.2 [RIC10] Analysis of the extreme dust events

[RIC11] The six dust events occurred between February 2020 and March 2022. Four of these episodes affected both the Canary Islands (off NW Africa) and mainland Spain (SW Europe), whereas two of them only affected to the Canaries.

The first two events occurred in February 2020 (Fig.4): 3-5 February 2020 (dx-01; Fig.4A1) and 22-29 February 2020 (dx-02; Fig.4A2). Throughout six weeks (from mid-January to ending February) a blocking anticyclone established over Iberia, i.e. the Iberian Peninsula (Fig. 4F and 4H), resulting in anomalous easterly wind over central Algeria (wind anomaly not shown for the sake of brevity), a scenario favourable to dust events (Alonso-Pérez et al., 2011a). On 3-5 February 2020 a cyclone reached Cape Verde, the low (over Cabo Verde) to high (over Iberia) L-to-H dipole configuration (Fig.4F) resulted in a strong pressure/geopotential gradient and winds that prompted dust emissions and a dense plume of Saharan dust that expanded over the Atlantic to the Canary Islands and toward the Azores (Fig.4G). Across the Canary Islands the dx-01 event resulted in (i) 1h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 300-2100  $\mu\text{g}/\text{m}^3$  (Fig.2C1) and 100-400  $\mu\text{g}/\text{m}^3$  (Fig.2B1; Fig.S1 [improvement]), respectively, and (ii) 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 100-535<sup>x</sup>  $\mu\text{g}/\text{m}^3$  (Fig.4A1; Fig.5A1; <sup>x</sup> = maximum at Tenerife, San Miguel Tajao AQMS) and 50-165<sup>x</sup>  $\mu\text{g}/\text{m}^3$  (Fig.4A2; Fig.5A2) (<sup>x</sup>Tenerife, Tomé Cano AQMS), respectively. In Madeira the dust impact was smoother, with 24h average PM<sub>10</sub> concentrations within the range 50-115  $\mu\text{g}/\text{m}^3$  (Fig.4C), due to this island remained aside the core of the dust plume (Fig.4G). Dust events prompted by the summer North African dipole (NAFDI) was were [improvement] originally introduced by Rodríguez et al. (2015). This concept of meteorological L-to-H dipoles has also been found to drive dust events in the Middle East (Kaskaoutis et al., 2015, 2017) and in [improvement] the June 2020 Godzilla dust event (Francis et al., 2020).

On 22 [R1C9] February 2020, a new cyclone reached again the region of Cape Verde, resulting in a similar L-to-H dipole meteorology (Fig.4G) which prompted a dusty jet stream in the subtropical North Atlantic, impacting the Canary Islands (Fig.4A2, 4H and 4I) [improvement]. This scenario caused the dark oranges skies, record breaking temperatures, wildfires linked to strong dry winds, closure of maritime and air navigation space and the massive (thousands) flight cancellations described above (Fig.1)(Cuevas et al., 2021). During this dx-02 event, extreme PM<sub>x</sub> concentrations were recorded in the Canary Islands, with: (i) 1h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 2000-5254<sup>x</sup> μg/m<sup>3</sup> (Fig.2C1; Fig.S1) (<sup>x</sup>Gran Canaria, Arinaga AQMS) and 400-1129<sup>x</sup> μg/m<sup>3</sup> (<sup>x</sup>Gran Canaria, Mercado Central AQMS) (Fig.2B1; Fig.S1), respectively, and (ii) 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 600-1840<sup>x</sup> μg/m<sup>3</sup> (Fig.4B1, 5B1) (<sup>x</sup>Gran Canaria, Playa del Inglés AQMS) and 200-404<sup>x</sup> μg/m<sup>3</sup> (Fig.4B2, 5B2) (<sup>x</sup>Gran Canaria, Playa del Inglés AQMS), respectively. Concentrations (24h average) of PM<sub>10</sub> during this dx-02 event were up to 6 times higher than the upper limit of PM<sub>10</sub> during the regular dust events in the Canary Islands (≈300 μg/m<sup>3</sup>) and also much higher than the extraordinary 680 μg/m<sup>3</sup> recorded during the dust event of 26 January 2000 (Viana et al., 2002). During this dx-02 episode, the highest 1-hour PM<sub>10</sub> (3500-5254 μg/m<sup>3</sup>) and PM<sub>2.5</sub> (800-1129 μg/m<sup>3</sup>) and 24-h average PM<sub>10</sub> (1200-1840 μg/m<sup>3</sup>) and PM<sub>2.5</sub> (230-404 μg/m<sup>3</sup>) concentrations were recorded in the AQMS located in the central part of the dust plume, in Gran Canaria and Tenerife islands (Fig.5B1-5B2)(Fig.S1). After 24 [R1C9] February 2020 the Saharan dust plume shifted northward over the Atlantic reaching northern mainland [improvement] Spain (Fig.4J), resulting in (24h average) PM<sub>10</sub> concentrations within the range 70-155 μg/m<sup>3</sup> (Fig.4D1) in central Spain (Madrid, Extremadura and Castilla La Mancha regions), 70-75 μg/m<sup>3</sup> (Fig.4D2) in central Portugal [R1C4] and 80-200 μg/m<sup>3</sup> in eastern Spain (Comunidad Valenciana region) (Fig.4E), and PM<sub>2.5</sub> concentrations within the range 20-33 μg/m<sup>3</sup> in central Spain and central Portugal [R1C4] (Fig.4D2), respectively. The re-analysis of MERRA-2 properly tracked these two dx-01 & dx-02 events, with dust and dust<sub>2.5</sub> (i.e. dust in the PM<sub>2.5</sub> fraction) within the range of the PM<sub>10</sub> and PM<sub>2.5</sub> concentrations recorded in the AQMS (see orange circles in Figure 4B-4D).

The third dust event (dx-03: 15-21 Feb 2021) (Fig.6D) was also caused by the intense wind (Fig.6C) linked a dipole L-to-H meteorology (Fig.6B), with the associated blocking anticyclone located over Iberia and a cyclone over the Sahel (Fig.6B). The impact on the Canary Islands occurred during 15-19 February [R1C9] and in Madeira during 16-18 [R1C4] February 2021 (Fig.6C). In the Canaries, the highest 1-hour average PM<sub>10</sub> (1000-1352 μg/m<sup>3</sup>) and PM<sub>2.5</sub> (200-326 μg/m<sup>3</sup>) were recorded in Gran Canaria, Tenerife, La Gomera and La Palma (Fig.S1). The 16 [R1C9] February 2021 resulted in 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 400-711<sup>x</sup> μg/m<sup>3</sup> (Fig.5; Fig.6A1)(<sup>x</sup>Las Galletas AQMS, Tenerife) and 80-205<sup>x</sup> μg/m<sup>3</sup> (Fig.5; Fig.6A2)(<sup>x</sup>Las Galanas, La Gomera), respectively. On 16 February 2021, the highest PM<sub>10</sub> (> 480 μg/m<sup>3</sup>) and PM<sub>2.5</sub> (>120 μg/m<sup>3</sup>) concentrations were recorded on the western side of the Canary Islands (Fig. 5C1-5C2) [improvement]. Subsequently, the dusty air mass tracked the northward anticyclonic circulation, resulting in (24h average) PM<sub>10</sub> concentrations within 80-180 μg/m<sup>3</sup> in Madeira (Fig. 6A3) [R1C4], reaching central mainland Portugal [R1C4] and Spain (18-21 Feb 2021; Fig.6E), and resulting in 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 75-150 μg/m<sup>3</sup> (Fig.6A4) and 20-55

$\mu\text{g}/\text{m}^3$  (Fig. 6A4-6A5), respectively. The MERRA-2 reanalysis properly tracked the dx-03 event, except during 16-17 of February in Madeira [RIC9] and 21 of [RIC9] February in central mainland Spain, when it clearly over estimated dust concentrations (see orange circles in Figure 6A1-6A4). A few days later, 23-28 [RIC9] February 2021, another dust event impacted across central to northern and eastern Europe due to eastward shift the L-to-H dipole and the resulting northward dust transport across the central Mediterranean (Meinander et al., 2023; Peshev et al., 2023).




Figure 4. Events dx-01 and dx-02. Satellite view (NOAA-20 VIIRS) of the dust plume during the first day of dx-01 (A1) and dx-02 (A2) events. Time series of (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> data recorded in AQMS of the Canary Islands (B1-B2), Madeira island (C) [RIC4], central Portugal and Spain (D1-D2) and eastern Spain (E), which includes dust and dust<sub>2.5</sub> concentrations (μg/m<sup>3</sup>) obtained with MERRA-2 model (orange circle) in each region (27-29°N, 15-17.5°W domain for the Canary Islands, 32-34°N, 16-18°W for Madeira island [RIC4], 39-41°N, 9.2-4.3°W for central Portugal and Spain and 37-41.5°N, 6.8°W-1.2°E for eastern Spain). The geopotential height (GPH) of the 850hPa is shown for the first day of dx-01 (E) and dx-2 (G) events. Wind vector at 850hPa for the 23 Feb 2020 dx-02 (E). MERRA-2 column dust load for the 04 Feb 2020 (F), 23 Feb 2020 (I) and 28 Feb 2020 (J).





Figure 5. Surface dust and dust<sub>2.5</sub> concentrations of MERRA-2 reanalysis (26.5-30.0°N, 19.3-12.0°W) and observed (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> measured in AQMS during specific days of dx-01, dx-02, dx-03, dx-04 and dx-05 events in the Canary Islands.





5 **Figure 6. Event dx-03. Time series (A1-A4): (1) of (24h average)  $PM_{10}$  and  $PM_{2.5}$  in AQMS of the Canary Islands, Madeira, central mainland Portugal [RIC4] and central mainland Spain, and (2) of dust and dust<sub>2.5</sub> concentrations ( $\mu\text{g}/\text{m}^3$ ) obtained with MERRA-2 in the Canary Islands (27-29°N, 15-17.5°W), Madeira island (32-34°N, 16-18°W) [RIC4], central mainland Portugal and Spain (39-41°N, 9.2-4.3°W). The geopotential height (GPH) of the 850hPa level (B, [RIC13] 15 feb 2021), wind at 1000hPa (C, [RIC13] 16 feb 2021), the satellite image (NOAA-20 VIIRS)(D, [RIC13] 16 feb 2021) and the column dust load (E, [RIC13] 18 feb 2021) are included.**

Another two extreme dust events occurred in January 2022 (Fig.7). The dx-04 event impacted on the Canary Islands during 14-17 [RIC9] January 2022, resulting in (24h average)  $PM_{10}$  and  $PM_{2.5}$  concentrations within the range 275-883 $\times$   $\mu\text{g}/\text{m}^3$  (Fig.7A1; Fig.5D1)( $\times$ Tenerife, Casa Cuna AQMS) and 60-136 $\times$   $\mu\text{g}/\text{m}^3$  (Fig.7A2; Fig.5D2) ( $\times$ Tenerife, Caletilla AQMS) and 1-hour average  $PM_{10}$  and  $PM_{2.5}$  concentrations reaching values within 1200-2170  $\mu\text{g}/\text{m}^3$  and 240-550  $\mu\text{g}/\text{m}^3$  across the Canaries (Fig.S1) [improvement], respectively. A few days later, the dx-05 occurred (29 Jan - 1 Feb 2022) impacting again the Canary Islands, resulting in (24h average)  $PM_{10}$  and  $PM_{2.5}$  concentrations within the range 314-1055 $\times$   $\mu\text{g}/\text{m}^3$  (Fig.7A1; Fig.5E1)( $\times$ Fuerteventura, El Charco AQMS) and 70-199 $\times$   $\mu\text{g}/\text{m}^3$  (Fig.7A2; Fig.5E2) ( $\times$ El Charco AQMS) and 1-hour average

15

PM<sub>10</sub> and PM<sub>2.5</sub> concentrations reaching values within 1000-2520 µg/m<sup>3</sup> and 400-545 µg/m<sup>3</sup> in the eastern islands (Fig.S1 [improvement]), respectively. In Madeira, PM<sub>10</sub> concentrations ranged within 80-225 µg/m<sup>3</sup> during dx-04 and dx-05 (Fig.7B3 [RIC4]). In both cases a massive dust plume was transported northward, approaching to Spain and Portugal (Fig.7D1-7D2). As in the previous cases, these dust episodes were caused by a L-to-H dipole meteorology, with an anticyclonic core over Europe expanding to North Africa and a cyclone south of the Canary Islands (Fig.7C1-7C2). In the two events, MERRA-2 clearly overestimated dust concentrations (Fig.7B1-7B2).




Figure 7. Events dx-04 and dx-05. Satellite view (NOAA-20 VIIRS) of the dust plumes (A1-A3). Time series of (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> in AQMS of the Canary Islands (B1-B2) and Madeira island (B3) [RIC4] and MERRA-2 surface dust and dust<sub>2.5</sub> concentrations in the Canary Islands (27-29°N, 15-17.5°W) and Madeira (32-34°N, 16-18°W) [RIC4]. The geopotential height (GPH) of the 850hPa level (C1-C2) and column dust load (D1-D2) are included.

5

Finally, the sixth dust event (dx-06: 15-20 March 2022) first impacted mainland Spain and Portugal [RIC4] and subsequently the Canary Islands (Fig.8). The event was also prompted by a meteorological L-to-H dipole, linked to the location of the cyclone *Celia* over Morocco and an anticyclonic core over the central Mediterranean (Fig. 8B). The resulting dusty jet (Fig. 8C-8E) expanded from southeast to northwest mainland Spain and Portugal [RIC4] on the 15 and 16 March 2022. Once in northern Spain, the dust plume split in two branches, a branch travelled eastward across central Europe tracking the

10

anticyclonic circulation at the north of the high-H (Qor-El-Aine et al., 2022), the other branch travelled southward over mainland Portugal [R1C4] to the Canary Islands tracking the cyclonic L circulation (Fig.8E-8F).




Figure 8. Event dx-06. Time series of (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> in AQMS of the Canary Islands (blue time series) and different regions of mainland Spain (green time series) and mainland Portugal (black time series): A1-A2) Northern (N) Spain (Cantabria and Galicia regions) and Portugal (Norte). A3-A4) Central-Northern (CN) Spain (Castilla y León region) and Portugal (Norte); A5-A6) Central (C) Spain (Madrid + Extremadura + Castilla La Mancha) and Portugal (Centro + north Alentejo); A7-A8) Southern (S) Spain (Andalucía) and Portugal (south Alentejo and Algarve); A9-A10) Canary Islands. [RIC4] The plots also include surface dust concentrations in these regions obtained with MERRA-2 reanalysis (A1-A10). The geopotential height (GPH) (B) and wind (C) in the 850hPa level, the satellite NOAA-20 VIIRS image and column dust load (E-G) is also included.




Figure 9. Surface concentrations of dust (A1-A3) and dust particles smaller than 2.5 microns (dust<sub>2.5</sub>) (B1-B3) from 14 to 16 March 2022 in mainland Spain according to MERRA-2 reanalysis. Daily (24h) average concentrations of PM<sub>10</sub> (A1-A2) and PM<sub>2.5</sub> (B1-B2) measured in AQMS are shown with black numbers.

In Iberia mainland Spain [improvement], PM<sub>x</sub> concentrations experienced a sharp increase, from their regular background levels (10-30 µg/m<sup>3</sup>) to 24h average PM<sub>10</sub> and PM<sub>2.5</sub> values within the range (i) 500-3070 µg/m<sup>3</sup> and 100-700 µg/m<sup>3</sup> in southern regions of Spain and Portugal (Murcia, Andalucía, Algarve and Alentejo) [R1C4] (Fig.8A7, 8A8, 9), (ii) 200-1000 µg/m<sup>3</sup> and 60-160 µg/m<sup>3</sup> in central parts of Spain and Portugal (Castilla La Mancha, Madrid, Extremadura, Vale do Tejo and Lisboa) [R1C4] (Fig.8A5, 8A6, 9), (iii) 200-1000 µg/m<sup>3</sup> and 60-260 µg/m<sup>3</sup> in central northern Spain (Centro, Castilla y León) (Fig.8A3, 8A4, 9), and (iv) 150-500 µg/m<sup>3</sup> and 75-130 µg/m<sup>3</sup> in northern Portugal and Spain (Norte, Cantabria and Galicia) (Fig.8A1, 8A2, 9) during the 15-16 March 2022, respectively. In Lisboa, these extremely high PM<sub>10</sub> and PM<sub>2.5</sub> concentrations were even registered in the indoors environment (Gomes et al., 2022) [R1C4]. After several days traveling across thousands of kilometres, the dust plume impacted in the Canary Islands during 17-20 March 2022 (Fig.8F), resulting in (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> values within the range 150-430 µg/m<sup>3</sup> (Fig.8A9) and 30-80 µg/m<sup>3</sup> (Fig.8A10). A regular (no extreme) dust event impacted southern Spain 24-25 March 2022 (PM<sub>10</sub> and PM<sub>2.5</sub>: 50-420 µg/m<sup>3</sup> and 30-120 µg/m<sup>3</sup>; respectively) (Fig.8A7-8A8) and the Canary Islands 24-25 March 2022 (PM<sub>10</sub> and PM<sub>2.5</sub>: 40-80 and 20-35 µg/m<sup>3</sup>, respectively).

This historic dx-06 event (Fig.1D-1E) started the evening of 14 March 2022 (>21h), with a dust inflow in south-eastern Spain that led to 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 70-260 µg/m<sup>3</sup> and 25-43 µg/m<sup>3</sup> (Fig.9A1-9B1). On 15 March 2022, the massive dust plume moved from south-eastern Spain, where it led to 24h average PM<sub>10</sub> values within the range 580-3070 µg/m<sup>3</sup>, toward the west and northwest of Iberia, resulting in (24h average) PM<sub>10</sub> concentrations within the ranges 825-950 µg/m<sup>3</sup> in central Spain, 600-650 µg/m<sup>3</sup> in central Portugal and 440-810 µg/m<sup>3</sup> in northern Portugal and northwest Spain (Fig.9A2). Concentrations of PM<sub>2.5</sub> (24h average) were within the range 95-160 µg/m<sup>3</sup> in south-eastern Spain, 40-70 µg/m<sup>3</sup> southern Portugal and south-western Spain and 60-145 µg/m<sup>3</sup> in central-north Portugal and north-western Spain (Fig.9B3). These extremely high PM<sub>x</sub> concentrations were also recorded indoor (Gomes et al., 2022). During the 16 March 2022, high PM<sub>10</sub> and PM<sub>2.5</sub> values were still recorded (Fig.9A3-9B3), with the highest PM<sub>10</sub> concentrations linked to the still ongoing dust inflow by south-eastern Spain (800 µg/m<sup>3</sup>) and the southward transport of dust in southern Portugal (300-330 µg/m<sup>3</sup>). Because of the massive dust load the solar energy production in Spain dropped in a 50% (Micheli et al., 2024). Details on this event, as the maximum hourly PM<sub>10</sub> concentrations (Fig.S2) and the names of the AQMS, are provided in the Supplement [R2C5]

This historic dx-06 event (Fig.1D-1E) started the evening of 14 March 2022 (>21h), with a dust inflow in south-eastern Spain that led to 24h average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the range 70-260 µg/m<sup>3</sup> and 25-43 µg/m<sup>3</sup> (Fig.9A1-9B1). On 15 March 2022, mainland Spain was blanketed by the massive dusty air mass (Fig.9A2-9B2), which led to 24h average PM<sub>10</sub> values of: (i) 3069 µg/m<sup>3</sup> (Mediterraneo AQMS; Almería province), 1586 µg/m<sup>3</sup> (El Ejido; Almería), 848 µg/m<sup>3</sup>

(Ronda del Valle; Jaén), 621  $\mu\text{g}/\text{m}^3$  (Palacio de Congresos; Granada) and 587  $\mu\text{g}/\text{m}^3$  (Mompean; Murcia) in south-eastern Spain and 518  $\mu\text{g}/\text{m}^3$  in central-southern Spain (Sierra Norte; Seville), (ii) 949  $\mu\text{g}/\text{m}^3$  (Villa del Prado; Madrid), 944\*  $\mu\text{g}/\text{m}^3$  (Segovia 2; Segovia), 928  $\mu\text{g}/\text{m}^3$  (Medina del Campo; Valladolid), 841\*  $\mu\text{g}/\text{m}^3$  (Avila II; Avila) and 825  $\mu\text{g}/\text{m}^3$  (Salamanca-6; Salamanca) in central Spain, and (iii) 808\*  $\mu\text{g}/\text{m}^3$  (Lario; León), 562  $\mu\text{g}/\text{m}^3$  (León-01; León), 472  $\mu\text{g}/\text{m}^3$  (Laza; Orense) and 371  $\mu\text{g}/\text{m}^3$  (Los Tojos; Cantabria) in Norwest Spain (Fig.9A2). Some of these PM<sub>10</sub> concentrations (highlighted with asterisk\*) are actually slightly underestimated due to the automatic PM<sub>10</sub> monitor reached the saturation level (1000  $\mu\text{g}/\text{m}^3$ ) during a few hours and the *duxt r* reconstruction method could not be applied to the lack of parallel PM<sub>2.5</sub> data. On 15 March 2022, the 24h average PM<sub>2.5</sub> values reached values of (Fig.9B2): (i) 688  $\mu\text{g}/\text{m}^3$  (Mediterraneo; Almería), 164  $\mu\text{g}/\text{m}^3$  (Ronda del Valle; Jaén), 158  $\mu\text{g}/\text{m}^3$  (Palacio de Congresos; Granada) in south-eastern Spain, (ii) 252  $\mu\text{g}/\text{m}^3$  (Salamanca-6), 153  $\mu\text{g}/\text{m}^3$  (Medina del Campo; Valladolid) and 119  $\mu\text{g}/\text{m}^3$  (Villa del Prado; Madrid) in central Spain and (iii) 129  $\mu\text{g}/\text{m}^3$  (Laza; Orense) in north west Spain. Because of the massive dust load the national photovoltaic capacity of Spain dropped in a 50%. (Michel et al., 2023 <https://doi.org/10.1016/j.seta.2024.103607>). During the 16 March 2022, high PM<sub>10</sub> and PM<sub>2.5</sub> values were still recorded (Fig.9A3-9B3), with the highest PM<sub>10</sub> concentrations linked to the still ongoing dust inflow by south-eastern Spain: 800  $\mu\text{g}/\text{m}^3$  (Bailén; Jaén), 774  $\mu\text{g}/\text{m}^3$  (Ronda del Valle; Jaén), 535  $\mu\text{g}/\text{m}^3$  (Mediterraneo; Almería) and 432  $\mu\text{g}/\text{m}^3$  (El Ejido; Almería). During the dx-06 event, the hourly PM<sub>10</sub> concentrations exceeded 2000  $\mu\text{g}/\text{m}^3$  at four sites which reached maximum PM<sub>10</sub> values: >4573\*  $\mu\text{g}/\text{m}^3$  (Mediterraneo; Almería), 3502  $\mu\text{g}/\text{m}^3$  (El Ejido; Almería), 2813  $\mu\text{g}/\text{m}^3$  (Mompean; Murcia) and 2154  $\mu\text{g}/\text{m}^3$  (Ronda del Valle; Jaén) (Fig.2C3). Similarly, the hourly values of PM<sub>2.5</sub> exceeded 400  $\mu\text{g}/\text{m}^3$  at four sites, where the maximum 1h PM<sub>2.5</sub> concentration were: 1000\*  $\mu\text{g}/\text{m}^3$  (Mediterraneo; Almería, PM<sub>2.5</sub> analyzer saturated), 471  $\mu\text{g}/\text{m}^3$  (Mompean; Murcia), 443  $\mu\text{g}/\text{m}^3$  (Salamanca-6), 400 (Ronda del Valle; Jaén). On 15 March, hourly PM<sub>2.5</sub> concentrations at Mediterraneo AQMS saturated at 1000  $\mu\text{g}/\text{m}^3$  during 6 hours (08-13h), so the daily average PM<sub>2.5</sub> value = 688  $\mu\text{g}/\text{m}^3$  at this site is actually somewhat underestimated and consequently also the daily average PM<sub>10</sub> value = 3069  $\mu\text{g}/\text{m}^3$  (obtained with the *duxt r* method). [R2C5] this paragraph was transferred to the Supplement.

### 3.3 Record breaking events

25 The analysis of the 2000-2022 time series of (24h average) PM<sub>10</sub> (Fig.10) and PM<sub>2.5</sub> (Fig.11) data, evidence that the *duxt r* events we report here are record beating episodes in mainland Spain, continental Portugal [R1C4] and the Canary Islands. The massive dusty airmass that blackened the Iberian Peninsula mainland Spain during the 15 and 16 of March 2022 (dx-06; Fig.10 and 11) resulted in the highest PM<sub>10</sub> and PM<sub>2.5</sub> concentrations ever recorded in the regional scale across northern Spain (Cantabria region; Fig.10A), central - northern Spain (Castilla y León region; Fig. 10C & 11A), central Spain (Castilla La Mancha, Extremadura and Madrid region; Fig.10D & 10A), southern Spain (Andalucía region; Fig. 10E, 10F & 11C) and continental Portugal (Fig.10G) [R1C4]. For example, [improvement] In central Spain (Castilla La Mancha + Extremadura +



Madrid), regular Saharan dust events typically induce (24h average)  $PM_{10}$  concentrations within the range 40-140  $\mu\text{g}/\text{m}^3$  (highlighted with black arrows in Fig.10D) (Pey et al., 2013; Rodríguez et al., 2001), **anomalous intense dust events as that occurred 22 February 2016 (Sorribas et al., 2017) resulted in  $PM_{10}$  concentrations 200-380  $\mu\text{g}/\text{m}^3$  [improvement]**, whereas during the dx-06 event 33 AQMS of this region recorded (24h)  $PM_{10}$  concentration within 300-949<sup>x</sup>  $\mu\text{g}/\text{m}^3$  (<sup>x</sup>Villa del Prado AQMS in Madrid region)(white arrow in Fig. 10D). After analysing the 2001-2011 time series, Pey et al.(2013) concluded that in the Western Mediterranean, Saharan dust events inducing  $PM_{10} > 100 \mu\text{g}/\text{m}^3$  (24h average) are actually rare. The impact of the dx-06 event is not observed in Cataluña since it did not reach Northeast Spain (Fig.10B).





Figure 10. Time series (2000-2022) of (24h average)  $PM_{10}$  concentrations in 123 AQMS distributed across Portugal (5 AQMS), northern (N; 3 AQMS), North-East (NE; 8 AQMS), Central-North (CN; 30 AQMS), Central (C; 35 AQMS), Southwest (SW; 16 AQMS) and Southeast (SE; 11 AQMS) mainland Spain and in the Canary Islands (15 AQMS). Black arrows indicate regular dust events. White arrows indicate the dust events. The asterisk indicates the intense event occurred 22 Feb 2016.



**Figure 11. Time series (2000-2022) of (24h average)  $PM_{2.5}$  concentrations of in a total of 74 AQMS distributed across the Central-North (CN; 4 AQMS), Central (C; 16 AQMS) and Southeast (S; 10 AQMS) mainland Spain and in the Canary Islands (44 stations). Black arrows indicate regular dust events. White arrows indicate the duxt events.**

In the Canary Islands, the most intense Saharan dust events ever recorded occurred in the period 2020-2022 linked to the dust events described in this study. Since 2005 to 2020, intense Saharan dust events have regularly been associated with (24h average) PM<sub>10</sub> and PM<sub>2.5</sub> values with 200-400 µg/m<sup>3</sup> (Fig.10H) and 100-200 µg/m<sup>3</sup> (Fig.11D), respectively. This PM<sub>10</sub> range is similar to (1) that of total suspended particles during Saharan dust events of the period 1998-2003, based on AQMS data not yet normalised to the European standards (Alonso-Pérez et al., 2007, 2012; Viana et al., 2002), and to (2) that of total dust at Izaña Observatory (Tenerife) during the 1987-2014 Saharan dust events (Rodríguez et al., 2015). In contrast, in the period 2020-2022, the dust events described here led to PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the ranges (24h average) 600-1840 µg/m<sup>3</sup> (Fig.10G) and 200-404 µg/m<sup>3</sup> (Fig.11D), respectively. The three most intense dust events ever recorded in the Canary Islands, exceeding the threshold of 600 µg/m<sup>3</sup> of PM<sub>10</sub> as 24h average, in descending order of magnitude, are:

1. dx-02 event: the 23 and 24 [R1C9] February 2020 the (24h average) PM<sub>10</sub> averaged in all AQMS were 531 and 930 µg/m<sup>3</sup> (averages of 34 AQMS distributed in the 7 Canary Islands), respectively. The 22, 23 and 24 [R1C9] February 2020, a total of 6, 25 and 12 AQMS recorded a (24h average) PM<sub>10</sub> concentration between 600 and 1840<sup>x</sup> µg/m<sup>3</sup> (<sup>x</sup>Gran Canaria, Playa del Inglés AQMS). Previous to this event, the (24h average) PM<sub>10</sub> concentrations had only exceeded 600 µg/m<sup>3</sup> in just one AQMS (618 µg/m<sup>3</sup> at Las Galanas during the dust event 28 March 2018; Fig.10G).
2. dx-05 event: 29 and 30 [R1C9] January 2022 the (24h average) PM<sub>10</sub> averaged in all the AQMS was 463 and 501 µg/m<sup>3</sup> (averages of 44 AQMS distributed in the 7 islands). The 29 and 30 of Jan 2022, a total of 9 and 10 AQMS recorded a (24h average) PM<sub>10</sub> concentration between 600 and 1055<sup>x</sup> µg/m<sup>3</sup> (<sup>x</sup>Fuerteventura, El Charco AQMS).
3. dx-03 event: the 16 [R1C9] February 2021 the (24h average) PM<sub>10</sub> averaged in all AQMS was 463 µg/m<sup>3</sup> (average of 36 AQMS distributed in the 7 islands). The 16 [R1C9] February 2021, 4 AQMS recorded a (24h average) PM<sub>10</sub> concentration between 600 and 638<sup>x</sup> µg/m<sup>3</sup> (<sup>x</sup>Las Galletas AQMS, Tenerife).

In mainland Spain and continental Portugal [R1C4], the dx-06 is the most intense dust event ever recorded. In Spain total of 20 AQMS, distributed across south-eastern, central to central-northern Spain, registered (24h average) PM<sub>10</sub> concentrations within the range 600 to 3070<sup>x</sup> µg/m<sup>3</sup> (<sup>x</sup>Mediterraneo AQMS in Almería province). In Portugal, 4 AQMS located in the central regions registered (24h average) PM<sub>10</sub> concentrations within the range 600 to 648<sup>x</sup> µg/m<sup>3</sup> (<sup>x</sup>Chamusca AQMS in Lisboa - Vale do Tejo) [R1C4]. The 2 decades time series of PM<sub>x</sub> also offer other interesting data. In many of regions of Spain there is a clear 2000-2020 decreasing trend of PM<sub>10</sub> concentrations linked to the reduction of emissions following air quality policies (Fig.10A, 10C, 10E) (Li et al., 2018; Querol et al., 2014), suggesting that desert dust may have an increasing relative contribution to PM<sub>10</sub> concentrations as also pointed by recent projections (Gomez et al., 2023). [improvement]

### 3.4 Meteorological anomalies linked to dust events

In winter, North African dust is regularly transported southward to tropical latitudes (Merdji et al., 2023). In this season, dust transport northward, to the subtropical North Atlantic and southern Europe, occurs during rather short periods, under specific meteorological scenarios described in previous studies (Flaounas et al., 2015; Fluck and Raveh-Rubin, 2023a; Rodriguez et al., 2001). Winter extreme Saharan dust events have been observed in the southern Sahara – Sahel region, from Mauritania to Niger along the Sahel, associated with PM<sub>10</sub> and PM<sub>2.5</sub> concentrations within the ranges 800 – 5000 µg/m<sup>3</sup> and 600 – 1300 µg/m<sup>3</sup> (Fluck and Raveh-Rubin, 2023b; Marticorena et al., 2010), respectively, induced by H-L dipoles, i.e. an anticyclonic high pressure H core over Atlantic and western North Africa and a cut of low L over the Mediterranean, a meteorological configuration that results in strong southern winds (Fluck and Raveh-Rubin, 2023b). [R2C2][R2C3]

The 2020-2022 six dust events we report here were induced by the L-to-H meteorological dipoles formed by cyclones located at the southwest of a blocking anticyclone over western Europe. Fig. 12 (A1 to F1) show the anomaly of the 500 hPa geopotential height during the onset of the six dust events studied here [improvement] (Fig.12 A2 to F2). All dust events occurred during northern-hemisphere meteorological anomalies that resemble the anomalies of the atmospheric circulation that have been linked, in previous studies, to the anthropogenic global warming [Reply to R2C1] (Fig.12): (i) subtropical anticyclones intensified and expanded and shifted [R2C1] to higher latitudes (Cherchi et al., 2018; Cresswell-Clay et al., 2022), (ii) anomalous low pressures expanding northward beyond the tropical belt, resembling the tropical expansion (Seidel et al., 2008; Yang et al., 2020, 2023) (e.g. dx-01, dx-02, dx-03 and dx-04) and (iii) mid-latitudes amplified undulation of the [improvement] [R2C2] Rossby waves due to the concatenation of cut-off lows cyclones and anticyclones pointing to a weakening of the polar vortex (e.g. dx-03, dx-05, dx-06 and 23-Mar-2022) (Mann et al., 2017; Screen and Simmonds, 2013). The dust events observed at the Eastern Mediterranean in March 2018 (Solomos et al., 2018) and in March 2020 (Mifka et al., 2023) [improvement], at North Africa in June 2020 (Bi et al., 2023; Francis et al., 2020), at Uzbekistan in November 2021 (Xi et al., 2023) and in China in March 2021 (Gui et al., 2022; Liu et al., 2023) [improvement], occurred in this context of cyclones, blocking anticyclones and dipoles linked to mid-latitudes amplified Rossby waves.

The winter blocking anticyclone that we observed over western Europe/Western Mediterranean during the dust events (Fig.12A1-G1) fits with the picture of the industrial-era eastward expansion and shift of the North Atlantic anticyclone starting in the 1850s and accelerating in the last few decades (Alonso-Pérez et al., 2011b; Cresswell-Clay et al., 2022), a trend that is expected to continue as the concentrations of greenhouse gases increase, according to the CMIP5 multi-model simulations (Cherchi et al., 2018); the low pressures at the southwest of the blocking anticyclone that we observe during the dust events (Fig.12B1-12E1) are expected to follow as the tropic expands northward off North Africa in the forthcoming decades (e.g. see Fig 1F of Cherchi et al. 2018 for the 2075-2100 period).

The anomalies linked to the dust events are also evident in the trajectory of the cyclones that finally form the L-to-H dipoles leading to the extreme dust emissions. Red circles in Fig.12A1-12G1 indicate the location of the cyclones from days before

(negative number) to days after (positive number) of the dust event. Due to the anticyclonic blocking over western Europe, the mid-latitudes North Atlantic cyclones didn't follow their regular path across Europe or the Mediterranean. The cut-off lows forming the L-to-H dipoles reached this region (Canary Islands, Cape Verde or inner Sahara) by two main paths:




Figure 12. Anomaly (A1-G1) of the 500 hPa geopotential height (GPH) (respect to the 1991-2020 climatology) and column dust in the onset of the dx-01 to dx-06 and 24-26 March 2022 events (A2-G2). Red circles (A1-G1) indicate the location of the cyclones days before (negative number) to days after (positive number) the first day of the dust event.

1. they deviated southward from the regular mid-latitudes westerly circulation in the North Atlantic mid-latitudes [improvement] as results of the blocking anticyclone over western Europe; subsequently these cut-off lows may stay during long periods (up to 12 days) in the subtropical and tropical north-east Atlantic (near the Canary Islands and Cape Verde). This is the case of the dipoles of events dx-02, dx-04, dx-05 and dx-06, and also of the intense dust event of 23 March 2022. The cyclones of the events dx-02, dx-04 and dx-05 stayed (blocked by the anticyclonic situation) near the Canary Islands and Cape Verde during 10, 12 and 8 days (Fig. 12B1, 12D1 and 12E1), respectively.
2. they deviated northward from the tropical belt. This is the case of the events dx-01 and dx-03, associated with cyclones which had moved from Cape Verde to the west of the Canary Islands (Fig.12A1, dx-01) and to the inner Sahara (Fig.12C1, dx-03), respectively, across a tropical band anomalously shifted to northward.

The observed sharp increase in dust transport to the western Euro-Mediterranean region in the 2020-2022 winters has been also been associated to the meteorology linked to dipoles and blocking anticyclones (Cuevas et al., 2023).

#### 15 4 Summary and conclusions

In winter 2020, 2021 and 2022 a set of six extreme dust events expanded northward from NW Africa to the Atlantic and Europe, causing extremely high concentrations of PM<sub>10</sub> and PM<sub>2.5</sub> in the Governmental Air Quality Monitoring Stations of the Canary Islands, mainland Spain and continental Portugal [improvement], up to [improvement] exceeding the upper operation limit of many PM<sub>10</sub> monitors. We developed the *duxt-r* methodology for assessing the consistency of the PM<sub>10</sub> and PM<sub>2.5</sub> data and re-construct the underestimated PM<sub>10</sub> concentrations. During these extreme dust events, the 1-hour average PM<sub>10</sub> and PM<sub>2.5</sub> concentrations were within the range 1000-6000 µg/m<sup>3</sup> and 400-1200 µg/m<sup>3</sup>, whereas the 24-h average PM<sub>10</sub> and PM<sub>2.5</sub> data were within the range 500-3070 µg/m<sup>3</sup> and 200-690 µg/m<sup>3</sup>, respectively. These extreme dust episodes are were caused by the intense winds associated with the meteorological dipoles formed by a blocking anticyclone over western Europe and a cut-off low cyclone [improvement] located at the southwest, near the Canary Islands, Cape Verde or into the Sahara. The analysis of the 2000-2022 time series of PM<sub>10</sub> and PM<sub>2.5</sub> shows that these events have no precedent in Spain [improvement]. Record beating breaking PM<sub>10</sub> and PM<sub>2.5</sub> (24h average) concentrations were measured in the Canary Islands during the 22-24 February 2020, 1840 µg/m<sup>3</sup> and 404 µg/m<sup>3</sup>, and in mainland Spain during 15-16 March 2022, 3069 µg/m<sup>3</sup> and 688 µg/m<sup>3</sup> (Almeria province), respectively. All dust events occurred during northern-hemisphere meteorological anomalies associated with subtropical anticyclones shifted intensified [improvement] and expanded to higher latitudes, anomalous low pressures expanding beyond the tropical belt and a concatenation of cut-off lows and anticyclones suggesting to a weakening of the polar vortex, with features that resemble the change in the atmospheric circulation linked to climate change [R2C1]. These events



occurs in a the context in which Climate projections forecast the expansion of the North African drylands toward the northwest, increasing the risk of desertification of Spain and Portugal, with an associated increase in regional dust loads. The air quality monitoring networks need to adapt the strategy and operation range of the PM<sub>10</sub> and PM<sub>2.5</sub> monitoring programs to ensure accurate measurements can be done [improvement] during these extreme dust events due to the importance of having suitable data in the public data sets for health effects studies, modelling, etc... New studies have reported on recent record beating PM<sub>10</sub> and PM<sub>2.5</sub> episodes linked to dipoles induced extreme dust events from North Africa and Asia, in a paradoxical context of multidecadal decrease of dust emissions, a topic that will require further investigations. [R2C1]

### Data availability

10 The data used in the manuscript are available in public data bases, including the PM<sub>10</sub> data reconstructed as result of this study. Data are also available at DIGITAL CSIC (<http://hdl.handle.net/10261/364553>) and by request to the first author at [sergio.rodriguez@csic.es](mailto:sergio.rodriguez@csic.es) [improvement]

### Author contributions

15 SR and JLD performed the conceptualisation, investigation, data collection, treatment and formal analysis. SR wrote the original version of the manuscript, which was subsequently revised and edited by SR and JLD.

### Competing interests

~~The authors declare that they have no conflict of interest.~~ One of the authors [SR] is editor in the journal Atmospheric Chemistry and Physics.

### 20 Acknowledgments

We thanks to the 17 Autonomous Regions of Spain and the Ministry for the Ecological Transition and the Demographic Challenge for the kind supply to the data of the air quality monitoring stations. [R1C4] We also thanks to the Agência Portuguesa do Ambiente for supplying the air quality data of Portugal. We acknowledgement the access to modelling reanalysis data by the portal web sites of NASA - Giovanni and NOAA – Physical Science Laboratory. [R1C5] Satellite images were

provided by the NASA World View website. We also thanks to Jon Vilches (air quality management in the Government of the Canary Islands) for the interesting talks on the PM<sub>x</sub> monitoring devices.

### **Financial support**

This study is part of the project AERO-EXTREME (PID2021-125669NB-I00) funded by the ~~funded by the~~ State Research  
5 Agency/Agencia Estatal de Investigación of Spain and the European Regional Development Funds.

## References

- Alonso-Pérez, S., Cuevas, E., Querol, X., Viana, M. and Guerra, J. C.: Impact of the Saharan dust outbreaks on the ambient levels of total suspended particles (TSP) in the marine boundary layer (MBL) of the Subtropical Eastern North Atlantic Ocean, *Atmos. Environ.*, 41(40), 9468–9480, doi:10.1016/j.atmosenv.2007.08.049, 2007.
- Alonso-Pérez, S., Cuevas, E. and Querol, X.: Objective identification of synoptic meteorological patterns favouring African dust intrusions into the marine boundary layer of the subtropical eastern north Atlantic region, *Meteorol. Atmos. Phys.*, 113(3–4), 109–124, doi:10.1007/s00703-011-0150-z, 2011a.
- Alonso-Pérez, S., Cuevas, E., Perez, C., Querol, X., Baldasano, J. M., Draxler, R. and De Bustos, J. J.: Trend changes of African air mass intrusions in the marine boundary layer over the subtropical Eastern North Atlantic region in winter, *Tellus B*, 63(2), 255–265, doi:10.1111/j.1600-0889.2010.00524.x, 2011b.
- Alonso-Pérez, S., Cuevas, E., Querol, X., Guerra, J. C. and Pérez, C.: African dust source regions for observed dust outbreaks over the Subtropical Eastern North Atlantic region, above 25°N, *J. Arid Environ.*, 78, 100–109, doi:10.1016/j.jaridenv.2011.11.013, 2012.
- Bi, H., Chen, S., Zhang, D., Wang, Y., Kang, L., Alam, K., Tang, M., Chen, Y., Zhang, Y. and Wang, D.: The Circum-global Transport of Massive African Dust and its Impacts on the Regional Circulation in Remote Atmosphere, *Bull. Am. Meteorol. Soc.*, doi:10.1175/BAMS-D-23-0072.1, 2023.
- Cañadillas-Ramallo, D., Moutaoikil, A., Shephard, L. E. and Guerrero-Lemus, R.: The influence of extreme dust events in the current and future 100% renewable power scenarios in Tenerife, *Renew. Energy*, 184, 948–959, doi:10.1016/j.renene.2021.12.013, 2022.
- Cherchi, A., Ambrizzi, T., Behera, S., Freitas, A. C. V., Morioka, Y. and Zhou, T.: The Response of Subtropical Highs to Climate Change, *Curr. Clim. Chang. Reports*, 4(4), 371–382, doi:10.1007/s40641-018-0114-1, 2018.
- Cresswell-Clay, N., Ummenhofer, C. C., Thatcher, D. L., Wanamaker, A. D., Denniston, R. F., Asmerom, Y. and Polyak, V. J.: Twentieth-century Azores High expansion unprecedented in the past 1,200 years, *Nat. Geosci.*, 15(7), 548–553, doi:10.1038/s41561-022-00971-w, 2022.
- Cuevas, E., Milford, C., Barreto, A., Bustos, J. J., García, R. D., Marrero, C. L., Prats, N., Bayo, C., Ramos, R., Terradellas, E., Suárez, D., Rodríguez, S., de la Rosa, J., Vilches, J., Basart, S., Werner, E., López-Villarrubia, E., Rodríguez-Mireles, S., Pita Toledo, M. L., González, O., Belmonte, J., Puigdemunt, R., Lorenzo, J. A., Oromí, P. and del Campo-Hernández, R.: Desert Dust Outbreak in the Canary Islands (February 2020): Assessment and Impacts. (Eds. Cuevas, E., Milford, C. and

- Basart, S.), State Meteorological Agency (AEMET), Madrid, Spain and World Meteorological Organization, Geneva, Switzerland, WMO Global At. [online] Available from: <http://hdl.handle.net/20.500.11765/12703>, 2021.
- Cuevas, E., Barriopedro, D., Delia García, R., Alonso-Pérez, S., González-Alemán, J., Werner, E., Suárez, D., Bustos, J., García-Castrillo, G., García, O., Barreto, A. and Basart, S.: Sharp increase of Saharan dust intrusions over the Western Mediterranean and Euro-Atlantic region in winters 2020-2022 and associated atmospheric circulation, *Atmos. Chem. Phys. Discuss.*, doi:<https://doi.org/10.5194/egusphere-2023-1749>, 2023.
- Domínguez-Rodríguez, A., Báez-Ferrer, N., Abreu-González, P., Rodríguez, S., Díaz, R., Avanzas, P. and Hernández-Vaquero, D.: Impact of Desert Dust Events on the Cardiovascular Disease: A Systematic Review and Meta-Analysis, *J. Clin. Med.*, 10(4), 727, doi:10.3390/jcm10040727, 2021.
- 10 Dumont, M., Tuzet, F., Gascoïn, S., Picard, G., Kutuzov, S., Lafaysse, M., Cluzet, B., Nheili, R. and Painter, T. H.: Accelerated Snow Melt in the Russian Caucasus Mountains After the Saharan Dust Outbreak in March 2018, *J. Geophys. Res. Earth Surf.*, 125(9), doi:10.1029/2020JF005641, 2020.
- Evan, A. T., Flamant, C., Gaetani, M. and Guichard, F.: The past, present and future of African dust, *Nature*, 531(7595), 493–495, doi:10.1038/nature17149, 2016.
- 15 Filonchik, M. and Peterson, M.: Development, progression, and impact on urban air quality of the dust storm in Asia in March 15–18, 2021, *Urban Clim.*, 41, 101080, doi:10.1016/j.uclim.2021.101080, 2022.
- Flaounas, E., Kotroni, V., Lagouvardos, K., Kazadzis, S., Gkikas, A. and Hatzianastassiou, N.: Cyclone contribution to dust transport over the Mediterranean region, *Atmos. Sci. Lett.*, 16(4), 473–478, doi:10.1002/asl.584, 2015.
- Fluck, E. and Raveh-Rubin, S.: A 16-year climatology of the link between dry air intrusions and large-scale dust storms in North Africa, *Atmos. Res.*, 292, 106844, doi:10.1016/j.atmosres.2023.106844, 2023a.
- 20 Fluck, E. and Raveh-Rubin, S.: Dry air intrusions link Rossby wave breaking to large-scale dust storms in Northwest Africa: Four extreme cases, *Atmos. Res.*, 286, 106663, doi:10.1016/j.atmosres.2023.106663, 2023b.
- Francis, D., Fonseca, R., Nelli, N., Cuesta, J., Weston, M., Evan, A. and Temimi, M.: The Atmospheric Drivers of the Major Saharan Dust Storm in June 2020, *Geophys. Res. Lett.*, 47(24), doi:10.1029/2020GL090102, 2020.
- 25 Francis, D., Nelli, N., Fonseca, R., Weston, M., Flamant, C. and Cherif, C.: The dust load and radiative impact associated with the June 2020 historical Saharan dust storm, *Atmos. Environ.*, 268, 118808, doi:10.1016/j.atmosenv.2021.118808, 2022.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.

- D., Sienkiewicz, M. and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Clim.*, 30(14), 5419–5454, doi:10.1175/JCLI-D-16-0758.1, 2017.
- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C. and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, *Rev. Geophys.*, 50(3),  
5 doi:10.1029/2012RG000388, 2012.
- Gomes, J., Esteves, H. and Rente, L.: Influence of an Extreme Saharan Dust Event on the Air Quality of the West Region of Portugal, *Gases*, 2(3), 74–84, doi:10.3390/gases2030005, 2022.
- Gomez, J., Allen, R. J., Turnock, S. T., Horowitz, L. W., Tsigaridis, K., Bauer, S. E., Oliv  , D., Thomson, E. S. and Ginoux, P.: The projected future degradation in air quality is caused by more abundant natural aerosols in a warmer world, *Commun. Earth Environ.*, 4(1), 22, doi:10.1038/s43247-023-00688-7, 2023.  
10
- Govarchin-Ghale, Y. A., Tayanc, M. and Unal, A.: Dried bottom of Urmia Lake as a new source of dust in the northwestern Iran: Understanding the impacts on local and regional air quality, *Atmos. Environ.*, 262, 118635, doi:10.1016/j.atmosenv.2021.118635, 2021.
- Gui, K., Yao, W., Che, H., An, L., Zheng, Y., Li, L., Zhao, H., Zhang, L., Zhong, J., Wang, Y. and Zhang, X.: Record-breaking  
15 dust loading during two mega dust storm events over northern China in March 2021: aerosol optical and radiative properties and meteorological drivers, *Atmos. Chem. Phys.*, 22(12), 7905–7932, doi:10.5194/acp-22-7905-2022, 2022.
- Guiot, J. and Cramer, W.: Climate change: The 2015 Paris Agreement thresholds and Mediterranean basin ecosystems, *Science (80-. )*, 354(6311), 465–468, doi:10.1126/science.aah5015, 2016.
- Jiang, Y., Miao, Y., Zhao, Y., Liu, J. and Gao, Y.: Extreme-wind events in China in the past 50 years and their impacts on  
20 sandstorm variations, *Front. Earth Sci.*, 10, doi:10.3389/feart.2022.1058275, 2023.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Hig-gins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R. and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.
- Kaskaoutis, D. G., Rashki, A., Francois, P., Dumka, U. C., Houssos, E. E. and Legrand, M.: Meteorological regimes  
25 modulating dust outbreaks in southwest Asia: The role of pressure anomaly and Inter-Tropical Convergence Zone on the 1–3 July 2014 case, *Aeolian Res.*, 18, 83–97, doi:10.1016/j.aeolia.2015.06.006, 2015.
- Kaskaoutis, D. G., Houssos, E. E., Rashki, A., Bartzokas, A., Legrand, M., Francois, P. and Kambezidis, H. D.: Modulation of Atmospheric Dynamics and Dust Emissions in Southwest Asia by the Caspian Sea—Hindu Kush Index, pp. 941–947., 2017.

- Kaskaoutis, D. G., Rashki, A., Dumka, U. C., Mofidi, A., Kambezidis, H. D., Psiloglou, B. E., Karagiannis, D., Petrinoli, K. and Gavriil, A.: Atmospheric dynamics associated with exceptionally dusty conditions over the eastern Mediterranean and Greece in March 2018, *Atmos. Res.*, 218, 269–284, doi:10.1016/j.atmosres.2018.12.009, 2019.
- Katra, I.: Soil Erosion by Wind and Dust Emission in Semi-Arid Soils Due to Agricultural Activities, *Agronomy*, 10(1), 89, 5 doi:10.3390/agronomy10010089, 2020.
- Kok, J. F., Storelvmo, T., Karydis, V. A., Adebisi, A. A., Mahowald, N. M., Evan, A. T., He, C. and Leung, D. M.: Mineral dust aerosol impacts on global climate and climate change, *Nat. Rev. Earth Environ.*, 4(2), 71–86, doi:10.1038/s43017-022-00379-5, 2023.
- Kutuzov, S., Legrand, M., Preunkert, S., Ginot, P., Mikhalevko, V., Shukurov, K., Poliukhov, A. and Toropov, P.: The Elbrus 10 (Caucasus, Russia) ice core record – Part 2: history of desert dust deposition, *Atmos. Chem. Phys.*, 19(22), 14133–14148, doi:10.5194/acp-19-14133-2019, 2019.
- Lambert, A., Hallar, A. G., Garcia, M., Strong, C., Andrews, E. and Hand, J. L.: Dust Impacts of Rapid Agricultural Expansion on the Great Plains, *Geophys. Res. Lett.*, 47(20), doi:10.1029/2020GL090347, 2020.
- Li, J., Chen, B., de la Campa, A. M. S., Alastuey, A., Querol, X. and de la Rosa, J. D.: 2005–2014 trends of PM10 source 15 contributions in an industrialized area of southern Spain, *Environ. Pollut.*, 236, 570–579, doi:10.1016/j.envpol.2018.01.101, 2018.
- Liu, J., Wu, D., Liu, G., Mao, R., Chen, S., Ji, M., Fu, P., Sun, Y., Pan, X., Jin, H., Zhou, Y. and Wang, X.: Impact of Arctic amplification on declining spring dust events in East Asia, *Clim. Dyn.*, 54(3–4), 1913–1935, doi:10.1007/s00382-019-05094-4, 2020.
- 20 Liu, J., Wang, X., Wu, D., Wei, H., Li, Y. and Ji, M.: Historical footprints and future projections of global dust burden from bias-corrected CMIP6 models, *npj Clim. Atmos. Sci.*, 7(1), 1, doi:10.1038/s41612-023-00550-9, 2024.
- Liu, X., Zhang, Y., Yao, H., Lian, Q. and Xu, J.: Analysis of the Severe Dust Process and Its Impact on Air Quality in Northern China, *Atmosphere (Basel)*, 14(7), 1071, doi:10.3390/atmos14071071, 2023.
- Lorentzou, C., Kouvarakis, G., Kozyrakis, G. V., Kampanis, N. A., Trahanatzi, I., Fraidakis, O., Tzanakis, N., Kanakidou, M., 25 Agouridakis, P. and Notas, G.: Extreme desert dust storms and COPD morbidity on the island of Crete, *Int. J. Chron. Obstruct. Pulmon. Dis.*, Volume 14, 1763–1768, doi:10.2147/COPD.S208108, 2019.
- Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K. and Coumou, D.: Influence of Anthropogenic Climate Change on Planetary Wave Resonance and Extreme Weather Events, *Sci. Rep.*, 7(1), 45242, doi:10.1038/srep45242, 2017.

- Marticorena, B., Chatenet, B., Rajot, J. L., Traoré, S., Coulibaly, M., Diallo, A., Koné, I., Maman, A., NDiaye, T. and Zakou, A.: Temporal variability of mineral dust concentrations over West Africa: analyses of a pluriannual monitoring from the AMMA Sahelian Dust Transect, *Atmos. Chem. Phys.*, 10(18), 8899–8915, doi:10.5194/acp-10-8899-2010, 2010.
- Meinander, O., Kouznetsov, R., Uppstu, A., Sofiev, M., Kaakinen, A., Salminen, J., Rontu, L., Welti, A., Francis, D., Piedehierro, A. A., Heikkilä, P., Heikkinen, E. and Laaksonen, A.: African dust transport and deposition modelling verified through a citizen science campaign in Finland, *Sci. Rep.*, 13(1), 21379, doi:10.1038/s41598-023-46321-7, 2023.
- Merdji, A. B., Lu, C., Xu, X. and Mhawish, A.: Long-term three-dimensional distribution and transport of Saharan dust: Observation from CALIPSO, MODIS, and reanalysis data, *Atmos. Res.*, 286, 106658, doi:10.1016/j.atmosres.2023.106658, 2023.
- 10 Micheli, L., Almonacid, F., Bessa, J. G., Fernández-Solas, Á. and Fernández, E. F.: The impact of extreme dust storms on the national photovoltaic energy supply, *Sustain. Energy Technol. Assessments*, 62, 103607, doi:10.1016/j.seta.2024.103607, 2024.
- Middleton, N.: Variability and Trends in Dust Storm Frequency on Decadal Timescales: Climatic Drivers and Human Impacts, *Geosciences*, 9(6), 261, doi:10.3390/geosciences9060261, 2019.
- 15 Middleton, N., Kashani, S. S., Attarchi, S., Rahnama, M. and Mosalman, S. T.: Synoptic Causes and Socio-Economic Consequences of a Severe Dust Storm in the Middle East, *Atmosphere (Basel)*, 12(11), 1435, doi:10.3390/atmos12111435, 2021.
- Mifka, B., Telišman Prtenjak, M., Kavre Piltaver, I., Mekterović, D., Kuzmić, J., Marciuš, M. and Ciglencečki, I.: Intense Desert Dust Event in the Northern Adriatic (March 2020); Insights From the Numerical Model Application and Chemical  
20 Characterization Results, *Earth Sp. Sci.*, 10(7), doi:10.1029/2023EA002879, 2023.
- Millán-Martínez, M., Sánchez-Rodas, D., Sánchez de la Campa, A. M. and de la Rosa, J.: Contribution of anthropogenic and natural sources in PM10 during North African dust events in Southern Europe, *Environ. Pollut.*, 290, 118065, doi:10.1016/j.envpol.2021.118065, 2021.
- Miri, A. and Middleton, N.: Long-term impacts of dust storms on transport systems in south-eastern Iran, *Nat. Hazards*, 114(1),  
25 291–312, doi:10.1007/s11069-022-05390-z, 2022.
- Mona, L., Amiridis, V., Cuevas, E., Gkikas, A., Trippetta, S., Vandenbussche, S., Benedetti, A., Dagsson-Waldhauserova, P., Formenti, P., Haeefe, A., Kazadzis, S., Knippertz, P., Laurent, B., Madonna, F., Nickovic, S., Papagiannopoulos, N., Pappalardo, G., García-Pando, C. P., Popp, T., Rodríguez, S., Sealy, A., Sugimoto, N., Terradellas, E., Vimic, A. V., Weinzierl, B. and Basart, S.: OBSERVING MINERAL DUST IN NORTHERN AFRICA, THE MIDDLE EAST AND EUROPE:  
30 CURRENT CAPABILITIES AND CHALLENGES AHEAD FOR THE DEVELOPMENT OF DUST SERVICES, *Bull. Am.*

- Meteorol. Soc., doi:10.1175/BAMS-D-23-0005.1, 2023.
- Monteiro, A., Basart, S., Kazadzis, S., Votsis, A., Gkikas, A., Vandenbussche, S., Tobias, A., Gama, C., García-Pando, C. P., Terradellas, E., Notas, G., Middleton, N., Kushta, J., Amiridis, V., Lagouvardos, K., Kosmopoulos, P., Kotroni, V., Kanakidou, M., Mihalopoulos, N., Kalivitis, N., Dagsson-Waldhauserová, P., El-Askary, H., Sievers, K., Giannaros, T., Mona, L., Hirtl, M., Skomorowski, P., Virtanen, T. H., Christoudias, T., Di Mauro, B., Trippetta, S., Kutuzov, S., Meinander, O. and Nickovic, S.: Multi-sectoral impact assessment of an extreme African dust episode in the Eastern Mediterranean in March 2018, *Sci. Total Environ.*, 843, 156861, doi:10.1016/j.scitotenv.2022.156861, 2022.
- Mulitza, S., Heslop, D., Pittauerova, D., Fischer, H. W., Meyer, I., Stuu, J.-B., Zabel, M., Mollenhauer, G., Collins, J. A., Kuhnert, H. and Schulz, M.: Increase in African dust flux at the onset of commercial agriculture in the Sahel region, *Nature*, 466(7303), 226–228, doi:10.1038/nature09213, 2010.
- Nishonov, B. E., Kholmatjanov, B. M., Labzovskii, L. D., Rakhmatova, N., Shardakova, L., Abdulakhatov, E. I., Yarashev, D. U., Toderich, K. N., Khujanazarov, T. and Belikov, D. A.: Study of the strongest dust storm occurred in Uzbekistan in November 2021, *Sci. Rep.*, 13(1), 20042, doi:10.1038/s41598-023-42256-1, 2023.
- Peshev, Z., Chaikovskiy, A., Evgenieva, T., Pescherenkov, V., Vulkova, L., Deleva, A. and Dreischuh, T.: Combined Characterization of Airborne Saharan Dust above Sofia, Bulgaria, during Blocking-Pattern Conditioned Dust Episode in February 2021, *Remote Sens.*, 15(15), 3833, doi:10.3390/rs15153833, 2023.
- Pey, J., Querol, X., Alastuey, A., Forastiere, F. and Stafoggia, M.: African dust outbreaks over the Mediterranean Basin during 2001–2011: PM<sub>10</sub> concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology, *Atmos. Chem. Phys.*, 13(3), 1395–1410, doi:10.5194/acp-13-1395-2013, 2013.
- Pi, H., Huggins, D. R., Abatzoglou, J. T. and Sharratt, B.: Modeling Soil Wind Erosion From Agroecological Classes of the Pacific Northwest in Response to Current Climate, *J. Geophys. Res. Atmos.*, 125(2), doi:10.1029/2019JD031104, 2020.
- Preunkert, S., Legrand, M., Kutuzov, S., Ginot, P., Mikhalenko, V. and Friedrich, R.: The Elbrus (Caucasus, Russia) ice core record – Part 1: reconstruction of past anthropogenic sulfur emissions in south-eastern Europe, *Atmos. Chem. Phys.*, 19(22), 14119–14132, doi:10.5194/acp-19-14119-2019, 2019.
- Prospero, J. M. ., Ginoux, P., Torres, O., Nicholson, S. E. . and Gill, T. E. .: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40(1), 2-1-2–31, doi:10.1029/2000RG000095, 2002.
- Pu, B. and Jin, Q.: A Record-Breaking Trans-Atlantic African Dust Plume Associated with Atmospheric Circulation Extremes in June 2020, *Bull. Am. Meteorol. Soc.*, 102(7), E1340–E1356, doi:10.1175/BAMS-D-21-0014.1, 2021.
- Qor-El-Aine, A., Béres, A. and Géczi, G.: Case Study of the Saharan Dust Effects on PM<sub>10</sub> and PM<sub>2.5</sub> Concentrations in



- Budapest in March 2022, *J. Cent. Eur. Green Innov.*, 10(Suppl 1), 67–78, doi:10.33038/jcegi.3500, 2022.
- Querol, X., Alastuey, A., Pandolfi, M., Reche, C., Pérez, N., Minguillón, M. C., Moreno, T., Viana, M., Escudero, M., Orío, A., Pallarés, M. and Reina, F.: 2001–2012 trends on air quality in Spain, *Sci. Total Environ.*, 490, 957–969, doi:10.1016/j.scitotenv.2014.05.074, 2014.
- 5 Ridley, D. A., Heald, C. L. and Prospero, J. M.: What controls the recent changes in African mineral dust aerosol across the Atlantic?, *Atmos. Chem. Phys.*, 14(11), 5735–5747, doi:10.5194/acp-14-5735-2014, 2014.
- Rodríguez-Caballero, E., Stanelle, T., Egerer, S., Cheng, Y., Su, H., Canton, Y., Belnap, J., Andreae, M. O., Tegen, I., Reick, C. H., Pöschl, U. and Weber, B.: Global cycling and climate effects of aeolian dust controlled by biological soil crusts, *Nat. Geosci.*, 15(6), 458–463, doi:10.1038/s41561-022-00942-1, 2022.
- 10 Rodríguez, S. and López-Darias, J.: Dust and tropical PM<sub>x</sub> aerosols in Cape Verde: Sources, vertical distributions and stratified transport from North Africa, *Atmos. Res.*, 263, 105793, doi:10.1016/j.atmosres.2021.105793, 2021.
- Rodríguez, S., Querol, X., Alastuey, A., Kallos, G. and Kakaliagou, O.: Saharan dust contributions to PM<sub>10</sub> and TSP levels in Southern and Eastern Spain, *Atmos. Environ.*, 35(14), 2433–2447, doi:10.1016/S1352-2310(00)00496-9, 2001.
- Rodríguez, S., Alastuey, A., Alonso-Pérez, S., Querol, X., Cuevas, E., Abreu-Afonso, J., Viana, M., Pérez, N., Pandolfi, M.  
15 and de la Rosa, J.: Transport of desert dust mixed with North African industrial pollutants in the subtropical Saharan Air Layer, *Atmos. Chem. Phys.*, 11(13), 6663–6685, doi:10.5194/acp-11-6663-2011, 2011.
- Rodríguez, S., Alastuey, A. and Querol, X.: A review of methods for long term in situ characterization of aerosol dust, *Aeolian Res.*, 6, 55–74, doi:10.1016/j.aeolia.2012.07.004, 2012.
- Rodríguez, S., Cuevas, E., Prospero, J. M., Alastuey, A., Querol, X., López-Solano, J., García, M. I. and Alonso-Pérez, S.:  
20 Modulation of Saharan dust export by the North African dipole, *Atmos. Chem. Phys.*, 15(13), 7471–7486, doi:10.5194/acp-15-7471-2015, 2015.
- Rodríguez, S., Riera, R., Fonteneau, A., Alonso-Pérez, S. and López-Darias, J.: African desert dust influences migrations and fisheries of the Atlantic skipjack-tuna, *Atmos. Environ.*, 312, 120022, doi:10.1016/j.atmosenv.2023.120022, 2023.
- Screen, J. A. and Simmonds, I.: Exploring links between Arctic amplification and mid-latitude weather, *Geophys. Res. Lett.*,  
25 40(5), 959–964, doi:10.1002/grl.50174, 2013.
- Seidel, D. J., Fu, Q., Randel, W. J. and Reichler, T. J.: Widening of the tropical belt in a changing climate, *Nat. Geosci.*, 1(1), 21–24, doi:10.1038/ngeo.2007.38, 2008.
- Solomos, S., Kalivitis, N., Mihalopoulos, N., Amiridis, V., Kouvarakis, G., Gkikas, A., Biniotoglou, I., Tsekeri, A., Kazadzis, S., Kottas, M., Pradhan, Y., Proestakis, E., Nastos, P. and Marengo, F.: From Tropospheric Folding to Khamsin and Foehn

- Winds: How Atmospheric Dynamics Advanced a Record-Breaking Dust Episode in Crete, *Atmosphere (Basel)*, 9(7), 240, doi:10.3390/atmos9070240, 2018.
- Sorribas, M., Adame, J. A., Andrews, E. and Yela, M.: An anomalous African dust event and its impact on aerosol radiative forcing on the Southwest Atlantic coast of Europe in February 2016, *Sci. Total Environ.*, 583, 269–279, doi:10.1016/j.scitotenv.2017.01.064, 2017.
- 5 Tong, D. Q., Gill, T. E., Sprigg, W. A., Van Pelt, R. S., Baklanov, A. A., Barker, B. M., Bell, J. E., Castillo, J., Gassó, S., Gaston, C. J., Griffin, D. W., Huneus, N., Kahn, R. A., Kuciauskas, A. P., Ladino, L. A., Li, J., Mayol-Bracero, O. L., McCotter, O. Z., Méndez-Lázaro, P. A., Mudu, P., Nickovic, S., Oyarzun, D., Prospero, J., Raga, G. B., Raysoni, A. U., Ren, L., Sarafoglou, N., Sealy, A., Sun, Z. and Vimic, A. V.: Health and Safety Effects of Airborne Soil Dust in the Americas and Beyond, *Rev. Geophys.*, 61(2), doi:10.1029/2021RG000763, 2023.
- 10 Viana, M., Querol, X., Alastuey, A., Cuevas, E. and Rodríguez, S.: Influence of African dust on the levels of atmospheric particulates in the Canary Islands air quality network, *Atmos. Environ.*, 36(38), 5861–5875, doi:10.1016/S1352-2310(02)00463-6, 2002.
- Vukovic, A., Cvetkovic, B., Giannaros, T., Shahbazi, R., Sehat Kashani, S., Prieto, J., Kotroni, V., Lagouvardos, K., Pejanovic, G., Petkovic, S., Nickovic, S., Vujadinovic, M., Basart, S., Darvishi, A. and Terradellas, E.: Numerical Simulation of Tehran Dust Storm on 2 June 2014: A Case Study of Agricultural Abandoned Lands as Emission Sources, *Atmosphere (Basel)*, 12(8), 1054, doi:10.3390/atmos12081054, 2021.
- 15 Xi, X., Steinfeld, D., Cavallo, S., Wang, J., Chen, J., Zulpykharov, K. and Henebry, G. M.: What caused the unseasonal extreme dust storm in Uzbekistan during November 2021?, *Environ. Res. Lett.*, doi:10.1088/1748-9326/ad02af, 2023.
- 20 Xie, X., Liu, X., Shi, Z., Li, X., Xie, X., Sun, H., He, J., Che, H., Zhang, X.-Y., Zhisheng, A., Wang, D. and Liu, Y.: Sharp decline of dust events induces regional wetting over arid and semi-arid Northwest China in the NCAR Community Atmosphere model, *Environ. Res. Lett.*, doi:10.1088/1748-9326/ad16a5, 2023.
- Yang, H., Lohmann, G., Lu, J., Gowan, E. J., Shi, X., Liu, J. and Wang, Q.: Tropical Expansion Driven by Poleward Advancing Midlatitude Meridional Temperature Gradients, *J. Geophys. Res. Atmos.*, 125(16), doi:10.1029/2020JD033158, 2020.
- 25 Yang, H., Lohmann, G., Shi, X. and Müller, J.: Evaluating the Mechanism of Tropical Expansion Using Idealized Numerical Experiments, *Ocean. Res.*, 2, doi:10.34133/olar.0004, 2023.
- Yu, H., Chin, M., Yuan, T., Bian, H., Remer, L. A., Prospero, J. M., Omar, A., Winker, D., Yang, Y., Zhang, Y., Zhang, Z. and Zhao, C.: The fertilizing role of African dust in the Amazon rainforest: A first multiyear assessment based on data from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations, *Geophys. Res. Lett.*, 42(6), 1984–1991, doi:10.1002/2015GL063040, 2015.
- 30

- Yu, H., Tan, Q., Zhou, L., Zhou, Y., Bian, H., Chin, M., Ryder, C. L., Levy, R. C., Pradhan, Y., Shi, Y., Song, Q., Zhang, Z., Colarco, P. R., Kim, D., Remer, L. A., Yuan, T., Mayol-Bracero, O. and Holben, B. N.: Observation and modeling of the historic “Godzilla” African dust intrusion into the Caribbean Basin and the southern US in June 2020, *Atmos. Chem. Phys.*, 21(16), 12359–12383, doi:10.5194/acp-21-12359-2021, 2021.
- 5 Yu, Y. and Ginoux, P.: Enhanced dust emission following large wildfires due to vegetation disturbance, *Nat. Geosci.*, 15(11), 878–884, doi:10.1038/s41561-022-01046-6, 2022.
- Zafra-Pérez, A., Boente, C., García-Díaz, M., Gómez-Galán, J. A., de la Campa, A. S. and de la Rosa, J. D.: Aerial monitoring of atmospheric particulate matter produced by open-pit mining using low-cost airborne sensors, *Sci. Total Environ.*, 904, 166743, doi:10.1016/j.scitotenv.2023.166743, 2023.
- 10 Zhang, T., Zheng, M., Sun, X., Chen, H., Wang, Y., Fan, X., Pan, Y., Quan, J., Liu, J., Wang, Y., Lyu, D., Chen, S., Zhu, T. and Chai, F.: Environmental impacts of three Asian dust events in the northern China and the northwestern Pacific in spring 2021, *Sci. Total Environ.*, 859, 160230, doi:10.1016/j.scitotenv.2022.160230, 2023.