

Measurement report: The promotion of low-level jet and thermal-effect on development of deep convective boundary layer at the southern edge of the Taklimakan Desert

5 Lian Su¹, Chunsong Lu^{2,3}, Jinlong Yuan^{2,3}, Xiaofei Wang⁴, Qing He⁵, Haiyun Xia^{1,2,3,*}

¹School of Earth and Space Science, University of Science and Technology of China, Hefei 230026, China

²School of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing 210044, China

³Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Key Laboratory of Meteorological Disaster of Ministry of Education,

10 NUIST, Nanjing 210044, China

⁴Xinjiang Uygur Autonomous Region Meteorological Service, Urumqi 830002, China

⁵Institute of Desert Meteorology, China Meteorological Administration, Urumqi 830002, China

Correspondence to: Haiyun Xia (hsia@ustc.edu.cn)

Abstract. A vigorous development process of the deep convective boundary layer (CBL) was observed at the southern edge
15 of the Taklimakan Desert on 6 June, 2022. Based on coherent Doppler wind lidar and ERA5 data, the formation mechanism
of the deep CBL exceeding 5 km was analyzed, which was mainly driven by the low-level jet (LLJ) and thermal-effects.
During the stage of LLJ prior to the formation of the deep CBL, the LLJ had made sufficient preparations for the development
of the deep CBL in terms of momentum, energy, and material. Firstly, the cold downhill airflow from the Tibet Plateau leading
to LLJ weakens the height and intensity of the temperature inversion layer, which reduces the energy demand for the broken
20 of the temperature inversion layer. Secondly, the LLJ not only supplements the material and energy in the residual layer, but
also suppresses the exchange with the lower atmosphere. In addition, the LLJ provides a driving force for the development of
the deep CBL. During the stage of thermal effects, the sensible heat driven air-pump from the Tibet Plateau and cold front
transit provide additional impetus for the development of the deep CBL. Finally, the formation of deep CBL was catalyzed by
25 extreme thermal effects of the underlying surface, such as the furnace effect and the atmospheric superadiabatic expansion
process. The study of the development of the deep CBL is important for revealing the land-air exchange process of momentum,
energy, and material between the Taklimakan Desert and the Tibetan Plateau.

1 Introduction

The atmospheric boundary layer (ABL) is the place where the earth's surface exchanges momentum, energy and material with the free atmosphere (Stull, 1988; Garratt, 1994). The boundary layer height (BLH) is an important meteorological reference variable in the vertical direction, which indicates the atmospheric environmental capacity of the region and the vertical diffusion degree of pollutants (Holtslag and Boville, 1993). The study in the temporal and spatial distribution of BLH, which is closely related to human life, plays an important auxiliary role in monitoring air pollution and formulating pollution control policies according to local conditions.

The convective boundary layer (CBL) belongs to an unstable ABL, and the height of CBL should usually be lower than 2-3 km. However, under certain conditions, such as arid region and monsoon climate, the height of CBL can continue to develop upward and may exceed 5 km (Garratt, 1994). At present, a large number of scholars have found and analyzed the deep CBL phenomenon in the subcontinent of India (Basha and Ratnam, 2009; Raman et al., 1990), Sahara desert (Birch et al., 2012; Marsham et al., 2008), Mongolia (Han et al., 2015), Tibet Plateau (Che and Zhao, 2021; Lai et al., 2023), Badain Jaran Desert (Han et al., 2012), and Gobi desert (Zhang et al., 2002). These studies also revealed that the deep CBL exerts an influence on the local pollutant transmission and diffusion, cloud formation processes, strong convective weather, rainfall, drought and so on. The Taklimakan Desert (TD), which plays an important role in global climate change, has also carried out corresponding research work. For example, in the hinterland of the TD, the intense surface heating is not the primary reason for the formation of deep CBL, whereas the presence of weak temperature inversion and near-neutral residual layer (RL) above the CBL are crucial (Zhang et al., 2022; Xu et al., 2018); The low-level jet (LLJ) can trigger significant air accumulation and dynamic convergence in the lower-level atmosphere, while the deep CBL is usually accompanied by the LLJ on the following night (Wang et al., 2019); The deep CBL enables the formation of clouds in the late afternoon, the formation of clouds will not only lead to a significant cooling of surface, but also make the momentum in the upper part of the boundary layer to transport downward and cause dust emissions (Zhang et al., 2024). Due to the fact that the TaZhong station is located in the center of the TD, and is equipped with the most comprehensive observation equipment, most studies of deep CBL are concentrated here. However, the MinFeng station, which is located on the northern slope of the Tibet Plateau (TP) and has severe wind-sand activities (Yang et al., 2016; Xiao et al., 2008), was established in 2018 (Yang et al., 2020). On the one hand, there is a lack of sufficient study results of deep CBL, and the particularity of geographical locations (TD, slope terrain, Kunlun Mountains, TP) further complicates the formation mechanism of deep CBL. On the other hand, the deep CBL plays an important role in the regional circulation and weather, its study not only helps to reveal the mechanism of local dust emission and transport (Jia et al., 2015; Meng et al., 2019), but also promotes the understanding of the land-atmosphere interaction between the TD and the TP.

When experiments are carried out in the desert, the harsh desert climate environment will increase the performance requirements and the maintenance costs of meteorological equipment, thus increasing the difficulty of environmental monitoring. The coherent Doppler wind lidar (CDWL) needs to operate under the conditions of strict sealing and precise


temperature control. In the detection of the BLH, the CDWL has the characteristics of low blind area, high radial spatial resolution and temporal resolution, long detection distance and little influence by ground clutter. These characteristics of CDWL enable it to obtain the air flow conditions of the atmosphere from the calculated wind field information, monitor the change of the BLH in real time and more accurately, and help to understand the diffusion and retention of dust pollutants.

5 Overall, the CDWL can be used for long-term continuous and stable detection in desert areas, and it is one of the effective ways to estimate the BLH in the desert (Li et al., 2017; Zhang et al., 2020; Collis, 1966; Zhang et al., 2021).

In this paper, the CDWL was used to conduct a long-term stable observation experiment in the MinFeng area of the TD, and a representative formation process of the deep CBL was observed on 6 June, 2022, local time (UTC+8). The CDWL data and ERA5 data were used to analyze the causes of the formation of the deep CBL. This paper is organized as follows: the study
10 site, datasets and methods are described in Sect. 2. The CDWL observation results are presented and analyzed in Sect. 3. From the perspective of the whole desert region, the ERA5 reanalysis data were also analyzed in Sect. 4. Finally, a conclusion is drawn in Sect. 5.

2 Site, data resources, methods

2.1 Study site



15

Figure 1. Elevation map of the Taklimakan Desert. The red circle represents the study site of MinFeng.

The Taklimakan Desert is the second largest shifting desert in the world and the largest desert of China. Due to the blocking effect of the Tibet Plateau on the warm and humid airflow, the TD has become a typical extreme arid climate zone. The study site of MinFeng (37.06° N, 82.69° E, elevation 1418 m), located on the southern edge of TD and adjacent to the
20 northern foot of Kunlun Mountains, which is significantly influenced by the TP. The area has the characteristics of long sunshine hours, strong radiation, scarce precipitation, and the convergence of east-west airflows here, making it the most frequent site for wind-sand disasters in China. The average number of dusty days in this area exceeds 200 days per year, and

the frequent wind-sand weather has a significant impact on human activities and health (Zhou et al., 2020; Zhou et al., 2022; Yang et al., 2016; Wu et al., 2016).

2.2 Instruments and dataset

5 A compact CDWL working at eye-safe wavelength of 1.5 μm is used in this study. The pulse energy and repetition frequency of the laser are 130 μJ and 10 kHz, respectively. The temporal resolution and radial spatial resolution are 1 minute and 30 m, respectively. During the experiment, the lidar operates in the velocity azimuth display (VAD) scanning mode with an elevation angle of 70°. The key parameters of the CDWL are listed in Table 1.

Local meteorological data are provided by the MinFeng County Meteorological Bureau, including air temperature, ground temperature, relative humidity, horizontal visibility, etc.

10 ERA5 (Hersbach et al., 2020) is the fifth-generation global climate reanalysis dataset from the European Centre for Medium-Range Weather Forecasts (ECMWF). In this paper, ERA5 reanalysis data, such as $\mu\text{-v-}\omega$ wind vector, atmospheric temperature, relative humidity, mean sea level pressure, surface latent heat flux, boundary layer height, etc. are applied to analyze the regional variation of the whole desert. The spatial resolution of the reanalysis data is $0.25^\circ \times 0.25^\circ$, and the temporal resolution is 1 hour.

15 **Table 1.** Key Parameters of the CDWL

Parameter	Value
Wavelength	1.5 μm
Pulse energy	130 μJ
Pulse repetition frequency	10 kHz
Diameter of telescope	100 mm
Radial spatial resolution	30 m
Azimuth scanning range	0-360°
Zenith angle	70°
Sample rate of ADC	500 MS/s

2.3 Methods

The data from CDWL has the characteristics of high radial spatial and temporal resolution, and can be used to estimate turbulent kinetic energy dissipation rate (TKEDR) at different heights. The backscatter signal and turbulence intensity detected by CDWL are sharply reduced due to the temperature inversion characteristic of the atmospheric boundary layer top (Hooper and Eloranta, 1986). Based on this, the TKEDR threshold method can effectively estimate the BLH (Wang et al., 2021; Banakh et al., 2021).

The calculation formula of TKEDR is as follows (Banakh and Smalikho, 2018):

$$TKEDR = \left[\frac{\bar{D}_L(\varphi_l) - \bar{D}_L(\varphi_1)}{A(l\Delta y_k) - A(\Delta y_k)} \right]^{\frac{3}{2}} \quad (1)$$

where $\bar{D}_L(\varphi_l)$ is azimuth structure function. L is the serial number for the laser beam's line of sight. $\varphi_l = l\Delta\theta$, $\Delta\theta$ is the azimuth angle resolution, and $l=1,2,3,\dots$. The $A(l\Delta y_k)$ is calculated theoretically for the Kolmogorov model of the two-dimensional turbulence spectrum (Banakh et al., 2017), Δy_k is the transverse dimension of the probed volume, and k is the range gate number, $k=1,2,3,\dots$. The error analysis for calculating TKEDR and BLH was conducted by Viktor A. Banakh (Banakh et al., 2017; Banakh et al., 2021).

In this experiment, the value of l is set to 2, and the threshold of TKEDR is set to $10^{-4} \text{ m}^2 \text{ s}^{-3}$. When the location is at the height of $H_n = \Delta R * N$ (N is the index number of bins, and ΔR is radial spatial resolution), if all TKEDR values within the range $[\Delta R * (N+1), \Delta R * (N+5)]$ are less than the threshold, then H_n is used as the BLH.

The same type of CDWL also realizes the calculation and verification of TKEDR and BLH in various application scenarios (Wang et al., 2021; Wang et al., 2022; Jiang et al., 2022; Yuan et al., 2020; Yuan et al., 2021; Wu et al., 2023; Li et al., 2023).

3 Lidar results and local analysis

The underlying surface of the desert makes the night temperature drop greatly, so that the inversion layer (IL) is more likely to be formed at night. Compared with the night, the change of the mixed boundary layer height during the daytime can better reflect the development of local dust pollution. In this paper, the statistics have been collected on the probability of monthly occurrence of boundary layer at different heights during the daytime (8:00 LT~21:00 LT) from September 2021 to August 2022, and the number of samples is 50663. It can be clearly seen from Figure 2 that a small number of boundary layer heights developed more than 5 km in June 2022. In order to explain this phenomenon, the typical boundary layer data on 6 June, 2022 were selected for analysis.




Figure 2. The monthly probability distribution of the boundary layer height of MinFeng in the Taklimakan Desert during the daytime from September 2021 to August 2022.

Fig. 3 shows the continuous observation results of the CDWL, the local meteorological equipment, and the ERA5 on 6 June, 2022, local time (UTC+8). The radial spatial resolution of the CDWL is 30 m. By analyzing the original data of power spectrum, the data such as CNR, TKEDR, vertical wind speed, horizontal wind speed and horizontal wind direction were obtained. The CNR can be used as an indicator of aerosol concentration (Pea et al., 2013). When the CNR value is below -17 dB, the

calculated products are eliminated in the preprocessing. In this paper, 0:00 LT~12:00 LT is divided into the LLJ stage, and 12:00 LT~24:00 LT is divided into the thermal effect stage.

The LLJ generally refers to the strong and narrow airflow zone with wind speed greater than 12 m s^{-1} within 3 km height (Bonner, 1968). It can be seen from Fig. 3b that there is an obvious LLJ phenomenon in the study area before 12:00 LT, and the central axis of the LLJ is located at about 1 km.

At 0:00 LT~6:00 LT, before the formation of the LLJ. It can be clearly found in Fig. 3b that the horizontal wind speed has subsided from 4 km to 2 km, and the wind is the downhill wind blowing from the TP to the desert (Fig. 3e, $>3 \text{ km}$, $225^\circ\text{-}295^\circ$). When the downhill airflow was superimposed on the desert background wind field, the horizontal wind speed was increased to a maximum of 16.62 m s^{-1} at 5:41 LT, which reached a critical sand-raising wind speed of $3.5\text{-}10.9 \text{ m s}^{-1}$ (Yang et al., 2017) and promoted the formation of LLJ (Matsumoto and Ninomiya, 1971; Mcnider and Pielke, 1981). The enhanced turbulence activity near the surface was also observed, and the BLH was stabilized at about 1 km (Fig. 3c). In addition, according to the local meteorological data, the study site had dust weather during this period (Fig. A1).

At 6:00 LT~12:00 LT, this period is the maintenance stage of the LLJ. In Fig. 3(f-g), it can be found that during the period of 6:00 LT~10:00 LT, with the cold downhill airflow and the upstream cold airflow traveling to the desert basin where the study site is located, the near-surface temperature dropped sharply and the relative humidity increased significantly. Meanwhile, the surface weather station also recorded that the difference of the ground-air temperature reached a minimum of $-0.8 \text{ }^\circ\text{C}$ at 8:00 LT. At 6:00 LT~8:00 LT, firstly, as shown in Fig. 3a, the downward transfer of momentum may have caused the dust aerosol to lift into the residual layer at about 3 km (Washington et al., 2006; Fiedler et al., 2013; Ge et al., 2016), and promoted the dust aerosol in the residual layer to be supplemented from the desert hinterland. Secondly, the height of the inversion layer in the meteorology also often refers to the height of the ABL. Therefore, it can be seen from Fig. 3c that the downhill cold airflow weakened the intensity of the IL, reduced its height, and formed a near-surface IL (also analyzed in Fig. 5(p-t)). At 7:58 LT, the maximum wind speed of 15.04 m s^{-1} was reached, and the strong wind shear effect beneath the LLJ can provide a momentum source for turbulent activity and cause intermittent pulsation of turbulence (Ohya et al., 2006; Mathieu et al., 2005), so that the BLH can be raised to more than 2 km and the material in the residual layer may be partially supplemented.

The LLJ and IL can provide sufficient momentum, energy and material preparation for the development of the deep CBL during the period from 8:00 LT to 11:00 LT. Firstly, it can be seen from Fig. 3(a-b), due to the existence of the LLJ and the IL, the atmosphere exhibited a stratified state (Blackadar, 1957), so that the high concentration of CNR values was distributed below 1 km, which can provide a material basis for boundary layer development. Secondly, the LLJ and the IL can play a role in maintaining the balance of atmospheric thermal structure, inhibiting the internal turbulent motion and energy exchange with the upper atmosphere, and storing enough momentum and energy for the development of deep CBL. Finally, the weakening of the height and stability of the IL enables the developing boundary layer to reduce the energy of the broken IL, which is conducive to the vigorous development of the subsequent boundary layer.

At 12:00 LT~19:00 LT, with the shortwave radiation of the sun gradually heating the atmosphere and the driving force provided by the LLJ, the stable stratification at low altitude was broken, and the turbulent mixing process began to be reactivated. In

Fig.3 (c-d), it can be seen that the values of TKEDR were consistently maintained at a high level, indicating a significant enhancement in the vertical transport capacity of the atmosphere (Wang et al., 2020). Consequently, the underlying CNR value below 1 km in Fig. 3a increased significantly, and the height of the CBL in Fig. 3c was rapidly developed and exceeded 5 km (Meng et al., 2019). Due to the existence of the LLJ and the IL in the early stage, the dust aerosol was concentrated below 1 km. The lower dust aerosol layer can quickly absorb solar radiation and heat the atmosphere, which can form a “furnace effect” (Ma et al., 2021; Ma et al., 2020) to make the material conditions of the desert boundary layer develop rapidly into thermal conditions. As shown in Fig. 3a and Fig. 3c, the temperature difference of ground-air reached a maximum of 25 °C at 15:00 LT, and the height of the CBL was rapidly increasing around 13:00 LT. This strong thermal effect rapidly promoted the growth of the CBL. The LLJ with large wind speed can also provide a basic dynamic condition for the development of the CBL, and the lifting process of the horizontal wind can be obviously found at 12:00 LT in Fig. 3b. The TP, with its characteristics of high altitude, thin air, and little weakening effect on solar radiation, enables the sensible heat heating of its surface to suck up the surrounding atmosphere, forming a “sensible heat driven air-pump” of the TP (TP-SHAP) (Wu et al., 2012; Wu et al., 2007). The study site of this paper is located at the northern foot of the TP, where the TP-SHAP has a significant effect on lifting dust aerosols (Tan et al., 2021). The TP-SHAP can lift dust aerosols from the bottom of the desert along the northern slope of the Kunlun Mountains to the TP (Ge et al., 2014), thus further promoting the development of the local boundary layer (analyzed in Fig. 7). Additionally, the role of continuous hot weather and the entrainment process of the residual layer cannot be ignored (Zhang et al., 2011; Marsham et al., 2008). The local meteorological data show that the hot weather, with temperatures greater than 30 °C, has persisted for 6 days.

At 17:00 LT~24:00 LT, the study site began to be covered by clouds (Fig. 3c, Fig. A3). Before 18:00 LT, the study site was covered by scattered clouds, and after 18:00 LT, the cloud completely covered the study site. The presence of clouds can greatly weaken the solar radiation reaching the surface, causing the surface temperature to decrease rapidly (Fig. 3a), and gradually weakening the turbulence intensity of the atmosphere (Fig. 3c). At 17:00 LT~20:00 LT, the atmospheric turbulence remained active during the initial stages of cloud formation. On the one hand, as the surface temperature was still much higher than the atmospheric temperature, and the heat preservation effect of dust on the atmosphere continued to provide energy for the upper atmosphere. On the other hand, the cold clouds moved towards the warm air mass over the desert, promoting the formation of an upper-level cold front and causing strong convective motion in the lower atmosphere (Fig. 3d), and the height of deep CBL reached its peak at 18:00 LT (Fig. 3c). When the cloud completely covered the study site, the surface radiation further cooled the near-surface air, greatly weakening the atmospheric turbulence intensity and significantly reducing the CBL height before sunset, and the ground-air temperature difference changed to -0.6 °C at 22:00 LT. After the boundary layer developed into a nocturnal stable boundary layer (SBL), the airflow began to recover into a relatively strong and narrow airflow zone (Hoecker, 1963).

The local surface meteorological observation data of the day are shown in Fig. A1. At the experimental site, the representative CNR, horizontal wind speed, TKEDR, BLH, vertical wind speed and wind direction in different seasons are also presented in Fig. A2. The cloud coverage over the Taklimakan Desert is shown in Fig. A3.





Figure 3. The continuous observation results of the CDWL, the local meteorological equipment, and the ERA5 on 6 June, 2022, local time (UTC+8). (a) CNR, the local air and ground temperatures are also shown in this subgraph. (b) horizontal wind speed. (c) $\log_{10}(\text{TKEDR})$ and boundary layer height, the time of sunrise and sunset are marked with arrow symbols, and the cloud base height is marked with a purple dotted line. (d) vertical wind speed, the positive vertical wind speed represents the descending speed, and vice versa. (e) horizontal wind direction, 0° represents the wind blows to the north. (f) atmospheric temperature. (g) relative humidity. The height represents the height above the ground of the lidar site. The temperature and relative humidity contours in Fig. 3f and Fig. 3g are denoted by white dashed lines.

5

4 ERA5 results and regional analysis

4.1 Low-level jet



5 **Figure 4.** The distributions of ERA5 data over the Taklimakan Desert at 750 hPa (about 1.05 km above the ground) from 2:00 LT to 10:00 LT, on 6 June, 2022, local time (UTC+8). (a-e) wind vector. (f-j) geopotential height. (k-o) air temperature. (p-t) relative humidity. The purple circle represents the study site of MinFeng. The contours are denoted by dashed lines.

The LLJ is closely related to air pollution, dust storm, heavy rainfall and many other aspects. Studying the LLJ can partially reveal the dust emission and transmission process in the study area. Fig. 4 shows the variations in wind vector, geopotential height, atmospheric temperature, and relative humidity of the TD at 750 hPa (about 1.05 km above the ground) from 2:00 LT to 10:00 LT. As seen in the wind vector subgraphs of Fig. 4(a-e), the study site is located in the LLJ region, thus confirming the validity of observing LLJ using CDWL. In Fig. 4(a-c), the upstream and downstream of the horizontal airflow correspond to the divergence region (with lower temperature and relative humidity) and the convergence region (with higher temperature and relative humidity), respectively. The horizontal airflow can cause the divergence of the upstream region to sink and the convergence of the downstream region to rise, which is conducive to the development of the subsequent LLJ and the transport of dust aerosols (Bonner, 1968; Han et al., 2022). The TD is surrounded by three mountains, forming a unique horseshoe-shaped terrain structure. At the northeast of the study site, the northwest wind with lower wind speed deflects to the west after

10

15

encountering the blocking of the TP. In order to ensure the conservation of the potential vorticity, an east wind with higher wind speed is formed. The blocking of the mountains further catalyzes the formation of the LLJ (Wexler, 1961) and may also play a role in the transport of dust aerosols (Caton Harrison et al., 2021). Compared with the northern side of the desert, the southern side has higher temperature, lower geopotential height, and lower relative humidity, which is conducive to the formation of more active airflow. Under the background conditions of thermal and potential difference, it is helpful to the formation and enhancement of the desert background wind field, such as gradient wind and thermal wind, and to promotes the formation of LLJ (Stensrud, 1996; Rife et al., 2010). At night in the TD, the wind field of the LLJ rotates clockwise and has a typical inertial oscillation phenomenon (Blackadar, 1957). In summary, the LLJ formed a water vapor convergence area in front of the study site and maintained the temperature within a relatively high range (16°C isotherm), which enhanced the potential instability (Fig. 4(g-i)), thereby strengthening the convective potential of the atmosphere and providing the necessary energy and water vapor conditions for the subsequent development of the boundary layer.




Figure 5. The distributions of ERA5 data over the Taklimakan Desert from 2:00 LT to 10:00 LT, on 6 June, 2022, local time (UTC+8). (a-e) mean sea level pressure. (f-j) surface sensible heat flux. (k-o) surface latent heat flux. (p-t) temperature inversion distribution, obtained by subtracting 850 hPa (38 m above the ground) from the temperature data of 825 hPa (232 m above the ground). (u-x) boundary layer height. The purple circle represents the study site of MinFeng. The contours are denoted by dashed lines.

Fig. 5 shows the distribution of mean sea level pressure, surface sensible heat flux, surface latent heat flux, temperature inversion distribution, and BLH. At 2:00 LT~8:00 LT, it can be observed from Fig. 5(a-d) that the pressure of the TP (1025 hPa) is much higher than that near the study site in the desert basin (1007 hPa). Therefore, a large pressure gradient formed between the TP and the TD, which can promote the formation of the downhill airflow under the combined action of gravity and pressure gradient force. When the airflow of the downhill is superimposed on the background wind field of the TD, the wind speed of the TD can be enhanced (similar to Fig. 3b), which also indicates that the influence of topographic baroclinicity is significant (Jones, 2019). During this period, the surface sensible heat flux at the study site is generally greater than -10 W m^{-2} , and the surface latent heat flux is positive. The distribution of these fluxes will inhibit the convection and turbulence activities in the boundary layer, thereby facilitating the development of the nocturnal SBL (Zhang et al., 2017). At the study site, the atmospheric temperature change caused by the LLJ promoted the formation of the IL, and the most obvious IL phenomenon was observed at 8:00 LT. The IL can weaken the convective motion of atmosphere, resulting in the boundary layer height near the study site being constrained within 0.25 km, thereby limiting the diffusion and mixing of dust pollutants, and serving as a source of dust material for further boundary layer development.

At 8:00 LT~10:00 LT, with the increase of solar radiation, the average sea level pressure on the southeast TP gradually decreased, the inversion layer started to dissipate, and the sensible and latent heat fluxes of the desert increased rapidly.

4.2 Heat factors

In June, the precipitation in the study region is scarce, the underlying surface is dominated by fine sand, and the soil is dry with strong evaporation capacity. Fig. 6 shows the distribution of wind vector, air temperature, surface sensible heat flux, surface latent heat flux and boundary layer height over the TD from 12:00 LT to 22:00 LT. At 12:00 LT~18:00 LT, with the dissipation of the LLJ, the study site gradually shifted from the east wind to the northeast wind blowing towards the TP. The continuous high solar radiation made the surface sensible heat flux near the study site greater than 300 W m^{-2} (at 16:00 LT), so the efficiency of heating the atmosphere was very high (Zhang et al., 2002). Firstly, heating the atmosphere with such a high surface sensible heat flux can promote the generation of thermal convection, thus strengthening the atmospheric turbulence and making the boundary layer develop rapidly. Secondly, the atmospheric temperature of the study region is also always located in the relative highest value region at each moment ($30 \text{ }^{\circ}\text{C}$ isotherm), so that the region can form a lower low-pressure center to gather dust from the relatively cold area. Therefore, at 18:00 LT, a deep CBL with a maximum height of 4546 m was formed near the study site. At 20:00 LT~22:00 LT, the surface sensible heat flux turned to a negative value rapidly and the BLH decreased to less than 1 km.

Compared with the CDWL data, the temporal and spatial resolution of the ERA5 reanalysis data is more coarse, and the calculation method of the BLH is also different, leading to a lower maximum height of the deep CBL compared to the CDWL data.




Figure 6. The distributions of ERA5 data over the Taklimakan Desert from 12:00 LT to 22:00 LT, on 6 June, 2022, local time (UTC+8). (1-6) wind vectors (850 hPa, 38 m above the ground). (7-12) air temperature (850 hPa, 38 m above the ground). (13-18) surface sensible heat flux. (19-24) surface latent heat flux. (25-30) boundary layer height. The purple circle represents the study site of MinFeng. The contours are denoted by dashed lines.

5

Fig. 7 shows the variation of meteorological data in the vertical section of the TD at the nearest study site of MinFeng. In Fig. 7(a-e), the wind vector is synthesized by v and scaled ω (ω scaled by 10), which depicts the atmospheric vertical motion of the TD. The elevation of the study site is 1418 m. The Richardson number (Ri) can reflect the influence of vertical shear of horizontal wind on atmospheric stability and the state of atmospheric turbulence (Stull, 1988). There is a certain deviation in selecting different Ri values to identify the BLH (Guo et al., 2016). Generally, Ri values less than or equal to 0.25 are used to represent the turbulent motion state of the atmosphere, with a critical value of 0.25 often chosen to identify the BLH (Zhang et al., 2013). At 12:00 LT, the near-surface potential temperature on the left side is higher than that on the right side, and the 20 % relative humidity contour line, as well as the Ri critical value 0.25 contour line, are maintained at a lower elevation.

10




Figure 7. The meteorological elements in the vertical section of the Taklimakan Desert at the study site from 12:00 LT to 20:00 LT, on 6 June, 2022, local time (UTC+8). (a-e) wind vectors (synthesized by v and scaled ω , ω scaled by 10). (f-g) potential temperature. (k-o) relative humidity. (p-t) Richardson number. The purple circle represents the study site of MinFeng. The leftmost of the study site is the Tibet Plateau.

5

At 14:00 LT~18:00 LT, in the elevation range of 1.4 km to 5 km, firstly, the potential temperature on the north side of the TP is almost constant with elevation (Fig. 7(h-i), 316K). This indicates that the rate of temperature reduction during the rapid heating and expansion of the air is greater than the rate of dry adiabatic cooling, forming an atmospheric superadiabatic expansion process. In this atmospheric state, a strong updraft can be formed (Fig. 7(b-c)), which makes the atmosphere in an unstable state. The obvious uplift process can be seen in the 0.25 Ri contour in Fig. 7(q-s) and the 20% relative humidity contour in Fig. 7(l-n). This atmospheric superadiabatic expansion process is very conducive to the diffusion and transport of dust aerosols and the development of boundary layer (Arnette et al., 1998; Nilsson et al., 2001). This phenomenon also provides evidence that TKEDR is maintained within a large numerical range in Fig. 3c. Secondly, the topography and potential temperature of the south side of the desert basin are higher than those of the north side, forming an airflow from the northern side to the southern side (Fig. 7(b-c)). This allows the dust aerosol can climb and transport along the Kunlun Mountains to the TP. Above the elevation of 5 km, the higher potential temperature area is distributed in the left TP, which can form the TP-SHAP. Firstly, the gradient of the potential temperature of the TP is much higher than that of the desert basin, making the

15

atmosphere unstable and easier to form atmospheric convective motion, and vertically suck dust aerosols around the TP (Jia et al., 2015; Wu et al., 2017; Feng et al., 2020). Secondly, at the same height, the potential temperature on the left side is higher than that on the right side, and the potential temperature slope on the left side is also larger, which can form a horizontal suction of dust aerosols over the desert.

5 As shown in Fig. 7(m-o), the relative humidity sinks from top to bottom above 4 km, while over the study site it gradually increases. Fig. A4n, on the other hand, shows the rapid drop in temperature over the desert caused by the invasion of cold air. The cold air invaded the TD and intersected with the warm air over the desert to form an upper-level cold front, and the cold air sank to force the desert basin to produce a strong convective motion (Fig. 7d, Fig. 7e, also analyzed in Sect. 3). At about 18:00 LT, with the participation of the upper-level cold front, the height of deep CBL reached its peak (Fig. 3c, Fig. 6(28), Fig. 10 7s).


5 Conclusion

In this study, the CDWL data and ERA5 reanalysis data were used to comprehensively analyze the development process of a representative deep convective boundary layer on the southern edge of the Taklimakan Desert and the northern foot of the Tibet Plateau on 6 June, 2022. The results indicate that the formation of this deep convective boundary layer stems from the 15 combined effects of multiple factors under complex terrain, including the Taklimakan Desert, slope terrain, the Kunlun Mountains, and the Tibet Plateau. The primary factor is the low-level jet and inversion layer, which provide sufficient momentum, energy, and material prerequisites for the development of the atmospheric boundary layer. Furthermore, the thermal effect facilitates the formation of the deep convective boundary layer. The schematic diagram of the development of the low-level jet and the deep convective boundary layer is shown in Fig. 8.

20 The formation of low-level jet in the study site is the combined effect of multiple factors. The first is the pressure gradient force and thermal difference between the north and south of the desert, which can form a background wind field from north to south. Secondly, the terrain blocking effect of the Tibet Plateau makes the northwest wind with lower wind speed deflect into the east wind with higher wind speed. Then, the terrain baroclinicity of the Tibet Plateau is easy for downhill airflow to form at night, and when the downhill airflow is superimposed on the background wind field, the wind speed of the background wind 25 field will be enhanced. Finally, the divergence structure of the upstream airflow and the convergence structure of the downstream airflow promote the development of the low-level jet.

The low-level jet and the inversion layer also play an important role in the development of the deep convective boundary layer. Firstly, the cold downhill wind is one of the factors contributing to the formation of the low-level jet, which can weaken the height and strength of the inversion layer, thereby reducing the energy demand for the breaking the inversion layer. Secondly, 30 the low-level jet can not only cause intermittent turbulence pulsations, but also inhibit the exchange of energy and material with the lower atmosphere, thus providing key material and energy supplements for the development of the deep convective boundary layer. Furthermore, the low-level jet can provide a dynamic basis for the subsequent development of the boundary

layer. Finally, the existence of the inversion layer and the low-level jet can inhibit the internal turbulent motion in the lower atmosphere, causing dust aerosols to accumulate near the ground, thus providing material basis and storing sufficient momentum and energy for the subsequent development of the deep convective boundary layer.



5 **Figure 8.** A schematic diagram of the development of low-level jet and deep convective boundary layer. (a) before sunrise. (b) before sunset.

The underlying surface of the desert itself has a strong heating effect on the atmosphere, and the corresponding thermal factors can catalyze the formation of the deep convective boundary layer. First of all, the accumulation of dust aerosols at low altitudes in the desert can form a furnace effect, rapidly transforming the material conditions for the development of the desert atmosphere boundary layer into thermal conditions, and promote the formation of the atmospheric superadiabatic expansion process. Secondly, the thermal effect of the Tibet Plateau sensible heat driven air-pump can suck up the atmosphere around the plateau and lift dust aerosols. Finally, the passage of a cold front can produce strong convective motion in the desert area. Overall, the results reveal the formation process of a typical deep convective boundary layer in the Taklimakan Desert, and also reflect the process of land-atmosphere momentum, energy and material exchange and transport between the Taklimakan Desert and the Tibetan Plateau. However, the data range of CDWL only covers a point area in MinFeng, and lacks the observation results of the mountains along the northern side of the Tibet Plateau. The follow-up work will combine with multi-site observations of Raman lidar and weather radar to study the transmission characteristics of dust to the Tibet Plateau, as well as conduct long-term statistical analysis of the effects of extreme weather such as drought and dust storm on the boundary layer height.

Data availability

The ERA5 data sets are publicly available from ECMWF website at <https://cds.climate.copernicus.eu>. The two datasets used in ERA5 are “ERA5 hourly data on pressure levels from 1940 to present” and “ERA5 hourly data on single levels from 1940 to present”. The CDWL data can be downloaded from https://figshare.com/articles/dataset/deep_CBL_lidar_datos/25434556 (Su et al., 2024). The Fengyun-4A meteorological satellite data of China can be downloaded from <http://www.nsmc.org.cn>.

Appendix A: The results of other observational data

The surface meteorological observation data on 6 June, 2022 (UTC + 8) are shown in Fig. A1. The CNR, horizontal wind speed, $\log_{10}(\text{TKEDR})$ and boundary layer height, vertical wind speed, wind direction during the field experiment in different seasons are shown in Fig. A2. The cloud coverage over the Taklimakan Desert is shown in Fig. A3. Fig. A4 shows temperature variations at different altitudes over the Taklimakan Desert.




Figure A1. The surface meteorological observation data on 6 June, 2022, (UTC+8). (a) temperature. (b) dust weather. (c) horizontal visibility. (d) relative humidity. (e) local atmospheric pressure.




Figure A2. The continuous observation results of CDWL in different seasons. (a-e) 10 Sep., 2021. (f-j) 12 Dec., 2021. (k-o) 3 Mar., 2022.




Figure A3. The cloud coverage over the Taklimakan Desert recorded by the Fengyun-4A meteorological satellite of China on 6 June, 2022 (UTC+8).

5




Figure A4. Temperature variations over the Taklimakan Desert during the period from 12:00 LT to 18:00 LT (UTC+8), observed at different atmospheric pressures or heights. The height represents the height from the ground at the study site. The purple circle represents the study site.

5 Appendix B: Methods

The CNR is obtained by the ratio of the signal area to the noise area of the power spectrum (Fujii and Fukuchi, 2005):

$$CNR = A_s/A_n \quad (B1)$$

where A_s is the signal area of the power spectrum, A_n is the noise area of the power spectrum.

The line of sight velocity (V_{los}) of CDWL is given by the following formula:

$$10 \quad V_{los} = \lambda f_i/2 \quad (B2)$$

where λ is the central wavelength of the emitted laser, f_i is the Doppler frequency shift for aerosols.

The wind vector in the atmosphere can be expressed by \vec{V} :

$$\vec{V} = (u, v, w) \quad (B3)$$

u, v, w represent the north-south velocity, east-west velocity, and vertical velocity, respectively.

15 When using Velocity Azimuth Display (VAD) scanning mode, the direction vector \vec{S} can be expressed as:

$$\vec{S} = (\cos\theta\cos\varphi, \sin\theta\cos\varphi, \sin\varphi) \quad (B4)$$

Where θ is the azimuth angle of the laser beam, and φ is the elevation of the laser beam.

From formula B2, B3, and B4, it can be concluded that (Browning and Wexler, 1968):

$$V_{los} = \vec{V} \cdot \vec{S}_m \quad (B5)$$

20 From formula B5, u, v, w can be calculated. The horizontal wind direction is calculated as follows:

Author contribution

HX conceived, designed the study. LS and JY performed the lidar experiments. LS performed the analysis of lidar data and ERA5 data. XW and QH provide the field experiment site and the local meteorological data. LS carried out the analysis and prepared the figures, with comments from other co-authors. LS, LC and JY wrote the manuscript. All authors read and approved the final manuscript.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements.

Thanks for the support of the Xinjiang Uygur Autonomous Region Meteorological Service and the Desert Meteorological Institute of China Meteorological Administration for this experiment and local meteorological data; The European Centre for Medium-Range Weather Forecasts for providing support with atmospheric reanalysis data.

Financial support.

This research has been supported by the National Natural Science Foundation of China (grant no. 42030612).

References

- Arnette, S. A., Samimy, M., and Elliott, G. S.: The effects of expansion on the turbulence structure of compressible boundary layers, *Journal of Fluid Mechanics*, 367, 67-105, 10.1017/S0022112098001475, 1998.
- Banakh, V., Smalikho, I., and Falits, A.: Estimation of the turbulence energy dissipation rate in the atmospheric boundary layer from measurements of the radial wind velocity by micropulse coherent Doppler lidar, *Optics Express*, 25, 22679-22692, 10.1364/OE.25.022679, 2017.
- Banakh, V. A. and Smalikho, I. N.: Lidar studies of wind turbulence in the stable atmospheric boundary layer, *Remote Sensing*, 10, 1219, 10.3390/rs10081219, 2018.
- Banakh, V. A., Smalikho, I. N., and Falits, A. V.: Estimation of the height of the turbulent mixing layer from data of Doppler lidar measurements using conical scanning by a probe beam, *Atmospheric Measurement Techniques*, 14, 1511-1524, 10.5194/amt-14-1511-2021, 2021.

- Basha, G. and Ratnam, M. V.: Identification of atmospheric boundary layer height over a tropical station using high-resolution radiosonde refractivity profiles: Comparison with GPS radio occultation measurements, *Journal of Geophysical Research*, 114, D16101, 10.1029/2008JD011692, 2009.
- Birch, C. E., Parker, D. J., Marsham, J. H., and Devine, G.: The effect of orography and surface albedo on stratification in the summertime Saharan boundary layer: Dynamics and implications for dust transport, *Journal of Geophysical Research*, 117, 10.1029/2011JD015965., 2012.
- Blackadar, A. K.: Boundary Layer Wind Maxima and Their Significance for the Growth of Nocturnal Inversions, *Bulletin of the American Meteorological Society*, 38, 283-290, 10.1175/1520-0477-38.5.283, 1957.
- Bonner, W. D.: Climatology of the low level jet, *Monthly Weather Review*, 96, 833-850, 10.1175/1520-0493(1968)096<0833:COTLLJ>2.0.CO;2, 1968.
- Browning, K. and Wexler, R.: The determination of kinematic properties of a wind field using Doppler radar, *Journal of Applied meteorology and climatology*, 7, 105-113, 10.1175/1520-0450(1968)007<0105:TDOKPO>2.0.CO;2, 1968.
- Caton Harrison, T., Washington, R., Engelstaedter, S., Jones, R., and Savage, N. H.: Influence of Orography Upon Summertime Low-Level Jet Dust Emission in the Central and Western Sahara, *Journal of Geophysical Research: Atmospheres*, 126, 1-24, 10.1029/2021JD035025, 2021.
- Che, J. and Zhao, P.: Characteristics of the summer atmospheric boundary layer height over the Tibetan Plateau and influential factors, *Atmos Chem Phys*, 21, 5253-5268, 10.5194/acp-21-5253-2021, 2021.
- Collis, R.: Lidar: a new atmospheric probe, *Quarterly Journal of the Royal Meteorological Society*, 92, 220-230, 10.1002/qj.49709239205open, 1966.
- Feng, X., Mao, R., Gong, D., Zhao, C., Wu, C., Zhao, C., Wu, G.-Q., Lin, Z., Liu, X., Wang, K., and Sun, Y.: Increased Dust Aerosols in the High Troposphere Over the Tibetan Plateau From 1990s to 2000s, *Journal of Geophysical Research: Atmospheres*, 125, 1-11, 10.1029/2020JD032807, 2020.
- Fiedler, S., Schepanski, K., Heinold, B., Knippertz, P., and Tegen, I.: Climatology of nocturnal low-level jets over North Africa and implications for modeling mineral dust emission, *Journal of Geophysical Research. Atmospheres*, 118, 6100 - 6121, 10.1002/jgrd.50394, 2013.
- Fujii, T. and Fukuchi, T.: *Laser remote sensing*, CRC press 2005.
- Garratt, J. R.: The atmospheric boundary layer, *Earth-Science Reviews*, 37, 89-134, 10.1016/0012-8252(94)90026-4, 1994.
- Ge, J., Liu, H., Huang, J., and Fu, Q.: Taklimakan Desert nocturnal low-level jet: climatology and dust activity, *Atmos Chem Phys*, 16, 7773-7783, 10.5194/ACP-16-7773-2016, 2016.
- Ge, J., Huang, J., Xu, C., Qi, Y., and Liu, H.: Characteristics of Taklimakan dust emission and distribution: A satellite and reanalysis field perspective, *Journal of Geophysical Research: Atmospheres*, 119, 11,772-711,783, 10.1002/2014JD022280, 2014.

- Guo, J., Miao, Y., Zhang, Y., Liu, H., Li, Z., Zhang, W., He, J., Lou, M., Yan, Y., Bian, L., and Zhai, P.: The climatology of planetary boundary layer height in China derived from radiosonde and reanalysis data, *Atmos Chem Phys*, 16, 13309-13319, 10.5194/acp-16-13309-2016, 2016.
- Han, B., Lü, S., and Ao, Y.: Development of the convective boundary layer capping with a thick neutral layer in Badanjilin: Observations and simulations, *Advances in Atmospheric Sciences*, 29, 177-192, 10.1007/s00376-011-0207-4, 2012.
- Han, B., Zhao, C., Lü, S., and Wang, X.: A diagnostic analysis on the effect of the residual layer in convective boundary layer development near Mongolia using 20th century reanalysis data, *Advances in Atmospheric Sciences*, 32, 807-820, 10.1007/s00376-014-4164-6, 2015.
- Han, Z., Ge, J., Chen, X., Hu, X., Yang, X., and Du, J.: Dust Activities Induced by Nocturnal Low-Level Jet Over the Taklimakan Desert From WRF-Chem Simulation, *Journal of Geophysical Research: Atmospheres*, 127, 10.1029/2021JD036114, 2022.
- Hoecker, W. H.: Three southerly low-level jet systems delineated by the Weather Bureau special pibal network of 1961, *Monthly Weather Review*, 91, 573-582, 10.1175/1520-0493(1963)091<0573:TSLJSD>2.3.CO;2, 1963.
- Holtstlag, A. and Boville, B.: Local versus nonlocal boundary-layer diffusion in a global climate model, *Journal of climate*, 6, 1825-1842, 10.1175/1520-0442(1993)006<1825:LVNBLD>2.0.CO;2, 1993.
- Hooper, W. P. and Eloranta, E. W.: Lidar Measurements of Wind in the Planetary Boundary-Layer - the Method, Accuracy and Results from Joint Measurements with Radiosonde and Kytoon, *J Clim Appl Meteorol*, 25, 990-1001, 10.1175/1520-0450(1986)025<0990:Lmowit>2.0.Co;2, 1986.
- Jia, R., Liu, Y., Chen, B., Zhang, Z., and Huang, J.: Source and transportation of summer dust over the Tibetan Plateau, *Atmospheric Environment*, 123, 210-219, 10.1016/J.ATMOENV.2015.10.038, 2015.
- Jiang, P., Yuan, J., Wu, K., Wang, L., and Xia, H.: Turbulence detection in the atmospheric boundary layer using coherent Doppler wind lidar and microwave radiometer, *Remote Sensing*, 14, 2951, 10.5194/amt-2021-288, 2022.
- Jones, C.: Recent changes in the South America low-level jet, *npj Climate and Atmospheric Science*, 2, 1-8, 10.1038/s41612-019-0077-5, 2019.
- Lai, Y., Chen, X., Ma, Y., Sun, F., Zhou, D., and Xie, Z.: Variation of Atmospheric Boundary Layer Height Over the Northern, Central, and Southern Parts of the Tibetan Plateau During Three Monsoon Seasons, *Journal of Geophysical Research: Atmospheres*, 128, 1-14, 10.1029/2022JD038000, 2023.
- Li, H., Yang, Y., Hu, X. M., Huang, Z., Wang, G., Zhang, B., and Zhang, T.: Evaluation of retrieval methods of daytime convective boundary layer height based on lidar data, *Journal of Geophysical Research: Atmospheres*, 122, 4578-4593, 10.1002/2016JD025620, 2017.
- Li, M., Xia, H., Su, L., Han, H., Wang, X., and Yuan, J.: The Detection of Desert Aerosol Incorporating Coherent Doppler Wind Lidar and Rayleigh–Mie–Raman Lidar, *Remote Sensing*, 15, 5453, 10.3390/rs15235453, 2023.

- Ma, Y., Xin, J., Wang, Z., Tian, Y., Wu, L., Tang, G., Zhang, W., de Arellano, J. V.-G., Zhao, D., Jia, D., Ren, Y., Gao, Z., Shen, P., Ye, J., and Martin, S. T.: How do aerosols above the residual layer affect the planetary boundary layer height?, *The Science of the total environment*, 151953, 10.1016/j.scitotenv.2021.151953, 2021.
- Ma, Y., Ye, J., Xin, J., Zhang, W., Vilà-Guerau de Arellano, J., Wang, S., Zhao, D., Dai, L., Ma, Y., Wu, X., Xia, X. a., Tang, G., Wang, Y., Shen, P., Lei, Y., and Martin, S. T.: The Stove, Dome, and Umbrella Effects of Atmospheric Aerosol on the Development of the Planetary Boundary Layer in Hazy Regions, *Geophysical Research Letters*, 47, 1-10, 10.1029/2020GL087373, 2020.
- Marshall, J. H., Parker, D. J., Grams, C. M., Grey, W. M. F., and Johnson, B. T. T.: Observations of mesoscale and boundary-layer circulations affecting dust uplift and transport in the Saharan boundary layer, *Atmos Chem Phys*, 8, 8817-8846, 10.5194/ACPD-8-8817-2008, 2008.
- Mathieu, N., Strachan, I. B., Leclerc, M., Karipot, A., and Pattey, E.: Role of low-level jets and boundary-layer properties on the NBL budget technique, *Agricultural and Forest Meteorology*, 135, 35-43, 10.1016/j.agrformet.2005.10.001, 2005.
- Matsumoto, S. and Ninomiya, K.: On the Mesoscale and Medium-scale Structure of a Cold Front and the Relevant Vertical Circulation, *Journal of the Meteorological Society of Japan*, 648-662, 1971.
- McNider, R. T. and Pielke, R. A.: Diurnal Boundary-Layer Development over Sloping Terrain, *Journal of the Atmospheric Sciences*, 38, 2198-2212, 10.1175/1520-0469(1981)038<2198:DBLDOS>2.0.CO;2, 1981.
- Meng, L., Yang, X.-h., Zhao, T., He, Q., Lu, H., Mamtimin, A., Huo, W., Yang, F., and Liu, C.: Modeling study on three-dimensional distribution of dust aerosols during a dust storm over the Tarim Basin, Northwest China, *Atmospheric Research*, 218, 285-295, 10.1016/J.ATMOSRES.2018.12.006, 2019.
- Nilsson, E. D., Rannik, Ü., Kulmala, M., Buzorius, G., O'Dowd, C. D., and O'Dowd, C. D.: Effects of continental boundary layer evolution, convection, turbulence and entrainment, on aerosol formation, *Tellus B: Chemical and Physical Meteorology*, 53, 441 - 461, 10.3402/tellusb.v53i4.16617, 2001.
- Ohya, Y., Nakamura, R., and Uchida, T.: Intermittent Bursting of Turbulence in a Stable Boundary Layer with Low-level Jet, *Bound-Lay Meteorol*, 126, 349-363, 10.1007/S10546-007-9245-Y, 2006.
- Pea, Gryning, S.-E., Hahmann, and A., N.: Observations of the atmospheric boundary layer height under marine upstream flow conditions at a coastal site, *Journal of Geophysical Research: Atmospheres*, 118, 1924-1940, 10.1002/jgrd.50175, 2013.
- Raman, S., Templeman, B., Templeman, S., Holt, T. R., Murthy, A., Singh, M. P., Agarwal, P., Nigam, S., Prabhu, A., and Ameenullah, S.: Structure of the Indian southwesterly pre-monsoon and monsoon boundary layers: Observations and numerical simulation, *Atmospheric Environment. Part A. General Topics*, 24, 723-734, 10.1016/0960-1686(90)90273-P, 1990.
- Rife, D. L., Pinto, J. O., Monaghan, A. J., Davis, C. A., and Hannan, J. R.: Global Distribution and Characteristics of Diurnally Varying Low-Level Jets, *Journal of Climate*, 23, 5041-5064, 10.1175/2010JCLI3514.1, 2010.
- Stensrud, D. J.: Importance of Low-Level Jets to Climate: A Review, *Journal of Climate*, 9, 1698-1711, 10.1175/1520-0442(1996)009<1698:IOLLJT>2.0.CO;2, 1996.

- Stull, R. B.: An Introduction to Boundary Layer Meteorology, Springer Science & Business Media, 10.1007/978-94-009-3027-8, 1988.
- Su, L., Lu, C., Yuan, J., Wang, X., He, Q., and Xia, H.: Measurement report: The promotion of low-level jet and thermal-effect on development of deep convective boundary layer at the southern edge of the Taklimakan Desert, Figshare [dataset], 10.6084/m9.figshare.25434556.v1, 2024.
- Tan, Z., Liu, Y., Zhu, Q., and Shao, T.: Impact of massive topography on the dust cycle surrounding the Tibetan Plateau, *Atmospheric Environment*, 264, 118703, 10.1016/J.ATMOSENV.2021.118703, 2021.
- Wang, L., Yuan, J., Xia, H., Zhao, L., and Wu, Y.: Marine mixed layer height detection using ship-borne coherent Doppler wind lidar based on constant turbulence threshold, *Remote Sensing*, 14, 745, 10.3390/rs14030745, 2022.
- Wang, L., Qiang, W., Xia, H., Wei, T., Yuan, J., and Jiang, P.: Robust Solution for Boundary Layer Height Detections with Coherent Doppler Wind Lidar, *Advances in Atmospheric Sciences*, 38, 1920-1928, 10.1007/s00376-021-1068-0, 2021.
- Wang, M., Xu, X., Xu, H., Lenschow, D. H., Zhou, M., Zhang, J., and Wang, Y.: Features of the deep atmospheric boundary layer over the Taklimakan Desert in the summertime and its influence on regional circulation, *Journal of Geophysical Research: Atmospheres*, 124, 12755-12772, 10.1029/2019JD030714, 2019.
- Wang, T., Han, Y., Huang, J., Sun, M., Jian, B., Huang, Z., and Yan, H.: Climatology of Dust-Forced Radiative Heating Over the Tibetan Plateau and Its Surroundings, *Journal of Geophysical Research: Atmospheres*, 125, 1-14, 10.1029/2020JD032942, 2020.
- Washington, R., Todd, M. C., Engelstaedter, S., M'bainayel, S., and Mitchell, F.: Dust and the low-level circulation over the Bodélé Depression, Chad: Observations from BoDEX 2005, *Journal of Geophysical Research*, 111, D03201, 10.1029/2005JD006502, 2006.
- Wexler, H.: A Boundary Layer Interpretation of the Low-level Jet, *Tellus A*, 13, 368-378, 10.1111/J.2153-3490.1961.TB00098.X, 1961.
- Wu, Wu, G., Guoxiong, Liu, Liu, Y., Yimin, Wang, Wang, T., Tongmei, Wan, Rj, Rijin, Liu, X.-d., Li, Li, W., Weiping, Zz, Zaizhi, Zhang, Zhang, Q., Qiong, Duan, Duan, A., Anmin, Liang, Xy, and Xiaoyun: The Influence of Mechanical and Thermal Forcing by the Tibetan Plateau on Asian Climate, *Journal of Hydrometeorology*, 8, 770-789, 10.1175/JHM609.1, 2007.
- Wu, G., He, B., Duan, A., Liu, Y., and Yu, W.: Formation and variation of the atmospheric heat source over the Tibetan Plateau and its climate effects, *Advances in Atmospheric Sciences*, 34, 1169-1184, 10.1007/s00376-017-7014-5, 2017.
- Wu, G., Liu, Y., He, B., Bao, Q., Duan, A., and Jin, F. F.: Thermal Controls on the Asian Summer Monsoon, *Scientific Reports*, 2, 1-6, 10.1038/srep00404, 2012.
- Wu, K., Wei, T., Yuan, J., Xia, H., Huang, X., Lu, G., Zhang, Y., Liu, F., Zhu, B., and Ding, W.: Thundercloud structures detected and analyzed based on coherent Doppler wind lidar, *Atmospheric Measurement Techniques Discussions*, 2023, 1-22, 10.5194/amt-2023-73, 2023.
- Wu, Y.-P., Feng, G.-L., and Li, B.-L.: Interactions of multiple atmospheric circulation drive the drought in Tarim River Basin, *Scientific Reports*, 6, 26470, 10.1038/srep26470, 2016.

- Xiao, F., Zhou, C., and Liao, Y.: Dust storms evolution in Taklimakan Desert and its correlation with climatic parameters, *Journal of Geographical Sciences*, 18, 415-424, 10.1007/s11442-008-0415-8, 2008.
- Xu, H., Wang, M., Wang, Y., and Cai, W.: Performance of WRF Large Eddy Simulations in Modeling the Convective Boundary Layer over the Taklimakan Desert, China, *Journal of Meteorological Research*, 32, 1011-1025, 10.1007/s13351-5 018-8001-1, 2018.
- Yang, F., He, Q., Huang, J., Mamtimin, A., Yang, X., Huo, W., Zhou, C., Liu, X., Wei, W., Cui, C., Wang, M., Li, H., Yang, L., Zhang, H., Liu, Y., Zheng, X., Pan, H., Jin, L., Zou, H., Zhou, L., Liu, Y., Zhang, J., Meng, L., Wang, Y., Qin, X., Yao, Y., Liu, H., Xue, F., and Zheng, W.: Desert Environment and Climate Observation Network over the Taklimakan Desert, *Bulletin of the American Meteorological Society*, 102, E1172-E1191, 10.1175/BAMS-D-20-0236.1, 2020.
- 10 Yang, X., Shen, S., Yang, F., He, Q., Ali, M., Huo, W., and Liu, X.: Spatial and temporal variations of blowing dust events in the Taklimakan Desert, *Theoretical and Applied Climatology*, 125, 669-677, 10.1007/s00704-015-1537-4, 2016.
- Yang, X., He, Q., Matimin, A., Yang, F., Huo, W., Liu, X., Zhao, T., and Shen, S.: Threshold velocity for saltation activity in the Taklimakan Desert, *Pure and Applied Geophysics*, 174, 4459-4470, 10.1007/s00024-017-1644-5, 2017.
- 15 Yuan, J., Xia, H., Wei, T., Wang, L., Yue, B., and Wu, Y.: Identifying cloud, precipitation, windshear, and turbulence by deep analysis of the power spectrum of coherent Doppler wind lidar, *Optics Express*, 28, 37406-37418, 10.1364/OE.412809, 2020.
- Yuan, J., Wu, K., Wei, T., Wang, L., Shu, Z., Yang, Y., and Xia, H.: Cloud seeding evidenced by coherent Doppler wind Lidar, *Remote Sensing*, 13, 3815, 10.3390/rs13193815, 2021.
- Zhang, H., Zhang, X., Li, Q., Cai, X., Fan, S., Song, Y., Hu, F., Che, H., Quan, J., Kang, L., and Zhu, T.: Research Progress on Estimation of the Atmospheric Boundary Layer Height, *Journal of Meteorological Research*, 34, 482-498, 10.1007/s13351-20 020-9910-3, 2020.
- Zhang, L., Zhang, H., Cai, X., Song, Y., Mamtimin, A., and He, Q.: Physical mechanisms of deep convective boundary layer leading to dust emission in the Taklimakan Desert, *Geophysical Research Letters*, 51, e2024GL108521, 10.1029/2024GL108521, 2024.
- Zhang, L., Zhang, H., Li, Q., Wei, W., Cai, X., Song, Y., Mamtimin, A., Wang, M., Yang, F., and Wang, Y.: Turbulent 25 mechanisms for the deep convective boundary layer in the Taklimakan Desert, *Geophysical Research Letters*, 49, e2022GL099447, 10.1029/2022GL099447, 2022.
- Zhang, Q., Cao, X., Wei, G., and Huang, R.: Observation and Study of Land Surface Parameters over Gobi in Typical Arid Region, *Advances in Atmospheric Sciences*, 19, 121-135, 10.1007/s00376-002-0039-3, 2002.
- Zhang, Q., Zhang, J., Qiao, J., and Wang, S.: Relationship of atmospheric boundary layer depth with thermodynamic processes 30 at the land surface in arid regions of China, *Science China Earth Sciences*, 54, 1586-1594, 10.1007/S11430-011-4207-0, 2011.
- Zhang, W., Guo, J., Miao, Y., Liu, H., Song, Y., Fang, Z., He, J., Lou, M., Yan, Y., Li, Y., and Zhai, P.: On the Summertime Planetary Boundary Layer with Different Thermodynamic Stability in China: A Radiosonde Perspective, *Journal of Climate*, 31, 1451-1465, 10.1175/JCLI-D-17-0231.1, 2017.

Zhang, Y., Seidel, D. J., and Zhang, S.: Trends in Planetary Boundary Layer Height over Europe, *Journal of Climate*, 26, 10071-10076, 10.1175/JCLI-D-13-00108.1, 2013.

Zhang, Y., Wu, Y., and Xia, H.: Spatial resolution enhancement of coherent Doppler wind lidar using differential correlation pair technique, *Optics Letters*, 46, 5550-5553, 10.1109/jlt.2022.3160294, 2021.

- 5 Zhou, C., Yang, F., Mamtimin, A., Huo, W., Liu, X., He, Q., Zhang, J., and Yang, X.: Wind erosion events at different wind speed levels in the Tarim Basin, *Geomorphology*, 369, 107386, 10.1016/j.geomorph.2020.107386, 2020.

Zhou, C., Liu, Y., He, Q., Zhong, X., Zhu, Q., Yang, F., Huo, W., Mamtimin, A., Yang, X., and Wang, Y.: Dust characteristics observed by unmanned aerial vehicle over the Taklimakan Desert, *Remote Sensing*, 14, 990, 10.3390/rs14040990, 2022.