

Remote Sensing detectability of airborne Arctic dust

Norman T. O'Neill¹, Keyvan Ranjbar², Liviu Ivănescu³, Yann Blanchard⁴, Seyed Ali Sayedain¹, Yasmin AboEl-Fetouh⁵

5 ¹Centre d'Applications et de Recherches en Télédétection, Université de Sherbrooke, Sherbrooke, Canada

²Flight Research Laboratory, National Research Council Canada, Ottawa, Canada

³Metrology Research Centre, National Research Council Canada, Ottawa, Canada

⁴Centre pour l'Étude et la Simulation du Climat à l'Échelle Régionale, Département des sciences de la Terre et de l'atmosphère, Université du Québec à Montréal, Montréal, Canada

10 ⁵Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

Correspondence to: N. T. O'Neill (Norman.T.ONeill@USherbrooke.ca)

Abstract. Remote sensing (RS) based estimates of Arctic dust are oftentimes overestimated due to a failure in separating out the dust contribution from that of spatially homogeneous clouds or low-altitude cloud-like plumes. A variety of illustrations are given with a particular emphasis on [questionable](#) claims of using brightness temperature differences (BTDs) as a signature indicator of Arctic dust transported from mid-latitude deserts or generated by local Arctic sources. While there is little dispute about the presence of both Asian and local dust across the Arctic, the direct RS detectability of airborne dust, as ascribed to satellite (MODIS and AVHRR) measurements of significantly negative brightness-temperature differences at 11 and 12 μm (BTD₁₁₋₁₂) has been misrepresented in certain cases. While it is difficult to account for all examples of strongly negative BTD₁₁₋₁₂ values in the Arctic, it is unlikely that airborne dust plays a significant role. One, much more likely contributor would be water [plumeclouds](#) in the Arctic inversion layer.

The RS detectability of the impact of Arctic dust (notably due to Arctic dust from local sources) can, however, be of significance. Sustained dust deposition can substantially decrease (visible to shortwave IR) snow and ice reflectance albedo (pan-chromatic reflectance) and the signal measured by satellite sensors. Significantly negative BTD₁₁₋₁₂ values would however only represent a limited area near the drainage basin sources according to our event-level case studies. The enhanced INP (Ice Nucleating Particle) role of local Arctic dust can, for example, induce significant changes in the properties of low-level mixed-phase clouds (cloud optical depth changes $\ll 1$) that can be readily detected by active and passive RS instruments. It is critical that the distinction between the RS detectability of airborne Arctic dust versus the RS detectability of the impacts of that dust be understood if we are to appropriately parameterize, for example, the radiative forcing influence of dust in this climate sensitive region.

30 **1 Introduction**

Vincent (2018) (VCT) reported on the use of MODIS and AVHRR thermal infrared (TIR) brightness temperature differences (BTDs) in the western Canadian Arctic (Beaufort Sea and Amundsen Gulf region) to detect the presence of “persistent low-level dust clouds” and dust deposited on ice, snow and water. A later publication (Bowen & Vincent, 2021) (B&V) argued that negative BTD_{11-12} ($BT_{11\ \mu m} - BT_{12\ \mu m}$) values were a unique signature of dust (without explicitly distinguishing between airborne and surface deposited dust) and that this measure could be directly used to estimate the relative spatial extent of dust in the Arctic. Those two water bodies (along with other place names and geographic features that are discussed below) appear in the Figure 1 map of the grey-shaded Canadian Arctic Archipelago (CAA) and associated Arctic and sub-Arctic regions.




Figure 1 – Map of the (grey-shaded) islands of the Canadian Arctic Archipelago (CAA). The map also includes Arctic and subarctic research sites in Alaska, northern Canada and Greenland (indicated with golden stars) and geographic features that are discussed in the main text.

Formatted: Font: 9 pt, Complex Script Font: 9 pt

40

We acknowledge the robustness of negative BTD_{11-12} values as a potential indicator of optically thick, airborne dust embedded in a normal-lapse-rate atmospheric layer or as a sign of accumulated dust deposition. We disagree with the affirmation that airborne dust clouds of Asian origin were commonly detected using passive, satellite-based remote sensing (RS). Springtime Asian dust, while representing a robust pan-Arctic seasonal event, yields, on average, very weak coarse mode (CM), roughly submicron, aerosol optical depths (AOD) at 550 nm^1 . The multi-year, six-station, pan-Arctic, AERONET/AEROCAN monthly-binned (geometric means) climatology of AboEl-Fetouh et al. (2020) (AeF) indicate (their Figure 7) that the Resolute Bay CM AODs are largely of the greatest amplitude across the six stations. Using that site as a reference, their CM AODs will were generally be $< \sim$ the Resolute Bay value ($0.016 \times 1.5^{\pm 1}$ during the Asian-dust dominated April, May springtime and $< \sim 0.0023 \times 1.2^{\pm 1}$ during the June, July, August (likely local dust) summertime). DODs (Dust Optical Depths) will be \sim CM AODs in the absence of any other significant CM source such as sea-salt or volcanic ash (or CM cloud in the absence of proper cloud screening).

Springtime Asian dust aerosols at such small CM AOD (DOD) values are difficult if not impossible to detect using passive satellite-based RS techniques at any wavelength. The BTD_{11-12} variation per unit change in DOD is $\sim -0.3 \text{ }^\circ\text{K}$ (see the discussion of Figure A3 Figure B1 for details). The corresponding change in BTD_{11-12} for a springtime Asian dust DOD of 0.016 (the Resolute Bay maximum) would be an undetectable $-0.005 \text{ }^\circ\text{K}$ (an amplitude that is significantly less than the nominal MODIS BTD_{11-12} noise figure of $0.07 \text{ }^\circ\text{K}$ (again, see the discussion of Figure A3 Figure B1).

While the monthly-averaged springtime Asian dust DODs are $< \sim 0.016$ there are springtime (mid-tropospheric) Asian dust events that do lead to more substantive DODs over the Arctic. In general these are limited to a few notable days in a given year, with individual DODs being $< \sim 0.4$: see Appendix A.4 for details on the roughly week-long dust intrusion over the Mount Logan (Yukon territory of Canada) in April of 2001 (DODs $< \sim 0.3$)² (associated with DODs of $< \sim 0.3$), Stone et al. (2007) for the roughly week long, mid-tropospheric dust intrusions of April 2002 over Barrow, Alaska (DODs $< \sim 0.4$), Zhao et al. (2022) for moderate DODs ($< \sim 0.1$) associated with single-day intrusions over Barrow in April of 2015, and Thulasiraman et al. (2002) for evidence that the sub-0.4 DODs of the April, 2001 event were arguably a broad west coast phenomenon stretching from (at least) low-Arctic Canada to the southern United States. DODs ~ 0.4 could incite a marginally detectable BTD_{11-12} signal ($-0.3 \text{ }^\circ\text{K} \times 0.4 \sim -0.12 \text{ }^\circ\text{K}$) and would be more easily identified in true-color and AOD imagery (at least over water).

VCT noted that a second dust source could have been locally generated dust storms (although dust from southern latitudes was claimed to be the major source). Indeed, Meinander et al. (2022) recently reviewed the importance of high latitude dust generated from high latitude local sources. However, even strong local dust plumes will likely not induce large DODs beyond the short temporal and spatial window associated with their detectable plume presence. Outside this window, the monthly averaged DOD upper limit reported by AeF make it very unlikely that DODs could be detected using passive, satellite-based RS in the thermal infrared red (TIR).

¹ Note that, unless otherwise stated, our AODs and DODs will always be referenced to a wavelength of 550 nm

² see Appendix A.4 for details on this well documented event

Empirical support for this affirmation is provided, for example, by Ranjbar et al.'s (2021) detection of a high-Arctic dust plume near its drainage-basin source: their MODIS BTD_{11-12} values (amplitude ~ 1.5 °K) are \sim the amplitude of the most extreme negative values (~ -1.0 °K) reported by VCT. However, Ranjbar's visible (532 nm) DODs were ~ 0.5 (a BTD_{11-12} to DOD sensitivity of $\text{dBTD}_{11-12}/\text{dDOD} \sim 1.5 / 0.5 = 3$ °K per unit DOD). The AeF Resolute Bay summertime³ CM AOD maximum ($0.0023 \times 1.2^{\pm 1}$) would, assuming approximate proportionality, produce generally undetectable BTD_{11-12} changes ($|\text{BTD}_{11-12}| < \sim 3 \times 0.0023 = 0.007$ °K).

~~We also~~As another source of independent support for the general weakness of Arctic DODs, we note that AeF's summertime DOD statistics are \sim DOD computations derived from the simulated local dust ("Arctic dust") polar map (Figure 1 of Kawai et al., 2023 [KA]). ~~Their~~The KA multi-year (2010-2019) "annual-mean vertically integrated mass concentrations" ("Arctic dust mass" with units of $\text{mg}\cdot\text{m}^{-2}$) yield DODs that \ll AeF's summertime CM AOD (see Figure A1 of Appendix A.3.2 where we compare the KA DOD simulations for the four AeF sites in or near the ~~Canadian Arctic Archipelago~~CAA). ~~The summertime constraint on their comparison comes from the Asian dust domination of AeF's springtime CM AOD (Asian dust is a dust component that is not modelled by the KA simulations).~~KA's Resolute Bay and Thule DODs are \sim AeF's summertime CM AOD analogues while being significantly less in the case of the dual PEARL and OPAL (Eureka) sites. ~~In Appendix A.3.3 we argue that one can aspire to make up the difference in the latter case inasmuch as the KA simulations appear to significantly underestimate local dust DOD relative to CALIOP derived estimates (based on a graphic provided by KA in their supplementary material). However, we also point out that this argument is inconclusive since CALIOP estimates of local dust DOD are unvalidated. In the end, the partial validation of KA's DOD simulations relative to the AeF CM AOD climatology is limited to a statement that the amplitude of their absolute DOD difference is $\ll 0.002$.~~

Having argued that Arctic DODs are, in general, at the margins of RS detectability, we must also take issue with B&V's affirmation that: "While it is possible that a substance other than mineral dust is causing large-scale negative BTD_{11-12} signatures in the polar environment, there is nothing in the literature to support this conjecture." We will present an alternative mechanism below involving inversion-layer liquid (droplet) clouds whose cloud optical depths (CODs) are sufficiently large to induce significant negative BTD_{11-12} signatures.

In general, there is often a tendency in the literature to significantly overestimate DOD magnitudes of Asian dust. It is no trivial matter to decouple such relatively small DODs from very large CM CODs ~~for~~of clouds that are often in the neighbourhood of those dust plumes. Such clouds may indeed result from dust nucleation: see for e.g., Hildner et al. (2010) and their discussion involving a high-altitude Asian dust plume that apparently nucleates into a highly-depolarizing cloud (captured by the AHSRL lidar above our PEARL [Polar Environment Atmospheric Research Lab] observatory at Eureka, Nunavut). In more general terms, Eck et al. (2009) noted the shortcomings in applying the AERONET (temporally-based) V2 cloud screening algorithm in the presence of spatially homogeneous clouds at Barrow, AK: spatially homogeneous (insufficiently variable) clouds such

³ More typical of average spring to-autumn-local DODs if one excludes the springtime Asian-dust dominated DODs (see Appendix A.3.2)

Formatted: Superscript

as thin cirrus are erroneously classified as dust (false-positive “dust”). More recently, Stone et al. (2014) underscored the potential for the same false-positive problem in their Barrow-based climatology of Arctic aerosols. Ranjbar et al. (2022) argued that the authors of a case study involving the transport of Asian dust into the high Arctic likely confused DOD with nearby COD and thereby significantly overestimated the DOD of a thin Asian dust plume about 7 km above the PEARL observatory. Analogous problems plague polar winter data. O’Neill et al. (2016) used lidar profiles and a spectral cloud-screening approach to estimate the large (starphotometer-derived) CM AOD errors that would be associated with the application of frequently inadequate (temporally-based) cloud screening paradigms to polar winter AODs-optical depths acquired at the PEARL-OPAL, Eureka site. The authors concluded that: “Spatially homogeneous clouds and low altitude ice clouds that remain after temporal cloud screening represent an inevitable systematic error in the estimation of AOD [more so for CM AOD]: the [positive bias] AOD error was estimated to vary from 78 to 210% at Eureka and from 2 to 157% at Ny-Ålesund.” In a not unrelated finding, Zamora et al. (2022) pointed out that the CALIPSO (CALIOP) classification algorithm was likely misclassifying wintertime “diamond dust” as mineral dust in their pan-Arctic analysis.

In terms of satellite-based estimates of DOD, B&V claimed that “average ~~aerosol optical depth (AOD)~~ is ~~was~~ “a proxy for dust aerosol concentration” and employed the 1998 to 2010 SeaWiFS AOD climatology of Hsu et al. (2012) to report a slight increase in AOD over the global oceans (and given their dust proxy assertion, a slight increase in DOD) in an apparent effort to support an increasing trend in their average RSED (“relative spatial extent of dust”) parameter over the Arctic and Antarctic.

This is yet another instance of DOD overestimation in the literature: AOD is almost universally dominated by ~~(non-dust)~~ fine mode (roughly submicron) particles and cannot be viewed as a proxy for “dust aerosol concentration” ~~(while there is evidence that fine mode dust exists, there is little evidence that it dominates other types of fine mode aerosols)~~. The proxy assumption is especially questionable when claiming to report a trend involving a minor AOD species (dust) coupled with a satellite AOD product that is less accurate than the AOD generated from ground-based AERONET data (for which a DOD trend analysis would be a challenge on its own merits): the bias error (amplitude $\gg 0.01$) between the ~~SeaWiFS-SeaWiFS~~ AODs and AERONET AODs (Figure 2 of Hsu et al., 2012) ~~are~~ (for example) ~~is~~ \gg the spread of AeFs spring and summer geometric standard deviation envelope for Resolute Bay ($0.0051 \times 2.53.0^{\pm 1}$).

A more realistic DOD satellite product over the Arctic is the MIDAS (ModIs Dust AeroSol) data set (Gkikas et al., 2021). The MIDAS reanalysis system is based on MODIS AODs coupled with MDFs (mineral dust fraction; a semi-intensive parameter of ~~DOD~~ \div AOD) derived from MERRA-2 (Modern-Era Retrospective analysis for Research and Applications version 2) whose key components are the Goddard GOCART aerosol model and GEOS (Goddard Earth Observing System). The MIDAS annual DOD (arithmetic) mean for the 2003 – 2017 period (for the more accurate retrievals over water around Resolute Bay) is $\sim 0.01 \pm 0.02$ (where 0.02 is a computed estimate of retrieval uncertainty rather than a standard deviation). The AeF Resolute Bay value of $[0.0051 \times 3.0^{\pm 1}]$ is contained within the MIDAS uncertainty envelope.

Commented [SAS1]: Why is this value different from what see at line 110?

Commented [NTO2R1]: Good catch (I corrected the value above)

(water mixed with ice) clouds have been reported in the literature: de Boers et al. (2009) and Shupe et al. (2015) provide lidar / radar supported illustrations of ~~explicit~~ mixed-phase events at Eureka and Barrow respectively⁴. The former paper reported a 4-year (2004-2007) frequency-of-occurrence (%) series (three-month-wide bins) of combined Barrow and Eureka results showing a general predominance of “SON” autumn bins (~ 10%) at <~ 1.5 km mean cloud-base-height⁵ for single-layer, mixed-phase stratiform clouds. The latter paper provided a 2-year Barrow climatology (which is more relevant to the Barrow-region focus of the analyses that follows) showing that the monthly occurrence (%) was highest in October (~ 40%) while being only moderate and of lower altitude from March (~ 10%) to ~ 25% in April and May (with a strong preponderance of sub-1-km liquid occurrence). Yi et al., 2019 reported comprehensive satellite-based (MODIS and CALIOP) water cloud (“Arctic fog”), March-to-October statistics for a large Arctic Ocean region roughly centered north of Barrow. Their water cloud limitation to fog (cloud base height = cloud height – cloud thickness being < 1000 feet [300 meters]) was, however, rather restrictive with respect to the types of liquid cloud events that we investigate below (events requiring that the water cloud be imbedded in significant temperature slices of the Arctic inversion layer).

Nearly all our negative BTD_{11-12} Barrow examples shared one feature that is rarely mentioned in typical BTD_{11-12} literature; the ubiquitous and strong Arctic temperature-inversion up to altitudes ~ 1 km that occurs during the polar winter and summer (see, again, the Supplementary material “BTDBarrowSummer.xlsx” and “BTDBarrowWinter.xlsx” ~~X~~ for details on our analysis ~~of BTD_{11-12} results~~ near Barrow and Bradley et al., 1993 and Palo et al., 2017 for statistical summaries of the Arctic inversion layer). Inversion-layer cloud events are the most easily detectable instances of a fundamental principle; that a “warm-cloud” in the Arctic inversion layer (in an atmosphere clear of higher altitude clouds) transforms the more common negative-lapse-rate BTD_{11-12} signature from a generally positive to negative dependency with the degree of negativity being dependent (for a given surface emissivity, inversion-layer strength and water vapour load) on the COD and effective particle radius (see, for example, Key, 2002; and Liu et al., 2004).

We generated MODTRAN-simulated BTD_{11-12} vs BT_{11} graphics ~~employing a variety of whose~~ input parameters ~~to encompassed~~ a wide variety of COD and particle size conditions about ~~the a~~ specific March 22, 2015 event at Barrow (see the discussion ~~of that event of associated with~~ Fig. 1 below). The resulting BTD_{11-12} vs BT_{11} patterns are shown in ~~Figure A3~~ Figure B1 while Table 1 below presents a descriptive summary of ~~both empirical and simulated patterns~~ those simulations⁶ (the optical details and boundary layer conditions associated with each BTD_{11-12} vs BT_{11} pattern are given in Appendix ~~B.1~~ B.1). The “convex downward” shape of a large-COD water cloud in an inversion layer will produce almost exclusively negative values that fundamentally depend on the COD = 0 and ∞ singularities on the BTD_{11-12} vs BT_{11} patterns of ~~Figure A3~~ Figure B1

⁴ The reader will note that for the ~~specific~~ purposes of this paper we do not distinguish between liquid clouds and mixed phase clouds. Below we argue that the ice COD in mixed phase clouds is typically negligible compared to the liquid water COD

⁵ With a plume thickness of <~ 600 m. These are the 3-year results for Eureka: the single Barrow year of 2004, with a SON occurrence of 26% was more coherent with the Shupe et al. results.

⁶ as well a simulations and empirical evidence from the literature

(while a high-altitude ice or liquid cloud will produce, as per the upper left graphic of [Figure A3](#)[Figure B1](#), generally positive BTD_{11-12} values).

Table 1 - Empirical (E) & simulated (S) BTD_{11-12} vs BT_{11} results for a variety of cloud-~~or~~-dust plumes embedded in positive temperature-lapse-rate (inversion layer) or negative lapse-rate regions. Blue, red and ~~gold-brown~~ text refer respectively to water clouds, ice clouds and dust plumes.

Temperature lapse rate (dT/dz)	Emissivity Slope ⁺ ($d\epsilon/d\lambda$)	BTD_{11-12} vs BT_{11} pattern	Cloud or dust plume & surface scenarios	Citations / comments
Negative	Negative	Convex upwards ¹	Low-altitude to mid-altitude water cloud	Baum et al. (2000) (S) ^a , Key (2002) (S) ^b
Positive	Negative	Convex downwards ²	Low-altitude (inversion layer) water clouds	This study ^c (S & E) Key (2002) ^d (S)
Negative	Negative	Convex upwards	High altitude ice clouds	This study (S ^e & E).
Positive	Negative	Convex downwards	Low-altitude (inversion layer) ice clouds	This study (S ^f)
Negative	Positive ^g	Convex downwards	High altitude Asian dust plumes	Various (S & E) This study (S) ^h
Positive	Negative	Convex upwards	Low altitude (inversion layer) Asian dust plume parameters	This study (S-)

¹ Otherwise known as concave downward. These curves generally (but not always) consist of positive BTD_{11-12} values

² Otherwise known as concave upward. These curves generally (but not always) consist of negative BTD_{11-12} values

^a Surface to top-of-water-cloud temperature decrease of $\cong 297$ to $\cong 284$ °K (low altitude water cloud)

^b Surface to top-of-water-cloud temperature decrease of 263.16 to 238.65 °K (mid altitude water cloud)

^c Surface, top-of-inversion-layer ~~plumecloud~~ and top of high-altitude ~~plumecloud~~ temperatures $\cong 256$, 262 and 213 °K. See the captions of Figures S6 and the discussion of [Figure A3](#)[Figure B1](#) for details

^d Surface to top-of-water-cloud temperature increase of 235.66 to 253.15 °K

^e Same approximate values as footnote c. The high altitude ice-cloud literature is resplendent with convex-upward BTD_{11-12} vs BT_{11} examples (see, for example, Figure 3 of the tropical atmosphere simulation of Ackerman et al., 2010)

^f Same approximate values as footnote c

^g Han et al. (2012) point out that quartz is a major component of Asian dust. Quartz-dominated dust is generally characterized by positive $dk/d\lambda$ (resulting in positive $d\epsilon/d\lambda$ according to our calculations)

^h The dust parameterizations that we employed in our BTD_{11-12} vs BT_{11} simulations (see our dust refractive index survey in Appendix B) were sometimes referenced to results for African dust

ⁱ Many There are many S & E examples in the literature of convex-upward-downward BTD_{11-12} vs BT_{11} patterns (see, for e.g., Figures 2 and 4 of Zhang et al., 2006). The generally negative nature of those patterns are a well-known high altitude signature of Asian dust plumes.

Table glossary


E – empirical results, S – simulated results

ϵ – emissivity spectrum of a surface composed of water, ice or dust soil

$dk/d\lambda$ & $d\epsilon/d\lambda$ – 11 to 12 μm spectral slope of respectively, the complex part of the refractive index & emissivity

Figure [4a-2a](#) shows a radar profile for the specific inversion-layer (Figure S6) Barrow illustration of the March 22, 2015 event (with Figure [4b-2b](#) providing a zoom of the radar profile in the inversion layer between -0 and 2 km). Figure [4c-2c](#) shows

both the MODIS-measured moderate-amplitude BTD_{11-12} values and the MODTRAN-simulated values (details in the figure caption). This demonstrates how (i) a warm, liquid-water, inversion-layer (negative BTD_{11-12}) cloud (whose lower and upper boundaries are explicitly shown in Figure S6 and Figure 2b) coupled with the positive BTD_{11-12} presence of a cold (negative or normal lapse rate) ice-cloud around 6-km-altitude (during roughly the first half of the displayed time period) produces systematically varying BTD_{11-12} values that oscillate between the negative to positive extremes of the two phenomena. This demonstrates explicitly how (i) a warm liquid-water, inversion layer plume (whose lower and upper boundaries are explicitly shown in Figure S6 and Figure 1b) produces systematically-varying BTD_{11-12} values that oscillate from negative to positive as driven by the negative BTD_{11-12} predominance of the inversion layer plume and positive BTD_{11-12} predominance of a cold (negative or normal lapse rate) ice cloud around 6 km altitude and (ii) that MODTRAN radiative transfer simulations were largely successful in capturing the BTD_{11-12} oscillations.



Formatted: Subscript

Formatted: Subscript

Commented [NTO6]: We'll keep the legend as is (with the possibility that Yann might agree that "MPL" should be "NSHRL")

215 Figure ~~4-2~~ – Radar backscatter coefficient profiles acquired at the ARM Barrow site on Mar. 22, 2015; ~~from~~ (a) 0 to 10 km altitude ~~and~~ (b)
a zoom ~~of (a)~~ from 0 to 2 km altitude. ~~The green and red curves (as measured by the combination of lidar and radar backscatter profiles)~~
~~indicate, respectively, the bottom and top of what we inferred to be a water-dominated, mixed-phase cloud that was partially contained in an~~
~~inversion layer. The dynamical details of this event, including radiosonde temperature profiles, –including the juxtaposition of what we~~
~~inferred to be a water-dominated, mixed-phase, inversion-layer plume (as measured by the combination of radar backscatter profile and~~
220 ~~NSHSRL backscatter and depolarization profiles) are given in the caption of Figure S6 and its caption. The solid black curves of Figure (c)~~
~~shows, for the MODIS pixel nearest to the Barrow site, – both the MODIS-measured BT₁₁ time series (upper graph) and MODIS-measured~~
~~BTD₁₁₋₁₂ values time series (lower graph). – (for the pixel nearest to the Barrow site) and the~~ The solid red curves show their MODTRAN-
~~simulated values analogues. These simulations employing employed representative input parameters representative of during~~ that day (see
the captions of Figure S6 and S7 for further details).

Specific details on the vertical extent of the mixed phase backscatter coefficient profile, the ~~ir~~ ~~respective water vs ice~~ COD
225 contributions and their relationship to the temperature (inversion layer) profile provided by the Barrow radiosonde profiles are
presented in the discussion of Figure S6. In those details, we argue that the radar profiles provide key information about the
upper boundary of the water / mixed phase ~~plumecloud~~ (beyond the upper bound defined by the extinction limit of the lidar)
and its attendant extension to altitudes where there was actually a strong temperature inversion.

The same (Table 1) inversion layer BTD₁₁₋₁₂ convexity reversal should apply to warm, low-level ice-clouds in an Arctic
230 inversion layer: however as shown in ~~the lower left graphic of Figure A3~~ Figure B1(b), the amplitude of the convex downward
pattern can be insufficient to move the pattern into the negative BTD₁₁₋₁₂ range. In any case, we did not, in our survey of
significantly negative, near-Barrow BTD₁₁₋₁₂ (MODIS) values, find any obvious lidar / radar retrievals dominated by
synchronous, low-level, optically-thick ice-clouds in the inversion layer. This is not unexpected since the CODs of near-surface
ice clouds are substantially smaller than those of water clouds (see Shonk et al., 2019 for a general statement and specific
235 examples in the Dec. 29, 2006 Eureka case study of de Boer et al., 2009 as well as Sections 3d vs 4c of Zuidema et al., 2005
for a May 1 – 18, 1998 case study at a floating ice camp ~ 600 km northwest of Barrow). Morrison et al. (2012) also point out
the dominance of water CODs over ice CODs across a ~~5-5~~ day (May 11 – 15, 2011) Eureka event and underscore the
persistence of Arctic mixed-phase clouds in general. This dominance of water COD over ice COD was also found in our Figure
~~4-2~~ case study (see the caption of Figure S6 for details).

240 We would note that a convex-down to convex-up pattern reversal would occur ~~for when comparing high altitude dust clouds~~
~~with~~ warm dust clouds located in an Arctic inversion layer: the generally negative contours for ~~cold dust clouds~~ the former
would be transformed to positive contours (see ~~the bottom right graphic of Figure A3~~ Figure B1(d)) that would actually
confound the classic negative BTD₁₁₋₁₂ signature of cold, high-altitude dust clouds (see for example, Figures 3 (MODIS BTD₁₁₋
245 ₁₂) and Figure 4 (simulations) of Zhang et al., 2006). Support for this affirmation ~~for the case of local dust~~ comes indirectly
from Ranjbar et al. (2021): the lapse rate in the Lake Hazen case was, in all likelihood, an “~~classical~~” (inversion-layer-free)
rate of decreasing temperature with increasing altitude. This results in the negative BTD₁₁₋₁₂ plume values reported in that
paper (the defining BTD₁₁₋₁₂ vs BTD₁₁ pattern is more in the nature of the convex-down shapes of ~~the upper right graphic of~~
Figure A3 Figure B1(c)).

The overarching message of this section is that negative BTD₁₁₋₁₂ values in the Arctic are not a unique signature of the
250 pervasiveness of dust across the Arctic. The BTD₁₁₋₁₂ signature of airborne dust in the inversion layer would generally be too

Commented [NT07]: This change was necessitated by what
perceived to be a less than clear linkage between this figure and
Figure S6

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

weak to detect and of the wrong sign with respect to the classical negative signature of desert dust plumes in a normal lapse rate environment. DODs in the Arctic are generally too small to induce significant BTD_{11-12} amplitudes. The BTD_{11-12} signature of deposited dust can be significantly negative but, as suggested by our case study on snow deposition of local dust (see the following section), tend to be spatially limited to dust dominated regions of the drainage basin source.

255 4 RS detectability of dust impacts

An affirmation of the general marginality of airborne-dust RS detectability in the Arctic is not to say that the impacts of Asian or local dust are necessarily marginal in terms of satellite-based RS. The cumulative deposition of local dust associated with weak DODs can (as also noted by VCT) be substantial over seasonal or longer time scales with significant changes in surface reflectance (and attendant impacts on early snow melt coupled with a feedback effect of even greater reflectance changes).

260 ~~AVHRR remote sensing imagery dating back as far as 1991 was employed by Woo et al. (1991) to argue that dust covered areas on the Fosheim Peninsula (region of– Eureka) were the first to experience snow melt (Ranjbar et al., (2021) showed a true color May 18, 2019 reflectance image of northern Ellesmere Island with significant browning of the generally white ice/snow surface near known drainage basins (including Lake Hazen and the Fosheim Penninsula region of Eureka) showed image browning regions on MODIS “Corrected Reflectance (True Color)” (RGB) images that corresponded to Woo’s “dark spot” regions).~~ Figure S8 shows, what we argue, are examples of dust deposition on snow or ice in the neighbourhood of
265 ~~southern-CAA~~ drainage basins whose flow dynamics have induced local dust plumes. This illustrates how the accumulation of local dust deposition by dust plumes produces, (i) true-color images of significantly modified snow reflectance in the visible spectral region ($< 60\%$ average reflectance change⁷; ~~c.f. the “Dusty Snow” reflectance changes in the Figure 1a spectra of Painter et al., 2007~~) but only (ii) weakly positive BTD_{11-12} dust-deposition signatures near their drainage basin sources (while
270 also showing that significantly negative BTD_{11-12} signatures do occur in what are likely the very localized pure dust regions of the drainage basins). This, as indicated in the legend of Figure S8, is likely a BTD_{11-12} difference ~~that can be ascribed to a pure~~ ~~resulting from the greater~~ dust-surface emissivity ~~difference between bands of band 11-12 and relative to band 1211~~. These illustrations strongly suggest that, significantly negative signatures of dust on snow or ice are likely to be ~~extremely-very~~ limited in their spatial extent.

275 The reflectance effects associated with the deposition of Asian dust on snow are less evident. Asian dust deposition was detected by ground teams at higher altitudes (where sources of local dust would be unlikely) in the Mount Logan (Yukon) region of the St. Elias range during the strong April 2001 Asian dust event (Zdanowicz et al., 2006). The authors suggested that up to 45% of the airborne dust mass abundance was deposited in the snow (over a 9 day period) and that the mechanism for deposition was scavenging by snow flakes. MODIS ~~corrected true color reflectance~~ RGB images show no obvious impact:
280 this effort to determine an impact is not helped by these agents of deposition arguably confounding / camouflaging the

⁷c.f. the “Dusty Snow” reflectance changes in the Figure 1a spectra of Painter et al., 2007

Formatted: Font: (Default) Times New Roman, Complex
Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Complex
Script Font: Times New Roman

darkening impact of dust. Zhao et al. (2022) employed a variety of ground- and satellite-based, passive and active RS data as well as surface nephelometer measurements of CM scattering coefficient to investigate the albedo (spectrally integrated reflectance) impact of dust deposition on snow during March 14, 2013 and April 20, 2015 Asian dust events over Alert (Nunavut) and Barrow, Alaska respectively⁸. The CM scattering coefficients coupled with estimates of the dust plume (mid-tropospheric) altitude over each site suggested direct deposition links between the dust plumes and the surface dust (they did not attempt to elaborate on any explanation of the deposition dynamics). ~~A radiative transfer model was then employed by~~ The authors employed a radiative transfer model to argue that daily dust deposition events could reduce snow surface (panchromatic) albedo by as much as 2.3% at Barrow and 1.9% at Alert. These albedo reductions would be quite substantial if dust depositions (in relatively unperturbed snow conditions) were allowed to accumulate over, for example the 9-day period of the April 2001 event. However, the simulations of Groot Zwaafink et al. (2016) on the substantially greater contribution of local Arctic dust (versus Asian or African dust) to dust deposition suggests that ~~the mid-latitude (Asian and African)~~ dust would, in general, play a secondary reflectance perturbation role compared to local dust.

A second substantial impact of Arctic dust particles is associated with their role as INP (ice nucleation particles) and their indirect effect on cloud dynamics. The core message of Kawai et al. (2023) was not a statement about the weak optical influence of local Arctic dust, but rather a simulation-based affirmation that local dust was the dominant INP source in the lower Arctic troposphere during summer and fall ~~(much more dominant than Asian dust in general)~~. A similar statement concerning the dominance of local dust over Asian dust as INPs was made by Xi et al. (2022) based on INP (droplet freezing) measurements made near the source of local dust plumes at the sub-Arctic Lhù'ään Mán' (Kluane Lake) site in the Canadian Yukon territory. Barr et al. (2023) reported on the greater INP activity of local dust (from glacial drainage basins on the southern Alaskan coast) relative to Arctic dust from low-latitude sources. Tobo et al. (2019) described the important role of local dust as an INP source in the Svalbard region and noted that the high ice nucleating ability of the local dust was likely governed by the presence of organic matter. Shi et al. (2022) analyzed the radiative forcing impacts of local Arctic dust (what they called "HLD" for high-latitude dust ~~as an acronym for Arctic dust~~ whose source is in the Arctic ~~or sub-Arctic~~). Their simulations (roughly supported by INP comparisons with measured INPs carried out at 9 stations) show, for example, that HLD INPs likely instigated a maximum depletion in the liquid water path (LWP) of mixed phase clouds in the fall season ~~over the Canadian Arctic Archipelago~~ (and lesser but still significant LWP changes during the summer and winter seasons). Those LWP depletions ($\sim 8 \text{ g}\cdot\text{m}^{-2}$) amount to water COD reductions of $< \sim 1.5$ (at any wavelength for which the Mie extinction efficiency (Q) is \sim the optically large-particle asymptotic value of ~ 2). Such COD changes (along with their associated extinction coefficient profile change) would be readily detected using standard passive & active, satellite-based sensors (from the visible to the thermal IR).

⁸ those two intensive-analysis days were supplemented by neighbouring days for which CM nephelometer measurements suggested the dust-event extended beyond those two core days.

310 5 Conclusions

We presented a variety of examples showing how direct RS-based estimates of CM Arctic dust were oftentimes excessively large due to a failure in separating out the contribution of CM clouds (or cloud-like optical contributions). A particular emphasis was placed on a paper by Vincent (2018) who reported an optically strong airborne dust presence in the western Canadian Arctic that was ascribed to dust of Asian origin or dust from local sources. While we do not dispute the presence of both Asian and local dust in the Arctic, the direct RS detectability of airborne dust (attributed to satellite-measured values of significantly negative BTD_{11-12} values) was likely-almost surely misrepresented. While it is difficult to account for all examples of strongly negative BTD_{11-12} , it is very unlikely that airborne dust plays a major RS role in any case other than plumes of strong DOD ($> \sim 0.5$). One, much more likely contributor would be water plumeclouds (or, more generally stated, water dominated, mixed phase clouds) in the Arctic inversion layer.

320 The RS detectability of the impact of Arctic dust and notably Arctic dust from local drainage basin sources can, however, be of significance. Sustained dust deposition can substantially decrease the (visible to shortwave IR) snow and ice reflectance and the attendant signal measured by satellite sensors (while significantly negative BTD_{11-12} values represent an extremely limited area according to our event level case studies). The reportedly-strongersubstantial INP (Ice Nucleating Particle) role of local Arctic dust can, for example, induce significant changes in the properties of low-level mixed phase clouds (optical depth changes $< \sim$ unity) that can be readily detected by active and passive RS instruments. It is clearly critical that the distinction between the RS detectability of Arctic dust versus the RS detectability of the impacts of Arctic dust be understood if we are to properly account for and model the radiative forcing impacts of dust in this-the climate sensitive Arctic region.

6 Appendices

Appendix A – Intensive and extensive microphysical and optical parameters of local and Asian dust

330 A.1 Effective radius relationships for spherical particles

The effective radius for spherical particles is defined by Hansen & Travis (1974) (HT) as:

$$r_{eff} = \frac{\int r^3 \frac{dn}{d \ln r} d \ln r}{\int r^2 \frac{dn}{d \ln r} d \ln r} = \frac{\int (D/2)^3 \frac{dn}{d \ln D} d \ln D}{\int (D/2)^2 \frac{dn}{d \ln D} d \ln D} = \frac{1}{2} \frac{\int D^3 \frac{dn}{d \ln D} d \ln D}{\int D^2 \frac{dn}{d \ln D} d \ln D} = \frac{1}{2} D_{eff} \quad (A1)$$

where the very last relation amounts to a definition of D_{eff} . Equation (A1) can then be recast in terms of total particle-surface and particle-volume concentration:

$$335 \quad D_{eff} = \frac{\frac{2^3}{3\pi} \int \pi \left(\frac{D}{2}\right)^3 \frac{dn}{d \ln D} d \ln D}{\frac{2^2}{\pi} \int \pi \left(\frac{D}{2}\right)^2 \frac{dn}{d \ln D} d \ln D} = \frac{3 \int \frac{dv}{d \ln D} d \ln D}{2 \int \frac{ds}{d \ln D} d \ln D} = \frac{3v}{2s} \quad (A2a)$$

From equation (A1) the effective diameter can be recast as:

$$D_{eff} = \frac{3 \int \frac{4}{3} \left(\frac{D}{2}\right) \frac{ds}{d \ln D} d \ln D}{2 \int \frac{ds}{d \ln D} d \ln D} = \frac{\int D \frac{ds}{d \ln D} d \ln D}{\int \frac{ds}{d \ln D} d \ln D} = \frac{\int D \frac{ds}{d \ln D} d \ln D}{\int \frac{ds}{d \ln D} d \ln D} = \langle D \rangle_{\omega} = ds/d \ln D \quad (A2b)$$

the weighted mean of D where the weight $\omega = ds/d \ln D$. Ginoux (2003) argues that the shape of dust particles are, in general, better represented by prolate ellipsoids (see the following section).

340 A.2 Computation of D_{eff}

Kawai et al. (2023) (KA) employed Kok's particle-volume [size](#) distribution as the basis of their multi-year simulations (ultimately it was the starting point⁹ of their computations of seasonally averaged particle-mass columnar densities). Kok's particle-volume size distribution (his equation (6)) is related to his particle-number size distribution (his equation (5) by $dV_d/d\ln D_d = C_N/C_V D_d^3 dN_d/d\ln D_d$. We can¹⁰ recast this as;

$$\begin{aligned}
 345 \quad d\tilde{v}/d\ln D &= C_N/C_V [3/(4\pi) 2^3] [(4/3)\pi (D/2)^3] d\tilde{n}/d\ln D \\
 &= (C_N/C_V 6/\pi)v_{sp}(D) d\tilde{n}/d\ln D \\
 &= C_{Kok}v_{sp}(D) d\tilde{n}/d\ln D \quad (A3)
 \end{aligned}$$

where $v_{sp}(D) = (4/3)\pi (D/2)^3$ is the volume of a spherical particle of radius $D/2$ and $C_{Kok} = C_N/C_V \times 6/\pi$. However C_{Kok} ($= 0.144$ for Kok's C_N and C_V values of 0.9539 and $12.62 \mu\text{m}$ respectively) is not close to unity as would be expected for small dust particles. Equation (A3) is apparently the correct inter-distribution relationship between Kok's (Figure 6) "normalized" number and "normalized" volume size distributions¹¹.

However equation (A2a) applied to Kok's normalized distributions gives unrealistic estimates of the effective radius ($0.78 \mu\text{m}$)¹². Those normalized distributions were not defined and so we have to tentatively conclude that the normalization precluded the application of equation (A2a)¹³. Dust particles are not sufficiently large to have substantial non-sphericity effects and so one expects the departure of C_{Kok} from unity to be relatively small. Ginoux (2003) cited Okada et al. (2001) to indicate that dust particles near their source (Chinese desert sites) displayed an ellipsoid aspect ratio (λ) of ~ 1.5 and that moderately higher values of 2 showed no significant departure from sphericity (their Figure 5, for example, shows that simulated particle volume distributions for $\lambda = 2$ were quite close to the spherical-particle AERONET inversions for 6 sites near or in the desert sources of dust). Accordingly we can, in general, treat dust particles as being approximately spherical ($d\tilde{v}/d\ln D \sim v_{sp}(D) d\tilde{n}/d\ln D$) and the light-grey broken-line open-circles in the Supplementary material ("Local_dust_PSDs.xlsx") represent the appropriate distribution¹⁴ for the employment of equation (A2a). This yields a D_{eff} value of $5.40 \mu\text{m}$ ($r_{eff} = 2.70 \mu\text{m}$).

⁹ The emission (source) particle-volume size distribution

¹⁰ -dropping his "d" (dust) subscript, using lower case letters for these point-volume parameters and \tilde{n} and \tilde{v} for their "normalized" distributions

¹¹ as verified by the fact that the black, solid-line, open-circle ($C_{Kok}v_{sp}(D) d\tilde{n}/d\ln D$) curve is very close to Kok's gold, solid-line full-circle curve ($d\tilde{v}/d\ln D$) in the Supplementary material ("Local_dust_PSDs.xlsx")

¹² Versus, for example, a volume-weighted geometric mean diameter (VMD) of $6.51 \mu\text{m}$ (the AERONET-inversion type of calculation)

¹³ Meaning, that Kok's normalized distributions were not equally proportional to their physical representations (the physical representations being symbolized by the hatless variables in his section)

¹⁴ Meaning that we treat the distributions as being spherical-particle distributions

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

A.3 DOD computations for KA's local dust particles

A.3.1 DOD mass efficiency (DOD_m)

If V is the columnar, particle-volume abundance ρ is the dust particle density and A is the particle-number abundance then the particle-mass abundance (A_m) in the case of the KA local dust simulations¹⁵ (or any unimodal particle-volume or particle-mass distribution) is given by;

$$A_m = \rho V \cong \rho \frac{4}{3} \pi r_{eff}^3 A \quad (A4a)$$

$$A \cong \frac{A_m}{\rho \frac{4}{3} \pi r_{eff}^3} \cong \frac{A_m}{m} \quad (A4b)$$

where the concept of intensive parameters averaged over a unimodal particle size distribution is discussed, for e.g., in O'Neill et al. (2005). If the dust extinction cross section is σ and Q is the dust extinction efficiency, then the dust optical depth (DOD) is;

$$\tau \cong \sigma A \cong \sigma \frac{A_m}{\rho \frac{4}{3} \pi r_{eff}^3} = Q \pi r_{eff}^2 \frac{A_m}{\rho \frac{4}{3} \pi r_{eff}^3} = \frac{Q A_m}{\frac{4}{3} \rho r_{eff}} \quad (A5)$$

Employing the mean r_{eff} of 2.7 μm from the Kok distributions (Appendix A.2 above) yields $x_{eff} = 2 \pi r_{eff} / \lambda = 33$ for $\lambda = 0.5 \mu\text{m}$. Q approaches an asymptote ~ 2.3 for values of the product $x_{eff}(1 - m_r) > \sim 10$ (Figure 16.3 of Hinds, 1999 with m_r being the real part of the refractive index) and refractive indices representative of dust¹⁶. Employing the MITR¹⁷ density of 2.6 $\text{g}\cdot\text{cm}^{-3} \rightarrow 2.6 \times 10^9 \text{mg}/\text{m}^3$ yields;

$$\tau \sim \frac{2.3 A_m}{\frac{4}{3} (2.6 \times 10^9) \text{mg}/\text{m}^3 \times (2.65 \times 10^{-6}) \text{m}} \sim 0.250 \times 10^{-3} (\text{mg}/\text{m}^2)^{-1} A_m \quad (A6)$$

One can define a "DOD mass efficiency" $\times 10^4$ (DOD per unit columnar mass abundance) as;

$$DOD_m \times 10^4 = \frac{\tau}{A_m} \sim 2.5 (\text{mg}/\text{m}^2)^{-1} \quad (A7)$$

A.3.2 DOD extracted from KA's particle-mass abundances

KA's multi-year A_m averages derived from their Figure 1, at the position of AeF's sites that are in or near the ~~Canadian Arctic Archipelago (CAA)~~, were employed to compute the DOD estimates of Table A.1 below. We note that "Eureka" is meant to represent the similar environments of two Eureka sites (the sea-level OPAL site and the 610-meter elevation "PEARL" (Ridge lab) site): the resolution of KA's A_m values allows no such distinction to be made for the KA simulations.

Table A1 – Local dust DODs derived from the A_m (mass abundances) of KA's Figure 1 (at the position of the four AeF sites within or near the CAA).

¹⁵ The parameter that they call "vertically integrated ... mass concentrations"

¹⁶ 1.53 – 0.0078i for the MITR ("Mineral transported") class of dust (the "opdat" directory of the OPAC package). Hess et al.'s MITR density transforms as $2.6 \text{g}/\text{cm}^3 = 2.6 \times 10^3 \text{mg}/(10^{-2} \text{m})^3 = 2.6 \times 10^9 \text{mg}/\text{m}^3$

¹⁷ Optical and μ physical parameters are listed in Table 1c of Hess et al. (1998).

Formatted: Font: 10 pt, Complex Script Font: 10 pt

Formatted: Font: 10 pt, Complex Script Font: 10 pt

Site	Lat., Long.	A_m [mg/m ²]	Computed DOD (τ)
Resolute Bay	75°N, 95°W	14.68	0.0037
Eureka (PEARL & OPAL)	80°N, 86°W	0.32	0.000079
Thule	77°N, 69°W	3.16	0.00079

Figure A1 shows a comparison between AeF's CM AOD (geometric mean) summertime (JJA) climatology and the KA simulations. The KA simulations include no Asian dust component: their yearly means ~~are expected to~~ should, in principle be ~ the summertime AeF values (if the latter can be assumed to be dominated by local dust). The AeF CM AODs of the April & May springtime period are (as per AeF), largely dominated by Asian dust with values $> \sim 0.0036-06$ (and thus largely off-scale in Figure A1 ~~only one springtime point shows up~~). We also note that PEARL and OPAL are separated by a distance of only 15-km distance with their elevations being, respectively, 610 and 10 m. The simple subtraction labelled "OPAL – PEARL" in Figure A1 is arguably a better measure of local dust DOD than their individual summertime means (assuming the local dust is largely limited to altitudes less than that of the PEARL (Ridge lab) site: ~~–~~) in this case the ~~The KA precision~~ KA underestimation is for the Resolute Bay, Thule and "OPAL – PEARL" (Eureka) site would then nominally be $\ll 0.001$. This is a number that one hesitates to quote given the preponderance of uncertainties that plague both the simulations and the measurements (for e.g., the quality, respectively of local dust emissions about a given AeF site given the coarse KA spatial resolution of $1.9^\circ \times 2.5^\circ$ and the nominal AERONET AOD error ~ 0.01). The most optimistic affirmation is arguably that the summertime dust estimates are the same order of magnitude.




Figure A1 – Local dust DODs derived from the A_m (mass abundances) of KA's Figure 1 (at the position of the four AeF (AERONET) sites in or near the CAA) along with summertime (JJA-averaged) CM AODs from AeF's four sites (Canadian Arctic Archipelago).

A.3.3 KA-model “underestimation” of local DOD

Figure S4 of KA's supplementary material suggests that their simulated 550 nm “annual-mean zonally averaged dust AOD” significantly underestimates the local DOD relative to the CALIOP 532 nm estimate of local DOD¹⁸. The 3rd column of Table A2 (corresponding to the 2nd column of Table A1) shows the DODs (τ) computed from KA's Figure 1 at AeF's AERONET sites (the last column of Table A1 has been inserted into the 4th column of Table A2). The computations include a “correction” of the simulated KA's DODs ($DOD - \Delta DOD$) to yield values that would account for the gap between the simulated KA DODs and the CALIOP DODs. The result, relative to AeF's summertime estimate is a better comparison for Eureka and Thule and a rather large overestimate for Resolute Bay. The limitation of such a “correction” is the credibility of

¹⁸ Underestimates by a bias which we label ΔDOD . There are, for example, biases of $\Delta DOD \sim -0.005$ at the 75°N lat. of Resolute Bay and ~ -0.003 at the near 80°N lat. of Eureka and Thule: these values were estimated from KA's Figure S4 ($(-1) \times [CALIOP \text{ curve} - \text{red KA simulation curve}]$).

Formatted: Centered

Formatted: Line spacing: single

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Font:

CALIOP in classifying and estimating local DOD in the Arctic. There is no validation in the literature of CALIOP’s capability in classifying local dust plumes: ~~the The conclusion of this exercise has to be limited to an argument that the KA simulations succeed in producing a correct order of magnitude at the three CAA sites simply that there is no reason to change the order of magnitude conclusion of the previous section.~~

420

Table A2 – Computed DOD values (DOD (τ)) employing the nominal **DOD_m** value of equation (A4A7) ~~above~~ and corrected DOD values (DOD - Δ DOD) ~~computed by increasing the nominal DOD values of Table A1 to values suggested by the bias CALIOP DOD values from KA’s Figure S4.~~

Site	Lat., long.	Computed DOD	“Corrected DOD” (DOD - Δ DOD ^a)
Resolute Bay	75°N, 95°W	0.0037	0.0087
Eureka (PEARL & OPAL)	80°N, 86°W	0.000079	0.0031
Thule	77°N, 69°W	0.00079	0.0038

^a The simulations underestimate the CALIOP “truth”: their Δ DOD bias is accordingly negative

A.4 Estimation of Mount Logan DODs during the Asian dust event of April 2001

425

Table A3 shows visually extracted DODs from NAAPS simulations over the region of Mount Logan (Yukon) during the April 11 to 19, 2001 Asian dust event. The DOD values are the midpoints of the standard NAAPS color-scale bins. If there is no NAAPS DOD (no NAAPS dust at the position of Mount Logan) then the bin is assigned a value of 0.0¹⁹. The arithmetic average of all the DOD values below is $\langle DOD \rangle = 0.13$. This table supports the discussion ~~above~~ surrounding the well documented Asian dust event of 2001 and the dust deposition consequences in the Mount Logan region.

Commented [NT09]: Deleted this text because of its excessive (obvious) detail

430

Table A3 – Visually determined NAAPS DODs²⁰ at Mount Logan, YK (60° 34’ N, 140° 24’ W) during the April, 11 to 19, 2001 Asian dust event.

ddhh (UT)	DOD _{bin center}	ddhh (UT)	DOD _{bin center}	ddhh (UT)	DOD _{bin center}
1100	0.0	1400	0.0	1700	0.15
1106	0.0	1406	0.15	1706	0.15
1112	0.0	1412	0.3	1712	0.15
1118	0.0	1418	0.3	1718	0.15
1200	0.0	1500	0.3	1800	0.15
1206	0.15	1506	0.3	1806	0.15
1212	0.15	1512	0.3	1812	0.0
1218	0.15	1518	0.3	1818	0.0
1300	0.15	1600	0.15	1900	0.15
1306	0.0	1606	0.15	1906	0.15

¹⁹ or a value of 0.15 if the edge of the 0.15-valued (yellow) colored plume cannot be visually separated from the position of Mount Logan

²⁰ https://www.nrlmry.navy.mil/aerosol-bin/aerosol/display_directory_all_t.cgi?DIR=/web/aerosol/public_html/globaer/ops_01/noramer/200104&TYPE=


1312	0.15	1612	0.15	1912	0.0
1318	0.15	1618	0.15	1918	0.0

Appendix B – Computational details in support of Table 1

B.1 MODTRAN simulations of BT_{11-12} vs BT_{11} patterns for liquid water, ice and dust (Optical constants and radiative transfer computations at 11 and 12 μm)

B.1 MODTRAN simulations of BT_{11-12} vs BT_{11} patterns for liquid water, ice and dust

Figure B1 shows MODTRAN BTD₁₁₋₁₂ vs BT_{11} patterns over a “Snow/Ice” surface for water and ice clouds/plumes (left hand graphs) and dust plumes (right hand graphs) over a “Snow/Ice” surface at high altitude (top graphs) and within an a low-altitude inversion layer (bottom graphs) are shown in Figure B1. The general atmospheric conditions and cloud/plume parameterization represent a range of values that include the specific conditions of Figures 4a2a, 4b2b, S6 and S7 (conditions of Barrow and its surroundings on March 22, 2015). The nominal temperatures employed in the MODTRAN simulations at a the (snow/ice) surface, inversion-layer–cloud/plume–top and high–altitude cloud–top were, respectively 255.56, 262.05 and 212.66 °K. These graphs provide support for all the simulation (S) based BT_{11-12} vs BT_{11} classifications (BTD_{11-12} vs BT_{11} pattern characterization) of Table 1.







Formatted: Heading 3

Formatted: Indent: Before: -0.01", After: -0.01", Line spacing: 1.5 lines

Formatted: Font: Italic, Complex Script Font: 10 pt, Italic English (United Kingdom), Subscript

Formatted: Left, After: -0.01", Line spacing: single

Formatted: Left, Indent: Before: -0.01", After: -0.01"



470

Figure B1 – $BTD_{11.12}$ vs BT_{11} simulations for different types of clouds/plumes over a “Snow/Ice” surface (surface of Feldman, 2014). Graphs Left-hand graphs, top & bottom respectively (a) and (b) respectively: — ice and liquid cloud at a high altitude and within a low altitude temperature inversion layer. Graphs Right-hand graphs, top & bottom (c) and (d) respectively: Asian dust plume-cloud at a high altitude and within a low altitude inversion layer. The wavelength-dependent optical depth (τ) is reported at a wavelength of 550 nm in order to make a link with optical effects in the visible wavelength region.

475

The eccentric nature/significant curvature of the low level “Reff = 3 μm” red curve in the bottom right graph of Figure (d) B1 appears to be an approach to suggest a balanced radiative transfer condition wherein there is little change in BTD_{11} with increasing plume-cloud optical depth (an approach to an idealistic singularity of a straight vertical line). We determined that this effect was largely due to non-linearities in the spectra of the extinction efficiency and the fact that 550 nm referencing of the cloud/plume optical depth; (τ) is fixed, in the MODTRAN simulations, at a single wavelength of 550 nm.

480

Formatted: Line spacing: single

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Line spacing: single

Formatted: Line spacing: single

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Font: 9 pt, Complex Script Font: 9 pt

Formatted: Font: 9 pt, Complex Script Font: 9 pt

The optical properties of the liquid and dust particles were generated with a Mie Code (MiePlot4621, written by Philip Laven²¹) (<http://www.philiplaven.com/mieplot.htm>), using the refractive index of water (Hale and Query, 1973) and dust²² (Volz, 1973), for monodisperse particles. The optical properties of the ice crystals were extracted from Ping Yang's database (Yang et al., 2013) and correspond to a modified-gamma distribution with effective variance of 0.1 (Petty and Huang, 2011) of severely roughened column aggregates (Yang et al., 2013). This is the same distribution that is assumed in the Collection 6 MODIS cloud product (Holz et al., 2016).

Weak DODs associated with high-altitude Asian dust, the BTD_{11-12} to DOD sensitivity ($dBTD_{11-12} / dDOD$) would be best represented by a slope near $DOD = 0$ ($\tau = 0$ on the graphs) for the case of the near $1.5 \mu\text{m}$ peak radius of springtime Asian dust (see, for example, the right panel of Figure 16 of Burton et al. (2012) and Figure 3 of AeF for the springtime Asian dust particle size distribution). This yields a value (from the detailed numerical results employed in generating these graphs) of $-0.30 \text{ }^\circ\text{K}$ per unit change in DOD.

The brightness temperatures correspond to the EOS-1 TERRA MODIS spectral response functions for bands 31 (max. at $11.0 \mu\text{m}$) and 32 (max. at $11.9 \mu\text{m}$), downloaded from the Satellite Application Facility for Numerical Weather Prediction (NWP SAF) website²³, this website:

<https://nwp-saf.eumetsat.int/site/software/rttov/download/coefficients/spectral-response-functions/>

A nominal noise figure for MODIS Band 31 (the $11 \mu\text{m}$ band) is $0.05 \text{ }^\circ\text{K}$ (the cloud-discrimination ATBD of Team et al., 2010). Given a roughly equivalent (incoherent) noise for band 32 (the $12 \mu\text{m}$ band) yields a BTD_{11-12} noise value of $\sqrt{2} \times 0.05 = 0.07 \text{ }^\circ\text{K}$

B.2 Choice of refractive indices at 11 and 12 μm

The refractive indices of water droplets and ice crystals are, as per the previous section, relatively well constrained and known. The observed dust refractive indices in the literature are principally dependent on dust composition (see, for e.g., Volz, 1972; Koepke et al., 1997; Rothman et al., 2009 and Sadrian et al., 2023); this dependence impacts the behavior of the BTD_{11-12} vs BTD_{11} patterns. A unique choice of refractive index based on dust composition is not possible given the diversity of dust types that characterize Asian and local dust over the Arctic (coupled with the often incomplete information on their composition). Figure B2 illustrates the infra-red refractive index and (simulated) surface emissivity (ϵ) spectrum of water and ice particles as well as two distinctly different complex refractive indices of dust for two frequently referenced citations (Koepke, 1997

²¹ <http://www.philiplaven.com/mieplot.htm>

²² see Appendix B.2 for a discussion of our choice of dust refractive index.

²³ <https://nwp-saf.eumetsat.int/nwp-saf.eumetsat.int/site/software/rttov/download/coefficients/spectral-response-functions/>

<https://nwp-saf.eumetsat.int/site/software/rttov/download/coefficients/spectral-response-functions/>

Formatted: Font color: Text 1

Field Code Changed

Formatted: Font: Symbol

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Default Paragraph Font, Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Default Paragraph Font, Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) Times New Roman, Font color: Text 1, Complex Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Left

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font

Formatted: Default Paragraph Font


Formatted: Default Paragraph Font

Formatted: Normal



510 with principal components of quartz and clay and Volz, 1973 with principal components of clay, illite, and kaolinite). Those different refractive indices result in significantly different emissivity spectral slopes between 11 and 12 μm .

515 We chose the Volz (1973) refractive indices essentially because the 11 to 12 μm spectral slope of the derived ϵ values were of the same sign as the ϵ slopes presented in VCT. This choice was underpinned by different levels of empirical and simulated evidence: the BTD_{11-12} vs BT_{11} “convex downward” pattern (generally indicating negative BTD_{11-12} values) for a normal ($dT / dz < 0$) lapse rate (Figure B1 and Table 1) is coherent with satellite-based Asian dust measurements (as per footnote g of Table 1) as well as the negative lapse rate and attendant negative BTD_{11-12} across the optically thick local-dust Lake Hazen plume that was discussed ~~above~~. Another level of empirical evidence in the Arctic ~~is~~ was the significantly negative BTD_{11-12} values in the dust emission regions of the drainage basins ~~snow near the sources of dust~~ on Eglinton and Bank’s Island (see Figure S8 and its caption). We also found moderately negative BTD_{11-12} values in areas around Lake Hazen that were clearly the result of dust deposition on snow and ice.

Formatted: Font: Symbol



Formatted: Centered






Figure B2 – Real (a) and imaginary (b) parts of the refractive index (n and k) and surface emissivity (ϵ) spectra (c) employed for the MODTRAN simulations of Appendix B1. The ϵ spectra were approximately computed using the formulations of Masuda et al. (1988).

525

B.2 Survey of dust refractive indices (11 and 12 μm)

This survey was carried out to support the refractive index decisions employed in the generation of Figure B1. Table B1 is a concise summary of the survey results.

530 Table B1: Survey of published dust / sand refractive indices ($n - ik$) and complementary data at 11 and 12 μm

Source	Users	n (11 μm) ^a	n (12 μm) ^a	k (11 μm) ^a	k (12 μm) ^a	Comment
Koepke et al. (1997) ^{a,24}	Li et al., (2018)	1.829	1.774	0.209	0.428	“Substance”: “Sand” “Type”: “Quartz and clay”
Rothman et al. (2009)		1.6 ^d	2.0 ^d	0.02 ^d	0.1 ^d	“HITRAN Quartz” ^b Rothman citation is from Han et al. (2012)
Sadriani et al. (2023)		1.595	1.510	0.085	0.005	Taklimakan, China (sample S1014 for e.g.) Carbonate absorption was identified in S1014

²⁴ [https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Sand/Quartz_and_clay_\(Koepke_et_al_1997\)/sand_koepke_1997.r](https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Sand/Quartz_and_clay_(Koepke_et_al_1997)/sand_koepke_1997.r)

Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Space After: 8 pt, Line spacing: Multiple 1.0
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Not Italic, Complex Script Font: 9 Not Italic
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Complex Script Font: 9 pt
 Formatted: Font: 9 pt, Font color: Accent 1, Complex Script Font: +Headings CS (Times New Roman), 9 pt, English (United Kingdom)

Volz (1972) ²⁵	Table 3 of Zhang et al. (2006) ^e	1.618	1.548	0.105	0.095	“Substance”: “Dust” “Sample”: “Rainout dust” No comment about the composition of the dust
Volz (1973) ²⁶	Gu et al. (2003)	1.825	1.778	0.302	0.184	“Substance”: “Sand” “Type”: “Saharan sand”; “Sample”: “Barbados” From Volz (1973): “clay, illite, and kaolinite along with traces of quartz”
Wald et al. (1998)				no <i>k</i> data but $\varepsilon = 0.92$	no <i>k</i> data but $\varepsilon = 0.90$	Case of “pure quartz”. Smaller particles (0 – 70 μm diameter ²⁷) show a significantly greater positive $d\varepsilon/d\lambda$ (emissivity (ε)) ^e slope from 11 to 12 μm

^a the Koepke & Volz links: EODG (Earth Obs. Data Group), Aerosol Refractive Index Archive (consulted Oct. 4, 2023); The italicized text are the labels from that site. These numbers are very close to the OPAC numbers (Hess et al., 1998)

^b “HITRAN is also tested because quartz is one of the major components of Asian dust [Jeong, 2008]” (Han et al., 2012). This quote is a statement about the % weight of components in bulk samples of Asian dust (see Jeong, 2008).

^c Precisely 11.03 and 12.02 μm respectively for all table entries except the OPAC and HITRAN values.

^d These are coarse visual estimates from Figure 2 of Han et al. (2012).

^e The *n* and *k* values of their Table 3 “Dust like” component are those of Volz (1972) and not, as claimed, Koepke et al. (1997). Their reference to Gu et al. (2003) as a data source is more directly a reference to Volz et al. (1973)

^f The authors state (in the discussion of their Figure 1 *c* spectra) that: “Because the samples were made by grinding, the 0.75 μm sample is numerically dominated by < 5 μm diameter particles”

^g and thus, perhaps, a greater absorption coefficient and thus *k* slope.

Appendix C – Acronym and symbol glossary

AERONET	World-wide NASA network of combined sunphotometer / sky-scanning radiometers manufactured by CIMEL Électronique. See AERONET website²⁷ for documentation and data downloads
AHSRL	Arctic High Spectral Resolution Lidar
AOD	The community uses “AOD” to represent anything from nominal aerosol optical depth which hasn’t been cloud-screened to the conceptual (theoretical) interpretation of aerosol optical depth. In this paper we use it in the latter sense and apply adjectives as required.
AquaQUA	Polar orbiting NASA satellite whose payload includes the MODIS-Aqua multi-band imager. Aqua passes south to north over the equator in the afternoon (originally known as EOS PM-1)
ARM	Atmospheric Radiation Measurements
β	Backscatter cross section ($\text{m}^{-1} \text{sr}^{-1}$)
BT, BTD	Brightness Temperature, Brightness Temperature Difference
CAA	Canadian Arctic Archipelago
CM	Coarse mode (generally referring to particles of super-micrometer radii)

²⁵ [https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Dust/Rainout_dust_\(Volz_1972\)/dust_volz_1972.ri](https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Dust/Rainout_dust_(Volz_1972)/dust_volz_1972.ri)

²⁶ [https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Sand/Saharan_sand/Barbados_\(Volz_1973\)/sand_volz_1973.ri](https://eodg.atm.ox.ac.uk/ARIA/data?Sand_and_Dust/Sand/Saharan_sand/Barbados_(Volz_1973)/sand_volz_1973.ri)

²⁷ <https://aeronet.gsfc.nasa.gov/>

Commented [SAS10]: As we removed Table B1 in the last version of the paper, I’m wondering if we should ignore the sentence “These numbers are very close to the OPAC numbers” not?

Commented [NTO11R10]: I don’t understand your question Ali. We didn’t remove Table B1 ... it is here and in the reviewer’s file. I agreed with one thing though: I deleted “(see section “Row removed from Table B1” below).”

Formatted: No underline

Formatted: Heading 2

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Commented [NTO12]: I don’t know how you formatted this text (it should be just a regular table ... I can’t figure out how to insert a new entry!). Please make a regular Word table and then insert the definition of the CAA (Canadian Arctic Archipelago)

Formatted: Indent: Before: 0.01"

COD	Cloud Optical Depth
DOD	Dust Optical Depth
FM	Fine mode (generally referring to particles of sub-micrometer radii)
HLD	High Latitude Dust
HYSPLIT	HYbrid Single-Particle Lagrangian Integrated Trajectory
INP	Ice Nucleation Particle
KAZRGE	Ka ARM Zenith Radar (KAZR)²⁸ GEneral mode. Zenith pointing Doppler radar operating at 35 GHz (8.6 mm)
LWP	Liquid Water Path
MISR	Multi-angle Imaging SpectroRadiometer
MODIS	Moderate Resolution Imaging Spectroradiometer
NSHSRL	North Slope High Spectral Resolution Lidar
Terra	Terra passes from north to south across the equator in the morning
TIR	Thermal InfraRed

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

Formatted: Indent: Before: 0.01"

7 Author contribution

Norm T. O’Neill: Writing – original draft preparation – review & editing, Conceptualization, Methodology, Investigation, Formal analysis, Visualization, Validation, Project administration, Data curation, Funding acquisition, Resources. **Keyvan Ranjbar:** Writing – review & editing, Conceptualization, Investigation, Software, Formal analysis, Visualization, Validation. **Liviu Ivănescu:** Writing – review & editing, Conceptualization, Investigation, Software, Formal analysis, Visualization, Validation. **Yann Blanchard:** Writing – review & editing, Conceptualization, Investigation, Software, Formal analysis, Visualization, Validation. **Seyed Ali Sayedain:** Writing – review & editing, Conceptualization, Validation. **Yasmin AboEl-Fetouh:** Writing – review & editing, Conceptualization.

540 8 Competing interests

The authors declare that they have no conflict of interest.

²⁸ <https://adc.arm.gov/metadata/html/nsakazrgeC1.b1.html>

9 Acknowledgements

This work was supported by the ESS-DA program of the Canadian Space Agency (CSA). We also gratefully acknowledge the longstanding research support provided by Natural Sciences and Engineering Research Council of Canada (NSERC). With respect to the Barrow (Utqiagvik) North Slope Alaska (NSA) analysis: data were obtained from the Atmospheric Radiation Measurement (ARM) user facility, a U.S. Department of Energy (DOE) office of science user facility managed by the Biological and Environment Research program. Valuable in-kind support was provided respectively by the AEROCAN network of Environment and Climate Change Canada (ECCC) and the NASA AERONET network. We also acknowledge the efforts of Antonis Gkikas of the National Observatory of Athens (NOA) Institute for Astronomy, Astrophysics, Space Applications and Remote Sensing (IAASARS) in providing detailed MIDAS DOD retrievals and uncertainties over the Arctic.

10 Financial support

This research has been supported by the Canadian Space Agency (grant nos. 16SUASACIA and 21SUASACOA), and the Natural Sciences and Engineering Research Council of Canada (grant nos. RGPIN-2017-05531 and RGPIN-2023-04943).

9 Acknowledgements

This work was supported by CANDAC (the Canadian Network for the Detection of Atmospheric Change) via the Canadian NSERC (National Sciences and Engineering Research Council) CREATE Training Program in Arctic Atmospheric Science, as well as by grant 21SUASACOA from the ESS-DA program of the Canadian Space Agency (CSA). Financial support was also provided by O'Neill's NSERC Discovery Grant. With respect to the Barrow (Utqiagvik) North Slope Alaska (NSA) analysis: data were obtained from the Atmospheric Radiation Measurement (ARM) user facility, a U.S. Department of Energy (DOE) office of science user facility managed by the Biological and Environment Research program. Valuable in-kind support was provided respectively by the AEROCAN network of Environment and Climate Change Canada (ECCC) and the NASA AERONET network. We also acknowledge the efforts of Antonis Gkikas of the National Observatory of Athens (NOA) Institute for Astronomy, Astrophysics, Space Applications and Remote Sensing (IAASARS) in providing detailed MIDAS DOD retrievals and uncertainties over the Arctic.

10 Financial support

This research has been supported by the National Aeronautics and Space Administration, Goddard Space Flight Center (grant no. SURA-GSTR-4100-NASA), the Canadian Space Agency (grant nos. 500353/15FASTA12, 19ATORA07, 16SUASACIA and 21SUASACOA), and the Natural Sciences and Engineering Research Council of Canada (grant nos. CREATE 384996-10, RGPCC 433842-2012 and RGPIN 2017-05531).

Commented [SAS13]: If there is any number associated to NSERC grant, please let me know to ask ACP guys to add to the funding section of the paper

Commented [NTO14R13]: All the following "Financial support" text (from my 2023 SMF vs FMF AMT paper) applies; "Financial support. This research has been supported by the National Aeronautics and Space Administration, Goddard Space Flight Center (grant no. SURA-GSTR-4100-NASA), the Canadian Space Agency (grant nos. 500353/15FASTA12, 19ATORA07, 16SUASACIA and 21SUASACOA), and the Natural Sciences and Engineering Research Council of Canada (grant nos. CREATE 384996-10, RGP 433842-2012 and RGPIN-2017-05531)."

Formatted: Heading 1

570 10 Acronym and symbol glossary

AERONET	World-wide NASA network of combined sunphotometer / sky-scanning radiometers manufactured by CIMEL Électronique. See AERONET website ²⁹ for documentation and data downloads
AHSRL	Arctic High Spectral Resolution Lidar
AOD	The community uses "AOD" to represent anything from nominal aerosol optical depth which hasn't been cloud-screened to the conceptual (theoretical) interpretation of aerosol optical depth. In this paper we use it in the latter sense and apply adjectives as required.
Aqua	<u>Polar orbiting NASA satellite whose payload includes the MODIS-Aqua multi-band imager. Aqua passes south to north over the equator in the afternoon (originally known as EOS PM-1)</u>
ARM	Atmospheric Radiation Measurements
β	Backscatter cross section ($m^{-1}sr^{-1}$)
BT, BTd	Brightness Temperature, Brightness Temperature Difference
CM	Coarse mode (generally referring to particles of super-micronmicrometer radii)
COD	Cloud Optical Depth
DOD	Dust Optical Depth
FM	Fine mode (generally referring to particles of sub-micrometer radii generally sub-micron)
HLD	High Latitude Dust
HYSPPLIT	HYbrid Single Particle Lagrangian Integrated Trajectory
INP	Ice Nucleation Particle
KAZRGE	Ka-ARM Zenith Radar (KAZR) ³⁰ General mode. Zenith pointing Doppler radar operating at 35 GHz (8.6 mm)
LWP	Liquid Water Path
MISR	Multi-angle Imaging SpectroRadiometer
MODIS	Moderate Resolution Imaging Spectroradiometer
NSHSRL	North Slope High Spectral Resolution Lidar
Terra	Terra passes from north to south across the equator in the morning
TIR	Thermal InfraRed

Commented [NT015]: I don't know how you formatted this text (it should be just a regular table ... I can't figure out how to insert a new entry!). Please make a regular Word table and then insert the definition of the CAA (Canadian Arctic Archipelago)

²⁹ <https://aeronet.gsfc.nasa.gov/>

³⁰ <https://adc.arm.gov/metadata/html/nsakazrgeC1.b1.html>

References

- AboEl-Fetouh, Y., O'Neill, N. T., Ranjbar, K., Hesaraki, S., Abboud, I., & Sobolewski, P. S.: Climatological-Scale Analysis of Intensive and Semi-intensive Aerosol Parameters Derived From AERONET Retrievals Over the Arctic. *Journal of Geophysical Research: Atmospheres*, 125(10). <https://doi.org/10.1029/2019jd031569>, 2020.
- 575 Barr, S. L., Wyld, B., McQuaid, J. B., Neely, R. R., & Murray, B. J.: Southern Alaska as a source of atmospheric mineral dust and ice-nucleating particles. *Science Advances*, 9(33), 1–12. <https://doi.org/10.1126/sciadv.adg3708>, 2023.
- Bowen, M., & Vincent, R. F.: An assessment of the spatial extent of polar dust using satellite thermal data. *Scientific Reports*, 11(1), 1–9. <https://doi.org/10.1038/s41598-020-79825-7>, 2021.
- Bradley, R. S., Keimig, F. T., & Diaz, H. F.: Recent changes in the North American Arctic boundary layer in winter. 580 *Journal of Geophysical Research*, 98(D5), 8851–8858. <https://doi.org/10.1029/93JD00311>, 1993.
- [Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., & Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements—methodology and examples. *Atmospheric Measurement Techniques*, 5\(1\), 73–98. <https://doi.org/10.5194/amt-5-73-2012>, 2012.](https://doi.org/10.5194/amt-5-73-2012)
- 585 ~~[Burton SP, Ferrare RA, Hostetler CA, Hair JW, Rogers RR, Obland MD, Butler CF, Cook AL, Harper DB, Froyd KD: Aerosol classification using airborne High Spectral Resolution Lidar measurements—methodology and examples. *Atmospheric Measurement Techniques*, 2012 Jan 10;5\(1\):73–98](https://doi.org/10.5194/amt-5-73-2012)~~
- de Boer, G., Eloranta, E. W., & Shupe, M. D.: Arctic mixed-phase stratiform cloud properties from multiple years of surface-based measurements at two high-latitude locations. *Journal of the Atmospheric Sciences*, 66(9), 2874–2887. <https://doi.org/10.1175/2009JAS3029.1>, 2009.
- 590 Eck, T. F., Holben, B. N., Reid, J. S., Sinyuk, A., Hyer, E. J., O'Neill, N. T., Shaw, G. E., Vande Castle, J. R., Chapin, F. S., Dubovik, O., Smirnov, A., Vermote, E., Schafer, J. S., Giles, D., Slutsker, I., Sorokine, M., & Newcomb, W. W.: Optical properties of boreal region biomass burning aerosols in central Alaska and seasonal variation of aerosol optical depth at an Arctic coastal site. *Journal of Geophysical Research Atmospheres*, 114(11), 1–14. <https://doi.org/10.1029/2008JD010870>, 2009.
- 595 Feldman, D. R., W. D. Collins, R. Pincus, X. Huang, and X. Chen (2014), Far-infrared surface emissivity and climate, *Proc. Natl. Acad. Sci. U.S.A.*, 111(46), 16,297–16,302, <https://www.pnas.org/doi/10.1073/pnas.1413640111>
- Ginoux, P.: Effects of nonsphericity on mineral dust modeling. *Journal of Geophysical Research: Atmospheres*, 108(2). <https://doi.org/10.1029/2002jd002516>, 2003.
- Gkikas, A., Proestakis, E., Amiridis, V., Kazadzis, S., Di Tomaso, E., Tsekeri, A., Marinou, E., Hatzianastassiou, N., & 600 Pérez García-Pando, C.: ModIs Dust AeroSol (MIDAS): A global fine-resolution dust optical depth data set. *Atmospheric Measurement Techniques*, 14(1), 309–334. <https://doi.org/10.5194/amt-14-309-2021>, 2021.

- Gong, S. L., Zhang, X. Y., Zhao, T. L., McKendry, I. G., Jaffe, D. A., & Lu, N. M.: Characterization of soil dust aerosol in China and its transport and distribution during 2001 ACE-Asia: 2. Model simulation and validation. *Journal of Geophysical Research: Atmospheres*, 108(9), 1–19. <https://doi.org/10.1029/2002jd002633>, 2003.
- 605 Groot Zwaafink, C. D., Grythe, H., Skov, H., & Stohl, A.: Substantial contribution of northern high-latitude sources to mineral dust in the Arctic. *Journal of Geophysical Research*, 121(22), 13,678–13,697. <https://doi.org/10.1002/2016JD025482>, 2016.
- Gu, Y., Rose, W. I., & Bluth, G. J. S.: Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of April 7, 2001 sandstorm in China. *Geophysical Research Letters*, 30(15), 1998–2001. <https://doi.org/10.1029/2003GL017405>, 2003.
- 610 [Hale, G. M., & Querry, M. R.: Optical Constants of Water in the 200-nm to 200- \$\mu\$ m Wavelength Region. *Applied Optics*, 12\(3\), 555. <https://doi.org/10.1364/ao.12.000555>, 1973.](https://doi.org/10.1364/ao.12.000555)
- [Hale, G. M. and Querry, M. R. \(1973\) Optical constants of water in the 200-nm to 200- \$\mu\$ m wavelength region, *Appl. Opt.* 12, 555–563.](https://doi.org/10.1364/ao.12.000555)
- 615 Han, H. J., Sohn, B. J., Huang, H. L., Weisz, E., Saunders, R., & Takamura, T.: An improved radiance simulation for hyperspectral infrared remote sensing of Asian dust. *Journal of Geophysical Research Atmospheres*, 117(9), 1–10. <https://doi.org/10.1029/2012JD017466>, 2012.
- Hansen, J.E., Travis, L.D.: Light scattering in planetary atmospheres, *Space Sci Rev* 16, 527–610, <https://doi.org/10.1007/BF00168069>, 1974.
- 620 Hess, M., Koepke, P., & Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC. *Bulletin of the American Meteorological Society*, 79(5), 831–844. [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), 1998.
- Hildner, R. D.: Understanding Aerosol-Cloud Interactions in Ice Saturated Environments using AHSRL, CALIOP and Trajectory Cluster Analysis, 2nd Symposium on Aerosol-Cloud-Climate Interactions, Jan 20, 2010.
- 625 Hinds, W. C.: *Aerosol technology : properties, behavior, and measurement of airborne particles*. John Wiley & Sons; 1999, 483 pp., 1999.
- [Holz, R. E., Platnick, S., Meyer, K., Vaughan, M., Heidinger, A., Yang, P., Wind, G., Dutcher, S., Ackerman, S., Amarasinghe, N., Nagle, F., & Wang, C.: Resolving ice cloud optical thickness biases between CALIOP and MODIS using infrared retrievals. *Atmospheric Chemistry and Physics*, 16\(8\), 5075–5090. <https://doi.org/10.5194/acp-16-5075-2016>, 2016.](https://doi.org/10.5194/acp-16-5075-2016)
- 630 [Holz, R. E., et al. \(2016\), Resolving ice cloud optical thickness biases between CALIOP and MODIS using infrared retrievals, *Atmos. Chem. Phys.*, 16\(8\), 5075–5090, doi:10.5194/acp-16-5075-2016.](https://doi.org/10.5194/acp-16-5075-2016)
- Hsu, N. C., Gautam, R., Sayer, A. M., Bettenhausen, C., Li, C., Jeong, M. J., Tsay, S. C., & Holben, B. N.: Global and regional trends of aerosol optical depth over land and ocean using SeaWiFS measurements from 1997 to 2010. *Atmospheric Chemistry and Physics*, 12(17), 8037–8053. <https://doi.org/10.5194/acp-12-8037-2012>, 2012.

- 635 Jeong, G. Y.: Bulk and single-particle mineralogy of Asian dust and a comparison with its source soils, *J. Geophys. Res.*, 113, D02208. <https://doi.org/10.1029/2007JD008606>, 2008.
- Kawai, K., Matsui, H., & Tobo, Y.: Dominant Role of Arctic Dust With High Ice Nucleating Ability in the Arctic Lower Troposphere. *Geophysical Research Letters*, 50(8), 1–10. <https://doi.org/10.1029/2022GL102470>, 2023.
- Key, J.: The Cloud and Surface Parameter Retrieval (CASPR) System for Polar AVHRR: Users Guide. *Cooperative Institute for Meteorological Satellite Studies, University of Wisconsin*, 1225, 33–69, 2002.
- 640 Koepke, P., Hess, M., Schult, I., Shettle, E.P.: Global Aerosol Data Set. Report No. 243, ISSN: 0937-1060. Max-Planck-Institut für Meteorologie, Hamburg, 1997.
- Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle. *Proceedings of the National Academy of Sciences of the United States of America*, 108(3), 1016–
- 645 1021. <https://doi.org/10.1073/pnas.1014798108>, 2011.
- Li, J., Wong, M. S., Wai, K. M., & Lee, K. H.: Characterization of Asian Dust Storms by MTSAT Satellite Retrievals. *5th International Workshop on Earth Observation and Remote Sensing Applications, EORSA 2018 - Proceedings*, 1–5. <https://doi.org/10.1109/EORSA.2018.8598641>, 2018.
- Liu, Y., Key, J. R., Frey, R. A., Ackerman, S. A., & Menzel, W. P.: Nighttime polar cloud detection with MODIS. *Remote Sensing of Environment*, 92(2), 181–194. <https://doi.org/10.1016/j.rse.2004.06.004>, 2004.
- 650 Meinander, O., Dagsson-Waldhauserova, P., Amosov, P., Aseyeva, E., Atkins, C., Baklanov, A., Baldo, C., Barr, S. L., Barzycka, B., Benning, L. G., Cvetkovic, B., Enchilik, P., Frolov, D., Gassó, S., Kandler, K., Kasimov, N., Kavan, J., King, J., Koroleva, T., Krupskaya, V., Kulmala, M., Kusiak, M., Lappalainen, H. K., Laska, M., Lasne, J., Lewandowski, M., Luks, B., McQuaid, J. B., Moroni, B., Murray, B., Möhler, O., Nawrot, A., Nickovic, S., O'Neill, N. T., Pejanovic, G., Popovicheva,
- 655 O., Ranjbar, K., Romanias, M., Samonova, O., Sanchez-Marroquin, A., Schepanski, K., Semenkov, I., Sharapova, A., Shevnina, E., Shi, Z., Sofiev, M., Thevenet, F., Thorsteinsson, T., Timofeev, M., Silas Umo, N., Uppstu, A., Urupina, D., Varga, G., Werner, T., Arnalds, O., Vukovic Vimic, A.: Newly identified climatically and environmentally significant high-latitude dust sources. *Atmospheric Chemistry and Physics*, 22(17), 11889–11930, <https://doi.org/10.5194/acp-22-11889-2022>, 2022.
- 660 [Masuda, K., Takashima, T., & Takayama, Y.: Emissivity of pure and sea waters for the model sea surface in the infrared window regions. *Remote Sensing of Environment*, 24\(2\), 313–329. https://doi.org/10.1016/0034-4257\(88\)90032-6, 1998.](https://doi.org/10.1016/0034-4257(88)90032-6)
- [Masuda, K., Takashima, T., & Takayama, Y. \(1988\). Emissivity of pure and sea waters for the model sea surface in the infrared window regions. *Remote Sensing of Environment*, 24\(2\), 313–329. doi:10.1016/0034-4257\(88\)90032-6](https://doi.org/10.1016/0034-4257(88)90032-6)
- Morrison, H., De Boer, G., Feingold, G., Harrington, J., Shupe, M. D., & Sulia, K.: Resilience of persistent Arctic mixed-phase clouds. *Nature Geoscience*, 5(1), 11–17. <https://doi.org/10.1038/ngeo1332>, 2012.
- 665 Okada, K., Heintzenberg, J., Kai, K., & Qin, Y.: Shape of atmospheric mineral particles collected in three Chinese arid regions. *Geophysical Research Letters*, 28(16), 3123–3126. <https://doi.org/10.1029/2000GL012798>, 2001.

- O'Neill, N. T., Baibakov, K., Hesaraki, S., Ivanescu, L., Martin, R. V., Perro, C., Chaubey, J. P., Herber, A., & Duck, T. J.: Temporal and spectral cloud screening of polar winter aerosol optical depth (AOD): Impact of homogeneous and inhomogeneous clouds and crystal layers on climatological-scale AODs. *Atmospheric Chemistry and Physics*, 16(19), 12753–12765. <https://doi.org/10.5194/acp-16-12753-2016>, 2016.
- O'Neill, N. T., Ranjbar, K., Ivanescu, L., Eck, T. F., Reid, J. S., Giles, D. M., Pérez-Ramírez, D., & Chaubey, J. P.: Relationship between the sub-micron fraction (SMF) and fine-mode fraction (FMF) in the context of AERONET retrievals. *Atmospheric Measurement Techniques*, 16(4), 1103–1120. <https://doi.org/10.5194/amt-16-1103-2023>, 2023.
- O'Neill, N. T., Thulasiraman, S., Eck, T. F., & Reid, J. S.: Robust optical features of fine mode size distributions: Application to the Québec smoke event of 2002. *Journal of Geophysical Research: Atmospheres*, 110(11), 1–16. <https://doi.org/10.1029/2004JD005157>, 2005.
- Painter, T. H., Barrett, A. P., Landry, C. C., Neff, J. C., Cassidy, M. P., Lawrence, C. R., McBride, K. E., & Farmer, G. L.: Impact of disturbed desert soils on duration of mountain snow cover. *Geophysical Research Letters*, 34(12), 1–6. <https://doi.org/10.1029/2007GL030284>, 2007.
- Palo, T., Vihma, T., Jaagus, J., & Jakobson, E.: Observations of temperature inversions over central Arctic sea ice in summer. *Quarterly Journal of the Royal Meteorological Society*, 143(708), 2741–2754. <https://doi.org/10.1002/qj.3123>, 2017.
- Petty, G. W., & Huang, W.: The modified gamma size distribution applied to inhomogeneous and nonspherical particles: Key relationships and conversions. *Journal of the Atmospheric Sciences*, 68(7), 1460–1473. <https://doi.org/10.1175/2011JAS3645.1>, 2011.
- Petty, G. and Huang, W. (2011). The Modified Gamma Size Distribution Applied to Inhomogeneous and Nonspherical Particles: Key Relationships and Conversions. *Journal of the Atmospheric Sciences* 68(7) pp. 1460–1473.
- Ranjbar, K., O'Neill, N. T., & Aboel-Fetouh, Y.: Comment on “Short-cut transport path for Asian dust directly to the Arctic: A case Study” by Huang et al. (2015) in *Environ. Res. Lett. Atmospheric Chemistry and Physics*, 22(3), 1757–1760. <https://doi.org/10.5194/acp-22-1757-2022>, 2022.
- Ranjbar, K., O'Neill, N. T., Ivanescu, L., King, J., & Hayes, P. L.: Remote sensing of a high-Arctic, local dust event over Lake Hazen (Ellesmere Island, Nunavut, Canada). *Atmospheric Environment*, 246, 118102. <https://doi.org/10.1016/j.atmosenv.2020.118102>, 2021.
- Rothman, L. S., Gordon, I. E., Barbe, A., Benner, D. C., Bernath, P. F., Birk, M., Boudon, V., Brown, L. R., Campargue, A., Champion, J. P., Chance, K., Coudert, L. H., Dana, V., Devi, V. M., Fally, S., Flaud, J. M., Gamache, R. R., Goldman, A., Jacquemart, D., ... Vander Auwera, J.: The HITRAN 2008 molecular spectroscopic database. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 110(9–10), 533–572. <https://doi.org/10.1016/j.jqsrt.2009.02.013>, 2009.
- Sadrian, M. R., Calvin, W. M., Perrin, A. E., Engelbrecht, J. P., & Moosmüller, H.: Variations in Infrared Complex Refractive Index Spectra of Surface Soils from Global Dust Entrainment Regions. *Atmosphere*, 14(4), 1–14. <https://doi.org/10.3390/atmos14040675>, 2023.

- Shi, Y., Liu, X., Wu, M., Zhao, X., Ke, Z., & Brown, H.: Relative importance of high-latitude local and long-range-transported dust for Arctic ice-nucleating particles and impacts on Arctic mixed-phase clouds. *Atmospheric Chemistry and Physics*, 22(4), 2909–2935. <https://doi.org/10.5194/acp-22-2909-2022>, 2022.
- Shonk, J. K. P., Chiu, J. Y. C., Marshak, A., Giles, D. M., Huang, C. H., MacE, G. G., Benson, S., Slutsker, I., & Holben,
705 B. N.: The impact of neglecting ice phase on cloud optical depth retrievals from AERONET cloud mode observations. *Atmospheric Measurement Techniques*, 12(9), 5087–5099. <https://doi.org/10.5194/amt-12-5087-2019>, 2019.
- Shupe, M. D., Turner, D. D., Zwink, A., Thieman, M. M., Mlawer, E. J., & Shippert, T.: Deriving arctic cloud microphysics at Barrow, Alaska: Algorithms, results, and radiative closure. *Journal of Applied Meteorology and Climatology*, 54(7), 1675–1689. <https://doi.org/10.1175/JAMC-D-15-0054.1>, 2015.
- 710 Stone, R. S., Anderson, G. P., Andrews, E., Dutton, E. G., Shettle, E. P., & Berk, A.: Incursions and radiative impact of Asian dust in northern Alaska. *Geophysical Research Letters*, 34(14), 1–5. <https://doi.org/10.1029/2007GL029878>, 2007.
- Stone, R. S., Sharma, S., Herber, A., Eleftheriadis, K., & Nelson, D. W.: A characterization of Arctic aerosols on the basis of aerosol optical depth and black carbon measurements. *Elementa*, 2, 1–22. <https://doi.org/10.12952/journal.elementa.000027>, 2014.
- 715 Tobo, Y., Adachi, K., DeMott, P. J., Hill, T. C. J., Hamilton, D. S., Mahowald, N. M., Nagatsuka, N., Ohata, S., Uetake, J., Kondo, Y., & Koike, M.: Glacially sourced dust as a potentially significant source of ice nucleating particles. *Nature Geoscience*, 12(4), 253–258. <https://doi.org/10.1038/s41561-019-0314-x>, 2019.
- Vincent, R. F.: The Effect of Arctic Dust on the Retrieval of Satellite Derived Sea and Ice Surface Temperatures. *Scientific Reports*, 8(1), 6–15. <https://doi.org/10.1038/s41598-018-28024-6>, 2018.
- 720 Volz, F. E.: Infrared Optical Constants of Ammonium Sulfate, Sahara Dust, Volcanic Pumice, and Flyash. *Applied Optics*, 12(3), 564. <https://doi.org/10.1364/ao.12.000564>, 1973.
- Volz, F. E.: Infrared Refractive Index of Atmospheric Aerosol Substances. *Applied Optics*, 11(4), 755. <https://doi.org/10.1364/ao.11.000755>, 1972.
- Wald, A. E., Kaufman, Y. J., Tanré, D., & Gao, B. C.: Daytime and nighttime detection of mineral dust over desert using
725 infrared spectral contrast. *Journal of Geophysical Research Atmospheres*, 103(D24), 32307–32313. <https://doi.org/10.1029/98JD01454>, 1998.
- [Woo, M. K., Edlund, S. A., Young, K. L.: Occurrence of early snow-free zones on Fosheim peninsula, Ellesmere Island, northwest territories. Curr. Res. Part B, Geol. Surv. Canada Pap. 91, 9–14. https://publications.gc.ca/collections/collection_2017/rncan-nrcan/M44-91-1B.pdf, 1991.](https://publications.gc.ca/collections/collection_2017/rncan-nrcan/M44-91-1B.pdf)
- 730 [Woo, M.K., Edlund, S.A., Young, K.L., 1991. Occurrence of early snow-free zones on Fosheim peninsula, Ellesmere Island, northwest territories. Curr. Res. Part B, Geol. Surv. Canada Pap. 91, 9–14.](https://publications.gc.ca/collections/collection_2017/rncan-nrcan/M44-91-1B.pdf)
- Xi, Y., Xu, C., Downey, A., Stevens, R., Bachelder, J. O., King, J., Hayes, P. L., & Bertram, A. K.: Ice nucleating properties of airborne dust from an actively retreating glacier in Yukon, Canada. *Environmental Science: Atmospheres*, 2(4), 714–726. <https://doi.org/10.1039/d1ea00101a>, 2022.

- 735 Xian, P., Zhang, J., O'Neill, N.T., Toth, T.D., Sorenson, B., Colarco, P.R., Kipling, Z., Hyer, E.J., Campbell, J.R., Reid, J.S. and Ranjbar, K., 2022. Arctic spring and summertime aerosol optical depth baseline from long-term observations and model reanalyses–Part 1: Climatology and trend. *Atmospheric Chemistry and Physics*, 22(15), pp.9915–9947, <https://doi.org/10.5194/acp-22-9915-2022>.
- 740 [Yang, P., Bi, L., Baum, B. A., Liou, K. N., Kattawar, G. W., Mishchenko, M. I., & Cole, B.: Spectrally consistent scattering, absorption, and polarization properties of atmospheric ice crystals at wavelengths from 0.2 to 100 \$\mu\text{m}\$. *Journal of the Atmospheric Sciences*, 70\(1\), 330–347. <https://doi.org/10.1175/JAS-D-12-039.1>, 2013.](https://doi.org/10.1175/JAS-D-12-039.1)
- ~~Yang, P., Bi, L., Baum, B., Liou, K., Kattawar, G., Mishchenko, M. and Cole, B. (2013). Spectrally Consistent Scattering, Absorption, and Polarization Properties of Atmospheric Ice Crystals at Wavelengths from 0.2 to 100 μm . *Journal of the Atmospheric Sciences* 70(1) pp. 330–347. Available at: <https://journals.ametsoc.org/view/journals/atsc/70/1/jas-d-12-039.1.xml>~~
- 745 Yi, L., Li, K. F., Chen, X., & Tung, K. K.: Arctic fog detection using infrared spectral measurements. *Journal of Atmospheric and Oceanic Technology*, 36(8), 1643–1656. <https://doi.org/10.1175/JTECH-D-18-0100.1>, 2019.
- Zamora, L. M., Kahn, R. A., Evangelio, N., Groot Zwaafink, C. D., & Huebert, K. B.: Comparisons between the distributions of dust and combustion aerosols in MERRA-2, FLEXPART, and CALIPSO and implications for deposition freezing over wintertime Siberia. *Atmospheric Chemistry and Physics*, 22(18), 12269–12285. <https://doi.org/10.5194/acp-22-12269-2022>, 2022.
- Zdanowicz, C., Hall, G., Vaive, J., Amelin, Y., Percival, J., Girard, I., Biscaye, P., & Bory, A.: Asian dustfall in the St. Elias Mountains, Yukon, Canada. *Geochimica et Cosmochimica Acta*, 70(14), 3493–3507. <https://doi.org/10.1016/j.gca.2006.05.005>, 2006.
- 755 Zhang, P., Lu, N. meng, Hu, X. qing, & Dong, C. hua.: Identification and physical retrieval of dust storm using three MODIS thermal IR channels. *Global and Planetary Change*, 52(1–4), 197–206. <https://doi.org/10.1016/j.gloplacha.2006.02.014>, 2006.
- Zhao, X., Huang, K., Fu, J. S., & Abdullaev, S. F.: Long-range transport of Asian dust to the Arctic: identification of transport pathways, evolution of aerosol optical properties, and impact assessment on surface albedo changes. *Atmospheric Chemistry and Physics*, 22(15), 10389–10407. <https://doi.org/10.5194/acp-22-10389-2022>, 2022.
- 760 Zuidema, P., Baker, B., Han, Y., Intrieri, J., Key, J., Lawson, P., Matrosov, S., Shupe, M., Stone, R., & Uttal, T.: An Arctic springtime mixed-phase cloudy boundary layer observed during SHEBA. *Journal of the Atmospheric Sciences*, 62(1), 160–176. <https://doi.org/10.1175/JAS-3368.1>, 2005.