

How does the lifetime of detrained cirrus impact the high cloud radiative effect in the tropics?

George Horner¹ and Edward Gryspeerdt^{1,2}

¹Department of Physics, Imperial College London, London, UK

²Grantham Institute - Climate Change and the Environment, Imperial College London, London, UK

Correspondence: George Horner (g.horner20@imperial.ac.uk)

Abstract. The lifetime of cirrus clouds from deep convection plays an important role in determining their overall cloud radiative effect (CRE). The net CRE of cirrus clouds from deep convection is close to zero over their whole lifetime. This CRE is the result of a near-cancellation of a large shortwave (SW) cooling and large longwave (LW) warming, such that small changes in cirrus properties have the potential to produce a ~~large~~ significant net radiative effect. Changes in the atmospheric and sea surface temperature structure, along with changes in anthropogenic aerosol, have been hypothesised to impact the lifetime of detrained cirrus clouds, altering this radiative balance. Constraining the potential CRE response to changes in cirrus lifetime is therefore vital to understand the strength of these proposed climate forcings and feedbacks.

This paper tracks the evolution of detrained cirrus clouds along trajectories from deep convection. The total cirrus CRE in the tropics is found to be ~~3.6~~ warming, at 5.0 $\pm 0.4 \text{ Wm}^{-2}$. It's found that cirrus clouds along trajectories from oceanic origin convection have a ~~CRE of 2.9~~ warming CRE of 4.4 $\pm 0.4 \text{ Wm}^{-2}$. In contrast, cirrus clouds along trajectories from land convection have a warming of ~~6.3~~ 7.7 $\pm 0.6 \text{ Wm}^{-2}$ throughout their lifetime. This contrast is predominantly due to differences in the diurnal cycle of the initial convection over land and ocean.

A proposed extension to the lifetime of the detrained cirrus leads to changes in the total cirrus CRE in the tropics. In all cases, doubling the lifetime of the detrained cirrus leads to an increase in the total cirrus CRE of ~~1.2~~ 0.7 $\pm 0.1 \text{ Wm}^{-2}$. ~~This result~~ Whilst there is uncertainty in the strength of mechanisms responsible for a change in cirrus lifetime, this work provides an important constraint on the impact ~~of changes in the lifetime of detrained cirrus in a future climate or in response to aerosol perturbations on the total tropical CRE~~ that any potential lifetime extension may have.

1 Introduction

Deep convective clouds and their associated thin cirrus outflows occur predominantly in the tropical and subtropical regions. Deep convection is characterised by optically thick, high-altitude, and relatively short-lived convective cores. When these cores reach their maximum altitude, large anvil cirrus spread out from them around the areas of convection (Fan et al., 2013). The radiative ~~forcing~~ effect of these convective cores and anvil cirrus clouds are strongly dependent on their optical thickness and ice water path (Berry and Mace, 2014; Hartmann and Berry, 2017), which change over time (Horner and Gryspeerdt, 2023). An important feature of the tropics is the cancellation between large longwave (LW) and shortwave (SW) components of the

25 total cloud radiative effect (CRE) (Wielicki et al., 1996). This cancellation occurs due to the significant LW warming from high altitude, thin cirrus clouds (typically defined as optical thickness below 0.05 (Lee et al., 2009a)), and the high albedo of optically thick clouds occurring in the same region. This leads to a CRE in cloudy regions that is very similar to non-cloudy regions in the tropics. (Ramanathan et al., 1989; Hartmann and Berry, 2017; Harrison et al., 1990). It is unclear whether this cancellation in tropical CRE is a coincidence or due to feedbacks (Kiehl, 1994; Hartmann et al., 2001). Since the LW and SW CREs are so large, any small percentage change in the magnitudes of either of these components in a future climate could lead to significant changes in the overall CRE in the tropics. It is therefore vital to understand how changes in properties and lifetime of cirrus clouds from convection affect the overall tropical CRE.

Cirrus clouds cover approximately 60-80% of the tropics (Wylie and Menzel, 1999; Stubenrauch et al., 2006; Sassen et al., 2008; Nazaryan et al., 2008; Lee et al., 2009b). Luo and Rossow (2004) found using observational data that 50% of tropical cirrus were detrained cirrus, i.e. they originated out of deep convective cores. Using ground-based data and satellites, Mace et al. (2006) found that 47% of cirrus observed over Manus Island in the western Pacific originated from deep convection occurring within the past 12 h. Many other modelling and observational studies also show that at least 50% of tropical cirrus originate from convection (Massie et al., 2002; Gehlot and Quaas, 2012; Riihimaki et al., 2012). Given that detrained cirrus therefore cover approximately 30-40% of the tropics, their radiative effects are an important contributor to the overall CRE balance in the tropics. Luo and Rossow (2004) found that detrained cirrus had longer average lifetimes than in situ cirrus, at around 30 h compared to 19 h for in situ, ~~suggesting some mechanism for sustaining water vapour at the detrained cirrus height that allows for longer lifetimes~~ due to the fact that the detrained cirrus are thicker than in situ cirrus, and thus take more time to dissipate.

In situ cirrus, whilst not formed directly from convective cores, do not exist independently from their detrained counterparts, given that the formation of in situ cirrus relies on water vapour in the upper troposphere, transported via convection. The lifetime and properties of detrained cirrus are affected by any factor that affects the distribution of deep convection (Gasparini et al., 2023). These ~~large-scale~~ large-scale factors range from variable sea surface temperatures (SSTs) to the Madden-Julian Oscillation (MJO) and El-Niño Southern Oscillation (ENSO) (Riley et al., 2011; Sweeney et al., 2023). Not only will these factors affect the distribution and properties of the detrained cirrus, but the abundance of deep convection which lifts moisture into the free troposphere and generates detrained cirrus is likely to also affect the distribution of in situ cirrus (Massie et al., 2002). Conversely, the existence of in situ cirrus will also affect the impact that convective air and moisture has on the free troposphere and lower stratosphere, and therefore the development of detrained cirrus (Jensen et al., 2020). For these reasons, the total cirrus CRE in the tropics must be considered when investigating lifetime changes to detrained cirrus. Any change in the properties or amount of detrained cirrus will ultimately affect the average properties of all cirrus in the tropics.

55 Detrained cirrus and in situ cirrus have different radiative and microphysical properties. Of particular importance is the time dependence of the properties of detrained cirrus compared to in situ cirrus. The properties of detrained cirrus strongly depend on the time since the cirrus cloud formed out of deep convection (Horner and Gryspeerd, 2023). Close to convection, the detrained anvil cirrus will be optically thick, with a SW cooling dominating the net CRE. As time since convection increases, the detrained cirrus have a tendency to thin and spread out, decreasing the SW cooling as the albedo decreases (Luo and

60 Rossow, 2004; Hartmann et al., 2018; Gasparini et al., 2019, 2021). The albedo decreases faster than the LW emissivity, contributing to this switch from cooling to warming in the net CRE Berry and Mace (2014). The properties of in situ cirrus will vary, depending on the optical thickness and ice water path, as well as the altitude of the clouds (Hartmann and Berry, 2017). In contrast to in situ cirrus, where there is little time dependence on their radiative properties, the time dependence of detrained cirrus is an important factor in determining their CRE. Therefore when calculating the radiative balance in the
65 tropics, any changes in the lifetime of detrained cirrus would have a significant impact on the delicate radiative balance in the tropics, given that their properties are such a strong function of time since convection.

In addition to the overall lifetime of the detrained cirrus, the timing of the initial convection plays an important role in determining the total cirrus cloud CRE over the detrained cirrus' lifetime. Jones et al. (2023) showed that the timing of the initial convection has a large effect on the total CRE across the detrained cirrus' lifetime. This is due to the dependence of SW
70 cooling on incoming solar radiation. If convection occurs later in the day when solar insolation is lower, the SW cooling of the younger, thicker anvil cirrus will be less than if the convection occurs in the morning or middle of the day.

The diurnal dependence on the total lifetime CRE of detrained cirrus means that the location where the convection occurs has a large effect on the CRE across its lifetime. Many studies have investigated the diurnal cycle of regional convection (Chen and Houze Jr, 1997; Lopez-Bravo et al., 2023; Krishna et al., 2021). Fewer studies have investigated large-scale differences in
75 the diurnal cycle of deep convection (Ruppert and Hohenegger, 2018). The difference in the diurnal cycle from terrestrial and oceanic convection ~~, where the differences in the diurnal cycle of convection are significant, is~~ has been thoroughly studied (Dai et al., 1999; Dai, 2001; Albright et al., 1985; Zuidema, 2003). Of particular importance in this paper is not the mechanisms that govern the behaviour of terrestrial and oceanic convection but the effect that the difference in timing of this convection has on the subsequent radiative properties of the detrained cirrus.

80 The temperature of anvil clouds will remain the same in a warming climate for a given increase in altitude under the fixed anvil temperature (FAT) hypothesis (Hartmann and Larson, 2002). However, many modelling studies have predicted changes in the cloud fraction of detrained anvil clouds in a warming climate (Bony et al., 2016; Cronin and Wing, 2017; Su et al., 2017). ~~There are disagreements with this theory (Bony et al., 2015), with some criticism of the original methodology (Del Genio and Kovari, 2002; Hartmann and Michelsen, 2002). Nevertheless, studies find evidence for a decreasing cloud area with warming SSTs (Zelinka and Hartmann, 2011; Choi et al., 2017).~~ Changes in cloud area can be thought of as changes in the lifetime from a Lagrangian viewpoint, assuming that the advection of the anvil cirrus remains the same (which under the stability iris hypothesis may not be true), where 'time since convection' is a function of distance from a convective core (Horner and Gryspeerdt, 2023). Raghuraman et al. (2024) used observational evidence to investigate these feedbacks, finding a near zero high-cloud amount feedback, providing evidence against an iris feedback.

90 Although there is some observational evidence for a decrease in cloud area/lifetime with warming SSTs, there is also a large spread in model responses to warming SSTs (Wing et al., 2020), with some models showing an increase in high cloud amount with warming (Ohno and Satoh, 2018; Satoh et al., 2012; Tsushima et al., 2014). Gasparini et al. (2021) used Lagrangian trajectories and model studies to investigate the lifetime of anvil cirrus in present and future climate. They found the lifetime of the anvil cloud to be 15 h, significantly shorter to that of Luo and Rossow (2004) who found anvil lifetimes of 30 h. These studies

95 are sensitive to the choice of methods and thresholds for cirrus detection used. For example, Luo and Rossow (2004) determine cirrus lifetime to be the point at which the cloud fraction along a trajectory first drops below 20% of the maximum cloud fraction along the trajectory (nominally 100%). There is no robust definition of when detrained cirrus have dissipated, therefore there is some spread in the given lifetimes of detrained cirrus in the literature.

Crucially, Gasparini et al. (2021) found no difference in the lifetime of anvil cirrus in the present climate compared to a 4K warming scenario. ~~Clearly,~~ Beydoun et al. (2021) also found a similar negligible change in anvil lifetime using RCEMIP-style models. It should be noted that neither of these studies represent all the processes that would impact cirrus lifetime, particularly the lack of diurnal cycle. More observational studies are needed to assess the impact that these particular processes may have on the lifetime of detrained cirrus clouds. Overall, there is much uncertainty in the response of high cloud area and lifetime to a warming climate.

105 Drivers of lifetime changes in detrained cirrus relate to the properties of the deep convection itself. Studies have investigated the aerosol invigoration hypothesis (Abbott and Cronin, 2021; Fan et al., 2013; Koren et al., 2010, 2005; Rosenfeld et al., 2008), which states that an increase in aerosol concentration should lead to increased updrafts in deep convection, leading to higher altitude, and potentially, longer lived anvil cirrus. ~~A full~~ Herbert and Stier (2023) provide an observational study investigating the impact of AOD on deep convection over the Amazon, however this is limited to short timescales, close to the convection.

110 A much earlier study by Lin et al. (2006) also shows, using observations, a correlation between AOD and convective strength (via higher cloud tops). There is still more work needed to understand the full impact that aerosol invigoration may have on the lifecycle of detrained cirrus as they evolve beyond the deep convection itself. A full critical evaluation of aerosol invigoration is given by Varble et al. (2023).

There are many factors that impact the decay, and therefore lifetime, of detrained cirrus clouds. (Gasparini et al., 2022) used

115 idealised cloud resolving models to investigate the impact that the diurnal cycle has on the cirrus lifetime. They found that tropical anvil clouds forming in the day are more widespread and longer lasting than those formed out night, due to the shortwave radiative heating during the day which lofts and spreads the anvil clouds. During night there is an increase in ice nucleation, however this doesn't compensate for the entrainment of dry air that also acts to reduce anvil lifetime. (Sokol and Hartmann, 2020) also found that anvils thin more rapidly during the night due the net radiative cooling eroding

120 the cloud top. In both cases, it is found that the timing of convection is an important control on the lifetime of the detrained cirrus cloud.

This paper seeks to understand the extent to which changing the lifetime of detrained cirrus impacts the overall tropical CRE. If the effect is small, then many of the drivers of anvil development mentioned above may not be important when considering changes in total cirrus CRE in a warming tropics. This question is answered by first characterising the overall high cloud CRE

125 in the tropics as a combination of the CRE of the detrained cirrus and in situ cirrus distributions. The distribution of detrained cirrus is ~~is~~ then modified by artificially altering its lifetime, and ~~investigate what impact the impact that~~ this has on the overall CRE is investigated. How the CRE of detrained cirrus from land convection differs to oceanic convection is also investigated, and how their lifetime extensions change the overall tropical CRE. Answering these questions will allow us to investigate how

sensitive the overall CRE in the tropics is to changes in the lifetime of the detrained cirrus clouds, enabling us to constrain
130 future changes due to climate change or anthropogenic aerosol emissions.

2 Methods

2.1 Characterising cirrus evolution

The Time Since Convection (TSC) algorithm (Horner and Gryspeerdt, 2023) assigns each grid box in the tropics a time in
hours since that grid box last experienced a convective event, as defined by the presence of a deep convective core. The
135 deep convective cores are identified from the 3-hourly International Satellite Cloud Climatology Project (ISCCP)-H dataset
(Rossow et al., 2017) based on their cloud fraction, cloud optical thickness, and cloud top pressure data. This characterises
each gridbox as belonging to a distinct cloud regime, including a deep convective regime. The deep convective gridboxes
are then advected forward in time following the ERA5 reanalysis wind field data ([Hersbach et al., 2018](#)) averaged between
200hPa and 300hPa([Hersbach et al., 2018](#)), with the associated TSC increasing as the air parcels are advected. [As shown in](#)
140 [Horner and Gryspeerdt \(2023\), these trajectories typically remain at this pressure level for the entirety of the detrained cirrus](#)
[lifetime](#). Where new deep convection occurs ([locations updated every three hours](#)), the TSC is reset to zero and advected as
new convection. For a full description of the algorithm see Horner and Gryspeerdt (2023). This method is used to characterise
cirrus properties as a function of TSC to see how these cloud properties change as they move from deep convection through
anvil cirrus to thin cirrus and eventually dissipate. The study period for this paper is between 1984-2016, however any results
145 involving CERES data only go back to 2000.

2.2 Cirrus Type Flag: Land/Ocean and In situ/Detrained

Air parcels along trajectories from deep convection contain detrained cirrus clouds, or they may contain in situ cirrus. Detrained
cirrus, [also known as liquid-origin cirrus \(Luebke et al., 2016\)](#), are defined as cirrus that have detrained directly from a deep
convective core, such as thick anvil cirrus, with the air parcel remaining cloudy along the entire trajectory up to that point. In
150 contrast, in situ cirrus were not directly detrained from a convective core, but appear along the trajectories after the detrained
cirrus has dissipated.

The TSC algorithm assigns a flag to each air parcel depending on properties at the origin of the convection. At TSC=0, all
air parcels are considered detrained, with the transition to an in situ state occurring when the ISCCP-H high cloud fraction
along a trajectory becomes less than 10%. After this point, the air parcel is then considered in situ and remains as such until it
155 is replaced by newer convection. These in situ air parcels may contain a cirrus cloud, which by definition would be an in situ
cirrus cloud, or they may not contain any cirrus cloud at all. While the detrained air parcels by definition must be cloudy (and
so contain a detrained cirrus cloud), the in situ air parcels could be cloud free and hence don't necessarily contain an in situ
cirrus cloud. Instead, it indicates the origin of any potential cirrus cloud in a grid box.




Figure 1. Sample TSC snapshot from April 9th 2008, 00:30UTC. (a) Time since convection (TSC) (b) In situ versus detrained cirrus flag values (c) land versus ocean convective origin flag. (d-f) ~~Annual-2008 annual~~ mean for (a-c).

It is worth noting that the results are not hugely sensitive to the definition of lifetime used (Figure S1). There is no strict definition for how detrained lifetime should be defined, therefore a choice for when to cut off detrained cirrus has to be made, in this work that cut off is 10%.

When two trajectories enter the same gridbox, the air that has more recently experienced convection (with a lower TSC) is deemed to be a more important indicator of cloud properties. As a result, the properties of the younger convection define that gridbox.

When two trajectories diverge and a gridbox is left empty, the value for that gridbox is interpolated as the average of the surrounding eight gridboxes. In practice, the standard deviation of the flag values of the air parcels in the eight surrounding gridboxes is usually small. However, where there is a mixture of air parcel types in the surrounding boxes, this is represented in the flag value associated with the air parcel. The value of a detrained cirrus flag is 100, and an in situ flag is -100, with mixtures taking a value in-between. For example, if six of the eight surrounding gridboxes contain detrained cirrus flags with a value of 100, and two contain in situ flags with a value of -100, the flag value of the divergent gridbox will become 50. I.e., it is more likely to be a detrained cirrus air parcel, but with decreased confidence due to the interpolation in the TSC algorithm. In all subsequent analyses, only air parcels with a high confidence (a magnitude larger than 50) are considered.

This method is also adapted to store properties associated with the initial convection. The convective origin flag marks each air parcel with either a land-origin or an ocean-origin flag. This flag remains on the air parcel until it is replaced by a flag from

175 newer convection. It is important to note that this flag doesn't just indicate whether a cirrus cloud is currently over land or ocean, but rather whether the convection it originated from occurred over land or ocean. For this reason, it is often the case that land-origin convection exists over the ocean and vice versa (Fig. 1c).

Figure 1 shows (a) a snapshot TSC map from 00:30UTC on the 9th April 2008, a snapshot of the associated (b) cirrus type flag and (c) convective origin flag. As values are interpolated during the advection algorithm generating the TSC and associated
 180 flags, the flag values are not a binary classification. Instead, the positive or negative sign indicates the flag value (i.e., positive cirrus type flag indicates detrained, negative in situ), and the magnitude can be considered an indication of the confidence in that flag following the merging of trajectories over time.

2.3 High Cloud Radiative Effect

The high cloud CRE is calculated using the CERES SYN1deg L3 1-hourly dataset (NASA/LARC/SD/ASDC, 2017). It is nec-
 185 essary to isolate the CRE of the cirrus clouds from the effects of low clouds in order to investigate how the radiative properties of the high clouds alone change with TSC. To isolate the radiative effect of the high clouds, the monthly mean combined low cloud and clear sky albedo (hereafter referred to as the 'background albedo') are subtracted from the instantaneous all sky albedo to give an estimated high cloud albedo. To identify the background albedo, the ISCCP histograms are used to isolate gridboxes where the high cloud fraction is less than <1%, thereby attributing the albedo to the low cloud and surface only. The
 190 monthly mean background is used in order to give sufficient coverage to co-locate regions of high cloud and background cloud. Equation 1 is used to isolate the albedo of the high clouds only:

$$\alpha_{high} = \alpha_{all_sky} - \bar{\alpha}_{bkg} \quad (1)$$

where α_{high} is the high cloud albedo, α_{all_sky} is the all sky albedo and $\bar{\alpha}_{bkg}$ is the monthly mean background albedo. The background albedo is defined as the α_{all_sky} in regions with a <1% high cloud fraction. The SW high cloud CRE can then be
 195 calculated using:

$$SW_CRE_{high} = F_{\downarrow} \times \alpha_{high} \quad (2)$$

where F_{\downarrow} is the incoming solar radiation. One shortcoming in using the monthly mean background albedo, and subtracting this from the instantaneous all sky albedo, is that this introduces regions where there is a negative albedo, and therefore an apparent positive instantaneous high cloud SW CRE. This occurs in regions where the all sky is less than the monthly mean
 200 background albedo. Figure 2 shows (a) an instantaneous all sky CRE, (b) the monthly mean background albedo (in this case the month of March), and (c) the instantaneous high cloud CRE calculated from these.




Figure 2. Snapshot of (a) All sky SW CRE for 12:00UTC 02/03/2008, (b) Monthly mean background SW CRE (March), (c) High cloud SW CRE for 12:00UTC 02/03/2008.

As seen in Figure 2c), in regions with little to no high cloud, the SW CRE can be positive, since the all sky instantaneous albedo is less than the annual mean background albedo. These un-physical regions disappear in the annual mean, but are required to avoid biasing the mean high cloud albedo. The LW CRE is calculated as the all-sky LW minus the clear-sky LW, due to the small LW CRE of low clouds.

3 Results

3.1 TSC distribution

Figure 1b) shows the instantaneous detrained versus in situ flag map, and e) shows the yearly mean flag values. The detrained cirrus in red are centered around regions of active convection. These extend beyond the original convective cores, much like
210 anvil cirrus would extend beyond the active convection, All other regions contain in situ air parcels, i.e. if cirrus were to occur in these regions they would be classed as in situ in origin.

Any region in Figure 1e) where the mean value is above zero are regions where detrained cirrus clouds are more often found. ~~Comparing~~ Compared to Fig 1d), these occur at low TSC values, close to convection. The likelihood of finding detrained cirrus air parcels is highest over the Maritime Continent. This is because there is much more sustained convection over the ocean,
215 which means convection here has a smaller diurnal variation. This allows for a constant supply of detrained air from convection over the course of 24 hours. This compares to terrestrial regions over South America and central Africa where the convective cycle is more diurnal. This means there are large periods in the day where there is little convection, and the detrained origin air parcels are advected away and replaced by in situ air parcels from the ocean.

Figure 3a) shows the fraction of grid boxes containing detrained or in situ air parcels, as a function of TSC. Figure 3b
220 shows the total counts of both origin types. Note, the scallop-like pattern along the lines in Figure 3a) is due to the temporal resolution of ISCCP data meaning that convection is only initiated every three hours. At low TSC values, most air parcels along trajectories contain detrained cirrus clouds. At ≈ 30 hours, the number of gridboxes containing detrained cirrus or in situ air parcels is approximately equal. For example, at 30 hours from convection, approximately 50% of the detrained cirrus have dissipated. At higher TSC values than this, new cirrus clouds that occurs along the trajectory are more likely to be in situ in
225 origin.

The median lifetime, that is the TSC at the point a gridbox switches from containing a detrained cirrus air parcel to an in situ air parcel, of a detrained cirrus cloud is 36 hours. ~~This is~~ The modal lifetime is 25 hours. These values are in agreement with Luo and Rossow (2004) who found the average detrained cirrus lifetime to be 30 ± 16 hours.

The standard deviation of the lifetime is found to be 30 hours. This is much larger than that found by Luo and Rossow (2004) since
230 it doesn't track specific convective events but rather tracks all air from convection.

Figure 1d) shows the instantaneous snapshot of the land and ocean origin air parcels. The majority of the tropics contain ocean origin air parcels (blue), as the majority of convection that occurs in the tropics occurs over the ocean. Therefore for a given detrained cirrus cloud, it is more likely to be oceanic than terrestrial in origin.

The land origin air parcels occur predominantly over land or near the coast. Figure 1f) shows the mean land-ocean origin
235 value. There are large regions over the southern Atlantic ocean that contain more frequent land origin air parcels than average. This is due to wind fields that carry air from land origin convection over the ocean, making it more likely to find land-origin detrained cirrus in this region compared to the rest of the global oceans.

The average air parcel origin map (Figure 1f) shows which type of air parcel is most common for a particular region. For example, land air parcels are most likely seen over oceans in the SE Atlantic off the coast of Africa. Figure 1c) shows that these instances of land origin air parcels can occur in many regions over ocean, and can travel long distances from any land source. Figure 4 shows the distribution of the lifetime of the land and ocean-origin detrained cirrus. The modal lifetime for land-origin detrained cirrus is 15 hours, compared to 25 hours for ocean-origin detrained cirrus. It should be noted that many of the very low lifetimes occur when convection replaces itself between timesteps. However, these instances have a negligible impact on the median lifetime. Removing these very low lifetimes (lifetimes less than 5 hours), increases the median lifetime by less than 1 hour.




Figure 3. Distribution of detrained and in situ cirrus flags as a function of TSC. (a) The count of each flag as a fraction of the total TSC distribution. (b) The total distribution of each flag.

The difference in land and ocean origin detrained cirrus lifetime is significant. The land origin lifetimes are much shorter than the ocean origin lifetimes likely due to the timing of the initial convection. Recalling that most land convection occurs closer to the evening, this means that the thicker anvil exists for most of its life during solar night. This is compared to the oceanic convection that sees a small morning peak, followed by a much greater proportion occurring during the day than at night. This means that for ocean, most of the thick anvil will spend its evolution in the solar day. Detrained cirrus is likely to dissipate much faster during the night due to, among other factors, the entrainment of dry air acting to reduce anvil lifetime (Gasparini et al., 2022). These lifetime differences also support findings by Sokol and Hartmann (2020) who found that anvils thin more rapidly during the night due to the net radiative cooling which erodes the cloud top. As well as the lifetimes being different between land-origin and ocean-origin-detrained cirrus, the distribution shown in Figure 4 shows an oscillation in the rate of decay for both origin cirrus. Further work is needed, however this may provide some extra evidence of a diurnal cycle effect on the lifetime.




Figure 4. Total distribution of the lifetimes of ocean-origin and land-origin.

3.2 Cloud Radiative Effect - Detrained versus In Situ

Figure 5 shows the LW, SW and net high cloud CRE for detrained ~~air parcels and in situ air parcels, and the mean of both,~~
~~weighted by relative distribution of the detrained cirrus air parcels and in situ air parcels,~~ in situ and net of all air parcels as
 260 a function of ~~time since convection-~~ TSC. They are weighted in each TSC bin by the relative occurrence of each bin. The total
 net CRE is negative (a cooling) at low TSC values becoming warming after ≈ 6 hours, reflecting the shift from optically thick
 deep convective core and anvil cirrus to thinner cirrus. For the first 10 hours, the mean high CRE is the same as the detrained
 cirrus CRE, as almost all air parcels at this point are considered detrained (Fig. 35).


The low occurrence of in situ air parcels at low TSC accounts for the erratic jumps in the weighted mean of the in situ CRE
 265 close to convection (Fig.5). As these in situ air parcels become more common, the mean high-cloud CRE approaches the in
 situ CRE, although some differences remain even at several days from convection. This shift in the high-cloud CRE is most
 evident when considering the LW and SW CRE separately. The detrained clouds have a larger SW cooling, LW warming, and
 stronger net positive CRE than the in situ air parcels. There is still a small trend in the CRE of the in situ air parcels even when
their occurrence is low close to convection. This is likely due to shallower, mid-tropospheric convection that is not identified
 270 as deep convection, but still contributing to the CRE along trajectories.

After the initial cooling, the net high cloud CRE peaks at $\approx 10 \text{ Wm}^{-2}$ at 10 hours since convection. There are some small
 oscillations in the CRE due to the diurnal cycle of convection formation (see Figure 7). As the proportion of in situ air parcels
 increases as a function of TSC, the mean net CRE decreases towards zero, as the in situ air parcels have a CRE closer to zero
 than the detrained cirrus clouds. At low TSC values, the change in high cloud CRE is dominated by the changing properties
 275 of the detrained cirrus clouds. As TSC increases, the change in total cirrus CRE reflects the change in distribution away from
 detrained cirrus clouds, towards in situ cirrus air parcels.

The total lifetime CRE of the high clouds for the entire tropics is calculated as the ~~sum of average~~ the CRE along trajectories,
 weighted according to the distribution of TSC values in the tropics and the latitude that the values occur at, (see Fig 3b inset).
 The total lifetime ~~forcing-radiative effect~~ of high clouds in the tropics along trajectories from deep convection is $3.6-5.1 \pm$

280 $0.3-0.4 \text{ Wm}^{-2}$. The total radiative effect of the detrained cirrus is found to be $1.4 \pm 0.1 \text{ Wm}^{-2}$, whereas the total radiative effect from the in situ is found to be $9.2 \pm 0.5 \text{ Wm}^{-2}$. Although when summed over their whole lifetime detrained cirrus are more cooling than in situ, extending their lifetime still increases the overall warming because they have a greater instantaneous warming at longer TSC values than the in situ.

The change in the total lifetime forcing of the tropical cirrus for an increase in the lifetime of detrained cirrus. The uncertainties here are the interannual variability from



2000-2016.

Figure 5. The cloud radiative effects (Net, LW, SW) as a function of the TSC for both detrained, in situ air parcels, and the weighted sum of both. The dotted line shows the modal lifetime of the detrained cirrus.

3.3 Land versus Ocean TSC distribution

285 Figure 1d) shows the instantaneous snapshot of the land and ocean origin air parcels. The majority of the tropics contain ocean origin air parcels (blue), as the majority of convection that occurs in the tropics occurs over the ocean. Therefore for a given detrained cirrus cloud, it is more likely to be oceanic than terrestrial in origin.

The land origin air parcels occur predominantly over land or near the coast. Figure 1f) shows the mean land-ocean origin value. There are large regions over the southern Atlantic ocean that contain more frequent land origin air parcels than average.

290 This is due to wind fields that carry air from land origin convection over the ocean, making it more likely to find land-origin detrained cirrus in this region compared to the rest of the global oceans.

The average air parcel origin map (Figure 1f) hides details that can be seen in the instantaneous picture (Figure 1e). There are many instances where air from terrestrial convection is carried long distances over the ocean, and vice versa for air from ocean convection.

295 3.3 Cloud Radiative Effect - Land versus Ocean

There are significant differences in the CRE of air parcels that originate from convection over land, versus those that originate over the ocean. Figure 6 shows the LW, SW and net CRE for ocean-origin air parcels and land-origin air parcels. It is found that 20% of air parcels are land-origin, and 80% are ocean-origin. The ocean-origin evolution is very similar to the all-origin evolution since the majority of the air parcels from convection are oceanic in origin. The largest differences in the land versus ocean CRE occur at low TSC values, driven predominantly by differences in the SW cooling in the early stages of the evolution. These differences arise due to the differences in timings of convection and the strong diurnal cycle in convection over land. Figure 7 shows the timing of convection from land-origin versus ocean-origin convection. There is a significant peak in convective activity over land in the late afternoon into the early evening. This is in contrast to oceanic convective which occurs comparatively uniformly throughout the day, with a smaller peak in the morning, which adds to the contrasting SW diurnal cycle between land and ocean.

Some of the difference is also due to the contrast in surface albedo between land and ocean. Further work is needed to quantify the contribution of both the albedo contrast, and the diurnal cycle difference, by focusing on land origin convection that exists over land and over ocean.




Figure 6. Total LW, SW and Net CRE as a function of TSC. Dotted lines show the land-origin CRE and solid lines show the ocean-origin CRE. The vertical lines show the modal lifetimes of the ocean and land detrained cirrus.

The timing of this convection over land means that the SW cooling from the optically thick cirrus is minimised compared to oceanic convection. Therefore the total net CRE is positive in the first hours from convection. By the time the incoming solar radiation is maximised, the cirrus has already thinned sufficiently that the SW cooling is significantly smaller than the LW warming. As a result, the land-origin air parcels have a positive CRE throughout its entire lifecycle. The timing of terrestrial convection is shown to play a vital role in determining the total CRE of the detrained air parcels (Jones et al., 2023).

The differences in the LW warming between land and ocean-origin air parcels are less clear. In the early stages of the convection, they are similar. After 20 hours, the land-origin air parcels appears to have a slightly larger LW warming than the ocean-origin air parcels. This could be ~~de~~ due to stronger convection occurring over land, leading to higher altitude cirrus producing a larger LW warming (Takahashi et al., 2017). The land-origin parcels also have a stronger SW cooling after the initial stage of convection, suggesting that the detrained cirrus from land-origin convection are optically thicker than their ocean-origin counterparts.

Over the entire range of lifetimes, the weighted ~~sum~~ average of the land-origin CRE is ~~6.3~~ 7.7 \pm ~~0.5~~ 0.6 Wm^{-2} , the ocean-origin has a total CRE of ~~2.9~~ 4.3 \pm 0.4 Wm^{-2} . This difference in the lifetime CRE is almost entirely due to the differences in the




Figure 7. ~~The normalised diurnal~~ Diurnal cycle of tropical deep convection, normalised such that the maxima of each cycle is one, for (a) the entire globe, (b) land origin convection (c) ocean origin convection. The error bars correspond to the standard deviation of the interannual variability between 1984 and 2016.

timing of the initial convection, and as such the lack of SW cooling that occurs in land-origin cirrus compared to ocean-origin cirrus.

3.4 Lifetime adjustments

325 Given that the CRE of detrained cirrus clouds is very different to the CRE of in situ cirrus air parcels (Fig. 5), the lifetime of the detrained cirrus clouds are very important when determining the total tropical CRE. As mentioned above, the median lifetime of the detrained cirrus is approximately 36 hours, with a modal lifetime of 25 hours. This is shown separately for land and ocean origin cirrus in Figure 4. This is defined as the mean TSC when the air parcel changes from detrained to in situ.

The impact of extensions to detrained cirrus lifetime on the tropical CRE can be estimated through an artificial modification
 330 of the detrained cirrus distribution. The lifetime of the detrained cirrus can be artificially extended by modifying the fractional distribution of the detrained cirrus in Figure 3, a novel method introduced in this work. In this work, a simple approach to extend the distribution of the detrained cirrus is used. The original distribution is stretched along the TSC-axis by a given percentage, and this new distribution can be thought to represent the distribution of detrained cirrus in this new extended lifetime regime. This is shown in Figure 8. By stretching the distribution along the x-axis by some factor the amount of detrained cirrus is
 335 artificially increased for a given TSC.

It is important to note that the TSC distribution is not changed when the lifetime of detrained cirrus are extended, just the radiative properties of the cirrus as a function of TSC. This decouples the cloud evolution from the convective occurrence and

Total distribution of the lifetimes of ocean-origin and land-origin.




Figure 8. The fractional distribution of the detrainment versus in situ air parcels. Dotted lines represent various distributions when the detrainment cirrus are extended by a percentage.

large scale flow. This change ~~therefore~~, therefore, maintains the overall shape in the distribution of TSC, whilst increasing the proportion of detrainment cirrus air parcels for a given TSC. This increases the mean detrainment cirrus lifetime by the prescribed factor.

Increasing the distribution is analogous to increasing the mean lifetime of each detrainment cirrus by the same percentage. Figure 8 shows that for a 50% increase in the detrainment cirrus lifetime, this shifts the point at which the number of in situ and detrainment parcels are equal from around 30 hours to just under 50 hours.

~~Figure 4 shows the distribution of the lifetime of the land and ocean-origin detrainment cirrus. The mean lifetime for land-origin detrainment cirrus is 31 hours, compared to 39 hours for ocean-origin detrainment cirrus. Therefore a 50% increase in the lifetime of the detrainment cirrus corresponds approximately to a 16-hour and 20-hour increase in the lifetime of the land-origin and ocean-origin detrainment cirrus respectively.~~

As the all-cirrus LW, SW and net CREs along trajectories from convection are a weighted average of the CRE contributions of detrainment cirrus and in situ air parcels, changing the lifetime of the detrainment cirrus changes the weighting and thus total CRE along the trajectories. With detrainment cirrus having a stronger warming effect, extending their lifetime increases the overall warming of tropical cirrus clouds.

Figure 9 shows how the total tropical CRE changes when the lifetime of the detrainment cirrus is increased. In all cases, the the tropical CRE increases almost linearly for a given increase in the lifetime of the detrainment cirrus. For a doubling in the lifetime of all-origin detrainment cirrus, the tropical CRE increases from a baseline of ~~3.6~~ 5.0 ± 0.4 Wm^{-2} to ~~4.8~~ 5.7 ± 0.5 Wm^{-2} . ~~This is an increase of over 50% in the high cloud CRE.~~ The uncertainties given represent the interannual variability in the increase in CRE for a given increase in detrainment cirrus lifetime. In practice these uncertainties are likely larger due to the inherent uncertainties in the initial calculation of CRE, however these are normalised to a baseline, therefore there is no uncertainty in the lifetime extension when the lifetime extension is zero hours.




Figure 9. ~~Total LW~~ Change in CRE for all, ~~SW~~ land and ~~Net CRE~~ ocean origin air parcels for high-clouds as a function of TSC. ~~Dotted lines show given percentage increase in the weighted contributions lifetime of the detrained and in situ flagged grid boxes to the total CRE cirrus clouds.~~

The increase in the high cloud CRE is similar for an increase in the lifetime of ocean-origin detrained cirrus. This is to
 360 be expected given the greater proportion of convection originating over ocean versus land. A doubling in the ocean-origin detrained lifetime leads to an increase in the tropical CRE of ~~0.9~~ $0.3 \pm 0.1 \text{ Wm}^{-2}$, from a baseline CRE of ~~2.9~~ $4.4 \pm 0.4 \text{ Wm}^{-2}$, to ~~3.8~~ $4.7 \pm 0.4 \text{ Wm}^{-2}$.

Interestingly the increase is not linear over time, with the first 50% lifetime extension increasing the all-origin CRE by
 0.06 Wm^{-2} more than the final 50% increase. One reason for this is the decreasing difference between the in situ CRE and
 365 detrained CRE at longer TSC values. Therefore any adjustment to the relative distributions at longer TSC will have a slightly smaller impact than initial adjustments to the shorter lifetimes.

For land-origin cirrus the increase in tropical CRE is much greater for a given increase in the detrained cirrus lifetime. A
 doubling the land-origin detrained cirrus lifetime increases the tropical CRE from a baseline of ~~6.3~~ $7.7 \pm 0.5 \text{ Wm}^{-2}$ to ~~8.6~~ $9.7 \pm 0.5 \text{ Wm}^{-2}$. This is because land-origin detrained cirrus are much more warming than ocean-origin cirrus, particularly
 370 at low TSC values (Figure 6). This is due in part to a smaller SW cooling in land-origin detrained cirrus because of differences in the timing of convection. This means that increasing the proportion of detrained cirrus for a given TSC, increases the overall CRE by a larger amount than the same increase for ocean-origin detrained cirrus.

Figure 10 shows the LW, SW and Net CRE as a function of TSC, for a baseline, as well as a ~~25~~ 50% and a 100% increase in the lifetime of detrained cirrus. This shows an increase in the Net CRE at all TSC values given an increase in the lifetime
 375 of detrained cirrus. At very low TSC values there is little change in the net CRE for an increase in lifetime. This is because at these low TSC values, detrained air parcels account for almost all the air parcels. Therefore there is no change in the relative distributions of detrained versus in situ air parcels (see Figure 3), and no change in the overall CRE at low TSC values.




Figure 10. Total net (left axis) and LW and SW (right axis) CRE as a function of TSC. The dashed and dotted lines represent a 25% and 50% and 100% increase in the lifetime of detrained cirrus respectively. Note the separate axis for Net, LW and SW ~~foreign~~ radiative effects, introduced here to show the changes in CRE for each lifetime extension more clearly.

4 Discussion

In this work, the overall high cloud CRE, including both detrained origin cirrus and in situ origin air parcels (which may contain in situ cirrus) was found to be ~~3.6~~ 5.0 $\pm 0.4 \text{ Wm}^{-2}$. However, there are still a number of uncertainties that need addressing. These uncertainties fall into three main categories: a) isolating the ~~foreign~~ radiative effects from high clouds, b) the distinction between detrained and in situ air parcels, and c) the method for extending the lifetime.

In order to isolate the ~~foreign~~ radiative effects of high and low clouds, this paper attempts to consider the contributing radiative effect of the low clouds by calculating the monthly mean ‘background albedo’, which is then subtracted from the instantaneous all-sky albedo to give the ‘high cloud’ albedo. The background albedo is the all-sky albedo in regions with no high cloud. This method gives an approximate high cloud SW CRE. However, it relies on the assumption that ~~that~~ the monthly mean background albedo is approximately representative of the instantaneous background albedo, if it were possible to be calculated for everywhere in the tropics at each timestep.

As the annual mean background albedo is subtracted from the instantaneous all-sky, there are occasions where the annual background is greater than the all-sky Figure 2. This produces a negative high cloud albedo and thus a positive SW cooling value. It can be assumed in these regions this occurs when the all-sky is close to the clear-sky albedo value, with total cloud fraction being lower than the average low cloud fraction. ~~These-~~This assumes that there is little correlation between the low and high cloud fraction.For a further discussion of cloud overlap statistics, see Tompkins and Giuseppe (2015). The impact of these cases averages to zero when calculating the annual mean SW high cloud CRE, and are less important for ~~our~~this work's analysis as they occur at large TSC values in regions where there is little high cloud. This situation occurs in regions where the average low cloud cover is high, and therefore the average background albedo is high. Therefore, if a trajectory is observed at this location that contains an optically thin high cloud, and little to no low cloud, the observed albedo will be lower than the average albedo, thus resulting in a 'negative' high cloud SW CRE. It is also known that the ISCCP cloud type histograms may over attribute the TOA cooling to low clouds (Stephens et al., 2018). This may lead to a positive (warming) bias in the results in this work, therefore it should be understood that these results represent an upper bound. However this bias is likely small in comparison to other uncertainties mentioned above.

The second area of uncertainty in this work surrounds the definition of detrained cirrus. This work defines the end of a detrained cirrus lifetime, and the beginning of the in situ air parcels, as the point at which a cloud along a trajectory from deep convection moves below 10% cirrus cloud fraction for the first time. Any cirrus that then appears along a trajectory after this point is defined as in situ in origin. This is similar to Luo and Rossow (2004) who define the end of their cirrus lifetime as the point at which the cirrus cloud reaches 1/5 of the maximum cloud fraction along the trajectory. Changing the definition of detrained cirrus would not change the overall high cloud CRE. However, it ~~would change our~~does change the calculated lifetime of detrained cirrus, which is shown in Figure S1 in the supplementary. There is no universal definition for “detrained” or “anvil” cirrus, and as such the lifetimes of these clouds vary depending on how they are defined. Nevertheless, ~~our lifetimes~~the lifetimes in this work fall within the expected ranges given in the literature ~~Luo and Rossow (2004)~~(Luo and Rossow, 2004) and as shown in Figure S2, the final values for the change in CRE for a given lifetime extension are not particularly sensitive to the threshold used to define the convection.

The method used to extend the lifetime of the detrained cirrus is relatively idealised, insofar as it models a lifetime extension as a change in the distribution of detrained cirrus at the expense of in situ cirrus. Moreover, the extension in the distribution modifies the distribution mostly at the tail end of the detrained cirrus lifetimes, meaning that the oldest detrained cirrus are the ones that see the largest increase. The purpose of this work was not to assess the methods through which a lifetime extension would occur. Instead, the aim was to provide a bound on the impact that increasing the lifetime of the detrained cirrus would have on the tropical high cloud CRE. By modifying the distribution to represent an increase in lifetime, particularly in a way that may impact the longer lived detrained cirrus more than the short lived cirrus, we do provide such an upper bound, since any modification to the shorter lived cirrus would not increase the CRE by as much, as they are already more cooling. In reality, any physical routes through which a lifetime extension will likely increase the total CRE by less than the values we provide here. Further work is needed to assess the mechanisms through which lifetime extensions might occur, and what the range of impacts this may have on the CRE. For example, the lifetime could change due to a stronger clustering of convective cores

(Jones et al., 2024) or increased updrafts via aerosol invigoration (Abbott and Cronin, 2021). Each of these mechanisms may impact the lifetime in a distinct way from the idealised set up in this work. Investigating these mechanisms and the specific impacts they had on the lifetime would make for an interesting comparison study to the idealised extension proposed in this work, and would be a necessary addition to put these results into context, as well as developing a stronger constraint on the potential changes of the CRE.

As well as showing the difference in detrained cirrus CRE due to the origin of the convection, this work shows how extending the lifetime of the detrained cirrus changes the overall lifetime CRE. In this work, the lifetime is a simple distribution, and extending the lifetime is merely a case of extending the distribution at the expense of in situ air parcels. This assumes that every cirrus cloud in the tropics is either detrained or in situ and these are distinct from each other, i.e. an increase in detrained air parcel amount should correlate with a decrease in in situ air parcel amount. The purpose of extending the lifetime isn't to obtain a concrete value of the CRE for a given lifetime extension, but instead to indicate the trend in CRE for an increase in lifetime. This helps to bound estimates of how much the lifetime of a detrained cirrus would need to change to significantly affect the tropical high cloud CRE.

This work finds that increasing the lifetime of the detrained cirrus increases the overall high cloud CRE in the tropics, resulting in an overall warming. This is because on average the detrained cirrus have a larger net CRE than the in situ air parcels. For a 50% increase in the lifetime, the average CRE of tropical cirrus increases by $0.6-0.3 \pm 0.1 \text{ Wm}^{-2}$. Comparing this to aerosol invigoration studies which have shown lifetime extensions of detrained cirrus on the order of 30% (Zang et al., 2023), it may be the case that aerosol invigoration may not have a large impact on the total tropical CRE, however more work is needed to constrain the lifetime extension from aerosol invigoration, which is currently highly uncertain.

The average lifetime of detrained cirrus clouds is found to be approximately 36 hours, similar to that found by Luo and Rossow (2004). This implies that increasing the average lifetime of detrained cirrus clouds by 15 hours would increase the overall high cloud CRE by $0.6-0.4 \pm 0.1 \text{ Wm}^{-2}$.

This increase is larger for land-origin convection, which increases by $1.2-1.1 \pm 0.1 \text{ Wm}^{-2}$ for a 15 hour increase in lifetime, and smaller for ocean-origin convection which increases by $0.4-0.2 \pm 0.1 \text{ Wm}^{-2}$.

5 Conclusions

The lifetime of detrained cirrus clouds from convection plays an important role in determining the overall CRE of high clouds detrained CRE in the tropics, and this may impact the total high cloud CRE in the tropics. Small changes in the evolution of these clouds, either driven by their initial conditions or meteorological environment have been hypothesised to produce large responses in the high cloud CRE, making understanding this evolution essential to constrain both climate forcings and feedbacks. Tracking the evolution of cirrus clouds throughout the tropics as they move away from convective events and isolating the high cloud CRE from the background albedo variation caused by low clouds, this work characterises the lifecycle of the cirrus CRE using satellite observations. Defining the lifetime of detrained cirrus based on the point along a trajectory following convection where the high cloud fraction drops below 10% (representing the time when the detrained

460 cirrus has dissipated), the impact of cirrus formation mechanisms on CRE is isolated. By tracking the properties of the initial convection, this method also allows the impact of the initial convection on the CRE to be investigated. The tracking approach used in this work differs from previous studies, such as (Jones et al., 2023), by using ISCCP data with a much longer time record, as well as covering the entirety of the tropics, without explicitly tracking individual clouds at all. This allows for regional variations such as the land ocean contrast to be thoroughly investigated.

Most detrained air parcels remain in the central tropical belt, close to active convection, while in situ air parcels exist throughout the tropics (Figs. 1b,e). The high cloud CRE of the detrained cirrus air parcels is found to be considerably more warming than the CRE of in situ air parcels (Figure 5). The high cloud radiative effect is calculated by subtracting the variability 465 in the background albedo due to changes in low clouds from the overall CRE (Figure 2). Calculating the total high cloud CRE as the weighted mean CRE along trajectories from deep convection gives a total high cloud topical CRE as $3.6\text{--}5.0 \pm 0.4 \text{ Wm}^{-2}$.

Air parcels along trajectories from oceanic convection are compared to those from detrained convection (1c) and f)). It is found that air from land convection exists over large regions of the oceans, and air from oceanic convection often exists over 470 land. The high cloud CRE along trajectories from land-origin convection is $6.3\text{--}8.8 \pm 0.5\text{--}0.4 \text{ Wm}^{-2}$, compared to $2.9\text{--}3.4 \pm 0.4 \text{ Wm}^{-2}$ for cirrus along trajectories from ocean-origin convection. Strong differences in the timing of the initial convection over land and ocean (Figure 7) lead to these differences in the net CRE of detrained cirrus clouds as a function of TSC. Detrained cirrus clouds from land are found to have a net warming across their entire lifetime, as land convection occurs much later in the day than oceanic convection (Figure 7). This reduces the SW cooling from the thick anvils at low TSC values compared to 475 oceanic convection, which detrained from convection much earlier in the day when the SW cooling is still significant. This is similar to the results of Jones et al. (2023), who showed the importance of the diurnal cycle in determining the lifetime CRE of detrained cirrus clouds.

The lifetime of detrained cirrus clouds is artificially increased by stretching the distribution of the detrained cirrus air parcels at the expense of the in situ air parcels (Figure 8). These extended distributions are then used as the weights when calculating 480 the total cirrus CRE in the tropics. It is found that increasing the lifetime of the detrained cirrus clouds increases the total CRE in the tropics almost linearly. For example, a 50% increase in the detrained cirrus lifetime, from an average of 36 hours to 54 hours, increases the total tropical CRE from $3.6\text{--}4.4 \pm 0.4 \text{ Wm}^{-2}$ to $4.8 \pm 0.4 \text{ Wm}^{-2}$.

The impacts of lifetime extension of detrained cirrus air parcels is calculated for both land-origin and ocean-origin cirrus. Increasing the land-origin detrained cirrus lifetime by 50% increases the total land-origin CRE in the tropics by $1.2\text{--}1.1 \pm 0.1$ 485 Wm^{-2} . On the other hand, increasing the ocean-origin detrained cirrus lifetime by 50% increases the total ocean-origin CRE in the tropics by only $0.5\text{--}0.2 \pm 0.1 \text{ Wm}^{-2}$. These contrasts arise because of the differences in the CRE of land-origin and ocean-origin detrained cirrus in the early stages of their development. Ocean-origin detrained cirrus have a negative (cooling) CRE at low TSC, therefore any increase in the lifetime of the detrained cirrus also increases the time spent cooling, thus dampening the warming impact they may also have at longer TSC. Land-origin detrained cirrus are warming at all stages of 490 their development, therefore any increase in their lifetime will amplify their warming impact (see Figure 6).

While this work shows the potential for cirrus lifetime changes to modify high cloud CRE in the tropics, further work is required to assess and constrain these potential changes in detrained cirrus lifetime. In particular, to assess how specific mechanisms through which detrained cirrus lifetimes may be extended (such as aerosol invigoration), impact the overall CRE, within the bounds provided in this work. Building on this estimate of how the radiative effect of these clouds responds to lifetime extensions, the impact of meteorological or anthropogenic drivers (such as changes in SSTs or aerosol invigoration) have on the lifetime of anvil cirrus, and therefore the overall tropical cirrus CRE must be bounded to fully constrain the magnitude of this effect.

Data availability. The data used is highlighted in Sect. 2.1. The ISCCP-H data were obtained from the NCEI/NOAA Climate Data Records (<https://doi.org/10.7289/V5QZ281S>, Rossow et al. (2017))

The ERA5 data were obtained from the Climate Data Store (<https://doi.org/10.24381/cds.bd0915c6>, Hersbach et al. (2018)).

The CERES data were obtained from the NASA EARTHDATA ASDC (https://doi.org/10.5067/TERRA+AQUA/CERES/SYN1DEG-1HOUR_L3.004A, NASA/LARC/SD/ASDC (2017)).

Author contributions. Both authors contributed to study design and interpretation of results. GH performed the analysis and prepared the manuscript with comments from EG

Competing interests. The contact author has declared that neither of the authors has any competing interests.

Acknowledgements. This research has been supported by the Royal Society (grant no. URF/R1/191602) and the Horizon Europe programme under Grant Agreement No 101137680 via project CERTAINTY (Cloud-aERosol inTeractions & their impActs IN The earth sYstem).

References

- Abbott, T. H. and Cronin, T. W.: Aerosol invigoration of atmospheric convection through increases in humidity, *Science*, 371, 83–85, <https://doi.org/10.1126/science.abc5181>, 2021.
- Albright, M. D., Recker, E. E., Reed, R. J., and Dang, R.: The diurnal variation of deep convection and inferred precipitation in the central tropical Pacific during January–February 1979, *Monthly weather review*, 113, 1663–1680, 1985.
- Berry, E. and Mace, G. G.: Cloud properties and radiative effects of the Asian summer monsoon derived from A-Train data, *Journal of Geophysical Research: Atmospheres*, 119, 9492–9508, <https://doi.org/10.1002/2014JD021458>, 2014.
- Beydoun, H., Caldwell, P. M., Hannah, W. M., and Donahue, A. S.: Dissecting Anvil Cloud Response to Sea Surface Warming, *Geophysical Research Letters*, 48, e2021GL094049, <https://doi.org/https://doi.org/10.1029/2021GL094049>, e2021GL094049 2021GL094049, 2021.
- Bony, S., Stevens, B., Frierson, D. M., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., et al.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261–268, 2015.
- Bony, S., Stevens, B., Coppin, D., Becker, T., Reed, K. A., Voigt, A., and Medeiros, B.: Thermodynamic control of anvil cloud amount, *Proceedings of the National Academy of Sciences*, 113, 8927–8932, 2016.
- Chen, S. S. and Houze Jr, R. A.: Diurnal variation and life-cycle of deep convective systems over the tropical pacific warm pool, *Quarterly Journal of the Royal Meteorological Society*, 123, 357–388, <https://doi.org/https://doi.org/10.1002/qj.49712353806>, 1997.
- Choi, Y.-S., Kim, W., Yeh, S.-W., Masunaga, H., Kwon, M.-J., Jo, H.-S., and Huang, L.: Revisiting the iris effect of tropical cirrus clouds with TRMM and A-Train satellite data, *Journal of Geophysical Research: Atmospheres*, 122, 5917–5931, 2017.
- Cronin, T. W. and Wing, A. A.: Clouds, circulation, and climate sensitivity in a radiative-convective equilibrium channel model, *Journal of Advances in Modeling Earth Systems*, 9, 2883–2905, 2017.
- Dai, A.: Global precipitation and thunderstorm frequencies. Part II: Diurnal variations, *Journal of Climate*, 14, 1112–1128, 2001.
- Dai, A., Giorgi, F., and Trenberth, K.: Observed and model simulated precipitation diurnal cycle over the contiguous United States, *J. Geophys. Res.*, 104, 6377–6402, 1999.
- Del Genio, A. D. and Kovari, W.: Climatic properties of tropical precipitating convection under varying environmental conditions, *Journal of Climate*, 15, 2597–2615, 2002.
- Fan, J., Leung, L. R., Rosenfeld, D., Chen, Q., Li, Z., Zhang, J., and Yan, H.: Microphysical effects determine macrophysical response for aerosol impacts on deep convective clouds, *Proceedings of the National Academy of Sciences*, 110, E4581–E4590, <https://doi.org/10.1073/pnas.1316830110>, 2013.
- Gasparini, B., Blossey, P. N., Hartmann, D. L., Lin, G., and Fan, J.: What Drives the Life Cycle of Tropical Anvil Clouds?, *Journal of Advances in Modeling Earth Systems*, 11, 2586–2605, <https://doi.org/10.1029/2019MS001736>, 2019.
- Gasparini, B., Rasch, P. J., Hartmann, D. L., Wall, C. J., and Dütsch, M.: A Lagrangian Perspective on Tropical Anvil Cloud Lifecycle in Present and Future Climate, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033487, <https://doi.org/10.1029/2020JD033487>, e2020JD033487 2020JD033487, 2021.
- Gasparini, B., Sokol, A. B., Wall, C. J., Hartmann, D. L., and Blossey, P. N.: Diurnal Differences in Tropical Maritime Anvil Cloud Evolution, *Journal of Climate*, 35, 1655 – 1677, <https://doi.org/10.1175/JCLI-D-21-0211.1>, 2022.
- Gasparini, B., Sullivan, S. C., Sokol, A. B., Kärcher, B., Jensen, E., and Hartmann, D. L.: Opinion: Tropical cirrus – from micro-scale processes to climate-scale impacts, *Atmospheric Chemistry and Physics*, 23, 15 413–15 444, <https://doi.org/10.5194/acp-23-15413-2023>, 2023.

- 545 Gehlot, S. and Quaas, J.: Convection-Climate Feedbacks in the ECHAM5 General Circulation Model: Evaluation of Cirrus Cloud Life Cycles with ISCCP Satellite Data from a Lagrangian Trajectory Perspective, *Journal of Climate*, 25, 5241 – 5259, <https://doi.org/10.1175/JCLI-D-11-00345.1>, 2012.
- Harrison, E. F., Minnis, P., Barkstrom, B. R., Ramanathan, V., Cess, R. D., and Gibson, G. G.: Seasonal variation of cloud radiative forcing derived from the Earth Radiation Budget Experiment, *Journal of Geophysical Research: Atmospheres*, 95, 18 687–18 703, <https://doi.org/10.1029/JD095iD11p18687>, 1990.
- 550 Hartmann, D. L. and Berry, S. E.: The balanced radiative effect of tropical anvil clouds, *Journal of Geophysical Research: Atmospheres*, 122, 5003–5020, <https://doi.org/10.1002/2017JD026460>, 2017.
- Hartmann, D. L. and Larson, K.: An important constraint on tropical cloud-climate feedback, *Geophysical research letters*, 29, 12–1, 2002.
- Hartmann, D. L. and Michelsen, M. L.: No evidence for iris, *Bulletin of the American Meteorological Society*, 83, 249–254, 2002.
- 555 Hartmann, D. L., Moy, L. A., and Fu, Q.: Tropical Convection and the Energy Balance at the Top of the Atmosphere, *Journal of Climate*, 14, 4495 – 4511, [https://doi.org/10.1175/1520-0442\(2001\)014<4495:TCATEB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<4495:TCATEB>2.0.CO;2), 2001.
- Hartmann, D. L., Gasparini, B., Berry, S. E., and Blossey, P. N.: The Life Cycle and Net Radiative Effect of Tropical Anvil Clouds, *Journal of Advances in Modeling Earth Systems*, 10, 3012–3029, <https://doi.org/https://doi.org/10.1029/2018MS001484>, 2018.
- Herbert, R. and Stier, P.: Satellite observations of smoke–cloud–radiation interactions over the Amazon rainforest, *Atmospheric Chemistry and Physics*, 23, 4595–4616, <https://doi.org/10.5194/acp-23-4595-2023>, 2023.
- 560 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1959 to present., <https://doi.org/10.24381/cds.bd0915c6>, 2018.
- Horner, G. and Gryspeerdt, E.: The evolution of deep convective systems and their associated cirrus outflows, *Atmospheric Chemistry and Physics*, 23, 14 239–14 253, <https://doi.org/10.5194/acp-23-14239-2023>, 2023.
- 565 Jensen, E. J., Pan, L. L., Honomichl, S., Diskin, G. S., KrÄmer, M., Spelten, N., GÄEnther, G., Hurst, D. F., Fujiwara, M., Vamel, H., Selkirk, H. B., Suzuki, J., Schwartz, M. J., and Smith, J. B.: Assessment of Observational Evidence for Direct Convective Hydration of the Lower Stratosphere, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032 793, <https://doi.org/https://doi.org/10.1029/2020JD032793>, e2020JD032793 10.1029/2020JD032793, 2020.
- 570 Jones, W. K., Stengel, M., and Stier, P.: A Lagrangian Perspective on the Lifecycle and Cloud Radiative Effect of Deep Convective Clouds Over Africa, *EGUsphere*, 2023, 1–25, <https://doi.org/10.5194/egusphere-2023-2059>, 2023.
- Jones, W. K., Stengel, M., and Stier, P.: A Lagrangian perspective on the lifecycle and cloud radiative effect of deep convective clouds over Africa, *Atmospheric Chemistry and Physics*, 24, 5165–5180, <https://doi.org/10.5194/acp-24-5165-2024>, 2024.
- Kiehl, J. T.: On the Observed Near Cancellation between Longwave and Shortwave Cloud Forcing in Tropical Regions, *Journal of Climate*, 7, 559–565, [https://doi.org/10.1175/1520-0442\(1994\)007<0559:OTONCB>2.0.CO;2](https://doi.org/10.1175/1520-0442(1994)007<0559:OTONCB>2.0.CO;2), 1994.
- 575 Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A., and Rudich, Y.: Aerosol invigoration and restructuring of Atlantic convective clouds, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005GL023187>, 2005.
- Koren, I., Feingold, G., and Remer, L. A.: The invigoration of deep convective clouds over the Atlantic: aerosol effect, meteorology or retrieval artifact?, *Atmospheric Chemistry and Physics*, 10, 8855–8872, <https://doi.org/10.5194/acp-10-8855-2010>, 2010.
- 580 Krishna, U. V. M., Das, S. K., Deshpande, S. M., and Pandithurai, G.: Physical processes controlling the diurnal cycle of convective storms in the Western Ghats, *Scientific Reports*, 11, 14 103, <https://doi.org/10.1038/s41598-021-93173-0>, 2021.

- Lee, J., Yang, P., Dessler, A. E., Gao, B.-C., and Platnick, S.: Distribution and Radiative Forcing of Tropical Thin Cirrus Clouds, *Journal of the Atmospheric Sciences*, 66, 3721 – 3731, <https://doi.org/10.1175/2009JAS3183.1>, 2009a.
- Lee, J., Yang, P., Dessler, A. E., Gao, B.-C., and Platnick, S.: Distribution and radiative forcing of tropical thin cirrus clouds, *Journal of the atmospheric sciences*, 66, 3721–3731, 2009b.
- Lin, J. C., Matsui, T., Pielke Sr., R. A., and Kummerow, C.: Effects of biomass-burning-derived aerosols on precipitation and clouds in the Amazon Basin: a satellite-based empirical study, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006884>, 2006.
- Lopez-Bravo, C., Vincent, C. L., Huang, Y., and Lane, T. P.: The Diurnal Cycle of Rainfall and Deep Convective Clouds Around Sumatra and the Associated MJO-Induced Variability During Austral Summer in Himawari-8, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD039132, <https://doi.org/https://doi.org/10.1029/2023JD039132>, e2023JD039132 2023JD039132, 2023.
- Luebke, A. E., Afchine, A., Costa, A., Groß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmospheric Chemistry and Physics*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- Luo, Z. and Rossow, W. B.: Characterizing Tropical Cirrus Life Cycle, Evolution, and Interaction with Upper-Tropospheric Water Vapor Using Lagrangian Trajectory Analysis of Satellite Observations, *Journal of Climate*, 17, 4541 – 4563, <https://doi.org/10.1175/3222.1>, 2004.
- Mace, G. G., Deng, M., Soden, B., and Zipser, E.: Association of Tropical Cirrus in the 10–15-km Layer with Deep Convective Sources: An Observational Study Combining Millimeter Radar Data and Satellite-Derived Trajectories, *Journal of the Atmospheric Sciences*, 63, 480–503, <https://doi.org/10.1175/JAS3627.1>, 2006.
- Massie, S., Gettelman, A., Randel, W., and Baumgardner, D.: Distribution of tropical cirrus in relation to convection, *Journal of Geophysical Research: Atmospheres*, 107, AAC 19–1–AAC 19–16, <https://doi.org/10.1029/2001JD001293>, 2002.
- NASA/LARC/SD/ASDC: CERES and GEO-Enhanced TOA, Within-Atmosphere and Surface Fluxes, Clouds and Aerosols 1-Hourly Terra-Aqua Edition4A, 10.5067/TERRA+AQUA/CERES/SYN1DEG-1HOUR_L3.004A, 2017.
- Nazaryan, H., McCormick, M. P., and Menzel, W. P.: Global characterization of cirrus clouds using CALIPSO data, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Ohno, T. and Satoh, M.: Roles of cloud microphysics on cloud responses to sea surface temperatures in radiative-convective equilibrium experiments using a high-resolution global nonhydrostatic model, *Journal of Advances in Modeling Earth Systems*, 10, 1970–1989, 2018.
- Raghuraman, S. P., Medeiros, B., and Gettelman, A.: Observational Quantification of Tropical High Cloud Changes and Feedbacks, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD039364, <https://doi.org/10.1029/2023JD039364>, e2023JD039364, 2024.
- Ramanathan, V., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., and Hartmann, D.: Cloud-Radiative Forcing and Climate: Results from the Earth Radiation Budget Experiment, *Science*, 243, 57–63, <https://doi.org/10.1126/science.243.4887.57>, 1989.
- Riihimäki, L. D., McFarlane, S. A., Liang, C., Massie, S. T., Beagley, N., and Toth, T. D.: Comparison of methods to determine Tropical Tropopause Layer cirrus formation mechanisms, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016832>, 2012.
- Riley, E. M., Mapes, B. E., and Tulich, S. N.: Clouds Associated with the Madden-Julian Oscillation: A New Perspective from CloudSat, *Journal of the Atmospheric Sciences*, 68, 3032 – 3051, <https://doi.org/https://doi.org/10.1175/JAS-D-11-030.1>, 2011.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or Drought: How Do Aerosols Affect Precipitation?, *Science*, 321, 1309–1313, <https://doi.org/10.1126/science.1160606>, 2008.

- 620 Rossow, W., Golea, V., Walker, A., Knapp, K., Young, A., Hankins, B., and Inamdar, A.: International Satellite Cloud Climatology Project (ISCCP) Climate Data Record, H-Series, <https://doi.org/10.7289/V5QZ281S>, 2017.
- Ruppert, J. H. and Hohenegger, C.: Diurnal Circulation Adjustment and Organized Deep Convection, *Journal of Climate*, 31, 4899 – 4916, <https://doi.org/https://doi.org/10.1175/JCLI-D-17-0693.1>, 2018.
- Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol lidar and infrared pathfinder satellite
625 observations (CALIPSO) measurements, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Satoh, M., Iga, S.-i., Tomita, H., Tsushima, Y., and Noda, A. T.: Response of upper clouds in global warming experiments obtained using a global nonhydrostatic model with explicit cloud processes, *Journal of Climate*, 25, 2178–2191, 2012.
- Sokol, A. B. and Hartmann, D. L.: Tropical Anvil Clouds: Radiative Driving Toward a Preferred State, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD033 107, <https://doi.org/https://doi.org/10.1029/2020JD033107>, e2020JD033107 10.1029/2020JD033107, 2020.
- 630 Stephens, G., Winker, D., Pelon, J., Trepte, C., Vane, D., Yuhas, C., L  cuyer, T., and Lebsock, M.: CloudSat and CALIPSO within the A-Train: Ten Years of Actively Observing the Earth System, *Bulletin of the American Meteorological Society*, 99, 569 – 581, <https://doi.org/10.1175/BAMS-D-16-0324.1>, 2018.
- Stubenrauch, C., Ch  din, A., R  del, G., Scott, N., and Serrar, S.: Cloud properties and their seasonal and diurnal variability from TOVS Path-B, *Journal of climate*, 19, 5531–5553, 2006.
- 635 Su, H., Jiang, J. H., Neelin, J. D., Shen, T. J., Zhai, C., Yue, Q., Wang, Z., Huang, L., Choi, Y.-S., Stephens, G. L., et al.: Tightening of tropical ascent and high clouds key to precipitation change in a warmer climate, *Nature communications*, 8, 15 771, 2017.
- Sweeney, A., Fu, Q., Pahlavan, H. A., and Haynes, P.: Seasonality of the QBO Impact on Equatorial Clouds, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 737, <https://doi.org/https://doi.org/10.1029/2022JD037737>, e2022JD037737 2022JD037737, 2023.
- Takahashi, H., Luo, Z. J., and Stephens, G. L.: Level of neutral buoyancy, deep convective outflow, and convective core:
640 New perspectives based on 5 years of CloudSat data, *Journal of Geophysical Research: Atmospheres*, 122, 2958–2969, <https://doi.org/https://doi.org/10.1002/2016JD025969>, 2017.
- Tompkins, A. M. and Giuseppe, F. D.: An Interpretation of Cloud Overlap Statistics, *Journal of the Atmospheric Sciences*, 72, 2877 – 2889, <https://doi.org/10.1175/JAS-D-14-0278.1>, 2015.
- Tsushima, Y., Iga, S.-i., Tomita, H., Satoh, M., Noda, A. T., and Webb, M. J.: High cloud increase in a perturbed SST experiment with a
645 global nonhydrostatic model including explicit convective processes, *Journal of Advances in Modeling Earth Systems*, 6, 571–585, 2014.
- Varble, A. C., Igel, A. L., Morrison, H., Grabowski, W. W., and Lebo, Z. J.: Opinion: A critical evaluation of the evidence for aerosol invigoration of deep convection, *Atmospheric Chemistry and Physics*, 23, 13 791–13 808, <https://doi.org/10.5194/acp-23-13791-2023>, 2023.
- Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee, R. B., Smith, G. L., and Cooper, J. E.: Clouds and the Earth’s Radiant Energy System (CERES): An Earth Observing System Experiment, *Bulletin of the American Meteorological Society*, 77, 853–868, [https://doi.org/10.1175/1520-0477\(1996\)077<0853:CATERE>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0853:CATERE>2.0.CO;2), 1996.
- Wing, A. A., Stauffer, C. L., Becker, T., Reed, K. A., Ahn, M.-S., Arnold, N. P., Bony, S., Branson, M., Bryan, G. H., Chaboureau, J.-P., et al.: Clouds and convective self-aggregation in a multimodel ensemble of radiative-convective equilibrium simulations, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002 138, 2020.
- 655 Wylie, D. P. and Menzel, W. P.: Eight years of high cloud statistics using HIRS, *Journal of Climate*, 12, 170–184, 1999.

Zang, L., Rosenfeld, D., Pan, Z., Mao, F., Zhu, Y., Lu, X., and Gong, W.: Observing Aerosol Primary Convective Invigoration and Its Meteorological Feedback, *Geophysical Research Letters*, 50, e2023GL104151, <https://doi.org/10.1029/2023GL104151>, e2023GL104151 2023GL104151, 2023.

Zelinka, M. D. and Hartmann, D. L.: The observed sensitivity of high clouds to mean surface temperature anomalies in the tropics, *Journal of Geophysical Research: Atmospheres*, 116, 2011.

Zuidema, P.: Convective clouds over the Bay of Bengal, *Monthly Weather Review*, 131, 780–798, 2003.