

Sensitivity of Totten Glacier dynamics to sliding parameterizations and ice shelf basal melt rates

Yiliang Ma¹, Liyun Zhao¹, Rupert Gladstone², Thomas Zwinger³, Michael Wolovick⁴, JunShun Wang¹, John C. Moore²

¹State Key Laboratory of Earth Surface Processes and Hazards Risk Governance (ESPHR), Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

²Arctic Centre, University of Lapland, Rovaniemi, Finland

³CSC – IT Center for Science Ltd., Espoo, Finland

⁴Glaciology Section, Alfred Wegener Institute, Bremerhaven, Germany

Correspondence to: Liyun Zhao (zhaoliyun@bnu.edu.cn) and John C. Moore (john.moore.bnu@gmail.com)

Abstract.

Totten Glacier in East Antarctica holds a sea level potential of 3.85 m and is mostly grounded below sea level. It features very large discharge rates among the highest for East Antarctic outlet glaciers and has been losing mass over recent decades.

Recent thinning of the Totten ice shelf is likely to be due to high basal melt rates driven by increasing intrusion of warm Circumpolar Deep Water. Here we simulate the evolution of the Totten Glacier subregion using a full-Stokes model with different basal sliding parameterizations (linear Weertman, nonlinear Weertman, and regularised Coulomb) as well as sub-shelf melt rates to quantify their effect on the projections. The modelled grounding line retreat and decline in ice volume above floatation using the linear Weertman and the regularised Coulomb sliding parameterizations are close, and both larger than that using the nonlinear Weertman sliding parameterization. The simulated grounding line retreats mainly on the eastern and southern grounding zone of Totten Glacier. The change of sub-shelf cavity thickness is dominated by sub-shelf melt rates, yielding strong volume above floatation dependence on melting through the mechanism of reduced buttressing.

1. Introduction

Totten Glacier (TG), located in the Aurora Basin, contains 3.85 m of sea-level change equivalent, had an annual ice discharge of 71.4 ± 2.6 Gt yr^{-1} from the year 2009 to 2017 which is the third highest ice discharge from the East Antarctic Ice Sheet (EAIS), and shows on-going dynamic change (Rignot et al., 2019; Greenbaum et al., 2015). TG speed slowed down from 1989 to 2001, then accelerated and stayed at relatively constant speed over the 2005-2015 period (Li et al., 2016). The rate of TG mass loss increased by 28% to 7.3 Gt yr^{-1} between 2003-2017 relative to its 1979-2003 rate (Rignot et al., 2019). Bedrock elevations in the TG subbasin are mostly below sea level, with the deepest region nearly 2000 m below sea level (Morlighem et al., 2020). The grounding line retreated by 1-3 km during the period 1996-2013, and the ice at the grounding line thinned by 0.7 ± 0.1 m (Li et al., 2015). A further grounding line retreat of 0.5-2 km along the ice plains beneath the

glacier central trunk occurred from 2013 to 2021 (Li et al., 2023). The mass imbalance of TG and the existence of deepening topography extending far inland have raised concerns that this subbasin has the threat of notable future grounded ice loss caused by marine ice sheet instability if the grounding line were to continue retreating (Schoof, 2012; Li et al., 2016; Morlighem et al., 2020).

Warm modified Circumpolar Deep Water (mCDW) invades the Totten ice shelf cavity through a deep channel incised in the sea floor at a rate of 0.22 ± 0.07 Sv (Rintoul et al., 2016; Silvano et al., 2017). This inflow of heat induces rapid basal melting, that may thin the ice shelf, remove grounding on sea floor topographic high points and hence reduce the ice shelf buttressing force on the grounded ice. Observations suggest that episodic grounding line retreats are coincident with high thinning rates and high ice velocities (Li et al., 2023), and that TG may be very sensitive to deep ocean temperature (Li et al., 2016).

The future evolution of TG has been studied with various numerical ice sheet models employing different relationships between mass loss and ocean thermal forcing. Some studies used sub-shelf melt rate parameterizations (Sun et al., 2016; Pelle et al., 2021), and some used coupled ice-ocean modelling (McCormack et al., 2021; Pelle et al., 2021). These studies project that TG will experience continuous and notable retreat on its eastern sectors by 10 km at 2200 under the A1B scenario (Sun et al., 2016) and 10 km by 2100 under the SSP585 scenario (Pelle et al., 2021). Upstream of the grounding line, ice flow is mostly controlled by basal sliding and vertical shear stress. In contrast with the many studies on the impact of sub-shelf melt rates on TG dynamics, there has not been an investigation of TG dynamics sensitivity to basal sliding parameterization.

Several types of basal sliding parameterization have been proposed and improved over the past 70 years (Weertman 1957; Nye 1970; Kamb, 1970; Fowler, 1981; Budd et al., 1979; Schoof, 2005; Gagliardini et al., 2007; Tsai and others, 2015). The Weertman sliding parameterization is formulated assuming that temperate ice slides perfectly on a hard bed, and the effective pressure (the difference between ice overburden pressure and basal water pressure) is sufficiently high that no cavities form as the ice flows over the bed, so the basal drag does not depend on effective pressure. The linear relation between basal friction and basal sliding velocity is only valid when regelation dominates or under the simplifying assumption that ice deforms like a Newtonian viscous material (Nye 1970; Kamb, 1970; Cuffey and Paterson, 2010). But this linear Weertman sliding is used very widely in ice sheet dynamic simulations mainly due to its simplicity (e.g. Sun et al., 2016; Nowicki et al., 2016; Pelle et al., 2020). If the obstacle size is not large enough to induce regelation processes, sliding is dominated by ice-deformation on the largest occurring roughness features, and a nonlinear Weertman sliding parametrization with an exponent of $1/3$ applies (Fowler, 1981). In the case of fast sliding on a hard bed, cavity formation increases with sliding rate, the basal drag attains a constant value or even decreases as sliding rate increases, showing a strong dependence on effective pressure (Cuffey and Paterson, 2010). Effective pressure has been included in several basal

sliding parameterizations (Budd et al., 1979, 1984; Schoof, 2005; Tsai and others, 2015; Gagliardini et al., 2007). In the Budd sliding parameterization (Budd et al., 1979, 1984), basal drag is proportional to both basal sliding velocity and effective pressure, and can reach arbitrarily high values. In the Coulomb sliding parameterization, basal drag is only proportional to effective pressure, and thus has upper limit; this Coulomb plastic rheology is able to describe sliding over tills (Iverson and others, 1998). Tsai et al. (2015) proposed a sliding parameterization which accommodates the Weertman type friction and the Coulomb friction regimes, and the basal drag instantaneously switches from Coulomb friction at low effective pressure to Weertman type friction at high effective pressure. Schoof (2005) and Gagliardini et al. (2007) developed a regularized Coulomb sliding parameterization, which naturally has a continuous transition between Coulomb friction regime and Weertman type friction regime. It has been suggested that regularized Coulomb sliding parameterization is better suited for fast flowing areas such as Pine Island Glacier in the Amundsen Sea Embayment (Joughin et al., 2019).

The choice of basal sliding parameterizations used by models can have an important impact on the simulated grounding line retreat rate and glacier mass loss (Gladstone et al., 2014, 2017; Brondex et al., 2017, 2019). Brondex et al. (2017) found that simulations on an idealised two-dimensional ice sheet geometry with different basal sliding parameterizations (the nonlinear Weertman with exponent of 1/3, Schoof, and Budd sliding parameterization) show notably different transient behaviour although they start from identical initial states, and the nonlinear Weertman sliding parameterization forecasts the least volume above floatation (VAF) loss and Budd sliding parameterization the most. Yu et al. (2018) showed that the use of the Budd sliding parameterization produces more grounding line retreat and more VAF loss than with the linear Weertman sliding parameterization with sub-shelf melt rate linearly dependent of ice shelf basal elevation and the maximum basal melt rate is larger than 80 m a⁻¹ in simulations of Thwaites Glacier. Brondex et al. (2019) simulated the evolution of Amundsen Sea Embayment for 100 years and found that grounding line retreat and VAF loss with the linear and nonlinear (with exponent of 1/3) Weertman sliding parameterizations are less than that with a regularised Coulomb sliding parameterization, which are less than that with Budd sliding parameterization. They also found that simulations using the non-linear versions (with exponent of 1/3) of the Budd and Weertman sliding parameterizations produce more VAF losses than those using the linear versions although they lead to similar or less grounded ice area loss.

In this study, we conduct a series of simulations to assess the dynamic response of TG under different basal sliding parameterizations from the year 2015 to 2100 and sub-shelf melt rates from the year 2015 to 2050. We use the full-Stokes ice-flow model, solved by Elmer/Ice (Gagliardini et al., 2013). Full-Stokes models consider all the components of the 3D deviatoric stress tensor. Furthermore, full-Stokes models have been suggested as essential to infer a correct pattern of basal drag and to capture all higher order stresses near the grounding line of ice streams (Morlighem et al., 2010). Therefore, we expect to gain more insight into basal processes using a full-Stokes model. We specify three types of basal sliding parameterizations (linear and nonlinear Weertman, and regularised Coulomb) and a range of sub-shelf melt rates with the same parameterization but different maximal basal melt rate of 0, 20, 40, 80 and 160 m yr⁻¹. We compare the results from the

different simulations and investigate the impact of different basal sliding parameterizations and sub-shelf melt rates on TG evolution.

2. Data, model domain and mesh refinement

2.1 Data

The Totten subregion is part of the Aurora basin of the EAIS. We take its 2D domain boundary outline and ice front location from the MEaSUREs Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version 2 dataset (Mouginot et al., 2017). The bed elevation, surface elevation (Fig. 1) and ice thickness are derived from MEaSUREs BedMachine Antarctica, version 2, at a resolution of 500 m (Morlighem et al., 2020). The surface ice velocity data (Fig. 1), with a resolution of 450 m, are obtained from MEaSUREs InSAR-based Antarctic ice velocity Map, version 2 (Rignot et al., 2017).




Figure 1. Bed elevation (a), surface elevation (b), surface ice flow speed (c), modelled basal temperature (g) and ice thickness (i) of Totten subregion and its surroundings. The solid black curve is the outline of Totten subregion, from MEaSUREs Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version 2 dataset (Mouginot et al., 2017). The inland black curve is the grounding line. The bed and surface elevations are from MEaSUREs BedMachine Antarctica, version 2 (Morlighem et al., 2020). The surface ice flow speed is from MEaSUREs InSAR-based Antarctic ice velocity Map, version 2 (Rignot et al., 2017). Subglacial lake Vostok and Law Dome are marked in plot (a). The inset plot in plot (b) shows drainage basin divisions and the location of our domain (red curve) in Antarctica. The prescribed ice temperatures are taken from the output of the ice sheet model SICOPOLIS (Greve et al., 2020). (d)-(f) show the enlarged coastal region of (a)-(c), and (h) shows the enlarged coastal region of (g).

Ice sheet temperature data, at 8 km resolution (Fig. 1d), are sourced from the output files of the ice sheet model SICOPOLIS (Greve et al., 2020), which is initialized by a paleoclimatic spin-up over 140 000 years until 1990, when the topography is nudged towards that of the present-day (Greve et al., 2020). The geothermal heat flux applied in their study is from Martos et al. (2017), at a resolution of 15 km. The surface mass balance (SMB) data applied for the period 2015-2100 consists of repeated 1995-2030 time-varying output of the Community Earth System Model Version 2 (CESM2) of the Coupled Model Intercomparison Project Phase 6 (CMIP6) under the SSP5-8.5 scenario at 32 km resolution (Nowicki et al., 2016; Fig. S1), which is close to the RACMO2.4p1 SMB data (Van Dalum et al., 2024). The ocean forcing is simplified to a parameterization scheme based on the Whole Antarctic Ocean Model (WAOM v1.0) output for modelling the basal melt rate of the ice shelves (see Section 3.3). See Table 1 for the dataset details.


Table 1. Datasets used in this study.

Variable name	Dataset	Resolution	Period	Reference
Subbasin boundary	MEaSUREs Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version 2	25-250 m	1992-2015	Mouginot et al., 2017
Surface elevation, bedrock elevation, and ice thickness	MEaSUREs BedMachine Antarctica version 2	500 m	1970-2019	Morlighem et al., 2020
Surface ice velocity	MEaSUREs InSAR-based Antarctic ice velocity Map, version 2	450 m	1996-2016	Rignot et al., 2017
Ice temperature	Output of SICOPOLIS	8 km	2015	Greve et al., 2020
SMB	CESM2 output under SSP5-8.5	32 km	2015-2100	Nowicki et al., 2016
ice shelf basal melt rate	Whole Antarctic Ocean Model (WAOM v1.0) output	2 km	2007	Richter et al., 2022

2.2 Mesh generation and refinement

We use Gmsh software (Geuzaine and Remacle, 2009) to generate an initial 2D horizontal footprint mesh within the domain boundary. The initial grid was then optimised by the anisotropic mesh adaptation code Mmg (Dapogny et al., 2014), which

135 helps to refine the mesh where ice velocity and thickness gradients identify regions needing increased resolution. The resulting mesh is shown in Fig. 2 and has minimum and maximum element sizes of approximately 900 m and 20 km. The 2D mesh contains 31 434 elements and 15 996 nodes. It is then vertically extruded using 20 terrain-following layers, with a vertically refined mesh toward the base.



140 **Figure 2.** The refined 2D horizontal domain footprint mesh (a). Boxes outlined in panel (a) are shown in detail and overlain with surface ice velocity in panel (b). The solid black curves in (b) represent the positions of the grounding line.

3. Method

3.1 Ice Flow Model

We carried out all the simulations using the ice flow model Elmer/Ice version 9.0 (Gagliardini et al., 2013). We solve the Stokes equation and use an variational inverse method (Morlighem et al., 2010) as implemented in Elmer/Ice (Gagliardini et al., 2013; Gillet-Chaulet et al., 2012) to determine the basal friction coefficient in the linear Weertman sliding parameterization and a stress enhancement factor. The Stokes equation includes conservation equations for the momentum and mass:

$$\nabla \cdot \boldsymbol{\tau} - \nabla p = \rho_i \mathbf{g}, \quad (1)$$

150 $\nabla \cdot \mathbf{v} = 0, \quad (2)$

where $\boldsymbol{\tau}$ is the deviatoric stress tensor, p is the isotropic pressure, ρ_i is the ice density, $\mathbf{g}$ is the acceleration due to gravity (0, 0, -9.81) m s^{-2} , and $\mathbf{v}$ is ice velocity. The effective viscosity, η , or the ice-rheology,

$$\boldsymbol{\tau} = 2\eta \dot{\boldsymbol{\epsilon}}, \quad (3)$$

given by Glen's flow law (Cuffey and Paterson, 2010),

$$155 \quad \eta = \frac{1}{2} (E \cdot A(T))^{-\frac{1}{n}} \dot{\epsilon}_e^{\frac{1-n}{n}} \quad (4)$$

is dependent on ice temperature via the flow rate factor $A(T)$, defined by an Arrhenius equation. We use an exponent of $n=3$. $\dot{\epsilon}$ denotes strain rate and the term $\dot{\epsilon}_e = \sqrt{\text{tr}(\dot{\epsilon}^2)}/2$ is the effective strain rate. The ice temperature distribution comes from the output of ice sheet model SICOPOLIS (Greve et al., 2020). We introduce $E_\eta = E^{-1/(2n)}$, to obtain

$$\eta = \frac{1}{2} E_\eta^2 \cdot A(T)^{-\frac{1}{n}} \dot{\epsilon}_e^{\frac{1-n}{n}} \quad (5)$$

160 which ensures $\eta > 0$ for any value of E_η . We refer E_η to as the 2D vertically constant stress enhancement factor which is set initially to unity everywhere, indicating no stress enhancement, before performing the viscosity inversion. The enhancement factor is often used to account for anisotropy effects (Ma et al., 2010). Values of $E_\eta < 1$ indicate softer and faster deforming ice, whereas higher values, $E_\eta > 1$, mean stiffer ice.

165 The ice surface is assumed to have a vanishing Cauchy-stress vector. At the ice front under the sea level and the bottom of ice shelf, the normal Cauchy-stress is set equal to the hydrostatic water pressure. The so-called sea spring (Gagliardini et al., 2013) stabilization method is applied to the bottom free surface in the case of a temporal evolution away from the hydrostatic equilibrium. On the lateral boundary, the normal stress is equal to the hydrostatic ice pressure and the normal velocity is set to zero, the ice is free to slip in the tangential direction. The bed for grounded ice is assumed to be rigid, impenetrable, and
 170 fixed over time since the ice sheet geometry change over decades is too small to affect lithosphere deformation (de Boer et al., 2014; Coulon et al., 2021). The normal component of the velocity is set to zero at the ice–bed interface. We implement alternative basal sliding relations (Sect. 3.2) for the grounded ice in Elmer/Ice. For transient simulations, we prescribe SMB at the ice surface (Sect. 2.1), and sub-shelf basal melt rate parameterization along the surface in contact with the ocean (Sect. 3.3).

175

3.2 Basal sliding parameterizations and conversion between them

We use three basal sliding parameterizations in this study: linear Weertman sliding (Eq. (6)), nonlinear Weertman sliding (Eq. (7); Weertman, 1957) and regularised Coulomb sliding parameterization (Eq. (8-9); Schoof, 2005; Gagliardini et al., 2007) given by

$$180 \quad \tau_b = C_{\text{LW}} \mathbf{u}_b, \quad (6)$$

$$\tau_b = C_{\text{NW}} \mathbf{u}_b^{1/m}, \quad (7)$$

$$\tau_b = C_s N \left[\frac{\chi \cdot u_b^{-m}}{1 + a \cdot \chi^q} \right]^{\frac{1}{m}} u_b, \quad (8)$$

$$\chi = \frac{u_b}{C_s^m N^m A_s}, \quad (9)$$

where τ_b is the basal shear stress, $\mathbf{u}_b$ is the vector of basal sliding velocity, u_b is basal sliding speed, C_{LW} and C_{NW} are
 185 friction coefficients of linear and nonlinear Weertman sliding parameterization. We set $m=3$. The regularised Coulomb
 sliding parameterization considers the existence of water cavities which may be present between ice and soft sediment layers.
 The parameter A_s is the sliding parameter in the absence of cavitation and C_s is the maximum value reached by
 τ_b / N (Gagliardini et al., 2007). In Eq. (8), q is a post-peak exponent satisfying $q \geq 1$; $a = (q-1)^{q-1} / q^q$. We assume $q=1$,
 obtaining $a=1$. N represents the effective pressure on the ice bottom and is defined as $N = \sigma_{nn} - p_w$, where σ_{nn} is the
 190 normal stress at the bed, and p_w is the water pressure. The parameters and their units are listed in Table 2.

Lower effective pressure reduces the basal friction and enhances the basal sliding speed. This effect has been included in
 several sliding parameterizations (Budd et al., 1984; Schoof, 2005; Tsai et al., 2015). Hydrology models determining the
 effective pressure distribution rely on a set of poorly constrained parameters such as sheet and channel conductivities and
 195 basal melt water input that are hard to measure (Werder et al., 2013). Therefore, many authors take effective pressure as the
 hydrostatic pressure imposed by an assumed perfect connection between the subglacial drainage system and the ocean
 (Morlighem et al., 2010; Tsai et al., 2015; Gladstone et al., 2017). For TG, Dow et al. (2020) simulated the basal hydrology
 using GlaDS subglacial hydrology model to best match the specularly content, and found the modelled water pressure is
 $\sim 95\%$ of overburden in most of the region. Motivated by their findings, in this study, we take the effective pressure as 5% of
 200 overburden. This assumption decreases the effective pressure by an order of magnitude in most inland regions and halves it
 near the GL, compared with that under the assumption of perfect hydrostatic connection (Fig. S2).

We apply the basal shear stress and basal sliding velocity field obtained from the inversion with the linear Weertman sliding
 parameterization to the other two sliding parameterizations, and estimate the sliding parameters there. In other words, we
 205 convert the local friction coefficient in linear Weertman sliding parameterization to that in other sliding parameterizations
 (Brondex et al., 2017). In Eq. (6), we set $C_{LW} = 10^\beta$ to make sure C_{LW} is positive. The conversion from C_{LW} to C_{NW} then is
 trivially given by $C_{NW} = 10^\beta u_b^{1-1/m}$. To convert the friction coefficient from a linear Weertman to a regularised Coulomb
 sliding parameterization, as given in Eq. (8), we rearrange the latter

$$\left(\frac{\tau_b}{C_s N}\right)^m = \frac{\chi}{1+\chi}, \quad (10)$$

210 The limit forms of Eq. (10) with small $\chi \rightarrow 0$ or large $\chi > 10$ correspond to slow-flowing regimes with large effective pressure and fast-flowing regimes with small effective pressure, respectively.

In the limit of small χ , we recover a non-linear Weertman law

$$\tau_b = A_s^{-1/m} u_b^{1/m}. \quad (11)$$

Combining Eq. (11) and Eq. (6) with $C_{LW} = 10^\beta$, we get

$$215 \quad A_s = (10^\beta \cdot u_b^{1-1/m})^{-m}. \quad (12)$$

In the limit of Eq. (10) for large χ (i.e., N is relatively small), we obtain a Coulomb-type of friction

$$\tau_b = C_s N. \quad (13)$$

We modify Eq. (12) by multiplying a smooth transition term from small N region to large N region as below, such that it can
220 be applied to the whole region,

$$A_s = \tanh(2N) \cdot (10^\beta \cdot u_b^{1-1/m})^{-m}. \quad (14)$$

Combining Eq. (6), (8) and (9) with $C_{LW} = 10^\beta$, then using Eq. (14), we can get C_s in regularised Coulomb sliding parameterization as

$$C_s = \frac{10^\beta \cdot u_b}{N \cdot (1 - 10^{m\beta} \cdot A_s \cdot u_b^{m-1})^{1/m}} = \frac{10^\beta \cdot u_b}{N \cdot (1 - \tanh(2N))^{1/m}}. \quad (15)$$

225 Eq. (14) and Eq. (15) determine the conversion of the inverted friction coefficients to the regularised Coulomb sliding parameterization. We keep the converted friction coefficients A_s and C_s fixed in all the prognostic simulations with regularised Coulomb sliding parameterization.

Eq. (10) is equivalent to Coulomb-type friction with a factor:

$$230 \quad \tau_b = \left(\frac{\chi}{1+\chi}\right)^{1/m} C_s N. \quad (16)$$

This factor $\left(\frac{\chi}{1+\chi}\right)^{1/m}$ is about 0.6 if $\chi=0.275$, about 0.8 if $\chi=1$, and about 0.97 if $\chi=10$.

Table 2. List of parameters used in Section 3.2 and their units.

Parameter	Description	Unit
C_{LW}	friction coefficients of linear Weertman	$\text{m}^{-1} \text{ yr Pa}$
C_{NW}	friction coefficients of nonlinear Weertman	$\text{m}^{-1/3} \text{ yr}^{1/3} \text{ Pa}$
C_s	the maximum value reached by τ_b / N	No unit
A_s	the sliding parameter in the absence of cavitation	$\text{m yr}^{-1} \text{ Pa}^{-3}$
q	friction law exponent, $q \geq 1$, we take $q=1$	No unit

235

3.3 Ice-shelf basal melt parameterization

Sub-shelf melting of TG is influenced by ocean heat intrusions imposed by mCDW, which may be modified by several processes such as the Antarctic Slope Current and upstream coastal freshening (Nakayama et al., 2021). But we do not consider the complex ocean circulation here. Instead, we use a simplified parameterization to mimic the spatial pattern and magnitude inferred from satellites (Rignot et al., 2013; Depoorter et al., 2013; Liu et al., 2015; Adusumilli et al., 2020) and ocean model simulations (Richter et al., 2022). We consider sub-shelf melt as a sum of low melt which plays a dominant role in the shallow water (<500 m) and high melt which plays a dominant role in the deep water (>1000 m) near the grounding line. We express mass balance in ice equivalent units.

245 We parameterise the sub-shelf melt rate in the shallow water using ice shelf bottom depth and ice shelf basal melting rate as modelled by the 2 km spatial resolution, tide and eddy-resolving Whole Antarctic Ocean Model (WAOM v1.0; Richter et al., 2022). We did linear regression between sub-shelf melt rate and $1/(d+d_0)$, where d is ice shelf bottom depth (unit: m), and d_0 is a tuned parameter. We found the best fit as follows with $d_0=100$ m (Fig. S3):

$$M_d = 450.284 \times \frac{1}{d + d_0} + 1.1917, \quad (17)$$

250 where M_d represents the basal melting rate of the ice shelf (unit: m yr^{-1} i.e.).

The WAOM v1.0 estimates of Antarctic ice shelf basal mass loss agree with satellite observations in many regions (Richter et al., 2022). However, for the TG subregion, the satellite estimates (Rignot et al., 2013; Depoorter et al., 2013; Liu et al., 2015; Adusumilli et al., 2020) suggest higher basal melting (area average melt $\geq 10 \text{ m yr}^{-1}$) than that obtained with WAOM v1.0 (Richter et al., 2022). To account for the high melt rate in the deep water, we add an extra melt rate term as

$$M_e = \begin{cases} M_{\max} \times \left(\frac{d}{2000} \right), & d < 2000 \text{ m}; \\ M_{\max}, & d \geq 2000 \text{ m}. \end{cases} \quad (18)$$

where $M_{\max}$ is the maximum basal melt rate, and d is the depth of the bottom surface of the ice shelf (unit: m).

In addition, we use two scaling factors S_w and S_i to reflect the influence of cavity geometry and avoid numerical
 260 instability caused by a vanishingly thin ice shelf, following Gladstone et al. (2017).

$$S_w = \tanh \left(e \frac{H_w}{H_{w0}} \right), \quad (19)$$

where H_w is water column thickness, given by $H_w = z_i - b$ where z_i is the ice shelf bottom elevation and b is the bed
 elevation. We set $H_{w0} = 200$ m as a reference water column thickness. S_w approaches 1 with thick water column
 ($H_w > H_{w0}$). And S_w approaches 0 when the water column thickness goes to zero near the grounding line, capturing the
 265 influence of cavity geometry on melt rate as a very thin water column restricts circulation. A high step change in forcing
 across the grounding line (basal drag or ocean induced melt) causes problems for stability (Gladstone et al., 2017). S_w
 prevents high melting right next to the grounding line, so reduces that step change.

A vanishingly thin ice shelf causes the tetrahedral elements to have a very high horizontal to vertical length scale, which
 270 increases the likelihood of instability. S_i is an ice-shelf depth-scaling parameter, used to prevent melting the ice shelf once
 the draft passes $|z_{i0}|$ given by

$$S_i = \max \left[\tanh \left(e \frac{z_{i0} - z_i}{z_s} \right), 0 \right], \quad (20)$$

where we set $z_{i0} = -80$ m as a reference ice base elevation relative to sea level (positive upwards) and $z_s = 100$ m as a
 (directionless) scaling depth. The use of S_i gives zero melting for $z_i > z_{i0}$.

275 In a combination of Eq. (17) – (20), the sub-shelf melt rate then is parameterised as

$$M = S_w \cdot S_i (M_d + M_e). \quad (21)$$

The parameterizations of M_d , M_e with $M_{max} = 40$ m yr⁻¹ and the total sub-shelf melt with water column thickness along
 a flowline are illustrated in Fig. S3. The spatial distribution of estimated sub-shelf melt rates using different values of M_{max}
 280 (0, 20, 40, 80 and 160 m yr⁻¹) are shown in Fig. 3. The modelled sub-shelf melt rates, with varying maximal values, cover
 most of the estimates from previous studies.

Flux gate calculations using satellite data have reported steady-state area-averaged basal melt rates of 10.5 ± 0.7 m yr⁻¹
 (2003-2008; Rignot et al., 2013), 9.89 ± 1.92 m yr⁻¹ (2003-2009; Depoorter et al., 2013) and 11.5 ± 2.0 m yr⁻¹ (2010-2018;
 285 Adusumilli et al., 2020). Notably, Liu et al. (2015) used a similar method without assuming steady-state calving front, and
 derived a higher melt rate of 17.9 ± 1.2 m yr⁻¹ during the period 2003–2011. This elevated value aligns better with Rintoul et

al.'s (2016) synoptic cavity water exchange estimates. In situ measurements by Vaňková et al. (2023) using autonomous phase-sensitive radar along grounding lines revealed large spatial melt variability, correlating with water column thickness gradients. Our model simulations for sub-shelf melt rate with the maximum value of 20, 40, 80 and 160 m yr⁻¹ (Fig. 3) yield area-averaged melt rates of 7.8, 14.0, 26.6 and 51.6 m yr⁻¹ respectively for the present day ice shelf geometry .




Figure 3. Spatial distribution of modelled sub-shelf melt rate with the maximum value of (a) 20 m yr⁻¹, (b) 40 m yr⁻¹, (c) 80 m yr⁻¹, (e) 160 m yr⁻¹ with coloured contours indicating water column thickness under the ice shelf with 100 m intervals. Plot (d) shows the satellite-based estimates (Adusumilli et al., 2020).

3.4 Experiment design

The following simulations are carried out in series as part of the initialization procedure. We firstly do a steady state simulation with $E_\eta = 1$, and the linear Weertman sliding parameterization, expressing the sliding coefficient by $C_{\text{LW}} = 10^\beta$, where β is from the output of a whole Antarctic ice sheet inversion for the year 2015 (Gladstone et al., 2019), and linearly interpolated from their mesh to the finer mesh in this study. Then we relax the free surface of the domain by a short transient run of 1 year with a small time-step size of 0.1 year without any sub-shelf melt rate to reduce the non-physical spikes in the initial surface geometry (Zhao et al., 2018; Wang et al., 2020). Taking the results from surface relaxation, we use the

variational inverse method (Morlighem et al., 2010) to adjust the spatial distribution of basal friction coefficient and stress enhancement factor to minimize the mismatch between the magnitudes of the simulated and observed (Fig. 1c; Table 1) surface velocities, following the same approach as in Gladstone and Wang (2022) and Wang et al. (2020). The inversions make use of Tikhonov regularization (Gillet-Chaulet et al., 2012). Two separate regularization parameters, λ_β and λ_η , are used for the drag inversions and stress enhancement factor inversions. The total cost function, J_{tot} , is the sum of misfit, J_0 , and weighted regularization term for the drag inversion:

$$J_{tot} = J_0 + \lambda_\beta J_{reg}.$$

λ_β would be replaced by λ_η for the viscosity enhancement factor inversion. We take $\lambda_\beta = 10^3$ and $\lambda_\eta = 10^4$. We firstly obtain the optimal spatial field of β , an updated modelled ice velocity field, and the stress field. We then perform a viscosity inversion to obtain the optimal spatial field of E_η , and further updated modelled ice velocity and stress fields. We take this state as the one for the initial year 2015.

With the modelled basal shear stress and basal sliding velocity, we convert the basal drag coefficient from the linear Weertman parameterization to those representing non-linear Weertman and regularised Coulomb parameterizations using the method described in Section 3.2. Using the three different sliding parameterizations and different sub-shelf melt rates, we carry out two sets of sensitivity tests (Table 3). We run simulations in the sliding parameterization group from 2015 to 2100, and melt rate group from 2015 to 2050, all with a time-step size of 0.1 year. A so-called ‘reference run’ is chosen to illustrate more dynamic process during the 85 years simulation. We chose regularised Coulomb parameterization as the basal sliding in the reference run because it is arguably the most physically sound (Tsai et al., 2015; Joughin et al., 2019). Given the expectation of a warming climate, we chose for our reference run a melt rate parameterization with a maximum melt that approximates to an estimate for the temporal maximum over recent decades (i.e. 80 m yr⁻¹; Adusumilli et al., 2020; Nakayama et al., 2021). We use maximum sub-shelf melt rates of 0, 20, 40, 80 and 160 m yr⁻¹ (Fig. 3) in the sensitivity tests.

We did simulations of melt rate group to the year 2050, because solving the Stokes equation for Totten glacier subregion is computationally expensive and the sensitivity of Totten glacier dynamics to sub-shelf melt rate has been shown from 2015 to 2050. Since the modelled sub-shelf melt rate with the maximum value of 40 m yr⁻¹ is closest to the satellite-based estimates (Section 3.3), the experiment C_m40 (which belongs to the melt rate group) was selected as the control experiment to evaluate the model drift, and was done from 2015 to 2100. In the drift control experiment C_m40, the modelled annual mean thinning rate from 2015 to 2020 is -0.03 m yr⁻¹, larger than the observed thinning rate of -0.02 m yr⁻¹ (Nilsson et al., 2023), indicating higher model sensitivity to the present day climate forcing.

Table 3. Summary of sensitivity experiments.

Experiment group	Experiment	Sliding parameterization	Maximum melt rate (m yr ⁻¹)	Simulation period
Reference run	C_m80	regularised Coulomb	80	2015-2100
Sliding parameterization group	LW_m80	linear Weertman	80	2015-2100
	NW_m80	nonlinear Weertman	80	2015-2100
Melt rate group	C_m160	regularised Coulomb	160	2015-2050
	C_m40 (drift control)	regularised Coulomb	40	2015-2100
	C_m20	regularised Coulomb	20	2015-2050
	C_m0	regularised Coulomb	0	2015-2050

4. Results

4.1 Initial state and reference run result

We obtain the inverted optimal spatial distribution of basal sliding coefficient exponent β (Fig. 4a) in the linear Weertman parameterization and the stress enhancement factor (Fig. 4b) for the initial year 2015, and keep them fixed in all the prognostic simulations. The basal sliding coefficient exponent β is smaller in the fast flow regions where ice speed >10 m yr⁻¹, and larger in the slow ice flow region. The inversion results show a rib-like pattern of basal sliding coefficient and basal shear stress in fast-flowing region (Fig. 4e, g). But it does not directly give us the cause. Ripples, oscillations, or ribs have been found in other inverse models of both Greenland and Antarctica (Sergienko and Hindmarsh, 2013; Sergienko et al., 2014; Wolovick et al., 2023). E_η remains at 1 over most of the grounded region; $E_\eta > 1$ in the vicinity of the grounding line indicates relatively stiffer ice, and $E_\eta < 1$ on the ice shelf indicating more rapid deformation likely due to shear-band damage.

The converted friction coefficients A_s and C_s in the regularised Coulomb sliding parameterization are also kept fixed in the prognostic simulations. The value of χ (Eq. (9)) in the initial year is from 0.3 to 0.5 near the grounding line with spatial variability, mainly on the eastern branch, which means that the basal friction in the fast-flowing region is 60%-70% of that in a true Coulomb regime (Eq. (16)). As the effective pressure is reduced as the ice thins, χ increases and varies from 0.3 to 5 near the grounding line in the year 2100, which is closer to a true regularised Coulomb sliding parameterization (Fig. S4).

For the initial year 2015, the basal shear stress (Fig. 4c) for the initial state is concentrated in the range of 4-400 kPa. The basal effective viscosity (Fig. 4d) is smaller in regions with relatively high ice temperature (Fig. 1) and fast flow speed (>10 m yr⁻¹), and larger in the upstream regions with low ice temperature and slow flow speed. Shear margins of lower viscosity exist on both sides of the TIS.




Figure 4. Spatial distribution of modelled (a) basal sliding coefficient β , (b) stress enhancement factor E_η , (c) basal shear stress τ_b , (d) basal effective viscosity, and their corresponding enlarged ones (e-h). The solid yellow (a-c; e-g) or cyan (d, h) curves represent the grounding line and the solid white curves in (a-d) show modelled basal speed contours of 10 m yr⁻¹.

In the reference run (C_m80), ice thickness upstream of the grounding line decreases notably, and the ice velocity speeds up upstream of the grounding line but decelerates near the ice front of the ice shelf over the 85-year simulation (Fig. 5). But there are large spatial differences in ice thickness change of the grounded ice. The fast-flowing regime in the grounded ice becomes larger and the ice shows dynamic thinning at the eastern and southern grounding line (Fig. 5). The western side of TIS is grounded on Law Dome (Fig. 1). The grounding line retreats primarily along the southern and eastern grounding zones, which is consistent with previous studies (e.g. Pelle et al., 2021). We select two flowlines for analysis: FL1 along the main trunk, and FL2 on the eastern branch (Fig. 5) of TG. Modelled annual averaged grounded ice thinning rate from 2015 to 2100 measured from the 2100 grounding line position and extending 20 km further upstream is 2.14~2.35 m yr⁻¹ along FL1 and 2.88~5.06 m yr⁻¹ along FL2. The modelled grounding line retreats 12.20 km along FL1, and 23.29 km along FL2 by the year 2100. The basal topography along FL1 is prograde sloping bed with large slope upstream of grounding line, but the

basal topography along FL2 has more variations in slope and the slopes are smaller than that along FL1. Hence the ice is more stable along FL1 than along FL2.




Figure 5. Surface velocity at the beginning of the initial year 2015 (a) and the end of the year 2100 (b) in the reference run. The surface velocity difference (2100 minus 2015) is shown in (c). The grounding line positions in the year 2015, 2040, 2060, 2080 and 2100 in the reference run C_m80 are shown in (a-c). The grounding line position in the year 2100 in the drift control run C_m40 is shown in (b-g) with black lines. Pink and purple solid lines in (a) and (b) represent flowlines FL1 and FL2 as labelled. The solid color portions of the figures show the ice flow velocity (upper colorbar) profiles along FL1 (c, d) and FL2 (e, f) in the reference run in the initial year 2015 (d, f) and the end year 2050 (e, g), with bedrock in dark grey and seawater in blue. The geometry change of TG is marked with colored solid lines for the years 2015, 2040, 2060, 2080 and 2100. The vertical elevations are exaggerated by a factor of 25.

4.2 Sensitivity experiments results

The grounding line at the year 2050 and 2100 from the sliding parameterization and at the year 2050 from the melt rate groups are shown in Fig. 6. The grounding line mainly retreats in the southern and eastern grounding zones, and grounding line evolution is sensitive to both sliding parameterizations and sub-shelf melt rates (Fig. 6). The modelled grounding line

390 retreats most using the linear Weertman sliding parameterization, slightly less distance using the regularised Coulomb sliding parameterization, and least using the nonlinear Weertman sliding parameterization (Fig. 6). The grounding line retreats more using the regularised Coulomb sliding parameterization than with the nonlinear Weertman sliding parameterization, which might be because the effective pressure used in the regularised Coulomb is reduced as the ice thins, leading to more basal sliding and faster ice speed. Moreover, the value of χ (Eq. 9) is below 1 in the fast flowing region from our posterior estimate, showing a reduced basal friction and enhanced basal sliding (Eq. 16) compared with a true regularised Coulomb sliding parameterization.

395

Along the main trunk FL1 both the regularized Coulomb (C_{m80}) and linear Weertman (LW_{m80}) sliding parameterizations produce comparable grounding line retreat rates of 0.14 km yr^{-1} over 2015-2100 (Fig. 7). These rates corresponds to retreats of 8.08 km by 2050 and 12.20 km by 2100. In contrast, with nonlinear Weertman sliding, retreats are 5.94 km by 2050 along FL1 followed by then stabilization (Fig. 6). Meanwhile, the drift control experiment (C_{m40}) shows a grounding line retreat of 8.08 km along FL1 by 2100. Along FL2, the grounding line retreats 12.09 km by 2050 and 35.85 km by 2100 ($\sim 0.42 \text{ km yr}^{-1}$) using linear Weertman; 10.83 km by 2050 and 23.29 km by 2100 ($\sim 0.27 \text{ km yr}^{-1}$) using regularised Coulomb; and only 1.32 km by 2050 ($\sim 0.04 \text{ km yr}^{-1}$) followed by near-stabilization using nonlinear Weertman sliding parameterizations (Fig. 6). Meanwhile, the drift control experiment (C_{m40}) shows a grounding line retreat of 12.33 km along FL2 by 2100.

405 Between 2015 and 2100, when applying the regularised Coulomb and linear Weertman sliding parameterizations, the surface ice flow speed exhibits an acceleration upstream of the grounding line, and the ice thickness upstream of the grounding line experiences a gradual decrease. This ice flow speed acceleration and ice thinning are more pronounced in the eastern grounding zones compared to the southern ones (Fig. S5, Fig. S6). Conversely, when the nonlinear Weertman sliding parameterization is employed, the surface ice flow velocity undergoes a slight deceleration in the vicinity of the grounding line, while it experiences a slight acceleration at locations further upstream (Fig. S5). The surface lowering is much smaller using nonlinear Weertman sliding parameterization than that using the regularised Coulomb and linear Weertman sliding parameterizations. The modelled grounded ice thinning rate in the reference run (C_{m80}) near the projected 2100 grounding line is double that of the drift control run (C_{m40}). Furthermore, surface ice velocity acceleration in C_{m80} at the projected 2100 grounding line position along the main trunk FL1 exceeds that in the drift control run by a factor of three over the 2015-2100 period (Table 4). More specifically, the surface ice speed change and ice thickness change along FL1 and FL2 from experiments in the sliding parameterization group and the drift control experiment are shown in Table 4, Figure 7 and 8.

420 The oscillations in the difference between the surface and basal ice speeds near the grounding line and upstream are caused by the rib-like pattern of basal friction and basal velocity. The difference between surface and basal speeds drops to zero, which means the basal friction approaches zero in the fast flowing trunks upstream of the grounding line (Figs. 7 and 8). The

fast-flowing ice streams are supported by shear margins or isolated sticky regions through long-distance stress transmission (Wolovick et al., 2023).

Table 4. Surface ice speed change, Δv , and ice thickness change rate, Δh , from the year 2015 to 2100 (2100 minus 2015) at the projected 2100 grounding line position and 15 km further upstream, along FL1 and FL2 from the drift control experiment and experiments in the sliding parameterization group.

		Along FL1		Along FL2	
	Experiment	Grounding Line 2100	15 km Upstream	Grounding Line 2100	15 km Upstream
Δv (m yr ⁻¹)	C_m40 (drift control)	87	21	210	145
	C_m80	262	195	265	272
	LW_m80	240	165	537	200
	NW_m80	18	−32	−34	−39
Δh (m yr ⁻¹)	C_m40 (drift control)	−1.19	−1.10	−3.36	−2.79
	C_m80	−2.35	−2.14	−5.06	−2.88
	LW_m80	−2.65	−2.29	−6.42	−2.83
	NW_m80	−0.91	−0.59	−0.79	−0.70


The simulation grounded area loss rate between 2015 to 2100 from the sliding parameterization group experiments are heterogeneous in both time and space (Fig. 5, 6, 9a). The grounded area loss is 3 149 km², 2 611 km², 1 105 km² by the year 2050, and 6 582 km², 5 601 km², 1 360 km² by the year 2100 with linear Weertman (LW_m80), regularised Coulomb (C_m80) and nonlinear Weertman (NW_m80) sliding parameterizations respectively (Fig. 9a). The grounded area loss in LW_m80 and C_m80 experiments by 2100 significantly exceed the 3 413 km² from the drift control experiment. The modelled grounded area loss using nonlinear Weertman after the year 2050 is much less than that before 2050. The modelled grounded area loss rate using regularised Coulomb (C_m80) becomes slower from 2050 to 2075 while the glacier retreats over rumpled terrain, and re-accelerates after the year 2075. The modelled grounded area using linear Weertman decreases at a nearly constant rate becoming gradually slower in the last two decades.

However, the modelled VAF using the three basal sliding parameterizations decreases almost linearly over time (Fig. 9c), due mainly to the dynamic ice flow. The modelled VAF loss of the TG sub-basin from the sliding parameterization group experiments is equivalent to global sea level rise of 5.67 mm, 5.48 mm and 3.29 mm over the period 2015-2050 and 16.35 mm, 15.97 mm and 7.34 mm over the period 2015-2100 using linear Weertman, regularised Coulomb and non-linear Weertman sliding parameterizations, respectively, compared with 11.29 mm in the drift control experiment (Fig. 9c).

445 In the melt rate group of experiments (Table 3), higher basal melt rates lead to more grounding line retreat, thinner ice in both the ice shelf and the grounded ice near the grounding line (Fig. S7), smaller effective pressure, hence faster ice flow speeds, and larger basal drag (Fig. 7, 8). Ice thickness change of the ice shelf near the grounding line is highly dependent on basal melt rates. The ice becomes thicker using maximal basal melt rate smaller than 40 m yr^{-1} , and thinner when basal melt exceeds that value (Fig. S7).

450 The experiments C_{m160} , C_{m80} , C_{m40} in melt rate group (Table 3) show that the glacier grounding line retreats 11.29 km, 8.08 km and 5.94 km over the period 2015-2050 along FL1, while C_{m20} and C_{m0} do not yield any retreat. The experiments C_{m160} , C_{m80} , C_{m40} , C_{m20} , C_{m0} result in grounding line retreat of 12.33 km, 10.83 km, 7.89 km, 5.14 km and 2.83 km along FL2 by the end of the year 2050, and the grounded area is reduced by about 3405 km^2 , 2612 km^2 , 1519 km^2 , 916 km^2 and 289 km^2 , respectively (Fig. 8).

The change of sub-shelf cavity thickness is dominated by sub-shelf melt rates (Fig. 7, Fig. 8), although different basal sliding parameterizations could yield different retreat of grounding line position, hence different sub-shelf cavity thickness near the grounding line. The influence of different basal sliding parameterizations on sub-shelf cavity thickness is negligible in
 460 prograde sloping bed with large slope upstream of the grounding line such as FL1 (Fig. 7), and obvious at locations with relatively small and variable slope such as FL2 (Fig. 8). The ice speed near the grounding line is slower with nonlinear Weertman sliding parameterization than with the other two parameterizations. This causes less grounding line retreat, hence thicker ice shelf near the grounding line (Fig. 8).



465 **Figure 6.** Grounding line positions (coloured curves) at the year 2050 from experiments in the sliding parameterization group (a), at the year 2100 from experiments in the sliding parameterization group and drift control experiment (b), at the year 2050 from experiments in the melt rate group (c). The solid black line represents the initial grounding line position in the year 2015. The pink and purple solid lines are FL1 and FL2, respectively. The grey background shows the bed elevation.




Figure 7. Surface ice speed (a, b), ice speed difference (c, d; surface minus basal), basal drag (e, f), ice thickness change (g, h), and ice-sheet profiles (i, j) along FL1 in the initial year 2015 (black solid line) and the year 2100 (coloured solid line) from experiments (Table 3) in the sliding parameterization group and drift control experiment (**left column**) and the year 2050 (coloured solid line) in the melt rate group (**right column**). Grounding line positions are marked with black vertical dashed line for the year 2015 and coloured vertical dashed line for the year 2100 in the sliding parameterization group (**left column**) and the year 2050 (coloured solid line) in the melt rate group (**right column**). The figure is shaded to show the geometry of the TG along the flow line in 2015, where dark grey is the bedrock, light blue the ice shelf and dark blue seawater. The elevations are exaggerated by a factor of 25.




Figure 8. The same as Figure 7 but for FL2.

VAF change is sensitive to the choice in both basal sliding parameterizations and the sub-shelf melt rates. The modelled VAF loss of TG sub-basin from the melt rate group experiments over the period 2015-2050 is equivalent to global sea level rise of 2.8 mm, 3.58 mm, 4.22 mm, 5.48 mm and 7.47 mm, using sub-shelf melt rate parameterizations with maximum values of 0, 20, 40, 80 and 160 m yr⁻¹ (Fig. 9d).




Figure 9. Grounded area change (**a**, **b**) and VAF change (**c**, **d**) of sliding parameterization group experiments and drift control experiment (**a**, **c**) over the period 2015-2100 and melt rate group experiments (**b**, **d**) over the period 2015-2050.

5. Discussion

The effect of the exponent value, m in Eqn. (2), in the Weertman sliding parameterization has been investigated in the literature (Brondex et al., 2017, 2019; Nias et al., 2018). However, the relationship between the value of m and the rates of retreat and ice loss is unclear. Some simulations using the nonlinear Weertman sliding parameterization with $m=3$ respond faster to the changes in forcing (Sun et al., 2016), or lead to more loss of VAF (Brondex et al., 2019) than linear Weertman sliding parameterization. For Pine Island Glacier, Nias et al. (2018) show that $m=3$ can produce higher or lower changes in VAF depending on the bathymetry product being used. Barnes and Gudmundsson (2022) claimed that this relationship contains complexities and can be affected by regional differences, so there is no general statement to be made about what the general effect of increasing m will be on a system. In our study, the nonlinear Weertman sliding parameterization yields

500 generally less grounding line retreat and less ice loss than linear Weertman in the western side of TIS, and the effect of larger m has regional differences.

We ran a steady state simulation using the Blatter-Pattyn model, restarting from the steady state result of the reference run using the full-Stokes model and using the same basal sliding parameterization as in the reference run. We found that both
505 models reveal similar surface velocity patterns (Fig. S8). The ice speed simulated using Blatter-Pattyn is several hundred meters per year faster than that simulated using full-Stokes near the grounding line, but several hundred meters per year slower at the ice shelf, especially near the ice front. This is consistent with the finding in Rückamp et al. (2022), which compares the full-Stokes and Blatter-Pattyn models applied to the Northeast Greenland Ice Stream. They also found that the discrepancies between full-Stokes and Blatter-Pattyn increase with higher velocities, and are stronger when using a power-law friction than a linear friction law. The speed difference in the vicinity of the grounding line is significant because marine
510 ice sheet behaviour is largely controlled by feedbacks (involving both ocean-induced melt and basal resistance) close to the grounding line. The Blatter-Pattyn simulation predicting faster flow speed in this region would cause differences in the evolution over time of the system, hence we chose to focus on full-Stokes simulations for the current study. A comprehensive investigation into the time evolving impacts of the various approximations to the Stokes equations on
515 evolution of the Totten Glacier would be valuable, but is not the focus of the current study where we use a complete representation of stresses. Restarting from the steady state result using full-Stokes model in the reference run, the steady state simulation using Blatter-Pattyn model is 5 times faster than one timestep forward iteration using full-Stokes model with linear Weertman sliding parameterization .

520 Rib-like patterns in basal friction have been found in inversions in both Antarctica and Greenland (Sergienko and Hindmarsh, 2013; Sergienko et al., 2014; Wolovick et al., 2023). These ribs in the basal friction mimic, but do not exactly follow, similar rib-like patterns in the gravitational driving stress (Wolovick et al., 2023). Sergienko and Hindmarsh (2013) posited that basal traction ribs form from coupled instabilities in the till-water-ice system, and supported that supposition with process modeling which produced broadly similar rib-like patterns as those seen in their inverse model. The rib-like pattern may also
525 arise as compensating errors to other missing processes or uncertainties in datasets (Berends et al., 2023). However, strictly speaking, an inversion cannot tell us the definitive cause of any given structure in the inverted field, it can only inform us that the structure exists. Wolovick et al. (2023) found that rib-like structure in the basal drag was sensitive to the choice of regularization in the inversion, which should be expected for short-wavelength features. However, they also found that the ribs were present in both their base case inversion with the best optimal corner lambda value, as well as in their best
530 combined basal drag map. We did not perform a full L-curve analysis in this paper, but we did test values of 10^3 , 10^4 , 10^5 , 10^6 , 10^8 , 10^{10} , for the regularization parameter λ_β in the inversion, choosing 10^3 , which is relatively small, and hence amenable to more variable basal friction fields. If we had chosen a larger lambda value, it is likely we would have seen fewer ribs in our inverted result. However, the fact that similar ribbed structure has been seen in many different inverse

models applied to many different geographic regions - and that ribbed structure is also seen in the gravitational driving stress,
535 which does not depend on any inverse model results - suggests that at least some of this structure must be genuine.

The sub-shelf melt parameterization used here is mainly controlled by the maximum melt rate near the grounding line. We do not consider the circulation processes such as plumes that occur in the ice shelf cavity, especially the invasion process of mCDW and changes in ocean temperature, salinity, and their variability after the expansion of the ice shelf cavity. Some
540 studies obtain the distribution of sub-shelf melt rate by coupling ice sheet with ocean models. Pelle et al. (2021) used an asynchronously coupled ice-ocean model to project the evolution of TG to 2100. The maximum in their modelled sub-shelf melt rate is along the western TIS and exceed 80 m yr^{-1} under the SSP5-8.5 scenario. They also projected patent retreat on TG's eastern and southern sectors. McCormack et al. (2021) investigate Totten's response to variable ocean forcing, using the Ice-sheet and Sea-level System Model (ISSM; Larour et al., 2012) coupled to the Potsdam Ice-shelf Cavity mOdel combined
545 with a Plume model (PICOP; Lazeroms et al., 2018). They found the grounding line is steady and close to present-day only for current ocean temperatures and retreats for temperature increases $\geq 0.2^\circ\text{C}$ mainly on the eastern and southern sectors. Both ocean temperature variability and topography control grounding line stability. Our modelled grounding line retreat occurs with relatively higher basal melt rate ($\geq 40 \text{ m yr}^{-1}$), and mainly on the eastern and southern grounding zones, and is qualitatively consistent with these previous studies (Pelle et al., 2021; McCormack et al., 2021) though the magnitude and
550 spatial pattern differ from ours.

We compared the modelled surface velocity and grounding line position after 35 years of simulation using two choices of effective pressures: 1) assuming 5% of the overburden and 2) assuming perfect hydrostatic connection (Fig. S9). The surface speed differs by $\pm 400 \text{ m a}^{-1}$ mainly on the ice shelf and near the grounding line (Fig. S9). There are significant differences
555 in grounding line position after 35 years. The grounding line retreats faster assuming perfect hydrostatic connection. This grounding line position difference (Fig. S9) is similar to that between use of regularised Coulomb sliding law and linear Weertman sliding law after 35 years (Fig. 6).

Effective pressure has been included in several basal sliding parameterizations (e.g. Budd et al., 1984; Schoof, 2005; Tsai et al., 2015; Brondex et al., 2017, 2019) to represent the effect of water at the ice/bed interface. These studies with effective pressure-dependent basal sliding parameterizations usually adopt the assumption of perfect hydrological connectivity to the ocean (Morlighem et al., 2010; Tsai et al., 2015; Gladstone et al., 2017; Brondex et al., 2017, 2019). Reducing the effective pressure allows more water to flow through the basal network and enhances basal slip (Zwally et al., 2002; Stearns et al., 2008). Simulations using Weertman sliding parameterization produce less VAF loss than effective pressure-dependent laws
565 like Budd and regularised Coulomb sliding parameterization assuming a perfect hydrological connection between the subglacial drainage system and the ocean (Brondex et al., 2017, 2019). Therefore, the choice of effective pressure in basal sliding parameterizations has marked effect on the modelled ice dynamic evolution. In this study we take the effective

pressure as 5% of overburden pressure from subglacial hydrology modelling (Dow et al., 2020). The regularised Coulomb sliding parameterization yields more VAF loss than nonlinear Weertman sliding parameterization. This is consistent with previous studies (Brondex et al., 2017, 2019), despite our effective pressure values taken from a subglacial hydrology modelling is less responsive to ice thickness changes than the widely used assumption of perfect hydrological connection to the ocean. Leguy et al. (2014) proposed a function for the effective pressure, $N(p)$, depending on a parameter p , that accounts for connectivity between the subglacial drainage system and the ocean. It could produce a smooth transition between finite basal friction in the ice sheet and zero basal friction in the ice shelf, but it faces a key challenge of choosing realistic values of p for the study domain.

6. Conclusions

We convert the friction coefficient from the linear Weertman sliding parameterization to the regularised Coulomb sliding parameterization, and prescribe the two parameters used in regularised Coulomb sliding parameterization with spatial variability and smooth transitions from slow-flowing to fast-flowing regions. The defect in the method is that the resulting basal friction in the fast-flowing region is only 60% - 80% of a true regularised Coulomb sliding parameterization. We find that our simulated grounding line retreat and VAF loss of Totten glacier using regularised Coulomb sliding parameterization is less than that using linear Weertman sliding parameterization, and more than that using nonlinear Weertman sliding parameterization. This agrees with previous findings in the literature that effective pressure-dependent sliding parameterizations usually produce more VAF loss than nonlinear Weertman-type sliding parameterizations. This is despite the fact that our effective pressure parameterization is only 5% as responsive to ice thickness change as the widely used assumption of perfect hydrological connection to the ocean. There are no general statements to be made about the effect of different exponents used in Weertman sliding parameterizations, because regional bathymetry and basal velocity play large roles as well as sliding parameterization.

The modelled grounding line retreats when the maximal basal melt rates are $\geq 40 \text{ m yr}^{-1}$, and mainly along the eastern and southern grounding zones. This is consistent with previous studies, although with large differences in local retreat pattern and magnitude. The differences in pattern are mainly due to the differences in ocean thermal forcing and hence the details of basal melt rate parameterizations and model setup. More long-term ocean state observations would reduce the uncertainty in future projections by allowing both spatial and temporally realistic variability in sub ice shelf melt rates. It seems that TG does not face the risk of marine ice sheet instability in the near future because of the prograde slope in the first tens of kilometres inland of the grounding line. Our simulations do not retreat as past this slope before the year 2100. Once the grounding line retreats further onto the retrograde bed slope then we would expect far faster grounding line retreats in the long term. The change of sub-shelf cavity thickness is dominated by sub-shelf melt rates, although different basal sliding parameterizations could yield different retreat of grounding line position, hence different sub-shelf cavity thickness near the

600 grounding line. This indicates the need to consider geometry change of sub-shelf cavity and water circulation inside the sub-shelf cavity when using ocean model or plume model for long term projections.

Data availability. MEaSURES Antarctic Boundaries for IPY 2007–2009 from Satellite Radar, version 2, is available at <https://doi.org/10.5067/AXE4121732AD> (Mouginot et al., 2017). MEaSURES BedMachine Antarctica, version 2, is
605 available at <https://doi.org/10.5067/E1QL9HFQ7A8M> (Morlighem, 2020). MEaSURES InSAR-based Antarctic ice velocity Map, version 2, is available at <https://doi.org/10.5067/D7GK8F5J8M8R> (Rignot et al., 2017). Ice temperature of SICOPOLIS output (Greve et al., 2020) is available upon request to the author. SMB of CESM2 under the SSP5-8.5 scenario is available from the Globus ISMIP6_Forc endpoint, at https://app.globus.org/file-manager?origin_id=ad1a6ed8-4de0-4490-93a9-8258931766c7&origin_path=%2F (Nowicki et al., 2016). Elmer/Ice is a free and open-source software,
610 published on Zenodo (Elmer/Ice Authors, 2022, <https://doi.org/10.5281/zenodo.7892181>).

Author contribution. LZ and JCM conceived the study. LZ, RG, TZ, MW and JM designed the methodology. MY and LZ carried out the experiments and produced the estimates. MY and LZ wrote the original draft, and all the authors revised the paper.

615

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgments. This work was supported by National Key Research and Development Program of China (grant no. 2021YFB3900105), Academy of Finland (grant no. 322430 and 355572), and the Finnish Ministry of Education and Culture
620 and CSC-IT Center for Science (Decision diary number OKM/10/524/2022). The authors would like to thank the anonymous referees for their valuable suggestions.

Financial support. This research has been supported by National Key Research and Development Program of China (grant no. 2021YFB3900105).

625 **References**

- Adusumilli, S., Fricker, H.A., Medley, B., , Padman, L., & Siegfried M. R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves. *Nat. Geosci.* 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- Barnes, J. M. and Gudmundsson, G. H.: The predictive power of ice sheet models and the regional sensitivity of ice loss to
630 basal sliding parameterisations: a case study of Pine Island and Thwaites glaciers, West Antarctica, *The Cryosphere*, 16, 4291–4304, <https://doi.org/10.5194/tc-16-4291-2022>, 2022.

- Berends, C. J., Van De Wal, R. S., Van Den Akker, T., & Lipscomb, W. H.: Compensating errors in inversions for subglacial bed roughness: same steady state, different dynamic response, *The Cryosphere*, 17(4), 1585-1600, 2023.
- Budd, W., Keage, P., and Blundy, N.: Empirical studies of ice sliding. *J. Glaciol.*, 23(89), 157–170, 1979.
- 635 Budd, W., Jenssen, D., and Smith, I.: A three-dimensional time-dependent model of the West Antarctic ice sheet. *Ann. Glaciol.*, 5, 29–36, 1984.
- Brondex, J., Gagliardini, O., Gillet-Chaulet, F., and Durand, G.: Sensitivity of grounding line dynamics to the choice of the friction law, *J. Glaciol.*, 63, 854–866, 2017.
- Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to
640 the friction law, *The Cryosphere*, 13, 177–195, 2019.
- Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, fourth edition, Academic Press, Amsterdam, 2010.
- Coulon, V., Bulthuis, K., Whitehouse, P. L., Sun, S., Haubner, K., Zipf, L., & Pattyn, F.: Contrasting response of West and East Antarctic ice sheets to glacial isostatic adjustment, *J. Geophys. Res.-Earth Surf.*, 126, e2020JF006003, <https://doi.org/10.1029/2020JF006003>, 2021.
- 645 Dapogny, C., Dobrzynski, C., and Frey, P.: Three-dimensional adaptive domain remeshing, implicit domain meshing, and applications to free and moving boundary problems, *J. Comput. Phys.*, 262, 358–378, <https://doi.org/10.1016/j.jcp.2014.01.005>, 2014.
- de Boer, B., Stocchi, P., and van de Wal, R. S. W.: A fully coupled 3-D ice-sheet-sea-level model: algorithm and applications, *Geosci. Model Dev.*, 7, 2141-2156, <https://doi.org/10.5194/gmd-7-2141-2014>, 2014.
- 650 Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502, 89–92, <https://doi.org/10.1038/nature12567>, 2013.
- Dow, C. F., McCormack, F. S., Young, D. A., Greenbaum, J. S., Roberts, J. L., and Blankenship, D. D.: Totten Glacier subglacial hydrology determined from geophysics and modeling, *Earth Planet. Sci. Lett.*, 531, 115961, <https://doi.org/10.1016/j.epsl.2019.115961>, 2020.
- 655 Fowler, A. C.: A Theoretical Treatment of the Sliding of Glaciers in the Absence of Cavitation, *Philos. Trans. R. Soc. Lond. Ser. A-Math. Phys. Eng. Sci.*, 298(1445): 637-681, 1981.
- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., de Fleurian, B., Greve, R., Malinen, M., Martín, C., Råback, P., Ruokolainen, J., Sacchettini, M., Schäfer, M., Seddik, H., and Thies, J.: Capabilities and performance of
660 Elmer/Ice, a new-generation ice sheet model, *Geosci. Model Dev.*, 6, 1299–1318, <https://doi.org/10.5194/gmd-6-1299-2013>, 2013.
- Gagliardini, O., Cohen, D., Raback, P., Zwinger, T.: Finite-element modeling of subglacial cavities and related friction law, *J. Geophys. Res.*, 112 (F02027), 1–11, 2007.
- Geuzaine, C. and Remacle, J.-F.: Gmsh: A 3-D finite element mesh generator with built-in pre- and post-processing facilities,
665 *Int. J. Numer. Methods Eng.*, 79, 1309–1331, <https://doi.org/10.1002/nme.2579>, 2009.

- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.
- 670 Gladstone, R., Zhao, C., and Zwinger, T.: ISMIP6 Projections-Antarctica read me file, Zenodo, <https://doi.org/10.5281/zenodo.3484635>, 2019.
- Gladstone, R. M., Warner, R. C., Galton-Fenzi, B. K., Gagliardini, O., Zwinger, T., and Greve, R.: Marine ice sheet model performance depends on basal sliding physics and sub-shelf melting, *The Cryosphere*, 11, 319–329, <https://doi.org/10.5194/tc-11-319-2017>, 2017.
- 675 Gladstone, R. and Wang, Y.: Antarctic regional inversions using Elmer/Ice: methodology, Zenodo, <https://doi.org/10.5281/zenodo.5862046>, 2022.
- Greenbaum, J. S., Blankenship, D. D., Young, D. A., Richter, T. G., Roberts, J. L., Aitken, A. R. A., Legresy, B., Schroeder, D. M., Warner, R. C., van Ommen, T. D., and Siegert, M. J.: Ocean access to a cavity beneath Totten Glacier in East Antarctica, *Nat. Geosci.*, 8, 294–298, <https://doi.org/10.1038/ngeo2388>, 2015.
- 680 Greene, C. A., Young, D. A., Gwyther, D. E., Galton-Fenzi, B. K., and Blankenship, D. D.: Seasonal dynamics of Totten Ice Shelf controlled by sea ice buttressing, *The Cryosphere*, 12, 2869–2882, <https://doi.org/10.5194/tc-12-2869-2018>, 2018.
- Greve, R., Calov, R., Obase, T., Saito, F., Tsutaki, S. and Abe-Ouchi, A.: ISMIP6 future projections for the Antarctic ice sheet with the model SICOPOLIS, Zenodo, <https://doi.org/10.5281/zenodo.4035932>, 2020.
- Greve, R. and Blatter, H.: *Dynamics of Ice Sheets and Glaciers*, Springer Berlin Heidelberg, Berlin, Heidelberg, <https://doi.org/10.1007/978-3-642-03415-2>, 2009.
- 685 Gudmundsson, G. H., Krug, J., Durand, G., Favier, L., and Gagliardini, O.: The stability of grounding lines on retrograde slopes, *The Cryosphere*, 6, 1497–1505, <https://doi.org/10.5194/tc-6-1497-2012>, 2012.
- Gwyther, D. E., O’Kane, T. J., Galton-Fenzi, B. K., Monselesan, D. P., and Greenbaum, J. S.: Intrinsic processes drive variability in basal melting of the Totten Glacier Ice Shelf. *Nat. Commun.*, 9(3141). <https://doi.org/10.1038/s41467-018-05618-2>, 2018.
- 690 Hansen, P. C.: The L-curve and its use in the numerical treatment of inverse problems, in: *Computational Inverse Problems in Electrocardiology*, WIT Press, Southampton, 119–142, 2001.
- Iverson, N.R., T.S. Hooyer and R.W. Baker: Ring-shear studies of till deformation: Coulomb-plastic behavior and distributed strain in glacier beds, *J. Glaciol.*, 44, 634–642, 1998.
- Joughin, I., Smith, B.E., and Schoof, C.G.: Regularized coulomb friction laws for ice sheet sliding: application to pine island glacier, antarctica. *Geophys. Res. Lett.*, 46(9), 4764–4771, 2019. doi: 10.1029/2019GL082526.
- 695 Roberts, J., Galton-Fenzi, B. K., Paolo, F. S., Donnelly, C., Gwyther, D. E., Padman, L., et al. (2018b). Ocean forced variability of Totten Glacier mass loss. *Geological Society, London, Special Publications*, 461(1), 175–186. <https://doi.org/10.1144/SP461.6>

- Kang, H., Zhao, L., Wolovick, M., and Moore, J. C.: Evaluation of six geothermal heat flux maps for the Antarctic Lambert–Amery glacial system, *The Cryosphere*, 16, 3619–3633, <https://doi.org/10.5194/tc-16-3619-2022>, 2022.
- Kamb, W. B.: Sliding motion of glaciers: theory and observation. *Rev. Geophys.*, 8(4): 673–728, 1970.
- Leguy, G. R., Asay-Davis, X. S., and Lipscomb, W. H.: Parameterization of basal friction near grounding lines in a one-dimensional ice sheet model, *The Cryosphere*, 8, 1239–1259, <https://doi.org/10.5194/tc-8-1239-2014>, 2014
- Leguy, G. R., Lipscomb, W. H., and Asay-Davis, X. S.: Marine ice sheet experiments with the Community Ice Sheet Model, *The Cryosphere*, 15, 3229–3253, <https://doi.org/10.5194/tc-15-3229-2021>, 2021.
- Li, T., Dawson, G. J., Chuter, S. J., & Bamber, J. L.: Grounding line retreat and tide-modulated ocean channels at Moscow University and Totten Glacier ice shelves, East Antarctica. *The Cryosphere*, 17(2), 1003–1022, 2023. <https://doi.org/10.5194/tc-17-1003-2023>
- Li, X., Rignot, E., Morlighem, M., Mouginot, J., and Scheuchl, B.: Grounding line retreat of Totten Glacier, East Antarctica, 1996 to 2013, *Geophys. Res. Lett.*, 42, 8049–8056, <https://doi.org/10.1002/2015GL065701>, 2015.
- Li, X., Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow dynamics and mass loss of Totten Glacier, East Antarctica, from 1989 to 2015, *Geophys. Res. Lett.*, 43, 6366–6373, doi:10.1002/2016GL069173, 2016.
- Liu, Y., Moore, J. C., Cheng, X., Gladstone, R. M., Bassis, J. N., Liu, H., Wen, J., and Hui, F.: Ocean-driven thinning enhances iceberg calving and retreat of Antarctic ice shelves, *P. Natl. Acad. Sci. USA*, 112, 3263–3268, <https://doi.org/10.1073/pnas.1415137112>, 2015.
- Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *J. Geophys. Res.*, 117, F01022, <https://doi.org/10.1029/2011JF002140>, 2012.
- Lazeroms, W. M. J., Jenkins, A., Gudmundsson, G. H., and van de Wal, R. S. W.: Modelling present-day basal melt rates for Antarctic ice shelves using a parametrization of buoyant meltwater plumes. *The Cryosphere*, 12, 49–70. <https://doi.org/10.5194/tc-12-49-2018>, 2018.
- Martos, Y. M., Catalán, M., Jordan, T. A., Golynsky, A., Golynsky, D., Eagles, G., and Vaughan, D. G.: Heat Flux Distribution of Antarctica Unveiled, *Geophys. Res. Lett.*, 44, 11,417–11,426, <https://doi.org/10.1002/2017GL075609>, 2017.
- McCormack, F. S., Roberts, J. L., Gwyther, D. E., Morlighem, M., Pelle, T., and Galton-Fenzi, B. K.: The impact of variable ocean temperatures on Totten Glacier stability and discharge. *Geophysical Research Letters*, 48, e2020GL091790. <https://doi.org/10.1029/2020GL091790>, 2021.
- Morlighem, M.: MEaSUREs BedMachine Antarctica, Version 2, <https://doi.org/10.5067/E1QL9HFQ7A8M>, 2020.
- Morlighem, M., Rignot, E., Seroussi, H., Larour, E., Ben Dhia, H., and Aubry, D.: Spatial patterns of basal drag inferred using control methods from a full-Stokes and simpler models for Pine Island Glacier, West Antarctica, *Geophys. Res. Lett.*, 37, L14502, <https://doi.org/10.1029/2010GL043853>, 2010.

- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. van den, Ommen, T. D. van, Wessem, M. van, and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nat. Geosci.*, 13, 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- Mouginot, J., Scheuchl B. and Rignot, E.: MEaSURES Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version 2, <https://doi.org/10.5067/AXE4121732AD>, 2017.
- Nilsson, J., Gardner, A. S. & Paolo, F.: MEaSURES ITS_LIVE Antarctic Grounded Ice Sheet Elevation Change. (NSIDC-0782, Version 1). [Data Set]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/L3LSVDZS15ZV>, 2023.
- Nye, J. F.: Glacier sliding without cavitation in a linear viscous approximation. *Proceedings of the Royal Society of London, Ser. A*, 315, (1522): 381-403, 1970.
- Nakayama, Y., Greene, C. A., Paolo, F. S., Mensah, V., Zhang, H., Kashiwase, H., Simizu, D., Greenbaum, J. S., Blankenship, D. D., Abe-Ouchi, A., and Shigeru Aoki, S.: Antarctic Slope Current modulates ocean heat intrusions towards Totten Glacier. *Geophysical Research Letters*, 48, e2021GL094149, <https://doi.org/10.1029/2021GL094149>, 2021.
- Nowicki, S. M. J., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geosci. Model Dev.*, 9, 4521–4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.
- Nias, I., Cornford, S., and Payne, A.: New mass-conserving bedrock topography for Pine Island Glacier impacts simulated decadal rates of mass loss, *Geophys. Res. Lett.*, 45, 3173–3181, 2018.
- Pelle, T., Morlighem, M., and McCormack, F. S.: Aurora Basin, the weak underbelly of East Antarctica, *Geophys. Res. Lett.*, 47, e2019GL086821, <https://doi.org/10.1029/2019GL086821>, 2020.
- Pelle, T., Morlighem, M., Nakayama, Y., and Seroussi, H.: Widespread grounding line retreat of Totten Glacier, East Antarctica, over the 21st century. *Geophys. Res. Lett.*, 48, e2021GL093213, <https://doi.org/10.1029/2021GL093213>, 2021.
- Ribeiro, N., Herraiz-Borreguero, L., Rintoul, S. R., McMahon, C. R., Hindell, M., Harcourt, R., and Williams, G.: Warm modified Circumpolar Deep Water intrusions drive ice shelf melt and inhibit Dense Shelf Water formation in Vincennes Bay, East Antarctica. *J. Geophys. Res.-Oceans*, 126, e2020JC016998. <https://doi.org/10.1029/2020JC016998>, 2021.
- Richter, O., Gwyther, D. E., Galton-Fenzi, B. K., and Naughten, K. A.: The Whole Antarctic Ocean Model (WAOM v1.0): development and evaluation, *Geosci. Model Dev.*, 15, 617–647, <https://doi.org/10.5194/gmd-15-617-2022>, 2022.
- Rignot, E., Jacobs, S. S., Mouginot, J., and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341, 266–270, <https://doi.org/10.1126/science.1235798>, 2013.

- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proc. Natl. Acad. Sci.*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- 770 Rignot, E., J. Mouginot and Scheuchl, B.: MEaSUREs InSAR-Based Antarctica Ice Velocity Map, Version 2, <https://doi.org/10.5067/D7GK8F5J8M8R>, 2017.
- Rintoul, S. R., Silvano, A., Pena-Molino, B., van Wijk, E., Rosenberg, M., Greenbaum, J. S., and Blankenship, D. D.: Ocean heat drives rapid basal melt of the Totten Ice Shelf, *Sci. Adv.*, 2, e1601610, <https://doi.org/10.1126/sciadv.1601610>, 2016.
- 775 Rückamp, M., Kleiner, T., and Humbert, A.: Comparison of ice dynamics using full-Stokes and Blatter-Pattyn approximation: application to the Northeast Greenland Ice Stream, *The Cryosphere*, 16, 1675–1696, <https://doi.org/10.5194/tc-16-1675-2022>, 2022.
- Shechepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Model.*, 9, 347–404, 2005.
- 780 Schoof, C.: The effect of cavitation on glacier sliding, *P. Roy. Soc. A-Math. Phys.*, 461, 609–627, [doi:10.1098/rspa.2004.1350](https://doi.org/10.1098/rspa.2004.1350), 2005.
- Schoof, C.: Marine ice sheet stability. *Journal of Fluid Mechanics*, 698, 62–72, 2012.
- Sergienko, O. V., & Hindmarsh, R. C. A.: Regular patterns in frictional resistance of ice-stream beds seen by surface data inversion. *Science*, 342(6162), 1086–1089, 2013.
- 785 Sergienko, O. V., Creyts, T. T., & Hindmarsh, R. C. A.: Similarity of organized patterns in driving and basal stresses of Antarctic and Greenland ice sheets over extensive areas of basal sliding. *Geophysical Research Letters*, 41(11), 3925–3932, 2014.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, 790 G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N.-J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere*, 14, 3033–3070, <https://doi.org/10.5194/tc-14-3033-2020>, 2020.
- 795 Stearns, L.A., Smith, B.E. and Hamilton, G.S.: Increased flow speed on a large east Antarctic outlet glacier caused by subglacial floods. *Nat. Geosci.*, 1(12), 827–831, 2008.
- Silvano, A., Rintoul, S. R., Peña-Molino, B., and Williams, G. D.: Distribution of water masses and meltwater on the continental shelf near the Totten and Moscow University ice shelves, *J. Geophys. Res.-Oceans*, 122, 2050–2068, <https://doi.org/10.1002/2016JC012115>, 2017.

- Sun, S., Cornford, S. L., Moore, J. C., Gwyther, D. E., Gladstone, R. M., Galton-Fenzi, B. K., Zhao, L.: Impact of ocean forcing on the Aurora Basin in the 21st and 22nd centuries, *Ann. Glaciol.*, 57(73), 1–8, 2016.
- Tsai, V. C., Stewart, A. L., and Thompson, A. F.: Marine ice-sheet profiles and stability under Coulomb basal conditions, *J. Glaciol.*, 61, 205–215, 2015.
- Van Dalum, C., Van de Berg, W. J., and Van den Broeke, M.: Monthly RACMO2.4p1 data for Antarctica (11 km) for SMB, SEB and near-surface variables (1979-2023), <https://doi.org/10.5281/zenodo.14217232>, Zenodo [data set], 2024.
- 805 Vaňková, I., Winberry, J. P., Cook, S., Nicholls, K. W., Greene, C. A., and Galton-Fenzi, B. K.: High spatial melt rate variability near the Totten Glacier grounding zone explained by new bathymetry inversion. *Geophys. Res. Lett.*, 50, e2023GL102960, <https://doi.org/10.1029/2023GL102960>, 2023.
- Wang, Y., Zhao, C., Gladstone, R., Galton-Fenzi, B., and Warner, R.: Thermal structure of the Amery Ice Shelf from borehole observations and simulations, *The Cryosphere*, 16, 1221–1245, 2022.
- 810 Weertman, J.: On the Sliding of Glaciers, *J. Glaciol.*, 3, 33–38, <https://doi.org/10.3189/S0022143000024709>, 1957.
- Werder, M. A., Hewitt, I. J., Schoof, C. G., Flowers, G.E., 2013. Modeling channelized and distributed subglacial drainage in two dimensions. *J. Geophys. Res.- Earth Surf.*, 118, 2140–2158. <https://doi.org/10.1002/jgrf.20146>
- Wolovick, M., Humbert, A., Kleiner, T., and Rückamp, M.: Regularization and L-curves in ice sheet inverse models: a case study in the Filchner–Ronne catchment, *The Cryosphere*, 17, 5027–5060, <https://doi.org/10.5194/tc-17-5027-2023>, 2023.
- 815 Yu, H., Rignot, E., Seroussi, H., and Morlighem, M.: Retreat of Thwaites Glacier, West Antarctica, over the next 100 years using various ice flow models, ice shelf melt scenarios and basal friction laws. *The Cryosphere*, 12(12), 3861–3876. <https://doi.org/10.5194/tc-12-3861-2018>, 2018.
- Zhao, C., Gladstone, R. M., Warner, R. C., King, M. A., Zwinger, T., and Morlighem, M.: Basal friction of Fleming Glacier, Antarctica – Part 1: Sensitivity of inversion to temperature and bedrock uncertainty, *The Cryosphere*, 12, 2637–2652, <https://doi.org/10.5194/tc-12-2637-2018>, 2018.
- 820 Zoet, L. K., Iverson, N. R., Andrews, L., and Helanow, C.: Transient evolution of basal drag during glacier slip. *J. Glacio.*, 68(270), 741–750. <https://doi.org/10.1017/jog.2021.13>, 2022.
- Zwally, H. J., Abdalati, W., Herring, T., Larson, K., Saba, J., Steffen, K.: Surface melt-induced acceleration of Greenland ice-sheet flow. *Science*, 297, 218–222, 2002.