

Importance of basement faulting and salt decoupling for the structural evolution of the Fars Arc, Zagros fold-and-thrust belt: A numerical modeling approach

Fatemeh Gomar¹, Jonas B. Ruh², Mahdi Najafi³, Farhad Sobouti¹

¹Department of Earth Sciences, Institute for Advanced Studies in Basic Sciences, Zanjan, 45137-66731, Iran

²Institute of Marine Sciences, Consejo Superior de Investigaciones Científicas, 08003, Barcelona, Spain

³Geosciences Barcelona, Consejo Superior de Investigaciones Científicas, GEO3BCN-CSIC, 08028, Barcelona, Spain

Correspondence to: *Fatemeh Gomar (fatemehgomar@iasbs.ac.ir) and Jonas B. Ruh (jruh@icm.csic.es)*

Formatted: Spanish (Spain)

Abstract. Understanding the tectonic evolution and crustal-scale structure of fold-and-thrust belts is crucial for exploring geological resources and evaluating seismic hazards. We conducted a series of two-dimensional finite-difference ~~two-dimensional~~ thermo-mechanical numerical models with a visco-elasto-plastic/brittle rheology to decipher how the interaction of inherited basement faults and salt décollement levels control the deformation process and structural style of the Fars Arc in the Zagros fold-and-thrust belt during tectonic inversion. Results~~Our results~~ indicate that strain accumulation pattern during initial rifting is controlled by the geometry of inherited faults. During the convergence phase, ~~fold-and-thrust belts display folding at two scales: the inverted basement faults form large-~~ wavelength ~~folds induced by basement deformation in the form of foreland-verging fault-propagation faults, and small~~ wavelength folds and thrust systems emerge above the anticlines in the sedimentary cover, while the thick salt layer ~~aspromotes the growth of second-order~~ detachment ~~folds; anticlines accompanied by both fore- and back-limb thrust~~ faults. Reactivated faults can serve as pathways for stress transfer, resulting in the emergence~~formation~~ of new faults ~~and thus triggering~~ seismic activity. The ~~tectonic events in structural evolution of~~ orogenic belts like the Zagros do not adhere to a fixed pattern; ~~they are it is~~ shaped by factors such as ~~the basement rock~~ properties ~~of basement rocks and the orientation of faults and inherited fault orientations~~. Shallow earthquakes predominantly occur along décollement anticlines in the sedimentary cover, while deeper ~~and~~ larger ones are associated with basement faults. Additionally, ~~we~~We also observe variations in resistance to deformation based on salt rheology and fault geometry, with listric faults minimizing resistance. The degree of basement involvement ~~in deformation~~ directly influences the model's resistance to deformation, with greater involvement facilitating easier deformation. Our results ~~showing related to~~ the temporal-and spatial relationship between thin- and thick-skinned tectonics can work as ~~an analogue~~analogues for similar orogenic belts worldwide, such as Taiwan, the Pyrenees, the Alps, the Appalachians, and the Kopet Dagh.

1 Introduction

Fold-and-thrust belts are complex tectonic domains formed in response to near- or far-field compressional stress fields in the Earth's crust. Multiple factors have been identified to control the structural styles in these belts, such as; the strength of the involved rocks (Davis et al., 1983), the involvement of the crystalline basement and its pre-existing faults (Pfiffner, 2017; Barchi and Tavarelli, 2022), the occurrence and rheology of décollement layers (Ruh et al.,

35 2012; Pla et al., 2019; Eslami Rezaei et al., 2023), and the intensity of surface processes (Talbot and Alavi, 1996; Cooper, 2007; Simpson, 2010; Malavieille and Konstantinov, 2010; Morley et al., 2011; McQuarrie and Ehlers, 2017). In particular, the presence of décollements, mechanically weak layers, composed of rocks that have lower mechanical strength than their surroundings, plays a significant role in the formation and evolution of fold-and-thrust belts. These layers are weak stratigraphic horizons that separate and mechanically decouple layers of greater strength and tend to accommodate and localize the deformation (Koyi and Mansurbeg, 2021). The relative weakness of décollement levels arises from variations in lithology (i.e., layer-parallel décollements) or structural features (i.e., inherited faults) that often promote localization of deformation as a result of increased stress contrasts and lower strength thresholds (Vogt et al., 2017; Broderie et al., 2018). The lithology of a décollement layer (salt, anhydrite, gypsum or shale), its stratigraphic position in the rock sequence (basal or middle), its thickness, and its fluid content control the distribution and geometry of faulting and folding during crustal convergence (Simpson, 2009; Ruh et al., 2012; Najafi et al., 2014; Santolaria et al, 2022). Décollement layers with a certain thickness may lead to the formation of single-layer detachment folds (Mitra, 2003; Wallace and Homza, 1998). Furthermore, multiple décollement horizons within a shortened sedimentary sequence may connect along thrust faults and form structural ramp-flat geometries (Boyer and Elliott, 1982; Dal Zilio, 2020).

50 Besides weak layers within the stratigraphy, pre-existing basement faults, inherited from older tectonic events, can influence the location and geometry of new faults and folds and their propagation and termination in the overlying sedimentary cover, by serving as zones of weakness during compressional deformation. This ultimately affects the overall structural architecture of a fold-and-thrust belt (Bonini, 2012; Granado and Ruh, 2019; Parizot et al., 2022). The effects of fault inheritance can be complex and dependent on various factors, such as the orientation, dip, geometry, and depth of the inherited structures (White et al., 1986), the timing and style of subsequent deformation (Zwaan et al., 2022), and the mechanical properties of the rocks involved (Ruh and Vergés, 2018).

60 The Fars Arc, located in the southeastern part of the Zagros Orogen, (Fig. 1), offers an ideal area to study the combined influence of cover mechanical stratigraphy and basement inheritance. It is a typical example of a fold-and-thrust belt that contains multiple mechanically weak stratigraphic layers in the Phanerozoic sedimentary cover, as well as pre-existing inherited faults in the Precambrian crustal basement (Jackson and Fitch, 1981; Berberian, 1995; Talebian and Jackson, 2004; Mouthereau et al., 2006; Yamato et al., 2011; Karasözen et al., 2019; Najafi et al., 2021). According to seismic, well-logging, and surface geological data, the Ediacaran–Lower Cambrian Hormuz series constitutes the main décollement in the Fars Arc. It consists of ~2 km of salt-bearing evaporites with minor carbonate and shale layers and decouples the crustal basement from the overlying sedimentary cover (Kent, 1958; Sherkati et al., 2005; Callot et al., 2007; Leturmy et al., 2010). The presence of a thick salt layer at the base of the sedimentary cover in the Fars Arc is responsible for short-wavelength folds (Mouthereau et al., 2006). In some places, its basal décollement is spatially interrupted by basement faults related to the Cenozoic convergence (Kent, 1958; Talbot and Alavi, 1996; Sepehr and Cosgrove, 2004; Sherkati et al., 2005; Callot et al., 2007). The recent activity of these faults in the Fars Arc is documented by seismological data (Jackson and Fitch, 1981; Berberian and King, 1981; Berberian, 1995, Talebian and Jackson, 2004; Tatar and Hatzfeld, 2004; Karasözen et al., 2019; Nissen et al., 2019). Focal mechanisms of seismic events in the basement exhibit rather steep fault dip angles, suggesting the reactivation of inherited normal faults

(Jackson, 1980). ~~In the Fars Arc, the activity of inherited faults has affected the progression of deformation towards the foreland, with the Mountain Front Fault being related to basement thrusting (Bahroudi and Koyi, 2003; Mouthereau et al., 2006; Mouthereau et al., 2007a; Yamato et al., 2011; Ruh et al., 2014; Najafi et al., 2021).~~

Formatted: English (United Kingdom)

In the Fars Arc, the activity of inherited faults has influenced the progression of deformation towards the foreland, with the Mountain Front Fault being associated with basement thrusting (Bahroudi and Koyi, 2003; Mouthereau et al., 2006, 2007a; Yamato et al., 2011; Ruh et al., 2014; Najafi et al., 2021). ~~reported that~~ According to Mouthereau et al. (2006), basement deformation and thickening ~~is crucial~~ play a critical role in the Zagros Folded Belt, helping to ~~reproduce~~ explain the observed topographic growth ~~observed~~. They furthermore highlight that ~~also emphasize that the reactivation of pre-existing faults during the early stages~~ of compression in the Zagros foredeep ~~was controlled by the reactivation of pre-existing faults, suggesting that~~ suggests a significant influence of inherited structural features significantly affect current ~~present-day~~ deformation patterns. While ~~balanced~~, Balanced cross-sections demonstrate ~~support this by demonstrating~~ that basement involvement is ~~required~~ necessary to explain various ~~account~~ for varying base topographic elevations in the Paleozoic and Mesozoic formations (Blanc et al., 2003; Molinaro et al., 2005; Mouthereau et al., 2007a). However, other studies ~~suggest~~ propose that the ~~major effect~~ most substantial impact of basement deformation on surface structures ~~during tectonic inversion~~ occurred ~~only later~~ in the late stage of the region's tectonic history ~~of, particularly during the Zagros Pliocene and Pleistocene~~ (Molinaro et al., 2005; Sherkati et al., 2005; Tavani et al., 2018), ~~probably during the Pliocene and Pleistocene~~ (Vergés et al., 2011; Najafi et al., 2018; Etemed-Saeed et al., 2020). ~~There~~ It is a general agreement ~~widely accepted~~ that the high-angle reverse faults ~~originally~~ initially formed as normal faults in the Arabian basement during the Permian–Triassic rifting of the Neo-Tethys Ocean (Navabpour et al., 2010). ~~The faults were~~, remaining inactive ~~during~~ through the Jurassic- to Late Cretaceous passive margin phase ~~and were finally, before being~~ reactivated during the Cenozoic ~~continental~~ collision ~~of between~~ the Arabian and Eurasian plates.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: English (United Kingdom)

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Various 2D numerical modeling studies have investigated the evolution of fold-and-thrust belts and salt-bearing basins. Nilforoushan et al. (2013) demonstrated the influence of geothermal gradients and basement mineralogy on fault geometry and basement reactivation in the Fars arc, emphasizing the role of weak salt horizons in mechanical decoupling. Heydarzadeh et al. (2020) analysed factors such as sedimentation rates, erosion, and salt layer properties in Dehdasht basin, highlighting the importance of balanced surface processes and deformation rates. Humair et al. (2020) conducted simulations to study the interaction of folding and thrusting during Swiss Jura and Canadian Foothills fold-and-thrust belt evolution, focusing on the effects of layer-parallel shortening and initial geometrical perturbations. Their work showed that the magnitude of these perturbations influences whether folding or thrusting predominates, affecting the structural evolution and asymmetry of anticlines. Spitz et al. (2020) conducted 3D thermo-mechanical numerical simulations to investigate the influence of laterally variable inherited structures on fold-and-thrust belt evolution and nappe formation on Helvetic nappe system. The study demonstrated the fundamental importance of tectonic inheritance on fold-and-thrust belt evolution, with strain localization, folding, and nappe transport controlled by initial geometrical and mechanical heterogeneities. Almost all studies have focused on examining the collisional phase and deformation resulting from compression in the fold belts and the Fars Arc, while

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman


the earlier extensional history and its effect on later deformation have received less attention (e.g., Granado and Ruh, 2019).

~~On the other hand, the~~The interaction of the basal décollement level and the pre-existing basement faults and their combined influence on the distribution of thin- and thick-skinned tectonic styles in the Fars Arc, during both extensional and compressional phases, is not fully understood yet.

Understanding these processes is crucial for improving our geological models of the region, which has significant implications for hydrocarbon exploration. Although there is a general consensus on the importance of the décollement layer and basement faults, the detailed dynamics and their broader impact on regional tectonics require further investigation in during inversion tectonics.

The goal of our study is to test the impact of the Hormuz salt, and the mechanical properties and geometries of the basement faults inherited from the Permian continental rifting on the structural evolution of the Fars Arc. To achieve this goal, we conducted a series of two-dimensional thermo-mechanical numerical experiments to investigate how the presence or absence of fault inheritance (planar or listric) controls the deformation during rifting and subsequent tectonic shortening. Furthermore, we test the effect of a weak intermediate salt décollement of variable rheology on the structural evolution during convergence. ▲

Formatted: Font: (Default) Times New Roman, 10 pt, English (United Kingdom)



125 Figure 1: (a) Overview of the geotectonic situation of Iran. (b) Main tectonic features in the Zagros fold-and-thrust belt, located along the northeastern margin of the Arabian Plate. (c) Regional geological cross-section, showing the crustal geometry (modified from Etemad-Saeed et al. 2020; Najafi et al. 2021; Najafi and Lajmorak 2020).

2 Geotectonic and geological setting

130 The Zagros fold-and-thrust belt (ZFTB) is a NW-SE-trending orogenic belt that resulted from the convergence and continental collision between Arabian and Eurasian plates (Agard et al., 2011; Mouthereau et al. 2012; Vergés et al., 2024). The ZFTB extends over approximately 2000 km from the Taurus Mountains in Turkey in the northwest, to the Makran accretionary wedge in southeastern Iran (Fig. 1). Across strike, the ZFTB is bounded by the Main Zagros Thrust (i.e., the inherited suture of the old subduction zone) in the northeast, and by the Zagros Frontal Fault system in the southwest (Fig. 1b).

135 The NE margin of the Arabian Plate has been shaped by a series of tectonic events that ultimately resulted in the emergence of the ~~Zagros fold-and-thrust belt~~ZFTB (Agard et al., 2011; Vergés et al., 2024; Madanipour et al., 2024): 1) Permian–Triassic rifting and opening of the NW-SE-trending Neo-Tethys Ocean, 2) Jurassic–Late Cretaceous passive marginal stage, 3) Late Cretaceous ophiolite obduction, 4) Oligocene soft continental collision between Arabian and Eurasian plates, and 5) middle Miocene-to-recent folding propagation across the mountain range.

140 The Permian–Triassic opening of the Neo-Tethys Ocean separated the Arabian Plate to the southwest from the Iranian microplate to the northeast (Szabo and Kheradpir, 1978; Berberian and King, 1981; Agrad et al., 2005). Lithospheric extension related to continental rifting deformed the Arabian crystalline basement and produced a series of NW-trending half-grabens parallel to the current orientation of the Zagros orogen (e.g. Jackson and Fitch, 1981; Mouthereau et al., 2007a). Following the rifting episode, the Arabian margin became passive during the Jurassic and Cretaceous (e.g. Alavi, 2004; Agard et al., 2005). An early stage of contractional deformation occurred in the Late Cretaceous with the obduction of the Neo-Tethys ophiolite and radiolarite slices onto the Arabian margin, presently preserved in Kermanshah, Neyriz, and Hajiabad (Agard et al., 2005; Saura et al., 2011; Bernaola et al., 2011).

145 From the Oligocene onward, continental collision led to the formation of the ZFTB, age-constrained by recent thermochronometric and magnetostratigraphic data (Pirouz et al., 2017; Koshnaw et al., 2019; Barber et al., 2019). The main phase of folding in the Zagros took place during the Miocene to Pleistocene, and progressively propagated to the SW (Hessami et al., 2001; Khadivi et al., 2009; Mouthereau et al., 2012; Ruh et al., 2014; Vergés et al., 2019; Najafi et al., 2021). GPS measurements indicate that the present-day convergence rate is approximately 20–30 mm/yr in a roughly N-S direction, of which $\sim 6.5 \pm 2$ mm/yr is being consumed across the Zagros (Nilforoushan et al., 2003; Walpersdorf et al., 2006).

155 Based on its structural and stratigraphic characterizations, the Zagros is divided into the High Zagros imbricated zone in the NE, and the Zagros Simply Folded Belt (ZSFB) in the SW, separated by the High Zagros Fault (Fig. 1). The ZSFB extends to the Persian Gulf and the present-day Mesopotamian foredeep basin. It represents the deformed foreland of the orogeny, and displays elongated folds of regular wavelength (Falcon, 1974; Sepehr and Cosgrove, 2004; Mouthereau et al., 2006). The ZSFB consists of an 8–14 km-thick sedimentary sequence covering metamorphosed rocks of the Precambrian Arabian basement (Mouthereau et al., 2007b; Lacombe et al., 2011). Based on lateral stratigraphic and structural variations, the ZSFB is divided, from NW to SE, into the Lurestan Arc, the Izeh zone, the Dezful Embayment, and the Fars Arc (Fig. 1).

160 In this study, we focus on the Fars Arc, the largest tectonic domain of the Simply Folded Belt, limited by the Kazerun–Borazjan segmented dextral fault system to the west and the Minab–Zendan–Palami fault system to the east (Fig. 1b;

165 Regard et al., 2004; Lacombe et al., 2011). The Fars Arc extends over more than 300 km across strike, is ~200 km
wide from the High Zagros Fault to the deformation front; and developed as a result of folding of the thick sedimentary
cover (Stöcklin, 1974, Berberian and King, 1981). Its elongated folds show a distinctive periodic pattern with axial
lengths reaching to 200 km and wavelengths of ~15 km (Fig. 1c; Mouthereau et al., 2007b; Najafi et al., 2021). The
Fars Arc is scattered with salt diapirs, a majority of them located in the eastern part of the arc (Sherkati et al., 2006;
170 Callot et al., 2007).

The stratigraphy of the Fars Arc contains a major basal and several minor intermediate mechanically weak
décollement layers (Fig. 2). Near the base of the sedimentary cover, the ~2 km-thick salt-bearing evaporites of Hormuz
Formation of Ediacaran–Early Cambrian age overlay early Ediacaran sediments (Callot et al., 2007). The 3-km-thick
Paleozoic sequence is dominantly composed of sandstone, dolomite, and shale. The Dehram Group with a thickness
175 of more than 1 km was deposited during the Permian–Triassic. The Triassic Dashtak Formation of 550–850 m
thickness overlies the Dehram Group and plays as the minor intermediate décollement level in the frontal Fars Arc
(Motamedi et al., 2012; Najafi et al., 2014). It laterally grades into the dolomites of the Khanekkat Formation in the
interior of the range, where it loses its efficiency as a décollement level (Szabo and Kheradpir, 1978). During the
passive marginal stage in the Jurassic–Late Cretaceous, the Zagros basin was characterized by a shallow-marine
180 environment, when the 2-km-thick carbonate and minor detrital and evaporite successions of the Khami and Bangestan
groups were deposited (Sharp et al., 2010).

The Neo-Tethyan oceanic crust was obducted over the Arabian plate margin during the Late Cretaceous, and loading
resulted in the flexure of the Arabian lithosphere. This produced an early foreland basin in the Lurestan region of the
NW Zagros, referred to as the Amiran foreland basin by Homke et al. (2009) and Saura et al. (2011). The thick
successions of deep-water shales of the Paleocene to Eocene formations indicate regional subsidence. The passive
margin succession of the Paleocene to Miocene sediments has a thickness of ~3 km (Jahani et al., 2009; Najafi et al.,
185 2014). In the Oligocene, the Asmari Formation was deposited over the Pabdeh Formation in the Fars region. The post-
Asmari clastics, known as the Fars Group, including the Gachsaran, Mishan, and Aghajari formations (early Miocene
to Pliocene), have a total thickness of about 3 km. Migration of the Aghajari-Bakhtyari sedimentary system towards
190 the foreland and the propagation of folding were in sequence. They have migrated at a rate of 20 mm/yr in the Fars
Arc and 15 mm/yr in the Lurestan Arc during the last 20 Myr (Ruh et al., 2014; Vergés et al., 2019).

In the Fars Arc, two main décollement levels occur in the stratigraphy (Fig. 2); the Hormuz Formation represents the
basal and the major décollement level, and the evaporites of the Dashtak Formation constitute a minor intermediate
décollement level (Callot et al., 2012; Motamedi et al., 2012; Najafi et al., 2014).

195 Seismic activity at mid-crustal depths ~~is the main provides key~~ evidence for basement involvement in the Zagros. Most
~~of the earthquake~~ centroid depths ~~of earthquakes fall in the range from 4- to 25 km-depth range, in~~, affecting both the
basement and cover, ~~and most of them show with many exhibiting~~ reverse focal mechanisms (Jackson and Fitch, 1981;
Berberian, 1995; Talebian and Jackson, 2004; Karasözen et al., 2019). ~~The~~In the Fars Arc, the major inherited
basement reverse faults ~~in the Fars Arc are~~, from SW to NE, ~~include~~ the Mountain Front Fault, the Surmeh Fault, the
200 High Zagros Fault, and the Main Zagros Thrust (Fig. 41c). A geological cross-section ~~aeross~~of the Fars Arc
~~shows reveals~~ evidence of ~~eeeval~~both thin-skinned and thick-skinned tectonic deformation ~~styles occurring~~

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)


Formatted: English (United States)

simultaneously (Mouthereau et al., 2007b; Najafi et al., 2021; Fig. 1c). Deformation This deformation is manifested expressed through a combination of large-scale detachment folds and forced folds (Jackson, 1980; Lacombe et al., 2011; Mouthereau et al., 2012).

While numerous studies have focused on the Zagros fold-and-thrust belts and the Fars Arc, several crucial aspects of

- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)
- Formatted: English (United States)


205



the Fars Arc's geological evolution remain poorly understood. Specifically, the exact mechanisms and timing of basement involvement, the interaction between basement faults and salt décollements during tectonic inversion, and the relative influence of thin-skinned versus thick-skinned tectonics on the overall structural evolution are still unresolved (Mouthereau et al., 2006; 2012). To address these uncertainties, we employ a numerical model that simulates the full tectonic history of the Fars Arc, including both an initial extensional phase and a subsequent compressional phase.

- Formatted: English (United States)

210



215 Figure 2: Stratigraphic column of the sedimentary cover in the Fars Arc (modified from Callot et al., 2007; Jahani et al. 2009; Motamedi et al., 2012; Mouthereau et al. 2007; Najafi et al. 2014; Sepehr and Cosgrove 2004; Sherkati et al. 2006). Violet and red lithologies represent regional décollement horizons.

220 Many studies have been conducted in the Zagros fold-thrust belt and the Fars arc. However, several critical aspects of the geological evolution of the Fars Arc remain poorly understood. Specifically, the precise mechanisms and timing of basement involvement, the interaction between basement faults and salt décollements during tectonic inversion, and the relative contributions of thin-skinned versus thick-skinned tectonics to the overall structural evolution are not fully resolved (see Mouthereau et al., 2006, 2012). We employ a numerical model that simulates the complete tectonic history of the Fars Arc, including an initial extensional phase followed by a compressional phase.

Formatted: English (United States)

3 Numerical Model

225 In order to investigate the role of inherited extensional faults and mechanically weak décollement horizons on the structural evolution of the Fars Arc, a series of numerical experiments were conducted. We apply the two-dimensional finite-difference code “Norma” (Ruh et al., 2022), with a fully staggered Eulerian grid, freely moving Lagrangian markers (marker-in-cell technique; Gerya, 2019), and a temperature-dependent visco-elasto-plastic/brittle rheology.

3.1 Governing equations

230 The numerical code solves for the conservation of mass and momentum (the Stokes equation) on a Eulerian grid to calculate the velocity and pressure fields:

$$\frac{\partial v_i}{\partial x_i} = 0 \quad (1)$$

and

$$-\frac{\partial p}{\partial x_i} + \frac{\partial \tau_{ij}}{\partial x_j} = \rho g_i, \quad (2)$$

235 where v_i and x_i are velocities and spatial coordinates, p the dynamic pressure (mean stress), τ_{ij} the deviatoric stresses, ρ the density (constant), and g the gravitational acceleration. Temperature is solved by considering the energy conservation on the Eulerian grid:

$$\rho C_p \left(\frac{DT}{Dt} \right) = \kappa \frac{\partial^2 T}{\partial x_i^2}, \quad (3)$$

where C_p is isobaric heat capacity, $\frac{D}{Dt}$ is the total time derivative, T is temperature, t is time, κ is the thermal conductivity coefficient.

240 The pressure and temperature fields calculated from the governing equations are interpolated onto the Lagrangian markers based on a linear distance-weighted scheme (Gerya, 2019). The Lagrangian markers store all material properties and advect through the Eulerian grid based on a fourth-order Runge-Kutta interpolation of the calculated two-directional Eulerian velocity field.

245 3.2 Rheological model

The numerical code employs a visco-elasto-plastic/brittle rheology, where visco-elasticity is implemented by a Maxwell-type expression. The strain rate of a Maxwell body under stress consists of viscous and elastic components:

$$\dot{\epsilon}_{ij} = \frac{1}{2\eta} \tau_{ij} + \frac{1}{2G} \frac{D\tau_{ij}}{Dt}, \quad (4)$$

250 where G is the shear modulus and η is the effective viscosity. First-order finite differences in time are used to represent the objective co-rotational time derivatives of the visco-elastic stresses:

$$\frac{D\tau_{ij}}{Dt} = \frac{\tau_{ij} - \tau_{ij}^{old}}{\Delta t}. \quad (5)$$

Formatted: Kern at 16 pt

Formatted: Heading 1, Space Before: 24 pt

The effective viscosity η is calculated from the non-Newtonian dislocation creep equation:

$$\eta_{disl} = 0.5 \cdot \frac{1}{A_D} \cdot \sigma_{II}^{(1-n)} \cdot \exp\left(\frac{Q}{RT}\right), \quad (6)$$

where

$$255 \quad \sigma_{II} = \sqrt{\frac{1}{2} \tau_{ij}^2}, \quad (7)$$

and R , A_D , n , and Q are the gas constant, the pre-exponential factor, the power-law constant, and the thermal activation energy, respectively. Elastic behavior is achieved by updating the effective viscosity in response to an elastic time step (Maxwell time; $\Delta t_e = 1000$ years) and the stress history (Gerya and Yuen, 2007; Moresi et al., 2007; Moresi et al., 2003). The updated visco-elastic deviatoric stresses are defined by:

$$260 \quad \tau_{ij} = 2\eta \dot{\epsilon}_{ij} Z + \tau_{ij}^{old} (1 - Z), \quad (8)$$

where

$$Z = \frac{\Delta t_e G}{\eta_{disl} + \Delta t_e G}. \quad (9)$$

The effective viscosity is multiplied by this visco-elastic factor (Z) to obtain a numerical viscosity η_{num} to be used for solving the set of equations:

$$265 \quad \eta_{num} = \eta_{disl} \cdot Z = \frac{\eta_{disl} \Delta t_e G}{\eta_{disl} + \Delta t_e G}. \quad (10)$$

Plastic failure occurs if the second invariant of the visco-elastic stress tensor exceeds the yield stress, σ_y , following a pressure-dependent Drucker-Prager criterion:

$$\sigma_y = p \cdot (1 - \lambda) \cdot \sin \varphi + c \cdot \cos \varphi, \quad (11)$$

270 where p is the pressure (mean stress), and c , φ , and λ , are the cohesion, friction angle, and the fluid pressure ratio of the bulk material, respectively. The components of stress (σ_{xx} is the normal stress component, and σ_{xy} is the shear stress component) and the viscosity are then updated as:

$$\left. \begin{aligned} \sigma_{xx}^{new} = & \begin{cases} \frac{\sigma_x}{\sigma_H}, & \text{if } \sigma_H > \sigma_{yield} \\ \sigma_{xx} \frac{\sigma_y}{\sigma_{II}}, & \text{if } \sigma_H > \sigma_{yield} \\ \sigma_{xx}, & \text{if } \sigma_H \leq \sigma_{yield} \end{cases} \end{aligned} \right\} \sigma_{xx} \frac{\sigma_y}{\sigma_{II}}, \quad \text{if } \sigma_H > \sigma_{yield} \\ \left. \begin{aligned} \sigma_{xx}^{new} = & \begin{cases} \sigma_{xx}, & \text{if } \sigma_H \leq \sigma_{yield} \end{cases} \end{aligned} \right\} \sigma_{xx}, \quad \text{if } \sigma_H \leq \sigma_{yield} \end{aligned} \quad (12)$$

$$275 \quad \left. \begin{aligned} \sigma_{xy}^{new} = & \begin{cases} \frac{\sigma_{xy}}{\sigma_H}, & \text{if } \sigma_H > \sigma_{yield} \\ \sigma_{xy} \frac{\sigma_y}{\sigma_{II}}, & \text{if } \sigma_H > \sigma_{yield} \\ \sigma_{xy}, & \text{if } \sigma_H \leq \sigma_{yield} \end{cases} \end{aligned} \right\} \sigma_{xy} \frac{\sigma_y}{\sigma_{II}}, \quad \text{if } \sigma_H > \sigma_{yield} \\ \left. \begin{aligned} \sigma_{xy}^{new} = & \begin{cases} \sigma_{xy}, & \text{if } \sigma_H \leq \sigma_{yield} \end{cases} \end{aligned} \right\} \sigma_{xy}, \quad \text{if } \sigma_H \leq \sigma_{yield} \end{aligned} \quad (13)$$

$$\eta_{num} = \frac{\sigma_y}{2\dot{\epsilon}_{II}}, \quad \text{if } \sigma_{II} > \sigma_{yield}, \quad (14)$$

where

280 $\dot{\epsilon}_{II} = \sqrt{\frac{1}{2} \dot{\epsilon}_{ij}^2}$. (15)

The viscosities, including the visco-elastic effect and the plastic failure, are calculated on the Lagrangian markers and interpolated onto the Eulerian nodes using a distance-weighted scheme. To ensure numerical stability, they are capped by lower and upper cutoffs of 10^{17} and 10^{25} Pa·s, respectively.

3.3 Initial geometrical setup

285 The model domain is defined by a box of 500 km width and 60 km height (Fig. 3a). The Eulerian grid consists of 1001 × 121 nodes, with a nodal resolution of 500 × 500 m. The initial marker distribution defines, from bottom up, 1) a 30 km crustal basement layer, 2) a mechanically-weak salt horizon emplaced between 100 km < x < 450 km, 3) a 3-km-thick Palaeozoic sequence based on the stratigraphy of the Fars Arc, and 4) a 25-km-thick low-density and low-viscosity sticky-air layer to simulate a free surface, allowing for the vertical growth of the evolving fold-and-thrust belt (Crameri et al., 2012). The thickness of the salt layer varies in different models in order to examine its impact on the structural evolution. Some experiments include three inherited basement faults located at x = 100, 200, and 350 km (top of the basement), dipping at an angle of 50° at their top towards the hinterland (Mouthereau et al., 2006). These weak zones may either have a listric or planar geometry. The rheological parameters used in the experiments are provided in Table 1. The initial temperature field is characterized by a linear temperature increase with depth, starting from 0 °C at the interface between rock and sticky-air and reaching 600 °C at the bottom, in agreement with a constant geothermal gradient of 20 °/km inferred from paleothermal data (Aldega et al., 2018) and a relatively thick lithosphere (~200 km) beneath the Zagros (Jiménez-Munt et al. 2012; Priestley et al. 2012; Tunini et al., 2014). Each

Formatted: Kern at 16 pt

Formatted: Heading 1, Space Before: 24 pt

Table 1: Applied rheological parameter

Rock Type	$A_D(Pa^{-n})$	$Q(KJ mol^{-1})$	$n(-)$	$\rho(kgm^{-3})$	$\phi(^{\circ})^{\dagger}$	$c(MPa)^{\dagger}$	$\lambda(-)$	$k(Wm^{-1}K^{-1})$	$C_p(m^2K^{-1}s^{-2})$
Sticky-air	10^{-17}	-	1	1	-	-	-	200	3×10^6
Sediments ^a	5×10^{-18}	154	2.3	2500	10	1 (0.1)	0.4	2.5	1×10^3
Svn-rift sediments ^a	5×10^{-18}	154	2.3	2500	30 (20)	1 (0.1)	0.4	2.5	1×10^3
Paleozoic series ^a	5×10^{-18}	154	2.3	2700	30 (20)	1 (0.1)	0.4	2.5	1×10^3
Salt layer	Non-linear ^b	1.82×10^{-20}	32.4	5	2200	-	-	2.5	1×10^3
	Linear	10^{-18}	-	1	2200	-	-	2.5	1×10^3
Basement rock ^c	6.31×10^{-20}	276	3.05	2800	30 (20)	10 (0.1)	0.4	2.5	1×10^3
Basement fault ^c	6.31×10^{-20}	276	3.05	2800	10	0.1	0.4	2.5	1×10^3

^a Quartzite (Ranalli and Murphy, 1987; Stöckhert et al., 1999)

^b Rock salt (Li and Urai, 2012)

^c Diabase (Wilks and Carter, 1990)

[†] Values in brackets indicate strain weakened value.

Eulerian cell initially contains 16 randomly distributed Lagrangian markers carrying rock information and properties.

Table 1: Applied rheological parameter

Rock-Type	$A_D(\text{Pa}^{-n})$	$Q(\text{K} \cdot \text{mol}^{-1})$	$n(-)$	$\rho(\text{kgm}^{-3})$	$\varphi(^{\circ})^{\pm}$	$C(\text{MPs})^{\pm}$	$\lambda(-)$	$k(\text{wm}^{-1}\text{K}^{-1})$	$C_p(\text{m}^2\text{K}^{-1}\text{s}^{-2})$	
Sticky-air	10^{-17}	-	+	+	-	-	-	200	3×10^6	
Sediments*	5×10^{-16}	154	2.3	2500	10	1(0.1)	0.4	2.5	1×10^3	
Syn-rift sediments*	5×10^{-18}	154	2.3	2500	30(20)	1(0.1)	0.4	2.5	1×10^3	
Paleozoic series*	5×10^{-16}	154	2.3	2700	30(20)	1(0.1)	0.4	2.5	1×10^3	
Salt layer	Non-linear ^b	1.82×10^{-20}	32.4	5	2200	-	-	-	2.5	1×10^3
	Linear	10^{-18}	-	+	2200	-	-	-	2.5	1×10^3
Basement-rock ^c	6.31×10^{-20}	276	3.05	2800	30(20)	10(0.1)	0.4	2.5	1×10^3	
Basement-fault ^c	6.31×10^{-20}	276	3.05	2800	10	0.1	0.4	2.5	1×10^3	

*Quartzite (Ramalli and Murphy, 1987; Stöckhert et al., 1999)

† Rock salt (Li and Urni, 2012)

*Diabase (Wilks and Carter, 1990)

‡ Values in brackets indicate strain-weakened value.

3.4 Boundary conditions

The numerical experiments simulate the deformation of the NE margin of the Arabian Plate during Permian–Triassic rifting and Oligocene–recent continental collision. ~~The phase of tectonic quiescence between rifting and convergence is ignored here for simplification of the model setup. Tectonic quiescence affects the thermal state and, consequently, the rheological behaviour of the rocks. However, we assume that the temperature field reached a steady state by the onset of the collision phase to simplify the model setup. This implementation allows us to effectively capture the relevant thermal conditions while maintaining focus on the critical dynamics of the extension and collision phases.~~

The velocity boundary conditions are prescribed in a way to simulate the tectonic inversion of the rifted margin, including initial extension and subsequent convergence. During the rifting phase, an outward horizontal boundary velocity of $v_x = 5$ mm/yr is applied on the right side of the model domain, while the left boundary is kept fixed horizontally (Fig. 3a). The bottom boundary has zero vertical velocity, and the top boundary has an incoming vertical velocity of $v_y = 0.6$ mm/yr, ensuring the conservation of volume within the model domain (Fig. 3a). On all boundaries free-slip conditions (zero shear stress) are prescribed. The finite Eulerian domain impedes elastic bending of the lower boundary, affecting resulting surface taper angles compared to the ~~Zagros fold and thrust belt~~ZFTB (Mouthereau et al., 2006; McQuarrie, 2004). The extension phase is applied for a period of 5 Myr, resulting in a total extension of 25 km. During this period the rift basin is allowed to be filled with syn-rift deposits (equivalents of the Dashtak and Dehram formations). Following the extension phase, 4 km of sediments representing the depositional environment during the tectonic quiescence and subsidence period are added onto the existing stratigraphy (Fig. 3b). These sediments represent the Khami and Bangestan groups, and the Gurpi and Pabdeh formations (Fig. 2). The Late Cretaceous ophiolite obduction episode has affected only some parts of the Arabian margin, namely the Kermanshah, Neyriz, Hajiabad and Oman regions. We have not included the Late Cretaceous deformation in our modeling, as it accounts for just a few percent of the observed shortening in the ZSFB (e.g. Saura et al., 2011).

Crustal shortening in the experiments is simulated by imposing a horizontal velocity of 1 cm/yr at the bottom and the left side of the domain. The right-side acts as a partial backstop by allowing the basement part of the crust to escape the model but preventing the sedimentary cover from exiting. For the lower part of the right boundary (basement), different horizontal velocities between 0 and 1 cm/yr are tested (Fig. 3b). This arrangement permits us to vary the degree of basement involvement and to study the development of thin-skinned and thick-skinned tectonics in the model. The compressional phase ran for 15 Myr, accommodating a total shortening of 25% (i.e., 125 km). This amount offsets the 25 km extension during the extensional phase and the 20% shortening during the collisional phase, comparable to estimates for the SE Zagros (McQuarrie, 2004; Motamedi et al., 2012; Pirouz et al., 2017; Najafi et al., 2021).


3.5. Surface processes

335

The surface processes of sedimentation and erosion during the evolution of the models are simulated by applying diffusion of the rock/air interface. We used the diffusion equation:

$$\frac{\partial h_s}{\partial t} = \kappa K \frac{\partial^2 h_s}{\partial x^2}, \quad (16)$$

where h_s is the surface topography, t is time, κK denotes a diffusion constant ($\kappa K = 10^{-6} \frac{m^2}{s}$), and x is the horizontal coordinate. The left and right sides of the surface line prescribe free-slip boundaries.



340

Figure 3: Setup and boundary conditions of numerical modeling. (a) Initial compositional setup for extension phase with listric faults. (b) Fully extended model (5 Myr) with listric faults before convergence. v_x , v_y and v_{x-b} is horizontal velocity, vertical velocity, and basement horizontal velocity, respectively.

4. Results of modeling

345

A total of 11 experiments are presented, investigating the role of basement involvement and the impact of salt rheology and thickness on the structural evolution of the Fars Arc. Experiments are divided into four series varying in 1) the thickness of the salt layer, 2) the rheology of the salt layer, 3) the existence and geometry of pre-existing basement faults, and 4) the degree involvement of the basement during convergence (Table 2). First, we present the temporal evolution of the reference model, to which the rest of the models are compared. All experiments underwent 5 Myr of extension, followed by 15 Myr of compression. [A link to videos \(Graphics Interchange Format\) of all numerical experiments can be found in Data Availability.](#)

350

Table 2: List of the numerical models

Model	Salt thickness (km)	Salt rheology	Fault geometry	Basement velocity (mm s ⁻¹)	Figure
Model 1 (Reference model)	2	Non-linear	listric	0	4,5,8,11,12,14, S1
Model 2	0	Non-linear	listric	0	6,8,12, S2
Model 3	4	Non-linear	listric	0	6,8, 12,13, S3
Model 4	2	Linear (10 ¹⁸ Pa·s)	listric	0	7,8,12, S4
Model 5	2	Linear (10 ²⁰ Pa·s)	listric	0	7,8, 12, S5
Model 6	2	Non-linear	planar	0	9,12, S6
Model 7	2	Non-linear	no fault	0	9,12, S7
Model 8	2	Non-linear	listric	0.25	10,12, S8
Model 9	2	Non-linear	listric	0.5	10,12, S9
Model 10	2	Non-linear	listric	0.75	10,12, S10
Model 11	2	Non-linear	listric	1	10,12, S11

Formatted Table

360 4.1 Evolution of the reference model

The reference model (Model 1) has an initial 2-km-thick salt horizon with a power-law viscous rheology (Table 1 and 2). The basement exhibits three inherited listric faults and is shortened at the same rate as the sedimentary cover during convergence. The extensional phase is characterized by the development of half-graben basins forming along the listric basement faults that are filled by syn-extensional deposits (Fig. 4a-c [and S1](#)). The second invariant of the strain-rate tensor shows that the fastest deformation occurs along the pre-existing weak zones. During extension, all three inherited faults are simultaneously active, without any recognizable preference. Normal fault deformation along the basement faults propagates upward into the post-salt strata inducing typical half-grabens and bending of the hanging wall (Fig. 4c [and S1](#)).

365 After 5 Myr of compression, the cover sequence mainly deforms between the right backstop and the salt pinch-out at $x \approx 150$ km (Fig. 4d [and S1](#)). The basement fault closest to the backstop has experienced a significant amount of reverse motion and has caused a large step of several ~~kilometers~~[kilometres](#) in the salt décollement geometry. Faulting

of the sedimentary cover mainly develops where the syn-rift strata are the thinnest, i.e. in front of the basement thrusts. Strain localization occurs along the basement faults, and in the case of the frontal fault ($x \approx 120$ km) it cuts across the strata overlying the salt layer (Fig. 4d and S1). Towards the backstop, the deformation in the basement and the cover sequence is decoupled along the weak salt horizon.

As convergence progresses (10 Myr), the basement fault at $x \approx 300$ km undergoes inversion and creates a ramp-flat-ramp geometry (Fig. 4e and S1). Basement fault propagation deforms the syn-extensional strata and forms harpoon-like anticlines within the sedimentary cover. The syn-tectonic sedimentation concentrates within multiple structural basins between the developing anticlines. The salt layer decouples the deformation between the basement faulting and the sedimentary cover folding, allowing for more amplification within the sedimentary cover. Strain rates indicate intense deformation within the basement, which particularly localizes along the inverted normal faults, as well as the basal décollement level. The strain rate pattern starts to gradually diffuse below ~ 26 km depth, indicating the transition from brittle to ductile deformation in the lower crust (Fig. 4f) and S1).




Figure 4: Temporal evolution of the reference model during extensional (a-c) and compressional (d-f) phases. For each time panel, the compositional layers (top) derived from the Lagrangian markers, and the second invariant of the strain-rate tensor (bottom) are shown.

Patterns of the second invariant of the stress tensor after the extensional phase and after full convergence indicate an increase in stress with depth down to ~ 30 km ($y = 50$ km), where the brittle-to-ductile transition begins (Fig. 5a,d). The lower part of the basement displays low stresses given its lower viscosities and ductile nature (Fig. 5b,e). Viscosity plots furthermore indicate the position of the low-viscous décollement and basement thrusts.

Figure 5c illustrates the vertically averaged viscosity of the non-Newtonian salt horizon after 25 km of extension, influenced by both temperature and strain rate (Table 1; Li and Urai, 2012). During extension and the formation of half-grabens along pre-existing faults, the strain rate and temperature increase in these areas. Consequently, these

regions exhibit the lowest viscosities (10^{18} Pa·s), corresponding to the locations of the basement faults (Fig. 5c). In areas without strain localization within the cover sequence, the viscosity shows larger values (10^{19} - 10^{20} Pa·s). After 5 Myr of extension, the half-graben basins are covered by post-rift deposits, prior to the onset of convergence. After 15 Myr, the intensity of thin-skinned folding within the sedimentary cover increases (Fig. 5f). The development of long-wavelength folds is influenced by the reactivation of basement faults, which exert a significant impact on the deformation of the upper crustal region. The frontal basement fault is influenced by the salt pinch-out and over-thrusts the post-rift strata, whereas the other two faults undergo a transition to ramp-flat-ramp configuration. In the post-rift sediments, new shear zones are formed along the continuation of the pre-existing faults. These faults have been displaced due to the presence of basal salt, leading to deviation from their original dip. Strain rates indicate that the continuity of the salt décollement is disrupted due to the offset introduced by the inherited faults (Fig. 4f). The largest viscosities of salt are found in the anticlines with long wavelengths that form in the areas between two pre-existing faults (Fig. 5f). In contrast, lower viscosities are observed in areas where the strain rate reaches its maximum, specifically at the tip of the basement faults where the salt layer is thin.

410




Figure 5: Second invariant of the stress tensor and viscosity after full extension (a, b) and full convergence (d, e). c, f) Vertically-averaged viscosity of the nonlinear salt horizon after 5 Myr of extension (c) and 15 Myr of convergence (f), respectively.

415 **4.2 Effect of salt thickness**

Figure 6 shows the effect of the thickness of the salt layer on the deformation. In the absence of a salt layer (Model 2; Fig. 6a), no mechanical decoupling occurs between the sedimentary cover and the basement. The extension phase is dominated by the pre-existing faults and their geometry, and the majority of the deformation accumulates along the pre-existing normal faults. After 15 Myr of convergence, long-wavelength basement-cored anticlines develop in the sedimentary cover. The anticlines form asymmetrically and with low internal deformation of the sedimentary sequence (Fig. 6b and S2). The folding style within the sedimentary cover is determined by the pre-existing faults, resulting in the development of thrust faults that propagate upward through the overlying rocks. Basement faults cut the

420

sedimentary cover and reach the surface without any significant deviation from their initial orientation. Furthermore, backthrusts form in the hanging wall as conjugates to the reactivated inherited faults. Strain rate patterns reveal that the entire crustal package deforms uniformly, without mechanical decoupling along the different lithological layers. Increasing the salt layer thickness to 4 km (Model 3) does not significantly affect the structural development during the extension phase compared to the reference model (Fig. 4c and 6c). However, a thicker salt layer hinders the extensional faults from fully cross-cutting the salt after full extension (Fig. 6c and S3). During shortening, a thicker salt layer acts as a more efficient décollement, impeding small-scale deformation in the sedimentary cover relative to the reference model (Fig. 6d; compare to Fig. 4f). The sedimentary cover develops décollement folds cored by a thickened salt. The cover is pushed over the frontal basement fault on the left, which upon reactivation results in a salt-cored fault-propagation fold at $x \approx 210$ km. Furthermore, the increased amount of salt allows for vertical breakthroughs and diapir formation.




Figure 6: Composition inferred from Lagrangian markers and the second invariant of the strain-rate tensor, affected by salt thickness after 5 Myr of extension and 15 Myr of convergence. (a-b) In the absence of salt layer. (c-d) In the presence of a salt layer with 4 km thickness.

4.3 Effect of salt rheology

To investigate the impact of the rheology of the salt on the style of deformation, we considered two models with linear viscosity for the salt layer and compared the results with that of the reference model. For Models 4 and 5, we chose linear viscosities of 10^{18} and 10^{20} Pa·s, respectively, which are roughly equal to the lowest and largest viscosities observed in the reference model (Fig. 5c, f). After 5 Myr of extension, the low-viscosity Model 4 leads to the formation of symmetric ridges along the half-grabens and footwalls (Fig. 7a and S4). The majority of the deformation is accommodated by the salt layer and the inherited faults. After 15 Myr of compression, the pattern of folds within the sedimentary cover diverges from that of the reference model, as they display an inclination towards the hinterland (Fig. 7b and S4). Based on the strain rates, deformation is accommodated across the décollements.


In Model 5 with higher salt viscosity, the structural pattern after 5 Myr of extension resembles that of the reference model (Figs. 4c and 7c). Strain rates reveal significant deformation occurring within the basement, localizing along

450

the pre-existing weak zones. After 15 Myr of shortening, the style of folding shows significant differences from Model 4 (Figs. 7b, d; and S5). Several forethrusts develop above the imbricated basement blocks and backthrusts are generally absent. Strain rates indicate that the décollement remains laterally connected across the different basement blocks, whereas it is interrupted in the model with low linear viscosity (compare Fig. 7b and 7d).


455

Figure 8 presents the vertically-averaged strain rate within the salt layer across models with different décollement rheology. The influence of a power-law rheology on strain rate is characterized by a non-linear, accelerating response to stress. As stress levels increase, the strain rate increases more rapidly compared to a linear rheology, where the strain rate maintains a constant, linear relationship with stress. The mean strain rate in a salt layer with a power-law rheology exceeds that in a salt layer with linear rheology, as illustrated.



460

Figure 7: Composition inferred from Lagrangian markers and the second invariant of the strain-rate tensor, affected by different viscosity for linear salt layer after 5 Myr of extension and 15 Myr of convergence. (a-b) linear salt horizon with 10^{18} Pa·s viscosity. (c-d) linear salt horizon with 10^{20} Pa·s viscosity.



465

Figure 8: Vertically-averaged strain-rate over model width for models with variable salt rheology after 15 Myr of shortening.

4.4 Effect of basement fault geometry

Figure 9 shows two models, one with planar basement faults (Model 6) and one with no basement faults (Model 7). After 5 Myr of extension, the basement faults maintain a constant dip angle, resulting in the formation of conjugate normal faults rooting in the lower extent of the pre-existing faults (Fig. 9a and S6). Similar to the case with listric faults (Fig. 4c), the second invariant of the strain-rate tensor indicates high concentration of deformation through the development of fractures within the sedimentary layers above the salt horizon, specifically at the back of the footwall of the faults (Fig. 9a and S6). After 15 Myr of compression, multiple overturned forwards and backward verging folds develop within the sedimentary cover at $x \approx 350$ km (Fig. 9b and S6). These folds originate in relation to the activation of thrusts in the underlying basement. Notably, the planar faults play a pivotal role in the formation of backthrust faults. The principal strain rate is concentrated within vulnerable zones such as the salt layer, planar faults, and the newly formed backthrust faults within the basement (Fig. 9b and S6). Basement faults crosscut the salt décollement to form bypass thrusts.

In the absence of inherited faults (Fig. 9c and S7), no zone of significant strain accumulation is formed during the extension phase. The maximum strain rate accumulates primarily along the fractures caused by extension, while the basement is decoupled from the overlying sedimentary strata. After 15 Myr of shortening, the basement has undergone intense deformation by thrusting (Fig. 9d and S7). Near the backstop wall ($x = 350\text{--}500$ km), it is deformed into pop-up structures caused by conjugate thrusts, while forward verging thrusts dominate towards the foreland. The overlying sedimentary sequence is mainly shortened by décollement folding/faulting between $x = 200\text{--}350$ km. A large pop-up structure ($x \approx 230$ km) without mechanical decoupling between the basement and the sedimentary cover develops in front of the salt pinch-out (Fig. 9d and S7). The strain rates indicate that at the late stage of shortening, a thin-skinned style of deformation is taking place in front of the salt pinch-out and the associated basement thrust.





Figure 9: Composition inferred from Lagrangian markers and the second invariant of the strain-rate tensor, affected by different type of inherited faults after 5 Myr of extension and 15 Myr of convergence. (a-b) In the presence of planar inherited faults. (c-d) In the absence of any inherited faults.

4.5 Effect of basement shortening rate

The effect of basement involvement during convergence is implemented by varying the horizontal velocity of basement rocks exiting the right-side boundary, v_{x-b} (Fig. 3b). All of the previous models exhibit a rigid backstop over the entire right side that prevented the basement material from leaving the model domain ($v_{x-b} = 0$), resulting in 100% involvement of the basement in the deformation. In this section, we show a set of models, in which the basement is allowed to exit the model domain with a velocity that is a varying fraction of that on the left boundary (Fig 10). The basement will experience a slower rate of shortening with respect to the cover. As a result, the degree of thin-skinned deformation will increase.

With a basement involvement of 75% ($v_{x-b} = 0.25 \cdot v_x$), the two faults located on the right side of the model transform into flat-ramp-flat structures (Model 8; Fig. 10a and S8). Above the basement blocks, the sedimentary cover undergoes thrusting and the cover sequence overthrusts the frontal fault zone by ~50 km. If the basement shortening rate is 50% of the cover shortening rate ($v_{x-b} = 0.5 \cdot v_x$), the faults situated on the right side of the model remain continuous and undergo significant displacement (Model 9; Fig. 10b and S9). After 15 Myr of convergence, the fault on the right side partly exits the model domain. Additionally, strain rate indicates that most deformation localizes within the salt layer, the pre-existing basement faults, and faults within the sedimentary cover. If the basement is shortened by a rate that is involvement of 25% of the cover sequence shortening ($v_{x-b} = 0.75 \cdot v_x$), the fault on the right side almost entirely left the model domain, and the other two faults experience lesser degrees of involvement, roughly displaying 100% of tectonic inversion (Model 10; Fig. 10c and S10). In comparison to the previous cases, deformation in the sedimentary cover is more distributed laterally, not depending as strongly on the location of basement faults. Furthermore, the salt layer remains connected and forms a nearly-horizontal décollement horizon (Fig. 10c). In the case the basement exits the model domain with the same velocity as the bottom boundary ($v_{x-b} = v_x$), inherited basement faults are not reactivated, and the sedimentary cover deforms in a thin-skinned tectonic style (Model 11; Fig. 10d and S11). The second invariant of the strain-rate tensor demonstrates that the salt horizon acts as a décollement layer with basement steps that developed during the rifting phase. The resulting thin-skinned fold-and-thrust belt exhibits a flat surface taper given the weak salt rheology (Fig. 10d).



Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Figure 10: Composition inferred from Lagrangian markers and the second invariant of the strain-rate tensor, affected by different values of basement velocity after 15 Myr of convergence. (a) $v_{x-b} = 0.25$, (b) $v_{x-b} = 0.5$, (c) $v_{x-b} = 0.75$ and (d) $v_{x-b} = 1$.

520 5 Discussion

The models presented in the previous sections demonstrate the impact of weak zones, in the form of décollement layers and pre-existing basement faults, and the degree of involvement of the basement on the structural evolution and reactivation of inherited structures during tectonic inversion. In the following, we will provide insight into the effects of the implemented variables on the dynamic strength of the basement and strain localization within it. Furthermore, numerical results are compared to previous modeling attempts and natural examples from the Zagros fold-and-thrust belt (ZFTB).


5.1 Strength and localization of deformation within the basement


The mechanical strength of fold-and-thrust belts and the partitioning of strain during their growth has been a topic of interest for many decades (Chapple, 1978; Davis et al, 1983; Stockmal et al., 2007; Tavani et al., 2015). During continental collision and the emergence of mountain belts, plate convergence rates usually decrease step wise given the increasing strength of the affected plate boundary (Wortel et al., 2009). Weak zones ~~is within~~ the deformed rock ~~pile define~~ layers play a crucial role in shaping the structural evolution of ~~evolving~~ fold-and-thrust belts. ~~According to~~ In the presented case of the Zagros, numerical experiments ~~demonstrate that~~ both the sub-horizontal salt horizon and inherited weak basement faults ~~strongly affect strain~~ significantly influence the partitioning ~~in the Zagros,~~ eventually. ~~Independent of strain.~~ Regardless of whether the viscous décollement ~~is able to~~ can fully decouple the upper ~~from the~~ and lower crust mechanically, the basement ~~has~~ exerts a strong ~~effect~~ influence on the overall structural evolution of the Zagros fold-and-thrust belt, ~~which has to.~~ This must be taken into consideration ~~considered~~ when ~~attempting the construction of~~ constructing structural cross sections.

In fold-and-thrust belts with local weak zones in form of décollement layers and reactivated inherited faults, strength may vary across strike. For example, the reference model (Model 1) displays a stress distribution with maximum values in the brittle part of the basement, which is crosscut by hinterland-dipping faults (Fig. 11a). To illustrate across strike variations in strength, three vertical stress profiles are shown representing different segments of the evolving fold-and-thrust belt: i) At $x = 150$ km, basement faults and a salt décollement are absent, and the strength profile exhibits a simple form with a brittle ($y \approx 15$ – 40 km) and ductile ($y \approx 40$ – 60 km) part (Fig. 11b). ii) The second profile at $x = 320$ km crosscuts both salt décollement ($y \approx 18$ km) and a basement fault ($y \approx 40$ km) that strongly affect the strength profile (Fig. 11c). While the salt horizon shows very low stresses of only a few megapascal, stresses across the basement fault are reduced to ~ 150 MPa, in contrast to maximum stresses of ~ 600 MPa around the brittle-ductile transition. iii) Close to the backstop, at $x = 460$ km, maximal stresses increase to ~ 800 MPa at $y \approx 40$ km (Fig. 11d). At $y \approx 18$ km, the salt décollement displays an effective decoupling horizon, and an additional weak zone at $y \approx 27$ km is defined by flat ramp of a basement sliver stack. To infer the force acting across the fold-and-thrust belt related to the reference model, the differential stress profiles at each x -coordinate are integrated over the vertical distance. For the reference model after 15 Myr of convergence, values range in between ~ 7 – 15 TN/m, where the lowest values


coincide with the occurrence of basement faults (Fig. 11c). In general, there is no clear trend observable that would refer to a weakening or strengthening related to deformation, as crustal thickening (strengthening) goes hand in hand with basement thrusting (weakening).

Figure 12 illustrates shows the temporal evolution of the average force (e.g., average value of Fig. 11e for the reference model) for a variety of models to identify key parameters that affect the stress state of continental collision zones. All models show similar boundary forces during the extension phase of ~ 4 TN/m (Fig. 12), except Model 7 without pre-existing basement faults, which exhibits values of ~ 4.2 TN/m (Fig. 12b). Similar force values during extension for models with planar and listric inherited basement faults indicates that none of the tested fault geometries would preferably localize. During the convergence phase, boundary forces quickly increase, and most of the models show a general increasing trend from ~ 6 – 8 TN/m to ~ 10 – 12 TN/m over a time span of 15 Myr (Fig. 12). Minor increase and decrease in force are observed related to the absence (Model 2) and increased thickness of the salt horizon (Model 3), respectively, while the strength (viscosity) of the salt has no detectable effect (Fig. 12a). The absence and geometry of basement faults displays a significant importance during convergence, in contrast to the extension phase, where planar preexisting faults increase the necessary boundary force by ~ 1 – 2 TN/m, and no faults lead to another increase of ~ 1 – 2 TN/m during the first 10 Myr of convergence (Fig. 12b). More important is the effect of basement involvement in crustal shortening, with decreasing involvement resulting in decreasing boundary force (Fig. 12c). This illustrates the importance of an effective décollement level that is able to decouple an upper crust with distributed deformation from a basement that is underthrust towards the suture, where it deforms more intensely (Tavani et al., 2015; Pfiffner et al., 2002). A similar pattern of boundary force evolution was also observed from analogue models, however, with more acute variations for specific internal deformation of the compressed sand pile (McBeck et al., 2018).





575 **Figure 11:** (a) strain tensor superimposed on differential stress tensor. (b-d) Differential stress profiles of front of the deformation front, with decollement and basement fault, and without basement fault involved respectively. (e) The integrated stress profile for reference model.



580 **Figure 12: Horizontally-averaged boundary force over time. (a) Models with variable salt thickness and salt rheology. (b) Models with variable fault geometry. (c) Models with variable basement shortening rate. Model characteristics are listed in Table 2.**

5.2. Comparison with numerical modelling studies

585 The mechanics of fold-and-thrust belts has been investigated by means of numerical methods since the 1980's (e.g., Borja and Dreiss, 1989). Rapidly evolving computing capabilities led to a vast increase in geodynamic numerical modelling studies that provide an invaluable contribution towards our understanding of ~~geotectonic~~ **Geotectonic** systems (e.g., van Zelst et al., 2021; Gerya, 2022; Ismail-Zadeh and Tackley, 2010). Previous numerical models of fold-and-thrust belts ~~often have typically~~ focused on thin-skinned tectonic systems, investigating ~~a wide range of~~ parameters ~~and their effects on the structural evolution,~~ such as wedge and décollement strength, ~~wedge and décollement~~ rheology, surface processes, ~~and~~ mechanical stratigraphy, ~~beside others~~ (e.g., Stockmal et al., 2007; 590 Buiter et al., 2016; Simpson, 2011; Burbidge and Braun, 2002; Ruh et al., 2012). ~~The consideration~~ **However, the inclusion** of rifting prior to collision ~~represents adds~~ a new ~~aspect~~ **dimension** that ~~is commonly ignored~~ **has often been overlooked** in ~~previous these~~ studies ~~related to the Zagros fold-and-thrust belt.~~ This approach ~~provides allows for~~ a more comprehensive understanding of tectonic evolution by ~~allowing the investigation of~~ **examining how** pre-collisional structural configurations ~~and their influence on subsequent~~ **later** deformation patterns. Including rifting 595 before convergence is crucial as it sets the initial conditions that significantly impact structural evolution during collision (Buiter and Pfiffner 2003; Ruh et al., 2018; Granado and Ruh, 2019). This approach offers several advantages: It provides insight into structural inheritance, as rifting creates pre-existing weaknesses and fault systems that play a crucial role during subsequent compressional phases. Our model demonstrates how these inherited structures influence strain localization and deformation styles, providing insights into the evolution of complex geological features such as the Fars Arc. In the following, we discuss several numerical studies concerning tectonic 600 systems that are similar to the one presented in this study to set our results into perspective and to identify generally applicable observations.

Buiter and Pfiffner (2003) applied a two-dimensional, visco-plastic numerical model in order to evaluate the dynamics of tectonic inversion of a series of half-graben basins upon compression. They reported syn- and post-rift sediment 605 uplift, accompanied by basement block rotation, and the emergence of newly formed shear zones in the post-rift sequence, originating from basin-bounding faults. Weak sediments at the basin base contribute to the generation of basement shortcut faults. Furthermore, their study illustrates the predominant development of back thrusts as conjugates to listric basin-bounding faults. The structural evolution observed in our models aligns with their research findings, demonstrating a similar pattern of strain localization within syn-rift and post-rift layers, mainly initiated by deformation along pre-existing underlying faults. Additionally, to their findings, our results indicate that the formation 610 of back thrusts is mainly associated with the inversion of planar faults (Fig. 9; Model 6).

Nilforoushan et al. (2013) presented two-dimensional thermo-mechanical experiments of thick-skinned fold-and-thrust belts with salt décollements. They demonstrated that the geothermal gradient and the mineralogy of the basement (ductile flow law) strongly affect the geometry and reactivation of inherited basement faults. Furthermore, 615 they underlined the importance of a weak salt horizon at the base of the sedimentary succession for the mechanical decoupling and formation of a thin-skinned upper-crustal fold-and-thrust belt, resulting in a variability in the rate of

shortening between the cover and the basement. Their findings corroborate the hypothesis of Molinaro et al. (2005) that in the ~~Zagros Simply Folded Belt~~ZSFB, greater shortening occurs within the cover compared to the basement. From our results examining the effects of the shortening rate of the basement on structural development, we observed the emergence of long-wavelength folds arising directly from the movements of faults within the basement (Figs. 11a-d). These findings align with natural occurrences and are corroborated by geological cross-sectional data, reinforcing the principal characteristics of the Fars arc that are typified by folds of extensive wavelengths. This agrees with the suggestions by Mouthereau et al. (2006), who proposed that early activation of basement deformation dictates the formation of the large-wavelength folds. Furthermore, Mouthereau et al. (2006) determined that the viscosity of Hormuz salt decreases significantly with increasing temperature, exhibiting Newtonian fluid behaviour under low stress and strain-rate conditions. In our results, the average viscosity of non-Newtonian salt is affected by both temperature and strain rate, where weak zones with the lowest strain rates and temperatures exhibit the largest viscosity values. The resulting values are ~~slightly~~ larger than what was proposed for the Fars arc resulting from Critical Wedge Modelling (Mouthereau et al., 2006).

Ghazian and Buitier (2014) investigated the impact of salt for the southeast ~~Zagros fold and thrust belt~~ZFTB by 2D thermo-mechanical models. Their findings revealed that the presence of a thick basal Hormuz salt effectively decouples overlying sediments from the basement and promotes the localization of deformation in the sediments. Our results align with these findings, demonstrating a significant influence of the salt layer on controlling topographic height and folding patterns (Fig. 4 and 6). The deformation style in both the basement and the cover sequence decouples along the weak salt horizon. As the salt thickness increases, this layer functions more effectively as a décollement, impeding deformation in the sedimentary cover sequence (Fig. 6).

Bauville and Schmalholz (2015; 2017) and Kiss et al. (2020) conducted 2D numerical experiments to investigate the transition from basement-involved thin-skinned to thick-skinned tectonics and the effects of tectonic inheritance on the development of nappe systems. The key parameters controlling this transition were found to be the viscosity ratios within the basement and between the basement and the sedimentary cover above it. Specifically, a higher ratio within the basement favors thick-skinned deformation, while a higher ratio between the basement and cover leads to thin-skinned deformation. As indicated by our study, 100% participation of the basement in deformation leads to thick-skinned deformation (Fig. 4), while non-participation of the basement to thin-skinned deformation (Fig. 10; Model 11). Their findings show the tectonic and structural inheritance plays a significant role in controlling the tectonic evolution and resulting structures in the ~~fold-and-thrust-belt belts~~. The geometry and magnitude of mechanical heterogeneities like varying basement-cover interface characterized by half-grabens and horsts, as well as the vertical alternation of sedimentary layers with different mechanical strengths, influence nappe formation. Their observations align with our findings that the presence of pre-rift salt and inherited faults, which serve as structural inheritances, influence the shape and geometry of the basin (Fig. 4,6,7). Moreover, their simulations and ours revealed that both linear and power-law viscous rheologies demonstrated ~~exhibited~~ similar features to more complex simulations, indicating the robustness of these findings (Fig. 7; Model 4 and 5).

Eslamrezaei et al. (2023) used numerical discrete element models to investigate the impact of mechanical stratigraphy, décollement layers, and the number and thickness of cover sequences on the structural evolution and strain partitioning

of thin-skinned fold-and-thrust belts. Their modeling outcomes revealed that weak décollements played a crucial role in decoupled deformation within the fold-and-thrust belt under continuous shortening. The study underscored that shortening was primarily accommodated by thrust-related folds, resulting in notable variations in structural styles. In alignment with our findings (Fig. 4, 6 and 10), the weak décollement layer, due to its characteristics, exhibited an inability to retain stress and undergo deformation, as evidenced by the increasing strain rate along the basal detachment (Fig. 4). Both their study's conclusions and our results emphasized that the structural style and decoupling in thin-skinned fold-and-thrust belts and formation of structures such as box folds and harpoon structures, are influenced by factors such as rheology, the number, and thickness of décollements (Fig. 6 and 7).

5.3 Comparison with the structural evolution of the Fars Arc

The tectonic style of the Fars Arc is characterized by a combination of thin-skinned salt-related folding and thick-skinned basement thrusting (i.e., Mouthereau et al., 2007a). The structure of the sedimentary cover is shaped by a series of detachment faults and fault-propagation folds, involving several salt diapirs rooted in the Hormuz salt (e.g., Jahani et al., 2007; Motamedi and Gharabeigli, 2019; Vergés et al., 2024). In addition, at least three high-angle reverse basement faults (High Zagros, Surmeh, and Mountain Front faults) are constrained based on their seismicity, structural relief, magnetic anomalies, and abrupt topographic changes (e.g., Jackson and Fitch, 1981; Berberian, 1995; Talebian and Jackson, 2004; Mouthereau et al., 2007b; Teknik and Ghods, 2017; Karasözen et al., 2019). The dynamic importance of these faults is supported by our results, as numerical models without basement faults (Model 7 in Fig. 9) display a marked difference in structural style with geological observations in the Fars Arc. In the absence of basement faults in the models, deformation is concentrated only in the outer part of the arc, in the form of an imbricate thrust system and a large mushroom anticline near the salt pinch-out. Furthermore, the importance of the Hormuz salt for the distinct structural style of the Fars Arc is corroborated by the numerical model without a basal salt layer (Model 2 in Figs. 6a and 6b). In that model, four basement-involved anticlines form across the entire belt, each with wavelengths of about 100 km, while in contrast, in the Fars Arc, the anticlines can be grouped into short-wavelength detachment-fold anticlines and long-wavelength basement-involved anticlines (e.g. Mouthereau et al., 2006). In the model without salt, all structures verge towards the front without any back-thrusting. This type of fold and faults geometry is not representative of the observed structures in the Fars Arc (e.g. Motamedi et al., 2012; Najafi et al., 2014; Jahani et al., 2017). Observational evidence thus highlights the essential role that the basement and the salt layer play in determining the deformation style of the belt.

The results of the experiment with a 4-km-thick basal salt layer (Model 3; Fig. 6c-d) demonstrate that a thicker salt layer interacts with the basement faults to form symmetrical anticlines containing salt cores and discordant diapirs. The topography of the Fars Arc cannot be solely attributed to the presence of a salt layer, the basement's role in structural development is also significant. As our results indicate, processes involving the basement should be considered as an essential factor in reconstructing and assessing the geological development of the region. The models demonstrate that the basement faults ramp up through the upper part of the basement and eventually flatten along the Hormuz salt level at a depth of around 10 km (Model 1 in Fig. 4e4f). Similar structures of basement faults detaching along evaporitic levels in the middle of the sedimentary cover, or reaching the surface to cut the forelimbs of their

690 hanging wall anticlines, are documented in the Fars Arc (Vergés et al., 2024), such as the Kangan, Asaluyeh and
Tabnak anticlines, located to the south of the cross-section in the foreland vicinity and within the Mountain Front
Fault (Berberian, 1995; Mouthereau et al., 2007b; Figs. 13b,c). These anticlines are characterized by a pronounced
concentric surface geometry (Fig. 13a).


A well-studied example of syntectonic growth strata is the Dowlatabad syncline, located approximately 40 km
695 northeast of the Persian Gulf coastline (Figs. 13a, b) and limited by the Sefid anticline to the north and the NW segment
of the Pazan anticline to the south (Najafi et al., 2021). The Dowlatabad syncline is infilled by more than 2 km of syn-
folding sediments. Numerical results from Model 3, which has a thick salt layer, compare well with the formation of
the Dowlatabad syncline (Figs. 13a,b): i) The forelimbs of Pazan and Sefid anticlines were faulted by foreland-directed
thrusts rising from the basal décollement, ii) surface processes resulted in a smooth topography, and iii) the surface
700 has an asymmetric shape, while at depth, the structure has the form of a box fold.

In the Zagros belt, there is a noted relationship between faulting patterns and the distribution of salt diapirs (Talbot
and Alavi, 1996; Hessami et al., 2001; Bahroudi and Koyi, 2003; Sherhati and Letouzey, 2004; Sepehr and Cosgrove,
2005; Jahani et al., 2017). Our numerical results indicate that salt diapirism may be triggered at two distinct stages in
the case of a thick (4 km) salt décollement. During the rifting phase, small-scale diapirs start to grow above the tip of
705 normal faults, partially piercing the sedimentary cover (Model 3; Fig. 6c). This procedure can be considered as a short-
lasting and locally-developed reactive diapirism stage (e.g. Jackson and Vendeville, 1994; Jackson and Hudec, 2017).

More significantly, the growth of two salt diapirs are observed during the compression phase in the inner zone of the
fold-and-thrust belt (Model 3 in Fig. 6d). ~~Fernandez and Kaus (2014) showed with numerical modeling that pre-
existing salt diapirs can significantly influence the pattern and growth of three-dimensional folds and fold patterns,
710 accelerating fold formation and localizing deformation, highlighting the important role of diapirism in structural
evolution during tectonic processes.~~

In contrast to décollement folds, these salt bodies have pierced the overburden
strata and show discordant contact with overlying sediments. The process involves the buoyant rise of underground
salt when subjected to pressure from convergent tectonic forces. These early-growing diapirs, squeezed by further
shortening, lead to the development of secondary salt welds. This type of shortened diapirs has recently been
715 documented in the High Zagros zone in a field-based study by Taghikhani et al. (2024) and have been already known

in the SE Fars Arc (e.g., Callot et al., 2007; Jahani et al., 2007) and the Persian Gulf (e.g., Hassanpour et al., 2020,
2021; Snidero et al., 2020) through seismic interpretations. A well-exposed example of such a structure is the Hajiabad
diapir, located in the inner part of the Fars Arc, where an 8 km-long squeezed Hormuz salt diapir trends NW and dips
towards NE (Fig. 13d). The salt-bearing evaporites have been shortened between the two flanks of the precursor salt
720 wall, forming a secondary salt weld. Another well-exposed example in the inner Fars is the Kharman diapir (Fig. 13e),
which is in good correlation with observations from Model 3. Fernandez and Kaus (2014) showed with numerical
modeling that pre-existing salt diapirs can significantly influence the pattern and growth of three-dimensional folds
and fold patterns, accelerating fold formation and localizing deformation, highlighting the important role of diapirism
in structural evolution during tectonic processes.



725 **Figure 13:** (a) The result of numerical modeling with thick salt and basement involved (Model 3). (b) Geological cross section of the Dowlatabad syncline and Kangan anticline (modified from Najafi et al. 2021). (c) Aerial image of Asalouyeh and Kangan anticlines. (d) Satellite image of the Hajiabad Diapir (© Google Earth). (e) Satellite image of the Kharman Diapir (© Google Earth).


730 **5.4 Implication for seismic activity in the Fars Arc**

735 Figure 14a displays a seismicity map of the Zagros in the region of our profile. It shows events with magnitudes greater than $M_L = 4$, between 2000 and 2023 from the ISC global database (<http://www.ics.ac.uk>). We have excluded those events of the database with pre-fixed depths (usually set at 10 or 15 km) from the plot. Furthermore, a catalogue of small-magnitude earthquakes from Tatar et al. (2004) recorded by a local temporary network over a period of seven weeks is plotted. A local network with close station spacing allows for more precise location of earthquakes, and Tatar et al. (2004) state that these events have horizontal and vertical uncertainties below 2 km. The earthquakes are also plotted onto the geological cross-section of figure 1c and the compositional image of the final stage (15 Myr) of Model 1 (Fig. 14b, c). Several studies have asserted that the depths and focal mechanism solutions indicate that high-angle

reverse faults in the basement are responsible for major earthquakes in the SE Zagros (e.g., Jackson and Fitch, 1981; Berberian, 1995; Talebian and Jackson, 2004). Others have argued that most of the strongest earthquakes in the **SFBZ/SFB** probably lie in the lower sedimentary cover, but their aftershock activity mostly occurs deeper in the basement (e.g. Nissen et al., 2011; 2014). A more recent work by Karasözen et al. (2019) used calibrated earthquake relocation to show that the Zagros earthquakes nucleate in both the cover and the basement. According to our numerical results, the activation of basement structures localizes topographic features and affect thrust faults within the overlying sedimentary cover (Fig. 4g4f, 9b, 10). Notable examples of active basement faults in the Fars Arc include the Surmeh Fault and the Mountain Front Fault, which have distinct structural and topographic characteristics (Jackson, 1980; Jackson and McKenzie, 1984; Berberian, 1995; Mouthereau et al. 2007b). Critical wedge modeling confirms that basement-involved shortening is the primary factor influencing deformation and topography in the Fars Arc (Fig. 4g4f and 9b; Mouthereau et al., 2006). Balanced cross-sections provide additional evidence of the basement's involvement in shortening through major thrusts rooted in a deep décollement level within the lower crust (Molinaro et al. 2005; Mouthereau et al, 2007; Najafi et al; 2014; Najafi et al; 2021).

The majority of earthquakes are concentrated within the sedimentary cover, aligning with the décollement faults (Fig. 14b, c). This observation supports the theory of the sedimentary cover decoupling from the basement, influenced by the presence of the basal salt layer. The depth trend of earthquakes in the basement exhibits a notable correlation with the High Zagros Fault and the Surmeh Fault, which confirms the existence of pre-existing weak zones within the basement at these locations. Furthermore, the depth distribution of the earthquakes supports the idea of a listric geometry for the basement faults. By examining the deformation within the basement and the earthquake depths, we estimate the transition zone from brittle to ductile behaviour at around 30 km. While this depth is influenced by multiple factors, including material parameters, temperature, and strain rate (stress), our approximation is based on typical conditions relevant to the region. This suggests that diabase rheology, combined with the applied geotherm, is appropriate for modeling the basement under these specific conditions. Mouthereau et al. (2006) on the other hand concluded that diabase might be too weak to reproduce the observed topography in the Fars arc. However, they apply a crustal thickness of 45 km, which implies increased temperature conditions and thus a weaker diabase detachment. Our modeling results indicate that reverse faults cutting both flanks of décollement anticlines are likely responsible for the shallower earthquakes (Fig. 14c). This phenomenon was studied in the Shanul anticline, southeast of our study region, where earthquakes with reverse mechanisms occur on both flanks of the anticline at depths mostly ≤ 4 km (Jamalreyhani et al., 2021). However, basement faults are the source of deeper and larger earthquakes across the Zagros fold belt, generally concentrated in NE-dipping linear trends (Fig. 14) (e.g., Nissen et al., 2019; Fig. 14). Lacombe and Mouthereau (2002) discussed the relative chronology between mid-crustal décollement thrusting and shallow décollement folding of sedimentary cover in various orogenic belts, such as Taiwan, the Pyrenees, and the Alps. This relationship has been further investigated through analogue and numerical modeling in the Jura, Appalachians, Kopet Dagh, and Zagros fold-and-thrust belts (e.g., Pohn, 2000; Orjuela et al., 2021; Ruh and Vergés, 2018; as presented in this study). A comprehensive analysis of these studies, in alignment with our results, suggests that there is no definitive sequence for thin- and thick-skinned tectonics in all orogenic belts. This sequence appears to be controlled, at least in part, by the rheology and temperature of basement rocks, the thickness of evaporitic layers

in the sedimentary cover, the dip and flatness of inherited basement faults, as well as their orientation in relation to the tectonic transport direction.



780 Figure 14: (a) Seismicity of Zagros fold-and-thrust belt reported by ISC during the years 2000 to 2023 with magnitudes $M_t > 4$ and local temporary network data by Tatar et al (2004) with magnitudes $ML < 4$, superimposed on a shaded relief map derived from Global Topography. (b, c, d) Projection of earthquakes along profile A–B after removing fixed depths on geological cross section and the reference model (Model 1) after 15 Myr of convergence.

6 Conclusions

785 Our series of 2D finite-difference 2D thermo-mechanical numerical experiments ~~shed light on~~ provide insight into how the rheology of basal salt and the mechanical properties of inherited basement faults control the structural evolution and final style of fold-and-thrust belts. The models were designed based on the tectonic history (continental rifting and later collision) and mechanical stratigraphy of the Fars Arc in the Zagros orogenic belt. The comparison of numerical results with the actual structure of the Fars Arc has ~~yielded~~ led to the following conclusions.

- 790 1) Pre-existing weak zones, such as basement faults and décollement anticlines, serve as primary sites for deformation accumulation, particularly ~~evident~~ during extensional tectonic phases.
- 2) The basement faults form large-wavelength foreland-verging fault-propagation anticlines in the overlying sedimentary cover, while the thick salt layer ~~leads to~~ promotes the growth of second-order smaller-wavelength décollement anticlines accompanied by both fore- and back-limb thrust faults.
- 795 3) Reactivated faults play a crucial role in stress transfer, leading to the formation of new faults and seismic activity, at their tips propagating into the sedimentary cover, ~~with listric~~. Listric faults ~~proving~~ are more effective ~~in~~ accommodating strain rates.
- 4) The presence of a basal salt layer ~~at the base of the sedimentary cover influence~~ influences fault displacement, contributing to the development of back thrusts within the ~~sedimentary~~ cover.
- 800 5) The distribution of earthquakes is significantly influenced by the presence of weak zones, with shallow earthquakes predominantly occurring along décollement levels and larger, deeper earthquakes associated with basement faults. These findings ~~emphasize~~ highlight the importance of considering geological and structural complexities when assessing seismicity and tectonic events in tectonically active regions like the Fars Arc.
- 805 6) The degree of basement involvement directly influences the model's resistance, with greater involvement facilitating deformation process ~~through~~ over geologic time. In addition, variations in resistance to deformation based on salt rheology and fault geometry were observed, with listric faults minimizing resistance and therefore facilitating ~~the~~ deformation.
- 810 7) Intense basement deformation ~~apparent~~ observed in the ~~presented~~ numerical experiments indicates the importance of the lower crust for the construction of regional cross sections of the Zagros fold-and-thrust belt.

Author contribution

FG: modeling setup, experiments run, investigation, writing – original manuscript and handling, visualization, figure drafting. JR: hypothesis of work, numerical modeling code write up, interpretation of results, validation, supervision,

815 manuscript writing, and reviewing and editing. MN: hypothesis of work, interpretation of results, supervision,
fieldwork and cross-section construction, reviewing and editing. FS: hypothesis of work, modeling setup, supervision,
and reviewing and editing. All authors contributed and approved the submitted article.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

820 This manuscript is based on the Ph.D. dissertation of the corresponding author, Fatemeh Gomar, at the Institute for
Advanced Studies in Basic Sciences (IASBS), Zanjan, Iran. It was partly developed during her 6-month scientific stay
in the Structural Geology and Tectonics group at ETH Zurich, Switzerland.

Data availability

825 The GIF movies of all experiments ~~that support the findings of presented in~~ this study are available in Figshare with
the identifier: ~~<https://figshare.com/s/38141397f97519f7de31>~~ <https://figshare.com/s/cb0c7f5a9a6fda0e38ef>

References

- 830 Agard, P., Omrani, J., Jolivet, L., and Mouthereau, F.: Convergence history across Zagros (Iran): Constraints from
collisional and earlier deformation, *International Journal of Earth Sciences.*, 94, 401–419,
<https://doi.org/10.1007/s00531-005-0481-4>, 2005.
- Agard, P., Omrani, J., Jolivet, L., Whitechurch, H., Vrielynck, B., Spakman, W., Monié, P., Meyer, B., and Wortel, R.:
Zagros orogeny: A subduction-dominated process, *Geological Magazine.*, 148, 692–725,
<https://doi.org/10.1017/S001675681100046X>, 2011.
- 835 Alavi, M.: Regional stratigraphy of the Zagros fold-thrust belt of Iran and its proforeland evolution, *American Journal of*
Science., 304, 1–20, <https://doi.org/10.2475/ajs.304.1.1>, 2004.
- Alavi, M.: Structures of the Zagros fold-thrust belt in Iran, *American Journal of Science*, 307, 1064–1095,
<https://doi.org/10.2475/09.2007.02>, 2007.
- 840 Aldega, L., Bigi, S., Carminati, E., Trippetta, F., Corrado, S. and Kavooosi, M.A.: The Zagros fold-and-thrust belt in the
Fars province (Iran): II. Thermal evolution, *Marine and Petroleum Geology.*, 93, 376–390,
<https://doi.org/10.1016/j.marpetgeo.2018.03.022>, 2018.
- Anna, L.O., Roberts, L.N. and Potter, C.J.: Geologic assessment of undiscovered oil and gas in the Paleozoic-Tertiary
composite total petroleum system of the Eastern Great Basin, in: *Geologic Assessment of Undiscovered Oil and Gas*
Resources of the Eastern Great Basin Province, Nevada, Utah, Idaho, and Arizona, US Geological Survey Digital
Data Series, 50, 2007.

Formatted: Indent: Hanging: 0.51 cm, Don't swap
indents on facing pages

- 845 Bahroudi, A., and Koyi, H. A.: Effect of spatial distribution of Hormuz salt on deformation style in the Zagros fold and thrust belt: An analogue modelling approach, *Journal of the Geological Society.*, 160, 719–733, <https://doi.org/10.1144/0016-764902-135>, 2003.
- Barber, D. E., Stockli, D. F., and Galster, F.: The Proto-Zagros foreland basin in Lorestan, western Iran: Insights from multiminerale detrital geothermochronometric and trace elemental provenance analysis, *Geochemistry, Geophysics, Geosystems*, 20, 2657–2680, <https://doi.org/10.1029/2019GC008185>, 2019.
- 850 Berberian, M.: Master “blind” thrust faults hidden under the Zagros folds: active basement tectonics and surface morphotectonics, *Tectonophysics.*, 241, 193–224, [https://doi.org/10.1016/0040-1951\(94\)00185-C](https://doi.org/10.1016/0040-1951(94)00185-C), 1995.
- Berberian, M., and King, G. C. P.: Towards a Paleogeography and Tectonic Evolution of Iran, *Canadian Journal of Earth Sciences.*, 18, 210–265, <https://doi.org/10.1139/e81-019>, 1981.
- 855 Barchi, M. R., and Tavarnelli, E.: Thin vs. thick-skinned tectonics in the Umbria-Marche fold-and-thrust belt: Contrast or coexistence?, in: *From the Guajira Desert to the Apennines, and from Mediterranean Microplates to the Mexican Killer Asteroid: Honoring the Career of Walter Alvarez*, Geological Society of America, 0, [https://doi.org/10.1130/2022.2557\(05\)](https://doi.org/10.1130/2022.2557(05)), 2010.
- Bernaola, G., Sharp, I. R., Blanc, E., Saura, E., Hunt, D. W., Fernandez, N., Embry, J. C., Vergés, J., Serra-Kiel, J., Casini, G., Casciello, E., Romaine, I., and Homke, S.: Basin architecture and growth folding of the NW Zagros early foreland basin during the Late Cretaceous and early Tertiary, *Journal of the Geological Society.*, 168, 235–250, <https://doi.org/10.1144/0016-76492010-092>, 2011.
- 860 Blanc, E. J. P., Allen, M. B., Inger, S., and Hassani, H.: Structural styles in the Zagros Simple Folded Zone, Iran. *Journal of the Geological Society.*, 160, 401–412, <https://doi.org/10.1144/0016-764902-110>, 2003.
- 865 Bonini, M., Sani, F., and Antonielli, B.: Basin inversion and contractional reactivation of inherited normal faults: A review based on previous and new experimental models, *Tectonophysics.*, 522–523, 55–88, <https://doi.org/10.1016/j.tecto.2011.11.014>, 2012.
- Borderie, S., Graveleau, F., Witt, C., and Vendeville, B. C.: Impact of an interbedded viscous décollement on the structural and kinematic coupling in fold-and-thrust belts: Insights from analogue modeling, *Tectonophysics.*, 722, 118–137, <https://doi.org/10.1016/j.tecto.2017.10.019>, 2018.
- 870 Boyer, S. E., and Elliott, D.: Thrust systems, *AAPG bulletin.*, 66, 1196–1230, <https://doi.org/10.1029/2002tc001417>, 1982.
- Buiter, S. J. H., and Pfiffner, O. A.: Numerical models of the inversion of half-graben basins, *Tectonics.*, 22, 1–16, <https://doi.org/10.1029/2002tc001417>, 2003.
- 875 Buiter, S. J., Schreurs, G., Albertz, M., Gerya, T. V., Kaus, B., Landry, W., ... and Beaumont, C.: Benchmarking numerical models of brittle thrust wedges, *Journal of Structural Geology.*, 92, 140–177, <https://doi.org/10.1016/j.jsg.2016.03.003>, 2016.
- Burbidge, D. R., and Braun, J.: Numerical models of the evolution of accretionary wedges and fold-and-thrust belts using the distinct-element method, *Geophysical Journal International.*, 148, 542–561, <https://doi.org/10.1046/j.1365-246x.2002.01579.x>, 2002.
- 880

- Callot, J. P., Jahani, S., and Letouzey, J.: The Role of Pre-Existing Diapirs in Fold and Thrust Belt Development. Thrust Belts and Foreland Basins, in: Thrust Belts and Foreland Basins: from fold kinematics to hydrocarbon systems, Springer Berlin Heidelberg, 309–325, https://doi.org/10.1007/978-3-540-69426-7_16, 2007.
- 885 Callot, J. P., Trocmé, V., Letouzey, J., Albouy, E., Jahani, S., and Sherkat, S.: Pre-existing salt structures and the folding of the Zagros Mountains, Geological Society Special Publication., 363, 545–561, <https://doi.org/10.1144/SP363.27>, 2012.
- Crameri, F., Schmeling, H., Golabek, G. J., Duretz, T., Orendt, R., Buitert, S. J. H., ... and Tackley, P. J.: A comparison of numerical surface topography calculations in geodynamic modelling: an evaluation of the ‘sticky air’ method, Geophysical Journal International., 189, 38–54, <https://doi.org/10.1111/j.1365-246X.2012.05388.x>, 2012.
- 890 Chorowicz, J.: The east African rift system, Journal of African Earth Sciences., 43, 379–410, <https://doi.org/10.1016/j.jafrearsci.2005.07.019>, 2005.
- Cooper, M.: Structural style and hydrocarbon prospectivity in fold and thrust belts: A global review, Geological Society Special Publication., 272, 447–472, <https://doi.org/10.1144/GSL.SP.2007.272.01.23>, 2007.
- Cooper, M. A., Williams, G. D., de Graciansky, P. C., Murphy, R. W., Needham, T., de Paor, D., Stoneley, R., Todd, S. P., Turner, J. P., and Ziegler, P. A.: Inversion tectonics - a discussion, Geological Society Special Publication., 44, 335–347, <https://doi.org/10.1144/GSL.SP.1989.044.01.18>, 1989.
- 915 Dal Zilio, L., Ruh, J., and Avouac, J. P.: Structural evolution of orogenic wedges: interplay between erosion and weak décollements, Tectonics., 39, e2020TC006210, <https://doi.org/10.1029/2020TC006210>, 2020.
- Davis, D. M., and Engelder, T.: The role of salt in fold-and-thrust belts, Tectonophysics., 119, 67–88, [https://doi.org/10.1016/0040-1951\(85\)90033-2](https://doi.org/10.1016/0040-1951(85)90033-2), 1985.
- 900 Eslamirezai, N., Alavi, S. A., Nabavi, S. T., and Ghassemi, M. R.: Influence of multiple décollement and cover rock rheology on the structural evolution of thin-skinned fold-and-thrust belts: insights from discrete element modelling, Comptes Rendus, Géoscience., 355, 215–236, <https://doi.org/10.5802/crgeos.224>, 2023.
- Etamad-Saeed, N., Najafi, M. and Vergés, J.: Provenance evolution of Oligocene–Pliocene foreland deposits in the Dezful embayment to constrain Central Zagros exhumation history, Journal of the Geological Society., 177, 799–817, <https://doi.org/10.1144/jgs2019-206>, 2020.
- 905 Falcon, N. L.: Southern Iran: Zagros Mountains, Geological Society Special Publication., 4, 199–211, <https://doi.org/10.1144/GSL.SP.2005.004.01.11>, 1974.
- Gerya, T.: Introduction to numerical geodynamic modelling, Cambridge University Press, <https://doi.org/10.1017/9781316534243>, 2019.
- 910 Gerya, T.V., Yuen, D.A.: Robust characteristics method for modelling multiphase visco-elasto-plastic thermo-mechanical problems, Phys. Earth Planet. Inter., 163, 83–105, <https://doi.org/10.1016/j.pepi.2007.04.015>, 2007.
- Fernandez, N. and Kaus, B.J.: Influence of pre-existing salt diapirs on 3D folding patterns, Tectonophysics., 637, 354–369, <https://doi.org/10.1016/j.tecto.2014.10.021>, 2014.
- 915 Ghazian, R. K., and Buitert, S. J. H.: Numerical modelling of the role of salt in continental collision: An application to the southeast Zagros fold-and-thrust belt, Tectonophysics., 632, 96–110, <https://doi.org/10.1016/j.tecto.2014.06.006>, 2014.

- Granado, P., and Ruh, J. B.: Numerical modelling of inversion tectonics in fold-and-thrust belts, *Tectonophysics.*, 763, 14–29, <https://doi.org/10.1016/j.tecto.2019.04.033>, 2019.
- 920 Granado, P., Ruh, J.B., Santolaria, P., Strauss, P. and Muñoz, J.A.: Stretching and contraction of extensional basins with pre-rift salt: A numerical modeling approach, *Frontiers in Earth Science.*, 9, 648937, <https://doi.org/10.3389/feart.2021.648937>, 2021.
- Gutiérrez, F., Zarei, M., Hudec, M.R. and Deirmik, H.: Normal faulting and landsliding in morpho-structural domes related to buried salt stocks, Zagros Mountains, Iran. Insights into salt breakout, *Marine and Petroleum Geology.*, 155, 925 106376, <https://doi.org/10.1016/j.marpetgeo.2023.106376>, 2023.
- Hasterok, D., and Chapman, D. S.: Heat production and geotherms for the continental lithosphere, *Earth and Planetary Science Letters.*, 307, 59–70, <https://doi.org/10.1016/j.epsl.2011.04.034>, 2011.
- Hassanpour, J., Yassaghi, A., Muñoz, J. A., and Jahani, S.: Salt tectonics in a double salt-source layer setting (Eastern Persian Gulf, Iran): Insights from interpretation of seismic profiles and sequential cross-section restoration, *Basin Research.*, 33, 159-185, <https://doi.org/10.1111/bre.12459>, 2021.
- 930 Hessami, K., Koyi, H. A., Talbot, C. J., Tabasi, H., and Shabanian, E.: Progressive unconformities within an evolving foreland fold-thrust belt, Zagros Mountains, *Journal of the Geological Society.*, 158, 969–982, <https://doi.org/10.1144/0016-764901-007>, 2001.
- Holdsworth, R. E., Butler, C. A., and Roberts, A. M.: The recognition of reactivation during continental 935 deformation, *Journal of the Geological Society.*, 154, 73-78, <https://doi.org/10.1144/gsjgs.154.1.0073>, 1997.
- Homke, S., Vergés, J., Garcés, M., Emami, H., and Karpuz, R.: Magnetostratigraphy of Miocene–Pliocene Zagros foreland deposits in the front of the Push-e Kush arc (Lurestan Province, Iran), *Earth and Planetary Science Letters.*, 225, 397-410, <https://doi.org/10.1016/j.epsl.2004.07.002>, 2004.
- Humair, F., Bauville, A., Epard, J. L., and Schmalholz, S. M.: Interaction of folding and thrusting during fold-and-thrust- 940 belt evolution: Insights from numerical simulations and application to the Swiss Jura and the Canadian Foothills., *Tectonophysics.*, 789, 228474, <https://doi.org/10.1016/j.tecto.2020.228474>, 2020.
- Jackson, M.P. and Hudec, M.R.: Salt tectonics: Principles and practice, Cambridge University Press, 2017.
- Jackson, M.P.A. and Vendeville, B.C.: Regional extension as a geologic trigger for diapirism, *Geological Society of America Bulletin.*, 106, 57–73, [https://doi.org/10.1130/0016-7606\(1994\)106<0057:REAAGT>2.3.CO;2](https://doi.org/10.1130/0016-7606(1994)106<0057:REAAGT>2.3.CO;2), 1994.
- 945 Jackson, J. A.: Reactivation of basement faults and crustal shortening in orogenic belts, *Nature.*, 283, 343–346, <https://doi.org/10.1038/283343a0>, 1980.
- Jackson, J., and Fitch, T.: Basement faulting and the focal depths of the larger earthquakes in the Zagros mountains (Iran), *Geophysical Journal of the Royal Astronomical Society.*, 64, 561–586, <https://doi.org/10.1111/j.1365-246X.1981.tb02685.x>, 1981.
- 950 Jackson, J., and McKenzie, D.: The relationship between plate motions and seismic moment tensors, and the rates of active deformation in the Mediterranean and Middle East, *Geophysical Journal.*, 93, 45–73, <https://doi.org/10.1111/j.1365-246X.1988.tb01387.x>, 1988.

- Jahani, S., Callot, J. P., Letouzey, J., and De Lamotte, D. F.: The eastern termination of the Zagros Fold-and-Thrust Belt, Iran: Structures, evolution, and relationships between salt plugs, folding, and faulting, *Tectonics*, 28, 1–22, <https://doi.org/10.1029/2008TC002418>, 2009.
- 955 Jamalreyhani, M., Pousse-Beltran, L., Büyükakpınar, P., Cesca, S., Nissen, E., Ghods, A., López-Comino, J.Á., Rezapour, M. and Najafi, M.: The 2019–2020 Khalili (Iran) Earthquake Sequence—Anthropogenic Seismicity in the Zagros Simply Folded Belt?, *Journal of Geophysical Research: Solid Earth*, 126, 2021JB022797, <https://doi.org/10.1029/2021JB022797>, 2021.
- 960 Katumwehe, A. B., Abdelsalam, M. G., and Atekwana, E. A.: The role of pre-existing Precambrian structures in rift evolution: The Albertine and Rhino grabens, Uganda, *Tectonophysics*, 646, 117–129, <https://doi.org/10.1016/j.tecto.2015.01.022>, 2015.
- Karasözen, E., Nissen, E., Bergman, E. A., and Ghods, A.: Seismotectonics of the Zagros (Iran) From Orogen-Wide, Calibrated Earthquake Relocations, *Journal of Geophysical Research: Solid Earth*, 124, 9109–9129, <https://doi.org/10.1029/2019JB017336>, 2019.
- 965 Kendall, J., Vergés, J., Koshnaw, R. and Louterbach, M.: Petroleum tectonic comparison of fold and thrust belts: the Zagros of Iraq and Iran, the Pyrenees of Spain, the Sevier of Western USA and the Beni Sub-Andean of Bolivia, *Geological Society, London, Special Publications*, 490, 79-103, <https://doi.org/10.1144/SP490-2018-102>, 2020.
- 970 Khodabakhshnezhad, A., and Arian, M.: Salt tectonics in the Southern Iran, *International Journal of Geosciences*, 7, 367, <https://doi.org/10.4236/ijg.2016.73029>, 2016.
- Koshnaw, R. I., Stockli, D. F., and Schlunegger, F.: Timing of the Arabia-Eurasia continental collision—Evidence from detrital zircon U-Pb geochronology of the Red Bed Series strata of the northwest Zagros hinterland, Kurdistan region of Iraq, *Geology*, 47, 47-50, <https://doi.org/10.1130/G45499.1>, 2019.
- 975 Konstantinovskaya, E., and Malavieille, J.: Thrust wedges with décollement levels and syntectonic erosion: A view from analog models, *Tectonophysics*, 502, 336–350, <https://doi.org/10.1016/j.tecto.2011.01.020>, 2011.
- Koyi, H. A., and Mansurbeg, H.: The Role of Multiple Weak Lithologies in the Deformation of Cover Units in the Northwestern Segment of the Zagros FOLD-AND-THRUST Belt, *Journal of Petroleum Geology*, 44, 145-166, <https://doi.org/10.1111/jpg.12783>, 2021.
- 980 Lacombe, O. and Mouthereau, F.: Basement-involved shortening and deep décollement tectonics in forelands of orogens: Insights from recent collision belts (Taiwan, Western Alps, Pyrenees), *Tectonics*, 21, 12-1, <https://doi.org/10.1029/2001TC901018>, 2002.
- 985 Lacombe, O., Bellahsen, N., and Mouthereau, F.: Fracture patterns in the Zagros Simply Folded Belt (Fars, Iran): Constraints on early collisional tectonic history and role of basement faults, *Geological Magazine*, 148, 940–963, <https://doi.org/10.1017/S001675681100029X>, 2011.
- Lacombe, O., Ruh, J., Brown, D., and Nilfouroushan, F.: Introduction: Tectonic evolution and mechanics of basement-involved fold-and-thrust belts, *Geological Magazine*, 153, 759–762, Cambridge University Press, <https://doi.org/10.1017/S0016756816000650>, 2016.

- Leturmy, P., Molinaro, M., and de Lamotte, D. F.: Structure, timing and morphological signature of hidden reverse basement faults in the Fars Arc of the Zagros (Iran), *Geological Society Special Publication.*, 330, 121–138, <https://doi.org/10.1144/SP330.7>, 2010.
- Letouzey, J., and Sherkati, S.: Salt movement, tectonic events, and structural style in the central Zagros fold and thrust belt (Iran), in: *Salt Sediment Interactions and Hydrocarbon Prospectivity Concepts, Applications and Case Studies for the 21 century*, *Sedimentary geology*, <https://doi.org/10.5724/gcs.04.24.0753>, 2004.
- 995 Li, S., Abe, S., Reuning, L., Becker, S., Urai, J. L., and Kukla, P. A.: Numerical modelling of the displacement and deformation of embedded rock bodies during salt tectonics: A case study from the South Oman Salt Basin, *Geological Society, London, Special Publications.*, 363, 503-520, <https://doi.org/10.1144/SP363.24>, 2012.
- Li, S. Y., and Urai, J. L.: Rheology of rock salt for salt tectonics modeling, *Petroleum Science.*, 13, 712–724, <https://doi.org/10.1007/s12182-016-0121-6>, 2016.
- 1000 Madritsch, H., Schmid, S. M., and Fabbri, O.: Interactions between thin- and thick-skinned tectonics at the northwestern front of the Jura fold-and-thrust belt (eastern France), *Tectonics.*, 27, <https://doi.org/10.1029/2008TC002282>, 2008.
- [Madanipour S, Najafi M, Nozaem R, Vergés J, Yassaghi A, Heydari I, Khodaparast S, Soudmand Z, Aghajari L. The Arabia–Eurasia Collision Zone in Iran: Tectonostratigraphic and Structural Synthesis. *Journal of Petroleum Geology*. 2024 Apr;47\(2\):123-71. 2024.](#)
- 1005 Malavieille, J., and Konstantinovskaya, E.: Impact of surface processes on the growth of orogenic wedges: insights from analog models and case studies, *Geotectonics.*, 44, 541-558, <https://doi.org/10.1134/S0016852110060075>, 2010.
- McQuarrie, N.: Crustal scale geometry of the Zagros fold–thrust belt, Iran, *Journal of structural Geology.*, 26, 519-535, <https://doi.org/10.1016/j.jsg.2003.08.009>, 2004.
- McBeck, J., Cooke, M., Souloumiac, P., Maillot, B., and Mary, B.: The influence of detachment strength on the evolving deformational energy budget of physical accretionary prisms, *Solid Earth.*, 9, 1421–1436, <https://doi.org/10.5194/se-9-1421-2018>, 2018.
- McQuarrie, N., and Ehlers, T. A.: Techniques for understanding fold-and-thrust belt kinematics and thermal evolution, in *Linkages and Feedbacks in Orogenic Systems*, *Geological Society of America*, [https://doi.org/10.1130/2017.1213\(02\)](https://doi.org/10.1130/2017.1213(02)), 2017.
- 1015 Mitra, S.: A unified kinematic model for the evolution of detachment folds, *Journal of Structural Geology.*, 25, 1659-1673, [https://doi.org/10.1016/S0191-8141\(02\)00198-0](https://doi.org/10.1016/S0191-8141(02)00198-0), 2003.
- Mohajjel, M., and Fergusson, C. L.: Jurassic to cenozoic tectonics of the zagros orogen in northwestern Iran, *International Geology Review.*, 56, 263–287, <https://doi.org/10.1080/00206814.2013.853919>, 2014.
- Molinaro, M., Leturmy, P., Guezou, J. C., Frizon de Lamotte, D., and Eshraghi, S. A.: The structure and kinematics of the southeastern Zagros fold-thrust belt, Iran: From thin-skinned to thick-skinned tectonics, *Tectonics.*, 24, 1–19, <https://doi.org/10.1029/2004TC001633>, 2005.
- 1020 Morley, C. K., King, R., Hillis, R., Tingay, M., and Backe, G.: Deepwater fold and thrust belt classification, tectonics, structure and hydrocarbon prospectivity: A review, *Earth-Science Reviews.*, 104, 41–91, <https://doi.org/10.1016/j.earscirev.2010.09.010>, 2011.

Formatted: Indent: Hanging: 0.51 cm, Don't swap indents on facing pages

- 1025 Moresi, L., Dufour, F., Muhlhaus, H.B.: A Lagrangian integration point finite element method for large deformation modeling of viscoelastic geomaterials, *J. Comput. Phys.*, 184, 476–497, [https://doi.org/10.1016/S0021-9991\(02\)00031-1](https://doi.org/10.1016/S0021-9991(02)00031-1), 2003.
- Moresi, L., Quenette, S., Lemiale, V., Meriaux, C., Appelbe, B., and Muhlhaus, H.B.: Computational approaches to studying non-linear dynamics of the crust and mantle, *Phys. Earth Planet. Inter.*, 163, 69–82, <https://doi.org/10.1016/j.pepi.2007.06.009>, 2007.
- 1030 Motamedi, H., Sherkati, S., and Sepehr, M.: Structural style variation and its impact on hydrocarbon traps in central Fars, southern Zagros folded belt, Iran, *Journal of Structural Geology.*, 37, 124–133, <https://doi.org/10.1016/j.jsg.2012.01.021>, 2012.
- Motamedi, H., and Gharabeigli, G.: Structural style in the Fars geological province: Interaction of diapirism and multidécollement folding, In *Developments in Structural Geology and Tectonics.*, 3, 145–160, <https://doi.org/10.1016/B978-0-12-815048-1.00009-3>, 2019.
- Mouthereau, F., Lacombe, O., and Meyer, B.: The Zagros folded belt (Fars, Iran): Constraints from topography and critical wedge modelling, *Geophysical Journal International.*, 165, 336–356, <https://doi.org/10.1111/j.1365-246X.2006.02855.x>, 2006.
- 1040 Mouthereau, F., Tensi, J., Bellahsen, N., Lacombe, O., De Boisgrollier, T., and Kargar, S.: Tertiary sequence of deformation in a thin-skinned/thick-skinned collision belt: The Zagros Folded Belt (Fars, Iran), *Tectonics.*, 26, <https://doi.org/10.1029/2007TC002098>, 2007a.
- Mouthereau, F., Lacombe, O., Tensi, J., Bellahsen, N., Kargar, S., and Amrouch, K.: Mechanical Constraints on the Development of the Zagros Folded Belt (Fars), in: *Thrust belts and foreland basins: from fold kinematics to hydrocarbon systems*, Springer, 247–266, https://doi.org/10.1007/978-3-540-69426-7_13, 2007b.
- 1045 Mouthereau, F., Lacombe, O., and Vergés, J.: Building the Zagros collisional orogen: Timing, strain distribution and the dynamics of Arabia/Eurasia plate convergence, *Tectonophysics.*, 532–535, 27–60, <https://doi.org/10.1016/j.tecto.2012.01.022>, 2012.
- Najafi, M., Beamud, E., Ruh, J., Mouthereau, F., Tahmasbi, A., Bernalola, G., Yassaghi, A., Motamedi, H., Sherkati, S., Goodarzi, M.G.H. and Vergés, J.: Pliocene growth of the Dowlatabad syncline in Frontal Fars arc: Folding propagation across the Zagros Fold Belt, Iran, *Bulletin.*, 133, 1381–1403, <https://doi.org/10.1130/B35748.1>, 2021.
- 1050 Najafi, M., and Lajmorak, S.: Contractional salt-tectonic system in the south Dezful embayment, Zagros, *Journal of Structural Geology.*, 141, 104–204. <https://doi.org/10.1016/j.jsg.2020.104204>, 2020.
- Najafi, M., Vergés, J., Etemad-Saeed, N., and Karimnejad, H. R.: Folding, thrusting and diapirism: Competing mechanisms for shaping the structure of the north Dezful Embayment, Zagros, Iran, *Basin Research.*, 30, 1200–1229, <https://doi.org/10.1111/bre.12300>, 2018.
- 1055 Najafi, M., Yassaghi, A., Bahroudi, A., Vergés, J., and Sherkati, S.: Impact of the Late Triassic Dashtak intermediate décollement horizon on anticline geometry in the Central Frontal Fars, SE Zagros fold belt, Iran, *Marine and Petroleum Geology.*, 54, 23–36, <https://doi.org/10.1016/j.marpetgeo.2014.02.010>, 2014.
- 1060 Nilforoushan, F., Masson, F., Vernant, P., Vigny, C., Martinod, J., Abbassi, M., Nankali, H., Hatzfeld, D., Bayer, R., Tavakoli, F., Ashtiani, A., Doerflinger, E., Daignières, M., Collard, P., and Chéry, J.: GPS network monitors the

- Arabia-Eurasia collision deformation in Iran, *Journal of Geodesy.*, 77, 411–422, <https://doi.org/10.1007/s00190-003-0326-5>, 2003.
- 1065 Nissen, E., Ghods, A., Karasözen, E., Elliott, J.R., Barnhart, W.D., Bergman, E.A., Hayes, G.P., Jamal-Reyhani, M., Nemati, M., Tan, F. and Abdalnaby, W.: The 12 November 2017 M w 7.3 Ezgeleh-Sarpolzahab (Iran) earthquake and active tectonics of the Lurestan Arc, *Journal of Geophysical Research: Solid Earth.*, 124, 2124–2152, <https://doi.org/10.1029/2018JB016221>, 2019.
- 1070 Navabpour, P., Angelier, J., and Barrier, E.: Mesozoic extensional brittle tectonics of the Arabian passive margin, inverted in the Zagros collision (Iran, interior Fars), *Geological Society, London, Special Publications.*, 330, 65-96, <https://doi.org/10.1144/SP330.5>, 2010.
- Kent, P. E.: Recent Studies of South Persian Salt Plugs, *AAPG Bulletin.*, 42, 2951–2972, <https://doi.org/10.1306/0bda5c2d-16bd-11d7-8645000102c1865d>, 1985.
- Orjuela, M.Á., Martínez-Sánchez, D.A. and Jiménez, G.: Structural changes caused by dips in lateral ramps of fold and thrust belts, *Boletín de Geología.*, 43, 29-44, <https://doi.org/10.18273/revbol.v43n2-2021002>, 2021.
- 1075 Pla, O., Roca, E., Xie, H., Izquierdo-Llavall, E., Muñoz, J. A., Rowan, M. G., ... and Huang, S.: Influence of Syntectonic sedimentation and Décollement rheology on the geometry and evolution of Orogenic wedges: Analog modeling of the Kuqa fold-and-Thrust Belt (NW China), *Tectonics.*, 38, 2727-2755, <https://doi.org/10.1029/2018TC005386>, 2019.
- Pohn, H.A.: Lateral ramps in the folded Appalachians and in overthrust belts worldwide: a fundamental element of thrust-belt architecture, US Government Printing Office, 2000.
- 1080 Parizot, O., Missenard, Y., Barbarand, J., Blaise, T., Benedicto, A., Haurine, F., and Sarda, P.: How sensitive are intraplate inherited structures? Insight from the Cévennes Fault System (Languedoc, SE France), *Geological Magazine.*, 159, 2082-2094, <https://doi:10.1017/S0016756822000152>, 2022.
- Pfiffner, O. A., Schlunegger, F., and Buitter, S. J. H.: The Swiss Alps and their peripheral foreland basin: Stratigraphic response to deep crustal processes, *Tectonics.*, 21, 3-1, <https://doi.org/10.1029/2000TC900039>, 2002.
- 1085 Pfiffner, O. A.: Thick-skinned and thin-skinned styles of continental contraction, *Special Paper of the Geological Society of America.*, 414, 153–177, [https://doi.org/10.1130/2006.2414\(09\)](https://doi.org/10.1130/2006.2414(09)), 2006.
- Pfiffner, O. A.: Thick-skinned and thin-skinned tectonics: A global perspective, *Geosciences.*, 7, 71, <https://doi.org/10.3390/geosciences7030071>, 2017.
- 1090 Pirouz, M., Avouac, J. P., Hassanzadeh, J., Kirschvink, J. L., and Bahroudi, A.: Early Neogene foreland of the Zagros, implications for the initial closure of the Neo-Tethys and kinematics of crustal shortening, *Earth and Planetary Science Letters.*, 477, 168–182, <https://doi.org/10.1016/j.epsl.2017.07.046>, 2017.
- Ranalli, G., and Murphy, D. C.: Rheological stratification of the lithosphere, *Tectonophysics.*, 132, 281-295, [https://doi.org/10.1016/0040-1951\(87\)90348-9](https://doi.org/10.1016/0040-1951(87)90348-9), 1987.
- 1095 Ranalli, G.: Rheology of the crust and its role in tectonic reactivation, *Journal of geodynamics.*, 30, 3-15, [https://doi.org/10.1016/S0264-3707\(99\)00024-1](https://doi.org/10.1016/S0264-3707(99)00024-1), 2000.
- Regard, V., Bellier, O., Thomas, J. C., Abbassi, M. R., Mercier, J., Shabaniyan, E., Feghhi, K., and Soleymani, S.: Accommodation of Arabia-Eurasia convergence in the Zagros-Makran transfer zone, SE Iran: A transition between

- collision and subduction through a young deforming system, *Tectonics*, 23, 1–25, <https://doi.org/10.1029/2003TC001599>, 2004.
- 1100 Robert, A. M. M., Letouzey, J., Kavooosi, M. A., Sherkati, S., Müller, C., Vergés, J., and Aghababaei, A.: Structural evolution of the Kopeh Dagh fold-and-thrust belt (NE Iran) and interactions with the South Caspian Sea Basin and Amu Darya Basin, *Marine and Petroleum Geology*, 57, 68–87, <https://doi.org/10.1016/j.marpetgeo.2014.05.002>, 2014.
- 1105 Roustaei, M., Nissen, E., Abbassi, M., Gholamzadeh, A., Ghorashi, M., Tatar, M., ... and Parsons, B.: The 2006 March 25 Fin earthquakes (Iran)—insights into the vertical extents of faulting in the Zagros Simply Folded Belt, *Geophysical Journal International*, 181, 1275–1291, <https://doi.org/10.1111/j.1365-246X.2010.04601.x>, 2010.
- Ruh, J. B., Kaus, B. J. P., and Burg, J. P.: Numerical investigation of deformation mechanics in fold-and-thrust belts: Influence of rheology of single and multiple décollements, *Tectonics*, 31, 1–23, <https://doi.org/10.1029/2011TC003047>, 2012.
- 1110 Ruh, J. B., Hirt, A. M., Burg, J. P., and Mohammadi, A.: Forward propagation of the Zagros Simply Folded Belt constrained from magnetostratigraphy of growth strata, *Tectonics*, 33, 1534–1551, <https://doi.org/10.1002/2013TC003465>, 2014.
- Ruh, J. B., and Vergés, J.: Effects of reactivated extensional basement faults on structural evolution of fold-and-thrust belts: Insights from numerical modelling applied to the Kopet Dagh Mountains, *Tectonophysics*, 746, 493–511, <https://doi.org/10.1016/j.tecto.2017.05.020>, 2018.
- 1115 Ruh, J. B., Tokle, L., and Behr, W. M.: Grain-size-evolution controls on lithospheric weakening during continental rifting, *Nature Geoscience*, 15, 585–590, <https://doi.org/10.1038/s41561-022-00964-9>, 2022.
- Saura, E., Garcia-Castellanos, D., Casciello, E., Parravano, V., Urruela, A., and Vergés, J.: Modeling the flexural evolution of the Amiran and Mesopotamian foreland basins of NW Zagros (Iran-Iraq), *Tectonics*, 34, 377–395, <https://doi.org/10.1002/2014TC003660>, 2015.
- 1120 Santolaria, P., Harris, L. B., Casas, A. M., and Soto, R.: Influence of décollement-cover thickness variations in fold-and-thrust belts: Insights from centrifuge analog modeling, *Journal of Structural Geology*, 163, 104704, <https://doi.org/10.1016/j.jsg.2022.104704>, 2022.
- Sepehr, M., and Cosgrove, J. W.: Structural framework of the Zagros Fold-Thrust Belt, Iran, *Marine and Petroleum Geology*, 21, 829–843, <https://doi.org/10.1016/j.marpetgeo.2003.07.006>, 2004.
- 1125 Sharp, I., Gillespie, P., Morsalnezhad, D., Taberner, C., Karpuz, R., Vergés, J., ... and Hunt, D.: Stratigraphic architecture and fracture-controlled dolomitization of the Cretaceous Khami and Bangestan groups: an outcrop case study, Zagros Mountains, Iran, *Geological Society, London, Special Publications*, 329, 343–396, <https://doi.org/10.1144/SP329.14>, 2010.
- 1130 Sherkati, S., and Letouzey, J.: Variation of structural style and basin evolution in the central Zagros (Izeh zone and Dezful Embayment), Iran, *Marine and Petroleum Geology*, 21, 535–554, <https://doi.org/10.1016/j.marpetgeo.2004.01.007>, 2004.
- Sherkati, S., Letouzey, J., and De Lamotte, D. F.: Central Zagros fold-thrust belt (Iran): New insights from seismic data, field observation, and sandbox modeling, *Tectonics*, 25, 1–27, <https://doi.org/10.1029/2004TC001766>, 2006.

- 1135 Sherhati, S., Molinaro, M., Frizon de Lamotte, D., and Letouzey, J.: Décollement folding in the Central and Eastern Zagros fold-belt (Iran): Salt mobility, multiple décollements and late basement control, *Journal of Structural Geology*, 27, 1680–1696, <https://doi.org/10.1016/j.jsg.2005.05.010>, 2005.
- Simpson, G. D.: Mechanical modelling of folding versus faulting in brittle–ductile wedges, *Journal of Structural Geology*, 31, 369–381, <https://doi.org/10.1016/j.jsg.2009.01.011>, 2009.
- 1140 Simpson, G. D.: Formation of accretionary prisms influenced by sediment subduction and supplied by sediments from adjacent continents, *Geology*, 38, 131–134, <https://doi.org/10.1130/G30461.1>, 2010.
- Simpson, G.: Mechanics of non-critical fold–thrust belts based on finite element models, *Tectonophysics*, 499, 142–155, <https://doi.org/10.1016/j.tecto.2011.01.004>, 2011.
- Snidero, M., Carrera, N., Mencos, J., Butillé, M., Granado, P., Tavani, S., ... and Muñoz, J. A.: Diapir kinematics in a multi-layer salt system from the eastern Persian Gulf, *Marine and Petroleum Geology*, 117, 104402, <https://doi.org/10.1016/j.marpetgeo.2020.104402>, 2020.
- 1145 Spitz, R., Bauville, A., Epard, J. L., Kaus, B. J., Popov, A. A., and Schmalholz, S. M.: Control of 3-D tectonic inheritance on fold-and-thrust belts: insights from 3-D numerical models and application to the Helvetic nappe system., *Solid Earth*, 11(3), 999–1026, <https://doi.org/10.5194/se-11-999-2020>, 2020.
- 1150 Stöcklin, J.: Possible ancient continental margins in Iran, in: *The geology of continental margins*, Springer, Berlin, Heidelberg, 873–887, https://doi.org/10.1007/978-3-662-01141-6_64, 1974. Stockmal, G. S., Beaumont, C., Nguyen, M., and Lee, B.: Mechanics of thin-skinned fold-and-thrust belts: Insights from numerical models, in: *Whence the Mountains? Inquiries into the Evolution of Orogenic Systems: A Volume in Honor of Raymond A. Price*, Geological Society of America, 0, [https://doi.org/10.1130/2007.2433\(04\)](https://doi.org/10.1130/2007.2433(04)), 2007.
- 1155 Szabo, F., and Kheradpir, A.: Permian and Triassic stratigraphy, Zagros basin, south-west Iran, *Journal of Petroleum Geology*, 1, 57–82, <https://doi.org/10.1111/j.1747-5457.1978.tb00611.x>, 1978.
- Taghikhani, H., Yassaghi, A., Madanipour, S. and Najafi, M.: Salt tectonics along the High Zagros Fault in Iran, faulting through welded salt walls, *Journal of Asian Earth Sciences*, 259, 105928, <https://doi.org/10.1016/j.jseaes.2023.105928>, 2024.
- 1160 Talbot, C. J., and sepeh, M.: The past of a future syntaxis across the Zagros, *Geological Society Special Publication*, 100, 89–109, <https://doi.org/10.1144/GSL.SP.1996.100.01.08>, 1996.
- Talebian, M., and Jackson, J.: A reappraisal of earthquake focal mechanisms and active shortening in the Zagros mountains of Iran, *Geophysical Journal International*, 156, 506–526, <https://doi.org/10.1111/j.1365-246X.2004.02092.x>, 2004.
- 1165 Tatar, M., Hatzfeld, D. and Ghafory-Ashtiani, M.: Tectonics of the Central Zagros (Iran) deduced from microearthquake seismicity, *Geophysical Journal International*, 156, 255–266, <https://doi.org/10.1111/j.1365-246X.2003.02145.x>, 2004.
- Tavani, S., Storti, F., Lacombe, O., Corradetti, A., Muñoz, J. A., and Mazzoli, S.: A review of deformation pattern templates in foreland basin systems and fold-and-thrust belts: Implications for the state of stress in the frontal regions of thrust wedges, *Earth-Science Reviews*, 141, 82–104, <https://doi.org/10.1016/j.earscirev.2014.11.013>, 2015.
- 1170

- Tavani, S., Parente, M., Vitale, S., Iannace, A., Corradetti, A., Bottini, C., ... and Mazzoli, S.: Early Jurassic rifting of the Arabian passive continental margin of the Neo-Tethys. Field evidence from the Lurestan region of the Zagros fold-and-thrust belt, Iran, *Tectonics*, 37, 2586-2607, <https://doi.org/10.1029/2018TC005192>, 2018.
- 1175 Teknik, V., and Ghods, A.: Depth of magnetic basement in Iran based on fractal spectral analysis of aeromagnetic data, *Geophysical Journal International*, 209, 1878–1891, . <https://doi.org/10.1093/gji/ggx132>, 2017.
- Vergés, J., Goodarzi, M. G. H., Emami, H., Karpuz, R., Efstathiou, J., and Gillespie, P.: Multiple detachmnet folding in Pusht-e Kuh Arc, zagros: Role of mechanical stratigraphy, *AAPG Memoir*, 94, 69–94 <https://doi.org/10.1306/13251333M942899>, 2011.
- 1180 Tunini, L., Jiménez-Munt, I., Fernandez, M., Vergés, J. and Villasenor, A.: Lithospheric mantle heterogeneities beneath the Zagros Mountains and the Iranian Plateau: a petrological-geophysical study, *Geophysical Journal International*, 200(1), 596-614, <https://doi.org/10.1093/gji/ggu418>, 2014.
- Vergés, J. H. M. E. S. P. A., Emami, H., Garcés, M., Beamud, E., Homke, S., and Skott, P.: Zagros foreland fold belt timing across Lurestan to constrain Arabia–Iran collision, in: *Developments in structural geology and tectonics*, Elsevier, 29-52, <https://doi.org/10.1016/B978-0-12-815048-1.00003-2>, 2019.
- 1185 Vergés, J., Casini, G., Ruh, J., Cosgrove, J., Sherkat, S., Najafi, M., ... and Jahani, S.: STRUCTURAL STYLE AND TIMING OF NW-SE TRENDING ZAGROS FOLDS IN SW IRAN: INTERACTION WITH NORTH-SOUTH TRENDING ARABIAN FOLDS AND IMPLICATIONS FOR PETROLEUM GEOLOGY, *Journal of Petroleum Geology*, 47, 3-74, <https://doi.org/10.1111/jpg.12850>, 2024.
- 1190 Vogt, K., Matenco, L., and Cloetingh, S.: Crustal mechanics control the geometry of mountain belts. Insights from numerical modelling, *Earth and Planetary Science Letters*, 460, 12-21, <https://doi.org/10.1016/j.epsl.2016.11.016>, 2017.
- Wallace, W. K., and Homza, T. X.: Décollement folds with fixed hinges and variable décollement depth, northeastern Brooks Range, Alaska: Reply, *Journal of Structural Geology*, 20, 1591–1595, [https://doi.org/10.1016/S0191-8141\(98\)00040-6](https://doi.org/10.1016/S0191-8141(98)00040-6), 1998.
- 1195 Walpersdorf, A., Hatzfeld, D., Nankali, H., Tavakoli, F., Nilforoushan, F., Tatar, M., Vernant, P., Chéry, J., and Masson, F.: Difference in the GPS deformation pattern of North and Central Zagros (Iran), *Geophysical Journal International*, 167, 1077–1088, <https://doi.org/10.1111/j.1365-246X.2006.03147.x>, 2006.
- White, S. H., Bretan, P. G., and Rutter, E. H. (1986). Fault-zone reactivation: kinematics and mechanisms. *Philosophical Transactions of the Royal Society of London, Series A, Mathematical and Physical Sciences*, 317, 81-97, <https://doi.org/10.1098/rsta.1986.0026>, 1986.
- 1200 Wilks, K. R., and Carter, N. L.: Rheology of some continental lower crustal rocks, *Tectonophysics*, 182, 57-77, [https://doi.org/10.1016/0040-1951\(90\)90342-6](https://doi.org/10.1016/0040-1951(90)90342-6), 1990.
- Wortel, R., Govers, R., and Spakman, W.: Continental collision and the STEP-wise evolution of convergent plate boundaries: From structure to dynamics, in: *Subduction zone geodynamics*, Springer, Berlin, Heidelberg, 47-59, https://doi.org/10.1007/978-3-540-87974-9_3, 2009.
- 1205 Yamato, P., Kaus, B. J. P., Mouthereau, F., and Castelltort, S.: Dynamic constraints on the crustal-scale rheology of the Zagros fold belt, Iran, *Geology*, 39, 815–818, <https://doi.org/10.1130/G32136.1>, 2011.

Zwaan, F., Schreurs, G., Buitter, S. J., Ferrer, O., Reitano, R., Rudolf, M., and Willingshofer, E.: Analogue modelling of basin inversion: a review and future perspectives, *Solid Earth*, 13, 1859-1905, <https://doi.org/10.5194/se-13-1859-2022>, 2022.