

2001-2022 global gross primary productivity dataset using an ensemble model based on random forest

Xin Chen¹, Tiexi Chen^{1,2,3*}, Xiaodong Li⁴, Yuanfang Chai⁵, Shengjie Zhou¹, Renjie Guo⁶, Jie Dai¹

¹School of Geographical Sciences, Nanjing University of Information Science and Technology, Nanjing 210044, Jiangsu, China.

²Qinghai Provincial Key Laboratory of Plateau Climate Change and Corresponding Ecological and Environmental Effects, Qinghai University of Science and Technology, Xining 810016, China

³School of Geographical Sciences, Qinghai Normal University, Xining 810008, Qinghai, China.

⁴Qinghai Institute of Meteorological Science, Xining 810008, Qinghai, China.

⁵Department of Earth Sciences, Vrije Universiteit Amsterdam, Boelelaan 1085, 1081 HV, Amsterdam, the Netherlands

⁶Faculty of Geographical Science, Beijing Normal University, Beijing, China.

Correspondence to: Tiexi Chen (txchen@nuist.edu.cn)

Abstract. Advancements in remote sensing technology have significantly contributed to the improvement of models for estimating terrestrial gross primary productivity (GPP). However, discrepancies in the spatial distribution and interannual variability within GPP datasets pose challenges to a comprehensive understanding of the terrestrial carbon cycle. In contrast to previous models that rely on remote sensing and environmental variables, we developed an ensemble model based on the random forest (ERF model). This model used GPP outputs from established models (EC-LUE, GPP-kNDVI, GPP-NIRv, Revised-EC-LUE, VPM, MODIS) as inputs to estimate GPP. The ERF model demonstrated superior performance, explaining 85.1% of the monthly GPP variations at 170 sites, surpassing the performance of selected GPP models (67.7%-77.5%) and an independent random forest model using remote sensing and environmental variables (81.5%). Additionally, the ERF model improved accuracy across each month and various subvalues, mitigating the issue of "high value underestimation and low value overestimation" in GPP estimates. Over the period from 2001 to 2022, the global GPP estimated by the ERF model was 132.7 PgC yr⁻¹, with an increasing trend of 0.42 PgC yr⁻², which was comparable to or slightly better than the accuracy of other mainstream GPP datasets in term of validation results of GPP observations independent of FLUXNET (ChinaFlux). Importantly, for the growing number of GPP datasets, our study provides a way to integrate these GPP datasets, which may lead to a more reliable estimate of global GPP.

31 **1 Introduction**

32 Gross primary productivity (GPP) is the largest carbon flux in the global carbon cycle, and serves as the primary input of
33 carbon into the terrestrial carbon cycle. Uncertainties in GPP estimates can propagate to other carbon flux estimates, making
34 it crucial to clarify the spatio-temporal patterns of GPP (Ruehr et al., 2023; Xiao et al., 2019). However, global GPP is variously
35 estimated from 90 PgC yr⁻¹ to 160 PgC yr⁻¹ across different studies, with these variations becoming more pronounced when
36 scaled down to regional scales or specific ecosystem types (Anav et al., 2015; Jung et al., 2020; Ryu et al., 2019). This
37 variability underscores the necessity for innovative methods to reduce uncertainty in GPP estimates.

38 The light use efficiency (LUE) model is one of the most widely adopted methods for estimating GPP. It assumes that GPP is
39 proportional to the photosynthetically active radiation absorbed by vegetation, and optimizes the spatio-temporal pattern of
40 GPP through meteorological constraints such as temperature and moisture (Pei et al., 2022). However, variations in these
41 constraints varies significantly, leading to differences of over 10% in model explanatory power. (Yuan et al., 2014). Recent
42 studies have proposed some novel vegetation indices that have been shown to be effective proxies for GPP through theoretical
43 derivation and observed validation (Badgley et al., 2017; Camps-Valls et al., 2021). However, these vegetation indices often
44 use only remote sensing data as an input for estimating long-term GPP without considering meteorological factors, which has
45 led to some controversy (Chen et al., 2024; Dechant et al., 2022; Dechant et al., 2020). Both LUE and vegetation index models
46 use linear mathematical formulas to estimate GPP, but ecosystems are inherently complex, and the biases introduced by these
47 numerical models increase the uncertainty of GPP estimates. Machine learning models have shown great potential for
48 improving GPP estimates in previous studies (Guo et al., 2023; Jung et al., 2020). These models are trained by non-physical
49 means directly using GPP observations and selected environmental and vegetation variables, and the performance of the
50 models depends on the number and quality of observed data and the representativeness of input data. Nevertheless, direct
51 validation from flux towers of FLUXNET reveals that these models typically explain only about 70% of monthly GPP
52 variations, with similar performance to other GPP estimate models (Badgley et al., 2019; Jung et al., 2020; Wang et al., 2021;
53 Zheng et al., 2020). Due to deviations in the model structure, a common limitation across these models is the poor estimate of
54 monthly extreme GPP, leading to the phenomenon of "high value underestimation and low value overestimation " (Zheng et
55 al., 2020). Especially for extremely high values, which usually occur during the growing season and largely determine the
56 annual totals and interannual fluctuations of GPP, this underestimation may hinder our understanding of the global carbon
57 cycle.

58 It is challenging for a single model to provide accurate estimates for all global regions. Previous studies have shown that
59 ensemble models perform significantly better than single models and can handle some inherent issues in single models (Chen
60 et al., 2020; Yao et al., 2014). Traditional multi-model ensemble methods usually use a simple multi-model average or a
61 Bayesian model averaging. However, these methods typically assign fixed weights to each model and are essentially linear
62 combinations. Recent studies have incorporated machine learning techniques to multi-model ensembles to establish nonlinear
63 relationships between multiple simulated target variables and real target variable, improving simulation performance (Bai et

64 al., 2021; Tian et al., 2023; Yao et al., 2017). Whether this method can improve some common problems with individual GPP
65 estimate models, such as high value underestimation and low value overestimation, is not clear and needs to further
66 investigation.

67 In this study, we attempt to use an ensemble model based on the random forest (ERF model) to improve global GPP estimate.
68 Specifically, the work of this study includes the following: (1) Recalibrating parameters for each model, and comparing the
69 performance of six GPP estimate models and the ERF model; (2) Focusing on the phenomenon of “high value underestimation
70 and low value overestimation” in each model, and evaluating the performance of each model across different months,
71 vegetation types and subvalues (high value, median value, low value); (3) Developing a global GPP dataset using the ERF
72 model and validating its generalization using GPP observations from ChinaFlux.

73 **2 Method**

74 **2.1 Data at the global scale**

75 In this study, we selected remote sensing data from the Moderate Resolution Imaging Spectroradiometer (MODIS) and
76 meteorological data from ERA5 to estimate global GPP (Hersbach et al., 2020). For the remote sensing data, surface reflectance
77 (red band, near infrared band, blue band and shortwave infrared band), leaf area index (LAI) and fraction of photosynthetically
78 active radiation (FPAR) were used. For meteorological data, we selected average air temperature, dew point temperature,
79 minimum air temperature, total solar radiation and direct solar radiation. Dew point temperature and average air temperature
80 were used to calculate saturated vapor pressure difference (VPD) (Yuan et al., 2019), and diffuse solar radiation (DifSR) was
81 derived as the difference between total solar radiation and direct solar radiation. Minimum air temperature was obtained from
82 the hourly air temperature. CO₂ data were obtained from the monthly average carbon dioxide levels measured by the Mauna
83 Loa Observatory in Hawaii. Table 1 provides an overview of the datasets used in this study.

84

85 **Table 1.** Overview of the datasets used in this study.

Variable	Dataset	Spatial resolution	Temporal resolution	Temporal coverage
Surface reflectance (red band and near infrared band)	MCD43C4	0.05 °	daily	2001-2022
Surface reflectance (red band, near infrared band, blue band and shortwave infrared band)	MOD09CMG	0.05 °	daily	2001-2022
LAI	MOD15A2H	500m	8d	2001-2022
FPAR	MOD15A2H	500m	8d	2001-2022
Average air temperature (AT)	ERA5-land	0.1 °	Monthly	2001-2022

Dew point temperature (DPT)	ERA5-land	0.1 °	Monthly	2001-2022
Minimum air temperature (MINT)	ERA5-land	0.1 °	Monthly	2001-2022
Total solar radiation (TSR)	ERA5 monthly data on single levels	0.25 °	Monthly	2001-2022
Direct solar radiation (DirSR)	ERA5 monthly data on single levels	0.25 °	Monthly	2001-2022
CO ₂	NOAA's Earth System Research Laboratory	/	Monthly	2001-2022
Distribution map of C4 crops	Harvested Area and Yield for 175 Crops	1/12 °	Annual	2000
Land use	MCD12C1	0.05 °	Annual	2010

86

87 Previous studies have shown that the photosynthetic capacity of C4 crops is much higher than that of C3 crops (Chen et al.,
88 2014; Chen et al., 2011), so it is necessary to divide the cropland into C3 crops and C4 crops. To estimate the global GPP, we
89 used the "175 Crop harvested Area and yield" dataset, which describes the global harvested area and yield of 175 crops in
90 2000 (Monfreda et al., 2008). We extracted the sum of the area ratios of all C4 crops (corn, corn feed, sorghum, sorghum feed,
91 sugarcane, millet) at each grid as the coverage of C4 crops (Figure S1). Consequently, the estimated value of cropland GPP
92 can be expressed as: coverage of C3 crops × simulated GPP value of C3 crops + coverage of C4 crops × simulated GPP value
93 of C4 crops, which has been used in a previous study (Guo et al., 2023).

94 The land use map was derived from the IGBP classification of MCD12C1, and 2010 was chosen as the reference year (that is,
95 land use data is unchanged in the simulation of global GPP). In order to meet the requirements of subsequent research, land
96 use types were grouped into 9 categories: Deciduous Broadleaf Forest (DBF), Evergreen Needleleaved Forest (ENF),
97 Evergreen Broadleaf Forest (EBF), Mixed Forest (MF), Grassland (GRA), Cropland (including CRO-C3 and CRO-C4),
98 Savannah (SAV), Shrub (SHR), Wetland (WET).

99 Finally, for higher resolution data, we gridded the dataset to 0.05 ° by averaging all pixels whose center fell within each 0.05 °
100 grid cell for upscaling. For lower resolution data, we used the nearest neighbor resampling method to 0.05 °. In addition,
101 MODIS data were aggregated to a monthly scale to ensure spatio-temporal consistency.

102 **2.2 Observation data at the site scale**

103 GPP observations were sourced from the FLUXNET 2015 dataset, which includes carbon fluxes and meteorological variables
104 from more than 200 flux sites around the world (Pastorello et al., 2020). GPP cannot be obtained directly from flux sites and
105 usually needs to be obtained by dismantling the net ecosystem exchange. We chose a monthly level GPP based on the nighttime
106 partitioning method and retained only high quality data ($NEE_VUT_REF_QC > 0.8$) for every year, ultimately selecting 170
107 sites with 10932 monthly values for this study (Figure S2). In addition, we selected monthly average air temperature, total
108 solar radiation and VPD. The site observations do not provide direct solar radiation, so we extracted data from ERA5 covering
109 the flux tower. Monthly minimum air temperature was derived from hourly air temperature. Since some required data in GPP
110 simulation are not directly available at flux sites, LAI and FPAR were extracted from MOD15A2H (500 m), and surface
111 reflectance data (red band, near infrared band, blue band and shortwave infrared band) were derived from MCD43A4 (500 m)
112 and MOD09A1 (500 m). These data are roughly similar to the footprint of the flux site and can represent the land surface of
113 the flux site (Chu et al., 2021).

114 **2.3 GPP estimate model**

115 We selected six independent models to estimate GPP in this study. These models are widely used with few model parameters
116 and have demonstrated reliable accuracy in previous studies (Badgley et al., 2017; Zhang et al., 2017; Zheng et al., 2020). The
117 six models are EC-LUE, Revised-EC-LUE, NIRv-based linear model, kNDVI-based linear model, VPM, MODIS. The VPM,
118 MODIS and EC-LUE are LUE models based on remote sensing data and meteorological data (Running et al., 2004; Xiao et
119 al., 2004; Yuan et al., 2007). Zheng et al., (2020) proposed the Revised-EC-LUE model, which divides the canopy into sunlit
120 and shaded leaves, improving the estimate of global GPP (Zheng et al., 2020). The NIRv and kNDVI are novel vegetation
121 indices calculated from the red and near-infrared bands of the reflectance spectrum (Badgley et al., 2017; Camps-Valls et al.,
122 2021). Similar to solar induced chlorophyll fluorescence, they exhibit a linear relationship with GPP and are considered
123 effective proxies for GPP. Detailed descriptions of all models can be found in Text S1.

124 To reduce uncertainty in GPP estimates from a single model, we used the ERF model, the basic idea of which is to restructure
125 the simulated values of multiple models. In this study, we directly used the ERF model to establish the relationship between
126 the GPP simulated by the above six models and GPP observations. In addition, for comparison with the ERF model, we also
127 used the random forest (RF) method for modeling. In this study, we used average air temperature, minimum air temperature,
128 VPD, direct solar radiation, diffuse solar radiation, FPAR and LAI as explanatory variables. Both models used the random
129 forest method, which has been widely used in previous studies of GPP estimate (Guo et al., 2023; Jung et al., 2020). Random
130 forest is an ensemble learning algorithm that combines the outputs of multiple decision trees to produce a single result, and is
131 commonly used for classification and regression problems (Belgiu and Drăguț, 2016). In the regression problem, the output
132 result of each decision tree is a continuous value, and the average of all decision tree outputs is taken as the final result. An
133 overview of all models used can be found in Table 2.

134 **Table 2.** Overview of the models used in this study.

ID	Model	Input data	Output
1	EC-LUE	FPAR, VPD, AT, SRAD, CO ₂	GPP _{EC}
2	Revised-EC-LUE	LAI, VPD, AT, DifSR, DirSR, CO ₂	GPP _{REC}
3	kNDVI-GPP	Red band and near infrared band (MCD43)	GPP _{kNDVI}
4	NIRv-GPP	Red band and near infrared band (MCD43)	GPP _{NIRv}
5	VPM	Red band, near infrared band, blue band, shortwave infrared band (MOD09), AT, SRAD	GPP _{VPM}
6	MODIS	FPAR, SRAD, MINT, VPD	GPP _{MODIS}
7	Random forest model (RF)	LAI, FPAR, AT, MINT, VPD, DifSR, DirSR	GPP _{RF}
8	Ensemble model based on random forest (ERF)	GPP _{EC} , GPP _{REC} , GPP _{kNDVI} , GPP _{NIRv} , GPP _{MODIS} , GPP _{VPM}	GPP _{ERF}

135

136 **2.4 Model parameter calibration and validation**

137 FLUXNET only provides GPP observations and meteorological data, lacking direct measurements for LAI, FPAR, and surface
 138 reflectance, so remote sensing data is needed. Considering the variety of remote sensing data sources, such as MODIS and
 139 AVHRR, it is evident that calibrating the same GPP estimate model with different remote sensing data can yield varied
 140 parameters. In addition, the number of sites used to calibrate model parameters is also an important influencing factor for
 141 model parameters. The original parameters of these models were calibrated with only a limited number of sites (e.g., 95 sites
 142 for Revised EC-LUE model and 104 for NIRv-GPP) (Wang et al., 2021; Zheng et al., 2020). Therefore, to reduce the impact
 143 of the uncertainty of model parameters on simulation results, we did not use original parameters and conducted parameter
 144 calibration for GPP estimate models across different vegetation types. For EC-LUE, Revised EC-LUE, VPM and MODIS, the
 145 Markov chain Monte Carlo method was used to calibrate model parameters. Traditionally, the mean of the posterior distribution
 146 of parameters is taken as the optimal value. However, previous studies have indicated that some model parameters are not well
 147 constrained when calibrating multiple model parameters (Wang et al., 2017; Xu et al., 2006), so we selected the parameter
 148 with the smallest root-mean-square error (RMSE) as the optimal parameter in each iteration. For each vegetation type, we
 149 randomly selected 70% of the data for parameter calibration, and repeated the process 200 times. In order to avoid overfitting,
 150 we adopted the mean of the 200 calibrated parameters as the final model parameters. Similarly, for the two vegetation index
 151 models, we randomly selected 70% of the data in each vegetation type for parameter calibration, repeating the process 200
 152 times and using the mean of the 200 calibrated parameters as the final model parameters.

153 After obtaining GPP estimates from the six GPP models, we evaluated the simulation performance of the RF model and the
 154 ERF model respectively. For both models, we evaluated the model performance using 5-fold-cross-validation, where the

155 process was repeated 200 times, and the mean of the 200 GPP estimates was considered the final GPP estimate. In addition,
156 we used a second validation method in which all data from 70% of the sites were selected for modeling and only all data from
157 the remaining 30% of the sites were validated, a process that was repeated 200 times. This validation will further illustrate the
158 generalization of the model, i.e. its potential for estimating GPP without local observations. We utilized the determination
159 coefficient (R^2) and RMSE as metrics to evaluate the simulation performance of all models. Additionally, we used the ratio of
160 GPP simulations to GPP observations (Sim/Obs) to measure whether the model overestimates or underestimates.

161 **2.5 Global GPP estimate based on ERF model and its uncertainty.**

162 Based on the ERF model, we estimated global GPP for 2001-2022 (ERF_GPP). It is important to note that in this process, we
163 used all the site data to build the model. The uncertainties of ERF_GPP can be attributed to two primary factors: the influence
164 of the number of GPP observations and the influence of the number of features (that is, the simulated GPP). For the first type
165 of uncertainty, we randomly selected 80% of the data to build a model and simulated the multi-year average of global GPP.
166 The process was repeated 100 times, yielding 100 sets of multi-year averages of ERF_GPP. Their standard deviations were
167 considered as the uncertainty of ERF_GPP caused by the number of GPP observations. For the second type of uncertainty, we
168 selected different number of features to build a model and simulated the multi-year average of global GPP. A total of 56 sets
169 of multi-year averages of ERF_GPP were obtained. The standard deviation of different combinations was considered to be the
170 uncertainty of ERF_GPP caused by the number of features.

171 **2.6 Evaluation of the generalization of different GPP datasets**

172 The majority of flux sites in FLUXNET are concentrated in Europe and North America, it is unclear whether the different GPP
173 estimate methods are suitable for regions with sparse flux sites. Recently, ChinaFlux has published GPP observations from
174 several sites, offering an opportunity to evaluate the generalization of different GPP datasets. However, the spatial resolution
175 of most GPP datasets is 0.05° , and a direct comparison with GPP observations at flux sites is challenging. Therefore, we
176 extracted 0.05° MODIS land use covering the flux sites. If the vegetation type of the flux site matched the MODIS land use,
177 the site was used for the analysis. Finally, a total of 12 flux sites were selected (Figure S2), and Table S1 shows the information
178 of these sites. The same procedure was applied to FLUXNET, resulting in the selection of 52 sites. It should be noted that due
179 to the absence of meteorological data from some sites in Chinaflux, we did not validate all GPP estimate models at the site
180 scale (500 m).

181 We evaluated the generalization of ERF_GPP at 12 ChinaFlux sites and 52 FLUXNET sites. In addition, we selected a number
182 of widely used GPP datasets for comparison, including BESS (Li et al., 2023), GOSIF (Li and Xiao, 2019), FLUXCOM:
183 random forest-based version (FLUXCOM-RF) and ensemble version (FLUXCOM-ENS) (Jung et al., 2020), NIRv (Wang et
184 al., 2021), Revise-EC-LUE (Zheng et al., 2020), MODIS (Running et al., 2004), VPM (Zhang et al., 2017), which were
185 generated using different GPP estimate methods. These GPP datasets all have a spatial resolution of $500\text{ m}-0.5^\circ$, similar to the



186 resampling process in section 2.1, we have unified them to 0.05 °. The common time range for these datasets spanned from
187 2001 to 2018, and the temporal resolution was unified to monthly to match the GPP observations.

188 **3 Result**

189 **3.1 Performance of GPP estimate models at site scale**

190 Table S2-S7 show the optimization results of the six GPP estimate model parameters. Consistent with previous study, in the
191 Revised EC-LUE model, the light use efficiency parameter of shade leaves was significantly higher than that of sunlit leaves
192 (Zheng et al., 2020). It is necessary to divide cropland into C3 crops and C4 crops. In all models, the light use efficiency
193 parameters of C4 crops were significantly higher than those of C3 crops, which was particularly reflected in the two vegetation
194 index models of GPP_{KNDVI} and GPP_{NIRv} , the slope of the linear regression directly reflected the difference in photosynthetic
195 capacity of the different crops.

196 Figure 1 shows the performance of all models across different vegetation types. Overall, the performance of the ERF model
197 was better than that of the other GPP estimate models. GPP_{ERF} had the higher accuracy among all models, with R^2 between
198 0.61-0.91 and RMSE between 0.72-2.78 $gC\ m^{-2}\ d^{-1}$. In contrast, the LUE and vegetation index models performed slightly
199 weaker, especially in EBF, where R^2 was both below 0.5. It is worth noting that compared to other vegetation types, the RMSE
200 was highest for cropland, with 6 out of 8 models for C4 crops exceeding 3 $gC\ m^{-2}\ d^{-1}$, suggesting that these existing GPP
201 estimate models may not properly capture the seasonal changes in cropland GPP. The six models with calibrated parameters
202 and ERF model were found to have no significant deviation across vegetation types. However, GPP_{RF} was significantly
203 underestimated for C4 crops and overestimated for SHR.

a**b****c**

205 **Figure 1.** The performance of the eight models on different vegetation types. a, b and c represent R^2 , RMSE, and Sim/Obs respectively.

206 Combining the results of all flux sites, GPP_{ERF} explained 85.1% of the monthly GPP variations, while the seven GPP estimate

207 models only explained 67.7%-81.5% of the monthly GPP variations (Figure 2). Another validation method in which the

208 validation data were not selected randomly but instead sites were entirely used for either training or validation also showed

209 similar results, the average R^2 and RMSE of 200 validation results of ERF model were 0.822 and 1.68 $gC\ m^{-2}\ d^{-1}$, which were

210 obviously better than other models (Figure S3). In order to further prove the robustness of the ERF model, we also used GPP

211 estimate models with original parameters for modeling and validation. As shown in Figure S4, the performance of these GPP

212 estimate models decreased significantly, with R^2 ranging from 0.570 to 0.719 and RMSE ranging from 2.29 to 3.81 $gC\ m^{-2}\ d^{-1}$.

213 The phenomenon of "high value underestimation and low value overestimation" was also pronounced. However, the ERF

214 model maintained a consistent advantage, with R^2 significantly higher than other GPP estimate models (0.856). In addition,

215 we tested the effect of the number of GPP estimate models on the accuracy of the ERF model. As shown in Table S8, as the

216 number of GPP in the ERF model increased, the performance gain of the model gradually decreased.

217 In summary, GPP_{ERF} showed high accuracy in terms of vegetation type and the ability to interpret monthly variations in GPP,


218 which also illustrates the potential of the ERF model to improve GPP estimate. However, it was observed that most GPP

219 simulations exhibited the phenomenon of "high value underestimation and low value overestimation". For example, GPP_{EC} ,

220 GPP_{REC} , GPP_{MODIS} and GPP_{RF} showed obvious underestimation in the months when the monthly GPP value surpassed 15 gC

221 $m^{-2}\ d^{-1}$ (Figure 2). Therefore, it is necessary to evaluate the performance of different models in each month and different

222 subvalues.



223




224 **Figure 2.** Comparison between the GPP simulations of the eight models and the GPP observations. a-h represents GPP_{EC}, GPP_{NIRv}, GPP_{KNDVI},
 225 GPP_{REC}, GPP_{VPM}, GPP_{MODIS}, GPP_{RF}, GPP_{ERF}, respectively.

226

227 3.2 Performance of GPP estimate models in each month and different subvalues


228 Figure 3 shows the simulation accuracy of the eight models in each month. The ERF model maintained a higher accuracy than
 229 other GPP estimate models, with GPP_{ERF} consistently achieving higher R^2 and lower RMSE in most months, and no evident
 230 phenomena of "high value underestimation and low value overestimation". In contrast, the accuracy of other GPP estimate
 231 models was less satisfactory accuracy, especially during winter (most flux sites are concentrated in the Northern Hemisphere),

232 the LUE models tended to underestimate GPP, and the Sim/Obs remained at 0.72-1.01, although R^2 were above 0.7. Meanwhile,
233 the vegetation index models overestimated GPP, Sim/Obs remained at 1.34-1.73, and R^2 were relatively low, mostly around
234 0.6.



236 **Figure 3.** Performance of the eight models in each month. a, b and c represent R^2 , RMSE, and Sim/Obs respectively.

237 We further compared the performance of all models in different subvalues, including high value ($GPP > 15 \text{ gC m}^{-2} \text{ d}^{-1}$), median
 238 value ($15 \text{ gC m}^{-2} \text{ d}^{-1} > GPP > 2 \text{ gC m}^{-2} \text{ d}^{-1}$), low value ($GPP < 2 \text{ gC m}^{-2} \text{ d}^{-1}$). For extreme values, most models performed poorly
 239 (Figure 4), with R^2 for GPP estimate models falling below 0.3, and only GPP_{VPM} showing better performance in the high value.
 240 GPP_{ERF} demonstrated some improvement in both low and high values, with R^2 0.32 and 0.43, RMSE of 0.89 and 4.73 gC m^{-2}
 241 d^{-1} , and Sim/Obs closer to 1, respectively. In the median value, all models performed better, with no significant bias in the
 242 GPP estimate. The R^2 of GPP estimate models ranged from 0.44 to 0.68, and the RMSE remained between 1.82 and 2.54 gC m^{-2}
 243 d^{-1} . Further analysis was made at two typical sites, it was obvious that GPP_{EC} , GPP_{REC} and GPP_{MODIS} on CN-Qia exhibited
 244 obvious underestimation during the growing season (Figure S5). On CH_Lae, GPP_{KNDVI} and GPP_{VPM} were significantly
 245 overestimated (Figure S6). In contrast, at both sites, GPP_{ERF} was more consistent with observations.



246


247 **Figure 4.** Performance of eight models in different subvalues.

248 3.3 Temporal and spatial characteristics of ERF_GPP and its generalization evaluation

249 Figure 5a shows the spatial distribution of the multi-year average of ERF_GPP. The high values of GPP were mainly
250 concentrated in tropical areas, exceeding $10 \text{ gC m}^{-2} \text{ d}^{-1}$, and relatively high in southeastern North America, Europe and southern
251 China, about $4\text{-}6 \text{ gC m}^{-2} \text{ d}^{-1}$. From 2001-2022, China and India showed the fastest increase in GPP, mostly at $0.1 \text{ gC m}^{-2} \text{ d}^{-1}$
252 (Figure 5b), similar to a previous study that reported that China and India led the global greening (Chen et al., 2019). We
253 further investigated the annual maximum GPP, as shown in Figure 5c, and the North American corn belt was the global leader
254 in GPP at more than $15 \text{ gC m}^{-2} \text{ d}^{-1}$, compared to only $10 \text{ gC m}^{-2} \text{ d}^{-1}$ in most tropical forests. In 2001-2022, the global GPP was
255 $132.7 \pm 2.8 \text{ PgC yr}^{-1}$, with an increasing trend of 0.42 PgC yr^{-2} (Figure 5d). The lowest value was $128.6 \text{ PgC yr}^{-1}$ in 2001, and
256 the highest value was $136.2 \text{ PgC yr}^{-1}$ in 2020.

257 The results of the two uncertainty analyses consistently indicated that ERF_GPP exhibited higher uncertainty in tropical
258 regions (Figures S7 and S8), and the uncertainty of ERF_GPP caused by the number of GPP observations was relatively small,
259 the standard deviation of 100 simulations was about $0.3 \text{ gC m}^{-2} \text{ d}^{-1}$ in the tropics and lower in other regions, below 0.1 gC m^{-2}
260 d^{-1} . In contrast, the uncertainty of ERF_GPP caused by the number of features was more pronounced, especially when fewer
261 features were included in the model. It is worth noting that when the number of features was five, the uncertainty was already
262 substantially less, and the standard deviation was generally lower than $0.5 \text{ gC m}^{-2} \text{ d}^{-1}$.

263



264

265 **Figure 5.** Spatial and temporal characteristics of ERF_GPP during 2001-2022. a represents the multi-year average, b represents the trend, c
266 represents the multi-year average of the annual maximum, and d represents the interannual change of GPP.


267

268 As shown in Figure 6, ERF_GPP and other GPP datasets were validated using GPP observations from ChinaFlux. Among all
269 models, VPM demonstrated the best performance, with R^2 of 0.86 and RMSE of $1.34 \text{ gC m}^{-2} \text{ d}^{-1}$. ERF_GPP also exhibited
270 high generalization, with R^2 of 0.75, RMSE of $1.72 \text{ gC m}^{-2} \text{ d}^{-1}$, there was no “high value underestimation and low value
271 overestimation”, which was comparable to the accuracy of BESS and GOSIF. However, the simulation accuracy of the other
272 GPP datasets in Chinaflux was relatively poor, with the R^2 of NIRv being only 0.64, while FLUXCOM-ENS, FLUXCOM-
273 RF, MODIS and Revised EC-LUE were significantly underestimated, with the Sim/Obs being only 0.71-0.89. In the validation
274 of FLUXNET, the R^2 of FLUXCOM-ENS, MODIS, and Revised EC-LUE ranged from 0.57 to 0.67, and the RMSE ranged
275 from 2.67 to $3.30 \text{ gC m}^{-2} \text{ d}^{-1}$, and exhibited different degrees of underestimation (Figure S9). Other GPP datasets demonstrated
276 similar performance, with ERF_GPP being the best ($R^2 = 0.74$, $\text{RMSE} = 2.26 \text{ gC m}^{-2} \text{ d}^{-1}$).

277

278

279



280

281 **Figure 6.** Comparison between the GPP datasets and the GPP observations from ChinaFlux. a-i represents BESS, FLUXCOM-ENS,
 282 FLUXCOM-RF, GOSIF, MODIS, NIRv, VPM, Revise-EC-LUE, ERF_GPP, respectively.

283 **4 Discussion**

284 **4.1 Performance analysis of different models**

285 After parameter calibration, both LUE and vegetation index models obtained reliable model accuracy. However, noticeable
 286 errors persist in different months and subvalues, indicating the prevalent phenomenon of "high value underestimation and low
 287 value overestimation" (Figures 1-4). In addition to MODIS, the GPP simulated by the other three LUE models is generally
 288 underestimated in winter (Figure 3), which may be caused by biases in the parameters used in meteorological constraints. In

289 the expression form of the temperature constraint adopted by LUE models, the maximum temperature, minimum temperature
290 and optimum temperature for limiting photosynthesis are all constants, however these values may not be fixed (Grossiord et
291 al., 2020; Huang et al., 2019). A previous study has demonstrated that the GPP estimate could be effectively improved by
292 using dynamic temperature parameters (Chang et al., 2021). Moreover, the form of meteorological constraint is also an
293 important influencing factor. Compared with other LUE models, VPM does not use VPD constraints, but incorporates land
294 surface water index from satellite observations as constraints (Xiao et al., 2004), which may be the reason why the model
295 performs better than other models at high value (Figure 4). Conversely, the two vegetation index models overestimated GPP
296 in winter, and even overestimated by 70% in December. The vegetation index model does not consider meteorological
297 constraints that believe that all environmental impacts on vegetation have been included in the vegetation indices (kNDVI,
298 NIRv) (Badgley et al., 2017; Camps-Valls et al., 2021). However, it is a fact that under high temperature or low radiation, the
299 vegetation index may still maintain the appearance of high photosynthesis (greening), while in fact the GPP is low (Chen et
300 al., 2024; Doughty et al., 2021; Yang et al., 2018). Furthermore, the relationship between these vegetation indices and GPP is
301 not robust, and the vegetation indices based on reflectance may have hysteresis (Wang et al., 2022).

302 Compared to other GPP estimate models, the ERF model demonstrated better performance ($R^2 = 851$). Since there are no
303 physical constraints, the machine learning model needs to find the relationship between explanatory variables and target
304 variable from a large amount of training data (such as $GPP=f(LAI, T, P, \text{etc.})$) (Guo et al., 2023; Jung et al., 2020). Therefore,
305 the reliability of the model usually depends on the representativeness of the training data. For example, LAI can explain GPP
306 to a large extent, while complex modeling relationships are still needed from LAI to GPP. The difference between the ERF
307 model and the RF model lies in the explanatory variables. The ERF model uses multiple GPP simulations that are more
308 representative and aligned with the target variable, thus making the GPP simulations more accurate. In other words, the ERF
309 model does not need to take into account the uncertainties of the model structure (such as meteorological constraints) and
310 model parameters (such as maximum light use efficiency), but rather focuses on the uncertainties inherent in the simulated
311 GPP. To further clarify the impact of explanatory variables on the ERF model, we conducted a feature importance analysis
312 (Figure S10). From an average of 200 times, the results of the ERF model did not depend on a single GPP simulation. Even
313 GPP_{MODIS} , with the highest relative importance, accounted for no more than 25%, suggesting that the ERF model behaves
314 more like a weighted average of multiple GPP simulations. In addition, it is important to emphasize that the accuracy of the
315 ERF model is still robust even for GPP simulations of original parameters (Figure S4), which means that we can try to use this
316 method to integrate the currently published GPP data sets to obtain a more accurate global GPP estimate.

317 It is worth noting that in the study of Tian et al. (2023), the ERF model was also used to improve the GPP estimate. Our study
318 extends this work in several ways. Firstly, parameter calibration was carried out in our study so that the final validation results
319 are comparable, that is, differences in model performance are mainly due to the uncertainty of the model structure. Secondly,
320 our study focused on the phenomenon of "high value underestimation and low value overestimation" of GPP estimate models,
321 with results indicating that the ERF model performed well across various vegetation types, months, and subvalues. Finally, we

322 generated the ERF_GPP dataset and validated it on different observational datasets, further confirming the robustness of the
323 ERF model in GPP estimate.

324 **4.2 Robustness of ERF_GPP**

325 Due to the inherent advantages of the RF method, the accuracy of the model was comparable to that of the ERF model, even
326 if a very simple model that used longitude, latitude, month, and year as explanatory variables (Figure S11 a). However, the
327 global GPP estimated by this model was not reliable (Figure S11 b). This illustrates that an excellent model performance based
328 on the FLUXNET sites does not necessarily imply an equivalent prediction skill in other regions. ERF model can overcome
329 this limitation to some extent. On the one hand, the explanatory variables used in the model are derived from GPP simulation
330 in which contain a lot of remote sensing information, which can ensure that the global GPP estimated by the model is reliable.
331 On the other hand, the second validation method also further shows that the ERF model has good generalization and has greater
332 potential than other models in estimating global GPP.

333 Since the current GPP datasets are generated based on remote sensing data and FLUXNET GPP observations, there is a strong
334 similarity in spatial distribution among all GPP datasets. Therefore, the validation of GPP observations independent of
335 FLUXNET is crucial. Validation results from GPP observations of ChinaFlux indicated that ERF_GPP exhibited good
336 generalization in China ($R^2=0.75$), which was slightly lower than the accuracy of 5-fold-cross-validation during modeling,
337 possibly due to the mismatch between the 0.05° GPP estimate and the footprint of the flux tower (Chu et al., 2021). In addition,
338 the validation of FLUXNET further confirms the reliability of ERF_GPP. Overall, this is comparable to or slightly better than
339 the simulation accuracy of current mainstream GPP datasets. We also observed a clear improvement in the spatial maximum
340 value of ERF_GPP in some corn growing regions, such as the North American Corn Belt (Figure 5c), which is supported by
341 previous studies showing that C4 crops have much higher GPP peaks than other vegetation types (Chen et al., 2011; Yuan et
342 al., 2015).

343 Due to the increasing drought trend, the constraining effect of water on vegetation is gradually increasing, and some studies
344 have reported the decoupling phenomenon of LAI and GPP under some specific conditions (Hu et al., 2022; Jiao et al., 2021).
345 However, in China and India with significant greening, GPP continues to increase in most datasets, and ERF_GPP supports
346 this view. This phenomenon may be attributed to the low drought pressure on croplands in China and India due to irrigation,
347 which poses less constraint on GPP (Ai et al., 2020; Ambika and Mishra, 2020). The global estimate of ERF_GPP is $132.7 \pm$
348 2.8 PgC yr^{-1} , which is close to estimates from most previous studies (Badgley et al., 2019; Wang et al., 2021). A study have
349 suggested that global GPP may reach 150-175 PgC yr^{-1} (Welp et al., 2011), however, there is no further evidence to support
350 this view.

351 ERF_GPP exhibited higher uncertainty in tropical regions, similar reports have been made in previously published GPP
352 datasets (Badgley et al., 2019; Guo et al., 2023). The scarcity of flux observations in these regions (Pastorello et al., 2020),
353 coupled with the well-known issue of cloud pollution and saturation in remote sensing data (Badgley et al., 2019), exacerbates
354 the uncertainty in GPP estimates for these regions. Therefore, in future studies, on the one hand, more flux observations in

355 tropical regions are needed, and on the other hand, attempts can be made to combine optical and microwave data to improve
356 GPP estimate.

357 **4.3 Limitations and uncertainties**

358 In this study, we improved GPP estimate based on the ERF model. Nonetheless, there are still some limitations and
359 uncertainties due to the availability of data and methods. First, C4 crop distribution maps were used in our study to improve
360 estimates of cropland GPP. However, this dataset only represents the spatial distribution of crops around the year 2000, which
361 introduce uncertainty into GPP simulations of cropland in a few C3 and C4 alternating areas. Secondly, the ERF model
362 considers six GPP simulations, and it is not clear whether adding more GPP simulations to the model can further improve the
363 GPP estimate. Finally, our model did not consider the effect of soil moisture on GPP, and some previous studies have
364 highlighted the importance of incorporating soil moisture in GPP estimates, especially for dry years (Stocker et al., 2018;
365 Stocker et al., 2019).

366 **5 Conclusion**

367 In this study, we compared the performance of the ERF model with other GPP estimate models at the site scale, especially for
368 the phenomenon of "high value underestimation and low value overestimation", and further developed the ERF_GPP dataset.
369 Overall, GPP_{ERF} had higher model accuracy, explaining 85.1% of the monthly GPP variations, and demonstrated reliable
370 accuracy in different months, vegetation types and subvalues. Over the period from 2001 to 2022, the global estimate of
371 ERF_GPP was $132.7 \pm 2.8 \text{ PgC yr}^{-1}$, corresponding to an increasing trend of 0.42 PgC yr^{-2} . Validation results from ChinaFlux
372 indicated that ERF_GPP had good generalization. For the current emerging GPP estimate models, the ERF model provides an
373 alternative method that lead to better model accuracy.

374 **Data and code availability**

375 The ERF_GPP for 2001-2022 is available at <https://doi.org/10.6084/m9.figshare.24417649> (Chen et al., 2023). The spatial
376 resolution of ERF_GPP is 0.05° and the temporal resolution is monthly. Code is available from the author upon reasonable
377 request.

378 **Author contributions**

379 X.C. and T.X.C. conceived the scientific ideas and designed this research framework. X.C. compiled the data, conducted
380 analysis, prepared figures. X.C., T.X.C. and Y.F.C. wrote the manuscript. D.X.L., R.J.G., J.D., and S.J.Z. gave constructive
381 suggestions for improving the manuscript.

382 **Financial support**

383 This study was supported by the National Natural Science Foundation of China (No. 42130506, 42161144003 and 31570464)
384 and the Postgraduate Research & Practice Innovation Program of Jiangsu Province (No. KYCX23_1322).

385 **Acknowledgments**

386 We thank the two anonymous reviewers and the associate editor (Anja Rammig) for their thoughtful comments and help with
387 this study.

388 **Declaration of interests**

389 The authors have not disclosed any competing interests.

390 **References**

- 391 Ai, Z. et al., 2020. Variation of gross primary production, evapotranspiration and water use efficiency for global croplands.
392 *Agricultural and Forest Meteorology*, 287.
- 393 Ambika, A.K. and Mishra, V., 2020. Substantial decline in atmospheric aridity due to irrigation in India. *Environmental*
394 *Research Letters*, 15(12).
- 395 Anav, A. et al., 2015. Spatiotemporal patterns of terrestrial gross primary production: A review. *Reviews of Geophysics*, 53(3):
396 785-818.
- 397 Badgley, G., Anderegg, L.D., Berry, J.A. and Field, C.B., 2019. Terrestrial gross primary production: Using NIRV to scale
398 from site to globe. *Global change biology*, 25(11): 3731-3740.
- 399 Badgley, G., Field, C.B. and Berry, J.A., 2017. Canopy near-infrared reflectance and terrestrial photosynthesis. *Science*
400 *advances*, 3(3): e1602244.
- 401 Bai, Y. et al., 2021. On the use of machine learning based ensemble approaches to improve evapotranspiration estimates from
402 croplands across a wide environmental gradient. *Agricultural and Forest Meteorology*, 298: 108308.
- 403 Belgiu, M. and Drăguț, L., 2016. Random forest in remote sensing: A review of applications and future directions. *ISPRS*
404 *journal of photogrammetry and remote sensing*, 114: 24-31.
- 405 Camps-Valls, G. et al., 2021. A unified vegetation index for quantifying the terrestrial biosphere. *Science Advances*, 7(9):
406 eabc7447.
- 407 Chang, Q. et al., 2021. Assessing variability of optimum air temperature for photosynthesis across site-years, sites and biomes
408 and their effects on photosynthesis estimation. *Agricultural and Forest Meteorology*, 298.
- 409 Chen, C. et al., 2019. China and India lead in greening of the world through land-use management. *Nature Sustainability*, 2(2):
410 122-129.
- 411 Chen, T., Van Der Werf, G., Gobron, N., Moors, E. and Dolman, A., 2014. Global cropland monthly gross primary production
412 in the year 2000. *Biogeosciences*, 11(14): 3871-3880.
- 413 Chen, T., van der Werf, G.R., Dolman, A.J. and Groenendijk, M., 2011. Evaluation of cropland maximum light use efficiency
414 using eddy flux measurements in North America and Europe. *Geophysical Research Letters*, 38.
- 415 Chen, X. et al., 2024. Vegetation Index-Based Models Without Meteorological Constraints Underestimate the Impact of
416 Drought on Gross Primary Productivity. *Journal of Geophysical Research: Biogeosciences*, 129(1): e2023JG007499.
- 417 Chen, Y., Yuan, H., Yang, Y. and Sun, R., 2020. Sub-daily soil moisture estimate using dynamic Bayesian model averaging.
418 *Journal of Hydrology*, 590: 125445.

419 Chu, H. et al., 2021. Representativeness of Eddy-Covariance flux footprints for areas surrounding AmeriFlux sites.
420 Agricultural and Forest Meteorology, 301: 108350.

421 Dechant, B. et al., 2022. NIRVP: A robust structural proxy for sun-induced chlorophyll fluorescence and photosynthesis across
422 scales. Remote Sensing of Environment, 268: 112763.

423 Dechant, B. et al., 2020. Canopy structure explains the relationship between photosynthesis and sun-induced chlorophyll
424 fluorescence in crops. Remote Sensing of Environment, 241: 111733.

425 Doughty, R. et al., 2021. Small anomalies in dry-season greenness and chlorophyll fluorescence for Amazon moist tropical
426 forests during El Nino and La Nina. Remote Sensing of Environment, 253.

427 Grossiord, C. et al., 2020. Plant responses to rising vapor pressure deficit. New Phytologist, 226(6): 1550-1566.

428 Guo, R. et al., 2023. Estimating Global GPP From the Plant Functional Type Perspective Using a Machine Learning Approach.
429 Journal of Geophysical Research-Biogeosciences, 128(4).

430 Hersbach, H. et al., 2020. The ERA5 global reanalysis. Quarterly Journal of the Royal Meteorological Society, 146(730):
431 1999-2049.

432 Hu, Z. et al., 2022. Decoupling of greenness and gross primary productivity as aridity decreases. Remote Sensing of
433 Environment, 279: 113120.

434 Huang, M. et al., 2019. Air temperature optima of vegetation productivity across global biomes. Nature ecology & evolution,
435 3(5): 772-779.

436 Jiao, W. et al., 2021. Observed increasing water constraint on vegetation growth over the last three decades. Nature
437 Communications, 12(1).

438 Jung, M. et al., 2020. Scaling carbon fluxes from eddy covariance sites to globe: synthesis and evaluation of the FLUXCOM
439 approach. Biogeosciences, 17(5): 1343-1365.

440 Li, B. et al., 2023. BESSv2.0: A satellite-based and coupled-process model for quantifying long-term global land-atmosphere
441 fluxes. Remote Sensing of Environment, 295.

442 Li, X. and Xiao, J., 2019. A Global, 0.05-Degree Product of Solar-Induced Chlorophyll Fluorescence Derived from OCO-2,
443 MODIS, and Reanalysis Data. Remote Sensing, 11(5).

444 Monfreda, C., Ramankutty, N. and Foley, J.A., 2008. Farming the planet: 2. Geographic distribution of crop areas, yields,
445 physiological types, and net primary production in the year 2000. Global Biogeochemical Cycles, 22(1).

446 Pastorello, G. et al., 2020. The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data.
447 Scientific data, 7(1): 1-27.

448 Pei, Y. et al., 2022. Evolution of light use efficiency models: Improvement, uncertainties, and implications. Agricultural and
449 Forest Meteorology, 317: 108905.

450 Ruehr, S. et al., 2023. Evidence and attribution of the enhanced land carbon sink. Nature Reviews Earth & Environment, 4(8):
451 518-534.

452 Running, S.W. et al., 2004. A continuous satellite-derived measure of global terrestrial primary production. Bioscience, 54(6):
453 547-560.

454 Ryu, Y., Berry, J.A. and Baldocchi, D.D., 2019. What is global photosynthesis? History, uncertainties and opportunities.
455 Remote sensing of environment, 223: 95-114.

456 Stocker, B.D. et al., 2018. Quantifying soil moisture impacts on light use efficiency across biomes. New Phytologist, 218(4):
457 1430-1449.

458 Stocker, B.D. et al., 2019. Drought impacts on terrestrial primary production underestimated by satellite monitoring. Nature
459 Geoscience, 12(4): 264+.

460 Tian, Z. et al., 2023. Fusion of Multiple Models for Improving Gross Primary Production Estimation With Eddy Covariance
461 Data Based on Machine Learning. Journal of Geophysical Research: Biogeosciences, 128(3): e2022JG007122.

462 Wang, J. et al., 2017. Decreasing net primary production due to drought and slight decreases in solar radiation in China from
463 2000 to 2012. Journal of Geophysical Research: Biogeosciences, 122(1): 261-278.

464 Wang, S., Zhang, Y., Ju, W., Qiu, B. and Zhang, Z., 2021. Tracking the seasonal and inter-annual variations of global gross
465 primary production during last four decades using satellite near-infrared reflectance data. Science of the Total
466 Environment, 755: 142569.

467 Wang, X. et al., 2022. Satellite solar-induced chlorophyll fluorescence and near-infrared reflectance capture complementary
468 aspects of dryland vegetation productivity dynamics. Remote Sensing of Environment, 270: 112858.

469 Welp, L.R. et al., 2011. Interannual variability in the oxygen isotopes of atmospheric CO₂ driven by El Nino.
470 Nature, 477(7366): 579-582.

471 Xiao, J. et al., 2019. Remote sensing of the terrestrial carbon cycle: A review of advances over 50 years. Remote Sensing of
472 Environment, 233: 111383.

473 Xiao, X. et al., 2004. Modeling gross primary production of temperate deciduous broadleaf forest using satellite images and
474 climate data. Remote sensing of environment, 91(2): 256-270.

475 Xu, T., White, L., Hui, D. and Luo, Y., 2006. Probabilistic inversion of a terrestrial ecosystem model: Analysis of uncertainty
476 in parameter estimation and model prediction. Global Biogeochemical Cycles, 20(2).

477 Yang, J. et al., 2018. Amazon drought and forest response: Largely reduced forest photosynthesis but slightly increased canopy
478 greenness during the extreme drought of 2015/2016. Global Change Biology, 24(5): 1919-1934.

479 Yao, Y. et al., 2017. Improving global terrestrial evapotranspiration estimation using support vector machine by integrating
480 three process-based algorithms. Agricultural and Forest Meteorology, 242: 55-74.

481 Yao, Y. et al., 2014. Bayesian multimodel estimation of global terrestrial latent heat flux from eddy covariance, meteorological,
482 and satellite observations. Journal of Geophysical Research: Atmospheres, 119(8): 4521-4545.

483 Yuan, W. et al., 2015. Uncertainty in simulating gross primary production of cropland ecosystem from satellite-based models.
484 Agricultural and Forest Meteorology, 207: 48-57.

485 Yuan, W. et al., 2014. Global comparison of light use efficiency models for simulating terrestrial vegetation gross primary
486 production based on the LaThuile database. Agricultural and Forest Meteorology, 192: 108-120.

487 Yuan, W. et al., 2007. Deriving a light use efficiency model from eddy covariance flux data for predicting daily gross primary
488 production across biomes. Agricultural and Forest Meteorology, 143(3-4): 189-207.

489 Yuan, W. et al., 2019. Increased atmospheric vapor pressure deficit reduces global vegetation growth. Science advances, 5(8):
490 eaax1396.

491 Zhang, Y. et al., 2017. A global moderate resolution dataset of gross primary production of vegetation for 2000–2016.
492 Scientific data, 4(1): 1-13.

493 Zheng, Y. et al., 2020. Improved estimate of global gross primary production for reproducing its long-term variation, 1982–
494 2017. Earth System Science Data, 12(4): 2725-2746.

495

496