

Petrogenesis of Early Paleozoic I-type granitoids in the Longshoushan and implications for the tectonic affinity and evolution of the southwestern Alxa Block

Renyu Zeng^{1,2,3}, Hui Su¹, Mark B. Allen³, Haiyan Shi⁴, Houfa Dua,^{1,2} Chenguang Zhang⁵, Jie Yan^{1,}

5

¹ School of Earth Sciences, East China University of Technology, Nanchang, 330013, China

² State Key Laboratory of Nuclear Resources and Environment, East China University of Technology, Nanchang 330013, China

³ Department of Earth Sciences, Durham University, Durham DH1 3LE, UK

10 ⁴ Qinghai Geological Survey, Technology Innovation Center for Exploration and Exploitation of Strategic Mineral Resources in Plateau Desert Region, Ministry of Natural Resources, Xining 810000, China

⁵ School of Geographic Sciences, Xinyang Normal University, Xinyang, 464000, China

Correspondence to: Renyu Zeng (zengrenyu@126.com)

15 **Abstract.** In the Paleozoic, the Alxa Block was situated between the Central Asian Orogenic Belt and the North Qilian Orogenic Belt, and experienced intense magmatic activity. Thus, the Alxa Block is an important area for understanding the tectonic framework and evolution of these two orogenic belts. However, there has long been debate regarding the tectonic affinity and tectonic evolution of the Longshoushan, located in southwestern margin of the Alxa Block, during the Paleozoic. In this study, we present zircon U-Pb ages, whole-rock major and trace elements, and Hf isotopic data for the granitoids from

20 the east of the Longshoushan to investigate these issues. Bulk-rock analyses show that these granitoids are weakly peraluminous, with high SiO₂ and K₂O, but low MgO, TFe₂O₃ and P₂O₅. They are also characterized by enrichment in LREE and LILE, depletion in HREE and HFSE, and a large range of εHf(t) values (monzogranite: -0.3 to -16.2; K-feldspar granite: 3.5 to -7.7). These geochemical features indicate that these granitoids are highly fractionated I-type granites, which were formed by crust and mantle-derived magma mixing. LA-ICP-MS zircon U-Pb dating constrains that the monzogranite and K-

25 feldspar granite formed at 440.8 ± 2.1 Ma and 439.4 ± 2.0 Ma, respectively. Combining these results with previous chronological data, the geochronology framework of Paleozoic magmatic events in the Longshoushan is consistent with the North Qilian Orogenic Belt to the south, but significantly differs from other parts of the Alxa Block and the Central Asian Orogenic Belt to the north. This result indicates that the Longshoushan was primarily influenced by the North Qilian Orogenic Belt during the Early Paleozoic. Integrated with previous studies, a three-stage tectonic model is proposed of Early Paleozoic

30 accretion and arc magmatism leading to collision in the Longshoushan. (1) 460-445 Ma: Arc magmatism on an active continental margin with the northward subduction of the North Qilian back-arc basins (NQ bab). (2) 445-435 Ma: Magmatic rocks, dominated by I-type granites, were formed in a continent-continent collision setting. Significant crustal thickening is interpreted to result from compressional stress and/or magmatic additions. (3) 435-410 Ma: The development of abundant A-

type granites and mafic dikes in response to extension, supported by a change in trace element chemistry indicating crustal thinning at this stage. This sequence of events and their timings is similar to other parts of the Central China Orogenic Belt, and requires either a coincidence of several oceanic plates closing at the same time, or an along-strike repetition of the same system.

1 Introduction

The Alxa Block is a region of Precambrian basement adjacent to Phanerozoic orogenic belts (Fig. 1a). Its northern and southern sides are adjacent to the Central Asian Orogenic Belt and the North Qilian Orogenic Belt, respectively (Song et al., 2013; Xue et al., 2017; Zeng et al., 2016, 2021; Hui et al., 2021; Allen et al., 2023). The eastern margin may have undergone collision and amalgamation events with the North China Craton, although the timing of the amalgamation is in dispute (Li et al., 2012; Wang et al., 2015; Yuan et al., 2015; Dan et al., 2016). The Alxa Block therefore documents the tectonic evolution of the Proto-Tethys and Paleo-Asian oceans.

The Longshoushan is a mountain range located in the south part of the Alxa Block (Fig. 1b) and is characterized by Early Paleozoic magmatic activity. It is generally believed that the region to the north part of the Longshoushan, known as the Beidashan, is mainly influenced by the Central Asian Orogenic Belt (Liu et al., 2016; Zhou et al., 2016), while the region to the south, the Hexi Corridor, is influenced by the North Qilian Orogenic Belt (Wei et al., 2013; Zhang et al., 2017). However, the tectonic background of the Longshoushan itself is still debated. Some researchers suggest that the Early Paleozoic magmatic rocks in the Longshoushan are related to the subduction-accretionary orogenesis of the North Qilian Orogenic Belt (Liu et al., 2021; Zeng et al., 2021), while others propose a connection between these magmatic events and the roll-back of the Paleo-Asian Oceanic slab (Liu et al., 2016; Zhou et al., 2016; Xue et al., 2017). Furthermore, it is widely believed that the Longshoushan underwent a stress transition from a compressional to an extensional environment during the Early Paleozoic (Zeng et al., 2016; Zhang et al., 2017; Wang et al., 2018; Liu et al., 2021). However, there are different viewpoints regarding the age of this transition, with some suggesting it occurred in the Late Ordovician (~ 450-440 Ma, Zhang et al., 2017; Liu et al., 2021) and others proposing it took place in the Early Silurian (~ 433-430 Ma, Yu et al., 2015; Zeng et al., 2016).

As an important component of the continental crust, granitoid is of great significance in studying crustal properties, tectonic framework and tectonic evolution (e.g. Pearce et al., 1984; Li et al., 2014; Wang et al., 2017; Wu et al., 2017; Zeng et al., 2022). Extensive research has been conducted on granitoids in the Longshoushan, including the Zhimengou, Taohualashan, Jiling, Jinchuan and Qingshanbao plutons, and the western continuation of the Helishan area. However, research on the granitoids in the east of the Longshoushan has not yet been undertaken to the same extent. In this contribution, whole-rock geochemistry, zircon U-Pb ages, and Hf isotopes of the granitoids in the east of the Longshoushan are systematically studied. In conjunction with published data on Paleozoic magmatic rocks in the Longshoushan, this study also aims to investigate the tectonic background, crustal thickness variations, and tectonic evolution of the Longshoushan during the Early Paleozoic.

2 Geological setting

The Alxa Block is separated from the Hexi Corridor and the Qilian Orogenic Belt by the Longshoushan Fault to the south, from the Tarim Craton by the Altyn Fault to the west (Zhang and Gong, 2018), from the North China Craton by either the Bayanwulashan Fault, the western marginal fault of the Ordos Basin, or the Helanshan Fault to the east (Hui et al., 2021) (Fig. 1). On the northern margin of the Alxa Block, there are two important ophiolite belts: the Engger Us Ophiolite Belt in the north and the Qagan Qulu Ophiolite Belt in the south. Hf isotope data show that the magma sources of the Paleozoic magmatic rocks on both sides of the Qagan Qulu Ophiolite Belt are different (Zhang et al., 2015). Hence, recent studies have indicated that the Qagan Qulu Ophiolite Belt was most likely a Late Paleozoic suture related to the closure of a back-arc basin, representing the tectonic boundary between the Alxa Block and the Central Asian Orogenic Belt (Shi et al., 2014; Zhang et al., 2015; Hui et al., 2021; Zhang et al., 2023).




Figure 1 (a) Sketch geological map showing the tectonic location of the Alxa Block (modified from Song et al., 2013). (b) Simplified geological map of the Alxa Block Phanerozoic granitoids (modified from Zheng et al., 2014; Hui et al., 2021). AB- Alxa Block; CQB- Central Qilian Block; NCC- North China Craton.

The tectono-thermal events in the Alxa Block mainly occurred during the Paleoproterozoic, and during Paleozoic to early Mesozoic. During the Paleoproterozoic, the Alxa Block experienced a ~2.5 Ga magmatic-metamorphic event, ~2.3 Ga and 2.05-2.0 Ga magmatic events, as well as a 1.95-1.80 Ga magmatic-metamorphic event (Zhang et al., 2013; Gong et al., 2016; Zeng et al., 2018; Qi et al., 2019; Wang et al., 2019). The ~2.5 Ga magmatic-metamorphic event is relatively limited in

distribution and is mainly reported in the Beidashan (Zhang et al., 2013) and the Longshoushan (Qi et al., 2019). The ~2.3 Ga
85 and 2.05-2.0 Ga magmatic events are primarily found in the Bayanwulashan, Diebusuge and Longshoushan areas, and they
are generally believed to be related to an extensional tectonic setting (Dong et al., 2007; Dan et al., 2012; Zeng et al., 2018).
The 1.95-1.80 Ga metamorphic events are widely documented in the metamorphic basement throughout the Alxa Block (Zhang
et al., 2013; Gong et al., 2016; Zeng et al., 2018). It is generally believed to be associated with a late Paleoproterozoic orogeny
(Gong et al., 2016; Zeng et al., 2018), possibly recording the amalgamation of the North China Craton (Zhang et al., 2013;
90 Zeng et al., 2018).


During the Paleozoic to Early Mesozoic, the Alxa Block was mainly influenced by the Central Asian Orogenic Belt to
the north and the North Qilian Orogenic Belt to the south, resulting in extensive magmatic activities. Zhang and Gong (2018)
divided the magmatic activities during this period into three stages: 460-390 Ma, 360-300 Ma, and 299-230 Ma. These
magmatic activities in the Alxa Block generally exhibited a progressive younging trend from south to north. Compared to the
95 other two stages, the 460-390 Ma magmatic events were relatively limited in distribution within the Alxa Block. They were
mainly developed in the central-southern parts of the Alxa Block, such as the Beidashan and Longshoushan areas, as well as
in the eastern regions like the Yamatu and Langshan areas. The tectonic settings of the 460-390 Ma magmatism in the Alxa
Block are not fully understood. Some researchers believe that 460-390 Ma magmatic events are related to the subduction of
oceanic crust related to the Central Asian Orogenic Belt (Liu et al., 2016; Zhou et al., 2016; Xue et al., 2017). For example,
100 Xue et al. (2017) suggest that these rocks are a result of the slab rollback of the Paleo-Asian Oceanic slab, while Zhou et al.
(2016) propose a connection with post-arc rifting. However, some work suggests that the 460-390 Ma magmatic events in the
southern margin of the Alxa Block, particularly in the Longshoushan, are related to the North Qilian Orogenic Belt (Zeng et
al., 2016, 2021; Zhang et al., 2017; Liu et al., 2021). Additionally, some researchers suggest that the 460-390 Ma magmatic
events in the eastern part of the Alxa Block are the product of collision and accretion between the Alxa Block and the North
105 China Craton (Wang et al., 2015; Dan et al., 2016).

The magmatic events during the periods of 360-300 Ma and 299-230 Ma are widely distributed in the Alxa Block,
especially in the Yabulai-Nuorigong-Honggueryulin zone and the Beidashan. The predominant rock type is granite, with
subordinate gabbroic rocks (e.g., Zhang and Gong, 2018; Dan et al., 2014; Zhang et al., 2016). These two periods of magmatic
events are generally believed to be associated with the subduction-collision and post-collisional evolution of the Central Asian
110 Orogenic Belt (Zhang and Gong, 2018; Liu et al., 2017).

3 Samples and petrography

In this study we investigated two plutons located in the east of the Longshoushan (Fig. 1b), including monzogranite and K-
feldspar granite. These two plutons occupy an area of approximately 12 km². In the study area, the early Precambrian
Longshoushan Complex, Neoproterozoic Dunzigou Formation, Hanmushan Formation and Neogene strata are well exposed
115 (Fig. 2). The Longshoushan Complex is mainly composed of biotite quartz schist and migmatite, while the Dunzigou

Formation and Hanmushan Formation are marine clastic and carbonate sedimentary rocks. The monzogranite and K-feldspar granite both intruded into the Early Precambrian Longshoushan Complex (Fig. 2). Four samples of monzogranite and four samples of K-feldspar granite were collected at locations shown in Fig. 2b.



120 **Figure 2 (a) Simplified geological map of the southwestern Alxa Block (Wang et al., 2020); (b) Simplified geological map of the east**
of the Longshoushan area. Data are in Supplementary materials Table S1 (All age data were obtained using the zircon U-Pb
method).

3.1 Monzogranite

The monzogranite is generally pale red in colour with fine to medium grained texture (Fig. 3a). The mineral assemblage
 125 contains K-feldspar (~32 %), plagioclase (~28 %), quartz (~35 %) and biotite (~5 %) as well as accessory minerals such as
 magnetite, zircon and apatite. K-feldspar grains are euhedral or subhedral, always exhibit grid twinning (Fig. 3b) and can
 contain rounded quartz and plagioclase inclusions (Fig. 3c). Plagioclase commonly show polysynthetic twinning, and are
 altered to sericite and carbonate, with rims surrounded by newly formed K-feldspar (Fig. 3b). Biotite is the main dark mineral,
 occurring in flaky shapes and filling in the interstices between the other minerals, with embayment features and chloritization
 130 (Fig. 3b).




Figure 3 Field photographs and photomicrographs of monzogranite (a, b, c) and K-feldspar granite (d, e, f) (b, c, e, f- perpendicular polarized light). Q-quartz; Kfs-feldspar; Pl- plagioclase; Mc-Microcline; Bt-biotite.

3.2 K-feldspar granite

135 The K-feldspar granite is generally pale red in colour with medium grained texture (Fig. 3d). The mineral assemblage contains K-feldspar (~55 %), quartz (~30 %), plagioclase (~10 %) and biotite (~5 %) as well as accessory minerals such as apatite, magnetite and zircon. K-feldspar has a grain size range between 3mm to 4 mm, mainly composed of microcline with grid twinning, and containing numerous plagioclase and quartz inclusions (Fig. 3e). Plagioclase usually exhibits polysynthetic twinning and has sericitization and carbonatization. Euhedral to subhedral crystals are tabular-shaped. The main dark mineral is biotite; grains have embayed crystal margins and chloritization (Fig. 3f).

140

4 Analytical methods

Representative rock samples were selected for geochemical analysis. The analyses of whole-rock major and trace elemental compositions were conducted at Analytical Chemistry & Testing Services (ALS) Chemex (Guangzhou) Ltd, by using X-ray fluorescence (XRF) (ME-XRF26) and ICP-MS (ME-MS81), respectively. The detailed methods were given by Zeng et al.

145 (2022).

Cathodoluminescence (CL) imaging of zircon were performed at Chengpu Geological Testing Co. Ltd, Langfang, China using the TIMA analysis. The LA-ICP-MS zircon U-Pb dating and trace element analyses were carried out at Wuhan Sample Solution Analytical Technology Co., Ltd., Wuhan, China. Laser ablation system is the COMPexPro 102 ArF excimer laser (wavelength of 193 nm and maximum energy of 200 mJ) with a spot size of 35 μm, repetition rate at 8 Hz. ICP-MS is Agilent

150 7900.Zircon 91500 and glass NIST610 were used as external standards for U-Pb dating and trace element calibration, respectively. GJ-1 and Plešovice were used for quality control. In the experiment, the weighted mean age of GJ-1 and Plešovice were 600.3 ± 3.7 Ma (n=7, MSWD = 0.04) and 337.5 ± 3.0 Ma (n=4, MSWD = 0.03), respectively. In this paper, the $^{206}\text{Pb}/^{238}\text{U}$ age and $^{207}\text{Pb}/^{206}\text{U}$ ages are determined for younger zircons (<1000 Ma) and older grains (>1000 Ma). The detailed methods were given by Zong et al. (2017).

155 Zircon in-situ Lu–Hf analyses were undertaken at the Wuhan Sample Solution Analytical Technology Co., Ltd, Hubei, China using a Neptune Plus MC-ICP-MS. An ArF excimer laser ablation system of GeoLas HD was used with 44 μm spot size. The detailed analytical program is the same as outlined by Hu et al. (2012). In order to ensure the reliability of the analysis data, three international zircon standards of Plešovice, 91500 and GJ-1 are analyzed simultaneously with the actual samples. During our analyses, the value of Plešovice, 91500 and GJ-1 were 0.282472–0.282495, 0.282302–0.282314 and 0.282024–
160 0.282032 respectively, consistent with their recommended values (Plešovice: 0.282482 ± 23 ; 91500: 0.282308 ± 106 ; GJ-1: 0.282010 ± 89 , Zhang et al., 2020).

5 Analytical results

The data for major and trace elements, zircon U-Pb ages, zircon trace elements, and zircon Hf isotopes are shown in Tables S2, S3, S4 and S5, respectively.

165 5.1 Geochemical characteristics

5.1.1 Monzogranite

The samples of monzogranite have SiO_2 contents ranging from 71.59 wt.% to 72.05 wt.%. They have low Na_2O (3.75–3.88 wt.%) and high K_2O (4.40–4.52 wt.%). All samples fall in the granite area in the TAS classification (Fig.4a), and the Alkali - calcic to Calc-alkaline series areas in the $\text{SiO}_2\text{-N}_2\text{O}+\text{K}_2\text{O}$ diagram (Fig.4b). Contents of Al_2O_3 are 14.40–14.47 wt.%, A/KNC
170 are 1.05–1.07. All compositions fall in the weakly peraluminous field of the A/NC-A/KNC diagram (Fig.4c).




Figure 4 Geochemical classification diagrams. (a) TAS diagram (after Middlemost, 1994); (b) A/CNK-A/NK diagram (after Maniar and Piccoli, 1989); (c) $\text{Na}_2\text{O}+\text{K}_2\text{O}-\text{CaO}$ vs. SiO_2 diagram (after Frost et al., 2001). The source of the published data can be found in Supplementary materials Table S1.

The La_N/Yb_N values of monzogranite range from 22.26 to 27.84, showing enrichment of LREE and depletion of HREE. On chondrite-normalized REE patterns (Fig. 5c), the monzogranite has relatively inclined LREE patterns and flat HREE patterns. All the samples display negative Eu anomalies ($\delta\text{Eu}=0.73\text{--}0.75$) and slight positive Ce anomalies ($\delta\text{Ce}=0.99\text{--}1.22$). In the primitive mantle-normalized trace element diagram (Fig. 5d), these samples show enrichment of LILEs (e.g., Rb and K), depletion of HFSEs (e.g., Nb, Ta, Ti and P), and no depletion of Hf and Zr.




Figure 5 Chondrite-normalized REE patterns and primitive mantle-normalized trace element patterns for Early Paleozoic magmatic rocks (chondrite and primitive mantle values are from Sun and McDonough, 1989). The source of the published data can be found in Supplementary materials Table S1.

5.1.2 K-feldspar granite

185 The samples of K-feldspar granite have SiO₂ contents from 75.77 wt.% to 76.18 wt.%. They have low Na₂O (3.09–3.13 wt.%) and high K₂O (5.54–6.18 wt.%). All samples fall in the granite area in the TAS classification (Fig.4a), and the Alkali -calcic series and Calc-alkaline series areas in the SiO₂-N₂O+K₂O diagram (Fig.4b). Contents of Al₂O₃ are 12.33–12.36 wt.%, A/KNC are 1.01–1.03, and all samples are weakly peraluminous in the A/NC-A/KNC diagram (Fig.4c).

190 The La_N/Yb_N values of K-feldspar granite samples are from 6.38 to 11.94, showing enrichment of LREE and depletion of HREE. On chondrite-normalized REE patterns (Fig. 5c), the K-feldspar granite has relatively inclined LREE patterns and flat HREE patterns. All samples display clear negative Eu anomalies ($\delta\text{Eu}=0.19\text{--}0.41$) and slightly positive Ce anomalies ($\delta\text{Ce}=1.08\text{--}1.35$). In the primitive mantle-normalized trace element diagram (Fig. 5d), these samples show enrichment of LILEs (e.g., Rb, Ba and K), strong depletion of HFSEs (e.g., Nb, Ta, Ti and P), and no depletion of Hf and Zr.

5.2 Zircon U-Pb geochronology, trace elements and Hf isotopes

195 5.2.1 Monzogranite

Zircons from monzogranite (JC-20-9) are mostly euhedral, transparent, colorless, and 80–160 μm in length. The zircons have bright CL intensity and oscillatory zoning in the CL images (Fig. 6a), with Th/U ratios of 0.10–1.27. 25 spots cluster on the concordia curve with ²⁰⁶Pb/²³⁸U ages of 429–449 Ma, defining a weighted mean age of 440.8±2.1 Ma (MSWD=1.6) (Fig. 6a, b).




Figure 6 Concordia diagrams for zircon LA-ICP-MS U-Pb analyses.

Sixteen zircons were analyzed for REE. The spots have Σ REE of 704–2660 ppm (average 1270 ppm), Σ LREE of 18–212 ppm (average 89 ppm) and Σ HREE of 623–2550 ppm (average 1181 ppm), Σ LREE/ Σ HREE of 0.02–0.17, δ Eu of 0.12–0.38 (average 0.24) and δ Ce of 1.16–125.04 (average 18.38). On chondrite-normalized REE patterns (Fig. S1), these spots show depletion of LREEs, enrichment of HREEs, and clear negative Eu anomalies and positive Ce anomalies.

Fifteen zircons were analyzed for Lu-Hf isotopes. The zircons have $^{176}\text{Hf}/^{177}\text{Hf}$ ranging from 0.282295 to 0.282617, with $\epsilon\text{Hf}(t)$ of 3.72 to -7.83 and two-stage Hf isotope depleted mantle model age ($T_{\text{DM}2}$) of 1187 Ma to 1917 Ma by using the weighted mean age.

5.2.2 K-feldspar granite

Zircons from K-feldspar granite (JC-20-19) are mostly euhedral, transparent, colorless, and 80–130 μm in length, and have bright CL intensity and oscillatory zoning (Fig. 6c) with Th/U ratios of 0.29–1.01. Among these, 17 spots cluster on the concordia curve with $^{206}\text{Pb}/^{238}\text{U}$ ages of 432–446 Ma, defining a weighted mean age of 439.4 ± 2.0 Ma (MSWD=0.88) (Fig. 6c, d). Seven spots have ages of 825–2344 Ma. Among them, three zircons are weakly luminescent and lacking discernible internal structure with Th/U ratios of 0.03–0.09. They have $^{207}\text{Pb}/^{206}\text{Pb}$ ages of 1857 Ma, 1847 Ma and 1894 Ma, respectively. Three zircons have bright CL intensity and oscillatory zoning with Th/U ratios of 0.48–1.00. They have $^{207}\text{Pb}/^{206}\text{Pb}$ ages of 2216 Ma, 2344 Ma and 2307 Ma, respectively.

Thirteen zircon spots, which define the weighted mean age of 439.4 ± 2.0 Ma, have ΣREE of 542–1414 ppm (average 862 ppm), ΣLREE of 26–260 ppm (average 80 ppm) and ΣHREE of 502–1154 ppm (average 783 ppm), $\Sigma\text{LREE}/\Sigma\text{HREE}$ of 0.04–0.23, δEu of 0.15–0.58 (average 0.34) and δCe of 1.78–129.07 (average 20.97). All the spots show depletion of LREEs, enrichment of HREEs, and clear negative Eu anomalies and positive Ce anomalies.

Thirteen zircons were analyzed for Lu-Hf isotopes from sample JC-20-19. Twelve spots, which define the weighted mean age of 439.4 ± 2.0 Ma, show $^{176}\text{Hf}/^{177}\text{Hf}$ and $\epsilon\text{Hf}(t)$ values of 0.282494 to 0.282056 and -0.49 to -16.27, respectively, with $T_{\text{DM}2}$ of 1453 Ma to 2446 Ma by using the weighted mean age. One older zircon (825 Ma) has a $^{176}\text{Hf}/^{177}\text{Hf}$ ratio of 0.281812, with $\epsilon\text{Hf}(t)$ of -16.07, and $T_{\text{DM}2}$ of 2722.

225 6 Discussion

6.1 Formation age

In this study, zircons from the monzogranite and K-feldspar granite can be divided into two groups (I-type and II-type, Fig. 7a). The I-type zircons (the zircons from the monzogranite and the majority of zircons from the K-feldspar granite) have high Th/U ratio values (0.10-1.27), as well as bright intensity and oscillatory zoning in the CL images. The above characteristics are consistent with those of igneous zircon (Hoskin and Schaltegger, 2003). Hence, the weighted mean ages of 440.8 ± 2.1 Ma (MSWD=1.6) and 439.4 ± 2.0 Ma (MSWD=0.88) are taken to represent the magmatic emplacement age of monzogranite and K-feldspar granite, respectively. Four analyses from sample JC-20-9 have $^{207}\text{Pb}/^{206}\text{Pb}$ ages of 825 Ma, 2216 Ma, 2307 Ma and 2344 Ma, which are much older than the emplacement ages of the K-feldspar granite. Hence, these zircons are captured or inherited magmatic zircons. The II-type zircons (spots #2, #20 and #22 from JC-20-9) have low Th/U ratio values (0.03-0.09), as well as being weakly luminescent and lacking discernible internal structure in the CL images, which suggests that their ages most likely represent the timing of metamorphism (Hoskin and Schaltegger, 2003). The spots from the II-type zircons have ages of 1847 – 1894 Ma, which is consistent with the age of the metamorphic events in the Longshoushan area during the Paleoproterozoic (Gong et al., 2016; Zeng et al., 2018).




Figure 7 Age-Th/U (a), T- $\log(f_{O_2})$ (b) and T- δ_{Ce} (c) diagrams of zircon

6.2 Crystallization conditions

Temperature, oxygen fugacity, and pressure are important indicators for the crystallization conditions of magmatic rocks. They play a significant role in investigating the origin and evolution of magmas, as well as their relationship with mineralization processes (Xiao et al., 2017; Zeng et al., 2022).


Zircon saturation thermometry (T_{Zm}) and the Ti-in-zircon thermometer (T_{Zr-Ti}) are commonly used methods for estimating magma temperatures. Based on the calculation method introduced by Watson and Harrison (1983), the T_{Zm} values for the monzogranite and K-feldspar granite are in the range of 839-842 °C (with an average value of 840 °C) and 745-817 °C (with an average value of 782 °C), respectively. The solubility of Ti in zircon is related to the temperature during zircon formation, and the activities of TiO_2 (a_{TiO_2}) and SiO_2 (a_{SiO_2}) in the melt (Watson and Harrison, 2005; Ferry and Watson, 2007). Since both monzogranite and K-feldspar granite contain abundant quartz and zircon, but rare ilmenite and xenotime, the a_{SiO_2} and a_{TiO_2} are 1 and 0.5, respectively (Schiller and Finger, 2019). According to the calculation method proposed by Ferry and Watson (2007), the Ti temperatures of 16 autocrystic zircons from the monzogranite range from 727 to 883 °C (with an average of 791 °C), while the Ti temperatures of 13 autocrystic zircons from the K-feldspar granite range from 687 to 851 °C (with an average of 756 °C). In addition, the Al_2O_3/TiO_2 ratio of the whole rock can be used as an indicator of the formation temperature of granites (Hui et al., 2021). Hot granites typically have lower Al_2O_3/TiO_2 ratios, while cold granites have higher Al_2O_3/TiO_2 ratios (Sylvester, 1998). The samples of monzogranite have lower Al_2O_3/TiO_2 ratios (72.0 to 72.4) than that of the samples of K-feldspar granite (154.5 to 246.6), implying that the melts of monzogranite are hotter than those of K-feldspar granite. This result is in consistent with the outcomes obtained from zircon saturation thermometry (T_{Zm}) and the Ti-in-zircon thermometer.

Due to the distinct partitioning behavior of Ce^{4+} and Ce^{3+} in zircon, the cerium in zircon can reflect the oxygen fugacity condition of the magmatic systems (Trail et al., 2012). Using the formulas proposed by Trail et al. (2012), the $\log fO_2$ for the monzogranite and K-feldspar granite falls within the range of -27 to -10 (with an average of -18) and -26 to -6 (with an average of -19), respectively. In the $\log fO_2$ -T (°C) and Ce/Ce^*-T (K) diagrams (Fig. 7b and c), the monzogranite and the K-feldspar granite mostly fall below the FMQ buffer, indicating low oxygen fugacity conditions.

Generally, when garnet is the dominant residual phase in the source region, the melt that forms will exhibit a significant inclined HREE distribution pattern, with $Y/Yb > 10$ and $Ho/Yb > 1.2$ characteristics (Tang et al., 2021). In this study, the HREE distribution patterns of the monzogranite and K-feldspar granite are relatively flat, with Y/Yb ranging from 7.91 to 10.93, and Ho/Yb ranging from 0.33 to 0.38, indicating that garnet is not the dominant residual phase. Therefore, the pressure of the granite source region should be less than 1.7 GPa (mostly less than 1.5 GPa, Tang et al., 2021). The Yb contents of the monzogranite and the K-feldspar granite are both less than 2 ppm (except for one sample), and the Sr contents of the two plutons are both less than 400 ppm, indicating the characteristics of low Yb and low Sr granite (Zhang et al., 2006). In addition, these two plutons have the characteristic low Al_2O_3 contents and a distinct negative Eu anomaly. These features are consistent with the low Sr and low Yb granites in North Hebei, implying that the monzogranite and the K-feldspar granite are partial melts of high-pressure granulites containing plagioclase, clinopyroxene, garnet, and amphibole, at the bottom of the thickened lower crust (Li et al., 2004; Zhang et al., 2006), with source pressures estimated to be between 0.8 and 1.5 GPa (Zhang et al., 2010).

6.3 Granitoid petrogenesis

The monzogranite and K-feldspar granite have low Zr (58–175 ppm), Ce (37.6–94.5 ppm), Zr + Nb + Ce + Y (129.6–296.9 ppm) and FeO^*/MgO (4.3– 12.2), distinct from typical A-type granites ($\text{Zr}>250$ ppm, $\text{Ce}>100$ ppm, $\text{Zr} + \text{Nb} + \text{Ce} + \text{Y}>350$ ppm, $\text{FeO}^*/\text{MgO}>16$; Whalen et al., 1987). In addition, the magma temperature of the monzogranite ($T_{\text{Zr}}: 839\text{--}842$ °C; $T_{\text{Zr-Ti}}: 728$ to 883 °C) and K-feldspar granite ($T_{\text{Zr}}: 745\text{--}817$ °C; $T_{\text{Zr-Ti}}: 687$ to 851 °C) is significantly lower than that of typical A-type granite (> 900 °C, Skjerlie and Johnston, 1992). Furthermore, all compositions fall in the region of I- and S- type granite rather than the A-type granite in the discrimination diagrams (Fig. 8a-c). Thus, the monzogranite and K-feldspar granite are not A-type granites and are more compatible with being I or S-type granites. The samples of monzogranite and K-feldspar granite have $\text{A/CNK} < 1.1$, normative corundum $< 1\%$ in the CIPW norm calculation and there is no presence of peraluminous minerals, such as cordierite, andalusite, muscovite and garnet, which is distinctly different from the mineralogy of S-type granites (Chappell and White, 1992). As shown in Th-Rb and SiO_2 - P_2O_5 diagrams (Fig. 8d and e), the positive trend between Th and Rb and the negative trend between P_2O_5 and SiO_2 also demonstrate the evolutionary trend of typical I-type granite (Wolf and London, 1994; Kong et al., 2018; Liu et al., 2021). Moreover, In the P-(REE+Y) diagram (Fig. 8f), autocrystic zircons in both the monzogranite and the K-feldspar granite have lower P concentrations, and the concentration of P and REE+Y does not show an obvious positive correlation, showing the characteristics of zircons in I- type granite rather than S-type granite.



295 **Figure 8** Discrimination diagram of granitoid genetic types. (a, b, c) 10000 Ga/Al vs. Zr, 10000 Ga/Al vs. Y and Zr+Nb+Ce+Y vs. TFeO/MgO (after Whalen et al., 1987); (d) Rb vs. Th; (e) SiO₂ vs. P₂O₅; (f) P vs. REE+Y (after Burnham and Berry, 2017).

As noted, the monzogranite and K-feldspar granite have close temporal and spatial relationship, similar Na₂O+K₂O contents (Fig. 4a), uniform REE and trace element patterns (Fig 5c, d). These characteristics suggest a co-magmatic origin. The genesis of highly fractionated I-type granite has been proposed to two primary mechanisms: (1) fractional crystallization

(FC) or assimilation fractional crystallization (AFC) of mantle-derived mafic magmas (He et al., 2019), or (2) partial melting
 300 of crustal materials with involvement of some mantle-derived components, accompanied by subsequent fractional
 crystallization (Karsli et al., 2010; Liu et al., 2021).




Figure 9. SiO₂ vs. major element oxides (a–e), SiO₂ vs. Sr (f), Sr vs. Rb/Sr (h), Sr vs. Ba(g), and La vs. (La/Yb)_N (i) for the Dakouzi pluton (the partition coefficients of Rb, Sr, and Ba are from Philpotts and Schnetzler, 1970; the partition coefficients of La and Yb are from Ewart and Griffin, 1994)

305

Pl- plagioclase; **Kfs-** K-feldspar; **Bt-** biotite; **Zr-** zircon; **Sph-** sphene; **Ap-** apatite; **Mon-** monazite; **Allan-** allanite.

Mantle-derived primary melt would be expected to have high MgO, TFe₂O₃, CaO and Cr contents (Ma et al., 2013), which are obviously inconsistent with the characteristics of the monzogranite and K-feldspar granite in this study. Furthermore, mafic melts can only form a minor volume of granitic melts through crystal differentiation, with a proportion of about 9:1
 310 (Zeng et al., 2016). However, in the Longshoushan, most of the ~440 Ma magmatic rocks are granitic rocks, and no large-

scale exposures of mafic-ultramafic rocks have been reported. Therefore, monzogranite and K-feldspar granite are unlikely to have been formed by the FC or AFC processes of mantle-derived magma.

The Hf isotopes of zircon are not affected by fractional crystallization, and thus can provide precise constraints on the origin of magma (Griffin et al., 2002). The zircon $\epsilon\text{Hf}(t)$ values of the monzogranite range from -0.37 to -16.28, while those of the K-feldspar granite range from 3.53 to -7.74, both exhibiting a large range of $\epsilon\text{Hf}(t)$ (Fig. 10). This phenomenon indicates the existence of different Hf isotopic end-members in the source regions of these two plutons, which could be attributed to the mixing of mafic and felsic melts (Zhang et al., 2015; He et al., 2019), heterogeneous composition of source crust materials (Zeng et al., 2016), or variable degrees of melting of source minerals (Zeng et al., 2021). In addition, some quartz and biotite boundaries have been melted and transformed into irregular embayments, and plagioclase inclusions occur within the quartz. These observations suggest the injection of high-temperature magma during the magma evolution process (Duan et al., 2021). On the Rb/V-1/V and La/Cr-1/Cr diagrams (Fig. 11a and b), samples from monzogranite and K-feldspar granite exhibit a linear trend, indicating magma mixing (Liu et al., 2021). On the MgO-FeO_T and SiO₂/MgO-Al₂O₃/MgO diagrams (Fig. 11c and d), our samples also show an obvious magma mingling trend (Tang et al., 2021). In addition, in the Longshoushan, Zhang et al. (2021) reported an age of ~441 Ma for the Xijing clinopyroxene diorite vein, and Wang et al. (2019) reported crystallization ages of ~440 Ma for the mantle-derived mafic microgranular enclaves in the Jiling granite. Therefore, there was mantle-derived magmatism in the Longshoushan at ~440 Ma, which could provide the mantle material for the granitic magmatism. We use the ~441 Ma Xijing clinopyroxene diorite (Zhang et al., 2021) and our sample of K-feldspar granite (JC-20-20) to assess magma mixing simply in terms of two end-member compositions. Our simulation results show a curvilinear evolutionary relationship on the Th/Nd-Th and Rb/Nd-Rb diagrams (Fig. 11e and f), which is also consistent with magma mixing (Schiano et al., 2010; Tang et al., 2021). To conclude, the monzogranite and K-feldspar granite in this study were derived by magma mixing of mantle and crust-derived magmas.




Figure 10 Zircon $\epsilon_{\text{Hf}}(t)$ -age (Ma) diagram for samples in this study and published data for the region. The source of the published data can be found in Supplementary materials Table S1 (The red arrow represents the variation trend of $\epsilon_{\text{Hf}}(t)$).

335




Figure 11 Diagram of magma mixing discrimination. (a) Rb/V vs. 1/V diagram; (b) La/Cr vs. 1/Cr diagram; (c) MgO vs. FeO^T diagram; (d) SiO₂/MgO vs. Al₂O₃/MgO diagram; (e) Th vs. Th/Nd diagram; (f) Rb vs. Rb/Nd diagram (a and b after Schiano et al., 2010; e after Zorpi et al, 1989)

340 In crust-mantle mixing processes, crust-derived magmas typically have lower $\epsilon Hf(t)$ values. Therefore, lower $\epsilon Hf(t)$ values can roughly reflect the composition of the crustal source. As shown in Fig. 10, some spots with lower $\epsilon Hf(t)$ values from both rock bodies fall within the evolutionary trend line of the Early Precambrian basement strata, known as the Longshoushan Complex, into which the studied plutons are intruded. Furthermore, abundant 1.85-1.90 Ga metamorphic zircons and ~2.3 Ga plutons have been identified in the Longshoushan Complex (Gong et al., 2016; Zeng et al., 2018), which

345 corresponds to the inherited zircon ages found in the K-feldspar granite. Therefore, the Longshoushan Complex is mostly likely the crustal source of the granitoids in this study. Compared with the K-feldspar granite, the monzogranite has higher $\epsilon Hf(t)$ values and is closer to the mantle end-member in the mixing diagrams, indicating a relatively lower proportion of crust-derived magmas.

All the samples of the monzogranite and the K-feldspar granite are characterized by relatively high SiO₂, total alkali and differentiation index, low TFe₂O₃, MgO and CaO contents, low Nb/Ta (9.5–14.5) and Zr/Hf values (26.3–40.2), and enrichment of Rb, Th and U, indicating that these rocks are highly fractionated (Xiao et al., 2014; Wu et al., 2017). This is consistent with the samples of these two granites being in the fractionated felsic granites (FG) area in the FeO^T/MgO vs. Zr+Nb+Ce+Y diagram (Fig. 8c). In this study, the granitoids have negative Eu-Ba-Sr anomalies (Fig. 5c and d), indicating plagioclase and K-feldspar fractionation (Harris et al., 1990). On the Rb/Sr vs. Sr diagrams (Fig. 9g), the granitoid samples show a separation trend of plagioclase. The negative correlation of SiO₂ with TFeO and MgO suggests fractionation of amphibole and biotite during magmatic evolution (Fig. 9a and d). As shown in the Sr-Ba diagram (Fig. 9h), amphibole is likely to dominate the fractional phase. The increasing TiO₂ content with decreasing SiO₂ (Fig. 9e), along with the clear Ti, Nb and Ta anomalies in the primitive-mantle-normalized trace element diagrams (Fig. 5b), implies the fractionation of rutile (Foley et al., 2000). In a diagram of (La/Yb)_N vs. La (Fig. 9i), the variation of REE contents suggests the fractionation of monazite and allanite. Hence, the monzogranite and K-feldspar granite have experienced extensive fractional crystallization of plagioclase, K-feldspar, amphibole, biotite, rutile, monazite and allanite.

6.4 Tectonic implications

6.4.1 Tectonic affinity of the Longshoushan in the Paleozoic

During the Paleozoic-Mesozoic, the Alxa Block was situated between multiple tectonic domains, with its northern and southwestern sides experiencing subduction-accretionary orogenesis of the Central Asian Orogenic Belt and the North Qilian Orogenic Belt, respectively (Song et al., 2013; Liu et al., 2017; Xue et al., 2017; Zhang and Gong et al., 2018; Hui et al., 2021). The North Qilian Orogenic Belt is a component of the much larger Central China Orogenic Belt, which extends ~2000 km east and west of the Alxa Block.

It is widely believed that the Yabulai-Nuoergong-Honggueryulin zone and Zongnaishan-Shalazhashan zone at the northern side of the Alxa Block were mainly influenced by the Central Asian Orogenic Belt (Li et al., 2017; Liu et al., 2017; Sun et al., 2017; Zhang and Gong et al., 2018; Hui et al., 2021). As shown in Fig.13a, magmatic activities in these areas were mainly concentrated in the late Paleozoic-Mesozoic, with a peak age of ~270 Ma. The magmatic activity in the Longshoushan, which lies at the southern margin of the Alxa Block, mainly occurred in the Early Paleozoic with a peak age of ~440 Ma (Fig. 12d), which is significantly different from the magmatic events in the northern side of the Alxa Block and basically consistent with the Hexi Corridor, North Qilian Orogenic Belt and Central Qilian Block (Fig. 12e-g), as well as the wider Central China Orogenic Belt (Allen et al., 2023). Therefore, the Longshoushan was most likely part of the evolution of the Central China Orogenic Belt during the Paleozoic, and specifically adjacent regions of the North Qilian Orogenic Belt and the Hexi Corridor. There is a small amount of Late Paleozoic magmatic rocks in the Longshoushan, such as the Taohualashan pluton (Xue et al., 2017). These plutons are mainly distributed on the northwest side of the Longshoushan (Fig. 1b), close to the northern boundary of the Alxa Block (near the Central Asian Orogenic Belt), and are considered to related to the Central Asian Orogenic Belt

(Xue et al., 2017; Zhang et al., 2021). The Beidashan is located to the north of the Longshoushan, and both Early Paleozoic and Late Paleozoic-Mesozoic magmatic rocks are extensively exposed there (Fig. 12c). This feature indicates that the Beidashan is located at the intersection of two tectonic domains, and was influenced by both the Central Asian Orogenic Belt and the North Qilian Orogenic Belt. In addition, similar to the Longshoushan, the Early Paleozoic magmatic rocks in the Beidashan are mainly distributed on the southeastern side, while the Paleozoic-Mesozoic magmatic rocks are mainly distributed on the northwest side (Fig. 1b). Therefore, in the Alxa Block, the Longshoushan and the southeast side of the Beidashan are mainly influenced by the North Qilian Orogenic Belt, while other areas to the north are mainly influenced by the Central Asian Orogenic Belt. Apart from Beidashan and Longshoushan, Early Paleozoic magmatic rocks are mainly distributed in the eastern part of the Alxa Block, including Bayan Nuru, Bayanwulashan, and Langshan areas. Various authors suggested that these magmatic events are related to the convergence between the Alxa Block and the North China Craton (Dan et al., 2016; Wang et al., 2015).




Figure 12 Kernel density estimate (KDE) plots for Palaeozoic and Mesozoic magmatic ages in the Alxa Block, North Qilian Belt and Central Qilian Block. Ophiolitic rocks are not included. Data are in Supplementary materials Table S1. All dates are zircon U-Pb ages (histogram bin width is 20).

395

Figure 12 indicates that magmatism in the Alxa Block started later than 500 Ma which is later than in other parts of the Central China Orogenic Belt to the south. In addition, the Longshoushan area lacks the twin peaks in Early Paleozoic ages for magmatism found in other areas such as the North Qilian Orogenic Belt and Central Qilian Block, at ~500 Ma and 430 Ma (Allen et al., 2023). We suggest that the reason for these difference in age spectrum is that the Longshoushan was initially far enough away from the Proto-Tethyan subduction zone that it was not affected by arc magmatism, which was focused on the North Qilian Orogenic Belt and Central Qilian Block to the south. Detailed discussion can be found in Section 6.4.3.

400

6.4.2 Evaluation of crustal thickness

Hu et al. (2017) designed an empirical formula between the median Sr/Y ratio of intermediate-acid rocks and the crustal thickness in continental collisional orogens. In addition, Mantle and Collins (2008) designed the empirical relationship between the maximum Ce/Y ratio of basalt and crustal thickness. Therefore, Sr/Y and Ce/Y ratios of magmatic rocks can be used as proxies for crustal thickness. The Longshoushan experienced the convergence and collision of the Alxa Block and the basement of the North Qilian Orogenic Belt in the Early Paleozoic, so the intermediate-acid rocks and basic-ultrabasic rocks formed in this process can be used to estimate the thickness of the paleo-crust by using the formulas designed by Hu et al. (2017) and Mantle and Collins (2008), respectively.

In this study, we collected published data for 153 magmatic rock samples of Early Paleozoic age in the Longshoushan, including 8 ultrabasic rock samples, 31 basic rock samples, 45 intermediate rock samples and 88 acidic rock samples. According to the location, lithology and formation age of these rocks, the above samples are divided into 31 subsets in this study. Among them, 26 subsets of intermediate-acid rocks were run through the data processing scheme recommended by Hu et al., (2017) to process data. The first step is to eliminate the samples with $\text{SiO}_2 > 72\%$ and $\text{MgO} < 0.5\%$ or $> 6.0\%$ in each subset, thereby removing those that exceed the range of original data fitting the formula; The second step is to remove individual data points with Sr/Y outliers from each data subset using the modified Thompson tau statistical method. The third step is to eliminate the subset whose average Rb/Sr value is greater than 0.35, in order to filter out samples that have been strongly affected by fractionation. After the above process, there are 6 subsets of 32 samples left. According to the calculation formula of Hu et al., (2017), the crustal thickness is calculated by the median ratio of Sr/Y in each subset of samples. On the other hand, 5 subsets of basic-ultrabasic rock samples were processed by the Mantle and Collins (2008) method. The first step is to eliminate the samples with $\text{SiO}_2 < 44\%$, $\text{MgO} < 4\%$ and $\text{LOI} > 4\%$. In the second step, samples of alkaline and shoshonitic basalts derived from the enriched lithospheric mantle are excluded, as these rocks typically have abnormally high Ce/Y ratios (> 4), rendering the results unusable. After the above process, there are 4 groups of 28 samples left. According to the calculation formula of Mantle and Collins (2008), the corresponding crustal thickness is obtained through the maximum Ce/Y ratio in each subset. See Table S6 the calculation results. As shown in Fig. 13, the thickness of the crust underwent a transition from thickening to thinning during the period of 460-410 Ma, with the initial timing of the transition being ~435 Ma.




Figure 13 Plot of crustal thickness versus age of magmatic rocks for each data subset in the Longshoushan, southwestern Alxa Block. Data are in Supplementary materials Table S1.

430 6.4.3 Tectonic evolution of the Longshoushan in the Early Paleozoic

A notable feature of the Longshoushan is the large volume of Late Ordovician-Silurian magmatic rocks (Fig. 2). Based on the statistics and analysis of U-Pb ages and geochemical data of these rocks, three stages of magmatic activities can be defined (Fig. 14).








Figure 14 Integrated column of magmatism evolution of Early Paleozoic in the Longshoushan area. On the left is the histogram of age distribution of Early Paleozoic magmatic rocks in the Longshoushan area. Data are in Supplementary materials Table S1

The first stage (460 Ma–444 Ma): The magmatic rocks in this period are all acidic rocks (except one mafic rock, Fig. 14), calc-alkaline (alkali-calcic) granites (Fig. 4), which is consistent with the magmatic rock assemblage formed in a subduction environment (Song et al., 2015; Wang et al., 2020). Additionally, the 461–444 Ma magmatic rocks are enriched in LREEs and
440 LILEs, depleted in HREEs and HFSEs, and exhibit significant negative anomalies of Nb, Ta, P and Ti (Fig. 5a and b), which are consistent with the characteristics of arc-derived magmas (Jahn et al., 1999; Li et al., 2016). Hence, the magmatic events during this period are related to slab subduction. During this period, there are some A-type granites, which are possibly related to the extension caused by the rollback of the subducting slab. This phenomenon has been observed in the Central Asian Orogenic Belt as well as in eastern Australia (Cawood et al., 2011; Wu et al., 2021).

445 In the North Qilian Orogenic Belt, there are two ophiolite belts, the Northern Ophiolite Belt (NOB) and the Southern Ophiolite Belt (SOB), separated by the North Qilian Volcanic Arc (NQVA). Allen et al. (2023) identified the oceans represented by these two ophiolite belts as the main Proto-Tethys ocean and North Qilian back-arc basin (NQ bab), respectively. The Dachadaban tholeiite-boninite suite (517-487 Ma) and Laohushan IAB-type boninite (492-488 Ma) are considered to have been formed by the northward subduction of the Proto-Tethys on the south side of the North Qilian Volcanic Arc (NQVA)
450 (Xia et al., 2012; Fu et al., 2022). Their formation ages indicate that this subduction was already underway in the mid-late Cambrian (Fig. 15a). Under this tectonic setting, intense magmatic activity occurred in the North Qilian Volcanic Arc (NQVA) and the Southern Ophiolite Belt (SOB) during the mid-late Cambrian and early Ordovician. In comparison, the initiation age of the northward subduction of the North Qilian back-arc basin (NQ bab) is later. The formation age of SSZ-type ophiolites (gabbro, basalt, and plagiogranite) formed in the northward subduction of the North Qilian back-arc basin (NQ bab) is 454-
455 448 Ma (Fu et al., 2020), while the age of metamorphism of the Laohushan boninite related to this subduction event is ~455 Ma (Fu et al., 2022). Therefore, there is no intense Cambrian magmatic activity in the Northern Ophiolite Belt (NOB), Hexi Corridor (HC), and Longshoushan. The magmatic activity in these areas gradually intensified in the Middle-Late Ordovician, which is consistent with the North Qilian back-arc basin (NQ bab) undergoing northward subduction by this time (Fig. 15b).



Abbreviations are as follows. CEKT: Central East Kunlun Terrane. CQB: Central Qilian Block. HC: Hexi Corridor. KC: Kuangping Complex. NOB: Northern Ophiolite Belt. NQgB: North Qinling Belt. NQUB: North Qaidam Ultra High-pressure Belt. NQVA: North Qilian Volcanic Arc. OB: Oulongbuluke Block. SKT: South Kunlun Terrane. SOB: Southern Ophiolite Belt. SQgB: South Qinling Block. QB: Qaidam Block.

The second stage (444–435 Ma): Compared to the first stage of magmatic activity, the intensity of the 444–435 Ma magmatic activity significantly increased, and the rocks have higher total alkali content (Fig. 4a and b). In this stage, although calc-alkaline granitic magmatism was still dominant, there are also some alkaline rocks. In addition, as previously mentioned, the ~440 Ma monzogranite and ~439 Ma K-feldspar granite were formed by crust and mantle-derived magma mixing. During this period, mantle-derived intermediate-basic rocks also existed in the Longshoushan (Figure 7). Therefore, there was mantle-derived magmatic activity in the Longshoushan from 444 to 435 Ma. The above magmatic rock assemblage is consistent with the post-collisional magmatic rock assemblage formed after a major ocean closed (Zhang et al., 2019; Wang et al., 2020). Noting that the meaning of post-collisional has multiple interpretations, the post-collisional stage refers to the stage after the initial collision between continental plates or between continental plates and island arcs in this study (Liegeois et al., 1998). In this study, the zircon $\epsilon\text{Hf}(t)$ values of the monzogranite and K-feldspar granite are primarily negative, with a minimum value reaching -16.27. As shown in Fig. 10, the Hf isotope characteristics indicate a significant decrease in zircon $\epsilon\text{Hf}(t)$ during the 444–435 Ma period compared to the previous stage (Fig. 10), reflecting the increasing contribution of ancient continental crust components in the magma source region. This phenomenon is also consistent with the variations in magmatic rocks during the transition from subduction to collision (Li et al., 2023). As shown in Fig. 13, significant crustal thickening occurred in the Longshoushan during the period of 444–437 Ma. This is consistent with that the monzogranite and K-feldspar granite in this study were formed through partial melting near the base of the thickened lower crust, with source pressures approximately between 0.8 and 1.5 GPa. In addition, research has shown that thickened crust also occurred in the Hexi Corridor (HC), the Northern Ophiolite Belt (NOB) and the North Qilian Volcanic Arc (NQVA) on the south side of the Longshoushan during 450–430 Ma (Zhang et al., 2017; Yang et al., 2019). The above phenomena are consistent with the shortening and thickening of continental crust caused by the closure of ocean basins and subsequent continent-continent collision (Li et al., 2023). In addition, the latest arc-type magmatic rocks in the North Qilian Orogenic Belt were found to have formed at 446 Ma (Wang et al., 2005). Silurian molasse rocks are widely distributed and unconformably overlie pre-Silurian strata in this orogenic belt (Song et al., 2013; Xia et al., 2016; Zeng et al., 2021). Hence, it is generally believed that the closure of the NQ bab occurred at the end of the Ordovician (~444 Ma, Song et al., 2013; Zeng et al., 2021). Therefore, the Longshoushan transitioned from a subduction environment to a post-collisional environment from ~444 Ma (Fig. 15c).

Collectively, the characteristics of the Longshoushan rocks from 444–435 Ma are similar to the signatures of the magmatism after 440 Ma in the North Qaidam Ultra High-Pressure Belt (NQUB) (Allen et al, 2023), presently ~500 km to the southwest of the Longshoushan. This coincidence in timing and evolution requires either a chance juxtaposition of separate

arcs and continents across separate subduction zones, or, and more likely in our view, the duplication of sections of what was
495 once a single subduction and collision system.

The third stage (435–410 Ma): During the evolution of orogenic events, calc-alkaline granites are commonly replaced
gradually by high-silica alkaline to peralkaline magmatic rocks (Fig. 4), representing the transition from the post-collisional
compressional environment to an extensional environment (Liégeois et al., 1998; Bonin, 2004). In the Longshoushan, the 444-
435 Ma magmatic rocks are mainly high potassium calc-alkaline I-type granite (Fig. 14). However, the proportion of alkaline
500 rock series significantly increased and multiple A-type granites appeared during 435–410 Ma, indicating the characteristics of
an intracontinental extensional environment. During this period, a significant number of mantle-derived mafic dykes were
intruded, which also reflects an extensional setting (Zeng et al., 2021). Therefore, the Longshoushan transitioned from a post-
collisional compressional environment to an extensional setting at ~435 Ma. As discussed earlier, the transition in crustal
thickness from thickening to thinning in the Longshoushan also occurred at ~435 Ma, which is consistent with the inferred
505 stress changes in the study area (although we cannot rule out efficient erosion as the cause of crustal thinning). The driver for
the proposed extension is not clear, but it would have been enhanced by a crust with initially greater thickness than normal.

There are two different views regarding the mechanism of the transition from a compressional environment to an
extensional environment in the North Qilian orogenic belt: slab break-off (Xia et al., 2016; Zeng et al., 2021) and lithospheric
delamination (Yu et al., 2015; Zhang et al., 2017). Studies have shown that the mantle-derived magmas in this stage mainly
510 derived from the enriched lithospheric mantle, indicating that the lithospheric mantle in the Longshoushan has not been
completely delaminated (Zeng et al., 2021). Small-scale delamination of hydrated lithosphere is a possible generation
mechanism (Kaisalaniemäki et al., 2014) (Fig. 15d). A complementary mechanism is adiabatic melting during extension and
lithospheric thinning, supported by the observations of crustal thinning over time. These mechanisms are not mutually
exclusive. While we cannot fully rule out slab break-off, we note modelling studies that suggest it does not normally generate
515 significant magmatism (Freeburn et al., 2017).

7 Conclusions

- (1) LA-ICP-MS zircon U-Pb dating indicates that the monzogranite and the K-feldspar granite in the east of Longshoushan
formed at 440.8 ± 2.1 Ma (MSWD=1.6) and 439.4 ± 2.0 Ma (MSWD=0.88), respectively.
- (2) The monzogranite and K-feldspar granite in the east of the Longshoushan were generated through various degrees of
520 mixing between crustal-derived magma from the thickened lower crust and mantle-derived magma, and then underwent
extensive fractional crystallization with removal of plagioclase, K-feldspar, amphibole, biotite, rutile, monazite and allanite.
- (3) Based on the temporal and spatial distribution of the magmatic rocks in the Longshoushan, we propose a three-stage
tectonic evolution model. ① 461 to 445 Ma: The northward subduction of NQ bab resulted in the develop of arc-type magmatic
rocks. ② 444 to 435 Ma: The closure of the NQ bab and subsequent continent-continent collision resulted in significant crustal

525 thickening in the study area. ③ 435 to 410 Ma: Tectonics transitioned from a collisional compressional environment to an extensional environment, evidenced by crustal thinning at this time.

Data availability. The data presented in the supplement.

Supplement. The supplement related to this article (Tables S1, S2, S3, S4, S5 and S6) is available online at:

530 *Author contributions.* Renyu Zeng: Conceptualization, Methodology, Writing - original draft. Hui Su: Writing- original draft, visualization. Mark B. Allen: Writing - review & editing; Haiyan Shi: Visualization; Chenguang Zhang: Visualization; Jie Yan: Funding acquisition.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests

Acknowledgements

535 We thank Prof. M. Santosh, Dr. Zhijie Song and Dr. Liangpeng Deng for constructive reviews and useful suggestions. We are also grateful to Qixing Ai and Yuhua Wang for their help with the field work. This research was funded by the Jiangxi Provincial Natural Science Foundation (No. 20232BAB213061), the China Uranium Industry Co. LTD. - East China University of Technology Innovation Partnership Foundation (2023NRE-LH-12), National Nature Science Foundation of China (Grants No. 42030809, 42262017, 42002095; 42162013), China Scholarship Council (No. 202008360018), Open
540 Research Fund Program of Key Laboratory of Metallogenic Prediction of Nonferrous Metals and Geological Environment Monitoring (Central South University), Ministry of Education (2022YSJS10)

References

- Allen, M.B., Song, S.G., Wang, C., Zeng, R.Y., and Wen, T.: An oblique subduction model for closure of the Proto-Tethys and Palaeo-Tethys oceans and creation of the Central China Orogenic Belt, *Earth-Sci. Rev.*, 104385, <https://doi.org/10.1016/j.earscirev.2023.104385>, 2023.
- 545 Bonin, B.: Do coeval mafic and felsic magmas in post-collisional to within-plate regimes necessarily imply two contrasting, mantle and crustal, sources? *Lithos*, 78, 1-24, <https://doi.org/10.1016/j.lithos.2004.04.042>, 2004.
- Burnham, A.D., and Berry, A.J.: Formation of Hadean granites by melting of igneous crust, *Nat. Geosci.*, 10, 457-461, <https://doi.org/10.1038/ngeo2942>, 2017.
- 550 Cawood, A.P., Pisarevsky, A.S., and Leitch, C.E.: Unraveling the New England orocline, east Gondwana accretionary margin, *Tectonics*, 30, 1-15, <https://doi.org/10.1029/2011TC002864>, 2011.
- Chappell, B.W., and White, A.J.R.: I- and S-type granites in the Lachlan Fold Belt, *Geol. Soc. Am. Spec. Pap.*, 272, 1-26, <https://doi.org/10.1130/SPE272-p1>, 1992.

- 555 Dan, W., Li, X.H., Guo, J.H., Liu, Y., Wang, X.C.: Paleoproterozoic evolution of the eastern Alxa Block, westernmost North China: Evidence from in situ zircon U-Pb dating and Hf-O isotopes, *Gondwana Res.*, 21, 838-864, <https://doi.org/10.1016/j.gr.2011.09.004>, 2012.
- Dan, W., Li X.H., Wang, Q., Wang, X.C., and Liu, Y.: Neoproterozoic S-type granites in the Alxa Block, westernmost North China and tectonic implications: in situ zircon U-Pb-Hf-O isotopic and geochemical constraints, *Am. J. Sci.*, 14, 110-153, <https://doi.org/10.2475/01.2014.04>, 2014.
- 560 Dan, W., Li, X.H., Wang, Q., Wang, X.C., Wyman, D.A., and Liu, Y.: Phanerozoic amalgamation of the Alxa Block and North China Craton: Evidence from Paleozoic granitoids, U-Pb geochronology and Sr-Nd-Pb-Hf-O isotope geochemistry, *Gondwana Res.*, 32, 105-121, <https://doi.org/10.1016/j.gr.2015.02.011>, 2016.
- Dong, C.Y., Liu, D.Y., Li, J.J., Wan, Y.S., Zhou, H.Y., Li, C.D., Yang, Y.H., and Xie, L.W.: New evidence for the timing of the formation of the western segment of the Khondalite Belt in the North China Craton: SHRIMP dating Hf isotopic composition of zircons from the Bayanwula Helan Shan region, *Chinese Science Bulletin*, 16, 1913-1922, [10.3321/j.issn:0023-074x.2007.16.012](https://doi.org/10.3321/j.issn:0023-074x.2007.16.012), 2007.
- Duan, F.H., Zhi, Q., Li, Y.J., Xiao, H., Wang P.L., and Gao J.P.: Petrogenesis and geodynamic setting of the Late Carboniferous granodioriteporphyry in Miaoergou pluton, southern West Junggar, *Acta Petrol. Sin.*, 37, 1159-1176, <https://doi.org/10.18654/1000-0569/2021.04.12>, 2021.
- 570 Ferry, J.M., and Watson, E.B.: New thermodynamic models and revised calibrations for the Ti-in-zircon and Zr-in-rutile thermometers, *Contrib. Mineral. Petrol.*, 154, 429-437, <https://doi.org/10.1007/s00410-007-0201-0>, 2007.
- Foley, S.F., Barth, M.G., and Jenner, G.A.: Rutile/Melt partition coefficients for trace elements and an assessment of the influence of rutile on the trace element characteristics of subduction zone magmas, *Geochim. Cosmochim. Ac.*, 64, 933-938, [https://doi.org/10.1016/S0016-7037\(99\)00355-5](https://doi.org/10.1016/S0016-7037(99)00355-5), 2000.
- 575 Freeburn, R., Bouilhol, P., Maunder, B., Magni, V., and Hunen, J.V.: Numerical models of the magmatic processes induced by slab break off, *Earth Planet. Sci. Lett.*, 478, 203-213, <https://doi.org/10.1016/j.epsl.2017.09.008>, 2017.
- Frost, C.D., and Frost, B.R.: On ferroan (A-type) granitoids: Their compositional variability and modes of origin, *J. Petrol.*, 52, 39-53, <https://doi.org/10.1093/petrology/egq070>, 2011.
- Fu, D., Kusky, T.M., Wilde, S.A., Windley, B.F., Polat, A., Huang, B., and Zhou, Z.P.: Boninitic blueschists record subduction initiation and subsequent accretion of an arc-forearc in the northeast Proto-Tethys Ocean, *Geology*, 50, 10-15, <https://doi.org/10.1130/G49457.1>, 2022.
- 580 Fu, D., Kusky, T.M., Wilde, S.A., Windley, B.F., Polat, A., Huang, B., and Zhou, Z.P.: Structural anatomy of the early Paleozoic Laohushan ophiolite and subduction complex: Implications for accretionary tectonics of the Proto-Tethyan North Qilian orogenic belt, northeastern Tibet, *Geol. Soc. Am. Bull.*, 132, 2175-2201, <https://doi.org/10.1130/B35442.1>, 2020.
- 585

- Gong, J.H., Zhang, J.X., Wang, Z.Q., Yu, S.Y., Li, H.K., and Li, Y.S.: Origin of the Alxa Block, western China: New evidence from zircon U-Pb geochronology and Hf isotopes of the Longshoushan Complex, *Gondwana Res.*, 36, 359-375, <https://doi.org/10.1016/j.gr.2015.06.014>, 2016.
- Griffin, W.L., Wang, X., Jackson, S.E., Pearson, N.J., O'Reilly, S.Y., Xu, X.X., and Zhou, X.M.: Zircon chemistry and magma mixing, SE China: In-situ analysis of Hf isotopes, Tonglu and Pingtan igneous complexes, *Lithos*, 61, 237-269, [https://doi.org/10.1016/S0024-4937\(02\)00082-8](https://doi.org/10.1016/S0024-4937(02)00082-8), 2002.
- Harris, N.B.W., Inger, S., and Xu, R.H.: Cretaceous plutonism in Central Tibet: an example of post-collision magmatism? *J. Volcanol. Geotherm. Res.*, 44, 21-32, [https://doi.org/10.1016/0377-0273\(90\)90009-5](https://doi.org/10.1016/0377-0273(90)90009-5), 1990.
- He, H.Y., Li, Y.L., Wang, C.S., Han, Z.P., Ma, P.F., and Xiao, S.Q.: Petrogenesis and tectonic implications of Late Cretaceous highly fractionated I-type granites from the Qiangtang block, central Tibet, *J. Asian Earth Sci.*, 176, 337-352, <https://doi.org/10.1016/j.jseaes.2019.02.022>, 2019.
- Hoskin, P.W.O., and Schaltegger, U.: The composition of zircon and igneous and metamorphic petrogenesis, *Rev. Mineral. Geochem.*, 53, 27-62, <https://doi.org/10.2113/0530027>, 2003.
- Hu, F.Y., Ducea, M.N., Liu, S.W., and Chapman, J.B.: Quantifying Crustal Thickness in Continental Collisional Belts: Global Perspective and a Geologic Application, *Sci. Rep.*, 7, 1-10, <https://doi.org/10.1038/s41598-017-07849-7>, 2017.
- Hu, Z.C., Liu, Y.S., Gao, S., Liu, W.G., Zhang, W., and Tong, X.R.: Improved in situ Hf isotope ratio analysis of zircon using newly designed X skimmer cone and jet sample cone in combination with the addition of nitrogen by laser ablation multiple collector ICP-MS, *J. Anal. At. Spectrom.*, 27, 1391-1399, <https://doi.org/10.1039/C2JA30078H>, 2012.
- Hui, J., Zhang, K.J., Zhang, J., Qu, J.F., Zhang, B.H., Zhao, H., and Niu, P.F.: Middle-late Permian high-K adakitic granitoids in the NE Alxa block, northern China: Orogenic record following the closure of a Paleo-Asian oceanic branch? *Lithos*, 400-401, 106379, <https://doi.org/10.1016/j.lithos.2021.106379>, 2021.
- Jahn, B.M., Wu, F.Y., Lo, C.H., and Tsai, C.H.: Crust-mantle interaction induced by deep subduction of the continental crust: geochemical and Sr-Nd isotopic evidence from post-collisional mafic-ultramafic intrusions of the Northern Dabie complex, central China, *Chem. Geol.*, 157, 119-146, [https://doi.org/10.1016/S0009-2541\(98\)00197-1](https://doi.org/10.1016/S0009-2541(98)00197-1), 1999.
- Karsli, O., Dokuz, A., Uysal, İ., Aydin, F., Chen, B., Kandemir, R., and Kandemir, R.: Relative contributions of crust and mantle to generation of Campanian high-K calc-alkaline I-type granitoids in a subduction setting, with special reference to the Harşit Pluton, Eastern Turkey, *Contrib. Mineral. Petrol.*, 160, 467-487, [10.1007/s00410-010-0489-z](https://doi.org/10.1007/s00410-010-0489-z), 2010.
- Kong, H., Li, H., Wu, Q.H., Xi, X.S., Dick, J.M., and Gabo-Ratio, J.A.S.: Co-development of Jurassic I-type and A-type granites in southern Hunan, South China: Dual control by plate subduction and intraplate mantle upwelling, *Chemie der Erde-Geochemistry*, 78, 500-520, <https://doi.org/10.1016/j.chemer.2018.08.002>, 2018.
- Li, B., Jiang, S.Y., Lu, A.H., Zhao, H.X., Yang, T.L., and Hou, M.L.: Zircon U-Pb dating, geochemical and Sr-Nd-Hf isotopic characteristics of the Jintonghu monzonitic rocks in western Fujian Province, South China: Implication for Cretaceous

- crust-mantle interactions and lithospheric extension, *Lithos*, 260, 413-428, <https://doi.org/10.1016/j.lithos.2016.05.002>, 2016.
- 620 Li, C.D., Zhang, Q., Miao, L.C., and Meng, X.F.: Mesozoic high-Sr, low-Y and low-Sr, low-Y types granitoids in the northern Hebeiprovince: Geochemistry and petrogenesis and its relation to mineralization of gold deposits, *Acta Petrol. Sin.*, 20, 269-284, <https://doi.org/10.1016/j.sedgeo.2004.01.002>, 2004.
- Li, H., Sun, H.S., Wu, J.H., Evans, N.J., Xi, X.S., Peng, N.L., Cao, J.Y., and Gabo-Ratio, J.A.S.: Re-Os and U-Pb geochronology of the Shazigou Mo polymetallic ore field, Inner Mongolia: Implications for Permian-Triassic mineralization at the northern margin of the North China Craton, *Ore Geology Reviews*, 83, 287-299, <https://doi.org/10.1016/j.oregeorev.2016.12.010>, 2017.
- 625 Li, J.Y., Zhang, J., and Qu, J.F.: Amalgamation of North China Craton with Alxa Block in the Late of Early Paleozoic: Evidengce from Sedimentary Sequences in the Niushou Mountain, Ningxia Hui Autonomous Region, NW China, *Geological Review.*, 58, 208-214, [10.16509/j.georeview.2012.02.014](https://doi.org/10.16509/j.georeview.2012.02.014), 2012.
- 630 Li, S., Wang, T., Xiao, W.J., and Ho, Q.L.: Tectono-magmatic evolution from accretion to collision in the southern margin of the Central Asian Orogenic Belt, *Acta Petrol. Sin.*, 39, 1261-1275, <https://doi.org/10.18654/1000-0569/2023.05.03>, 2023.
- Liégeois, J.P., Navez, J., Hertogen, J., and Black, R.: Contrasting origin of post-collisional high-K calc-alkaline and shoshonitic versus alkaline and peralkaline granitoids, The use of sliding normalization, *Lithos*, 45, 1-28, [https://doi.org/10.1016/S0024-4937\(98\)00023-1](https://doi.org/10.1016/S0024-4937(98)00023-1), 1998.
- 635 Liu, Q., Zhao, G.C., Han, Y.G., Eizenhöfer, P.R., Zhu, Y.L., Hou, W.Z., and Zhang, X.R.: Timing of the final closure of the Paleo-Asian Ocean in the Alxa Terrane: Constraints from geochronology and geochemistry of Late Carboniferous to Permian gabbros and diorites, *Lithos*, 274-275, 19-30, <https://doi.org/10.1016/j.lithos.2016.12.029>, 2017.
- Liu, Q., Zhao, GC., Sun, M., Han, Y.G., Eizenhofer, P.R., Hou, W.Z., Zhang, X.R., Zhu, YL., Wang, B., and Liu, D.X.: Early Paleozoic subduction processes of the Paleo-Asian Ocean: Insights from geochronology and geochemistry of Paleozoic plutons in the Alxa Terrane, *Lithos*, 262, 546-560, <https://doi.org/10.1016/j.lithos.2016.07.041>, 2016.
- 640 Liu, Q.Q., Li, H., Shao, Y.J., Girei, M.B., Jiang, W.C., Yuan, H.M., and Zhang, X.: Age, genesis, and tectonic setting of the Qiushuwan Cu-Mo deposit in East Qinling (Central China): constraints from Sr-Nd-Hf isotopes, zircon U-Pb and molybdenite Re-Os dating, *Ore Geology Reviews*, 132, 103998, <https://doi.org/10.1016/j.oregeorev.2021.103998>, 2021.
- Liu, W.Heng., Spier, C., Niu, Y.B., Liu, X.D., Yan, J., Wang, K.X., and Pan, J.Y.: Magma mixing in the genesis of the Qingshanbao granitoids in the Longshoushan area: Implications for the tectonic evolution of the North Qilian orogenic belt, NW China, *Geol. J.*, 56, 4594-4617, <https://doi.org/10.1002/gj.4195>, 2021.
- 645 Li, Y.L., Zhou, H.W., Brouwer, F.M., Xiao, W.J., Wijbrans, J.R., and Zhong, Z.Q.: Early Paleozoic to Middle Triassic bivergent accretion in the Central Asian Orogenic Belt: insights from zircon U-Pb dating of ductile shear zones in central Inner Mongolia, China, *Lithos*, 205, 84-111, <https://doi.org/10.1016/j.lithos.2014.06.017>, 2014.

- 650 Ma, L., Jiang, S.Y., Dai, B.Z., Jiang, Y.H., Hou, M.L., Pu, W., and Xu, B.: Multiple sources for the origin of Late Jurassic Linglong adakitic granite in the Shandong Peninsula, eastern China: Zircon U-Pb geochronological, geochemical and Sr-Nd-Hf isotopic evidence, *Lithos*, 162-163, 251-263, <https://doi.org/10.1016/j.lithos.2013.01.009>, 2013.
- Maniar, P.D., and Piccoli, P.M.: Tectonic discrimination of granitoids, *Geol. Soc. Am. Bull.*, 101, 635-643, [https://doi.org/10.1130/0016-7606\(1989\)101<0635:TDOG>2.3.CO;2](https://doi.org/10.1130/0016-7606(1989)101<0635:TDOG>2.3.CO;2), 1989.
- 655 Mantle, G.W., and Collins, W.J.: Quantifying crustal thickness variations in evolving orogens: Correlation between arc basalt composition and Moho depth, *Geology*, 36, 87-90, <https://doi.org/10.1130/G24095A.1>, 2008.
- Middlemost, E.A.K.: Naming materials in the magma/igneous rock system, *Earth-Sci. Rev.*, 37, 215-224, [https://doi.org/10.1016/0012-8252\(94\)90029-9](https://doi.org/10.1016/0012-8252(94)90029-9), 1994.
- Pearce, J.A., Harris, N.B.W., and Tindle, A.G.: Trace Element Discrimination Diagrams for the Tectonic Interpretation of
660 Granitic Rocks, *J. Petrol.*, 25, 956-983, <https://doi.org/10.1093/petrology/25.4.956>, 1984.
- Qi, J.W., Zhang, S.M., Yang, C.S., Lan, D.C., and Wang, L.L.: The LA-ICP-MS zircon U-Pb age of the pegmatoidal alaskite and its relationship with uranium mineralization in Hongshiquan area, Gansu Province, *Geol. Bull.*, 38, 562-572, <https://doi.org/CNKI:SUN:ZQYD.0.2019-04-009>, 2019.
- Schiano, P., Monzier, M., Eissen, J.P., Martin, H., and Koga, K.T.: Simple mixing as the major control of the evolution of
665 volcanic suites in the Ecuadorian Andes, *Contrib. Mineral. Petrol.*, 160, 297-312, <https://doi.org/10.1007/s00410-009-0478-2>, 2010.
- Schiller, D., and Finger, F.: Application of Ti-in-zircon thermometry to granite studies: problems and possible solutions, *Contrib. Mineral. Petrol.*, 174, 51, <https://doi.org/10.1007/s00410-019-1585-3>, 2019.
- Shi, X.J., Wang, T., Zhang, L., Castro, A., Xiao, X.C., Tong, Y., Zhang, J.X., Guo, L., and Yang, Q.D.: Timing, petrogenesis
670 and tectonic setting of the Late Paleozoic gabbro-granodiorite-granite intrusions in the Shalazhashan of northern Alxa: Constraints on the southernmost boundary of the Central Asian Orogenic Belt, *Lithos*, 208-209, 158-177, <https://doi.org/10.1016/j.lithos.2014.08.024>, 2014.
- Skjerlie, K.P., and Johnston, A.D.: Vapor-absent melting at 10 kbar of a biotite-and amphibole-bearing tonalitic gneiss: implications for the generation of A-type granites, *Geology*, 20, 263-266, [https://doi.org/10.1130/0091-7613\(1993\)021<0089:VAMAKO>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0089:VAMAKO>2.3.CO;2), 1992.
- 675 Song, S.G., Niu, Y.L., Su, L., and Xia, X.H.: Tectonics of the North Qilian Orogen, NW China, *Gondwana Res.*, 23, 1378-1401, <https://doi.org/10.1016/j.gr.2012.02.004>, 2013.
- Song, S.G., Wang, M.J., Wang, C., and Niu, Y.L.: Magmatism during continental collision, subduction, exhumation and mountain collapse in collisional orogenic belts and continental net growth: A perspective, *Science China Earth Sciences*,
680 58(8), 1284-1304, <https://doi.org/10.1007/s11430-015-5102-x>, 2015.

- Sun, H.S., Li, H., Danišik, M., Xia, Q.L., Jiang, C.L., Wu, P., Yang, H., Fan, Q.R., and Zhu, D.S.: U-Pb and Re-Os geochronology and geochemistry of the Donggebi Mo deposit, Eastern Tianshan, NW China: Insights into mineralization and tectonic setting, *Ore Geology Reviews*, 86, 584-599, <https://doi.org/10.1016/j.oregeorev.2017.03.020>, 2017.
- 685 Sun, S.S., and McDonough, W.F.: Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes, *Geol. Soc. Lond. Spec. Publ.*, 42, 313-345, <https://doi.org/10.1144/GSL.SP.1989.042.01.19>, 1989.
- Sylvester, P.J.: Post-collisional strongly peraluminous granites, *Lithos*, 45, 29-44, [https://doi.org/10.1016/S0024-4937\(98\)00024-3](https://doi.org/10.1016/S0024-4937(98)00024-3), 1998.
- Tang, Y.L., Shi, Y., Hu, X.M., Liu, X.J., and Huang, C.W.: Petrogenesis of Early Paleozoic I-type granitoids in the Wuyi-Yunkai Orogen, South China: Implications for the tectono-magmatic evolution of the Cathaysia Block, *J. Asian Earth Sci.*, 220, 104906, <https://doi.org/10.1016/j.jseaes.2021.104906>, 2021.
- 690 Trail, D., Watson, E.B., and Tailby, N.D.: Ce and Eu anomalies in zircon as proxies for the oxidation state of magmas, *Geochim. Cosmochim. Ac.*, 97, 70-87, <https://doi.org/10.1016/j.gca.2012.08.032>, 2012.
- Wang, C.Y., Zhang Q., Qian Q., and Zhou, M.F.: Geochemistry of the Early Paleozoic Baiyin Volcanic Rocks (NW China): Implications for the Tectonic Evolution of the North Qilian Orogenic Belt, *J. Geol.*, 113, 83-94, <https://doi.org/10.1086/425970>, 2005.
- 695 Wang, T., Tong, Y., Zhang, L., Li, S., Huang, H., Zhang, J.J., Guo, L., Yang, Q.D., Hong, D.W., Donskaya, T., Gladkochub, D., and Tserendash, N.: Phanerozoic granitoids in the middle and eastern parts of CentralAsia and their tectonic significance, *J. Asian Earth Sci.*, 145, 368-392, <https://doi.org/10.1016/j.jseaes.2017.06.029>, 2017.
- Wang, Z.Z., Chen, X.H., Li, B., Zhang, Y.P., and Xu, S.L.: The discovery of the Paleoproterozoic syenite in Helishan, Gansu Province, and its implications for the tectonic attribution of the Alxa Block, *Geology in China*, 46, 1094-1104. [10.12029/gc20190510](https://doi.org/10.12029/gc20190510), 2019.
- 700 Wang, Z.Z., Chen, X.H., Shao, Z.G., Li, B., Ding, W.C., Zhang, Y.P., Wang, Y.C., Zhang Y.Y., Xu, S.L., and Qin, X.: Petrogenesis of the Late Silurian-Early Devonian granites in the Longshoushan-Helishan area, Gansu Province, and its tectonic implications for the Early Paleozoic evolution of the southwestern Alxa Block, *Acta Geol. Sin.*, 94, 2243-2261, <https://doi.org/10.19762/j.cnki.dizhixuebao.2020027>, 2020.
- 705 Wang, Z.Z., Han, B.F., Feng, L.X., and Liu, B.: Geochronology, geochemistry and origins of the Paleozoic-Triassic plutons in the Langshan area, western Inner Mongolia, China, *J. Asian Earth Sci.*, 97, 337-351, <https://doi.org/10.1016/j.jseaes.2014.08.005>, 2015.
- Watson, E.B., and Harrison, T.M.: Zircon saturation revisited: temperature and composition effects in variety of crustal magma types, *Earth Planet. Sci. Lett.*, 64, 295-304, [https://doi.org/10.1016/0012-821X\(83\)90211-X](https://doi.org/10.1016/0012-821X(83)90211-X), 1983.
- 710 Watson, E.B., and Harrison, T.M.: Zircon saturation revisited: temperature and composition effects in a variety of crustal magma types, *Earth Planet. Sci. Lett.*, 64, 295-304, [https://doi.org/10.1016/0012-821X\(83\)90211-X](https://doi.org/10.1016/0012-821X(83)90211-X), 1983.
- Watson, E.B., and Harrison, T.M.: Zircon Thermometer Reveals Minimum Melting Conditions on Earliest Earth, *Science*, 308, 841-844, <https://doi.org/10.1126/science.1120977>, 2005.

- 715 Wei, Q.Q., Hao, L.B., Lu, J.L., Zhong, Y.Y., Zhao, X.Y., and Shi, H.L.: LA-MC-ICP-MS Zircon U-Pb Dating of Hexipu Granite and Its Geological Implications, *Bull. Mineral., Petrol. Geochem.*, 32, 729-735, <https://doi.org/10.3969/j.issn.1007-2802.2013.06.007>, 2013.
- Whalen, J.B., Currie, Kenneth L., and Chappell, B.W.: A-type granites: geochemical characteristics, discrimination and petrogenesis, *Contrib. Mineral. Petrol.*, 95, 407-419, <https://doi.org/10.1007/BF00402202>, 1987.
- 720 Wolf, M.B., and London, D.: Apatite dissolution into peraluminous haplogranitic melts: An experimental study of solubilities and mechanism, *Geochim. Cosmochim. Ac.*, 58, 4127-4145, [https://doi.org/10.1016/0016-7037\(94\)90269-0](https://doi.org/10.1016/0016-7037(94)90269-0), 1994.
- Wu, D.D., Li, S., Chew, D., Liu, T.X., and Guo, D.H.: Permian-Triassic magmatic evolution of granitoids from the southeastern Central Asian Orogenic Belt: Implications for accretion leading to collision, *Science China Earth Sciences*, 64(5): 788-806, <https://doi.org/10.1007/S11430-020-9714-5>, 2021.
- 725 Wu, F.Y., Liu, X.C., Ji, W.Q., Wang, J.M., and Yang, L.: Highly fractionated granites: recognition and research, *Science China Earth Sciences*, 60, 1201-1219, [10.1007/s11430-016-5139-1](https://doi.org/10.1007/s11430-016-5139-1), 2017.
- Xia, L.Q., Li, X.M., Yu, J.Y., and Wang, G.Q.: Mid-late neoproterozoic to early paleozoic volcanism and tectonic evolution of the Qilianshan, NW China, *Georesj*, 9-12, 1-41, <https://doi.org/10.1016/j.grj.2016.06.001>, 2016.
- Xia, X., Song, S., and Niu, Y.: Tholeiite-Boninite terrane in the North Qilian suture zone: Implications for subduction initiation and back-arc basin development, *Chem. Geol.*, 328, 259-277, <https://doi.org/10.1016/j.chemgeo.2011.12.001>, 2012.
- 730 Xiao, B., Chen, H.Y., Hollings, P., Han, J.S., Wang, Y.F., Yang, J.T., and Cai, K.D.: Magmatic evolution of the Tuwu-Yandong porphyry Cu belt, NW China: Constraints from geochronology, geochemistry and Sr-Nd-Hf isotopes, *Gondwana Res.*, 43, 74-91, <https://doi.org/10.1016/j.gr.2015.09.003>, 2017.
- Xiao, W.J., Windley, B.F., Sun, S., Li, J.L., Huang, B.C., Han, C.M., Yuan, C., Sun, M., and Chen, H.L.: A tale of amalgamation of three Permo-Triassic collage systems in Central Asia: oroclines, sutures, and terminal accretion, *Annu. Rev. Earth Planet. Sci.*, 43, 477-507, <https://doi.org/10.1146/annurev-earth-060614-105254>, 2015.
- 735 Xue, S., Ling, M.X., Liu, Y.L., Zhang, H., and Sun, W.D.: The genesis of early Carboniferous adakitic rocks at the southern margin of the Alxa Block, North China, *Lithos*, 278-281, 181-194, <https://doi.org/10.1016/j.lithos.2017.01.012>, 2017.
- Yang, H., Zhang, H.F., Xiao, W.J., Luo, B.J., Gao, Z., Tao, L., Zhang, L.Q., and Guo, L.: Petrogenesis of Early Paleozoic high Sr/Y intrusive rocks from the North Qilian orogen: Implication for diachronous continental collision, *Lithosphere*, 12, 53-73, <https://doi.org/10.1130/L1129.1>, 2019.
- 740 Yu, S.Y., Zhang, J.X., Qin, H.P., Sun, D.Y., Zhao, X.L., Cong, F., and Li, Y.S.: Petrogenesis of the early Paleozoic low-Mg and high-Mg adakitic rocks in the North Qilian orogenic belt, NW China: Implications for transition from crustal thickening to extension thinning, *J. Asian Earth Sci.*, 107, 122-139, <https://doi.org/10.1016/j.jseaes.2015.04.018>, 2015.
- 745 Yuan, W., and Yang, Z.Y.: The Alashan Terrane did not amalgamate with North China block by the Late Permian: Evidence from Carboniferous and Permian paleomagnetic results, *J. Asian Earth Sci.*, 104, 145-159, [10.1016/j.jseaes.2014.02.010](https://doi.org/10.1016/j.jseaes.2014.02.010), 2015.

- Zeng, R.Y., Allen, M.B., Mao, X.C., Lai, J.Q., Yan, J., and Wan, J.J.: Whole-rock and zircon evidence for evolution of the Late Jurassic high Sr/Y Zhoujiapuzi granite, Liaodong Peninsula, North China Craton, *Solid Earth*, 13, 1259-1280, <https://doi.org/10.5194/se-2021-129>, 2022.
- 750
- Zeng, R.Y., Lai, J.Q., Mao, X.C., Li, B., Ju, P.J., and Tao, S.L.: Geochemistry, zircon U-Pb dating and Hf isotopes composition of Paleozoic granitoids in Jinchuan, NW China: Constraints on their petrogenesis, source characteristics and tectonic implication, *J. Asian Earth Sci.*, 121, 20-33, <https://doi.org/10.1016/j.jseaes.2016.02.009>, 2016.
- Zeng, R.Y., Lai, J.Q., Mao, X.C., Li, B., Zhang, J.D., Bayless, R., and Yang, L.Z.: Paleoproterozoic multiple tectonothermal events in the Longshoushan area, Western North China Craton and their geological implication: evidence from geochemistry, zircon U-Pb geochronology and Hf isotopes, *Minerals*, 8, 361, <https://doi.org/10.3390/MIN8090361>, 2018.
- 755
- Zeng, R.Y., Lai, J.Q., Mao, X.C., Xiao, W.Z., Yan, J., Zhang, C.G., and Xiao, W.Z.: Geochemistry and zircon ages of the Yushigou diabase in the Longshoushan area, Alxa Block: implications for crust-mantle interaction and tectonic evolution, *Geol. Mag.*, 158, 685-700, <https://doi.org/10.1017/S0016756820000783>, 2021.
- 760
- Zhang, B., Guo, F., Zhang, X.B., Wu, Y.M., Wang, G.Q., and Zhao, L.: Early Cretaceous subduction of Paleo-Pacific Ocean in the coastal region of SE China: Petrological and geochemical constraints from the mafic intrusions, *Lithos*, 334-335, 8-24, <https://doi.org/10.1016/j.lithos.2019.03.010>, 2019.
- Zhang, D.H., Wei, J.H., Fu, L.B., Chen, H.Y., Tan, J., Li, Y.J., Shi, W.J., and Tian, N.: Formation of the Jurassic Changboshan-Xieni-qishan highly fractionated I-type granites, northeastern China: implication for the partial melting of juvenile crust induced by asthenospheric mantle upwelling, *Geol. J.*, 50, 122-138, <https://doi.org/10.1002/gj.2531>, 2015.
- 765
- Zhang, J.J., Wang, T., Castro, A., Zhang, L., Shi, X.J., Tong, Y., Zhang, Z.C., Guo, L., Yang, Q.D., and Iaccheri, L.M.: Multiple Mixing and Hybridization from Magma Source to Final Emplacement in the Permian Yamatu Pluton, the Northern Alxa Block, China, *J. Petrol.*, 157, 183-197, <https://doi.org/10.1007/BF01765318>, 2016.
- Zhang, J.J., Wang, T., Zhang, L., Tong, Y., Zhang, Z.C., Shi, X.J., Guo, L., Huang, H., Yang, Q.D., Huang, W., Zhao, J.X., Ye, K., and Hou, J.Y.: Tracking deep crust by zircon xenocrysts within igneous rocks from the northern Alxa, China: Constraints on the southern boundary of the Central Asian Orogenic Belt, *J. Asian Earth Sci.*, 108, 150-169, <https://doi.org/10.1016/j.jseaes.2015.04.019>, 2015.
- 770
- Zhang, J.X., and Gong, J.H.: Revisiting the nature and affinity of the Alxa Block, *Acta Petrol. Sin.*, 34, 940-962, <https://doi.org/CNKI:SUN:YSXB.0.2018-04-008>, 2018.
- 775
- Zhang, J.X., Gong, J.H., Yu, S.Y., Li, H.K., and Hou, K.J.: Neoproterozoic multiple tectonothermal events in the western Alxa block, North China Craton and their geological implication: Evidence from zircon U-Pb ages and Hf isotopic composition, *Precambrian Res.*, 235, 36-57, <https://doi.org/10.1016/j.precamres.2013.05.002>, 2013.
- Zhang, L., Zhang, H., Hawkesworth, C., Luo, B., and Yang, H.: Mafic rocks from the southern Alxa block of Northwest China and its geodynamic evolution in the Paleozoic, *J. Geol. Soc.*, 178, 2020-038, <https://doi.org/10.1144/jgs2020-038>, 2021.

- 780 Zhang, L., Wang, T., Zhang, J.J., Shi, X.J., Ren, H.D., Yang, Q.D., Tong, Y., Guo, L.: Revisiting the boundary between the Central Asian Orogenic Belt and North China Craton in Alxa area, China: Insights from zircon U-Pb ages and Hf isotopes of Phanerozoic granitoids, *Gondwana Res*, 119: 119-137, <https://doi.org/10.1016/j.gr.2023.03.011>, 2023.
- Zhang, L.Q., Zhang, H.F., Zhang, S.S., Xiong, Z.L., Luo, B.J., Yang, H., Pan, F.B., Zhou, X.C., Xu, W.C., and Guo, L.: Lithospheric delamination in post-collisional setting: Evidence from intrusive magmatism from the North Qilian orogen to southern margin of the Alxa block, NW China, *Lithos*, 288-289, 20-34, <https://doi.org/10.1016/j.lithos.2017.07.009>, 2017.
- 785 Zhang, Q., Jin, W.J., Li, C.D., and Wang, Y.L.: Revisiting the new classification of granitic rocks based on whole-rock Sr and Yb contents: Index, *Acta Petrol. Sin.*, 26, 985-1015, <https://doi.org/CNKI:SUN:YSXB.0.2010-04-002>, 2010.
- Zhang, Q., Wang, Y., Li, C.D., Wang, Y.L., Jin, W.J., and Jia, X.Q.: Granite classification on the basis of Sr and Yb contents and its implications, *Acta Petrol. Sin.*, 22, 2249-2269, <https://doi.org/10.3321/j.issn:1000-0569.2006.09.001>, 2006.
- 790 Zhang, W., and Hu Z.C.: Estimation of Isotopic Reference Values for Pure Materials and Geological Reference Materials, *At. Spectrosc.*, 41, 93-102, <https://doi.org/10.46770/AS.2020.03.001>, 2020.
- Zhou, X.C., Zhang, H.F., Luo, B.J., Pan, F.B., Zhang, S.S., and Guo, L.: Origin of high Sr/Y-type granitic magmatism in the southwestern of the Alxa Block, Northwest China, *Lithos*, 256-257, 211-227, <https://doi.org/10.1016/j.lithos.2016.04.021>, 2016.
- 795 Zong, K.Q., Klemd, R., Yuan, Y., He, Z.Y., Guo, J.L., Shi, X.L., Liu, Y.S., Hu, Z.C., and Zhang, Z.M.: The assembly of Rodinia: The correlation of early Neoproterozoic (ca. 900 Ma) high-grade metamorphism and continental arc formation in the southern Beishan Orogen, southern Central Asian Orogenic Belt (CAOB), *Precambrian Res.*, 290, 32-48, <https://doi.org/10.1016/j.precamres.2016.12.010>, 2017.
- 800 Zorpi, M.J., Coulon, C., Orsini, J.B., and Cocirta, C.: Magma mingling, zoning and emplacement in calc-alkaline granitoid plutons, *Tectonophysics*, 157, 315-329, [https://doi.org/10.1016/0040-1951\(89\)90147-9](https://doi.org/10.1016/0040-1951(89)90147-9), 1989.