



Exploring drought hazard, vulnerability, and related impacts to agriculture in Brandenburg

Fabio Brill¹, Pedro Henrique Lima Alencar², Huihui Zhang¹, Friedrich Boeing^{3,4}, Silke Hüttel^{5,6}, Tobia Lakes^{1,7}

5 ¹ Geography Department, Humboldt-Universität zu Berlin, Berlin, 10099, Germany

² Department of Ecohydrology and Landscape Assessment, Technical University Berlin, Berlin, 10623, Germany

³ Department Computational Hydrosystems, Helmholtz Centre for Environmental Research (UFZ), Leipzig, 04318, Germany

⁴ Institute for Environmental Science and Geography, University of Potsdam, Potsdam-Golm, 14476, Germany

⁵ Department of Agricultural Economics and Rural Development, University of Göttingen, Göttingen, 37073, Germany

10 ⁶ Faculty of Agriculture, University of Bonn, Bonn, 53115, Germany

⁷ Integrative Research Institute on Transformations of Human-Environment Systems (IRI THESys), Humboldt-Universität zu Berlin, Berlin, 10099, Germany

Correspondence to: Fabio Brill (fabio.brill@hu-berlin.de)

Abstract.

15 Adaptation to an increasingly dry regional climate requires spatially explicit information about current and future risks. Existing drought risk studies often rely on expert-weighted composite indicators, while empirical evidence on impact-relevant factors is still scarce. The aim of this study is to investigate to what extent hazard and vulnerability indicators can explain observed agricultural drought impacts via data-driven methods. We focus on the German federal state of Brandenburg, 2013-2022, including several consecutive drought years. As impact indicators we use thermal-spectral anomalies (LST/NDVI) on
20 field level, and empirical yield gaps from reported statistics on county level. Empirical associations to the impact indicators on both spatial levels are compared. Non-linear models explain up to about 60% variance in the yield gap data, with lumped models for all crops being more stable than models for individual crops, and models for the drought years performing better than for the pre-drought years. Meteorological drought in June and soil quality are selected as strongest impact-relevant factors. Rye is found less vulnerable than wheat, despite growing on poorer soils. LST/NDVI only weakly relates to our empirical
25 yield gaps. We recommend comparing different impact indicators on multiple scales to proceed with the development of empirically grounded risk maps.



1 Introduction

Agricultural drought risk mapping is essential for spatial prioritization of adaptation actions and measures, and particularly to raise awareness of stakeholders throughout the social-ecological system (Mishra and Singh, 2011; Blauhut, 2020; Kim et al., 2021). In the light of climate change, droughts are expected to occur in higher frequency and unprecedented magnitudes, which poses a major challenge for risk management (Hanel et al., 2018; Hari et al., 2020; Satoh et al., 2022; Kreibich et al., 2022). Risk in this context can be conceptualized as potential for negative impacts, assembled from the components hazard, exposure, and vulnerability – while definitions of terms have shifted over the years, the recent guideline by the Intergovernmental Panel on Climate Change (IPCC) is very clear on that matter (Reisinger et al., 2020). A sound understanding of hazard thresholds and vulnerability conditions associated with impacts under droughts (hereinafter “impact-relevant factors”) is thus urgently needed to provide reliable risk maps and move towards impact-based forecasting (Sutanto et al., 2019). However, many drought risk maps are still being produced by more or less arbitrary weighting of indicators to a composite score (Kim et al., 2015; Dabanli, 2018; Kim et al., 2021; Khoshnazar, 2023), sometimes based on expert opinion (Frischen et al., 2020; Abdullah et al., 2021; Stephan et al., 2023), or by process-based models for individual agricultural crops (Söder et al., 2022). A review of international examples found that drought studies in particular do often neither define their target of investigation in sufficient detail, nor include any sort of validation, thereby making the results difficult to interpret and use (Hagenlocher et al., 2019). Such aggregated indicators could harm more than they help by masking important differences between areas (Jhan et al., 2020). For Brandenburg, our study region, Ihinegbu and Ogunwumi (2022) produced a drought event map based on weighting of the normalized difference vegetation index (NDVI), land surface temperature (LST), and rainfall, without considering vulnerability or impacts. We suggest that drought risk mapping should be more closely related to investigations of actual hazard-impact relationships.

Droughts are natural hazards with a relatively slow-onset character, although there is recently more scientific attention towards flash droughts (Alencar and Paton, 2022). Distinguished are purely meteorological droughts, soil moisture droughts, hydrological low flow in rivers, as well as socio-economic droughts that impose consequences on the broader population and might lead to water conflicts (Wilhite and Glantz, 1985). For agriculture, the direct biophysical drought impacts arguably start once water availability restricts plant growth. Depending on the drought intensity, duration, and timing within the plant phenological stage, crop health is affected, which translates into yield levels, product quality and ultimately prices (Santini et al., 2022). Historically, droughts are associated with famine and high death tolls (Mishra et al., 2019; Contreras, 2019). With modern disaster response, the impacts usually stay on the economic level, but also monetary loss can have severe consequences for individuals, businesses, and entire regions, that are to be anticipated and managed proactively (Erfurt et al., 2019; Krishnamurthy et al., 2022). While there are mechanisms to partially compensate losses due to extreme events (European Commission, 2023), a notable residual business risk remains with the farms – potentially leading to stress and anxiety experienced by farmers (Austin et al., 2018; Abunyewah et al., 2024). Indirect effects are then propagated along the value



65 chain and within the affected region. More than 100 billion euro have been attributed to drought events between 1986 and
2016 in the European Union (Blauhut et al., 2016). In the German federal state of Brandenburg, our study region, the local
government spent 72 million euro of compensations to farmers for drought-related losses in 2018 alone, accounting for about
45% of the actual claims of that year (MLUK, 2019). This, however, was only the beginning of a prolonged multi-year drought
(Boeing et al., 2022). As an area that was historically water-rich, Brandenburg now needs to prepare for a dryer future
(Kahlenborn et al., 2021; MLUK, 2023), making it an interesting case for an empirical study.

70

Methods for empirically investigating impact-relevant factors for natural hazards range from simple regression to state-of-the-
art data-mining algorithms. Investigated impacts include for example damage to buildings from river floods (Merz et al., 2013),
debris flows (Jakob et al., 2012), or compound events (Brill et al., 2020), as well as casualties from floods (Tellman et al.,
2020) and heat (Şalap-Ayça and Goto, 2023), or the occurrence of wildfires (Kondylatos et al., 2022). There have been similar
75 attempts to uncover impact-relevant factors from text reports of past droughts (Stahl et al., 2016; Blauhut et al., 2016; de Brito
et al., 2020; Sodge et al., 2023; Stephan et al. 2023b), and from yield anomalies for selected crops (Sutanto et al., 2019; Peichl
et al., 2021; Tanguy et al., 2023). Despite these recent efforts, empirical evidence on regional impact-relevant factors and non-
linearities of actual observed drought impacts is still rather scarce (Bachmair et al., 2016; Sutanto et al., 2019; Peichl et al.,
2021; Tanguy et al., 2023). A key bottleneck of such data-driven studies is the availability of impact data.

80

One potential solution to solve the data availability issue is the use of remote sensing data products, from which indicators of
crop health can be derived. The ratio between LST and NDVI is a well-established observable indicator for that purpose
(McVicar and Bierwirth, 2001; Karnieli et al., 2010; Crocetti et al., 2020). Mid growing season is generally regarded as the
most decisive time of observation (Ghazaryan et al., 2020). Reinermann et al. (2019) used remote-sensing time series from
85 2000 to 2018 and detected negative vegetation anomalies in Germany during summer months, particularly in the drought year
2018. The correlation strength of drought indicators to yields was found to increase over time (Lüttger and Feike, 2018).
However, most data-driven studies using earth observation merely model the occurrence of drought or treat anomalies of
spectral indicators as “observed impact” without proper comparison to yields (Houmma et al., 2022).

90

Based on these identified gaps, the aim of this study is twofold: (1) we investigate the recent drought years in Brandenburg by
combining indicators on hazard, vulnerability, and impacts from multiple data sources, and (2) we derive empirical
relationships of hazard and vulnerability indicators to the different impact indicators by data-driven methods. Additionally, an
interactive web map was developed to assist on the exploration of the components of regional drought risk. The findings
provide new insights on the complexity of the impact-hazard-vulnerability relationship of agricultural droughts for our study
95 region in Brandenburg, as well as on limitations of currently available datasets. This has implications for modelling and
monitoring.



2 Material & Methods

2.1 Approach & Study Area

To achieve our two objectives, we select a set of indicators based on a literature analysis, including impact indicators on two different levels: field and county level. Spatiotemporal patterns are investigated by visual inspection. We then conduct data-driven analyses to identify hazard and vulnerability indicators empirically associated to the impact indicators on both levels (Fig. 1). These data-driven analyses consist of correlation checks, machine learning regression and model inspection techniques. In addition to this paper, we provide an interactive web-based visualization tool to foster the exploration of data beyond the printed figures.

105




Figure 1. Workflow of the presented study

As study region we choose the German federal state of Brandenburg, which has a relevant agricultural sector that has been affected by drought in recent years, and where reported yields as well as high spatial resolution data on grown crops are available. Brandenburg is characterized by flat topography, sandy soils and lakes stemming from the latest ice age, as well as

110



former peatland areas that have been drained over centuries for the purpose of obtaining arable land (LBGR, 2010). The climate is continental and comparably dry for German standards, with averaged precipitation around 600 mm/a, and evapotranspiration around 500 mm/a, including smaller subregions with negative water balances (Germer et al., 2011). Regional climate
115 projections indicate a further reduction in precipitation during the crop growing season, i.e. harsher conditions for agriculture (Kahlenborn et al., 2021; MLUK, 2023). Soil water is generally expected to decrease in the region (Holsten et al., 2009). Agriculture in Brandenburg is primarily rainfed, though, and current priorities of the regional water management suggest that the uptake of large-scale irrigation will not be a realistic option in the near future (MLUK, 2023). Despite this setting, the agricultural sector is very important for the region and its population in the 18 counties (in German: Landkreise, correspond
120 to NUTS-3 regions), with about 1 million ha, one third of the state, used for arable farming (MLUK, 2023). The agricultural sector of Brandenburg has also been identified as highly vulnerable to drought in European- scale studies (de Stefano et al., 2015; Blauhut et al., 2016).


2.2 Exposure & Vulnerability Indicators

125 Spatially explicit information about exposure, i.e. cropped agricultural land, is derived from the Integrated Administration and Control System (IACS), that provides the field-level data on crops for farms which have applied for annual payments within the EU's Common Agricultural Policy (CAP) (Leonhardt et al., 2023). These shapes provide the basis of our field-level analysis. We selected 12 of the most important crop types in Brandenburg in terms of area of production, for which matching information in the yield reports and average values per LBG are available (Table A1). In some cases we only used the winter
130 variety, in other cases we had to merge summer and winter varieties to match the yield reports (Table A2). The 12 crops used in this study are: winter wheat, rye, triticale, oat, winter barley, winter canola, grain maize, sunflower, potatoes, lupines, peas, and sugar beet. The total cropped area covered by our 12 selected crop types is fluctuating in the investigated time period (2013-2022) between about 638.000 to 686.000 ha, with no clear trend. The largest unconsidered fraction is silage maize, which is mostly used as fodder and thus not consistently covered in the reported statistics. Rye is among the most commonly
135 found crops in the region, and regarded as reliable source of income on sandy soils even with little precipitation (LBV, 2024). Wheat is considered to be more demanding but also to realize higher prices. Cultivation of potatoes and sugar beet has been drastically reduced over the last decades, partially owing to the increasingly dry climate (LBV, 2024). Farm level product prices were purchased from the company Agrarmarkt Informations-Gesellschaft (AMI) (cf. LELF, 2021 for publicly available data until 2020).

140 Vulnerability, a characteristic of the exposure, conceptually provides the link between hazard and impacts. Vulnerability indicators attempt to capture the relevant characteristics. We compiled a list of environmental and socio-economic indicators and their assumed direction of influence on agricultural drought vulnerability (cf. Walz et al., 2018; Meza et al., 2019; Frischen et al., 2020; Zhou et al., 2022; Stephan et al., 2023). A gridded estimate of agricultural soil quality (in German: Ackerzahl,



145 AZL) is available in 5 m resolution (Schmitz and Müller, 2020). Based on the AZL, 5 different agricultural production areas
 (in German: Landbaugebiete, LBG), are classified, for which average yields for the most important crops are published (LELF,
 2016). As a specific water-related indicator we include the plant-available water capacity (in German: nutzbare Feldkapazität,
 NFK) (BGR, 2015). To capture potential water accumulation in the landscape, we further derived the topographic wetness
 index (TWI) from a digital elevation model (BKG, 2017). We extracted mean values of AZL, NFK, and TWI per agricultural
 150 field for the available point in time, assuming that they do not change over time. Other indicators, in particular the socio-
 economic datasets, were only available per county for Brandenburg. This restricted their use to simple correlation analysis
 with impact indicators on the same spatial level. Large parts of Brandenburg are classified as “disadvantaged area” due to
 rather poor soils – the exception here being the northeastern counties Uckermark and Märkisch-Oderland. These two counties
 also exhibit the highest scores for secured succession (along with Potsdam), despite long-known problems with general
 155 unemployment in the Uckermark (10.7% in the year 2022). Smaller strips and patches of high quality LBG-1 soils are found
 in the West (Fig. 2). The spatial distribution of crop types partially reflects these patterns, e.g. winter wheat is typically grown
 on high-quality soils, making it the dominant crop type in the abovementioned areas, while rye is most common throughout
 the rest of Brandenburg on poorer, sandy soils.



160

Figure 2. Spatial distribution of agricultural soil quality (LBG). Distribution of winter wheat and rye in the year 2022.



2.3 Hazard indicators: SPEI and SMI

The Standardized Precipitation Evaporation Index (SPEI) captures both precipitation and potential evaporation and has evolved as one of the most commonly used meteorological drought indicators in recent years (Vicente-Serrano et al. 2010, Rossi et al. 2023, Tanguy et al. 2023). Monthly values of SPEI-1 (one-month accumulation SPEI) were provided by Zhang et al. (2024) on 10 km grid resolution from 2013 to 2022. As the harvest of main crops in the region typically starts in July, we used data for the months March to July. In addition to the monthly SPEI, a metric of growing season drought magnitude was computed as the sum of $\text{SPEI-1} < -0.5$ over the period between March and July (SPEI-Magnitude). Regarding soil moisture droughts, the model-based German drought monitor developed at the Helmholtz Centre for Environmental Research (UFZ) is the most established regional product (Samaniego et al., 2013; Zink et al., 2016; Boeing et al., 2022) and has already been used for similar purpose (Peichl et al., 2021). We use the monthly soil moisture index (SMI) and aggregated soil drought magnitude (SMI-Magnitude) in the top soil (25 cm), again from March to July. To add some information on slower long-term drought processes, we further include the annual drought magnitude for the total soil (up to 1.8 m depth), which is temporally aggregated from April to October (SMI-Total). Negative SPEI values indicate meteorological drought while positive SMI values indicate soil drought.

2.4 Impact indicators: crop health observations and empirical yield gaps


As an indicator of crop health, the ratio of LST and NDVI between May and June, i.e. roughly mid growing season, of each year (2013-2022) was obtained from Landsat-8 satellite imagery, using all images of the T1_L2 collection. Pre-processing and cloud-masking were conducted within the Google Earth Engine (Gorelik et al., 2017). The temporal aggregation of the satellite data is necessarily a compromise: a comparison between years gets more precise when the interval is shorter, but to smooth out potential variations in overpass and cloud cover, as well as disturbances on individual pixels, mean values across several weeks are generally more trusted (Ghazaryan et al., 2020). Images were downloaded in 30 m spatial resolution and then aggregated on individual fields. A small fraction of fields had to be discarded due to missing data, e.g. because of cloud cover, and we continued the statistical analysis with the remaining ones. As different crop types exhibit characteristic spectra, we further computed the anomalies of LST/NDVI over the entire observation period stratified by crop type. By doing so, the resulting anomalies (LST/NDVI-anom.) are comparable among different crops.

We further calculated empirical yield gaps per county for 12 crops for the last 10 years (2013-2022), by subtracting actual reported yields (total production in tonnes) by the regional statistical authority (Amt für Statistik Berlin-Brandenburg, 2022; Alencar, 2022) from an estimate of expected yields under non-drought conditions. We refer to expected yields as the product of cropped area (per crop type in a given year) and the respective 5-year average yield (tonnes per hectare) per LBG from the time 2010-2014. The expected yields are computed on field level and then aggregated on the level of counties, to be comparable to the reported yield data. We refer to relative gap as the fraction of this empirical yield gap and the expected yield. A relative



195 gap value of 1 thus implies that all yield was lost, while a value of -1 implies that double the expected amount was reported, and a value of 0 indicates a perfect match between expected and reported numbers. To correct for differences in the total area reported in IACS as compared to the yield reports, we added the difference in area per crop, multiplied with the average yield per hectare of that crop within the respective county as derived from the data. Some minor assumptions had to be made to merge crop types reported in IACS with the reports like neglecting spelt in the statistics for wheat (details in Appendix A). Multiplication of the empirical yield gaps and prices of the respective year results in a total estimate of monetary loss in euro. 200 As not all of the 12 considered crops are grown in all regions in every year, the total monetary loss estimate can be based on partially different crops per region. We assume this reflects the real agroeconomic situation in each region.

For a plausibility check, we compared the resulting empirical yield gaps and loss estimates to regional newspaper reports. For individual crops (rye, wheat, maize, barley) we were able to also calculate the potential production (PP) and water-limited 205 production (WLP) by the process model WOFOST on a 2 km grid resolution (Jänicke et al., 2017; de Wit et al., 2019). Crop growth is modelled from irradiation, temperature, CO₂ concentration, plant characteristics, seeding date, and availability of water. The physically modelled potential production from WOFOST matches very well with the expected yields derived by our empirical approach for soil quality range LBG-2 in the case of wheat and barley, and LBG-1 in the case of rye (Fig. 3). We are thus confident that our choice of reference period (2010-2014) is reasonable and can be used to produce estimates in a 210 realistic range. Only for maize the modelled potential production is higher than the average values for Brandenburg suggest on any soil type. For further comparison we use the newspaper reported impact score by (Sodoge et al., 2023), for the category “agriculture”. All data used is summarised in Table 1.



215 **Figure 3.** Comparison on field level (a) wheat (b) rye (c) barley (d) maize. The original resolution of the crop model is 2 km



Table 1. Indicators and data sources

Category	Abbreviation	Indicator Description	Spatial Res.	Data source and references	
Hazard	SPEI (monthly)	Standardized Precipitation-Evaporation Index,	10 km	Zhang et al. (2024)	
	SPEI Magnitude	Sum of SPEI < -0.5, March–July			
	SMI (monthly)	Soil Moisture Index, top soil (25 cm)	4 km	Boeing et al. (2022), UFZ Drought Monitor / Helmholtz Centre for Environmental Research https://www.ufz.de/index.php?en=37937	
SMI Magnitude	Top soil, March–July				
SMI Total	Total soil (max. 1.8 m), April–October				
Exposure	-	Agricultural land, on which one of the 12 selected crops is reported in the IACS dataset	Fields (vector)	Integrated Administration and Control System (IACS) MLUK (2022c), Leonhardt et al. (2023)	
Impact	LST/NDVI	Land Surface Temperature / Normalized Difference Vegetation Index. Mean of May–June	30 m	Landsat-8, collection: Landsat/LC08/C02/T1_L2 Courtesy of the U.S. Geological Survey (USGS) accessed via Google Earth Engine	
	LST/NDVI-anom.	Anomalies per crop			
Environmental Vulnerability	Empirical yield gap	Expected - Reported	County	5-year average hectare yield per LBG: LELF (2016) Reported: Amt für Statistik Berlin-Brandenburg (2022) Compiled by Alencar (2022) https://github.com/pedroalencar1/CropYield_BBr	
	Relative yield gap	(Expected - Reported) / Expected where Expected is based on 5-year average hectare yields per LBG and the annual area in ha			
	Loss estimate	Sum (empirical yield gap * farm level price), for all crops reported in a county per year	County	Farm level prices: AMI, cf. LELF (2021) for publicly available data until 2020	
	PP	Potential production from a crop model	2 km	WOFOST: de Wit et al. (2019) Forcing: Jänicke et al., (2017)	
	WLP	Water limited production			
Modelled Gap	PP-WLP	County	Sodoge et al. (2023)		
Environmental Vulnerability	Newspaper reported-impacts	Number of newspaper articles reporting agricultural drought impacts (text-mining based)	County	Sodoge et al. (2023)	
	AZL	Agricultural soil quality (“Ackerzahl”),	5 m	Schmitz & Müller (2020) LELF (2021)	
	LBG	5-class ordinal range (“Landbaugebiet”)			
	TWI	Topographic wetness index	200 m	BKG (2017)	
	NFK	Plant available water (“nutzbare Feldkapazität”)	250 m	BGR (2015a)	
	-	Soil depth	County	BGR (2015b)	
	-	Soil water erosion	County	BGR (2014a)	
	-	Soil wind erosion	County	BGR (2014b)	
	-	Water exchange frequency	County	BGR (2015c)	
	-	Forest ratio	County	Statistische Ämter des Bundes und der Länder (2020a)	
	-	Farmland ratio	County	Statistische Ämter des Bundes und der Länder (2020a)	
	-	Protected area	County	LfU (2020)	
	-	Disadvantaged area	County	MLUK (2022b)	
	-	Livestock health	County	Statistische Ämter des Bundes und der Länder (2020b)	
	Socio-economic Vulnerability	-	Secured succession	County	Statistische Ämter des Bundes und der Länder (2020b)
		-	Poverty	County	Amt für Statistik Berlin Brandenburg (2019b)
		-	Education	County	Statistische Ämter des Bundes und der Länder (2021)
-		Unemployment	County	Statistische Ämter des Bundes und der Länder (2022)	
-		Social dependency	County	Eurostat (2021)	
-		Agricultural population density	County	Eurostat (2021), Statistische Ämter des Bundes und der Länder (2010)	
-		GDP per farmer	County	Eurostat (2022), Statistische Ämter des Bundes und der Länder (2010)	
-		GDP per capita	County	Eurostat (2021, 2022)	
-		Agricultural dependency for livelihood	County	Statistische Ämter des Bundes und der Länder (2020d)	
-		Public participation (voting)	County	Amt für Statistik Berlin- Brandenburg (2019a)	
-	Investments in DRR (less favoured areas)	County	MLUK (2022a)		

2.5 Statistical procedures and algorithms

Exploratory analysis is conducted by calculating descriptive statistics, correlation matrices and by visual inspection of spatiotemporal patterns in the data. For the latter, we additionally provide a simple web app in R-Shiny. Changes over the investigated years are analysed by plotting the shift of statistical distributions and the temporal evolution of regional mean values. To investigate empirical relations between our hazard and vulnerability indicators and the to our impact indicators, we apply the statistical learning algorithm extreme gradient boosting (XGBoost) (Chen and Guestrin, 2016) combined with the model inspection technique Shapley additive explanations (SHAP) (Shapley, 1953; Lundberg and Lee, 2017). This



230 combination is widely used in the field of explainable artificial intelligence (XAI) and has recently been successfully applied
in many different scientific studies to derive insights from complex non-linear and interacting datasets (Yang et al., 2021; Jena
et al., 2023; Raihan et al., 2023; Li et al., 2024). XGBoost is an ensemble method based on boosting, i.e. consecutive models
are trained on the residuals of the predecessor, thereby increasing the fit step-by-step (as opposed to bagging like in Random
Forest, where an ensemble is trained in parallel fashion and aggregated via majority voting). We use a common tree-based
235 model variant to allow for a hierarchical structure. As sampling scheme, we implemented a nested cross-validation, with an
inner loop for hyperparameter optimization and an outer loop to assess the skill on independent holdout sets (not used in
parametrization). SHAP values were computed for the best model of each nested iteration, selected by the highest R^2 score on
the holdout set. The SHAP values represent a game-theoretic estimate of effect size, where the feature values are treated as
players that can join a coalition game (model). The resulting values give the expected marginal contribution for each feature
240 value across all possible coalitions, in the unit of the model target, and fulfill the efficiency property, meaning that they sum
up to the difference between the overall expected value and the specific model prediction for a set of feature values. By
computing these SHAP values for all samples used to construct a model, it is possible to visualize the effect each feature has
within the inspected model. Note that this does not necessarily imply insights into processes in nature, but rather into empirical
relations in the data as learned by the specific model.

245 In total our dataset contains 437.476 agricultural fields across 18 counties. With 12 crop types and 10 years the theoretical
maximum number of data points on county level is 2160, of which missing entries have to be removed (not all crops grown in
all counties in all years). Predictive features on field level are the indicator values. Some feature engineering is necessary to
convert the field-level data into features on county level. It is reasonable to assume that damaging processes are more dependent
250 on extreme conditions than on the mean value over a large area. To retain as much information about the hazard distributions,
we computed the relative affected area (non-)exceeding specified thresholds (in regular intervals of 0.5 for SPEI, 0.25 for
LST/NDVI-anom., 0.05 for SMI, 5 for SMI-Total, and using the LBG class limits for AZL). A total of 68 features were created
this way on county level.

3 Results & Discussion


255 3.1 Spatiotemporal patterns of hazard, vulnerability and impact indicators

3.1.1 Temporal evolution on country level

The temporal evolution of mean indicator values for entire Brandenburg suggests that the investigated decade can be divided
into a pre-drought phase (2013-2017) and a drought phase (2018-2022) (Fig. 4). In 2013 and 2014 the SMI-Total is close to 0,
observed vegetation health is at its maximum (i.e. negative LST/NDVI-anom.), essentially no impact-related statements are



260 captured in the newspaper text-mining data, and our economic calculation even estimates a plus of about 100 million euro
compared to the expectations. Especially 2014 indeed made headlines with record-breaking (positive) yields (Agrarheute,
2014). However, the crop model WOFOST still estimates a gap between potential production and water-limited production in
that year, and also in the SPEI magnitude there is some drought signal visible. We interpret this as locally and temporally
constrained meteorological effects that did not propagate to the soil and consequently did not have a negative effect on crop
265 health and yields. The year 2017 was then rather wet, which is reflected in SPEI, LST/NDVI, and the media impact statements.
However, the soil drought did only decline slightly according to the SMI. From 2018 a multi-year drought started. There seems
to be a temporal lag of 1 year between meteorological and soil moisture drought indicators, likely reflecting the propagation
from atmospheric conditions to the deeper soil layers. This is also visible in data for the year 2021, where SPEI-Magnitude
indicates a good meteorological water balance, but soil moisture drought stayed. Interesting to note though is that the satellite
270 observations of crop health peak in the same year as the SMI-Total, 2019, while the estimated economic loss (12 crops), as
well as the crop model (for wheat) and newspaper reported impacts exhibit peaks at the meteorological drought maximum in
2018. The distribution of LST/NDVI-Anomalies has been shifting towards higher values in recent years – not only the median,
but also the upper tail of the distribution became heavier (Fig. 5). This upper part of the distribution is where we expect impacts
like reduced yields. The most notable exposure changes over the decade are decreasing trends for rye (-30%), triticale (-22%),
275 winter canola (-26%), sugar beet (-33%), and lupines (-38%), increase of winter wheat (+19%), winter barley (+28%), oat
(+44%), peas (+132%), and sunflower (+145%) (Fig. 6). Changes in crop choice may partially reflect a response to experienced
crop-damaging conditions, but are also driven by unconsidered factors such as fertilizer or market prices (e.g. Albers et al.,
2017). Our total loss estimate from the 12 crops for Brandenburg 2018 is 132 million euro, which comes close to the official
numbers: 72 million euro of compensations have been issued by the state, and this sum was considered to account for about
280 45% of actual claims (which would translate to a loss of 160 million euro when taken at face value) (MLUK 2019).



285 **Figure 4.** Evolution of indicator values over entire Brandenburg. Our loss estimate is given in the original values on the left axis. Bars
 indicate the number of agricultural impact statements in newspapers on the original scale (left axis). All other indicators were extracted on
 agricultural fields, area weighted, and scaled to fit the same axis. SPEI-Magnitude has only negative values and is thus scaled to [-1, 0].
 SMI-Total and crop model-based gap (PP-WLP) have only positive values and are thus scaled to [0, 1]. LST/NDVI-Anom. has positive
 and negative values and is thus scaled to [-1, 1].

290




Figure 5. Shifting distribution of LST/NDVI anomalies by year.





Figure 6. Area covered by 12 selected crop types in Brandenburg

295



300

Our empirical yield gaps peak in the year 2018 for most crops in most regions, but the variability between counties is high for most crops (Fig. 7). Only winter wheat, winter canola, and winter barley exhibit low to moderate variability between counties. Sugar beet is only reported in a few cases. Plausibility checks against newspaper articles suggest that our relative gap estimates are in a reasonable range: yield reduction for individual crops from 25% to more than 50% have been reported in 2018 and 2019, with winter canola performing worse in 2019 (Agrarheute, 2018; DLF 2019). Grain crops did better in 2022 than 2021, but maize much worse (Tagesschau, 2022). The year 2014 on the other hand is remembered for record-breaking yields with “+24% compared to the previous 5-year average and 11% higher than the previous year”, indicating that 2013 was still well above average (Agrarheute, 2014), which is captured in our estimates.



305


Figure 7. Relative yield gaps per county in percent for the 12 investigated crops.



3.1.2 Spatiotemporal Patterns

310 A more faceted picture appears when comparing the spatial distributions of hazard and impact indicators alongside each other
for consecutive years (Fig. 8). Essentially the entire state of Brandenburg was affected by meteorological drought in 2018,
with the SPEI-Magnitude minimum registered in the South-West. Soils in the South were already dry by then, but severe soil
moisture drought throughout the country developed a year later. Contrarily, during the rather rainy year 2021 the accumulated
soil drought persisted. When another intense meteorological drought struck in 2022, only the soils in the North had moderately
315 hazard indicators. Patches of high anomalies (i.e. supposedly damaged fields) are found scattered across the country, while
low anomalies (i.e. supposedly healthy crops) appear to dominate in the areas of good soil quality (cf. Fig. 2). The highest
economic loss per hectare is mapped in the southern areas Spree-Neiße and Oberspreewald-Lausitz (the highest absolute loss
in the Uckermark, due to the large fraction of agricultural land). While the exceptional years 2018 and 2019 also caused severe
losses in the North and West of Brandenburg, the South-East ranks high in the relative loss estimates throughout all of the
320 investigated years. Loss per hectare from our empirical approach is higher than the crop model estimates by Söder et al. (2022),
who report separate numbers of around 90 euro per hectare from summer drought plus 60 euro per hectare from spring drought
in 2018 in the region. Our estimates refer to the sum of all damaging processes.

325 A number of socio-economic vulnerability indicators are particularly concerning in the North-Western areas Prignitz and
Ostprignitz-Ruppin, as well as in the Southern county Oberspreewald-Lausitz: those regions rank above average on agricultural
dependency for livelihood and below average on secured succession, while Prignitz has a particularly high agricultural
population density on top. All three exhibit low scores for the coping capacity indicators education and participation in local
politics. The socioeconomic vulnerability indicators and low resolution maps for all investigated years can be viewed at
<https://fabiobrill.shinyapps.io/agrdrought-explorer-brandenburg/>, while the high resolution data can be obtained from the
330 GitHub repository <https://github.com/fabiobrill/brandenburg-drought-study>.



335 **Figure 8.** Spatiotemporal patterns of aggregated meteorological and soil moisture drought hazard indicators, crop health anomalies, and county-scale loss estimates per hectare.



3.2 Empirical investigation of impact-relevant factors

3.2.1 Relations between indicators on field level

340 The spatiotemporal patterns suggest non-trivial and multi-way interactive relationships between our chosen hazard, vulnerability and impact indicators. This is further supported by a correlation analysis, which shows that the bivariate linear relations in the data are mostly weak (Fig. 9). Correlations slightly increase when subdividing the data by crop, presumably because the relationships are more linear for individual crops, however the effect is almost negligible (not shown). The meteorological and soil moisture hazard indicators SPEI and SMI are correlated among each other. Monthly SPEI and SMI are essentially uncorrelated to LST/NDVI-anom. in March, very weakly correlated in April, and moderately correlated in May and June. As the LST/NDVI measurements are also from May and June, the additional correlation in July has to be a spurious effect stemming from the collinearity in the SPEI layers (almost 0.5 between June and July). Raw NDVI – and therefore also LST/NDVI – is clearly related to AZL, meaning that crops grown on better soils tend to be “greener”, with or without drought. This effect is reduced in the anomalies. TWI and NFK exhibit no relation except to AZL. The modelled water-limited production from WOFOST only weakly relates to LST/NDVI (not shown).

350




Figure 9. Pearson’s correlation coefficient on field level data. Almost all correlations are statistically significant due to the high number of samples (n=437.245 complete observations, 474.966 in total).

355



An XGBoost model trained to predict LST/NDVI-anom. from monthly hazard indicators SPEI and SMI, aggregated SMI-Total, environmental vulnerability factors AZL, TWI, and NFK, as well as crop type, obtains R^2 scores around 0.5 (Appendix B). AZL is chosen by the models as most important feature, followed by the categorical variable crop type, while the other environmental vulnerability factors, TWI and NFK, have little influence (Fig. 10). Each dot in these plots corresponds to a sample, and the SHAP values represent the feature effects (conditional expectation) on the predicted quantity, i.e. LST/NDVI-anom. in this case. Interaction plots for crop types highlight that wheat, canola, and barely are grown on relatively good soils, lupines on bad soils, and rye on both (Fig. 11). While the absolute effect of AZL on the predictions is higher than the effect of crop type, particularly wheat is modelled to be impacted more likely than other crops despite growing on better soils (higher AZL).

360

365

370




Figure 10. SHAP summary plot.




Figure 11. Interaction plot for crop type and AZL.

375




Figure 12. SHAP dependence plots for selected features: (a) AZL, (b) TWI (c) SMI Total, (d) SPEI March, (e) SMI May, (f) SPEI June. Centre line derived by a loess regression on the SHAP values. Colour visualizes interaction with a second feature.

380 From a methodological point of view, it is worth to mention that SHAP plots based on the full dataset exhibit far larger variance
 on the y-axis than preliminary experiments with only 10% of the data. One reason for this might be the spatial resolution of
 the features, but we assume that it is also related to the complexity of the regression task. While there are some clear effects in
 the centre lines, it also becomes obvious that no single feature explains the full data. Several steps in our analysis include
 simplifications, e.g. calculations using mean values per field imply that an entire field is treated as a unit. For larger fields it
 385 might be realistic that only parts are affected, however such effects are below the credible resolution of input data. We
 acknowledge that particularly for maize, which is typically harvested from September on, a longer observation window might
 be better suited. Adjusting the remote sensing data to the actual sowing and harvest dates of each crop might improve the
 results – however, doing so would further complicate the data pre-processing and was considered out of scope of this study.
 Although agriculture in Brandenburg is predominantly rainfed, a future study could also benefit from spatially explicit
 390 information on irrigated areas (Ghazaryan et al. 2022).


3.2.2 Relations between indicators on county level

When arranging the 12 crops by correlation among the relative gaps (i.e. each sample referring to a county in a given year), it
 appears that almost all crops are positively correlated over time, while spatially (and thus spatiotemporally) several groups
 395 emerge (not shown). Correlation between the newspaper-based “agriculture” impact score by Sodge et al. (2023) and our
 relative economic impact measure (in euro per hectare) over all 12 crops for the years 2013-2022 is 0.75 for entire Brandenburg



and 0.53 on county level. When compressing the data to mean values over the entire timespan to merge them with the socio-economic vulnerability indicators, the highest correlation of newspapers reported impacts is to participation in local politics (0.69). From our data and analysis, we see no meaningful correlation between the vulnerability indicators to reported impacts or calculated losses. A major drawback is the resolution of the indicators. For these reasons they were not included in the following XGBoost regression analysis.

The statistical learning models trained to predict the empirical yield gaps on county level obtain R^2 scores around 0.6 when using all features and all data (Fig. 13). Models using only LST/NDVI features as predictors perform poorly ($R^2 \sim 0.2$). It is quite remarkable that a field-level (i.e. high spatial resolution) observation of crop health does not provide more useful information for predicting yield. Models using hazard indicators as predictors perform better. Monthly values of SPEI are clearly to be preferred over seasonally aggregated magnitude, and the same is true for SMI. However, we observe that models using only SPEI perform slightly better than those using only SMI. One potential reason for this might be that the SMI is itself model-based, which introduces further uncertainty. We find a minor improvement when using both SPEI and SMI, where SMI-total is more relevant than the monthly top soil layers (as complementary information to SPEI). The additional improvement when adding LST/NDVI features on top is almost negligible. Our predictive features explain much more variance for the drought years 2018-2022 than for the pre-drought years 2013-2017, as expected. Models trained on the full dataset exhibit both higher skill and less variance. A similar effect is observed when training separate models for the different crop types: individual models for winter wheat perform better than individual models for rye, but a lumped model using all crops is much more stable. We explain this by the higher number of training samples in combination with a tree-based model structure that exploits similarities between crops. The R^2 skill score of the final model used for inspection via SHAP plots is 0.62 on the holdout set, i.e. about 60% of the variance in the empirical yield gaps can be explained by our drought-related features, while about 40% remain unexplained. Agricultural crops are highly managed and face numerous threats, not only droughts. It would be unreasonable to assume that drought indicators alone could fully explain real observed yield data. In a similar published attempt, Peichel et al. (2021) report that their best model for winter wheat obtained an R^2 of 0.68, which is very close to our best models – however, they do not report any details on the variability of this score. Empirical damage models, such as used for floods, typically report rather weak model fits (e.g. Wagenaar et al. 2017, Sieg et al. 2017). In the European Drought Risk Atlas, Rossi et al. (2023) do not even report model fit at all, but still uncover plausible impact-relevant factors for droughts.



425 **Figure 13.** Distributions of the R² skill score based on 10 repetitions for each setup. (a) Separate models for pre-drought and drought
430 years. (b) Separate models for individual crop types.

435 Model inspection identifies SPEI below -1 in June as the most relevant condition in the lumped model for all crops (Fig. 14) and also in crop-specific models for wheat and rye (Appendix C). Note that the features on county level always refer to the relative affected area above or below a threshold, e.g. the value of “SPEI June < -1” indicates the relative area per crop per county affected by SPEI in June below -1. However, a large fraction of the data indicates that non-exceeding -1 coincided with negative empirical yield gaps, i.e. higher than expected yields. To investigate this in more detail, we run another model setup using only data with positive empirical yield gap (n=827). Data on county level always includes mixed effects, i.e. the constraint “empirical yield gap > 0” on county level does not imply that there are no damaged fields in the data, but rather that damaged fields are outweighed by fields with higher than expected yields within the same county. Features based on SPEI in June are still among the most important predictors for such a subset, with thresholds of -0.5, and -1 ranked high (Fig. 14b). Even more severe meteorological drought conditions (SPEI < -2) are apparently just too rare in this dataset to be influential on county level. In March the threshold of 0 is again selected in reverse direction, i.e. indicating damage from too wet conditions (cf. Fig. 15d). Multiple AZL features are selected, confirming once more that soil quality is a relevant drought vulnerability factor (the regression target is already based on expected yield estimates that account for AZL, so this effect is on top). LST/NDVI as predictive feature for the empirical yield gaps is of low relevance when using all data, but ranks higher when restricting the training data to positive yield gaps. In the comparison of crops (Fig. 15a), lupines clearly stick out, which is explained by the high losses in the yield data (cf. Fig. 7). The interaction of crop type with AZL < 36 shows once more that rye is growing on worse soils than wheat, but still has lower SHAP values with respect to the regression on impacts. Triticale is on a similar level as wheat, canola even higher. From all these crops, rye is thus empirically found the most robust.

445 To check the stability of the SHAP values, we repeated the model fitting several times and inspected the resulting summary plots. The first features are always crop type and SPEI in June. Beyond the first few ranks, feature effects get very similar and the exact ranks can shift in repeated model runs (depending on the random data subset and respective model parameters). The



455

effects for the different crop types and shapes of the dependence plots are also stable results, confirmed in multiple setups. Focussing the models on positive empirical yield gaps can make the feature effects more linear (Fig. 15b and 15c). Nonlinear responses in the dependence plots for single features on county level are likely empirical artefacts, as the definition of a feature as relative affected area should more or less linearize the physical response. Although spatial neighbourhood effects, like water lacking in a hydrologically connected area, could introduce nonlinearities, we assume in general that more affected area should lead to more impact, regardless of the criterion.




Figure 14. SHAP summary plots for the best model trained on (a) all data, and (b) empirical yield gap > 0

460




Figure 15. SHAP dependence plots for selected features on county level. (a) Effect of crop type and interaction with relative area of AZL < 36 from a model trained on all data (b) Effect of SPEI in June < -1 from a model trained on all data and (c) from a model trained only on positive empirical yield gaps. (d) Effect of SPEI in March < 0 from a model trained only on positive empirical yield gaps. (e) Effect of LST/NDVI anomaly > 0.5 from a model trained only on positive empirical yield gaps.

465

4. Conclusion

Our analysis of spatiotemporal patterns of agricultural drought hazard, exposure, vulnerability, and impact indicators for Brandenburg, 2013-2022, empirically shows that the links between these components are complex and, consequently, risk mapping and monitoring need to be supported by thorough investigations from multiple datasets. We present agricultural impact indicators on two spatial levels – the crop health indicator LST/NDVI on individual fields, and empirical yield gaps on county level – and relate them. Anomalies of LST/NDVI are shifted to higher values during the drought years, but spatial patterns are rather scattered. The South-East ranks high in our per-hectare economic loss estimates throughout all of the investigated years, although in the exceptional years 2018 and 2019 high losses are also registered in the North and West of Brandenburg. It is not immediately obvious how the spatial patterns of the individual hazard and vulnerability indicators relate to both impact indicators. Statistical learning algorithms can be used to establish the missing link between the indicators.

470

475

We apply XGBoost regression to model the impact indicators on both scales, and compare the SHAP values to derive impact-relevant factors from the data. We find the importance of SPEI in June for regressing the observed impacts substantiated by multiple model setups: (1) On field level, regressing LST/NDVI-anom., the SHAP values of SPEI in June strongly increase below -1. (2) On county level, regressing empirical yield gaps, the relative area affected by SPEI < -1 is selected as most important predictive feature for a model trained on all data, as well as for crop-specific models (both wheat and rye). (3) Even

480



when removing all data where empirical yield gap < 0 , i.e. more yield reported than expected, SPEI features from June still top the ranking, although several thresholds are selected (mainly -0.5 and -1). This is of particular concern as current regional climate simulations for Brandenburg project a shift in seasonality of rainfall: more in winter, and less during summer months.

485 Too wet conditions in March are found to be an impact-relevant factor, in agreement with Peichl et al. (2021). SMI-Total adds complementary information to monthly SPEI. No real model improvement obtained when using both SPEI and SMI monthly values, though. From the considered vulnerability factors, AZL (i.e. agricultural soil quality) is by far the most relevant one. There is a clear influence of AZL on LST/NDIV-anom., with vulnerability rising at AZL below about 35. LST/NDVI is, somewhat surprisingly, not a good predictor for the empirical yield gaps in our study. We thus advise caution when interpreting

490 empirical results from a single impact indicator. AZL is also related to selected crop types. Most notably, wheat is grown on high quality soils, while rye predominantly on low to medium quality soils. Rye is empirically found more robust under drought conditions in the region – based on both impact datasets. The cropped area of rye decreased by about 30% between 2013 and 2022 in Brandenburg, though, and the area for winter wheat increased by 19% in the same time. Such choices of crop types simultaneously affect exposure and vulnerability, and thus risk.

495 Data-driven techniques can capture complex interactions in human-environments such as agriculture. SHAP plots uncover which factors drive the prediction of impact indicators in the models. This does not necessarily relate to causal effects in nature, though. We thus suggest to cross-check results obtained from different model setups and different regression targets. Model inspection in this study shows that features are generally used in a physically meaningful direction, which is a prerequisite if

500 data-driven models are to be trusted. No single feature explains the full data, though, and in fact such simplified interpretations are against the logic of using a strongly nonlinear ML algorithm to tackle complex regression problems. Rather than attempting to weight indicators manually, empirical impact data should be the benchmark to evaluate hazard and vulnerability indicators for the purpose of risk mapping. Further improvements in modelling observed impacts likely require more detailed spatially explicit data on vulnerability and management, e.g. irrigation. Monitoring and impact-based forecasting are needed to prepare

505 for future hazards, which can hardly be mitigated. Stronger remote sensing indicators on drought impacts seem necessary in that context. To enable farmers and the general population to evaluate local adaptation options, i.e. changes in exposure and vulnerability, risk-related data needs to be made generally available – not only to policy-makers. Interactive visualization tools allow for exploring and foster a more open communication of risk-related geoinformation.

510 **Acknowledgements:**

This research was supported by the Einstein Research Unit “Climate and Water under Change” from the Einstein Foundation Berlin and Berlin University Alliance (ERU-2020-609), and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – SFB 1502/1–2022 - project number: 450058266. We thank Jan Sodoge for providing the newspaper text-mining data for the full investigated period of time, and acknowledge the preliminary work conducted by Thomas

515 Hoffmann and Marlen Laudien during their theses at HU Berlin.



Author contribution

520 FB(1): Conceptualization, Methodology, Software, Formal analysis, Validation, Visualization, Writing–original draft. PA: Resources, Writing–review & editing. HZ: Resources, Writing–review & editing. FB(2): Resources, Writing–review & editing. SH: Resources, Writing–review & editing. TL: Conceptualization, Resources, Writing–review & editing, Funding acquisition. All authors have read and agreed to the submitted version of the manuscript.

Competing interests

525 PA is member of the editorial board of this special issue. All other authors declare that they have no conflict of interest.

Code and data availability

530 All data and scripts needed to reproduce the figures are publicly available via GitHub: <https://github.com/fabiobrill/brandenburg-drought-study/>. The processed dataset is stored on GitHub LFS and will be transferred to a permanent repository with a DOI during the publication process. Scripts used to conduct the preprocessing and analysis are publicly available via the same repository. Except for the crop prices, which were obtained from AMI under a commercial license, all other raw data used in this study are either open (see Table 1) or can be made available upon reasonable request to the authors. The interactive data exploration app in R-Shiny is also available via the GitHub repository and can be run locally. An independent publicly hosted version is accessible online: <https://fabiobrill.shinyapps.io/agrdrought-explorer-brandenburg/>

535 References

- Abdullah, M. F., Siraj, S., and Hodgett, R. E.: An Overview of Multi-Criteria Decision Analysis (MCDA) Application in Managing Water-Related Disaster Events: Analyzing 20 Years of Literature for Flood and Drought Events, *Water-Sui*, 13, 1–27, <https://doi.org/10.3390/w13101358>, 2021.
- 540 Abunywah, M., Okyere, S. A., Opoku Mensah, S., Erdiaw-Kwasie, M., Gajendran, T. and Byrne, M. K.: Drought impact on peri-urban farmers' mental health in semi-arid Ghana: The moderating role of personal social capital, *Environmental Development*, 49, 1-18, <https://doi.org/10.1016/j.envdev.2023.100960>, 2024.
- Agrarheute, <https://www.agrarheute.com/pflanze/brandenburg-rekordernte-gerste-raps-446957>, last accessed: 06 March 2024, 2014.
- Agrarheute, <https://www.agrarheute.com/markt/marktfruechte/erste-bilanzen-neue-prognosen-katastrophale-ernte-norden-545961>, last accessed: 06 March 2024, 2018.
- 545 Albers, H., Gornott, C., and Hüttel, S.: How do inputs and weather drive wheat yield volatility? The example of Germany, *Food Policy*, 70, 50-61, [10.1016/j.foodpol.2017.05.001](https://doi.org/10.1016/j.foodpol.2017.05.001), 2017.
- Alencar, P. H. L. and Paton, E. N.: How do we identify flash droughts? A case study in Central European Croplands, *Hydrol. Res.*, 53, 1150–1165, <https://doi.org/10.2166/nh.2022.003>, 2022.
- 550 Amt für Statistik Berlin-Brandenburg: Kommunalwahlen Im Land Brandenburg: Endgültiges Ergebnis Der Wahlen Zu Den Kreistagen Der Landkreise Und Stadtverordnetenversammlungen Der Kreisfreien Städte, <https://www.statistik-berlin-brandenburg.de/kommunalwahlen-brandenburg>, last accessed: 29 March 2024, 2019a.



- Amt für Statistik Berlin-Brandenburg: Regionaler Sozialbericht Berlin Und Brandenburg 2019, <https://web.statistik-berlin-brandenburg.de/instantatlas/interaktivekarten/sozialbericht/atlas.html>, last accessed: 29 March 2024, 2019b.
- Amt für Statistik Berlin-Brandenburg: Ernteberichterstattung über Feldfrüchte und Grünland in Brandenburg, <https://www.statistik-berlin-brandenburg.de/c-ii-1-m>, last accessed: 29 March 2024, 2022
- 555 Austin, E. K., Handley, T., Kiem, A. S., Rich, J. L., Lewin, T. J., Askland, H. H., Askarimarnani, S. S., Perkins, D. A. and Kelly, B. J.: Drought-related stress among farmers: findings from the Australian Rural Mental Health Study, *The Medical Journal of Australia*, 209, 159–165, <https://doi.org/10.5694/mja17.01200>, 2018.
- Bachmair, S., Stahl, K., Collins, K., Hannaford, J., Acreman, M., Svoboda, M., Knutson, C., Smith, K. H., Wall, N., Fuchs, B., Crossman, N. D., and Overton, I. C.: Drought indicators revisited: the need for a wider consideration of environment and society, *WIREs Water*, 3, 516–536, <https://doi.org/10.1002/wat2.1154>, 2016.
- 560 BGR (Bundesanstalt für Geowissenschaften und Rohstoffe): Potentielle Erosionsgefährdung Der Ackerböden Durch Wasser in Deutschland Auf Basis von Klimaszenarien, <https://geoportal.bgr.de/mapapps/resources/apps/geoportal/index.html?lang=de#/datasets/portal/35d9601a-84f3-4e33-9436-69509bfd48c4>, last accessed: 29 March 2024, 2014a
- BGR (Bundesanstalt für Geowissenschaften und Rohstoffe): Potentielle Erosionsgefährdung Der Ackerböden Durch Wind in Deutschland. https://www.bgr.bund.de/DE/Themen/Boden/Ressourcenbewertung/Bodenerosion/Wind/PEG_wind_node.html, last accessed: 29 March 2024, 2014b.
- 565 BGR (Bundesanstalt für Geowissenschaften und Rohstoffe): Nutzbare Feldkapazität Im Effektiven Wurzelraum in Deutschland, https://www.bgr.bund.de/DE/Themen/Boden/Produkte/produktkatalog_node.html?jsessionid=78D8CD03B9D125DBDF2B089280CFCB6E.inter.net942, last accessed: 29 March 2024, 2015a.
- 570 BGR (Bundesanstalt für Geowissenschaften und Rohstoffe). Physiologische Gründigkeit Der Böden Deutschlands (PhysGru1000_250 V1.0), <https://www.govdata.de/daten/-/details/physiologische-grundigkeit-der-boden-deutschlands>, last accessed: 29 March 2024, 2015b.
- BGR (Bundesanstalt für Geowissenschaften und Rohstoffe): Austauschhäufigkeit Des Bodenwassers in Landwirtschaftlich Genutzten Böden Deutschlands, <https://geoportal.bgr.de/mapapps/resources/apps/geoportal/index.html?lang=de#/datasets/portal/3bfef3bf-855a-435f-9d88-553b10000a4c>, last accessed: 29 March 2024, 2015c.
- 575 BKG (Bundesanstalt für Kartographie und Geodäsie): Digitales Geländemodell Gitterweite 200 M, <https://mis.bkg.bund.de/trefferanzeige?docuuid=eaaa67a1-5ecb-4e57-af38-b5f1d6d57e2a>, last accessed: 29 March 2024, 2017.
- Blauhut, V.: The triple complexity of drought risk analysis and its visualisation via mapping: a review across scales and sectors, *Earth-Sci. Rev.*, 210, 1–22, <https://doi.org/10.1016/j.earscirev.2020.103345>, 2020.
- 580 Blauhut, V., Stahl, K., Stagge, J. H., Tallaksen, L. M., de Stefano, L., and Vogt, J.: Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors, *Hydrol. Earth Syst. Sci.*, 20, 2779–2800, <https://doi.org/10.5194/hess-20-2779-2016>, 2016.
- Brill, F., Passuni Pineda, S., Espichán Cuya, B., and Kreibich, H.: A data-mining approach towards damage modelling for El Niño events in Peru, *Geomat. Nat. Haz. Risk*, 11, 1966–1990, <https://doi.org/10.1080/19475705.2020.1818636>, 2020.
- Boeing, F., Rakovec, O., Kumar, R., Samaniego, L., Schrön, M., Hildebrandt, A., Rebmann, C., Thober, S., Müller, S., and Zacharias, S.: High-resolution drought simulations and comparison to soil moisture observations in Germany, *Hydrol. Earth Syst. Sci.*, 26, 5137–5161, <https://doi.org/10.5194/hess-26-5137-2022>, 2022.
- 585 Chen, T. and Guestrin, C.: XGBoost: A Scalable Tree Boosting System, 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, San Francisco, USA, 13 – 17 August, 785–794, <http://dx.doi.org/10.1145/2939672.2939785>, 2016.
- Contreras, D.: The Integrated Spatial Pattern of Child Mortality during the 2012–2016 Drought in La Guajira, Colombia, *Sustainability-Basel*, 11, 1–23, <https://doi.org/10.3390/su11247190>, 2019.
- 590 Crocetti, L., Forkel, M., Fischer, M., Jurečka, F., Grlj, A., Salentinig, A., Trnka, M., Anderson, M., Ng, W.-T., Kokalj, Ž., Bucur, A., and Dorigo, W.: Earth Observation for agricultural drought monitoring in the Pannonian Basin (southeastern Europe): current state and future directions, *Reg. Environ. Change*, 20, 1–17, <https://doi.org/10.1007/s10113-020-01710-w>, 2020.



- 595 Dabanli, I.: Drought hazard, vulnerability, and risk assessment in Turkey, *Arab. J. Geosci.*, 11, 1–12, <https://doi.org/10.1007/s12517-018-3867-x>, 2018.
- De Sherbinin, A., Bukvic, A., Rohat, G., Gall, M., McCusker, B., Preston, B., Apotsos, A., Fish, C., Kienberger, S., Muhonda, P., Wilhelm, O., Macharia, D., Shubert, W., Sliuzas, R., Tomaszewski, B., and Zhang, S.: Climate vulnerability mapping: A systematic review and future prospects, *WIREs Clim. Change*, 10, 1–23, <https://doi.org/10.1002/wcc.600>, 2019.
- De Stefano, L., González Tánago, I., Ballesteros, M., Urquijo, J., Blauhut, V., Stagge, J. H. and Stahl, K.: Methodological approach considering different factors influencing vulnerability - pan-European scale, Technical Report No. 26, 2015.
- 600 De Wit, A., Boogaard, H., Fumagalli, D., Janssen, S., Knapen, R., van Kraalingen, D., Supit, I., van der Wijngaart, R., and van Diepen, K.: 25 years of the WOFOST cropping systems model, *Agr. Syst.*, 168, 154–167, <https://doi.org/10.1016/j.agsy.2018.06.018>, 2019.
- DLF (Deutschlandfunk), <https://www.deutschlandfunkkultur.de/duerre-in-brandenburg-noch-schlimmer-als-letztes-jahr-100.html>, last accessed: 06 March 2024, 2019.
- Eurostat: Population on 1 January by Age Group, Sex and NUTS 3 Region (demo_r_pjangrp3), [https://ec.europa.eu/eurostat/databrowser/product/page/DEMO_R_PJANGRP3\\$DEFAULTVIEW](https://ec.europa.eu/eurostat/databrowser/product/page/DEMO_R_PJANGRP3$DEFAULTVIEW), last accessed: 29 March 2024, 2021.
- 605 Eurostat: Gross Domestic Product (GDP) at Current Market Prices by NUTS 3 Regions (nama_10r_3gdp), https://ec.europa.eu/eurostat/databrowser/product/page/NAMA_10R_3GDP_custom_2867669, last accessed: 29 March 2024, 2022.
- Erfurt, M., Glaser, R., and Blauhut, V.: Changing impacts and societal responses to drought in southwestern Germany since 1800, *Reg. Environ. Change*, 19, 2311–2323, <https://doi.org/10.1007/s10113-019-01522-7>, 2019.
- 610 European Commission: €430 million of EU funds to support the EU agricultural sector, press release, Brussels, 1-3 https://ec.europa.eu/commission/presscorner/detail/en/IP_23_3189, 2023.
- Frischen, J., Meza, I., Rupp, D., Wietler, K., and Hagenlocher, M.: Drought Risk to Agricultural Systems in Zimbabwe: A Spatial Analysis of Hazard, Exposure, and Vulnerability, *Sustainability-Basel*, 12, 1–23, <https://doi.org/10.3390/su12030752>, 2020.
- Germer, S., Kaiser, K., Bens, O., and Hüttel, R. F.: Water Balance Changes and Responses of Ecosystems and Society in the Berlin-Brandenburg Region – a Review. *DIE ERDE – Journal of the Geographical Society of Berlin*, 142(1-2), 65–95, <https://www.die-erde.org/index.php/die-erde/article/view/43>, 2011.
- 615 Ghazaryan, G., König, S., Rezaei, E. E., Siebert, S., and Dubovyk, O.: Analysis of Drought Impact on Croplands from Global to Regional Scale: A Remote Sensing Approach, *Remote Sens.-Basel*, 12, 1–17, <https://doi.org/10.3390/rs12244030>, 2020.
- Ghazaryan, G., Ernst, S., Sempel, F., and Nendel, C.: Field-Level Irrigation Monitoring with Integrated Use of Optical and Radar Time Series in Temperate Regions, in: *Proceedings of the International Geoscience and Remote Sensing Symposium (IGARSS)*, Kuala Lumpur, Malaysia, 17–22 July 2022, 5448-5451, <https://doi.org/10.1109/IGARSS46834.2022.9884067>, 2022.
- 620 Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sens. Environ.*, 202, 18–27, <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- Hagenlocher, M., Meza, I., Anderson, C. C., Min, A., Renaud, F. G., Walz, Y., Siebert, S., and Sebesvari, Z.: Drought vulnerability and risk assessments: state of the art, persistent gaps, and research agenda, *Environ. Res. Lett.*, 14, 1–13, <https://doi.org/10.1088/1748-9326/ab225d>, 2019.
- 625 Hanel, M., Rakovec, O., Markonis, Y., Máca, P., Samaniego, L., Kysely, J., and Kumar, R.: Revisiting the recent European droughts from a long-term perspective, *Sci. Rep.-UK*, 8, 1–11, <https://doi.org/10.1038/s41598-018-27464-4>, 2018.
- Hari, V., Rakovec, O., Markonis, Y., Hanel, M., and Kumar, R.: Increased future occurrences of the exceptional 2018–2019 Central European drought under global warming, *Sci. Rep.-UK*, 10, 1–10, <https://doi.org/10.1038/s41598-020-68872-9>, 2020.
- 630 Holsten, A., Vetter, T., Vohland, K. and Krysanova, V.: Impact of climate change on soil moisture dynamics in Brandenburg with a focus on nature conservation areas, *Ecol. Model.*, 220, 2076-2087, <https://doi.org/10.1016/j.ecolmodel.2009.04.038>, 2009.
- Houmma, I. H., El Mansouri, L., Gadal, S., Garba, M., and Hadria, R.: Modelling agricultural drought: a review of latest advances in big data technologies, *Geomat. Nat. Haz. Risk*, 13, 2737–2776, <https://doi.org/10.1080/19475705.2022.2131471>, 2022.
- Ihinegbu, C. and Ogunwumi, T.: Multi-criteria modelling of drought: a study of Brandenburg Federal State, Germany, *Modeling Earth Systems and Environment*, 8, 2035–2049, <https://doi.org/10.1007/s40808-021-01197-2>, 2022.



- 635 Jakob, M., Stein, D., and Ulmi, M.: Vulnerability of buildings to debris flow impact, *Nat. Hazards*, 60, 241–261, <https://doi.org/10.1007/s11069-011-0007-2>, 2012.
- Jänicke, B., Meier, F., Fenner, D., Fehrenbach, U., Holtmann, A., and Scherer, D.: Urban–rural differences in near-surface air temperature as resolved by the Central Europe Refined analysis (CER): sensitivity to planetary boundary layer schemes and urban canopy models, *Int. J. Climatol.*, 37, 2063–2079, <https://doi.org/10.1002/joc.4835>, 2017.
- 640 Jena, R., Shanableh, A., Al-Ruzouq, R., Pradhan, B., Gibril, M. B. A., Khalil, M. A., Ghorbanzadeh, O., Ganapathy, G. P., Ghamisi, P.: Explainable Artificial Intelligence (XAI) Model for Earthquake Spatial Probability Assessment in Arabian Peninsula. *Remote Sensing*, 15 (9), 2248, <https://doi.org/10.3390/rs15092248>, 2023.
- Jhan, H.-T., Ballinger, R., Jaleel, A., and Ting, K.-H.: Development and application of a Socioeconomic Vulnerability Indicator Framework (SVIF) for Local Climate Change Adaptation in Taiwan, *Sustainability-Basel*, 12, 1–27, <https://doi.org/10.3390/su12041585>, 2020.
- 645 Kahlenborn, W., Porst, L., Voss, M., Fritsch, U., Renner, K., Zebisch, M., Wolf, M., Schönthaler, K. and Schauer, I.: Climate Impact and Risk Assessment 2021 for Germany – Summary, Umweltbundesamt, Dessau-Roßlau, https://www.umweltbundesamt.de/sites/default/files/medien/479/publikationen/cc_27-2021_climate_impact_and_risk_assessment_2021_for_germany_english_summary_bf.pdf, 2021.
- Karnieli, A., Agam, N., Pinker, R. T., Anderson, M., Imhoff, M. L., Gutman, G. G., Panov, N., and Goldberg, A.: Use of NDVI and Land Surface and Temperature for Drought and Assessment: and Merits and Limitations, *J. Climate*, 23, 618–633, <https://doi.org/10.1175/2009JCLI2900.1>, 2010.
- 650 Khoshnazar, A., Perez, G. C., and Sajjad, M.: Characterizing spatial-temporal drought risk heterogeneities: A hazard, vulnerability and resilience-based modeling, *J. Hydrol.*, 619, 1–16, <https://doi.org/10.1016/j.jhydrol.2023.129321>, 2023.
- Kim, H., Park, J., Yoo, J., and Kim, T.-W.: Assessment of drought hazard, vulnerability, and risk: A case study for administrative districts in South Korea, *J. Hydro-environ. Res.*, 9, 28–35, <https://doi.org/10.1016/j.jher.2013.07.003>, 2015.
- 655 Kim, S. J., Park, S., Lee, S. J., Shaimerdenova, A., Kim, J., Park, E., Lee, W., Kim, G. S., Kim, N., Kim, T. H., Lim, C.-H., Choi, Y., and Lee, W.-K.: Developing spatial agricultural drought risk index with controllable geo-spatial indicators: A case study for South Korea and Kazakhstan, *Int. J. Disast. Risk Re.*, 54, 1–12, <https://doi.org/10.1016/j.ijdr.2021.102056>, 2021.
- Kondylatos, S., Prapas, I., Ronco, M., Papoutsis, I., Camps-Valls, G., Piles, M., Fernández-Torres, M.-Á., and Carvalhais, N.: Wildfire Danger Prediction and Understanding With Deep Learning, *Geophys. Res. Lett.*, 49, 1–11, <https://doi.org/10.1029/2022gl099368>, 2022.
- 660 Kreibich, H., van Loon, A.F., Schröter K., Ward P.J., Mazzoleni, M., Sairam, N., et. al.: The challenge of unprecedented floods and droughts in risk management, *Nature*, 608, 80–86, <https://doi.org/10.1038/s41586-022-04917-5>, 2022.
- Krishnamurthy R, P. K., Fisher, J. B., Choularton, R. J., and Kareiva, P. M.: Anticipating drought-related food security changes, *Nature Sustainability*, 5, 956–964, <https://doi.org/10.1038/s41893-022-00962-0>, 2022.
- LBGR (Landesamt für Bergbau, Geologie und Rohstoffe Brandenburg): Atlas zur Geologie von Brandenburg, Cottbus, 1-69, ISBN 978-3-9808157-4-1, https://lbgr.brandenburg.de/sixcms/media.php/9/4_Geoatlas_1-69.pdf, 2010.
- 665 LBV (Landesbauernverband Brandenburg e.V.): Ackerbau in Brandenburg, <https://www.lbv-brandenburg.de/50-themen/ackerbau/209-ackerbau-in-Brandenburg>, last access: 27 February 2024.
- LELF (Landesamt für Ländliche Entwicklung, Landwirtschaft und Flurneuordnung): Richtwerte zur Bewertung von Aufwuchsschäden an landwirtschaftlichen Kulturen im Land Brandenburg, http://www.lenzburger.de/downloads/richtwerte_aufwuchsschaeden_bb_07_11_2016.pdf, last accessed: 06 March 2024, 2016.
- 670 LELF (Landesamt für Ländliche Entwicklung, Landwirtschaft und Flurneuordnung): Datensammlung für die betriebswirtschaftliche Bewertung landwirtschaftlicher Produktionsverfahren im Land Brandenburg, <https://lelf.brandenburg.de/sixcms/media.php/9/Datensammlung-2021-web.pdf>, last accessed: 06 March 2024, 2021
- Leonhardt, H., Hüttel, S., Lakes, T., Wesemeyer, M., and Wolff, S.: Use Cases of the Integrated Administration and Control System’s Plot-Level Data: Protocol and Pilot Analysis for a Systematic Mapping Review, *Ger. J. Agr. Econ.*, 72, 168–184, <https://doi.org/10.30430/gjae.2023.0385>, 2023.
- 675



- LFU (Landesamt für Umwelt Brandenburg), Schutzgebiete Nach Naturschutzrecht Des Landes Brandenburg, <https://geobroker.geobasis-bb.de/gbss.php?MODE=GetProductInformation&PRODUCTID=AB2F53A4-A68E-413F-84C4-A972D2A2DA0B>, last accessed: 29 March 2024, 2020.
- 680 Li, H., Vulova, S., Rocha, A. D., and Kleinschmit, B.: Spatio-temporal feature attribution of European summer wildfires with Explainable Artificial Intelligence (XAI), *Sci. Total Environ.*, 916, 1–13, <https://doi.org/10.1016/j.scitotenv.2024.170330>, 2024.
- Lundberg, S. M. and Lee, S.-I.: A Unified Approach to Interpreting Model Predictions, 1-9, arXiv [preprint], <http://arxiv.org/pdf/1705.07874.pdf>, 2017.
- Lüttger, A. B. and Feike, T.: Development of heat and drought related extreme weather events and their effect on winter wheat yields in Germany, *Theor. Appl. Climatol.*, 132, 15–29, <https://doi.org/10.1007/s00704-017-2076-y>, 2018.
- 685 McVicar, T. R. and Bierwirth, P. N.: Rapidly assessing the 1997 drought in Papua New Guinea using composite AVHRR imagery, *Int. J. Remote Sens.*, 22, 2109–2128, <https://doi.org/10.1080/01431160120728>, 2001.
- Merz, B., Kreibich, H., and Lall, U.: Multi-variate flood damage assessment: a tree-based data-mining approach, *Nat. Hazard Earth Sys.*, 13, 53–64, <https://doi.org/10.5194/nhess-13-53-2013>, 2013.
- Meza, I., Hagenlocher, M., Naumann, G., Vogt, J. V., and Frischen, J.: Drought vulnerability indicators for global-scale drought risk assessments: global expert survey results report, Publications Office of the European Union, Luxembourg, 1-56, <https://doi.org/10.2760/738544>, 2019.
- 690 Mishra, A. K. and Singh, V. P.: Drought modeling - A review, *J. Hydrol.*, 403, 157–175, <https://doi.org/10.1016/j.jhydrol.2011.03.049>, 2011.
- Mishra, V., Tiwari, A. D., Aadhar, S., Shah, R., Xiao, M., Pai, D. S., and Lettenmaier, D.: Drought and famine in India, 1870–2016. *Geophysical Research Letters*, 46, 2075–2083, <https://doi.org/10.1029/2018GL081477>, 2019.
- MLUK (Ministerium für Landwirtschaft, Umwelt und Klimaschutz): Berechenbar: Dürrehilfen 2018 abgeschlossen - rund 72 Millionen Euro für notleidende Agrarbetriebe, Pressemappe, agrar presseportal, Potsdam, <https://www.agrar-presseportal.de/landwirtschaft/agrarpolitik/berechenbar-duerrehilfen-2018-abgeschlossen-rund-72-millionen-euro-fuer-notleidende-agrarbetriebe-27692.pdf>, 2019.
- 695 MLUK (Ministerium für Landwirtschaft, Umwelt und Klimaschutz): Agrarbericht Des Ministeriums Für Landwirtschaft, Umwelt Und Klimaschutz Des Landes Brandenburg, <https://agrarbericht.brandenburg.de/abo/de/start/agrarstruktur/naturliche-bedingungen/>, last accessed: 29 March 2024, 2022a.
- MLUK (Ministerium für Landwirtschaft, Umwelt und Klimaschutz): Benachteiligtes Gebiet, <https://geoportal.brandenburg.de/detailansichtdienst/render?view=gdiib&url=https://geoportal.brandenburg.de/gs-json/xml?fileid=f901b82c-54b7-4ef1-8365-2205da79c79b>, last accessed: 29 March 2024, 2022b.
- 700 MLUK (Ministerium für Landwirtschaft, Umwelt und Klimaschutz): Daten aus dem Agrarförderantrag, <https://geoportal.brandenburg.de/detailansichtdienst/render?view=gdiib&url=https://geoportal.brandenburg.de/gs-json/xml?fileid=996f8fd1-c662-4975-b680-3b611fcb5d1f>, last accessed: 29 March 2024, 2022c.
- 705 MLUK (Ministerium für Landwirtschaft, Umwelt und Klimaschutz): Strategie des Landes Brandenburg zur Anpassung an die Folgen des Klimawandels, 1-177, <https://mluk.brandenburg.de/sixcms/media.php/9/Klimaanpassungsstrategie-Brandenburg-LF.pdf>, 2023.
- Peichl, M., Thober, S., Samaniego, L., Hansjürgens, B., and Marx, A.: Machine-learning methods to assess the effects of a non-linear and damage spectrum taking into account soil moisture on and winter wheat yields in Germany, *Hydrol. Earth Syst. Sc.*, 25, 6523–6545, <https://doi.org/10.5194/hess-25-6523-2021>, 2021.
- 710 Raihan, M. J., Khan, M. AM., Kee, S.-H., and Al Nahid, A.: Detection of the chronic kidney disease using XGBoost classifier and explaining the influence of the attributes on the model using SHAP, *Scientific Reports*, 13 (1), <https://doi.org/10.1038/s41598-023-33525-0>, 2023
- Reinermann, S., Gessner, U., Asam, S., Kuenzer, C., and Dech, S.: The Effect of Droughts on Vegetation Condition in Germany: An Analysis Based on Two Decades of Satellite Earth Observation Time Series and Crop Yield Statistics, *Remote Sens.-Basel*, 11, 1–21, <https://doi.org/10.3390/rs11151783>, 2019.
- 715 Reisinger, A., Howden, M., Vera, C., Garschagen, M., Hurlbert, M., Kreibich, S., Mach, K. J., Mintenbeck, K., O'Neill, B., Pathak, M., Pedace, R., Pörtner, H.-O., Poloczanska, E., Corradi, M. R., Sillmann, J., van Aalst, M., Viner, D., Jones, R., Ruane, A. C., and Ranasinghe, R.: The concept of risk in



the IPCC Sixth Assessment Report: A Summary of Cross-Working Group Discussions, Intergovernmental Panel on Climate Change, Geneva, Switzerland, 15 pp., 2020.

720 Rossi, L., Wens, M., De Moel, H., Cotti, D., Sabino Siemons, A., Toreti, A., Maetens, W., Masante, D., van Loon, A., Hagenlocher, M., Rudari, R., Naumann, G., Meroni, M., Avanzi, F., Isabellon, M. and Barbosa, P.: European Drought Risk Atlas, Publications Office of the European Union, Luxembourg, 1-101, <https://doi.org/10.2760/33211>, 2023.

725 Şalap-Ayça, S. and Goto, E. A.: Beware the Rise of Models When They Are Wrong: A Look at Heat Vulnerability Modeling Through the Lens of Sensitivity (Short Paper), in: 12th International Conference on Geographic Information Science (GIScience 2023), Schloss-Dagstuhl-Leibniz Zentrum für Informatik, Germany, Leibniz International Proceedings in Informatics (LIPIcs), 277, 64:1-64:6, <https://doi.org/10.4230/LIPICS.GISCIENCE.2023.64>, 2023.

Samaniego, L., Kumar, R., and Zink, M.: Implications of Parameter Uncertainty on Soil Moisture Drought Analysis in Germany, *Journal of Hydrometeorology*, 14(1), 47-68, <https://doi.org/10.1175/JHM-D-12-075.1>, 2013

Santini, M., Noce, S., Antonelli, M., and Caporaso, L.: Complex drought patterns robustly explain global yield loss for major crops, *Sci. Rep.-UK*, 12, 1–17, <https://doi.org/10.1038/s41598-022-09611-0>, 2022.

730 Satoh, Y., Yoshimura, K., Pokhrel, Y., Kim, H., Shiogama, H., Yokohata, T., Hanasaki, N., Wada, Y., Burek, P., Byers, E., Müller Schmied, H., Gerten, D., Ostberg, S., Newland Gosling, S., Stanslas Boulange, J. E., and Oki, T.: The timing of unprecedented hydrological drought under climate change, *Nat. Commun.*, 13, 1–11, <https://doi.org/10.1038/s41467-022-30729-2>, 2022.

Schmitz, T. and Müller, D.: Digitale Karte der Bodenwertzahlen für Brandenburg, FORLand Technisches Papier 01, AgEcon Search, Germany, 1-13, <https://doi.org/10.22004/ag.econ.308812>, 2020.

735 Schymanski, E. L. and Schymanski, S. J.: Water science must be Open Science, *Nat. Water*, 1, 4-6, <https://doi.org/10.1038/s44221-022-00014-z>, 2023.

Shapley, L. S.: A value for n-person games, in: *Contributions to the Theory of Games, Volume II, Annals of Mathematics Studies*, edited by: Kuhn, H. and Tucker, A. W., Princeton University Press, Princeton, NJ, 307–317, <https://doi.org/10.1073/pnas.39.10.1095>, 1953.

Sieg, T., Vogel, K., Merz, B., and Kreibich, H.: Tree-based flood damage modeling of companies: Damage processes and model performance, *Water Resources Research*, 53 (7), 6050-6068, <https://doi.org/10.1002/2017wr020784>, 2017.

740 Söder, M., Berg-Mohnicke, M., Bittner, M., Ernst, S., Feike, T., Fröhlich, C., Golla, B., Jänicke, C., Jorzig, C., Leppelt, T., Liedtke, M., Möller, M., Nendel, C., Offermann, F., Riedesel, L., Romanova, V., Schmitt, J., Schulz, S., Seserman, D.-M., and Shawon, A. R.: Klimawandelbedingte Ertragsveränderungen und Flächennutzung (KlimErtrag), Thünen Working Paper 198, AgEcon Search, Johann Heinrich von Thünen-Institut, Braunschweig, Germany, 234 pp., <https://doi.org/10.22004/ag.econ.324625>, 2022.

745 Sodoge, J., Kuhlicke, C., and de Brito, M. M.: Automated spatio-temporal detection of drought impacts from newspaper articles using natural language processing and machine learning, *Weather and Climate Extremes*, 41, 1–9, <https://doi.org/10.1016/j.wace.2023.100574>, 2023.

Stahl, K., Kohn, I., Blauhut, V., Urquijo, J., de Stefano, L., Acácio, V., Dias, S., Stagge, J. H., Tallaksen, L. M., Kampragou, E., van Loon, A. F., Barker, L. J., Melsen, L. A., Bifulco, C., Musolino, D., de Carli, A., Massarutto, A., Assimacopoulos, D., and van Lanen, H. A. J.: Impacts of European drought events: insights from an international database of text-based reports, *Nat. Hazard Earth Sys.*, 16, 801–819, <https://doi.org/10.5194/nhess-16-801-2016>, 2016.

750 Statistische Ämter des Bundes und der Länder: Arbeitskräfte Und Deren Arbeitsleistung in Landwirtschaftlichen Betrieben, <https://www.regionalstatistik.de/genesis/online?operation=table&code=41141-08-02-4-B&byypass=true&levelindex=1&levelid=1657801475190#abreadcrumb>, last accessed: 29 March 2024, 2010.

Statistische Ämter des Bundes und der Länder: Bodenfläche Nach Art Der Tatsächlichen Nutzung, <https://www.regionalstatistik.de/genesis/online?operation=table&code=33111-01-02-4&byypass=true&levelindex=0&levelid=1659007696599#abreadcrumb>, last accessed: 29 March 2024, 2020a.

755 Statistische Ämter des Bundes und der Länder: Landwirtschaftliche Betriebe Insgesamt Sowie Mit Ökologischem Landbau Und Deren Landwirtschaftlich Genutzte Fläche (LF) Und Viehbestand, <https://www.regionalstatistik.de/genesis/online?operation=table&code=41141-04-02-4&byypass=true&levelindex=0&levelid=1660746074321#abreadcrumb>, last accessed: 29 March 2024, 2020b.



- 760 Statistische Ämter des Bundes und der Länder: Landwirtschaftliche Betriebe Mit Hofnachfolge,
<https://www.regionalstatistik.de/genesis//online?operation=table&code=41141-09-01-4&bypass=true&levelindex=0&levelid=1663767785139#abreadcrumb>, last accessed: 29 March 2024, 2020c.
- Statistische Ämter des Bundes und der Länder: Landwirtschaftliche Betriebe Nach Rechtsform Und Sozialökonomische Betriebstypen,
<https://www.regionalstatistik.de/genesis//online?operation=table&code=41141-07-01-4&bypass=true&levelindex=0&levelid=1661699406158#abreadcrumb>, last accessed: 29 March 2024, 2020d.
- 765 Statistische Ämter des Bundes und der Länder: Schulabgangsquote an Allgemeinbildenden Schulen VIII,
<https://www.bildungsmonitoring.de/bildung/online?operation=table&code=BBD15.1i&bypass=true&levelindex=0&levelid=1660318442403#abreadcrumb>, last accessed: August 12 2022, 2021.
- Statistische Ämter des Bundes und der Länder: Arbeitslosenquote Regionalatlas Deutschland,
<https://regionalatlas.statistikportal.de/?BL=DE&TCCode=AI008-1-5&ICode=AI0801>, last accessed: August 11, 2022, 2022.
- 770 Stephan, R., Terzi, S., Erfurt, M., Cocuccioni, S., Stahl, K., and Zebisch, M.: Assessing agriculture’s vulnerability to drought in European pre-Alpine regions, *Nat. Hazard Earth Sys.*, 23, 45–64, <https://doi.org/10.5194/nhess-23-45-2023>, 2023.
- Stephan, R., Stahl, K., and Dormann, C.F.: Drought impact prediction across time and space: limits and potentials of text reports. *Environmental Research Letters*, 18 (2023) 074004, <https://doi.org/10.1088/1748-9326/acd8da>, 2023b.
- Sutanto, S. J., van der Weert, M., Wanders, N., Blauhut, V., and van Lanen, H. A. J.: Moving from drought hazard to impact forecasts, *Nat. Commun.*, 10, 1–7, <https://doi.org/10.1038/s41467-019-12840-z>, 2019.
- 775 Tagesschau, <https://www.tagesschau.de/wirtschaft/unternehmen/landwirtschaft-erntebilanz-bauern-duerre-101.html>, last accessed: 06 March 2024, 2022.
- Tanguy, M., Eastman, M., Magee, E., Barker, L. J., Chitson, T., Ekkawatpanit, C., Goodwin, D., Hannaford, J., Holman, I., Pardthaisong, L., Parry, S., Rey Vicario, D., and Visessri, S.: Indicator-to-impact links to help improve agricultural drought preparedness in Thailand, *Nat. Hazards Earth Syst. Sci.*, 23, 2419–2441, <https://doi.org/10.5194/nhess-23-2419-2023>, 2023.
- 780 Tellman, B., Schank, C., Schwarz, B., Howe, P. D., and de Sherbinin, A.: Using Disaster Outcomes to Validate Components of Social Vulnerability to Floods: Flood Deaths and Property Damage across the USA, *Sustainability-Basel*, 12, 1–28, <https://doi.org/10.3390/su12156006>, 2020.
- Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index. *Journal of Climate*, 23 (7), 1696–1718, <https://doi.org/10.1175/2009jcli2909.1>, 2010.
- Wagenaar, D., de Jong, J., and Bouwer, L. M.: Multi-variable flood damage modelling with limited data using supervised learning approaches, *Nat. Hazards Earth Syst. Sci.*, 17(9), 1683–1696, [10.5194/nhess-17-1683-2017](https://doi.org/10.5194/nhess-17-1683-2017), 2017.
- 785 Walz, Y., Dall, K., Graw, V., Villagran de Leon, J.-C., Haas, S., Kussul, N. and Jordaan, A.: Understanding and reducing agricultural drought risk: Examples from South Africa and Ukraine, *Policy Report No. 3*, United Nations University – Institute for Environment and Human Security (UNU-EHS), Bonn, 1-29, 2018.
- Wilhite, D. A. and Glantz, M. H.: Understanding: the Drought Phenomenon: The Role of Definitions, *Water Int.*, 10, 111–120, <https://doi.org/10.1080/02508068508686328>, 1985.
- 790 Yang, C., Chen, M., and Yuan, Q.: The application of XGBoost and SHAP to examining the factors in freight truck-related crashes: An exploratory analysis, *Accid Anal Prev*, 158, 106153, <https://doi.org/10.1016/j.aap.2021.106153>, 2021
- Zhang H, Loaiciga H.A., and Sauter, T.: A Novel Fusion-Based Methodology for Drought Forecasting, *Remote Sensing*, 16(5), 828, <https://doi.org/10.3390/rs16050828>, 2024.
- 795 Zhou, R., Jin, J., Cui, Y., Ning, S., Bai, X., Zhang, L., Zhou, Y., Wu, C., and Tong, F.: Agricultural drought vulnerability assessment and diagnosis based on entropy fuzzy pattern recognition and subtraction set pair potential, *Alexandria Engineering Journal*, 61, 51–63, <https://doi.org/10.1016/j.aej.2021.04.090>, 2022.
- Zink, M., Samaniego, L., Kumar, R., Thober, S., Mai, J., Schäfer, D., and Marx, A.: The German drought monitor, *Environ. Res. Lett.*, 11, 1–9, <https://doi.org/10.1088/1748-9326/11/7/074002>, 2016.



Appendix A

Table A1. Average yields [dt/ha] per LBG 2010–2014, as used to estimate expected yields. Compiled from LELF (2016)

Crop	LBG-1	LBG-2	LBG-3	LBG-4	LBG-5
Winter wheat	77	65	50	38	23
Winter rye	63	55	43	35	25
Summer rye	37*	33*	25.8*	21*	15*
Winter barley	75	63	50	36	25
Oat	55	45	35	27	18
Winter triticale	66	60	48	37	23
Summer triticale	39.6*	36*	28.8*	22.2*	13.8*
Grain maize	90	80	70	60	50
Peas	35	30	25	20	NA
Lupines	NA	25	21	18	15
Potatoes	370	350	320	250	220
Potatoes (starch)	450	420	390	320	250
Sugar beet	650	620	580	NA	NA
Winter canola	43	38	32	25	20
Summer canola	23	18	14	11	NA
Sunflower	28	25	20	17	15

*Assumption, based on 60% of winter variety.

805

Table A2. Merging of the crop types between the three datasets IACS, yield reports, and average yields per LBG. Silage maize has been discarded later, and also for sugar beet we did not find prices 2021-2022

Crop	LBG average yields	IACS data	Yield reports	Assumptions made
Grain maize	Grain maize	Grain maize	Grain maize	-
Sunflower	Sunflower	Sunflower	Sunflower	-
Sugar beet	Sugar beet	Sugar beet	Sugar beet	-
Lupines	Lupines	Lupines	Lupines	-
Peas	Peas	Peas	Peas	-
Winter barley	Winter barley	Winter barley	Winter barley	-
Winter canola	Winter canola	Winter canola	Winter canola	-
Oat	Oat	Winter oat, Summer oat	Oat	Merge IACS to “Oat”
Potatoes	Potatoes Potatoes (starch)	Potatoes (various) Potatoes (starch)	Potatoes combined	Merge to “Potatoes”
Winter wheat	Winter wheat	Winter wheat	Winter wheat + spelt	Neglect spelt
Rye	Winter rye	Winter rye Summer rye	Rye + winter mix	Assume LBG values for summer rye as 60% of winter rye; Merge IACS to “Rye”;
Triticale	Winter triticale	Winter triticale Summer triticale	Triticale	Neglect winter mix Assume LBG values for summer triticale as 60% of winter triticale; Merge IACS to “Triticale”

810



815 **Appendix B**

Table B1. Model setups on field level ($y = \text{LST/NDVI-anom.}$). The indicators denoted with an ‘x’ are included in the respective setup. Performance initially assessed on 10% of the data to check the relative differences.

Setup	Crop type	SPEI Magnitude	SPEI Monthly	SMI Magnitude	SMI Monthly	Total Soil Magnitude	Vulnerability AZL, TWI, nFK	R ² (mean of 10 repetitions)
F1		x		x		x		0.09
F2	x	x		x		x	x	0.17
F3	x		x					0.20
F4	x				x			0.15
F5	x		x		x		x	0.26
F6	x*		x		x		x	0.25
								0.48**
F7	x*		x		x	x	x	0.25
								0.51**

*as categorical feature rather than one-hot encoded, ** re-trained on the full dataset

820

Table B2. Model setups on county level (target = relative empirical yield gap) using all available samples per setup (scores on holdout data). The indicators denoted with an ‘x’ are included in the respective setup.

Setup	Crop type	LST/NDVI	SPEI Magnitude	SPEI Monthly	SMI Magnitude	SMI Monthly	SMI Total	Vulnerability AZL	R ² (mean of 10 repetitions)
LK1	x	x							0.22
LK2	x		x						0.41
LK3	x		x		x		x	x	0.52
LK4	x			x				x	0.54
LK5	x					x		x	0.48
LK6	x			x		x		x	0.53
LK7	x			x		x	x	x	0.56
LK8	x	x		x		x	x	x	0.57
LK9	x*	x		x		x	x	x	0.53
LK9b	x*	x		x		x	x	x	0.40**

*as categorical feature rather than one-hot encoded, **trained only on samples where empirical yield gap > 0

825

830

835

840

845



Appendix C




Figure C1. SHAP summary plots for models trained only on (a) wheat (b) rye




Figure C2. SHAP values for all features of the best model trained on (a) all data, and (b) empirical yield gap > 0. Fig. 14 in the main paper only displays the first 15 of these.