

Contribution of blowing snow sublimation to the surface mass balance of Antarctica

Srinidhi Gadde ^{1,2} and Willem Jan van de Berg ¹

¹Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

²Faculty of Geo-Information and Earth Observation (ITC), Univeristy of Twente, Enschede, The Netherlands

Correspondence: s.nagaradagadde@utwente.nl, w.j.vandenberg@uu.nl

Abstract. Blowing snow sublimation is ~~an important a key~~ boundary layer process in polar regions and is the major ablation term in the surface mass balance (SMB) of the Antarctic ice sheet. ~~In this study, we update~~ This study updates the blowing snow model in the Regional Atmospheric Climate Model (RACMO) ~~version 2.3p3, to include, among other things, the effect of incorporating~~ blowing snow sublimation ~~in into~~ the prognostic equations for temperature and water vapour. These updates ~~are necessary to remove the undesired address~~ numerical artefacts in ~~this version's modelled blowing snow flux. Specifically, instead of a~~ the previous model version by replacing the uniformly discretised ice particle radius distribution ~~used in the previous version of the model~~, which limited the maximum ice particle radius to $\leq 50 \mu\text{m}$, ~~we use a non-uniformly discretised ice particle radius distribution to include all relevant radius between~~ with a non-uniform distribution covering radii from 2 to 300 μm without ~~any~~ additional computational overhead. The ~~updated model results are compared against the improved model is validated against~~ meteorological observations from site D47 in Adélie Land, East Antarctica. The updates ~~alleviate fix~~ the numerical artefacts ~~observed in the previous model results and successfully predict~~, successfully predicting the power-law variation of the blowing snow flux with wind speed while improving the prediction of ~~the magnitude of the blowing snow flux. Furthermore, we obtained an average blowing snow layer depth of $230 \pm 116 \text{ m}$ at the observation site D47, which matches the typical values obtained from satellite observations. A qualitative comparison of the blowing snow frequency from updated RACMO~~ its magnitude. Additionally, a qualitative comparison with CALIPSO satellite ~~observations data~~ shows that RACMO ~~successfully predicts accurately forecasts~~ the spatial pattern of monthly blowing snow frequencies. ~~We also compared updated RACMO results with CRYOWRF model results. While blowing snow sublimation dominates the total sublimation in RACMO, surface sublimation is the major contributor to total sublimation in CRYOWRF. Model results show that in the absence of~~ The model also yields an average blowing snow layer depth of $230 \pm 116 \text{ m}$ at D47, matching typical satellite observation values. Results reveal that, without blowing snow, sublimation in Antarctica ~~is limited to summer months (October–March) as there is negligible~~ mainly occurs in summer (October–March), with minimal surface sublimation in ~~Antarctic winter (April–September)~~ winter (April–September). Introducing the blowing snow model creates an additional sublimation mechanism ~~with its major contribution in the Antarctic winter months. For~~ primarily contributing in winter. From 2000 ~~—~~ 2012, ~~the model-integrated blowing snow sublimation is averaged~~ 175 \pm 7 Gt yr⁻¹, an increase of a 52% compared ~~to the results of the previous model version. The updated total sublimation, the sum of blowing snow sublimation increase from the previous version. Total sublimation, summing blowing snow~~ and surface sublimation, ~~is reached~~ 234 \pm 10 Gt yr⁻¹, which

is 47% higher compared to a simulation without a blowing snow model. Increased sublimation contributes This increase leads to a 1.2% reduction in the integrated SMB of the Antarctic ice sheet compared to the previous model results. In addition, we observe changes in the 's integrated SMB. Additionally, changes in sublimation in coastal and lower escarpment zones , indicating underscore the importance of the model updates to the climatology of blowing snow in Antarctica.
for Antarctic climatology.

1 Introduction

In the coastal regions of Antarctica, strong katabatic winds lift loose snow off the ground, causing drifting snow (e.g. Kodama et al. (1985)). When this snow rises further and is suspended in the atmospheric boundary layer, it is called blowing snow. This wind-driven transport can be categorised as drifting (< 1.8 m a.g.l) and blowing (> 1.8 m a.g.l) snow (Serreze and Barry, 2005, p. 54). It redistributes the snow on the surface of an ice sheet and can also give rise to black-ice areas, affecting the local surface energy balance (SEB) (van den Broeke and Bintanja, 1995). Furthermore, it is well known that the suspended snow particles are more prone to sublimation than surface snow (Schmidt, 1972; Bintanja, 2001). Therefore, drifting and blowing snow transport and sublimation are important factors contributing to Antarctica's surface mass balance (SMB), particularly in coastal regions (Bintanja, 1998). For brevity, from hereon, both drifting and blowing snow are combined and referred to as blowing snow.

Blowing snow is a significant contributor to the (local) SMB of the polar regions and plays a crucial role in the climate system of Antarctica. While there have been automatic weather station (AWS) observations of blowing snow-related processes from Antarctica (van den Broeke et al., 2004; Thiery et al., 2012; Barral et al., 2014; Amory, 2020), continent-wide estimates of blowing snow are difficult to obtain from such observations. Though continent-wide estimates derived from satellite-based products are available (Palm et al., 2017), they are restricted to optically thin cloud conditions and snow suspended in upper layers of the boundary layer (> 30 m a.g.l.) (Palm et al., 2011) and therefore are not suitable for estimates of near-surface blowing snow and its contribution to SMB. Hence, the continent-wide estimates can only be obtained by parameterising blowing snow processes and embedding these parameterisations in regional climate models (RCMs) (Bintanja, 1998; Déry and Yau, 2001; Lenaerts and van den Broeke, 2012; Amory et al., 2021; Toumelin et al., 2021). However, the representation of blowing snow in RCMs is challenging due to the complex and dynamic nature of the phenomenon involving multiple feedbacks with related processes such as snow precipitation and surface sublimation.

Including a blowing snow model in RCMs has been found to improve the SMB estimates in the regions where katabatic winds form (Mottram et al., 2021). Specifically, without modelling blowing snow processes, it is difficult to capture the spatial gradients in the sublimation of snow accurately (Agosta et al., 2019), which is particularly important in the escarpment regions of Antarctica. To improve our understanding of the Antarctic climate, it is crucial to accurately model the occurrence and impacts of blowing snow in RCMs. However, due to the coupled nature of blowing snow and the high sensitivity of the model to parameters, it is difficult to obtain a perfect agreement between observed and RCM estimates of blowing snow flux (Lenaerts




Figure 1. (a) Yearly average (2000-2012) 10-m wind speed (m s^{-1}). Location of observational site D47 in Adélie Land, East Antarctica, IL represents an interior location (71.1°S , 111.7°E), and dashed lines represent the latitude 82°S , North of which CALIPSO satellite data is available. (b) Variation of near surface blowing snow flux TR_{ds} ($\text{kg m}^{-1} \text{s}^{-1}$) vs 2-m wind speed (m s^{-1}). Solid and dashed lines represent the variation of observed and simulated (RACMO2.3p3) near-surface snow-drift fluxes, respectively.

60 et al., 2014; van Wessem et al., 2018; Amory et al., 2015, 2021).

The polar version of the regional atmospheric climate model (RACMO) (van Wessem et al., 2018; van Dalum et al., 2022) is coupled with a blowing snow scheme based on the PIEKTUK model (Déry and Yau, 2001; Lenaerts et al., 2012) to represent snow transport in polar regions. Evaluation of RACMO against snow particle counter (SPC) observations from Greenland showed that RACMO2.3p1 (hereafter Rp1) overestimated the snow particle transport (Lenaerts et al., 2014). Therefore, in
 65 RACMO2.3p2 (hereafter Rp2) (van Wessem et al., 2018), the linear saltation coefficient was subsequently halved to match the SPC observations from Greenland.

Recently, we evaluated blowing snow fluxes from RACMO2.3p3 (hereafter Rp3) against the SPC observational data at site D47 (location: 67.4°S , 138.7°E), Adélie Land, East Antarctica (Amory, 2020). Figure 1(a) shows the yearly (2000-2012) average 10-m wind speed obtained by Rp3 and the location of the observation site D47. Since the coastal regions of Antarctica
 70 witness very high wind speeds (Fig. 1(a)) and the concentration of blowing snow particles increases with the wind speed (Radok, 1977; Budd, 1966; Amory, 2020), the blowing snow transport (TR_{ds} $\text{kg m}^{-1} \text{s}^{-1}$) is expected to increase in a power-law fashion with velocity. However, Figure 1(b) shows that TR_{ds} from Rp3 does not show a rapid increase with velocity as expected. Evaluation of Rp3 blowing snow flux with observations also revealed that it consistently underestimates the magnitude of the observed flux. The evaluation shows the need to improve the blowing snow model in Rp3 and systematic
 75 comparison of blowing snow fluxes against observations to obtain reliable estimates of Antarctic SMB.

In this study, several updates to the blowing snow scheme in Rp3 are presented. The updates aim to improve the coupling

of the blowing snow processes with Rp3 atmospheric physics. Next, near-surface blowing snow fluxes obtained from Rp3 are compared against the observed fluxes from site D47, Adélie Land, East Antarctica (Amory, 2020). The observations from site D47 are particularly suitable for evaluations since the region experiences frequent blowing snow, and the observations employ 80 2nd generation FlowCapt™ sensors, which have been found to predict the blowing snow fluxes with reasonable accuracy (Amory, 2020). The details of RACMO and the modifications to the blowing snow scheme in Rp3 are presented in Section 2, and details of the observational site and available data are presented in Section 3. Blowing snow frequency and fluxes from the updated RACMO are evaluated against the observations in Section 4, followed by comparing results against version Rp3. Furthermore, we discuss the impact of the snow drift updates on the continent-wide estimates of SMB for Antarctica by 85 comparing the modelled SMB for 2000-2012 with a no-blowing snow case and model results from CRYOWRF (Gerber et al., 2023), followed by conclusions in Section 5.

2 Model descriptions

2.1 Regional Atmospheric Climate Model (RACMO)

RACMO is built on the semi-implicit semi-Lagrangian dynamics kernel of the numerical weather prediction model HIRLAM 90 (High-resolution limited area model; Undén et al. (2002)), version 5.0.3, with the European Center for Medium-Range Weather Forecasts (ECMWF) physics package, including both surface and atmospheric processes, from cycle 33r1 (ECMWF, 2009). The model assumes hydrostatic equilibrium, and the operational polar version, Rp2, has been verified to produce realistic results at the resolutions used in this study (van Wessem et al., 2015, 2016). This polar (p) version of RACMO2 includes a multilayer snow model that calculates the snow albedo evolution, melt, refreezing, percolation and run-off of meltwater 95 (Greuell and Konzelmann, 1994; Ettema et al., 2010; Kuipers Munneke et al., 2011). It also includes a blowing snow scheme based on the PIEKTUK model (Déry and Yau, 1999; Lenaerts et al., 2012).

In the newer version Rp3, the snow and ice albedo parameterisations were updated using Two-streAm Radiative TransfEr in Snow model (TARTES; Libois et al. (2013)) coupled with the Spectral-to-NarROWBand ALbedo (SNOWBAL) module, version 1.2 (van Dalum et al., 2019). Rp3 has produced results that compare well with both in-situ and remote sensing observations of 100 SMB of Antarctica (van Dalum et al., 2022). Rp2 and Rp3 are introduced in detail in Noël et al. (2018) and van Dalum et al. (2019), respectively. At the lateral boundaries, the simulations presented here are forced with ECMWF ERA5 reanalysis data (Hersbach et al., 2020) with an update interval of 3 hours.

2.2 Blowing snow model

In Rp3, we use the bulk (non-spectral) version of the PIEKTUK model (Déry and Yau, 1999), which employs an evolution 105 equation for the mixing ratio of blowing snow q_b (kg kg^{-1}) and an additional equation for the evolution of snow particle number concentration N , which is the double-moment version of the model (hereafter PIEKTUK-D, Déry and Yau (2001)). Here, we introduce only the essential features of the PIEKTUK model, and the additional details can be found in Déry and Yau




Figure 2. Schematic showing the blowing snow model levels, Rp3 model levels, and key processes involving blowing snow. Q_T represents the snow transport due to blowing snow. The figure shows that the model level of blowing snow is above the saltation height. In the schematic, q_b (mixing ratio) and N (number concentration) represent the boundary conditions for the blowing snow model calculated from q_{salt} and N_{salt} using the classical equation for suspended particle concentration.

(2001). PIEKTUK is an Inuktitut word for blowing snow (Déry et al., 1998). Figure 2 shows the blowing snow processes and the coupling between PIEKTUK-D and Rp3, presenting the important snow transport mechanisms over an ice sheet. When the friction velocity, a measure of the wind shear at the surface, exceeds the threshold friction velocity, the snow particles perform a downwind motion of a series of jumps or skips, a process called saltation. When the saltating snow particles get suspended in the boundary layer due to turbulent mixing, they form the blowing snow. In PIEKTUK-D, this transition from saltation to suspension, governed by different physical mechanisms, is assumed to happen at the elevation h_{salt} .

To calculate the sublimation and transport of blowing snow, the evolution equation for the blowing snow mixing ratio q_b (kg kg^{-1}) is written as:

$$\frac{\partial q_b}{\partial t} = \frac{\partial}{\partial z} \left(K_b \frac{\partial q_b}{\partial z} + v_b q_b \right) + S_b \quad (1)$$

where t (s) denotes time, z (m) the vertical coordinate, v_b (m s^{-1}) is the bulk terminal velocity, K_b ($\text{m}^2 \text{s}^{-1}$) represents turbulent eddy diffusivity for blowing snow, and S_b ($\text{kg kg}^{-1} \text{s}^{-1}$) the bulk sublimation rate. The lowest blowing snow model level is set to 0.1 m, at the top of the saltation layer at height h_{salt} . The boundary condition for solving Eq. (1) is given by relating the blowing snow mixing ratio q_b at the lowest model level with the mixing ratio at the top of the saltation layer, q_{salt} . There exist several empirical formulations for erosion of the snow particles in the saltation layer, q_{salt} (kg kg^{-1}). In Rp1, saltation flux was parameterised using Déry and Yau (1999),

$$q_{salt} = c_{salt} (1 - u_{thr}/U_{fml})^{2.59} / u_*, \quad (2)$$

in which c_{salt} is a constant, initially set to 0.385 and returned to 0.192 in Rp2 and Rp3, respectively. Furthermore, u_{thr} represents the threshold wind velocity (m s^{-1}), and U_{fml} represents wind speed at the first model level (m s^{-1}). The threshold wind velocity is defined by $u_{thr} = 9.43 + 0.18 T_{2m} + 0.0033 T_{2m}^2$ with T_{2m} in $^\circ\text{C}$ (Déry and Yau, 1999).

In Rp3, we update the saltation parameterisation Pomeroy (1989),

$$q_{\text{salt}} = \frac{e_{\text{salt}}}{gh_{\text{salt}}} (u_*^2 - u_{*t}^2), \quad (3)$$

where e_{salt} , the saltation efficiency, is set to $1/(3.25u_*)$, u_* (ms^{-1}) is the friction velocity, $h_{\text{salt}} = 0.08436u_*^{1.27}$ represents
 130 the thickness of the saltation layer (m) according to the relation of Pomeroy and Male (1992), $g = 9.81$ is the gravitational
 acceleration (ms^{-2}), and u_{*t} represents the threshold friction velocity (ms^{-1}).

The parameterisation of the threshold friction velocity in Eq. (3) is given by Gallée et al. (2001):

$$u_{*t} = u_{*t0} \exp\left(\frac{-n}{1-n} + \frac{n_0}{1-n_0}\right), \quad (4)$$

where, $n = (1 - \rho_s/\rho_i)$ is the snow porosity, $n_0 = (1 - \rho_0/\rho_i)$, with ρ_s as the actual mean snow density of the upper 5 cm,
 135 and ρ_i is the density of ice. n_0 is the porosity of fresh snow with $\rho_0 = 300 \text{ kg m}^{-3}$. The reference threshold friction velocity
 u_{*t0} is calculated based on the potential for snow erosion by the wind. u_{*t0} is characterised by a snow mobility index, $\text{Mo} =$
 $0.75d - 0.5s + 0.5$, where the variables d and s represent the snow grain dendricity, and sphericity, respectively. However,
 dendricity and sphericity are not modelled in RACMO. Therefore, we take $d = s = 0.5$, hence set Mo to 0.625. The threshold
 friction velocity based on (Gallée et al., 2001), depend on the snow mobility index which denotes the potential for snow erosion
 140 by the wind, with $\text{Mo} = 0.75d - 0.5s + 0.5$, where d and s represent dendricity and sphericity of fresh snow. Gallée et al. (2001)
 mention that the crystal shape of freshly fallen snow does not allow a large grain cohesion in the snowpack. Therefore, this
 allows relatively high snow mobility index Mo values for large d . Sintering is enhanced when the number of rounded shapes
 increases so that Mo decreases when s decreases. Explicitly modelling the snow mobility index requires solving prognostic
 equations for snow particle characteristics. Without sophisticated models for snow particle characteristics, the snow mobility
 145 index was set to 0.625 ($d = 0.5$ and $s = 0.5$) to match blowing snow frequency observations. A detailed discussion is available
 in Lenaerts et al. (2012). A parametric study of the snow particle characteristics is out of the scope of the present study.

Finally, u_{*t0} is defined by Gallée et al. (2001) as

$$u_{*t0} = \frac{\log(2.868) - \log(1 + \text{Mo})}{0.085} C_D^{0.5}, \quad (5)$$

where $C_D = u_*^2/U_{\text{fml}}^2$ represents the drag coefficient of momentum.

150 The governing equation for the evolution the concentration of particles (N) is

$$\frac{\partial N}{\partial t} = \frac{\partial}{\partial z} \left(K_N \frac{\partial N}{\partial z} + v_b N \right) + S_N. \quad (6)$$

Here, K_N ($\text{m}^2 \text{s}^{-2}$) is the eddy diffusivity for N , and S_N ($\text{m}^3 \text{s}^{-1}$) denotes the rate of change of particle numbers due to the
 sublimation process. The lower boundary condition for solving Eq. (6) is here also the particle concentration at the top of the
 saltation layer (N_{salt}) (Déry and Yau, 1999), which will be defined below.

155 In PIEKTUK-D, the bulk blowing snow mixing ratio q_b is related to N via the spectral number density $F(r)$, following Schmidt
 (1982):

$$q_b = \frac{4\pi\rho_{\text{ice}}}{3\rho} \int_0^\infty r^3 F(r) \text{d}r, \quad (7)$$

where the distribution of $F(r)$ follows two-parameter gamma distribution (Budd, 1966; Schmidt, 1982) by the relation:

$$F(r) = \frac{Nr^{(\alpha-1)} \exp^{-r/\beta}}{\beta^\alpha \Gamma(\alpha)}, \quad (8)$$

160 where, r represents the radius of ice particles, α (dimensionless) and β (m) the shape and scale parameters of the gamma distribution Γ . Substituting, Eq. (8) in (7), we obtain the particle number concentration N_{salt} at the saltation layer:

$$N_{\text{salt}} = \frac{3\rho q_{\text{salt}}\Gamma(\alpha)}{4\pi\rho_{\text{ice}}\Gamma(\alpha+3)\beta^3}, \quad (9)$$

with $\alpha = 4.0$, $\beta = 100/\alpha$ (μm), and density of ice $\rho_{\text{ice}} = 917 \text{ kg m}^{-3}$. Eq. (7) is discretised with the hypothesis that ice particle size follows a two-parameter gamma distribution, with particle size bins covering particles of radius 2 to 300 μm (Déry et al.,
165 1998).

Finally, in the blowing snow model, the mass change of an ice particle due to the blowing snow sublimation is given by the model of Thorpe and Mason (1966):

$$\frac{dm}{dt} = \left(2\pi r \sigma - \frac{Q_r}{KN_{\text{Nu}}T_a} \left[\frac{L_s}{R_v T_a} - 1 \right] \right) / \left(\frac{L_s}{KN_{\text{Nu}}T_a} \left[\frac{L_s}{R_v T_a} - 1 \right] + R_v \frac{T_a}{N_{\text{Sh}} D e_i} \right), \quad (10)$$

where σ (dimensionless and negative) is the water vapour deficit with respect to ice $(e - e_i)/e_i$, where e and e_i are the
170 vapour pressure and its value at saturation over ice. T_a is the ambient air temperature (K), K the thermal conductivity of air ($\text{W m}^{-1} \text{K}^{-1}$), L_s the latent heat of sublimation ($\text{J kg}^{-1} \text{K}^{-1}$), R_v the gas constant for water vapour, and D the molecular diffusivity of water vapour in air ($\text{m}^2 \text{s}^{-1}$), Q_r the net radiation transferred to the ice particle (W), and N_{Nu} and N_{Sh} being the Nusselt and Sherwood numbers.

2.3 Major changes to blowing snow model in RACMO

175 Six major updates in the implementation of PIEKTUK-D in Rp3 are summarised below :

1. In Rp3, uniformly discretised 12-particle size bins were used, with a constant particle bin size $\Delta r = 4 \mu\text{m}$. Therefore, size bins with a mean particle radius greater than 50 μm were excluded, which caused the unexpected variation of TR_{ds} observed in Fig. 1(b). To solve the issue, we use a grid stretching technique similar to Direct Numerical Simulations (DNS) of channel flows to obtain non-uniform distribution with smooth stretching following a tangent hyperbolic function (Vinokur, 1983). We now use 16-particle size bins with varying Δr to include all relevant particle size classes with
180 particles of mean radius for each bin from 2 to 300 μm , while keeping the computational overhead the same as before. Déry et al. (1998) report convincing results by including particles of mean radius for each bin from 2 to 254 μm .
2. Previously, in the blowing snow model of Rp3, 32 vertical levels equidistant on a logarithmic scale were used. The blowing snow model was not fully coupled to the boundary-layer model as the blowing snow grid levels did not match
185 the model atmospheric levels. Specifically, instead of the actual velocity profiles, temperature and velocity profiles were reconstructed using logarithmic relations from the first model atmospheric level. In addition, the friction velocity (u_*)

was recalculated in the blowing-snow model, assuming near-neutral conditions. These inconsistencies have now been resolved, and actual velocity and temperature profiles and friction velocities from the boundary layer model are used, which constitutes another major improvement. We have reduced the vertical levels to 16 to reduce computational expenses, with eight logarithmically varying levels up to the lowest model level (dashed lines in Figure 2). Furthermore, above the lowest atmospheric level, the PIEKTUK-D model levels coincide with the model atmospheric levels, and this facilitates easier coupling of blowing snow sublimation as tendencies in the prognostic equations.

190

195

200

205

210

215

3. We found that PIEKTUK-D, when coupled to Rp3, is highly sensitive to the model time step. In PIEKTUK-D, the integrated blowing snow flux quickly reaches equilibrium, and it depends on the time step used to solve the evolution equations. While Déry and Yau (1999) specify a model time step of 2 seconds for PIEKTUK-D; in Rp3, the model time step was of the order of 300 – 600 seconds. This time step was too large to predict the drift fluxes reliably. To overcome this, we introduce sub-stepping in the blowing snow model. We ran the model with different Δt , with different time step sizes. We found large difference between the values of integrated blowing snow flux for $\Delta t = 600, 300, 100, 50, 20, 10,$ and 5 seconds. For $\Delta t = 10$ and $\Delta t = 5$, the magnitude of blowing snow flux was nearly the same, so we choose $\Delta t = 10$ s. Furthermore, the model quickly reaches a steady state in 5 sub-steps. Therefore, we use five sub-steps with a time step of 10 seconds and the fluxes from the last sub-step are taken as the representative flux for the full Rp3 model step.
4. In the original PIEKTUK model implementation by Déry and Yau (1999), the blowing snow mixing ratios are reset to zero only if the friction velocity is lower than the threshold friction velocity in two consecutive time steps, providing a realistic initial approximation of blowing snow quantities in each time step. However, previously in Rp3, N and q_b were reset to zero after every model time step, though in reality, the blowing snow events last for hours. Resetting the flux to zero is unrealistic and calls for a proper initialisation of the variables. Therefore, we now initialise N and q_b from the previous time step if two consecutive time steps satisfy the condition $u_* > u_{*t}$; otherwise, the values are reset to zero, indicating the end of the blowing snow event.
5. In Rp3, the bulk sublimation rate S_b was used to calculate an integrated blowing snow sublimation flux, and this integrated moisture flux was added to the surface. While this approach works reasonably in obtaining SMB estimates, it is not realistic since it limits the effect of blowing snow sublimation to the surface. To rectify this error in representation, we now add blowing snow sublimation rate ($-S_b$) and latent heat due to blowing snow ($L_s S_b / c_p$) as tendencies to the prognostic equations of atmospheric water vapour and temperature, respectively.
6. In Rp3, snowdrift was modelled if $u_* > u_{*t}$ and Eq. (2) was used to estimate the saltation flux. This parameterisation caused sharp variations in the saltation flux in Rp3 and was not optimal. Therefore, the saltation flux is now derived with Eq. (3), which produces smooth variations of q_{salt} . Furthermore, Eq. (3) is also used in the MAR model to parameterise saltation flux (Amory et al., 2021). Finally, the formula to derive the vapour saturation pressure to ice (e_i) in Eq. (10) has been updated to the AERKi formula (Alduchov and Eskridge, 1996; CY45R1—Part IV, 2018), as this formula is used in IFS code in which the blowing snow module is embedded.

3.1 In situ snowdrift observations

The in situ observations used for evaluation are presented and discussed in detail by Amory (2020) and Amory et al. (2020a); here, we summarise the key information. The observational site D47 (location: 67.4°S, 138.7°E, Fig. 1(a)), is located at an elevation of 1560 m, and at a distance of 105 km from the shore. Due to its topographical situation, the site experiences strong katabatic winds with a strong directional consistency Amory (2020). Due to the high surface winds, the site experiences frequent blowing snow events and is ideally suited for evaluating RACMO results. For evaluation, observations of near-surface quantities such as 2-m wind speed, temperature, and air relative humidity are used, complemented with half-hourly drifting-snow transport fluxes. These observations are available for 2010–2012 with half-hourly temporal resolution. The drifting-snow transport fluxes are measured with second-generation FlowCapt™ sensors. The sensors convert the acoustic vibration caused by blowing snow particles into integrated snow mass flux. The equipment consists of two 1-m length acoustic tubes, superimposed vertically to measure snow flux in the first 2 m above the ground.

The blowing snow scheme in Rp3 has multiple levels, with the lowest vertical level set at 0.1 m. For comparison with observations, we obtain an average, vertically integrated, blowing snow flux, $Q_{T,RACMO}$ ($\text{kg m}^{-2} \text{s}^{-1}$), from the lowest model level up to 2-m height. Following Amory et al. (2021), since there are two acoustic tubes for measurement, we combine snow mass flux from both the tubes into an average, near-surface, mass flux $Q_{T,OBS}$ ($\text{kg m}^{-2} \text{s}^{-1}$), as :

$$Q_{T,OBS} = \frac{Q_{T,1}h_1 + Q_{T,2}h_2}{h_1 + h_2}, \quad (11)$$

where, $Q_{T,i}$ is the observed snow mass flux integrated over the exposed length of h_i of the corresponding 2G-FlowCapt™ sensor. The height of the sensor at D47 is 2.8 m. However, Amory (2020) mention that, due to harsh weather conditions at D47, it was difficult to reset the height of the sensors owing to the elevation changes due to snow. As a result, by late December 2012, the measurement heights decreased from their initial values to 1.5 m for wind speed and direction and 0.9 m for temperature and relative humidity. Therefore, we compare the instantaneous fluxes for the year 2011. The data is made available on Zenodo as quantities at 2-m height. Since the first atmospheric level in RACMO (approximately 8–10 m) is above this height, we obtain the 2-m wind speed using the Monin-Obukhov similarity theory.

3.2 Satellite data for evaluating monthly blowing snow frequency

We compare model results with lidar data from CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation, Palm et al. (2017)), which measures blowing snow quantities for the Antarctic Ice Sheet north of 82 °S (Palm et al., 2017, 2018). These satellite observations include only those blowing snow layers deeper than 30 m and only those events without clouds. Making an accurate one-to-one comparison of the model results with the satellite observations requires filtering of the model results to layers deeper than 30 m and for cases with no or optically thin cloud conditions, of which the former is not possible with the data exported from the current RACMO simulations. Therefore, we use the satellite observations to look at the seasonal patterns and only qualitatively compare the model results.




Figure 3. Comparison of simulated instantaneous near-surface blowing snow flux Q_T ($\text{kgm}^{-2}\text{s}^{-1}$) with observations for the year 2011: (a) Rp3, (b) RpNew (c) Variation of near-surface blowing snow flux Q_T with 2-m wind speed U_{2m} (ms^{-1}). Open circular, filled red diamond, and filled circular markers represent data from RpNew, Rp3, and AWS data, respectively. Observed and simulated blowing snow fluxes $\text{kgm}^{-2}\text{s}^{-1}$ (d) Rp3, and (e) RpNew. Solid lines represent the 1:1 line and the dashed lines represent the best-fit line. The colours represent the normalised point density from low (0.0, black) to high (1.0, red).

4 Results and discussion

Three Rp3 simulations for 2000–2012, forced by ERA5 reanalysis data, were run for the evaluation presented here. The first one, hereon referred to as RpNew, employed all updates listed in Section 2.3. A second simulation was carried out with the blowing snow scheme switched off, hereon referred to as NO-DRIFT, to study the effects of blowing snow compared to the no-blowing snow scenario. Finally, a simulation with the original blowing snow code of Rp3 has been carried out to compare the change in SMB estimates and related quantities.

4.1 Model evaluation with observations at site D47

4.1.1 Blowing snow flux and near-surface relative humidity

260 Figure 3(a) presents the instantaneous blowing snow mass flux obtained from Rp3 compared with the observations for the year 2011. As can be seen in the figure Rp3 fails to predict the peaks of blowing snow magnitude when compared with observations and as such does not reliably predict the magnitude of the blowing snow. In Rp3, the linear saltation coefficient c_{salt} (Eq. (2)) was reduced (van Wessem et al., 2018) which resulted in the low, limited snow transport flux seen in Figure 3(a). As halving c_{salt} roughly led to halving the snow drift flux, the RACMO versions preceding version 2.3p2, with doubled c_{salt} , overestimated 265 Q_T for most of the time (not shown).

Figure 3(b) presents the instantaneous blowing snow mass flux obtained with the RpNew. We observe that RpNew works well in predicting the magnitude of the blowing snow flux. Specifically, the magnitude of the blowing snow flux matches the observations reasonably well in the Antarctic winter (April–September). However, it is underestimated in the Antarctic summer (October–March). The underestimation might be related to the amount of loose snow available at the surface, possibly due 270 to inaccuracies of the modelled surface snow compaction in summer or the direct interaction between precipitation and snow drift, which Rp3 does not resolve. As we found no clear cause for the underestimation of snow drift during summer, further study is necessary to uncover the seasonal differences in the blowing snow flux.

Figure 3(c) presents the variation of blowing snow mass flux with the near-surface wind speed. Flux from Rp3 fails to produce power-law variation with the wind speed; however, RpNew successfully predicts it. The primary reason for this improvement is the non-uniform ice particle radius distribution, allowing us to include all relevant ice particles in the range 275 between 2 to 300 μm . Coupled with the better coupling with RACMO prognostic variables and sub-stepping, the behaviour of the flux follows the expected power-law variation seen in Figure 3(c).

In Figure 3(d) and (e), we present the comparison of simulated near-surface blowing snow mass flux with observed flux for Rp3 and RpNew, respectively. Simulated flux from Rp3 has a positive bias, with a very low R^2 (p-value < 0.01) indicating that 280 Rp3 fails to capture the variability in the blowing snow flux observations. The predictions with Rp3 also have a higher RMSE of $0.035 \text{ kg m}^{-2} \text{ s}^{-1}$. Also, it is apparent from Figure 3(d) that Rp3 fails to predict the blowing snow fluxes reliably when compared with the observations. In contrast, with RpNew, we have a reasonable agreement between the observed and simulated fluxes (Fig. 3(e)) with $R^2 = 0.56$ (p-value < 0.01). The agreement indicates that the changes introduced significantly improve its ability to predict the blowing snow fluxes. Though the predictions are considerably improved compared to the observations, 285 both Rp3 and RpNew underestimate the blowing snow fluxes. The underestimation is mostly due to the underestimation of velocities reported in Table A1 and the model sensitivity to the chosen parameters. Since the snow transport flux varies in a power-law fashion with the wind speed (Radok, 1977; Budd, 1966; Amory, 2020), the flux is highly sensitive to the wind-speed predictions; even a slight underestimation in the velocity introduces a significant difference in the blowing snow mass flux.

Though RpNew results show the desired behaviour, it fails to capture the spread in the observational data (Fig. 3(c)). Through 290 sensitivity analysis of the data, we found that the spread in the data depends on the modelling choices made, e.g. the parameter α in two-parameter gamma distribution (Eq. 8) and the threshold friction velocity. Budd (1966) and Schmidt (1982) report




Figure 4. Density scatter plots of observed and simulated near-surface relative humidity w.r.t ice at site D47, for cases (a) NO-DRIFT, (b) Rp3, (c) RpNew. Solid lines represent the 1:1 line and the dashed lines represent the best-fit line. The colours represent the normalised point density from low (0.0, black) to high (1.0, red).

that the distribution of ice particle diameters follows a two-parameter gamma function that varies with height from the ground, with α value varying between 2 and 14. However, for simplified implementation, following Déry and Yau (2002), we used a constant $\alpha = 4$, which does not vary with height; this influences the modelled snow mass flux at different heights. Furthermore, the use of constant snow grain properties in the calculation of snow mobility index used in the calculation of the threshold friction velocity (Eq. (4)) can influence the spread in the data. Irrespective of these simplifications, RpNew reasonably accurately predicts the blowing snow fluxes.

Improving the blowing snow prediction is expected to improve the near-surface humidity predictions. Figure 4(a), (b), and (c) present a comparison of observed relative humidity with respect to ice against the simulated relative humidity for the three experiments. Figure 4(a) shows that the NO-DRIFT case shows a negative bias in the moisture, with low $R^2 = 0.07$ (p-value < 0.01) and a high error indicated by an RMSE of 18.84%. With Rp3, the results are slightly improved with a lower negative bias and a higher $R^2 = 0.35$ (p-value < 0.01). However, Figure 4(c) shows that with RpNew, the modelled results show an improved correlation with the observations ($R^2 = 0.49$, p-value < 0.01). Though the data has a large spread, the RMSE is 6.6%, and the figure shows an improved match between the observed and simulated data. It is evident from Figure 4(a), (b), and (c) that the updates improve the moisture prediction when compared with the observations.

4.1.2 Blowing snow events at site D47

To quantify the ability to model the blowing snow events, we follow Amory et al. (2017, 2021) and classify blowing snow events as the occurrences when the blowing snow mass flux is greater than $10^{-3} \text{ kg m}^{-2} \text{ s}^{-1}$. Subsequently, we create confusion matrices comparing the blowing snow events from observed and simulated data. The diagonal entries in the confusion matrix represent the events correctly predicted by the simulations, and the off-diagonal entries represent the remaining events.

Table 1. Confusion matrix presenting the comparison between observed and simulated blowing snow events. Table (a) considers all snow drift events, Table (b) on strong snow drift events. In (a), DRIFT represents the events where $Q_T > 10^{-3} \text{ kg m}^{-2} \text{ s}^{-1}$ and NO-DRIFT represents the remaining events. In the tables below, first number in each cell represents events from RpNew and the second number from Rp3.

(a) Blowing snow events from calculated as
 $Q_T > 10^{-3} \text{ kg m}^{-2} \text{ s}^{-1}$.

	SIM	NO-DRIFT	DRIFT
OBS			
NO-DRIFT		18%,16%	2%,4%
DRIFT		26%,17%	54%,63%

(b) High blowing snow flux events calculated as
 $Q_T > 0.05 \text{ kg m}^{-2} \text{ s}^{-1}$.

	SIM	$Q_T \leq 0.05$	$Q_T > 0.05$
OBS			
$Q_T \leq 0.05$		84%, 86%	2%, 0%
$Q_T > 0.05$		9%, 14%	5%, 0%

Table 1(a) represents the confusion matrix presenting the percentage of blowing snow events observed and simulated by RpNew and Rp3, respectively. In Table 1(a), we see that out of the total observations, there are 80% of observed blowing snow events, and RpNew manages to predict 54% of these events. In contrast, Rp3 predicts 63% of the total blowing snow events. We calculate the blowing snow frequency as the ratio of correctly simulated blowing snow events and the total number of observed blowing snow events. For RpNew, we obtain a blowing snow frequency of 0.68, and Rp3 has a blowing snow frequency of 0.79. Rp3 performs comparatively better in predicting the blowing snow events as the threshold friction velocity calculated in Rp3 is lower than RpNew. As mentioned previously, in Rp3, u_* was recalculated in every step with a simple logarithmic-law assumption, which reduced the threshold friction velocity used in the model to trigger a blowing snow event. Since the assumption was not correct, we changed it and used the friction velocity calculated from the physics module. With RpNew, the model's friction velocity is consistently higher than Rp3 (not shown here). This can be tuned in future versions to match the observations better. Furthermore, we do not observe any seasonality in the underestimation of blowing snow events, we observe only marginal differences in the blowing snow frequency of RpNew over Antarctica summer (October–March) and winter (April–September) months. Clearly, Rp3 does a better job in identifying the blowing snow events, however Fig. 3(a) shows that it does not capture any peaks in the blowing snow fluxes.

To evaluate the performance of RpNew in identifying the higher magnitude blowing snow fluxes, we create another confusion matrix where we compare the blowing snow events with blowing snow mass flux $Q_T > 0.05 \text{ kg m}^{-2} \text{ s}^{-1}$. Table 1(b) presents a comparison of the observed and simulated blowing snow fluxes for events with $Q_T > 0.05 \text{ kg m}^{-2} \text{ s}^{-1}$. The tables show that 14% of the observed events account for events with high blowing snow mass flux. While RpNew captures 5% out of the 14% high blowing snow events, Rp3 does not capture any of these events. Therefore, RpNew shows a marked improvement in predicting blowing snow peaks compared to Rp3. However, RpNew still underestimates the number of strong blowing snow events which is closely related to the underestimation of the wind speed (Table A1) in RpNew and Rp3. The results show that RpNew provides reasonable estimates of low- and high-magnitude blowing snow events while future improvements are needed.




Figure 5. (a) Blowing snow sublimation rate for the year 2011 in mm s^{-1} , inset shows the histogram of blowing snow layer depth (m). (b) Yearly averaged of blowing snow sublimation rate with height (mm s^{-1}).

335 4.1.3 Blowing snow sublimation at site D47

Figure 5(a) shows the modelled instantaneous profiles of blowing snow sublimation rate for 2011 at site D47. In the Antarctic winter (April–September), deep blowing snow layers are modelled, with a typical range of blowing snow layer heights and snow sublimation between 100 and 500 m. A shallower blowing snow layer is observed in Antarctic summer (October–March). The figure shows multiple events with continuous blowing snow storms in winter, indicating a significant contribution of blowing snow to Antarctic sublimation. Although sublimation over a thick layer coincides with blowing snow events (Fig. 3b), we do not see a direct relation between the near-surface snow drift flux and the intensity or total magnitude of blowing snow sublimation. This shows the necessity to explicitly couple the blowing snow model to the atmospheric model layers, as the modelled temperatures, humidities, and wind speeds of the lowermost model level are unlikely representative of the whole boundary layer.

345 Figure 5(b) presents the yearly averaged blowing snow sublimation rate profile for 2011 at site D47. The average blowing snow layer depth is 230 ± 116 m. As the air is saturated at the surface, the sublimation at the surface is negligible, with sublimation increasing away from the ground and maximum sublimation above. Déry and Yau (2002) and Toumelin et al. (2021) have reported a similar variation of the blowing snow sublimation. It is worth noting here that both the drifting snow concentration and horizontally drifting snow transport are peaking even close to the ground (not shown). As depicted in Figure 350 5(a), blowing snow sublimation starts below the model’s first atmospheric level (approximately 8–10 m). Previously in Rp3, this vertical profile of sublimation was not represented as the blowing snow sublimation was added to the surface.

Based on lidar data from CALIPSO observations, Palm et al. (2017) report for the Antarctic Ice Sheet north of 82°S an average snow layer depth of 120 m, with typical blowing snow layers of 200 m all along the coastal katabatic wind regions (see Fig. 5 in Palm et al. (2018)). For the site D47, RpNew shows a similar mean layer depth of 230 ± 116 m, and a similar typical range (Inset Fig. 5(a)). This analysis shows that RpNew satisfactorily reproduces all the necessary features of the blowing snow sublimation and can be used to obtain continent-wide estimates. However, it is worth mentioning that total blowing snow sublimation is sensitive to horizontal resolution. At the 27 km resolution employed in the study, strong spatial gradients near the coast would not be accurately captured. Subsequently, the impact of blowing snow on sublimation and horizontal transport of mass can be underestimated.




Figure 6. Blowing snow frequency visualised to provide a qualitative comparison with satellite measurements of Palm et al. (2018). The panels show average blowing snow frequency over the decade 2001–2010 and use the same colour scale as Palm et al. (2018). A colour blind friendly version of this figure is given in the appendix (Figure B1).

360 4.2 Continental blowing snow frequency

Figure 6 gives the monthly variation of mean blowing snow frequency [from RpNew](#) over Antarctica for the decade 2001–2010. Blowing snow frequency is obtained by calculating all the blowing snow events with the blowing snow mixing ratio $q_b > 10^{-6} \text{ kg kg}^{-1}$. The cutoff q_b , the limits, and the colourmap in Figure 6 are chosen to facilitate a qualitative comparison with the satellite observations presented in Figure 3 in Palm et al. (2018). We observe that the monthly blowing snow frequency largely follows the seasonal trend in the surface wind patterns over Antarctica, with high-frequency blowing snow in winter compared to summer. Despite the differences between the two approaches, the simulated blowing snow frequency is qualitatively similar to that obtained from the CALIPSO satellite observations (Palm et al., 2017, 2018). The results show a persistent blowing snow hotspot in East Antarctica near Adélie Land, observed in satellite observations and our simulations. We can also infer that the satellite observations slightly underestimate the frequencies compared with the simulations due to the reasons mentioned in

365
370 Section 3.2.

Our results are also qualitatively similar to the simulations with the CRYOWRF model carried out by Gerber et al. (2023), although the simulated periods are different. [Specifically, Gerber et al. \(2023\) report a zone of strongest blowing snow along the coast of East Antarctica with highest values of blowing snow frequency slightly inland. Zones with reduced blowing snow are found over the Amery ice shelf, toward the western Queen Maud land. RpNew shows similar qualitative trends.](#) Most of the blowing snow hotspots observed in our simulations also correspond to the ‘wind glaze’ areas in East Antarctica reported by Scambos et al. (2012). Scarchilli et al. (2010) report blowing snow frequencies of 80% at the wind convergence zone of Terra Nova Bay (East Antarctica); we observe approximately 80–90% blowing snow frequency in the area during the Antarctic winter months.

4.3 Continent-wide estimates of blowing snow climate over Antarctica

Figure 7 presents the updated continent-wide estimates of the blowing snow climate of Antarctica by comparing the yearly average (2000-2012) quantities of RpNew and NO-DRIFT simulations. Similar to previous model results (Lenaerts and van den Broeke (2012)), we observe negligible blowing snow sublimation (Figure 7(a)) in the interior parts of Antarctica with maximum sublimation towards the coast. Model results show blowing snow sublimation hotspots ($SU_{ds} > 100 \text{ mm w.e. yr}^{-1}$) in George V Land, Adélie Land, Wilkes Land, and Queen Mary Land in Eastern Antarctica with non-zero sublimation all along the coast of Antarctica. RACMO shows negligible sublimation over Dome Fuji, Dome Argus, and Dome C, which form the interior parts of Eastern Antarctica, due to the lower wind speed and low temperatures in these regions (Fig. 1(a)). Similarly, we observe negligible sublimation over Ronne and Ross ice shelves, which also experience low wind speeds. This shows that blowing snow sublimation is mostly limited to the katabatic wind regions of Antarctica. Maximum blowing snow sublimation of $335 \pm 30 \text{ mm w.e. yr}^{-1}$ occurs in Adélie Land at the location 66.9°S , 130.4°E . Palm et al. (2017) based on the CALIPSO lidar observations report a maximum blowing snow sublimation of $250 \pm 125 \text{ mm w.e. yr}^{-1}$ near the coast between longitudes 140 and 150°E . A slight shift in the location of the maximum blowing snow sublimation between the satellite and model results can be related to the fact that CALIPSO observes moderate and strong events without thick cloud cover, while Figure 7 displays all snowdrift events (see Section 3.2). Both spatial distribution and the magnitude of blowing snow sublimation from RpNew match reasonably well with CALIPSO observations of Palm et al. (2017).

Figure 7(b) provides the difference between the 2-meter temperature for the RpNew and NO-DRIFT cases. The figure shows that blowing snow sublimation reduces the near-surface temperature. At blowing snow sublimation hotspots, we observe a cooling of $0.1 - 0.3 \text{ K}$, with negligible change in the temperature over most of interior Antarctica. It is worth mentioning here that with Rp3, we observed a slight warm bias compared to the case with NO-DRIFT (not shown); this shows that the coupling was previously incorrect. The results have appreciably improved with RpNew. However, the overall effect of blowing snow sublimation on the yearly average near-surface temperature in Antarctica seems marginal, similar to previous model results.

Higher sublimation due to blowing snow in RpNew leads to higher near-surface relative humidity (Fig. 7(c)) when compared to NO-DRIFT simulations. We observe higher relative humidity along the Antarctic coast with a maximum of 10% in the coastal George V Land and Adélie Land. This increase in relative humidity is higher than what was previously observed with Rp3. Similar to sublimation, blowing snow transport TR_{ds} ($\text{kg m}^{-1} \text{ yr}^{-1}$) (Fig. 7(d)) is negligible over interior Antarctica.

405 We observe a strong blowing snow transport near coastal George V Land with maximum transport of $9 \times 10^6 \text{ kg m}^{-1} \text{ yr}^{-1}$. Along the rest of the Antarctic coast, blowing snow transport is approximately $2 \times 10^6 - 3 \times 10^6 \text{ kg m}^{-1} \text{ yr}^{-1}$. Blowing snow erosion ER_{ds} (mm w.e. yr^{-1}) (Fig. 7(e)) which is a contributor to Antarctic SMB, shows complex convergence and divergence patterns all along the Antarctic coast. Similar to Bromwich et al. (2004) and Lenaerts and van den Broeke (2012), we observe large blowing snow divergence near escarpment areas with significant katabatic wind acceleration. Furthermore, 410 areas with blowing snow convergence are near blowing snow divergence, which indicates that blowing snow is important for redistributing the precipitation in the coastal areas of Antarctica. However, the magnitude of ER_{ds} is not significant enough for a major contribution to SMB, as only the snow blown off Antarctica counts for the integrated SMB.

Total sublimation SU_{total} (mm w.e. yr^{-1}), the sum of blowing snow and surface sublimation ($SU_{ds} + SU_s$), follows the spatial distribution of blowing snow sublimation. Maximum total sublimation of $396 \text{ mm w.e. yr}^{-1}$ is observed at the same 415 location as the maximum blowing snow sublimation, indicating the leading contribution of blowing snow sublimation to the total sublimation. Total sublimation is higher in the RpNew simulations when compared to NO-DRIFT simulations (Figure 7(f)); in the regions near Adélie Land, the difference in total sublimation is as high as $200 \text{ mm w.e. yr}^{-1}$.

4.4 Seasonal variation of integrated sublimation

In Figure 8 we present the different components of sublimation from RpNew, NO-DRIFT, Rp3, and CRYOWRF (Gerber et al., 420 2023). Monthly contribution to the yearly average integrated blowing snow sublimation (Fig. 8(a)) shows that the blowing snow sublimation in the Antarctic summer (October–March) is lower than in Antarctic winter (April–September) due to higher temperatures and summer snow densities, making it difficult for the snow to lift off from the ground. Blowing snow sublimation SU_{ds} increases with the onset of winter and remains relatively constant over winter with an approximate contribution of $15 - 20 \text{ Gt mo}^{-1}$ in winter. [For RpNew blowing snow sublimation contributes 108 Gt during winter out of the total sublimation of 175 Gt yr⁻¹. This amounts to 62% of the total blowing snow sublimation.](#) 425 Constant blowing snow indicates that blowing snow sublimation is a major contributor to total sublimation in winter. Due to the updates made in RpNew, the blowing snow sublimation has nearly doubled throughout the winter compared to Rp3. CRYOWRF produces blowing sublimation which is comparable in magnitudes to Rp3. Surface sublimation SU_s dominates the sublimation in summer due to higher temperatures (Fig. 8(b)) and reaches a relatively constant value in winter. We observe that surface sublimation in Antarctic summer with 430 CRYOWRF is nearly 1.5 times the surface sublimation observed with RpNew, while winter sublimation is comparable. SU_s is negligible in Antarctic winter and it is nearly zero. Interestingly, with the introduction of blowing snow sublimation in Rp3 and RpNew, we observed negative surface sublimation, indicating the deposition of water vapour onto the snow surface in winter. This deposition agrees with the measurements of King et al. (1996), who measured small, downward water vapour fluxes in the winter of 1991 at Halley station, East Antarctica. A similar seasonal cycle in the surface sublimation with a negative surface 435 sublimation in winter has been reported by King et al. (2001). While the blowing snow sublimation is increased in RpNew, the negative surface sublimation also increases, balancing the net change in total sublimation. The deposition follows the same spatial and seasonal pattern as the blowing snow sublimation. Since the condensation is directly proportional to the difference between the vapour pressure of water at the surface and above the surface, with RpNew, which has better coupling with the




Figure 7. Yearly mean (2000-2012): (a) Blowing snow sublimation SU_{ds} in mm water equivalent, (b) Difference in near-surface temperature, ΔT_{2m} between RpNew and NO-DRIFT simulations, (c) Difference in relative humidity ΔRH_{2m} , (d) Blowing snow flux $[kgm^{-1}yr^{-1}]$, (e) Erosion due to blowing snow, ER_{ds} or the divergence due to blowing snow $[kgm^{-1}yr^{-1}]$, and (f) Total sublimation SU_{total} , including surface- and blowing-snow sublimation $[kgm^{-1}yr^{-1}]$.




Figure 8. Monthly contribution to the yearly mean (2000–2012) (all in $Gtmo^{-1}$) (a) integrated blowing snow sublimation over total ice sheet, (b) integrated surface sublimation, and (c) integrated total sublimation ($SU_{tot} = SU_{ds} + SU_s$).

atmosphere, there is more condensation in winter compared to Rp3. Specifically, condensation in winter is nearly doubled with
 440 RpNew compared to Rp3. However, we do not see such a negative surface sublimation with CRYOWRF.

Total sublimation SU_{total} (Fig. 8(c)) follows a similar pattern as the surface sublimation, with higher values during Antarctic
 summer and relatively constant values in winter. It is clear from the NO-DRIFT case that the surface sublimation is negligible
 in Antarctic winter. The magnitude of total sublimation is comparable between RpNew and CRYOWRF in the Antarctic winter
 (March–September); the difference in the sublimation between different models is mostly a summer phenomenon. CRYOWRF
 445 consistently produces higher summer sublimation values than RpNew. While the seasonal trends of the sublimation remain
 unaltered between Rp3 and RpNew, interestingly, we observe that an increase in blowing snow sublimation in RpNew in
 winter leads to an increase in deposition, leading to limited overall changes to total sublimation.

4.5 Changes in integrated SMB

Table 2. Total ice sheet, including ice shelves, integrated SMB mean 2000–2012 values (Gtyr^{-1}) with interannual variability σ : total (snow
 and rain) precipitation (P_{tot}), total sublimation (SU_{tot}), surface sublimation (SU_{s}), blowing snow sublimation (SU_{ds}), blowing snow erosion
 (ER_{ds}), run-off (RU). ER_{ds} only considers the transport aspect of blowing snow. ER_{ds} is positive in case of erosion due to divergence of the
 blowing snow flux, and negative if convergence of the blowing snow flux brings snow to a grid box. Furthermore, as ER_{ds} only considers
 the snow redistribution, the spatially integrated impact on the SMB is zero as long as drifting snow is not blown off the ice sheet. Integrated
 surface mass balance is given by: $\text{SMB} = P_{\text{tot}} - SU_{\text{ds}} - SU_{\text{s}} - ER_{\text{ds}} - \text{RU}$. (a) Change between RpNew and NO-DRIFT. Percentage change
 is calculated as $(\text{RpNew} - \text{NO-DRIFT})/\text{NO-DRIFT}$, and (b) SMB difference between RpNew (2000–2012) with CRYOWRF (2010–2020)
 (Gerber et al., 2023)

(a) RpNew (2000–2012) and NO-DRIFT (2000–2012)						(b) RpNew (2000–2012) and CRYOWRF (2010–2020)					
	NO-DRIFT		RpNew		RpNew - NO-DRIFT	RpNew		CRYOWRF		RpNew - CRYOWRF	
	mean	σ	mean	σ	mean (% change)	mean	σ	mean	σ	mean	
P_{tot}	2622	96	2678	96	+56 (2%)	2678	96	3101	-	-423	
SU_{tot}	161	8	234	10	+76 (47%)	234	10	335	-	-101	
SU_{s}	161	8	59	8	-102 (-63%)	59	8	234	-	-175	
SU_{ds}	-	-	175	7	-	175	7	101	-	+74	
ER_{ds}	-	-	8	0.5	-	8	0.5	31	-	-23	
RU	7	3	7	3	0	7	3	5	-	+2	
SMB	2454	95	2428	96	-26 (-1%)	2428	96	2730	-	-302	

Table 2(a) presents the SMB and its components integrated over the whole ice sheet (including ice shelves) for the years 2000
 450 – 2012 in Gtyr^{-1} along with their inter-annual variability. Compared to NO-DRIFT, RpNew has an increased precipitation of
 56 Gtyr^{-1} caused by the higher moisture content in the atmosphere due to blowing snow sublimation. The total sublimation
 is increased by 76 Gtyr^{-1} with blowing snow sublimation being the major contributor. There is a decrease in surface sub-
 limation of 102 Gtyr^{-1} as blowing snow sublimation is now the dominant mechanism of sublimation. With blowing snow

transport fluxes, we have a snow erosion increase of 8 Gtyr^{-1} . Overall, the integrated SMB is reduced by 26 Gtyr^{-1} , due to
455 a net increase in blowing snow sublimation. The change amounts to only a 1% decrease in SMB compared to the NO-DRIFT
case. Since the change in SMB with the updates is minor and the SMB results from RACMO have been previously evaluated
against several in-situ and remote sensing observations, we refer to Noël et al. (2018); van Wessem et al. (2018) for the SMB
evaluation. Though there is negligible change in the overall SMB, blowing snow sublimation is highly important to local SMB,
especially in the escarpment areas in Eastern Antarctica.

460 Table 2(b) compares the integrated quantities obtained from RpNew with the results from CRYOWRF from 2010–2020.
While the period is different, Gerber et al. (2023) is the only other study (other than RACMO studies) that reports SMB results
of the entire Antarctica with a blowing snow model, making these results interesting to compare. Our experience with RACMO
runs suggest that the total sublimation from Rp3 does not vary much in the decade between 2000–2020. Therefore, we do not
expect a large difference in sublimation during this time period, and therefore, the results are comparable. Table 2(b) shows that
465 there is a large difference in SMB (11%) and precipitation (14%) between RpNew and CRYOWRF. While precipitation and
SMB are comparably higher in CRYOWRF, the ablation terms of CRYOWRF, especially sublimation, are more interesting.
Specifically, CRYOWRF produces higher total sublimation ($+101 \text{ Gtyr}^{-1}$) when compared to RpNew. From monthly average
sublimation (Fig. 8) we observed that the CRYOWRF produces higher surface sublimation in Antarctic summer (October–
March), when compared to winter.

470

5 Summary and conclusions

In this study, we updated the blowing snow model in the regional climate model RACMO, version 2.3p3 (Rp3), to better
represent the blowing snow phenomenon, the major ablation term in the SMB of the Antarctic ice sheet. As observed in the
limited available observations, the unaltered version of the model Rp3 failed to accurately predict the power-law variation of
475 blowing snow mass flux with wind speed. Furthermore, choices made in the unaltered version to reduce the blowing snow
model's computational expenses led to simplifications and assumptions that affected the model results. In the present work,
we updated the empirical formulation of saltation flux used as the boundary condition for the blowing snow model. We in-
creased the number and distribution of ice-particle radius classes to cover all the relevant blowing snow radius classes. We
also improved the coupling of the blowing snow model with RACMO by providing velocity, temperature profiles and friction
480 velocities from RACMO to the blowing snow model, which was previously being modelled as a logarithmic-law velocity and
the friction velocity was based on the first model level velocities. In addition, we found that the blowing snow model was very
sensitive to its time step and introduced sub-stepping for the blowing snow model, which significantly improved the results.

We ran the original blowing snow model (Rp3) and the updated code (RpNew) for Antarctica on a 27 km grid laterally
forced by 3-hourly ERA5 data. We performed three experiments for 2000–2012: NO-DRIFT, RpNew, and Rp3. In the exper-
485 iment NO-DRIFT, RACMO was run without the blowing snow model. The results from the updated model were evaluated
against in-situ observations from site D47, Adélie Land, Antarctica (Amory, 2020). Important surface quantities such as the

near-surface wind, temperature, humidity and blowing snow fluxes were compared. We found that RpNew results compared well against the blowing snow observations, successfully predicting both blowing snow frequency and magnitude. Furthermore, RpNew also successfully predicts the power-law variation of the blowing snow transport fluxes with wind speed. Comparison
490 of continental blowing snow frequency obtained from RpNew with CALIPSO satellite observations (Palm et al., 2018) shows that qualitatively, RpNew predicts the blowing snow frequency over Antarctica reasonably well.

The updated estimates of blowing snow sublimation from RpNew also agree well with the continent-wide estimates of blowing snow sublimation from satellite observations. Average blowing snow depth of 230 ± 116 m obtained from RpNew matches reasonably well with the satellite observations from Palm et al. (2017). Furthermore, Palm et al. (2017) from CALIPSO lidar
495 observations report a maximum blowing snow sublimation of 250 ± 125 mm w.e. yr^{-1} near the Antarctic coast around 140°E longitude. We observe a maximum blowing snow sublimation of 335 ± 30 mm w.e. yr^{-1} at the location: 66.9°S , 130.4°E . CALIPSO satellite observations indicate blowing snow sublimation could be as high as 393 ± 196 Gt yr^{-1} . We observe a blowing snow sublimation of 176 ± 10 Gt yr^{-1} with RpNew which shows there is a significant difference between model results and satellite observations. Palm et al. (2018) attribute the high blowing snow sublimation estimates to the errors associated with
500 MERRA-2 reanalysis data (Gelaro et al., 2017) used for calculating sublimation, particle radius error, and extinction errors and therefore, the satellite estimates involve a large error. However, without other continental-scale estimates of blowing snow sublimation, future studies must properly document the differences between different methods.

In the absence of blowing snow sublimation, the sublimation in Antarctica is mostly a summertime phenomenon (October–March), shown by NO-DRIFT experiment with negligible surface sublimation in winter. However, with the introduction of the
505 blowing snow model, total sublimation increases with a large contribution of blowing snow sublimation in the Antarctic winter (April–September). We observe an interesting self-limiting nature of total sublimation from RpNew model results. Specifically, while the RpNew leads to an increase in the blowing snow sublimation, we observed a corresponding decrease in the surface sublimation and a non-negligible increase in deposition, balancing the total sublimation in Antarctic winter. In RpNew, sublimation in Antarctica is a self-limiting mechanism where large blowing snow sublimation saturates the near-surface layers,
510 limiting the potential for surface sublimation. Future intercomparison studies with other models are necessary to test the hypothesis. We also compared RpNew result with the simulation results from CRYOWRF (Gerber et al., 2023). While blowing snow sublimation is the major contributor to the total sublimation in RpNew, surface sublimation is the dominant contributor to total sublimation in CRYOWRF. Furthermore, sublimation in CRYOWRF is nearly four times higher than RpNew surface sublimation. The difference shows that future intercomparison studies are necessary to identify the major contributor to total
515 sublimation.

In conclusion, the updates introduced to the regional climate model RACMO in this study significantly improve the representation of blowing snow physics within the model. Blowing snow and surface sublimation are the major mass loss terms in ~~the SMB of Antarctica, leading locally~~Antarctica's SMB, and locally, this leads to a negative SMB, which results in the formation of blue ice areas. This study presents a step forward in modelling blowing snow to produce a physically sound and
520 reliable estimate of ~~the SMB of Antarctica~~Antarctica's SMB. However, some limitations and simplifications exist in the model, which could be further improved. Currently, the snow properties such as dendricity and helicity are assumed to be constant.

525 As a result, the model does not capture variance in the observational data accurately; fixing this simplification will be a major improvement. In addition, the blowing snow model is not coupled with the radiation model, and as such, the effect of blowing snow on radiative transfer is not considered, which could be further improved. Finally, modelling of surface snow density and wind packing should be improved, as we have observed that the poor representation of the temporal and spatial variability of the surface snow density gives the largest uncertainties and model deviations in the modelled firn densification and firn air content.

Data availability. Monthly-accumulated sublimation and Yearly accumulated SMB components for NO-DRIFT, Rp3, and RpNew are publicly available on Zenodo (<https://zenodo.org/doi/10.5281/zenodo.12509004>) for the years 2000-2012. Observational data was downloaded from Amory et al. (2020b). CRYOWRF results were downloaded from Gerber et al. (2022).

Author contributions. SG and WJB conceived this study, decided on the new model settings. SG performed the code development and performed the model simulations and led the writing of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Appendix A: Near surface climate

Table A1. Root-mean-squared-error (RMSE), slope, intercept, bias, and coefficient of determination (R^2) of comparison of NO-DRIFT, Rp3, and RpNew simulations against observations at site D47. Statistics are reported for 2-m wind speed in ms^{-1} , 2-m temperature in $^{\circ}\text{C}$, 2-m relative humidity w.r.t ice in %, and the near surface blowing snow flux in $\text{kg m}^{-2} \text{s}^{-1}$.

	NO-DRIFT				Rp3				RpNew			
	U_{2m}	T_{2m}	RH_{2m}	Q_T	U_{2m}	T_{2m}	RH_{2m}	Q_T	U_{2m}	T_{2m}	RH_{2m}	Q_T
Slope	0.76	1.01	0.44	-	0.78	1.01	0.67	0.24	0.75	1.01	0.82	0.5
RMSE	3.84	3.09	18.84	-	3.69	3.17	9.39	0.04	3.88	3.04	6.64	0.03
R^2	0.77	0.91	0.07	-	0.76	0.91	0.35	0.24	0.76	0.91	0.49	0.57
Bias	-3.34	1.61	-13.88	-	-3.17	1.75	-5.61	0.003	-2.72	1.45	-0.87	-0.01

535 We evaluate the performance of RpNew in predicting the near-surface wind speed, temperature, relative humidity, and snow transport fluxes for 2010–2012 compared to the Rp3 and NO-DRIFT experiments. Table A1 presents the statistics comparing observed near-surface quantities against simulated results from the three experiments. We observe that the model underestimates the near-surface wind speed in all three experiments however with the current updates, the model bias is slightly

decreased from -3.34 m s^{-1} in the NO-DRIFT case to -2.72 m s^{-1} in the case of RpNew. Model captures the variability in
540 the data reasonably well, with negligible differences between the three experiments. The coefficient of determination (R^2)
is approximately 0.76, indicating that model results resemble the synoptic evolution of the wind strength well. A RMSE of
approximately 3.88 m s^{-1} indicates that there are still significant differences between the model results and observations. As
all three simulations underestimate the wind speed, we also performed tests with dual mass flux–TKE scheme (van Meijgaard
et al., 2012), which allows better modelling of the turbulent boundary-layer processes. However, it did not improve the wind-
545 speed predictions appreciably (not shown). Therefore, this scheme was not used further. The under-prediction of simulated
wind speed is likely due to the lower vertical resolution of the model, wherein the first atmospheric level is approximately 8 to
10 m above the surface, and the 2-m wind speed is calculated based on the similarity theory and is not simulated.

In the blowing snow model, the mass change of an ice particle due to the blowing snow sublimation is given by the model
of Thorpe and Mason (1966) (Eq. (10)). Since the mass change depends on water vapour deficit and air temperature, accurate
550 prediction of these quantities is necessary to obtain reliable estimates of blowing snow sublimation. Table A1 shows that the
near-surface temperature is overpredicted for all three experiments, and all simulations have a slight positive temperature bias.
However, with the updates to the model, the bias in the model is improved from $1.61 \text{ }^\circ\text{C}$ for the NO-DRIFT case to $1.45 \text{ }^\circ\text{C}$ for
RpNew. RpNew also shows an improved temperature prediction with a lower bias of $0.3 \text{ }^\circ\text{C}$ compared to Rp3. The variability
is modelled well, with an RMSE of 3°C , and a high $R^2 = 0.91$. The numbers show that RpNew predicts the near-surface wind
555 and temperature better than the other two experiments.

Appendix B: Blowing snow frequency

Figure B1 provides a colour-blind friendly version of the blowing snow frequency.

Appendix C: Difference between RpNew and Rp3

560 In Figure C1, we present the difference in some important variables between RpNew and Rp3 to quantify the magnitude of
change between the two versions. The blowing snow transport TR_{ds} (Fig. C1(b)) decreased somewhat over most of Antarctica
with significant but localised increases in transport along George V Land, Adélie Land, and Dronning Maud Land. At these
locations, the blowing snow transport is increased by 2 - 3 times compared to Rp3 due to better modelling of snow particle
distribution, which includes more particles with well-distributed ice particle radii and particle initialisation. As visualised in
565 Figure C1(g) as an example, however, for most of Antarctica, most blowing snow events are reduced in intensity by the model
updates. Only a few instances per year does the wind speed exceed the threshold for which the updated blowing snow model
simulates higher blowing snow transport.

Conversely, for most of Antarctica, we observe higher blowing snow sublimation (Fig. C1(a)) due to the ability of RpNew to
capture the peaks in blowing snow fluxes and the change in initialisation employed for the blowing snow model. This increase




Figure B1. Blowing snow frequency visualised to provide a qualitative comparison with satellite measurements of Palm et al. (2018). Figures show the average blowing snow frequency from 2000 to 2012.

570 indicates the necessity of a direct two-way coupling of the atmosphere with the blowing snow model. In RpNew, the snow particles are lifted into the warmer and drier air of the upper part of the stable boundary layer. In Rp3, particles were not lifted that high - due to errors in the particle size distribution. The larger ice shelves are the only regions of Antarctica where blowing snow sublimation decreases. Here, the stable boundary layer is generally very thick (e.g. van den Broeke and Van Lipzig (2003), Fig. 10), inhibiting the blowing snow from reaching the warmer air above the surface layer.

575 Results show RpNew is slightly colder by 0.3–0.4 K (Fig. C1(c)) along the coastal areas when compared to Rp3. This results from a better coupling of blowing snow sublimation to the tendencies of temperature, which allows the removal of latent heat from upper vertical levels of Rp3. Compared to Rp3, RpNew has higher relative humidity (Fig. C1(d)) also, due to better coupling of blowing snow moisture tendencies, the change in moisture leads to an increase in the dew-point temperature of 2 – 4 K (not shown here) in the first few vertical layers of Rp3. Furthermore, the total sublimation is higher in RpNew (Fig. C1(e))
 580 when compared to Rp3. Along the coast, the difference is as high as $100 \text{ mm w.e. yr}^{-1}$. Overall, the averaged surface mass balance (Fig. C1(f)) is changed mostly along the coastal Antarctica with a reduction of approximately $30 - 40 \text{ mm w.e. yr}^{-1}$.




Figure C1. Yearly averaged (2000-2012) difference between RpNew and Rp3 quantities: (a) Blowing snow sublimation (mm yr^{-1}), (b) Blowing snow transport ($\text{kg m}^{-1}\text{yr}^{-1}$), (c) Near-surface temperature T_{2m} in $^{\circ}\text{C}$, (d) Relative humidity in percentage, (e) Total sublimation (mm yr^{-1}), and (f) SMB (mm yr^{-1}). SU_{ds} , SU_{total} , and SMB are in mm_m water equivalent. (g) Instantaneous drifting snow flux at an interior region of East Antarctica (71.1°S , 111.7°E).

Since there is an increase in the moisture availability, there is relatively higher precipitation over Ronne and Ross ice shelves with a corresponding increase in SMB of approximately $20 \text{ mm w.e. yr}^{-1}$.

In conclusion, changes introduced in RpNew greatly influence the overall sublimation pattern in Antarctica and moisture content in lower levels of the atmosphere. In the RpNew, blowing snow's impact is more regional than Rp3. However, the overall impact on SMB is limited, with a decrease in SMB on the Eastern Antarctic coast and a slight increase in SMB in Western Antarctica due to higher moisture content created by blowing snow sublimation.

Table D1. Total ice sheet, including ice shelves, integrated SMB mean 2000–2012 values (Gt yr^{-1}) with interannual variability σ : total (snow and rain) precipitation (P_{tot}), total sublimation (SU_{tot}), surface sublimation (SU_{s}), blowing snow sublimation (SU_{ds}), blowing snow erosion (ER_{ds}), run-off (RU). Integrated surface mass balance is given by: $\text{SMB} = P_{\text{tot}} - SU_{\text{ds}} - SU_{\text{s}} - ER_{\text{ds}} - \text{RU}$.

	RpNew		Rp3		RpNew - Rp3
	mean	σ	mean	σ	mean (%change)
P_{tot}	2678	96	2655	98	+23 (0.9%)
SU_{tot}	234	10	186	6	+48 (26%)
SU_{s}	59	8	71	5	-12 (17%)
SU_{ds}	175	7	115	4	+60 (52%)
ER_{ds}	8	0.5	5	0.2	+3 (60%)
RU	7	3	7	3	0 (0%)
SMB	2428	97	2458	96	-30 (1.2%)

Appendix D: Changes in integrated SMB between RpNew and Rp3

590 Table D1 presents the SMB and its components integrated over the whole ice sheet (including ice shelves) for the years 2000
– 2010 in Gt yr^{-1} along with their inter-annual variability. Compared to Rp3, RpNew has an increased precipitation of 23
 Gt yr^{-1} caused by the higher moisture content in the atmosphere due to higher blowing snow sublimation. The total sublima-
tion is increased by 48 Gt yr^{-1} with blowing snow sublimation being the major contributor. There is a slight decrease in surface
sublimation (12 Gt yr^{-1}) as air in the boundary layer is saturated more efficiently with RpNew compared to Rp3, which causes
595 a reduction in the potential for the surface sublimation. With higher blowing snow transport fluxes, we have a higher snow
erosion increase of 3 Gt yr^{-1} . This number remained small as snow erosion only influences the integrated SMB once the snow
is blown off the ice sheet. Overall, the integrated SMB is reduced by 30 Gt yr^{-1} , due to a net increase in blowing snow subli-
mation. The change amounts to only a 1.2% decrease in SMB compared to Rp3. Since the SMB changes with the updates are
minor and the SMB results from Rp2 have been previously evaluated against several in-situ and remote sensing observations,
600 we refer to Noël et al. (2018); van Wessem et al. (2018) for the SMB evaluation. Though there is negligible change in the over-
all SMB, blowing snow sublimation is highly important to local SMB, especially in the escarpment areas in Eastern Antarctica.

Acknowledgements. We would like to thank Charles Amory for the discussion about observational dataset from site D47, East Antarctica. We
would also like to thank Melchior van Wessem and Christiaan van Dalum for discussions about RACMO model development. This project
605 has received funding from the European Union’s Horizon 2020 research and innovation program under grant agreement no. 101003590
(PolarRES). We also acknowledge the ECMWF for archiving facilities and computational time on their supercomputers.

References

- Agosta, C., Amory, C., Kittel, C., Orsi, A., Favier, V., Gallée, H., van den Broeke, M. R., Lenaerts, J. T. M., van Wessem, M. J., van de Berg, W. J., et al.: Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979–2015) and identification of dominant processes, *The Cryosphere*, 13, 281–296, <https://doi.org/10.5194/tc-13-281-2019>, 2019.
- 610 Alduchov, O. A. and Eskridge, R. E.: Improved Magnus form approximation of saturation vapor pressure, *Journal of Applied Meteorology and Climatology*, 35, 601–609, [https://doi.org/https://doi.org/10.1175/1520-0450\(1996\)035%3C0601:IMFAOS%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0450(1996)035%3C0601:IMFAOS%3E2.0.CO;2), 1996.
- Amory, C.: Drifting-snow statistics from multiple-year autonomous measurements in Adélie Land, East Antarctica, *The Cryosphere*, 14, 1713–1725, <https://doi.org/10.5194/tc-14-1713-2020>, 2020.
- 615 Amory, C., Trouvilliez, A., Gallée, H., Favier, V., Naaim-Bouvet, F., Genthon, C., Agosta, C., Piard, L., and Bellot, H.: Comparison between observed and simulated aeolian snow mass fluxes in Adélie Land, East Antarctica, *The Cryosphere*, 9, 1373–1383, <https://doi.org/10.5194/tc-9-1373-2015>, 2015.
- Amory, C., Gallée, H., Naaim-Bouvet, F., Favier, V., Vignon, E., Picard, G., Trouvilliez, A., Piard, L., Genthon, C., and Bellot, H.: Seasonal variations in drag coefficient over a sastrugi-covered snowfield in coastal East Antarctica, *Boundary-Layer Meteorology*, 164, 107–133, <https://doi.org/10.1007/s10546-017-0242-5>, 2017.
- 620 Amory, C., Genthon, C., and Favier, V.: A drifting snow data set (2010–2018) from coastal Adélie Land, eastern Antarctica, Zenodo [data set], 10, <https://doi.org/10.5281/zenodo.3630496>, 2020a.
- Amory, C., Genthon, C., and Favier, V.: A drifting snow data set (2010–2018) from coastal Adélie Land, Eastern Antarctica, Data, <https://doi.org/10.5281/zenodo.3630497>, 2020b.
- 625 Amory, C., Kittel, C., Toumelin, L. L., Agosta, C., Delhasse, A., Favier, A., and Fettweis, X.: Performance of MAR (v3.11) in simulating the drifting-snow climate and surface mass balance of Adélie Land, East Antarctica, *Geoscientific Model Development*, 14, 3487–3510, <https://doi.org/10.5194/gmd-14-3487-2021>, 2021.
- Barral, H., Genthon, C., Trouvilliez, A., Brun, C., and Amory, C.: Blowing snow in coastal Adélie Land, Antarctica: three atmospheric-moisture issues, *The Cryosphere*, 8, 1905–1919, <https://doi.org/10.5194/tc-8-1905-2014>, 2014.
- 630 Bintanja, R.: The contribution of snowdrift sublimation to the surface mass balance of Antarctica, *Annals of Glaciology*, 27, 251–259, <https://doi.org/10.3189/1998AoG27-1-251-259>, 1998.
- Bintanja, R.: Snowdrift sublimation in a katabatic wind region of the Antarctic ice sheet, *Journal of Applied Meteorology and Climatology*, 40, 1952–1966, [https://doi.org/10.1175/1520-0450\(2001\)040<1952:SSIAKW>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1952:SSIAKW>2.0.CO;2), 2001.
- Bromwich, D. H., Guo, Z., Bai, L., and Chen, Q.-S.: Modeled Antarctic precipitation. Part I: Spatial and temporal variability, *Journal of Climate*, 17, 427–447, [https://doi.org/https://doi.org/10.1175/1520-0442\(2004\)017%3C0427:MAPPIS%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2004)017%3C0427:MAPPIS%3E2.0.CO;2), 2004.
- 635 Budd, W. F.: The drifting of nonuniform snow particles, *Studies in Antarctic meteorology*, 9, 59–70, <https://doi.org/10.1029/AR009p0059>, 1966.
- CY45R1—Part IV, I. D.: Physical processes, IFS Documentation CY45R1, 2018.
- Déry, S. J. and Yau, M. K.: A bulk blowing snow model, *Boundary-Layer Meteorology*, 93, 237–251, <https://doi.org/10.1023/A:1002065615856>, 1999.
- 640 Déry, S. J. and Yau, M. K.: Simulation of blowing snow in the Canadian Arctic using a double-moment model, *Boundary-Layer Meteorology*, 99, 297–316, <https://doi.org/10.1023/A:1018965008049>, 2001.

- Déry, S. J. and Yau, M. K.: Large-scale mass balance effects of blowing snow and surface sublimation, *Journal of Geophysical Research: Atmospheres*, 107, ACL–8, <https://doi.org/10.1029/2001JD001251>, 2002.
- 645 Déry, S. J., Taylor, P. A., and Xiao, J.: The thermodynamic effects of sublimating, blowing snow in the atmospheric boundary layer, *Boundary-Layer Meteorology*, 89, 251–283, <https://doi.org/10.1023/A:1001712111718>, 1998.
- ECMWF: IFS Documentation CY33R1 - Part IV: Physical Processes, 4, ECMWF, <https://doi.org/10.21957/8o7vwlbdr>, 2009.
- Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Box, J. E., and Steffen, K.: Climate of the Greenland ice sheet using a high-resolution climate model—Part 1: Evaluation, *The Cryosphere*, 4, 511–527, <https://doi.org/10.5194/tc-4-511-2010>, 2010.
- 650 Gallée, H., Guyomaréh, G., and Brun, E.: Impact of snow drift on the Antarctic ice sheet surface mass balance: possible sensitivity to snow-surface properties, *Boundary-Layer Meteorology*, 99, 1–19, <https://doi.org/10.1023/A:1018776422809>, 2001.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., et al.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *Journal of climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 655 Gerber, F., Sharma, V., and Lehning, M.: Reproducibility dataset for CRYOWRF validation, <https://doi.org/http://dx.doi.org/10.16904/envidat.347>, 2022.
- Gerber, F., Sharma, V., and Lehning, M.: CRYOWRF—Model Evaluation and the Effect of Blowing Snow on the Antarctic Surface Mass Balance, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 744, <https://doi.org/10.1029/2022JD037744>, 2023.
- Greuell, W. and Konzelmann, T.: Numerical modelling of the energy balance and the englacial temperature of the Greenland Ice Sheet. Calculations for the ETH-Camp location (West Greenland, 1155 m asl), *Global and Planetary change*, 9, 91–114, [https://doi.org/10.1016/0921-8181\(94\)90010-8](https://doi.org/10.1016/0921-8181(94)90010-8), 1994.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- King, J. C., Anderson, P. S., Smith, M. C., and Mobbs, S. D.: The surface energy and mass balance at Halley, Antarctica during winter, 665 *Journal of Geophysical Research: Atmospheres*, 101, 19 119–19 128, <https://doi.org/10.1029/96JD01714>, 1996.
- King, J. C., Anderson, P. S., and Mann, G. W.: The seasonal cycle of sublimation at Halley, Antarctica, *Journal of Glaciology*, 47, 1–8, <https://doi.org/10.3189/172756501781832548>, 2001.
- Kodama, Y., Wendler, G., and Gosink, J.: The effect of blowing snow on katabatic winds in Antarctica, *Annals of Glaciology*, 6, 59–62, <https://doi.org/10.3189/1985AoG6-1-59-62>, 1985.
- 670 Kuipers Munneke, P., Van den Broeke, M., Lenaerts, J., Flanner, M., Gardner, A., and Van de Berg, W.: A new albedo parameterization for use in climate models over the Antarctic ice sheet, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2010JD015113>, 2011.
- Lenaerts, J. T. M. and van den Broeke, M. R.: Modeling drifting snow in Antarctica with a regional climate model: 2. Results, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2010JD015419>, 2012.
- 675 Lenaerts, J. T. M., van den Broeke, M. R., Déry, S. J., van Meijgaard, E., van de Berg, W. J., Palm, S. P., and Rodrigo, J. S.: Modeling drifting snow in Antarctica with a regional climate model: 1. Methods and model evaluation, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016145>, 2012.
- Lenaerts, J. T. M., Smeets, C. J. P. P., Nishimura, K., Eijkelboom, M., Boot, W., van den Broeke, M. R., and van de Berg, W. J.: Drifting snow measurements on the Greenland Ice Sheet and their application for model evaluation, *The Cryosphere*, 8, 801–814, 680 <https://doi.org/10.5194/tc-8-801-2014>, 2014.

- Libois, Q., Picard, G., France, J. L., Arnaud, L., Dumont, M., Carmagnola, C. M., and King, M. D.: Influence of grain shape on light penetration in snow, *The Cryosphere*, 7, 1803–1818, <https://doi.org/10.5194/tc-7-1803-2013>, 2013.
- Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., Boberg, F., van de Berg, W. J., Fettweis, X., Gossart, A., et al.: What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates, *The Cryosphere*, 15, 3751–3784, <https://doi.org/10.5194/tc-15-3751-2021>, 2021.
- Noël, B., van de Berg, W. J., van Wessem, M. J., van Meijgaard, E., van As, D., Lenaerts, J., Lhermitte, S., Munneke, P. K., Smeets, C. J. P., van Ulft, L. H., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.
- Palm, S. P., Yang, Y., Spinhirne, J. D., and Marshak, A.: Satellite remote sensing of blowing snow properties over Antarctica, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD015828>, 2011.
- Palm, S. P., Kayetha, V., Yang, Y., and Pauly, R.: Blowing snow sublimation and transport over Antarctica from 11 years of CALIPSO observations, *The Cryosphere*, 11, 2555–2569, <https://doi.org/10.5194/tc-11-2555-2017>, 2017.
- Palm, S. P., Kayetha, V., and Yang, Y.: Toward a satellite-derived climatology of blowing snow over Antarctica, *Journal of Geophysical Research: Atmospheres*, 123, 10–301, <https://doi.org/10.1029/2018JD028632>, 2018.
- Pomeroy, J. W.: A process-based model of snow drifting, *Annals of Glaciology*, 13, 237–240, <https://doi.org/10.3189/S0260305500007965>, 1989.
- Pomeroy, J. W. and Male, D. H.: Steady-state suspension of snow, *Journal of hydrology*, 136, 275–301, [https://doi.org/10.1016/0022-1694\(92\)90015-N](https://doi.org/10.1016/0022-1694(92)90015-N), 1992.
- Radok, U.: Snow drift, *Journal of Glaciology*, 19, 123–139, <https://doi.org/10.3189/S0022143000215591>, 1977.
- Scambos, T. A., Frezzotti, M., Haran, T., Bohlander, J., Lenaerts, J., Van Den Broeke, M., Jezek, K., Long, D., Urbini, S., Farness, K., et al.: Extent of low-accumulation ‘wind glaze’ areas on the East Antarctic plateau: implications for continental ice mass balance, *Journal of glaciology*, 58, 633–647, <https://doi.org/10.3189/2012JoG11J232>, 2012.
- Scarchilli, C., Frezzotti, M., Grigioni, P., Silvestri, L. D., Agnoletto, L., and Dolci, S.: Extraordinary blowing snow transport events in East Antarctica, *Climate Dynamics*, 34, 1195–1206, <https://doi.org/10.1007/s00382-009-0601-0>, 2010.
- Schmidt, R. A.: Sublimation of wind-transported snow: a model, vol. 90, Rocky Mountain Forest and Range Experiment Station, Forest Service, US . . . , 1972.
- Schmidt, R. A.: Vertical profiles of wind speed, snow concentration, and humidity in blowing snow, *Boundary-Layer Meteorology*, 23, 223–246, <https://doi.org/10.1007/BF00123299>, 1982.
- Serreze, M. C. and Barry, R. G.: *The Arctic climate system*, Cambridge University Press, 2005.
- Thiery, W., Gorodetskaya, I. V., Bintanja, R., van Lipzig, N. P. M., van den Broeke, M. R., Reijmer, C. H., and Munneke, P. K.: Surface and snowdrift sublimation at Princess Elisabeth station, East Antarctica, *The Cryosphere*, 6, 841–857, <https://doi.org/10.5194/tc-6-841-2012>, 2012.
- Thorpe, A. and Mason, B.: The evaporation of ice spheres and ice crystals, *British Journal of Applied Physics*, 17, 541, <https://doi.org/10.1088/0508-3443/17/4/316>, 1966.
- Toumelin, L. L., Amory, C., Favier, V., Kittel, C., Hofer, S., Fettweis, X., Gallée, H., and Kayetha, V.: Sensitivity of the surface energy budget to drifting snow as simulated by MAR in coastal Adelie Land, Antarctica, *The Cryosphere*, 15, 3595–3614, <https://doi.org/10.5194/tc-15-3595-2021>, 2021.

- Undén, P., Rontu, L., Jarvinen, H., Lynch, P., Sánchez, F. J. C., Cats, G., Cuxart, J., Eerola, K., Fortelius, C., García-Moya, J. A., et al.: HIRLAM-5 scientific documentation, <http://hdl.handle.net/20.500.11765/6323>, 2002.
- 720 van Dalum, C. T., van de Berg, W. J., Libois, Q., Picard, G., and van den Broeke, M. R.: A module to convert spectral to narrowband snow albedo for use in climate models: SNOWBAL v1. 2, *Geoscientific Model Development*, 12, 5157–5175, <https://doi.org/10.5194/gmd-12-5157-2019>, 2019.
- van Dalum, C. T., van de Berg, W. J., and van den Broeke, M. R.: Sensitivity of Antarctic surface climate to a new spectral snow albedo and radiative transfer scheme in RACMO2.3p3, *The Cryosphere*, 16, 1071–1089, <https://doi.org/10.5194/tc-16-1071-2022>, 2022.
- 725 van den Broeke, M. and Bintanja, R.: The interaction of katabatic winds and the formation of blue-ice areas in East Antarctica, *Journal of Glaciology*, 41, 395–407, <https://doi.org/10.3189/S0022143000016269>, 1995.
- van den Broeke, M. and Van Lipzig, N.: Factors controlling the near-surface wind field in Antarctica, *Monthly Weather Review*, 131, 733–743, 2003.
- van den Broeke, M. R., Reijmer, C. H., and van de Wal, R. S. W.: A study of the surface mass balance in Dronning Maud Land, Antarctica, using automatic weather stations, *Journal of Glaciology*, 50, 565–582, <https://doi.org/10.3189/172756504781829756>, 2004.
- 730 van Meijgaard, E., van Ulft, L., Lenderink, G., de Roode, S. R., Wipfler, E. L., Boers, R., and van Timmermans, R. M. A.: Refinement and application of a regional atmospheric model for climate scenario calculations of Western Europe, KVR 054/12, KvR, <https://library.wur.nl/WebQuery/wurpubs/427097>, 2012.
- van Wessem, J. M., Ligtenberg, S. R. M., Reijmer, C. H., van de Berg, W. J., van den Broeke, M. R., Barrand, N. E., Thomas, E. R., Turner, J., Wuite, J., Scambos, T. A., et al.: The modelled surface mass balance of the Antarctic Peninsula at 5.5 km horizontal resolution, *The Cryosphere*, 10, 271–285, <https://doi.org/10.5194/tc-10-271-2016>, 2016.
- 735 van Wessem, M. J., Reijmer, C. H., van de Berg, W. J., van den Broeke, M. R., Cook, A. J., van Ulft, L. H., and van Meijgaard, E.: Temperature and wind climate of the Antarctic Peninsula as simulated by a high-resolution regional atmospheric climate model, *Journal of Climate*, 28, 7306–7326, <https://doi.org/10.1175/JCLI-D-15-0060.1>, 2015.
- 740 van Wessem, M. J., van de Berg, W. J., Noël, P. Y. B., van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 2: Antarctica (1979–2016), *The Cryosphere*, 12, 1479–1498, <https://doi.org/10.5194/tc-12-1479-2018>, 2018.
- Vinokur, M.: On one-dimensional stretching functions for finite-difference calculations, *Journal of Computational Physics*, 50, 215–234, [https://doi.org/https://doi.org/10.1016/0021-9991\(83\)90065-7](https://doi.org/https://doi.org/10.1016/0021-9991(83)90065-7), 1983.