



Examining ENSO related variability in tropical tropospheric ozone in the RAQMS-Aura chemical reanalysis

Maggie Bruckner¹, R. Bradley Pierce^{1,2}, Allen Lenzen²

¹Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, Madison, WI, 53706, USA

5 ²Space Science and Engineering Center, University of Wisconsin–Madison, Madison, WI, 53706, USA

Correspondence to: Maggie Bruckner (mebruckner@wisc.edu)

Abstract

The El Niño-Southern Oscillation (ENSO) is a major driver of interannual variability in both tropical and mid-latitudes and has been found to have a strong impact on the distribution of tropospheric ozone in the tropical Pacific in satellite
10 observational datasets, chemical transport models, and chemistry-climate simulations. Here we analyze interannual variability in tropical tropospheric ozone by applying composite analysis, empirical orthogonal function (EOF) analysis and multiple linear regression to the Real-time Air Quality Modeling System (RAQMS) Aura chemical reanalysis. As shown in similar studies, the dominant mode of interannual variability in tropical tropospheric ozone is driven by ENSO. ENSO composites show that the ENSO signature in tropospheric ozone is strongest near the tropopause. We also show an
15 enhancement in tropical ozone over the maritime continent below 700 hPa during El Niño that is dependent on the magnitude of the biomass burning emissions in the region. We reconstruct the ENSO variability in tropical tropospheric ozone through a multiple linear regression of principal components for precipitation and CO. The multiple linear regression quantifies that variability in biomass burning contributes to ENSO variability in tropical tropospheric ozone though the dominant driver is convection.

20 1 Introduction

The development of methods to calculate tropospheric ozone residuals (TOR) from satellite total column observations (eg. Fishman and Balok, 1999; Fishman et al., 1990; Fishman and Larsen, 1987) provided the first global view of tropospheric ozone and showed a systematic zonal wave one structure in the tropics. This zonal wave one structure is consistent with the climatological average state of tropical atmosphere which is dominated by the Pacific Walker circulation, defined by
25 ascending motion over warm SSTs near the maritime continent and descending over cooler SSTs in the eastern Pacific, with easterlies at surface and westerlies aloft. Climatologically, tropospheric ozone columns are lowest over the Pacific and highest downwind of western Africa (Fishman et al., 1990, 1996, 2003). The enhancement downwind of western Africa is strongest during September-October-November (SON) and is associated with photochemical production of ozone from biomass burning emissions (Fishman et al., 2003, 1996, 2005). Tropospheric ozone concentrations over Africa and South



30 America are lowest in March-April-May (MAM) (Fishman et al., 2003, 1990). The Fishman, Wozniak, and Creilson 2003
TOR seasonal climatology also shows a variance of 5-10 DU over the maritime continent from Decenber-January-February
(DJF) to June-July-August (JJA). The El Niño-Southern Oscillation (ENSO) is a major driver of interannual variability in
both tropical and mid-latitudes (eg. McPhaden et al., 2006; Trenberth, 1997), and has been found to have a strong impact on
the distribution of tropospheric ozone in the tropical Pacific (Doherty et al., 2006; Peters et al., 2001; Sekiya and Sudo, 2012;
35 Ziemke et al., 2010).

ENSO phases of El Niño and La Niña are tracked using a variety of indexes including the Niño 3.4 index (Bamston et al.,
1997; Trenberth, 1997) and the Ozone ENSO Index (Ziemke et al., 2010). El Niño events occur when a warm SST anomaly
develops in the eastern Pacific and reduces the east-west temperature gradient across the equatorial Pacific. In response to
the SST anomaly, the trade winds weaken. Convection is enhanced over the eastern Pacific, leading to increased
40 precipitation in the region and an eastward shift of the Walker Circulation. Correspondingly convection is suppressed over
the maritime continent and leads to drier than usual conditions During El Niño events, tropospheric ozone is lower over the
Pacific as the enhanced convection lofts low ozone air masses from near the ocean surface higher into the column, and
higher over the maritime continent as high upper tropospheric ozone descends (eg. Doherty et al., 2006; Hou et al., 2016;
Sudo and Takahashi, 2001). Variability in the location of the maximum SST anomaly during the El Niño phase has led to a
45 distinction between canonical (eastern Pacific) El Niño events and El Niño Modoki (central Pacific) events (eg. Larkin and
Harrison, 2005; Kim and Yu, 2012; Santoso et al., 2017). In the canonical El Niño, the maximum SST anomaly extends into
the eastern tropical Pacific cold pool while during El Niño Modoki the maximum SST anomaly is in the central Pacific. The
ascending branches of the Walker circulation are over the central Pacific during El Niño Modoki (Ashok et al., 2007).
Following from the differences in the Walker circulation, the pattern of the ENSO response in tropical tropospheric ozone
50 depends on the type of El Niño (Hou et al., 2016).

La Niña events occur when the eastern Pacific is cooler than average, and the atmosphere responds in an opposite manner to
El Niño as enhanced vertical motion and convection occurs over the maritime continent, suppression of convection occurs
over the east Pacific, and enhanced downwelling over the east Pacific. Tropical tropospheric ozone columns reflect the
impacts of higher concentration upper tropospheric ozone descending over the Pacific and comparatively lower
55 concentration lower tropospheric ozone ascending near the maritime continent during La Niña (eg. Ziemke and Chandra,
2003; Doherty et al., 2006).

The influence of ENSO on tropospheric ozone has previously been investigated in observational datasets, chemical transport
models, and chemistry-climate models. Application of statistical techniques (regression, correlation, and empirical
orthogonal functions) to TOR data revealed that interannual variability in measurements over the tropical Pacific is
60 dominated by ENSO (eg. Ziemke et al., 1998, 2010; Doherty et al., 2006). ENSO variability in tropical tropospheric ozone
columns has been reproduced in chemical transport models and climate models (eg. Sudo and Takahashi, 2001; Chandra et
al., 2002; Peters et al., 2001; Doherty et al., 2006; Sekiya and Sudo, 2014). ENSO variability in equatorial Pacific
tropospheric ozone was initially thought to be equally due to shifts in biomass burning emissions and meteorological



65 conditions (Sudo and Takahashi, 2001; Chandra et al., 2002). More contemporary studies indicate enhancement in biomass
burning during El Niño results in regional enhancement of ozone with little contribution to global tropospheric ozone
variability and that the response of tropospheric ozone to ENSO is primarily due to dynamical processes (Doherty et al.,
2006; Inness et al., 2015).

In this study we will investigate the interannual variability of tropical tropospheric ozone in a chemical re-analysis extending
from 2006 through 2016. A chemical re-analysis produces a long-term data record by cycling a model forecast and data
70 assimilation system to combine forecasts and observations in a statistically consistent manner that accounts for forecast and
observation error (Miyazaki et al., 2020; Yumimoto et al., 2017). The data record obtained is a best-estimate of the real
composition of the atmosphere, as analyses produced are constrained by observations of a limited number of species and the
evolution of those species by model physics (Miyazaki et al., 2020). A comparison of several recent chemical reanalyses
including the Copernicus Atmospheric Monitoring Service (CAMS) reanalysis (Inness et al., 2019), and the Tropospheric
75 Chemistry Reanalysis version 2 (TCR-2) (Miyazaki et al., 2020) found that these analyses are suitable for generating ozone
climatologies and looking at trends, though individual reanalyses will differ due to model configuration (Huijnen et al.,
2020). While reanalysis has been used to look at the ENSO signal in CO, O₃, NO_x, and smoke aerosols (Inness et al., 2015),
our analysis will make use of the chemical production and loss terms, convective mass flux, and diabatic heating from the
reanalysis to examine variability in tropospheric ozone. We also focus on the 2006-2016 period, which includes significant
80 biomass burning events during the 2015/2016 El Niño event.

This study seeks to: 1) evaluate the tropical tropospheric ozone column variability associated with ENSO in a 1x1 degree
chemical reanalysis using the Real-time Air Quality Modeling System (RAQMS, Pierce et al., 2007) and satellite
measurements from the NASA Aura satellite (Pierce et al., 2016) and 2) investigate how the 2015/2016 extreme El Niño
event impacts the ENSO response.

85 **2 Methods**

2.1 RAQMS-Aura

The Real-time Air Quality Modeling System (RAQMS) Aura Reanalysis, hereafter RAQMS-Aura, is a chemical reanalysis
using RAQMS (Pierce et al, 2007), a global chemical transport model with full stratospheric and tropospheric chemistry, and
satellite trace gas and aerosol retrievals from the NASA satellites (Terra, Aqua, and Aura) covering 2006 through 2016.
90 RAQMS-Aura provides 1°x1° global chemical analyses, on 35 hybrid model levels from the surface to approximately 60 km
above ground level, at 3-hour time steps. The operational grid point statistical interpolation (GSI) 3d var system (Wu et al.,
2002) is used to assimilate retrievals from the following Aura instruments: Aura Ozone Monitoring Instrument (OMI) cloud
cleared total column ozone (McPeters et al., 2008), Microwave Limb Sounder (MLS) (Froidevaux et al., 2008) stratospheric
ozone profiles, and OMI tropospheric column NO₂ (Boersma et al., 2007; Bucsela et al., 2013). NASA Terra and Aqua
95 Moderate Resolution Imaging Spectrometer (MODIS) aerosol optical depth (AOD) (Remer et al., 2005) and Atmospheric



Infrared Sounder (AIRS) carbon monoxide (Yurganov et al., 2008; McMillan et al., 2005) are also assimilated at three-hour intervals. Analysis increments from the OMI tropospheric column NO₂ retrievals are used for off-line adjustment of a priori 2010 Hemispheric Transport of Air Pollution (HTAP, 2010) anthropogenic emission inventories following an offline mass balance approach similar to East et al. 2022. Biomass burning emissions in RAQMS-Aura use Terra and Aqua MODIS fire 100 detections and are calculated using a bottom-up approach developed by Soja et al. 2004 and compared to other approaches in Al-Saadi et al. 2008. This approach estimates total carbon emissions at MODIS fire detections with the US Forest Service Haines Index (Haines, 1989) to determine fire weather severity and gridded, ecosystem-dependent estimates of carbon consumption for low, medium, and high fire severity fires. Emission ratios are then used to estimate emissions of CO, NO_x, and hydrocarbons from the calculated total carbon emissions.

105 The dynamical core of RAQMS is the UW hybrid model (Schaack et al., 2004). The UW hybrid model utilizes physical parameterizations from the NCAR Community Climate Model (CCM3) (Kiehl et al., 1998), including the moist convection scheme. The CCM3 moist convection scheme combines the Zhang and McFarlane (1995) deep convection scheme with shallow and midlevel convection following Hack (1994). The deep convection scheme treats convection as an ensemble of updrafts and downdrafts, and the shallow convection scheme treats convection as separate plumes within 3 successive layers 110 whereby mass is detrained from one layer into the next (Kiehl et al., 1998; Zhang et al., 1998). RAQMS-Aura initializes its meteorological fields with archived analyses from the National Center for Environmental Prediction (NCEP) Global Data Assimilation System (GDAS) (Kleist et al., 2009; Wang et al., 2013). These fields are impacted by updates to physics, resolution, and data assimilation used in the GDAS system (MODEL CHANGES SINCE 1991, 2023).

2.2 ENSO Composites

115 Anomaly composites are used to evaluate how well RAQMS-Aura reproduces observed ENSO variability. El Niño and La Niña periods are determined by use of the Niño 3.4 index. ENSO events are defined as occurring when the index is at least 0.4°C greater (El Niño) or less (La Niña) than average for 5 consecutive months (eg. Trenberth, 1997; Ziemke et al., 2015). The anomaly is defined as the deviation from the average annual cycle during the RAQMS-Aura analysis period (2006-2016). Anomaly composites for El Niño and La Niña periods are generated for precipitation, convective mass flux, diabatic 120 heating, ozone concentration, carbon monoxide, and net ozone production from monthly mean RAQMS-Aura analyses. Anomaly composites are also generated for satellite observations of tropospheric ozone column, total column carbon monoxide, and total precipitation. To investigate the vertical structure of ENSO variability in RAQMS-Aura, anomaly cross section composites are calculated between 7.5°S to 2.5°N for convective mass flux, diabatic heating, ozone, carbon monoxide, and net ozone production.

125 2.3 Empirical Orthogonal Function (EOF) Analysis

EOF analysis has been used previously by Peters et al. 2001 and Doherty et al. 2006 to identify ENSO variability in modeled tropospheric ozone concentrations. EOF analysis is performed on de-seasonalized and detrended precipitation, CO



column, and tropical tropospheric ozone column (TTOC) monthly mean anomalies to determine the dominant modes of tropical variability in RAQMS-Aura analyses.

130 Following Doherty et al 2006 the resulting EOF patterns for each RAQMS-Aura variable are multiplied by the standard deviation of the associated principal component (PC) to produce the physical magnitude of change associated with the mode. The PCs are correlated against the Niño 3.4 index to assess whether the mode captured by the EOF accounts for ENSO variability. A multiple linear regression is constructed using the precipitation and CO PCs to investigate how variability in convection and biomass burning emissions drive the ozone ENSO signal.

135 **3 Results**

3.1 Validation of RAQMS-Aura Precipitation

Prior to investigating variability of the RAQMS-Aura chemical fields, we evaluate RAQMS-Aura convection and precipitation processes through comparisons with observations. In RAQMS-Aura, sub-grid-scale mass flux between model layers occurs through shallow and deep convective schemes. Diabatic heating is generated by the sub-grid-scale convective parameterizations and influences the grid-scale thermodynamics. Convective mass flux and diabatic heating will be used in the composite analysis to look at the impact of ENSO on vertical transport and tropical tropospheric ozone concentrations.

140 Monthly mean total and convective precipitation from RAQMS-Aura is compared to estimates of precipitation from the Tropical Rainfall Measuring Mission (TRMM) Multi-satellite precipitation Analysis (TMPA) 3B43 product (Huffman et al., 2007). TRMM 3B43 merges satellite IR and microwave precipitation estimates with rain gauge data to produce a best estimate of monthly mean precipitation rate from 50°S to 50°N at 0.25x0.25 degree resolution, which in this study is averaged onto the RAQMS 1x1 degree grid. Our analysis is focused on meridional structure and seasonal maps to look at average regional biases, and time-series of the maritime continent and Pacific Intertropical convergence zone (ITCZ) regions to look at longer-term trends.

3.1.1 Meridional Structure

150 Figure 1 displays the meridional averaged convective, large-scale, and total precipitation for RAQMS-Aura and total precipitation from TRMM 3B43 for each season. The seasonal average meridional precipitation maxima in RAQMS-Aura are broader than observed in TRMM 3B43. During DJF and MAM, observed tropical precipitation peaks in both the northern hemisphere (NH) and southern hemisphere (SH). During JJA and SON, observed tropical precipitation peaks only in the NH.

155 In DJF the observed hemispheric peaks are of similar magnitude with the NH peaking at 0.247 mm/hour and the SH peaking at 0.233 mm/hour. TRMM 3B43 MAM indicates that the NH branch is more active during this season than the SH branch, as the NH peak is 0.293 mm/hour and the SH peak is 0.229 mm/hour. RAQMS-Aura reproduces the observed double peaks for DJF and MAM, though the magnitude is overestimated in RAQMS-Aura by 0.08-0.12 mm/hour, and the DJF SH peak is



larger than the NH peak and 5 degrees to the south of the observed peak. In JJA and SON, the reanalysis reproduces the
160 observed single maxima, though it is broader by more than 15 degrees latitude, and the absolute maximum is displaced
approximately 2.5 degrees to the north.

Between 40°N and 40°S the total precipitation in RAQMS-Aura is predominately convective precipitation, with ratios of
convective precipitation to total precipitation exceeding 0.6 on average. It is common for tropical precipitation to be
predominately convective precipitation in global models, leading to a “drizzling bias”. This “drizzling bias” is the result of
165 convective parameterizations producing convective precipitation that is too frequent and long-lasting but not as intense as
observed while the total precipitation amount is realistic (Chen et al., 2021; Chen and Dai, 2019).




Figure 1. Zonally and seasonally averaged precipitation from RAQMS-Aura and TRMM 3B43 for a) DJF, b) MAM, c) JJA, and d) SON.



170 3.1.2 Horizontal Structure

While RAQMS-Aura reasonably reproduces the seasonality of the observed meridional structure, the distributions are broader than in observations. Seasonal maps of precipitation allow us to examine the reasons for this in more detail. Figure 2 shows seasonal maps of precipitation from the TRMM 3B43 observations and RAQMS-Aura. TRMM 3B43 and RAQMS-Aura are well correlated for all seasons, with DJF displaying a correlation of 0.86, MAM a correlation of 0.75, JJA a correlation of 0.71, and SON a correlation of 0.77. These correlations show that the RAQMS-Aura reanalysis broadly captures the seasonal changes in the spatial pattern of tropical precipitation.




Figure 2. Seasonal mean precipitation for TRMM 3B43 (a, c, e, g) and RAQMS-Aura (b, d, f, h).

Precipitation over land in South America and Africa is consistently overestimated relative to TRMM 3B43 by 0.2-0.3 mm/hour. This overestimation over land is a long-standing bias of the dynamical component of RAQMS (Schaack et al, 2006). RAQMS-Aura overestimates precipitation in the Gulf of Mexico and Caribbean by >0.3 mm/hour during JJA and SON. During DJF and MAM, the average bias over the Gulf of Mexico is less than +/- 0.1 mm/hour. RAQMS-Aura overestimates precipitation over the Caribbean by ~0.14 mm/hour during DJF and by ~0.16 during MAM. RAQMS-Aura overestimates precipitation near India by >0.3 mm/hour during MAM and JJA. In the northwest Pacific, RAQMS-Aura



185 shows larger overestimates of precipitation in JJA and SON relative to DJF and JJA, with overestimates relative to TRMM of 0.05 mm/hour in DJF, >0.3 mm/hour in JJA, 0.15 mm/hour in MAM, and >0.3 mm/hour in SON.

RAQMS-Aura does capture precipitation features like the ITCZ and western North Atlantic storm track well, though there is bias in the precipitation amount. RAQMS-Aura underestimates precipitation in the western North Atlantic off the east coast of the US along a storm track region by 0.17 mm/hour in DJF, ~0.15 mm/hour in JJA, ~0.15 mm/hour in MAM, and ~0.17 mm/hour in SON. During DJF, precipitation is overestimated by 0.2-0.3 mm/hour in RAQMS-Aura in the Southern Hemisphere maximum over the Pacific and off the northern coast of Australia. The strength of the SH maximum is consistently overestimated by RAQMS-Aura, as it is higher than TRMM 3B43 by ~0.1 mm/hour in JJA, 0.25-0.3 mm/hour in MAM, and ~0.1 mm/hour in SON. RAQMS-Aura tends to underestimate the strength of the ITCZ in all seasons, with a small underestimate of ~0.05mm/hour in MAM and ~0.15 mm/hour in DJF. RAQMS-Aura underestimates the ITCZ over the east and central Pacific by a max of ~0.25 mm/hour in SON and JJA.

3.1.3 Time Series

The comparison of TRMM 3B43 precipitation and RAQMS-Aura indicates that RAQMS-Aura captures the expected seasonality in the ITCZ and over landmasses though tends to overestimate convective precipitation. Following this characterization of regional biases in RAQMS-Aura, we look closer at how the RAQMS-Aura represents precipitation within the tropics by evaluating the time series for 3 key regions, which are defined in figure 3. The region over the maritime continent is defined by broadscale ascent in the average Walker Circulation. Time series for the maritime continent, NH ITCZ, and SH maximum regions are displayed in figures 4, 5, and 6 respectively.





Figure 3. Regions for timeseries overlaid on mean 2006-2016 TRMM precipitation.

205 Over the maritime continent, RAQMS-Aura has a correlation of 0.619 with TRMM and a mean bias of 0.064 mm/hour (22.27%). The bias between TRMM and RAQMS-Aura is initially higher, ~0.2 mm/hour at a max, then decreases after 2010 within this region. There is also an increased bias in 2015 and late 2016 over the maritime continent. Across the ITCZ in the



northern hemisphere RAQMS-Aura has a correlation of 0.715 and bias of -0.0115 mm/hour (-4.90%) with TRMM. Prior to 2010 RAQMS-Aura displays a small bias relative to TRMM 3B43. Post 2010 RAQMS-Aura underestimates peak
210 precipitation, though the correlation of the measurements with TRMM 3B43 slightly increases to 0.774 within this region. Within a section of the SH precipitation maximum, RAQMS-Aura has a correlation of 0.599 and bias of 0.038 mm/hour (13.53%) with TRMM. The good correlation and bias of less than 25% for each region indicate that RAQMS-Aura has skill in reproducing the observed precipitation in the regions of interest for this study. Shifts in bias observed between 2009 and 2011 appear tied to upgrades to the GDAS system. Changes to GDAS implemented in 2009 included use of variational
215 quality control in the assimilation system and flow dependent reweighting of background error variance (MODEL CHANGES SINCE 1991, 2023).



220 **Figure 4. Mean precipitation over the Maritime continent for TRMM 3B43 and RAQMS-Aura Precipitation. RAQMS-Aura precipitation is on average biased 0.064 mm/hour (22.27%) higher than TRMM 3B43. RAQMS-Aura and TRMM 3B43 have a correlation of 0.619.**





Figure 5. Mean precipitation in NH ITCZ region RAQMS-Aura precipitation is on average biased 0.012 mm/hour (4.90%) lower than TRMM 3B43. RAQMS-Aura and TRMM 3B43 have a correlation of 0.715.



225 Figure 6. Mean precipitation in SH maximum precipitation region. RAQMS-Aura precipitation is on average biased 0.038 mm/hour (13.53%) higher than TRMM 3B43. RAQMS-Aura and TRMM 3B43 have a correlation of 0.599.

3.2 ENSO Composites


Based on comparison of RAQMS-Aura total precipitation with TRMM 3B43 we conclude that RAQMS-Aura reasonably reproduces convection over the Pacific Ocean, particularly within the ITCZ. To characterize the anomaly associated with



230 ENSO, composites for El Niño and La Niña periods are generated for precipitation, convective mass flux, diabatic heating, ozone concentration, carbon monoxide, and net ozone production from monthly mean RAQMS-Aura analyses.

3.2.1 Precipitation

Composites of the de-seasonalized anomaly in precipitation for TRMM and RAQMS-Aura for positive ENSO and negative ENSO are given in figure 7. The TRMM and RAQMS-Aura composites are strongly correlated, with a correlation of 0.77
 235 for El Niño composites and 0.739 for the La Niña composites. The dominant feature of the El Niño phase in the TRMM data and RAQMS-Aura reanalysis is an enhancement of precipitation in the tropics east from 150°E to the western coast of Central America and suppressed precipitation over the maritime continent. RAQMS-Aura however diverges from observations by displaying suppression of precipitation in regions around 7.5°S-39°S, 150°W-120°W and 7.5°N-20°N, 150°E-180°E where precipitation is enhanced in TRMM. During the La Niña phase, precipitation is suppressed over the
 240 central Pacific and enhanced over the maritime continent. For both TRMM and RAQMS-Aura the El Niño and La Niña composites are near mirrors of one another, with the location of the maximum change shifted west during the negative phase from the positive phase.



245 **Figure 7. Composited precipitation anomalies for El Niño in RAQMS-Aura (a) and TRMM 3B43 (b) and La Niña in RAQMS-Aura (c) and TRMM 3B43 (d).**


3.2.2 Response of Tropospheric Total Column Ozone and Carbon Monoxide Column to ENSO

ENSO composites for OMI-MLS TOR (Ziemke et al., 2006) and Measurements of Pollution in the Troposphere (MOPITT) CO (Emmons et al., 2004) are used to confirm the representativeness of RAQMS-Aura ENSO chemical signals. The OMI-



250 MLS TOR is a satellite residual product where total ozone columns from the OMI instrument and stratospheric columns from MLS instrument (both on-board the Aura satellite) are combined to infer the tropospheric ozone column. Both the OMI-MLS TOR and the MOPITT CO data used are monthly mean Level 3 products.

Tropical tropospheric ozone column (TTOC) anomalies in RAQMS-Aura and the OMI-MLS TOR for the positive and negative phases of ENSO are shown in figure 8. TTOC anomalies are 1-2 DU larger during the positive phase of ENSO than in the negative phase. Within both the RAQMS-Aura TTOC and OMI-MLS TOR, El Niño is associated with an increase over the maritime continent and a decrease over the central and eastern Pacific Ocean. The decrease over the Pacific Ocean is flanked by increased concentrations to the north and south. Outside of the Pacific region, the tropospheric column anomaly associated with the ENSO phase is less than 1 DU. During La Niña, a small decrease in tropospheric ozone occurs over the maritime continent while an increase occurs over the central-eastern Pacific. The location of the peak decrease in TTOC in the eastern Pacific depicted in the El Niño composite is comparable to that found by Oman et al. 2011 and Olsen, Wargan, and Pawson 2016, while in the earlier studies of Peters et al., 2001, Doherty et al., 2006, and Ziemke and Chandra, 2003 this peak decrease in TTOC is more towards the southeast. As our analysis is consistent with observations, the differences from earlier analyses are likely due to variability in ENSO and the influence of the large 2015 El Niño event during the 2006-2016 period under consideration in this study.



265 **Figure 8. Composited TTOC anomalies associated with El Niño in RAQMS-Aura (a) and OMI-MLS TOR (c) and La Niña in RAQMS-Aura(b) and OMI-MLS TOR(d)**

CO column anomalies for RAQMS-Aura and MOPITT are presented in figure 9. MOPITT CO anomalies appear noisier due to the sparse spatial sampling of the MOPITT instrument. RAQMS-Aura reproduces ENSO-related variability in CO as observed by MOPITT with both El Niño and La Niña composites having a correlation of 0.850. RAQMS-Aura CO column is on average increased across the tropics during El Niño, with stronger enhancements of $0.4 \times 10^{18} \text{ mol/cm}^2$ observed over the maritime continent. Enhanced CO over the maritime continent is tied to enhanced biomass burning during El Niño as precipitation is suppressed, increasing fuel aridity, and thereby increasing susceptibility to fire (Yin et al., 2016; Reid et al.,



2013; van der Werf et al., 2017). RAQMS-Aura CO column decreases over the maritime continent during La Niña and is enhanced over South America. During La Niña, rainfall is enhanced over the maritime continent, resulting in CO decreases as fires are suppressed.




Figure 9. Composit ed CO column anomalies associated with El Niño in RAQMS-Aura (a) and MOPITT (c) and La Niña in RAQMS-Aura(b) and MOPITT(d).


3.2.3 Vertical structure of tropospheric response to ENSO

280 As this study utilizes reanalysis data, we can provide further context to the patterns in TTOC and CO columns. In particular, we explore how the vertical structure of convective mass flux, large-scale diabatic heating, and ozone production/loss terms respond to ENSO. Meridionally averaged vertical profile cross sections are calculated between 7.5°S and 2.5°N. This latitude band was selected as it cuts across the maximum and minimum precipitation anomalies associated with ENSO (fig 7) and for consistency with the cross-sections analyzed by Doherty et al. 2006.

285 Convective mass flux anomalies between 7.5°S and 2.5°N for the positive and negative phases of ENSO are presented in Figure 10. The strongest convective mass flux anomaly is over the Pacific Ocean during both the positive and negative phase of ENSO. This strong convective mass flux anomaly is also where the absolute maximum precipitation anomaly occurs, which is expected given the dominance in convective precipitation in this region. Diabatic heating anomalies presented in figure 11 are qualitatively similar to the convective mass flux ENSO anomalies. This is because the majority of the diabatic heating in this region is associated with the large-scale response to sub-grid-scale convective precipitation. The convective mass flux and diabatic heating anomalies during El Niño indicate decreased upward vertical transport over the maritime continent where precipitation is suppressed and increased upward vertical transport over the central Pacific where precipitation is enhanced. Conversely, the convective mass flux and diabatic heating anomalies during La Niña both indicate enhanced vertical transport over the maritime continent and increased downward vertical transport over the central Pacific.



295 In Doherty et al 2006 and Sudo and Takahashi 2001 the positive and negative mass flux anomalies are of similar magnitudes while here the negative flux anomaly over Micronesia is $\frac{1}{2}$ - $\frac{1}{3}$ the strength of the anomaly over the central-eastern Pacific. This may be a consequence of the high bias in precipitation over Micronesia in the RAQMS-Aura reanalysis, as the precipitation anomaly El Niño composite indicates that precipitation is not suppressed as much as in observations over the region. However, these differences in the strength of the vertical motion anomalies are consistent with the ENSO precipitation anomaly over the central Pacific being larger than that of the anomaly over the maritime continent in TRMM observations and RAQMS-Aura analyses. The precipitation and mass flux anomaly patterns display suppressed (enhanced) vertical motion over the Pacific and enhanced (suppressed) vertical motion over the maritime continent during the negative (positive) phase.



305 **Figure 10. RAQMS-Aura convective mass flux (CMFLX) anomalies for a) positive and b) negative ENSO phases.**





Figure 11. RAQMS-Aura diabatic heating anomalies (colors) and theta (contours) for a) positive and b) negative ENSO phases.




Ozone anomaly cross-sections associated with ENSO are presented in Figure 12. During El Niño the tropospheric ozone anomaly extends across the depth of the troposphere over the maritime continent, with two distinct stronger (>3 ppbv) enhancements above 550 hPa and below 700 hPa. Over the central Pacific (from 160°E to 140°W) where the convective mass flux is enhanced in the El Niño composite through the depth of the troposphere, a decrease in the ozone concentration of 3-5 ppbv occurs. The lower troposphere enhancement over the maritime continent is accompanied by a positive anomaly in net O₃ production (fig 14a), indicating that some of the enhancement in TTOC over the maritime continent during El Niño is due to enhancement in chemical production and not solely due to shifts in the circulation pattern. The El Niño ozone anomaly cross-section is <1 ppbv throughout the majority of the troposphere off the South American Coast, indicating that the TTOC decrease is due to the decreased (>9 ppbv) concentrations near the tropopause, above 200 hPa. The La Niña ozone anomaly cross-section section shows enhancement in ozone over the central Pacific and decrease over the maritime continent. Over the maritime continent a distinct stronger (>2 ppbv) decrease is seen below 700 hPa and above 350 hPa.



320 **Figure 12. Anomalies in RAQMS-Aura ozone profiles below the tropopause associated with a) El Niño and b) La Niña.**

CO anomaly cross-sections for each ENSO phase are presented in figure 13. Tropical CO is anomalously high during El Niño and anomalously low during La Niña. Tropical CO is enhanced over the maritime continent during El Niño throughout the tropical troposphere, with the strongest enhancement near the surface indicative of a strong increase in biomass burning emissions. The near-surface enhancements in CO over South America and Africa during El Niño are also likely tied to CO emissions from biomass burning, though these enhancements are not spread through the depth of the troposphere as occurs over the maritime continent. The negative CO anomalies associated with La Niña are largest over the maritime continent and present through the depth of the troposphere. The enhancement in CO Column over South America associated with La Niña is not present in the La Niña vertical cross-section as it is to the south of the latitudes used to generate the cross-section composite.



330

Figure 13. Anomalies in RAQMS-Aura CO profiles below the tropopause associated with a) El Niño and b) La Niña.

Net ozone production (production-loss terms) anomalies are presented in Figure 14. The largest anomalies are closest to the surface and below 700 hPa. The change in net ozone production is smaller in La Niña than El Niño. Enhanced production of 2-3 ppbv/day is found over central Africa, Indonesia, and the Amazon rainforest in Brazil. These regions show reductions of ~ 1.3 ppbv/day in ozone production in the La Niña composite. El Niño is known to increase fire emissions in Indonesia as a consequence of the decreased rainfall over the region (Field et al., 2016; Park et al., 2021), and so the increased production of ozone during El Niño captured by RAQMS-Aura is likely to be partially due to enhanced chemical production of ozone in biomass burning plumes. Enhanced production during El Niño occurs over all 3 biomass burning regions but only the maritime continent shows a significant (>4 ppbv) enhancement in O₃ below 700 hPa. In contrast, the enhanced production over South America and Africa is associated with weak (<2 ppbv) ozone enhancement. The average winds below 800hPa during El Niño over South America (not shown) are northeasterly, resulting in transport of the ozone associated with biomass burning to the south and out of the latitudes included in the cross-section (7.5°S to 2.5°N). Over the maritime continent, the average winds below 750 hPa are southerly and decline in strength through the cross-section. Based on these wind patterns, ozone associated with biomass burning over the maritime continent experiences less meridional transport and influences the ozone profile within this meridional cross-section.

335

340

345





Figure 14. Anomalies in RAQMS-Aura net O₃ production associated with a) El Niño and b) La Niña.

3.3 EOF Analysis

In addition to composite analysis, Empirical Orthogonal Function (EOF) analysis is used to investigate the role played by ENSO in TTOC variability. The first EOF of TTOC has been previously found to be associated with ENSO, while TTOC EOFs 2 and 3 are uncorrelated with ENSO (Doherty et al., 2006; Sekiya and Sudo, 2012). ENSO positive and negative phases are near opposites of each other, and so it is reasonable to expect that much of the variability associated with ENSO can be captured with a single EOF. The EOF spatial patterns are displayed for TTOC, precipitation, and CO column in figures 15, 16, and 17. PC time series are presented in figure 18, alongside the Niño 3.4 index for reference.

3.3.1 EOFs

EOF patterns for TTOC are displayed in figure 15. The TTOC PC₁ has a correlation of 0.747 with the Niño 3.4. The associated EOF indicates a 2-2.5 DU enhancement over the maritime continent and a 1.6-2 DU decrease over the Pacific (figure 15a). EOF₁ captures similar features to those in the El Niño TTOC composite, though the enhancement in TTOC near Vietnam is weaker relative to the enhancement near Indonesia in the EOF compared to the composite. TTOC PC₂ and PC₃ are weakly correlated with the Niño 3.4 index, with correlations of -0.144 and -0.209 respectively. TTOC EOF₂ explains around half as much variance as TTOC EOF₁ and shows a wave 1 like pattern with a peak in the northeast Pacific. TTOC EOF₃ accounts for changes of less than 1 DU on average, and a maximum near 3 DU. At the most, this is ~10% of the mean TTOC and less than 1% on average. TTOC EOF₃ captures an increase across the equatorial Pacific and decreases elsewhere.




365 **Figure 15. Patterns for RAQMS-Aura TTOC EOF 1-3, scaled by 1 standard deviation of the associated PC. EOF₁ explains 17.20% of the non-seasonal variance in TTOC, EOF₂ explains 8.70% and EOF₃ explains 6.00%.**

370 EOF patterns for total precipitation are displayed in figure 16. The precipitation PC₁ is strongly correlated with the Niño 3.4 index, with a correlation of 0.870, as well as a strong correlation with the TTOC PC₁ (0.818). The associated EOF pattern is similar to the El Niño precipitation composite in figure 7a, though the magnitude of the decreased precipitation in the western Pacific relative to the enhancement in the central Pacific is smaller than in the composite. Precipitation EOFs 2 and 3 combined capture a similar amount of the variability in precipitation as EOF₁ alone. Their PCs are not correlated with the Niño 3.4 index, with a PC₂ correlation of -0.02, and a PC₃ correlation of -0.093. The EOF₂ pattern depicts a small, localized enhancement in the central southern Pacific ocean, slightly stronger enhancements of ~0.06 mm/hour in the Caribbean and



375 NW equatorial Pacific, and decreased precipitation in the remainder of the northern hemisphere Pacific. The EOF₃ pattern accounts for changes of <math><0.03</math> mm/hour on average. The largest of these small changes are a decrease in precipitation in the central Pacific to the east of where the maximum precipitation anomaly associated with ENSO is located. Precipitation PC₃ has a correlation of 0.695 with TTOC PC₂, indicating there is some co-variability between the two.




380 **Figure 16. Patterns for RAQMS-Aura total precipitation EOF 1-3, scaled by 1 standard deviation of the associated PC. EOF1 explains 8.33% of the non-seasonal variance in total precipitation, EOF2 explains 4.73% and EOF3 explains 4.46%.**

EOF patterns for CO column are displayed in figure 17. Interannual variability in tropical CO has been shown to be predominately influenced by biomass burning emissions (Rowlinson et al., 2019). All 3 CO column EOF patterns appear to be heavily influenced by extreme biomass burning events, as the strongest changes are over the maritime continent and



385 South America and the peaks in the PCs correspond with years with enhanced biomass burning in the regions highlighted by the largest values in the EOF (eg. van der Werf et al., 2017). CO PC amplitude peaks are larger than 2 for PC₁ in late 2015; PC₂ in 2006, 2007, 2010, and 2015; and PC₃ in 2006, 2015, and 2016 (figure 18). EOF₁ explains 46.96% of the non-seasonal variance in CO, while EOF₂ explains 9.46% and EOF₃ explains 6.48%.



390 **Figure 17.** Patterns for RAQMS-Aura CO Column EOF 1-3, scaled by 1 standard deviation of the associated PC. EOF₁ explains 46.96% of the non-seasonal variance in CO column, EOF₂ explains 9.46% and EOF₃ explains 6.48%.




Figure 18. Timeseries of PC₁ (a), PC₂ (b), and PC₃ (c) for TTOC, total precipitation, and CO Column. Niño 3.4 Index time series included for reference.

Most variability in CO columns from 2006-2016 is explained by EOF₁. The physical pattern is indicative of a tropics-wide decrease (increase) in CO, with the peak change of $\sim 0.3 \times 10^{18}$ mol/cm² centered over the maritime continent. CO PC₁ has a correlation of -0.399 with the Niño 3.4 index, which indicates an ENSO influence on CO variability. Additionally, CO PC₁ is correlated with precipitation PC₁ (-0.435), suggesting that ENSO related changes in precipitation contribute to the ENSO driven CO variability. This is consistent with precipitation influences on biomass burning. The CO EOF₂ pattern shows CO column enhancements over Brazil and decreases over the maritime continent. CO PC₂ has a correlation of -0.297 with the Niño 3.4 index, and correlation of -0.435 with TTOC PC₁, suggesting that ENSO related changes in CO contribute to ENSO



driven TTOC variability. EOF₃ pattern again highlights the maritime continent and Brazil varying together, with an opposing change in CO across the Pacific. CO PC₃ displays a correlation of -0.145 with the Niño 3.4 index.

3.3.2 Multiple Linear Regression reconstruction of TTOC PC₁

405 From the composite analyses we are able to show that ENSO related shifts in precipitation correspond with changes in vertical motion, CO concentration, net ozone production, and tropospheric ozone concentrations. The composite analysis also indicates that some of the enhancement in TTOC over the maritime continent during El Niño is due to enhanced production of ozone from biomass burning emissions. The EOF analysis further links variation in biomass burning to the TTOC variation as CO PCs 1 and 2 are moderately correlated with TTOC and precipitation PC₁. To quantify the relative importance of dynamical and biomass burning variability on ENSO related variability in TTOC, a multiple linear regression analysis is constructed using the principal components. The regression equation is shown in equation (1).

$$PC1_{TTOC} = w_1 PC1_{CO} + w_2 PC2_{CO} + w_3 PC3_{CO} + w_4 PC1_{precip} + e \quad (1)$$

The principal components are from the EOF analysis; w_1 , w_2 , w_3 , w_4 , and e are regression coefficients as determined using a least squares fit. The resulting regression model is shown in equation (2).

$$PC1_{TTOC} = 0.11 * PC1_{CO} - 0.2 * PC2_{CO} + 0.004 * PC3_{CO} + 0.8 * PC1_{precip} - 3.3 \times 10^{-10} \quad (2)$$

415 This multiple PC regression reproduces the PC_{1TTOC} very well, with the regression-based estimate correlating with the original PC_{1TTOC} at 0.85 (fig 19a).




Figure 19. a) TTOC PC₁ from EOF analysis and reconstructed from multiple linear regression. b) Contribution to regression of Precipitation PC₁ and combined contribution of CO PCs 1-3.

420 The strongest weighted PC in the regression is the precipitation PC₁, which is expected given its strong correlation with TTOC PC₁. This supports the result from Doherty et al. 2006 and Inness et al. 2015 that ENSO variability in TTOC is primarily driven by convective transport. The weights for CO PC₁ and PC₂ are also significant, indicating that CO, as a proxy for biomass burning, also contributes to TTOC variability.


A timeseries breaking down the contributions of precipitation PC₁ and the combined CO PCs to the TTOC PC₁ predicted by the regression is shown in figure 19b. The precipitation PC₁ regression contribution is positive during El Niño periods and negative during La Niña periods. The combined regression contribution of the CO PCs shows that variability in CO contributes to ENSO variability in TTOC in an episodic way. As the CO column anomaly is linked to anomalous biomass burning emissions and net ozone production near the surface, it can be concluded that a portion of the ENSO variability in TTOC is due to biomass burning though it is a smaller portion than that linked to the dynamical effects of ENSO.



430 As inferred from the regression, El Niño increases in TTOC over the maritime continent are linked to CO PC₁ enhancements
in CO over the maritime continent while CO PC₂ is associated with enhancements in CO over South America and Africa and
decreases over Indonesia. Timeseries of the CO column and TTOC anomalies (not shown) have a correlation of 0.668 over
the maritime continent and 0.566 over South America. The TTOC and CO anomalies over the maritime continent are
positive during El Niño events and negative during La Niña events. Over South America, the sign of the TTOC and CO
435 anomalies are less consistent with ENSO phase.

3.4 2015/2016 extreme El Niño

Through the satellite era, extreme El Niño events in 1982/1983, 1997/1998, and 2015/2016 have been observed alongside
weak and moderate events. These extreme events have a larger impact on the distribution of TTOC and have a larger
contribution from biomass burning emissions than weaker El Niño events (Doherty et al., 2006; Inness et al., 2015). The
440 2015/2016 extreme El Niño was the strongest El Niño since the 1997/1998 event (Santoso et al., 2017). 2015 and 1997 are
also among the most extreme maritime continent biomass burning events, with 1997 ranking first followed by 2015 in an
analysis of surface visibility at airports in Sumatra and Kalimantan from 1990-2015 (Field et al., 2016). Here we investigate
how the inclusion of the 2015 extreme El Niño influences our interpretation of the importance of biomass burning on TTOC
ENSO variability. As in prior analyses (Chandra et al., 1998, 2009; Sudo and Takahashi, 2001), we focus on October as
445 biomass burning in the maritime continent peaks around October and would have its greatest impact on TTOC around the
same time (Field et al., 2016). In RAQMS-Aura, the CO PC amplitudes have the largest variability in October and the largest
contributions of the CO PCs to the TTOC PC₁ regression occur in October.



450 **Figure 20. RAQMS-Aura October 2015 a) TTOC anomaly, b) convective mass flux anomaly, and c) tropospheric ozone profile anomaly, d) P-L, e) CO.**



The RAQMS-Aura 2015 October TTOC anomaly is shown in figure 20a. This pattern is similar to the October 1997 anomaly in TTOC modeled by Sudo and Takahashi 2001 with an increase over the maritime continent that is 2-3 times stronger than the decrease over the eastern Pacific. However, the peak decrease over the eastern Pacific is more towards the central Pacific during 2015 than in 1997. The maximum increase over the maritime continent is 10-15 DU in October 2015, less than the maximum 20-24 DU increase in October 1997. RAQMS-Aura TTOC increases over South America in October 2015 by 1-4 DU, while the Sudo and Takahashi simulated October 1997 changes by less than 2 DU over South America. These differences over Africa and South America in 2015 versus 1997 are consistent the differences in patterns of convective mass flux. In 2015 mass flux is decreased aloft over Brazil and Africa (fig 20b), while in 1997 changes in mass flux over Brazil and Africa are weaker and are slightly positive (Sudo and Takahashi, 2001). The core of the upward mass flux anomaly over the Pacific is ~30-40 degrees closer to the dateline in 2015.

Over the maritime continent, the ozone concentration anomaly below 650 hPa is stronger than in the 2006-2016 El Niño average (fig 20c). This is linked to stronger ozone production in October 2015 (fig 20d). This enhancement in O₃ production in 2015 is likely due to increased fire activity, as CO column is increased throughout the tropics in 2015 (fig 20e) and the CO anomaly over the maritime continent is more widespread and stronger by $\sim 0.2 \times 10^{18}$ mol/cm² than the 2006-2016 El Niño average. There is also an enhancement in CO, ozone, and net ozone production over South America in October 2015 relative to the 2006-2016 El Niño composite. This shows that the biomass burning activity in 2015 was anomalous compared to the other El Niño years included in the RAQMS-Aura reanalysis, with significant burning occurring over both South America and the maritime continent.

4 Conclusions

The RAQMS-Aura reanalysis captures observed ENSO variability in TTOC, CO, and precipitation. ENSO composites of tropospheric ozone, carbon monoxide, convective mass flux, diabatic heating, and ozone net chemical production show that the observed ENSO signatures in TTOC result from a combination of convective redistribution and variability in production of ozone from biomass burning emissions, which are modulated by ENSO variability in precipitation. The location of the peak decrease in TTOC resulting from increased vertical motion in the eastern Pacific depicted in the El Niño composite found by this study is comparable to other studies of TTOC variability in the 2000s and 2010s (Oman et al., 2011; Olsen et al., 2016). The location of the peak decrease in TTOC contrasts with that found by analyses of 1970s-2000 where it is more towards the southeast and near the South American coast (Peters et al., 2001; Doherty et al., 2006; Ziemke and Chandra, 2003). The RAQMS-Aura El Niño TTOC composite is in agreement with the El Niño composite OMI-MLS TOR observations, and the analysis of convective flux indicates that the ozone decreases over the central Pacific are due to enhanced vertical motion. Therefore, we believe the difference in position of the peak decrease in TTOC is due to characteristics of El Niño during our analysis period. El Niño events from 2006-2016 were predominately El Niño Modoki events, while El Niño events between 1979 and 2002 display greater variability in type of El Niño and includes more



canonical ENSO events (Hou et al., 2016; Santoso et al., 2017; Lee and McPhaden, 2010). The ascending branches of Walker circulation cell is over the central Pacific during El Niño Modoki (Ashok et al., 2007), while during canonical El Niño the ascending branch is over the eastern Pacific. Since TTOC is decreased where vertical motion is enhanced during ENSO and increased where vertical motion is suppressed, it is expected that under El Niño Modoki conditions the largest decrease in TTOC will be in the central Pacific with TTOC increases in the western and eastern Pacific. This response of TTOC to El Niño Modoki is shown by Hou et al. 2016 and is in-line with the El Niño RAQMS-Aura TTOC anomaly composite calculated by this study (Fig 9a).

The strongest ENSO variability in tropospheric ozone is shown to occur near the tropopause. An enhancement in ozone below 700 hPa during El Niño occurs over the maritime continent that is dependent on the magnitude of the biomass burning emissions. The EOF analyses and multiple linear regression further indicate that ENSO variability in TTOC is driven by shifts in the location of the ascending and descending branches of the Walker circulation. The EOF and multiple linear regression analyses also indicate that variability in biomass burning, as inferred from CO anomalies, contributes to ENSO variability in TTOC. During the 2015/2016 strong El Niño event TTOC, CO, and convective mass flux anomalies were stronger than in the weaker ENSO events captured by the RAQMS-Aura reanalysis. The 2015 CO concentrations align with the mode captured by CO EOF₁ while the other El Niño years in our analysis align with the mode in CO EOF₂. Biomass burning enhanced TTOC and CO anomalies occurred over both South America and the maritime continent in October 2015 in contrast to the other El Niño years between 2006 and 2016 where biomass burning enhanced TTOC and CO was only found over the maritime continent.

Code and data availability

The software used to generate the figures in this work and all raw data can be provided upon request by the corresponding author.

Author Contributions

Bruckner and Pierce planned the study and collaborated on the analysis. Lenzen performed the RAQMS-Aura reanalysis. Bruckner wrote the paper. Pierce reviewed and edited the paper.

Competing Interests

The contact author has declared that none of the authors has any competing interests.



5 References

- 510 Al-Saadi, J., Soja, A., Pierce, R. B., Szykman, J., Wiedinmyer, C., Emmons, L., Kondragunta, S., Zhang, X., Kittaka, C., Schaack, T., and Bowman, K.: Intercomparison of near-real-time biomass burning emissions estimates constrained by satellite fire data, <https://doi.org/10.1117/1.2948785>, 2008.
- MODEL CHANGES SINCE 1991: https://www.emc.ncep.noaa.gov/gmb/STATS/html/model_changes.html, last access: 11 October 2023.
- 515 Ashok, K., Behera, S. K., Rao, S. A., Weng, H., and Yamagata, T.: El Niño Modoki and its possible teleconnection, *J. Geophys. Res. Oceans*, 112, <https://doi.org/10.1029/2006JC003798>, 2007.
- Bamston, A. G., Chelliah, M., and Goldenberg, S. B.: Documentation of a highly ENSO-related sst region in the equatorial pacific: Research note, *Atmosphere-Ocean*, 35, 367–383, <https://doi.org/10.1080/07055900.1997.9649597>, 1997.
- Boersma, K. F., Eskes, H. J., Veeffkind, J. P., Brinksma, E. J., van der A, R. J., Sneep, M., van den Oord, G. H. J., Levelt, P. F., Stammes, P., Gleason, J. F., and Bucsela, E. J.: Near-real time retrieval of tropospheric NO₂ from OMI, *Atmospheric Chem. Phys.*, 7, 2103–2118, <https://doi.org/10.5194/acp-7-2103-2007>, 2007.
- 520 Bucsela, E. J., Krotkov, N. A., Celarier, E. A., Lamsal, L. N., Swartz, W. H., Bhartia, P. K., Boersma, K. F., Veeffkind, J. P., Gleason, J. F., and Pickering, K. E.: A new stratospheric and tropospheric NO₂ retrieval algorithm for nadir-viewing satellite instruments: Applications to OMI, *Atmospheric Meas. Tech.*, 6, 2607–2626, <https://doi.org/10.5194/AMT-6-2607-2013>, 2013.
- 525 Chandra, S., Ziemke, J. R., Min, W., and Read, W. G.: Effects of 1997–1998 El Niño on tropospheric ozone and water vapor, *Geophys. Res. Lett.*, 25, 3867–3870, <https://doi.org/10.1029/98GL02695>, 1998.
- Chandra, S., Ziemke, J. R., Bhartia, P. K., and Martin, R. V.: Tropical tropospheric ozone: Implications for dynamics and biomass burning, *J. Geophys. Res. Atmospheres*, 107, ACH 3-1-ACH 3-17, <https://doi.org/10.1029/2001JD000447>, 2002.
- 530 Chandra, S., Ziemke, J. R., Duncan, B. N., Diehl, T. L., Livesey, N. J., and Froidevaux, L.: Effects of the 2006 El Niño on tropospheric ozone and carbon monoxide: implications for dynamics and biomass burning, *Atmospheric Chem. Phys.*, 9, 4239–4249, <https://doi.org/10.5194/acp-9-4239-2009>, 2009.
- Chen, D. and Dai, A.: Precipitation Characteristics in the Community Atmosphere Model and Their Dependence on Model Physics and Resolution, *J. Adv. Model. Earth Syst.*, 11, 2352–2374, <https://doi.org/10.1029/2018MS001536>, 2019.
- 535 Chen, D., Dai, A., and Hall, A.: The Convective-To-Total Precipitation Ratio and the “Drizzling” Bias in Climate Models, *J. Geophys. Res. Atmospheres*, 126, e2020JD034198, <https://doi.org/10.1029/2020JD034198>, 2021.
- Dentener, F., Keating, T., and Akimoto, H. (Eds.): in: Hemispheric Transport of Air Pollution 2010: Part A: Ozone and Particulate Matter, Air Pollution Studies No.17, United Nations, New York, NY, 2010.
- 540 Doherty, R. M., Stevenson, D. S., Johnson, C. E., Collins, C. E., and Sanderson, M. G.: Tropospheric ozone and El Niño–Southern Oscillation: Influence of atmospheric dynamics, biomass burning emissions, and future climate change, *J. Geophys. Res. Atmospheres*, 111, 19304, <https://doi.org/10.1029/2005JD006849>, 2006.
- East, J. D., Henderson, B. H., Napelenok, S. L., Koplitz, S. N., Sarwar, G., Gilliam, R., Lenzen, A., Tong, D. Q., Pierce, R. B., and Garcia-Menendez, F.: Inferring and evaluating satellite-based constraints on NO_x emissions estimates in air quality simulations, *Atmospheric Chem. Phys.*, 22, 15981–16001, <https://doi.org/10.5194/acp-22-15981-2022>, 2022.



- 545 Emmons, L. K., Deeter, M. N., Gille, J. C., Edwards, D. P., Attié, J.-L., Warner, J., Ziskin, D., Francis, G., Khattatov, B., Yudin, V., Lamarque, J.-F., Ho, S.-P., Mao, D., Chen, J. S., Drummond, J., Novelli, P., Sachse, G., Coffey, M. T., Hannigan, J. W., Gerbig, C., Kawakami, S., Kondo, Y., Takegawa, N., Schlager, H., Baehr, J., and Ziereis, H.: Validation of Measurements of Pollution in the Troposphere (MOPITT) CO retrievals with aircraft in situ profiles, *J. Geophys. Res. Atmospheres*, 109, <https://doi.org/10.1029/2003JD004101>, 2004.
- 550 Field, R. D., Werf, G. R. van der, Fanin, T., Fetzner, E. J., Fuller, R., Jethva, H., Levy, R., Livesey, N. J., Luo, M., Torres, O., and Worden, H. M.: Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought, *Proc. Natl. Acad. Sci.*, 113, 9204–9209, <https://doi.org/10.1073/PNAS.1524888113>, 2016.
- Fishman, J. and Balok, A. E.: Calculation of daily tropospheric ozone residuals using TOMS and empirically improved SBUV measurements: Application to an ozone pollution episode over the eastern United States, *J. Geophys. Res. Atmospheres*, 104, 30319–30340, <https://doi.org/10.1029/1999JD900875>, 1999.
- 555 Fishman, J. and Larsen, J. C.: Distribution of total ozone and stratospheric ozone in the tropics: Implications for the distribution of tropospheric ozone, *J. Geophys. Res. Atmospheres*, 92, 6627–6634, <https://doi.org/10.1029/JD092iD06p06627>, 1987.
- Fishman, J., Watson, C. E., Larsen, J. C., and Logan, J. A.: Distribution of tropospheric ozone determined from satellite data, *J. Geophys. Res. Atmospheres*, 95, 3599–3617, <https://doi.org/10.1029/JD095iD04p03599>, 1990.
- 560 Fishman, J., Hoell Jr., J. M., Bendura, R. D., McNeal, R. J., and Kirchhoff, V. W. J. H.: NASA GTE TRACE A experiment (September–October 1992): Overview, *J. Geophys. Res. Atmospheres*, 101, 23865–23879, <https://doi.org/10.1029/96JD00123>, 1996.
- Fishman, J., Wozniak, A. E., and Creilson, J. K.: Global distribution of tropospheric ozone from satellite measurements using the empirically corrected tropospheric ozone residual technique: Identification of the regional aspects of air pollution, *Atmospheric Chem. Phys.*, 3, 893–907, <https://doi.org/10.5194/acp-3-893-2003>, 2003.
- 565 Fishman, J., Creilson, J. K., Wozniak, A. E., and Crutzen, P. J.: Interannual variability of stratospheric and tropospheric ozone determined from satellite measurements, *J. Geophys. Res. Atmospheres*, 110, <https://doi.org/10.1029/2005JD005868>, 2005.
- 570 Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sunnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113, D15S20, <https://doi.org/10.1029/2007JD008771>, 2008.
- 575 Hack, J. J.: Parameterization of moist convection in the National Center for Atmospheric Research community climate model (CCM2), *J. Geophys. Res. Atmospheres*, 99, 5551–5568, <https://doi.org/10.1029/93JD03478>, 1994.
- Haines, D. A.: A lower atmosphere severity index for wildlife fires, *Natl. Weather Dig.*, 13, 23–27, 1989.
- 580 Hou, X., Zhu, B., Fei, D., Zhu, X., Kang, H., and Wang, D.: Simulation of tropical tropospheric ozone variation from 1982 to 2010: The meteorological impact of two types of ENSO event, *J. Geophys. Res. Atmospheres*, 121, 9220–9236, <https://doi.org/10.1002/2016JD024945>, 2016.



- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P., and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, *J. Hydrometeorol.*, 8, 38–55, <https://doi.org/10.1175/JHM560.1>, 2007.
- 585 Huijnen, V., Miyazaki, K., Flemming, J., Inness, A., Sekiya, T., and Schultz, M. G.: An intercomparison of tropospheric ozone reanalysis products from CAMS, CAMS interim, TCR-1, and TCR-2, *Geosci. Model Dev.*, 13, 1513–1544, <https://doi.org/10.5194/gmd-13-1513-2020>, 2020.
- Inness, A., Benedetti, A., Flemming, J., Huijnen, V., Kaiser, J. W., Parrington, M., and Remy, S.: The ENSO signal in atmospheric composition fields: emission-driven versus dynamically induced changes, *Atmospheric Chem. Phys.*, 15, 9083–9097, <https://doi.org/10.5194/acp-15-9083-2015>, 2015.
- 590 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmospheric Chem. Phys.*, 19, 3515–3556, <https://doi.org/10.5194/acp-19-3515-2019>, 2019.
- 595 Kiehl, J. T., Hack, J. J., Bonan, G. B., Boville, B. A., Williamson, D. L., and Rasch, P. J.: The National Center for Atmospheric Research Community Climate Model: CCM3, *J. Clim.*, 11, 1131–1149, [https://doi.org/10.1175/1520-0442\(1998\)011<1131:TNCFAR>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<1131:TNCFAR>2.0.CO;2), 1998.
- Kim, S. T. and Yu, J.-Y.: The two types of ENSO in CMIP5 models, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL052006>, 2012.
- 600 Kleist, D. T., Parrish, D. F., Derber, J. C., Treadon, R., Wu, W.-S., and Lord, S.: Introduction of the GSI into the NCEP Global Data Assimilation System, *Weather Forecast.*, 24, 1691–1705, <https://doi.org/10.1175/2009WAF2222201.1>, 2009.
- Larkin, N. K. and Harrison, D. E.: On the definition of El Niño and associated seasonal average U.S. weather anomalies, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2005GL022738>, 2005.
- Lee, T. and McPhaden, M. J.: Increasing intensity of El Niño in the central-equatorial Pacific, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL044007>, 2010.
- 605 McMillan, W. W., Barnet, C., Strow, L., Chahine, M. T., McCourt, M. L., Warner, J. X., Novelli, P. C., Korontzi, S., Maddy, E. S., and Datta, S.: Daily global maps of carbon monoxide from NASA's Atmospheric Infrared Sounder, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2004GL021821>, 2005.
- McPeters, R., Kroon, M., Labow, G., Brinksma, E., Balis, D., Petropavlovskikh, I., Veefkind, J. P., Bhartia, P. K., and Levelt, P. F.: Validation of the Aura Ozone Monitoring Instrument total column ozone product, *J. Geophys. Res. Atmospheres*, 113, <https://doi.org/10.1029/2007JD008802>, 2008.
- 610 McPhaden, M. J., Zebiak, S. E., and Glantz, M. H.: ENSO as an Integrating Concept in Earth Science, *Science*, 314, 1740–1745, <https://doi.org/10.1126/science.1132588>, 2006.
- Miyazaki, K., Bowman, K., Sekiya, T., Eskes, H., Boersma, F., Worden, H., Livesey, N., Payne, V. H., Sudo, K., Kanaya, Y., Takigawa, M., and Ogochi, K.: Updated tropospheric chemistry reanalysis and emission estimates, TCR-2, for 2005–2018, *Earth Syst. Sci. Data*, 12, 2223–2259, <https://doi.org/10.5194/essd-12-2223-2020>, 2020.
- Olsen, M. A., Wargan, K., and Pawson, S.: Tropospheric column ozone response to ENSO in GEOS-5 assimilation of OMI and MLS ozone data, *Atmospheric Chem. Phys.*, 16, 7091–7103, <https://doi.org/10.5194/acp-16-7091-2016>, 2016.



- 620 Oman, L. D., Ziemke, J. R., Douglass, A. R., Waugh, D. W., Lang, C., Rodriguez, J. M., and Nielsen, J. E.: The response of tropical tropospheric ozone to ENSO, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL047865>, 2011.
- Park, M., Worden, H. M., Kinnison, D. E., Gaubert, B., Tilmes, S., Emmons, L. K., Santee, M. L., Froidevaux, L., and Boone, C. D.: Fate of Pollution Emitted During the 2015 Indonesian Fire Season, *J. Geophys. Res. Atmospheres*, 126, e2020JD033474, <https://doi.org/10.1029/2020JD033474>, 2021.
- 625 Peters, W., Krol, M., Dentener, F., and Lelieveld, J.: Identification of an El Niño-Southern Oscillation signal in a multiyear global simulation of tropospheric ozone, *JOURNAL OF GEOPHYSICAL RESEARCH*, <https://doi.org/10.1029/2000JD900658>, 2001.
- Pierce, R. B., Schaack, T., Al-Saadi, J. A., Fairlie, T. D., Kittaka, C., Lingenfelter, G. S., Natarajan, M., Olson, J. R., Soja, A. J., Zapotocny, T., Lenzen, A., Stobie, J., Johnson, D., Avery, M. A., Sachse, G. W., Thompson, A., Cohen, R., Dibb, J. E., Crawford, J. H., Rault, D. F., Martin, R., Szykman, J., and Fishman, J.: Chemical data assimilation estimates of continental U.S. ozone and nitrogen budgets during the Intercontinental Chemical Transport Experiment-North America, *J. Geophys. Res. Atmospheres*, 112, <https://doi.org/10.1029/2006JD007722>, 2007.
- 630 Pierce, R. B., Lenzen, A., and Harkey, M.: Aura Chemical Reanalysis in support Air Quality Applications, https://weather.ndc.nasa.gov/conference/public_health_av/Lenzen_NASA_2016_Applied_Science_Program_Review_Pierce_Final.pdf, 2016.
- 635 Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J. R., Christopher, S. A., Di Girolamo, L., Giglio, L., Holz, R. E., Kearney, C., Miettinen, J., Reid, E. A., Turk, F. J., Wang, J., Xian, P., Zhao, G., Balasubramanian, R., Chew, B. N., Janjai, S., Lagrosas, N., Lestari, P., Lin, N.-H., Mahmud, M., Nguyen, A. X., Norris, B., Oanh, N. T. K., Oo, M., Salinas, S. V., Welton, E. J., and Liew, S. C.: Observing and understanding the Southeast Asian aerosol system by remote sensing: An initial review and analysis for the Seven Southeast Asian Studies (7SEAS) program, *Atmospheric Res.*, 122, 403–468, <https://doi.org/10.1016/j.atmosres.2012.06.005>, 2013.
- 640 Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS Aerosol Algorithm, Products, and Validation, *J. Atmospheric Sci.*, 62, 947–973, <https://doi.org/10.1175/JAS3385.1>, 2005.
- 645 Rowlinson, M. J., Rap, A., Arnold, S. R., Pope, R. J., Chipperfield, M. P., McNorton, J., Forster, P., Gordon, H., Pringle, K. J., Feng, W., Kerridge, B. J., Latter, B. L., and Siddans, R.: Impact of El Niño–Southern Oscillation on the interannual variability of methane and tropospheric ozone, *Atmospheric Chem. Phys.*, 19, 8669–8686, <https://doi.org/10.5194/acp-19-8669-2019>, 2019.
- 650 Santoso, A., Mcphaden, M. J., and Cai, W.: The Defining Characteristics of ENSO Extremes and the Strong 2015/2016 El Niño, *Rev. Geophys.*, 55, 1079–1129, <https://doi.org/10.1002/2017RG000560>, 2017.
- Schaack, T. K., Zapotocny, T. H., Lenzen, A. J., and Johnson, D. R.: Global Climate Simulation with the University of Wisconsin Global Hybrid Isentropic Coordinate Model, *J. Clim.*, 17, 2998–3016, [https://doi.org/10.1175/1520-0442\(2004\)017<2998:GCSWTU>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2998:GCSWTU>2.0.CO;2), 2004.
- 655 Sekiya, T. and Sudo, K.: Role of meteorological variability in global tropospheric ozone during 1970–2008, *J. Geophys. Res. Atmospheres*, 117, <https://doi.org/10.1029/2012JD018054>, 2012.
- Sekiya, T. and Sudo, K.: Roles of transport and chemistry processes in global ozone change on interannual and multidecadal time scales, *J. Geophys. Res. Atmospheres*, 119, 4903–4921, <https://doi.org/10.1002/2013JD020838>, 2014.



- 660 Soja, A. J., Cofer, W. R., Shugart, H. H., Sukhinin, A. I., Stackhouse Jr., P. W., McRae, D. J., and Conard, S. G.: Estimating fire emissions and disparities in boreal Siberia (1998–2002), *J. Geophys. Res. Atmospheres*, 109, <https://doi.org/10.1029/2004JD004570>, 2004.
- Sudo, K. and Takahashi, M.: Simulation of tropospheric ozone changes during 1997–1998 El Niño: Meteorological impact on tropospheric photochemistry, *Geophys. Res. Lett.*, 28, 4091–4094, <https://doi.org/10.1029/2001GL013335>, 2001.
- Trenberth, K. E.: The Definition of El Niño, *Bull. Am. Meteorol. Soc.*, 78, 2771–2778, [https://doi.org/10.1175/1520-0477\(1997\)078<2771:TDOENO>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<2771:TDOENO>2.0.CO;2), 1997.
- 665 Wang, X., Parrish, D., Kleist, D., and Whitaker, J.: GSI 3DVar-Based Ensemble–Variational Hybrid Data Assimilation for NCEP Global Forecast System: Single-Resolution Experiments, *Mon. Weather Rev.*, 141, 4098–4117, <https://doi.org/10.1175/MWR-D-12-00141.1>, 2013.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 670 Wu, W.-S., Purser, R., and Parrish, D.: Three-Dimensional Variational Analysis with Spatially Inhomogeneous Covariances, *Mon. Weather Rev. - MON WEATHER REV.*, 130, [https://doi.org/10.1175/1520-0493\(2002\)130<2905:TDAVWS>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<2905:TDAVWS>2.0.CO;2), 2002.
- Yin, Y., Ciais, P., Chevallier, F., van der Werf, G. R., Fanin, T., Broquet, G., Boesch, H., Cozic, A., Hauglustaine, D., Szopa, S., and Wang, Y.: Variability of fire carbon emissions in equatorial Asia and its nonlinear sensitivity to El Niño, *Geophys. Res. Lett.*, 43, 10,472–10,479, <https://doi.org/10.1002/2016GL070971>, 2016.
- 675 Yumimoto, K., Tanaka, T. Y., Oshima, N., and Maki, T.: JRAero: the Japanese Reanalysis for Aerosol v1.0, *Geosci. Model Dev.*, 10, 3225–3253, <https://doi.org/10.5194/gmd-10-3225-2017>, 2017.
- Yurganov, L. N., McMillan, W. W., Dzhola, A. V., Grechko, E. I., Jones, N. B., and van der Werf, G. R.: Global AIRS and MOPITT CO measurements: Validation, comparison, and links to biomass burning variations and carbon cycle, *J. Geophys. Res. Atmospheres*, 113, <https://doi.org/10.1029/2007JD009229>, 2008.
- 680 Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian climate centre general circulation model, *Atmosphere-Ocean*, 33, 407–446, <https://doi.org/10.1080/07055900.1995.9649539>, 1995.
- 685 Zhang, G. J., Kiehl, J. T., and Rasch, P. J.: Response of Climate Simulation to a New Convective Parameterization in the National Center for Atmospheric Research Community Climate Model (CCM3), *J. Clim.*, 11, 2097–2115, [https://doi.org/10.1175/1520-0442\(1998\)011<2097:ROCSTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2097:ROCSTA>2.0.CO;2), 1998.
- Ziemke, J. R. and Chandra, S.: La Nina and El Nino—induced variabilities of ozone in the tropical lower atmosphere during 1970–2001, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2002GL016387>, 2003.
- 690 Ziemke, J. R., Chandra, S., and Bhartia, P. K.: Two new methods for deriving tropospheric column ozone from TOMS measurements: Assimilated UARS MLS/HALOE and convective-cloud differential techniques, *J. Geophys. Res. Atmospheres*, 103, 22115–22127, <https://doi.org/10.1029/98JD01567>, 1998.



695 Ziemke, J. R., Chandra, S., Duncan, B. N., Froidevaux, L., Bhartia, P. K., Levelt, P. F., and Waters, J. W.: Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's Chemical Transport Model, *J. Geophys. Res. Atmospheres*, 111, <https://doi.org/10.1029/2006JD007089>, 2006.

Ziemke, J. R., Chandra, S., Oman, L. D., and Bhartia, P. K.: A new ENSO index derived from satellite measurements of column ozone, *Atmospheric Chem. Phys.*, 10, 3711–3721, <https://doi.org/10.5194/acp-10-3711-2010>, 2010.

700 Ziemke, J. R., Douglass, A. R., Oman, L. D., Strahan, S. E., and Duncan, B. N.: Tropospheric ozone variability in the tropics from ENSO to MJO and shorter timescales, *Atmospheric Chem. Phys.*, 15, 8037–8049, <https://doi.org/10.5194/acp-15-8037-2015>, 2015.