

# Transient Attracting Profiles in the Great Pacific Garbage Patch

Luca Kunz<sup>1</sup>, Alexa Griesel<sup>1</sup>, Carsten Eden<sup>1</sup>, Rodrigo Duran<sup>2,3</sup>, and Bruno Sainte-Rose<sup>4</sup>

<sup>1</sup>Institute of Oceanography, Universität Hamburg, Hamburg, Germany

<sup>2</sup>National Energy Technology Laboratory, U.S. Department of Energy, Albany OR, USA

<sup>3</sup>[Theiss Research Planetary Science Institute, La Jolla CA 1700 East Fort Lowell, Tucson, AZ 85719, USA](#)

<sup>4</sup>The Ocean Cleanup, Rotterdam, The Netherlands

**Correspondence:** Luca Kunz (luca.kunz@[uni-hamburg.orac.deearth](mailto:uni-hamburg.orac.deearth))

**Abstract.** A major challenge for cleanup operations in the Great Pacific Garbage Patch is the daily prediction of plastic concentrations that allows to identify hotspots of marine debris. Lagrangian simulations of large particle ensembles are the method in use and effectively reproduce observed particle distributions at synoptic scales  $\mathcal{O}(1000\text{km})$   $\mathcal{O}(1000\text{ km})$ . However, they lose accuracy at operational scales  $\mathcal{O}(1-10\text{km})$   $\mathcal{O}(1-10\text{ km})$  and operators regularly experience differences between predicted and encountered debris accumulations within the garbage patch. Instead of asking *Where do objects go as they follow the current?* as in Lagrangian methods, we here take a different approach and question *Which locations attract material?*. The recently developed concept of Transient Attracting Profiles (TRAPs) provides answers to this ~~since it allows to detect the most attractive because it uncovers the most attracting~~ regions of the flow. TRAPs are the ~~attractive-attracting~~ form of hyperbolic Objective Eulerian Coherent Structures and ~~can be computed are computable~~ from the instantaneous strain field on the ocean surface. They describe flow features ~~that which~~ attract drifting objects and could facilitate offshore cleanups ~~which that~~ are currently taking place in the Great Pacific Garbage Patch. However, the concept remains unapplied since little is known about the persistence and ~~attractive properties attraction~~ of these features ~~in~~, ~~specifically within~~ the Pacific. Therefore, we compute a ~~20-years~~ ~~20-year~~ dataset of daily TRAP detections from satellite-derived mesoscale velocities within the North Pacific subtropical gyre. We are the first to track these instantaneous flow features as they propagate through space and time. ~~This It~~ allows us to study the life cycle of TRAPs, which can range from days to seasons and ~~on average lasts for~~  $\bar{A} \approx (6 \pm 12)$  ~~days~~ ~~lasts an average of~~ ~~six days~~. We show how long-living TRAPs with lifetimes ~~of~~  $A > 30$  ~~days~~ ~~beyond 30 days~~ intensify and weaken over their life cycle ~~and~~, ~~and we~~ demonstrate that the evolution stage of TRAPs affects the motion of nearby surface drifters. Our findings indicate that, ~~at the mesoscale~~, operators in the Great Pacific Garbage Patch should search for long-living TRAPs that are at an advanced stage of their life cycle. These TRAPs are ~~the~~ most likely to induce ~~a~~ large-scale confluence of drifting objects and their streamlining into hyperbolic pathways. Such a streamlined bypass takes ~~on average~~  $\bar{\varphi} \approx (5.3 \pm 3.8)$  ~~days and could be exploited~~, ~~on average, five days and creates an opportunity~~ to filter the flow around TRAPs. But we also find TRAPs that retain material over multiple weeks where we suspect material clustering at the submesoscale, ~~prospective~~. ~~Prospective~~ research could investigate this ~~with soon available~~ ~~further by applying our algorithms to soon available~~ high-resolution observations of the flow. ~~Eventually, our research may also benefit other challenges that are related to the search at sea~~ ~~Our analysis contributes~~ ~~to a better understanding of TRAPs and can benefit even more offshore operations~~, such as optimal drifter deployment, ~~the identification of foraging hotspots~~ ~~oil spill containment~~ or humanitarian search and rescue.




## Short summary

TRansient Attracting Profiles (TRAPs) indicate the most ~~attractive~~ attracting regions of the flow and have the potential to  
30 facilitate offshore cleanup operations in the Great Pacific Garbage Patch. We study the characteristics of TRAPs and the  
prospects for predicting debris transport from a mesoscale permitting dataset. Our findings provide an advanced understanding  
of TRAPs in this ~~particular~~ specific region and demonstrate the importance of TRAP lifetime estimations to an operational  
application. Our ~~algorithm to track TRAPs~~ TRAPs tracking algorithm complements the recently published TRAPs concept  
and prepares its use with high-resolution observations from the SWOT mission. Our findings may also benefit research in other  
35 fields like ~~e.g.~~ optimal drifter deployment, sargassum removal, ~~the identification of foraging hotspots~~ oil spill containment or  
search and rescue.

## 1 Introduction

The horizontal long-term flow at the ocean surface is understood ~~as-to be~~ the main forcing that transports floating material over large distances (van Sebille et al., 2020) and can be well-described by the combination of geostrophic and Ekman currents (Röhrs et al., 2021). Floating marine debris follows the large-scale convergence within each of the five subtropical gyres and forms basin-scale accumulation zones (van Sebille et al., 2020) ~~which-that~~ exhibit elevated levels of plastic concentration. In this context, the North Pacific subtropical gyre ~~is-to-date-~~ to date, the area of highest scientific and public concern. First initiatives to clean up ocean plastic pollution at global scale are taking place in this particular gyre (Slat, 2022), and a variety of experiments has been dedicated to estimating the limits of this accumulation zone, which is colloquially termed the *Great Pacific Garbage Patch* (Onink et al., 2019; Lebreton et al., 2018; Law et al., 2014). Figure 1 highlights this large-scale convergence zone and the horizontal long-term flow at the surface of the northeast Pacific ~~ocean~~ Ocean.



**Figure 1.** Mean circulation and EKE in the eastern North Pacific. ~~Quivers-Vectors~~ indicate the direction and magnitude of ~~20-years-20-year~~ average near-surface geostrophic + Ekman current velocities for the period 2000-2019 (CMEMS, 2022a). Arrow size scales with increasing velocity magnitude, ~~the-~~ The red box defines our study domain and highlights the area of large-scale convergence due to Ekman transport. ~~In this region-~~ The boundaries approximate the estimated limits of the Great Pacific Garbage Patch ~~is-expected-to-be-located-~~ (Law et al., 2014; Lebreton et al., 2018; Onink et al., 2019). The ocean surface is coloured by the magnitude of eddy kinetic energy (EKE) w.r.t. to the same period, derived from geostrophic currents only (CMEMS, 2022b).

A constant challenge for cleanup operations in this region is to predict ~~day-to-day~~ day-to-day variabilities of plastic ~~concentrations~~ ~~and-with-this-concentration~~ and to identify hotspots of marine debris. The common method is to release large ensembles of virtual particles at the ocean surface and to derive their trajectories and distribution as they follow the time-evolving surface flow (van Sebille et al., 2018; van Sebille et al., 2020; Duran et al., 2021). Measurements from altimetry or estimations from



numerical ocean models typically provide the velocity fields that drive these Lagrangian particle simulations, which eventually allow to derive particle concentrations at the ocean surface. At synoptic scales  $\mathcal{O}(1000\text{km})$   $\mathcal{O}(1000\text{ km})$ , Lagrangian simulations succeed in predicting the limits of the Great Pacific Garbage Patch (Onink et al., 2019; Lebreton et al., 2018; Law et al., 2014). However, within the garbage patch and at operational scales  $\mathcal{O}(1-10\text{km})$   $\mathcal{O}(1-10\text{ km})$ . The Ocean Cleanup reports that Lagrangian simulations often fail in accurately predicting the particle distributions they observe at sea. This deficiency ~~can be a consequence of~~ may result from the combination of Lagrangian methods with submesoscale velocity estimations from numerical models. At operational scales and within this region, numerical models currently represent the only source for now- and forecast estimations of the surface flow, but these simulations can only approximate the true dynamics at sea. On the other hand, Lagrangian simulations can produce significant trajectory errors if the underlying velocity data or the trajectory ~~modeling itself are~~ modelling itself is missing important physics ~~and errors~~, specifically because errors can accumulate quickly during the integration process (~~Duran et al., 2021~~) (Duran et al., 2021; Serra et al., 2020).

Instead of asking *Where do objects go as they follow the current?* as in Lagrangian methods, we here take a fundamentally different approach and question *Which locations attract material?*. The recently published concept of Transient Attracting Profiles (TRAPs, Serra et al. (2020)) provides answers to this since it allows to detect the most ~~attractive~~ attracting regions of the flow. TRAPs are the ~~attractive~~ attracting form of hyperbolic Objective Eulerian Coherent Structures (OECSs, Serra and Haller (2016)) and ~~can be computed~~ are computable from the symmetric part of the velocity gradient. They indicate regions of maximal compression and stretching on the ocean surface ~~which are expected to~~ that translate into the attraction and hyperbolic transport of nearby floating objects.

Serra et al. (2020) and Duran et al. (2021) provide experiments that show the capability of TRAPs to attract drifting objects. They demonstrate how TRAPs uncover the stretching of tracer patterns that ~~is hidden to~~ remains undetected by conventional diagnostics like streamlines or divergence. Their experiments also indicate that TRAPs are insensitive to the shape, submergence level, release time and release position of drifting objects. These parameters are generally uncertain in applications but must be considered in Lagrangian simulations. Moreover, Serra and Haller (2016), Serra et al. (2020) and Duran et al. (2021) argue that TRAPs are more robust to moderate errors in the underlying velocity field while trajectory-based methods are ~~sensitively dependent to inevitable errors~~ susceptible to error accumulation during the velocity integration, see Table B1 for more ~~benefits~~ details and benefits of the TRAPs method.

So, it stands to reason that TRAPs could facilitate offshore cleanups ~~which that~~ are currently taking place in the Great Pacific Garbage Patch. However, the concept remains unapplied since the persistence and attractive properties of TRAPs have not been characterised in this particular region. Therefore, we create a ~~20-years~~ 20-year dataset of daily TRAP detections and study these features within the North Pacific subtropical gyre.

We compute TRAPs from near-surface geostrophic + Ekman current velocities since geostrophic velocity from altimetry is the only large-scale observation that resolves flow features at the mesoscale (Abernathey and Haller, 2018). Many studies have established that altimetry-derived velocity products are accurate for Lagrangian transport applications, see ~~e.g.~~ Sect. 3.1.3 of Duran et al. (2021) and references therein. Moreover, ~~the contributions we make~~ our methods can be applied to higher-  
90 resolution observations that ~~soon will~~ will soon be available from the SWOT mission (International Altimetry Team, 2021).

Since altimetry ~~can be seen as a filter for~~ acts like a filter on the ocean circulation that ~~seperates~~ separates out all small-scale, short-term features of the flow, our study ~~is naturally embedded in~~ will focus on the low-frequency circulation. ~~This motivates us to~~ Therefore, we will locate TRAPs within the mesoscale eddy field by comparing our dataset to corresponding records of  
95 mesoscale eddy detections. We investigate how these coherent structures relate in order to advance our understanding of strain between eddies and ~~how it can be utilised to predict debris transport~~ its potential for the prediction of debris transport.

Serra et al. (2020) mention that TRAPs "necessarily persist over short times" with examples of TRAPs existing for several hours and attracting nearby objects within two to three hours. These time scales for persistence and impact derive from TRAPs computed upon submesoscale velocity fields with a high tempo-spatial resolution. However, the concept is scale-invariant and can be applied to velocity data of any resolution. TRAP characteristics will depend on the velocity data used for their computation, and the lifetime and impact of TRAPs will be relative to the time scales of the oceanic structures that give rise to the hyperbolic-type Lagrangian motion that TRAPs identify. For this reason, Duran et al. (2021) find persistent TRAPs that predict transport patterns eight days in advance. They compute TRAPs from mesoscale surface velocities with daily frequency and consequently study these structures at larger scales than Serra et al. (2020). With our choice of altimetry data, we follow Duran et al. (2021) and expect mesoscale TRAPs to exist and impact on time scales comparable to those of mesoscale flow features. Indeed, some examples from altimetry data in Serra and Haller (2016) show that different types of OECSSs, including the attracting hyperbolic type studied here, can last for at least six days.  
100  
105

110 TRAPs that persist over several days will then highlight permanent ~~features of the flow~~ flow features where we might find large-scale confluence of material. ~~The detection of~~ Therefore, identifying persistent TRAPs can ~~therefore~~ help to point cleanup operations in the right direction, which motivates us to design a tracking algorithm that follows TRAPs through space and time. We are the first to track these Eulerian flow features, ~~to~~ determine their lifetimes and ~~to~~ describe their propagation through the domain. We further combine these ~~findings~~ methods with observations of surface drifters to investigate the TRAP properties  
115 ~~that are~~ relevant for an offshore cleanup in the Great Pacific Garbage Patch. ~~But~~ Eventually, the findings we make ~~also~~ have the potential to facilitate even more maritime search operations that are taking place in other contexts and regions.

The paper is organised as follows. In Section 2, we review the theoretical aspects of Transient Attracting Profiles, outline the design of the experiment and go through the methods we use ~~to give a comprehensive~~ for our analysis of mesoscale TRAPs.  
120 Section 3 presents ~~the results of our analysis~~ our results in four parts - the spatial distribution of TRAPs, their life cycle and

propagation, vorticity patterns around TRAPs and the impact of TRAPs on nearby drifters. In Section 4, we discuss our findings and the directions they offer for future research.

## 2 Methodology

### 125 2.1 Transient Attracting Profiles

Serra et al. (2020) derive TRAPs from the instantaneous strain field ~~of-on~~ the ocean surface using snapshots of the two-dimensional surface velocity field  $\mathbf{u}(\mathbf{x}, t)$ , with  $\mathbf{u}$  being dependent on position  $\mathbf{x}$  and time  $t$ . The symmetric part of the velocity gradient represents the time-dependent strain tensor  $\mathbf{S}(\mathbf{x}, t) = \frac{1}{2}(\nabla\mathbf{u}(\mathbf{x}, t) + [\nabla\mathbf{u}(\mathbf{x}, t)]^T)$  with ~~eigenvalues-the eigenvalue~~ fields  $s_i(\mathbf{x}, t)$  and eigenvector fields  $\mathbf{e}_i(\mathbf{x}, t)$ .  ~~$\mathbf{S}$ ,  $s_i$  and eigenvectors- $\mathbf{e}_i$ .~~ We denote the respective quantities at a fixed position  
 130  $\mathbf{x}_0$  and time  $t_0$  and we apply the notation for the diagonal form of  ~~$\mathbf{S}(\mathbf{x}, t)$~~   $\mathbf{S}$  from Serra and Haller (2016):

$$\mathbf{S}\mathbf{e}_i = s_i\mathbf{e}_i, \quad |\mathbf{e}_i| = 1, \quad i = 1, 2; \quad s_1 \leq s_2, \quad \mathbf{e}_2 = \mathbf{R}\mathbf{e}_1, \quad \mathbf{R} := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad (1)$$


The deformation of any fluid's surface element  $A$  is determined by the local strain rates  $s_{i_2}$ , which specify the rates of stretching ( $s_i > 0$ ) or compression ( $s_i < 0$ ) of  $A$  along the principle axes indicated by the local eigenvectors  $\mathbf{e}_i$ , see ~~(Olbers et al., 2012)~~ Olbers et al. (2012) for details. Due to the condition  $s_1 \leq s_2$ , the local eigenvector  $\mathbf{e}_1$  ~~lines describe lines of minimal identifies~~  
 135 the direction of compression and  $\mathbf{e}_2$  ~~lines lines of maximal~~ the direction of stretching for a non-uniform deformation with  
 $s_1 < 0$  and  $s_2 > 0$ . The compression and stretching of surface elements ~~translates translate~~ into the attraction and repulsion of material ~~and negative~~. Negative local minima of  ~~$s_1(\mathbf{x}, t)$  therefore  $s_1(\mathbf{x}, t_0)$  at snapshot time  $t_0$  then~~ describe the most  
 attractive ~~attracting~~ regions of the flow, maximising attraction normal to  $\mathbf{e}_2$  at the respective position. For incompressible conditions,  $s_1 = -s_2$  further holds and local minima of  ~~$s_1 s_2(\mathbf{x}, t_0)$~~  simultaneously indicate local maxima of  ~~$s_2$ , so that the~~  
 140  $s_2(\mathbf{x}, t_0)$ . The strongest attraction and strongest repulsion then occur at the same position and in orthogonal directions.

TRAPs ~~represent  $\mathbf{e}_2$  lines starting indicate the most attracting regions of the flow as they start~~ at negative local minima of the  ~~$s_1 s_2(\mathbf{x}, t_0)$~~  strain field and extend until tangent to the local eigenvectors  $\mathbf{e}_2$  until the strain rate  $s_1$  along the  $\mathbf{e}_2$ -line tangent  
 ceases to be monotonically increasing. Consequently, TRAPs contain one minimum value of  ~~$s_1 s_2(\mathbf{x}, t_0)$~~ , i.e. the point of  
 145 strongest attraction perpendicular towards the TRAP. The position of this minimum value-local minimum is called the TRAP core, which represents an objective saddle-type stagnation point of the unsteady flow (Serra and Haller, 2016). ~~The TRAP~~  
~~itself, as it is at every point tangent~~ A TRAP describes the direction of maximal stretching because it is tangent at each point to the unit ~~eigenvector~~ eigenvectors  $\mathbf{e}_2$  ~~describes the direction of maximum stretching and will,~~ and in the following ~~also,~~ it will  
 be referred to as a TRAP curve.

150

Figure 2 ~~shows presents~~ an example of a Transient Attracting Profile. The TRAP is displayed as a red curve with the TRAP core as a red dot in the middle. The image ~~displays shows~~ this structure upon a colourmap of the underlying  ~~$s_1$  strain fields~~  
~~well as velocity quivers  $s_1(\mathbf{x}, t_0)$  strain field, superimposed by velocity vectors of the surrounding flow, all~~ derived from the  
 same snapshot of surface geostrophic + Ekman current velocity at time  $t_0$ . The TRAP core ~~is positioned in coincides with~~ the  
 155 point of highest attraction, i.e. a negative local minimum of the  ~~$s_1 s_2(\mathbf{x}, t_0)$~~  field, and the velocity ~~vectors indicate the flow~~

around the TRAP, i.e. indicates water motion normal towards the TRAP and subsequent transport to both ends of the structure. The figure also displays the positions of this TRAP after 10 and 20 days.



**Figure 2.** Example Transient Attracting Profile. The red structure represents one TRAP from our 20-years-20-year record of daily TRAPs, computed from snapshots of surface geostrophic + Ekman current velocity (CMEMS, 2022a). The red dot in the middle indicates the TRAP core, the red line represents the TRAP curve. Quivers-Vectors illustrate the surrounding flow around the structure and the colourmap indicates the  $s_1$  strain field, derived from the same velocity snapshot at time  $t_0$ . The geometries in salmon colour indicate, with increasing transparency, the position of the same TRAP after 10 and 20 days.

## 2.2 TRAPs computation

160 Serra et al. (2020) have published a programme to compute TRAPs from two-dimensional snapshots of an Eulerian velocity field  $u(x, t)$  (Serra, 2020), see Table B2 for details of the algorithm and Kunz (2024a), Kunz (2024b) for our post-processing of the output. We set-truncate TRAP curves wherever the attraction rate along the curve falls below 30 % of the attraction at the respective core. This cutoff criterion makes sense physically because the attraction of nearby parcels becomes negligible as distance increases away from the core. Without cutoff, TRAPs can become indefinitely long and merge with nearby structures, 165 which makes them hard to distinguish. Their converging ends then put wrong emphasis on regions between TRAP cores where the attraction rate is comparably low, see Fig. S1 in the Supplementary Material for details. In addition, we limit TRAPs to a maximal arclength of  $1^\circ$  to represent a reasonable length scale for the regions between mesoscale eddies that is at the same time representative for cleanup operations since TRAP lengths are of minor importance to our analysis and because the statistics will refer to the position and attraction of the TRAP core. This choice does not affect our main diagnostics but is optional and 170 can be neglected in future studies. We call the points that discretise the TRAP curve curve points and the programme, and the algorithm gives the positions of all curve points and corresponding TRAP cores in the domain. It further also outputs the

normal attraction rate  $s_1$  at every TRAP core, which we name *core attraction*. We set the boundaries of our study domain to [22.5° N, 42.5° N] in latitudes-latitude and [-160° E, -125° E] in longitudes-longitude and compute TRAPs within this domain. We choose these limits, highlighted by the red box in Fig. 1, ~~in order~~ to enclose the North Pacific accumulation zone  
175 between Hawaii and California as defined by Onink et al. (2019). These boundaries also warrant no intersection with any land mass. We compute TRAPs from daily snapshots of near-surface geostrophic + Ekman currents and therefore extract the velocity fields  $u_0$  and  $v_0$  from the product *Global Total Surface and 15m Current (COPERNICUS-GLOBCURRENT)* from *Altimetric Geostrophic Current and Modeled Ekman Current Reprocessing* that is provided by the E.U. Copernicus Marine Service (CMEMS, 2022a). The velocity fields are available at three-hourly instantaneous time steps, from which we select data  
180 at UTC midnight to sample snapshots with a daily frequency. ~~Velocity fields are provided on a~~ The latitude-longitude grid of the velocity field has a resolution of 0.25° ~~resolution~~.

### 2.3 Tracking ~~Algorithm~~algorithm

Our tracking algorithm runs on the full TRAPs record and finds spatially proximate detections at consecutive timestamps,  
185 which can be identified as one single feature of the flow. The only free parameter  $\epsilon$  defines the size of the search area around a current-given TRAP to look for a detection in the ~~next snapshot and is set following~~ snapshot, and we set it to  $\epsilon = 0.25^\circ$ , ~~see Kunz (2024a) for more details.~~ For a higher value, the algorithm creates "jumps" from a current to an unrealistically far future TRAP detection and overestimates trajectory lengths, see Sect. S2 in the Supplementary Material for a detailed explanation and motivation for this choice. The algorithm assigns a unique label to each TRAP trajectory and its associated instances ~~and it~~  
190 ~~It~~ derives metrics like ~~e.g.~~ the lifetime  $\Lambda$  of TRAPs and their age  $\tau$  at a particular snapshot. ~~We can then~~ The algorithm only captures the time spent inside the study domain and period. Therefore, it gives rise to potential bias in the lifetime estimation of TRAPs that reach beyond the tempo-spatial limits of the domain. However, we find that only 5.4 % of all TRAP trajectories are adjacent to these limits and might not entirely occur within the study domain. Our conclusions and the distribution of TRAP lifetimes don't change if those biased trajectories are excluded, see Sect. S3 in the Supplementary Material, where we  
195 analyse this in detail. With the trajectory estimation, we can now derive the zonal and meridional translation speeds  $c_x$  and  $c_y$  for every instance of a TRAP trajectory. Therefore, we choose all TRAPs that persist for at least three days and ~~average the forward and backward shifted velocity at a current timestamp. The forward/backward shifted velocity is the distance to its succeeding/preceding position divided by the time lapsed between both instances, respectively. This way we deliberately create~~ compute propagation speeds using centred differences. We derive no velocities at the start and end of a trajectory ~~and~~  
200 ~~do not gain propagation speeds for trajectories of two days lifetime. In turn, we obtain translation speeds of individual TRAP instances which we consider more accurate than taking the full distance travelled by a TRAP and dividing it by the respective lifetime where the TRAP forms and decays. At these stages,  $c_x$  and  $c_y$  could be estimated by following the underlying oceanic structure that creates the TRAP.~~

## 2.4 Mesoscale eddy data

205 We compare TRAPs against eddy detections from ~~The altimetric Mesoscale Eddy Trajectory Atlas (META3.2 DT) which~~ The altimetric Mesoscale Eddy Trajectories Atlas (META3.2 DT) that is produced by SSALTO/DUACS and distributed by AVISO+ with support from CNES, in collaboration with IMEDEA (AVISO+ et al., 2022). This dataset provides Eulerian detections of eddies derived from sea surface height (SSH) contours, is at frequency with our TRAPs record and includes estimations of the eddy contour speed  $U$ . The eddy contour speed  $U$  is the average geostrophic speed of the contour of maximum circum-  
210 average geostrophic speed for the detected eddy. We filter the dataset for eddy detections within the study domain and period and retrieve 28,645 cyclonic and 24,193 anticyclonic eddy trajectories from 689,460 cyclonic and 686,720 anticyclonic eddy detections. Since we ~~simply~~ cut off eddy trajectories beyond the domain boundaries, ~~4.2% of all eddy trajectories will show discontinuities since trajectories can show discontinuities if~~ they leave and return ~~into to~~ the domain. This effect occurs for 4.2 % of all eddy trajectories. We do not correct for this ~~cut-off~~ cut-off since the impact on our aggregate statistics should be  
215 negligible. We estimate the eddy lifetime by taking the time difference between the first and the last occurrence within our domain, added by one day. ~~As a consequence~~ Accordingly, the estimated eddy lifetime can also include times spent outside the domain, given that the eddy returns ~~back into to~~ it again. We further derive eddy propagation speeds as we do for TRAPs.

## 2.5 Vorticity curve


We use the relative vorticity field to characterise the flow around TRAPs. Many TRAPs seem to be surrounded by four vortices  
220 of alternating polarity. In this case, two vortices on each side of a TRAP curve exhibit perpendicular flow towards the TRAP core and tangential flow away from it ~~;~~ with respect to the TRAP curve. We call this vorticity pattern a *quadrupole* and classify variations of it. To detect vorticity patterns without a coordinate transformation, we draw a circle in the horizontal plane around every TRAP core with a radius equal to the distance between the core and the furthest curve point. Starting from the position of the furthest curve point, we ~~parameterize~~ parameterise the circle with angles increasing counter-clockwise ~~;~~ and bilinearly  
225 interpolate the vorticity field to the points on this circle. The *vorticity curve*  $\zeta(\alpha)$  then describes the vorticity along this circle with respect to the angle of parameterisation, i.e. the phase  $\alpha$ .

By visualising the vorticity curve, we ~~are able to directly~~ can see polarity changes in the surrounding vorticity field and their spatial orientation towards the TRAP. Figure 3a illustrates an example of such a vorticity measurement around a TRAP. For  
230 a quadrupole in the northern hemisphere, the polarity pattern along the vorticity curve then results in cyclonic ( $\oplus$ , positive vorticity), anticyclonic ( $\ominus$ , negative vorticity), cyclonic, anticyclonic and the vorticity curve reveals four zero crossings.

Since the ensemble mean of all vorticity curves  $\zeta(\alpha)$  indicates a quadrupole pattern, we filter the ensemble for specific combinations of four vortices ~~of with~~ either cyclonic or anticyclonic rotation. We do not explicitly resolve patterns with less than  
235 four vortices because we can identify them from this classification if needed. In total, there are  $2^4$  possible vortex combinations, 10 of which exert distinct dynamics, i.e. are unique under rotations of  $180^\circ$  around a TRAP core. To isolate each pattern, we



240 ~~further~~ remove a constant average background vorticity from every vorticity curve and then filter the ensemble for these 10 vortex combinations. To detect the vortex configuration within a given vorticity curve, we divide the curve into four phase intervals of  $\alpha_I = [0, \pi/2)$ ,  $\alpha_{II} = [\pi/2, \pi)$ ,  $\alpha_{III} = [\pi, 3\pi/2)$ ,  $\alpha_{IV} = [3\pi/2, 2\pi)$   $\alpha_I = [0, \pi/2)$ ,  $\alpha_{II} = [\pi/2, \pi)$ ,  $\alpha_{III} = [\pi, 3\pi/2)$ ,  $\alpha_{IV} = [3\pi/2, 2\pi)$  and compute the average vorticity within each interval, see Fig. S2-S4 for more details on all 10 vorticity patterns.



**Figure 3.** Vorticity circle and drifter transport around a TRAP. (a) For each TRAP, Blue dots indicate the vorticity field  $\zeta$  is interpolated to interpolation points on a circle around of the core (dotted line) vorticity curve  $\zeta(\alpha)$ . The circle parameterisation starts at blue triangle highlights the position starting point of the furthest curverpoint w.r.t. the core (triangle) and with a phase of parameterisation at  $\alpha = 0^\circ$ .  $\alpha$  increases counter-clockwise and as we sample the vorticity  $\zeta(\alpha)$  along the curve as vorticity curve  $\zeta(\alpha)$  circle. (b) Filled and empty big red circle markers indicate the current position of one drogued and one undrogued drifter two drifters around the same TRAP instance, respectively. Red tails with small markers indicate their show the respective drifter positions throughout the preceding preceding seven days. Quivers One drifter is drogued and indicated by filled markers, the other drifter is undrogued and represented by empty markers. In both panels, vectors illustrate the surrounding flow and the colourmap indicates the relative vorticity field  $\zeta(\mathbf{x}, t_0)$  at snapshot time  $t_0$ .

## 2.6 Impact on drifters

245 Since TRAPs highlight the most attractive-attracting regions and their cores represent objective saddle points of the unsteady surface flow, we expect them to attract and disperse drifting objects in a hyperbolic pattern. We study their transient impact on drifting objects and therefore look by looking at surface drifter trajectories around TRAPs.

250 To compare drifters and TRAPs at simultaneous timestamps, we consult 24-hourly drifter positions at UTC midnight from the Global Drifter Program (Lumpkin and Centurioni, 2019). For our study domain and period, this dataset provides 842, i.e. 328 drogued and 514 undrogued, drifter trajectories distributed over 221,979, i.e. 67,885 drogued and 154,094 undrogued, positions. We call these daily drifter positions *drifter days*. See Fig. S2 in the Supplementary Material for an overview of all drifter positions. Figure 3b illustrates an example for-of hyperbolic drifter motion that we find in from this dataset - drifters are first attracted perpendicular towards the TRAP and then transported along one of its branches towards the end. This is how TRAPs are



~~expected to organise nearby floating objects. See Fig. S1 in the Supplementary Material for an overview on all drifter positions.~~

255 We want to see how drifters behave in the surroundings of a TRAP and ~~therefore~~ detect pairs of drifters and nearby TRAPs. In Kunz (2024a), we provide a comprehensive description of our pair algorithm that works from a drifter's perspective and searches for the closest TRAP within a ~~distance radius~~ of 75 kilometres. ~~For every instance of a~~ We choose this limit since it represents the average distance  $\bar{d}$  plus 1 standard deviation between a drifter and its closest TRAP core, i.e.  $\bar{d} \approx (51 \pm 25)$  km for drogued and  $\bar{d} \approx (49 \pm 24)$  km for undrogued drifters. 86 % of drifter days occur within a 75 km distance to their closest  
260 TRAP core, see Fig. S3 in the Supplementary Material for the respective distribution. For every drifter-TRAP pair, the algorithm records the drifter's current distance to the ~~closest~~-TRAP. It ~~also~~-saves attributes like ~~e.g.~~-the TRAP age  $\tau$  at first encounter, the TRAP lifetime  $\Lambda$  and its attraction rate  $s_1$  at every instance of the pair. Moreover, we ~~will~~-know the daily vorticity pattern in ~~which the surroundings of~~ a drifter-TRAP pair ~~is embedded~~, and we measure the pair's duration, i.e. the retention time  $\varphi$  of a drifter around its closest TRAP.

265 A lot of pairings will only last for ~~a one~~ day due to ephemeral TRAPs with lifetimes of  $\Lambda = 1$  day, due to drifters passing by in the periphery of a TRAP or due to ~~the drifter finding another TRAP that is closer~~ a drifter meeting another structure in the way. We exclude these one-day pairs from our analysis since we cannot infer any motion statistics from them. ~~But the scope of our remaining measurements will still~~, and we note that we observe a lot of one-day pairings for drifters beyond the 75 km limit.

270 Because our pair algorithm searches for the closest TRAP around a drifter, its detection is insensitive to the individual attraction strength or impact range of surrounding TRAPs. The definition of a dynamic impact range, however, would be a valuable contribution to the TRAPs concept that we propose for future research. Our approach here will allow us to ~~carry out a statistical analysis about drifter motion around TRAPs~~ show the aggregate effect of TRAPs on drifters and to provide a first  
275 estimate of the retention times that drifters can spend around a TRAP.

### 3 Results


#### 3.1 Spatial distribution of TRAPs

We first look at important circulation features of the study area. Figure 1 shows the distribution of eddy kinetic energy (EKE) over the domain with respect to the period 2000-2019. The lowest values of EKE occur in the northwest corner of the domain. This subregion is part of an *eddy desert* in the northeast Pacific where mesoscale eddies are low in amplitude and short in lifetime, if present at all (Chelton et al., 2011). ~~There are~~ We find two distinct regions of high EKE in the northeast and ~~the~~ southwest of the domain, indicating frequent turbulence and mesoscale eddy activity. These two regions neighbour the California Upwelling System (CALUS) and the North Hawaiian Ridge Current (NHRC), respectively, which are known for the production of energetic mesoscale eddies (Pegliasco et al., 2015; Lindo-Atichati et al., 2020).

Within our domain and period, we detect 4,076,065 TRAP instances from which we identify 720,391 TRAP trajectories. TRAPs occur everywhere ~~in the domain~~ but with distinct patterns in quantity, persistence and attraction strength. In panel (a) of Fig. 4, we separate the domain into bins of  $0.25^\circ \times 0.25^\circ$  and show the 20-year bin averages of instantaneous TRAP core attraction rates  $s_1$ . A comparison with Fig. 1 reveals that TRAPs are particularly strong in regions of high EKE close to the CALUS and the NHRC. In the central-north, ~~the~~ middle and southeast of the domain, moderate to low EKE prevails, and TRAPs are ~~on average~~, on average, moderately to weakly attracting. The eddy desert in the northwest corner remains with a clear preference for weak TRAPs. We derive a mean attraction rate and standard deviation of  $\bar{s}_1 \approx (-0.23 \pm 0.11) \text{ s}^{-1}$  over all TRAP instances. The most ~~attractive~~ attracting TRAP that we find is located within the eastern hotspot of Fig. 4a and exhibits an attraction rate of  $s_1 = -1.73 \text{ s}^{-1}$ . We correlate the average attraction rates  $s_1$  from Fig. 4a with the EKE field given in Fig. 1 and find a ~~pearson~~ Pearson correlation coefficient of  $r = -0.93$  with a p-value of  $p < 0.001$ . ~~This~~ It indicates a strong and significant negative correlation between both variables. ~~It suggests that weak /strong~~ We infer from this that weak TRAPs, i.e. with a less ~~/more~~ negative attraction rate  $s_1$ , ~~are located~~ occur in regions of low ~~/EKE~~, while strong TRAPs occur in regions of high EKE.

In panel (b), we visualise our tracking results and show the trajectories of TRAPs with lifetimes  $\Lambda > 30$  days. We call this subset *long-living* TRAPs. From ~~time-lapse~~ time-lapse animations of the TRAPs field, we suspect that the trajectories here indicate westward propagation with a tendency towards the equator, see Videos ~~SV1, SV2 and SV3 in~~ (Kunz, 2024c) S1, S2 and S3 in Kunz (2024c). Pegliasco et al. (2015) find similar propagation characteristics for anticyclonic mesoscale eddies ~~that~~ originate originating in the California Upwelling System east of our domain. A comparison to panel (a) suggests that attraction strength may vary along the trajectories of long-living TRAPs ~~which however~~, which, however, will exclude very weak attraction rates. Long-living TRAPs are indeed stronger than TRAPs with lifetimes  $\Lambda \leq 30$  days. We find mean attraction rates of  $\bar{s}_{1,\Lambda > 30} \approx (-0.28 \pm 0.11) \text{ s}^{-1}$  and  $\bar{s}_{1,\Lambda \leq 30} \approx (-0.20 \pm 0.09) \text{ s}^{-1}$  for TRAP instances associated with these groups, respectively.

In panel (c), we separate the domain into bins of  $1^\circ \times 1^\circ$  and count the number of TRAP trajectories that pass through each histogram bin. We find more TRAPs passing through the northwest than through the southeast half of the domain. TRAP trajectories are especially abundant around the eddy desert, i.e. in the northwest and central-north of the study region. We complete the picture by deriving the average lifetime  $\bar{\Lambda}$  of all trajectories that pass through a histogram bin in panel (d). We find a clear preference for ephemeral TRAPs in the northwest corner and for persistent TRAPs within the southeast half of the domain. We summarise that TRAP trajectories are very abundant but only remain for a few days around the eddy desert while they become less abundant but more persistent towards the equator and the eastern boundary. It suggests that the underlying oceanic structures that create TRAPs show different characteristics in these two regions. Our observations are therefore consistent with the sparse occurrence of primarily weak mesoscale eddies in the eddy desert around the northern domain boundary (Chelton et al., 2011), and with the generation of energetic mesoscale eddies around the CALUS and the NHRC (Pegliasco et al., 2015; Lindo-Atichati et al., 2020) which eventually propagate through the southeastern part of the domain.



**Figure 4.** Distribution of TRAP characteristics over the domain and in the period 2000-2019. (a) TRAP core attraction strength  $s_1$  averaged over 20 years and all instances within bins of  $0.25^\circ \times 0.25^\circ$  size. Red colours indicate high attraction. (b) Trajectories of long-living TRAPs with lifetimes  $\Lambda > 30$  days. Trajectories are coloured by the associated TRAP lifetime  $\Lambda$ , using the respective colour scale on the right. The most persistent TRAP is indicated by a thick line. (c) Number of identified TRAP trajectories that pass through histogram bins of  $1^\circ \times 1^\circ$  size. (d) Average lifetime  $\bar{\Lambda}$  of TRAP trajectories passing through each bin.

### 3.2 Life cycle and propagation

325 We find that TRAPs typically persist for a ~~view-few~~ days. However, the life cycle of some profiles can also span several seasons. Figure 5a presents the distribution of TRAP lifetimes  $\Lambda$  over all TRAP trajectories. We find a mean lifetime of  $\bar{\Lambda} \approx (6 \pm 12)$  days. The most persistent TRAP counts a lifetime of  $\Lambda = 294$  days and is indicated by the thick red-brown trajectory in Fig. 4b. Only ~~4% of all~~ 4 % of TRAP trajectories exhibit lifetimes of  $\Lambda > 30$  days, but they include around ~~41%~~ 41 % of all instantaneous detections.


330

In Fig. 5a, we also present the lifetime distribution for all mesoscale eddy detections that AVISO+ et al. (2022) find in our domain. We consult the eddy census product by AVISO+ et al. (2022) to provide a first overview of three comparable features between TRAPs and mesoscale eddies. AVISO+ et al. (2022) identify 28,645 cyclonic and 24,193 anticyclonic mesoscale eddy trajectories within our study domain and period. On average, these cyclonic eddies persist for  $\bar{\Lambda}_{\oplus} \approx (24 \pm 40)$  days and anti-  
335 cyclonic ones for  $\bar{\Lambda}_{\ominus} \approx (29 \pm 56)$  days. The distributions in panel (a) resemble one another, but there are considerably more TRAP than eddy trajectories in our domain, and their lifetimes ~~are shifted~~ shift towards smaller values. Over their lifetime, ~~approx-~~ approximately 25 % of ~~all~~ TRAPs find no eddy detection, 54 % find one and 21 % find multiple eddy detections within a radius of  $0.5^\circ$  arclength. ~~We find that this~~ This cumulative number of ~~encountered~~ encountered eddy detections depends on TRAP lifetime with a correlation coefficient of  $r = 0.65$  and a p-value of  $p < 0.001$ . Long-living TRAPs ~~encounter~~ on average  
340 , on average, four eddy detections over their life cycle ~~while some of them,~~ while some long-living TRAPs can be related to up to 18 different eddy detections. From this, we understand that the detections from AVISO+ et al. (2022) are not ~~really suitable~~ to explain suitable for explaining individual TRAP detections.

We expect high rotation speeds of mesoscale eddies to create high strain between them, which should reflect in a relation  
345 between  $s_1$  and the eddy contour speed  $U$ . We investigate this relation by recasting panel (a) from Fig. 4 but for the mean eddy contour speeds  $U$ . We find a correlation coefficient between both histograms of  $r = -0.94$  with a p-value of  $p < 0.001$ . It confirms that, on average, TRAP attraction strength scales with eddy contour speed. Panel (b) of Fig. 5 then views ~~on~~ the evolution of  $s_1$  and  $U$  over the lifetime of long-living TRAPs and eddies. We adopt this approach from Pegliasco et al. (2015),  
who study the evolution of eddy radii and amplitudes. On average, both TRAP attraction and eddy contour speed intensify in  
350 the first half and decrease in the second half of a respective life cycle. We conclude that the life cycles of TRAPs and eddies relate.

Panel (c) presents the latitudinal dependence of the zonal translation speed  $c_x$  for TRAPs and eddies. We find that ~~for both,~~  $c_x$  is primarily negative and thus westward for both features. The mean values  $\bar{c}_x$  vary with latitude and decrease towards the  
355 equator, which indicates that both phenomena propagate faster towards the west at lower latitudes. ~~This~~ The average westward propagation ranges between ~~0 to 4 cm s<sup>-1</sup> on average~~ 0 and 4 cm s<sup>-1</sup>. The close agreement between the latitudinal means of  $c_x$  provides evidence that, on average, TRAPs move along with mesoscale eddies. We find similar coincidence for the latitudinal

and longitudinal distribution of ~~both~~ zonal and meridional propagation speeds  $c_x$  and  $c_y$ , see Fig. A1. However, we find that mesoscale eddies exhibit more extreme propagation speeds, suggesting that eddies might only create persistent strain within a  
 360 specific dynamic range or ~~a specific~~ around a certain combination of eddies.




**Figure 5.** Comparisons between TRAPs and mesoscale eddies. (a) ~~Distribution~~ Distribution of lifetime  $\Lambda$  over all TRAP trajectories, cyclonic and anticyclonic eddy trajectories in the domain. The distributions of eddy lifetimes are cut off at 300 days for clarity. (b) Evolution of attraction rate  $s_1$  over TRAP lifetime and contour speed  $U$  over eddy lifetime for phenomena with lifetimes  $\Lambda > 30$  days. (c) Latitudinal dependence of zonal propagation speed  $c_x$  for 2,951,028 TRAP, 604,296 cyclonic and 608,650 anticyclonic eddy instances. Lines in (b) and (c) indicate bin means, shaded bands their errors w.r.t. a confidence level of ~~95%~~ 95%. Mesoscale eddies as detected by AVISO+ et al. (2022).

Our results indicate that TRAPs are ~~certainly~~ located within the mesoscale eddy field, but we wonder where exactly. While TRAPs exert maximum normal attraction and induce significant deformation, i.e. strain-dominated regions, coherent eddies are characterised by water parcels rotating about a common axis and closed transport barriers. Hence, coherent eddies represent vorticity-dominated regions which should exclude TRAPs. As a consequence, TRAPs should emerge at the eddy periphery.  
 365 However, we find that the eddy detections from AVISO+ et al. (2022) ~~are not able to~~ cannot explain all TRAP occurrences. We often ~~find~~ observe TRAPs at eddy boundaries, ~~but we also frequently observe~~. But we also frequently find TRAPs in regions with no eddy detections or even inside eddy contours, see Fig. A2 or Video [SV3-S3](#) for details. The latter occurs for 15 % of all TRAP detections and suggests the presence of multiple eddies within one eddy detection. Even though the eddy dataset allows us to reveal average relations between TRAPs and mesoscale eddies, it is not suited ~~to describe~~ for describing the actual  
 370 dynamics around individual TRAPs. Instead, we use the relative vorticity field to characterise the flow around TRAPs.

### 3.3 Vorticity patterns around TRAPs

A considerable number of TRAPs seems to be surrounded by four vortices of alternating polarity, which exhibit perpendicular flow towards the TRAP core and tangential flow away from it w.r.t. the TRAP curve. We call this pattern a *quadrupole*. Here, we unravel this quadrupole and variations of it to demonstrate the driving mechanisms behind the formation of mesoscale  
 375 TRAPs. We compute the vorticity curves  $\zeta(\alpha)$  around all available TRAPs and normalise each by its maximum absolute value to obtain  $\hat{\zeta}(\alpha)$ . First, we take this large ensemble of curves and show its mean and standard deviation in Fig. 6. ~~This~~ It reveals the mean pattern in the vorticity field around a TRAP.




**Figure 6.** Ensemble of 3,568,850 vorticity curves  $\hat{\zeta}(\alpha)$  around TRAPs. Every curve  $\zeta(\alpha)$  is normalised by its maximum absolute value.  $\hat{\zeta} > 0 \Rightarrow \hat{\zeta}(\alpha) > 0$  indicates cyclonic,  $\hat{\zeta} < 0 \Rightarrow \hat{\zeta}(\alpha) < 0$  anticyclonic rotation in the surrounding flow field of a TRAP. The blue line indicates the ensemble mean, the shaded band renders its standard deviation. Black lines represent an arbitrary 0.1%–0.1% subset from this ensemble. The panels on the right sketch illustrate two TRAPs, one surrounded by a quadrupole as it can be interpreted from suggested by the ensemble mean, the other surrounded by vortices of unknown polarity: What vorticity patterns will appear around individual samples?

The ensemble of all vorticity curves clearly resembles a sine wave. The similarity between the ensemble mean and a harmonic function, as well as its smoothness, is remarkable. Even though individual curves might not follow this shape, the entirety of the approx. approximately 3.5 million curves generates a robust signal. Since a vorticity curve with four zero crossings and a polarity pattern of ~~eyelonie~~, ~~anticyelonie~~, ~~eyelonie~~, ~~anticyelonie~~ cyclonic-anticyclonic-cyclonic-anticyclonic sequence indicates a quadrupole, the above signal gives reason to believe that the mean pattern in the vorticity field around a TRAP is a quadrupole. Figure 3 illustrates such a typical quadrupole situation.

We then remove the background vorticity from each vorticity curve and filter the ensemble of all vorticity curves  $\zeta(\alpha)$  for the 10 distinct vortex combinations. The quadrupole pattern in Fig. 6 serves as our reference pattern, and we define the other nine patterns in terms of variation from it. Therefore, we introduce the *quadrupole order*  $q$ . It describes the number of vortices in a given pattern that need to change polarity in order to obtain the reference quadrupole pattern. We group the vorticity patterns of all the above 3,568,850 TRAP instances by  $q$  and illustrate the most frequent groups in Fig. 7, ~~see Fig. S2 for more~~. See Fig. S4 for details on all 10 vorticity patterns and their attributed quadrupole order.





**Figure 7.** Quadrupole orders and their frequency. The quadrupole order  $q$  describes the number of vortices in a given pattern that need to change polarity in order to obtain the reference quadrupole pattern. After removing the background vorticity from every vorticity curve  $\hat{\zeta}(\alpha)$ , 99.97 % of the ensemble can be explained by the five patterns illustrated here. Figure S2-S4 provides an overview on all vorticity patterns.


We shorthand name TRAP instances within a specific quadrupole surrounding  $q$ -th-order  $q$ -th-order quadrupoles. Zero-, first and second-order Zero-, first- and second-order quadrupoles constitute 99.97 % of the signal in Fig. 6. We identify approximately 57 % of all TRAPs as zero-order-zero-order quadrupoles, 34 % as first-order-first-order quadrupoles and 8 % as second-order-second-order quadrupoles. As we expect, the most prevalent group is the reference quadrupole indicated by the ensemble mean. However, a significant share of TRAPs is also surrounded by first order quadrupoles surrounded by a first order quadrupole. A quick look into the ensemble means in Fig. S2 reveals that first order S4 reveals that first-order quadrupoles can also include dipoles with one dominating polarity and some second-order, and some second-order quadrupoles might represent symmetric dipoles. Higher-order Higher-order quadrupoles also exist and. They involve the lowest attraction rates, but we neglect them since they rarely occur.

400

We now Next, we want to know how quadrupole patterns evolve over the TRAP lifetime  $\Lambda$  and how they relate to the attraction rate  $s_1$ . Panel (a) in Fig. 8 illustrates the shares of quadrupole orders zero, one and two at different evolution stages of long-living TRAPs. We observe a gradual change in the vorticity field around long-living TRAPs towards the reference quadrupole state during the first half of a life cycle, and away from the reference quadrupole state during the second half. Zero-order Zero-order quadrupoles are the most important vortex group throughout the entire lifetime, and they are especially abundant during the mature phase of TRAPs, i.e. between 20 % and 80 % of lifetime. However, the first-order quadrupole is also a likely first-order quadrupole is a comparably probable state at the formation and decay phase of this cycle.

405

A comparison between Figs. 8a and 5b suggests that the evolutions of quadrupole order  $q$  and TRAP attraction rate  $s_1$  go along with each other, i.e. TRAPs intensify towards their mid-life while their vorticity surroundings approach a **lower-order** **lower-order** quadrupole state. From this, we expect that **zero-, first and second-order zero-, first- and second-order** quadrupoles create different kinds of strain. Panel (b) in Fig. 8 displays the **distributions-distribution** of the instantaneous attraction rates  $s_1$  w.r.t. these quadrupole orders. The weakest TRAPs, i.e. with the **largest-highest** strain rates  $s_1$ , are only surrounded by **first order quadrupoles. Zero- and second-order first-order quadrupoles. Zero- and second-order** quadrupole environments appear for slightly higher attractions, i.e. smaller **values-of-strain rates**  $s_1$ . All three distributions peak within the same niche **but with** **but with a** decreasing attraction rate, the probability **density-for-second-and-first-order densities for second- and first-order** quadrupoles decline faster than for the reference quadrupole. It shows that the probability **to-find-of-finding** strongly attracting TRAPs is higher among quadrupoles of order zero. The average attraction rates  $\bar{s}_1$  confirm this tendency, **and** the p-values for the respective differences in mean values result in  $p < 0.001$ . However, the strongest TRAPs are attributed to both the **zero-and** **first-order quadrupole environments and zero- and first-order quadrupole environments, and we note that** in panel (b), we are looking at slight differences at the **very** extremes. Therefore, we conclude that there is a trend towards the reference quadrupole environment for increasingly attractive TRAPs. **Nevertheless, first-order, while first-order** quadrupoles are likewise able to induce high strain. We now investigate which of these groups organises tracer patterns particularly well.




**Figure 8.** Distribution of quadrupole orders over TRAP lifetime and attraction. (a) Shares of **zero-zero-, first-first- and second-order second-order** quadrupoles over the lifetime of 26,675 long-living TRAPs with  $\Lambda > 30$  days. (b) Distribution of TRAP attraction **rates-rate**  $s_1$  for **zeroall zero-, first-first- and second-order second-order** quadrupoles **within the ensemble of vorticity curves  $\hat{c}(\alpha)$ .**

### 3.4 Impact on drifters

We identify **33.878-33.878** drifter-TRAP pairs with retention times of  $\varphi > 1$  day. These pairs cover 73 % of all drifter days and exhibit a mean retention time of  $\bar{\varphi} \approx (4.8 \pm 3.7)$  days which reflects the transient impact of TRAPs, i.e. drifters are **accumulated-attracted** and dispersed again within a few days. However, we also find a few drifters that spend multiple weeks around a TRAP. The highest retention time we measure counts  $\varphi = 46$  days. Pairs with retention times  $\varphi > 7$  days represent only 9 % of all pairs but cover 28 % of all drifter days. **In general, we We generally** observe similar behaviours of drogued and undrogued drifters around TRAPs, **but. But** we emphasise a subtle difference in retention times to explain why we find 2.6 times more undrogued than drogued drifter-TRAP pairs while there are only 1.6 times more undrogued than drogued drifters



in our domain. Figure 9 presents the distribution of retention times  $\varphi$  over these drifter-TRAP pairs w.r.t. a pair's drogue state. The probability density function illustrates, irrespective of sample size, which drogue type is more likely to enter into a long retention. With increasing retention time  $\varphi$ , the probability density for undrogued drifters declines faster than for drogued ones. In general, drogued drifters show a higher susceptibility to stay around TRAPs for 8-eight days and longer. ~~A-Eventually, a~~ longer retention of drogued drifters ~~eventually~~ makes them less available for transient pairings.



**Figure 9.** Retention times of drifters around TRAPs. ~~Probability~~ Filled bars indicate the probability density function of retention times  $\varphi$  with respect to a drifter's drogue type for drogued drifters. Empty bars present the probability density for undrogued drifters. Drogued drifters remain on average  $\bar{\varphi} \approx (5.3 \pm 4.3)$  days, undrogued ones  $\bar{\varphi} \approx (4.6 \pm 3.4)$  days around a TRAP. The p-value for the difference in means results in  $p < 0.001$ . Pairs with retention times of ~~±one~~ day are excluded.

We study the average drifter motion around TRAPs and find that transport patterns depend on the evolution stage of the respective TRAP. Thus, we divide all drifter-TRAP pairs into four groups. The first group includes pairs for which the respective TRAP forms and decays during the drifter visit. Technically, this means that a pair ~~includes covers~~ the first and ~~the~~ last detection of a TRAP. The second group consists of pairs which ~~occur during start at~~ the formation of a TRAP but end before its decay. The third group ~~entails represents~~ pairs which begin after the formation of a TRAP but end with its decay. ~~And, and~~ the fourth group defines pairs that exist throughout the lifetime of a TRAP but exclude its formation and decay. Table 1 summarises the average retention times, attraction rates and TRAP lifetimes related to these groups. We find the longest average retention time for pairs that occur throughout the life cycle of a TRAP without formation or decay. These pairs ~~show to involve~~ TRAPs of significantly stronger attraction and higher persistency compared to pairs that ~~occur during TRAP formation and/~~ ~~experience~~ TRAP formation or decay.


**Table 1.** Characteristics of drifter-TRAP pairs at specific TRAP evolution stages.  $\bar{\varphi}$  indicates the average retention ~~times-time~~ over all respective pairs,  $\bar{\Lambda}$  the average lifetime over all associated TRAPs and  $\bar{s}_1$  the average attraction rate over all associated TRAP instances. TRAP formation is given if a pair includes the first detection of a TRAP, ~~and~~ TRAP decay if a pair includes the last detection.

	TRAP formation	TRAP decay	$\bar{\varphi}$ [days]	$\bar{\Lambda}$ [days]	$\bar{s}_1$ [s <sup>-1</sup> ]	# pairs	share in drifter days
Group I	yes	yes	(3.5 ± 2.6)	(3.5 ± 2.6)	(-0.17 ± 0.07)	<del>5456</del> <u>5,456</u>	8.5 %
Group II	yes	no	(4.8 ± 3.8)	(19.6 ± 20.8)	(-0.21 ± 0.08)	<del>5521</del> <u>5,521</u>	11.8 %
Group III	no	yes	(4.7 ± 3.7)	(19.6 ± 20.3)	(-0.21 ± 0.08)	<del>5609</del> <u>5,609</u>	11.8 %
Group IV	no	no	(5.3 ± 3.8)	(47.9 ± 35.4)	(-0.27 ± 0.11)	<del>17292</del> <u>17,292</u>	40.9 %

In Fig. 10, we illustrate the average drifter motion around TRAPs with respect to this grouping. We rotate drifter tracks around TRAPs towards the zonal axis and allocate their respective drifter positions to hexagonal bins. For every bin, we average the elapsed retention time  $\varphi$  ~~and~~, the eastward and ~~the~~ northward velocity components over all binned drifter positions. 450 Bin colours indicate the average elapsed retention time  $\varphi$  per bin, ~~quivers-vectors~~ the average velocity components.

Panel (a) highlights the average drifter motion around TRAPs that form and decay. The situation appears ~~rather chaotic somewhat disorganised~~, with no specific motion pattern to detect. Velocity ~~quivers-vectors~~ hardly show any preferred direction of ~~transport-drifter transport~~, and green colours dominate the picture. The colour map indicates that the retention times of 455 the underlying drifter-TRAP pairs begin and end anywhere throughout the search circle. Panel (b) gives insight into drifter motion during the formation of TRAPs ~~which overlast that persist beyond~~ the retention time. We find that drifters ~~are majorly transported-mostly move~~ away from the TRAP and in parallel to the TRAP curve. Panel (c) describes the opposite situation in which decaying TRAPs ~~rather attract-attract rather~~ than disperse drifters. This attraction is ~~generally directed-directed mainly~~ towards the TRAP core. Panels (a) to (c) reveal that drifters around TRAPs that ~~either form-and/form~~ or decay do not tend to 460 follow a hyperbolic pattern. But TRAPs that are at ~~a-the~~ final stage of their life cycle can at least indicate the confluence of material.

In contrast, the velocity ~~quivers-vectors~~ in panel (d), where TRAP formation and decay are excluded, describe hyperbolic transport as expected, ~~i.e. drifters~~. ~~Drifters~~ flow perpendicularly towards the TRAP core and tangentially away from it, with 465 respect to the TRAP curve. We also observe distinct regions of ~~light-blue and red bin colours~~ ~~which indicate~~, ~~indicating~~ that many drifters enter the zone perpendicular towards the TRAP and leave it at one end of the structure. A comparison with Figs. 5b, 8a and Table 1 clarifies that hyperbolic transport primarily occurs throughout the mature phase of a long-living TRAP because, at this stage, the surrounding flow ~~is particularly organised-organises~~ into a quadrupole pattern that generates high strain.



**Figure 10.** Drifter motion around TRAPs. For the first TRAP detection of every drifter-TRAP pair, we determine the angle between the longer branch of vector pointing from the first TRAP instance core to the furthest curve point and the zonal axis, with  $0^\circ$  pointing eastward and angles increasing counterclockwise. We use this angle to rotate all drifter positions towards the zonal axis. We allocate all rotated drifter positions to hexagonal bins and for. For every bin, we average the elapsed retention time  $\varphi$  as well as the eastward and northward velocity components over all binned drifter instances. The colour mapping indicates bin averages of  $\varphi$ , quivers-vectors the average velocity components. A transparent TRAP in the middle indicates-represents a generic profile, a the black circle draws the limits of the drifter search zone around it. (a) TRAP forms and decays, (b) TRAP only forms, (c) TRAP only decays and (d) TRAP neither forms nor decays during the drifter visit.

#### 470 4 Summary, discussion and conclusion

We studied the characteristics of TRAPs and the prospects for predicting debris transport from this mesoscale permitting dataset. Our findings provide an advanced understanding of TRAPs in the Great Pacific Garbage Patch and demonstrate the importance of TRAP lifetime estimations to an operational application. We find that the life cycle of TRAPs can range from days to seasons with an average lifetime of  $\bar{\Lambda} \approx (6 \pm 12)$  days. However, 41 % of aH-TRAP detections relate to profiles with lifetimes of  $\Lambda > 30$  days. Such long-living TRAPs exhibit a distinct evolution of attraction strength, ~~they~~. They intensify during the first and weaken during the second half of their life cycle. At the same time, the vorticity field around TRAPs gradually changes towards and away from a ~~particularly~~ specifically ordered state, i.e. a group of four vortices with alternating polarity. Therefore, the life cycle of TRAPs can explain why we observe hyperbolic drifter transport primarily throughout the mature phase of long-living TRAPs. At this stage, the surrounding flow ~~is particularly organised and generates high~~ creates the optimal vorticity pattern for generating high, hyperbolic strain. We find that hyperbolic transport around TRAPs takes on average  $\bar{\varphi} \approx (5.3 \pm 3.8)$  days ~~but also detect~~. In general, retention times can be very short, and especially, strong TRAPs quickly attract and disperse material. But we also detect a few drifters that spend multiple weeks around a TRAP. The highest retention time we measure counts  $\varphi = 46$  days.

485 We identify the evolution stage of TRAPs as the most significant predictor for drifter motion. ~~But~~ However, the coherence of surrounding vortices might further explain our observations of hyperbolic transport and long retentions. We propose to investigate this with higher-resolution measurements of the flow, ~~which~~ we see around many TRAPs in [SV3Video S3](#). The mesoscale eddy detections by AVISO+ et al. (2022) neither capture such vortices at the limit between the meso- and submesoscale ~~which reflects in~~, which explains the high number of TRAPs without close-by eddy detections. Moreover, we find 15 % of aH-TRAPs inside these eddy detections, ~~although coherent eddies should exclude strain-dominated regions. It is known that eddy detection algorithms based on closed contours of sea surface height often yield false-positive identifications (Beron-Vera et al., 2013; Andrade-Canto and Beron-Vera, 2022). Liu and Abernathey (2023) present an alternative algorithm which detects eddies from Lagrangian-averaged vorticity deviations (LAVD, Haller et al. (2016)). They claim the absence of false-positive detections, however~~ false-positive detections. However, we still find TRAPs that intersect these eddy detections, possibly due to the discrepancy between the ~~time-scale~~ time scale over which Lagrangian averaging takes place and the instantaneous nature of TRAPs, see Fig. A2 for details. We expect true coherence and no inclusion of TRAPs when we use a detection framework that descends from the deformation tensor  $\mathbf{S}(\mathbf{x}, t)$ . Serra and Haller (2016) introduce elliptic OECSs, ~~which can be derived from singularities of  $\mathbf{S}(\mathbf{x}, t)$  and build a complement to the strain-dominated regions uncovered by TRAPs, with the additional benefit of both~~ computations being instantaneous methods being applicable to Eulerian snapshots of velocity. A computational implementation of elliptic OECSs would be subject to future research.

There is an ongoing debate on whether mesoscale eddies accumulate and transport floating material. van Sebille et al. (2020) discuss confirming examples such as Brach et al. (2018), Budyansky et al. (2015) and Dong et al. (2014) ~~while~~ 505 ~~Abernathy and Haller (2018) and Wang et al. (2015) show that transport by coherent eddies is negligible. This controversy rises from the use of different concepts to identify eddy boundaries.~~ Abernathy and Haller (2018) ~~find~~ Early et al. (2011) use the zero contour of relative vorticity to define eddy boundaries in an idealised flow. They show how an anticyclonic eddy core perfectly transports floats and tracers over large distances, but they also explain why fluid from the outside cannot be entrained by the core. The authors illustrate that the ring of fluid around an eddy core simultaneously entrains and sheds fluid from 510 and into the environment and can disperse material over different scales. Abernathy and Haller (2018), however, argue that transport by coherent eddies is negligible and that material transport is caused by stirring and filamentation at the periphery of strictly coherent eddies, rather than by the coherent motion within eddy cores. They emphasise the lack of need for objective methods to identify such peripheral regions. ~~But TRAPs describe the limit between eddy peripheries and should be able to contribute to this investigation. We find that on large time scales, there is no preference of drifter positions around TRAPs or~~ 515 ~~within eddy boundaries defined by AVISO+ et al. (2022).~~ The Froyland et al. (2015) demonstrate an objective approach using finite-time coherent sets from a transfer operator that minimises mass loss from eddy boundaries. They track the long-term decay of an observed Agulhas ring and estimate the proportion of surface water leaking from this coherent structure. Using their theory, Denes et al. (2022) derive finite-time coherent sets from a dynamic Laplace operator and estimate the material transport provided by the periphery of a modelled Agulhas ring. They show that the quasi-coherent outer ring of this eddy 520 significantly contributes to the entrainment and retention of fluid. TRAPs are intrinsic to these peripheral regions, and the concept should facilitate further understanding of these processes.

As a first approach to this, we derived a 20-year time series of the number of daily drifter positions spent around TRAPs and within mesoscale eddies detected by AVISO+ et al. (2022), and compared the retention of drifters by both features, see Fig. 525 S5 for details. We find that, on average, the share of drifter days spent around TRAPs or within eddies approx. approximately equals the proportion of surface area that is covered by these structures. However, there are temporary preferences on short time scales which motivate a follow-up analysis, see ~~This result suggests no preference of drifter positions for any of these features. However, the number of drifters within our domain barely supports the analysis, and at times with few drifters, we observe high standard deviations in the respective shares of drifter days. Drifters become more abundant towards the end of~~ 530 ~~the time series in Fig. S3 for details,~~ where the share of drifter days around TRAPs tends to surpass the respective proportion of covered surface area, contrary to eddy detections. More studies are needed to clarify if a preference of drifter positions can be observed in experiments with more drifters, a larger domain, and on what time scale this might occur.

There are obvious limits to the application of this mesoscale permitting dataset. The effective resolution in space and time 535 ~~can be expected to be greater~~ should be coarser than the  $0.25^\circ$  latitude-longitude grid and the daily frequency. We observe the effects of this in animations where TRAPs reappear after a one-day gap. At such a gap, our tracking algorithm defines a new trajectory ~~and therefore TRAP lifetimes might be underestimated. In a similar way,~~ and therefore, we might underestimate

TRAP lifetimes. Similarly, detection gaps affect the identification of drifter-TRAP pairs, leading to an underestimation of retention times. Nevertheless, we find a remarkable consistency between Duran et al. (2021) and our study. They find persistent mesoscale TRAPs that predict the spread of surface oil at least eight days in advance ~~and confirm~~, in agreement with our finding of retention times of  $\varphi = (5.3 \pm 3.8)$  days for hyperbolic drifter motion. ~~The agreement between their study and ours, as well as~~ These comparable time scales and the overall similar behaviour we observe for drogued and undrogued drifters ; ~~further underlines~~ further underline the concept's robustness against differences in tracer properties.

545 Our results for drifter-TRAP pairs are observations of the hyperbolic-type Lagrangian motion induced by TRAPs, and therefore, our findings confirm the persistence of TRAPs over periods considerably longer than a few hours, which is the lifetime of a TRAP that is mathematically guaranteed to exist. We know from our drifter-TRAP statistics that, over about a week, drifters are attracted normally to a TRAP to then accelerate and leave the TRAP in a tangential direction. Given that the drifter and altimetry datasets are independent oceanic observations, these statistics show that TRAPs often persist for at  
550 least a week. Importantly, we note that the behaviour of drifters in the vicinity of forming or decaying TRAPs is distinct from the hyperbolic behaviour observed near TRAPs that are neither forming nor decaying. This result shows that we are following the same TRAP and not following different TRAPs that quickly form and decay at locations that coincide with the path of propagating eddies. Due to the relatively short transport time of a drifter in the vicinity of a TRAP, we would expect that a TRAP can persist for considerably longer than a week. Indeed, the persistent relation between TRAPs and mesoscale  
555 vorticity structures that we report, including the similarity in their propagation speed, suggests that the lifetimes of mesoscale TRAPs are often related to the lifetime of long-lived mesoscale structures. Duran et al. (2021) present another example of the temporal continuity of TRAPs and their influence on hyperbolic tracer deformation, again from independent observations. The hyperbolic nature of this latter deformation pattern is established in Olascoaga and Haller (2012) and Duran et al. (2018).


560 Even though our study cannot resolve important submesoscale processes like ~~e.g.~~ filamentation, Langmuir cells or submesoscale vortices, it demonstrates the capability of TRAPs to predict material transport from ~~first-order~~ first-order mesoscale observations. We observed the effect of TRAPs on drifters, both drogued and undrogued, that sample any oceanic structures found along their path, including submesoscales. Thus, our results include the aggregate effect of such flow components on the daily drifter positions that we use in the analysis, even though these flow components are not resolved by the satellite altimetry  
565 itself. Our algorithms offer a great chance to reapply the TRAPs concept to future high-resolution observations that will be provided by the current SWOT mission (International Altimetry Team, 2021). TRAPs computed from submesoscale velocities will probably show different characteristics than their mesoscale counterparts but are crucial to an application of the concept during offshore cleanups. An interesting approach would be to study the flow around drifter-TRAP pairs with long retention times. These long retention times might be due to drifter trapping within submesoscale vortices and filaments that result from  
570 instabilities at mesoscale fronts (van Sebille et al., 2020; Zhang et al., 2019). Mesoscale TRAPs indicate these mesoscale fronts and might provide a window to enhanced material clustering at the submesoscale. ~~We propose to recast~~, where we also expect higher numbers of small-scale TRAPs. However, it remains to be investigated whether mesoscale TRAPs from

575 large-scale observations of sea surface height  $\mathcal{O}(10 - 100 \text{ km})$  can be used as a proxy for material accumulation at operational scales  $\mathcal{O}(1 - 10 \text{ km})$ . Subject to current research is the recast of our TRAPs record (Kunz, 2024b) ~~for~~ to periods with available SWOT measurements~~to investigate whether TRAPs can indicate material accumulation at operational scales  $\mathcal{O}(1 - 10 \text{ km})$  but from large-scale observations of sea surface height  $\mathcal{O}(10 - 100 \text{ km})$ , even if,~~ with resolution about 1 order of magnitude higher than the data used here. Since the geostrophic assumption is needed to obtain a sea-surface velocity from SWOT measurements. ~~Such,~~ such exploration can be complemented with additional observations in coastal regions from high-frequency radar, which gives the full and not only the geostrophic sea-surface velocity. These surveys can further improve the applicability of TRAPs  
580 to offshore cleanups.

Our results can ~~already support offshore cleanup operations~~ support cleanup operations in the Great Pacific Garbage Patch since they reveal which TRAPs are the most likely to indicate a large-scale confluence of drifting objects. ~~Operators~~ At the mesoscale, operators should search for long-living TRAPs that are at an advanced stage of their life cycle. These TRAPs  
585 streamline floating objects into hyperbolic pathways. Such a streamlined bypass involves a short but strong attraction~~which could,~~ which can be exploited to *filter* the flow around a TRAP. ~~But we also find TRAPs that retain material over multiple weeks. These long-term retentions can start at any stage of a TRAP's life cycle and drifters remain closer to the TRAP core than for streamlined bypasses. An investigation of the conditions causing this behaviour could further improve the prospects of a cleanup operation, especially since long retentions of  $\varphi > 7$  days are rare but cover up to 28% of all drifter days. And~~ The  
590 state-of-the-art cleanup system consists of a two-kilometres-long surface barrier, towed by two vessels (The Ocean Cleanup, 2023) , and it could move along TRAPs to act like a filter on the through-flowing water. The large-scale navigation along mesoscale TRAPs could then be complemented by vessel-based methods that enable the detection of debris and attracting flow features on the small scales, for instance, through automated object recognition using cameras (de Vries et al., 2021) or shipboard marine X-band radar (Lund et al., 2018). More knowledge about the relation between submesoscale and mesoscale TRAPs could  
595 further contribute to bridging observational gaps between these scales.


Eventually, this research is not limited to the subject of marine debris. ~~Our findings may also benefit other challenges like e.g. optimal drifter deployment, sargassum removal, the identification of foraging hotspots or,~~ and various offshore applications can benefit from the detection and tracking of these hyperbolic structures. For instance, authorities might use TRAPs to mitigate  
600 sargassum transport towards ports and coastal areas where beaching events cause limited accessibility. In case TRAPs indicate enhanced small-scale clustering of organic compounds, biologists could use them to monitor and protect the foraging of marine species. Oceanographers can apply the TRAPs concept to optimise drifter deployments whenever drifters should separate fast or remain in a specific region. Furthermore, TRAPs are suitable for estimating oil transport at the ocean surface, making them a considerable tool for the emergency response to oil spills. And finally, a better understanding of TRAPs will help establish  
605 their use in the essential search and rescue operations that are ever-saving lives at sea.

## Appendix A: Figures



**Figure A1.** Comparison of propagation speeds between TRAPs and mesoscale eddies. Latitudinal dependence of the (a) zonal propagation speed  $c_x$  and (b) meridional propagation speed  $c_y$ . Longitudinal dependence of the (c) zonal propagation speed  $c_x$  and (d) meridional propagation speed  $c_y$ . Lines indicate bin means, shaded bands the respective standard deviations from 2,951,028 TRAP, 604,296 cyclonic and 608,650 anticyclonic eddy instances. Mesoscale eddies as detected by AVISO+ et al. (2022).





**Figure A2.** Snapshot of mesoscale TRAPs and mesoscale eddy detections within the study domain. Black lines represent TRAPs. Cyan/pink lines depict the speed contours of cyclonic/anticyclonic mesoscale eddies as detected by AVISO+ et al. (2022). Blue/red patches depict Lagrangian particles trapped by cyclonic/anticyclonic coherent eddy boundaries as detected by Liu and Abernathey (2022) for the same snapshot. The colour scale on the right indicates the relative vorticity  $\zeta(\mathbf{x}_\ell, t_0)$  of Lagrangian particles at position  $\mathbf{x}_\ell$ , with  $\ell \in [1, n]$  being the index for each of the  $n$  Lagrangian particles at snapshot time  $t_0$ .

## Appendix B: Tables

**Table B1.** Benefits of TRAPs. Adopted from Serra and Haller (2016), Serra et al. (2020) and Duran et al. (2021).

<b>use velocity snapshots</b>	TRAPs can be directly inferred from snapshots of the surface velocity field and thus are light in computation. They allow for input data with time gaps, and predictions can be made from a nowcast of the velocity field.
<b>avoid pitfalls of trajectory integration</b>	TRAPs do not require costly trajectory integrations, tracer release times, locations or the length of the observation period. Velocity errors do not accumulate, which should make TRAPs less sensitive to uncertainties in the underlying velocity field. <del>They TRAPs</del> also provide full domain coverage <del>while Lagrangian methods. Lagrangian methods, on the other hand,</del> become computationally demanding with increasing resolution of the initial particle field where particles may eventually leave a finite-size domain.
<b>indicate direction of transport</b>	What makes TRAPs special is that they indicate the directions of transport while other structures like <del>e.g. <math>s_T s_1(x, t)</math></del> minima, divergence minima, or ridges of the Finite-Time Lyapunov Exponent (FTLE) field don't. FTLE ridges do not necessarily indicate the direction of largest stretching (Haller, 2015).
<b>easy to interpret</b>	TRAPs describe localised, one-dimensional curves of attraction in contrast to large open sets of <del>e.g. horizontal divergence or particle density.</del>
<b>robust</b>	TRAPs are robust to different inertia and windage effects and thus are insensitive to <del>the</del> varying shape or submergence level of drifting objects.
<b>uncover hidden flow structures</b>	TRAPs can be perpendicular to streamlines and exist in divergence-free flows.
<b>observer-independent</b>	The objective nature of TRAPs leads to the same conclusions on different platforms. Classic Eulerian quantities like streamlines, velocity magnitude, velocity gradient, energy, <del>or</del> relative vorticity are not objective and will lead to different results in different frames of reference.
<b>scale-invariant</b>	TRAPs can be computed for velocity fields of any temporal and spatial resolution. The concept can be applied to the scales of the confluence phenomena of interest.

**Table B2.** Algorithm to compute Transient Attracting Profiles in a two-dimensional flow. Adopted from Serra and Haller (2016) and Serra et al. (2020).

---

**Input:** A two-dimensional Eulerian velocity field  $\mathbf{u}(\mathbf{x}, t)$  **Compute**

the rate-of-strain tensor  $\mathbf{S}(\mathbf{x}, t) = \frac{1}{2}(\nabla\mathbf{u}(\mathbf{x}, t) + [\nabla\mathbf{u}(\mathbf{x}, t)]^\top)$  at the current time  $t = t_0$ , defined on a rectangular grid over the  $(x_1, x_2)$  coordinates.  $\mathbf{x} = (x_1, x_2)$  coordinates.

**Compute**

1. the instantaneous rate-of-strain tensor  $\mathbf{S}(\mathbf{x}, t_0) = \frac{1}{2}(\nabla\mathbf{u}(\mathbf{x}, t_0) + [\nabla\mathbf{u}(\mathbf{x}, t_0)]^\top)$
2. the eigenvalue fields  $s_1(\mathbf{x}, t) \leq s_2(\mathbf{x}, t)$  and the associated unit eigenvector fields  $\mathbf{e}_i(\mathbf{x}, t)$  of  $\mathbf{S}(\mathbf{x}, t)$  for  $i = 1, 2$ .
3. the set  $S_{1m}(t)$  of negative local minima of  $s_1(\mathbf{x}, t)$ .
4. TRAPs ( $e_2$ -lines) as solutions of the ODE

$$\begin{cases} \frac{d\mathbf{x}}{ds} = \text{sign}\langle \mathbf{e}_2(\mathbf{x}(s), t_0), \mathbf{e}_2(\mathbf{x}(s - \Delta), t_0) \rangle \cdot \mathbf{e}_2(\mathbf{x}(s), t_0) \\ \mathbf{x}(0) \in S_{1m}(t_0). \end{cases}$$

where  $s$  is the independent variable of the TRAP curve parameterisation  $\mathbf{x}(s)$  at time  $t_0$ ,  $\Delta$  denotes the integration step in the arclength parameter  $s$  and  $\mathbf{x}(0)$  indicates the position of a TRAP core. The sign term in step 4 guarantees the local smoothness of the direction field  $\mathbf{e}_2(\mathbf{x}, t_0)$ . Stop integration when  $s_1(\mathbf{x}(s)) - s_1(\mathbf{x}(s), t_0)$  ceases to be monotone increasing or when  $s_1(\mathbf{x}(s)) > 0.3 \cdot s_1(\mathbf{x}(0))$ . TRAP curves are shortened such that the attraction rate along the curve is everywhere at least 30 % of the attraction at the respective core in order to ensure that TRAPs indicate a distinguished attraction relative to nearby structures (Serra et al., 2020).

**Output:** Transient Attracting Profiles at time  $t_0$  and their normal attraction rate field  $s_1(\mathbf{x}, t)$ .

---

*Code and data availability.* We provide source code at Kunz (2024a) to post-process TRAP computations, ~~to~~ track TRAPs through the domain and ~~to~~ identify drifter-TRAP pairs. We ~~further also~~ provide a ~~20-years-20-year~~ dataset of TRAP detections with lifetime estimations ~~and~~ vorticity pattern detections and our dataset of drifter-TRAP pair detections at Kunz (2024b).

*Video supplement.* We provide animations of the TRAPs tracking algorithm in [SV1-Video S1](#), of the evolution of TRAPs, drifter positions and the relative vorticity field in [SV2-Video S2](#), and of the evolution of TRAPs, mesoscale eddy detections and the relative vorticity field in [SV3-Video S3](#). [SV1-Videos S1](#), [SV2-S2](#) and [SV3-S3](#) are available at Kunz (2024c), and Fig. [S4-S6](#) summarises the details.

*Author contributions.* LK programmed the algorithms, did the analyses and wrote much of the text. AG supervised the writing process, contributed to interpreting the results and wrote text. AG and CE acquired the funding. CE contributed to editing the paper, interpreting the results and wrote text. RD contributed to developing the methodology, editing [the paper](#), ~~interpretation of~~ [interpreting the](#) results and wrote text. BSR contributed to [developing the](#) methodology, ~~interpretation~~ [interpreting the results](#) and acquired computational resources. All authors contributed to the design of the experiment. [LK, AG and RD prepared the revised version of the manuscript.](#)

*Competing interests.* The authors declare that they have no conflict of interest.

620 *Acknowledgements.* This paper is a contribution to the project L3 (Meso- to submesoscale turbulence in the ocean) of the Collaborative Research Centre TRR 181 “Energy Transfer in Atmosphere and Ocean” funded by the German Research Foundation (DFG) and [it](#) has been written in collaboration with The Ocean Cleanup. [The authors thank the anonymous referees for their careful review and helpful comments on the initial manuscript. LK has consulted the free versions of the AI tools DeepL and Grammarly for assistance in English writing.](#)

## References

- 625 Abernathey, R. and Haller, G.: Transport by Lagrangian Vortices in the Eastern Pacific, *Journal of Physical Oceanography*, 48, 667–685, <https://doi.org/10.1175/JPO-D-17-0102.1>, 2018.
- Andrade-Canto, F. and Beron-Vera, F. J.: Do Eddies Connect the Tropical Atlantic Ocean and the Gulf of Mexico?, *Geophysical Research Letters*, 49, 20, <https://doi.org/10.1029/2022GL099637>, 2022.
- AVISO+, CNES, SSALTO/DUACS, and IMEDEA: The altimetric Mesoscale Eddy Trajectories Atlas 3.2 Delayed-Time (META3.2 DT) Allsat Version, AVISO+ [dataset], <https://doi.org/10.24400/527896/a01-2022.005.220209>, 2022.
- 630 Beron-Vera, F. J., Wang, Y., Olascoaga, M. J., Goni, G. J., and Haller, G.: Objective Detection of Oceanic Eddies and the Agulhas Leakage, *Journal of Physical Oceanography*, 43, 1426–1438, <https://doi.org/10.1175/JPO-D-12-0171.1>, 2013.
- Brach, L., Deixonne, P., Bernard, M.-F., Durand, E., Desjean, M.-C., Perez, E., van Sebille, E., and ter Halle, A.: Anticyclonic eddies increase accumulation of microplastic in the North Atlantic subtropical gyre, *Marine Pollution Bulletin*, 126, 191–196, <https://doi.org/10.1016/j.marpolbul.2017.10.077>, 2018.
- 635 Budyansky, M., Goryachev, V., Kaplunenko, D., Lobanov, V., Prants, S., Sergeev, A., Shlyk, N., and Uleysky, M.: Role of mesoscale eddies in transport of Fukushima-derived cesium isotopes in the ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, 96, 15–27, <https://doi.org/10.1016/j.dsr.2014.09.007>, 2015.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91, 640 167–216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- CMEMS: Global Total Surface and 15m Current (COPERNICUS-GLOBCURRENT) from Altimetric Geostrophic Current and Modeled Ekman Current Reprocessing, E.U. Copernicus Marine Service Information (CMEMS) [dataset], <https://doi.org/10.48670/moi-00050>, 2022a.
- CMEMS: Global Ocean Gridded L 4 Sea Surface Heights And Derived Variables Reprocessed 1993 Ongoing, E.U. Copernicus Marine Service Information (CMEMS) [dataset], <https://doi.org/10.48670/moi-00148>, 2022b.
- 645 de Vries, R., Egger, M., Mani, T., and Lebreton, L.: Quantifying Floating Plastic Debris at Sea Using Vessel-Based Optical Data and Artificial Intelligence, *Remote Sensing*, 13, 3401, <https://doi.org/10.3390/rs13173401>, 2021.
- Denes, M. C., Froyland, G., and Keating, S. R.: Persistence and material coherence of a mesoscale ocean eddy, *Physical Review Fluids*, 7, 034 501, <https://doi.org/10.1103/PhysRevFluids.7.034501>, 2022.
- 650 Dong, C., McWilliams, J., Liu, Y., and Chen, D.: Global heat and salt transports by eddy movement, *Nature Communications*, 5, 3294, <https://doi.org/10.1038/ncomms4294>, 2014.
- Duran, R., Beron-Vera, F., and Olascoaga, M.: Extracting quasi-steady Lagrangian transport patterns from the ocean circulation: An application to the Gulf of Mexico, *Scientific Reports*, 8, 5218, <https://doi.org/10.1038/s41598-018-23121-y>, 2018.
- Duran, R., Nordam, T., Serra, M., and H. Barker, C.: Chapter 3 - Horizontal transport in oil-spill modeling, in: *Marine Hydrocarbon Spill Assessments*, edited by Makarynskyy, O., pp. 59–96, Elsevier, <https://doi.org/10.1016/B978-0-12-819354-9.00004-1>, <https://doi.org/10.48550/arXiv.2009.12954>, 2021.
- 655 Early, J. J., Samelson, R. M., and Chelton, D. B.: The Evolution and Propagation of Quasigeostrophic Ocean Eddies, *Journal of Physical Oceanography*, 41, 1535–1555, <https://doi.org/10.1175/2011JPO4601.1>, 2011.
- Froyland, G., Horenkamp, C., Rossi, V., and van Sebille, E.: Studying an Agulhas ring’s long-term pathway and decay with finite-time coherent sets, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 25, 083 119, <https://doi.org/10.1063/1.4927830>, 2015.
- 660

- Haller, G.: Lagrangian Coherent Structures, *Annual Review of Fluid Mechanics*, 47, 137–162, <https://doi.org/10.1146/annurev-fluid-010313-141322>, 2015.
- Haller, G., Hadjighasem, A., Farazmand, M., and Huhn, F.: Defining coherent vortices objectively from the vorticity, *Journal of Fluid Mechanics*, 795, 136–173, <https://doi.org/10.1017/jfm.2016.151>, 2016.
- 665 International Altimetry Team: Altimetry for the future: Building on 25 years of progress, *Advances in Space Research*, 68, 319–363, <https://doi.org/10.1016/j.asr.2021.01.022>, 2021.
- Kunz, L.: Track and analyse Transient Attracting Profiles in the Great Pacific Garbage Patch, Zenodo [code], <https://doi.org/10.5281/zenodo.12761097>, 2024a.
- Kunz, L.: Datasets to Transient Attracting Profiles in the Great Pacific Garbage Patch, Zenodo [dataset],  
670 <https://doi.org/10.5281/zenodo.10993736>, 2024b.
- Kunz, L.: Supplementary Videos to Transient Attracting Profiles in the Great Pacific Garbage Patch, Zenodo [dataset], <https://doi.org/10.5281/zenodo.10943728>, 2024c.
- Law, K. L., Morét-Ferguson, S. E., Goodwin, D. S., Zettler, E. R., DeForce, E., Kukulka, T., and Proskurowski, G.: Distribution of Surface Plastic Debris in the Eastern Pacific Ocean from an 11-Year Data Set, *Environmental Science & Technology*, 48, 4732–4738,  
675 <https://doi.org/10.1021/es4053076>, 2014.
- Lebreton, L., Slat, B., Ferrari, F., Sainte-Rose, B., Aitken, J., Marthouse, R., Hajbane, S., Cunsolo, S., Schwarz, A., Levivier, A., et al.: Evidence that the Great Pacific Garbage Patch is rapidly accumulating plastic, *Scientific Reports*, 8, 1–15, <https://doi.org/10.1038/s41598-018-22939-w>, 2018.
- Lindo-Atichati, D., Jia, Y., Wren, J. L. K., Antoniadis, A., and Kobayashi, D. R.: Eddies in the Hawaiian Archipelago Region: Formation, Characterization, and Potential Implications on Larval Retention of Reef Fish, *Journal of Geophysical Research: Oceans*, 125, 5,  
680 <https://doi.org/10.1029/2019JC015348>, 2020.
- Liu, T. and Abernathey, R.: A global Lagrangian eddy dataset based on satellite altimetry (GLED v1.0), Zenodo [dataset], <https://doi.org/10.5281/zenodo.7349753>, 2022.
- Liu, T. and Abernathey, R.: A global Lagrangian eddy dataset based on satellite altimetry, *Earth System Science Data*, 15, 1765–1778,  
685 <https://doi.org/10.5194/essd-15-1765-2023>, 2023.
- Lumpkin, R. and Centurioni, L.: Global Drifter Program quality-controlled 6-hour interpolated data from ocean surface drifting buoys, NOAA National Centers for Environmental Information [dataset], <https://doi.org/10.25921/7ntx-z961>, 2019.
- Lund, B., Haus, B. K., Horstmann, J., Graber, H. C., Carrasco, R., Laxague, N. J. M., Novelli, G., Guigand, C. M., and Özgökmen, T. M.: Near-Surface Current Mapping by Shipboard Marine X-Band Radar: A Validation, *Journal of Atmospheric and Oceanic Technology*, 35,  
690 1077–1090, <https://doi.org/10.1175/JTECH-D-17-0154.1>, 2018.
- Olascoaga, M. J. and Haller, G.: Forecasting sudden changes in environmental pollution patterns, *Proceedings of the National Academy of Sciences*, 109, 4738–4743, <https://doi.org/10.1073/pnas.1118574109>, 2012.
- Olbers, D., Willebrand, J., and Eden, C.: *Ocean Dynamics*, Springer Berlin, Heidelberg, <https://doi.org/10.1007/978-3-642-23450-7>, 2012.
- Onink, V., Wichmann, D., Delandmeter, P., and van Sebille, E.: The Role of Ekman Currents, Geostrophy, and Stokes Drift in the Accumulation of Floating Microplastic, *Journal of Geophysical Research: Oceans*, 124, 1474–1490, <https://doi.org/10.1029/2018JC014547>,  
695 2019.
- Pegliasco, C., Chaigneau, A., and Morrow, R.: Main eddy vertical structures observed in the four major Eastern Boundary Upwelling Systems, *Journal of Geophysical Research: Oceans*, 120, 6008–6033, <https://doi.org/10.1002/2015JC010950>, 2015.

- Röhrs, J., Sutherland, G., Jeans, G., Bedington, M., Sperrevik, A. K., Dagestad, K.-F., Gusdal, Y., Mauritzen, C., Dale, A., and LaCasce, J. H.: Surface currents in operational oceanography: Key applications, mechanisms, and methods, *Journal of Operational Oceanography*, 16, 60–88, <https://doi.org/10.1080/1755876X.2021.1903221>, 2021.
- Serra, M.: Compute Transient Attracting Profiles (TRAPs), GitHub [code], <https://github.com/MattiaSerra/TRAPs>, 2020.
- Serra, M. and Haller, G.: Objective Eulerian coherent structures, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 26, 053 110, <https://doi.org/10.1063/1.4951720>, 2016.
- 705 Serra, M., Sathe, P., Rypina, I., Kirincich, A., Ross, S. D., Lermusiaux, P., Allen, A., Peacock, T., and Haller, G.: Search and rescue at sea aided by hidden flow structures, *Nature Communications*, 11, 2525, <https://doi.org/10.1038/s41467-020-16281-x>, 2020.
- Slat, B.: First 100,000 KG Removed From the Great Pacific Garbage Patch, <https://theoceancleanup.com/updates/first-100000-kg-removed-from-the-great-pacific-garbage-patch/>, last access: 25 July 2024, 2022.
- The Ocean Cleanup: System 03: A Beginner’s Guide, <https://theoceancleanup.com/updates/system-03-a-beginners-guide/>, last access: 25  
710 July 2024, 2023.
- van Sebille, E., Griffies, S. M., Abernathy, R., Adams, T. P., Berloff, P., Biastoch, A., Blanke, B., Chassignet, E. P., Cheng, Y., Cotter, C. J., Deleersnijder, E., Döös, K., Drake, H. F., Drijfhout, S., Gary, S. F., Heemink, A. W., Kjellsson, J., Koszalka, I. M., Lange, M., Lique, C., MacGilchrist, G. A., Marsh, R., Mayorga Adame, C. G., McAdam, R., Nencioli, F., Paris, C. B., Piggott, M. D., Polton, J. A., Rühls, S., Shah, S. H., Thomas, M. D., Wang, J., Wolfram, P. J., Zanna, L., and Zika, J. D.: Lagrangian ocean analysis: Fundamentals and practices, *Ocean Modelling*, 121, 49–75, <https://doi.org/10.1016/j.ocemod.2017.11.008>, 2018.
- 715 van Sebille, E., Aliani, S., Law, K. L., Maximenko, N., Alsina, J. M., Bagaev, A., Bergmann, M., Chapron, B., Chubarenko, I., Cózar, A., et al.: The physical oceanography of the transport of floating marine debris, *Environmental Research Letters*, 15, 023 003, <https://doi.org/10.1088/1748-9326/ab6d7d>, 2020.
- Wang, Y., Olascoaga, M. J., and Beron-Vera, F. J.: Coherent water transport across the South Atlantic, *Geophysical Research Letters*, 42, 4072–4079, <https://doi.org/10.1002/2015GL064089>, 2015.
- 720 Zhang, Z., Qiu, B., Klein, P., and Travis, S.: The influence of geostrophic strain on oceanic ageostrophic motion and surface chlorophyll, *Nature Communications*, 10, 2838, <https://doi.org/10.1038/s41467-019-10883-w>, 2019.