

NO₃ reactivity during a summer period in a temperate forest below and above the canopy

Patrick Dewald¹, Tobias Seubert¹, Simone T. Andersen¹, Gunther N. T. E. Türk¹, Jan Schuladen¹, Max R. McGillen², Cyrielle Denjean³, Jean-Claude Etienne³, Olivier Garrouste³, Marina Jamar⁴, Sergio Harb⁵,
5 Manuela Cirtog⁵, Vincent Michoud⁶, Mathieu Cazaunau⁵, Antonin Bergé⁵, Christopher Cantrell⁵,
Sebastien Dusanter⁴, Bénédicte Picquet-Varrault⁵, Alexandre Kukui⁷, Chaoyang Xue^{1,7}, Abdelwahid Mellouki^{2,8}, Jos Lelieveld¹, and John N. Crowley¹

¹Atmospheric Chemistry Department, Max Planck Institute for Chemistry, 55128 Mainz, Germany

²Institut de Combustion, Aérothermique, Réactivité Environnement (ICARE), CNRS, 1C Avenue de la Recherche
10 Scientifique, CEDEX 2, 45071 Orléans, France

³CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

⁴IMT Nord Europe, Institut Mines-Télécom, Université de Lille, Center for Energy and Environment, 59000 Lille, France

⁵Université Paris Est Créteil and Université de Paris Cité, CNRS, LISA, F-94010 Créteil, France

⁶Université Paris Cité and Université Paris Est Créteil, CNRS, LISA, F-75013 Paris, France

15 ⁷Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS, Orléans, France

⁸University Mohammed VI Polytechnic (UM6P), Lot 660, Hay Moulay Rachid Ben Guerir, 43150, Morocco

Correspondence to: John N. Crowley (john.crowley@mpic.de)

Abstract. We present direct measurements of BVOC-induced nitrate radical (NO₃) reactivity (k^{VOC}) through the diel cycle in the suburban, temperate forest of Rambouillet near Paris (France). The data were obtained in a six-week summer period in
20 2022 as part of the ACROSS campaign (Atmospheric ChemistRy Of the Suburban foreSt). k^{VOC} was measured in a small (700 m²) clearing mainly at a height of 5.5 m above ground level, but also at 40 m (for 5 days/nights). At nighttime, mean values (and 25th-75th percentile ranges) of $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m}) = (0.24_{-0.06}^{+0.32}) \text{ s}^{-1}$ and $k_{\text{night}}^{\text{VOC}}(40 \text{ m}) = (0.016_{-0.007}^{+0.018}) \text{ s}^{-1}$ indicate a significant vertical gradient and low NO₃ reactivity above the canopy, whereas $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ showed peak values of up to 2 s⁻¹ close to the ground. The strong vertical gradient in NO₃ reactivity could be confirmed by measurements between 0 and 24 m on one
25 particular night characterised by a strong temperature inversion, and is a result of the decoupling of air masses aloft from the ground- and canopy-level sources of BVOCs (and NO). No strong vertical gradient was observed in the mean daytime NO₃ reactivity with $k_{\text{day}}^{\text{VOC}}(5.5 \text{ m}) = (0.12 \pm 0.04) \text{ s}^{-1}$ for the entire campaign and $k_{\text{day}}^{\text{VOC}}(40 \text{ m}) = (0.07 \pm 0.02) \text{ s}^{-1}$ during the 5-day period.

Within the clearing, the fractional contribution of VOCs to the total NO₃ loss rate coefficient (k^{tot} , determined by photolysis,
30 reaction with NO and VOCs) was 80-90 % during the night and ~50 % during the day. In terms of chemical losses of α -pinene below canopy height in the clearing, we find that at nighttime OH and O₃ dominate with NO₃ contributing “only” 17 %, which decreases further to 8.5 % during the day. Based on measured OH, O₃ and calculated NO₃ concentrations, the chemical lifetime of BVOCs at noon is about one hour and is likely to be longer than timescales of transport out of the canopy (typically in the order of minutes), thus significantly reducing the importance of daytime, in-canopy processing. Clearly, in forested regions

35 where sufficient NO_x is available, the role of NO₃ and OH as initiators of BVOC oxidation are not strictly limited to the night and to the day, respectively, as often implied in e.g. atmospheric chemistry text-books.

1 Introduction

Forests emit great quantities (~ 1000 Tg yr⁻¹) of a variety of biogenic volatile organic compounds (BVOCs) such as isoprene and monoterpenes into the atmosphere (Guenther et al., 2012; Hakola et al., 2012; Vermeuel et al., 2023). The transport of
40 combustion-related emissions from urban and industrialised regions results in the presence of NO_x (the sum of nitric oxide, NO and nitrogen dioxide, NO₂) in forested regions, as does microbial activity in soils (Ludwig et al., 2001; Barger et al., 2005; Pilegaard, 2013). An important step in photochemical ozone (O₃) generation is the oxidation of VOCs, which may be initiated by hydroxyl (OH) and nitrate radicals (NO₃) or O₃ itself (Geyer et al., 2001; Lelieveld et al., 2008; Peräkylä et al., 2014). The interaction of largely anthropogenic NO_x with BVOCs is thus a key component of tropospheric ozone production in many
45 regions (Pusede et al., 2015). Here we focus on NO₃, which is formed from the reaction between NO₂ (e.g. from R1) and O₃ (R2) and is in a thermal equilibrium with NO₂ and dinitrogen pentoxide (N₂O₅, R3 and R4) (Wayne et al., 1991).



In the troposphere, NO₃ reacts efficiently with NO to re-form NO₂ (R5), reacts with unsaturated VOCs (R6) and is photolysed rapidly with a lifetime that is often only a few seconds (R7a and R7b) (Finlayson-Pitts and Pitts, 2000). NO is reduced in concentration at night owing to its reaction with O₃ and since both NO and sunlight drastically reduce the lifetime of NO₃, the latter is often thought of as a “nighttime only” oxidant (Wayne et al., 1991; Platt and Heintz, 1994; Martinez et al., 2000;
55 Brown and Stutz, 2012). Note that, in some environments, direct heterogeneous losses of NO₃ can also be important as can indirect losses via N₂O₅ uptake (Saathoff et al., 2001; Bertram and Thornton, 2009; Phillips et al., 2016).



While both the reaction with NO (R5) and photolysis (R7) recycle either NO or NO₂, reactions between NO₃ and VOCs result in a variety of gas-phase products including organic nitrates (RONO₂) and nitric acid (HNO₃) (Hallquist et al., 1999; Ayres et al., 2015; Ng et al., 2017) which may be lost by deposition and/or transferred to the condensed phase, forming e.g. secondary organic aerosols (SOA) (Bates et al., 2022; Day et al., 2022; DeVault et al., 2022). The interaction of NO₃ with BVOCs can
65 represent an efficient process for the removal of NO_x from the gas phase and a mechanism for SOA generation (Fry et al., 2014; Romer Present et al., 2020), making the fractional contribution of R6 to the overall loss rate of NO₃ of particular interest.

As unsaturated BVOCs such as isoprene and monoterpenes are often present at parts per billion by volume (ppbv) levels in the forest (Kesselmeier and Staudt, 1999; Hakola et al., 2009) the local NO₃ lifetimes are typically short not only during the day, but also at night (McLaren et al., 2004; Liebmann et al., 2018a; Liebmann et al., 2018b). NO₃ mixing ratios are often
70 below 1 part per trillion by volume (pptv), making its detection in highly reactive air masses very challenging (Liebmann et al., 2018a). Measuring the NO₃ reactivity (together with NO₂ and O₃ to calculate the NO₃ production rate) provides a means to assess the atmospheric fate of the nitrate radical even when its mixing ratio is too low (< 0.5 pptv) to be detected (Dewald et al., 2022).

There are also meteorological effects that induce differences in the fate of NO₃ during the day and night. While daytime
75 insolation at ground level can result in efficient (turbulent) mixing of the boundary layer, the radiative cooling of the ground at lower temperatures at night means that the nocturnal boundary layer can be highly stratified (Stull, 1988). This results in strong gradients in the mixing ratios of e.g. BVOCs (Fish et al., 1999) and the below-canopy reactivity of NO₃ can be very different to that above (Mogensen et al., 2015; Liebmann et al., 2018a). To date, NO₃ vertical profiles are available for high altitudes (i.e. above the boundary layer) in non-forested environments (Smith et al., 1993; von Friedeburg et al., 2002; Stutz
80 et al., 2004; Brown et al., 2007a; Yan et al., 2021), yet highly-resolved vertical profiles of NO₃ (or its reactivity) for low altitudes at nighttime are sparse (Brown et al., 2007b; Liebmann et al., 2018a).

In this study, we present and analyse nighttime and daytime NO₃ reactivity measurements above and below the canopy in a temperate forest ca. 50 km from Paris (France) during the summer of 2022 as part of the ACROSS campaign (Atmospheric ChemistRy Of the Suburban foreSt) (Cantrell and Michoud, 2022).

85 **2 Experimental**

2.1 Site description and meteorology

The ACROSS campaign took place from mid-June until the end of July 2022 at a clearing (ca. 700 m²) in the suburban forest of Rambouillet in France (N48.687, E1.704). Rambouillet forest covers an area of about 150 km² and is located about 50 km to the south-west of Paris. The surrounding trees are mainly oaks (~68 %) and pine (up to 25 %) (Marchant et al., 2017), with
90 an average height of ~25 m. Daytime maximum temperatures during ACROSS were ~20–40 °C (Fig.1, panel b), while nighttime temperatures decreased to 10–25 °C, often with a significant (positive) gradient in height that started to develop in the late afternoon when tree-induced shadowing of the ground led to radiative cooling (Andersen et al, 2024).

48h-back trajectories show that air originated from the Atlantic Ocean between June 25 and July 2, whereas air passed predominantly over industrialized regions including Paris, the UK, Benelux states and the Ruhr area from July 2 to July 18
95 (Andersen et al., 2024). Local wind directions and speeds are shown as a wind-rose in the Supplement (Fig. S1).

The clearing housed several instrumented containers and also a tower, which enabled measurements at 40 m to be made, either via instruments located at the top of the tower or via sampling from a gas-manifold attached to a high-flow inlet at the top of the tower.

2.2 NO₃ reactivity

100 The flow-tube cavity-ring-down spectrometer (FT-CRDS, Liebmann et al. (2017)) used to quantify NO₃ reactivity was installed in the MPIC container and sampled mainly from the centre of a high volume-flow (10000 standard L min⁻¹, SLPM) stainless steel tube ($\varnothing = 15$ cm) the top of which was at 5.5 m above ground level. We present measurements of k^{VOC} , which is defined as the VOC-induced pseudo first-order NO₃ loss rate coefficient (in units of s⁻¹) and is equal to $\Sigma k_i[\text{VOC}]_i$, with [VOC]_i and k_i being the VOC concentrations and the corresponding rate coefficient for R6, respectively. The FT-CRDS was

105 connected to the high-flow inlet with a 1.5 m long piece of ¼ inch outer diameter (OD) PFA tubing (overall residence time of 0.4 s) equipped with a Teflon membrane filter (2 µm pore, 47 mm diameter, Pall Corp.) to prevent particles entering the cavity. The filter was replaced every three days. From July 18, the NO₃ reactivity setup sampled air alternately from the high-flow inlet and from the manifold taking air from 40 m. The instrument was attached to the manifold with ca. 20 m ¼ inch PFA tubing (ca. 5 s residence time).

110 NO₃ was generated by mixing 3–5 standard cm³ min⁻¹ (sccm) of NO (1 part per million by volume (ppmv) in N₂, Air Liquide) with O₃ in a thermostated (30 °C), Teflon-coated (FEPD-121, Chemours) reactor (ca. 5 min residence time) at a pressure of 1100 Torr. O₃ was generated by the 185 nm photolysis of O₂ in a flow of 400 sccm dry synthetic air, which was provided from a commercial zero-air generator (CAP-180, Fuhr GmbH). In order to convert the resultant N₂O₅ quantitatively to NO₃ (R4), the flow was heated to 140 °C in ca. 15 cm of ¼ inch OD PFA tubing. The flow containing NO₃ was then mixed with 2.8

115 SLPM of either synthetic air (to define zero reactivity) or ambient air within a ¼ inch PFA T-piece and directed to the FEP-coated, darkened and thermostated (20°C) flow-tube where NO₃ had 11 s to react. NO₃ surviving the reactor was quantified on-line by cavity-ring-down spectroscopy at 662 nm. Zeroing (“baseline measurement”) was achieved by adding an excess of NO (3 sccm of 100 ppmv in N₂, Air Liquide) to titrate NO₃ (R5). In synthetic air, the NO₃ mixing ratios were typically in the range of 30–50 pptv. As the presence of NO₃ and N₂O₅ in ambient air would bias the measurement, at nighttime the air was

120 sampled through a 2 L glass flask (heated to 35 °C, ~ 40 s residence time) to ensure that ambient N₂O₅ is converted to NO₃. All radicals including NO₃, OH, RO₂ and HO₂ are lost on the glass walls and thus prevented from reaching the flowtube. From July 18, air was sampled through the glass flask throughout the diel cycle. No difference in k^{VOC} was observed directly after switching between “daytime mode” (no flask) and “nighttime mode” (sampling through flask), indicating that no compounds significantly contributing to k^{VOC} are lost in the glass flask. Accurate quantification of NO₃ reactivity requires that the synthetic

125 air is humidified to ambient level (monitored with a commercial sensor, IST, HYT393), which was achieved with a permeation tube (MH-070-24F-4, PermaPure LLC) filled with de-ionized water (LiChrosolv, Merck GmbH). Dynamic dilution of ambient air with synthetic air extended the dynamic range of the instrument to NO₃ reactivities of up to ca. 2.1 s⁻¹. Since reactions R1 to R5 and wall losses (0.001 s⁻¹) take place in addition to the reaction of interest (R6), a numerical simulation procedure was used to separate contributions of NO_x in the flowtube and thus extract the NO₃ reactivity towards VOCs as detailed in

130 Liebmann et al. (2017). When sampling through the glass flask, ambient NO mixing ratios used for the simulation were corrected for the impact of R1 with correction factors between 1 and 40 %.

The uncertainty in k^{VOC} is determined by the stability of the NO_3 source, the cavity stability (i.e. noise level and baseline stability) and by the numerical simulation corrections. The uncertainty induced by the simulation is dependent on the ratio between NO_2 and k^{VOC} (Liebmann et al., 2017). During the campaign source and cavity stability, numerical simulation and
135 uncertainties in the NO correction (impact of R5) contributed on average ca. 11, 9 and 26 % to the average overall uncertainty of ca. 30 %. The limit of detection (LOD) is derived from the variability (two standard deviations, 2σ) in consecutive baselines and NO_3 source measurements and was on average 0.006 s^{-1} during this campaign. In this manuscript we index the NO_3 reactivity according to the following scheme: $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$, $k_{\text{day}}^{\text{VOC}}(5.5 \text{ m})$, $k_{\text{night}}^{\text{VOC}}(40 \text{ m})$ and $k_{\text{day}}^{\text{VOC}}(40 \text{ m})$ which are the measured reactivities towards VOCs during the day/night at the two different heights. Note that reactivity measurements at 40
140 m was limited to the last 5 days of the campaign. We also refer to k^{tot} which is the total NO_3 loss term including reaction with NO and photolysis as well as reaction with VOCs.

2.3 NO, NO_2 and O_3

NO_2 mixing ratios at 5.5 m were measured by sampling via 1.5 m $\frac{1}{4}$ inch (OD) PFA tubing and a membrane filter (2 μm pore, 47 mm diameter, Pall Corp.) through the second (405 nm) cavity of the NO_3 reactivity instrument (Liebmann et al., 2018b).
145 The instrument's LOD was 87 pptv (2σ , 4 s) and the measurement was associated with a total uncertainty of 7 %. On top of the tower (40 m), NO_2 was measured via a cavity attenuated phase shift (CAPS) setup (precision of 6 %, LOD of 40 pptv) which was zeroed on an hourly basis.

NO was measured using a commercial instrument based on chemiluminescence detection (Ecophysics, CLD 780 TR, LOD of 10 pptv for 1 min averaging time) which was installed in a container ca. 17 m distance from the MPIC container and sampled
150 at a height of 3.2 m above the ground. The NO mixing ratios were corrected for a change in sensitivity during the campaign (Andersen et al., 2024). Measurement of NO at 40 m height was carried out using another CLD (Teledyne, T200UP) that sampled from the tower-manifold. This measurement was corrected for losses from R1, with corrections ranging from 1–28 %. The instrument's LOD was 30 pptv and its total uncertainty was 3.2 %.

A commercial ozone monitor (2B Technologies, model 205) based on UV absorption was installed in the MPIC container and
155 measured ozone mixing ratios at 5.5 m height with an LOD of 2 ppbv and an associated uncertainty of 5 %. O_3 from top of the tower was quantified with means of a second ozone monitor with an LOD of 2.5 ppbv (HORIBA, APOA370).

2.4 Photolysis frequencies and temperatures

Actinic flux was measured on top of the tower (41 m) as well as above the roof of the MPIC container (5 m), in both cases using spectral radiometers (Metcon GmbH). Actinic fluxes were converted to photolysis frequencies of NO_3 and of other
160 compounds using IUPAC- and NASA-evaluated absorption cross sections (Burkholder et al., 2016; IUPAC, 2024). Commercial temperature sensors (Atexis PT1000 and Thermostat PT100) monitored ambient air temperature simultaneously at 5 m, 13 m, 21 m and 41 m.

3 Results and discussion

An overview of the measurements relevant for analysis of NO₃ reactivity is given in Fig. 1 where data obtained at 5.5 m (orange) and 40 m above ground level (blue) are plotted.

The VOC-induced NO₃ reactivity at a height of 5.5 m ($k^{\text{VOC}}(5.5 \text{ m})$, panel f, orange triangles) was generally high with values between ~ 0.1 and 0.5 s^{-1} and also highly variable with nighttime peak values of up to 2 s^{-1} . In contrast, $k_{\text{night}}^{\text{VOC}}(40 \text{ m})$ was often close to or below the LOD ($\leq 0.006 \text{ s}^{-1}$) when sampling from top of the tower between July 18 – July 23 (panel f, blue circles). NO mixing ratios (panel e) at 3.2 m a.g.l. (orange circles) were on average between 0.1 and 0.3 ppbv, but occasionally peaked at 1–4 ppbv mostly in the morning during the continental phase. As detailed in Andersen et al. (2024), nighttime NO mixing ratios were close to or below the LOD when sampling air masses which, according to two-day back trajectories, were largely of continental origin (July 2 – July 18). In contrast, up to several ppbv of NO were observed at night during a period dominated by air with its origin over the Atlantic (June 26 – July 2) when O₃ levels (at 5.5 m) were below the LOD (panel c). NO₂ mixing ratios (panel d) were similar at both heights and, in the absence of anthropogenic influence, mostly between 0.5 and 2 ppbv. The large diel variation in O₃ mixing ratios at 5.5 m (40–80 ppbv during the day and as low as zero at nighttime, panel c) results from its net daytime photochemical production through reactions involving OH, NO_x and VOCs (Crutzen and Lelieveld, 2001) and its nocturnal losses via reactions (e.g. with NO and BVOC) and deposition. Nocturnal O₃ mixing ratios at 40 m (20–40 ppbv) are higher than at 5.5 m (0–20 ppbv) and its diel cycle at 40 m is weaker than at 5.5 m. This results from a combination of removal processes of O₃ at lower levels (reaction with NO released from soil, reaction with unsaturated, biogenic VOCs released from vegetation and deposition to soil and foliar surfaces) and weak vertical mixing at nighttime (Andersen et al., 2024).

3.1 k^{VOC} : Variability and controlling factors close to the ground

At a height of 5.5 m, k^{VOC} shows large variability across the diel cycle and also from night-to-night (Fig. 1, panel f), which are driven by variability in emissions of reactive BVOCs and in the height of the nocturnal boundary layer (NBL). In Fig. 2, $k^{\text{VOC}}(5.5 \text{ m})$ is plotted together with the difference in temperature (ΔT) measured at 5 and 41 m where $\Delta T = T(5 \text{ m}) - T(41 \text{ m})$ and shows that the largest values of $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ occur on nights when a temperature inversion ($\Delta T < 0 \text{ K}$) evolves. The impact of temperature inversion on NO₃ reactivity at 5.5 m is illustrated by comparing a night with no temperature inversion (July 24 – July 25, period A) to a night with moderate temperature inversion (30 June – 1 July, period B). In the absence of a temperature inversion we see a roughly constant value $< 0.1 \text{ s}^{-1}$ for $k^{\text{VOC}}(5.5 \text{ m})$ through the diel cycle (period A in the inset to Fig. 2). In contrast, when a temperature inversion developed (period B in the inset to Fig. 2), $k^{\text{VOC}}(5.5 \text{ m})$ was relatively low ($< 0.1 \text{ s}^{-1}$) until ca. 19:30 UTC. Over the next 2 hours, it gradually increased to reach peak values as large as 0.4 s^{-1} , which were associated with larger variability. Over the same period O₃ levels decreased from ~ 30 –40 ppbv to < 10 ppbv, which has been rationalised in terms of suppression of entrainment of above-canopy air (with higher levels of O₃) into air masses close

to the ground during temperature inversions (Andersen et al., 2024). The temperature inversion and associated reduction in vertical mixing impedes upward transport of both NO (emitted from the soil) and BVOCs (emitted from vegetation) so that both may accumulate within and below the canopy after sunset. The association of high values of $k^{\text{VOC}}(5.5 \text{ m})$ with low values of O_3 during nighttime temperature inversions has previously been reported for the boreal forest (Liebmann et al., 2018a). In addition, elevated NO_3 reactivity at night is also aided by the fact that the nocturnal mixing ratios of O_3 and OH are diminished due to deposition and/or lack of photochemistry, so that the lifetime and mixing ratios of monoterpenes increase (see section 3.6).

The hypothesis that temperature inversions partially drive the observed NO_3 reactivity within the canopy is reinforced by close inspection of the temperature profiles in period B in the inset to Fig. 2. At 18:00 UTC, no gradient in temperature ($16.5 \text{ }^\circ\text{C}$) between 5 and 41 m was observed. At 20:30 UTC, a positive gradient in temperature was observed at heights $> 20 \text{ m}$, becoming more distinct at 22:00 UTC. Under these conditions, vertical mixing from ground level to above-canopy-levels (ca. 20 m) is suppressed, whereas some mixing may still take place between 5 and 13 m where no temperature difference was observed. Weak (but non-zero) vertical mixing at the lower levels may be the cause of the high variability in $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ whereby instabilities in the stratification at lower levels of the NBL allow sampling (at 5.5 m) of air with different (variable) time spent at lower/higher levels and thus with highly variable reactivity. The resulting high variability in NO_3 mixing ratios has been documented for NO_3 measurements made close to the ground (Brown et al., 2003; Crowley et al., 2010).

In forests, the most abundant BVOCs are typically isoprene and monoterpenes (Hakola et al., 2012; Vermeuel et al., 2023). Since the corresponding rate coefficients for the reaction of NO_3 with monoterpenes are up to two orders of magnitude larger than that of isoprene (IUPAC, 2024), monoterpenes are expected to be the main contributor to k^{VOC} during the ACROSS campaign. Relative monoterpene emission factors are temperature-dependent and described by $\exp(\beta(T-297 \text{ K}))$ with $\beta = 0.1 \text{ K}^{-1}$ in forested environments (Guenther et al., 1993), resulting in a strong seasonal variation (Hakola et al., 2006; Vermeuel et al., 2023). As a consequence, correlations between air temperature and VOC mixing ratios and OH reactivity have been reported (Kalogridis et al., 2014; Pfannerstill et al., 2021). Figure 3a shows that, during the day (black data points), with temperatures varying from 297 to 311 K, an increase in $k^{\text{VOC}}(5.5 \text{ m})$ is observed. The inset contains daytime values measured between 10:00 and 14:00 UTC against air temperature and a linear regression suggests a fair correlation between the two (Pearson correlation coefficient $r = 0.66$). The expected factor of 4 increase in the emission rate over this range as reported in Guenther et al. (1993) is much larger than the observed change in $k^{\text{VOC}}(5.5 \text{ m})$. This is however expected as the daytime concentrations of monoterpenes (MTs) will be determined not only by emission rates but also by their lifetime, which, in the clearing, will be reduced by reactions with daytime oxidants such as the OH and O_3 and (possibly more importantly) transport out of the canopy (Bohn, 2006). At nighttime (blue and red dots in Fig. 3a), there is no clear correlation between $k^{\text{VOC}}(5.5 \text{ m})$ and temperature, although the highest values of $k^{\text{VOC}}(5.5 \text{ m})$ are generally observed at lower temperatures. A plot of $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ versus ΔT and coloured according to the temperature at 5 m (Fig. 3b) reveals that higher NO_3 reactivity is accompanied by large (negative) values of ΔT . The maximum value of $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m}) \approx 2 \text{ s}^{-1}$ was observed when a strong temperature inversion

($\Delta T < -6$ K) coincided with a high nocturnal air temperature. During cooler nights ($T(5\text{ m}) < 12$ °C), $k^{\text{VOC}}(5.5\text{ m})$ was < 0.5 s⁻¹ even during periods with very strong temperature inversions. Our analysis thus shows that while temperature is an important factor influencing NO₃ reactivity at 5.5 m through enhanced rates of emission of BVOCs, the effect (at least at sub-canopy levels) is greatly amplified by temperature inversions which favour accumulation of MTs in a shallow boundary layer close to the surface. In an upcoming publication we use monoterpene measurements to analyse NO₃ (and N₂O₅) mixing ratios and lifetimes at both 5.5 and 40 m heights during ACROSS and draw comparison between directly measured NO₃ reactivity and that attributed to various BVOC.

3.2 Nitrate radical lifetime within and above the canopy

For the last five days of the ACROSS campaign (July 18 – July 23), the NO₃ reactivity was measured at a height of 40 m every even hour, with data obtained at 5.5 m every odd hour. The corresponding time-series of this measurement with a discussion of the vertical gradients observed during this period is appended in the Supplement (S2). With the data from this period, we could derive the total (chemical) loss rate coefficient of NO₃ not only at 5.5 m but also at 40 m, which is given by

$$k^{\text{tot}} \approx k^{\text{VOC}} + J_{\text{NO}_3} + k_5[\text{NO}] \quad (1)$$

where k^{VOC} is our directly measured VOC-induced loss, $J_{\text{NO}_3} = k_{7a} + k_{7b}$ is the NO₃ photolysis frequency (Fig. 1, panel a) and k_5 is the rate coefficient (IUPAC, 2024) for the reaction between NO and NO₃. Equation (1) neglects NO₃ loss processes resulting from direct (and indirect) heterogeneous reactions of NO₃ (and N₂O₅) as well as reactions with HO_x and organic radicals, which, for the ACROSS environment, is justified in the Supplement (S3). The NO₃ lifetime τ_{NO_3} is the inverse of the total NO₃ loss rate coefficient:

$$\tau_{\text{NO}_3} = 1/k^{\text{tot}} \quad (2)$$

We plot τ_{NO_3} through the diel cycle (5.5 and 40 m) in Fig. 4. For calculating $k^{\text{tot}}(5.5\text{ m})$, the whole campaign period (as in Fig. 1) was considered since data gaps in both NO and J_{NO_3} close to the ground and on top of the tower reduced the availability of quasi-simultaneous values of $k^{\text{tot}}(5.5\text{ m})$ and $k^{\text{tot}}(40\text{ m})$ (see Supplement S4). At a height of 5.5 m, the nocturnal lifetime of NO₃ is 2–3 s and (counterintuitively) somewhat longer (4–5 s) during the day despite the photolysis of NO₃. At a height of 40 m, above the canopy, the daytime lifetime of NO₃ is close to 3–4 s, whereas at night it increases to 10–12 s. The similar lifetimes at both heights during the day imply a vertically well-mixed layer. On the other hand, the longer nocturnal lifetime at 40 m compared to 5.5 m can be attributed to decoupling from direct (ground-level and within canopy) sources of NO and BVOCs (see section 3.1). The longer nocturnal lifetime of NO₃ at 40 m enabled it to be detected on some nights of the campaign, whereas NO₃ measurements were, unlike N₂O₅, always below LOD close to the ground. A detailed analysis of the NO₃ (and N₂O₅) measurements will be presented in a future publication.

3.3 Fate of the nitrate radical within the canopy

A mean diel cycle of k^{VOC} (5.5 m) for the whole campaign is depicted in Fig. 5. The higher mean nighttime value ($k_{\text{night}}^{\text{VOC}}$ (5.5 m) = $(0.24_{-0.06}^{+0.32}) \text{ s}^{-1}$) compared to $k_{\text{day}}^{\text{VOC}}$ (5.5 m) = $(0.12 \pm 0.04) \text{ s}^{-1}$) is a result of the accumulation of BVOCs in a shallow, sub canopy layer with reduced rates of canopy-venting owing to temperature inversions. The observation of a daytime minimum and nighttime maximum in k^{VOC} (5.5 m) is consistent with measurements in the boreal forest in Finland (Liebmann et al., 2018a) where terpene emissions dominated the fate of NO_3 and strong temperature-inversions were present at night. Values of k^{VOC} in the boreal forest in autumn were a factor of 2–3 lower than those we measured in the temperate forest during ACROSS, which is presumably related to the lower temperatures as well as other factors such as leaf area index, vegetation-type and availability of oxidizing agents which affect the abundance of monoterpenes.

The fractional contribution F^{VOC} , of the reaction of NO_3 with VOCs to its total loss rate coefficient k^{tot} , is given by Eq. (3)

$$F^{\text{VOC}} = k^{\text{VOC}}/k^{\text{tot}} \quad (3)$$

and shown in Fig. 5. At night, F^{VOC} (grey shaded area, blue line) is ca. 0.7–0.8, with the remaining 20 to 30 % assigned to reaction with NO. During the day, while photolysis (20 %) and reaction with NO (30 %) gain in importance, VOCs still account for 50 % of the NO_3 reactivity. The large daytime contribution of k^{VOC} is partly related to the fact that actinic flux (and thus the photolysis frequencies of NO_3) within the clearing at 5.5 m height is reduced compared to above canopy levels (Fig. 1, panel a). In addition, the photolysis frequency for NO_2 is also reduced, so that the distribution of NO_x between NO and NO_2 is shifted towards NO_2 and away from NO, resulting in a reduction in $k_5[\text{NO}]$. In order to assess the impact of this effect, we calculated F^{VOC} with above-canopy values of J_{NO_3} and NO. In this scenario, daytime F^{VOC} increases to 33 %, i.e. the reduction of photolysis frequencies increases F^{VOC} by a factor of 1.5. Liebmann et al. (2018a) reported a daytime average for F^{VOC} of only 20 % in a boreal forest. Despite the fact that both sites are similarly affected by low NO and high monoterpene levels, this value is still significantly lower than 33 %, which can be reconciled with lower daytime values of k^{VOC} in the boreal forest. The comparatively high daytime contribution of VOCs to NO_3 consumption below the canopy thus stems from both reduction in J_{NO_3} and higher values of k^{VOC} , latter most likely due to higher concentrations of monoterpenes than in the boreal forest. This observation is consistent with even higher daytime VOC contributions to NO_3 loss of >97 % reported for sunlit wildfire plumes by Decker et al. (2021) who reconciled their result with VOC mixing ratios that were sufficiently high to outcompete photolysis and NO.

We note at this point, that the photolysis frequencies measured in the clearing will be substantially larger than in the non-cleared forest. By comparing J_{NO_3} (5.5 m) with values observed above the canopy in the early morning and afternoon when the integrating-dome of the spectral-radiometer experienced only diffuse sunlight, we observe a reduction in J_{NO_3} (5.5 m) by a factor > 10 (Supplement S5). A substantial reduction in in-forest actinic flux compared to in a clearing and above the canopy has been reported by Bohn (2006) for a temperate, deciduous forest, albeit of different tree type and density of foliage. As discussed by Bohn (2006), a reduction in NO_3 photolysis frequency implies that the relative importance of NO_3 (compared to

OH and O₃) as a daytime oxidant in the forest canopy is even larger than that (~50 %) derived above. A caveat to this is that the formation of NO₃ requires the presence of O₃ and NO₂, both of which are likely to have substantial deposition terms in dense foliage in the non-cleared forest. We also recognise that, during the daytime, the in-canopy chemical-lifetimes of BVOCs may be much longer (see section 3.6) than the average residence time with respect to canopy-outflow, so that most BVOC daytime oxidation may take place above the forest.

3.4 Fate of the nitrate radical above the canopy

In Fig. 6, we plot the diel cycle of $k^{\text{VOC}}(40 \text{ m})$ which shows a daytime maximum and nighttime minimum, which is the opposite of that measured at 5.5 m (see Fig. 5) but is typical for measurement sites that are decoupled from near-ground emissions during the night (Crowley et al., 2011; Liebmann et al., 2018b; Dewald et al., 2022). The daytime mean value of $k_{\text{day}}^{\text{VOC}}(40 \text{ m}) = (0.07 \pm 0.02) \text{ s}^{-1}$ is similar to those measured at 5.5 m, which presumably results from vertical mixing that is rapid compared to chemical lifetimes of BVOCs and NO. In contrast, the average value (\pm 25th and 75th percentile, respectively) of $k_{\text{night}}^{\text{VOC}}(40 \text{ m}) = (0.016_{-0.007}^{+0.018}) \text{ s}^{-1}$ is approximately one order of magnitude lower than $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$. Recall however, that due to its poor data coverage during the 5-day-period, $k^{\text{VOC}}(5.5 \text{ m})$ from the whole campaign is used for this comparison.

The daytime, chemical, gas-phase NO₃ loss processes at 40 m (see pie chart in Fig. 6) also differ significantly from those at 5.5 m (see Fig. 5) with the contribution of BVOCs almost halved (25.5 %) in favour of both NO (48 %) and photolysis (26.5 %). Reaction with NO is thus the main daytime loss process for NO₃ at 40 m, which is a result of greater NO mixing ratios at this height during the 5-day-period. But even at night, NO mixing ratios during this period were still high enough (40-150 pptv) to compete with VOCs. In addition, some days between July 18 and July 23 were cloudy, which is why NO₃ loss via reaction with NO prevails even at a height of 40 m. A time series comparing J_{NO_3} , $k_5[\text{NO}]$ and k^{tot} at both heights during this period is appended in the Supplement (S4).

3.5 Vertical gradient (0-24 m) in NO₃ reactivity

During the night from July 17 to July 18 (a night with a strong temperature inversion with ΔT between -4 and -6 K), gas-phase NO₃ reactivity (resulting from reaction with both BVOCs and NO), NO₂ and O₃ were measured at heights between 0 and 24 m above ground level in 4 m steps. Measurements of NO₂ and O₃ at 40 m during the same night are also plotted in Fig. 7. As NO mixing ratios were not available at all heights, the NO₃ reactivity was not corrected for this and thus includes NO₃ removal via reaction with NO ($k^{\text{NO}} = k_5[\text{NO}]$), i.e. k^{VOC} becomes $k^{\text{VOC}} + k^{\text{NO}}$.

Between 20:00 and 00:00 UTC five profiles of $k^{\text{VOC}} + k^{\text{NO}}$ were measured, each taking < 20 min. The first profile was measured after an increase in $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ which was associated with the onset of the temperature inversion. $k^{\text{VOC}} + k^{\text{NO}}$ (averaged over the 5 single profiles) is plotted versus height in Fig. 7c along with the data obtained at 40 m in the following nights (see section 3.2). The data reveal a strong trend in NO₃ reactivity with the highest values (0.34 s⁻¹) measured at ground

level, where NO is expected to have a greater impact, gradually decreasing with height to a value of ca. 0.08 s⁻¹ at 24 m. The data at 40 m (uncorrected for NO) are broadly consistent with this trend and, together with Fig. 6, imply that NO is the main contributor to NO₃ reactivity above canopy level. The gradient in NO₂ (Fig. 7b) shows no clear trend for heights below 24 m and is determined by its nighttime production (mainly the reaction between NO and O₃) which depends on the availability of NO and the availability of O₃ (which has a positive gradient). As modelled in Stutz et al. (2004), a negative nocturnal gradient in NO with height is expected, notably due to soil emission and reaction with O₃ (Andersen et al., 2024). In addition, NO₂ loss via deposition is expected to be more important at the lower levels. These processes appear to roughly counterbalance each other on this night, resulting in an almost constant mixing ratio between ground level and 24 m. This observation is consistent with those in Stutz et al. (2004), who could, if at all, only find very weak gradients. Fig. 7a also includes the vertical profiles of O₃ which increases from ca. 27 ppbv at ground-level to 58 ppbv at 40 m presumably a result of near-ground loss processes such as deposition and reaction with NO / BVOCs and lack of entrainment of O₃ from above the canopy, where higher mixing ratios were measured (Fig. 1, panel c) (Brown et al., 2007a). The measured vertical profiles of O₃ were much less distinct in Stutz et al. (2004), which contradicted their model calculations that suggested a monotonic increase within the first 100 m, broadly consistent with our case study.

3.6 Fractional contribution of NO₃ to BVOC oxidation below the canopy

Our results show that, within the canopy, the nitrate radical is lost by reactions with BVOCs not only during the night but also during the day. Here we calculate the fractional contribution of NO₃ to the oxidation of a dominant BVOC, α -pinene (Guenther et al., 2012; Liebmann et al., 2018a; Vermeuel et al., 2023). As detailed in Eq. (4) the overall loss rate coefficient for α -pinene ($k_{\alpha\text{-pinene}}$) can be calculated from the concentrations of each oxidant (OH, NO₃ and O₃) and the corresponding rate coefficients $k_{x+\alpha\text{-pin}}$ of 5.3×10^{-11} , 6.2×10^{-12} and 9.6×10^{-17} cm³ molecule⁻¹ s⁻¹ for the reaction of OH, NO₃ and O₃ with α -pinene at 298 K, respectively (IUPAC, 2024).

$$k_{\alpha\text{-pinene}} = k_{\text{OH}+\alpha\text{-pinene}}[\text{OH}] + k_{\text{O}_3+\alpha\text{-pinene}}[\text{O}_3] + k_{\text{NO}_3+\alpha\text{-pinene}}[\text{NO}_3]_{\text{ss}} \quad (4)$$

As measured NO₃ mixing ratios near ground-level were below the LOD of 2 pptv throughout the whole campaign, we have derived steady-state mixing ratios, [NO₃]_{ss}, from the ratio of production ($k_2[\text{NO}_2][\text{O}_3]$) and overall NO₃ loss rate coefficients k^{tot} (Eq. 1) (Heintz et al., 1996; Brown et al., 2003; McLaren et al., 2010; Crowley et al., 2011). Note that the OH measurements were carried out with a chemical ionization mass spectrometer (CIMS, Kukui et al. (2008), see Supplement S3) ca. 17 m apart from the MPIC container. Figure 8 (upper panel) depicts the campaign-averaged, median mixing ratios of OH, O₃ and NO₃ (steady-state) throughout the diel cycle. Nighttime, steady-state NO₃ mixing ratios were 0.1-0.2 pptv during late evening until midnight, decreasing to < 0.1 ppt between midnight and dawn. This is not only related to the reduced availability of O₃ but also to the increase in k^{VOC} , both of which were usually accompanied by temperature inversions. Remarkably, daytime NO₃ values were similar and occasionally even higher than nighttime values, emphasising the fact that photolysis was not the major daytime loss process for NO₃. As expected, OH mixing ratios were highest during the daytime (peak value of 0.14 pptv),

350 but were not zero at night (0.01–0.03 pptv). Nighttime OH is formed in the oxidation of terpenes by O₃ and in secondary reactions of RO₂ and HO₂ (formed in same process) with NO. Median O₃ mixing ratios reached ca. 44 ppbv during the day and continuously decreased to ca. 15 ppbv during the night and the early morning.

The fractional contribution of each oxidant to $k_{\alpha\text{-pinene}}$ is presented as a median diel profile for the whole campaign in the lower panel of Fig. 8. Based on the above analysis, in the summertime forested environment probed during ACROSS, NO₃ contributed *only* ca. 17 % to the nighttime oxidation of α -pinene, a value that is exceeded by both OH (ca. 24 %) and O₃ (60 %). The daytime dominance of OH and O₃ (on average ca. 50 and 41.5 %, respectively) is expected, whilst, with a contribution of 8.5 % the diel- and campaign-averaged contribution of NO₃ is still significant, which is in agreement with recent publications (Schulze et al., 2017; Liebmann et al., 2018b; Mermet et al., 2021; Dewald et al., 2022) and model calculations showing that NO₃ oxidizes BVOCs below the canopy level during daytime in both coniferous and deciduous forests (Forkel et al., 2006; 360 Fuentes et al., 2007). We further note that the oxidation of BVOCs by OH, O₃ and NO₃ results in greatly different products, so that in terms of formation of organic nitrates, NO₃-initiated oxidation can still dominate (Liebmann et al., 2019). A detailed analysis on the role of NO₃-initiated organic nitrate formation will be presented in a further publication from the ACROSS campaign. Recall that the fractional contributions are highly dependent on the ratios between the rate coefficients of the oxidants and are thus different for other monoterpenes. As shown in the Supplement (S6), this is reflected in lower fractional 365 contributions of NO₃ to the oxidation of β -pinene and limonene with values of 12.4 and 13.5 % at night and 4.4 and 6.2 % during the day.

Our analysis is not consistent with the generally accepted text-book paradigm that OH-initiated oxidation processes are predominant during the day and NO₃-initiated oxidation prevails at night. However, an important caveat to our analysis is the neglect of daytime transport (venting) of BVOCs out of the forest. If this proceeds at time-scales that are short compared to 370 the chemical lifetime of BVOCs, then in-canopy oxidation will be of reduced importance compared to oxidation once BVOCs have been transported to higher levels within the boundary layer and free troposphere (Bohn, 2006).

4 Conclusions and summary

During a field campaign in a peri-urban, temperate (oak and pine) forest in France during a summer period in 2022, the reactivity of the NO₃ radical towards VOCs was measured both within the canopy (height of 5.5 m), above the canopy (height 375 of 40 m) and on one night at several heights between 0 and 24 m. NO₃ lifetimes were generally short (1–3 s) which were driven mainly by the abundance of BVOCs. Diel cycles of NO₃ at 5.5 m and 40 m were distinct, with the highest reactivity at 40 m occurring during the day and the highest reactivity at 5.5 m measured at nighttime. The highest nighttime reactivities were associated with high temperatures (driving the BVOC emissions) and with strong nighttime temperature inversions (preventing mixing of BVOCs and NO out of the nocturnal surface layer). At 5.5 m, BVOCs represented the dominant loss term for NO₃ 380 both during the night (70–80 %) and during the day (~50 %), which is partially a result of reduced NO₃ (and NO₂) photolysis frequencies at sub-canopy heights. NO₃ reactivity decreased rapidly with height above the ground with nocturnal lifetimes

(with respect to reaction with VOCs) of > 100 s at 40 m and as low as 2.5 s at ground-level. This gradient is driven largely by BVOC and NO emissions into a shallow, stratified near-surface layer under canopy height.

385 The conventional wisdom, that OH is a daytime oxidant and NO₃ a nighttime oxidant appears not to apply to forested regions with significant BVOC emissions where both NO₃ and OH have important roles throughout the diel cycle. Hence, NO₃-initiated organic nitrate formation could become significant during the day, whereas OH-initiated nocturnal chemistry would be enhanced in such environments.

Data Availability. All data can be found on <https://across.aeris-data.fr/catalogue/>.

390

Author contributions. PD analysed the data and wrote the original draft of the manuscript and, together with JNC, revised it. CC and VM were responsible for the campaign organization with contributions from individual group leads. All authors provided measurements and commented on the manuscript.

395 *Competing Interests.* At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements.

JC acknowledges Chemours for the provision of a FEPD-121 sample used to coat the flowtube and the Deutsche Forschungsgemeinschaft (project “MONOTONS”, project number: 522970430). STA thanks the Alexander von Humboldt foundation for funding her stay at the MPIC. The ACROSS project has received funding from the French National Research Agency (ANR) under the investment program integrated into France 2030, with the reference ANR-17-MPGA-0002, and it was supported by the French National program LEFE (Les Enveloppes Fluides et l'Environnement) of the CNRS/INSU (Centre National de la Recherche Scientifique/Institut National des Sciences de l'Univers). IMT Nord Europe acknowledges financial support from the CaPPA project, which is funded by the French National Research Agency (ANR) through the PIA
405 (Programme d'Investissement d'Avenir) under contract ANR-11-LABX-0005-01, the Regional Council “Hauts-de-France” and the European Regional Development Fund (ERDF). Data from the ACROSS campaign are hosted by the French national centre for Atmospheric data and services AERIS. AK acknowledges financial support from the VOLTAIRE project (ANR-10-LABX-100-01) funded by ANR through the PIA (Programme d'Investissement d'Avenir). The scientific colour map “Berlin” (Crameri, 2023) is used in this study to prevent visual distortion of the data and exclusion of readers with colour-
410 deficiencies (Crameri et al., 2020).

References

Andersen, S. T., McGillen, M. R., Xue, C., Seubert, T., Dewald, P., Türk, G. N. T. E., Schuladen, J., Denjean, C., Etienne, J. C., Garrouste, O., Jamar, M., Harb, S., Cirtog, M., Michoud, V., Cazaunau, M., Bergé, A., Cantrell, C., Dusanter, S., Picquet-Varrault, B., Kukui, A., Mellouki, A., Carpenter, L. J., Lelieveld, J., and Crowley, J. N.: Measurement report: Sources, sinks

- 415 and lifetime of NO_x in a sub-urban temperate forest at night, *EGUsphere*, 2024, 1-29, doi:10.5194/egusphere-2023-2848, 2024.
- Ayres, B. R., Allen, H. M., Draper, D. C., Brown, S. S., Wild, R. J., Jimenez, J. L., Day, D. A., Campuzano-Jost, P., Hu, W., de Gouw, J., Koss, A., Cohen, R. C., Duffey, K. C., Romer, P., Baumann, K., Edgerton, E., Takahama, S., Thornton, J. A., Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Wennberg, P. O., Nguyen, T. B., Teng, A., Goldstein, A. H., Olson, K., and Fry, J. L.: Organic nitrate aerosol formation via NO₃ + biogenic volatile organic compounds in the southeastern United States, *Atmos. Chem. Phys.*, 15, 13377-13392, doi:10.5194/acp-15-13377-2015, 2015.
- 420 Barger, N. N., Belnap, J., Ojima, D. S., and Mosier, A.: NO gas loss from biologically crusted soils in Canyonlands National Park, Utah, *Biogeochemistry*, 75, 373-391, doi:10.1007/s10533-005-1378-9, 2005.
- Bates, K. H., Burke, G. J. P., Cope, J. D., and Nguyen, T. B.: Secondary organic aerosol and organic nitrogen yields from the nitrate radical (NO₃) oxidation of alpha-pinene from various RO₂ fates, *Atmos. Chem. Phys.*, 22, 1467-1482, doi:10.5194/acp-22-1467-2022, 2022.
- 425 Bertram, T. H., and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351-8363, doi:10.5194/acp-9-8351-2009, 2009.
- Bohn, B.: Solar spectral actinic flux and photolysis frequency measurements in a deciduous forest, *J. Geophys. Res.-Atmos.*, 430 111, D15303, doi:10.1029/2005JD006902, 2006.
- Brown, S. S., Stark, H., Ryerson, T. B., Williams, E. J., Nicks, D. K., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Nitrogen oxides in the nocturnal boundary layer: Simultaneous in situ measurements of NO₃, N₂O₅, NO₂, NO, and O₃, *J. Geophys. Res.-Atmos.*, 108, 4299, doi:10.1029/2002JD002917, 2003.
- Brown, S. S., Dube, W. P., Osthoff, H. D., Stutz, J., Ryerson, T. B., Wollny, A. G., Brock, C. A., Warneke, C., De Gouw, J., 435 A., Atlas, E., Neuman, J. A., Holloway, J. S., Lerner, B. M., Williams, E. J., Kuster, W. C., Goldan, P. D., Angevine, W. M., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Vertical profiles in NO₃ and N₂O₅ measured from an aircraft: Results from the NOAA P-3 and surface platforms during the New England Air Quality Study 2004, *J. Geophys. Res.-Atmos.*, 112, D22304, doi: 10.1029/2007jd008893, 2007a.
- Brown, S. S., Dube, W. P., Osthoff, H. D., Wolfe, D. E., Angevine, W. M., and Ravishankara, A. R.: High resolution vertical 440 distributions of NO₃ and N₂O₅ through the nocturnal boundary layer, *Atmos. Chem. Phys.*, 7, 139-149, doi:10.5194/acp-7-139-2007, 2007b.
- Brown, S. S., and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405-6447, doi:10.1039/C2CS35181A, 2012.
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D., 445 M., and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18," JPL Publication 15-10, Jet Propulsion Laboratory, Pasadena: <https://jpldataeval.jpl.nasa.gov>, access: 23 July 2023, 2016.
- Cantrell, C., and Michoud, V.: An Experiment to Study Atmospheric Oxidation Chemistry and Physics of Mixed Anthropogenic-Biogenic Air Masses in the Greater Paris Area, *Bulletin of the American Meteorological Society*, 103, 599-603, doi:10.1175/BAMS-D-21-0115.1, 2022.
- 450 Crameri, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nat. Commun.*, 11, 5444, doi:10.1038/s41467-020-19160-7, 2020.

- Crameri, F.: Scientific colour maps (8.0.1), Zenodo, doi:10.5281/zenodo.8409685, 2023.
- Crowley, J. N., Schuster, G., Pouvesle, N., Parchatka, U., Fischer, H., Bonn, B., Bingemer, H., and Lelieveld, J.: Nocturnal nitrogen oxides at a rural mountain site in south-western Germany, *Atmos. Chem. Phys.*, 10, 2795-2812, doi:10.5194/acp-10-2795-2010, 2010.
- Crowley, J. N., Thieser, J., Tang, M. J., Schuster, G., Bozem, H., Hasaynali Beygi, Z., Fischer, H., Diesch, J.-M., Drewnick, F., Borrmann, S., Song, W., Yassaa, N., Williams, J., Pöhler, D., Platt, U., and Lelieveld, J.: Variable lifetimes and loss mechanisms for NO₃ and N₂O₅ during the DOMINO campaign: Contrast between marine, urban and continental air, *Atmos. Chem. Phys.*, 11, 10863-10870, doi:10.5194/acp-11-10853-2011, 2011.
- 460 Crutzen, P. J., and Lelieveld, J.: Human impacts on atmospheric chemistry, *Annu. Rev. Earth Planet. Sci.*, 29, 17-45, doi:10.1146/annurev.earth.29.1.17, 2001.
- Day, D. A., Fry, J. L., Kang, H. G., Krechmer, J. E., Ayres, B. R., Keehan, N. I., Thompson, S. L., Hu, W. W., Campuzano-Jost, P., Schroder, J. C., Stark, H., DeVault, M. P., Ziemann, P. J., Zarzana, K. J., Wild, R. J., Dube, W. P., Brown, S. S., and Jimenez, J. L.: Secondary Organic Aerosol Mass Yields from NO₃ Oxidation of alpha-Pinene and Delta-Carene: Effect of RO₂ Radical Fate, *J. Phys. Chem. A*, doi:10.1021/acs.jpca.2c04419, 2022.
- 465 Decker, Z. C. J., Robinson, M. A., Barsanti, K. C., Bourgeois, I., Coggon, M. M., DiGangi, J. P., Diskin, G. S., Flocke, F. M., Franchin, A., Fredrickson, C. D., Gkatzelis, G. I., Hall, S. R., Halliday, H., Holmes, C. D., Huey, L. G., Lee, Y. R., Lindaas, J., Middlebrook, A. M., Montzka, D. D., Moore, R., Neuman, J. A., Nowak, J. B., Palm, B. B., Peischl, J., Piel, F., Rickly, P. S., Rollins, A. W., Ryerson, T. B., Schwantes, R. H., Sekimoto, K., Thornhill, L., Thornton, J. A., Tyndall, G. S., Ullmann, K., Van Rooy, P., Veres, P. R., Warneke, C., Washenfelder, R. A., Weinheimer, A. J., Wiggins, E., Winstead, E., Wisthaler, A., Womack, C., and Brown, S. S.: Nighttime and daytime dark oxidation chemistry in wildfire plumes: an observation and model analysis of FIREX-AQ aircraft data, *Atmos. Chem. Phys.*, 21, 16293-16317, doi:10.5194/acp-21-16293-2021, 2021.
- DeVault, M. P., Ziola, A. C., and Ziemann, P. J.: Products and Mechanisms of Secondary Organic Aerosol Formation from the NO₃ Radical-Initiated Oxidation of Cyclic and Acyclic Monoterpenes, *ACS Earth Space Chem.*, 6, 2076-2092, doi:10.1021/acsearthspacechem.2c00130, 2022.
- 475 Dewald, P., Nussbaumer, C. M., Schuladen, J., Ringsdorf, A., Edtbauer, A., Fischer, H., Williams, J., Lelieveld, J., and Crowley, J. N.: Fate of the nitrate radical at the summit of a semi-rural mountain site in Germany assessed with direct reactivity measurements, *Atmos. Chem. Phys.*, 22, 7051-7069, doi:10.5194/acp-22-7051-2022, 2022.
- Finlayson-Pitts, B. J., and Pitts, J. N.: CHAPTER 7 - Chemistry of Inorganic Nitrogen Compounds: Chemistry of the Upper and Lower Atmosphere, edited by: Finlayson-Pitts, B. J., and Pitts, J. N., Academic Press, San Diego, 264-293, 2000.
- 480 Fish, D. J., Shallcross, D. E., and Jones, R. L.: The vertical distribution of NO₃ in the atmospheric boundary layer, *Atmos. Environ.*, 33, 687-691, doi:10.1016/S1352-2310(98)00332-X, 1999.
- Forkel, R., Klemm, O., Graus, M., Rappenglück, B., Stockwell, W. R., Grabmer, W., Held, A., Hansel, A., and Steinbrecher, R.: Trace gas exchange and gas phase chemistry in a Norway spruce forest: A study with a coupled 1-dimensional canopy atmospheric chemistry emission model, *Atmos. Environ.*, 40, S28-S42, doi:10.1016/j.atmosenv.2005.11.070, 2006.
- 485 Fry, J. L., Draper, D. C., Barsanti, K. C., Smith, J. N., Ortega, J., Winkle, P. M., Lawler, M. J., Brown, S. S., Edwards, P. M., Cohen, R. C., and Lee, L.: Secondary Organic Aerosol Formation and Organic Nitrate Yield from NO₃ Oxidation of Biogenic Hydrocarbons, *Environ. Sci. Technol.*, 48, 11944-11953, doi:10.1021/es502204x, 2014.

- 490 Fuentes, J. D., Wang, D., Bowling, D. R., Potosnak, M., Monson, R. K., Goliff, W. S., and Stockwell, W. R.: Biogenic hydrocarbon chemistry within and above a mixed deciduous forest, *J. Atmos. Chem.*, 56, 165-185, doi:10.1175/1520-0477(2000)081<1537:BHITAB>2.3.CO;2, 2007.
- Geyer, A., Alicke, B., Konrad, S., Schmitz, T., Stutz, J., and Platt, U.: Chemistry and oxidation capacity of the nitrate radical in the continental boundary layer near Berlin, *J. Geophys. Res.-Atmos.*, 106, 8013-8025, doi:10.1029/2000jd900681, 2001.
- 495 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and Monoterpene Emission Rate Variability - Model Evaluations and Sensitivity Analyses, *J. Geophys. Res.-Atmos.*, 98, 12609-12617, doi:10.1029/93jd00527, 1993.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model. Dev.*, 5, 1471-1492, doi:10.5194/gmd-5-1471-2012, 2012.
- 500 Hakola, H., Tarvainen, V., Back, J., Ranta, H., Bonn, B., Rinne, J., and Kulmala, M.: Seasonal variation of mono- and sesquiterpene emission rates of Scots pine, *Biogeosciences*, 3, 93-101, doi:10.5194/bg-3-93-2006, 2006.
- Hakola, H., Hellén, H., Tarvainen, V., Bäck, J., Patokoski, J., and Rinne, J.: Annual variations of atmospheric VOC concentrations in a boreal forest, *Boreal Env. Res.*, 14, 722-730, 2009.
- 505 Hakola, H., Hellén, H., Hemmilä, M., Rinne, J., and Kulmala, M.: In situ measurements of volatile organic compounds in a boreal forest, *Atmos. Chem. Phys.*, 12, 11665-11678, doi:10.5194/acp-12-11665-2012, 2012.
- Hallquist, M., Wangberg, I., Ljungstrom, E., Barnes, I., and Becker, K. H.: Aerosol and product yields from NO₃ radical-initiated oxidation of selected monoterpenes, *Environ. Sci. Technol.*, 33, 553-559, doi:10.1021/es980292s, 1999.
- Heintz, F., Platt, U., Flentje, H., and Dubois, R.: Long-term observation of nitrate radicals at the tor station, Kap Arkona (Rügen), *J. Geophys. Res.-Atmos.*, 101, 22891-22910, doi:10.1029/96JD01549, 1996.
- 510 IUPAC: Task Group on Atmospheric Chemical Kinetic Data Evaluation, edited by: Ammann, M., Cox, R.A., Crowley, J.N., Herrmann, H., Jenkin, M.E., McNeill, V.F., Mellouki, A., Rossi, M. J., Troe, J. and Wallington, T. J.: <https://iupac.aeris-data.fr/en/home-english/>, access: 29 January, 2024.
- 515 Kalogridis, C., Gros, V., Sarda-Esteve, R., Langford, B., Loubet, B., Bonsang, B., Bonnaire, N., Nemitz, E., Genard, A. C., Boissard, C., Fernandez, C., Ormeño, E., Baisnée, D., Reiter, I., and Lathièrre, J.: Concentrations and fluxes of isoprene and oxygenated VOCs at a French Mediterranean oak forest, *Atmos. Chem. Phys.*, 14, 10085-10102, doi:10.5194/acp-14-10085-2014, 2014.
- Kesselmeier, J., and Staudt, M.: Biogenic volatile organic compounds (VOC): An overview on emission, physiology and ecology, *J. Atmos. Chem.*, 33, 23-88, doi:10.1023/a:1006127516791, 1999.
- 520 Kukui, A., Ancellet, G., and Le Bras, G.: Chemical ionisation mass spectrometer for measurements of OH and Peroxy radical concentrations in moderately polluted atmospheres, *J. Atmos. Chem.*, 61, 133-154, doi:10.1007/s10874-009-9130-9, 2008.
- Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737-740, doi:10.1038/nature06870, 2008.

- 525 Liebmann, J., Karu, E., Sobanski, N., Schuladen, J., Ehn, M., Schallhart, S., Quéléver, L., Hellen, H., Hakola, H., Hoffmann, T., Williams, J., Fischer, H., Lelieveld, J., and Crowley, J. N.: Direct measurement of NO₃ radical reactivity in a boreal forest, *Atmos. Chem. Phys.*, 18, 3799-3815, doi:10.5194/acp-18-3799-2018, 2018a.
- 530 Liebmann, J., Sobanski, N., Schuladen, J., Karu, E., Hellen, H., Hakola, H., Zha, Q., Ehn, M., Riva, M., Heikkinen, L., Williams, J., Fischer, H., Lelieveld, J., and Crowley, J. N.: Alkyl nitrates in the boreal forest: formation via the NO₃-, OH- and O₃-induced oxidation of biogenic volatile organic compounds and ambient lifetimes, *Atmos. Chem. Phys.*, 19, 10391-10403, doi:10.5194/acp-19-10391-2019, 2019.
- Liebmann, J. M., Schuster, G., Schuladen, J. B., Sobanski, N., Lelieveld, J., and Crowley, J. N.: Measurement of ambient NO₃ reactivity: Design, characterization and first deployment of a new instrument, *Atmos. Meas. Tech.*, 10, 1241-1258, doi:10.5194/amt-2016-381, 2017.
- 535 Liebmann, J. M., Muller, J. B. A., Kubistin, D., Claude, A., Holla, R., Plaß-Dülmer, C., Lelieveld, J., and Crowley, J. N.: Direct measurements of NO₃-reactivity in and above the boundary layer of a mountain-top site: Identification of reactive trace gases and comparison with OH-reactivity, *Atmos. Chem. Phys.*, 18, 12045-12059, doi:10.5194/acp-18-12045-2018, 2018b.
- Ludwig, J., Meixner, F. X., Vogel, B., and Förstner, J.: Soil-air exchange of nitric oxide: An overview of processes, environmental factors and modeling studies, *Biogeochemistry*, 52, 225-257, 2001.
- 540 Marchant, A., Le Coupanec, A., Joly, C., Perthame, E., Sertour, N., Garnier, M., Godard, V., Ferquel, E., and Choumet, V.: Infection of *Ixodes ricinus* by *Borrelia sensu lato* in peri-urban forests of France, *PloS One*, 12, e0183543, doi:10.1371/journal.pone.0183543, 2017.
- Martinez, M., Perner, D., Hackenthal, E. M., Kulzer, S., and Schutz, L.: NO₃ at Helgoland during the NORDEX campaign in October 1996, *J. Geophys. Res.-Atmos.*, 105, 22685-22695, doi:10.1029/2000JD900255, 2000.
- 545 McLaren, R., Salmon, R. A., Liggio, J., Hayden, K. L., Anlauf, K. G., and Leaitch, W. R.: Nighttime chemistry at a rural site in the Lower Fraser Valley, *Atmos. Environ.*, 38, 5837-5848, doi:10.1016/j.atmosenv.2004.03.074, 2004.
- McLaren, R., Wojtal, P., Majonis, D., McCourt, J., Halla, J. D., and Brook, J.: NO₃ radical measurements in a polluted marine environment: links to ozone formation, *Atmos. Chem. Phys.*, 10, 4187-4206, doi:10.5194/acp-10-4187-2010, 2010.
- 550 Mermet, K., Perraudin, E., Dusanter, S., Sauvage, S., Léonardis, T., Flaud, P.-M., Bsaibes, S., Kammer, J., Michoud, V., Gratién, A., Cirtog, M., Al Ajami, M., Truong, F., Batut, S., Hecquet, C., Doussin, J.-F., Schoemaeker, C., Gros, V., Locoge, N., and Villenave, E.: Atmospheric reactivity of biogenic volatile organic compounds in a maritime pine forest during the LANDEX episode 1 field campaign, *Sci. Total Environ.*, 756, 144129, doi:10.1016/j.scitotenv.2020.144129, 2021.
- Mogensen, D., Gierens, R., Crowley, J. N., Keronen, P., Smolander, S., Sogachev, A., Nölscher, A. C., Zhou, L., Kulmala, M., Tang, M. J., Williams, J., and Boy, M.: Simulations of atmospheric OH, O₃ and NO₃ reactivities within and above the boreal forest, *Atmos. Chem. Phys.*, 15, 3909-3932, doi:10.5194/acp-15-3909-2015, 2015.
- 555 Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Inuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrault, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos. Chem. Phys.*, 17, 2103-2162, doi:10.5194/acp-17-2103-2017, 2017.
- 560

- Peräkylä, O., Vogt, M., Tikkanen, O. P., Laurila, T., Kajos, M. K., Rantala, P. A., Patokoski, J., Aalto, J., Yli-Juuti, T., Ehn, M., Sipilä, M., Paasonen, P., Rissanen, M., Nieminen, T., Taipale, R., Keronen, P., Lappalainen, H. K., Ruuskanen, T. M., Rinne, J., Kerminen, V. M., Kulmala, M., Back, J., and Petaja, T.: Monoterpenes' oxidation capacity and rate over a boreal forest: temporal variation and connection to growth of newly formed particles, *Boreal Environ. Res.*, 19, 293-310, 2014.
- 565 Pfannerstill, E. Y., Reijrink, N. G., Edtbauer, A., Ringsdorf, A., Zannoni, N., Araújo, A., Ditas, F., Holanda, B. A., Sá, M. O., Tsokankunku, A., Walter, D., Wolff, S., Lavrič, J. V., Pöhlker, C., Sörgel, M., and Williams, J.: Total OH reactivity over the Amazon rainforest: variability with temperature, wind, rain, altitude, time of day, season, and an overall budget closure, *Atmos. Chem. Phys.*, 21, 6231-6256, doi:10.5194/acp-21-6231-2021, 2021.
- 570 Phillips, G. J., Thieser, J., Tang, M. J., Sobanski, N., Schuster, G., Fachinger, J., Drewnick, F., Borrmann, S., Bingemer, H., Lelieveld, J., and Crowley, J. N.: Estimating N₂O₅ uptake coefficients using ambient measurements of NO₃, N₂O₅, ClNO₂ and particle-phase nitrate, *Atmos. Chem. Phys.*, 16, 13231-13249, doi:10.5194/acp-16-13231-2016, 2016.
- Pilegaard, K.: Processes regulating nitric oxide emissions from soils, *Philos. Trans. R. Soc. London, Ser. B*, 368, 20130126, doi:10.1098/rstb.2013.0126, 2013.
- 575 Platt, U., and Heintz, F.: Nitrate radicals in tropospheric chemistry, *Isr. J. Chem.*, 34, 289-300, doi:10.1002/ijch.199400033, 1994.
- Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and Recent Trends in the Chemistry of Continental Surface Ozone, *Chem. Rev.*, 115, 3898-3918, doi:10.1021/cr5006815, 2015.
- Romer Present, P. S., Zare, A., and Cohen, R. C.: The changing role of organic nitrates in the removal and transport of NO_x, *Atmos. Chem. Phys.*, 20, 267-279, doi:10.5194/acp-20-267-2020, 2020.
- 580 Saathoff, H., Naumann, K. H., Riemer, N., Kamm, S., Mohler, O., Schurath, U., Vogel, H., and Vogel, B.: The loss of NO₂, HNO₃, NO₃/N₂O₅, and HO₂/HOONO₂ on soot aerosol: A chamber and modeling study, *Geophys. Res. Lett.*, 28, 1957-1960, doi:10.1029/2000GL012619, 2001.
- 585 Schulze, B. C., Wallace, H. W., Flynn, J. H., Lefer, B. L., Erickson, M. H., Jobson, B. T., Dusanter, S., Griffith, S. M., Hansen, R. F., Stevens, P. S., VanReken, T., and Griffin, R. J.: Differences in BVOC oxidation and SOA formation above and below the forest canopy, *Atmos. Chem. Phys.*, 17, 1805-1828, doi:10.5194/acp-17-1805-2017, 2017.
- Smith, J. P., Solomon, S., Sanders, R. W., Miller, H. L., Perliski, L. M., Keys, J. G., and Schmeltekopf, A. L.: Atmospheric NO₃: 4. Vertical Profiles at Middle and Polar Latitudes at Sunrise, *J. Geophys. Res.-Atmos.*, 98, 8983-8989, doi:10.1029/93JD00041, 1993.
- 590 Stull, R. B.: Stable Boundary Layer: An Introduction to Boundary Layer Meteorology, edited by: Stull, R. B., Springer Netherlands, Dordrecht, 499-543, 1988.
- Stutz, J., Alicke, B., Ackermann, R., Geyer, A., White, A., and Williams, E.: Vertical profiles of NO₃, N₂O₅, O₃, and NO_x in the nocturnal boundary layer: 1. Observations during the Texas Air Quality Study 2000 J. *Geophys. Res.-Atmos.*, 109, D12306, doi:10.1029/2003JD004209, 2004.
- 595 Vermeuel, M. P., Novak, G. A., Kilgour, D. B., Claffin, M. S., Lerner, B. M., Trowbridge, A. M., Thom, J., Cleary, P. A., Desai, A. R., and Bertram, T. H.: Observations of biogenic volatile organic compounds over a mixed temperate forest during the summer to autumn transition, *Atmos. Chem. Phys.*, 23, 4123-4148, doi:10.5194/acp-23-4123-2023, 2023.


von Friedeburg, C., Wagner, T., Geyer, A., Kaiser, N., Vogel, B., Vogel, H., and Platt, U.: Derivation of tropospheric NO₃ profiles using off-axis differential optical absorption spectroscopy measurements during sunrise and comparison with simulations, *J. Geophys. Res.-Atmos.*, 107, 4168, doi:10.1029/2001JD000481, 2002.

600 Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, *Atmos. Env. A*, 25A, 1-206, doi:10.1016/0960-1686(91)90192-A, 1991.


Yan, Y. H., Wang, S. S., Zhu, J., Guo, Y. L., Tang, G. Q., Liu, B. X., An, X. X., Wang, Y. S., and Zhou, B.: Vertically increased NO₃ radical in the nocturnal boundary layer, *Sci. Total Environ.*, 763, 142969, doi:10.1016/j.scitotenv.2020.142969, 2021.

605

Figures



610 Figure 1: Time series of NO_3 photolysis frequencies (J_{NO_3} , panel a), temperature (T, panel b), ozone (O_3 , panel c), nitrogen dioxide (NO_2 , panel d), nitric oxide (NO, panel e) and VOC-induced NO_3 reactivity (k^{VOC} , panel f) sampled at 5.5 m (orange triangles, 3.2 m and 5 m for NO and T, respectively) and 40 m (blue circles) above ground level. Major and minor ticks on the x-axis represent 00:00 UTC of the corresponding date. Dashed lines separate periods with air of Atlantic and continental origin. Nighttime periods are grey-shaded.



615 **Figure 2: Time series of k^{VOC} (second panel, logarithmic scale) near ground level (5.5 m) coloured by O_3 mixing ratios measured at the same height (colour scale on the right). Nighttime periods are grey-shaded. The difference between temperatures measured at 5 m and 41 m (ΔT) is plotted in the first panel. Temperature inversions are marked by dashed, vertical lines. Periods A (lower left panel) and B (lower right panel) exemplify daytime-nighttime transitions both with (right) and without (left) clear temperature inversions, with NO_3 reactivity plotted along with temperature profiles at selected times. A vertical profile of k^{VOC} was measured in period P (see section 3.5).**

620




Figure 3: (a) NO₃ reactivity at 5.5 m ($k^{\text{VOC}}(5.5 \text{ m})$) plotted against the expected change in the monoterpane emission factor (E_{MT}) relative to 297 K using $E_{\text{MT}} \propto \exp(0.1\text{K}^{-1}(T - 297\text{K}))$ (Guenther et al., 1993). Data points are coloured according to time of the day (UTC) with night/morning (blue), daytime (black) and afternoon/night (red). The inset shows $k^{\text{VOC}}(5.5 \text{ m})$ between 10:00 and 14:00 UTC plotted against air temperature (5 m). Linear regression (blue solid line) yields a Pearson correlation coefficient (r) of 0.66. (b) Nocturnal ground-level NO₃ reactivity $k_{\text{night}}^{\text{VOC}}(5.5 \text{ m})$ plotted versus the temperature difference (ΔT) between 41 m and 5 m. The data points are coloured according to the ground-level temperature $T(5 \text{ m})$.

625

630







Figure 4: Mean diel cycle of overall NO₃ lifetime (τ_{NO_3}) calculated from Eq. (2) at 5.5 m (dark orange triangles with solid line, June 17 – July 23) and 40 m (blue circles with dashed line, every 2h, July 18 – July 23). Shaded areas represent the standard deviation (1σ). The nighttime period (19:30 to 03:30 UTC) is shaded grey.



640 **Figure 5: Mean diel profiles (10 min averages, June 17 – July 23) of VOC-induced NO_3 reactivity (k^{VOC} , right y-axis) at 5.5 m along with its fractional contribution (F^{VOC} , left y-axis) to the overall NO_3 loss. Shaded areas represent the standard deviation (1σ). The nighttime period is grey-shaded and separated by black solid lines. The pie chart shows the fractional contribution of each process in Eq. (1) to the overall NO_3 loss term during the day.**



645 Figure 6: Mean diel profiles (1h averages, every two hours, July 18 – July 23) of VOC-induced NO_3 reactivity (k^{VOC} , orange triangles) at 40 m along with its fractional contribution (F^{VOC} , blue circles) to the overall NO_3 loss. Shaded areas represent the standard deviation (1σ). The nighttime period is grey-shaded. The pie chart shows the fractional contribution of each process in Eq. (1) to the overall NO_3 loss term during the day.



655 **Figure 7: Box-and-whisker plots (full range, 25th, 50th and 75th percentiles) of vertical profiles of (a) O₃ (b) NO₂ and (c) total gas-phase NO₃ reactivity ($k^{\text{VOC}} + k^{\text{NO}}$) measured during the temperature-inverted night of Jul 17 to Jul 18 between 20:00 and 00:00 UTC. NO₃ reactivities measured on top of the tower (40 m) during the nights between July 18 to July 23 are also shown.**




Figure 8: Campaign-averaged median diel profiles (1h) of the oxidants (upper panel) $[\text{NO}_3]_{\text{ss}}$ (violet circles), OH (blue triangles) and O_3 (dark yellow squares) and their contribution to the overall loss rate coefficient of α -pinene according to Eq. (4) close to the ground (lower panel). The nighttime period is grey-shaded. Shaded areas represent the standard deviation (1σ).