

Peering into the heart of thunderstorm clouds: Insights from cloud radar and spectral polarimetry

Ho Yi Lydia Mak and Christine Unal

Department of Geoscience and Remote Sensing, Delft University of Technology, Delft, The Netherlands

Correspondence: Ho Yi Lydia Mak (hylmak@connect.ust.hk), Christine Unal (c.m.h.unal@tudelft.nl)

Abstract. Lightning is a natural phenomenon that can be dangerous to humans. It is ~~however~~ therefore challenging to study thunderstorm clouds using direct observations since it can be dangerous to fly into thunderstorm clouds. In this study, a cloud radar at 35 GHz with 45° elevation is used to study the properties and dynamics of thunderstorm clouds. It is based on a ~~case of thunderstorm~~ thunderstorm case on 18 June 2021 from 16:10 to 17:45 UTC near Cabauw, the Netherlands. The ~~technique of spectral polarimetric analysis at millimeter wavelength, which has not been used in previous studies about thunderstorm clouds so far, is used to help understand the behaviours of different types of particles~~ time and location of individual lightning strikes are determined using the BLIDS system, operated by SIEMENS, which is based on the Time-Of-Arrival principle. Concurrently, spectral polarimetry in the millimeter band—an innovative technique not previously applied in thunderstorm cloud studies—is employed to elucidate the behavior of various particle types within a radar resolution volume. Spectral polarimetric radar variables are ~~used to investigate possible hydrometeors in the clouds and also used to~~ look for vertical alignment of ice crystals that is expected due to electric torque. Due to challenges posed by ~~Mie~~ non-Rayleigh scattering, scattering simulations are carried out to aid the interpretation of spectral polarimetric variables. It is shown that the start of the Mie regime in the Doppler spectrum can be clearly identified by the use of the spectral differential phase. From the results, there is a high chance that supercooled liquid water and conical graupel are present in the investigated thunderstorm clouds. There is also a possibility of ice crystals arranged in chains at the cloud top. Ice crystals become vertically aligned a few seconds before lightning and return to their usual horizontal alignment afterwards. However, this phenomenon has been witnessed in only a few cases, specifically when the lightning strike is in close proximity to the radar's line of sight or when the lightning is ~~exceptionally~~ strong. Doppler analyses show that updrafts are found near the core of the thunderstorm cloud, while downdrafts are observed at the edges. Strong turbulence is also observed as ~~reflected~~ shown by the large Doppler spectrum width.

20 1 Introduction

Lightning is ~~a natural phenomena that is dangerous to humans. It is~~ the electric discharge caused by an electrical breakdown of charges built up in a cloud. Scientists began investigating atmospheric electrification and lightning several hundred years ago. Many studies have shown that the charge distribution in most thunderclouds follow a tripole structure, with positive charges in the upper and lower levels and negative charges in the middle level (Wang, 2013). The positive charge center near the cloud base is relatively small, thus is sometimes ignored. Typically, a breakdown can occur when the environmental electric

field established by the charges is around $100\text{-}300\text{ V m}^{-1}$, though the critical field at the point of breakdown is likely much higher (Wang, 2013). During a thunderstorm, the electric field builds up and breaks down continuously. The time needed to accumulate large enough electric fields for lightning to occur ranges from less than a minute to several minutes (Gunn, 1954; Marshall and Winn, 1982). For active thunderstorm clouds with tens of kV m^{-1} in the interior, the magnitude of the electric field decreases to 3 kV m^{-1} within 5 km away from the cloud edge on average (Merceret et al., 2008).

Over the years, numerous charging mechanisms were proposed to account for charge separation in thunderstorm clouds. These can be divided into three major categories: convective charging, inductive charging and non-inductive charging. According to the convective charging mechanism proposed by Vonnegut (1955), updrafts carry fair-weather positive charges into the cloud to form a positive charge center. Negative charges are then attracted to the top and edges of the cloud, which are subsequently brought to the lower level by downdrafts. However, numerous investigators such as Chiu and Klett (1976) have found inconsistencies between this mechanism and observations, such as opposite cloud polarity if the cloud forms close to the ground. Inductive charging includes different charge separation mechanisms that involve charges induced by the external fair-weather electric field, such as charging by selective ion capture (Wilson et al., 1929), drop breakup charging and particle rebound charging. However, many studies have shown that these mechanisms are quantitatively unrealistic or ineffective as they are only applicable when the electric field strength is below the typical thresholds required for lightning initiation in thunderstorms (Pruppacher and Klett, 1980; Wang, 2013). For non-inductive charging, charge separation occurs without the presence of an external electric field. Under this category, the most widely accepted mechanism is charging due to the collision of ice crystals with riming graupel ~~pellet~~pellets, which was first studied in the laboratory by Reynolds et al. (1957). It was found that graupel pellets that are growing by the accretion of supercooled droplets ~~acquires~~acquire negative charges as they collide with ice crystals. Takahashi (1978) further investigated this phenomenon and found that the magnitude and sign of the electrification depend largely on temperature and cloud water content. The optimal cloud water content for graupel to become highly charged is $1\text{ to }2\text{ g m}^{-3}$. Graupel will become positively charged if the temperature is above the charge reversal temperature T_R , which ranges from $-20\text{ }^\circ\text{C}$ to $-10\text{ }^\circ\text{C}$, and negatively charged otherwise (Takahashi, 1978). Within the updraft column in a thundercloud where temperature is ~~low~~below T_R , negatively charged graupel and positively charged ice crystals will be formed. The negatively charged graupel will fall at the periphery of the column where the updraft is weak, while the positively charged ice crystals with negligible fall velocity will be thrown upwards. As the graupel reach a region warmer than T_R , they become positively charged, forming the tripole structure of most thunderclouds. Although non-inductive charging due to the collision of graupel and ice crystals best explains tripolar cloud structure, it should be noted that all charging mechanisms above could contribute to certain extent at some time to cloud charging even though these mechanisms alone would produce inadequate or reversed charges (Pruppacher and Klett, 1980).

To know what could be observed in thunderstorm clouds, it is important to first identify the ingredients of thunderstorms. A wide variety of ice particles can be found in thunderclouds. Ice crystals of different shapes and sizes can be formed at different temperatures and ice supersaturation (Bailey and Hallett, 2009). These crystals can grow within clouds through three major processes (Pruppacher and Klett, 1980; Lamb and Verlinde, 2011): riming, water vapor diffusional growth and aggregation. Riming occurs when supercooled water droplets collide with ice crystals and freeze on them, ~~forming large, dense and near~~

~~spherical particles~~generally resulting in increased particle size, density and sphericity. Conical graupel can be formed if riming occurs while particles fall through strong updrafts containing water droplets. Since the bottom windward side of the particle grows faster than the top leeward side, the particle develops a conical shape (Tang et al., 2017). Scattering simulations carried out by Oue et al. (2015) and ~~Lu et al. (2016)~~ showed data from the scattering database created by Lu et al. (2016) indicate that conical graupel can produce negative differential reflectivity (Z_{DR}) values at X-, Ka- and W-band. Diffusional growth takes place when water vapor diffuses towards ice crystals from gas phase. During this process, crystals keep their characteristic shape (Lamb and Verlinde, 2011). Aggregation occurs when ice crystals collide with each other and form larger crystals that are particles that tend to be more spherical in shape. ~~When~~ Various lab measurements have demonstrated that when an electric field of more than around 50 kV m⁻¹ is present, aggregation of ice crystals may be enhanced due to attractive electrical forces induced between neighbouring conducting crystals (Connolly et al., 2005), forming elongated chains rather than spherical clusters almost spherical clusters (Connolly et al., 2005). The efficiency of this process is the highest at approximately -10 °C according to laboratory studies, but these studies are only conducted at temperatures higher than -20 °C (Connolly et al., 2005). In the atmosphere, chain-like aggregates are observed in convective storms at temperatures below -40 °C (Connolly et al., 2005; Stith et al., 2002). Figure 1(a) shows some examples of plate crystals arranged into chains in anvil clouds, i.e. the region of convective cloud detraining from the main cell of the thunderstorm cloud, captured by a cloud particle imager taken by Connolly et al. (2005) at an altitude of around 12 km where the temperature is below -40 °C. Chain-like aggregates can also be formed from frozen droplets, such as those observed by Gayet et al. (2012) near the top of an overshooting convective cloud at 11080 m where the temperature is -58 °C as shown in Fig. 1(b). The enhancement of aggregation starts to decrease when the electric field exceeds 150 kV m⁻¹ since the strong electric field would fragment the ice particle (Connolly et al., 2005). Meanwhile, laboratory experiments have found that electric field enhanced aggregation does not occur when the ice particle number concentration is below 2 cm⁻³ (Wahab, 1974). High concentration concentrations of ice particles could be present in convective clouds if strong updrafts carry supercooled droplets to a level of -37 °C where they freeze rapidly by the process of homogeneous nucleation (Gayet et al., 2012).

~~Evidences~~ Evidence of the presence of graupel and ice crystals in thunderstorm clouds were found using polarimetric and Doppler measurements. Mattos et al. (2016) used X-band radar to compare storms with and without lightning activities and analysed the vertical distribution of hydrometeors within the clouds. They found that in the lower layer of thunderclouds (from 0 to -15 °C), there is an enhanced positive specific differential phase shift (K_{DP}) probably associated with supercooled oblate raindrops lofted by updraft; in the middle layer (from -15 to -40 °C), there is negative Z_{DR} and K_{DP} and moderate horizontal reflectivity, which are possibly associated with the presence of conical graupel. With Ka-band cloud radar, Sokol et al. (2020) identified a mixture of hydrometeors at an elevation of 4-7 km (from -6.6 to -27°C) with a predominance of ice and snow particles and graupel based on the terminal velocities of different hydrometeors. The coexistence of different types of hydrometeors is supported by the measured high Doppler spectrum width, which also implies the existence of collisions of hydrometeors.

In addition to the existence of a variety of hydrometeors in thunderstorm clouds, it was first suggested by Vonnegut (1965) based on changes in cloud brightness observed during lightning that ice crystals would align under strong electric field. Wein-




Figure 1. Examples of (a) plate crystals arranged into chains in anvil clouds taken by Connolly et al. (2005) (chain lengths from left to right are 381, 632 and 721 μm respectively) and (b) frozen drops arranged into chains near the top of an overshooting convective cloud taken by Gayet et al. (2012).

heimer and Few (1987) studied the magnitude of electric field needed to align particles of different sizes and shapes. They compared the magnitudes of electrical torques that try to align particles' long axis with the electric field, and aerodynamic torques that attempt to align particles with their long axes perpendicular to their direction of motion. They estimated that for an electric field of 100 kV m^{-1} , plates with a major dimension of less than 0.6 mm can be aligned, while the threshold is 1 mm for dendrites and 0.2 mm for thick plates. Columns of all sizes can be aligned by such a field. Meanwhile, only particles smaller than 0.05 mm can be aligned by an electric field of 10 kV m^{-1} . Such alignment of ice crystals is observed in various thunderstorm cases using polarimetric radar measurements. For example, Lund et al. (2009) observed negative Z_{DR} in or near clusters of lightning initiations using S-band radar, while Mattos et al. (2016), using X-band radar, found that in the upper layer (above $-40 \text{ }^\circ\text{C}$) of thunderclouds, K_{DP} becomes more negative with increasing lightning density. These are likely due to ice particles being aligned vertically by a large vertical electric field. Meanwhile, only one study that used cloud radar to study the alignment of ice crystals during thunderstorms is found. Using a Ka-band radar, Sokol et al. (2020) observed high linear depolarisation ratio (L_{DR}) in clouds that produce lightning in the vicinity, which is likely caused by the **alignment-canting** of ice crystals in an electric field.

Another important ingredient for lightning is strong updraft. According to Zipser and Lutz (1994), lightning is highly unlikely if the mean updraft speed is less than around 6 to 7 m s^{-1} , or the peak updraft speed is less than around 10 to 12 m s^{-1} . Deierling and Petersen (2008) found that time series of updraft volume in the charging zone where the temperature is below $-5 \text{ }^\circ\text{C}$ with vertical velocities exceeding 10 m s^{-1} is highly correlated to total lightning activity. In general, it is common to find updrafts of more than 10 m s^{-1} and up to 30 m s^{-1} in thunderstorms [Stith et al. \(2016\)](#); [Marshall et al. \(1995\)](#) ([Stith et al., 2016](#); [Marshall et al., 1995](#)).

Up to this date, most research about thunderstorms made use of S-band (2-4 GHz), C-band (4-8 GHz) and X-band (9-12 GHz) radar, while limited studies were conducted using cloud radar with millimeter wavelength. Radars at lower fre-

quencies are common choices for investigating thunderstorms as they have larger ranges and suffer from less attenuation, but high frequency cloud radars could bring new insights on thunderstorm clouds before precipitation starts given their higher spatial resolution. Moreover, existing studies ~~have only analysed integrated~~ of thunderstorms have generally analysed integrated polarimetric radar variables that include the contribution of all particles within each radar resolution volume. ~~There~~ Polarimetric Doppler spectra are investigated at C-band in the context of RELAMPAGO field experiment in Argentina in (Aiswarya Lakshmi et al., 2024). However, there have been no attempts to utilise the polarimetric Doppler spectra at millimeter wavelength to disentangle the contributions of different types of particles ~~Nonetheless, in thunderstorm clouds. At millimeter wavelength, complications occur because~~ variations in the Doppler spectra not only can indicate another type of particles, but also the presence of Mie scattering regime when the particles grow. This study explores new ways to study thunderstorm events by using cloud radar observations and polarimetric Doppler spectra. The goal is to establish links between radar observations and physical processes in thunderstorms to enhance our understanding about lightning.

The manuscript is organized as follows: Section 2 provides essential details on the instruments, data, and case study. Section 3 outlines the methodology for conducting spectral polarimetric analysis of thunderclouds, focusing on key radar variables and the scattering simulation used (T-matrix method for spheroids and cylinders). Special attention is given to spectral differential reflectivity (sZ_{DR}) and spectral differential backscatter phase ($s\delta_{co}$). In Section 4, scattering simulation results illustrate how sZ_{DR} and $s\delta_{co}$ vary with ice particle radius, considering factors such as axis ratio, ice fraction, and canting angle. Section 5 applies this background to two thundercloud case studies, emphasizing ice particle alignment and notable microphysical properties. Finally, Section 6 presents the study's conclusions.

135 2 Instruments and data

The cloud radar used in this study is a dual-frequency scanning polarimetric frequency-modulated continuous-wave (FMCW) radar produced by Radiometer Physics GmbH located at Cabauw, the Netherlands (51.968° N 4.929° E). It operates at 35 GHz (Ka-band) and 94 GHz (W-band) in Simultaneous Transmission Simultaneous Reception (STSR) mode and measures at a constant elevation of 45° and constant azimuth of 282° (see Fig. A1) at some selected periods. Its half power beam width at 35 GHz is 0.84° and temporal sampling is 3.59 s. In this study of thunderstorm clouds, only the 35 GHz data is used since there are numerous issues associated with the 94 GHz data including significant attenuation, less sensitivity at large heights, Doppler aliasing and complications due to Mie-scattering resonance. The configuration parameters for each chirp sequence is shown in Table 1. The ~~transmission-transmitted~~ power is continuously monitored, and the radar receiver (including the receiving antenna) undergoes calibration every six months using clear sky calibration. Short-term calibration is provided through periodic Dicke switching. Prior to the semiannual calibration procedure, the hydrophobic antenna radomes are replaced.

The cloud radar provides two types of output data. The Level 0 dataset contains the raw data, which includes the Doppler spectrum at horizontal and vertical polarizations (sZ_{hh} and sZ_{vv}), as well as the real and imaginary parts of the covariance spectrum between horizontal and vertical polarizations ($sC_{hh,vv}$). The Level 1 dataset contains processed data, including the equivalent radar reflectivity factor (Z_e , or Z_{hh}), mean Doppler velocity, Doppler spectrum width, differential reflectivity

Table 1. Configuration parameters of cloud radar at 35 GHz at 45° elevation for each chirp sequence.

Attributes	Chirp sequence		
	1	2	3
Integration time (s)	1.20	0.96	0.82
Range interval (m)	119.2-1192.5	1222.3-4889.1	4953.3-14969.9
Range resolution (m)	29.8	29.8	55.0
Nyquist velocity (\pm m s ⁻¹)	19.7	16.1	10.7
Doppler velocity resolution (m s ⁻¹)	0.15	0.13	0.17

150 (Z_{DR}), copolar correlation coefficient (ρ_{hv}), specific differential phase shift (K_{DP}), and slanted linear depolarization ratio (SL_{DR}). SL_{DR} is a proxy for L_{DR} , which can **only** be computed when the radar transmits horizontally and vertically polarized electromagnetic waves alternatively. Since the radar used in this study transmits them simultaneously, only SL_{DR} is available. ~~The radar also has~~ Compared to L_{DR} , SL_{DR} in the STSR mode loses the direct mean canting angle information due to the inability to acquire cross-polar measurements, but retains information on the variance of the canting angles and axis ratios. The

155 radar has also a passive broad band channel operated at a centre frequency of ~~31.4-89~~ GHz that provides information about the integrated liquid water path (LWP). A weather station is **also** attached, which provides rain rate, surface wind speed and wind direction, but does not provide the wind profile. Wind profile is obtained instead from European Centre for Medium-Range Weather Forecasting (ECMWF) Integrated Forecast System (IFS) output over Cabauw (O'Connor, 2022) available at <https://cloudnet.fmi.fi/>. This model provides hourly forecast of zonal (eastward) and meridional (northward) wind up to 80000

160 m with a horizontal resolution of 9 km. The model uses an eta-coordinate system, with vertical resolution of the first 10000 m ranges ranging from around 20 m near the surface to around 300 m at the top. A microwave radiometer beside the radar provides temperature and relative humidity profiles along the zenith. Lightning data is obtained from the online lightning map at meteologix.com provided by Siemens BLIDS. ~~Clicking on the lightning stroke gives its~~ The location, time, type, charge (positive or negative) and power of each lightning strike is given.

165 The thunderstorm case being studied took place on 18 June 2021 from 16:15 to 17:45 UTC near Cabauw. Four major thunderstorm clouds (numbered in Fig. 2, Fig. A1, Fig. A2 and Fig. A3) crossed the line of sight of the radar from southwest to northeast. The equivalent reflectivity factor, Z_e , and rain rate from 16:00 to 17:59 UTC are shown in Fig. 2, while Z_{DR} , K_{DP} , SL_{DR} and ρ_{hv} are shown in Fig. 3 and Fig. 4. Note that Z_e , SL_{DR} and ρ_{hv} are taken directly from the Level 1 files, while Z_{DR} and K_{DP} are re-calculated from Level 0 files and calibrated.

170 It is evident from Fig. 2 that due to significant attenuation, the top part of the second and fourth clouds which produced precipitation that reached the ground are missing. Some artefacts are observed, such as the noise from ground level to 2500 m over the entire period, and the ‘ghost’ signals between 2500 m and 3500 m from 16:10 to 16:25 UTC and from 17:30 to 17:40 UTC, which are likely due to signals from the top of the cloud being folded into the second chirp. These artefacts are also




Figure 2. Equivalent reflectivity factor on 18 June 2021 16:00-17:59 UTC. Black line shows the rain rate.

present in other variables, thus the data in the second chirp might not be reliable. From Fig. 2, no melting layer with high Z_e is visible even though the temperature was about 0°C at around 4000 m, which is likely due to convective mixing. However, after 17:15, a brief indication of a melting layer can be observed using the radar variables, Z_{DR} , SL_{DR} and ρ_{hv} .

From Fig. 3(a) and Fig. 4(a), negative Z_{DR} and high SL_{DR} values are observed from 16:42 to 16:48 UTC and from 17:24 to 17:30 UTC, which could be associated to-with the alignment of particles near lightning. From Fig. 4(b), lower ρ_{hv} values of 0.9 are found also from 16:42 to 16:48 UTC and from 17:24 to 17:30 UTC, which could suggest that there may be a mixture of hydrometeors in the cloud. However, at those times and locations, the decreasing ρ_{hv} and increasing SL_{DR} values could be due to a lower signal-to-noise ratio (SNR) because of the attenuated equivalent reflectivity factor, thus caution is required when interpreting these values. Also the differential reflectivity may be impacted by rain differential attenuation. Therefore, these times/locations will not be discussed further. Comparing Fig. 3(a) and (b), Z_{DR} and K_{DP} show different patterns in some areas, such as in the first high cloud and in the top part of the cloud from 17:20 to 17:25 UTC. These will be further investigated.

Fig. 5 shows the mean vertical velocity, vertical air velocity and Doppler spectrum width during the thunderstorm. The mean vertical velocity in Fig. 5(a) eliminates from the measured mean Doppler velocity the contribution of horizontal wind in the same hour obtained from ECMWF model data from the measured mean Doppler velocity forecast initialised at 17 June 2021 12:00 UTC. For such a complex system as thunderstorm, this leads to a first approximation of the mean vertical velocity of hydrometeors. In the first cloud from 16:10 to 16:30 UTC, particles are mainly falling, while in the other clouds, there are alternate regions where particles are falling and rising. The vertical air velocity is obtained from the smallest Doppler velocity bin. From Fig. 5(b), vertical air velocity varies a lot within the clouds. There are regions with upward velocity exceeding 20 m s^{-1} , which shows there may be strong updrafts in the thunderstorm clouds. There are also adjacent regions with upward and downward motion, such as near 16:22 and 17:20 UTC. These may represent convective motion in the clouds. Figure 5(c)




Figure 3. (a) Z_{DR} and (b) K_{DP} on 18 June 2021 16:00-17:59 UTC.

195 shows that some regions in the clouds have high Doppler spectrum width, such as within the first cloud and near the top of the fourth cloud. This could mean that there is a wide variety of particles within the radar resolution volume or the Doppler spectrum is broadened by turbulence or shear (Doviak and Zrnica, 2006; Feist et al., 2019).

For a better understanding of the cloud radar data, weather radar images from 16:15 to 17:40 UTC are shown in Fig. A1, Fig. A2 and Fig. A3 (Kachelmann GmbH). Lightning strikes within the 5 minutes prior to the labelled time are marked by yellow asterisks. The red triangle shows the cloud radar location and the red ruler shows the line of sight of the cloud radar with each mark equal to 1 km. Lightning occurred in all four major clouds labelled in Fig. 2. For the first cloud, lightning occurred at least near the line of sight at more than 10 km away from the radar. For the second cloud, lightning occurred at a perpendicular distance of around the ranges 3 to 8 km from the radar with a cross-range varying from 1 to 10 km. The third cloud only produced two lightning strikes after passing through the line of sight of the radar. The strikes were at a perpendicular distance of around 6 to 7 km from the radar. The fourth cloud produced a large number of lightning strikes near the radar line of sight from less than 1 km to more than 15 km away from the radar along range. Lightning was most active from 17:15 to 17:25 UTC, and became less active as the cloud passed through the line of sight of the radar and moved away.

205




Figure 4. (a) SL_{DR} and (b) ρ_{hv} on 18 June 2021 16:00-17:59 UTC.

3 Methodology

This section explains the steps required to analyse radar data to investigate thunderstorm events. First, the way to compute polarimetric and Doppler variables from raw data is explained in Sect. 3.1. Then, Sect. 3.2 explains how integrated variables and Doppler spectra were used to investigate properties of the thunderstorm clouds. Finally, Sect. 3.3 explains the motivation and method of performing scattering simulations.

3.1 Radar variables

3.1.1 Polarimetric variables calculation

~~Spectral~~ This research utilized spectral polarimetric radar variables which are estimated from the raw data (derived directly from the Level 0) were used in this research data. Consequently, most the majority of the integrated radar variables are were also computed from Level 0 data. This approach facilitates consistency checks between Level 0 and Level 1 data, enables spectral domain filtering when necessary, and allows for the dealiasing of Doppler spectra prior to the calculation of Doppler moments.




Figure 5. (a) Mean vertical velocity, (b) vertical air velocity and (c) Doppler spectrum width on 18 June 2021 16:00 - 17:59 UTC from 35 GHz radar with 45° elevation.

220 The integrated Z_{DR} and Ψ_{DP} ([differential phase shift](#)) can be computed by:

$$Z_{DR}(r, t) = 10 \log_{10} \left(\frac{\sum_v s Z_{hh}(r, v, t)}{\sum_v s Z_{vv}(r, v, t)} \right) \quad (1)$$

$$\Psi_{DP}(r, t) = \arctan \left(\frac{\sum_v \Im(-s C_{hh,vv}(r, v, t))}{\sum_v \Re(s C_{hh,vv}(r, v, t))} \right) \quad (2)$$

The covariance spectrum $sC_{hh,vv}$ corresponds to the Level 0 array CHVSpec. The minus sign in Eq. (2) is added in order to obtain the right trend for K_{DP} in rain, namely positive at 35 GHz and negative at 94 GHz. Here, r is the range, v is the Doppler velocity and t the time. Only data with signal-to-noise ratio above 10 dB were included in the summations to be consistent with the analysed spectral data.

The spectral differential reflectivity (sZ_{DR}) and spectral differential phase shift ($s\Psi_{DP}$) can be computed by:

$$sZ_{DR}(r, v, t) = 10 \log_{10} \left(\frac{sZ_{hh}(r, v, t)}{sZ_{vv}(r, v, t)} \right) \quad (3)$$

$$s\Psi_{DP}(r, v, t) = \arctan \left(\frac{-\Im(sC_{hh,vv}(r, v, t))}{\Re(sC_{hh,vv}(r, v, t))} \right) \quad (4)$$

Only the part of the spectra with signal-to-noise ratio above 10 dB were used to exclude the noisy edges of the spectra where values often fluctuate significantly (Yu et al., 2012). In addition, the spectra were smoothed using a 5-point moving average in Doppler bin to reduce noise. For this study, an extra polarimetric calibration was carried out using vertical profiles of precipitation involving high precipitating clouds. This procedure resulted in reducing the expected error associated with Z_{DR} and Ψ_{DP} from 0.18 dB to 0.05 dB and from 1.6° to 0.6° respectively.

The SL_{DR} and ρ_{hv} values were taken from the Level 1 dataset.

The specific differential phase shift (K_{DP}) was approximated from the calibrated Ψ_{DP} in degrees in two steps. First, Ψ_{DP} was smoothed using a 5-point moving average in range to reduce noise. Then, K_{DP} was computed by

$$K_{DP}(r, t) = \frac{\Delta \Psi_{DP}(r, t)}{2\Delta r} [^\circ \text{ km}^{-1}], \quad (5)$$

where Δr is the distance between adjacent range bins in km. Note that this quick estimation of the specific differential phase shift is meant for detecting areas of interest in thunderstorm cloud profiles. For quantitative values of K_{DP} , this processing may be too simple when large sized ice particles are present in the thunderstorm cloud and Mie-non-Rayleigh scattering occurs.

3.1.2 Doppler variables calculation

The measured Doppler velocity v of a particle, defined negative as the particle approaches the radar, is given by

$$v = (w - V_t) \sin \theta + v_H \cos \theta \cos(D - \pi - \phi), \quad (6)$$

where w is the vertical air velocity, v_H is the horizontal wind speed, V_t is the terminal fall velocity of the particle (positively defined), and θ is the elevation angle of the radar, D is the wind direction relative to North and ϕ is the azimuth angle of the radar beam relative to, both being relative to True North. The mean Doppler velocity can reflect the average motion of particles in a radar resolution volume along the line of sight of the radar. To extract it from Level 0 data, the first step is to unfold and dealias each Doppler spectrum. Then, the mean Doppler velocity ($\overline{v_D}$) can be computed by

$$\overline{v_D}(r, t) = \frac{1}{Z_{hh}(r, t)} \sum_{v_{\text{SNR}} > 10 \text{ dB}} v \times sZ_{hh}(r, v, t). \quad (7)$$

The Doppler spectrum width (σ_{v_D}) can also be computed by

$$\sigma_{v_D}(r, t) = \sqrt{\frac{1}{Z_{hh}(r, t)} \sum_{v_{SNR>10 \text{ dB}}} (v - \overline{v_D}(r, t))^2 \times sZ_{hh}(r, v, t)}. \quad (8)$$

The mean vertical velocity ($\overline{w - V_t}$) can give information about the vertical motion of hydrometeors in thunderstorm clouds. It can be estimated by solving Eq. (6) using the mean Doppler velocity ($\overline{v_D}$) together with v_H and D estimated from the
255 ECMWF model data.

It is also useful to extract the vertical air velocity, which can give information about the updraft and downdraft pattern in thunderstorm clouds. It can be estimated by assuming that the smallest particles in the Doppler spectra are so light that their fall velocity is very close to zero, thus their vertical velocity is equal to the vertical air velocity. Therefore, the first step is to identify the Doppler velocity of the ~~rightmost valid bin of~~ bin with the highest Doppler velocity value in the Doppler spectra
260 with a 10 dB SNR threshold. Then, the vertical air velocity w can be estimated by solving Eq. (6) with $V_t = 0$ and v_H and D estimated from the ECMWF model data.

3.2 Analysing radar variables

3.2.1 Analysing integrated variables

Integrated variables were used in this study to identify time instants and ranges where signals related to lightning activities
265 are present. During lightning, the electric field in clouds ~~would~~ can align ice crystals vertically, causing Z_{DR} and K_{DP} to become negative. When negative Z_{DR} or K_{DP} is observed in the integrated profile, more in depth analyses were carried out by investigating sZ_{DR} and $s\Psi_{DP}$ at those time instances to understand the causes of those negative values.

Other useful variables may be the linear depolarisation ratio (L_{DR}) and the copolar correlation coefficient (ρ_{hv}). High L_{DR} values may indicate canting of ice crystals in a specific direction due to cloud electrification (Sokol et al., 2020). With regard
270 to SL_{DR} , areas with low values may result from a reduction in canting angle variance caused by the alignment of ice particles. Regions with low ρ_{hv} could be regions where graupel and ice crystals co-exist, and they may collide with each other to produce an electric field. However, when SNR is low, SL_{DR} and ρ_{hv} values may become large and low, respectively, regardless of the characteristics of the particles. Therefore, analysis was made at sufficient SNR, which is above 10 dB.

3.2.2 Analysing Doppler ~~spectrums~~ spectra

275 While integrated variables contain information about all particles within a radar resolution volume, Doppler spectra separate the contributions of particles with different Doppler velocities, hence different sizes or densities. With spectral Z_{DR} , it would be possible to identify whether negative Z_{DR} is contributed by small particles that would appear on the right part of the Doppler spectrum, or by large particles that would appear on the left part of the Doppler spectrum. If negative Z_{DR} is observed for small particles, it is likely that an electric field is present that aligns the small particles. On the other hand,
280 based on the database described by (Lu et al., 2016), negative Z_{DR} for large particles only may indicate the presence of con-

ical graupel (Lu et al., 2016). However, the possible transition from Rayleigh to Mie scattering regime may complicate these interpretations of spectral Z_{DR} .

The vertical gradient of the differential phase shift (Ψ_{DP}) is related to K_{DP} . A positive gradient indicates positive K_{DP} and vice versa. With the use of $s\Psi_{DP}$ the Mie scattering regime can be identified. As mentioned before, fluctuations in sZ_{DR} values in the Mie scattering regime makes it difficult to interpret those values. It is therefore crucial to identify when the Mie scattering regime begins. ~~To do this, a useful fact is that the~~ This is done by making use of the following relationship between differential phase shift (Ψ_{DP}) ~~is the sum of the~~ two-way differential propagation phase (Φ_{DP}) and the differential backscatter phase (δ_{co}): $\Psi_{DP} = \Phi_{DP} + \delta_{co}$.

In the Rayleigh scattering regime, where δ_{co} is zero, the spectral differential phase shift ~~should be constant since at a fixed~~ range remains constant because the electromagnetic wave ~~that scatters from particles at a particular range has propagated at~~ both polarizations has passed through the same ~~set of~~ particles in all ~~previous ranges, and δ_{co} is zero~~ preceding ranges. This part of the spectrum is often referred to as the Rayleigh plateau (Unal and van den Brule, 2024). In the Mie scattering regime, δ_{co} is non-zero and depends on the particle properties, thus the differential phase shift spectrum is no longer flat. Therefore, the Mie scattering regime begins when the left part of the differential phase shift spectrum starts to increase or decrease. The effect of noise may sometimes affect the identification of the Mie scattering regime. It is useful to know that the maximum or minimum of spectral Ψ_{DP} are often aligned with the maximum or minimum of spectral Z_{DR} . Thus, if the maxima or minima of $s\Psi_{DP}$ and sZ_{DR} are aligned, one can be more confident that the fluctuations observed are due to Mie scattering resonance instead of noise.

The left column of Fig. 6 shows an example where the Mie scattering regime can be clearly identified using $s\Psi_{DP}$. The Rayleigh plateau is found from -1 to 3 m s^{-1} , while Mie-non-Rayleigh scattering occurs at Doppler velocity smaller than -1 m s^{-1} since $s\delta_{co}$ becomes non-zero. sZ_{DR} follows a similar trend, which strengthens the proof that Mie-non-Rayleigh scattering occurs. However, some cases can be more tricky, such as the one shown in the right column of Fig. 6. Here, the Rayleigh plateau ends at about -0.5 m s^{-1} , while sZ_{DR} only begins to decrease at about -4 m s^{-1} . To understand this better, scattering simulations are needed, which is discussed next.

3.3 Scattering simulations

Studying the Doppler spectrum of Z_{DR} is challenging when Mie scattering resonance is involved. This is because sZ_{DR} values fluctuate in the Mie scattering regime, so it will become difficult to determine whether the fluctuations in the observed Z_{DR} spectrum are due to changes in shape/density of hydrometeors or Mie scattering resonance. Therefore, scattering simulations were carried out to understand how Mie-non-Rayleigh scattering affects the Z_{DR} spectrum using the python code pyTmatrix (Waterman, 1965; Leinonen, 2014). The code is based on the T-matrix method (Waterman, 1965), which is a numerical model of electromagnetic and light scattering by non-spherical particles with sizes comparable to the wavelength of the incident radiation. The code supports simulations of spheroids or cylinders. The scattering matrix of a scatterer depends on several parameters, including the axis ratio, ice fraction and canting angle. The axis ratio is defined as the length along the scatterer's rotational axis to its width perpendicular to this axis. It is smaller than one-1 for oblate particles and larger than one-1 for




Figure 6. Two examples of Doppler spectra of Z_e , Z_{DR} and Ψ_{DP} at 35 GHz showing [Mie-non-Rayleigh](#) scattering.

315 prolate particles. Ice fraction (f_i) characterizes how much ice and air a scatterer is composed of, which affects the density of the particle. A value of 1 means pure ice, while a value of 0 means pure air. Ice fraction affects the complex effective relative permittivity of the scatterer (ϵ_{eff}). One approximation is given by the Maxwell-Garnett formula:

$$\frac{\epsilon_{eff} - 1}{\epsilon_{eff} + 2} = f_i \cdot \frac{\epsilon_i - 1}{\epsilon_i + 2}, \quad (9)$$

where ϵ_i is the complex relative permittivity of ice. The value of ϵ_i is $3.19015 + 0.00285i$ at 35 GHz at 266 K, and the temperature dependence is small for the part of the spectrum from ultraviolet (175 nm) to the microwave (1 cm) (Warren and Brandt, 2008). The complex effective refractive index of the scatterer (m_{eff}), which is a parameter that can be specified in the simulation code, can then be determined using

$$\epsilon_{eff} = m_{eff}^2 = \sqrt{\epsilon_{eff}}. \quad (10)$$

The canting angle refers to the Euler angle β of the scatterer defined in Fig. 7.

325 In the simulation, a scatterer object in the shape of a spheroid was defined, and the backscatter radar reflectivity (Z_e), differential reflectivity (Z_{DR}) and differential backscatter phase (δ_{co}) at 35 GHz, with 45° looking angle were retrieved. [The axis ratio and ice fraction of the particles in the simulation experiments were chosen according to the data given in Spek et al. \(2008\)](#)




Figure 7. Definition of Euler angles α and β . The xyz coordinate frame has the z axis aligned with the radar's zenith direction. The rotated frame is denoted as $x'y'z'$, corresponding to the particle's orientation. Starting from the xyz frame, a rotation by angle α around the z axis results in the intermediate frame $x'y_1z$. This is followed by a rotation by angle β around the x' axis to achieve the final $x'y'z'$ frame.

In the first experiment, the axis ratio of scatterers with spheroids with a zero mean canting angle was varied from 0.1 to 1.2, which covers the axis ratio range. This range encompasses the axis ratios of plates, dendrites, aggregates, and graupel. The ice fraction was fixed held constant at 0.6, which is representing the average ice fraction of different types of for the aforementioned ice particles. In the second experiment, the ice fraction of scatterers with spheroids with a zero mean canting was varied from 0.2 to 1, which covers the ice fraction range of plates, dendrites, aggregates and graupel. Simulations for both oblate and prolate particles were carried out, with an axis ratio of 0.8 or 1.2. In the third experiment, the canting angle was varied from 0° to 90° . Three sets of simulations were carried out to simulate different types of particles, including plates (axis ratio = 0.1, ice fraction = 0.98), slightly oblate aggregates (axis ratio = 0.8, ice fraction = 0.3) and graupel (axis ratio = 1.2, ice fraction = 0.6). For all simulations, the canting angles of the scatterers-spheroids follow a Gaussian distribution with a standard deviation of 0.1° . The Euler angle α of the scatterers (see Fig. 7) follows a uniform distribution from 0 to 360° .

Note that the T-matrix method (Leinonen, 2014) offers flexibility for simulating the radar spectral variables by varying different input parameters (axis ratio, ice fraction, Euler angles) for a first examination of trends at 35 GHz. Nonetheless, this method has limitations as it assumes that ice particles are spheroidal and have a fixed ice fraction or density. It ignores the non-homogeneity of ice particles, especially aggregates, which may result in a bias in the spectral polarimetric variables when frequency increases. This is another reason to carry out this study of thunderstorm clouds at 35 GHz but not at 94 GHz.

4 Scattering simulation results

This section gives an overview of the dependencies of spectral polarimetric radar variables of particles, sZ_{hh} , sZ_{DR} and $s\delta_{co}$, on axis ratio, ice fraction and canting angle in the Rayleigh and Mie scattering regimes based on scattering simulations.

4.1 Axis ratio

Figure 8 shows the simulation results for ~~horizontally aligned scatterers~~ spheroids with ice fraction 0.6 and zero mean canting angle with different axis ratios at 35 GHz. The radius refers to the maximum radius of the spheroid, i.e. half the length of its long axis. From Fig. 8(a), the first Mie minimum occurs at a maximum radius of around 2 mm for axis ratio 1.2, 2.6 mm for axis ratio 0.8, and 3.2 mm for axis ratio 0.4. Therefore, for oblate spheroidal particles, the position of the first Mie minimum goes towards larger radius when axis ratio decreases.

From Fig. 8(b), in the Rayleigh scattering regime, Z_{DR} decreases with increasing axis ratio, with positive values for oblate spheroids (axis ratio < 1) and negative values for prolate spheroids (axis ratio > 1). When entering the Mie scattering regime, Z_{DR} of oblate particles increases slightly, while that of prolate particles decreases. At the first Mie minimum, particles with axis ratio 0.1, 0.4 and 1.2 give a trough in Z_{DR} , but those with axis ratio 0.8 give a peak. In addition, the lines for different axis ratios cross over each other in the graph of Z_{DR} , meaning that the trend between Z_{DR} and axis ratio depends on particle size. From Fig. 8(c), δ_{co} of oblate particles increases when entering the Mie scattering regime and gives a peak at the first Mie minimum, while that of prolate particles decreases and gives a trough.




Figure 8. Simulated (a) radar reflectivity, (b) differential reflectivity and (c) differential backscatter phase for horizontally-aligned scatterers spheroids with different axis ratios as a function of maximum radius at 35 GHz with 45° looking angle. All scatterers-spheroids have ice fraction of 0.6 and zero mean canting angle.

4.2 Ice fraction

360 Figure 9 shows two sets of simulations for horizontally-aligned scatterers with spheroids with zero mean canting angle and different ice fractions. The Mie minima can be seen in the reflectivity plots (Fig. 9(a,d)). In the Rayleigh scattering regime, the magnitude of Z_{DR} increases with increasing ice fraction. The first extremum of Z_{DR} is attained at a smaller size for scatterers spheroids with higher ice fraction. For low ice fraction (0.2, 0.4 and 0.6), the sign of Z_{DR} does not change after entering the Mie scattering regime (except for radius larger than 3.2 mm for scatterers-spheroids with axis ratio 1.2 and ice fraction 0.6).

365 When ice fraction is large (0.8 and 1), the sign of Z_{DR} flips soon after reaching the first extremum, and the trend is rather unpredictable. For a radius of particles of this ice fraction with radius larger than 2.5 mm representing large aggregates such as- which could represent graupel, significant negative (positive) values could be obtained, which increases the interpretation challenge. The differential backscatter phase initially increases (decreases) for oblate (prolate) particles when entering the Mie scattering regime. The sign reverses afterwards, and the trend becomes less predictable especially if ice fraction is high.




Figure 9. Simulated radar variables for horizontally-aligned scatterers-spheroids with zero mean canting angle and different ice fractions as a function of maximum radius at 35 GHz with 45° looking angle. Panels (a-c) show the radar reflectivity, differential reflectivity and differential backscatter phase for scatterers-spheroids with fixed axis ratio of 0.8. Panels (d-f) show the same for scatterers-spheroids with fixed axis ratio of 1.2.

370 4.3 Canting angle

Figure 10 shows three sets of simulations for scatterers-spheroids with different canting angles. The Mie minima can be seen in the reflectivity plots (Fig. 10(a,d,g)). A zero mean canting angle corresponds to oblate spheroids being horizontally aligned and prolate spheroids being vertically aligned. To represent prolate particles as horizontally aligned, they are modeled with a mean canting angle of 90 degrees. For oblate particles (left and middle columns), Z_{DR} in the Rayleigh scattering regime is negative when the canting angle becomes larger than 45°. One can understand this as the effective axis ratio of an oblate scatterer-spheroid getting larger than one when it becomes vertically aligned. The opposite is true for prolate particles. However, in the Mie scattering regime, the relationship between the sign of Z_{DR} and the canting angle is not trivial. For scatterers-spheroids similar to plates with axis ratio 0.1 and ice fraction 0.98, the first extremum of Z_{DR} is positive for $\beta = 90^\circ$ but negative for

$\beta = 0^\circ$. There is no sharp extremum for $\beta = 30^\circ$ or 60° . For [scatterers-spheroids](#) similar to conical graupel with axis ratio 1.2 and ice fraction 0.6, the sign of Z_{DR} also changes when particle size becomes larger. The differential backscatter phase does not have a trend that can be easily summarised for different canting angles for all three cases. [In all instances, the most pronounced resonance patterns are found at canting angles \$\beta = 0^\circ\$ and \$\beta = 90^\circ\$.](#)




Figure 10. Simulated radar variables for [horizontally-aligned scatterers-spheroids](#) with different canting angles as a function of maximum radius at 35 GHz with 45° looking angle. [Panels \(a-c\)](#) show the radar reflectivity, differential reflectivity and differential backscatter phase for [scatterers-spheroids](#) similar to plates with fixed axis ratio of 0.1 and ice fraction of 0.98. [Panels \(d-f\)](#) show the same for [scatterers-spheroids](#) similar to [slightly oblate](#) aggregates with fixed axis ratio of 0.8 and ice fraction of 0.3. [Panels \(g-i\)](#) show the same for [scatterers-spheroids](#) similar to conical graupel with fixed axis ratio of 1.2 and ice fraction of 0.6.

4.4 Summary

In this section, the effects of axis ratio, ice fraction and canting angle of [scatterers-spheroids](#) on Z_{DR} and δ_{co} are investigated. [Table 2](#) summarises the key trends of Z_{DR} in the Rayleigh scattering regime and the trend of δ_{co} before the first Mie minimum for [horizontally-aligned scatterers-spheroids](#) with different axis ratios and ice fractions. [Their mean canting angle is zero.](#) Changing the canting angle has similar effect as altering the axis ratio of the [scatterers-spheroids](#) in terms of the initial trend

of Z_{DR} . In general, the sign of Z_{DR} is the same as the sign of δ_{co} before the first Mie minimum. However in some cases, δ_{co} shows a sign inversion at the first Mie minimum. The fluctuations of Z_{DR} and δ_{co} after the first Mie minimum are difficult to predict and often involve sign changes. The most unpredictable behaviours are found when ice fraction is high.

Table 2. Z_{DR} characteristics in Rayleigh scattering regime and trend of δ_{co} before first Mie minimum. The mean canting angle of the spheroids is zero.

	Z_{DR} in Rayleigh scattering regime	δ_{co} trend before first Mie minimum for horizontally aligned scatterers
Axis ratio < 1	positive, increase with decreasing axis ratio	increase
Axis ratio > 1	negative, more negative with increasing axis ratio	decrease
Ice fraction	magnitude increases with increasing ice fraction	same trend as Z_{DR} except for large ice fraction

From this first analysis, our investigation of spectral polarimetric variables in thunderstorm clouds will start by identifying the Rayleigh scattering part of the spectrum using the measurement of the spectral differential phase. In the Rayleigh scattering regime, the spectral differential backscatter phase is zero and the spectral differential phase equals the spectral differential propagation phase. This will prevent the misinterpretation of variations in spectral differential reflectivity caused by ~~Mie scattering resonance~~. Next, focus will be given to the sZ_{DR} signature in the Rayleigh scattering regime. ~~Afterwards~~ Subsequently, analysis can be ~~carried out considering~~ conducted using both sZ_{DR} and $s\delta_{co}$ ~~in within~~ the Mie scattering regime at least ~~until up to~~ the first Mie minimum. ~~Concerning second extrema, they are expected to be~~ Second extrema are challenging to interpret and ~~to measure, the latter in terms of signal-to-noise ratio~~ measure, especially at high altitudes where the signal-to-noise ratio is low.

For each sub-figure, simulations were conducted considering a single type of ice particle. However, in practice, a radar resolution volume may contain multiple types of ice particles, resulting in the final spectral polarimetric variables being composed of different modeled curves as a function of the radius range.

5 Case analysis

This section discusses interesting observations in the thunderstorm event on 18 June 2021 from 16:15 to 17:45 UTC near Cabauw. Focus has been given to the first and the fourth cloud that passed through the line of sight of the radar. The second cloud was not investigated as the radar suffered from significant attenuation due to the precipitation, while the third cloud was not studied as it only had two lightning ~~strokes~~ strikes after it passed through the line of sight of the radar.

5.1 First cloud

The first cloud came within the sight of the radar from 16:10 to 16:30 UTC. The centre of the cloud that contained lightning
410 activities was more than 10 km away from the radar, thus the radar could only see the edge of the cloud.




Figure 11. (a) Differential reflectivity, (b) specific differential phase shift and (c) vertical air velocity of the first thunderstorm cloud on 18 June 2021 from 16:09 to 16:30 UTC.

5.1.1 Alignment of particles

From Fig. 11(a) and (b), interesting-intriguing polarimetric signatures can be found-in-observed within the cloud. As-shown in Fig-Figure 11(a), the-illustrates that Z_{DR} values in the cloud is close to zero and do not show much variations. However, from-are near zero with minimal variation. Conversely, Fig. 11(b), there-is-reveals a cluster of negative K_{DP} values between 415 7600 m and 9300 m, which-may-indicate-suggesting the alignment of non-spherical small ice particles but-not-large-ones. If the- If these small ice particles have-sufficient-number-are present in sufficient concentration, K_{DP} would become negative. However, since Z_{DR} carries reflectivity weighting, The large ice particles that do, on the other hand, are expected to be slightly non-spherical, which leads to a small contribution to K_{DP} , and may not align with the electric field-an electric field unless it is sufficiently strong. Because Z_{DR} is reflectivity-weighted, large ice particles significantly influence Z_{DR} strongly. This could 420 explain, which likely explains why Z_{DR} does not show-exhibit significant negative values.

From Fig. 11(c), the first cloud mainly shows downdrafts from 16:15 to 16:18 UTC and after 16:22 UTC. Referring to Fig. A1, in these periods, the radar was looking at the edge of the thunderstorm cloud. Therefore, the radar did not see regions with strong updrafts which is normally found in the core of thunderstorm clouds, but observed downdrafts outside the core instead. From 16:18 to 16:22 UTC, updrafts of up to 12 m s^{-1} are observed, which could be because the core of the thunderstorm cloud 425 is closer to the line of sight of the radar. The estimated vertical air velocity is not uniform within the cloud, which suggests that there might be a lot of turbulence.




Figure 12. Spectral Z_{DR} on 18 June 2021 from 16:18:59 to 16:22:34 UTC.

Figure 12 shows the spectral Z_{DR} across the period when negative K_{DP} is observed. At 16:18:59 UTC, the right part of the spectrum, which corresponds to small ice particles, has positive sZ_{DR} , suggesting that the particles are horizontally aligned. However, at 16:21:05 UTC, the right part of the spectrum becomes slightly negative, suggesting that small ice particles are 430 vertically aligned. At 16:22:34 UTC, sZ_{DR} of the right part of the spectrum becomes positive again, which suggests that the particles return to being horizontally aligned. Figure 13 shows the mean sZ_{DR} of the highest-smallest 10% of the particles

in each radar resolution volume at the three time instants. This is achieved by averaging sZ_{DR} over the rightmost 10% of the Doppler bins. It is clear that from 7000 m to 9000 m, sZ_{DR} of the highest smallest 10% particles are positive at 16:18:59 UTC and 16:22:34 UTC, and is negative at 16:21:05 UTC. The question is: are these negative sZ_{DR} values associated with cloud electrification before lightning?
 435

Spectral Z_{DR} on 18 June 2021 from 16:18:59 to 16:22:34 UTC.




Figure 13. Mean sZ_{DR} of all particles and the highest smallest 10% of the particles in a radar resolution volume at (a) 18 June 2021 16:18:59 UTC, (b) 16:21:05 UTC and (c) 16:22:34 UTC.

Our expectation is that particles align vertically before a lightning strokestrike, and return to horizontal alignment afterwards. However, the lightning strokes strikes closest to the line of sight of the radar occurred at 16:20:17, 16:21:50 and 16:22:20 UTC (strokes strikes number 9, 11, 14-17 in Fig. B2), but negative K_{DP} is observed continuously from 16:20:11 to 16:21:37 UTC. Negative K_{DP} appears at a distance of 7600 m to 9300 m away from the radar, but the lightning strokes strikes occurred at
 440 least 13 km-13000 m away from the radar. If the electric field that caused these lightning strokes strikes is responsible for the alignment of particles observed, one would expect to observe negative K_{DP} also for ranges beyond 9000 m. Making a closer inspection with spectral Z_{DR} , negative sZ_{DR} smaller than -0.1 dB are found beyond 9000 m from 16:20:21 to 16:21:15 UTC (Fig. 16(b)), though more negative sZ_{DR} are found on the left side of the spectrum that corresponds to large particles instead of the right side as expected (e.g. 16:21:01 UTC in Fig. 16(b)).

445 The first inquiry is whether this could be caused by wind shear-

The first question is whether wind shear could be responsible for flipping the Doppler spectrum such that lighter particles, causing lighter particles to appear on the left side. The horizontal and vertical velocities (V_h and V_v) of particles are given by (Wang et al., 2019) described by Wang et al. (2019) as:

$$V_h = v_H + \frac{sV_t^2}{g}, \quad (11)$$

$$450 \quad V_v = -V_t + w \quad (12)$$

where v_H is the horizontal wind speed, w is the vertical wind, $s = \frac{dv_H}{dz}$ is the constant vertical wind shear, g is gravitational acceleration, and V_t is the terminal velocity of the particle which depends on its size. For a radar looking at an elevation angle θ and azimuth ϕ , the Doppler velocity is $V_v \sin \theta + V_h \cos \theta \cos(D - \pi - \phi)$. Without vertical wind shear, the Doppler spectrum will be shifted as a whole spectrum shifts uniformly by v_H and w , with lighter particles remaining leaving lighter particles on the right side of the spectrum. If s is negative, since V_t increases with particle size, the left side of the spectrum would shift to the left a negative shear s causes the spectrum to widen as the left side shifts more than the right side of the spectrum, thus the spectrum widens right (Fig. 14(b)). On the other hand, if s is positive, the left side of the spectrum would shift to the right more than the right side of the spectrum, while positive shear narrows it (Fig. 14(c)). If the right shift of the left side of the spectrum rightward shift on the left due to the term $\frac{sV_t^2}{g}$ is larger than exceeds the original spectrum width, the spectrum could be flipped flip (Fig. 14(d)). Assuming a spectrum width of For a 10 m s⁻¹ and taking spectrum width and $V_t = 2$ m s⁻¹ corresponding to the upper bound of the terminal velocity of plates ; i.e. $V_t = 2$ m s⁻¹ (Spek et al., 2008), for a radar looking at 45°, the vertical wind shear required to flip the Doppler spectrum is approximately 35 (Spek et al., 2008), a shear of approximately 25000 m s⁻¹ km⁻¹. However, according to the ECMWF Integrated Forecast System output over Cabauw (O'Connor, 2022) shown in Fig. 15(e), this is would be required to flip the spectrum, much larger than the strongest vertical wind shear of 4 m s⁻¹ km⁻¹ observed from 7500 m to 10000 m. Therefore, it is unlikely that in ECMWF data shown in Fig. 15(c). Thus, wind shear is unlikely to explain the negative sZ_{DR} on the left side of the spectrum is due to flipping of the spectrum caused by wind shear.




Figure 14. A figure to illustrate effects of the sign of vertical wind shear s on the Doppler spectrum: (a) Doppler spectrum when there is no shear (b) Doppler spectrum widens when s is negative (c) Doppler spectrum may become narrow when s is positive (d) Doppler spectrum may flip when s is positive and $\frac{sV_t^2}{g}$ is larger than the original spectrum width.

470 ~~If the~~ Alternatively, electric fields could align particles vertically, particularly larger ice chains (Connolly et al., 2005), leading to negative sZ_{DR} ~~values are caused by vertical alignment of particles by the electric fields, one possible explanation for this on the left side of the Doppler spectrum. Here, the hypothesis~~ is that the axis ratios of small particles are close to one, thus their sZ_{DR} is close to zero. Meanwhile, large ice chains may be formed under an electric field (Connolly et al., 2005). sZ_{DR} could become negative if these chains are vertically aligned. Nonetheless, However, the most negative sZ_{DR} is found at 16:21:01 UTC, which is not exactly the moment before any lightning strokes. This could be because the region with chains and sufficiently strong does not coincide directly with lightning, suggesting the electric field had ~~moved away from the line of sight of the radar when lightning occurred, or that the electric field was not strong enough to trigger lightning either weakened or moved out of radar view by the time of the strikes.~~




Figure 15. (a) Mean horizontal wind, (b) horizontal wind direction relative to North and (c) vertical wind shear at 18 June 2021 16:00 and 17:00 UTC. [These data are outputs of the ECMWF Integrated Forecast System \(IFS\) over Cabauw \(O'Connor, 2022\).](#)

480 The negative sZ_{DR} values on the right edge of the spectra at around 7500 m to 9000 m also do not occur right before any lightning strokes. Although there is a lightning at a perpendicular distance of between 7500-9000 m similarly do not align with lightning events occurring at cross-ranges larger or equal to 13 km away from the radar at 16:21:09 UTC (stroke number 5 in Fig. B2), from the literature, electric field in thunderstorm clouds reduces significantly over, as significant electric fields extend only about 5 km (Merceret et al., 2008), thus it is unlikely that the alignment of particles observed is caused by the build up of electric field that generated the lightning stroke. There is another lightning at a perpendicular distance of around 11 km away from the radar in thunderstorms (Merceret et al., 2008). Though a strike at 16:21:50 UTC (stroke number strike 7 in Fig. B2). It is possible that the electric field associated with this stroke caused the vertical alignment of particles observed, but the relationship is hard to prove since the spatial and temporal variation of the electric field is unknown. The next lightning, 485 cross-range 11 km) may have contributed, this is difficult to confirm due to the unknown electric field variation. The subsequent strike at 16:29:08 UTC at a perpendicular distance of around 11.5 km (stroke (strike number 8 in Fig. B3) occurred 8 minutes after negative sZ_{DR} is observed. This is a rather long period of time compared to common durations, cross-range 11.5 km) is too delayed considering common duration of charging cycles (Gunn, 1954; Marshall and Winn, 1982), thus it is unlikely that 490 it is associated with the observed to be connected to earlier negative sZ_{DR} values.

Meanwhile, one should not rule out other possible causes of vertical alignment of particles. As suggested by Brussaard (1976), canting of particles may occur when there is vertical wind shear. He derived the Wind shear-induced particle canting (Brussaard, 1976) is another potential cause. The canting angle of particles due to vertical wind shear, i.e. difference in hori-

zonal wind speed in vertical direction, ~~by assuming is given by~~

495
$$\tan \beta = -\frac{sV_t}{g}$$
 (13)

~~The equation holds assuming a linear wind profile, no updraft and that the mean orientation of their-the particles rotational symmetric axes are-always-is parallel to the direction of the airflow around them. If there is no updraft and the particle is falling at its terminal velocity, for a linear wind profile, the canting angle (γ) of the particle is given by-~~

$$\tan \gamma = -\frac{sV_t}{g}$$

500 ~~To estimate the largest possible canting angle in the current cloud, terminal velocity V_t is taken to be 2 (Brussaard, 1976). Using the vertical shear $s = \frac{dv_H}{dz} = 4 \text{ m s}^{-1}$ (Spek et al., 2008) and $s = \frac{dv_H}{dz} = 4 \text{ km}^{-1} = 0.004 \text{ s}^{-1}$ and terminal velocity of 2 m s^{-1} according to the ECMWF Integrated Forecast System output (O'Connor, 2022) shown in Fig. 15(c). Using Eq. (13), the corresponding, the canting angle is estimated at 0.05° , which is insignificant. Therefore, wind shear is not likely the cause of negligible. Even considering underestimation due to model resolution, achieving significant canting would require a~~
505 ~~much higher shear of 4.9 s^{-1} , making wind shear an unlikely cause of the observed negative sZ_{DR} in this case. On the other hand, it is found by Cho et al. (1981) that turbulence is unable to destroy the preferred orientation of falling ice crystals values. Additionally, turbulence is not expected to disrupt ice crystal orientation in cumulonimbus clouds (Cho et al., 1981).~~

In conclusion, the vertical alignment of particles observed in the first cloud could be due to electric field, though the electric field may not be strong enough to trigger lightning, or there are ~~lightnings-lightning strikes~~ that are not measured by the
510 lightning sensor.

5.1.2 ~~Supercooled liquid water~~ Interesting microphysical properties

5.1.2.1 ~~Supercooled liquid water~~

Another interesting feature observed in this cloud is the possible presence of supercooled liquid water. From 16:20:21 to 16:21:15 UTC, ~~spectograms spectrograms~~ of reflectivity show a separate mode of particles on the right side of the spectrum at
515 around 6000 m (see Fig. 16(a)), ~~where air temperature measured by the microwave radiometer is around -12.5°C .~~ From Fig. 16(b), sZ_{DR} of this mode of particles is close to zero.




Figure 16. (a) Spectral reflectivity and (b) spectral Z_{DR} on 18 June 2021 from 16:20:21 to 16:21:15 UTC showing presence of supercooled liquid water near 6000 m.

Figure 17 shows the time series of spectral reflectivity and spectral Z_{DR} at 5916 m. A small peak at Doppler velocity of around -4 to -3 m s^{-1} is consistently present. The sZ_{DR} of this mode of particles is lower than the left part of the spectrum, with values of around -0.1 to 0 dB. By manually identifying the part of the Doppler spectrum that may contain supercooled liquid water for 139 range bins over 16 time steps, it was found that the average sZ_{DR} is -0.0370 dB. Since the error of Z_{DR} after calibration is 0.05 dB and small supercooled liquid water droplets are nearly spherical and have a differential reflectivity of 0 dB, there is a high chance that supercooled liquid water is indeed present in the cloud. This is further supported by the liquid water path measured by the cloud radar with a passive channel that has the same looking direction as the radar. From 16:20:21 to 16:21:15 UTC (marked by the red lines in Fig. 18), there is indeed a peak in liquid water path, which agrees with the hypothesis that supercooled liquid water may be present in the cloud. Supercooled liquid water plays a role in the non-inductive charging mechanism as it is needed for riming to occur, which in turn forms graupel that collide with ice crystals to produce charges. Nonetheless, the radar was not able to look at the lower part of the cloud, thus it is unknown whether graupel is formed in this case.

(a) Spectral reflectivity and (b) spectral Z_{DR} on 18 June 2021 from 16:20:21 to 16:21:15 UTC showing presence of supercooled liquid water near 6000 m.




Figure 17. (a) Spectral reflectivity and (b) spectral Z_{DR} on 18 June 2021 from 16:20:32 to 16:21:08 UTC at 5916 m.




Figure 18. Liquid water path of the first cloud measured at 94.89 GHz. Time interval from 16:20:21 to 16:21:15 UTC is marked by red lines.

5.2 Fourth cloud

530 The fourth cloud came within the sight of the radar from 17:15 to 17:40 UTC. The part of the cloud that passed through the line of sight of the radar from 17:15 to 17:20 UTC did not contain active lightning activities. From 17:20 to 17:35 UTC, the part of the cloud with the most active lightning activities passed through the line of sight of the radar. Afterwards, lightning activities ceased and the cloud moved away from the line of sight of the radar. For an overview of the cloud including the radar images showing its motion, see Appendix A.




Figure 19. (a) Differential reflectivity, (b) specific differential phase shift, (c) slanted linear depolarisation ratio and (d) copolar correlation coefficient of the fourth thunderstorm cloud on 18 June 2021 from 17:14 to 17:26 UTC. Vertical black lines indicate time instants 17:20:26, 17:21:31, 17:22:25, 17:22:57 and 17:23:47 UTC.




Figure 20. (a) Vertical air velocity and (b) Doppler spectrum width of the fourth thunderstorm cloud on 18 June 2021 from 17:14 to 17:26 UTC. Vertical black lines indicate time instants 17:20:26, 17:21:31, 17:22:25, 17:22:57 and 17:23:47 UTC.

535 5.2.1 Evidence Alignment of conical graupel particles

At 17:21:32 UTC, a lightning strike of 5 kA occurred around 8500 m away on the line of sight of the radar (strike number 7 in Fig. B4). This is a cloud-to-cloud lightning strike with medium strength. One second before that, negative sZ_{DR} values are observed for large and small particles from 8000 m to 9000 m as shown in Fig. 21(a). The minimum value is around -0.40 dB on the left side of the spectrum and -0.36 dB on the right side of the spectrum. Negative sZ_{DR} values disappeared at 17:21:38 UTC. Note that the timestamps of the cloud radar correspond to the end of the measurement after all chirp sequences have been transmitted, therefore the spectrum at 17:21:34 UTC may contain backscattered signals before the lightning, which could explain why negative sZ_{DR} is still observed. Since the location and time of negative sZ_{DR} agree well with that of the lightning strike and there are no other strikes close to this one in time and space, what is observed here is likely the vertical alignment and relaxation of particles right before and after a lightning strike.

545 The SL_{DR} across this lightning strike also shows interesting signature. As shown in Fig. 21(b), at 17:21:31 UTC, SL_{DR} from 8000 m to 9000 m suddenly decreases significantly and only recovered at 17:21:38 UTC. During this period, ρ_{hv} does not change significantly and is high (Fig. 21(c)), which means that the decrease in SL_{DR} is not due to low SNR. One possible cause is that almost all crystals are vertically aligned right before the lightning close to the location of lightning, which leads to low canting variance. As a result, there is a sudden decrease in SL_{DR} .




Figure 21. (a) Spectral differential reflectivity (b) slanted linear depolarisation ratio and (c) copolar correlation coefficient before and after lightning strike (5 kA) at 17:21:32 UTC on the line-of-sight of the radar between 8000 m and 9000 m (strike number 7 in Fig. B4).




Figure 22. Spectral differential reflectivity before and after a strong lightning strike (-18 kA) at 17:20:27 UTC at 5 km cross-range (strike number 92 in Fig. B5).

550 At 17:20:27 UTC, a strong cloud-to-cloud lightning discharge with a peak current of -18 kA occurred at a perpendicular distance of the range 3000 m (strike number 92 in Fig. B5), placing it at a distance of around 5500 m from the radar's line of sight. Despite being quite distant from the line of sight of the radar, negative sZ_{DR} values are observed for small particles from 5200 m to 5700 m before the lightning as shown in Fig. 22, which is probably due to the large magnitude of the electric field that generated the strong lightning. The minimum sZ_{DR} observed is around -0.36 dB, which is similar to that observed

555 in the previous case. Also similar to the previous case is that sZ_{DR} values returned to the level before the lightning about 4 seconds after the lightning from 17:20:31 UTC onward. However, unlike the previous case, negative sZ_{DR} is only found for small particles, which may be because electric field strength reduces with distance from the lightning strike, thus it is not strong enough to align larger and heavier particles vertically. It is difficult to pinpoint when negative sZ_{DR} first emerged due to this particular lightning strike. Slightly negative sZ_{DR} of about -0.16 dB can be found for light particles as early as 17:19:39

560 UTC, which could be due to other lightning strike in the same cloud. Although the same part of the spectrum shows positive sZ_{DR} instead one timestamp before at 17:19:09 UTC, a closer inspection reveals that it may be due to a different population of particles.

Comparing the spectral reflectivity at 17:19:09 and 17:19:39 UTC in Fig. 23 with the spectral differential reflectivity in Fig. 22, negative sZ_{DR} from 5000 m to 6000 m at 17:19:39 UTC likely belongs to particles associated with the upper part of the

565 spectrum, while positive sZ_{DR} at the same height at 17:19:09 UTC belongs to particles associated with the lower part of the spectrum. Since the two populations of particles may overlap, small negative sZ_{DR} values may be masked by positive values. Therefore, it is not possible to conclude that no negative sZ_{DR} values or no vertical alignment of particles are found before 17:19:39 UTC.

Also, unlike the previous case, right before the lightning at 17:20:27 UTC, SL_{DR} does not show a sudden decrease. This

570 could be because the lightning occurred some distance away from the line of sight of the radar. Therefore, not all particles are aligned, thus SL_{DR} did not decrease significantly.




Figure 23. Spectral reflectivity at 17:19:09 and 17:39:39 UTC.

In summary, the cloud-to-cloud lightning discharge within the radar's line-of-sight resulted in the vertical alignment of all ice particles within the radar resolution volume, whereas a discharge occurring cross-range led to the vertical alignment of only small ice particles. It should be noted that this effect may be influenced by the peak current magnitude of the lightning discharge.

From 17:23:40 UTC, sZ_{DR} becomes negative for the entire Doppler spectrum above 7000 m, such as the spectrum at 17:23:47 UTC shown in Fig. 27(b). This could be due to vertical alignment of all particles by strong cloud electric field. However, from Fig. 2, most of the thunderstorm cloud above 4000 m from 17:24 to 17:29 UTC was not visible to the radar due to large attenuation. There is also significant amount of liquid water below the cloud, leading to differential attenuation of horizontal and vertical polarizations, which may cause Z_{DR} values to be negatively biased. An evidence of differential attenuation is that Z_{DR} values become more negative as the thickness of the layer that contains liquid water with oblate particles increases. Also, many lightnings occurred close to each other in time during this period, so it is impossible to isolate each lightning strike and analyse the changes before and after each strike. These limit the investigation on the period with the most intense lightning activities.

5.2.2 Interesting microphysical properties

5.2.2.1 Evidence of conical graupel

According to Fig. 19(a), from around 17:22 UTC, a region with large negative differential reflectivity appears at around 4000 m to 6000 m. From Fig. 19(c) and (d), this region has enhanced slanted linear depolarisation ratio and reduced copolar correlation coefficient. Inspecting the spectograms during this period, it is found that from 17:21:24 UTC, a separate particle mode with negative sZ_{DR} is present on the left side of the Doppler spectrum at around 6000 m as shown in Fig. 21(a). The reflectivity

of this mode grew with time and it descended to around 4300 m near 17:24 UTC. The spectral reflectivity and sZ_{DR} of one instant where at a specific moment when this mode is present is shown in Fig. 24(a) and (b). When negative sZ_{DR} appears on the left part of the spectrum, the sZ_{DR} of on the right part of the spectrum is close to zero. Since small particles are The observed negative sZ_{DR} values on the left part of the spectrum may suggest the presence of conical graupel (Lu et al., 2016), as smaller particles, which are typically more easily aligned by an electric field and they are not, do not appear to be aligned in this case, the as indicated by the absence of slightly negative sZ_{DR} in the left part of the spectrum might be due to the presence of conical graupel (Lu et al., 2016) values.




Figure 24. Spectrograms of (a) equivalent reflectivity, (b) differential reflectivity, and profiles of (c) slanted linear depolarisation ratio, (d) copolar correlation coefficient at 17:22:25 UTC. Spectra at 5021 m indicated by black horizontal line in (a) and (b) are shown in Fig. 25(d-f-g-i).

The Figure 25(g-i) presents the Doppler spectra of reflectivity, Z_{DR} , and Ψ_{DP} at 5021 m at the instant shown for the time instant depicted in Fig. 24 are shown in Fig. 25(d-f). The spectral differential phase shift deviates from the Rayleigh plateau at about -2 m s^{-1} , signifying Mie non-Rayleigh scattering. To ensure correct interpretation of sZ_{DR} , scattering simulation is-simulations are carried out using typical parameters of conical graupel. From the literature, the theoretical axis ratio of conical graupel is 1.05 and the density, while measurements of mean axial ratios of conical graupels show values ranging from 1.1 to 1.3 for sizes in excess of 1 mm (Spek et al., 2008). The density of conical graupel is 0.55 g cm^{-3} (Spek et al., 2008), which is equivalent to an ice fraction of 0.6. The diameter of conical graupel, while the diameter is typically 2 to 8 mm (Pruppacher and Klett, 1980). The canting angle follows a Gaussian distribution with a zero mean and a standard deviation of 300.1° (Bringi and Chandrasekar, 2004). Conical shape is not supported by the simulation code used, thus the shape is assumed to be spheroidal. The simulated Since the T-matrix method can only simulate spheroidal but not conical particles, the simulation results are also compared to the results from the database created by Lu et al. (2016) for conical graupel with density 0.55 g

cm^{-3} and cone angles 40° and 50° . The cone angles were selected to match the trend of the observations. The reflectivity, Z_{DR} and δ_{co} obtained from the database are shown in Fig. 25(b-e). a-c), while those obtained from the T-matrix simulations are shown in Fig. 25(d-f).

The trends of differential reflectivity and differential backscatter phase obtained from the database are similar to those obtained by the T-matrix method. In Rayleigh scattering regime, the differential reflectivity of the simulated conical graupel is mostly negative. Both Z_{DR} and δ_{co} decrease when the Mie scattering regime is reached. δ_{co} reaches a minimum earlier than Z_{DR} . As particle size increases further, δ_{co} increases sharply and becomes positive, during which Z_{DR} reaches its minimum. Afterwards, δ_{co} reaches a local maximum and decreases slightly before increasing again. Z_{DR} increases and continues to oscillate. Similar behaviours are observed patterns are evident in the Doppler spectra observed at 5021 m at 17:22:25 UTC (Fig. 25(e-fg-i)). $s\Psi_{DP}$ reaches a minimum at -3.9 m s^{-1} and increases sharply as particle size further increases. sZ_{DR} reaches a minimum at -5.0 m s^{-1} while $s\Psi_{DP}$ is still increasing. Afterwards, $s\Psi_{DP}$ reaches a maximum and decreases slightly, while sZ_{DR} continues to increase. The Summarizing, the measurements of sZ_{DR} and $s\Psi_{DP}$ exhibit similar characteristics to both simulations, with $s\Psi_{DP}$ displaying a trough at smaller graupel sizes compared to sZ_{DR} .




Figure 25. Simulated reflectivity, differential reflectivity and differential backscatter phase of conical graupel by Lu et al. (2016) (a-c) and T-matrix method (d-f). (g-i) Spectral reflectivity, differential reflectivity and differential phase shift at 5021 m at 17:22:25 UTC. Note that the Doppler velocity decreases towards more negative values when the radius increases.

Although the results obtained from the database of Lu et al. (2016) and the T-matrix method show similar trends, the magnitudes of Z_{DR} are different, which suggests the limitation of T-matrix in simulating conical species. However, based on the similarity of the shapes of the curves, it is likely that the species observed has a shape between prolate spheroids simulated by the T-matrix method and conical graupel modelled by Lu et al. with cone angle of about 40-50°. It is also worth noting that the minimum of reflectivity in Fig. 25(dg) is not located at the Mie minimum according to the simulation (Fig. 25(ad)). Also the sZ_{DR} values on the small particles side is slightly positive. This suggests that the two peaks in spectral reflectivity represent two particle populations, the left peak corresponding to conical graupel and the right peak relating to nearly spherical smaller ice particles. Comparing the simulation results and the observed spectra, the conical graupel present is about 3 to 5.2 mm in diameter. This hypothesis is supported by a lower co-polar correlation coefficient.

~~(a-c) Simulated reflectivity, differential reflectivity and differential backscatter phase of conical graupel. (d-f) Spectral reflectivity, differential reflectivity and differential phase shift at 5021 m at 17:22:25 UTC. Note that the Doppler velocity decreases towards more negative values when the radius increases.~~

From 4400 m to 5600 m where the negative sZ_{DR} signature of graupel is the most prominent, SL_{DR} increases and ρ_{hv} decreases as shown in Fig. 24(c) and (d). This is likely because the radar resolution volume contains a variety of hydrometeors, including conical graupel and other small ice particles.

Unfortunately, it is challenging to look for supercooled liquid water in this case since there is liquid water at the bottom of the cloud below the 0 °C level at around 4000 m, which means it is impossible to identify supercooled liquid water using liquid water path. The presence of liquid water introduces an additional challenge, namely differential attenuation, which influences the sZ_{DR} values. While no direct measurements of the Rain Drop Size Distribution (RDSD) are available, a simulation can provide an estimate of the differential attenuation. For this purpose, the convective RDSD typical of the Netherlands based on disdrometer data Gatidis et al. (2024), is considered. The corresponding intercept parameter N_w equals $1300 \text{ mm}^{-1} \text{ m}^{-3}$ and the mass-weighted mean diameter D_m is 2.2 mm. The shape parameter, derived using the μ - λ relationship from the same study, along with the shape-size relationship used in Unal and van den Brule (2024), is applied. Consequently, in rainfall, the differential reflectivity is estimated at 0.15 dB, and the one-way differential attenuation at 0.06 dB km^{-1} . Except near the edges of the precipitation, Z_{DR} measurements show an increase from 0 dB to 0.2 dB as height decreases from 3000 m to 2200 m. Thus, the two-way differential attenuation contribution is expected to be low, at less than 0.12 dB, and does not significantly affect the interpretation of the results discussed.

It is worth noting from Fig. 24(a) and (b) that the population of graupel ends at around 4000 m. Since the radar is looking at an elevation angle of 45°, this suggests that graupel is not present closer than 4000 m from the radar, which means the region with graupel is localised in the thunderstorm cloud.

In Fig. 24(a) and (b), the spectrograms are plotted with vertical velocity instead of Doppler velocity as in other spectrograms in this article. The vertical velocity is estimated by assuming uniform horizontal wind predicted by the ECMWF model in the same hour. By plotting with vertical velocity, it is clear that the graupel are falling, while smaller ice particles on the right with positive vertical velocities are brought upwards by updrafts. As the falling graupel collide with the rising ice particles, charges can be produced. According to Takahashi (1978), if temperature is below -10 °C , graupel will become negatively

charged and vice versa. From the temperature profile measured by the microwave radiometer coupled to the cloud radar, the temperature is $-10\text{ }^{\circ}\text{C}$ at around 5550 m. This means that above 5550 m, falling graupel that collides with rising ice particles becomes negatively charged, forming a negative charge region in the cloud. Meanwhile, small ice particles that gained positive charges due to collisions are brought upwards by updrafts, so the upper part of the cloud is positively charged. Below 5550 m where temperature is above $-10\text{ }^{\circ}\text{C}$, falling graupel acquires positive charge, causing the cloud base to become positively charged. This could result in the typical tripolar structure of thunderstorm clouds. Nonetheless, the temperature profile inside the thunderstorm cloud may be different from the temperature profile measured by the microwave radiometer looking towards the zenith, so the actual charge distribution in the cloud may be different.

665 5.2.3 Alignment of particles

At 17:21:32 UTC, a lightning of 5 kA occurred around 8.5 km away on the line of sight of the radar (stroke number 7 in Fig. B4). This is a cloud-to-cloud lightning with medium strength. One second before that, negative sZ_{DR} values are observed for large and small particles from 8000 m to 9000 m as shown in Fig. 21(a). The minimum value is around -0.40 dB on the left side of the spectrum and -0.36 dB on the right side of the spectrum. Negative sZ_{DR} values disappeared at 17:21:38 UTC. Note that the timestamps of the cloud radar correspond to the end of the measurement after all chirp sequences have been transmitted, therefore the spectrum at 17:21:34 UTC may contain backscattered signals before the lightning, which could explain why negative sZ_{DR} is still observed. Since the location and time of negative sZ_{DR} agree well with that of the lightning stroke and there are no other strokes close to this one in time and space, what is observed here is likely the vertical alignment and relaxation of particles right before and after a lightning stroke.

675 The SL_{DR} across this lightning stroke also shows interesting signature. As shown in Fig. 21(b), at 17:21:31 UTC, SL_{DR} from 8000 m to 9000 m suddenly decreases significantly to -100 dB and only recovered at 17:21:38 UTC. During this period, ρ_{nv} does not change significantly and is high (Fig. 21(c)), which means that the decrease in SL_{DR} is not due to low SNR. One possible cause is that almost all crystals are vertically aligned right before the lightning close to the location of lightning, which leads to low canting variance. As a result, there is a sudden decrease in SL_{DR} .

680 At 17:20:27 UTC, a strong cloud-to-cloud lightning of -18 kA occurred at a perpendicular distance of around 3 km away from the radar (stroke number 92 in Fig. B5), but it is about 5.5 km away from the line of sight of the radar. Despite being quite distant from the line of sight of the radar, negative sZ_{DR} values are observed for small particles from 5200 m to 5700 m before the lightning as shown in Fig. 22, which is probably due to the large magnitude of the electric field that generated the strong lightning. The minimum sZ_{DR} observed is around -0.36 dB , which is similar to that observed in the previous case. Also similar to the previous case is that sZ_{DR} values returned to the level before the lightning about 4 seconds after the lightning from 17:20:31 UTC onward. However, unlike the previous case, negative sZ_{DR} is only found for small particles, which may be because electric field strength reduces with distance from the lightning, thus it is not strong enough to align larger and heavier particles vertically. It is difficult to pinpoint when negative sZ_{DR} first emerged due to this particular lightning stroke. Slightly negative sZ_{DR} of about -0.16 dB can be found for light particles as early as 17:19:39 UTC, which could be due to

690 other lightning in the same cloud. Although the same part of the spectrum shows positive sZ_{DR} instead one timestamp before at 17:19:09 UTC, a closer inspection reveals that it may be due to a different population of particles.

(a) Spectral differential reflectivity (b) slanted linear depolarisation ratio and (c) copolar correlation coefficient before and after lightning stroke (5 kA) at 17:21:32 UTC (stroke number 7 in Fig. B4).

695 Spectral differential reflectivity before and after a strong lightning stroke (−18 kA) at 17:20:27 UTC (stroke number 92 in Fig. B5).

Spectral reflectivity at 17:19:09 and 17:19:39 UTC.

700 Comparing the spectral reflectivity at 17:19:09 and 17:19:39 UTC in Fig. 23 with the spectral differential reflectivity in Fig. 22, negative sZ_{DR} from 5000 m to 6000 m at 17:19:39 UTC likely belongs to particles associated with the upper part of the spectrum, while positive sZ_{DR} at the same height at 17:19:09 UTC belongs to particles associated with the lower part of the spectrum. Since the two populations of particles may overlap, small negative sZ_{DR} values may be masked by positive values. Therefore, it is not possible to conclude that no negative sZ_{DR} values or no vertical alignment of particles are found before 17:19:39 UTC.

705 Also, unlike the previous case, right before the lightning at 17:20:27 UTC, SL_{DR} does not show a sudden decrease. This could be because the lightning occurred some distance away from the line of sight of the radar. Therefore, not all particles are aligned, thus SL_{DR} did not decrease significantly.

710 From 17:23:40 UTC, sZ_{DR} becomes negative for the entire Doppler spectrum above 7000 m, such as the spectrum at 17:23:47 UTC shown in Fig. 27(b). This could be due to vertical alignment of all particles by strong cloud electric field. However, from Fig. 2, most of the thunderstorm cloud above 4000 m from 17:24 to 17:29 UTC was not visible to the radar due to large attenuation. There is also significant amount of liquid water below the cloud, leading to differential attenuation of horizontal and vertical polarizations, which may cause Z_{DR} values to be negatively biased. An evidence of differential attenuation is that Z_{DR} values become more negative as the thickness of the layer that contains liquid water increases. Also, many lightnings occurred close to each other in time during this period, so it is impossible to isolate each lightning stroke and analyse the changes before and after each stroke. These limit the investigation on the period with the most intense lightning activities.

715 5.2.2.2 Strong updraft and turbulence

5.2.3 Strong updraft and turbulence

720 As shown in Fig. 20(a), from 17:18 to 17:24 UTC, vertical air velocity is large and positive (15-30 m s^{−1}) above 7000 m, indicating strong updraft in the cloud. From Fig. 20(b), the top of the cloud above 6000 m has large Doppler spectrum width of 3 to 4 m s^{−1}. In stratiform rain, the cloud top usually has low spectrum width since small and light particles have a small range of fall velocities. The large spectrum width observed here might be due to strong turbulence in the thunderstorm cloud. Slanted linear depolarisation ratio is high and copolar correlation coefficient is low in this region, which could be the result of large canting variance of particles due to strong turbulence.

From 17:22:30 to 17:24:00 UTC from 5000 m to 7000 m, there are three co-locating peaks of SL_{DR} and troughs of ρ_{hv} . The lowest peak at around 5000 m is located just above the graupel layer, such as in the example shown in Fig. 26 where the peak is found at 5060 m. From Fig. 26(h), the vertical air velocity does not vary much near this height, so the sudden increase in SL_{DR} and decrease in ρ_{hv} may not be due to increased canting variance due to turbulence. Meanwhile, the spectral Z_{DR} where the peak of SL_{DR} and ρ_{hv} is located shows multiple peaks (Fig. 26(d)). This could be due to a variety of hydrometeors with different axis ratios that are the seeds for forming conical graupel. Therefore, the high SL_{DR} and low ρ_{hv} in this case are likely due to co-existence of different types of particles.

730 The middle and highest peaks of SL_{DR} and troughs of ρ_{hv} are found at around 5900 m and 6400 m, such as in the example shown in Fig. 27. From Fig. 27(h), vertical air velocity changes sharply at these heights, which can produce strong turbulence. Therefore, the sudden increase in SL_{DR} and decrease in ρ_{hv} may be due to increased canting variance under turbulence. With strong turbulence, the Doppler spectra is no longer ordered with small particles on the right and large particles on the left because particles with different sizes are mixed.




Figure 26. 18 June 2021 17:22:57 UTC where the lowest peak of SL_{DR} and trough of ρ_{hv} is observed. (a-b) Spectrograms of reflectivity and differential reflectivity (c-e) spectra of reflectivity, Z_{DR} and Ψ_{DP} at 5060 m (f-i) profiles of SL_{DR} , ρ_{hv} , vertical air velocity and Doppler spectrum width.




Figure 27. 18 June 2021 17:23:47 UTC where the three peaks of SL_{DR} and troughs of ρ_{hv} are observed. (a-b) Spectrograms of reflectivity and differential reflectivity (c-e) spectra of reflectivity, Z_{DR} and Ψ_{DP} at 5877 m (middle peak) and 6422 m (highest peak) (f-i) profiles of SL_{DR} , ρ_{hv} , vertical air velocity and Doppler spectrum width.

735 5.2.3 Possibility of chains

5.2.2.3 Possibility of chains

From Fig. 19(a), high Z_{DR} is observed at the top of the cloud from 17:22 UTC onward. The Doppler spectra at 10003 m at 18 June 2021 17:22:57 UTC is shown in Fig. 28(d-fa-c). The differential reflectivity of the Rayleigh plateau (Doppler velocity >

5 m s⁻¹) is around 0.2 dB, and the entire ~~spectrum of Z_{DR} spectrum~~ is positive. One hypothesis is that the large particles with
740 positive sZ_{DR} are chain-like aggregates that formed ~~under cloud electric field. According to Connolly et al. (2005), chains~~
~~can be observed when the temperature is lower than -40°C . In the case being studied, temperature earlier under a strong~~
~~E-field. The lower copolar correlation coefficient in Fig. 26(g) suggests a mixture of particles (small ice crystals and chains),~~
~~but currently, there is no high E-field to vertically align the small particles. At that moment, the temperature above 9600 m~~
is lower than -40°C ~~above 9600 m, which backs up the hypothesis that chains might be present. Mie scattering, and it is~~
745 ~~indeed possible for chains to be present at such temperatures, according to Connolly et al. (2005). Resonance occurs at around~~
5 m s⁻¹ as sZ_{DR} increases and $s\Psi_{DP}$ begins to fluctuate. Scattering ~~simulation is carried out to estimate the simulations for~~
~~models of chain aggregates that estimate the differential reflectivity and differential backscatter phase of these particles as a~~
~~function of size could be useful in supporting the existence of chains as well as estimating the size of the particles based on the~~
~~trend of Mie scattering. In the simulation, chains are modelled as prolate particles. From the literature, chain-like aggregates~~
750 ~~can be up to several tens of particles long (Gayet et al., 2012) and individual crystals in chains in the cloud. However, this~~
~~species is not included in the scattering database of Lu et al. (2016). Meanwhile, the T-matrix method that models particles as~~
~~spheroids is not suitable for simulating this type of particles. From Fig. 1a, the length-to-width ratio of the chain aggregates is~~
~~around 2-3. On the other hand, individual monomers, which are usually plates, are very thin, so the length-to-thickness ratio of~~
~~chains are aligned with their maximum dimension along the length of the chain (Connolly et al., 2005). The individual particles~~
755 ~~can be plates, with typical axis ratio of less than 0.3 (Spek et al., 2008), or quasi-spherical frozen droplets with diameters of~~
~~15-20 μm (Gayet et al., 2012). Based on these, the axis ratio of chains (ratio of long axis to short axis) should be much larger~~
~~than 1. However, the simulation code used cannot converge when particle size increases when the axis ratio is too large, thus~~
~~the maximum possible axis ratio that can be used to allow Mie scattering regime to be reached is 7. Since chains are a type~~
~~of aggregates, ice fraction of typical aggregates is used, which is 0.3. The chains are assumed to be oriented with their long~~
760 ~~axis parallel to the horizontal, thus the Euler angle $\beta = 90^{\circ}$. The orientation follows a Gaussian distribution with a standard~~
~~deviation of 0.1° . is an order of magnitude larger than its length-to-width ratio. Therefore, chains are not well-represented by~~
~~a spheroidal model, hence not suitable to be simulated by the T-matrix method. Therefore, the existence of chain aggregates in~~
~~top levels of the thunderstorm cloud remains a hypothesis only.~~

(a-c) Simulated reflectivity, differential reflectivity and differential backscatter phase of chains with axis ratio 7 and ice fraction 0.3 oriented with long axis parallel to the horizontal. (d-f) Spectral reflectivity, differential reflectivity and differential phase shift at 10003 m at 17:22:57 UTC. Note that the Doppler velocity decreases when the maximum dimension increases.




Figure 28. (a) Spectral reflectivity, (b) differential reflectivity and (c) differential phase shift at 10003 m at 17:22:57 UTC. Note that the Doppler velocity decreases when the maximum dimension increases.

765 The simulated scattering behaviour of chains are shown in Fig. 28(a-c). The maximum dimension refers to the length of
the long axis of the spheroid. Similar to the observed spectral differential phase shift in Fig. 28(f), the simulated differential
backscatter phase Fig. 28(c) shows a peak followed by a trough, while spectral differential reflectivity shows a peak. This
occurs when scatterers have a maximum dimension of around 1.5 mm, which suggests that the observed particles have sizes
reaching around 1.5 mm. However, from the literature, chains collected in thunderstorm clouds have lengths of at most 0.8 mm
(Connolly et al., 2005). There are several reasons for the discrepancy between results based on simulation and the literature. For
770 instance, the axis ratio used in the simulation may not be large enough for chains, and a spheroid model may not be sufficient
to describe the shape of chains. According to Fig. 1, chains can be irregular in shape, and the effect of the irregularities could
increase as their sizes become more comparable to the radar wavelength.

Other than using scattering simulations, the presence of chain-like aggregates could also be supported by checking whether the region concerned fulfils the conditions described in the literature for the enhancement of aggregation by electric fields to occur, which are an electric field of 50-150 kV m⁻¹ and ice particle number concentration of at least 2 cm⁻³. Unfortunately, these data are not available in this study.

6 Conclusions

In this study, two major thunderstorm clouds on 18 June 2021 from 16:10 to 17:45 UTC near Cabauw were studied in depth to explore how bulk and spectral cloud radar data at 35 GHz with 45° elevation can be used to help understand processes in thunderstorm clouds. Prior to the analysis of the spectral differential reflectivity, sZ_{DR} , the spectral differential phase, $s\Psi_{DP}$, which indicates in which range of Doppler velocities Rayleigh and Mie-non-Rayleigh scattering occur, was investigated. This prevents the misinterpretation of sZ_{DR} extrema as characteristics of different ice particle populations in the Mie scattering regime. Instead of L_{DR} , SL_{DR} is available in the STSR mode. Compared to L_{DR} , SL_{DR} in this mode loses direct information about the mean canting angle due to the inability to acquire cross-polar measurements, but still retains information on the variance of canting angles and axis ratios.

Several types of hydrometeors are observed in clouds that produced lightning. In the first cloud, supercooled liquid water is found at the edge of the cloud at around 6000 m, which is supported by the increased liquid water path and near zero differential reflectivity of a separate mode of particles on the right of the Doppler spectra. In the fourth cloud, comparison between scattering simulations and observations supports the presence of conical graupel with maximum dimension of 3-5.2 mm graupel with prolate spheroidal to conical shape. The falling graupel coexist with ice particles that are brought upwards by updrafts, which could lead to non-inductive charging. The temperature at the corresponding heights could give rise to a tripolar structure of the thunderstorm cloud. In both clouds discussed above, there is a possibility that chain-like aggregates of small ice particles are present in the top of the cloud, which is reflected by large magnitudes of sZ_{DR} on the left side of the Doppler spectrum. Nonetheless, no realistic scattering simulation could be carried out to confirm the size and characteristics of the chains.

Vertical alignment of ice particles can be observed right before lightning up to 4 seconds before lightning and disappears within 3 seconds after the lightning reflected by negative sZ_{DR} values as low as -0.4 dB at 45° elevation. When the lightning is close to the line of sight of the radar, particles of all sizes are vertically aligned with sZ_{DR} values all negative. At this juncture point, the bulk variable SL_{DR} tends to 0 because of the vertical alignment of all the particles in decreases significantly due to the reduction in the canting angle variance of all particles within the radar resolution volume. It is worth noting that negative sZ_{DR} can be observed if the lightning is close to the values are observed when lightning occurs either near the radar's line of sight of the radar or is far or at a greater distance (up to 5.5 km) but strong, but if the lightning is strong. However, a sudden decrease in SL_{DR} can only be observed if is only observed when the lightning is close but not when it is far nearby and not at greater distances. When the lightning is far away, only small particles on the right side of the Doppler spectra are vertically aligned and exhibit negative sZ_{DR} values, while the bulk variable Z_{DR} has positive values. However,

there are also some situations where negative sZ_{DR} is observed that suggests vertical alignment of particles by the electric field, yet there are no lightning ~~strokes~~ strikes measured nearby in space and time. This could be because the electric field is not strong enough to trigger lightning, or that some lightning ~~strokes~~ strikes were not recorded.

810 Updrafts and downdrafts can be observed at different parts of the thunderstorm cloud. Near the edge of the first cloud, downdrafts can be observed. At the top and near the core of the fourth cloud, strong updrafts of up to 30 m s^{-1} can be observed. In general, vertical air velocity is not uniform in thunderstorm clouds, which suggests that there is strong turbulence. This is also supported by large Doppler spectrum width of up to $3\text{-}4 \text{ m s}^{-1}$. When strong turbulence is present, slanted linear depolarisation ratio increases and copolar correlation coefficient decreases, which suggest that canting angle variance of particles within a radar resolution volume increases.

815 In the case being studied, only measurements with constant elevation and azimuth and zenith observation were available, but their drawback is that only a small part of the thunderstorm cloud along the radar's line of sight could be measured, which leads to a low number of thunderstorm events recorded by the radar. In addition, it is not possible to look at the whole thunderstorm cloud at the same time to analyse the spatial variations within the cloud. Also, each part of the thunderstorm cloud only passes over the line of sight of the radar once, thus it is impossible to analyse the evolution of different parts of the cloud. A more
820 appropriate radar measurement mode for studying thunderstorm clouds would be azimuth scan ~~with constant elevation~~-(PPI) with the constant elevation of 45° . With PPI mode, thunderstorm clouds in all directions can be measured by the radar, so there can be more cases to choose from for in-depth study or statistical analysis. Moreover, it may become possible to analyse differences between different parts of the thunderstorm cloud with different levels of lightning activities, as well as how the cloud evolves with time.




Figure A1. Radar images and location of lightning strokes-strikes (yellow asterisks) from 18 June 2021 16:10 to 16:40 UTC (© Open-StreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Red The different classes of the colour bar relate to precipitation rates. The red triangle shows-indicates the radar location, while the red ruler shows-line represents the radar's line of sight of radar, with each mark equal-along the line corresponding to a distance of 1 km.




Figure A2. Radar images and location of lightning ~~strokes~~ strikes from 18 June 2021 16:45 to 17:10 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Legend same as Fig. A1.





Figure A3. Radar images and location of lightning strokes-strikes from 18 June 2021 17:15 to 17:40 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Legend same as Fig. A1.

B Lightning maps

18 June 2021 16:20 UTC




	UTC	kA		UTC	kA
1	16:18:33	-4	14	16:17:10	-6
2	16:19:01	-5	15	16:19:53	-3
3	16:18:12	-3	16	16:17:10	-7
4	16:18:33	-4	17	16:17:11	-6
5	16:18:12	-4	18	16:18:44	-5
6	16:18:12	-3	19	16:16:42	-5
7	16:18:12	-4	20	16:15:12	-4
8	16:18:12	-3	21	16:16:01	-15
9	16:17:41	-5	22	16:15:12	-3
10	16:19:00	-4	23	16:18:05	-3
11	16:19:29	-3	24	16:17:29	-4
12	16:17:10	-3	25	16:17:29	-4
13	16:17:30	-4	26	16:17:30	-5

1 to 3 kA Sissy mutterer
 3 to 7 kA Medium roller
 > 7 kA Strong slammer

Figure B1. Location, time and power of lightning strokes-strikes from 18 June 2021 16:15 to 16:20 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Red triangle shows radar location, red ruler shows line of sight of radar with each mark equal to 1 km.

18 June 2021 16:25 UTC



	UTC	kA		UTC	kA
1	16:23:50		10	16:23:07	-3
2	16:21:14	-13	11	16:20:17	2
3	16:20:17	5	12	16:21:51	2
4	16:20:40	-8	13	16:20:17	4
5	16:21:09	-3	14	16:20:17	-3
6	16:20:40	-4	15	16:22:20	8
7	16:21:50	-3	16	16:21:50	4
8	16:21:00	4	17	16:22:20	-5
9	16:21:50	-3	18	16:20:37	-5

Figure B2. Location, time and power of lightning strokes-strikes from 18 June 2021 16:20 to 16:25 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Legend same as Fig. B1. Cloud-to-ground lightning in bold.

18 June 2021 16:30 UTC




Figure B3. Location, time and power of lightning ~~strokes~~ strikes from 18 June 2021 16:25 to 16:30 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Legend same as Fig. B1.

18 June 2021 17:25 UTC





Figure B4. Location, time and power of lightning ~~strokes~~ strikes from 18 June 2021 17:20 to 17:25 UTC (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). See Fig. B5 for lightning in green rectangle. Legend same as Fig. B1.

18 June 2021 17:25 UTC



UTC	kA	UTC	kA
36	17:21:28 -3	55	17:21:58 3
37	17:21:28 -6	56	17:21:37 -6
38	17:24:41 -3	57	17:23:15 2
39	17:21:37 -3	58	17:21:37 -4
40	17:24:41 -2	59	17:21:48 -2
41	17:24:23 -4	60	17:23:07 -3
42	17:24:03 4	61	17:23:15 -3
43	17:20:18 -5	62	17:21:01 -6
44	17:24:18 -2	63	17:22:30 -3
45	17:22:58 4	64	17:23:30 -4
46	17:23:16 -5	65	17:23:18 -5
47	17:24:16 -4	66	17:23:30 -4
48	17:24:16 -9	67	17:23:43 -3
49	17:22:49 -4	68	17:24:08 -4
50	17:22:49 -4	69	17:20:56 -2
51	17:21:37 -4	70	17:24:47 -4
52	17:21:37 -2	71	17:21:15 3
53	17:23:07 -13	72	17:20:24 -4
54	17:23:07 -5	73	17:23:50 -8

UTC	kA	UTC	kA	UTC	kA	UTC	kA	UTC	kA	UTC	kA	UTC	kA
36	17:21:28 -3	55	17:21:58 3	74	17:24:35 -4	93	17:20:27 -3	112	17:21:29 -2	131	17:23:36 -5	150	17:22:18 -4
37	17:21:28 -6	56	17:21:37 -6	75	17:21:15 -3	94	17:20:10 -3	113	17:20:30 -3	132	17:21:02 -5	151	17:22:23 -7
38	17:24:41 -3	57	17:23:15 2	76	17:23:43 -2	95	17:23:55 -3	114	17:21:35 -2	133	17:23:25 -5	152	17:22:23 -4
39	17:21:37 -3	58	17:21:37 -4	77	17:21:03 -5	96	17:20:27 -3	115	17:21:17 -6	134	17:22:27 -12	153	17:20:37 -5
40	17:24:41 -2	59	17:21:48 -2	78	17:20:27 -3	97	17:21:49 -3	116	17:21:42 -3	135	17:21:49 -4	154	17:20:30 -3
41	17:24:23 -4	60	17:23:07 -3	79	17:21:05 -3	98	17:23:55 -2	117	17:21:52 -4	136	17:21:14 -3	155	17:22:23 -4
42	17:24:03 4	61	17:23:15 -3	80	17:20:01 -3	99	17:20:18 -6	118	17:22:04 -6	137	17:22:41 -2	156	17:20:06 -3
43	17:20:18 -5	62	17:21:01 -6	81	17:22:30 -7	100	17:24:35 -3	119	17:21:45 -6	138	17:22:01 -4	157	17:21:05 -1
44	17:24:18 -2	63	17:22:30 -3	82	17:21:15 -2	101	17:21:01 -2	120	17:22:04 -4	139	17:21:21 -2	158	17:20:42 -4
45	17:22:58 4	64	17:23:30 -4	83	17:22:20 -4	102	17:21:08 -3	121	17:22:04 -3	140	17:20:49 -4	159	17:20:47 -2
46	17:23:16 -5	65	17:23:18 -5	84	17:22:20 7	103	17:22:07 -2	122	17:21:02 -3	141	17:22:33 -4	160	17:22:20 -5
47	17:24:16 -4	66	17:23:30 -4	85	17:23:25 -6	104	17:21:33 -3	123	17:23:01 -4	142	17:21:11 6	161	17:21:27 -2
48	17:24:16 -9	67	17:23:43 -3	86	17:23:55 -4	105	17:20:37 -3	124	17:22:55 -2	143	17:20:04 -3	162	17:21:31 -2
49	17:22:49 -4	68	17:24:08 -4	87	17:22:20 -3	106	17:23:18 -2	125	17:21:01 -3	144	17:21:15 -3		
50	17:22:49 -4	69	17:20:56 -2	88	17:22:22 -3	107	17:20:25 -2	126	17:21:26 7	145	17:21:29 -4		
51	17:21:37 -4	70	17:24:47 -4	89	17:21:57 -3	108	17:20:30 -7	127	17:23:18 -7	146	17:20:11 -2		
52	17:21:37 -2	71	17:21:15 3	90	17:21:05 -7	109	17:21:08 -2	128	17:23:50 3	147	17:21:49 -8		
53	17:23:07 -13	72	17:20:24 -4	91	17:22:55 -4	110	17:20:30 -8	129	17:23:18 -6	148	17:21:49 -5		
54	17:23:07 -5	73	17:23:50 -8	92	17:20:27 -18	111	17:22:49 -3	130	17:23:25 -8	149	17:24:35 -4		

Figure B5. Location, time and power of lightning strokes-strikes in green rectangle in Fig. B4 (© OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)(Kachelmann GmbH). Legend same as Fig. B1.

Data availability. These cloud radar data are not published yet but are available in hourly NetCDF files from co-author Christine Unal (c.m.h.unal@tudelft.nl) upon request.

Author contributions. HYLM conducted the simulations, generated the plots, analyzed the data, and wrote the manuscript. CU supervised
830 HYLM, providing guidance on the methodology and data analysis, and contributed to the writing of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors would like to thank (Rob Mackenzie, Mahaut Sourzac) and (Saverio Guzzo, André Castro) for their support for maintaining the cloud radar operational and for data management, respectively. This work was supported by the Ruisdael Observatory, a scientific research infrastructure co-financed by the Netherlands Organisation for Scientific Research (NWO), under grant number
835 184.034.015.

References

- Aiswarya Lakshmi, K., Sahoo, S., Biswas, S., and Chandrasekar, V.: Study of Microphysical Signatures Based on Spectral Polarimetry during the RELAMPAGO Field Experiment in Argentina, *Journal of atmospheric and oceanic technology*, 41, 235–260, 2024.
- Bailey, M. P. and Hallett, J.: A comprehensive habit diagram for atmospheric ice crystals: Confirmation from the laboratory, AIRS II, and
840 other field studies, *Journal of the Atmospheric Sciences*, 66, 2888–2899, 2009.
- Bringi, V. N. and Chandrasekar, V.: *Polarimetric Doppler weather radar: principles and applications*, Cambridge university press, 2001.
- Brussaard, G.: A meteorological model for rain-induced cross polarization, *IEEE Transactions on Antennas and Propagation*, 24, 5–11, 1976.
- Chiu, C.-S. and Klett, J. D.: Convective electrification of clouds, *Journal of Geophysical Research*, 81, 1111–1124, 1976.
- Cho, H., Iribarne, J., and Richards, W.: On the orientation of ice crystals in a cumulonimbus cloud, *Journal of the Atmospheric Sciences*, 38,
845 1111–1114, 1981.
- Connolly, P., Saunders, C., Gallagher, M., Bower, K., Flynn, M., Choullarton, T., Whiteway, J., and Lawson, R.: Aircraft observations of the influence of electric fields on the aggregation of ice crystals, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131, 1695–1712, 2005.
- Deierling, W. and Petersen, W. A.: Total lightning activity as an indicator of updraft characteristics, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
850
- Doviak, R. J. and Zrníc, D. S.: *Doppler radar and weather observations*, Dover publications, 2006.
- Feist, M., Westbrook, C., Clark, P., Stein, T., Lean, H., and Stirlingnolly, A.: Statistics of convective cloud turbulence from a comprehensive turbulence retrieval method for radar observations, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 145, 727–744, 2019.
- 855 Gatidis, C., Schleiss, M., and Unal, C.: A new power-law model for μ - Λ relationships in convective and stratiform rainfall, *Atmospheric Measurement Techniques*, 17, 235–245, 2024.
- Gayet, J.-F., Mioche, G., Bugliaro, L., Protat, A., Minikin, A., Wirth, M., Dörnbrack, A., Shcherbakov, V., Mayer, B., Garnier, A., et al.: On the observation of unusual high concentration of small chain-like aggregate ice crystals and large ice water contents near the top of a deep convective cloud during the CIRCLE-2 experiment, *Atmospheric Chemistry and Physics*, 12, 727–744, 2012.
- 860 Gunn, R.: Electric-field regeneration in thunderstorms, *Journal of Atmospheric Sciences*, 11, 130–138, 1954.
- Kachelmann GmbH: Lightning and radar Utrecht, <https://meteologix.com/nl/stormradar/utrecht/20210618-1615z.html>, accessed: 2023-05-12.
- Lamb, D. and Verlinde, J.: *Physics and chemistry of clouds*, Cambridge University Press, 2011.
- Leinonen, J.: High-level interface to T-matrix scattering calculations: architecture, capabilities and limitations, *Optics express*, 22, 1655–
865 1660, 2014.
- Lu, Y., Jiang, Z., Aydin, K., Verlinde, J., Clothiaux, E. E., and Botta, G.: A polarimetric scattering database for non-spherical ice particles at microwave wavelengths, *Atmospheric Measurement Techniques*, 9, 5119–5134, 2016.
- Lund, N. R., MacGorman, D. R., Schuur, T. J., Biggerstaff, M. I., and Rust, W. D.: Relationships between lightning location and polarimetric radar signatures in a small mesoscale convective system, *Monthly Weather Review*, 137, 4151–4170, 2009.
- 870 Marshall, T. C. and Winn, W. P.: Measurements of charged precipitation in a New Mexico thunderstorm: Lower positive charge centers, *Journal of Geophysical Research: Oceans*, 87, 7141–7157, 1982.

- Marshall, T. C., Rust, W. D., and Stolzenburg, M.: Electrical structure and updraft speeds in thunderstorms over the southern Great Plains, *Journal of Geophysical Research: Atmospheres*, 100, 1001–1015, 1995.
- 875 Mattos, E. V., Machado, L. A. T., Williams, E. R., and Albrecht, R. I.: Polarimetric radar characteristics of storms with and without lightning activity, *Journal of Geophysical Research: Atmospheres*, 121, 14–201, 2016.
- Merceret, F. J., Ward, J. G., Mach, D. M., Bateman, M. G., and Dye, J. E.: On the magnitude of the electric field near thunderstorm-associated clouds, *Journal of Applied Meteorology and Climatology*, 47, 240–248, 2008.
- O’Connor, E.: Model data from Cabauw on 18 June 2021, <https://hdl.handle.net/21.12132/1.009248a1fe7a4c17>, 2022.
- 880 Oue, M., Kumjian, M. R., Lu, Y., Jiang, Z., Clothiaux, E. E., Verlinde, J., and Aydin, K.: X-band polarimetric and Ka-band Doppler spectral radar observations of a graupel-producing Arctic mixed-phase cloud, *Journal of Applied Meteorology and Climatology*, 54, 1335–1351, 2015.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, Springer, 1980.
- Reynolds, S. E., Brook, M., and Gourley, M. F.: Thunderstorm charge separation, *Journal of Atmospheric Sciences*, 14, 426–436, 1957.
- 885 Sokol, Z., Minářová, J., and Fišer, O.: Hydrometeor distribution and linear depolarization ratio in thunderstorms, *Remote Sensing*, 12, 2144, 2020.
- Spek, A., Unal, C., Moisseev, D., Russchenberg, H., Chandrasekar, V., and Dufournet, Y.: A new technique to categorize and retrieve the microphysical properties of ice particles above the melting layer using radar dual-polarization spectral analysis, *Journal of atmospheric and oceanic technology*, 25, 482–497, 2008.
- 890 Stith, J. L., Dye, J. E., Bansemer, A., Heymsfield, A. J., Grainger, C. A., Petersen, W. A., and Cifelli, R.: Microphysical observations of tropical clouds, *Journal of Applied Meteorology and Climatology*, 41, 97–117, 2002.
- Stith, J. L., Basarab, B., Rutledge, S. A., and Weinheimer, A.: Anvil microphysical signatures associated with lightning-produced NO_x, *Atmospheric Chemistry and Physics*, 16, 2243–2254, 2016.
- Takahashi, T.: Riming electrification as a charge generation mechanism in thunderstorms, *Journal of Atmospheric Sciences*, 35, 1536–1548, 1978.
- 895 Tang, G., Yang, P., Stegmann, P. G., Panetta, R. L., Tsang, L., and Johnson, B.: Effect of particle shape, density, and inhomogeneity on the microwave optical properties of graupel and hailstones, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 6366–6378, 2017.
- Unal, C. and van den Brule, Y.: Exploring Millimeter-Wavelength Radar Capabilities for Raindrop Size Distribution Retrieval: Estimating Mass-Weighted Mean Diameter from the Differential Backscatter Phase, *Journal of atmospheric and oceanic technology*, 41, 583–603, 2024.
- 900 Vonnegut, B.: Possible mechanism for the formation of thunderstorm electricity, *Geophys. Res. Pap.*, 42, 169–181, 1955.
- Vonnegut, B.: Orientation of ice crystals in the electric field of a thunderstorm, *Weather*, 20, 310–312, 1965.
- Wahab, N. M. A.: *Ice crystal interactions in electric fields*, Ph.D. thesis, UMIST, 1974.
- Wang, P. K.: *Physics and dynamics of clouds and precipitation*, Cambridge University Press, 2013.
- 905 Wang, Y., Yu, T.-Y., Ryzhkov, A. V., and Kumjian, M. R.: Application of spectral polarimetry to a hailstorm at low elevation angle, *Journal of Atmospheric and Oceanic Technology*, 36, 567–583, 2019.
- Warren, S. G. and Brandt, R. E.: Optical constants of ice from the ultraviolet to the microwave: A revised compilation, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Waterman, P.: Matrix formulation of electromagnetic scattering, *Proceedings of the IEEE*, 53, 805–812, <https://doi.org/10.1109/PROC.1965.4058>, 1965.

- 910 Weinheimer, A. J. and Few, A. A.: The electric field alignment of ice particles in thunderstorms, *Journal of Geophysical Research: Atmospheres*, 92, 14 833–14 844, 1987.
- Wilson, C. et al.: Some thundercloud problems, *Journal of the Franklin Institute*, 208, 1–12, 1929.
- Yu, T.-Y., Xiao, X., and Wang, Y.: Statistical quality of spectral polarimetric variables for weather radar, *Journal of Atmospheric and Oceanic Technology*, 29, 1221–1235, 2012.
- 915 Zipser, E. J. and Lutz, K. R.: The vertical profile of radar reflectivity of convective cells: A strong indicator of storm intensity and lightning probability?, *Monthly Weather Review*, 122, 1751–1759, 1994.