



Quantifying radiative effects of light-absorbing particles deposition on snow at the SnowMIP sites

Enrico Zorzetto^{1,2}, Paul Ginoux³, Sergey Malyshev³, and Elena Shevliakova³

¹Earth and Environmental Science Department, New Mexico Institute of Mining and Technology, Socorro, NM, USA

²Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ, USA

³NOAA OAR Geophysical Fluid Dynamics Laboratory, Princeton, NJ, USA.

Correspondence: Enrico Zorzetto (enrico.zorzetto@nmt.edu)

Abstract. The deposition of light-absorbing particles (LAPs) leads to a decrease of surface albedo over snow covered surfaces. This effect, by increasing the energy absorbed by the snowpack, enhances snow melt and accelerates snow aging, process which in turn is responsible for further decreasing the snow albedo. Capturing this combined process is important in land surface modelling, as the change in surface reflectivity connected with the deposition of LAPs can modulate time and magnitude of snowmelt and runoff. These processes impact regional water resources, and can also lead to relevant feedbacks to the global climate system. We have recently developed a new numerical snowpack model for the GFDL land model (A Global Land Snow Scheme, or GLASS). GLASS provides a detailed description of snow mass and energy balance, as well as the evolution of snow microphysical properties (grain shape and size). We now extend this model to account for the presence of light-absorbing impurities, modelling their dry and wet deposition in the snowpack, the evolution of their vertical distribution in the snow due to precipitation and snow melt, and the effect of their concentration on snow optical properties. To test the effects of the resulting snow scheme, we force the GFDL land model with deposition of black carbon, mineral dust and organic carbon obtained from a general circulation model (GFDL AM4.0). We evaluate the new model configuration at a set of instrumented sites, including an alpine site (Col de Porte, France) where in-situ observations of snow (including spectral measurements of snow reflectivity and concentration of LAPs) allow for a comprehensive model evaluation. For the Col de Porte site, we show that GLASS reproduces the observed magnitudes of impurities concentration in the snowpack throughout a winter season. The seasonal evolution of the snow optical diameter is also qualitatively reproduced by the model, although the increase in snow grain diameter during the melt season appears to be underestimated. For a set of instrumented sites spanning a range of climates and LAP deposition rates (the ‘SnowMIP’ sites) we then evaluate the number of snow-days lost due to the deposition of dust and carbonaceous aerosols. We find that this loss ranges between 5 and 24 days depending on the site. The resulting snow model with LAP-aware snow reflectivity show a good agreement with measurements of broadband albedo and seasonal SWE over the study sites.



1 Introduction

The deposition of light absorbing particles (LAPs) on snow is known to reduce its surface reflectivity, in turn accelerating snow melt and snow aging through enhanced metamorphism (Warren, 1984; Doherty et al., 2010; Dumont et al., 2014; Hadley and Kirchstetter, 2012; Skiles et al., 2018; Sarangi et al., 2020). The impact of LAPs has been shown to lead to relevant impacts in the water cycle over extended regions of the world, such as the western US (Painter et al., 2012), the Alps (Di Mauro et al., 2015) and Pyrenees (Réveillet et al., 2022) and high mountain Asia (Ackroyd et al., 2021; He et al., 2018a), among other regions. The effects of the concentration of LAPs on snow optical properties is very nonlinear: While the deposition of LAPs directly affects snow albedo in the visible range, it additionally produces indirect effects. As the energy absorbed by snow increases, the evolution of snow grain size and shape with snow aging accelerates, driven both by thermal gradients and wet processes (Hadley and Kirchstetter, 2012). In turn, these metamorphic changes on snow grain properties modulate snow albedo, not only in the visible range but also in the near-infrared part of the spectrum (Skiles et al., 2018), and can enhance the surface albedo feedback, especially during spring (Huang et al., 2022). Accounting for LAP deposition and its effects on snow albedo is thus important in numerical snow schemes which are routinely used in hydrological and land surface models, as these processes impact the spatial variability of snow cover and can lead to long-term changes in snow at the ground. Furthermore, these changes in snow cover exert control both on surface and sub-surface hydrology (through changes in snow melt timing) and on land surface-atmosphere interactions, through changes in land surface temperature. Accurate snow predictions are thus of paramount importance for Earth System Models, especially as snow water equivalent and snow extent have both been shown to decline both in historical observations (Estilow et al., 2015; Kunkel et al., 2016) and in climate projections (Mudryk et al., 2020).

Multiple LAP species can affect snow albedo. Common species of LAPs found in snow include mineral dust (Painter et al., 2007; Sarangi et al., 2020), black carbon (Flanner et al., 2007; Réveillet et al., 2022; Flanner et al., 2012), and organic carbon (resulting from both natural and anthropogenic combustion processes), volcanic ash, and other biological elements such as algae (Cook et al., 2017). LAPs can be added to snow by wet deposition (i.e., LAPs contained in liquid or frozen precipitation) and by dry deposition, whereby LAPs are deposited on snow by gravitational settling or turbulent/laminar exchange with the atmosphere. Once deposited, the concentration of LAPs in a snow layer evolves as a result of snow melt and sublimation (Conway et al., 1996). In the case of sublimation, the concentration of LAPs near the snowpack surface increases due to the net loss of ice to the atmosphere. On the other hand, snow melt can remove LAPs from the snow through a phenomenon referred to as *scavenging*, depending on whether the particles are hydrophilic or hydrophobic. It has been observed (Sterle et al., 2013) that black carbon and dust tend to be retained in the snowpack even during the ablation season, leading to increased LAP concentrations in the uppermost layers following snow melt.

The most abundant LAP by mass is mineral dust, which generally originates from deserts and other poorly vegetated and dry regions due to wind-driven emission. Light absorptive properties of mineral dust can vary greatly, and are primarily controlled by its iron content. Black carbon has the largest absorption in the visible wavelength compared to organic carbon and other common LAPs.



In recent years, the growing recognition of i) the importance of LAPs in modulating snow processes, and ii) the interaction of these effects with the climate system and land hydrology has led to multiple modelling efforts focused on LAPs. Radiative transfer numerical models have been developed to represent the effect of LAPs in snow (Wiscombe and Warren, 1980; Liou et al., 2014; Flanner et al., 2007; Aoki et al., 2011; Libois et al., 2013; He et al., 2019), including simplified parameterizations tailored to global circulation models (Dang et al., 2015; He et al., 2018b), in which land surface schemes routinely employ a coarse 2-band representation of the solar spectrum. He et al. (2018a) and He et al. (2019) carried out an extensive evaluation of the radiative effects of LAPs in snow, focusing on the role of internal vs. external mixing state, and quantified the magnitude of their effects for snow grains of different shapes. However, all radiative transfer models rely on the knowledge of LAP concentrations and snow properties (e.g., snow optical diameter and grain shape) as well as the distribution of these properties within the snowpack. This presents a challenge, as most snow schemes employed in global Earth System models have a simplified representation of snowpack properties and their vertical distribution.

Detailed snow models accounting for the LAP deposition process have been developed. For instance, Tuzet et al. (2017, 2020) extended the high-detail snow model CROCUS (Vionnet et al., 2012) to include LAP processes, and used to evaluate the effect of LAPs over instrumented sites in the French Alps. Similarly, (Skiles and Painter, 2019) employed the radiative transfer model SNICAR (Flanner and Zender, 2005) to quantify LAPs effect in the snow model SNOWPACK (Lehning et al., 2002). However, generally highly detailed snow models are used in local or regional studies due to their complexity and computational cost. In comparison to these local studies, several Earth System Models still employ simple representation of snow processes such as LAP deposition, snow metamorphism, and a coarse vertical representation of the snowpack which does not allow to adequately characterize the vertical heterogeneity of the snowpack. These large differences in complexity and detail of snow schemes used in different ESMs have led to large inter-model differences being observed in previous model intercomparison efforts (Nijssen et al., 2003; Krinner et al., 2018; Menard et al., 2021) and are thought to contribute to the large spread still observed between model estimates of the surface albedo feedback (Flanner et al., 2011; Qu and Hall, 2014).

These persisting issues concerning snow predictions in have led to an increased attention towards the representation of LAP-on-snow processes both regional and global modelling efforts (Qian et al., 2015). The effect of LAPs has for example been implemented in the Community Land Model (CLM) using the SNICAR (Snow, Ice, and Aerosol Radiation Model) developed by Flanner and Zender (2005); Flanner et al. (2009), finding that the presence of LAPs leads to increased surface temperature and to relevant feedbacks in a global circulation model. More recently, a SNICAR-based parameterization for LAPs effect on snow optical properties was implemented in the DOE E3SM model (Golaz et al., 2022), and used to test the representation of snow processes over the Tibetan plateau (Hao et al., 2022) and over the Western United States (Hao et al., 2023).

Despite the recent progress, the global distribution of LAPs effects on snow remains characterized by large uncertainties, which include i) the magnitude of this effect, ii) the relative contributions of different LAP species, and iii) the interactions of LAPs-driven melt with snow metamorphic processes. To reduce this uncertainty, here we analyze the effect of LAPs at instrumented sites using a recent detailed snow scheme (Global Land Snow Scheme, GLASS) developed for Earth System Model simulations. In a companion manuscript (Zorzetto et al., 2024), we have presented this new snow scheme and discussed its implementation in the GFDL Earth System Model. GLASS includes a refined vertical structure of the snowpack and an



explicit description of snow metamorphism. GLASS is based on an implicit numerical solution of mass and energy balance for the snowpack, so that the model can be effectively employed in global-scale simulations of the land-atmosphere coupled system. In this manuscript, we extend GLASS to include the effect of light absorbing impurities, including their wet and dry
95 deposition, their mass balance within the snowpack, and their effects on snow optical properties, accounting for predicted snow microphysical structure (snow grain shape and optical diameter).

Harnessing this new model, here we focus on responding to two questions: How do LAPs affect snow melt in the spring? and, is the modelled snowpack in agreement with observations (bulk snow properties, grain properties, and LAP concentration near the snow surface?). While GLASS is designed for global applications, we focus our analysis on 10 "SnowMIP" sites (Ménard
100 et al., 2019), as i) all these sites include high-quality forcing data and validation data, which allow to reduce uncertainty compared to large-scale studies and are thus invaluable to evaluate model performance, and ii) offer a perfect opportunity to quantify LAP-driven snow melt for a set of site spanning a wide range of climate and terrain conditions, which can be used to further constrain large scale studies. Furthermore, this forcing and validation dataset was used in a recent SnowMIP ESM comparison effort (Krinner et al., 2018; Menard et al., 2021).

The paper is organized as follows: We first provide an overview of the GFDL land model and of the GLASS snow scheme. We then describe the new treatment of LAPs mass balance within the snowpack, and the revised snow albedo model used to capture LAP-driven snow darkening processes. The experimental setup is then discussed, along with the datasets used for forcing the model and for validation. A particular focus of the discussion is the Col de Porte site (France), where both
105 measurements of snow bulk properties and spectral measurements were carried out for the 2013-2014 snow season which allow us to evaluate model predictions of snow optical diameter and of LAP concentration at the snowpack surface. We then discuss the implications of our findings for land surface simulations over continental domains and in coupled land-atmosphere simulations.

2 Model Description

2.1 GFDL land model

115 The land model LM4.1 is the land component of the GFDL ESM4.1 (Dunne et al., 2020). The description of water and energy balance at the land surface and in subsurface soil layers is based on the Land Dynamics scheme (Milly et al., 2014). The land domain is modelled in a grid of cells composed of sub-units, termed tiles, which represent homogeneous areas of either soil, lake or glacier. In the present work we employ a "point model" version of LM4.1, whereby we assume that a single land model tile represent the sol-snow-atmosphere continuum at each site. Here we use the same physics time step routinely used
120 in global-scale simulations (30 minutes). Soil is modelled as a multi-layer medium coupled with snow and atmosphere above, with full representation of mass balance of liquid and frozen water, and vertical diffusion of heat. Exchanges of water and energy between land and atmosphere are computed according to the *Monin-Obukhov* similarity theory (Garratt, 1994).



2.2 Snow model

The physical description of snow processes is based on a snow scheme recently developed for LM4.1 (the Global LAnd Surface
125 Scheme, or GLASS, described by Zorzetto et al. (2024)). In GLASS, the snowpack is modelled as a 1D multi-layer medium.
Each layer k ($k = 1, \dots, n_L$ from the top), is characterized by ice ($w_{s,k}$) and liquid ($w_{l,k}$) contents (in kg m^{-2}), temperature,
density (i.e., layer thickness Δz_k), snow age, and a set of variables describing the snow microphysical structure, which evolve
based on dry and wet metamorphic processes, snow compaction, and wind drift effects. These variables are snow optical
diameter ($d_{opt,k}$), snow grain sphericity ($s_{p,k}$), and grain dendricity (δ_k). Snow metamorphism caused by dry processes driven
130 by temperature gradients is described according to (Flanner and Zender, 2006), while wet processes are modelled following
the parameterization by (Brun et al., 1992).

The evolution of snow grain shape is described based on the prognostic equations for snow grain dendricity and sphericity.
These are computed following the parameterization used in CROCUS (Brun et al., 1992; Vionnet et al., 2012; Carmagnola
et al., 2014). In GLASS, all three microphysical properties ($d_{opt,k}$, $s_{p,k}$, and δ_k) are prognostic variables. Wind drift and snow
135 compaction are also accounted for in the model following the approach by (Vionnet et al., 2012) and contribute to the evolution
of snow density and grain size size and shape. The vertical structure of the snowpack consists of a dynamic number n_L of snow
layers. New layers are created on top of the existing snowpack following snowfall events of large enough magnitude, so that
the vertical layering structure preserves snow physical properties in each layer. The vertical layers are also updated based on
computational considerations. At each time step, the snow vertical structure is compared to an optimal vertical discretization
140 defined for each given snow depth. If the layers are too coarse or too thin for a given snow depth, the layers undergo splitting or
merging. These operations are designed in order to strike a trade-off between computational cost and vertical detail, to satisfy
requirements of numerical efficiency (to avoid too large number of layers) and to ensure a proper description of the snowpack
vertical structure (too coarse a vertical discretization would hinder the representation of some physical processes, such as the
vertical heat diffusion). If two snow layers are characterized by values of density, optical diameter or impurities content which
145 are too different, merging of the two layers is not permitted in order to preserve the vertical heterogeneity of the snowpack. For
a more detailed description, see Zorzetto et al. (2024).

2.3 Tracer deposition in the snowpack

In this work GLASS was updated to include the mass balance of multiple LAP species. In addition to the set of variables
mentioned in Section 2.2, in this revised version each snow layer k is further characterized by the mass of each LAP specie
150 both internally mixed (IM) and externally mixed (EM) within the snow. These quantities ($w_{IM,i,k}$ and $w_{EM,i,k}$) in (kg m^{-2})
are tracked for each tracer specie i , so that in our current application we have 6 types of LAP in each layer (IM and EM, for
each specie: BC, OC and MD). A conceptual representation of the new processes included in GLASS are illustrated in Figure
1.

The dry flux $D_{c,i}$ (in $\text{mg/m}^2/\text{s}$) for each tracer specie ($i=\text{BC, MD, OC}$) is deposited in the uppermost snow layer. While in
155 general wind pumping can redistribute dry deposited LAPs within the snow, this process is limited to thicknesses smaller than



10 mm so that here we decided to assign all deposition to the uppermost snow layer (Clifton et al., 2008; Tuzet et al., 2017). At each model time step, we assume dry deposition is equal to the monthly average value from the forcing dataset.

The wet deposition flux in our forcing dataset was provided as monthly average. Since we force our snow model with in-situ observations, we adopt the following strategy to estimate wet deposition fluxes for each snowfall or rainfall event: We first
160 compute the monthly average concentration of each LAP specie in the precipitation (as the mass ratio of monthly average wet deposition to monthly precipitation, in [ppm]). We then assign in each model time step the total amount of wet-deposited LAPs as proportional to the rainfall and/or snowfall rate for that time step, so that the flux of tracer i due to liquid and solid precipitation is respectively $c_{l,i}f_l$ and $c_{s,i}f_s$. For the purpose of this study, we assume that rainfall and snowfall carry the same concentration of LAPs, and neglect any possible dependence of deposition fluxes e.g., on precipitation intensity.

165 In the case of large enough snowfall, the snow model creates a set of new snow layers on top of the existing snowpack. The newly deposited LAP mass is stored in these layers. In case of small snowfall events and snow already present at the ground, instead of creating new snow layers the model adds the fresh snow to the existing upper layer of the snowpack. In that case, the deposited LAPs are also added to the existing mass of each species in that layer.

2.4 LAPs in the snow model

170 The land model solves the energy balance at the ground using an implicit time stepping scheme. From this step, the snow temperature profile is updated and an estimate of the mass available for melt or freeze within the snowpack is computed. Sublimation leads to a thinning of the uppermost snow layer, or to its complete disappearance and to mass being removed from the underlying layers. In both cases, any existing mass of LAPs present in these layers is conserved. Thus, in general sublimation leads to an increase in concentration of LAPs at the top of the snowpack. If the entire snowpack disappears due
175 to sublimation, any existing mass of LAPs is lost. Sublimation does not lead to changes in the status of LAPs (internally vs externally mixed). Then, melt and freeze are applied to each layer of the snowpack, if the thermodynamic conditions and availability of liquid water or ice require it.

The mass of each LAP species is conserved by snowpack relayering operations. If a snow layer is split in two, the amount of each LAP species present in the layer is assigned to the new layers proportionally to their snow water masses. Similarly, in the
180 event of two snow layers being merged, the LAP mass in the newly formed layer will simply be the sum of that in the original layers, for each LAP species and LAP mixing state. When a snow layer disappears as a consequence of melt or sublimation, the entire LAP content of the layer accumulates in the layer immediately underneath, if any, or else is lost from the snowpack.

2.5 Snow melt

When a snow layer completely melts, the mass of LAPs present in that layer is deposited to the layer underneath. When there
185 is liquid water flow between a layer and the one underneath, a part of the LAPs present in the upper layer is scavenged by the water flow. The fraction of LAPs removed is based on a scavenging coefficient. Any LAPs carried by water flowing from snowpack to the substrate (glacier, lake or soil) are lost. For snow layer k and LAP species i the vertical flux (between layers



k and the underlying $k + 1$) of LAPs is given by:

$$M_{scav,i,k} = q_{L,i,k} C_{scav,i} \frac{w_{c,i,k}}{SWE_k} \quad (1)$$

190 As pointed out by Sterle et al. (2013), the magnitude of scavenging for dust and BC is highly uncertain, and appears to be low due to the tendency of these LAPs to remain in the snowpack throughout the ablation processes. Similar to previous studies (Tuzet et al., 2017; Ga Chan et al., 2022), here we set the scavenging coefficient to 0 for mineral dust (with particles generally too large for scavenging to occur) and organic carbon, and to 0.2 for black carbon, as suggested by (Flanner et al., 2007).

2.6 Snow albedo parameterization

195 In this work, the snow surface albedo is computed based on snow properties (optical diameter and shape) and on the concentration of LAPs near the snowpack surface. In this section, snow properties are averaged over a near surface layer of thickness set equal to up to 3 cm. Based on the work of Dang et al. (2015) and He et al. (2018a), snow albedo is expressed as a function of snow grain effective radius

$$\alpha_s = b_0 (\delta_{ns}, s_{p,ns}) + b_1 (\delta_{ns}, s_{p,ns}) R_n + b_2 (\delta_{ns}, s_{p,ns}) R_n^2 \quad (2)$$

200 with

$$R_n = \log_{10} \left(\frac{R_e}{R_0} \right) \quad (3)$$

here R_e represents the snow effective radius, defined as $R_e = 3V_s/(4A_s)$, and corresponds to $d_{opt,ns}/2$. The reference radius is $R_0 = 100 \mu\text{m}$. Here the optical diameter $d_{opt,ns}$ as well as the snow grain shape parameters δ_{ns} and $s_{p,ns}$ are averaged over the near surface layer of the snowpack, with thickness up to the upper 3 cm. For direct radiation, we account for the direction
205 of the incident radiation beam by computing an effective snow grain radius as proposed by Marshall (1989):

$$R'_e = R_e (1 + a_b \Delta\mu) \quad (4)$$

where $a_b = 0.781$ for visible band ($b = VIS$) and $a_b = 0.791$ for near infrared radiation ($b = NIR$). In the case of diffuse radiation, $\mu_D = 0.65$ ($\theta = 49.5^\circ$). For thick enough snowpack ($d > 0.02$ m), solar radiation penetrates within the snowpack and absorbed radiation is distributed exponentially

$$210 \quad Q_s(z) = \sum_{b=1}^2 (1 - \alpha_b) R_{s,b} e^{-\beta_b z} \quad (5)$$

where for each band b $R_{s,b}$ is the downward shortwave radiation at the surface, and β_b describes the penetration of light within the snowpack. The extinction coefficients for visible and near infrared light are estimated as in Jordan (1991) and Shrestha et al. (2010): $\beta_{NIR} = 400$, $\beta_{VIS} = 0.003759 \rho d_{opt}^{-0.5}$, with density and optical diameter averaged over the near-surface layer of the snowpack, up to a maximum depth of 3 cm.



215 2.7 Effect of LAPs on predicted albedo

The effect of LAPs is accounted for using the parameterization by He et al. (2018a), in which the albedo reduction is obtained as

$$\Delta\alpha_s = d_{0,b}(\delta_{ns}, s_{p,ns})(C_{eq,bc})^k \quad (6)$$

with

$$220 \quad k = d_{1,b}(\delta_{ns}, s_{p,ns}) \left(\frac{R_e}{R_0} \right)^{d_2} \quad (7)$$

Here the parameters $d_{0,b}(\delta, s_p)$ and $d_{1,b}(\delta, s_p)$ are given by He et al. (2018b) as a function of spectral band (visible and near-infrared bands are used here) and as a function of snow grain shape (sphere, spheroid, hexagon or Koch snowflake). Here we account for snow grain shape using the prognostic equation describing snow dendricity and sphericity. If the near surface layer is characterized by dendritic snow ($\delta_{ns} > 0.5$), optical parameters relative to Koch snowflakes are used. If this is not the case, non-dendritic snow is classified based on its sphericity, so that it can be idealized as a collections of spherical ($s_{p,ns} > 0.8$), spheroidals ($0.8 > s_{p,ns} > 0.2$) or hexagonal crystals ($s_{p,ns} < 0.2$).

To account for the radiative effects of multiple tracer species, we compute a BC-equivalent concentration as done by Ga Chan et al. (2022):

$$c_{eq,BC} = c_{BC} + c_{MD} \frac{\sigma_{abs,MD}}{\sigma_{abs,BC}} + c_{OC} \frac{\sigma_{abs,OC}}{\sigma_{abs,BC}} \quad (8)$$

230 Where $\sigma_{abs,t}$ is the absorption cross section of tracer t . These are set to $7330 \text{ m}^2\text{kg}^{-1}$ for BC, $67.8 \text{ m}^2\text{kg}^{-1}$ for MD, and $122 \text{ m}^2\text{kg}^{-1}$ for OC, values used by Ga Chan et al. (2022). All these quantities are averaged over the near-surface layer of the snowpack.

3 Experimental setup

3.1 Forcing and Validation Data

235 In this study, we use the land model driven by prescribed high-frequency meteorological data, including short-wave and long-wave downward radiation, rainfall and snowfall rates, air temperature, pressure, specific humidity, and wind speed. We first run a 100-year model spinup cycling through forcing data from the Global Soil Wetness Project Phase 3 (GSWP3) (1980-1990) with site-specific correction described in (Ménard et al., 2019). Then, we run the historical 1980-1996 years again using GSWP3 corrected data. Finally, we run the model for the entire duration of the in-situ meteorological observations available at each site (e.g., 1996-2014 for the Col de Porte site). As done in the companion manuscript, we have evaluated the memory of the soil temperature and ice content and found that equilibrium is reached in about 30 model years.

The dataset used for testing model performance at Col de Porte is described in Dumont et al. (2017); Morin et al. (2012); Lejeune et al. (2019) and was previously used to evaluate the effect of LAPs in snow schemes Tuzet et al. (2017); Ga Chan



et al. (2022). The data collected at this site is extensive and include long series of snow depth, runoff and temperature, as well
245 as daily average snow broadband albedo.

Additionally, spectral measurements were carried out in the snow year 2013-2014 at the Col de Porte site (Dumont et al.,
2017), which were then used to estimate snow specific surface area (SSA) and concentration of LAPs. In this study, we compare
daily averages of the model output with the daily averages of observational data.

3.2 LAP deposition data

250 For testing the new snow scheme, the land model is driven by aerosol deposition fluxes obtained by a fully coupled simulation
with the GFDL AM4.0 and LM4.0 model (Zhao et al., 2018a, b). In the following we give a brief description of the model, and
refer the reader to (Ga Chan et al., 2022) for an extensive description of the experimental setup.

AM4.0 tracks five types of aerosol, among which are the tracers used in this study (BC, MD and OC). Aerosols are charac-
terized by a log-normal size distribution, except in the case of dust for which 5 size bins are used, with characteristic particle
255 radii ranging from 0.1 to 10 μm . For each tracer species, the model simulates sources at the surface, atmospheric transport by
advection, turbulent diffusion and convection, and deposition by both wet and dry processes. The lifting of dust is computed
using the model by Ginoux et al. (2001), employing an empirical threshold for wind erosion, and using land cover data from
CMIP6 forcing. Natural and anthropogenic sources of BC and OC were also obtained from CMIP6 emission data. The wet de-
position process includes both effects of condensation within clouds, and below-cloud scavenging of aerosols by precipitation.
260 The dry deposition of aerosols is driven by gravitational settling and turbulent exchange in the atmospheric boundary layer.

4 Results

4.1 Predictions for the Col de Porte site

We start by evaluating the model results at the Col de Porte (*cdp*) site for the 2013-2014 snow season. This is a useful bench-
mark, as i) the same site and season were used in previous studies making this a useful case study for model intercomparison
265 (Tuzet et al., 2017; Ga Chan et al., 2022), and ii) at this site the measurements by Dumont et al. (2017) allow to evaluate not
only snow bulk properties but also surface snow properties, namely snow optical diameter and concentration of impurities.

For the *cdp* test site we find that GLASS reproduces daily snow albedo very well (Figure 2A). We compare three model con-
figurations, forcing the model with all three LAP species (GLASS-LAPS), forcing the model with mineral dust only (GLASS-
DUST) and with no impurities deposited on snow (GLASS-CLEAN). The analysis reveal that i) the effect of dust and car-
270 bonaceous aerosols on albedo are comparable in magnitude for this site, and ii) the best match to daily albedo observation is
provided by the GLASS-LAPS model configuration. Note that the model overestimates the albedo of underlying soil substrate.
The reason for this mismatch is that the value of soil reflectivity was not calibrated for this particular location, and there is
no vegetation in this configuration of the model. When considering snow water equivalent predictions (Figure 2B), the model
matches observations very closely. During the ablation phase, some difference can be observed between model configurations,




Figure 1. Conceptual scheme of the LAP deposition process implemented in GLASS and its effects on snow optical properties. The input of LAP mass in the snow is given by wet and dry deposition. Vertical exchange of LAPs with the snowpack is driven by vertical water flow and is proportional to a specie-specific scavenging coefficient. The resulting concentration in near surface layer is used, together with surface snow properties, to predict snow albedo.




Figure 2. Results for the Col de Porte site during the 2013-2014 snow year. Simulated and observed values of daily albedo (panel A), SWE (panel B), snow depth (panel C) and snow density (panel D). Observations are reported as black markers (circles for manual obs., triangles for automatic obs.). Model simulations are reported for model forced by all LAPs (GLASS-LAPS, red lines), model forced by dust only (GLASS-DUST, tan lines) and by model with no impurities in snow (GLASS-CLEAN, green lines).



275 with snow disappearing about 10 days later in the case of clean snow compared to the configuration forced by all impurities. However, there are also comparable differences between automatic and manual snow depth observations in this period, with manual observations being closest to the GLASS-LAPS model prediction.

All three model configurations tend to overestimate snow depth throughout the winter with minimal differences between LAP treatments (Figure 2C). During the melt season, similarly to the results obtained for SWE, the closest match to observations
280 is given by the models forced by impurities. The discrepancy between the excellent fit for SWE and the less good snow depth results can be explained by limitations in how snow density is represented in GLASS. However, the evolution of snow density simulated throughout the season exhibit a constant underestimation (Figure 2D), suggesting perhaps a discrepancy in initial density assigned to fresh snow. We note that because GLASS is designed for global scale climate predictions, these values are not calibrated for specific locations.

285 For the *cdp* site, spectral measurements were collected during 2014 and used to estimate snow SSA and impurities content (Dumont et al., 2017). The estimates of SSA vary significantly during the cold season in the range from 60 to 5 m²kg⁻¹. The snow model overall captures the magnitude and range of variations of SSA during the winter (Figure 3A). The decrease of SSA during the ablation season is qualitatively reproduced by the model, although observed values then to be somewhat lower than model predictions.

290 The concentration of LAPs in the near surface snow layer is also well captured by the model (Figure 3B), considering that forcing values are obtained from an atmospheric model climatology dataset. For this reason, we do not expect the model to closely match variations in $c_{eq,ns}$ throughout the entire snow season. The order of magnitude of observed LAP concentration is comparable with simulations, but intra-seasonal variations are not captured by the model. During spring, the snow ablation phase is characterized by a sharp increase in LAP concentration driven by the combined effect of sublimation and melt. During
295 this phase, the increase in $c_{eq,ns}$, is overall well represented in the model.

4.2 Predictions for the SnowMIP sites

For some SnowMIP sites, daily albedo observations are available for model evaluation (Figure 4). During the accumulation phase, some underestimation of daily albedo by the model can be observed at some of the sites (*snb*, *sap*, *wff*). During ablation phase, the results are dependent on the model's ability to correctly capture the timing of complete snow melt. However, in
300 general the decrease of albedo in the spring appears better reproduced by the model configurations with impurities, except in the case of *wff*.

SWE model predictions show a good fit to observations at most SnowMIP sites (Figure 5). However, for some stations overestimation (*rme*, *snb*) or underestimation of SWE (*swa*, *sap*, *wff*) are observed for the specific snow year examined. For the sites in Colorado (*swa* and *snb*) we find that dust dominates the LAP effect on snow melt when compared to carbonaceous
305 aerosols (Figure 5, panels A and B). The effect of LAPs is smallest for the Japanese site (*sap*) and for the one in Finland (*sod*). For the sites located in the Alps (*wff*, in Figure 5D, and the already examined Col de Porte, *cdp*, featured in Figure 2B), we find that i) the effect of carbonaceous particles (BC and OC) is larger than simulated for the North American sites, and has the same




Figure 3. Results for snow specific surface area (panel A) and near-surface concentration of impurities (panel B) for the Col de Porte site (2013-2014 snow year). Spectral measurements carried out by (Dumont et al., 2017) are reported as black marker for different value of retrieval parameter a . Model simulations by GLASS-LAPS are reported as blue lines.




Figure 4. Daily average albedo predictions for 6 SnowMIP sites for a single snow year. For each site, we compare model runs with all LAP species (GLASS-LAPS, red), with dust-only forcing (GLASS-DUST, tan), and with clean snow (GLASS-CLEAN, green). Where available, daily albedo observations are reported as reference (black circles).




Figure 5. SWE predictions for 6 SnowMIP sites for a single snow year. For each site, we compare model runs with all LAP species (GLASS-LAPS, red), with dust-only forcing (GLASS-DUST, tan), and with clean snow (GLASS-CLEAN, green). Where available, SWE observations are reported as reference (black circles for manual observations, black triangles for automatic observations).



		swa	snb	sap	wfj	sod	rme	cdp
CLEAN - LAPS	μ_t	23.6	19.1	7.2	25.9	4.4	8.3	9.8
	σ_t	5.0	3.2	3.2	7.7	1.2	3.4	4.5
CLEAN - DUST	μ_t	18.7	14.7	3.9	18.2	2.6	4.4	5.7
	σ_t	4.2	3.2	2.6	7.7	0.9	2.0	3.2

Table 1. Decrease in the number of snow days in the LAPS and DUST cases compared to CLEAN, including temporal mean (μ_t) and standard deviations (σ_t) of yearly values computed over the entire experiment period.

magnitude as the effect of dust. When the model is forced with dust only, the number of snow days lost decreases significantly (and is between half and two-thirds of that predicted in the all-LAPs scenario).

310 Similarly, the skill in snow depth predictions varies across sites (Figure 6). Good performance are observed for *swa*, *sap*, and *sod*, although in the latter case some overestimation is observed during the boreal winter. A considerable overestimation of snow is observed for *snb* and *rme* sites, while the melt occur earlier then observed in *wfj*, case in which clean snow model would have the best fit.

To quantify the effect of LAPs, we examine the number of snow days predicted by the different model configurations. The 315 decrease in number of snow days for the stations is reported in Table 1 (average and standard deviation across the years of the experiment) for the DUST only and all-LAPs scenarios. The largest average number of snow days lost is observed for *swa*, *snb* (dust-driven), and *wfj*. However, in the latter case the CLEAN scenario is actually the closest to the observations.

Figure 7 shows the differences in yearly maximum SWE values between CLEAN, LAPS and DUST runs. Here we find that the effect of LAPs on yearly maximum SWE peak is very small for all sites, as one would expect as the effect of LAPs is more 320 pronounced during the ablation rather than during the snow accumulation phase.

Figure 8 shows the differences in number of snow days values between CLEAN, LAPS and DUST runs. this number is quite significant for several of the sites. However, we find that the year-to-year variations in number of snow days lost to LAPs are generally contained.

5 Discussion

325 We found that the effect of dust is dominant over that of carbonaceous aerosols for the sites in North America (i.e., for the sites *snb*, *swa*, *rme*), where the combined effect of LAPs determine a significantly shorter snow season than would be in the case of clean snow (shorter by 24 and 19 days for the *swa* and *snb* sites, averaged over the entire time period of the observational records). In the case of the Senator Beck Basin site (*snb*), effects of similar magnitude have been reported by (Skiles and Painter, 2019), which found a dust-driven reduction of about 30 snow days, with previous estimates ranging between about 330 20 to 50 days depending on the LAP concentration in the spring (Skiles et al., 2012). Similarly, for the Swamp Angel site, in our analysis the melt date advances on average by 24 days, corresponding to the lower end of a previous study (Skiles et al., 2015). Our results on average indicate an effect of all LAPs on melt date which is the lower end of the range computed in these




Figure 6. Snow depth predictions for 6 SnowMIP sites for a single snow year. For each site, we compare model runs with all LAP species (GLASS-LAPS, red), with dust-only forcing (GLASS-DUST, tan), and with clean snow (GLASS-CLEAN, green). Where available, snow depth observations are reported as reference (black circles for manual observations, black triangles for automatic observations).




Figure 7. Snow water equivalent predictions for each year in different model configurations: With all LAP species (GLASS-LAPS, red), with dust-only forcing (GLASS-DUST, tan), and with clean snow (GLASS-CLEAN, blue).




Figure 8. Number of snow days simulated for each year in different model configurations: With all LAP species (GLASS-LAPS, red), with dust-only forcing (GLASS-DUST, tan), and with clean snow (GLASS-CLEAN, blue).



previous studies. While dust is less absorptive than BC, it is by far the most abundant by mass and at these sites appears to dominate the snow surface darkening. Similar results underlying the primary role played by mineral dust on snow melt have
335 been reported for other regions. For example, Sterle et al. (2013) reported a similar behavior for snow in the Sierra Nevada, with mineral dust exhibit a larger impact of snow compared to black carbon. Over high-mountain Asia, Sarangi et al. (2020) observed dust playing a fundamental role in the darkening of high-altitude snow.

On the other hand, we found that for the Alpine sites the effects of dust and carbonaceous particles are similar in magnitude. In the case of the Col de Porte site (*cdp*), the result we obtained (10 snow days lost due to all LAPs, on average) is slightly
340 larger than the results from a previous modelling studies, (Tuzet et al., 2017), which reported snow melt date advancing 6 to 9 days in the 2013-2014 season depending on LAP parameters used in the analysis. For the *cdp* site, we were able to compare modelled snow near-surface properties (snow specific surface area and concentration of BC-equivalent concentration of LAPs) with in-situ spectral measurements for one snow season. This analysis showed that significant scatter exist between modelled and observed values. However, the order of magnitude of LAP concentration was captured by the model, as well as the increase
345 in LAPs during the ablation season. We note here that since the snow scheme is forced by modelled deposition fluxes, a perfect match to observation was not necessarily expected. Similarly, the increase in snow optical diameter (i.e., decrease in SSA) during the snow ablation phase was also captured by the model, although with some overestimation of observed SSA values.

In future extensions of this work, we plan to explore the ability of GLASS-LAPS to reproduce spatial statistics of snow cover using sub-grid tiling schemes and accounting for topography (e.g., Chaney et al. (2018); Zorzetto et al. (2022)). Furthermore,
350 the model could be improved by using more detailed optical models such as SNICAR (Flanner and Zender, 2005). We note that while some discrepancies are observed for specific sites, the model was developed for global applications and was not tailored to terrain or climate conditions of these sites. Thus, multiple physical processes may be at the origin of these discrepancies, as also discussed by Zorzetto et al. (2024), and will be the subject of future model development efforts.

6 Conclusions

355 In this work we extended a recently developed numerical snow model (GLASS) to include the physical processes connected with the deposition of LAPs. GLASS, implemented in the GFDL land model, can be used in global scale, centuries-long climate simulations. We tested the new model configuration over a set of SnowMIP sites, forcing the model with in-situ meteorological data and with LAP deposition rates obtained from a general circulation model (AM4.0). We found that the model satisfactorily represents seasonal snow amounts over the test sites, although performance is to a certain extent location
360 dependent. This is not surprising given the large variability in climates, as well as surface properties and terrain types. Running the model with clean snow, dust only and forced by all LAPs allowed us to investigate the relative contribution of different aerosol species to snow melt. We found that the effect of LAPs on snowpack evolution is significant at all sites examined, with an average number of snow days lost due to LAPs varying between 5 and 24. For sites in the Western US, the effect of dust is predominant and is responsible for most of the LAP-driven melt. In other locations this is not the case: In the sites in the
365 Alps, for instance, carbonaceous aerosols play a larger role relative to dust. Our results support large-scale applications of the



new model configuration to simulate snowpack globally, under historical and projected climate condition. These analysis will be the objective of future work.

Code and data availability. The code and data used in this study are freely available online at <https://zenodo.org/records/10901373>

Author contributions. All authors contributed to research design and to writing the manuscript. E.Z. Developed software, performed analysis
370 of model results, and drafted a first version of the manuscript.

Competing interests. The authors declare no competing interest.

Acknowledgements. The authors acknowledge funding from the NOAA Climate Program Office (CPO) grant number NA18OAR4320123
375 "3D-Land Energy and Moisture Exchanges: Harnessing High Resolution Terrestrial Information to Refine Atmosphere-to-Land interactions
in Earth System Models". The authors thank Dr. Justin Perket and Dr. Nicole Schlegel at NOAA GFDL for reviewing a first draft of the
manuscript.



References

- Ackroyd, C., Skiles, S. M., Rittger, K., and Meyer, J.: Trends in snow cover duration across river basins in high mountain Asia from daily gap-filled MODIS fractional snow covered area, *Frontiers in Earth Science*, 9, 713 145, 2021.
- 380 Aoki, T., Kuchiki, K., Niwano, M., Kodama, Y., Hosaka, M., and Tanaka, T.: Physically based snow albedo model for calculating broadband albedos and the solar heating profile in snowpack for general circulation models, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *Journal of Glaciology*, 38, 13–22, 1992.
- 385 Carmagnola, C., Morin, S., Lafaysse, M., Domine, F., Lesaffre, B., Lejeune, Y., Picard, G., and Arnaud, L.: Implementation and evaluation of prognostic representations of the optical diameter of snow in the SURFEX/ISBA-Crocus detailed snowpack model, *The Cryosphere*, 8, 417–437, 2014.
- Chaney, N. W., Van Huijgevoort, M. H., Shevliakova, E., Malyshev, S., Milly, P. C., Gauthier, P. P., and Sulman, B. N.: Harnessing big data to rethink land heterogeneity in Earth system models, *Hydrology and Earth System Sciences*, 22, 3311–3330, 2018.
- 390 Clifton, A., Manes, C., Rüedi, J.-D., Guala, M., and Lehning, M.: On shear-driven ventilation of snow, *Boundary-layer meteorology*, 126, 249–261, 2008.
- Conway, H., Gades, A., and Raymond, C.: Albedo of dirty snow during conditions of melt, *Water resources research*, 32, 1713–1718, 1996.
- Cook, J., Hodson, A. J., Taggart, A., Mernild, S. H., and Tranter, M.: A predictive model for the spectral “bioalbedo” of snow, *Journal of Geophysical Research: Earth Surface*, 122, 434–454, 2017.
- 395 Dang, C., Brandt, R. E., and Warren, S. G.: Parameterizations for narrowband and broadband albedo of pure snow and snow containing mineral dust and black carbon, *Journal of Geophysical Research: Atmospheres*, 120, 5446–5468, 2015.
- Di Mauro, B., Fava, F., Ferrero, L., Garzonio, R., Baccolo, G., Delmonte, B., and Colombo, R.: Mineral dust impact on snow radiative properties in the European Alps combining ground, UAV, and satellite observations, *Journal of geophysical research: Atmospheres*, 120, 6080–6097, 2015.
- 400 Doherty, S., Warren, S., Grenfell, T., Clarke, A., and Brandt, R.: Light-absorbing impurities in Arctic snow, *Atmospheric Chemistry and Physics*, 10, 11 647–11 680, 2010.
- Dumont, M., Brun, E., Picard, G., Michou, M., Libois, Q., Petit, J., Geyer, M., Morin, S., and Josse, B.: Contribution of light-absorbing impurities in snow to Greenland’s darkening since 2009, *Nature Geoscience*, 7, 509–512, 2014.
- Dumont, M., Arnaud, L., Picard, G., Libois, Q., Lejeune, Y., Nabat, P., Voisin, D., and Morin, S.: In situ continuous visible and near-infrared
405 spectroscopy of an alpine snowpack, *The Cryosphere*, 11, 1091–1110, 2017.
- Dunne, J. P., Horowitz, L., Adcroft, A., Ginoux, P., Held, I., John, J., Krasting, J. P., Malyshev, S., Naik, V., Paulot, F., et al.: The GFDL Earth System Model version 4.1 (GFDL-ESM 4.1): Overall coupled model description and simulation characteristics, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 015, 2020.
- Estilow, T. W., Young, A. H., and Robinson, D. A.: A long-term Northern Hemisphere snow cover extent data record for climate studies and
410 monitoring, *Earth System Science Data*, 7, 137–142, 2015.
- Flanner, M. G. and Zender, C. S.: Snowpack radiative heating: Influence on Tibetan Plateau climate, *Geophysical research letters*, 32, 2005.
- Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *Journal of Geophysical Research: Atmospheres*, 111, 2006.



- 415 Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V., and Rasch, P.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmospheric Chemistry and Physics*, 9, 2481–2497, 2009.
- Flanner, M. G., Shell, K. M., Barlage, M., Perovich, D. K., and Tschudi, M.: Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008, *Nature Geoscience*, 4, 151–155, 2011.
- 420 Flanner, M. G., Liu, X., Zhou, C., Penner, J. E., and Jiao, C.: Enhanced solar energy absorption by internally-mixed black carbon in snow grains, *Atmospheric Chemistry and Physics*, 12, 4699–4721, 2012.
- Ga Chan, H., Ginoux, P., Malyshev, S., and Kapnick, S.: A parameterization of snowpack albedo reduction by light-absorbing impurities for use in large-scale models, NOAA Technical Report, 2022.
- Garratt, J. R.: The atmospheric boundary layer, *Earth-Science Reviews*, 37, 89–134, 1994.
- 425 Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research: Atmospheres*, 106, 20255–20273, 2001.
- Golaz, J.-C., Van Roekel, L. P., Zheng, X., Roberts, A. F., Wolfe, J. D., Lin, W., Bradley, A. M., Tang, Q., Maltrud, M. E., Forsyth, R. M., et al.: The DOE E3SM Model Version 2: Overview of the physical model and initial model evaluation, *Journal of Advances in Modeling Earth Systems*, 14, 2022.
- 430 Hadley, O. L. and Kirchstetter, T. W.: Black-carbon reduction of snow albedo, *Nature Climate Change*, 2, 437–440, 2012.
- Hao, D., Bisht, G., He, C., Bair, E., Huang, H., Dang, C., Rittger, K., Gu, Y., Wang, H., Qian, Y., et al.: Improving snow albedo modeling in E3SM land model (version 2.0) and assessing its impacts on snow and surface fluxes over the Tibetan Plateau, *Geoscientific Model Development Discussions*, pp. 1–31, 2022.
- Hao, D., Bisht, G., Rittger, K., Stillinger, T., Bair, E., Gu, Y., and Leung, L. R.: Evaluation of E3SM land model snow simulations over the western United States, *The Cryosphere*, 17, 673–697, 2023.
- 435 He, C., Flanner, M. G., Chen, F., Barlage, M., Liou, K.-N., Kang, S., Ming, J., and Qian, Y.: Black carbon-induced snow albedo reduction over the Tibetan Plateau: uncertainties from snow grain shape and aerosol–snow mixing state based on an updated SNICAR model, *Atmospheric Chemistry and Physics*, 18, 11 507–11 527, 2018a.
- He, C., Liou, K.-N., Takano, Y., Yang, P., Qi, L., and Chen, F.: Impact of grain shape and multiple black carbon internal mixing on snow albedo: Parameterization and radiative effect analysis, *Journal of Geophysical Research: Atmospheres*, 123, 1253–1268, 2018b.
- 440 He, C., Liou, K.-N., Takano, Y., Chen, F., and Barlage, M.: Enhanced snow absorption and albedo reduction by dust-snow internal mixing: modeling and parameterization, *Journal of Advances in Modeling Earth Systems*, 11, 3755–3776, 2019.
- Huang, H., Qian, Y., He, C., Bair, E. H., and Rittger, K.: Snow albedo feedbacks enhance snow impurity-induced radiative forcing in the Sierra Nevada, *Geophysical Research Letters*, 49, e2022GL098 102, 2022.
- 445 Jordan, R. E.: A one-dimensional temperature model for a snow cover: Technical documentation for SNTHERM. 89, 1991.
- Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., Hall, A., Rott, H., Brutel-Vuilmet, C., Kim, H., et al.: ESM-SnowMIP: assessing snow models and quantifying snow-related climate feedbacks, *Geoscientific Model Development*, 11, 5027–5049, 2018.
- Kunkel, K. E., Robinson, D. A., Champion, S., Yin, X., Estilow, T., and Frankson, R. M.: Trends and extremes in Northern Hemisphere snow characteristics, *Current Climate Change Reports*, 2, 65–73, 2016.
- 450



- Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning: Part II. Snow microstructure, *Cold regions science and technology*, 35, 147–167, 2002.
- Lejeune, Y., Dumont, M., Panel, J.-M., Lafaysse, M., Lapalus, P., Le Gac, E., Lesaffre, B., and Morin, S.: 57 years (1960–2017) of snow and meteorological observations from a mid-altitude mountain site (Col de Porte, France, 1325 m of altitude), *Earth System Science Data*, 11, 455 71–88, 2019.
- Libois, Q., Picard, G., France, J., Arnaud, L., Dumont, M., Carmagnola, C., and King, M.: Influence of grain shape on light penetration in snow, *The Cryosphere*, 7, 1803–1818, 2013.
- Liou, K., Takano, Y., He, C., Yang, P., Leung, L., Gu, Y., and Lee, W.: Stochastic parameterization for light absorption by internally mixed BC/dust in snow grains for application to climate models, *Journal of Geophysical Research: Atmospheres*, 119, 7616–7632, 2014.
- 460 Marshall, S. E.: A physical parameterization of snow albedo for use in climate models., NCAR Cooperative Thesis, 1989.
- Ménard, C. B., Essery, R., Barr, A., Bartlett, P., Derry, J., Dumont, M., Fierz, C., Kim, H., Kontu, A., Lejeune, Y., et al.: Meteorological and evaluation datasets for snow modelling at 10 reference sites: description of in situ and bias-corrected reanalysis data, *Earth System Science Data*, 11, 865–880, 2019.
- Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., Brutel-Vuilmet, C., Burke, E., Cuntz, M., Dai, Y., Decharme, 465 B., Dutra, E., Fang, X., Fierz, C., Gusev, Y., Hagemann, S., Haverd, V., Kim, H., Lafaysse, M., Marke, T., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Schädler, G., Semenov, V. A., Smirnova, T., Strasser, U., Swenson, S., Turkov, D., Wever, N., and Yuan, H.: Scientific and Human Errors in a Snow Model Intercomparison, *Bulletin of the American Meteorological Society*, 102, E61–E79, <https://doi.org/10.1175/bams-d-19-0329.1>, 2021.
- Milly, P. C., Malyshev, S. L., Shevliakova, E., Dunne, K. A., Findell, K. L., Gleeson, T., Liang, Z., Phillipps, P., Stouffer, R. J., and Swenson, 470 S.: An enhanced model of land water and energy for global hydrologic and earth-system studies, *Journal of Hydrometeorology*, 15, 1739–1761, 2014.
- Morin, S., Lejeune, Y., Lesaffre, B., Panel, J.-M., Poncet, D., David, P., and Sudul, M.: An 18-yr long (1993–2011) snow and meteorological dataset from a mid-altitude mountain site (Col de Porte, France, 1325 m alt.) for driving and evaluating snowpack models, *Earth system science data*, 4, 13–21, 2012.
- 475 Mudryk, L., Santolaria-Otín, M., Krinner, G., Ménégos, M., Derksen, C., Brutel-Vuilmet, C., Brady, M., and Essery, R.: Historical Northern Hemisphere snow cover trends and projected changes in the CMIP6 multi-model ensemble, *The Cryosphere*, 14, 2495–2514, 2020.
- Nijssen, B., Bowling, L. C., Lettenmaier, D. P., Clark, D. B., El Maayar, M., Essery, R., Goers, S., Gusev, Y. M., Habets, F., Van den Hurk, B., et al.: Simulation of high latitude hydrological processes in the Torne–Kalix basin: PILPS Phase 2 (e): 2: Comparison of model results with observations, *Global and Planetary Change*, 38, 31–53, 2003.
- 480 Painter, T. H., Barrett, A. P., Landry, C. C., Neff, J. C., Cassidy, M. P., Lawrence, C. R., McBride, K. E., and Farmer, G. L.: Impact of disturbed desert soils on duration of mountain snow cover, *Geophysical Research Letters*, 34, 2007.
- Painter, T. H., Skiles, S. M., Deems, J. S., Bryant, A. C., and Landry, C. C.: Dust radiative forcing in snow of the Upper Colorado River Basin: 1. A 6 year record of energy balance, radiation, and dust concentrations, *Water Resources Research*, 48, 2012.
- Qian, Y., Yasunari, T. J., Doherty, S. J., Flanner, M. G., Lau, W. K., Ming, J., Wang, H., Wang, M., Warren, S. G., and Zhang, R.: Light- 485 absorbing particles in snow and ice: Measurement and modeling of climatic and hydrological impact, *Advances in Atmospheric Sciences*, 32, 64–91, 2015.
- Qu, X. and Hall, A.: On the persistent spread in snow-albedo feedback, *Climate dynamics*, 42, 69–81, 2014.



- Réveillet, M., Dumont, M., Gascoïn, S., Lafaysse, M., Nabat, P., Ribes, A., Nheili, R., Tuzet, F., Ménégos, M., Morin, S., et al.: Black carbon and dust alter the response of mountain snow cover under climate change, *Nature communications*, 13, 5279, 2022.
- 490 Sarangi, C., Qian, Y., Rittger, K., Ruby Leung, L., Chand, D., Bormann, K. J., and Painter, T. H.: Dust dominates high-altitude snow darkening and melt over high-mountain Asia, *Nature Climate Change*, 10, 1045–1051, 2020.
- Shrestha, M., Wang, L., Koike, T., Xue, Y., and Hirabayashi, Y.: Improving the snow physics of WEB-DHM and its point evaluation at the SnowMIP sites, *Hydrology and Earth System Sciences*, 14, 2577–2594, 2010.
- Skiles, S. M. and Painter, T. H.: Toward understanding direct absorption and grain size feedbacks by dust radiative forcing in snow with
495 coupled snow physical and radiative transfer modeling, *Water Resources Research*, 55, 7362–7378, 2019.
- Skiles, S. M., Painter, T. H., Deems, J. S., Bryant, A. C., and Landry, C. C.: Dust radiative forcing in snow of the Upper Colorado River Basin: 2. Interannual variability in radiative forcing and snowmelt rates, *Water Resources Research*, 48, 2012.
- Skiles, S. M., Painter, T. H., Belnap, J., Holland, L., Reynolds, R. L., Goldstein, H. L., and Lin, J.: Regional variability in dust-on-snow processes and impacts in the Upper Colorado River Basin, *Hydrological Processes*, 29, 5397–5413, 2015.
- 500 Skiles, S. M., Flanner, M., Cook, J. M., Dumont, M., and Painter, T. H.: Radiative forcing by light-absorbing particles in snow, *Nature Climate Change*, 8, 964–971, 2018.
- Sterle, K. M., McConnell, J. R., Dozier, J., Edwards, R., and Flanner, M.: Retention and radiative forcing of black carbon in eastern Sierra Nevada snow, *The Cryosphere*, 7, 365–374, 2013.
- Tuzet, F., Dumont, M., Lafaysse, M., Picard, G., Arnaud, L., Voisin, D., Lejeune, Y., Charrois, L., Nabat, P., and Morin, S.: A multilayer
505 physically based snowpack model simulating direct and indirect radiative impacts of light-absorbing impurities in snow, *The Cryosphere*, 11, 2633–2653, 2017.
- Tuzet, F., Dumont, M., Picard, G., Lamare, M., Voisin, D., Nabat, P., Lafaysse, M., Larue, F., Revuelto, J., and Arnaud, L.: Quantification of the radiative impact of light-absorbing particles during two contrasted snow seasons at Col du Lautaret (2058 m asl, French Alps), *The Cryosphere*, 14, 4553–4579, 2020.
- 510 Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7. 2, *Geoscientific Model Development*, 5, 773–791, 2012.
- Warren, S. G.: Impurities in snow: Effects on albedo and snowmelt, *Annals of Glaciology*, 5, 177–179, 1984.
- Wiscombe, W. J. and Warren, S. G.: A model for the spectral albedo of snow. I: Pure snow, *Journal of Atmospheric Sciences*, 37, 2712–2733, 1980.
- 515 Zhao, M., Golaz, J.-C., Held, I., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L., Dunne, J., et al.: The GFDL global atmosphere and land model AM4. 0/LM4. 0: 1. Simulation characteristics with prescribed SSTs, *Journal of Advances in Modeling Earth Systems*, 10, 691–734, 2018a.
- Zhao, M., Golaz, J.-C., Held, I., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L., Dunne, J., et al.: The GFDL global atmosphere and land model AM4. 0/LM4. 0: 2. Model description, sensitivity studies, and tuning strategies, *Journal of Advances in
520 Modeling Earth Systems*, 10, 735–769, 2018b.
- Zorzetto, E., Malyshev, S., Chaney, N., Paynter, D., Menzel, R., and Shevliakova, E.: Effects of complex terrain on the shortwave radiative balance: A sub-grid-scale parameterization for the GFDL Land Model version 4.1, *Geoscientific Model Development*, pp. 1–28, 2022.
- Zorzetto, E., Malyshev, S., Ginoux, P., and Shevliakova, E.: A Global land snow scheme (GLASS) v1. 0 for the GFDL Earth System Model: Formulation and evaluation at instrumented sites, *EGUsphere*, 2024, 1–41, 2024.