

Evaluating adsorption isotherm models for determining the partitioning of ammonium between soil and soil-pore water in environmental soil samples

Matthew G. Davis*, Kevin Yan, Jennifer G. Murphy

5 Department of Chemistry, University of Toronto, Toronto, M5S 3H6, Canada

Correspondence to: Matthew G. Davis (mg.davis@mail.utoronto.ca)

Abstract: Ammonium in soil pore water is thought to participate in bidirectional exchange with the atmosphere; however, common soil nutrient analysis methods determine the bulk quantity of ammonium associated with the soil particles, rather than determining the aqueous ammonium concentration. Previous works have applied the Langmuir and Freundlich isotherm equations to ammonium-enriched soils to estimate partitioning, but this may not be representative of conditions in natural, unmanaged soils. In this work, environmental soil samples were collected from greenspaces in Toronto and used to evaluate several commonly used adsorption isotherm equations, including the Langmuir, Freundlich, Temkin and Toth equations, to determine their applicability in lightly managed and non-fertilized soils. We then compare ammonia emission potentials (a quantity predicting the propensity of ammonia to volatilize from a liquid reservoir) determined using a conventional high-salt extraction procedure to determine the soil ammonium content to that modelled using the Temkin and Langmuir equations, and demonstrate that conventional approaches may overestimate emission potentials from soils by a factor of 5 – 20.

Key words: ammonium soil adsorption; emission potential;

1. Introduction:

1.1. Contextualizing the significance of ammonium partitioning in soils

20 Globally, emissions of reduced nitrogen compounds (NH_3) make up as much as half of the global atmospheric reactive nitrogen sources (Flechar et al., 2013). Of the NH_3 budget, approximately two thirds of emissions are related to anthropogenic agricultural activities, with natural sources being responsible for only 15 – 20% of emissions (Bouwman et al., 1997; Sutton et al., 2008, 2013; Van Aardenne et al., 2001). Following emission, NH_3 tends to partition into fine particulate matter or deposit via wet or dry deposition on a timescale of hours to days. NH_3 is understood to engage in bidirectional exchange throughout the earth system, with NH_3 depositing, or volatilizing depending on local environmental conditions (Farquhar et al., 1980; Flechar et al., 1999, 2013; Guo et al., 2022a; Nemitz et al., 2000; Sutton et al., 1995; Walker et al., 2023; Wentworth et al., 2014; Zhang et al., 2010). The bidirectional exchange of NH_3 between soils and the atmosphere has been considered important to its overall budget, particularly in remote areas, but research on the mechanics of NH_3 partitioning in soils between adsorbed

(inaccessible) and aqueous (accessible) NH_3 often focuses on fertilized croplands with substantial concentrations of ammonia present (Venterea et al., 2015; Vogeler et al., 2011). Consequently, ammonia volatilization models may parameterize all or most of the soil ammonia as being readily able to exchange with the atmosphere, which may be reasonable for recently fertilized soils, but not for unmanaged soils (Massad et al., 2010a; Pleim et al., 2013a; Zhu et al., 2015a). However, cropland is estimated to make up less than 15% of the Earth's land area, while unfertilized or irregularly fertilized natural, semi-natural or pastoral land reflects nearly three quarters of terrestrial surfaces (Ellis et al., 2020). The short atmospheric lifetime of NH_3 makes it important to understand the exchange of NH_3 over all types of surfaces, despite agricultural cropland being the most globally significant source.

1.2. Importance of developing an ammonium adsorption partitioning model

Ammonium in soils is thought to be partitioned between NH_4^+ _(ads) adsorbed to soil particles and NH_4^+ _(aq) dissolved in soil pore water. Because soils tend to exist with both permanent and pH-dependent negative charges (Bache, 1976), NH_4^+ ions compete for adsorption against other cations in solution. The total quantity of cations that can adsorb to the soil particles is termed the cation exchange capacity (CEC, typically reported as centimoles positive charge/kg). Traditional methods for determining ammonium in soil—intended for nutrient analysis in which the total ammonia nitrogen is more important than the partitioning between NH_4^+ _(ads) and NH_4^+ _(aq)—use high concentration salt solutions to displace all cations from the soil. As a result, these approaches do not distinguish between NH_4^+ _(aq) and NH_4^+ _(ads) (Li et al., 2012) and therefore likely overestimate the emission potential of soils. In this manuscript, we explore a variety of adsorption isotherm models with the goal of identifying a simple approach to relate the total quantity of ammonium in soil to the aqueous fraction of ammonium that can participate in bidirectional exchange with the atmosphere. .

1.3. Adsorption Isotherm equations:

The adsorption behavior of molecules to surfaces is complex and dependent on the properties of both the surface and the adsorbed molecules. Numerous adsorption equations have been proposed, based on both theoretical and empirical models; however, as the partitioning of ammonium between soil pore water and soil particles is complex and influenced by many other simultaneous equilibria, we consider each of the examined equations as being empirically determined only, rather than based on a theoretical treatment of the system. Previous studies have sought to develop an adsorption model for NH_4^+ partitioning soils based on the Langmuir (Alnsour, 2020; Venterea et al., 2015) and Freundlich equations (Vogeler et al., 2011), but have focused on croplands and other agricultural soils. In this work, in addition to the Langmuir and Freundlich equations, we also investigate the application of the Temkin and Toth adsorption equations to NH_4^+ partitioning in soils. Each of these equations (as formulated in Table 1) represents the adsorbed NH_4^+ concentration (S , mg kg^{-1}) in terms of the NH_4^+ concentration in solution (C , mg L^{-1}). Except for the Freundlich equation, each of these equations incorporate a saturation point or maximum adsorption capacity (S_{max} , mg kg^{-1}), in this work, we investigate the potential to calculate these adsorption equations as a function of the measured CEC (converted to $\text{mg of NH}_4^+ \text{ kg}^{-1} \text{ soil}$), rather than treating S_{max} as a calculated fitting parameter.

Table 1: Adsorption isotherm equations applied in this work and their parameters

Adsorption Isotherm Model	Equation	Units	Reference
Langmuir	(1) $S = \frac{S_{max}k_L C}{1+k_L C}$	S _{max} (mg kg ⁻¹), K _L (L kg ⁻¹)	(Langmuir, 1916)
Freundlich	(2) $S = k_F C^{n_F}$	k _F , n _F (dimensionless)	(Freundlich, 1909)
Temkina	(3) $S = q_T \ln(1 + K_T C)$	q _T (product of S _{max} (mg kg ⁻¹) and $\frac{RT}{b}$ (dimensionless)), K _T (L mg ⁻¹)	(Temkin & Pyzhev, 1940)
Toth	(4) $S = \frac{bC}{(K_{To} + C^{n_T})^{\frac{1}{n_T}}}$	b (product of S _{max} (mg kg ⁻¹) and a dimensionless scaling factor), K _{To} (mg L ⁻¹), n _T (dimensionless)	(Tóth, 1995)

a The Temkin model is also given as $S = \frac{RT}{b} \ln K_T C$, see (Chu, 2021) for this formulation

2. Methods:

2.1. Soil Collection:

65 Soil samples were collected from several greenspace (parks, urban forest, roadside-sites, etc.) locations across Toronto. Samples were collected only on days preceded by at least two days without precipitation and were collected by inserting a 7.5 cm internal diameter steel tube into the ground to a depth of 5 – 10 cm and recovering a soil core by removing the tube from the ground. The soil cores were mixed, and sieved immediately, and transported back to the lab for analysis. Samples analyzed for NH₃ content were always analyzed immediately to avoid potential artefacts from freezing, samples analyzed for CEC or
70 adsorption curves were frozen for storage prior to analysis. Soil samples were collected from eight locations in Fall 2021 as a training set for developing the model. A subsequent 16 soil samples from across Toronto were collected in Spring/Summer 2023 to be used to evaluate the effectiveness of applying the model to uncharacterized soils, and to determine the impact on soil emission potentials.

2.2. Soil analysis:

2.2.1. Cation Exchange Capacity determination

75 The CEC was determined using the inductively-coupled plasma optical emission spectroscopy (ICP-OES) cation sum method (Bache, 1976). Briefly, 1 g of soil was measured out and mixed with 25 mL of 1 M NH₄Cl, shaken, and refrigerated for 36 - 48 hours to settle. The supernatant was filtered using 0.2 um syringe filters, diluted 50-fold using volumetric glassware and analyzed for the common soil-associated cations, Na⁺, K⁺, Mg²⁺, and Ca²⁺. The ICP-OES was calibrated using a commercially
80 available mixed standard of 6 cations (Li⁺, Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺) (Thermo Fisher Scientific). Another soil-associated cation that can contribute to CEC is Al³⁺, our initial measurements screened for Al, but we did not detect it in solution. The CEC was determined for all 24 collected samples and used to select three of the soil samples from the 16 collected in 2023 to be used as a test set for the developed model.

2.2.2. Adsorption curve characterization

85 The determination of the adsorption behavior of NH_4^+ was performed using a modified version of the procedure described by Venterea et al. (2015), combined with the ICP-OES cation sum method (Bache, 1976). Briefly, a series of batch equilibrium samples were prepared by mixing 1 g of soil with 25 mL aliquots of NH_4Cl solutions with concentrations ranging from 2.5 – 1000 mM. The samples were shaken, refrigerated for 36 – 48 hours to settle, filtered with 0.2 μm syringe filters, diluted 50-fold with volumetric glassware, and analyzed using ICP-OES. The quantity of NH_4^+ adsorbed onto each soil was inferred
90 based on the displaced Na^+ , Mg^{2+} , Ca^{2+} and K^+ ions measured in solution. In the cation sum method, the total quantity of adsorbable cations is determined by saturating the soil with an index cation (in this procedure, and generally, NH_4^+ from NH_4Cl), which displaces the original cations in the soil's adsorption sites. Thus the displaced cations measured for each NH_4Cl solution concentration are representative of the quantity of NH_4^+ adsorbing onto the soil. Adsorption curves were determined for the original eight soil samples in the training set, as well as the three selected soil samples for the test set.

95 2.2.3. pH and NH_4^+ determination

pH and NH_4^+ content was determined for the 16 soil samples collected in 2023. pH was determined for each soil sample using a typical approach of mixing soil with DIW in a 1:1 ratio, and then measuring the pH of the slurry by immersing a pH electrode (Hach) until a stable pH reading was obtained. To determine the pH using an alternative extraction solution, this process was repeated using a 0.01 M CaCl_2 solution.

100 NH_4^+ was extracted from the soil using a batch equilibrium method using both a 2 M KCl extraction solution and a DIW solution. In both cases, 2.5 g of soil was mixed with 25 mL of the extraction solution, shaken, and refrigerated for 36 hours to allow suspended solids to settle out of solution. Afterward, the supernatant was filtered using 0.2 μm PES membrane syringe filters. The soil NH_4^+ was quantified using the indophenol-blue salicylate method (Kempers & Zweers, 1986). Briefly, two reagent solutions, A and B were prepared: Reagent A consisted of a solution of 1 M sodium salicylate and 100 mg L^{-1}
105 sodium nitroprusside, while Reagent B consisted of a solution of 1 M NaOH and 0.12% by volume of 5% available chloride NaOCl. Soil extracts were prepared for analysis by adding 0.6 mL of reagent A to 2 mL of soil extract, followed by the addition of 1.4 mL of reagent B. The mixtures were then stored for 2 hours in the dark for color development, and then quantified using UV-VIS spectrometry (Lambda 365, Perkin-Elmer) at 649 nm.

2.3. Data analysis:

110 2.3.1. Characterizing adsorption parameters using a training and test set

Curve-fitting was done in R using the nls function to fit our experimental data for the eight samples in our training set to the Langmuir, Freundlich, Temkin, and Toth equations. Goodness of fit was evaluated by calculating the Akaike information criterion (AIC) using the AIC function from the R *stats* package. The AIC is calculated using equation (5):

$$(5) \text{ AIC} = 2\text{K} - 2\ln(\text{L})$$

115 Where K is the number of independent variables, and L is the log-likelihood estimate. The log-likelihood estimate can be extracted directly from nls objects fitted in R using the AIC function.

Fitting parameters for each equation were determined by pooling all the experimental data, standardizing each adsorption curve by the maximum adsorption achieved (i.e., all curves went from 0 to 1), and then fitting each equation from Table 1 to those
120 curves.

To validate the effectiveness of these equations when applied to uncharacterized soil samples, we selected three soil samples from the 16 soil samples collected in 2023 to form a test set, these samples were selected by choosing the soil samples with the lowest (10.9) and highest CECs (37.2), and a soil with an average CEC (25.3). As the original training set mostly consisted of samples with CECs from 20 – 30 (with two samples with CECs of 7.6 and 16), we chose two samples that were
125 significantly different than the average training set sample, as well as one similar sample to determine whether the fitting parameters could be used for ‘extreme’ samples, or only for samples similar to the training set. The test set was characterized in the same way as the training set, and was then modelled using three approaches:

- i. Using the average CEC for all the soil samples of 25 cmole kg⁻¹ (S_{max} of 4500 mg kg⁻¹), and the training set parameters.
- ii. Using the measured CEC for each soil sample to calculate S_{max}, and the training set parameters.
- 130 iii. Using the fitting algorithm as described in Sect.2.3.1 to determine the least squares fit for each equation to the experimental data.

2.3.2. Emission potential determination

The emission potential of ammonia is a quantity calculated as the ratio of aqueous NH₄⁺ to H⁺ (Eq.(6)).

$$(6) \Gamma = \frac{[NH_4^+]}{[H^+]}$$

135 Commonly, for soils this would be calculated using the total NH₄⁺ (determined using a salt solution extraction) and the pH measured using an extraction with deionized water. We denote this as Γ_{STD} , corresponding to $\frac{[NH_4^+]_{salt}}{[H^+]_{DIW}}$. The pH may also be measured in a (less concentrated) salt solution, which we denote as Γ_{SALT} , corresponding to $\frac{[NH_4^+]_{salt}}{[H^+]_{salt}}$. Similarly, though we are not aware of this as a common method, NH₄⁺ could be determined using a DIW extraction solution, resulting in a third parameterization of the emission potential as Γ_{DIW} , representing $\frac{[NH_4^+]_{DIW}}{[H^+]_{DIW}}$. Lastly, by applying one of the adsorption isotherm
140 models, the total soil NH₄⁺ can be partitioned into NH₄⁺_(ads) (S) and NH₄⁺_(aq) (C), and the emission potential can be calculated using only the NH₄⁺ in solution (C). These versions of the emission potential are denoted as Γ -sub-equation-name (e.g., as $\Gamma_{Langmuir}$, Γ_{Temkin} etc).

3. Results:

3.1. Performance of adsorption isotherm equations applied to an environmental soil training and test set:

145 We evaluated the ability of the Langmuir, Freundlich, Temkin and Toth equations to model the partitioning of NH_4^+ between
adsorbed and aqueous forms. An adsorption curve was characterized for each soil, the data from each soil adsorption
experiment was pooled, and then fit using the R nls function. The adsorption curves and fitting parameters are shown in Figure
1 and Table 2. Additionally, as our interest is ultimately in the performance of these adsorption equations at the lower
concentration limit, we refit each equation using only the extraction solutions $\leq 680 \text{ mg L}^{-1} \text{ NH}_4\text{Cl}$. The adsorption curves and
150 fitting parameters under those conditions are shown in Figure 2 and Table 3.

While the Langmuir (Alnsour, 2020; Guo et al., 2022b; Venterea et al., 2015) and Freundlich (Vogeler et al., 2011)
equations have been previously reported as being effective at modelling NH_4^+ partitioning in soils, we found them to be the
least effective of the equations we examined for the full adsorption curves, both over- and under-estimating the adsorbed NH_4^+
concentrations (Figure 1), while the Temkin and Toth equations better fit the experimental data. Computing the Akaike
155 information criterion for these equations results in an AIC of -138, -190, -222, and -249 for the Langmuir, Freundlich, Temkin,
and Toth equations respectively. The absolute value of the AIC is not important, but for a set of models, the model with the
lowest AIC is considered the best at fitting the experimental data, indicating an order of Toth>Temkin>Freundlich>Langmuir
for model effectiveness. However, fitting only the lower range of the adsorption curves (0 – 40 mM) slightly changed these
results, the Toth equation could not be fit by our algorithm, and the resulting AIC values were -93, -157, -152 for the Langmuir,
160 Freundlich, and Temkin equations respectively, indicating that the Freundlich equation best fit the experimental data. (Note
that AIC values for models fit to different datasets should not be directly compared to one another.)





Figure 1: Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, - · - ·), and Toth (orange, · · ·) equations. The curves are plotted using the experimental data from all eight soil adsorption experiments, the y-axis is normalized to the maximum adsorption achieved during each experiment.

165

Table 2: Comparison of goodness of fit and fitting parameters for the selected adsorption equations

Equation	Number of fitted parameters	AIC	Parameter 1	Mean \pm Standard error	Parameter 2	Mean \pm Standard error	Exponential Factor
Langmuir	1	-138	K_L	$9.29 \cdot 10^{-4} \pm 5.9 \cdot 10^{-5}$	S_{max}	-	-
Freundlich	2	-190	K_F	$S_{max} * 0.0520 \pm 3.4 \cdot 10^{-3}$	-	-	0.3050 ± 0.0074
Temkin	2	-222	K_T	$1.33 \cdot 10^{-2} \pm 1.2 \cdot 10^{-3}$	q_T	$S_{max} * 0.180 \pm 4.1 \cdot 10^{-3}$	-

Toth	3	-249	K_{To}	3.10 ± 0.65	b	$S_{max} * 2.45 \pm 0.35$	0.25 ± 0.027
------	---	------	----------	-----------------	---	---------------------------	------------------



170 **Figure 2:** Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, ---), and Temkin (green, ·-·-) equations. The curves are plotted using the experimental data from all eight soil adsorption experiments, but only using the extraction solutions of ≤ 40 mM for fitting, the y-axis is normalized to the maximum adsorption achieved during each experiment. The Toth equation could not be fit to the experimental data under these conditions.

Table 3: Comparison of goodness of fit and fitting parameters for the selected adsorption equations, data fit only for the ≤ 40 mM solutions


Equation	Number of fitted parameters	AIC	Parameter 1	Mean \pm Standard error	Parameter 2	Mean \pm Standard error	Exponential Factor
Langmuir	1	-93	K_L	$1.2 \cdot 10^{-3} \pm 8.0 \cdot 10^{-4}$	S_{max}	-	-

Freundlich	2	-157	K_F	$S_{max} * 0.025 \pm 3.0 \cdot 10^{-3}$	-	-	0.42 ± 0.019
Temkin	2	-152	K_T	$3.4 \cdot 10^{-2} \pm 6.5 \cdot 10^{-3}$	q_T	$S_{max} * 0.120 \pm 8.5 \cdot 10^{-3}$	-
Toth	3	Fit did not converge					

175

With our objective being to evaluate how well each equation can fit soils without going through the full characterization procedure, we analyzed the adsorption curves of the low-CEC, medium-CEC and high-CEC soils in our test set using: i) a “typical” CEC of 25, and the fitting parameters from Table 2, ii) the measured CEC and the fitting parameters from Table 2, and iii) by fitting the equations using the least-squares fitting algorithm. Using the first approach, we found that none of the equations could reasonably fit the experimental data when using an incorrect CEC and that each of the equations fit the experimental data reasonably well using the average parameters and the correct CEC (Figure 3). The relative goodness of fit for each equation was the same for the test set as for the training set, i.e. Toth>Temkin>Freundlich>Langmuir. We also tested an alternative approach for calculating each equation’s fitting parameters, in which rather than pooling the normalized data, and then fitting each curve, the curves were fit to each soil adsorption curve separately, and the resulting fitting parameters were then pooled. A full summary of the alternative fitting parameters and the test-set characterization is given in Figure A1 and Tables A1 – A3.

185



190 **Figure 3: Curve-fitting comparison between the unfitted Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, ·-·-), and Toth (orange, ···) equations using the fitting parameters from Table 3 applied to the test set of the a) Low CEC, b) Moderate CEC, c) High CEC samples (method ii)**

Overall, we find that the Toth and Temkin equations best fit the full adsorption curves, while the Freundlich and Temkin equations best fit the low-range adsorption curves. However, both the Freundlich and Toth equations seem to be less robust when applied to this system than the Temkin equation. Firstly, the Freundlich equation, as formulated in Table 2 and Table 3, has an intrinsic dependence on the CEC, which is contrary to the theoretical basis of the Freundlich equation. Using the

195

alternative fitting approach, the Freundlich equation does not require a CEC-dependence, but in that case, it has high error when fitting uncharacterized soils. Similarly, for the Toth equation, the alternative approach results in a significantly different set of fitting parameters, which resulted in the Toth equation quite poorly fitting the non-characterized soils. Additionally, while this may be a limitation of the fitting algorithm we selected, we found the Toth equation to be more difficult to consistently fit to our experimental data, with no solution being found for the low-range concentrations. Consequently, taking all of these factors into consideration, we find that the Temkin equation is most suitable for use evaluating NH_4^+ adsorption from uncharacterized soil samples. For comparison with previous studies, in which the Langmuir equation is the most frequently used adsorption equation, we will continue to analyze it in subsequent sections, despite it performing much worse than the Temkin equation for both the low-range and full-range of data.

3.2. Determining aqueous NH_4^+ concentrations using the Langmuir and Temkin equations

To relate the total ammonium measured (m_{NH_4}) to the adsorbed portion (S) and the aqueous portion (C), we define Eq.(7):

$$(7) m_{\text{NH}_4} = S + \frac{wC}{\rho}$$

Where m_{NH_4} is the total mass of ammonium per kg soil (mg kg^{-1}), w the volumetric moisture content of the soil (L water L^{-1} soil), and ρ is the bulk density of the dry soil (kg L^{-1}). By substituting the Langmuir (1) or Temkin (3) equations into Eq.(7), m_{NH_4} can be expressed in terms of C:

$$(8) m_{\text{NH}_4} = \frac{S_{\text{max}}k_L C}{1+k_L C} + \frac{wC}{\rho}$$

$$(9) m_{\text{NH}_4} = q_T \ln(1 + K_T C) + \frac{wC}{\rho}$$

To solve for C, we inverted these equations using *Wolfram Mathematica*, yielding Eq.(10) (Langmuir) and Eq.(11) (Temkin).

$$(10) C = \frac{-S_{\text{max}} \cdot K_L + K_L \cdot m_{\text{NH}_4} - Z + \sqrt{4K_L \cdot Z \cdot m_{\text{NH}_4} + (S_{\text{max}} \cdot K_L - K_L \cdot m_{\text{NH}_4} + Z)^2}}{2K_L \cdot Z}$$

$$(11) C = \frac{-Z + K_T \cdot q_T \cdot \mathbf{W}\left(\frac{Z}{e^{K_T \cdot q_T} + \frac{m_{\text{NH}_4}}{q_T} \cdot Z}\right)}{K_T \cdot Z}, \text{ where } \mathbf{W}(\mathbf{x}) \text{ is the Lambert W function, and } Z = \frac{w}{\rho}$$

If it is assumed that $m_{\text{NH}_4} \approx S$, then $\frac{wC}{\rho} \approx 0$, and these equations can be simplified to Eq.(12) and Eq.(13) respectively.

$$(12) C = \frac{m_{\text{NH}_4}}{k_L(S_{\text{max}} - m_{\text{NH}_4})}$$

$$(13) C = \frac{e^{\frac{m_{\text{NH}_4}}{q_T}} - 1}{K_T}$$

We tested this assumption and found that for our unfertilized soil samples, the simplified equations have a positive bias of only 0.57 – 1.5%, and that for most soils fertilized with the equivalent of up to 300 kg N ha^{-1} , the simplified equations should have a positive bias of less than 5%. Consequently, we used the simplified equations for our analysis.

3.3. Determining environmental soil emission potentials

225 Soil emission potentials were then calculated using the approaches described in Sect. 2.3.3, corresponding to the ‘standard’ approach, two approaches based on matching the NH_4^+ and pH extraction solutions, and two approaches based on applying the Langmuir and Temkin equations to the measured extract concentrations and pH values. The adsorption equations were calculated using both the low-range and full curve fitting parameters. For the Langmuir calculations, the fit parameters were $K_L = 9.29 \cdot 10^{-4} \text{ L kg}^{-1}$ (full) and $K_L = 1.20 \cdot 10^{-3} \text{ L kg}^{-1}$ (low-range), and S_{\max} calculated as the equivalent of the CEC in units of

230 mg kg^{-1} . For the Temkin calculations, the fit parameters were $K_T = 1.33 \cdot 10^{-2} \text{ L mg}^{-1}$, and $q_T (\text{mg kg}^{-1}) = 0.180 \cdot S_{\max}$ (full) and $K_T = 3.40 \cdot 10^{-2} \text{ L mg}^{-1}$, and $q_T (\text{mg kg}^{-1}) = 0.120 \cdot S_{\max}$. Among the approaches, Γ_{STD} results in the highest estimate for the emission potential ($17,000 \pm 19,000$), followed by Γ_{Langmuir} and Γ_{SALT} ($2730 - 3530$), and Γ_{Temkin} and Γ_{DIW} ($810 - 1450$) (Table 3). Emission potential is linearly related to equilibrium vapor pressure, at a temperature of 15°C , these emission potentials correspond to equilibrium concentrations of $2.4 - 50$ ppb. Although CEC is inversely related to the proportion of NH_4^+ present

235 in the aqueous phase, the emission potentials increase exponentially as a function of the CEC due to the strong positive correlation between soil pH and CEC (Figure 3).

Table 4: Comparison of emission potentials determined using the standard, Langmuir, Temkin, salt extraction and DIW extraction methods.

Method	Equation	Average Emission Potential (mol mol ⁻¹) (n = 16)	Equilibrium vapor pressure at 15°C (ppb)
Γ_{STD}	$\frac{[\text{NH}_4^+]_{\text{salt}}}{[\text{H}^+]_{\text{DIW}}}$	$17,000 \pm 19,000$	50 ± 55
Γ_{SALT}	$\frac{[\text{NH}_4^+]_{\text{salt}}}{[\text{H}^+]_{\text{salt}}}$	3300 ± 3100	9.6 ± 9.0
Γ_{DIW}	$\frac{[\text{NH}_4^+]_{\text{DIW}}}{[\text{H}^+]_{\text{DIW}}}$	1450 ± 1900	4.2 ± 5.5
$\Gamma_{\text{Temkin (Full)}}$	$\frac{e^{\frac{[\text{NH}_4^+]_{\text{salt}}}{q_T} - 1}}{K_T} \frac{1}{[\text{H}^+]_{\text{DIW}}}$	1370 ± 1300	4.0 ± 3.8
$\Gamma_{\text{Temkin (Low-range)}}$		810 ± 740	2.4 ± 2.2
$\Gamma_{\text{Langmuir (Full)}}$	$\frac{[\text{NH}_4^+]_{\text{salt}}}{k_L(S_{\max} - [\text{NH}_4^+]_{\text{salt}})} \frac{1}{[\text{H}^+]_{\text{DIW}}}$	3530 ± 3200	10 ± 9.3
$\Gamma_{\text{Langmuir (Low-Range)}}$		2730 ± 2500	7.7 ± 7.3

240




Figure 4: Soil emission potentials using the standard, salt extraction, Temkin, Langmuir and DIW extraction approaches, arranged by CEC.

245 **4. Discussion**

The conventional approach for calculating emission potentials in soils lacks a solid theoretical foundation, and ought to be applied with caution. Empirically, this approach has performed poorly, with several studies reporting that calculated values of Γ_{STD} are unrealistically high and do not correspond well to measured or modelled emission fluxes (Cooter et al., 2010; Flechard et al., 2013; Neftel et al., 1998; Nemitz et al., 2001; Xu et al., 2022). Our assessment that the conventional method overestimates the soil emission potential by at least a factor of 5 is similar to findings reported by Nemitz et al., (2001) and Cooter et al., (2010) who reported needing to reduce their soil emission potentials by a factor of 6.66 and 2–3 respectively for their modelled predictions to match their flux observations. We believe that our empirical treatment of this system with the Langmuir and Temkin equations provides a more reasonable approach to estimate the soil emission potential. Additionally, in our view, the ‘like-with-like’ extraction approach (e.g. Γ_{SALT} or Γ_{DIW}) has a more mechanistic basis than the conventional approach involving

255 dissimilar extraction solutions. A few recent studies have reported on a similar approach using a Langmuir adsorption model (Alnsour, 2020; Guo et al., 2022b), as well as by matching the NH_4^+ and pH extraction solutions (Wu et al., 2023), showing that these methods can be feasibly implemented into an environmental sampling campaign. Venterea et al., (2015) reported that soil ammonium partitioning could be effectively modelled using a modified version of the Langmuir isotherm equation of the form $S = \frac{S_{\max}C}{K+C}$, where K (mg L^{-1}) is the aqueous concentration at which $S = \frac{1}{2} S_{\max}$. The Venterea equation is equivalent
260 to the Langmuir isotherm when $K = \frac{1}{K_L}$; for the parameters determined in this study, $\frac{1}{2}$ the saturation capacity is reached at exactly $C = \frac{1}{K_L} = K$, indicating that for our analysis the Langmuir and Venterea equations are equivalent.

How does this approach compare to the implementation of bidirectional exchange of NH_3 in chemical transport models? Widely used models such as GEOS-CHEM and the EPA's CMAQ model soil emission potentials mechanistically, rather than being based on emission potentials prescribed using land-use categories (Bash et al., 2013; Pleim et al., 2013b;
265 Pleim, Ran, Appel, Shephard, et al., 2019; Zhu et al., 2015b). In both models, the exchangeable soil ammonia has been parameterized as the volumetric molar concentration of ammonia in the top 1 – 5 cm divided by the volumetric water content in the soil, or $[\text{NH}_4] = \frac{\text{NH}_4 \text{ mol m}^{-3}}{\theta \text{ m}^3 \text{ m}^{-3}}$ (Massad et al., 2010b). In this parameterization, the entire soil ammonia content of the top layer of soil is treated as being dissolved into the soil water, thus a lower water content would result in a higher concentration of ammonia, and consequently a higher emission potential. (The parameter used by these models for the
270 resistance to ammonia emissions from soil is inversely proportional to soil moisture, such that the actual emission of ammonia from soil would still be proportional to soil water content (Pleim et al., 2013b).) In our model of the system, most ammonium is present adsorbed to soils, with the aqueous concentration controlled by the equilibrium between adsorbed and aqueous ammonium, increasing the soil water content would thus allow more ammonium to enter the aqueous phase to maintain the equilibrium concentration. (Increased water content could also result in nutrient runoff, reducing the soil
275 ammonium content, or increased mineralization, increasing the soil ammonium content.) In the subsequent development of the bidirectional model coupled with an agricultural ecosystem model, Pleim et al., (2019) noted that reducing the ammonia available for emission by implementing the Langmuir adsorption isotherm approach proposed by Venterea et al. (2015) appeared to lead to unexpectedly low fractions of ammonia available for exchange. It may be that for recently fertilized soils, the applied fertilizer is not in contact with sufficient soil for this equilibrium relation to apply. An updated version of CMAQ
280 (v5.2.1) uses a Langmuir-derived approach described in Venterea et al. (2015) to predict NH_3 bi-directional exchange and Kelly et al., (2019) explores its indirect impact on $\text{PM}_{2.5}$ composition across the U.S., suggesting that the reduced emission potential from our approach is compatible with atmospheric agro-ecosystem modelling.

5. Conclusion

This work evaluated the Langmuir, Freundlich, Temkin, and Toth adsorption isotherm equations as applied to environmental soil samples and NH_4^+ partitioning. We determined that under ideal conditions the Toth equation was the most effective of these equations at fitting soil adsorption curves, but that the Temkin equation was most effective at predicting the adsorption behaviour of soil samples with minimal additional characterization required, and was effective over both low concentration ranges and the saturation concentration range. Applying this method to a series of environmental soil samples, we determined that the conventional method for directly measuring soil emission potentials may overestimate them by a factor of 5 – 20 (relative to using the Langmuir/Temkin equations respectively). Of the two adsorption equations, the Temkin equation fit the data better, with a significantly lower AIC (-222 vs -138 for the Langmuir equation). Comparing these empirical equations with an alternative approach for determining emission potential based on extracting NH_4^+ and pH with the same extraction solutions (i.e. DIW/DIW or Salt/Salt) showed that the Temkin equation fit using the full adsorption range agreed well with the DIW/DIW ratio method, but was significantly different when using the low-range fitting parameters.

295 Appendix A




Figure A1: Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, ·-·-), and Toth (orange, ···) equations for soil samples collected from a) King's College Circle, b) the University of Toronto Scarborough forest, c) High Park, and d) Queen's Park.

Table A1: Comparison of the residual standard errors, and the range of fitted parameters for adsorption curves fit to the un-pooled data. Parameters given in parentheses are reported for comparison with the parameters in Table 2. Goodness of fit is reported using the residual standard error (RSE) rather than AIC to avoid confusion between ranges of AICs for each equation.

Equation	Number of fitted parameters	RSE	Parameter 1	Mean \pm sd	Parameter 2	Mean \pm sd	Exponential Factor
Langmuir	1	17.5 – 25.7%	K_L	$9.33E-4 \pm 2.0E-4$	-	-	-
Freundlich	2	5.72 – 17.6%	K_F	197 ± 60 ($S_{max} * [0.053 \pm 0.012]$)	-	-	0.3064 ± 0.023
Temkin	2	3.0 – 16.1%	K_T	$1.34E-2 \pm 5.1E-3$	q_T	$S_{max} * [0.183 \pm 0.013]$	-
Toth	3	0.9 – 6.5%	K_{To}	3.75 ± 1.3	b	9960 ± 5590 ($S_{max} * [2.42 \pm 0.72]$)	0.276 ± 0.054

305

Table A2: Residual standard errors when applying average fit parameters from the pooled and (separately fit) training data to the test set soil samples, for i) a CEC of 25; ii) the experimentally measured CEC; iii) when the data is fit using the fitting algorithm

Sample Location	CEC (cmole kg^{-1})	NH_4^+ (mg kg^{-1})	pH	RSE (%) using estimated fit parameters			
				Langmuir	Freundlich	Temkin	Toth
High Park	10.95	2.906	7.04	280 (285)	330 (340)	320 (330)	330 (420)
				26 (26)	22 (26)	16 (18)	14 (28)
				25	19	14	8.8
Corktown	25.3	3.552	7.82	33 (33)	25 (31)	22 (26)	9.5 (39)
				34 (34)	28 (34)	25 (28)	9.7 (43)
				29	15	15	3.8
Riverdale Park East	37.25	2.835	7.42	64 (63)	62 (58)	66 (63)	84 (62)
				35 (35)	19 (25)	29 (31)	19 (39)
				31	11	24	23
Geometric Mean RSE		Estimated		84 (84)	80 (85)	77 (81)	64 (100)
		Estimated with measured CEC		31 (31)	21 (28)	21 (25)	14 (36)
		Fitted Equation		28.3	14.8	17.0	9.2

Table A3: Sample information for the adsorption test set

Sample	CEC (cmole kg ⁻¹)	Bulk NH ₄ ⁺ (mg kg ⁻¹)	pH	Extractant Solution (mg L ⁻¹)	Soil Mass (g)	Sum of displaced cations (cmole kg ⁻¹)	NH ₄ ⁺ adsorbed (mg kg ⁻¹)
High Park	10.95	2.906	7.04	18.04	1.006	0.989	178.4
				36.08	1.0124	0.949	171.2
				144.32	0.9953	1.951	351.9
				721.6	0.9976	4.028	726.6
				3608	0.9539	7.413	1337
				18040	1.0002	10.81	1950
Corktown	25.3	3.552	7.82	18.04	1.0081	1.793	323.5
				36.08	1.0095	2.222	400.9
				144.32	0.9944	4.228	762.8
				721.6	1.0013	9.064	1635
				3608	1.0131	15.54	2804
				18040	1.002	24.39	4400
Riverdale Park East	37.25	2.835	7.42	18.04	0.9948	4.447	802.2
				36.08	1.0193	4.861	876.8
				144.32	1.0151	6.690	1207
				721.6	1.0304	11.79	2127
				3608	0.9953	23.53	4244
				18040	1.027	36.26	6541

310 Author contribution statement

MGD: Investigation, Formal analysis, Writing – Original Draft; KY: Investigation, Methodology; JGM: Supervision, Funding acquisition, Writing – Review & Editing.

Competing interests statement

The authors declare that they have no conflict of interest.

315 **Funding statement**

This research was supported by a Natural Science and Engineering Research Council (NSERC) Discovery grant and a Grants and Contributions agreement GCXE19S016 with Environment and Climate Change Canada held by Jennifer Murphy. Matthew Davis held a Walter C. Sumner Memorial Fellowship while conducting this research. An Undergraduate Summer Research Award from NSERC supported Kevin Yan during this work.

320 **Acknowledgements**

We thank our colleagues Professor Myrna Simpson and Jenny Oh (University of Toronto) for their helpful discussions. We also thank the University of Toronto ANALEST facility staff for their technical assistance.

References

- Alnsour, N. I. (2020). *Bi-Directional Exchange of Ammonia from Soils in Row Crop Agro-ecosystems*. North Carolina State University.
- 325
- Bache, B. W. (1976). The measurement of cation exchange capacity of soils. *Journal of the Science of Food and Agriculture*, 27(3), 273–280. <https://doi.org/10.1002/jsfa.2740270313>
- Bash, J. O., Cooter, E. J., Dennis, R. L., Walker, J. T., & Pleim, J. E. (2013). Evaluation of a regional air-quality model with bidirectional NH₃ exchange coupled to an agroecosystem model. *Biogeosciences*, 10(3), 1635–1645.
- 330 <https://doi.org/10.5194/bg-10-1635-2013>
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Hoek, K. W. Van Der, Olivier, J. G. J., Tg, N., & Tg, N. (1997). A global high-resolution emission inventory for ammonia. *Global Biogeochemical Cycles*, 11(4), 561–587.
- Chu, K. H. (2021). Revisiting the Temkin Isotherm: Dimensional Inconsistency and Approximate Forms. *Industrial and Engineering Chemistry Research*, 60(35), 13140–13147. <https://doi.org/10.1021/acs.iecr.1c01788>
- 335 Cooter, E. J., Bash, J. O., Walker, J. T., Jones, M. R., & Robarge, W. (2010). Estimation of NH₃ bi-directional flux from managed agricultural soils. *Atmospheric Environment*, 44(17), 2107–2115. <https://doi.org/10.1016/j.atmosenv.2010.02.044>
- Ellis, E. C., Beusen, A. H. W., & Goldewijk, K. K. (2020). Anthropogenic biomes: 10,000 BCE to 2015 CE. *Land*, 9(5), 8–10. <https://doi.org/10.3390/LAND9050129>
- 340 Farquhar, G. D., Firth, P. M., Wetselaar, R., & Weir, B. (1980). On the Gaseous Exchange of Ammonia between Leaves and the Environment: Determination of the Ammonia Compensation Point. *Plant Physiology*, 66(4), 710–714. <https://doi.org/10.1104/pp.66.4.710>
- Flechard, C. R., Fowler, D., Sutton, M. A., & Cape, J. N. (1999). A dynamic chemical model of bi-directional ammonia exchange between semi-natural vegetation and the atmosphere. In *Quarterly Journal of the Royal Meteorological Society* (Vol. 125, Issue 559). <https://doi.org/10.1002/qj.49712555914>
- 345 Flechard, C. R., Massad, R.-S., Loubet, B., Personne, E., Simpson, D., Bash, J. O., Cooter, E. J., Nemitz, E., & Sutton, M. A. (2013). Advances in understanding, models and parameterizations of biosphere-atmosphere ammonia exchange. *Biogeosciences*, 10(7), 5183–5225. <https://doi.org/10.5194/bg-10-5183-2013>
- Freundlich, H. (1909). Kapillarchemie: eine Darstellung der Chemie der Kolloide und verwandter Gebiete. In *Akademische Verlagsgesellschaft*. Verlag der Akademischen Verlagsgesellschaft. <https://doi.org/10.1038/130866a0>
- 350 Guo, X., Pan, D., Daly, R. W., Chen, X., Walker, J. T., Tao, L., McSpirtt, J., & Zondlo, M. A. (2022a). Spatial heterogeneity of ammonia fluxes in a deciduous forest and adjacent grassland. *Agricultural and Forest Meteorology*, 326(July), 109128. <https://doi.org/10.1016/j.agrformet.2022.109128>

- 355 Guo, X., Pan, D., Daly, R. W., Chen, X., Walker, J. T., Tao, L., McSpirtt, J., & Zondlo, M. A. (2022b). Spatial heterogeneity of ammonia fluxes in a deciduous forest and adjacent grassland. *Agricultural and Forest Meteorology*, 326(July), 109128. <https://doi.org/10.1016/j.agrformet.2022.109128>
- Kelly, J. T., Koplitz, S. N., Baker, K. R., Holder, A. L., Pye, H. O. T., Murphy, B. N., Bash, J. O., Henderson, B. H., Possiel, N. C., Simon, H., Eyth, A. M., Jang, C., Phillips, S., & Timin, B. (2019). Assessing PM2.5 model performance for the conterminous U.S. with comparison to model performance statistics from 2007-2015. *Atmospheric Environment*, 360 214(May), 116872. <https://doi.org/10.1016/j.atmosenv.2019.116872>
- Kempers, A. J., & Zweepers, A. (1986). Ammonium determination in soil extracts by the salicylate method. *Communications in Soil Science and Plant Analysis*, 17(7), 715–723. <https://doi.org/10.1080/00103628609367745>
- Langmuir, I. (1916). The Constitution and Fundamental Properties of Solids and Liquids. Part I. Solids. *Journal of the American Chemical Society*, 38(11), 2221–2295.
- 365 Li, K. yi, Zhao, Y. yuan, Yuan, X. long, Zhao, H. bing, Wang, Z. hui, Li, S. xiu, & Malhi, S. S. (2012). Comparison of Factors Affecting Soil Nitrate Nitrogen and Ammonium Nitrogen Extraction. *Communications in Soil Science and Plant Analysis*, 43(3), 571–588. <https://doi.org/10.1080/00103624.2012.639108>
- Massad, R. S., Nemitz, E., & Sutton, M. A. (2010a). Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmospheric Chemistry and Physics*, 10(21), 10359–10386. [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-10-10359-2010)
- 370 10-10359-2010
- Massad, R. S., Nemitz, E., & Sutton, M. A. (2010b). Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmospheric Chemistry and Physics*, 10(21), 10359–10386. [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-10-10359-2010)
- 10-10359-2010
- Neftel, A., Blatter, A., Gut, A., Högger, D., Meixner, F., Ammann, C., & Nathaus, F. J. (1998). NH₃ soil and soil surface gas measurements in a triticale wheat field. *Atmospheric Environment*, 32(3), 499–505. [https://doi.org/10.1016/S1352-](https://doi.org/10.1016/S1352-2310(97)00162-3)
- 375 2310(97)00162-3
- Nemitz, E., Milford, C., & Sutton, M. A. (2001). A two layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia. *Q. J. R. Meteorol. Soc.*, 127, 815–833.
- Nemitz, E., Sutton, M. A., Schjoerring, J. K., Husted, S., & Paul Wyers, G. (2000). Resistance modelling of ammonia exchange over oilseed rape. *Agricultural and Forest Meteorology*, 105(4), 405–425. [https://doi.org/10.1016/S0168-](https://doi.org/10.1016/S0168-1923(00)00206-9)
- 380 1923(00)00206-9
- Pleim, J. E., Bash, J. O., Walker, J. T., & Cooter, E. J. (2013a). Development and evaluation of an ammonia bidirectional flux parameterization for air quality models. *Journal of Geophysical Research: Atmospheres*, 118(9), 3794–3806. <https://doi.org/10.1002/jgrd.50262>
- 385 Pleim, J. E., Bash, J. O., Walker, J. T., & Cooter, E. J. (2013b). Development and evaluation of an ammonia bidirectional flux parameterization for air quality models. *Journal of Geophysical Research: Atmospheres*, 118(9), 3794–3806. <https://doi.org/10.1002/jgrd.50262>

- 390 Pleim, J. E., Ran, L., Appel, W., & Shephard, M. W. (2019). *New Bidirectional Ammonia Flux Model in an Air Quality Model Coupled With an Agricultural Model* *Journal of Advances in Modeling Earth Systems*, 2934–2957. <https://doi.org/10.1029/2019MS001728>
- Pleim, J. E., Ran, L., Appel, W., Shephard, M. W., & Cady-Pereira, K. (2019). New Bidirectional Ammonia Flux Model in an Air Quality Model Coupled With an Agricultural Model. *Journal of Advances in Modeling Earth Systems*, 11(9), 2934–2957. <https://doi.org/10.1029/2019MS001728>
- 395 Spiess, A. N., & Neumeyer, N. (2010). An evaluation of R2 as an inadequate measure for nonlinear models in pharmacological and biochemical research: A Monte Carlo approach. *BMC Pharmacology*, 10, 1–11. <https://doi.org/10.1186/1471-2210-10-6>
- Sutton, M. A., Erisman, J. W., Dentener, F., & Möller, D. (2008). Ammonia in the environment: From ancient times to the present. *Environmental Pollution*, 156(3), 583–604. <https://doi.org/10.1016/j.envpol.2008.03.013>
- 400 Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R., ... de Vries, W. (2013). Towards a climate-dependent paradigm of ammonia emission and deposition. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621). <https://doi.org/10.1098/rstb.2013.0166>
- 405 Sutton, M. A., Schjørring, J. K., & Wyers, G. P. (1995). Plant-atmosphere exchange of ammonia. *Philos. T. R. Soc. Lond., A*, 351(29), 261–276.
- Temkin, M. I., & Pyzhev, V. (1940). Kinetics of ammonia synthesis on promoted iron catalyst. *Acta Phys. Chim. USSR*, 12(327).
- Tóth, J. (1995). Uniform interpretation of gas/solid adsorption. *Advances in Colloid and Interface Science*, 55(C), 1–239. [https://doi.org/10.1016/0001-8686\(94\)00226-3](https://doi.org/10.1016/0001-8686(94)00226-3)
- 410 Van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Goldewijk, C. G. M. K., & Lelieveld, J. (2001). A 1° × 1° resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990. *Global Biogeochemical Cycles*, 15(4), 909–928. <https://doi.org/10.1029/2000GB001265>
- Venterea, R. T., Clough, T. J., Coulter, J. A., & Breuillin-Sessoms, F. (2015). Ammonium sorption and ammonia inhibition of nitrite-oxidizing bacteria explain contrasting soil N2O production. *Scientific Reports*, 5(April), 1–15. <https://doi.org/10.1038/srep12153>
- 415 Vogeler, I., Cichota, R., Snow, V. O., Dutton, T., & Daly, B. (2011). Pedotransfer Functions for Estimating Ammonium Adsorption in Soils. *Soil Science Society of America Journal*, 75(1), 324–331. <https://doi.org/10.2136/sssaj2010.0192>
- Walker, J. T., Chen, X., Wu, Z., Schwede, D., Daly, R., Djurkovic, A., Oishi, A. C., Edgerton, E., Bash, J., Knoepp, J., Puchalski, M., Iiames, J., & Miniati, C. F. (2023). Atmospheric deposition of reactive nitrogen to a deciduous forest in 420 the southern Appalachian Mountains. *Biogeosciences*, 20(5), 971–995. <https://doi.org/10.5194/bg-20-971-2023>

- Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G., & Hems, R. (2014). Soil-atmosphere exchange of ammonia in a non-fertilized grassland: Measured emission potentials and inferred fluxes. *Biogeosciences*, *11*(20), 5675–5686. <https://doi.org/10.5194/bg-11-5675-2014>
- 425 Wu, Z., Walker, J. T., Oishi, A. C., Duman, T., Katul, G., Chen, X., Schwede, D., Bash, J., & Iiams, J. (2023). Estimating source-sink distributions and fluxes of reactive nitrogen and sulfur within a mixed forest canopy. *Agricultural and Forest Meteorology*, *333*(June 2022), 109386. <https://doi.org/10.1016/j.agrformet.2023.109386>
- Xu, M., Umehara, M., Sase, H., & Matsuda, K. (2022). Ammonia fluxes over an agricultural field in growing and fallow periods using relaxed eddy accumulation. *Atmospheric Environment*, *284*(April), 119195. <https://doi.org/10.1016/j.atmosenv.2022.119195>
- 430 Zhang, L., Wright, L. P., & Asman, W. A. H. (2010). Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models. *Journal of Geophysical Research Atmospheres*, *115*(20). <https://doi.org/10.1029/2009JD013589>
- Zhu, L., Henze, D., Bash, J., Jeong, G. R., Cady-Pereira, K., Shephard, M., Luo, M., Paulot, F., & Capps, S. (2015a). Global evaluation of ammonia bidirectional exchange and livestock diurnal variation schemes. *Atmospheric Chemistry and Physics*, *15*(22), 12823–12843. <https://doi.org/10.5194/acp-15-12823-2015>
- 435 Zhu, L., Henze, D., Bash, J., Jeong, G. R., Cady-Pereira, K., Shephard, M., Luo, M., Paulot, F., & Capps, S. (2015b). Global evaluation of ammonia bidirectional exchange and livestock diurnal variation schemes. *Atmospheric Chemistry and Physics*, *15*(22), 12823–12843. <https://doi.org/10.5194/acp-15-12823-2015>

440

