

Evaluating adsorption isotherm models for determining the partitioning of ammonium between soil and soil-pore water in environmental soil samples

Matthew G. Davis*, Kevin Yan, Jennifer G. Murphy

5 Department of Chemistry, University of Toronto, Toronto, M5S 3H6, Canada

Correspondence to: Matthew G. Davis (mg.davis@mail.utoronto.ca)

Abstract: Ammonium in soil pore water is thought to participate in bidirectional exchange with the atmosphere; however, common soil nutrient analysis methods determine the bulk quantity of ammonium associated with the soil particles, rather than determining the aqueous ammonium concentration. Previous works have applied the Langmuir and Freundlich isotherm equations to ammonium-enriched soils to estimate partitioning, but this may not be representative of conditions in natural, unmanaged soils. In this work, environmental soil samples were collected from greenspaces in Toronto and used to evaluate several commonly used adsorption isotherm equations, including the Langmuir, Freundlich, Temkin and Toth equations, to determine their applicability in lightly managed and non-fertilized soils. We then compare ammonia emission potentials (a quantity predicting the propensity of ammonia to volatilize from a liquid reservoir) determined using a conventional high-salt extraction procedure to determine the soil ammonium content to that modelled using the Temkin and Langmuir equations, and demonstrate that conventional approaches may overestimate emission potentials from soils by a factor of 5 – 20.

Key words: ammonium soil adsorption; emission potential;

1. Introduction:

1.1. Contextualizing the significance of ammonium partitioning in soils

20 Globally, emissions of reduced nitrogen compounds (NH_3) make up as much as half of the global atmospheric reactive nitrogen sources (Flechar et al., 2013). Of the NH_3 budget, approximately two thirds of emissions are related to anthropogenic agricultural activities, with natural sources being responsible for only 15 – 20% of emissions (Bouwman et al., 1997; Sutton et al., 2008, 2013; Van Aardenne et al., 2001). Following emission, NH_3 tends to partition into fine particulate matter or deposit via wet or dry deposition on a timescale of hours to days. NH_3 is understood to engage in bidirectional exchange throughout the earth system, with NH_3 depositing, or volatilizing depending on local environmental conditions (Farquhar et al., 1980; Flechar et al., 1999, 2013; Guo et al., 2022; Nemitz et al., 2000; Sutton et al., 1995; Walker et al., 2023; Wentworth et al., 2014; Zhang et al., 2010). The bidirectional exchange of NH_3 between soils and the atmosphere has been considered important to its overall budget, particularly in remote areas, but research on the mechanisms of NH_3 partitioning in soils between adsorbed

(inaccessible) and aqueous (accessible) NH_3 often focuses on fertilized croplands with substantial concentrations of NH_3 present (Venterea et al., 2015; Vogeler et al., 2011). Consequently, NH_3 volatilization models may parameterize all or most of the soil NH_3 as being readily able to exchange with the atmosphere, which may be reasonable for recently fertilized soils, but not for unmanaged soils (Massad et al., 2010; Pleim et al., 2013; Zhu et al., 2015). However, cropland is estimated to make up less than 15% of the Earth's land area, while unfertilized or irregularly fertilized natural, semi-natural or pastoral land reflects nearly three quarters of terrestrial surfaces (Ellis et al., 2020). The short atmospheric lifetime of NH_3 makes it important to understand the exchange of NH_3 over all types of surfaces, despite agricultural cropland being the most globally significant source.

1.2. Importance of developing an ammonium adsorption partitioning model

The propensity for ammonia to volatilize from liquid reservoirs is parameterized by the emission potential (Γ), which is calculated as the ratio of aqueous NH_4^+ to H^+ . However, NH_4^+ in soils is not only present as $\text{NH}_4^+_{(\text{aq})}$ in soil pore water, but also present as $\text{NH}_4^+_{(\text{ads})}$ adsorbed to soil particles. Because soils tend to exist with both permanent and pH-dependent negative charges (Bache, 1976), NH_4^+ ions compete for adsorption against other cations in solution. The total quantity of cations that can adsorb to the soil particles is termed the cation exchange capacity (CEC, typically reported as centimoles positive charge/kg). Traditional methods for determining NH_4^+ in soil—intended for nutrient analysis in which the total ammoniacal nitrogen is more important than the partitioning between $\text{NH}_4^+_{(\text{ads})}$ and $\text{NH}_4^+_{(\text{aq})}$ —use high concentration salt solutions to displace all cations from the soil. As a result, these approaches do not distinguish between $\text{NH}_4^+_{(\text{aq})}$ and $\text{NH}_4^+_{(\text{ads})}$ (Li et al., 2012) and therefore likely overestimate the emission potential of soils. In this manuscript, we explore a variety of adsorption isotherm models with the goal of identifying a simple approach to relate the total quantity of NH_4^+ in soil to the aqueous fraction of $\text{NH}_4^+_{(\text{aq})}$ that can participate in bidirectional exchange with the atmosphere.

1.3. Adsorption isotherm equations

The adsorption behavior of molecules to surfaces is complex and dependent on the properties of both the surface and the adsorbed molecules. Numerous adsorption equations have been proposed, based on both theoretical and empirical models; however, as the partitioning of NH_4^+ between soil pore water and soil particles is complex and influenced by many other simultaneous equilibria, we consider each of the examined equations as being empirically determined only, rather than based on a theoretical treatment of the system. Previous studies have sought to develop an adsorption model for NH_4^+ in soils based on the Langmuir (Alnsour, 2020; Venterea et al., 2015) and Freundlich equations (Vogeler et al., 2011), but have focused on croplands and other agricultural soils. In this work, in addition to the Langmuir and Freundlich equations, we also investigate the application of the Temkin and Toth adsorption equations to NH_4^+ sorption in soils. Each of these equations (as formulated in Table 1) represents the adsorbed NH_4^+ concentration (S , mg kg^{-1}) in terms of the NH_4^+ concentration in solution (C , mg L^{-1}). Except for the Freundlich equation, each of these equations incorporate a saturation point or maximum adsorption capacity

60 (S_{\max} , mg kg⁻¹), this work treats S_{\max} as an empirically measured property, equivalent to the CEC (converted to mg of NH₄⁺ kg⁻¹ soil), rather than as a calculated fitting parameter.

Table 1: Adsorption isotherm equations applied in this work and their parameters

Adsorption Isotherm Model	Equation [†]	Units	Reference
Langmuir	(1) $S = \frac{S_{\max}k_L C}{1+k_L C}$	S_{\max} (mg kg ⁻¹), K_L (L mg ⁻¹)	(Langmuir, 1916)
Freundlich	(2) $S = k_F C^{n_F}$	k_F , n_F (dimensionless)	(Freundlich, 1909)
Temkin*	(3) $S = q_T \ln(1 + K_T C)$	q_T (product of S_{\max} (mg kg ⁻¹) and $\frac{RT}{b}$ (dimensionless)), K_T (L mg ⁻¹)	(Temkin & Pyzhev, 1940)
Toth	(4) $S = \frac{bc}{(K_{To} + C^{n_T})^{1/n_T}}$	b (product of S_{\max} (mg kg ⁻¹) and a dimensionless scaling factor), K_{To} (mg L ⁻¹), n_T (dimensionless)	(Tóth, 1995)

*The Temkin model is also given as $S = \frac{RT}{b} \ln K_T C$, see (Chu, 2021) for this formulation

65 [†] Where S is the concentration of NH₄⁺ adsorbed to soil particles (mg kg⁻¹), C is the concentration of NH₄⁺ in solution (mg L⁻¹), and S_{\max} is the maximum adsorption capacity (mg kg⁻¹), which was determined empirically as equivalent to the measured cation exchange capacity of the soil. This work treats the remaining parameters as empirical constants only.

2. Methods:

2.1. Soil Collection:

70 Soil samples were collected from 24 greenspace (parks, urban forest, roadside-sites, etc.) locations across Toronto. Samples were collected only on days preceded by at least two days without precipitation and were collected by inserting a 7.5 cm internal diameter steel tube into the ground to a depth of 5 – 10 cm and recovering a soil core by removing the tube from the ground. The soil cores were mixed, and sieved immediately, and transported back to the lab for analysis. Samples analyzed for NH₄⁺ content were always analyzed immediately to avoid potential artefacts from freezing, samples analyzed for CEC or adsorption curves were frozen for storage prior to analysis. Soil samples were collected from eight locations in Fall 2021 as a training set for developing the model. A subsequent 16 soil samples from across Toronto were collected in Spring/Summer 2023 to be used to evaluate the effectiveness of applying the model to uncharacterized soils, and to determine the impact on soil emission potentials.

2.2. Soil analysis:

2.2.1. Cation Exchange Capacity determination

80 The CEC was determined using the inductively-coupled plasma optical emission spectroscopy (ICP-OES) cation sum method (Bache, 1976). Briefly, 1 g of soil was measured out and mixed with 25 mL of 1 M NH₄Cl, shaken, and refrigerated for 36 - 48 hours to settle. The supernatant was filtered using 0.2 µm syringe filters, diluted 50-fold using volumetric glassware and analyzed for the common soil-associated exchangeable cations, Na⁺, K⁺, Mg²⁺, and Ca²⁺. The ICP-OES (iCAP Pro,

ThermoFisher Scientific, Waltham, USA) was calibrated using a commercially available mixed standard of 6 cations (Li^+ , Na^+ , NH_4^+ , K^+ , Mg^{2+} , Ca^{2+}) (Dionex Cation II, ThermoFisher Scientific, Waltham, USA). Another soil-associated cation that can contribute to CEC is Al^{3+} , our initial measurements screened for Al, but we did not detect it in solution. The CEC was determined for all 24 collected samples and used to select three of the soil samples from the 16 collected in 2023 to be used as a test set for the developed model.

2.2.2. Adsorption curve characterization

The determination of the adsorption behavior of NH_4^+ was performed using a modified version of the procedure described by Venterea et al. (2015), combined with the ICP-OES cation sum method (Bache, 1976). Briefly, a series of batch equilibrium samples were prepared by mixing 1 g of soil with 25 mL aliquots of NH_4Cl solutions with concentrations ranging from 2.5 – 1000 mM. The samples were shaken, refrigerated for 36 – 48 hours to settle, filtered with 0.2 μm syringe filters, diluted 50-fold with volumetric glassware, and analyzed using ICP-OES. The quantity of NH_4^+ adsorbed onto each soil was inferred based on the displaced Na^+ , Mg^{2+} , Ca^{2+} and K^+ ions measured in solution. In the cation sum method, the total quantity of adsorbable cations is determined by saturating the soil with an index cation (in this procedure, and generally, NH_4^+ from NH_4Cl), which displaces the exchangeable cations on the soil's adsorption sites. Thus, the displaced cations measured for each NH_4Cl solution concentration are representative of the quantity of NH_4^+ adsorbing onto the soil. Adsorption curves were determined for the original eight soil samples in the training set, as well as the three selected soil samples for the test set.

2.2.3. pH and NH_4^+ determination

pH and NH_4^+ content was determined for the 16 soil samples collected in 2023. pH was determined for each soil by mixing soil with ultra-pure (18 $\text{M}\Omega$ cm) water (DIW) in a 1:2 ratio, and then measuring the pH of the slurry by immersing a pH electrode (Hach Company, Loveland, USA) until a stable pH reading was obtained. NH_4^+ was extracted from the soil using a 2 M KCl extraction solution. 2.5 g of soil was mixed with 25 mL of the extraction solution, shaken, and refrigerated for 36 hours to allow suspended solids to settle out of solution. Afterward, the supernatant was filtered using 0.2 μm PES membrane syringe filters. The soil NH_4^+ was quantified using the indophenol-blue salicylate method (Kempers & Zweers, 1986). Briefly, two reagent solutions, A and B were prepared: Reagent A consisted of a solution of 1 M sodium salicylate and 100 mg L^{-1} sodium nitroprusside, while Reagent B consisted of a solution of 1 M NaOH and 0.12% by volume of 5% available chloride NaOCl. Soil extracts were prepared for analysis by adding 0.6 mL of reagent A to 2 mL of soil extract, followed by the addition of 1.4 mL of reagent B. The mixtures were then stored for 2 hours in the dark for color development, and then quantified using UV-VIS spectrometry (Lambda 365, Perkin-Elmer, Waltham, USA) at 649 nm.

Conventionally, the soil emission potential is calculated as a function of the soil pH (H^+_{DIW}) and salt solution-extracted NH_4^+ ($\text{NH}_4^+_{\text{SALT}}$). In addition to investigating whether adsorption isotherm equations could be applied to estimate $\text{NH}_4^+_{\text{(aq)}}$ from the total soil NH_4^+ , we investigated the impact of calculating the emission potential using a 'like-with-like' ratio of H^+_{DIW} with NH_4^+ extracted using DIW as the solvent, as well as the ratio of $\text{NH}_4^+_{\text{SALT}}$ to H^+ determined from a salt-solution:soil

slurry. Consequently, the pH was also determined as described above, but using a 0.01 M CaCl₂ solution in place of DIW, while soil NH₄⁺ was also determined as described above, but using DIW as the solvent.

2.3. Data analysis

2.3.1. Characterizing adsorption parameters using a training and a test set

120 Curve-fitting was done in R using the nls function to fit our experimental data for the eight samples in our training set to the Langmuir, Freundlich, Temkin, and Toth equations. Goodness of fit was evaluated by calculating the Akaike information criterion (AIC) using the AIC function from the R *stats* package. The AIC is calculated using equation (5):

$$(5) \text{ AIC} = 2K - 2\ln(L)$$

Where K is the number of independent variables, and L is the log-likelihood estimate. The log-likelihood estimate can be
125 extracted directly from nls objects fitted in R using the AIC function.

Fitting parameters for each equation were determined by pooling all the experimental data, standardizing each adsorption curve by the maximum adsorption achieved (i.e., all curves went from 0 to 1), and then fitting each equation from Table 1 to those curves.

130 To validate the effectiveness of these equations when applied to uncharacterized soil samples, we selected three soil samples from the 16 soil samples collected in 2023 to form a test set, these samples were selected by choosing the soil samples with the lowest (10.9) and highest CECs (37.2), and a soil with an average CEC (25.3). As the original training set mostly consisted of samples with CECs from 20 – 30 (with two samples with CECs of 7.6 and 16), we chose two samples that were significantly different than the average training set sample, as well as one similar sample to determine whether the fitting
135 parameters could be used for ‘extreme’ samples, or only for samples similar to the training set. The test set was characterized in the same way as the training set, and was then modelled using three approaches:

- i. Using the average CEC for all the soil samples of 25 cmole kg⁻¹ (S_{max} of 4500 mg kg⁻¹), and the training set parameters.
- ii. Using the measured CEC for each soil sample to calculate S_{max}, and the training set parameters.
- iii. Using the fitting algorithm as described in Sect.2.3.1 to determine the least squares fit for each equation to the
140 experimental data.

2.3.2. Emission potential determination

The NH₃ emission potential is a quantity calculated as the ratio of aqueous NH₄⁺ to H⁺ (Eq.(6)).

$$(6) \Gamma = \frac{[NH_4^+]}{[H^+]}$$

Commonly, for soils this would be calculated using the total NH₄⁺ (determined using a salt solution extraction) and the pH
145 measured using an extraction with deionized water. We denote this as Γ_{STD} , corresponding to $\frac{[NH_4^+]_{salt}}{[H^+]_{DIW}}$. The pH may also be

measured in a (less concentrated) salt solution, which we denote as Γ_{SALT} , corresponding to $\frac{[\text{NH}_4^+]_{\text{salt}}}{[\text{H}^+]_{\text{salt}}}$. Similarly, though we are not aware of this as a common method, NH_4^+ could be determined using a DIW extraction solution, resulting in a third parameterization of the emission potential as Γ_{DIW} , representing $\frac{[\text{NH}_4^+]_{\text{DIW}}}{[\text{H}^+]_{\text{DIW}}}$. Lastly, by applying one of the adsorption isotherm models, the total soil NH_4^+ can be partitioned into $\text{NH}_4^+_{(\text{ads})}$ (S) and $\text{NH}_4^+_{(\text{aq})}$ (C), and the emission potential can be calculated using only the NH_4^+ in solution (C). These versions of the emission potential are denoted as Γ -sub-equation-name (e.g., as Γ_{Langmuir} , Γ_{Temkin} etc).

3. Results

3.1. Performance of adsorption isotherm equations applied to an environmental soil training and test set

We evaluated the ability of the Langmuir, Freundlich, Temkin and Toth equations to model the exchange of NH_4^+ between adsorbed and aqueous forms. An adsorption curve was characterized for each soil, the data from each soil adsorption experiment was pooled, and then fit using the R nls function. The adsorption curves and fitting parameters are shown in Figure 1 and Table 2. Additionally, as our interest is ultimately in the performance of these adsorption equations at the lower concentration limit, we refit each equation using only the extraction solutions $\leq 680 \text{ mg L}^{-1} \text{ NH}_4\text{Cl}$. The adsorption curves and fitting parameters under those conditions are shown in Figure 2 and Table 3.

While the Langmuir (Alnsour, 2020; Guo et al., 2022; Venterea et al., 2015) and Freundlich (Vogeler et al., 2011) equations have been previously reported as being effective at modelling NH_4^+ adsorption in soils, we found them to be the least effective of the equations we examined for the full adsorption curves, both over- and under-estimating the adsorbed NH_4^+ concentrations (Figure 1), while the Temkin and Toth equations better fit the experimental data. Computing the Akaike information criterion for these equations results in an AIC of -138, -190, -222, and -249 for the Langmuir, Freundlich, Temkin, and Toth equations, respectively. The absolute value of the AIC is not important, but for a set of models, the model with the lowest AIC is considered the best at fitting the experimental data, indicating an order of Toth>Temkin>Freundlich>Langmuir for model effectiveness. However, fitting only the lower range of the adsorption curves (0 – 40 mM) slightly changed these results, the Toth equation could not be fit by our algorithm, and the resulting AIC values were -93, -157, -152 for the Langmuir, Freundlich, and Temkin equations, respectively, indicating that the Freundlich equation best fit the experimental data. (Note that AIC values for models fit to different datasets should not be directly compared to one another.)




Figure 1: Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, - · - ·), and Toth (orange, · · ·) equations. The curves are plotted using the experimental data from all eight soil adsorption experiments, the y-axis is normalized to the maximum adsorption achieved during each experiment.

175

Table 2: Comparison of goodness of fit and fitting parameters for the selected adsorption equations

Equation	AIC	Parameter 1	Mean ± Standard error	Parameter 2	Mean ± Standard error	Exponential Factor
Langmuir	-138	K_L	$9.29 \cdot 10^{-4} \pm 5.9 \cdot 10^{-5}$	S_{max}	-	-
Freundlich	-190	K_F	$S_{max} * 0.0520 \pm 3.4 \cdot 10^{-3}$	-	-	0.3050 ± 0.0074
Temkin	-222	K_T	$1.33 \cdot 10^{-2} \pm 1.2 \cdot 10^{-3}$	q_T	$S_{max} * 0.180 \pm 4.1 \cdot 10^{-3}$	-

Toth	-249	K_{To}	3.10 ± 0.65	b	$S_{max} * 2.45 \pm 0.35$	0.25 ± 0.027
------	------	----------	-----------------	---	---------------------------	------------------




Figure 2: Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, -- --), and Temkin (green, · - · -) equations. The curves are plotted using the experimental data from all eight soil adsorption experiments, but only using the extraction solutions of ≤ 40 mM NH_4^+ for fitting, the y-axis is normalized to the maximum adsorption achieved during each experiment. The Toth equation could not be fit to the experimental data under these conditions.

180

Table 3: Comparison of goodness of fit and fitting parameters for the selected adsorption equations, data fit only for the ≤ 40 mM NH_4^+ solutions

Equation	AIC	Parameter 1	Mean \pm Standard error	Parameter 2	Mean \pm Standard error	Exponential Factor
Langmuir	-93	K_L	$1.2 \cdot 10^{-3} \pm 8.0 \cdot 10^{-4}$	S_{max}	-	-


Freundlich	-157	K_F	$S_{max} * 0.025 \pm 3.0 \cdot 10^{-3}$	-	-	0.42 ± 0.019
Temkin	-152	K_T	$3.4 \cdot 10^{-2} \pm 6.5 \cdot 10^{-3}$	q_T	$S_{max} * 0.120 \pm 8.5 \cdot 10^{-3}$	-
Toth	Fit did not converge					

185

With our objective being to evaluate how well each equation can fit soils without going through the full characterization procedure, we analyzed the adsorption curves of the low-CEC, medium-CEC and high-CEC soils in our test set using: i) a “typical” CEC of 25, and the fitting parameters from Table 2, ii) the measured CEC and the fitting parameters from Table 2, and iii) by fitting the equations using the least-squares fitting algorithm. Using the first approach, we found that none of the equations could reasonably fit the experimental data when using an incorrect CEC and that each of the equations fit the experimental data reasonably well using the average parameters and the correct CEC (Figure 3). The relative goodness of fit for each equation was the same for the test set as for the training set, i.e. Toth>Temkin>Freundlich>Langmuir. We also tested an alternative approach for calculating each equation’s fitting parameters, in which rather than pooling the normalized data, and then fitting each curve, the curves were fit to each soil adsorption curve separately, and the resulting fitting parameters were then pooled. A full summary of the alternative fitting parameters and the test-set characterization is given in Figure A1 and Tables A2 – A4.

190

195



200

Figure 3: Curve-fitting comparison between the unfitted Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, ·-·-), and Toth (orange, ···) equations using the fitting parameters from Table 3 applied to the test set of the a) Low CEC, b) Moderate CEC, c) High CEC samples (method ii)

Overall, we find that the Toth and Temkin equations best fit the full adsorption curves, while the Freundlich and Temkin equations best fit the low-range adsorption curves. However, both the Freundlich and Toth equations seem to be less robust when applied to this system than the Temkin equation. Firstly, the Freundlich equation, as formulated in Table 2 and Table 3, has an intrinsic dependence on the CEC, which is contrary to the theoretical basis of the Freundlich equation. Using the alternative fitting approach, the Freundlich equation does not require a CEC-dependence, but in that case, it has high error

205

when fitting uncharacterized soils. Similarly, for the Toth equation, the alternative approach results in a significantly different set of fitting parameters, which resulted in the Toth equation quite poorly fitting the non-characterized soils. Additionally, while this may be a limitation of the fitting algorithm we selected, we found the Toth equation to be more difficult to consistently fit to our experimental data, with no solution being found for the low-range concentrations. Consequently, taking all of these factors into consideration, we find that the Temkin equation is most suitable for evaluating NH_4^+ adsorption from uncharacterized soil samples. For comparison with previous studies, in which the Langmuir equation is the most frequently used adsorption equation, we will continue to analyze it in subsequent sections, despite it performing much worse than the Temkin equation for both the low-range and full-range of data.

3.2. Determining aqueous NH_4^+ concentrations using the Langmuir and Temkin equations

To relate the total NH_4^+ measured (m_{NH_4}) to the adsorbed portion (S) and the aqueous portion (C), we define Eq.(7):

$$(7) m_{\text{NH}_4} = S + \frac{wC}{\rho}$$

Where m_{NH_4} is the total mass of NH_4^+ per kg soil (mg kg^{-1}), w the volumetric moisture content of the soil (L water L^{-1} soil), and ρ is the bulk density of the dry soil (kg L^{-1}). By substituting the Langmuir (1) or Temkin (3) equations into Eq.(7), m_{NH_4} can be expressed in terms of C:

$$(8) m_{\text{NH}_4} = \frac{S_{\text{max}}k_L C}{1+k_L C} + \frac{wC}{\rho}$$

$$(9) m_{\text{NH}_4} = q_T \ln(1 + K_T C) + \frac{wC}{\rho}$$

Where S_{max} is the maximum adsorption capacity (mg kg^{-1}); q_T is the product of S_{max} and an empirical fitting constant (mg kg^{-1}); and K_L (L mg^{-1}) and K_T (L mg^{-1}) are empirical fitting constants.

To solve for C, we inverted these equations using *Wolfram Mathematica* (<https://www.wolfram.com/mathematica/>), yielding Eq.(10) (Langmuir) and Eq.(11) (Temkin).

$$(10) C = \frac{-S_{\text{max}} \cdot K_L + K_L \cdot m_{\text{NH}_4} - Z + \sqrt{4K_L \cdot Z \cdot m_{\text{NH}_4} + (S_{\text{max}} \cdot K_L - K_L \cdot m_{\text{NH}_4} + Z)^2}}{2K_L \cdot Z}$$

$$(11) C = \frac{-Z + K_T \cdot q_T \cdot \mathbf{W}\left(e^{K_T \cdot q_T + \frac{m_{\text{NH}_4}}{q_T} \cdot Z}\right)}{K_T \cdot Z}, \text{ where } \mathbf{W}(\mathbf{x}) \text{ is the Lambert W function, and } Z = \frac{w}{\rho}$$

If it is assumed that $m_{\text{NH}_4} \approx S$, then these equations can be simplified to Eq.(12) and Eq.(13) respectively.

$$(12) C = \frac{m_{\text{NH}_4}}{k_L(S_{\text{max}} - m_{\text{NH}_4})}$$

$$(13) C = \frac{e^{\frac{m_{\text{NH}_4}}{q_T}} - 1}{K_T}$$

We tested this assumption and found that for our unfertilized soil samples, the simplified equations have a positive bias of only 0.57 – 1.5% for the aqueous concentration (C), and that for most soils fertilized with the equivalent of up to 300 kg N ha⁻¹, the


235 simplified equations should have a positive bias of less than 5%. Consequently, we used the simplified equations for our analysis.

3.3. Determining environmental soil emission potentials

Soil emission potentials were then calculated using the approaches described in Sect. 2.3.3, corresponding to the ‘standard’ approach, two approaches based on matching the NH_4^+ and pH extraction solutions, and two approaches based on applying the Langmuir and Temkin equations to the measured extract concentrations and pH values. The adsorption equations were calculated using both the low-range and full curve fitting parameters. For the Langmuir calculations, the fit parameters were $K_L = 9.29 \cdot 10^{-4} \text{ L mg}^{-1}$ (full) and $K_L = 1.20 \cdot 10^{-3} \text{ L mg}^{-1}$ (low-range), and S_{max} calculated as the equivalent of the CEC in units of mg kg^{-1} . For the Temkin calculations, the fit parameters were $K_T = 1.33 \cdot 10^{-2} \text{ L mg}^{-1}$, and $q_T (\text{mg kg}^{-1}) = 0.180 \cdot S_{\text{max}}$ (full) and $K_T = 3.40 \cdot 10^{-2} \text{ L mg}^{-1}$, and $q_T (\text{mg kg}^{-1}) = 0.120 \cdot S_{\text{max}}$. Among the approaches, Γ_{STD} results in the highest estimate for the emission potential ($17,000 \pm 19,000 \text{ mol mol}^{-1}$), followed by Γ_{Langmuir} and Γ_{SALT} (2730 – 3530), and Γ_{Temkin} and Γ_{DIW} (810 – 1450) (Table 4). Emission potential is linearly related to equilibrium vapor pressure, at a temperature of 15°C, these emission potentials correspond to equilibrium concentrations of 2.4 – 50 ppb. Although CEC is inversely related to the proportion of NH_4^+ present in the aqueous phase, the emission potentials increase exponentially as a function of the CEC due to the strong positive correlation between soil pH and CEC (Figure 4).

250 **Table 4: Comparison of emission potentials determined using the standard, Langmuir, Temkin, salt extraction and DIW extraction methods. Standard deviations represent the underlying variability in the collected soil samples.**

Method	Equation	Average Emission Potential (mol mol^{-1}) (n = 16)	Equilibrium vapor pressure at 15°C (ppb)
Γ_{STD}	$\frac{[\text{NH}_4^+]_{\text{salt}}}{[\text{H}^+]_{\text{DIW}}}$	$17,000 \pm 19,000$	50 ± 55
Γ_{SALT}	$\frac{[\text{NH}_4^+]_{\text{salt}}}{[\text{H}^+]_{\text{salt}}}$	3300 ± 3100	9.6 ± 9.0
Γ_{DIW}	$\frac{[\text{NH}_4^+]_{\text{DIW}}}{[\text{H}^+]_{\text{DIW}}}$	1450 ± 1900	4.2 ± 5.5
Γ_{Temkin} (Full)	$\frac{e^{\frac{[\text{NH}_4^+]_{\text{salt}}}{q_T} - 1}}{K_T} \frac{1}{[\text{H}^+]_{\text{DIW}}}$	1370 ± 1300	4.0 ± 3.8
Γ_{Temkin} (Low-range)		810 ± 740	2.4 ± 2.2
Γ_{Langmuir} (Full)	$\frac{[\text{NH}_4^+]_{\text{salt}}}{k_L(S_{\text{max}} - [\text{NH}_4^+]_{\text{salt}})} \frac{1}{[\text{H}^+]_{\text{DIW}}}$	3530 ± 3200	10 ± 9.3
Γ_{Langmuir} (Low-Range)		2730 ± 2500	7.7 ± 7.3



255 **Figure 4: Soil emission potentials using the standard, salt extraction, Temkin, Langmuir and DIW extraction approaches, arranged by CEC.**

4. Discussion

The conventional approach for calculating emission potentials in soils lacks a solid theoretical foundation, and ought to be applied with caution. Empirically, this approach has performed poorly, with several studies reporting that calculated values of Γ_{STD} are unrealistically high and do not correspond well to measured or modelled emission fluxes (Cooter et al., 2010; Flechard et al., 2013; Neftel et al., 1998; Nemitz et al., 2001; Xu et al., 2022). Our assessment that the conventional method overestimates the soil emission potential by at least a factor of 5 is similar to findings reported by Nemitz et al., (2001) and Cooter et al., (2010) who reported needing to reduce their soil emission potentials by a factor of 6.66 and 2 – 3, respectively, for their modelled predictions to match their flux observations. We believe that our empirical treatment of this system with the Langmuir and Temkin equations provides a more reasonable approach to estimate the soil emission potential. Additionally, in our view, the ‘like-with-like’ extraction approach (e.g. Γ_{SALT} or Γ_{DIW}) has a more mechanistic basis than the conventional approach involving dissimilar extraction solutions. A few recent studies have reported on a similar approach using a Langmuir adsorption

model (Alnsour, 2020; Guo et al., 2022), as well as by matching the NH_4^+ and pH extraction solutions (Wu et al., 2023), showing that these methods can be feasibly implemented into an environmental sampling campaign. Venterea et al., (2015) reported that soil NH_4^+ partitioning could be effectively modelled using a modified version of the Langmuir isotherm equation of the form $S = \frac{S_{\max}C}{K+C}$, where K (mg L^{-1}) is the aqueous concentration at which $S = \frac{1}{2} S_{\max}$. The Venterea equation is equivalent to the Langmuir isotherm when $K = \frac{1}{K_L}$; for the parameters determined in this study, $\frac{1}{2}$ the saturation capacity is reached at exactly $C = \frac{1}{K_L} = K$, indicating that for our analysis the Langmuir and Venterea equations are equivalent.

How does this approach compare to the implementation of bidirectional exchange of NH_3 in chemical transport models? Widely used models such as GEOS-CHEM and the EPA's CMAQ model soil emission potentials mechanistically, rather than being based on emission potentials prescribed using land-use categories (Bash et al., 2013; Pleim et al., 2013; Pleim, et al., 2019; Zhu et al., 2015). In both models, the exchangeable soil NH_3 has been parameterized as the volumetric molar concentration of NH_4^+ in the top 1 – 5 cm divided by the volumetric water content in the soil, or $[\text{NH}_4^+] = \frac{\text{NH}_4^+ \text{ mol m}^{-3}}{\theta \text{ m}^3 \text{ m}^{-3}}$ (Massad et al., 2010). In this parameterization, the entire soil NH_4^+ content of the top layer of soil is treated as being dissolved into the soil water, thus a lower water content would result in a higher concentration of NH_4^+ , and consequently a higher emission potential. (The parameter used by these models for the resistance to NH_3 emissions from soil is inversely proportional to soil moisture, such that the actual emission of NH_3 from soil would still be proportional to soil water content (Pleim et al., 2013).) In our model of the system, most NH_4^+ is present adsorbed to soils, with the aqueous concentration controlled by the equilibrium between adsorbed and aqueous NH_4^+ , increasing the soil water content would thus allow more NH_4^+ to enter the aqueous phase to maintain the equilibrium concentration. (Increased water content could also result in nutrient runoff, reducing the soil NH_4^+ content, or increased mineralization, increasing the soil NH_4^+ content.) In the subsequent development of the bidirectional model coupled with an agricultural ecosystem model, Pleim et al. (2019) noted that reducing the NH_3 available for emission by implementing the Langmuir adsorption isotherm approach proposed by Venterea et al. (2015) appeared to lead to unexpectedly low fractions of NH_3 available for exchange. It may be that for recently fertilized soils, the applied fertilizer is not in contact with sufficient soil for this equilibrium relation to apply. An updated version of CMAQ (v5.2.1) uses a Langmuir-derived approach described in Venterea et al. (2015) to predict NH_3 bi-directional exchange, and Kelly et al. (2019) explore its indirect impact on $\text{PM}_{2.5}$ composition across the U.S., suggesting that the reduced emission potential from our approach is compatible with atmospheric agro-ecosystem modelling.

5. Conclusion

This work evaluated the Langmuir, Freundlich, Temkin, and Toth adsorption isotherm equations as applied to environmental soil samples and NH_4^+ partitioning. We determined that under ideal conditions the Toth equation was the most effective of these equations at fitting soil adsorption curves, but that the Temkin equation was most effective at predicting the adsorption

behaviour of soil samples with minimal additional characterization required and was effective over both low concentration ranges and the saturation concentration range. Applying this method to a series of environmental soil samples, we determined that the conventional method for directly measuring soil emission potentials may overestimate them by a factor of 5 – 20 (relative to using the Langmuir/Temkin equations respectively). Of the two adsorption equations, the Temkin equation fit the data better, with a significantly lower AIC (-222 vs -138 for the Langmuir equation). Comparing these empirical equations with an alternative approach for determining emission potential based on extracting NH_4^+ and pH with the same extraction solutions (i.e. DIW/DIW or Salt/Salt) showed that the Temkin equation fit using the full adsorption range agreed well with the DIW/DIW ratio method but was significantly different when using the low-range fitting parameters.

305 Appendix A




Figure A1: Curve-fitting comparison between the Langmuir (red, solid), Freundlich (blue, ---), Temkin (green, ·-·-), and Toth (orange, ···) equations for soil samples collected from a) King's College Circle, b) the University of Toronto Scarborough forest, c) High Park, and d) Queen's Park.

310

Table A2: Comparison of the residual standard errors, and the range of fitted parameters for adsorption curves fit to the un-pooled data. Parameters given in parentheses are reported for comparison with the parameters in Table 2. Goodness of fit is reported using the residual standard error (RSE) rather than AIC to avoid confusion between ranges of AICs for each equation.

Equation	RSE	Parameter 1	Mean ± sd	Parameter 2	Mean ± sd	Exponential Factor
Langmuir						
Freundlich						
Temkin						
Toth						

Langmuir	17.5 – 25.7%	K_L	$9.33 \cdot 10^{-4} \pm 2.0 \cdot 10^{-4}$	-	-	-
Freundlich	5.72 – 17.6%	K_F	197 ± 60 ($S_{max} * [0.053 \pm 0.012]$)	-	-	0.3064 ± 0.023
Temkin	3.0 – 16.1%	K_T	$1.34 \cdot 10^{-2} \pm 5.1 \cdot 10^{-3}$	q_T	$S_{max} * [0.183 \pm 0.013]$	-
Toth	0.9 – 6.5%	K_{To}	3.75 ± 1.3	b	9960 ± 5590 ($S_{max} * [2.42 \pm 0.72]$)	0.276 ± 0.054

Table A3: Residual standard errors when applying average fit parameters from the pooled and (separately fit) training data to the test set soil samples, for i) a CEC of 25; ii) the experimentally measured CEC; iii) when the data is fit using the fitting algorithm

Sample Location	CEC (cmole kg ⁻¹)	NH ₄ ⁺ (mg kg ⁻¹)	pH	RSE (%) using estimated fit parameters			
				Langmuir	Freundlich	Temkin	Toth
High Park	10.95	2.906	7.04	280 (285)	330 (340)	320 (330)	330 (420)
				26 (26)	22 (26)	16 (18)	14 (28)
				25	19	14	8.8
Corktown	25.3	3.552	7.82	33 (33)	25 (31)	22 (26)	9.5 (39)
				34 (34)	28 (34)	25 (28)	9.7 (43)
				29	15	15	3.8
Riverdale Park East	37.25	2.835	7.42	64 (63)	62 (58)	66 (63)	84 (62)
				35 (35)	19 (25)	29 (31)	19 (39)
				31	11	24	23
Geometric Mean RSE		Estimated		84 (84)	80 (85)	77 (81)	64 (100)
		Estimated with measured CEC		31 (31)	21 (28)	21 (25)	14 (36)
		Fitted Equation		28.3	14.8	17.0	9.2

Table A4: Sample information for the adsorption test set

Sample	CEC (cmole kg ⁻¹)	Bulk NH ₄ ⁺ (mg kg ⁻¹)	pH	Extractant Solution (mg L ⁻¹)	Soil Mass (g)	Sum of displaced cations (cmole kg ⁻¹)	NH ₄ ⁺ adsorbed (mg kg ⁻¹)
High Park	10.95	2.906	7.04	18.04	1.006	0.989	178.4
				36.08	1.0124	0.949	171.2
				144.32	0.9953	1.951	351.9

				721.6	0.9976	4.028	726.6
				3608	0.9539	7.413	1337
				18040	1.0002	10.81	1950
Corktown	25.3	3.552	7.82	18.04	1.0081	1.793	323.5
				36.08	1.0095	2.222	400.9
				144.32	0.9944	4.228	762.8
				721.6	1.0013	9.064	1635
				3608	1.0131	15.54	2804
				18040	1.002	24.39	4400
Riverdale Park East	37.25	2.835	7.42	18.04	0.9948	4.447	802.2
				36.08	1.0193	4.861	876.8
				144.32	1.0151	6.690	1207
				721.6	1.0304	11.79	2127
				3608	0.9953	23.53	4244
				18040	1.027	36.26	6541

320 **Author contribution statement**

MGD: Investigation, Formal analysis, Writing – Original Draft; KY: Investigation, Methodology; JGM: Supervision, Funding acquisition, Writing – Review & Editing.

Competing interests statement

The authors declare that they have no conflict of interest.

325 **Data Availability Statement**

The data used to construct the soil adsorption curves and the measurements of the properties of environmental soils are publicly archived here: <https://doi.org/10.5683/SP3/JGRIN0>

Funding statement

330 This research was supported by a Natural Science and Engineering Research Council (NSERC) Discovery grant and a Grants and Contributions agreement GCXE19S016 with Environment and Climate Change Canada held by Jennifer Murphy. Matthew Davis held a Walter C. Sumner Memorial Fellowship while conducting this research. An Undergraduate Summer Research Award from NSERC supported Kevin Yan during this work.

Acknowledgements

335 We thank our colleagues Professor Myrna Simpson and Jenny Oh (University of Toronto) for their helpful discussions. We also thank the University of Toronto ANALEST facility staff for their technical assistance.

References

- Alnsour, N. I.: Bi-Directional Exchange of Ammonia from Soils in Row Crop Agro-ecosystems, Ph.D. thesis, North Carolina State University, 2020.
- Bache, B. W.: The measurement of cation exchange capacity of soils. *Journal of the Science of Food and Agriculture*, 27(3), 273–280. <https://doi.org/10.1002/jsfa.2740270313>, 1976.
- Bash, J. O., Cooter, E. J., Dennis, R. L., Walker, J. T., & Pleim, J. E.: Evaluation of a regional air-quality model with bidirectional NH₃ exchange coupled to an agroecosystem model. *Biogeosciences*, 10(3), 1635–1645. <https://doi.org/10.5194/bg-10-1635-2013>, 2013.
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Hoek, K. W. Van Der, Olivier, J. G. J., Tg, N., & Tg, N.: A global high-resolution emission inventory for ammonia. *Global Biogeochemical Cycles*, 11(4), 561–587, 1997.
- Chu, K. H.: Revisiting the Temkin Isotherm: Dimensional Inconsistency and Approximate Forms. *Industrial and Engineering Chemistry Research*, 60(35), 13140–13147. <https://doi.org/10.1021/acs.iecr.1c01788>, 2021.
- Cooter, E. J., Bash, J. O., Walker, J. T., Jones, M. R., & Robarge, W.: Estimation of NH₃ bi-directional flux from managed agricultural soils. *Atmospheric Environment*, 44(17), 2107–2115. <https://doi.org/10.1016/j.atmosenv.2010.02.044>, 2010.
- Ellis, E. C., Beusen, A. H. W., & Goldewijk, K. K.: Anthropogenic biomes: 10,000 BCE to 2015 CE. *Land*, 9(5), 8–10. <https://doi.org/10.3390/LAND9050129>, 2020.
- Farquhar, G. D., Firth, P. M., Wetselaar, R., & Weir, B.: On the Gaseous Exchange of Ammonia between Leaves and the Environment: Determination of the Ammonia Compensation Point. *Plant Physiology*, 66(4), 710–714. <https://doi.org/10.1104/pp.66.4.710>, 1980.
- Flechard, C. R., Fowler, D., Sutton, M. A., & Cape, J. N.: A dynamic chemical model of bi-directional ammonia exchange between semi-natural vegetation and the atmosphere. In *Quarterly Journal of the Royal Meteorological Society* (Vol. 125, Issue 559). <https://doi.org/10.1002/qj.49712555914>, 1999.
- Flechard, C. R., Massad, R.-S., Loubet, B., Personne, E., Simpson, D., Bash, J. O., Cooter, E. J., Nemitz, E., & Sutton, M. A.: Advances in understanding, models and parameterizations of biosphere-atmosphere ammonia exchange. *Biogeosciences*, 10(7), 5183–5225. <https://doi.org/10.5194/bg-10-5183-2013>, 2013.
- Freundlich, H.: *Kapillarchemie: eine Darstellung der Chemie der Kolloide und verwandter Gebiete*. In Akademische Verlagsgesellschaft. Verlag der Akademischen Verlagsgesellschaft. <https://doi.org/10.1038/130866a0>, 1909.
- Guo, X., Pan, D., Daly, R. W., Chen, X., Walker, J. T., Tao, L., McSperritt, J., & Zondlo, M. A.: Spatial heterogeneity of ammonia fluxes in a deciduous forest and adjacent grassland. *Agricultural and Forest Meteorology*, 326(July), 109128. <https://doi.org/10.1016/j.agrformet.2022.109128>, 2022.
- Kelly, J. T., Koplitz, S. N., Baker, K. R., Holder, A. L., Pye, H. O. T., Murphy, B. N., Bash, J. O., Henderson, B. H., Possiel, N. C., Simon, H., Eyth, A. M., Jang, C., Phillips, S., & Timin, B.: Assessing PM_{2.5} model performance for the conterminous

- U.S. with comparison to model performance statistics from 2007-2015. *Atmospheric Environment*, 214(May), 116872. <https://doi.org/10.1016/j.atmosenv.2019.116872>, 2019.
- 370 Kempers, A. J., & Zweers, A.: Ammonium determination in soil extracts by the salicylate method. *Communications in Soil Science and Plant Analysis*, 17(7), 715–723. <https://doi.org/10.1080/00103628609367745>, 1986.
- Langmuir, I.: *The Constitution and Fundamental Properties of Solids and Liquids. Part I. Solids.* Journal of the American Chemical Society, 38(11), 2221–2295, 1916.
- Li, K. yi, Zhao, Y. yuan, Yuan, X. long, Zhao, H. bing, Wang, Z. hui, Li, S. xiu, & Malhi, S. S.: Comparison of Factors
375 Affecting Soil Nitrate Nitrogen and Ammonium Nitrogen Extraction. *Communications in Soil Science and Plant Analysis*, 43(3), 571–588. <https://doi.org/10.1080/00103624.2012.639108>, 2012.
- Massad, R. S., Nemitz, E., & Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmospheric Chemistry and Physics*, 10(21), 10359–10386. <https://doi.org/10.5194/acp-10-10359-2010>, 2010.
- 380 Neftel, A., Blatter, A., Gut, A., Högger, D., Meixner, F., Ammann, C., & Nathaus, F. J.: NH₃ soil and soil surface gas measurements in a triticale wheat field. *Atmospheric Environment*, 32(3), 499–505. [https://doi.org/10.1016/S1352-2310\(97\)00162-3](https://doi.org/10.1016/S1352-2310(97)00162-3), 1998.
- Nemitz, E., Milford, C., & Sutton, M. A.: A two layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia. *Q. J. R. Meteorol. Soc.*, 127, 815–833, 2001.
- 385 Nemitz, E., Sutton, M. A., Schjoerring, J. K., Husted, S., & Paul Wyers, G.: Resistance modelling of ammonia exchange over oilseed rape. *Agricultural and Forest Meteorology*, 105(4), 405–425. [https://doi.org/10.1016/S0168-1923\(00\)00206-9](https://doi.org/10.1016/S0168-1923(00)00206-9), 2000.
- Pleim, J. E., Bash, J. O., Walker, J. T., & Cooter, E. J.: Development and evaluation of an ammonia bidirectional flux parameterization for air quality models. *Journal of Geophysical Research: Atmospheres*, 118(9), 3794–3806. <https://doi.org/10.1002/jgrd.50262>, 2013.
- 390 Pleim, J. E., Ran, L., Appel, W., Shephard, M. W., & Cady-Pereira, K.: New Bidirectional Ammonia Flux Model in an Air Quality Model Coupled With an Agricultural Model. *Journal of Advances in Modeling Earth Systems*, 11(9), 2934–2957. <https://doi.org/10.1029/2019MS001728>, 2019.
- Spiess, A. N., & Neumeyer, N.: An evaluation of R² as an inadequate measure for nonlinear models in pharmacological and biochemical research: A Monte Carlo approach. *BMC Pharmacology*, 10, 1–11. <https://doi.org/10.1186/1471-2210-10-6>,
395 2010.
- Sutton, M. A., Erisman, J. W., Dentener, F., & Möller, D.: Ammonia in the environment: From ancient times to the present. *Environmental Pollution*, 156(3), 583–604. <https://doi.org/10.1016/j.envpol.2008.03.013>, 2008.
- Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B.,
400 Massad, R., ... de Vries, W.: Towards a climate-dependent paradigm of ammonia emission and deposition. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621). <https://doi.org/10.1098/rstb.2013.0166>, 2013.

- Sutton, M. A., Schjørring, J. K., & Wyers, G. P.: Plant-atmosphere exchange of ammonia. *Philos. T. R. Soc. Lond., A*, 351(29), 261–276, 1995.
- 405 Temkin, M. I., & Pyzhev, V.: Kinetics of ammonia synthesis on promoted iron catalyst. *Acta Phys. Chim. USSR*, 12(327), 1940.
- Tóth, J.: Uniform interpretation of gas/solid adsorption. *Advances in Colloid and Interface Science*, 55(C), 1–239. [https://doi.org/10.1016/0001-8686\(94\)00226-3](https://doi.org/10.1016/0001-8686(94)00226-3), 1995.
- 410 Van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Goldewijk, C. G. M. K., & Lelieveld, J.: A $1^\circ \times 1^\circ$ resolution data set of historical anthropogenic trace gas emissions for the period 1890-1990. *Global Biogeochemical Cycles*, 15(4), 909–928. <https://doi.org/10.1029/2000GB001265>, 2001.
- Venterea, R. T., Clough, T. J., Coulter, J. A., & Breuillin-Sessoms, F.: Ammonium sorption and ammonia inhibition of nitrite-oxidizing bacteria explain contrasting soil N₂O production. *Scientific Reports*, 5(April), 1–15. <https://doi.org/10.1038/srep12153>, 2015.
- 415 Vogeler, I., Cichota, R., Snow, V. O., Dutton, T., & Daly, B.: Pedotransfer Functions for Estimating Ammonium Adsorption in Soils. *Soil Science Society of America Journal*, 75(1), 324–331. <https://doi.org/10.2136/sssaj2010.0192>, 2011.
- Walker, J. T., Chen, X., Wu, Z., Schwede, D., Daly, R., Djurkovic, A., Oishi, A. C., Edgerton, E., Bash, J., Knoepp, J., Puchalski, M., Iiames, J., & Miniati, C. F.: Atmospheric deposition of reactive nitrogen to a deciduous forest in the southern Appalachian Mountains. *Biogeosciences*, 20(5), 971–995. <https://doi.org/10.5194/bg-20-971-2023>, 2023.
- 420 Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G., & Hems, R.: Soil-atmosphere exchange of ammonia in a non-fertilized grassland: Measured emission potentials and inferred fluxes. *Biogeosciences*, 11(20), 5675–5686. <https://doi.org/10.5194/bg-11-5675-2014>, 2014.
- Wu, Z., Walker, J. T., Oishi, A. C., Duman, T., Katul, G., Chen, X., Schwede, D., Bash, J., & Iiames, J.: Estimating source-sink distributions and fluxes of reactive nitrogen and sulfur within a mixed forest canopy. *Agricultural and Forest Meteorology*, 333(June 2022), 109386. <https://doi.org/10.1016/j.agrformet.2023.109386>, 2023.
- 425 Xu, M., Umehara, M., Sase, H., & Matsuda, K.: Ammonia fluxes over an agricultural field in growing and fallow periods using relaxed eddy accumulation. *Atmospheric Environment*, 284(April), 119195. <https://doi.org/10.1016/j.atmosenv.2022.119195>, 2022.
- Zhang, L., Wright, L. P., & Asman, W. A. H.: Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models. *Journal of Geophysical Research Atmospheres*, 115(20). <https://doi.org/10.1029/2009JD013589>, 2010.
- 430 Zhu, L., Henze, D., Bash, J., Jeong, G. R., Cady-Pereira, K., Shephard, M., Luo, M., Paulot, F., & Capps, S.: Global evaluation of ammonia bidirectional exchange and livestock diurnal variation schemes. *Atmospheric Chemistry and Physics*, 15(22), 12823–12843. <https://doi.org/10.5194/acp-15-12823-2015>, 2015.