

Uncertainty in the projected Antarctic contribution to sea level due to internal climate variability

Justine Caillet¹, Nicolas C. Jourdain¹, Pierre Mathiot¹, Fabien Gillet-Chaulet¹, Benoit Urruty¹, Clara Burgard^{1,2}, Charles Amory^{1,3}, Mondher Chekki¹, and Christoph Kittel^{1,4,5}

¹Univ. Grenoble Alpes, CNRS, INRAE, IRD, Grenoble INP, Institut des Geosciences de l'Environnement, Grenoble, France

²Laboratoire d'Océanographie et du Climat, LOCEAN-IPSL, Sorbonne Université, CNRS-IRD-MNHN, Paris, France

³Laboratoire de Sciences du Climat et de l'Environnement, LSCE-IPSL, CEA-CNRS-UVSQ-UMR8212, Université Paris Saclay, Gif-sur-Yvette, France

⁴Department of Geography, Laboratory of Climatology, SPHERES, University of Liège, Liège, Belgium

⁵Earth System Science and Departement Geografie, Vrije Universiteit Brussel, Brussels, Belgium

Correspondence: Justine Caillet (justine.caillet@univ-grenoble-alpes.fr)

Abstract. Identifying and quantifying irreducible and reducible uncertainties in the Antarctic Ice Sheet response to future climate change is essential for guiding mitigation and adaptation policy decision. However, the impact of the irreducible internal climate variability, resulting from processes intrinsic to the climate system, remains poorly understood and quantified. Here, we characterise both the atmospheric and oceanic internal climate variability in a selection of three CMIP6 models (UKESM1-0-LL, IPSL-CM6A-LR and MPI-ESM1.2-HR) and estimate their impact on the Antarctic contribution to [sea-level-sea-level change](#) over the 21st century under the SSP2-4.5 scenario. To achieve this, we use a standalone ice-sheet model driven by the ocean through parameterised basal melting and by the atmosphere through emulated surface mass balance estimates. [The atmospheric component of internal climate variability in Antarctica has a similar amplitude in the three CMIP6 models. In contrast, the amplitude of the oceanic component strongly depends on the climate model and its representation of convective mixing in the ocean. A low bias in sea-ice production and an overly stratified ocean lead to a lack of deep convective mixing which results in weak ocean variability near the entrance of ice-shelf cavities.](#) Internal climate variability affects the Antarctic contribution to [changes in sea-level-sea-level change](#) until 2100 by 45% to 93% depending on the CMIP6 model. This may be a low estimate as the internal climate variability in the CMIP models is likely underestimated. ~~For all the three climate models and for most Antarectic regions, the~~ [The](#) effect of atmospheric internal climate variability on the surface mass balance overwhelms the effect of oceanic internal climate variability on the dynamical ice-sheet mass loss by ~~more than~~ a factor of ~~3~~. ~~The atmospheric internal climate variability is similar in the three CMIP6 models analysed in this study. Conversely, the amplitude of oceanic internal climate variability around Antarctica strongly depends on the climate model as underestimated convection, due to either biases in the sea-ice behaviour or in the ocean stratification, leads to weak mid-depth ocean variability. We then issue recommendations for future~~ [o 5, except in Dronning Maud area and Amundsen, Getz and Aurora basins where both contributions may be similar depending on the CMIP model. Based on these results, we recommend that ice-sheet projections: use several members in the run and in its initialisation, favor 50-year averages to correct or weight simulations over the model projections consider \(i\) several climate models and several members of a single climate model to account for the impact of](#)

internal climate variability and (ii) longer temporal period when correcting historical climate forcing to match present-day period, and couple ice-sheet and climate models observations.

25 1 Introduction

The Antarctic Ice Sheet (AIS) is losing mass at an increasing rate (Rignot et al., 2019; Shepherd et al., 2019), particularly in the Amundsen and Totten/Moscow sectors, where ocean-induced melting under floating ice shelves is relatively high (Jenkins et al., 2018; Hirano et al., 2023). The AIS response to future climate change, including its potential instability (Garbe et al., 2020; Armstrong McKay et al., 2022), is one of the ~~major-main~~ sources of uncertainty in projections of global ~~sea-level~~
30 ~~sea-level~~ rise (Fox-Kemper et al., 2021), with an estimated contribution over 2015–2100 ranging from -5 to 43 cm under a high-end anthropogenic emission scenario (ISMIP6, Edwards et al., 2021).

~~The Estimates of the~~ AIS contribution to future ~~sea-level rise is sea-level rise are~~ currently mostly based on standalone ice-sheet models, ~~forced-driven~~ by atmospheric and oceanic data from the ~~Climate-Coupled~~ Model Intercomparison Project (CMIP, Eyring et al., 2016). The diversity of climate conditions across the CMIP models explains an important part of the uncertainty
35 in some of the drainage basins (Seroussi et al., 2023), despite the use of an anomaly method to reduce known biases in the CMIP models (Jourdain et al., 2020; Purich and England, 2021). Internal climate variability is usually not accounted for in the uncertainty ~~on-of~~ AIS projections. A single study, so far, has estimated that this uncertainty could be 18-21% higher due to internal climate variability (Tsai et al., 2020). This was estimated using a single ice-sheet model and two versions of the same climate model.

40 Climate variability is the combination of two components, ~~the variability resulting~~ on the one hand the variability resulting from external forcing of both natural (e.g., volcanoes or solar activity) and anthropogenic (e.g., CO₂ emissions) sources, and internal variability on the other hand. The latter results from processes intrinsic to the climate system, due to the chaotic nature of fluid dynamics and to non-linearities in the coupled interactions between the ocean, atmosphere, land and cryosphere (e.g., Kravtsov et al., 2007; Penduff et al., 2018; Gwyther et al., 2018; Hogg et al., 2022). For a given climate model, the impact of
45 internal climate variability can be isolated by considering several forced simulations with identical external forcing but slightly different initial conditions. For this reason, an increasing number of CMIP models include several members which differ only in their initial state.

A part of the internal climate variability can be characterised as modes such as the El Niño/Southern Oscillation (~~ENSO~~) and Interdecadal Pacific Oscillation (~~IPO~~), which have remote connections with the Amundsen Sea Low (ASL, Holland et al.,
50 2022; Dalaiden et al., 2023). The ASL is a low-pressure system located over the South Pacific sector of the Southern Ocean, which generates decadal wind anomalies that affect the oceanic undercurrent along the continental slope, thereby modulating the amount of warm water flowing towards the ice shelves of the Amundsen Sea Embayment (Silvano et al., 2022). The regional influence of these modes makes internal climate variability particularly strong in the Amundsen sector: internal climate variability is thought to be responsible for the retreat of Pine Island’s grounding line in the 1940s (Holland et al., 2022), and

55 mid-depth ocean warming trends over the 21st century can vary by a factor of two depending on the phasing of internal climate variability (Naughten et al., 2023).

In this paper, we first investigate atmospheric and oceanic internal climate variability of several CMIP6 models. Then, using a standalone ice-sheet model forced by CMIP6 model outputs, we quantify the impact of internal climate variability on [Antarctic](#) Sea Level Contribution (SLC) over the 21st century under the medium SSP2-4.5 scenario for both the whole ice sheet and the
60 main basins, especially the Amundsen basin which is expected to be particularly affected by internal climate variability.

2 Methods

2.1 CMIP6 ~~model selection~~ models

We choose to analyse three CMIP6 models to get a more general picture of the internal climate variability than [we would get](#) using a single model. ~~These three models were selected both for~~

65 [The selected models are UKESM1-0-LL \(19 members, Sellar et al., 2020\), MPI-ESM1.2-HR \(10 members, Müller et al., 2018\) and IPSL-CM6A-LR model \(33 members, Boucher et al., 2020\). This choice was made based on \(i\) the size of their ensemble and for \(at least 10 members\), \(ii\) the availability of 6-hourly outputs that were needed to run regional climate projections, and \(iii\) their representation of the present-day Southern Ocean and Antarctic seas, as basal melting is one of the main driver of the Antarctic contribution to sea level rise. The selection was done based on a review of three studies: the evaluation of the properties of Antarctic Shelf Bottom Water \(ASBW\) on the continental shelf and of Circumpolar Deep Water \(CDW\), located further offshore in the Antarctic circumpolar region and expected to intrude more on the continental shelf in a warmer climate, conducted by Purich and England \(2021\), the evaluation of dynamical features known to be important for the Southern Ocean, including the transport of the Antarctic Circumpolar Current \(ACC\), zonal wind stress properties, and meridional gradients of ocean properties, conducted by Beadling et al. \(2020\), the evaluation of the bottom properties of \[oceanic and atmospheric\]\(#\)
70 \[properties.\]\(#\)
75 \[properties.\]\(#\)](#)

[For the third point, the Southern Ocean, conducted by Heuzé \(2021\). A summary of this assessment is provided in Fig. A1. The selected models are: UKESM1-0-LL \(19 members, Sellar et al., 2020\), which is among \[three selected models are in the best half of the CMIP6 ensemble according to Agosta \\(2024\\) who ranked 45 models based on several atmospheric variables relevant for precipitation over Antarctica. These three models also have a high fidelity in the representation of the mean ocean\]\(#\)
80 \[properties, as detailed in Appendix A.\]\(#\)](#)

[Although their oceanic and atmospheric mean state are some of the ~~best models in the three studies;~~ closest to observations, the three selected models have distinct characteristics of their internal climate variabilities. As shown in Appendix B, the atmospheric variability at the scale of Antarctica is close to the multi-model median in IPSL-CM6A-LR \(33 members, Boucher et al., 2020\) , which represents reasonably well the deep ocean properties and to a lesser extent the main Southern Ocean dynamics, but produces overly cold CDWs; MPI-ESM1.2-HR \(10 members, Müller et al., 2018\), which was evaluated in only one of the three aforementioned studies, but has a higher ocean resolution \(0.4° vs 1°\) and a lower Equilibrium Climate Sensitivity \(3.0°C vs 5.3 and 4.6°C, Meehl et al., 2020\) than the two other models. It \[and MPI-ESM1.2-HR, while it is much higher in\]\(#\)](#)
85 [and MPI-ESM1.2-HR, while it is much higher in](#)

UKESM1-0-LL. The oceanic variability is also among the lowest in MPI-ESM1-2-HR, close to the multi-model median in UKESM1-0-LL, and much higher in IPSL-CM6A-LR.

90 It is interesting to note that both UKESM1-0-LL and IPSL-CM6A-LR have some-kind-of-prescribed ice-shelf melting at depth-to-ensure the conservation of the ice-sheet mass over the pre-industrial or the entire simulation that is vertically distributed to mimic the presence of unresolved ice-shelf cavities (Mathiot et al., 2017), which is known to be important for coastal ocean properties around Antarctica (Mathiot et al., 2017; Donat-Magnin et al., 2021).

~~Moreover, the present-day properties of the atmosphere of the three selected CMIP6 models compare well to ERA5.~~
95 ~~According to the evaluation of the CMIP5/6 models conducted in Agosta et al. (2022), UKESM1-0-LL and MPI-ESM1-2-HR are among the best performing models in the Antarctic region, while the IPSL-CM6A-LR model is between the first and second quartile.~~

~~Assessment of Southern Ocean and Antarctic seas properties in the CMIP6 models. In each panel, CMIP models are ranked by increasing RMSE. (a) Antarctic Shelf Bottom Water (ASBW) and Circumpolar Deep Water (CDW) temperature biases in the Ross, Amundsen/Bellingshausen, Weddell and Cosmonauts seas (Purich and England, 2021, their Fig. S16). (b) Biases in the Southern Ocean metrics defined in Tab. 1 of Beadling et al. (2020) and describing several characteristics of the Antarctic Circumpolar Current, zonal wind stress strength, location and curl, as well as meridional gradients of water mass properties. (c) RMSE of bottom ocean properties (density, salinity and temperature) in the Southern Ocean (Heuzé, 2021, their Figs. 1, A1, A2). For each panel, the metric is normalised by the CMIP6 multi-model standard deviation. A red triangle indicates the selected~~
100 ~~models. Most CMIP models prescribe meltwater fluxes at the surface, which tends to increase the ocean stratification (Mathiot et al., 2017) and reduce exchanges between the surface and deeper waters, thereby limiting variability at depth.~~
105

2.2 Ice-sheet model

We use the version v9.0 of the Elmer/Ice finite element model (Gagliardini et al., 2013), in a configuration of the entire Antarctic Ice Sheet adapted from Hill et al. (2023). The ice dynamics is computed by solving the Shallow Shelf Approximation (SSA)
110 of the Stokes equations (MacAyeal, 1989), assuming an isotropic rheology following Glen's flow law (Glen, 1955) and a linear friction law ~~-(i.e., $\tau_b = C u_b$ where τ_b is the basal shear stress, C the friction coefficient and u_b the basal velocity).~~ The location of the grounding line is determined using a flotation criterion and a sub-grid scheme is applied for the friction in partially floating elements (SEP3 in Seroussi et al., 2014).

The mesh is ~~preferentially~~ refined both close to the grounding line and in areas where observed surface velocities and
115 thickness show high curvatures ~~-,with-~~ ~~(i.e., high second derivative of the modelled field, Gillet-Chaulet et al., 2012).~~ ~~The mesh has~~ a maximum size of 50 km in the very interior of the ice sheet and a minimum size of 1 km in the refined areas. The model domain does not change over time, but the ice thickness is subject to a lower limit of 1 m and elements that reach this limit are considered deglaciated in the post-processing. For stability reasons, the domain boundary is slightly smoothed and isolated icebergs (ice-covered area disconnected from the ice sheet) with less than 7 elements, are removed if they appear during the
120 simulation (i.e, their thickness is set to the critical thickness of 1 m). Apart from these corrections, we assume a steady calving front.

Inverse methods (Gillet-Chaulet et al., 2012; Brondex et al., 2019) provide viscosity and friction parameters by minimising the misfit between modelled and observed velocities from Mouginot et al. (2019) using the ice thickness from BedMachine-Antarctica-v2 (Morlighem et al., 2020). Details of the inversion are available in Hill et al. (2023). Our model configuration does not represent a prognostic evolution of ice temperature and damage that may affect viscosity in transient simulations. From the inversion step, we run a 20-year “relaxation” under the present-day forcing described hereafter. This attenuates the artificial high surface elevation rate of change that occurs when we switch from a diagnostic to a prognostic simulation (Gillet-Chaulet et al., 2012).

The PICO box model (Reese et al., 2018) is used to parameterise ice-shelf basal melting, with a distinct calibration from Hill et al. (2023). Here, the parameters are those detailed in Reese et al. (2023), i.e., $C = 2 \text{ Sv m}^3 \text{ kg}^{-1}$ and $\gamma_T = 5.5 \times 10^{-5} \text{ m s}^{-1}$, which are based on the observed or ocean-modelled sensitivity of melt rates to ocean temperature changes. The present-day sea floor temperature and salinity for each of the 19 regions defined in Reese et al. (2018) are extracted from the ISMIP6 ocean climatology (Jourdain et al., 2020) and averaged within 50 km of the ice-shelf front as described in Burgard et al. (2022). A correction of temperature, ranging from -1.8°C to 0.6°C with respect to the ocean climatology, is added to match the 1994-2018 melt rates estimates from Adusumilli et al. (2020) (see Fig. 1). This correction differs from Reese et al. (2023) as the current ice-sheet geometry and the oceanic climatology used in this study are different from the one considered in Reese et al. (2023).

The present-day Surface Mass Balance (SMB) is based on the 1995-2014 climatology (a period of relatively stable SMB) of the RACMO-2.3.p2 regional climate model (Van Wessem et al., 2018). In contrast to Hill et al. (2023), we do not correct the surface mass balance to maintain a steady state, but we apply a 10% reduction of uniformly lower the inverted friction coefficients to be closer to the by 10% to reduce the model drift. For this, we minimise the RMSE between the modelled and the observed ice-sheet mass change (The IMBIE Team, 2018) for both West and East Antarctica. Our configuration correctly estimates change West Antarctica. The resulting model configuration overestimates the mass loss trend in the West Antarctica but still by only 6% but still largely overestimates mass gain trend in East Antarctica and in the Peninsula (Tab. 1). This mass change trend As a consequence, the simulated Antarctic Ice Sheet is currently gaining a little mass ($+36 \text{ Gt yr}^{-1}$, Tab. 1), instead of losing mass as observed ($-109 \pm 56 \text{ Gt yr}^{-1}$, Tab. 1). This growing bias is quite common in ice-sheet models (Seroussi et al., 2020; Aschwanden et al., 2021). This is why our results are primarily analysed in relative terms However, this bias should not impact most of the analyses presented here, as the projections in response to the CMIP6 climate models are analysed relatively to each other.

2.3 Ice-sheet projections to 2100

Antarctic future mass change The future mass imbalance of Antarctica results from combined effects of changes in surface mass balance and dynamics changes. Given that (SMB) and ice dynamics. In standalone ice-sheet simulations, variations in surface mass balance can be attributed to the atmosphere and dynamical mass loss can be attributed to the ocean as SMB changes have little impact on the dynamical contribution Antarctic dynamical contribution to sea level over a century (Seroussi et al., 2014, 2023), these two contributions. Thus, the effect of atmospheric and oceanic variations on Antarctic contribution to sea-level change can be analysed separately, and then summed to reconstruct the combined effect (Bindschadler et al., 2013).




Figure 1. Basal melt rate of main ice shelves over the period 1994-2014 before (lightblue) and after calibration (blue) compared to the melting estimates over the period 1994-2018 from Adusumilli et al. (2020, in red). Observed data uncertainties correspond to one standard deviation. Note that the data from Adusumilli et al. (2020) only cover the area northward of 81.5°S, which excludes part of the Filchner-Ronne and Ross ice shelves. Melting estimates over the period 2003-2008 from [Rignot et al. \(2013, in black\)](#) [Rignot et al. \(2013, in orange\)](#) are shown for comparison. [Numbers from 1 to 18 indicate the basin where ice shelves are located, as shown in the top right-hand corner.](#)

Table 1. Rates of ice-sheet mass change (Gt yr^{-1}) for the entire [Antarctica Antarctic Ice Sheet \(AIS\)](#) and ~~the main areas of its three major basins~~: East Antarctica (EAIS), West Antarctica (WAIS) and Peninsula (APIS). [The IMBIE data are from The IMBIE Team \(2018\).](#)

	IMBIE estimates (1992-2017)	Elmer/Ice (1994-2014)
AIS	-109 ± 56	+36
EAIS	$+5 \pm 46$	+107
WAIS	-94 ± 27	-127
APIS	-20 ± 15	+35

~~To investigate the impact of oceanic internal climate variability on the Antarctic Ice Sheet projections, we run~~ [In our study, the SMB contribution to sea level is directly deduced from the emulated SMB anomalies \(i.e., cumulative SMB – initial SMB\). The contribution of ice dynamics to sea level is estimated through Elmer/Ice simulations constrained by the driven by the SMB of the first member and the ocean of several members of the selected CMIP models from 2015 to 2100. We then remove the SMB contribution of the first member to deduced the dynamical contribution.](#)

We use the medium SSP2-4.5 scenario, which corresponds to a global warming of 1.4 to 3.0°C from 1995-2014 to 2081-2100 (90% confidence interval, Lee et al., 2021) ~~and seems the most representative of current efforts to tackle climate change (Riahi et al., 2017). As the choice of greenhouse gas emission scenario has only a limited impact on the projected Antarctic contribution to sea-level rise until 2100 (Seroussi et al., 2020), we have not repeated our calculations for other scenarios.~~ Because of the numerical cost of our simulations, we select a limited number of members. ~~The~~ In addition to the first member, the selection is made over the current period (1995-2014 means) to cover the widest range of values for both the ocean temperature on the continental shelf in the Amundsen Sea and the SMB over the whole of Antarctica (see next section. We focus on this region as (i) the largest mass loss is observed there and has been attributed to the ocean, and (ii) the amplitude of the standard deviation of the 1995-2014 mean potential temperature across all members is particularly high in this region (see section 3.1).

165
170 In total, we run 11 simulations, five with the IPSL-CM6A-LR model (r1i1p1f1, r3i1p1f1, r6i1p1f1, r11i1p1f1, r25i1p1f1, see the CMIP6 naming convention in <https://goo.gl/v1drZI>), four with the UKESM1-0-LL model (r1i1p1f2, r2i1p1f2, r4i1p1f2, r8i1p1f2) ~~and~~, and only two with the MPI-ESM1.2-HR model (r1i1p1f2, r2i1p1f2) given that its oceanic variability is very low (see section 3.1).

All the Elmer/Ice simulations start from the same state, corresponding to 2014, and yearly atmospheric and oceanic anomalies are added to the present-day atmospheric and oceanic forcing to drive future projections as previously done in ISMIP6 (Nowicki et al., 2020). All the anomalies are calculated with respect to the 1995-2014 period of a given member.

175

The annual ocean potential temperature and practical salinity from CMIP model outputs were interpolated to a stereographic (8 km × 8 km × 60 m) grid, then extrapolated to fill unrepresented areas as in Jourdain et al. (2020). The corresponding ocean anomalies were then added to the present-day temperature and salinity to feed the ice-shelf basal melt parameterisation.

180 Regional climate projections were not used to calculate the future SMB of ISMIP6-Antarctica (Nowicki et al., 2020; Seroussi et al., 2020), mostly because they were not available early enough in the intercomparison process. Since then, this kind of simulations have been used to refine SMB projections (Kittel et al., 2021, 2022). Using a dedicated regional climate model is particularly important for the IPSL-CM6A-LR model given that its snow physics over ice sheets is too simple to simulate firn saturation and runoff in a warmer climate. However, running the regional climate model driven by many members of the CMIP

185 ensemble would be computationally too expensive and practically not feasible due to the non availability of 6-hourly output for most members, which are needed to ~~constrain-drive~~ the regional model.

In this paper, we therefore use the approach developed by Jourdain et al. (2024) to emulate the behaviour of the *Modèle Atmosphérique Régional* (MAR, Kittel et al., 2021). This method uses exponential fits of accumulation and surface melting perturbations due to changes in surface air temperature, as well as simple physical relationships to derive runoff and

190 SMB. ~~The fit parameters are calculated from 1980-2100 MAR simulations forced by the corresponding CMIP model (either IPSL-CM6A-LR, UKESM1-0-LL or MPI-ESM1.2-HR). The SMB of a given member is estimated from the relevant MAR simulation of the first member, perturbed as a function of the annual temperature difference between the two CMIP model members.~~ This method is thoroughly evaluated in Jourdain et al. (2024) for the emulation of other CMIP models and scenarios based on a few existing regional simulations, and here we apply it to emulate other members based on existing regional

195 [simulations of the first member of each CMIP model](#). Similarly to the ocean forcing, we calculate annual anomalies (with respect to 1995-2014 mean SMB) and add them to the present-day SMB.

3 Results

We first characterize internal climate variability of the oceanic (subsect. 3.1) and atmospheric (subsect. 3.2) components in the selected CMIP6 models. For this, we use all available members and we describe the effect of internal climate variability on the present-day mean state, i.e., 1995-2014, which is used as a reference for the calculation of anomalies in ISMIP6 and in our Elmer/Ice simulations. Then, we estimate the importance of internal climate variability for [sea level](#) projections by examining transient Elmer/Ice simulations from 2015 to 2100 ([subsec. 3.3](#)), driven by [a the](#) subset of the CMIP6 ensemble used in the [two first subsections\(subsec. 3.3\)subsections 3.1 and 3.2](#).

3.1 Oceanic internal climate variability

205 Oceanic internal climate variability [is investigated through salinity and temperature variability](#). Oceanic internal climate variability at mid-depth is much weaker in MPI-ESM1.2-HR than in IPSL-CM6A-LR and UKESM1-0-LL (Fig. 2). MPI-ESM1.2-HR shows a relatively low and homogeneous internal climate variability on the continental shelf, with ~~typical~~ standard deviations of ~~0.017~~[0.02](#) g kg^{-1} and 0.06°C across the members (Fig. 2a,d). The mean salinity of this model is too low over the whole continental shelf (34.2 g kg^{-1}) compared to the World Ocean Atlas dataset ([WOA 2018 Boyer et al., 2018](#)) ([WOA 2018, Boyer et al., 2018](#)), particularly in front of the Ross and Filchner ice shelves (Fig. 3a,b). This suggests that the weak internal climate variability is related to an underestimation of dense water formation (Fig. 3b).

For IPSL-CM6A-LR and UKESM1-0-LL, salinity exhibits higher variability over the whole [continental](#) shelf (around ~~0.030~~[0.03-0.04](#) g kg^{-1} in Fig. 2b,c) but this does not systematically lead to a high variability in temperature (Fig. 2e,f). ~~The part that undergoes the largest~~ [A region that undergoes large](#) variability in mid-depth temperatures in both IPSL-CM6A-LR and UKESM1-0-LL is the region extending westward from the Bellingshausen Sea to the western Ross Sea. There are nonetheless noticeable differences between the two models.

For IPSL-CM6A-LR, the largest variability in mid-depth salinity is found in the western part of the Ross Sea, where High Salinity Shelf Water (HSSW) is formed (Fig. 2b). ~~As described by Mathiot and Jourdain (2023) and Siahaan et al. (2022), lower rates of HSSW formation in the eastern Ross Sea can favor the intrusion of~~ [The deepest part of the Ross Sea is occupied by the densest water mass, so that there is a competition between intrusions of relatively warm and salty](#) Circumpolar Deep Water (CDW) ~~in the western part, which may explain why~~ [advected from offshore and the production of cold dense water \(HSSW\) through sea-ice formation and associated convection \(Siahaan et al., 2022; Mathiot and Jourdain, 2023\). The variation between the occupation of these two water masses may explain the high](#) mid-depth temperature variability ~~is highest there in the Ross Sea~~ (Fig. 2e). In contrast, the ~~high~~ variability in salinity at the HSSW formation site of the eastern Weddell Sea (Fig. 2b) is probably too weak to be associated with any CDW intrusion (Fig. 2e).

For UKESM1-0-LL, the highest variability in mid-depth salinity is located around Prydz Bay in East Antarctica (Fig. 2c), which is an area of important dense shelf water formation (Williams et al., 2016). It nonetheless does not induce a strong temperature variability near the ice shelves (Fig. 2f). An interesting feature of UKESM1-0-LL (and IPSL-CM6A-LR to a lower extent) is the high salinity variability ~~in the deep ocean~~beyond the continental shelf, northward of the Amundsen Sea (Fig. 2c), which coincides with a region of high variability ~~of in~~ sea-ice concentration (not shown) and air temperature (Fig. 4f).

We now ~~examine focus on~~ the Amundsen Sea ~~more closely~~, as the region is ~~particularly important for the ice-sheet mass loss currently experiencing the largest mass loss in Antarctica~~. In MPI-ESM1.2-HR, the first 100 m are much fresher than observed and than in the two other models (Fig. 2g), and the entire water column is too warm with an overly strong and shallow thermocline (Fig. 2j). Sea-ice concentration is considerably lower than for the other two models and observations (Fig. 3i-l), which results in a lack of deep convection on the continental shelf. The low oceanic internal climate variability in MPI-ESM1.2-HR may result from this lack of convection ~~as~~, which prevents atmospheric internal climate variability ~~in the atmosphere does not easily propagate from propagating~~ into the deep ocean.

~~Both~~ The weaker stratification in IPSL-CM6A-LR and UKESM1-0-LL ~~seem to be prone to convection, with than in~~ MPI-ESM1.2-HR indicates the presence of more convective mixing, as convection mixes cold and salty water produced by sea-ice formation with warmer water at depth. Consequently, both IPSL-CM6A-LR and UKESM1-0-LL exhibit more realistic temperature profiles than MPI-ESM1.2-HR in the Amundsen Sea. All the IPSL-CM6A-LR members are nonetheless cold biased at depth (weakest bias of -0.75°C at 900 m depth in Fig. 2k), while all the UKESM1-0-LL members are warm biased (weakest bias of $+0.54^{\circ}\text{C}$ at 900 m depth in Fig. 2l). The spread across the ensemble is large for both models, with 0.79°C (IPSL-CM6A-LR) and 0.39°C (UKESM1-0-LL) difference in the 1995-2014 mean temperature at 900 m between the extreme members.

These conclusions remain valid for 60-year averages ~~instead of as well as~~ 20-year averages, albeit with attenuated internal climate variability. For example, there is still 0.43°C (IPSL-CM6A-LR) and 0.34°C (UKESM1-0-LL) difference in the 1955-2014 mean temperature at 900 m between the extreme members (~~not shown~~Fig. C1). This finding is consistent with a strong internal climate variability at multi-decadal time scales in the Amundsen Sea, as previously pointed out by Purich and England (2021) who identified typical periodicity of approximately 30 years for MPI-ESM1.2-HR, 70 years for IPSL-CM6A-LR and 120 years for UKESM1-0-LL (their Fig. S6). In comparison, paleoclimate reconstructions indicate a ~ 50 -year period for the wind variability at the Amundsen Sea shelf break (Holland et al., 2022).

3.2 Atmospheric internal climate variability

The SMB is defined as the difference between precipitation (liquid and solid, positive contribution) and evaporation, sublimation and ~~run-off~~runoff (negative contribution). The present-day Antarctic SMB mostly consists of snowfall, a small part of which ($<10\%$) is sublimated at the surface and in blowing snow (Van Wessem et al., 2018; Agosta et al., 2019; Mottram et al., 2021). ~~Run-off~~Runoff is currently negligible as most of the meltwater refreezes due to cold temperatures. By 2100 and for the SSP2-4.5 medium scenario, ~~run-off~~runoff is supposed to remain limited (Kittel et al., 2021), so the SMB is projected to increase largely due to the increased water vapour saturation in warmer air (~~e.g. Krinner et al., 2008; Agosta et al., 2013~~). ~~Here,~~




Figure 2. Comparison of the saline and thermal properties of the CMIP6 models MPI-ESM1.2-HR (left), IPSL-CM6A-LR (middle) and UKESM1-0-LL (right). (a-c) standard deviation of the 1995-2014 mean practical salinity across the ensemble relative to the multi-member mean, considering the salinity averaged from 200 m to 700 m depth. The 1500 m isobath (pink) delimits the continental shelf. (d-f) same as (a-c) but for potential temperature and with the 1500 m isobath in blue. (g-i) mean vertical profiles of practical salinity on the Amundsen Sea continental shelf (as defined in Cailliet et al., 2023). For each model, the blue curves represent the individual members (1995-2014 mean), and the red line the multi-member mean. The grey curve corresponds to the 2018 World Ocean Atlas data (WOA 2018, Boyer et al., 2018) over the period 1995-2017 and the black curve to observational climatology based on the WOA, EN4 and MEOP datasets and built for ISMIP6 (Jourdain et al., 2020). (j-l) same as (g-i) but for potential temperature.




Figure 3. Comparison of the saline, thermal and sea-ice properties of observations (left) and CMIP6 models MPI-ESM1.2-HR (middle left), IPSL-CM6A-LR (middle right) and UKESM1-0-LL (right). (a) 1995-2017 mean practical salinity from the 2018 World Ocean Atlas datasets (WOA 2018, Boyer et al., 2018), considering the salinity averaged from 200 m to 700 m depth. (b-d) 1995-2014 mean practical salinity across the climate model ensemble, considering the salinity averaged from 200 m to 700 m depth. The 1500 m isobath (pink) delimits the continental shelf. (e) same as (a) but for potential temperature. (f-h) same as (b-d) but for potential temperature and with the 1500 m isobath in blue. (f) 1995-2014 mean sea-ice concentration from NSIDC dataset (version 4.0) (Comiso, 2023). (j-l) same as (b-d) but for sea-ice concentration.

260 ~~we~~, resulting in more precipitation (e.g. Krinner et al., 2008; Agosta et al., 2013). We therefore focus on ~~SMB, precipitations and air temperatures~~ variability in emulated SMB and its main components such as precipitation and air temperature.

In contrast to the ocean, the atmospheric internal climate variability is relatively similar in the three selected CMIP6 models (Fig. 4). This is partly due to similar emulated present-day SMB: the integrated value over the whole ice sheet ranges between 2641 and 2892 Gt yr⁻¹ for all members of the three models. The present-day SMB internal climate variability is stronger in 265 coastal regions (Fig. 4a-c) where the average SMB is higher (~~not shown~~), consistently Fig. D1, consistent with the precipitation variability in the CMIP simulations (Fig. 4g-i).

The largest SMB ~~internal climate~~ variability is simulated along the coast of the Amundsen and Bellingshausen seas, which ~~is due both to the particularly high present-day mean SMB and to the high results from the high~~ internal climate variability of atmospheric circulation (e.g., Amundsen Sea Low position) and air temperature in these regions (Fig. 4d-i). UKESM1-0-LL 270 also exhibits significant variability in the Dronning Maud region. As previously reported by Marshall and Thompson (2016), the internal climate variability of ~~sea level~~ sea-level pressure and air temperature have the typical characteristics of the two Pacific-South American modes (usually referred to as PSA1 and PSA2), which are associated with wave trains originating in the tropical Pacific and possibly modulated by feedbacks with clouds and ~~sea-ice~~ sea ice (Wang et al., 2022).

3.3 Impact of internal climate variability on the Antarctic contribution to sea level

275 In our ice-sheet projections, Antarctica gains mass over the century for all members of the three CMIP models, with an estimated SLC in 2100 ranging from -1.34 to -8.46 cm (Fig. 5a). This contribution results from a compensation between (i) increased ice mass flux through the grounding line foreed-driven by the ocean (Fig. 5b), mainly occurring in West Antarctica (Fig. 5h) and (ii) increasing SMB (Fig. 5c), occurring in ~~each region~~ all regions for almost all members (Fig. 5f,i,l).

Regions behave in different ways. While the East Antarctica and the Peninsula gain mass (SLC in 2100 ranging respectively 280 from -3.80 to -6.32 cm in Fig. 5d and from -0.96 to -2.24 cm in Fig. 5j), West Antarctica loses mass (SLC in 2100 ranging from +0.11 to +3.78 cm in Fig. 5g). The West Antarctic positive SLC is mostly explained by the dynamical response of Pine Island and Thwaites ice shelves (~ 3 cm in Fig. 6c, basin 11) as well as Getz ice shelf (~ 1 cm, basin 10). The absolute trends in East Antarctica and the Peninsula regions are largely ~~contaminated~~ influenced by the unforced drift in Elmer/Ice (see section 2.2), but the simulations can still inform on the sensitivity to internal climate variability.

285 Internal climate variability affects the estimated SLC of Antarctica in 2100 by more than 45%, 79% and 93% for the ~~IPSL, UKESM and MPI~~ IPSL-CM6A-LR, UKESM1-0-LL and MPI-ESM1.2-HR models respectively (considering the difference between the lowest and highest member divided by the multi-member mean). Thus, the estimated SLC can vary by 1.64 cm, 4.35 cm and 2.33 cm, respectively (Fig. 5a) (considering the difference between the lowest and highest member). This uncertainty is comparable to that associated with the selection of the CMIP6 model (3 cm, Fig. 5a).

290 For the three climate models and in most Antarctic regions, the effects of atmospheric internal climate variability overwhelm the effects of oceanic internal climate variability (Figs. 5-6). ~~This is also true for West Antaretica where the SLC trend mostly results from the ocean-induced dynamics of the ice sheet but internal climate variability results from the atmosphere. On average~~ On average, by the end of the century, the amplitude of SLC variability ~~relative~~ related to the atmosphere (Fig. 5c) is




Figure 4. Comparison of the internal climate variability in surface mass balance, air temperature, precipitation and [sea-level-sea-level](#) pressure in MPI-ESM1.2-HR (left), IPSL-CM6A-LR (middle) and UKESM1-0-LL (right). (a-c) standard deviation of the 1995-2014 mean SMB across the ensemble relative to the multi-member mean, from the MAR-based reconstructions. (d-f) same as (a-c) but for air temperature at 2 m (directly from the CMIP6 outputs). (g-i) same as (a-c) but for total precipitation (shaded) and [sea-level-sea-level](#) pressure (contours every 5 hPa) from the CMIP6 outputs.

3.4 times higher than that ~~relative-related~~ to the ocean ~~-(Fig. 5b)~~. However, there are significant spatial variations across the individual basins and CMIP models.

The West Ross, Getz and Amundsen basins (No. 9,10,11 in Fig. 6) show the most significant atmospheric and oceanic variability in the WAIS region. For the IPSL-CM6A-LR model, internal oceanic variability even exceeds atmospheric variability in these basins (Fig. 6b-c). As described in the previous paragraphs, this variability results from competition of CDW intrusions and convective mixing on the continental shelf (subsect. 3.1), and from the atmospheric circulation, especially the varying position of the Amundsen Sea Low depending on the members (subsect. 3.2). It should be noted that the MPI-ESM1.2-HR model does not show any internal oceanic variability, as expected from the analyses carried out in subsect. 3.1.

In East Antarctica, the Totten basin, which is currently experiencing the highest melt rates in East Antarctica (Rignot et al., 2019), and the Dronning Maud basin (No. 5 and 1 in Fig. 6) show strong internal oceanic variability reaching or exceeding the internal atmospheric variability for the three CMIP6 models. The other basins, like those of the Peninsula, show low basal melting and are largely dominated by internal atmospheric variability, induced primarily by interconnections with the tropical Pacific (see subsect. 3.2).

4 Discussion

4.1 Robustness of internal climate variability in climate models

Since all the diagnoses we have done are based on CMIP models, the realism of their internal climate variability needs to be addressed.

Parsons et al. (2020) compared the distribution of standard deviation of global mean surface air temperature of CMIP piControl simulations to paleoclimate proxies representative of the 1450-1849 period (PAGES2k, 2019). While some of the CMIP6 models had a high-biased temperature variability, the three models used in this study are within the observational plausible range [0.03;0.15], with standard deviation (for variability beyond 25-year timescales) of 0.12°C in IPSL-CM6A-LR, 0.09°C in UKESM1-0-LL and 0.08°C in MPI-ESM1.2-HR.

However, based on ice core reconstructions of temperatures at the surface of Antarctica over the past 1,000 years, Casado et al. (2023) estimated that the internal climate variability was underestimated over Antarctica in the CMIP5 and CMIP6 models, although the three models used here were not part of the assessment. Previdi and Polvani (2016) suggested that the SMB interannual internal climate variability is well captured by the CMIP5 models, but this is only based on the reanalysis period and is therefore more relevant for the interannual variability than for the multi-decadal variability that is emphasized in our work. The higher fidelity of the internal climate variability in CMIP models at the interannual frequency than at multi-decadal frequencies was ~~also-indeed~~ reported by Cheung et al. (2017) for the main modes of variability in the Atlantic and Pacific oceans. Both IPSL-CM6A and MPI-ESM1.2-HR have an internal variability of their 20-year mean surface air temperature close to the CMIP6 multi-model median (Appendix B), so their atmospheric multi-decadal variability is possibly underestimated given the results of Casado et al. (2023). Nevertheless, this variability is significantly stronger in UKESM1-0-LL, which suggests that our study may cover realistic atmosphere internal variability.




Figure 5. Antarctic Sea Level Contribution (SLC) over the 21st century relative to year 2015 under the SSP2-4.5 scenario, for MPI-ESM1.2-HR (green), IPSL-CM6A-LR (purple) and UKESM1-0-LL (blue). Results are displayed for the whole ice sheet (upper row) and for the main sub-regions (as defined in The IMBIE Team, 2018). The left rows show the combination of the dynamical ice-sheet contribution (modulated by the oceanic internal climate variability, middle row) and the surface mass balance contribution (modulated by the atmospheric internal climate variability, right row). The dynamical contribution is calculated from the change in volume above flotation minus the accumulated SMB changes, using the method described in Goelzer et al. (2020) to convert to sea level sea level variations. The SMB contribution is calculated over the grounded ice area of BedMachine-Antarctica-v2, which is very close to Elmer/Ice’s initial state (difference of less than 0.1% in grounded area). The use of the grounded ice area from BedMachine-Antarctica-v2 instead of the one from Elmer/Ice (which takes into account the grounding line retreat), impacts the SLC due to the SMB by less than 1 mm. The solid line represents the multi-member mean, while the shaded area represents the range of values covered by the ensemble members. The number in bracket refers to the number




Figure 6. Regional Antarctic Sea Level Contribution (SLC) in 2100 relative to year 2015 for MPI-ESM1.2-HR (green), IPSL-CM6A-LR (purple) and UKESM1-0-LL (blue), integrated over the IMBIE drainage basins shown in (c) The IMBIE Team (2018). a) all contributions, b) dynamical ice-sheet contribution modulated by the ocean internal climate variability and d) the SMB contribution modulated by the atmospheric internal climate variability (see methods and definitions in the caption of Fig. 5). The box plots correspond to the ensemble median (line), interquartile range (box) and total range (whiskers) of each model.

Our results also show that the amplitude of oceanic internal climate variability around Antarctica strongly depends on the climate model. When compared with 12 other CMIP6 models (Appendix B), the three selected models cover the whole range of oceanic multi-decadal variability in the CMIP6 ensemble, with one of the lowest values (MPI-ESM1.2-HR), one close to the multi-model median (UKESM1-0-LL) and one of the highest values (IPSL-CM6A-LR). The low variability of the ~~MPI-MPI-ESM1.2-HR~~ model is inconsistent with ~~observations. This may result from a poor representation of some processes, in particular an underestimation of dense water formation and freshwater fluxes from basal melting, as well as underestimated sea ice production, which results in a lack of deep convection on the continental shelf and prevents any exchange between the mid-depth ocean and the atmosphere~~ the temperature and salinity profiles observed in the Amundsen Sea (Dutrieux et al., 2014; Jenkins et al., 2018), which likely results from model biases (see subsect. 3.1). Based on an eddy resolving 1/4° ocean model, Sérazin et al. (2017) have estimated that chaotic ~~We therefore consider that the plausible range of oceanic variability is covered by IPSL-CM6A-LR and UKESM1-0-LL. It is nonetheless important to keep in mind that these CMIP models do not resolve eddies, which have been suggested to generate substantial~~ low-frequency oceanic internal climate variability ~~overwhelms the atmospheric forced oceanic variability~~ variability in the Southern Ocean. ~~This suggests that the multi-decadal oceanic internal climate variability of coarser resolution models, like those considered here, could be underestimated. (Sérazin et al., 2017).~~

Furthermore, climate models do not explicitly include the ice sheet, ~~although even though~~ the non-linearities due to ice-sheet–ocean and ice-sheet–atmosphere interaction have the potential to generate internal climate variability (Kravtsov et al., 2007; Gwyther et al., 2018). To capture the full uncertainty due to internal climate variability, ice-sheet models would ideally be fully coupled to climate models and be run for multiple members. Although still challenging (Smith et al., 2021), this would enable a consistent representation of internal climate variability, including the effects of ice-sheet–ocean and ice-sheet–atmosphere feedbacks.

Therefore, the low-frequency internal climate variability that affects the ice-sheet mass through oceanic and atmospheric pathways is probably underestimated in current climate models and its impact on the Antarctic SLC as well.

350 **4.2 Internal climate variability as a source of uncertainty in ~~sea-level~~ sea-level projections**

~~Our simulations, involving three climate models, suggest~~ The comparison of the amplitude of SLC in 2100 due to internal variability (shaded area in Fig. 5a) with the one due to the choice of climate model (difference between extreme thick lines, Fig. 5a) shows that the choice of climate model and internal climate variability both have a similar impact on Antarctic SLC; ~~although there are disparities on a finer scale.~~ The relative importance of internal climate variability in our simulations (45-355 93%) is higher than the 18-21% reported by Tsai et al. (2020). However, its absolute importance is lower in our simulations, with a 2015-2100 SLC modulated by 1.6 to 4.4 cm, versus 8 cm in Tsai et al. (2020). This is likely due to the fact that the SLC projections of Tsai et al. (30 to 48 cm) are at the very high end of the ensemble of other ice-sheet projections (Seroussi et al., 2020; Payne et al., 2021; Coulon et al., 2023), which is partly due to the parameterised ice-shelf hydrofracturing and ice cliff failure in Tsai et al. (2020) as opposed to the aforementioned other models. In contrast, our simulations are ~~towards~~ towards ~~the at the very~~ low end of the ~~spectrum~~ ensemble of other ice-sheet projections (-8.5 to -1.3 cm), Fig. 5a). This is partly

due to the ~~moderate scenario (SSP2-4.5), although the scenario has only a limited impact on Antarctic projections to 2100 (Seroussi et al., 2020; Fox-Kemper et al., 2021), and to the~~ present-day drift in East Antarctica and Peninsula that we did not remove from our projected trends as opposed to the aforementioned other models.

365 The anomaly method used to build the ocean and atmospheric forcing in both our experiments and in ISMIP6 (Nowicki et al., 2020) was designed to correct biases in SMB and ~~ocean-ocean-induced~~ melting over the 1995-2014 period. However, given the ~~high variability of wide confidence interval on a~~ 20-year means ~~−~~ $[-0.06^{\circ}\text{C}; 0.24^{\circ}\text{C}]$ for air temperature and $[0.02^{\circ}\text{C}; 0.12^{\circ}\text{C}]$ for oceanic temperature, see Fig. B1. correcting a random phase of the historical CMIP simulations towards the actual 1995-2014 period may significantly shift the projections. For example, members with colder forcing over the present-day period become warmer throughout the 21st century due to the correction. Casado et al. (2023) recommend averaging over 50-years to be long enough to weaken internal climate variability and short enough not to dilute forced trends. This corresponds to the typical 370 period of internal climate variability in the paleoclimate reconstructions (Holland et al., 2022). As discussed in subsect. 3.1, some models like UKESM1-0-LL nonetheless have internal climate variability over longer periods, so that 50-year averages do not attenuate internal climate variability to a significant extent. Another issue with extending the period over which the correction is applied is that not so many observations were available 50 years ago in Antarctica.

375 Given the difficulties of correcting biases, it is tempting to select the members that are most in phase with observations and not ~~applying to apply~~ any bias correction, which is investigated in the next subsection. It is nonetheless important to consider that the anomaly method is only responsible for a part of the uncertainty associated with internal climate variability. Indeed, Tsai et al. (2020) highlighted important internal climate variability despite correcting the 1920–2012 period. As ice-sheet ~~modellers-modelers~~ sometimes run large ensemble simulations to select or weight the members that best fit observational 380 records (e.g., Coulon et al., 2023), it seems important that they either consider multiple climate model members or select the more realistic ones.

4.3 Identifying the best member

For greater confidence in the ice-sheet projections, the models have to be initialised and calibrated to match historical observations. Given the importance of internal ~~climate variability, it is tempting to select the variability, selecting the CMIP~~ member that is 385 most in phase with the observational record ~~−We here might be useful to achieve it. Such a member could also be primarily used for projections when running multiple members is too computationally expensive. Here, we~~ investigate this possibility with the example of the IPSL-CM6A-LR ~~model and its 33 members, and UKESM1-0-LL models.~~

The first challenge is to define metrics that can be used to quantify the phasing of individual members. Among the observations that are available over several decades, it is somehow an expert judgement to decide which metrics are most relevant for 390 the Antarctic mass variations.

~~Herewe propose to calculate the following metrics for individual members, in order to assess the complexity of identifying a best member~~ Here, we choose several metrics to ensure:

- 395 – a good representation of the mean atmospheric and oceanic states. We selected variables directly used to drive the ice-sheet model, such as SMB for the atmosphere and temperature for the ocean. For the ocean, we focused our analyses on the Amundsen sector as the region experiences the current main mass loss and CTD profile data are available for a relatively long period from 1994 to 2018.
- a good representation of the amplitude of oceanic variability using the same observational data described in the previous paragraph. We did not evaluate the variability of SMB since it has been relatively stable in recent years.
- 400 – a good representation of important modes of variability known to affect the ocean and atmosphere in/around Antarctica. We focus our analyses on the indices representative of the Southern Annular Model and the Interdecadal Pacific Oscillation.
- a good phasing of internal variability with observations, which could be important for future detection/attribution studies and for projected Antarctic contribution to sea-level rise. We chose two variables, sea-ice concentration and the presence of warm periods on the continental shelf of the Amundsen Sea to provide insights on the phasing of internal variability.

405 The metrics definition and the rank of all members are presented in Appendix E. Overall, ranks are not very consistent across the chosen metrics, and no member is best for all metrics. Although the perfect member does not exist, some members nevertheless seem more in phase with the observed climate variability than the other members. For the IPSL-CM6A-LR model, member 26 seems to be the most consistent with the observed variability despite a lack of variability in front of Pine Island and a sea-ice trend that is mostly negative as opposed to the positive observed trend. For the UKESM1-0-LL model, member 4
410 seems to be the most consistent with the observed variability despite an overestimated SAM trend and SMB but also a negative sea-ice trend.

However, the member selection appears very sensitive to the list of chosen metrics, and the phasing of the best member is only marginally better than for the other members. Given the number of degrees of freedom of climate models, it would probably be unrealistic to expect finding a member perfectly in phase with the observed variability among ensembles of a few
415 tens of members, even if models were not biased. For these reasons, it appears judicious to consider several climate model members in ice-sheet projections, to account for the substantial uncertainty related to internal climate variability.

For the same reason, the initialisation of ice-sheet models should account for internal climate variability, either by starting from various members and/or by including internal climate variability in the long initialisation of some ice-sheet models, as previously suggested by Robel et al. (2023).

420 5 Conclusions

In this study, we show that internal climate variability affects the Antarctic contribution to changes in sea level until 2100 by 45%-93%, i.e., a variation between 1.6 and 4.4 cm under the SSP2-4.5 scenario. This may be a low estimate as the internal climate variability of the CMIP models is likely underestimated. In our case, the uncertainty in Antarctic contribution to sea level due to internal climate variability is of comparable magnitude as the uncertainty related to the choice of the climate model.

425 The internal climate variability has a strong multi-decadal component, so that (i) it is not completely diluted over a century, and (ii) it strongly affects the 20-year averages used to build the forcing anomaly.

The effect of atmospheric internal climate variability on the surface mass balance overwhelms the effect of oceanic internal climate variability on the dynamical ice-sheet mass loss by a factor of 2 to 5, except in Amundsen, Getz and Aurora basins where both contributions may be similar depending on the CMIP model.

430 The atmospheric internal climate variability over Antarctica has similar amplitudes in the three CMIP6 models analysed in this study. Conversely, the amplitude of oceanic internal climate variability around Antarctica strongly depends on the climate model. The oceanic internal climate variability in the MPI-ESM1.2-HR model is very low, which may be explained by underestimated ocean convective mixing on the continental shelf, due to either biases in the sea-ice behaviour or in the ocean stratification.

435 From these results, we recommend using the following practices for future ice-sheet projections:

– A CMIP model selection based on the assessment of the model ability to produce a plausible multi-decadal variability in both the atmospheric and oceanic drivers of ice-sheet changes (in addition to the usual mean state assessment, which should ideally be done for multiple members).

440 – The consideration of several members for each climate model forcing given the difficulty or impossibility to identify a perfect member (2 or 3 members can already be very informative if running more is too computationally expensive).

– The use of longer reference period for the calculation of anomalies than that usually used (e.g., 20 years in ISMIP Nowicki et al., 2020) as climate models show important modes of variability longer than 20 years. Casado et al. (2023) recommend averaging over 50 years to be long enough to weaken internal climate variability and short enough not to dilute forced trends. Few observations were available 50 years ago in Antarctica, so the observational climatologies will likely remain representative of 20-30 years. This nonetheless likely remains a preferable approach than using the last 20 years.

445

Code and data availability. The ice-sheet model version and set of parameters used to run our experiments, the SMB reconstructions and the python scripts used to build the figures will be provided in a zenodo repository after the review.

Appendix A: Assessment of mean oceanic properties in the CMIP6 models

450 The present-day oceanic properties of multiple CMIP6 models are assessed through a review of three studies which evaluate
water masses properties in the Southern Ocean and Antarctic seas (Purich and England, 2021), oceanic and atmospheric metrics
relevant for the Southern Ocean dynamics (Beadling et al., 2020), and bottom properties in the Southern Ocean (Heuzé, 2021)
. Each of these studies provides the bias of several CMIP model variables. We normalise the bias of individual variables by
the multi-model standard deviation, and we rank the models based on the increasing RMSE calculated over the variables of a
given study (Fig. A1). Tab. A1 details the variables and observations used to estimate the model biases. The analysis is done
455 here for the first available member of each model.




Figure A1. Assessment of Southern Ocean and Antarctic seas properties in the CMIP6 models. In each panel, CMIP models are ranked by increasing RMSE. (a) Antarctic Shelf Bottom Water (ASBW) and Circumpolar Deep Water (CDW) temperature biases compared to temperatures from Schmidtko et al. (2014) in the Ross, Amundsen/Bellingshausen, Weddell and Cosmonauts seas (Purich and England, 2021, their Fig. S16). (b) Biases in the Southern Ocean metrics defined in Tab. 1 of Beadling et al. (2020) with respect to the observational estimates describing several characteristics of the Antarctic Circumpolar Current, zonal wind stress strength, location and curl, as well as meridional gradients of water mass properties. (c) RMSE of bottom ocean properties (density, salinity and temperature) in the Southern Ocean (Heuzé, 2021, their Figs. 1, A1, A2). For each panel, the metric is normalised by the CMIP6 multi-model standard deviation. Selected models are labelled with a star.

Table A1. List of metrics used for ranking CMIP6 models. Evaluating period and observational dataset to which CMIP6 models are compared are indicated in the two last columns.

<u>Metric</u>	<u>Name</u>	<u>Unit</u>	<u>Period</u>	<u>Observations</u>
Purich and England (2021)				
<u>- 22 CMIP6 models</u>				
<u>T_{ASBW}</u>	<u>Temperature of Antarctic Shelf Bottom Water (ASBW)</u>	<u>°C</u>	<u>1975-2012</u>	<u>Schmidtke et al. (2014)</u>
<u>T_{CDW}</u>	<u>Temperature of Circumpolar Deep Water (CDW)</u>	<u>°C</u>	<u>1975-2012</u>	<u>Schmidtke et al. (2014)</u>
Beadling et al. (2020) - 34				
<u>CMIP6 models</u>				
<u>ACC</u>	<u>Volume transport of the Antarctic Circumpolar Current (ACC) through the Drake Passage</u>	<u>Sv</u>	<u>1986-2005</u>	<u>Donohue et al. (2016)</u>
<u>τ_{max}</u>	<u>Maximum zonally averaged zonal wind stress</u>	<u>$N.m^{-2}$</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>lat τ_{max}</u>	<u>Position of the peak wind stress</u>	<u>°S</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>total $\tau^{(x)}$</u>	<u>Zonally averaged westerly wind stress</u>	<u>$10^{12} N$</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>total $\tau^{(x)}$ in DP</u>	<u>Zonally averaged westerly wind stress over the Drake Passage latitudes (55°S-64°S)</u>	<u>$10^{12} N$</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>total WSC in DP</u>	<u>Integrated Wind Stress Curl (WSC) over the Drake Passage latitudes (55°S-64°S)</u>	<u>$10^{12} N.m^{-1}$</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>min WSC</u>	<u>Minimum zonally integrated Wind Stress Curl</u>	<u>Nm^{-2}</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>lat of min WSC</u>	<u>Latitude of minimum zonally integrated Wind Stress Curl</u>	<u>°S</u>	<u>1986-2005</u>	<u>ERA5</u>
<u>$\Delta\rho$</u>	<u>Zonally and full-depth-averaged potential density difference between 65°S and 45°S</u>	<u>$kg.m^{-3}$</u>	<u>1986-2005</u>	<u>WOA18</u>
<u>ΔS</u>	<u>Zonally and full-depth-averaged salinity difference between 65°S and 45°S</u>	<u>(-)</u>	<u>1986-2005</u>	<u>WOA18</u>
<u>ΔT</u>	<u>Zonally and full-depth-averaged potential</u>	<u>°C</u>	<u>1986-2005</u>	<u>WOA18</u>

Appendix B: Atmospheric and oceanic components of the internal climate variability in multiple CMIP6 models

460

In Fig. B1 we briefly show where the three selected CMIP6 models seat in terms of internal climate variability. Based on the analysis of 15 CMIP6 models with more than 10 members, the multi-member standard deviation of 2 m air temperature over the whole Antarctica varies between 0.06°C and 0.24°C. Both IPSL-CM6A-LR and MPI-ESM1.2-HR have a variability close to the median (0.12 and 0.13°C), while the UKESM1-0-LL model is among the models with the highest variability (0.20°C). For the ocean, the multi-member standard deviation of the ocean temperature averaged between 200 and 700 m over the continental shelf varies between 0.02°C and 0.12°C. The MPI-ESM1.2-HR model is one of the models with the lowest ocean variability (0.02°C), UKESM1-0-LL is close to the median (0.04°C) and IPSL-CM6A-LR is one of the models with the highest variability (0.07°C).




Figure B1. Assessment of 1995-2014 multi-member mean and standard deviation of Antarctic air temperature at 2m (left) and circum-Antarctic ocean temperature between 200 and 700 m depth on the continental shelf (right) in 15 CMIP6 models. The number of members for each model is in brackets. When two numbers are indicated, they correspond to the available members for the atmosphere and ocean, respectively. The black dashed lines represent the observational means, from the ERA5 atmospheric reanalysis (Hersbach et al., 2020) and from the ISMIP6 observational ocean climatology (Jourdain et al., 2020).

465 **Appendix C: Assessment of oceanic internal climate variability for 60-year period**

In Fig. C1, we repeat the analysis that was presented in the main text and illustrated in Fig. 2, but analysing the multi-model variability of 60-year means (1955-2014) instead of 20-year means (1995-2014). This emphasises that substantial internal climate variability is still present in 60-year averages.




Figure C1. Comparison of the saline and thermal properties of the CMIP6 models MPI-ESM1.2-HR (left), IPSL-CM6A-LR (middle) and UKESM1-0-LL (right). (a-c) standard deviation of the 1955-2014 mean practical salinity across the ensemble relative to the multi-member mean, considering the salinity averaged from 200 m to 700 m depth. The 1500 m isobath (pink) delimits the continental shelf. (d-f) same as (a-c) but for potential temperature and with the 1500 m isobath in blue. (g-i) mean vertical profiles of practical salinity on the Amundsen Sea continental shelf (as defined in Caillet et al., 2023). For each model, the blue curves represent the individual members (1955-2014 mean), and the red line the multi-member mean. The grey curve corresponds to the 2018 World Ocean Atlas data (WOA 2018, Boyer et al., 2018) over the period 1955-2017. (j-l) same as (g-i) but for potential temperature.

Appendix D: Atmospheric mean properties

470 [Here we evaluate the mean atmospheric state of the three selected CMIP6 models in comparison to the ERA5 atmospheric reanalysis \(Hersbach et al., 2020\).](#)




Figure D1. [Comparison of mean surface mass balance, air temperature, precipitation and sea-level pressure from the ERA5 reanalysis \(left\) and the three CMIP6 models: MPI-ESM1.2-HR \(middle left\), IPSL-CM6A-LR \(middle right\) and UKESM1-0-LL \(right\).](#) (e) [1995-2014 mean air temperature at 2 m from ERA5 reanalysis dataset \(Hersbach et al., 2020\).](#) (i) [same as \(e\) but for total precipitation \(shaded\) and sea-level pressure \(contours every 5 hPa\).](#) (b-d) [1995-2014 mean emulated surface mass balance \(SMB\) across the climate model ensemble.](#) (f-h) [same as \(b-d\) but for air temperature at 2 m.](#) (j-l) [same as \(b-d\) but for total precipitation \(shaded\) and sea-level pressure \(contours every 5 hPa\).](#)

Appendix E: Best member ranking

Given the importance of internal climate variability, it is tempting to select the member that is most in phase with the observational record. Here we calculate the following metrics for individual members:

- 475 – Root mean squared difference between the multi-year mean ocean temperature profiles measured and modelled in front of Pine Island (years 1994, 2000, 2007, 2009, 2010, 2012, as given in Dutrieux et al., 2014) and Dotson (years 2000, 2006, 2007, 2009, 2011, 2012, 2014, as given in Jenkins et al., 2018) ice shelves. This is a proxy for the phase of multi-decadal variability in the region where the ocean has triggered the largest ice-sheet mass loss.
- Root mean squared difference between the standard deviation of multi-year ocean temperature profiles measured and
480 modelled in front of Pine Island and Dotson ice shelves. This is a proxy for the amplitude of multi-decadal variability in the region where the ocean has triggered the largest ice-sheet mass loss.
- Difference between observed and modelled trend in the Southern Annular Mode (SAM), estimated over 1965-2014 based on the index defined by Marshall (2003). SAM affects both the Antarctic SMB (Medley and Thomas, 2019) and ice-shelf basal melting (Verfaillie et al., 2022). Here we evaluate the phase of multi-decadal variability in individual
485 members by quantifying the modulation of the SAM 60-year trend by internal climate variability.
- Pearson correlation coefficient, root mean squared difference between and standard deviation of the observed and modelled Southern Annular Mode index (SAM, Marshall, 2003) with a 5-year running window on detrend data over 1965-2014. SAM index is based on the zonal pressure difference between the latitudes of 40°S ~~an~~ and 65°S and is a proxy
490 for the phase and amplitude of inter-annual variability over all Antarctica as SAM has a large impact on atmospheric and oceanic circulations. Taylor diagram combines these 3 metrics to quantify the degree of correspondence between modelled and observed SAM.
- Pearson correlation coefficient, root mean squared difference between and standard deviation for the observed and modelled Tripole Index for the Interdecadal Pacific Oscillation (TPI, Henley et al., 2015) with a 5-year running window over
495 1854-2014. The TPI is based on the difference between the Sea Surface Temperature Anomalies (SSTA) averaged over the central equatorial Pacific and the average of the SSTA in the Northwest and Southwest Pacific. The TPI describes decadal to interdecadal changes in the strength of the El Niño–Southern Oscillation (ENSO) and its teleconnections. ENSO affects the West Antarctic SMB (Genthon and Cosme, 2003; Scott et al., 2019) and ice-shelf basal melting in the Amundsen Sea (Steig et al., 2012; Holland et al., 2019) through the south-eastward propagation of atmospheric Rossby waves from the inter-tropical Pacific. Taylor diagram combines these 3 metrics to quantify the degree of correspondence
500 between modelled and observed TPI.
- Comparison of the mean ocean temperature at 750 m depth on the continental shelf in the Amundsen Sea between identified warm periods and the preceding cold periods. Three warm periods have been identified in observations: 1945±12 (1933-1957), 1970±4 (1966-1974) (based on sediment records, Smith et al., 2017) and 2006-2012 (based on Dotson and

505 Pine Island melt rates estimates, Dutrieux et al., 2014; Jenkins et al., 2018). These periods are compared respectively to 1850-1932, 1958-1965 and 1975-2005, supposed to be colder periods. For each member, we assume that the warm period exists if a 5-year mean, at least, within years that define the warm period is higher than the mean of the preceding cold period. This is a proxy for the phase of multi-decadal variability in the region where the ocean has triggered the largest ice-sheet mass loss.

- 510 – Root mean squared difference between observed and modelled sea-ice concentration trend around Antarctica over 1979-2014 (version 4, Meier et al., 2022). Here, we evaluate the phase of multi-decadal variability in individual members by quantifying the modulation of sea-ice concentration trend by internal climate variability (Zhang et al., 2019).
- Root mean squared difference between the 1995-2014 average SMB output of the MAR simulations forced by ERA5 described in Kittel et al. (2021) and the reconstructed SMB of each member of the IPSL-CM6A-LR model on the same period as described in Jourdain et al. (2024).

515 We then rank the performance of all the members by assigning them a rank, with lower rank for the member that best matches the observations):

- for root mean squared difference (RMSE) metric, the member with the lowest (respectively highest) RMSE value is assigned rank 1 (respectively rank 33).
- 520 – for metrics relative to Taylor Diagram (Pearson correlation coefficient and standard deviation), we first calculate a rank for each individual metric by assigning the lowest rank value to the lowest Pearson correlation coefficient and to the lowest difference between observed and modelled standard deviation. We then average all calculated ranks and finally assign the best (worst) final rank to the lowest (highest) average.
- for metric relative to warm and cold period alternation, we assume that the warm period exists if a 5-year mean, at least, within years that define the warm period is higher than the mean of the preceding cold period. If the condition is met, 525 the member is assigned the value 1, and 0 otherwise. This process is applied to each of the three warm periods and the values are then summed. Members with a value of 3 (of 0) are assigned the best (the worst) rankings.

The ranks of all members for all metrics are presented in Fig. E1 .~~Overall, ranks are not very consistent across the chosen metrics, and no member is best for all metrics. Although the perfect member does not exist, some members nevertheless seem more in phase with the observed climate variability than the other members. We have ranked the members according to the weighted average rank of each metric (weight of 1) (see top blue value on Fig. E1). Member 26 seems to be the most consistent with the observed variability despite a lack of variability in front of Pine Island and a sea-ice trend that is mostly negative as opposed to the positive observed trend. for IPSL-CM6A-LR and UKESM1-0-LL models.~~

530

~~However, the member selection appears very sensitive to the list of chosen metrics, and the phasing of the best member may be only marginally better than for the other members. Given the number of degrees of freedom of climate models, it would probably be unrealistic to expect finding a member perfectly in phase with the observed variability among ensembles of a few~~

535




Figure E1. Combination of relevant metrics for evaluating members of the [IPSL-CM6A-LR model \(top\)](#) and [UKESM1-0-LL model \(bottom\)](#) compared to various observations/index data. The y-axis represents the selected metrics while the x-axis represents individual members. A rank of 1 (respectively 33 [or 16](#)) refers to the member with the value closest to (respectively furthest from) the assessed observational value. The blue numbers at the top indicate the member’s overall ranking, with equal weight for all metrics.

~~tens of members, even if models were not biased. For these reasons, it appears judicious to consider several climate model members in ice-sheet projections, to account for the substantial uncertainty related to internal climate variability.~~

540 ~~In this study, we show that internal climate variability affects the Antarctic contribution to changes in sea level until 2100 by 45%-93%, i.e., a variation between 1.6 and 4.4 cm under the SSP2-4.5 scenario. This may be a low estimate as the internal climate variability of the CMIP models is likely underestimated. In our case, the uncertainty in Antarctic sea-level contribution due to internal climate variability is of comparable magnitude as the uncertainty related to the choice of the climate model. The internal climate variability has a strong multi-decadal component, so that (i) it is not completely diluted over a century, and (ii) it strongly affects the 20-year averages used to build the forcing anomaly.~~

545 ~~For all three climate models and for most Antarctic regions, the effect of atmospheric internal climate variability on the surface mass balance overwhelms the effect of oceanic internal climate variability on the dynamical ice-sheet mass loss by more than a factor of 3. The atmospheric internal climate variability is similar in the three CMIP6 models analysed in this study.~~

Conversely, the amplitude of oceanic internal climate variability around Antarctica strongly depends on the climate model. The oceanic internal climate variability of the MPI-ESM1.2-HR model is very low, which may be explained by underestimated convection, due to either biases in the sea-ice behaviour or in the ocean stratification.

550 From these results, we recommend changing a number of practices in ice-sheet projections:-

- Several members within each climate model should be considered as forcing: given the number of degrees of freedom of climate models, it would probably be unrealistic to expect finding a member perfectly in phase with the observed variability among ensembles of a few tens of members, even if models were not biased, so it is important to account for this uncertainty.

555 - The reference period for the calculation of anomalies or weights should be longer than the 20 years used here and in ISMIP6 (Nowicki et al., 2020). Casado et al. (2023) recommend averaging over 50 years to be long enough to weaken internal climate variability and short enough not to dilute forced trends. An issue with extending the period over which the correction is applied is that only few observations were available 50 years ago in Antarctica, so it is possible that this approach will reduce the expression of internal climate variability in the model but not in the observations. This
560 nonetheless likely remains a preferable approach than using the last 20 years.

- The initialisation of ice-sheet models should account for internal climate variability, either by starting from various members and/or by including internal climate variability in the long initialisation of some ice-sheet models, as previously suggested by Robel et al. (2023).

565 - Ice-sheet models should be fully coupled to climate models and be run for multiple members. Although still challenging (Smith et al., 2021), this would enable a consistent representation of internal climate variability, including the effects of ice-sheet-ocean and ice-sheet-atmosphere feedbacks.

Author contributions. JC and NCJ designed the overall study. JC wrote the initial draft and NCJ did the first review. JC ran the Elmer/Ice experiments with valuable inputs from PM, FGC, BU and MC. NCJ compiled the intercomparison of CMIP6 ocean models. NCJ, CA and CK calculated the multi-member surface mass balance. JC and CB worked on the basal melting parameterisation. All authors contributed to the final manuscript.

Competing interests. The authors declare no competing interests.

Acknowledgements. This study was funded by the French National Research Agency (ANR) under grants ANR-19-CE01-0015 (EIS) and ANR-22-CE01-0014 (AIAI). The work also benefited from the support of the French Government through the France 2030 program managed by ANR (ISCLIM, ANR-22-EXTR-0010). Some of the authors received funding from the European Union's Horizon 2020 research and innovation programme under Grant agreements 820575 (TiPACCs), 101003536 (ESM2025) and 101003826 (CRiceS).

This work was granted access to the high-performance computing (HPC) resources of TGCC under allocations A0140106066 and A0140106035 attributed by GENCI. The CMIP6 data were analysed on the ESPRI-MOD platform thanks to the CLIMERI-France infrastructure.

The authors would like to acknowledge David Ferreira for interesting discussion on sea-ice, Jérôme Servonnat for his valuable help on ESPRI-MOD as well as Hélène Seroussi, Gaël Durand, Jérémie Mougnot and Hartmut Hellmer for their constructive feedbacks.

References

- Adusumilli, S., Fricker, H. A., Medley, B., Padman, L., and Siegfried, M. R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nature geoscience*, 13, 616–620, 2020.
- Agosta, C., Favier, V., Krinner, G., Gallée, H., Fettweis, X., and Genthon, C.: High-resolution modelling of the Antarctic surface mass balance, application for the twentieth, twenty first and twenty second centuries, *Climate dynamics*, 41, 3247–3260, 2013.
- 585 Agosta, C., Amory, C., Kittel, C., Orsi, A., Favier, V., Gallée, H., van den Broeke, M. R., Lenaerts, J., van Wessem, J. M., van de Berg, W. J., et al.: Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979–2015) and identification of dominant processes, *The Cryosphere*, 13, 281–296, 2019.
- Agosta, C., Kittel, C., Amory, C., Edwards, T., and Davrinche, C.: Evaluation of CMIP5 and CMIP6 global climate models in the Arctic and Antarctic regions, atmosphere and surface ocean, in: EGU General Assembly Conference Abstracts, pp. EGU22–11 363, 2022.
- 590 Agosta, C., D. C. K. C. A. C. . E. T.: Evaluation of CMIP5 and CMIP6 global climate models in the Arctic and Antarctic regions, atmosphere and surface ocean, Zenodo, <https://doi.org/10.5281/zenodo.11595213>, 2024.
- Armstrong McKay, D. I., Staal, A., Abrams, J. F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S. E., Rockström, J., and Lenton, T. M.: Exceeding 1.5 C global warming could trigger multiple climate tipping points, *Science*, 377, eabn7950, 2022.
- 595 Aschwanden, A., Bartholomäus, T. C., Brinkerhoff, D. J., and Truffer, M.: Brief communication: A roadmap towards credible projections of ice sheet contribution to sea level, *The Cryosphere*, 15, 5705–5715, 2021.
- Beadling, R. L., Russell, J., Stouffer, R., Mazloff, M., Talley, L., Goodman, P., Sallée, J.-B., Hewitt, H., Hyder, P., and Pandde, A.: Representation of Southern Ocean properties across coupled model intercomparison project generations: CMIP3 to CMIP6, *Journal of Climate*, 33, 6555–6581, 2020.
- 600 Bindschadler, R. A., Nowicki, S., Abe-Ouchi, A., Aschwanden, A., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., et al.: Ice-sheet model sensitivities to environmental forcing and their use in projecting future sea level (the SeaRISE project), *Journal of Glaciology*, 59, 195–224, 2013.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., et al.: Presentation and evaluation of the IPSL-CM6A-LR climate model, *J. Adv. Model. Ea. Sys.*, 12, e2019MS002 010, 2020.
- 605 Boyer, T. P., García, H. E., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Weathers, K. A., Baranova, O. K., Paver, C. R., Seidov, D., et al.: World ocean atlas, 2018.
- Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to the friction law, *The Cryosphere*, 13, 177–195, 2019.
- Burgard, C., Jourdain, N. C., Reese, R., Jenkins, A., and Mathiot, P.: An assessment of basal melt parameterisations for Antarctic ice shelves, *The Cryosphere*, 16, 4931–4975, 2022.
- 610 Caillet, J., Jourdain, N. C., Mathiot, P., Hellmer, H. H., and Mougnot, J.: Drivers and reversibility of abrupt ocean state transitions in the Amundsen Sea, Antarctica, *Journal of Geophysical Research: Oceans*, 128, e2022JC018929, 2023.
- Casado, M., Hébert, R., Faranda, D., and Landais, A.: The quandary of detecting the signature of climate change in Antarctica, *Nature Climate Change*, pp. 1–7, 2023.
- 615 Cheung, A. H., Mann, M. E., Steinman, B. A., Frankcombe, L. M., England, M. H., and Miller, S. K.: Comparison of low-frequency internal climate variability in CMIP5 models and observations, *Journal of Climate*, 30, 4763–4776, 2017.

- Comiso, J. C.: Bootstrap Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS, Version 4, <https://doi.org/10.5067/X5LG68MH0130>, 2023.
- 620 Coulon, V., Klose, A. K., Kittel, C., Edwards, T., Turner, F., Winkelmann, R., and Pattyn, F.: Disentangling the drivers of future Antarctic ice loss with a historically-calibrated ice-sheet model, *EGUsphere*, 2023, 1–42, 2023.
- Dalaiden, Q., Abram, N., and Goosse, H.: Tropical Pacific variability and anthropogenic forcing are the key drivers of the West Antarctic atmospheric circulation variability over the 20th century, Tech. rep., Copernicus Meetings, 2023.
- Donat-Magnin, M., Jourdain, N. C., Kittel, C., Agosta, C., Amory, C., Gallée, H., Krinner, G., and Chekki, M.: Future surface mass balance and surface melt in the Amundsen sector of the West Antarctic Ice Sheet, *The Cryosphere*, 15, 571–593, 2021.
- 625 Donohue, K. A., Tracey, K., Watts, D. R., Chidichimo, M. P., and Chereskin, T.: Mean antarctic circumpolar current transport measured in drake passage, *Geophysical Research Letters*, 43, 11–760, 2016.
- Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., Steig, E. J., Ding, Q., Abrahamsen, E. P., and Schröder, M.: Strong sensitivity of Pine Island ice-shelf melting to climatic variability, *Science*, 343, 174–178, 2014.
- Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J.,
630 et al.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, 2021.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, 2016.
- Fox-Kemper, B., Hewitt, H., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., Slangen, A. B. A., and Yu, Y.: Ocean, Cryosphere and Sea Level
635 Change, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1211–1362, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.011>, 2021.
- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., De Fleurian, B., Greve, R., Malinen, M., Martín, C., Råback, P., et al.: Capabilities and performance of Elmer/Ice, a new-generation ice sheet model, *Geoscientific Model Development*, 6, 1299–1318, 2013.
- 640 Garbe, J., Albrecht, T., Levermann, A., Donges, J. F., and Winkelmann, R.: The hysteresis of the Antarctic ice sheet, *Nature*, 585, 538–544, 2020.
- Genthon, C. and Cosme, E.: Intermittent signature of ENSO in west-Antarctic precipitation, *Geophysical Research Letters*, 30, 2003.
- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, 2012.
- 645 Glen, J. W.: The creep of polycrystalline ice, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 228, 519–538, 1955.
- Goelzer, H., Coulon, V., Pattyn, F., De Boer, B., and Van De Wal, R.: Brief communication: On calculating the sea-level contribution in marine ice-sheet models, *The Cryosphere*, 14, 833–840, 2020.
- Gwyther, D. E., O’Kane, T. J., Galton-Fenzi, B. K., Monselesan, D. P., and Greenbaum, J. S.: Intrinsic processes drive variability in basal
650 melting of the Totten Glacier Ice Shelf, *Nature communications*, 9, 3141, 2018.
- Henley, B. J., Gergis, J., Karoly, D. J., Power, S., Kennedy, J., and Folland, C. K.: A tripole index for the interdecadal Pacific oscillation, *Climate dynamics*, 45, 3077–3090, 2015.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.

- 655 Heuzé, C.: Antarctic bottom water and North Atlantic deep water in CMIP6 models, *Ocean Science*, 17, 59–90, 2021.
- Hill, E. A., Urruty, B., Reese, R., Garbe, J., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Gudmundsson, G. H., Winkelmann, R., Chekki, M., et al.: The stability of present-day Antarctic grounding lines—Part 1: No indication of marine ice sheet instability in the current geometry, *The Cryosphere*, 17, 3739–3759, 2023.
- Hirano, D., Tamura, T., Kusahara, K., Fujii, M., Yamazaki, K., Nakayama, Y., Ono, K., Itaki, T., Aoyama, Y., Simizu, D., et al.: On-shelf
660 circulation of warm water toward the Totten Ice Shelf in East Antarctica, *Nature Communications*, 14, 4955, 2023.
- Hogg, A. M., Penduff, T., Close, S. E., Dewar, W. K., Constantinou, N. C., and Martínez-Moreno, J.: Circumpolar variations in the chaotic nature of Southern Ocean eddy dynamics, *Journal of Geophysical Research: Oceans*, 127, e2022JC018440, 2022.
- Holland, P. R., Bracegirdle, T. J., Dutrieux, P., Jenkins, A., and Steig, E. J.: West Antarctic ice loss influenced by internal climate variability and anthropogenic forcing, *Nature Geoscience*, 12, 718–724, 2019.
- 665 Holland, P. R., O’Connor, G. K., Bracegirdle, T. J., Dutrieux, P., Naughten, K. A., Steig, E. J., Schneider, D. P., Jenkins, A., and Smith, J. A.: Anthropogenic and internal drivers of wind changes over the Amundsen Sea, West Antarctica, during the 20th and 21st centuries, *The Cryosphere*, 16, 5085–5105, 2022.
- Jenkins, A., Shoosmith, D., Dutrieux, P., Jacobs, S., Kim, T. W., Lee, S. H., Ha, H. K., and Stammerjohn, S.: West Antarctic Ice Sheet retreat in the Amundsen Sea driven by decadal oceanic variability, *Nature Geoscience*, 11, 733–738, 2018.
- 670 Jourdain, N. C., Asay-Davis, X., Hattermann, T., Straneo, F., Seroussi, H., Little, C. M., and Nowicki, S.: A protocol for calculating basal melt rates in the ISMIP6 Antarctic ice sheet projections, *The Cryosphere*, 14, 3111–3134, 2020.
- Jourdain, N. C., Amory, C., Kittel, C., and Durand, G.: Changes in Antarctic surface conditions and potential for ice shelf hydrofracturing from 1850 to 2200, *The Cryosphere*, submitted, 2024.
- Kittel, C., Amory, C., Agosta, C., Jourdain, N. C., Hofer, S., Delhasse, A., Doutreloup, S., Huot, P.-V., Lang, C., Fichefet, T., et al.: Diverging
675 future surface mass balance between the Antarctic ice shelves and grounded ice sheet, *The Cryosphere*, 15, 1215–1236, 2021.
- Kittel, C., Amory, C., Hofer, S., Agosta, C., Jourdain, N. C., Gilbert, E., Le Toumelin, L., Vignon, É., Gallée, H., and Fettweis, X.: Clouds drive differences in future surface melt over the Antarctic ice shelves, *The Cryosphere*, 16, 2655–2669, 2022.
- Kravtsov, S., Dewar, W. K., Berloff, P., McWilliams, J. C., and Ghil, M.: A highly nonlinear coupled mode of decadal variability in a mid-latitude ocean–atmosphere model, *Dynamics of atmospheres and oceans*, 43, 123–150, 2007.
- 680 Krinner, G., Guicherd, B., Ox, K., Genthon, C., and Magand, O.: Influence of oceanic boundary conditions in simulations of Antarctic climate and surface mass balance change during the coming century, *Journal of Climate*, 21, 938–962, 2008.
- Lee, J.-Y., Dunne, J. P., Engelbrecht, F., Fischer, E., Fyfe, J. C., Jones, C. A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Chapter 4: Future global climate: scenario-based projections and near-term information, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 553–672,
685 [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research: Solid Earth*, 94, 4071–4087, 1989.
- 690 Marshall, G. J.: Trends in the Southern Annular Mode from observations and reanalyses, *Journal of climate*, 16, 4134–4143, 2003.
- Marshall, G. J. and Thompson, D. W.: The signatures of large-scale patterns of atmospheric variability in Antarctic surface temperatures, *Journal of Geophysical Research: Atmospheres*, 121, 3276–3289, 2016.

- Mathiot, P. and Jourdain, N. C.: Southern Ocean warming and Antarctic ice shelf melting in conditions plausible by late 23rd century in a high-end scenario, *Ocean Science*, 0, 1–27, 2023.
- 695 Mathiot, P., Jenkins, A., Harris, C., and Madec, G.: Explicit and parametrised representation of under ice shelf seas in az* coordinate ocean model NEMO 3.6, *Geosci. Model Dev.*, 10, 2849–2874, 2017.
- Medley, B. and Thomas, E.: Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise, *Nature Climate Change*, 9, 34–39, 2019.
- Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J.-F., Stouffer, R. J., Taylor, K. E., and Schlund, M.: Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models, *Science Advances*, 6, eaba1981, 2020.
- 700 Meier, W. N., Stewart, J. S., Windnagel, A., and Fetterer, F. M.: Comparison of hemispheric and regional sea ice extent and area trends from NOAA and NASA passive microwave-derived climate records, *Remote Sensing*, 14, 619, 2022.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, 2020.
- 705 Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., Boberg, F., van de Berg, W. J., Fettweis, X., Gossart, A., et al.: What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates, *The Cryosphere*, 15, 3751–3784, 2021.
- Mouginot, J., Rignot, E., and Scheuchl, B.: Continent-wide, interferometric SAR phase, mapping of Antarctic ice velocity, *Geophysical Research Letters*, 46, 9710–9718, 2019.
- 710 Müller, W. A., Jungclaus, J. H., Mauritzen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., et al.: A higher-resolution version of the max planck institute earth system model (MPI-ESM1. 2-HR), *J. Adv. Model. Ea. Sys.*, 10, 1383–1413, 2018.
- Naughten, K. A., Holland, P. R., and De Rydt, J.: Unavoidable future increase in West Antarctic ice-shelf melting over the twenty-first century, *Nature Climate Change*, pp. 1–7, 2023.
- 715 Nowicki, S., Goelzer, H., Seroussi, H., Payne, A. J., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis, X. S., Barthel, A., et al.: Experimental protocol for sea level projections from ISMIP6 stand-alone ice sheet models, *The Cryosphere*, 14, 2331–2368, 2020.
- PAGES2k: Consistent multidecadal variability in global temperature reconstructions and simulations over the Common Era, *Nature geoscience*, 12, 643–649, 2019.
- 720 Parsons, L. A., Brennan, M. K., Wills, R. C., and Proistosescu, C.: Magnitudes and spatial patterns of interdecadal temperature variability in CMIP6, *Geophysical Research Letters*, 47, e2019GL086 588, 2020.
- Payne, A. J., Nowicki, S., Abe-Ouchi, A., Agosta, C., Alexander, P., Albrecht, T., Asay-Davis, X., Aschwanden, A., Barthel, A., Bracegirdle, T. J., et al.: Future sea level change under coupled model intercomparison project phase 5 and phase 6 scenarios from the Greenland and Antarctic ice sheets, *Geophysical Research Letters*, 48, e2020GL091 741, 2021.
- 725 Penduff, T., Sérazin, G., Leroux, S., Close, S., Molines, J.-M., Barnier, B., Bessières, L., Terray, L., and Maze, G.: Chaotic variability of ocean heat content: climate-relevant features and observational implications, *Oceanography*, 31, 63–71, 2018.
- Previdi, M. and Polvani, L. M.: Anthropogenic impact on Antarctic surface mass balance, currently masked by natural variability, to emerge by mid-century, *Environmental Research Letters*, 11, 094 001, 2016.

- 730 Purich, A. and England, M. H.: Historical and future projected warming of Antarctic Shelf Bottom Water in CMIP6 models, *Geophysical Research Letters*, 48, e2021GL092752, 2021.
- Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., and Winkelmann, R.: Antarctic sub-shelf melt rates via PICO, *The Cryosphere*, 12, 1969–1985, 2018.
- Reese, R., Garbe, J., Hill, E. A., Urruty, B., Naughten, K. A., Gagliardini, O., Durand, G., Gillet-Chaulet, F., Gudmundsson, G. H., Chandler,
735 D., et al.: The stability of present-day Antarctic grounding lines–Part 2: Onset of irreversible retreat of Amundsen Sea glaciers under current climate on centennial timescales cannot be excluded, *The Cryosphere*, 17, 3761–3783, 2023.
- Riahi, K., Van Vuuren, D. P., Kriegler, E., Edmonds, J., O’neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., et al.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global environmental change*, 42, 153–168, 2017.
- 740 Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, 266–270, 2013.
- Rignot, E., Mouginot, J., Scheuchl, B., Van Den Broeke, M., Van Wessel, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, 2019.
- Robel, A., Verjans, V., and Ambelorun, A.: Biases in ice sheet models from missing noise-induced drift, *EGUosphere*, 2023, 1–14, 2023.
- Schmidtko, S., Heywood, K. J., Thompson, A. F., and Aoki, S.: Multidecadal warming of Antarctic waters, *Science*, 346, 1227–1231, 2014.
- 745 Scott, R. C., Nicolas, J. P., Bromwich, D. H., Norris, J. R., and Lubin, D.: Meteorological drivers and large-scale climate forcing of West Antarctic surface melt, *Journal of Climate*, 32, 665–684, 2019.
- Sellar, A. A., Walton, J., Jones, C. G., Wood, R., Abraham, N. L., Andrejczuk, M., Andrews, M. B., Andrews, T., Archibald, A. T., de Mora, L., et al.: Implementation of UK Earth system models for CMIP6, *J. Adv. Model. Ea. Sys.*, 12, e2019MS001946, 2020.
- Sérazin, G., Jaymond, A., Leroux, S., Penduff, T., Bessières, L., Llovel, W., Barnier, B., Molines, J.-M., and Terray, L.: A global probabilistic
750 study of the ocean heat content low-frequency variability: Atmospheric forcing versus oceanic chaos, *Geophysical Research Letters*, 44, 5580–5589, 2017.
- Seroussi, H., Morlighem, M., Larour, E., Rignot, E., and Khazendar, A.: Hydrostatic grounding line parameterization in ice sheet models, *The Cryosphere*, 8, 2075–2087, 2014.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel,
755 A., et al.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere*, 14, 3033–3070, 2020.
- Seroussi, H., Verjans, V., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., et al.: Insights on the vulnerability of Antarctic glaciers from the ISMIP6 ice sheet model ensemble and associated uncertainty, *The Cryosphere Discussions*, 2023, 1–28, 2023.
- 760 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet elevation and mass, *Geophysical Research Letters*, 46, 8174–8183, 2019.
- Siahaan, A., Smith, R. S., Holland, P. R., Jenkins, A., Gregory, J. M., Lee, V., Mathiot, P., Payne, A. J., Ridley, J. K., and Jones, C. G.: The Antarctic contribution to 21st-century sea-level rise predicted by the UK Earth System Model with an interactive ice sheet, *The Cryosphere*, 16, 4053–4086, 2022.
- 765 Silvano, A., Holland, P. R., Naughten, K. A., Dragomir, O., Dutrieux, P., Jenkins, A., Si, Y., Stewart, A. L., Peña Molino, B., Janzing, G. W., et al.: Baroclinic Ocean Response to Climate Forcing Regulates Decadal Variability of Ice-Shelf Melting in the Amundsen Sea, *Geophysical Research Letters*, 49, e2022GL100646, 2022.

- Smith, J. A., Andersen, T. J., Shortt, M., Gaffney, A. M., Truffer, M., Stanton, T. P., Bindschadler, R., Dutrieux, P., Jenkins, A., Hillenbrand, C. D., Ehrmann, W., Corr, H. F., Farley, N., Crowhurst, S., and Vaughan, D. G.: Sub-ice-shelf sediments record history of twentieth-century retreat of Pine Island Glacier, *Nature*, 541, <https://doi.org/10.1038/nature20136>, 2017.
- 770
- Smith, R. S., Mathiot, P., Siahhaan, A., Lee, V., Cornford, S. L., Gregory, J. M., Payne, A. J., Jenkins, A., Holland, P. R., Ridley, J. K., et al.: Coupling the UK Earth System Model to dynamic models of the Greenland and Antarctic ice sheets, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002520, 2021.
- Steig, E. J., Ding, Q., Battisti, D., and Jenkins, A.: Tropical forcing of Circumpolar Deep Water inflow and outlet glacier thinning in the Amundsen Sea Embayment, West Antarctica, *Annals of Glaciology*, 53, 19–28, 2012.
- 775
- The IMBIE Team: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- Tsai, C.-Y., Forest, C. E., and Pollard, D.: The role of internal climate variability in projecting Antarctica’s contribution to future sea-level rise, *Climate dynamics*, 55, 1875–1892, 2020.
- 780
- Van Wessem, J. M., Van De Berg, W. J., Noël, B. P., Van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T., Lhermitte, S., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2–Part 2: Antarctica (1979–2016), *The Cryosphere*, 12, 1479–1498, 2018.
- Verfaillie, D., Pelletier, C., Goosse, H., Jourdain, N. C., Bull, C. Y., Dalaiden, Q., Favier, V., Fichefet, T., and Wille, J. D.: The circum-Antarctic ice-shelves respond to a more positive Southern Annular Mode with regionally varied melting, *Communications Earth & Environment*, 3, 139, 2022.
- 785
- Wang, Y., Yuan, X., and Cane, M. A.: Coupled mode of cloud, atmospheric circulation, and sea ice controlled by wave-3 pattern in Antarctic winter, *Environmental Research Letters*, 17, 044053, 2022.
- Williams, G., Herraiz-Borreguero, L., Roquet, F., Tamura, T., Ohshima, K., Fukamachi, Y., Fraser, A., Gao, L., Chen, H., McMahon, C., et al.: The suppression of Antarctic bottom water formation by melting ice shelves in Prydz Bay, *Nature communications*, 7, 12577, 2016.
- 790
- Zhang, L., Delworth, T. L., Cooke, W., and Yang, X.: Natural variability of Southern Ocean convection as a driver of observed climate trends, *Nature Climate Change*, 9, 59–65, 2019.