

1 ~~Integrating palaeoecology and dendrochronology to explore the~~
2 ~~impact of climate and forest management on a peatland in Scots pine~~
3 ~~monoculture~~

4
5 **Assessing the impact of forest management and climate on a peatland**
6 **under Scots pine monoculture using a multidisciplinary approach**

7
8 Mariusz Bąk¹, Mariusz Lamentowicz¹, Piotr Kołaczek¹, Daria Wochal¹, Paweł Matulewski², Dominik
9 Kopec^{3,4}, Martyna Wietecha^{3,4}, Dominika Jaster², Katarzyna Marcisz¹

10
11 ¹Climate Change Ecology Research Unit, Faculty of Geographical and Geological Sciences, Adam
12 Mickiewicz University, Poznań, Poland

13 ²Anthropocene Research Unit, Faculty of Geographical and Geological Sciences, Adam Mickiewicz
14 University, Poznań, Poland

15 ³Department of Biogeography, Paleoecology and Nature Conservation, University of Lodz, Łódź, Poland

16 ⁴MGGP Aero Sp. z o. o., Tarnów, Poland

17
18 *Correspondence to:* Mariusz Bąk, mariusz.bak@amu.edu.pl

19
20 **Abstract:** Assessing the scale, rate, and consequences of climate change, manifested primarily by rising
21 average air temperatures and altered precipitation regimes, is a critical challenge in contemporary scientific
22 research. These changes are accompanied by various anomalies and extreme events that negatively impact
23 ecosystems worldwide. Monoculture forests, including Scots pine (*Pinus sylvestris* L.) monocultures, are
24 particularly vulnerable to these changes due to their homogeneous structure and simplified ecosystem
25 linkages compared to mixed forests, making them more sensitive to extreme events such as insect outbreaks,
26 droughts, fires, and strong winds. In the context of global warming, forest fires are becoming extremely
27 dangerous, and the risk of their occurrence increases as average temperatures rise. The situation becomes
28 even more dramatic when fire enters areas of peatlands, as these ecosystems effectively withdraw carbon
29 from the rapid carbon cycle and store it for up to thousands of years. Consequently, peatlands become
30 emitters of carbon dioxide into the atmosphere.

31 In this study, we aim to trace the last 300 years of historical development of a peatland situated in a Scots
32 pine monoculture. Our focus is on the Okoniny peatland located within the Tuchola Pinewoods in northern
33 Poland, one of the country's largest forest complexes. We delved into the phase when the peatland's
34 surroundings transitioned from a mixed forest to a pine monoculture and investigated the impact of changes
35 in forest management on the peatland vegetation and hydrology. Our reconstructions are based on a multi-
36 proxy approach using: pollen, plant macrofossils, micro- and macrocharcoal, and testate amoebae. We

Z komentarzem [MB1]: Based on the reviewers' comments, we proposed a new title that better represents the content of the article.

37 combine the peatland palaeoecological record with the dendrochronology of *Pinus sylvestris* to compare the
38 response of these two archives. Our results show that a change in forest management and progressive climate
39 warming affected the development of the peatland. We note an increase in acidity over the analyzed period
40 and a decrease in the water table over the last few decades that led to the lake-peatland transition. These
41 changes progressed with the strongest agricultural activity in the area in the 19th century. However, the 20th
42 century was a period of continuous decline in agriculture and an increase in the dominance of Scots pine in
43 the landscape as the effect of afforestation. Dendroclimatic data indicate a negative effect of temperature on
44 Scots pine and pressure from summer rainfall deficiency. Additional remote sensing analysis, using
45 hyperspectral, LiDAR, and thermal airborne data, provided information about the current condition of the
46 peatland vegetation. With the application of spectral indices and the analysis of land surface temperature,
47 spatial variations in peatland drying have been identified. Considering the context of forest management
48 and the protection of valuable ecosystems in monocultural forests, the conclusions are relevant for peatland
49 and forest ecology, palaeoecology, and forestry.

50
51 **Keywords:** palaeoecological data, palaeoecology, dendrochronology, dendroclimatic data, climate change,
52 monoculture forests, plantation, remote sensing, historical data, historical maps, multi-proxy, high-
53 resolution, airborne data, thermal data, vegetation indices, remote sensing

54 1. Introduction

55 ~~Recognizing how different ecosystems function under a changing climate and increasing human impact is~~
56 ~~crucial for their conservation and management.~~ Peatlands are vulnerable to various types of change, which
57 play an important role in the global carbon cycle and whose destabilization can create catastrophic positive
58 feedback for climate warming (Gallego-Sala et al., 2018; Wilson et al., 2016). Peatlands, although they only
59 cover about 3% of the Earth's total land area (Parish et al., 2008; Rydin and Jeglum, 2013), store more than
60 30% of the organic carbon (C) (Freeman et al., 2004; Gorham, 1991; Harenda et al., 2018), which is far
61 more carbon than the entire biomass of the world's forests (Beaulne et al., 2021b). Their advantage over
62 forests is not only due to their ability to accumulate C but also to the fact that they do not emit decomposed
63 carbon from the so-called rapid C cycle for up to thousands of years (Blodau, 2002; Gorham, 1991). The
64 estimation of C content accumulated in peatlands is challenging (Sanderson et al., 2023), although some
65 studies indicate ca. 600 Gt of C in the Northern Hemisphere alone (Yu et al., 2010). It has recently been
66 shown that even the smallest kettle-hole peatlands effectively accumulate ~~of~~C and serve as important C hot
67 spots (Karpińska-Kołaczek et al., 2024).

68
69 Insufficient awareness of the ecological importance of peatlands has led to them being treated as
70 wastelands and drained for hundreds of years to obtain land for agriculture, and forestry or exploited

Z komentarzem [MB2]: Changes in keywords

Z komentarzem [MB3]: We have removed as suggested by the reviewer.

Z komentarzem [MB4]: Editorial and linguistic corrections

Z komentarzem [MB5]: Editorial and linguistic corrections

71 commercially as an energy resource (Joosten et al., 2012; Łuców et al., 2022; Paavilainen and Päivänen,
72 1995). Many of these areas have also had to adapt to a changing environment resulting from the use of
73 various forest management techniques, e.g., the replacement of mixed forests with more easily managed
74 monoculture forests (plantations) (Lee et al., 2023; Łuców et al., 2021; Słowiński et al., 2019). Mixed
75 forests, through greater biodiversity, are more resilient and better able to adapt to environmental change
76 (Bauhus et al., 2017; Messier et al., 2022), providing a more comprehensive range of ecosystem services
77 (Felton et al., 2016; Huuskonen et al., 2021).

78 Despite being more straightforward to manage, forest monocultures are characterized by simplified
79 ecosystem linkages (Chapin et al., 2012). As a result, they are more susceptible to various extreme events
80 and disturbances, both natural and anthropogenic, including droughts, fires, strong winds, and pest
81 ~~gradations infestations~~ (Grondin et al., 2014). This is particularly important as disturbances of these types
82 of forests are becoming more common (Seidl et al., 2014; Westerling, 2016). Natural disturbance regimes
83 in forests are mainly a response to climate change (Hanson and Weltzin, 2000; Pureswaran et al., 2015;
84 Seidl et al., 2017; Trumbore et al., 2015), therefore they are expected to increase in frequency and severity
85 in the coming years (Gregow et al., 2017; Moritz et al., 2012; Wotton et al., 2010). Moreover, the problem
86 applies to all kinds of monoculture forests regardless of the dominant species and climate zones (Booth,
87 2013; Guariguata et al., 2008; McNulty et al., 2013; Spiecker, 2000), including pine plantations in the
88 temperate climate zone of Central and Eastern Europe (Łuców et al., 2021; Schüle et al., 2023). Thus,
89 peatlands, which are so crucial in terms of their impact on global climate change, located in the area of forest
90 monocultures are even more vulnerable to extreme phenomena and disturbance, despite the already high
91 climatic and anthropogenic pressure.

92 The history of peatlands' development can be traced using palaeoecological analyses, which allow
93 numerous reconstructions of past environmental conditions, including climate change (Lamentowicz et al.,
94 2015; Mauquoy and Yeloff, 2008). These include reconstructions of vegetation changes in the peatland and
95 its surroundings, changes in the water table, and reconstructions of past fire activity (Galka et al., 2022;
96 Kołaczek et al., 2018; Marcisz et al., 2020b, 2017; Mroczkowska et al., 2021). ~~This is because peat~~
97 ~~preserves, plant remains, Peat archive records contain a wide range of preserved micro- and macrofossils~~
98 ~~for example, pollen, spores, microbial remains, and charcoal are deposited in situ and brought in by wind or~~
99 ~~water, collectively called peat archives~~ (Godwin, 1981). While paleoenvironmental reconstructions based
100 on peat records have become common, few studies still integrate palaeoecological data with other methods.
101 For example, studies that combine palaeoecological and dendrochronological records, including
102 dendroclimatic reconstructions based on analysis of the annual growth of tree rings, are still relatively rare
103 (Ballesteros-Cánovas et al., 2022; Beaulne et al., 2021a; Dinella et al., 2021; Edvardsson et al., 2022, 2019,
104 2016; González de Andrés et al., 2022; Kuosmanen et al., 2020; Lamentowicz et al., 2009b). Yet, combining

Z komentarzem [MB6]: Editorial and linguistic corrections

Z komentarzem [MB7]: Editorial and linguistic corrections

105 peat records with dendrochronological data can benefit interpretations of trees and forest resilience and
106 resistance to disturbances compared to local environmental changes recorded in peat. Such a view of past
107 environmental changes through several proxies and other archive types is fundamental and will be helpful
108 for forest management and nature conservation in the future. To assess the current state of the peatland, we
109 also included remote sensing data in the analysis. Remote sensing methods have been applied to study
110 wetland conditions for over 50 years and are currently regarded as one of the most useful methods in this
111 research area (FAO, 2020; Guo et al., 2017). Remote sensing technologies enable the remote and non-
112 invasive acquisition of information about the research object using specialized sensors, typically mounted
113 aboard satellites or aircraft. In this study, data obtained from a multisensor aerial platform were used to
114 assess the extent of peatland, the identification of drainage ditches and the current vegetation condition.

115 ~~Our study aims to assess the impact of the introduction of pine monoculture on the development of~~
116 ~~*Sphagnum* peatlands in central and eastern Europe. We reconstructed hydrological conditions caused by~~
117 ~~changing climate and forest management, identified peat layers corresponding to the occurrence of extreme~~
118 ~~phenomena known from historical sources, and integrated palaeoecological and dendrochronological~~
119 ~~(dendroclimatic) data developed from annual growths of Scots pine (*Pinus sylvestris*). We also explored~~
120 ~~how peatland responded to extreme phenomena, such as outbreaks or fires, in situ and in the immediate~~
121 ~~environment.~~

122 ~~We have assumed that the introduction of pine monoculture has led to significant changes in peatland~~
123 ~~species composition in favour of peat mosses and a stabilization of the groundwater table. We also undertook~~
124 ~~to confirm that peatlands record and respond to extreme phenomena, both occurring in situ and in the~~
125 ~~immediate environment. We assumed that disturbance events that happened in the monoculture forest~~
126 ~~throughout the years would be recorded in the pine tree ring record and would validate and complete~~
127 ~~peatland reconstruction.~~


128 Our study aims to assess the impact of forest management (introduction of pine monoculture) and
129 changing climate on the vegetation, as well as hydrological, and trophic conditions of a peatland in CE
130 Europe by integrating various data sources - palaeoecology, dendrochronology, remote sensing, and
131 historical information. We assumed that the introduction of pine monoculture led to changes in the species
132 composition of peatlands in favor of *Sphagnum* mosses, as well as to the stabilization of the water table. We
133 also undertook to confirm whether peatlands register and respond to extreme events, both *in situ* and in the
134 immediate environment. We assumed that the disturbances that occurred in the monoculture forest would
135 be recorded in the tree rings (annual growths) record of Scots pine (*Pinus sylvestris* L.) and would confirm
136 and complement the palaeoecological reconstruction of the peatland. Thus, we have identified peat layers
137 corresponding to the occurrence of extremes known from historical sources and compared
138 dendrochronological (dendroclimatic) data with them.

Z komentarzem [MB8]: We have merged these paragraphs and rewritten the text to make the goals of our study clearer.

139 **2. Materials and methods**

140 **2.1. Study site**

141 The Okoniny peatland (53°40'52"–53°41'21"N 18°03'09"–18°03'40"E according to standard WGS
142 84) is located in northern Poland, about 60 km north of Bydgoszcz and about 20 km northeast of Tuchola
143 (Fig. 1). The study area is located within the Tuchola Pinewoods mesoregion (Kondracki, 2001), close to
144 the Pomeranian ice margin of the Vistulian Glaciation dated to ca. 17,000–16,000 cal. BP (Marks, 2012).
145 The entire area of the Tuchola Pinewoods is a young glacial landscape covered by glacial till, sand, and
146 numerous depressions and other forms originating from melting dead ice (Błaszkiwicz et al., 2015). Based
147 on the analysis of remote sensing data, it was determined that the surface area of the peatland is 27.08
148 hectares, with approximately 7.00 hectares designated as non-forested area. The direct catchment area of
149 the peatland covers a surface of 33.23 hectares. The current elevation of the peatland is around 119 m asl,
150 with the highest elevated area within the direct catchment reaching around 128 m asl. It is part of a protected
151 area (Regulation No. 64/97, 1997), included within the boundaries of the Tuchola Landscape Park (created
152 in 1985). Moreover, since 2008 the entire complex of the Tuchola Pinewoods has been included on the
153 Natura 2000 list as a Special Protection Area. Since 2010, it has been listed as a UNESCO Biosphere
154 Reserve (UNESCO, 2024).



155
 156 Figure 1. Location of the study area. (A) Location on a map of north-western Poland. (B) Location of the
 157 two study sites – dendrochronological sampling site and peat coring site. (C) Okoniny peatland sampling
 158 site with current peatland boundaries.

159
 160 The Okoniny peatland is located in a temperate latitude zone, with a transitional climate influenced
 161 by continental air masses from eastern Europe and oceanic air masses from the Atlantic Ocean (Beck et al.,
 162 2018). According to climate data obtained from the Institute of Meteorology and Water Management for the
 163 meteorological station in Chojnice (35 km west of the study area) for the thirty years period between 1991-
 164 2020, the coldest month is January with an average temperature of -1.5 °C, the warmest month is July with
 165 an average temperature of 18.0 °C. In the thirty years Between 1961-1990, both January and July were
 166 cooler by 1.6 °C compared to 1991-2020. The average annual temperature increased from 6.9 °C in 1951-

Z komentarzem [MB9]: Editorial and linguistic corrections

Z komentarzem [MB10]: Editorial and linguistic corrections

167 1990 to 8.1 °C in 1991-2020. In terms of precipitation, February has the least amount with an average of
168 31.1 mm for the period 1991-2020, and July has the most with an average of 80.7 mm for the period 1991-
169 2020. Compared to 1951-1990, the average precipitation for February increased by 7.7 mm, and for July
170 decreased by 4.1 mm. Mean annual rainfall increased from 558.1 mm for 1951-1990 to 612.4 mm for 1991-
171 2020.

172 Samples for dendroclimatic analysis were taken from forest division no. 91 in the Woziwoda
173 Forestry, Woziwoda Forest District, about 9.5 km west of the study site (Fig. 1). The oldest pine trees in the
174 forest district were selected for the study according to the indications of the forest survey and taxonomic
175 descriptions.

176

177 2.2. Peat and tree core sampling

178 A peat core was taken from the north-western part of the peatland in February 2022 using a Wardenaar
179 corer (chamber dimension: 10 cm × 10 cm × 100 cm) (Wardenaar, 1987). The entire length of the sampled
180 peat core – 96 cm-long monolith – was analyzed. The core was sampled continuously every 1 cm, except
181 for the top 10 cm, which contained a living *Sphagnum* layer. The first sample covered 4 cm of the surface
182 layer (0-4 cm), and the following three samples were taken every 2 cm (4-6, 6-8 and 8-10 cm). 90 samples
183 were obtained and analyzed for bulk density, ash content, peat and carbon accumulation rates, plant
184 macrofossils, testate amoebae, macroscopic and microscopic charcoal, and pollen.

185 The research tree stem material was taken in April 2023 from 23 living and healthy trees at the
186 Woziwoda site, ca. 9.5 km west of the Okoniny peatland. From each tree, a minimum of two cores were
187 taken (from the east and west sides) at a breast height (1.3 m) with a Pressler increment corer. In total, 50
188 cores were acquired from the Scots pine tree stems.

189

190 2.3. Radiocarbon dating and chronology

191 Ten samples containing *Sphagnum* stems and leaves were used for accelerator mass spectroscopy
192 (AMS) ¹⁴C dating of the entire length of the profile. The survey was conducted at the Poznan Radiocarbon
193 Laboratory in Poland (laboratory code marked Poz; Tab. 1). The IntCal20 (Reimer et al., 2020) and
194 Bomb21NH1 (Hua et al., 2021) atmospheric curves were used to calibrate the dates.

195

196 Table 1. The list of radiocarbon dates from Okoniny peatland with calibration in the OxCal v4.4.4 software
197 using [the](#) IntCal20 calibration curve for the atmospheric data and Bomb21NH1 curve for bomb series.

No	Laboratory code – number sample	Depth (cm)	¹⁴ C date (¹⁴ C BP)	Calibrated dates [cal. CE (2σ – 95.4%)	Dated material
----	------------------------------------	---------------	---	---	----------------

Z komentarzem [MB11]: Editorial and linguistic corrections

1	Poz-150386	10.5	100.86 ± 0.33 pMC	1952-1958 (33.9%) 2013-... (61.5%)	<i>Sphagnum</i> stems
2	Poz-150387	20.5	107.92 ± 0.34 pMC	1952-1958 (11.1%) 1996-2009 (84.4%)	<i>Sphagnum</i> stems
3	Poz-150388	30.5	132.8 ± 0.36 pMC	1958-1962 (20.8%) 1972-1984 (74.6%)	<i>Sphagnum</i> stems
4	Poz-150445	40.5	165 ± 30	1661-1706 (17.2%) 1720-1818 (44.0%) 1832-1892 (14.9%) 1906-... (19.5%)	<i>Sphagnum</i> stems
5	Poz-150446	50.5	85 ± 30	1688-1730 (26.1%) 1806-1924 (69.3%)	<i>Sphagnum</i> stems
6	Poz-150447	60.5	105 ± 30	1682-1736 (25.9%) 1802-1936 (69.5%)	<i>Sphagnum</i> stems
7	Poz-150449	70.5	135 ± 30	1674-1766 (32.8%) 1774-1776 (0.6%) 1798-1942 (62.0%)	<i>Sphagnum</i> stems
8	Poz-150450	80.5	165 ± 30	1661-1706 (17.2%) 1720-1818 (44.0%) 1832-1892 (14.9%) 1906-... (19.5%)	<i>Sphagnum</i> stems
9	Poz-150631	90.5	280 ± 30	1505-1596 (55.0%) 1616-1665 (37.8%) 1784-1794 (2.6%)	<i>Sphagnum</i> stems
10	Poz-150633	95.5	100 ± 30	1683-1735 (26.1%) 1802-1930 (69.3%)	<i>Sphagnum</i> stems

198
199 The absolute chronology of the entire core was based on a Bayesian age-depth model using OxCal
200 v4.4.4 (Bronk Ramsey, 2021). The *P_Sequence* command with a parameter *k* of 0.1 cm⁻¹ was used to
201 calculate the model, assuming $\log_{10}(k/k_0) = 2$, and interpolation = 1 cm. The most pronounced change in
202 peat composition, as manifested by changes in pollen concentration, testate amoeba species composition
203 and species composition of plant macrofossils, which may signal changes in peat accumulation rates, was
204 input using the *Boundary* command at a depth of 66 cm. For better readability of the age-depth model, mean
205 values (μ) were introduced and used to illustrate the modeled age.

206

207 2.4. Peat properties and carbon accumulation rate

208 Analyses of bulk density, loss on ignition, and peat carbon accumulation rate (PCAR) were carried out
209 for each of the ninety samples. Each sample's volume [cm³] was carefully ~~and cautiously~~ measured
210 beforehand using calipers to avoid compressing the material. Each sample was then placed in a separate
211 crucible and dried to determine the percentage of water content. The weighed and dried samples were
212 incinerated at 550 °C for 12 hours and reweighed according to the protocol of Heiri et al. (2001) to determine
213 the ash mass [g]. Bulk density [g/cm³] was obtained by dividing the dry sample mass by the volume of the
214 fresh sample according to Chambers et al. (2010). Loss on ignition [g] was obtained by subtracting the ash
215 mass from the dry sample mass. Accumulation rates obtained from the peat core chronologies were
216 multiplied by measuring the bulk density without ash and by 50% to obtain the PCAR, following the
217 protocol of Loisel et al. (2014). The top eleven centimeters of the core (0-11 cm) were discarded for PCAR
218 assessment due to the unrepresentative nature of the results obtained, as increased values of carbon
219 accumulation in near-surface peat cannot be used for inference (Young et al., 2019).

220

221 2.5. Plant macrofossil analysis

222 The analysis of plant macrofossils was carried out using the modified protocol of Mauquoy et al., 2010.
223 Each sample of approximately 5cm³ was wet sieved (mesh diameter: 200 µm). The generalized content of
224 the sample was estimated in percentage using a binocular microscope. Fruits, seeds, caryopses, achenes,
225 perigynia, bud scales, catkin scales, whole preserved leaves, whole preserved needles, cones, anthers,
226 sporangia, opercula, fungi sclerotia, and wood pieces were counted as total numbers in each sample. The
227 tissues of monocotyledon species and moss leaves (brown and *Sphagnum* mosses) were identified on slides
228 using a magnification of ×200 and ×400. The material was compared with the guides (Anderberg, 1994;
229 Berggren, 1969; Bojňanský and Fargašová, 2007; Mauquoy and van Geel, 2007). ~~The diagram for the~~
230 ~~analyzed proxy was plotted using the riojaPlot package for R (plant macrofossils) (Juggins, 2023).~~

231

232 2.6. Testate amoebae analysis

233 Samples for testate amoeba analysis (volume: ca. 5cm³) were washed under 300 µm sieves following
234 the method described by Booth et al. (2010). Testate amoebae were analyzed under a light microscope with
235 ×200 and ×400 magnifications until the sum of 100 tests per sample was reached (Payne and Mitchell,
236 2009). Several keys and taxonomic monographs (Clarke, 2003; Mazei and Tsyganov, 2006; Meisterfeld,
237 2001; Ogden and Hedley, 1980) as well as online resources (Siemensma, 2023) were used to achieve the
238 highest possible taxonomic resolution. The results of a testate amoebae analysis were used for the
239 quantitative depth-to-water table (DWT) and pH reconstructions. Both ~~the full diagram and the~~

Z komentarzem [MB12]: Editorial and linguistic corrections

Z komentarzem [MB13]: Editorial and linguistic corrections

Z komentarzem [MB14]: The section "Visualization of the palaeoecological results" has been removed, and information on software used for visualization has been placed next to the description of specific methods.

Z komentarzem [MB15]: The section "Visualization of the palaeoecological results" has been removed, and information on software used for visualization has been placed next to the description of specific methods.

240 reconstructions were performed in C2 software (Juggins, 2007) using the European training set (Amesbury
241 et al., 2016).

242

243 2.7. Pollen and non-pollen palynomorphs (NPPs)

244 Samples for palynological analysis (volume: 2 cm³) were prepared using standard laboratory
245 procedures (Berglund and Ralska-Jasiewiczowa, 1986). To remove the carbonates, samples were treated
246 with 10% hydrochloric acid. This step was followed by digestion in hot 10% potassium hydroxide (to
247 remove humic compounds) and soaking in 40% hydrofluoric acid for 24 h (to remove the mineral fraction).
248 Next, acetolysis was carried out. Three *Lycopodium* tablets (Batch 280521291, containing 18407 spores per
249 tablet; produced by Lund University) were added to each sample during the laboratory procedures for the
250 calculation of microfossil concentration (Stockmarr, 1971). Pollen, spores, and selected non-pollen
251 palynomorphs (NPPs) were counted under an upright microscope (Zeiss Axio SCOPE A1) until the number
252 of total pollen sum (TPS) grains in each sample reached at least 500, apart from 23 samples in which pollen
253 concentrations were very low. Sporomorphs were identified with the assistance of atlases, keys (Beug, 2004;
254 Moore et al., 1991), various publications, and the image database in the case of NPPs, for which there are
255 no atlases (Miola, 2012; Shumilovskikh et al., 2022; Shumilovskikh and van Geel, 2020). The results of the
256 palynological analysis were expressed as percentages, calculations are based on the ratio of an individual
257 taxon to the TPS, i.e., the sum of AP (arboreal pollen) and NAP (non-arboreal pollen), excluding aquatic
258 and wetland plants (together with Cyperaceae and Ericaceae), cryptogams, and fungi. [The diagram for the
259 analyzed proxy was plotted using Tilia/Tilia graph software \(pollen\) \(Grimm, 1992, 1991\).](#)

260

261 2.8. Macro- and microcharcoal analysis

262 Microscopic charcoal particles (size: > 10 µm) were counted from the same slides as pollen until the
263 number of charcoal particles and *Lycopodium* spores counted together, exceeded 200 (Finsinger and Tinner,
264 2005; Tinner and Hu, 2003). Microscopic charcoal influx or accumulation rates (MIC) were calculated by
265 multiplying charcoal concentrations by peat accumulation rates (PAR) (Davis and Deevey, 1964; Tinner and
266 Hu, 2003).

267 For macroscopic charcoal analysis, samples (volume: 2 cm³) were prepared by bleaching to create a
268 more visible contrast between the charcoal and the remaining organic matter following the method described
269 by Whitlock and Larsen (2001). Samples were sieved through a 500-µm mesh and only large charcoal
270 fragments > 600 µm were analyzed to obtain a local fire signal (Adolf et al., 2018). Samples were analyzed
271 with a binocular under 60× magnification. Macroscopic charcoal influx or accumulation rates (MAC,
272 particles/cm²/year) were calculated using the charcoal concentrations and PAR.

273

Z komentarzem [MB16]: Editorial and linguistic corrections

Z komentarzem [MB17]: The section “Visualization of the palaeoecological results” has been removed, and information on software used for visualization has been placed next to the description of specific methods.

274 **Visualization of the palaeoecological results**

275 Palaeoecological diagrams for the analyzed proxies were plotted using Tilia/Tilia graph software (pollen)
276 (Grimm, 1992, 1991), C2 software (testate amoebae) (Juggins, 2007) and riojaPlot package for R (plant
277 macrofossils) (Juggins, 2023). Quantitative reconstructions of testate amoebae-based depth to water table
278 (DWT) and pH changes were done in C2 software (Juggins, 2007), using the European training set
279 (Amesbury et al., 2016).

280

281 **2.9. Tree core chronology construction**

282 Tree cores underwent a standardized dendrochronological procedure (Zielski and Krąpiec, 2004).
283 Polished cores were scanned between 1200 - 2400 DPI using an Epson Perfection V700 Photo scanner.
284 Annual growth rings were measured on digital images with an accuracy of 0.01mm using Coorecorder.
285 This facilitated the selection of individual growth sequences, which were utilized to form a chronology for
286 each plot. Visual comparisons were made between individual sequences, and the significance of correlations
287 was assessed using Student's t-test (Baillie and Pilcher, 1973). Subsequently, cross-dating was conducted
288 using COFECHA software (Grissino-Mayer, 2001), which evaluates each data series concerning the
289 reference chronology created and compares the correlation coefficients obtained. Raw chronologies were
290 derived by employing an arithmetic mean. For climate-growth analysis standardized chronologies were
291 used, obtained by fitting a spline function (i.e., the "n-year spline" was set at 2/3 of the wavelength of n
292 years of single growth series) using the 'dplR' package (Bunn, 2008) package version 1.7.6 (2023) in the
293 software R version 4.3.0 (R Core Team, 2022). By using this standardization method, random variation in
294 the radial growth was removed (Cook et al., 1990). For the obtained chronologies i.e., raw (TRW) and
295 standardized (RWI), values for the following descriptive statistics were computed: the mean correlation
296 between series (inter-series correlation or Rbar), the GLK index (Gleichläufigkeit; Eckstein and Bauch,
297 1969), and EPS (express population signal) (McCarroll and Loader, 2004).

298

299 **2.10. Dendroclimatological and pointer years analysis**

300 The 'chron' function from 'dplr' package allowed for the making of a residual chronology, which was
301 used for climate-growth analysis. The 'dcc' function and its moving response (25-yrs window) function
302 method were used to determine the effects of climate conditions on the growth of Scots pine using the
303 'treeclim' package (Zang and Biondi, 2015) version 2.0.6.0 in R (R Core Team, 2022). This package allows
304 the use of the bootstrap procedure to test the significance and stability of the coefficients of determination
305 (r^2) over a set period (Guiot, 1991). Monthly mean air temperature (TEMP) and total monthly precipitation
306 (PREC) were used to analyze climate-growth for the period 1920-2022 (Klein Tank et al., 2002). Climate
307 data were acquired via Climate Explorer (Trouet and van Oldenborgh, 2013) and calculated from the

Z komentarzem [MB18]: The section "Visualization of the palaeoecological results" has been removed, and information on software used for visualization has been placed next to the description of specific methods.

Z komentarzem [MB19]: Renumbering

Z komentarzem [MB20]: Renumbering

308 monthly gridded observational dataset E OBS v. 25.0e (Haylock et al., 2008) obtained for the 17.75-18.00°E
309 and 53.50-53.75°N grid.

310 The Becker algorithm (Becker et al., 1994) was used to determine the pointer years in the Woziwoda
311 chronology. Calculations were made using the 'dplr' package in R and the 'pointer' function (Bunn, 2008).
312 Pointer years were calculated using adjustable thresholds of relative variation in radial growth set to a 10-
313 year time window and the number of series exhibiting a similar incremental growth pattern. The main
314 criterion for determining pointer years was the occurrence of unidirectional changes (i.e., a decrease or
315 increase in the number of annual rings) in a minimum of 85% of the tested sequences of annual increments
316 observed in a group of trees at the Woziwoda site.

317

318 2.11. Acquisition and post-processing of remote sensing data

319 The analysis of the current state of Okoniny peatland was conducted using airborne remote sensing
320 data. The data were acquired through a multisensor aerial platform by the MGGP Aero company on March
321 25, 2022 (leaf-off collection) and July 20, 2022, one of the warmest days of the year, which was particularly
322 important for acquiring thermal data (leaf-on collection). Multispectral images (acquired with the IXM-100
323 camera) and Airborne Laser Scanning data (ALS; acquired with the Riegl VQ780-II scanner) were obtained
324 in the leaf-off season. Subsequently, during the vegetation season, the dataset was enhanced by acquiring
325 hyperspectral data (collected using the HySpex VS-725 scanner) and thermal data (obtained with the
326 InfraTEC 9400 camera). Based on the multispectral images, an orthophotomap was generated with a Ground
327 Sampling Distance (GSD) of 10 cm. Hyperspectral data were used to create a mosaic consisting of 430
328 bands (in the range from 400 to 2500 nm), ALS data were applied for the development of a Digital Terrain
329 Model (DTM), and thermal data were used to produce a land surface temperature (LST) mosaic. Thermal
330 and hyperspectral mosaics and DTM were prepared with GSD = 1 m.

331 Photo interpretation was carried out to assess the extent of peatlands and the course of drainage ditches
332 using orthophotos and DTM as a base map. DTM was also used to delineate the catchment area of the
333 peatland. Hydrological modelling methods based on watershed analyses were employed for this purpose. A
334 hyperspectral mosaic was used to calculate spectral indices such as the Normalized Difference Vegetation
335 Index (NDVI; Rouse et al., 1974) and Moisture Stress Index (MSI; Hunt and Rock, 1989). Spectral indices
336 are mathematical formulas that enable the simultaneous analysis of reflectance across multiple spectral
337 ranges. The NDVI is a measure of healthy, green vegetation ranging from -1 to 1. Vegetation values
338 typically range from 0.2 to 0.8, with higher values indicating healthier and denser vegetation. The MSI index
339 is sensitive to increasing leaf water content. Its values range from 0 to more than 3, but the common values
340 for vegetation are from 0.4 to 2. Higher values indicate greater water stress and less water content in this
341 case. Thermal data was used for calculating Land Surface Temperature (LST), measured in degrees Celsius.

Z komentarzem [MB21]: Renumbering

342

343 **2.12. Historical maps and cartographic information**

344 Several historical cartographic studies were used to assess changes to the peatland and its surroundings.
345 The oldest of the materials used is the Schrötter-Engelhardt map of 1803. Work on creating the map began
346 in 1796 under the leadership of the Prussian government minister Friedrich Leopold von Schrötter (1743-
347 1815) and topographer Friedrich Bernhard Engelhardt (1768-1854). The manuscript was produced at a scale
348 of 1:50,000. Still, due to the concerns of the Prussian army command about the map being too detailed and
349 capable of being used by enemy armies, a generalized version was eventually published at a scale of
350 1:150,000. A larger-scale version of the map was not available until the 1920s (Jäger, 1982, 1981). In this
351 article, the generalized version of the map is interpreted.

352 The Prussian topographic map Messtischblatt of 1874 on a scale of 1:25 000, sheet No. 982, Zalesie
353 section, was also analyzed. and the Detailed Map of Poland issued by the Military Geographical Institute in
354 1933 at a scale of 1:25,000, PAN map sheet 34 - SLUP 26 - B (Linsk). In addition, a geological-agricultural
355 map compiled between 1899 and 1900 on the topographic Messtischblatt of 1874 was considered. The
356 Prussian Geological Survey produced the map (Königlich-Preußische Geologische Landesanstalt) and
357 provides information on alluvial and diluvial deposits covering the area under study. The maps show the
358 changes in the peat bog and its surroundings from the early 19th century to the 1930s. Aerial images from
359 1964, 1984, and 1997 obtained from the Central Office of Geodesy and Cartography were also used for the
360 same purpose (license no. DIO.7211.457.2023_PL_N).

361 **Insect outbreak data are based on the literature (Orłowicz, 1924; Schütte, 1893; Wilson, 2012).**

362

363 **3. Results and interpretation**

364 **3.1. Age-depth model and peat accumulation rate**

365 The age-depth model showed a model agreement index (A_{model}) of 60% (Fig. 2), precisely at the limit
366 of the recommended minimum for its reliability (60% according to Bronk Ramsey, 2008). The model
367 spanned the period of ca. 282 years, with a maximum uncertainty ca. 30 years (mostly in the section of ca.
368 1883-1783 cal. CE). Most of the core consisted of well-preserved *Sphagnum* peat, while the lower part
369 consisted of sedge peat. The peat accumulation rate averaged 3.6 mm/yr, with the highest values associated
370 with the undecomposed acrotelm zone. The upper layers located between 0 and 11 cm were excluded from
371 the analysis of peat accumulation rates. The fastest rate was 0.71 cm/yr (at 11.5 cm), and the slowest was
372 0.1 cm/yr (at 91.5 cm). The mean BD value across the core was 0.07 g/cm³. It was highest in the lower part
373 of the core with 0.10 g/cm³ between 96 and 70 cm, and lowest in the middle part - 0.05 g/cm³, between 69
374 and 30 cm. In the upper part between 29 and 0 cm, it was 0.06 g/cm³. **Similarly, this upper, undecomposed
375 layer was excluded from the peat carbon accumulation rate (PCAR) analysis. For the rest of the core (11-**


Z komentarzem [MB22]: Renumbering

Z komentarzem [MB23]: We added information on when the insect outbreak data originated.

376 96 cm), PCAR averaged 112 gC/m²/yr. The mean water content of the wet sample was 93.8%, and the mean
377 organic matter content of the dry sample was 95.5%.

378

Z komentarzem [MB24]: Completion of missing data for Fig. 2



379
380 Figure 2. Age-depth model of the peat profile in Okoniny based on ¹⁴C dating. ¹⁴C age-depth model of the
381 Okoniny peat profile. Water content, organic matter content, bulk density, and PCAR are also marked.

Z komentarzem [MB25]: Editorial and linguistic corrections

382

383 3.2. Palaeoecological analyses

384 3.2.1. Phase 1 (~1726–1838, 96–74 cm): wet conditions and low human impact

385 The high concentration of non-pollen palynomorphs (NPPs) such as cyanobacteria and the algae
386 *Tetraëdron minimum*, *Scenedesmus*, *Botryococcus*, and *Pediastrum* point to the presence of a shallow water
387 body in this time (Fig. 5). This was also confirmed by the plant macrofossils and pollen analyses. The plant
388 macrofossil and pollen analyses point to the presence of a shallow water body during this time interval.
389 Plant macrofossil analysis (Fig. 3) showed that the peatland vegetation in this phase was strongly dominated
390 by vascular vegetation, mainly monocotyledons with *Carex* spp. Shallow waters and edges of the water
391 body were overgrown by sedge communities (Cyperaceae pollen) (2.8-14.5%) (Fig. 5). Additionally, this
392 was indicated by the presence of macrophytes represented by pollen of *Potamogeton* subgen.
393 *Eupotamogeton* (0-0.9%), *Nymphaea* (0-0.4%), and *Utricularia* (0-0.3%) (Fig. 5). The high shares of

394 aquatic non-pollen palynomorphs (NPPs) such as cyanobacteria and the algae *Tetraëdron minimum*,
395 *Scenedesmus*, *Botryococcus*, and *Pediastrum* (Fig. 5) confirms results of plant macrofossil and pollen
396 analyses.

397 This phase was also characterized by the brown moss *Straminergon stramineum* (max. 9% of the
398 subsample content) (Fig. 3). This species occurs in a wide range of habitats (Hedenäs, 1993) but is most
399 common in wet, moderately acidic habitats (Blockeel, 2010). *Straminergon stramineum* is usually found as
400 scattered stems or small patches among other mosses but occasionally forms scattered mats, sometimes
401 partially submerged in water, next to lakes, on the edges of peat bogs or in lakeside marshes (Hill and
402 Blockeel, 2014).

403 This phase of peatland development was characterized by a very low concentration of testate amoebae
404 in the samples. *Centropyxis aculeata* was the most abundant species (Fig. 4). The dominance of plagiostomic
405 species from the genus *Centropyxis* may point to the presence of mineral input into the peatland
406 (Lamentowicz et al., 2009a; Marcisz et al., 2020a). The water level in the peatland was quite unstable and
407 fluctuated between 4.3 and 16.5 cm below the ground and the pH value ranged between 4.5 and 5.2, but due
408 to the low number of identified tests, these reconstructions should be taken suggestively viewed with caution
409 (Fig. 4).

410 The surrounding vegetation was characterized by the dominance of forests, as evidenced by the high
411 proportion of arboreal pollen (AP) (83.6-91.1%) in total pollen content (TP) (Fig. 5). The main species
412 recorded was were *Pinus sylvestris* (62.6-81.3% AP) and *Betula* (6.8-16.0% AP), with admixtures of *Alnus*
413 (2.5-7.7% AP), *Quercus* (1.8-8.1% AP), *Corylus avellana* (0.6-3.8% AP), *Carpinus betulus* (0-3.4% AP)
414 and *Fagus sylvatica* (0.4-3.3% AP). Values of Cerealia pollen sum (0-7.8% TP) with *Centaurea cyanus*, a
415 crop weed, indicated a stable presence of cultivated fields.

416

Z komentarzem [MB26]: Editorial corrections as suggested by the reviewer. We have changed the order of the information in this paragraph so that the information about Figure 3 appears first. This is with the idea that the information on lithology/plant composition forming the peat should appear before the results obtained by other proxies.

Z komentarzem [MB27]: Editorial and linguistic corrections

Z komentarzem [MB28]: Editorial and linguistic corrections

443 The local vegetation (Fig. 3) in this phase was dominated by *Sphagnum*, first by the subgenus
444 *Subsecunda*, then for most of this period by *Sphagnum papillosum*. *S. papillosum* occupies the more
445 oligotrophic lawns with a preference for open space (Clymo and Hayward, 1982; Laine et al., 2018). Along
446 with the appearance of *Sphagnum* from the subgenus *Subsecunda*, *Drosera intermedia* was also recorded.
447 Currently, in Poland, it is a very rare species, found in dispersed peatlands (Mirek et al., 2006). Individuals
448 often stand in the water even throughout the season. *Andromeda polifolia* also appeared in this phase.
449 Initially, the presence of *Sphagnum* was accompanied by *Straminergon stramineum* (max. 10%), but later it
450 disappeared completely. By the beginning of the twentieth century, a relatively high proportion of
451 monocotyledonous plants was also observed, represented in the samples by their epidermis, averaging about
452 20% in a sample, with a much higher proportion in the early stages. All these taxa indicate an intermediate
453 environment between a shallow lake and a moss peatland.

454 After an initial decline (from 9.2 cm at 73.5 cm, 1838 cal. CE, to 13.0 cm at 66.5 cm, 1862 cal. CE),
455 the water table level increased and stabilized at a high level, reaching a maximum of 6.8 cm at 47.5 cm,
456 1907 cal. CE (Fig. 4). The abundance of individual testate amoeba species also increased. Initially, *C.*
457 *aculeata* dominated, but later *Amphitrema wrightianum* and *Hyalosphenia papilo*, mixotrophic taxa that
458 contain endosymbiotic photosynthetic algae, begin to prevail (Lamentowicz and Mitchell, 2005a; Marcisz
459 et al., 2020a) (Fig. 4). Subsequently, the proportion of *A. wrightianum* and *H. papilo* began to decline in
460 favour of *Archerella flavum* and *Hyalosphenia elegans* (Fig. 4). All four species are associated with the
461 presence of *Sphagnum*, with *A. flavum* and *A. wrightianum* tolerating very wet or even submerged
462 *Sphagnum* habitats, which corresponds to a stably high-water table. Then, from the mid-1880s for another
463 ca. 20 years, *C. aculeata* again became dominant. After this period, species associated with *Sphagnum*–*A.*
464 *wrightianum*, *A. flavum* and *Heleopera sphagni* – began to dominate again. During this phase, further
465 acidification of the site was noted through a drop in the pH value from the initial 4.8 to 4.1 (Fig. 4).

466 The forests surrounding the peatland (55.1–92.7% TP) were still dominated by pine (64.5–92.8% AP),
467 although their percentage has decreased in comparison to phase 1, especially during the 1920s and 1930s
468 (Fig. 5). Deciduous taxa such as *Quercus*, *Corylus avellana*, *Carpinus betulus* and *Fagus sylvatica* retreated.
469 The percentage of Cerealia in the TP increased significantly, from 0–7.8% TP in the first phase to 2.8–19.8%
470 in the second phase, with a peak in the late 1910s and early 1920s, indicating the development of agriculture
471 in the vicinity of the peatland (Fig. 5). Around the same time, the proportion of *Rumex* also increases
472 significantly (0–11.5%). The low values of MAC (Fig. 3) and MIC (Fig. 5) indicate a low fire activity in the
473 studied area.

474
475 **3.2.3. Phase 3: (~1945–present, 37–0 cm): Lowering of the groundwater table as a result of climate**
476 **change, further afforestation with *Pinus sylvestris*, a succession of *Betula***

Z komentarzem [MB32]: Editorial corrections

477

478 The local vegetation (Fig. 3) underwent several changes during this phase. Although *Sphagnum*
479 dominated for the entire time, the subgenus *Sphagnum* receded in favour of first the subgenus *Cuspidata*
480 and then the subgenus *Acutifolia*. The beginning of the phase was marked by *Pohlia nutans*, which can win
481 the competition in unstable habitat conditions, such as during the dry season (Boulc'h et al., 2020). Its
482 occurrence correlated with the presence of Phryganella acropodia among testate amoebae (Fig. 4), which is
483 an indicator of low water levels in Sphagnum peatland (Diaconu et al., 2017; Lamentowicz and Mitchell,
484 2005b).

485 This was followed by *Alabasta militaris* (\bar{x} = 25.5%), *Galeripora discoides* (\bar{x} = 10.5%) and *Nebela*
486 *tincta* (\bar{x} = 8.2%) beginning to dominate (Fig. 4). *G. discoides* is typically present in acidic sites with
487 unstable hydrological conditions (Lamentowicz and Mitchell, 2005b; Sullivan and Booth, 2011). *N. tincta*
488 tolerates dry, highly acidic conditions with mineral matter supply (Booth, 2002; Koenig et al., 2018;
489 Lamentowicz et al., 2011). *A. militaris*, dominant in recent years, is indicative of dry and markedly acidic
490 conditions (Amesbury et al., 2016; Booth, 2002; Lamentowicz et al., 2011; Marcisz et al., 2020a; Sullivan
491 and Booth, 2011). Based on testate amoebae, this phase was distinguished by a significant drop in the
492 groundwater table, from an average level of 9.6 cm below the ground surface in the second phase to 15.7
493 cm. In the last decade, the most significant decline was observed, with an average level of 21.9 cm, with a
494 maximum of 27.5 cm, 1983 cal. CE. The pH continued to decrease – from 4.4 to 4.0 (Fig. 4).

495 On a regional scale, there is an increase in the relative abundance of *Pinus* pollen in the TP, from about
496 46% at the beginning of the phase to about 85% today as an effect of afforestation (Fig. 5). *Betula* pollen
497 concentration share has an apparent increase, from 0,7-11,3% in the second phase to 5,6-32,5%. The
498 increased concentration percentage of *Betula* pollen, combined with macroscopic remains in the form of
499 achenes and catkin scale scales, indicates the intensive succession of this species on the peatland surface.
500 The ruderal species *Urtica* and *Artemisia* were also more strongly manifested. The average proportion of
501 *Urtica* pollen in the TP TPS increased almost 8 times distinctly (from 0-0.7% to 0-2.9%). The percentage
502 of Cerealia in TP has decreased significantly, from nearly 20% in the early 1920s to just over 1% today.

503 Local (Fig. 3) and regional (Fig. 5) fire activity continued to be low, although two slightly more
504 intensive periods of regional fires were marked – ca. 1945-1963 and the early 2020s.

505

506 3.3. Dendrochronological and pointer years analysis

507 A total of 50 tree-ring series of 23 *Pinus sylvestris* L. trees from the Woziwoda site were successfully
508 cross-dated. Based on the well-synchronized tree-ring series TRW (Fig. 6) and RWI site chronologies upon
509 the TRW (Fig. 6) and RWI sites, well-synchronised tree-ring series spanning 222 years (1801-2022) were
510 was developed. The statistical characteristics of the ring-width series and the statistical parameters

Z komentarzem [MB33]: Editorial and linguistic corrections

Z komentarzem [MB34]: Editorial and linguistic corrections

Z komentarzem [MB35]: Editorial and linguistic corrections

Z komentarzem [MB36]: Editorial and linguistic corrections

Z komentarzem [MB37]: Editorial and linguistic corrections

Z komentarzem [MB38]: Editorial and linguistic corrections

511 indicating the signal strength of the regional RWI chronology are shown in Tab. 2. The mean EPS was 0.93,
 512 which is well above the threshold value (EPS = 0.85) required to produce a statistically robust RWI
 513 chronology. Mean series inter-correlation, MS, SNR, and other statistical parameters indicating the strength
 514 of chronology signals were also high, indicating the suitability of chronology for climate-growth analysis.
 515 Tab. 2 Descriptive statistics of standardized *Pinus sylvestris* L. (RWI) chronology for Woziwoda site


Chronology length	1801-2022
Mean tree age [yrs]	197
Number of tree/cores	23/50
Mean ring width (mm) ± SD	1.256 ± 0.702
Series intercorrelation	0.623
Average mean sensitivity	0.265
Expressed Population Signal (EPS)	0.93
Signal-to-noise ratio (SNR)	12.97
Rbar.eff (effective chronology signal)	0.361

516
 517 Across the study period (1920-2022) a significant positive relationship between growth and
 518 February mean temperature was identified (Fig. 7). The moving correlation analysis showed an increasing
 519 trend in the sensitivity of tree growth to climatic factors (Fig. 8). The positive response of tree growth to
 520 February mean temperature remained constant throughout the study period (1920-2022) (Fig. 8). However,
 521 the sensitivity of tree growth to summer temperature increased. The relationship between annual growth and
 522 summer temperature was not stable during the period 1920-2022. Nevertheless, in the last 30 years, a
 523 significant negative relationship between annual growth and June mean temperature was observed.

524 Climate-growth analysis for monthly data did not show a statistically significant relationship
 525 between growth and precipitation (Fig. 7). However, moving response analysis revealed significant short-
 526 term relationships between tree growth and precipitation. Furthermore, it was demonstrated that the
 527 influence of precipitation in the current year's months on tree growth calculated for the years 1960-2022
 528 was more significant than the relationships calculated for the years 1921-1959. In recent years, a particularly
 529 positive relationship between tree growth and early-year (February-April) precipitation as well as June
 530 precipitation has become apparent.


531 For Woziwoda site 8 positive and 13 negative pointer years were identified for the period 1814-
 532 2022 (with a minimum sample depth of 10 trees) (Fig. 6). The most pronounced positive pointer years with
 533 more than 90% tree response were as follows: 1847, 1863, 1912, 1941, 1945, 1957, and 1983. The most
 534 pronounced negative pointer years were: 1839, 1868, 1869, 1911, 1925, 1940, and 1950. Figure 6 provides
 535 marks of pointer years together with meteorological and ecological characteristics.

Z komentarzem [MB39]: Editorial and linguistic corrections



536
 537 Figure 6. The grey lines depict the individual tree ring series of each tree, while the green line represents
 538 the average raw chronology of *Pinus sylvestris* L. at the Woziwoda site. Identified within the Scots pine
 539 chronology from Woziwoda are pointer years, categorized as negative (NEG) (A) and positive (POS) (B).
 540 These pointer years are highlighted with colored asterisks: red for positive pointer years and blue for
 541 negative pointer years. The position of the asterisks refers to a scale of 0-100%. [Information on extreme](#)
 542 [phenomena is based on Orłowicz, 1924; Schütte, 1893, Broda 2000, Wilson 2012.](#)

Z komentarzem [MB40]: We added information on when the insect outbreak data originated.




543
 544 Figure 7. Response function coefficients between residual *Pinus sylvestris* L. chronology *residual Pinus*
 545 *sylvestris* L. chronology and climate variables: (A) mean air temperature (~~TEMP~~ Temp), and (B)
 546 precipitation (Prec) for the period 1920–2022. Names of the previous year’s months start with a lowercase
 547 letter. Solid lines represent significant coefficients at $p < 0.05$.

Z komentarzem [MB41]: Editorial corrections


Z komentarzem [MB42]: Editorial corrections

Z komentarzem [MB43]: Editorial corrections

(A) Temp



(B) Prec



548
549 Figure 8. Moving response correlations (25-year window) between residual *Pinus sylvestris* L. chronology
550 and climate variables: (A) mean air temperature (Temp), and (B) precipitation (Prec) for the period
551 1920–2022. The color code represents the correlation-coefficient response function coefficients. Significant
552 correlations are indicated by white asterisks.

553
554 **3.4. The current state of the peatland based on remote sensing data analysis**


- Z komentarzem [MB44]: Editorial corrections
- Z komentarzem [MB45]: Editorial corrections
- Z komentarzem [MB46]: Editorial and linguistic corrections

555 Presently, the ~~non-forest non-forested~~ part of the peatland is drained by two parallel ditches. One is
556 located in the northern, and the other is in the southern non-forested part of the peatland. The analysis of
557 thermal data obtained on a ~~torrid~~ midsummer day indicates that the average LST for the non-forested part
558 of the peatland is approximately 34.29 °C, with a temperature range extending from 19.22 °C to 46.37 °C.
559 There is a distinct internal variability in LST values within the studied area. Higher values, indicative of
560 more significant dehydration, were identified in the eastern part of the peatland, while lower values were
561 observed in the western part. A repeating spatial pattern of ~~their~~ values was observed in the analysis of
562 vegetation indices (NDVI and MSI). High NDVI values and low MSI values, indicative of good vegetation
563 condition and low water stress, were observed in the western and southwestern parts of the peatland (Fig.
564 9). The average NDVI value in these areas is 0.71, and MSI is 0.6. Conversely, low NDVI values and high
565 MSI values, indicative of significant dehydration of the peatland and low vegetation vigor, were observed
566 in the eastern part of the object (Fig. 9), where NDVI averages 0.63, and MSI is around 0.69. The overall
567 average NDVI for the object was 0.65, and for MSI, it was 0.68.

Z komentarzem [MB47]: Editorial and linguistic corrections


Z komentarzem [MB48]: Editorial and linguistic corrections

Z komentarzem [MB49]: Editorial and linguistic corrections



568
 569 Figure 9. Remote sensing characteristics of Okoniny peatland based on multisensorial airborne data acquired
 570 in 2022.

571
 572 **3.5. Historical maps and airborne images as confirmation of changes shown in palaeoecological data**



573
 574 Figure 10. Changes in the peatland and its surroundings since the beginning of the 19th century based on
 575 historical maps and aerial images. (A) Schrötter-Engelhardt map 1:150 000 (1802), (B) Messtischblatt map
 576 no. 982 1:25 000 (1874), (C) Prussian geological and agricultural map no. 2374 1:25 000 (1905), (D)
 577 Detailed Map of Poland 1:25 000 (1933), (E) Aerial photograph from 1964, (F) Aerial photograph from
 578 1984, (G) Aerial photograph from 1997. Maps no. A, B, C, and D are in the public domain. Aerial
 579 photographs were obtained from © Central Office of Geodesy and Cartography in Poland, license no.
 580 DIO.7211.457.2023_PL_N.

581
 582 Analysis of historical materials (Fig. 10), including maps and airborne images, confirms the results
 583 of the palaeoecological analysis. Both the Schrötter-Engelhardt map of 1802 and the Messtischblatt of 1874
 584 indicate the existence of a small lake in the coring area. Again, however, it should be noted that the Schrötter-
 585 Engelhardt map is a highly generalized study and does not give much information about the surroundings
 586 of today's peatland, other than that we are dealing with an area with the character of a dense forest complex

587 with wetlands in isolated places. Messtischblatt allows us to better interpret the surroundings of the analyzed
588 modern peatland at the time in which the map was prepared. A small lake named "Kolze See" is observed
589 in an advanced stage of development, i.e., progressive overgrowth. This lake is located in the surroundings
590 of wetlands (Bruch in German) somewhat distant heathland (Heide in German) and wasteland (Ödland in
591 German) (the original nomenclature of the map legend was adopted). This lake and two other lakes close
592 by are enclosed within a single catchment area. To the south, the area of the current peatland was adjacent
593 to an open, extensive meadow.

594 Even more information is provided by a 1905 geological-agricultural map prepared on the
595 topographic base Messtischblatt map of 1874. In addition to land use, it shows the type and thickness of
596 alluvial and diluvial deposits. According to this map, the area around the lake was covered by alluvial
597 sediments – humus with peat subsoil and shallow groundwater (org. in German: Humus (Peat) mit Torf-
598 Untergrund und nahem Grundwasser). The thickness of the peat was marked at two meters. However, it
599 should be noted that drilling surveys at that time only covered a maximum depth of two meters, so the maps
600 do not provide information on the total thickness of the sediments (Jasnowski, 1962). Places that were used
601 as heathland and wasteland on the topographic map are covered by sandy humus on a sandy substrate with
602 shallow groundwater (org. in German: Sandiger Humus mit Sand-Untergrund und nahem Grundwasser) and
603 by humic sands on a substrate of permeable sands with shallow groundwater (org. in German: Humoser
604 Sand mit durchlässigen Sand-Untergrund und nahem Grundwasser).

605 A Detailed Map of Poland from 1933 documents the change in an ecosystem from lake to land. The
606 area, which on Prussian topographic maps was a lake with a surrounding bog, is described as a meadow on
607 this map. Moreover, the meadows adjacent to the south were marked with drainage ditches, which were not
608 on the Prussian maps. The area's surroundings, as before, were dominated by coniferous forests.

609 Aerial photos document subsequent changes in the ecosystem. The 1964 photo shows the northern part
610 of today's peatland's agricultural use (regular surface layout). Lake Kały, located nearby, became completely
611 overgrown, and its area was later dug by a drainage ditch, brought to the studied peat bog. The surrounding
612 area of the peatland is dominated by dense forest with occasional open clear-cutting areas. A photo from
613 1984 documents the succession of trees in the north-central part of the peatland. In the surrounding area,
614 open forest areas have entirely disappeared. A photo from 1997 clearly shows the development of trees on
615 the peatland, which have formed a dense block in its north-central part. A distinct area of *Sphagnum*-
616 dominated peatland with a well-marked edge has also emerged. Currently, the northernmost part of the
617 peatland is overgrown by pine; it is almost impossible to identify the maximum extent of the peatland surface
618 in the field (Fig. 1).

619

620 4. Discussion

621 4.1. Exceptionally high peat accumulation rate

622 ~~Peat accumulates when vegetation production exceeds organic losses under high water levels and~~
623 ~~anaerobic conditions (Tobolski, 2000).~~ In the Okoniny peatland, a rapid rate of peat accumulation is
624 observed, averaging 3.56 mm/yr, with a maximum value of 7.1 mm/yr at a depth between 11 and 12 cm.
625 This accumulation rate is not commonly observed. There are only several peatlands in Poland for which
626 higher accumulation rates were reported. In the Tuchola Pinewoods, ~~the faster average rate of peat~~
627 ~~accumulation was recorded at these were~~ Dury – 10 mm/yr (Pawlyta and Lamentowicz, 2010), and Mukrza
628 – 4.6 mm/yr (Lamentowicz and Obremska, 2010). ~~At~~ Jelenia Wyspa mire where the accumulation rates
629 reached 0.4 mm/yr for the first 3000 years but accelerated to 3 mm/yr in the last 150 years (Lamentowicz
630 et al., 2007). ~~A much slower rate was on~~ and the Tuchola kettle-hole bog – 1.2 mm and after ca. 1320 cal.
631 Yr BP the accumulation rate dropped to 0.4 mm/yr (Lamentowicz et al., 2008b). In other pine monocultures,
632 such as the Noteć Forest, the Rzecin peatland stands out for its high accumulation rate – an average of 6.8
633 mm/yr in one profile and 7.5 mm/yr in the other one (Milecka et al., 2017). Peatlands in Tuchola Pinewoods,
634 including Okoniny peatland, generally have a faster accumulation rate than peatlands located in other parts
635 of Pomerania. ~~The especially small kettle-hole peatland peatlands characteristic of Tuchola Pinewoods~~
636 ~~accumulates that accumulate~~ carbon the fastest of all peatland types (Karpińska-Kołaczek et al., 2024). ~~For~~
637 ~~the Stążki mire~~ In Pomeranian peatlands, the highest accumulation rates were reported for the period
638 between ca. 150-1230 AD and reached 2.2 mm/yr in Stążki ~~(unfortunately, the more recent, topmost material~~
639 ~~was not analysed)~~ (Lamentowicz et al., 2008a). ~~Peat accumulation was even slower on Słowińskie Błota~~
640 ~~raised bog – 1.38 mm between 1830 and 2006, although the highest accumulation rate was 5 mm/yr (during~~
641 ~~AD 840-860) and 1.38 mm between 1830 and 2006, although the highest accumulation rate was 5 mm/yr~~
642 ~~(during AD 840-860) in Słowińskie Błota raised bog~~ (Lamentowicz et al., 2009b). At the Gołębiewo sites
643 the maximum accumulation rate was were 1.85 mm/yr for the first site and 0.36 mm/yr for the second site
644 (Pędziszewska and Latałowa, 2016). ~~The average 2 mm/yr accumulation rate for the Kusowskie Bagno bog~~
645 ~~was in its 4000-year history (Lamentowicz et al., 2015).~~ At Gązwa bog, the accumulation rate was estimated
646 at 1.46 mm/yr, more than twice as slow as at Okoniny peatland (Gałka et al., 2015). In other regions of
647 Poland, Jaczno bog (Suwałki Lakeland) stands out where peat accumulation rate in this peatland was very
648 rapid, averaging 2.76 mm/yr, with the highest values recorded in undecomposed and uncompact acrotelm
649 – up to 12.7 mm/yr (Marcisz et al., 2020b). At the Pawski Ług bog PAR was similar – 2.6 mm/yr
650 (Lamentowicz et al., 2020). For many *Sphagnum*-dominated peatlands in other parts of Poland, the average
651 PAR varied between 1.4-2.5 mm/yr (Gałka et al., 2015; Lamentowicz et al., 2020; M. Lamentowicz et al.,
652 2015; Marcisz et al., 2020b). Such high accumulation rate values are also rare in other parts of the temperate
653 climate zone of Europe. Teici bog (Latvia) showed similar accumulation rates - 3.5 mm/yr - from 1835 to
654 1965 AD and 10 mm/yr after 2000 (Stivirins et al., 2018). Okoniny peatland after 2000 (between 21.5 and

Z komentarzem [MB50]: We have removed the sentence as suggested.

655 11.5 cm) recorded an accumulation of 5.7 mm/yr. Saxnäs mosse in Sweden showed an almost linear peat
656 accumulation rate of 2-2.5 mm/yr (van der Linden et al., 2014). The maximum accumulation was recorded
657 at around 2310-2250 cal on the Estonian Hara bog. BP (31-15 cm) reaching 2.4 mm/yr (Łuców et al., 2022).
658 A comparison with other regions of Poland and Europe shows that the exceptionally high accumulation rates
659 at the analyzed site are worth highlighting.

Z komentarzem [MB51]: We shortened this part of the discussion and added information on why this comparison is important for our study.

661 4.2. Relationships between forest management and pollen analysis

662 4.2.1. The complex history of the Tuchola Pinewoods and its influence on the forest

663 The results of pollen analysis of the collected core enabled us to illustrate how the forest was managed
664 over the past 300 years. Due to political changes and several administrative decisions, the management
665 strategies of the Tuchola Pinewoods underwent vital changes. The consequences of the implementation of
666 forest management techniques were visible in the palaeoecological record.

667 In 1772, the area of Gdansk Pomerania with the Tuchola Pinewoods was included in the borders of
668 the Kingdom of Prussia as a result of the First Partition of Poland (Wilson, 2012). At that time, some of the
669 first legal regulations for planned forest management in the area appeared (Jaszczak, 2008a). Nevertheless,
670 in 1775, Frederick II the Great (1740-1786) issued a decree regarding government forests in Prussia, and
671 later state forests in Poland. It proposed the division of the forest into districts consisting of 50 man-made
672 clearings. However, this method of forest management worked well in the area of small forests in western
673 Prussia, but not for large forest complexes like the Tuchola Pinewoods. In 1782 Frederick II the Great issued
674 a special decree "On the development of the Tuchola Forest," in which it was written that the area of the
675 Tuchola Pinewoods was to be divided into eight districts of about 6000 hectares and 60 cutting areas in each
676 district. Only one man-made clearing in each district was provided for economic use, so no more than 100
677 hectares of forest per district (Jaszczak, 2008b). However, despite the introduction of many regulations
678 relating to forest management, the first decades of the 19th century brought devastating and predatory
679 deforestation on a large scale. For most of the 19th century, progressive deforestation was a problem in the
680 region, making the already poor conditions for agricultural development much worse (Wilson, 2012). With
681 each successive partition of Poland (1772, 1793, 1795), the Prussian government took over the state forests,
682 first the royal table estates (in Polish: dobra stolowe), and then also royal land (in Polish: królewsczyzny).
683 After the second partition, state forests were separated from agricultural land and transferred to separate
684 administrations (Nienartowicz, 2012). In 1810, the Prussian government issued the so-called Secularization
685 Act, under which forests were removed from churches and monasteries and attached to state forests. A law
686 had been in effect in Prussia since 1713, which prohibited the selling of state-owned property, but the
687 approach was different in Prussia.

688 To carry out the fastest possible Germanization of these lands, the local state property was sold off to
689 the Prussian nobilities, inviting them to settle in the area. Prussia's defeat of Napoleon's forces at the Battle
690 of Jena-Auerstedt in 1806 and the contribution Napoleon imposed in 1808 also contributed to the selling off
691 of the state forests. In December 1808, the government in Berlin passed an edict (published in November
692 1809) allowing the sale of state land, including forests, to cover the national debt (Broda, 2000; Kozikowski,
693 1911). Only large, compact forest complexes of a protective nature and economic importance were excluded
694 from this regulation (Broda, 2000). An additional reason for the loss of state forest area was the need to
695 redeem the servitude rights vested in peasants when selling property. The government compensated the
696 peasants for their rights to use the forest by transferring other forests without looking at area losses. The
697 peasants most often cut down all the forest given to them and turned the land into agricultural land. The
698 cause for such action was provided by the 1807 edict that removed state supervision of private forests, which
699 was later extended in 1811 (Broda, 2000; Kozikowski, 1911). Private forests could be freely managed from
700 then on, including dividing into smaller parcels, converting to agricultural land and selling.

701 This period was also a time of intense social and economic change, marked by the collapse of
702 feudalism in favour of a capitalist economy. The end of these transformations in Prussia was the
703 enfranchisement of peasants in 1811-1823. Economic development entailed a considerable demand for
704 timber, and this, in turn, became the basis for the robbery economy in forests. The selling off of state forests
705 slowed down only in the 1830s largely due to the efforts of G.L. Hartig, the general director of state forests
706 in Prussia, and stopped entirely in 1860.

707 Exploited forest areas were restored mainly with pine and spruce, either artificially or naturally.
708 Because of this, deciduous admixture species with entirely different life requirements began to disappear
709 over time (Broda, 2000). The introduction of easier-to-maintain coniferous species was driven by the
710 growing demand for wood in industry. The trend toward introducing pine monocultures intensified from the
711 1830s onward. Since forest management in Prussia's state forests served mainly fiscal purposes, the concept
712 of monoculture plantations did well for several more decades. This situation persisted until the 1860s when
713 a devastating pest gradation occurred (Broda, 2000). At that time, the first steps were taken regarding the
714 introduction of admixtures into restoration.

715 With the first partition of Poland in 1772 by Prussia, regulations for planned forest management began
716 to be introduced. The main planting species was Scots pine, which over time began to dominate the forest,
717 replacing deciduous admixture species. The region's forest cover and forest composition were also affected
718 by later political and administrative developments. For more information on the history of forest
719 management in the late 18th and early 19th centuries, see Supplementary File 1.

720 Our data confirm an increase in the proportion of pine pollen in the forest composition and a decrease
721 in the proportion of pollen of other species. From the 1730s to the mid-1860s, the share of pine pollen in

Z komentarzem [MB52]: We moved a large part of the text with historical information to Supplement No. 1, to reduce the length of the discussion. We left only a brief description of historical events that will be enough for the readers to understand the context of the study.

722 the pollen of all trees increased from about 60% to about 90%. Our pollen diagram shows the rapid increase
723 in *Pinus sylvestris* pollen **concentration percentage** after 1850. It can, therefore, be assumed that this resulted
724 from *Pinus sylvestris* introduced by mass monoculture plantings in the early 1830s reaching reproductive
725 capacity. Pine usually reaches sexual maturity between 10 and 15 years (Sullivan, 1993), although the
726 threshold age has been set at 25 years (Matthias and Giesecke, 2014). The decline in the share of deciduous
727 species and the increase in the share of Scots pine in the landscape began in Poland with the formation of
728 the state. However, at that time, it was associated with the expansion of agriculture and the harvesting of
729 preferred species such as *Carpinus betulus* (Czerwiński et al., 2021) Nevertheless, in the Prussian partition,
730 planned forest management permanently changed the composition of **Poland's Polish largest forest**
731 complexes, which were dominated by easy-to-grow pine (Broda, 1993) **(see Supplementary File 1)**. A
732 dynamic increase in the share of pine pollen until the 1860s in the Tuchola Pinewoods was also recorded at
733 the Czechowskie Lake (Słowiński et al., 2019). An increase in pine pollen **concentration percentage** since
734 the 19th century was also shown in pollen diagrams of other sites from Pomerania – Stążki (Lamentowicz
735 et al., 2008a), Słowińskie Błota (Lamentowicz et al., 2009b) – and in other monoculture plantation
736 complexes from **the** Prussian partitioning area – Rzecin peatland in the Noteć Forest (Milecka et al., 2017).

737 Although attempts were undertaken to correct earlier mistakes, this did not stop the massive
738 deforestation **(among other consequences of war events and administrative regulations on settlement, more**
739 **in Supplementary File 1)**. Until the 1870s, the feudal system was still mixed with capitalist components, but
740 from the 1870s onward, under monopoly capitalism, timber trade and processing began to reach a significant
741 size (Broda, 2000). However, it has been noted that forests regulate air temperature, store water in the soil
742 more efficiently, and reduce wind speed, preventing soil erosion, which can help local agriculture face
743 difficult environmental conditions (Wilson, 2012). For this reason, as early as the 1870s, the state
744 administration encouraged landowners to protect forest stands on their lands and establish forestry
745 cooperatives. The government also guaranteed funds for the reforestation of private and municipal lands. In
746 the mid-1870s, the Landtag set aside a budget for the purchase and reforestation of wasteland by the state.
747 However, these funds were used to a small extent, although this somewhat reduced the share of forested
748 private property (Broda, 2000; Wilson, 2012). In 1886, the Royal Settlement Commission (in German:
749 Königliche Ansiedlungskommission) was established to buy up the estates of impoverished Polish nobility
750 to acquire agricultural land for German settlers (Wilson, 2012).

751 At the end of the 19th century, Tuchola Pinewoods became the largest timber production hub in the
752 Prussian partition. The Bydgoszcz timber industry region also played a major role in wood processing. The
753 first steam sawmill in the Bydgoszcz region was built in 1873, and by 1913, there were 20 of them,
754 processing some 500,000 m³ of wood and employing more than 1,600 people (Broda, 2000). All this resulted
755 in a significant decline in the **concentration share** of tree pollen in the total pollen **concentration share** in our

Z komentarzem [MB53]: Editorial and linguistic corrections

Z komentarzem [MB54]: Editorial and linguistic corrections

Z komentarzem [MB55]: Editorial and linguistic corrections

Z komentarzem [MB56]: Editorial and linguistic corrections

Z komentarzem [MB57]: Editorial and linguistic corrections

Z komentarzem [MB58]: Editorial and linguistic corrections.
We explained shortly the reasons for the deforestation.

Z komentarzem [MB59]: Editorial and linguistic corrections

Z komentarzem [MB60]: Editorial and linguistic corrections

756 diagram, to less than 60% by the late 1920s and early 1930s. At the same time, we have seen intensive
757 agricultural development. At Okoniny, the proportion of Cerealia pollen doubled between ca. ~~1900-1920~~
758 ~~1900 and 1920~~. This trend is also confirmed by pollen data from the site in Okoniny Nadjeziorne, on the
759 other side of Okonińskie Lake (Tipton, 2023), as well as from Czechowskie Lake, about 25 km northeast of
760 our site (Słowiński et al., 2019). Despite intensive deforestation in general, further afforestation with pine
761 was also progressing. In 1893, pine forests accounted for 99% of all forests in Tuchola County
762 (Szwankowski, 2005). Intense changes in forest management (pine dominance) and agricultural
763 development (high ~~concentration percentage~~ of Cerealia pollen) in the 19th century are also evident in
764 records of profiles outside large, dense forest complexes – Kusowskie Bagno (Gałka et al., 2014), Linje
765 mire (Marcisz et al., 2015).

766

767 4.2.2. Impact of forest management on peatland vegetation

768 As a result of changes related to forest management, lake to peatland transition occurred rapidly. We
769 assume that this was primarily the result of drainage, which was undertaken in the area at the end of the 19th
770 century (see drainage ditches on the southern side and a dike in the middle part of the site on maps in Figure
771 6), and secondly, to a lesser extent, the transition from mixed forests to pine monoculture. These activities
772 contributed to an increase in the acidity of the peatland. Forest drainage is often associated with the
773 acidification of surface waters (Miller et al., 1990). The introduction of forest drainage, on or near peatlands,
774 to improve tree growth has been quite common in northern and northeastern Europe (Westman and Laiho,
775 2003). The oxidation of organic sediments and the detachment of ~~hydrogen ions H+ ions~~ increase acidity
776 (Ulrich, 1980). In addition, the supply of alkaline cations to the peat is impeded by drainage ditches
777 (Minkkinen et al., 2008). However, the long-term consequences of drainage are devastating to peatlands, as
778 they initiate vegetation succession, in which species typical of peatlands are replaced by forest vegetation
779 (Laine et al., 1995). In the example of our palaeoecological data, the dynamic succession of pine and birch
780 in the Okoniny peatland is evident, which is also supported by aerial imaging. As already mentioned, the
781 successive decline in pH is also the result of the impact of pine plantations growing in catchments. A drop
782 in pH in Okoniny has likely enabled the rapid growth and expansion of *Sphagnum* and the peatland
783 initiation. The crowns of forests, especially the needles, can increase the uptake of atmospheric pollutants
784 such as sulfur and nitrogen components, contributing to the acidification of surface waters (Nisbet, 2001;
785 Reynolds et al., 1994). Conifers also can capture ions of marine origin - Na and Mg cations. These, in turn,
786 displace hydrogen and aluminium cations from the soil, leading to acid runoff from the forests along with
787 surface runoff, which is known as the "sea-salt effect" (Drinan et al., 2013; Harriman et al., 2003; Reynolds
788 et al., 1994). We observed the presence of *Pinus* needles at the beginning of phase 2 (from 1838 cal. CE), at
789 the transition from pond to peatland ecosystem. Moreover, *Pinus* stomata were also present in palynological

Z komentarzem [MB61]: Editorial and linguistic corrections

Z komentarzem [MB62]: Editorial and linguistic corrections

Z komentarzem [MB63]: Editorial and linguistic corrections

790 samples at that time, pointing to more frequent needle falls. More pine trees in the Tuchola Pinewoods
791 resulted in much higher amounts of needles and other pine fragments accumulating on the forest ground,
792 leading to soil acidification. This, together with drier conditions, could quickly lead to acidification around
793 the pond, forming perfect conditions for *Sphagnum* to encroach – first as a floating mat that successively
794 overgrown the pond. We sampled the peat core close to the edge of the peatland, thus in the place where
795 moss encroachment on the open water body began; therefore, we were able to track this succession in our
796 record. This succession and disappearance of Lake Kolze are also clearly visible in historical maps (Figure
797 10). Other examples of quick encroachment of floating mats on the surface of the lake have been observed
798 and mapped in other open water bodies in the Tuchola Pinewoods (Kowalewski, 2003; Kowalewski and
799 Milecka, 2003) and other regions (Warner, 1993).

800

801 **4.3. Anomalies and extreme events**

802 **4.3.1. The impact of droughts and fires on the forest and peatland**

803 Historical sources indicate that in the 18th and 19th centuries, the Tuchola Pinewoods were relatively
804 often affected by droughts resulting in fires (Wilson, 2012). In 1781, there was a fire in Tuchola (ca. 16 km
805 SW), during which a large part of the city with the church and town hall burned down, and in 1792, Starogard
806 Gdański (ca. 42 km NE) burned almost to the ground (Orłowicz, 1924). Major fires also occurred in 1794,
807 and 1807, when more than 34,000 hectares of forest burned (Orłowicz, 1924; Schütte, 1893). Fires in 1809,
808 1810, 1812, 1813, and 1828 in the Świt forest district about 15 km from the study site were also recorded
809 (Cyzman, 2008). Palaeoecological data, especially MIC, confirm high fire activity in the first decades of the
810 19th century (a rapid increase). Słowiński et al. (2019) emphasized that data on fires before the 1830s,
811 especially regarding their area, should be treated with caution due to the lack of accurate measurement
812 techniques. In the Woziwoda Forest District, within which the Okoniny peatland is located, the forests of
813 the Biała and Barłogi forest districts also burned in 1842 (Cyzman, 2008). Intense fires also appeared in the
814 Tuchola Pinewoods between 1846 and 1848 (Orłowicz, 1924; Schütte, 1893).

815 Later, numerous fires were also reported in the Woziwoda Forest District. Between 1860 and 1889,
816 310 fires were observed, destroying 4206 hectares of the forest (Orłowicz, 1924; Schütte, 1893). The highest
817 number of fires in this period was registered in 1862-1864 and 1874-1875 when 3565 hectares of forest
818 burned; altogether, nearly 85% of the area burned in 1860-1889 (Schütte, 1893). The largest area burned in
819 1863 equaled 2333 hectares, including more than 1250 hectares in the Woziwoda forest district; altogether,
820 25% of all the forest burned in 1860-1889 (Orłowicz, 1924; Schütte, 1893). Meteorological data confirm
821 dry years in the period from 1862 to 1865. In 1862 and 1863, the annual precipitation in Bydgoszcz was
822 only a little over 450 mm (Kirschenstein, 2005), and it was then that the largest number of hectares of forest
823 in the known history of the Tuchola Forest burned (Dietze et al., 2019).

824 ~~However, contemporary linked the number of fires with political events and nationalist sentiment~~
825 ~~among the Polish population~~ The number of fires can also be linked to political events (Orłowicz, 1924;
826 Schütte, 1893; Wilson, 2012). In 1901, in the nearby Trzebciny and Gołębek Forest Districts, a fire
827 consumed 663 hectares of forest (there was a parallel children's strike in Września Province) (Orłowicz,
828 1924; Wilson, 2012). Fires could also be caused by agricultural activities and land preparation for crops
829 (Poraj-Górska et al., 2017). By the 1830s, charcoal production was widespread (McGrath et al., 2015), and
830 forest burning was used to create heathlands for beekeeping (Bienias, 2009).

Z komentarzem [MB64]: Editorial and linguistic corrections

831 Fires of the 1860s provide a regional signal at another site in the Tuchola Pinewoods – Czechowskie
832 Lake (Dietze et al., 2019). Increased fire activity in the mid-19th century was also observed at the Lake
833 Jaczno site (Poraj-Górska et al., 2017). At the Okoniny peatland, MIC and MAC values decreased after
834 1850, but at the same time, the water level stabilized and remained high. Fire activity remained low in areas
835 where wet conditions prevailed, such as southern Finland (Väliranta et al., 2007) and eastern Estonia
836 (Sillasoo et al., 2011).

Z komentarzem [MB65]: We added citations.

Z komentarzem [MB66]: We added citations.

837 In 1948, about 450 hectares of forest were burned near Osieczna, and in 1954, 80 hectares were burned
838 near Ocypel (Cherek, 2007). Palaeoecological data record an increased MIC supply during this period. The
839 first of these fires was also recorded in the sediments of Czechowskie Lake (Słowiński et al., 2019). The
840 summer drought of 1921 occurred over a larger area of Europe, from Poland and the Czech Republic to the
841 UK (van der Schrier et al., 2021). Summer droughts also affected the Tuchola Pinewoods in 1951 and 1959.
842 In 1959 Bydgoszcz received only 37 mm of precipitation from August to October (Mitosek, 1960), and from
843 1950 to 1958 Bydgoszcz received less than 500 mm of rain per year (Kirschenstein, 2005). Our
844 palaeoecological data confirm droughts in the 1950s. There is a sharp increase in the proportion of
845 *Phryganella acropodia* among the testate amoebae, an indicator of dry conditions (Diaconu et al., 2017);
846 ~~and a high concentration of coprophilous fungi~~ and an expansion of brown mosses in the form of *Pohlia*
847 *nutans* (up to 30% of the peat sample composition) are also marked. Dendroclimatic data recorded the
848 negative impact of climatic conditions on pine, especially strongly in 1950 and 1956.

Z komentarzem [MB67]: Editorial and linguistic corrections

849 Studies show that particle size illustrates the distance of the fire from the site, the heavier the particles,
850 the shorter distances they travel (Clark, 1988; Peters and Higuera, 2007). However, many factors determine
851 the particles' transport—the fire's intensity, the burning areas, and the wind direction. Adolf et al. (2018) point
852 out that the charcoal source area of occurrence of both MIC and MAC can reach a radius of 40 km. However,
853 it is often assumed that MAC indicates fires that occurred up to 1-3 km (Clark, 1990; Higuera et al., 2007;
854 Oris et al., 2014). The distances to which particles move are also determined by terrain and vegetation. They
855 move longer distances on flat terrain covered with grasses (Woodward and Haines, 2020), while they move
856 shorter distances in dense forests (Kelly et al., 2013; Oris et al., 2014). In this context, it should be assessed

857 that the local fire activity in the studied peatland was low, with an average of 0.36 particles/cm³/year,
858 although from historical sources, fires are known to have occurred nearby.

859

860 4.3.2. Insect outbreaks and their impact on pine monoculture

861 Palaeoecological studies based on the presence of insect head capsules and/or ~~feces faeces~~, as well as
862 other insect remains could be helpful, but these methods are rarely used (Bhiry and Fillion, 1996; Lavoie et
863 al., 2009; Simard et al., 2006; Waller, 2013). Often the main obstacle to performing this method is bad
864 preservation of insect remains in peat. In the Okoniny peatland, we ~~did not find~~ found no any insect remains,
865 even though quite a large sample volume has been analyzed for the plant macrofossil analysis. Therefore,
866 we can interpret the effect of insect outbreaks using other sources of evidence.

Z komentarzem [MB68]: Editorial and linguistic corrections

867 The earliest information on insect outbreaks from the forests of the Tuchola Pinewoods under planned
868 forest management dates back to 1836-1838. A gradation An infestation of the *Panolis flammea* occurred at
869 that time (Schütte, 1893). The insects also attacked between 1866 and 1868. As a result of this gradation
870 infestation, 1380 hectares of forest were destroyed in the Woziwoda forest district alone (Schütte, 1893).
871 The pollen diagram from the Okoniny peatland documents the phenomenon in the 1860s with a decrease in
872 *Pinus sylvestris* pollen and an increased presence of *Pinus stomata* that may indicate the event of the insect
873 outbreak (Barabach, 2015). The needles that fell were partially decomposed and carried downwind to the
874 peatland, where they were preserved (Słowiński et al., 2019). The same effect was noted in another closely
875 located peatland in Okoniny Nadjeziorne, where the 1866-1868 gradation infestation also corresponds with
876 increased numbers of *Pinus stomata* (Tipton, 2023). In 1855, *Lymantria monachal* appeared in large
877 numbers but damaged only some of the younger stands (Schütte, 1893).

Z komentarzem [MB69]: Editorial and linguistic corrections

Z komentarzem [MB70]: We moved the paragraph to the beginning of the chapter, as suggested. In the "Historical and cartographic information" chapter, we added information on when the insect outbreak data originated.

Z komentarzem [MB71]: Editorial and linguistic corrections

Z komentarzem [MB72]: Editorial and linguistic corrections

878 A serious incidence of *Panolis flammea* gradation infestation also occurred in 1922-1924
879 (Kielczewski, 1947; Mokrzecki, 1928). Between 1978 and 1985, with a peak in 1982, the forests of the
880 northern part of the country were overrun by *Lymantria monacha*, and this was the largest gradation
881 infestation since the establishment of the National Forests in 1924, with salvage treatments covering more
882 than 6.3 million hectares of forest over seven years (Broda, 2000; Jabłoński, 2015; Śliwa, 1989, 1987). Both
883 major gradations infestations are reflected in palynological data, manifested by declines in the pollen
884 percentage of trees, primarily *Pinus* and *Picea*. A decrease in conifer pollen during the gradation period has
885 also been shown by studies of other sites in the Tuchola Pinewoods (Łuców et al., 2021; Tipton, 2023).
886 Other pine monoculture in Poland, the Noteć Forest was also affected by gradation in 1922-1924, and this
887 event manifested itself in palaeoecological data (Barabach, 2015; Lamentowicz et al., 2015; Milecka et al.,
888 2017). Among other things, Barabach (2015) noted an increase in *Glomeromycota* fungal spores, which
889 according to this author may indicate intense soil erosion caused by the felling of dead trees, and a marked
890 increase in *Calluna* and Poaceae indicating an increase in the openness of the landscape. Lamentowicz et

Z komentarzem [MB73]: Editorial and linguistic corrections

Z komentarzem [MB74]: Editorial and linguistic corrections

Z komentarzem [MB75]: Editorial and linguistic corrections

Z komentarzem [MB76]: Editorial and linguistic corrections

Z komentarzem [MB77]: Editorial and linguistic corrections

891 al. (2015) noted an increase in mineral content in the sediment as indicated by *Centropyxis platystoma*,
892 which was confirmed by XMT analysis of the peat. Milecka et al. (2017) described higher ash ~~content and~~
893 ~~higher~~ charcoal content in the sediments. Although the Tuchola Pinewoods and the Noteć Forest are in the
894 region of highest risk of outbreaks, other areas of Poland were also affected, such as the Kampinos Forest
895 in 1972 (Śliwa, 1974), or over the last decade, the Białowieża Primeval Forest (Grodzki, 2016; Kamińska
896 et al., 2021).

Z komentarzem [MB78]: Editorial and linguistic corrections

897 ~~Palaeoecological studies based on the presence of insect head capsules and/or feces, as well as other~~
898 ~~insect remains could be helpful, but these methods are rarely used (Bhiry and Filion, 1996; Lavoie et al.,~~
899 ~~2009; Simard et al., 2006; Waller, 2013). Often the main obstacle to performing this method is bad~~
900 ~~preservation of insect remains in peat. In the Okoniny peatland, we did not find any insect remains, even~~
901 ~~though quite a large sample volume has been analyzed for the plant macrofossil analysis. Therefore, we can~~
902 ~~interpret the effect of insect outbreaks using other sources of evidence.~~

Z komentarzem [MB79]: We moved the paragraph to the beginning of the chapter, as suggested.

903 It's difficult to assess unequivocally whether the gradations affected the immediate vicinity of the
904 peatland, or whether this is a regional signal. ~~Historical~~ ~~Historic~~ maps could be helpful, but these usually do
905 not show the difference between old and new plantings (Barabach, 2012). However, dendrochronological
906 data obtained from pine trees could help to reconstruct the extent of the outbreak. The main problem in
907 monoculture forests though is that the forest is successively cut and new trees are planted regularly.
908 However, for our dendrochronological record, we were able to obtain samples from the oldest pine trees in
909 the area. The oldest trees in the region analyzed in this study were planted over 200 years ago in the close
910 vicinity of the Woziwoda Forest District, after the introduction of the Prussian forest management strategies,
911 and have been kept there by foresters for obtaining tree saplings and for monitoring. The influence of insect
912 outbreaks has been recorded in these pine trees and we were able to track all the outbreak events in the
913 wood. The first years after the gradations - 1839, 1869, 1925, and 1982 - manifested very strongly in the
914 dendrochronological data as negative indicator years.

Z komentarzem [MB80]: Editorial and linguistic corrections

916 4.4. Current condition of the peatland vs. remote sensing and dendroclimatic data

917 The assessed growth reactions of pine trees to climate factors at the Woziwoda site may be considered
918 typical. The effect of February air temperatures on Scots pine growth in northern Poland was previously
919 noted (Cedro, 2001; Cedro and Lamentowicz, 2011; Feliksik and Wilczyński, 2009; Koprowski et al., 2012,
920 2011; Matulewski et al., 2019; Zielski, 1996; Zielski et al., 2010; Zielski and Sygit, 1998). Although the
921 studied pines from Woziwoda showed a similar growth response to climate as other pines from northern
922 Poland, their climate sensitivity was greater. The highest negative correlation for pine radial growth from
923 the Woziwoda site was found with July's mean air temperature.

924 Another factor commonly affecting the radial growth of Scots pine, according to the literature, is pluvial
925 conditions in February. This linkage was identified by Cedro (2001), Feliksik and Wileczyński (2009),
926 Koprowski et al. (2011) in the Pomeranian region (Northern Poland). The present study confirmed a short-
927 term relationship between pine radial growth and precipitation sums in February (Fig. 7). Late February and
928 early March are when additional water is required due to the initiation of biochemical processes in trees
929 (Przybylski, 1993). Additionally, in our study, a stronger dependence of pine radial growth on precipitation
930 was demonstrated in June. A similar result for pine from northern Poland was obtained by Matulewski et al.
931 (2019), Zielski and Barankiewicz (2000), where pine growth was threatened by a water deficit in the summer
932 season. Increased pine demand for water occurs in June and July, **which are** the months of the most intense
933 growth (Obmiński, 1970). At the same time, these are the months when droughts have become more frequent
934 in recent years (Łabędzki, 2004; Spinoni et al., 2018). Our results confirm that within the temperature and
935 monthly precipitation values typically observed in Central Europe, the primary environmental factor
936 influencing the diversity of species growth in the near future will be the availability of water (Boczoń et al.,
937 2017; Taeger et al., 2013). This availability is determined by **both the precipitation level of precipitation** and
938 losses caused by evapotranspiration (Boczoń and Wróbel, 2015; Zajączkowski et al., 2013).

Z komentarzem [MB81]: Editorial and linguistic corrections

Z komentarzem [MB82]: Editorial and linguistic corrections

939 The higher climatic sensitivity of pines at the Woziwoda site was manifested also by a higher number
940 of pointer years. The pointer years identified in this study are confirmed by earlier studies performed on
941 pine trees in northern Poland for **the period** 1910-2014 (Matulewski et al., 2019; Zielski et al., 1998; Zielski
942 and Barankiewicz, 2000). The years 1911, 1940, 1950, and 1982 attract particular attention. These are years
943 in which dry and hot summers were recorded (Matulewski et al., 2019; Zielski, 1996). **Moreover, the years**
944 **1925 and 1982 are marked by insect outbreaks.**

Z komentarzem [MB83]: Editorial and linguistic corrections

Z komentarzem [MB84]: Editorial and linguistic corrections

945 Our data show that *Pinus sylvestris* has been under critical climatic pressure and is responding
946 negatively to a warming climate and changing precipitation regime. Models predict a severe decline in
947 coniferous species in the next 50 years, including *Pinus sylvestris* in the temperate zone of Europe (Dyderski
948 et al., 2018; Hanewinkel et al., 2013; Schueler et al., 2014). The disappearance of species currently dominant
949 in the forests of **Central and Eastern central and eastern** Europe will result in the profound disruption or
950 disappearance of ecosystems functionally related to them, such as peatlands (Dyderski et al., 2018).

Z komentarzem [MB85]: Editorial and linguistic corrections

951 Peatlands are also affected by accelerating climate change and on top of that they are at risk of losing
952 their favourable environment, especially in *Pinus sylvestris* monoculture forests, which are particularly
953 vulnerable to increasing extreme events. Studies conducted by various researchers confirm that remote
954 sensing data, provide a valuable source of information about peatlands and help in monitoring their
955 condition (Czapiewski and Szumińska, 2021; Kaplan et al., 2019; Lees et al., 2021; Rapinel et al., 2023).
956 The analyses conducted in this study have demonstrated that multisensor airborne data can be successfully
957 utilized to assess the current state of peatlands vegetation. **The application Applying** of simple remote

Z komentarzem [MB86]: Editorial and linguistic corrections

958 sensing indices enabled the detection of spatial differences in the condition and water stress of vegetation
959 in the Okoniny peatland. According to Rastogi et al. (2019), NDVI values for peatland vegetation may
960 decrease in areas affected by stress factors such as warming and reduced precipitation. Moreover, NDVI
961 values for healthy Sphagnum moss in peatland usually range from 0.8 to 0.9 during the summer, but they
962 are also species-dependent (Harris, 2008; Letendre et al., 2008; Péli et al., 2015). Consequently, the values
963 of NDVI observed in this study (averaging 0.65) may indicate a prevailing drought situation in certain areas
964 of the Okoniny peatland. Comparable findings can be drawn from the spatial variation of MSI values
965 presented in this study. Harris et al. (2006, 2005) demonstrated that MSI is significantly correlated with
966 near-surface moisture condition of Sphagnum moss. Despite the wide application of optical data and spectral
967 indices in assessing peatland conditions, Gerhards (2018) found that spectral indices may only be useful
968 under conditions of severe or prolonged water stress. For the pre-visual detection of initial vegetation water
969 stress symptoms, temperature-based indices are most suitable, exemplified by the LST index used in this
970 study. Although aerial thermal data has been previously applied in peatland research (Kopeć et al., 2016),
971 further research into the potential use of airborne thermal data in assessing peatland vegetation conditions
972 is recommended. To date, there have been few works in Poland using spectral data in peatland monitoring
973 (Bandopadhyay et al., 2021, 2019). However, none has attempted to collate palaeoecological,
974 dendrochronological, and remote sensing data.

975

976 5. Conclusions

977 Our data show that peatlands are highly sensitive to the progressive rise in Earth's temperatures and
978 changing precipitation regimes. Groundwater levels have dropped dramatically in recent years, causing
979 intense heating of the peatland surface in summer and stressing peat-forming vegetation to water scarcity.
980 The pine monocultures surrounding the peatlands are also sensitive to climate change. They are currently
981 responding very strongly to summer precipitation deficiency, and these data fit into dendrological predictive
982 models. Planned forest management has permanently changed the composition of the forest. Deciduous tree
983 species such as *Quercus*, *Fagus*, *Carpinus*, and *Corylus avellana* have almost **completely** disappeared.
984 Forest management has also contributed **to an increase to increased** in acidity in the peatland, and thus the
985 rapid development of *Sphagnum* specialized for life in acidic conditions. After the expansion of *Sphagnum*,
986 the water level in the peatland stabilized. Peatlands are also valuable archives of past climatic anomalies
987 and catastrophic events. Pest gradations are recorded, among other things, by the presence of *Pinus* stomata,
988 and periods of drought by an increase in the **concentration values** of coprophilous fungi. These events
989 correspond with dendrochronological records. There is a strong correlation between the first years after
990 hailstorms and smaller increments of tree rings. **Our study shows that palaeoecological and**
991 **dendrochronological data the combining of different data (palaeoecological, dendrochronological, remote**

Z komentarzem [MB87]: Editorial and linguistic corrections

Z komentarzem [MB88]: Editorial and linguistic corrections

Z komentarzem [MB89]: Editorial and linguistic corrections

992 **sensing, and historical)** can complement each other and create a more complete picture of past
993 environmental changes and expand knowledge of best practices for local (Konczal et al., 2024) and global
994 (Joosten, 2021) recommendations for peatland conservation in forests. Healthy wetlands could be key to
995 protecting forests and slowing the transformation of forests caused by climate change (Marcisz et al., 2024).
996 The results are **important essential** for peatland conservation in **the context of** planned forest management.

Z komentarzem [MB90]: Editorial correction as suggested by the reviewer. We added historical and remote sensing data to the text.

Z komentarzem [MB91]: Editorial and linguistic corrections

Z komentarzem [MB92]: Editorial and linguistic corrections

998 **Competing interests**

999 The contact author has declared that none of the authors has any competing interests.

1000

1001 **Acknowledgments**

1002 The study was funded by the National Science Centre, Poland, grant 2020/39/D/ST10/00641. Remote
1003 sensing data collection and visualization were done within the scope of the project "Protection of Valuable
1004 Ecosystems of Tuchola Forest" funded by the European Economic Area Financial Mechanism 2014-2021
1005 within the framework of the Environment, Energy and Climate Change Programme MF EEA 2014-2021
1006 "Implementation of Ecosystem Management Plans".

1007 We want to thank Stefan Konczal and other foresters from the Woziwoda Forestry Unit for their cooperation
1008 and help in the field, **as well as** for providing us with historical maps and sharing knowledge of the **forest's**
1009 **history and management of the forest**. We thank Małgorzata Suchorska (Adam Mickiewicz University,
1010 Poznań) for her help in the field.

Z komentarzem [MB93]: Editorial and linguistic corrections

Z komentarzem [MB94]: Editorial and linguistic corrections

1011

1012 **Data availability**

1013 **All data associated with this article are openly available on Mendeley Data repository under the DOI:**
1014 **10.17632/prdgmjcg69.1**

Z komentarzem [MB95]: Nee section

1015

1016 **Authors contribution**

1017 MB – fieldwork, laboratory analyses (bulk density, carbon accumulation, plant macrofossils, selection of
1018 plant macrofossils for AMS radiocarbon dating), age-depth modelling, data interpretation, visualization,
1019 writing (original draft)

1020 ML – fieldwork, support in plant macrofossil analysis, data interpretation, writing (commenting and editing)

1021 PK – fieldwork, laboratory analyses (pollen and spores), age-depth modelling, data interpretation,
1022 visualization, writing (commenting and editing)

1023 DW – laboratory analyses (testate amoebae), testate amoeba-based reconstructions, data interpretation

1024 PM – fieldwork, laboratory analyses (dendrochronology), data interpretation, visualization, writing
1025 (commenting and editing)

1026 DK, MW – fieldwork, remote sensing analyses and interpretation, writing (commenting and editing)
1027 DJ – laboratory analyses (dendrochronology), data interpretation
1028 KM – funding acquisition, conceptualization, fieldwork, laboratory analyses (charcoal), testate amoeba-
1029 based reconstructions, data interpretation, visualization, writing (commenting and editing)
1030

1031 References

- 1032
1033 Adolf, C., Wunderle, S., Colombaroli, D., Weber, H., Gobet, E., Heiri, O., van Leeuwen, J.F.N., Bigler, C.,
1034 Connor, S.E., Galka, M., La Mantia, T., Makhortykh, S., Svitavská-Svobodová, H., Vannière, B.,
1035 Tinner, W., 2018. The sedimentary and remote-sensing reflection of biomass burning in Europe.
1036 *Global Ecology and Biogeography* 27, 199–212. <https://doi.org/10.1111/geb.12682>
1037 Amesbury, M.J., Swindles, G.T., Bobrov, A., Charman, D.J., Lamentowicz, M., Mallon, G., Mazei, Y.,
1038 Mitchell, E.A.D., Payne, R.J., Roland, T.P., Turner, E.T., Warner, B.G., 2016. Development of a new
1039 pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quat Sci*
1040 *Rev* 152, 132–151. <https://doi.org/10.1016/j.quascirev.2016.09.024>
1041 Anderberg, A.-L., 1994. Atlas of seeds and small fruits of Northwest-European plant species with
1042 morphological descriptions. Part 4: Resedaceae - Umbelliferae. Risbergs Tryckeri AB, Uddevalla.
1043 Baillie, M.G.L., Pilcher, J., 1973. A simple cross-dating program for tree-ring research. *Tree-Ring Bull.*
1044 33, 7–14.
1045 Ballesteros-Cánovas, J.A., Edvardsson, J., Corona, C., Mažeika, J., Stoffel, M., 2022. Estimation of recent
1046 peat accumulation with tree saplings. *Progress in Physical Geography: Earth and Environment* 46,
1047 515–529. <https://doi.org/10.1177/03091333211073786>
1048 Bandopadhyay, S., Rastogi, A., Cogliati, S., Rascher, U., Gąbka, M., Juszczak, R., 2021. Can Vegetation
1049 Indices Serve as Proxies for Potential Sun-Induced Fluorescence (SIF)? A Fuzzy Simulation
1050 Approach on Airborne Imaging Spectroscopy Data. *Remote Sens (Basel)* 13, 2545.
1051 <https://doi.org/10.3390/rs13132545>
1052 Bandopadhyay, S., Rastogi, A., Rascher, U., Rademske, P., Schickling, A., Cogliati, S., Julitta, T., Mac
1053 Arthur, A., Hueni, A., Tomelleri, E., Celesti, M., Burkart, A., Stróżecki, M., Sakowska, K., Gąbka,
1054 M., Rosadziński, S., Sojka, M., Iordache, M.-D., Reusen, I., Van Der Tol, C., Damm, A.,
1055 Schuettemeyer, D., Juszczak, R., 2019. Hyplant-Derived Sun-Induced Fluorescence—A New
1056 Opportunity to Disentangle Complex Vegetation Signals from Diverse Vegetation Types. *Remote*
1057 *Sens (Basel)* 11, 1691. <https://doi.org/10.3390/rs11141691>
1058 Barabach, J., 2015. Zapis zdarzeń katastrofalnych na obszarze Puszczy Noteckiej w osadach Torfowiska
1059 Rzecin. *Wydział Nauk Geograficznych i Geologicznych*.
1060 Barabach, J., 2012. The history of Lake Rzecin and its surroundings drawn on maps as a background to
1061 palaeoecological reconstruction. *Limnological Review* 12, 103–114. [https://doi.org/10.2478/v10194-](https://doi.org/10.2478/v10194-011-0050-0)
1062 [011-0050-0](https://doi.org/10.2478/v10194-011-0050-0)
1063 Bauhus, J., Forrester, D.I., Gardiner, B., Jactel, H., Vallejo, R., Pretsch, H., 2017. Ecological Stability of
1064 Mixed-Species Forests, in: *Mixed-Species Forests*. Springer, Berlin, Heidelberg, pp. 337–382.
1065 https://doi.org/10.1007/978-3-662-54553-9_7
1066 Beaulne, J., Boucher, É., Garneau, M., Magnan, G., 2021a. Paludification reduces black spruce growth
1067 rate but does not alter tree water use efficiency in Canadian boreal forested peatlands. *For Ecosyst* 8,
1068 28. <https://doi.org/10.1186/s40663-021-00307-x>
1069 Beaulne, J., Garneau, M., Magnan, G., Boucher, É., 2021b. Peat deposits store more carbon than trees in
1070 forested peatlands of the boreal biome. *Sci Rep* 11, 2657. [https://doi.org/10.1038/s41598-021-82004-](https://doi.org/10.1038/s41598-021-82004-x)
1071 x

- 1072 Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. Present and
 1073 future Köppen-Geiger climate classification maps at 1-km resolution. *Sci Data* 5, 180214.
 1074 <https://doi.org/10.1038/sdata.2018.214>
- 1075 Becker, M., Nieminen, T., Gérémia, F., 1994. Short-term variations and long-term changes in oak
 1076 productivity in northeastern France. The role of climate and atmospheric CO₂. *Annales des Sciences*
 1077 *Forestières* 51, 477–492. <https://doi.org/10.1051/forest:19940504>
- 1078 Berggren, G., 1969. Atlas of seeds and small fruits of Northwest-European plant species (Sweden,
 1079 Norway, Denmark, East Fennoscandia and Iceland) with morphological descriptions. Part 2:
 1080 Cyperaceae. *Berlingska Boktryckeriet, Lund*.
- 1081 Berglund, B.E., Ralska-Jasiewiczowa, M., 1986. Pollen analysis and pollen diagrams, in: Berglund, B.E.
 1082 (Ed.), *Handbook of Holocene Palaeoecology and Palaeohydrology*. John Wiley & Sons, Chichester,
 1083 pp. 455–484.
- 1084 Beug, H.-J., 2004. *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*. Verlag Dr.
 1085 Friedrich Pfeil, München.
- 1086 Bhiry, N., Filion, L., 1996. Mid-Holocene Hemlock Decline in Eastern North America Linked with
 1087 Phytophagous Insect Activity. *Quat Res* 45, 312–320. <https://doi.org/10.1006/qres.1996.0032>
- 1088 Bienias, D., 2009. *Las i człowiek w Borach Tucholskich (uwagi o bartnictwie i smolarstwie w Borach*
 1089 *Tucholskich*, in: Woźny, J. (Ed.), *Dziedzictwo Techniczne Boro' w Tucholskich. Przedsiębiorstwo*
 1090 *Marketingowe LOGO, Bydgoszcz*, pp. 43–51.
- 1091 Błaszkiwicz, M., Piotrowski, J.A., Brauer, A., Gierszewski, P., Kordowski, J., Kramkowski, M.,
 1092 Lamparski, P., Lorenz, S., Noryskiewicz, A.M., Ott, F., Słowiński, M., Tyszkowski, S., 2015.
 1093 Climatic and morphological controls on diachronous postglacial lake and river valley evolution in
 1094 the area of Last Glaciation, northern Poland. *Quat Sci Rev* 109, 13–27.
 1095 <https://doi.org/10.1016/j.quascirev.2014.11.023>
- 1096 Blockeel, T., 2010. *Straminergon stramineum*, in: Atherton, I., Bosanquet, S., Lawley, M. (Eds.), *Mosses*
 1097 *and Liverworts of Britain and Ireland a Field Guide*. British Bryological Society, Plymouth, p. 720.
- 1098 Blodau, C., 2002. Carbon cycling in peatlands – A review of processes and controls. *Environmental*
 1099 *Reviews* 10, 111–134. <https://doi.org/10.1139/a02-004>
- 1100 Boczoń, A., Kowalska, A., Gawryś, R., 2017. Glebowo-wodne uwarunkowania prowadzenia gospodarki
 1101 leśnej w perspektywie zmian klimatu. *Sylvan* 161, 763–771.
- 1102 Boczoń, A., Wróbel, M., 2015. Wpływ suszy na pobór wody przez sosnę zwyczajną (*Pinus sylvestris* L.) o
 1103 różnej pozycji w drzewostanie. *Leśne Prace Badawcze* 76, 370–376.
- 1104 Bojňanský, V., Fargašová, A., 2007. *Atlas of seeds and fruits of central and east-european flora. The*
 1105 *Carpathian Mountains Region*. Springer, Dordrecht.
- 1106 Booth, R.K., 2002. Testate amoebae as paleoindicators of surface-moisture changes on Michigan
 1107 peatlands: modern ecology and hydrological calibration. *J Paleolimnol* 28, 329–348.
- 1108 Booth, R.K., Lamentowicz, M., Charman, D.J., 2010. Preparation and analysis of testate amoebae in
 1109 peatland paleoenvironmental studies. *Mires and Peat* 7 (2010/11), 1–7.
- 1110 Booth, T.H., 2013. Eucalypt plantations and climate change. *For Ecol Manage* 301, 28–34.
 1111 <https://doi.org/10.1016/j.foreco.2012.04.004>
- 1112 Boule'h, P.-N., Caullireau, E., Faucher, E., Gouerou, M., Guérin, A., Miray, R., Couée, I., 2020. Abiotic
 1113 stress signalling in extremophile land plants. *J Exp Bot* 71, 5771–5785.
 1114 <https://doi.org/10.1093/jxb/eraa336>
- 1115 Broda, J., 2000. *Historia leśnictwa w Polsce*. Wydawnictwo Akademii Rolniczej im. Augusta
 1116 Cieszkowskiego w Poznaniu, Poznań.
- 1117 Broda, J., 1993. Sosna w czasach historycznych, in: Białobok, S., Boratyński, A., Bugała, W. (Eds.),
 1118 *Biologia Sosny Zwyczajnej*. Instytut Dendrologii PAN, Poznań-Kórnik, pp. 17–31.
- 1119 Bronk Ramsey, C., 2021. OxCal v4.4.4 [WWW Document]. URL <https://c14.arch.ox.ac.uk/oxcal.html>
 1120 (accessed 11.21.23).
- 1121 Bronk Ramsey, C., 2008. Deposition models for chronological records. *Quat Sci Rev* 27, 42–60.

Z komentarzem [MB96]: We added citations.

1122 Bunn, A.G., 2008. A dendrochronology program library in R (dplR). *Dendrochronologia* (Verona) 26,
1123 115–124. <https://doi.org/10.1016/j.dendro.2008.01.002>

1124 Cedro, A., 2001. Próba oceny oddziaływania temperatury powietrza i opadów atmosferycznych na
1125 przyrost radialny sosny zwyczajnej (<s>Pinus sylvestris</s>) na Pomorzu Zachodnim. *Annales*
1126 *Universitatis Mariae Curie-Skłodowska. Sectio B, Geographia, Geologia, Mineralogia et*
1127 *Petrographia* 55/56, 105–112.

1128 Cedro, A., Lamentowicz, M., 2011. Contrasting responses to environmental changes by pine (*Pinus*
1129 *syvestris* L.) growing on peat and mineral soil: An example from a Polish Baltic bog.
1130 *Dendrochronologia* (Verona) 29, 211–217. <https://doi.org/10.1016/j.dendro.2010.12.004>

1131 Chambers, F.M., Beilman, D.W., Yu, Z., 2010. Methods for determining peat humification and for
1132 quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate and
1133 peatland carbon dynamics. *Mires and Peat* 7, 1–10.

1134 Chapin, F.S., Matson, P.A., Vitousek, P., 2012. Managing and Sustaining Ecosystems, in: Chapin, F.S.
1135 (Ed.), *Principles of Terrestrial Ecosystem Ecology*. Springer, p. 447.

1136 Cherek, E., 2007. Ochotnicza Straż Pożarna w Kasparusie 1932-2007. *Kasparus*.

1137 Clark, J.S., 1990. Fire and climate change during the last 750 yr in northwestern Minnesota. *Ecol.*
1138 *Monogr.* 60, 135–159. <https://doi.org/10.2307/1943042>

1139 Clark, J.S., 1988. Particle Motion and the Theory of Charcoal Analysis: Source Area, Transport,
1140 Deposition, and Sampling. *Quat Res* 30, 67–80. [https://doi.org/10.1016/0033-5894\(88\)90088-9](https://doi.org/10.1016/0033-5894(88)90088-9)

1141 Clarke, K.J., 2003. Guide to Identification of Soil Protozoa - Testate Amoebae, Soil Biodiversity
1142 Programme Research Report No. 4. Freshwater Biological Association, Ambleside, U.K.

1143 Clymo, R.S., Hayward, P.M., 1982. The Ecology of Sphagnum, in: Smith, A.J.E. (Ed.), *Bryophyte*
1144 *Ecology*. Chapman & Hall, London, New York, pp. 229–289.

1145 Cook, E.R., Briffa, K., Shiyatov, S., Mazepa, A., Jones, P.D., 1990. Data analysis, in: Cook, E.R.,
1146 Kairiukstis, L.A. (Eds.), *Methods of Dendrochronology: Applications in the Environmental Sciences*.
1147 Kluwer Academic Publ., Dordrecht, pp. 97–162.

1148 Cyzman, W., 2008. Jednolity Program Gospodarczo-Ochronny dla Leśnego Kompleksu Promocyjnego
1149 „Bory Tucholskie”.

1150 Czapiewski, S., Szumińska, D., 2021. An Overview of Remote Sensing Data Applications in Peatland
1151 Research Based on Works from the Period 2010–2021. *Land* (Basel) 11, 24.
1152 <https://doi.org/10.3390/land11010024>

1153 Czerwiński, S., Guzowski, P., Lamentowicz, M., Gałka, M., Karpińska-Kończak, M., Poniak, R., Łokas,
1154 E., Diaconu, A.-C., Schwarzer, J., Miecznik, M., Kończak, P., 2021. Environmental implications of
1155 past socioeconomic events in Greater Poland during the last 1200 years. Synthesis of paleoecological
1156 and historical data. *Quat Sci Rev* 259, 106902.
1157 <https://doi.org/https://doi.org/10.1016/j.quascirev.2021.106902>

1158 Davis, M.B., Deevey, E.S., 1964. Pollen Accumulation Rates: Estimates from Late-Glacial Sediment of
1159 Rogers Lake. *Science* (1979) 145, 1293–1295. <https://doi.org/10.1126/science.145.3638.1293>

1160 Diaconu, A.-C., Tóth, M., Lamentowicz, M., Heiri, O., Kuske, E., Tanțău, I., Panait, A.-M., Braun, M.,
1161 Feurdean, A., 2017. How warm? How wet? Hydroclimate reconstruction of the past 7500 years in
1162 northern Carpathians, Romania. *Palaeogeogr Palaeoclimatol Palaeoecol* 482, 1–12.
1163 <https://doi.org/10.1016/j.palaeo.2017.05.007>

1164 Dietze, E., Brykała, D., Schreuder, L.T., Jażdżewski, K., Blarquez, O., Brauer, A., Dietze, M., Obremska,
1165 M., Ott, F., Pieńczewska, A., Schouten, S., Hopmans, E.C., Słowiński, M., 2019. Human-induced
1166 fire regime shifts during 19th century industrialization: A robust fire regime reconstruction using
1167 northern Polish lake sediments. *PLoS One* 14, e0222011.
1168 <https://doi.org/10.1371/journal.pone.0222011>

1169 Dinella, A., Giammarchi, F., Prendin, A.L., Carrer, M., Tonon, G., 2021. Xylem traits of peatland Scots
1170 pines reveal a complex climatic signal: A study in the Eastern Italian Alps. *Dendrochronologia*
1171 (Verona) 67, 125824. <https://doi.org/10.1016/j.dendro.2021.125824>

1172 Drinan, T.J., Graham, C.T., O'Halloran, J., Harrison, S.S.C., 2013. The impact of catchment conifer
1173 plantation forestry on the hydrochemistry of peatland lakes. *Science of The Total Environment* 443,
1174 608–620. <https://doi.org/10.1016/j.scitotenv.2012.10.112>

1175 Dyderski, M.K., Paż, S., Frelich, L.E., Jagodziński, A.M., 2018. How much does climate change threaten
1176 European forest tree species distributions? *Glob Chang Biol* 24, 1150–1163.
1177 <https://doi.org/10.1111/gcb.13925>

1178 Eckstein, D., Bauch, J., 1969. Beitrag zur Rationalisierung eines dendrochronologischen Verfahrens und
1179 zur Analyse seiner Aussagesicherheit. *Forstwissenschaftliches Centralblatt* 88, 230–250.
1180 <https://doi.org/10.1007/BF02741777>

1181 Edvardsson, J., Baužienė, I., Lamentowicz, M., Šimanasienė, R., Tamkevičiūtė, M., Taminskas, J.,
1182 Linkevičienė, R., Skuratovič, Ž., Corona, C., Stoffel, M., 2019. A multi-proxy reconstruction of
1183 moisture dynamics in a peatland ecosystem: A case study from Čepkeliai, Lithuania. *Ecol Indic* 106,
1184 105484. <https://doi.org/https://doi.org/10.1016/j.ecolind.2019.105484>

1185 Edvardsson, J., Corona, C., Mažeika, J., Pukienė, R., Stoffel, M., 2016. Recent advances in long-term
1186 climate and moisture reconstructions from the Baltic region: Exploring the potential for a new multi-
1187 millennial tree-ring chronology. *Quat Sci Rev* 131, Part A, 118–126.
1188 <https://doi.org/https://doi.org/10.1016/j.quascirev.2015.11.005>

1189 Edvardsson, J., Helama, S., Rundgren, M., Nielsen, A.B., 2022. The Integrated Use of
1190 Dendrochronological Data and Paleoecological Records From Northwest European Peatlands and
1191 Lakes for Understanding Long-Term Ecological and Climatic Changes—A Review. *Front Ecol Evol*
1192 10. <https://doi.org/10.3389/fevo.2022.781882>

1193 FAO, 2020. Peatlands mapping and monitoring. Recommendations and technical overview. Rome.
1194 <https://doi.org/10.4060/ca8200en>

1195 Feliksik, E., Wilczyński, S., 2009. The Effect of Climate on Tree-Ring Chronologies of Native and
1196 Nonnative Tree Species Growing Under Homogenous Site Conditions. *Geochronometria* 33, 49–57.
1197 <https://doi.org/10.2478/v10003-009-0006-4>

1198 Felton, A., Gustafsson, L., Roberge, J.-M., Ranius, T., Hjältén, J., Rudolphi, J., Lindblad, M., Weslien, J.,
1199 Rist, L., Brunet, J., Felton, A.M., 2016. How climate change adaptation and mitigation strategies can
1200 threaten or enhance the biodiversity of production forests: Insights from Sweden. *Biol Conserv* 194,
1201 11–20. <https://doi.org/10.1016/j.biocon.2015.11.030>

1202 Finsinger, W., Tinner, W., 2005. Minimum count sums for charcoal-concentration estimates in pollen
1203 slides: accuracy and potential errors. *Holocene* 15, 293–297.

1204 Freeman, C., Fenner, N., Ostle, N., Kang, H., Dorwick, D.J., Reynolds, B., Lock, M.A., Sleep, D.,
1205 Hughes, S., Hudson, J., 2004. Export of dissolved organic carbon from peatlands under elevated
1206 carbon dioxide levels. *Nature* 430, 195–198.

1207 Gałka, M., Knorr, K.-H., Tobolski, K., Gallego-Sala, A., Kołaczek, P., Lamentowicz, M., Kajukalo-
1208 Drygalska, K., Marcisz, K., 2022. How far from a pristine state are the peatlands in the Białowieża
1209 Primeval Forest (CE Europe) – Palaeoecological insights on peatland and forest development from
1210 multi-proxy studies. *Ecol Indic* 143, 109421. <https://doi.org/10.1016/j.ecolind.2022.109421>

1211 Gałka, M., Miotk-Szpiganowicz, G., Marczevska, M., Barabach, J., van der Knaap, W.O., Lamentowicz,
1212 M., 2015. Palaeoenvironmental changes in Central Europe (NE Poland) during the last 6200 years
1213 reconstructed from a high-resolution multi-proxy peat archive. *Holocene* 25, 421–434.
1214 <https://doi.org/10.1177/0959683614561887>

1215 Gałka, M., Tobolski, K., Górska, A., Milecka, K., Fiałkiewicz-Kozielec, B., Lamentowicz, M., 2014.
1216 Disentangling the drivers for the development of a Baltic bog during the Little Ice Age in northern
1217 Poland. *Quaternary International* 328–329, 323–337.
1218 <https://doi.org/http://dx.doi.org/10.1016/j.quaint.2013.02.026>

1219 Gallego-Sala, A. V., Charman, D.J., Brewer, S., Page, S.E., Prentice, I.C., Friedlingstein, P., Moreton, S.,
1220 Amesbury, M.J., Beilman, D.W., Björck, S., Blyakharchuk, T., Bochicchio, C., Booth, R.K.,
1221 Bunbury, J., Camill, P., Carless, D., Chimner, R.A., Clifford, M., Cressey, E., Courtney-Mustaphi, C.,
1222 De Vleeschouwer, F., de Jong, R., Fiałkiewicz-Kozielec, B., Finkelstein, S.A., Garneau, M., Githumbi,

1223 E., Hribljan, J., Holmquist, J., Hughes, P.D.M., Jones, C., Jones, M.C., Karofeld, E., Klein, E.S.,
 1224 Kokfelt, U., Korhola, A., Lacourse, T., Le Roux, G., Lamentowicz, M., Large, D., Lavoie, M.,
 1225 Loisel, J., Mackay, H., MacDonald, G.M., Makila, M., Magnan, G., Marchant, R., Marcisz, K.,
 1226 Martínez Cortizas, A., Massa, C., Mathijssen, P., Mauquoy, D., Mighall, T., Mitchell, F.J.G., Moss,
 1227 P., Nichols, J., Oksanen, P.O., Orme, L., Packalen, M.S., Robinson, S., Roland, T.P., Sanderson,
 1228 N.K., Sannel, A.B.K., Silva-Sánchez, N., Steinberg, N., Swindles, G.T., Turner, T.E., Uglow, J.,
 1229 Välranta, M., van Bellen, S., van der Linden, M., van Geel, B., Wang, G., Yu, Z., Zaragoza-Castells,
 1230 J., Zhao, Y., 2018. Latitudinal limits to the predicted increase of the peatland carbon sink with
 1231 warming. *Nat Clim Chang* 8, 907–913. <https://doi.org/https://www.nature.com/articles/s41558-018-0271-1>
 1232
 1233 Gerhards, M., 2018. Advanced Thermal Remote Sensing for Water Stress Detection of Agricultural Crops.
 1234 Godwin, H., 1981. *Archives of the Peat Bogs*. Cambridge University Press, Cambridge.
 1235 González de Andrés, E., Shestakova, T.A., Scholten, R.C., Delcourt, C.J.F., Gorina, N. V., Camarero, J.J.,
 1236 2022. Changes in tree growth synchrony and resilience in Siberian *Pinus sylvestris* forests are
 1237 modulated by fire dynamics and ecohydrological conditions. *Agric For Meteorol* 312, 108712.
 1238 <https://doi.org/https://doi.org/10.1016/j.agrformet.2021.108712>
 1239 Gorham, E., 1991. Northern Peatlands: Role in the Carbon Cycle and Probable Responses to Climatic
 1240 Warming. *Ecological Applications* 1, 182–195. <https://doi.org/10.2307/1941811>
 1241 Gregow, H., Laaksonen, A., Alper, M.E., 2017. Increasing large scale windstorm damage in Western,
 1242 Central and Northern European forests, 1951–2010. *Sci Rep* 7, 46397.
 1243 <https://doi.org/10.1038/srep46397>
 1244 Grimm, E.C., 1992. *Tilia and Tilia-Graph. Pollen Spreadsheet and Graphics Programs*. 8th International
 1245 Palynological Congress (President: A. Pons) Aix-en-Provence, September 6-12, 1992. Program and
 1246 Abstracts, p. 56.
 1247 Grimm, E.C., 1991. *Tilia and Tilia Graph*. Illinois State Museum.
 1248 Grissino-Mayer, H.D., 2001. Evaluating crossdating accuracy: A manual and tutorial for the computer
 1249 program COFECHA. *Tree Ring Res* 57, 205–221.
 1250 Grodzki, W., 2016. Mass outbreaks of the spruce bark beetle *Ips typographus* in the context of the
 1251 controversies around the Białowieża Primeval Forest. *Forest Research Papers* 77, 324–331.
 1252 <https://doi.org/10.1515/frp-2016-0033>
 1253 Grondin, P., Gauthier, S., Borcard, D., Bergeron, Y., Noël, J., 2014. A new approach to ecological land
 1254 classification for the Canadian boreal forest that integrates disturbances. *Landsc Ecol* 29, 1–16.
 1255 <https://doi.org/10.1007/s10980-013-9961-2>
 1256 Guariguata, M.R., Cornelius, J.P., Locatelli, B., Forner, C., Sánchez-Azofeifa, G.A., 2008. Mitigation
 1257 needs adaptation: Tropical forestry and climate change. *Mitig Adapt Strateg Glob Chang* 13, 793–
 1258 808. <https://doi.org/10.1007/s11027-007-9141-2>
 1259 Guiot, J., 1991. The bootstrapped response function. *Tree-Ring Bulletin* 51, 39–41.
 1260 Guo, M., Li, J., Sheng, C., Xu, J., Wu, L., 2017. A Review of Wetland Remote Sensing. *Sensors* 17, 777.
 1261 <https://doi.org/10.3390/s17040777>
 1262 Hanewinkel, M., Cullmann, D.A., Schelhaas, M.-J., Nabuurs, G.-J., Zimmermann, N.E., 2013. Climate
 1263 change may cause severe loss in the economic value of European forest land. *Nat Clim Chang* 3,
 1264 203–207. <https://doi.org/10.1038/nclimate1687>
 1265 Hanson, P.J., Weltzin, J.F., 2000. Drought disturbance from climate change: response of United States
 1266 forests. *Science of The Total Environment* 262, 205–220. [https://doi.org/10.1016/S0048-9697\(00\)00523-4](https://doi.org/10.1016/S0048-9697(00)00523-4)
 1267
 1268 Harenda, K.M., Lamentowicz, M., Samson, M., Chojnicki, B.H., 2018. The Role of Peatlands and Their
 1269 Carbon Storage Function in the Context of Climate Change. pp. 169–187.
 1270 https://doi.org/10.1007/978-3-319-71788-3_12
 1271 Harriman, R., Watt, A.W., Christie, A.E.G., Moore, D.W., McCartney, A.G., Taylor, E.M., 2003.
 1272 Quantifying the effects of forestry practices on the recovery of upland streams and lochs from

1273 acidification. *Science of The Total Environment* 310, 101–111. <https://doi.org/10.1016/S0048->
1274 9697(02)00626-5

1275 Harris, A., 2008. Spectral reflectance and photosynthetic properties of *Sphagnum* mosses exposed to
1276 progressive drought. *Ecohydrology* 1, 35–42. <https://doi.org/10.1002/eco.5>

1277 Harris, A., Bryant, R., Baird, A., 2005. Detecting near-surface moisture stress in spp. *Remote Sens*
1278 *Environ* 97, 371–381. <https://doi.org/10.1016/j.rse.2005.05.001>

1279 Harris, A., Bryant, R.G., Baird, A.J., 2006. Mapping the effects of water stress on *Sphagnum*: Preliminary
1280 observations using airborne remote sensing. *Remote Sens Environ* 100, 363–378.
1281 <https://doi.org/10.1016/j.rse.2005.10.024>

1282 Haylock, M.R., Hofstra, N., Klein Tank, A.M.G., Klok, E.J., Jones, P.D., New, M., 2008. A European daily
1283 high-resolution gridded data set of surface temperature and precipitation for 1950–2006. *Journal of*
1284 *Geophysical Research: Atmospheres* 113. <https://doi.org/10.1029/2008JD010201>

1285 Hedenäs, L., 1993. A generic revision of the *Warnstorfia-Calliergon* group. *J Bryol* 17, 447–479.
1286 <https://doi.org/10.1179/jbr.1993.17.3.447>

1287 Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and
1288 carbonate content in sediments: Reproducibility and comparability of results. *J Paleolimnol* 25, 101–
1289 110. <https://doi.org/10.1023/A:1008119611481>

1290 Higuera, P., Peters, M., Brubaker, L., Gavin, D., 2007. Understanding the origin and analysis of sediment-
1291 charcoal records with a simulation model. *Quat Sci Rev* 26, 1790–1809.
1292 <https://doi.org/10.1016/j.quascirev.2007.03.010>

1293 Hill, M.O., Blockeel, T.L., 2014. *Straminergon stramineum*, in: Blockeel, T.L., Bosanquet, S.D.S., Hill,
1294 M.O., Preston, C.D. (Eds.), *Atlas of British and Irish Bryophytes*. British Bryological Society,
1295 Newbury, Berkshire, p. 464.

1296 Hua, Q., Turnbull, J.C., Santos, G.M., Rakowski, A.Z., Ancapichún, S., De Pol-Holz, R., Hammer, S.,
1297 Lehman, S.J., Levin, I., Miller, J.B., Palmer, J.G., Turney, C.S.M., 2021. ATMOSPHERIC
1298 RADIOCARBON FOR THE PERIOD 1950–2019. *Radiocarbon* 1–23.
1299 <https://doi.org/10.1017/RDC.2021.95>

1300 Hunt, E., Rock, B., 1989. Detection of changes in leaf water content using Near- and Middle-Infrared
1301 reflectances. *Remote Sens Environ* 30, 43–54. [https://doi.org/10.1016/0034-4257\(89\)90046-1](https://doi.org/10.1016/0034-4257(89)90046-1)

1302 Huuskonen, S., Domisch, T., Finér, L., Hantula, J., Hynynen, J., Matala, J., Miina, J., Neuvonen, S.,
1303 Nevalainen, S., Niemistö, P., Nikula, A., Piri, T., Siitonen, J., Smolander, A., Tonteri, T., Uotila, K.,
1304 Viiri, H., 2021. What is the potential for replacing monocultures with mixed-species stands to
1305 enhance ecosystem services in boreal forests in Fennoscandia? *For Ecol Manage* 479, 118558.
1306 <https://doi.org/10.1016/j.foreco.2020.118558>

1307 Jabłoński, T., 2015. Występowanie i zwalczanie leśnych foliofagów - trendy i prognozy. *Postępy Techniki*
1308 *w Leśnictwie* 132, 13–19.

1309 Jäger, E., 1982. Prussia-Karten 1542-1810: Geschichte der kartographischen Darstellung Ostpreussens
1310 vom 16. bis zum 19. Jahrhundert. Entstehung der Karten - Kosten - Vertrieb: bibliographischer
1311 Katalog. Anton H. Konrad Verlag, Weißenhorn.

1312 Jäger, E., 1981. Die Schroettersche Landesaufnahme von Ost- und Westpreußen (1796-1802).
1313 Entstehungsgeschichte, Herstellung und Vertrieb der Karte. *Z Ostforsch* 30, 359–389.

1314 Jasnowski, M., 1962. Budowa i roślinność torfowisk Pomorza Szczecińskiego. *Societas Scientiarum*
1315 *Stetinensis, Szczecin*.

1316 Jaszczak, R., 2008a. Urządzenie lasu w Polsce do 1939 roku. Część I — początki urządzania lasu na
1317 ziemiach polskich. *Sylvan* 152, 13–21. <https://doi.org/https://10.26202/sylvan.2006126>

1318 Jaszczak, R., 2008b. Urządzenie lasu w Polsce do 1939 roku. Część III — urządzenie lasu na ziemiach
1319 polskich w zaborze austriackim i pruskim. *Sylvan* 152, 3–10.
1320 <https://doi.org/https://10.26202/sylvan.2006180>

1321 Joosten, H., 2021. Global guidelines for peatland rewetting and restoration. Ramsar Technical Report No.
1322 11. Gland, Switzerland.

Z komentarzem [MB97]: We moved it to Supplementary File 1.

1323 Joosten, H., Tapio-Biström, M.-L., Tol, S., 2012. Peatlands - guidance for climate change mitigation
1324 through conservation, rehabilitation and sustainable use, 2nd ed. Food and Agriculture Organization
1325 of the United Nations, Rome.

1326 Juggins, S., 2023. Rioja: Analysis of Quaternary Science Data [WWW Document]. URL [https://cran.r-](https://cran.r-project.org/web/packages/rioja/index.html)
1327 [project.org/web/packages/rioja/index.html](https://cran.r-project.org/web/packages/rioja/index.html) (accessed 12.4.23).

1328 Juggins, S., 2007. C2 Version 1.5 User guide. Software for ecological and palaeoecological data analysis
1329 and visualisation. Newcastle University, Newcastle upon Tyne, UK, p. 73.

1330 Kamińska, A., Lisiewicz, M., Kraszewski, B., Stereńczak, K., 2021. Mass outbreaks and factors related to
1331 the spatial dynamics of spruce bark beetle (*Ips typographus*) dieback considering diverse
1332 management regimes in the Białowieża forest. *For Ecol Manage* 498, 119530.
1333 <https://doi.org/10.1016/j.foreco.2021.119530>

1334 Kaplan, G., Yigit Avdan, Z., Avdan, U., 2019. Mapping and Monitoring Wetland Dynamics Using
1335 Thermal, Optical, and SAR Remote Sensing Data, in: *Wetlands Management - Assessing Risk and*
1336 *Sustainable Solutions*. IntechOpen. <https://doi.org/10.5772/intechopen.80264>

1337 Karpińska-Kołaczek, M., Kołaczek, P., Marcisz, K., Gałka, M., Kajukało-Drygalska, K., Mauquoy, D.,
1338 Lamentowicz, M., 2024. Kettle-hole peatlands as carbon hot spots: Unveiling controls of carbon
1339 accumulation rates during the last two millennia. *Catena (Amst)* 237, 107764.
1340 <https://doi.org/10.1016/j.catena.2023.107764>

1341 Kelly, R., Chipman, M.L., Higuera, P.E., Stefanova, I., Brubaker, L.B., Hu, F.S., 2013. Recent burning of
1342 boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National*
1343 *Academy of Sciences* 110, 13055–13060. <https://doi.org/10.1073/pnas.1305069110>

1344 Kielczewski, B., 1947. Klęska sówki chojnówki jako zagadnienie biocenotyczne. *Prace Komisji*
1345 *Matematyczno-Przyrodniczej* 10, 167–171.

1346 Kirschenstein, M., 2005. Wieloletnie zmiany sum opadów atmosferycznych na wybranych stacjach
1347 północno-zachodniej Polski. *Śląskie Prace Geograficzne* 2, 199–214.

1348 Klein Tank, A.M.G., Wijngaard, J.B., Können, G.P., Böhm, R., Demarée, G., Gocheva, A., Mileta, M.,
1349 Pashiardis, S., Hejkrlik, L., Kern-Hansen, C., Heino, R., Bessemoulin, P., Müller-Westermeier, G.,
1350 Tzanakou, M., Szalai, S., Páldóttir, T., Fitzgerald, D., Rubin, S., Capaldo, M., Maugeri, M., Leitass,
1351 A., Bukantis, A., Aberfeld, R., van Engelen, A.F. V., Forland, E., Miletus, M., Coelho, F., Mares, C.,
1352 Razuvaev, V., Nieplova, E., Cegnar, T., Antonio López, J., Dahlström, B., Moberg, A., Kirchhofer,
1353 W., Ceylan, A., Pachaliuk, O., Alexander, L. V., Petrovic, P., 2002. Daily dataset of 20th-century
1354 surface air temperature and precipitation series for the European Climate Assessment. *International*
1355 *Journal of Climatology* 22, 1441–1453. <https://doi.org/10.1002/joc.773>

1356 Koenig, I., Mulot, M., Mitchell, E.A.D., 2018. Taxonomic and functional traits responses of Sphagnum
1357 peatland testate amoebae to experimentally manipulated water table. *Ecol Indic* 85, 342–351.
1358 <https://doi.org/https://doi.org/10.1016/j.ecolind.2017.10.017>

1359 Kołaczek, P., Karpińska-Kołaczek, M., Marcisz, K., Gałka, M., Lamentowicz, M., 2018. Palaeohydrology
1360 and the human impact on one of the largest raised bogs complex in the Western Carpathians (Central
1361 Europe) during the last two millennia. *Holocene* 28, 595–608.
1362 <https://doi.org/10.1177/0959683617735587>

1363 Konczal, S., Lamentowicz M, Bąk, M., Czerwiński, S., Kołaczek, P., Wochal, D., Marcisz, M., Chojnicki,
1364 B., Harenda, K., Poczta, P., Gąbka, M., Jaster, D., Matulewski, P., Jedliński, J., Niedzielko, J.,
1365 Wylązłowska, J., Żmuda, M., Żmuda, D., Kopec, D., Rosadziński, S., Wietecha, M., Landowska, J.,
1366 Landowski, J., 2024. Rekomendacje dla ochrony mokradeł w lasach, in: Lamentowicz, M., Konczal,
1367 S. (Eds.), *Jak Chronić Torfowiska w Lasach?* ArchaeGraph, Łódź, pp. 161–165.

1368 Kondracki, J., 2001. *Geografia regionalna Polski*. Wydawnictwo Naukowe PWN, Warszawa.

1369 Kopec, D., Michalska-Hejduk, D., Ślawik, Ł., Berezowski, T., Borowski, M., Rosadziński, S.,
1370 Chormański, J., 2016. Application of multisensoral remote sensing data in the mapping of alkaline
1371 fens Natura 2000 habitat. *Ecol Indic* 70, 196–208. <https://doi.org/10.1016/j.ecolind.2016.06.001>

- 1372 Koprowski, M., Przybylak, R., Zielski, A., Pospieszynska, A., 2012. Tree rings of Scots pine (*Pinus*
1373 *sylvestris* L.) as a source of information about past climate in northern Poland. *Int J Biometeorol* 56,
1374 1–10. <https://doi.org/10.1007/s00484-010-0390-5>
- 1375 Koprowski, M., Zielski, A., Skowronek, T., 2011. Analiza przyrostów rocznych dwóch sosen (*Pinus*
1376 *sylvestris*) o nietypowej budowie strzały na terenie Nadleśnictwa Borne Sulinowo. *Sylwan* 155, 555–
1377 562.
- 1378 Kowalewski, G., 2003. Shoreline changes of basins in the mire-lake reserves in s Tuchola Pinewoods.
1379 *Limnological Review* 3, 119–126.
- 1380 Kowalewski, G., Milecka, K., 2003. Palaeoecology of basins of organic sediment accumulation in the
1381 Reserve Dury. *Studia Quaternaria* 20, 73–82.
- 1382 ~~Kozikowski, A., 1911. Historia lasów Prus królewskich w świetle prawdy. *Sylwan* XXIX, 420–429.~~
- 1383 Kuosmanen, N., Čada, V., Halsall, K., Chiverrell, R.C., Schafstall, N., Kuneš, P., Boyle, J.F., Knížek, M.,
1384 Appleby, P.G., Svoboda, M., Clear, J.L., 2020. Integration of dendrochronological and
1385 palaeoecological disturbance reconstructions in temperate mountain forests. *For Ecol Manage* 475,
1386 118413. <https://doi.org/10.1016/j.foreco.2020.118413>
- 1387 Łabędzki, L., 2004. Problematyka susz w Polsce. *Woda-Środowisko-Obszary Wiejskie* 4, 47–66.
- 1388 Laine, J., Flatberg, K.I., Harju, P., Timonen, T., Minkinen, K., Laine, A., Tuittila, E.-S., Vasander, H.,
1389 2018. Sphagnum mosses. The Stars of European Mires . Department of Forest Sciences, University
1390 of Helsinki, Sphagna Ky, Helsinki.
- 1391 Laine, J., Vasander, H., Laiho, R., 1995. Long-Term Effects of Water Level Drawdown on the Vegetation
1392 of Drained Pine Mires in Southern Finland. *J Appl Ecol* 32, 785–802.
1393 <https://doi.org/10.2307/2404818>
- 1394 Lamentowicz, Ł., Gąbka, M., Rusińska, A., Sobczyński, T., Owsiany, P.M., Lamentowicz, M., 2011.
1395 Testate amoeba (Arcellinida, Euglyphida) ecology along a poor-rich gradient in fens of western
1396 Poland. *Int Rev Hydrobiol* 96, 256–380.
- 1397 Lamentowicz, M., Balwierz, Z., Forysiak, J., Płóciennik, M., Kittel, P., Kloss, M., Twardy, J., Żurek, S.,
1398 Pawlyta, J., 2009a. Multiproxy study of anthropogenic and climatic changes in the last two millennia
1399 from a small mire in central Poland. *Hydrobiologia* 631, 213–230. [https://doi.org/10.1007/s10750-](https://doi.org/10.1007/s10750-009-9812-y)
1400 [009-9812-y](https://doi.org/10.1007/s10750-009-9812-y)
- 1401 Lamentowicz, M., Cedro, A., Gałka, M., Goslar, T., Miotk-Szpiganowicz, G., Mitchell, E.A.D., Pawlyta,
1402 J., 2008a. Last millennium palaeoenvironmental changes from a Baltic bog (Poland) inferred from
1403 stable isotopes, pollen, plant macrofossils and testate amoebae. *Palaeogeogr Palaeoclimatol*
1404 *Palaeoecol* 265, 93–106.
- 1405 Lamentowicz, M., Gałka, M., Lamentowicz, Ł., Obremska, M., Kühl, N., Lücke, A., Jassey, V.E.J., 2015.
1406 Reconstructing climate change and ombrotrophic bog development during the last 4000years in
1407 northern Poland using biotic proxies, stable isotopes and trait-based approach. *Palaeogeogr*
1408 *Palaeoclimatol Palaeoecol* 418, 261–277. <https://doi.org/10.1016/j.palaeo.2014.11.015>
- 1409 Lamentowicz, M., Marcisz, K., Guzowski, P., Gałka, M., Diaconu, A.-C., Kołaczek, P., 2020. How
1410 Joannites' economy eradicated primeval forest and created anthroecosystems in medieval Central
1411 Europe. *Sci Rep* 10, 18775. <https://doi.org/10.1038/s41598-020-75692-4>
- 1412 Lamentowicz, M., Milecka, K., Gałka, M., Cedro, A., Pawlyta, J., Piotrowska, N., Lamentowicz, Ł., van
1413 der Knaap, W.O., 2009b. Climate and human induced hydrological change since AD 800 in an
1414 ombrotrophic mire in Pomerania (N Poland) tracked by testate amoebae, macro-fossils, pollen tree-
1415 rings of pine. *Boreas, An international journal of Quaternary research* 38, 214–229.
- 1416 Lamentowicz, M., Mitchell, E.A.D., 2005a. Testate amoebae (Protists) as palaeoenvironmental indicators
1417 in peatlands. *Polish Geological Institute Special Papers* 16, 58–64.
- 1418 Lamentowicz, M., Mitchell, E.A.D., 2005b. The ecology of testate amoebae (Protists) in Sphagnum in
1419 north-western Poland in relation to peatland ecology. *Microb Ecol* 50, 48–63.
- 1420 Lamentowicz, M., Mueller, M., Gałka, M., Barabach, J., Milecka, K., Goslar, T., Binkowski, M., 2015.
1421 Reconstructing human impact on peatland development during the past 200 years in CE

Z komentarzem [MB98]: We moved it to Supplementary File 1.

1422 Europethrough biotic proxies and X-ray tomography. *Quaternary International* 357, 282–294.
1423 <https://doi.org/http://dx.doi.org/10.1016/j.quaint.2014.07.045>

1424 Lamentowicz, M., Obremska, M., 2010. A rapid response of testate amoebae and vegetation to inundation
1425 of a kettle hole mire. *J Paleolimnol* 43, 499–511. <https://doi.org/10.1007/s10933-009-9347-2>

1426 Lamentowicz, M., Obremska, M., Mitchell, E.A.D., 2008b. Autogenic succession, land-use change, and
1427 climatic influences on the Holocene development of a kettle hole mire in Northern Poland. *Rev*
1428 *Palaeobot Palynol* 151, 21–40. <https://doi.org/10.1016/j.revpalbo.2008.01.009>

1429 Lamentowicz, M., Tobolski, K., Mitchell, E.A.D., 2007. Palaeoecological evidence for anthropogenic
1430 acidification of a kettle-hole peatland in northern Poland. *Holocene* 17, 1185–1196.

1431 Lavoie, M., Fillion, L., Robert, É.C., 2009. Boreal peatland margins as repository sites of long-term natural
1432 disturbances of balsam fir/spruce forests. *Quat Res* 71, 295–306.
1433 <https://doi.org/10.1016/j.yqres.2009.01.005>

1434 Lee, D., Holmström, E., Hynynen, J., Nilsson, U., Korhonen, K.T., Westerlund, B., Bianchi, S., Aldea, J.,
1435 Huuskonen, S., 2023. Current state of mixed forests available for wood supply in Finland and
1436 Sweden. *Scand J For Res* 1–11. <https://doi.org/10.1080/02827581.2023.2259797>

1437 Lees, K.J., Artz, R.R.E., Chandler, D., Aspinall, T., Boulton, C.A., Buxton, J., Cowie, N.R., Lenton, T.M.,
1438 2021. Using remote sensing to assess peatland resilience by estimating soil surface moisture and
1439 drought recovery. *Science of The Total Environment* 761, 143312.
1440 <https://doi.org/10.1016/j.scitotenv.2020.143312>

1441 Letendre, J., Poulin, M., Rochefort, L., 2008. Sensitivity of spectral indices to CO₂ fluxes for several
1442 plant communities in a *Sphagnum*-dominated peatland. *Canadian Journal of Remote Sensing* 34,
1443 S414–S425. <https://doi.org/10.5589/m08-053>

1444 Loisel, J., Yu, Z., Beilman, D.W., Camill, P., Alm, J., Amesbury, M.J., Anderson, D., Andersson, S.,
1445 Bochicchio, C., Barber, K., Belyea, L.R., Bunbury, J., Chambers, F.M., Charman, D.J., De
1446 Vleeschouwer, F., Fiałkiewicz-Kozieł, B., Finkelstein, S.A., Galka, M., Garneau, M., Hammarlund,
1447 D., Hinchcliff, W., Holmquist, J., Hughes, P., Jones, M.C., Klein, E.S., Kokfelt, U., Korhola, A.,
1448 Kuhry, P., Lamarre, A., Lamentowicz, M., Large, D., Lavoie, M., MacDonald, G., Magnan, G.,
1449 Mäkilä, M., Mallon, G., Mathijssen, P., Mauquoy, D., McCarroll, J., Moore, T.R., Nichols, J.,
1450 O'Reilly, B., Oksanen, P., Packalen, M., Peteet, D., Richard, P.J.H., Robinson, S., Ronkainen, T.,
1451 Rundgren, M., Sannel, A.B.K., Tarnocai, C., Thom, T., Tuittila, E.-S., Turetsky, M., Väliranta, M.,
1452 van der Linden, M., van Geel, B., van Bellen, S., Vitt, D., Zhao, Y., Zhou, W., 2014. A database and
1453 synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation.
1454 *Holocene* 24, 1028–1042. <https://doi.org/10.1177/0959683614538073>

1455 Łuców, D., Küttim, M., Słowiński, M., Kołaczek, P., Karpińska-Kołaczek, M., Küttim, L., Salme, M.,
1456 Lamentowicz, M., 2022. Searching for an ecological baseline: Long-term ecology of a post-
1457 extraction restored bog in Northern Estonia. *Quaternary International* 607, 65–78.
1458 <https://doi.org/10.1016/j.quaint.2021.08.017>

1459 Łuców, D., Lamentowicz, M., Kołaczek, P., Łokas, E., Marcisz, K., Obremska, M., Theuerkauf, M.,
1460 Tyszkowski, S., Słowiński, M., 2021. Pine Forest Management and Disturbance in Northern Poland:
1461 Combining High-Resolution 100-Year-Old Paleoeological and Remote Sensing Data. *Front Ecol*
1462 *Evol* 9. <https://doi.org/10.3389/fevo.2021.747976>

1463 Marcisz, K., Bąk, M., Kołaczek, P., Lamentowicz, M., Wochal, D., 2024. Historia lasu i mokradel
1464 zapisana w torfowiskach, in: Lamentowicz, M., Koneczal, S. (Eds.), *Jak Chronić Torfowiska w*
1465 *Lasach? ArchaeGraph, Łódź*, pp. 29–45.

1466 Marcisz, K., Galka, M., Pietrala, P., Miotk-Szpiganowicz, G., Obremska, M., Tobolski, K., Lamentowicz,
1467 M., 2017. Fire activity and hydrological dynamics in the past 5700 years reconstructed from
1468 *Sphagnum* peatlands along the oceanic–continental climatic gradient in northern Poland. *Quat Sci*
1469 *Rev* 177, 145–157. <https://doi.org/10.1016/j.quascirev.2017.10.018>

1470 Marcisz, K., Jassey, V.E.J., Kosakyan, A., Krashevskaya, V., Lahr, D.J.G., Lara, E., Lamentowicz, Ł.,
1471 Lamentowicz, M., Macumber, A., Mazei, Y., Mitchell, E.A.D., Nasser, N.A., Patterson, R.T., Roe,

- 1472 H.M., Singer, D., Tsyganov, A.N., Fournier, B., 2020a. Testate Amoeba Functional Traits and Their
 1473 Use in Paleoecology. *Front Ecol Evol* 8, 340. <https://doi.org/10.3389/fevo.2020.575966>
- 1474 Marcisz, K., Kołaczek, P., Gałka, M., Diaconu, A.-C., Lamentowicz, M., 2020b. Exceptional hydrological
 1475 stability of a Sphagnum-dominated peatland over the late Holocene. *Quat Sci Rev* 231, 106180.
 1476 <https://doi.org/https://doi.org/10.1016/j.quascirev.2020.106180>
- 1477 Marcisz, K., Tinner, W., Colombaroli, D., Kołaczek, P., Słowiński, M., Fiałkiewicz-Kozieł, B., Łokas, E.,
 1478 Lamentowicz, M., 2015. Long-term hydrological dynamics and fire history over the last 2000 years
 1479 in CE Europe reconstructed from a high-resolution peat archive. *Quat Sci Rev* 112, 138–152.
 1480 <https://doi.org/10.1016/j.quascirev.2015.01.019>
- 1481 Marks, L., 2012. Timing of the Late Vistulian (Weichselian) glacial phases in Poland. *Quat Sci Rev* 44,
 1482 81–88. <https://doi.org/https://doi.org/10.1016/j.quascirev.2010.08.008>
- 1483 Matthias, I., Giesecke, T., 2014. Insights into pollen source area, transport and deposition from modern
 1484 pollen accumulation rates in lake sediments. *Quat Sci Rev* 87, 12–23.
 1485 <https://doi.org/10.1016/j.quascirev.2013.12.015>
- 1486 Matulewski, P., Buchwal, A., Makohonienko, M., 2019. Higher climatic sensitivity of Scots pine (*Pinus*
 1487 *sylvestris* L.) subjected to tourist pressure on a hiking trail in the Brodnica Lakeland, NE Poland.
 1488 *Dendrochronologia* (Verona) 54, 78–86. <https://doi.org/10.1016/j.dendro.2019.02.008>
- 1489 Mauquoy, D., Hughes, P.D.M., van Geel, B., 2010. A protocol for plant macrofossil analysis of peat
 1490 deposits. *Mires and Peat* 7, 1–5.
- 1491 Mauquoy, D., van Geel, B., 2007. Mire and peat macros, in: *Encyclopedia of Quaternary Science*.
 1492 Elsevier, Heidelberg, pp. 2315–2336.
- 1493 Mauquoy, D., Yeloff, D., 2008. Raised peat bog development and possible responses to environmental
 1494 changes during the mid- to late-Holocene. Can the palaeoecological record be used to predict the
 1495 nature and response of raised peat bogs to future climate change? *Biodivers Conserv* 17, 2139–2151.
 1496 <https://doi.org/10.1007/s10531-007-9222-2>
- 1497 Mazei, Y., Tsyganov, A.N., 2006. Freshwater testate amoebae. KMK, Moscow.
- 1498 McCarroll, D., Loader, N.J., 2004. Stable isotopes in tree rings. *Quat Sci Rev* 23, 771–801.
 1499 <https://doi.org/10.1016/j.quascirev.2003.06.017>
- 1500 McGrath, M.J., Luyssaert, S., Meyfroidt, P., Kaplan, J.O., Bürgi, M., Chen, Y., Erb, K., Gimmi, U.,
 1501 McInerney, D., Naudts, K., Otto, J., Pasztor, F., Ryder, J., Schelhaas, M.-J., Valade, A., 2015.
 1502 Reconstructing European forest management from 1600 to 2010. *Biogeosciences* 12, 4291–4316.
 1503 <https://doi.org/10.5194/bg-12-4291-2015>
- 1504 McNulty, S., Caldwell, P., Doyle, T.W., Johnsen, K., Liu, Y., Mohan, J., Prestemon, J., Sun, G., 2013.
 1505 Forests and Climate Change in the Southeast USA, in: Ingram, K.T., Dow, K., Carter, L., Anderson,
 1506 J. (Eds.), *Climate of the Southeast United States*. Island Press, Washington, pp. 165–189.
- 1507 Meisterfeld, R., 2001. Testate amoebae, in: Costello, M.J., Emblow, C.S., White, R. (Eds.), *Patrimoines*
 1508 *Naturels*. Muséum National d’Histoire Naturelle - Institut d’Ecologie et de Gestion de la Biodiversité
 1509 (I.E.G.B.) - Service du Patrimoine Naturel (S.P.N.), Paris, pp. 54–57.
- 1510 Messier, C., Bauhus, J., Sousa-Silva, R., Auge, H., Baeten, L., Barsoum, N., Bruelheide, H., Caldwell, B.,
 1511 Cavender-Bares, J., Dhiedt, E., Eisenhauer, N., Ganade, G., Gravel, D., Guillemot, J., Hall, J.S.,
 1512 Hector, A., Hérault, B., Jactel, H., Koricheva, J., Kreft, H., Mereu, S., Muys, B., Nock, C.A.,
 1513 Paquette, A., Parker, J.D., Perring, M.P., Ponette, Q., Potvin, C., Reich, P.B., Scherer-Lorenzen, M.,
 1514 Schnabel, F., Verheyen, K., Weih, M., Wollni, M., Zemp, D.C., 2022. For the sake of resilience and
 1515 multifunctionality, let’s diversify planted forests! *Conserv Lett* 15. <https://doi.org/10.1111/conl.12829>
- 1516 Milecka, K., Kowalewski, G., Fiałkiewicz-Kozieł, B., Gałka, M., Lamentowicz, M., Chojnicki, B.H.,
 1517 Goslar, T., Barabach, J., 2017. Hydrological changes in the Rzecin peatland (Puszcza Notecka,
 1518 Poland) induced by anthropogenic factors: Implications for mire development and carbon
 1519 sequestration. *Holocene* 27, 651–664. <https://doi.org/10.1177/0959683616670468>
- 1520 Miller, J.D., Anderson, H.A., Ferrier, R.C., Walker, T.A.B., 1990. Hydrochemical Fluxes and their Effects
 1521 on Stream Acidity in Two Forested Catchments in Central Scotland. *Forestry* 63, 311–331.
 1522 <https://doi.org/10.1093/forestry/63.4.311>

Z komentarzem [MB99]: We added citations.

- 1523 Minkkinen, K., Byrne, K.A., Trettin, C., 2008. Climate impacts of peatland forestry, in: Strack, M. (Ed.),
 1524 Peatlands and Climate Change. International Peat Society, Saarijärvi, pp. 98–122.
- 1525 Miola, A., 2012. Tools for Non-Pollen Palynomorphs (NPPs) analysis: A list of Quaternary NPP types and
 1526 reference literature in English language (1972–2011). *Rev Palaeobot Palynol* 186, 142–161.
 1527 <https://doi.org/10.1016/j.revpalbo.2012.06.010>
- 1528 Mirek, Z., Zarzycki, K., Wojewoda, W., Szeląg, Z., 2006. Red list of plants and fungi in Poland. Czerwona
 1529 lista roślin i grzybów Polski. Instytut Botaniki im. W. Szafera, Polska Akademia Nauk, Kraków.
- 1530 Mitosek, H., 1960. Letnio-jesienna susza 1959 r. *Postępy Nauk Rolniczych* 7, 53–64.
- 1531 Mokrzejki, Z., 1928. *Strzygonia choinówka*. Związek Zawodowy Leśników w Rzeczypospolitej Polskiej,
 1532 Warszawa.
- 1533 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen Analysis*. Blackwell Scientific Publications,
 1534 Oxford.
- 1535 Moritz, M.A., Parisien, M.-A., Battlori, E., Krawchuk, M.A., Van Dorn, J., Ganz, D.J., Hayhoe, K., 2012.
 1536 Climate change and disruptions to global fire activity. *Ecosphere* 3, 1–22.
 1537 <https://doi.org/10.1890/ES11-00345.1>
- 1538 Mroczkowska, A., Kittel, P., Marcisz, K., Dolbunova, E., Gauthier, E., Lamentowicz, M., Mazurkevich,
 1539 A., Obremska, M., Płóciennik, M., Kramkowski, M., Łuców, D., Kublitskiy, Y., Słowiński, M., 2021.
 1540 Small peatland with a big story: 600-year paleoecological and historical data from a kettle-hole
 1541 peatland in Western Russia. *Holocene* 31, 1761–1776. <https://doi.org/10.1177/09596836211033224>
- 1542 Nienartowicz, A., 2012. Historia lasów na obszarze Parku Narodowego „Bory Tucholskie” i na terenach
 1543 sąsiednich, in: Matuszkiewicz, J. (Ed.), *Świat Roślin i Grzybów Parku Narodowego „Bory
 1544 Tucholskie”. Park Narodowy „Bory Tucholskie”, Charzykowy, pp. 47–62.*
- 1545 Nisbet, T.R., 2001. The role of forest management in controlling diffuse pollution in UK forestry. *For Ecol*
 1546 *Manage* 143, 215–226. [https://doi.org/10.1016/S0378-1127\(00\)00519-3](https://doi.org/10.1016/S0378-1127(00)00519-3)
- 1547 Obmiński, Z., 1970. Zarys ekologii, in: Białobok, S. (Ed.), *Sosna Zwyczajna. Nasze Drzewa Leśne*.
 1548 Warszawa-Poznań, pp. 152–231.
- 1549 Ogden, C.G., Hedley, R.H., 1980. *An Atlas of Freshwater Testate Amoebae*. Oxford University Press,
 1550 London.
- 1551 Oris, F., Ali, A.A., Asselin, H., Paradis, L., Bergeron, Y., Finsinger, W., 2014. Charcoal dispersion and
 1552 deposition in boreal lakes from 3 years of monitoring: Differences between local and regional fires.
 1553 *Geophys Res Lett* 41, 6743–6752. <https://doi.org/10.1002/2014GL060984>
- 1554 Orłowicz, M., 1924. *Ilustrowany przewodnik po województwie pomorskiem*. Książnica Polska, Lwów.
- 1555 Paavilainen, E., Päivänen, J., 1995. *Peatland Forestry: Ecology and Principles*, Ecological Studies.
 1556 Springer, Berlin.
- 1557 Parish, F., Sirin, A., Charman, D.J., Joosten, H., Minayeva, T., Silviu, M., Stringer, L., 2008. Assessment
 1558 on peatlands, biodiversity and climate change: main report, Global Environment Centre, Kuala
 1559 Lumpur and Wetlands International, Wageningen.
- 1560 Pawlyta, J., Lamentowicz, M., 2010. Age-depth model for modern peat core. Methodological approach,
 1561 methods of absolute chronology., in: 10th International Conference. GADAM Centre of Excellent.
 1562 Department of Radioisotopes, Institute of Physics, Silesian University of Technology, Gliwice, p.
 1563 109.
- 1564 Payne, R.J., Mitchell, E.A.D., 2009. How many is enough? Determining optimal count totals for
 1565 ecological and palaeoecological studies of testate amoebae. *J. Paleolimnol.* 42, 483–495.
 1566 <https://doi.org/10.1007/s10933-008-9299-y>
- 1567 Pędziszewska, A., Latałowa, M., 2016. Stand-scale reconstruction of late Holocene forest succession on
 1568 the Gdańsk Upland (N. Poland) based on integrated palynological and macrofossil data from paired
 1569 sites. *Veg Hist Archaeobot* 25, 239–254. <https://doi.org/10.1007/s00334-015-0546-7>
- 1570 Péli, E.R., Nagy, J.G., Cserhalmi, D., 2015. In situ measurements of seasonal productivity dynamics in
 1571 two sphagnum dominated mires in Hungary. *Carpathian Journal of Earth and Environmental
 1572 Sciences* 10, 231–240.

Z komentarzem [MB100]: We moved it to Supplementary File 1.

1573 Peters, M.E., Higuera, P.E., 2007. Quantifying the source area of macroscopic charcoal with a particle
1574 dispersal model. *Quat Res* 67, 304–310. <https://doi.org/10.1016/j.yqres.2006.10.004>

1575 Poraj-Górska, A.I., Żarczyński, M.J., Ahrens, A., Enters, D., Weisbrodt, D., Tylmann, W., 2017. Impact of
1576 historical land use changes on lacustrine sedimentation recorded in varved sediments of Lake Jaczno,
1577 northeastern Poland. *Catena (Amst)* 153, 182–193.
1578 <https://doi.org/http://dx.doi.org/10.1016/j.catena.2017.02.007>

1579 Przybylski, T., 1993. *Ekologia*, in: Białobok, S., Boratyński, A., Bugała, W. (Eds.), *Biologia Sosny*
1580 *Zwyczajnej*. Sorus, Poznań-Kórnik, pp. 255–300.

1581 Pureswaran, D.S., De Grandpré, L., Paré, D., Taylor, A., Barrette, M., Morin, H., Régnière, J., Kneeshaw,
1582 D.D., 2015. Climate-induced changes in host tree–insect phenology may drive ecological state-shift
1583 in boreal forests. *Ecology* 96, 1480–1491. <https://doi.org/10.1890/13-2366.1>

1584 R Core Team, 2022. R: A Language and Environment for Statistical Computing. R Foundation for
1585 Statistical Computing [WWW Document]. URL <https://www.R-project.org> (accessed 12.4.23).

1586 Rapinel, S., Panhelleux, L., Gayet, G., Vanacker, R., Lemerrier, B., Laroche, B., Chambaud, F.,
1587 Guelmami, A., Hubert-Moy, L., 2023. National wetland mapping using remote-sensing-derived
1588 environmental variables, archive field data, and artificial intelligence. *Heliyon* 9, e13482.
1589 <https://doi.org/10.1016/j.heliyon.2023.e13482>

1590 Rastogi, A., Stróżecki, M., Kalaji, H.M., Łuców, D., Lamentowicz, M., Juszczak, R., 2019. Impact of
1591 warming and reduced precipitation on photosynthetic and remote sensing properties of peatland
1592 vegetation. *Environ Exp Bot* 160, 71–80.
1593 <https://doi.org/https://doi.org/10.1016/j.envexpbot.2019.01.005>

1594 Regulation No. 64/97 of the Bydgoszcz Voivode of October 30, 1997, on the recognition as ecological
1595 sites of natural objects in the Bydgoszcz Voivodeship, 1997.

1596 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng,
1597 H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg,
1598 A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der
1599 Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L.,
1600 Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P.,
1601 Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The
1602 IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62,
1603 725–757. <https://doi.org/10.1017/RDC.2020.41>

1604 Reynolds, B., Ormerod, S.J., Gee, A.S., 1994. Spatial patterns concentrations in upland Wales in relation
1605 to catchment forest cover and forest age. *Environmental Pollution* 84, 27–33.
1606 [https://doi.org/10.1016/0269-7491\(94\)90067-1](https://doi.org/10.1016/0269-7491(94)90067-1)

1607 Rouse, J.W., Haas, R.H., Schell, J.A., Deering, D.W., 1974. *Monitoring vegetation systems in the Great*
1608 *Plains with ERTS, Third ERTS Symposium*. ed. NASA.

1609 Rydin, H., Jeglum, J.K., 2013. *The biology of peatlands (Second Edition)*. Oxford University Press.

1610 Sanderson, N., Loisel, J., Gallego-Sala, A., Anshari, G., Novita, N., Marcisz, K., Lamentowicz, M., Bąk,
1611 M., Wochal, D., 2023. Setting a new research agenda for tropical peatlands, recent carbon
1612 accumulation and ecosystem services. *Past Global Changes Magazine* 31, 121–121.
1613 <https://doi.org/10.22498/pages.31.2.121>

1614 Schueler, S., Falk, W., Koskela, J., Lefèvre, F., Bozzano, M., Hubert, J., Kraigher, H., Longauer, R., Olrik,
1615 D.C., 2014. Vulnerability of dynamic genetic conservation units of forest trees in Europe to climate
1616 change. *Glob Chang Biol* 20, 1498–1511. <https://doi.org/10.1111/gcb.12476>

1617 Schüle, M., Domes, G., Schwanz, C., Heinken, T., 2023. Early natural tree regeneration after wildfire in
1618 a Central European Scots pine forest: Forest management, fire severity and distance matters. *For*
1619 *Ecol Manage* 539, 120999. <https://doi.org/10.1016/j.foreco.2023.120999>

1620 Schütte, R., 1893. *Die Tucheler Haide vornehmlich in forstlicher Beziehung*. Bertling, Danzig.

1621 Seidl, R., Schelhaas, M.-J., Rammer, W., Verkerk, P.J., 2014. Increasing forest disturbances in Europe and
1622 their impact on carbon storage. *Nat Clim Chang* 4, 806. <https://doi.org/10.1038/nclimate2318>
1623 <https://www.nature.com/articles/nclimate2318#supplementary-information>

1624 Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D.,
1625 Petr, M., Honkaniemi, J., Lexer, M.J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel,
1626 T.A., Reyer, C.P.O., 2017. Forest disturbances under climate change. *Nat Clim Chang* 7, 395.
1627 <https://doi.org/10.1038/nclimate3303> [https://www.nature.com/articles/nclimate3303#supplementary-](https://www.nature.com/articles/nclimate3303#supplementary-information)
1628 [information](https://www.nature.com/articles/nclimate3303#supplementary-information)

1629 Shumilovskikh, L.S., Shumilovskikh, E.S., Schlütz, F., van Geel, B., 2022. NPP-ID: Non-Pollen
1630 Palynomorph Image Database as a research and educational platform. *Veg Hist Archaeobot* 31, 323–
1631 328. <https://doi.org/10.1007/s00334-021-00849-8>

1632 Shumilovskikh, L.S., van Geel, B., 2020. Non-Pollen Palynomorphs, in: Henry, E.G. (Ed.), *Handbook for*
1633 *the Analysis of Micro-Particles in Archaeological Samples*. Springer Cham, pp. 65–94.

1634 Siemensma, F., 2023. Microworld, world of amoeboid organisms [WWW Document]. URL
1635 <https://arcella.nl/> (accessed 11.23.23).

1636 Sillasoo, U., Väliaranta, M., Tuittila, E.S., 2011. Fire history and vegetation recovery in two raised bogs at
1637 the Baltic Sea. *Journal of Vegetation Science* 22, 1084–1093.

1638 Simard, I., Morin, H., Lavoie, C., 2006. A millennial-scale reconstruction of spruce budworm abundance
1639 in Saguenay, Québec, Canada. *Holocene* 16, 31–37. <https://doi.org/10.1191/0959683606hl904rp>

1640 Śliwa, E., 1989. Przebieg masowego pojawu brudnicy mniszki (*Lymantria monacha* L.) i jej zwalczania w
1641 Polsce w latach 1978-1985 oraz regeneracja aparatu asymilacyjnego w uszkodzonych
1642 drzewostanach. *Prace Instytutu Badawczego Leśnictwa* 710, 3–120.

1643 Śliwa, E., 1987. Występowanie i zwalczanie ważniejszych folio fagów w drzewostanach sosnowych w
1644 latach 1946-1985. *Sylvan* 11–12.

1645 Śliwa, E., 1974. Gradacja strzygoni choinówki (*Panolis flammea* Schiff.) w Kampinowskim Parku
1646 Narodowym. *Sylvan* 11, 61–66.

1647 Słowiński, M., Lamentowicz, M., Luców, D., Barabach, J., Brykała, D., Tyszkowski, S., Pieńczewska, A.,
1648 Śnieszko, Z., Dietze, E., Jażdżewski, K., Obremska, M., Ott, F., Brauer, A., Marcisz, K., 2019.
1649 Paleoeological and historical data as an important tool in ecosystem management. *J Environ*
1650 *Manage* 236, 755–768. <https://doi.org/10.1016/j.jenvman.2019.02.002>

1651 Spiecker, H., 2000. Growth of Norway Spruce (*Picea abies* [L.] Karst.) under Changing Environmental
1652 Conditions in Europe, in: Klimo, E., Hager, H., Kulhavý, J. (Eds.), *Spruce Monocultures in Central*
1653 *Europe – Problems and Prospects*. European Forest Institute, Joensuu, pp. 11–26.

1654 Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., Dosio, A., 2018. Will drought events become more
1655 frequent and severe in Europe? *International Journal of Climatology* 38, 1718–1736.
1656 <https://doi.org/10.1002/joc.5291>

1657 Stivrins, N., Liiv, M., Ozola, I., Reitalu, T., 2018. Carbon accumulation rate in a raised bog in Latvia, NE
1658 Europe, in relation to climate warming. *Estonian Journal of Earth Sciences* 67, 247.
1659 <https://doi.org/10.3176/earth.2018.20>

1660 Stockmarr, J., 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13, 615–621.

1661 Sullivan, J., 1993. <https://www.fs.usda.gov/database/feis/plants/tree/pinsyl/all.html> [WWW Document].
1662 *Pinus sylvestris*. In: *Fire Effects Information System*.

1663 Sullivan, M.E., Booth, R.K., 2011. The Potential Influence of Short-term Environmental Variability on the
1664 Composition of Testate Amoeba Communities in Sphagnum Peatlands. *Microb Ecol* 62, 80–93.
1665 <https://doi.org/10.1007/s00248-011-9875-y>

1666 Szwankowski, J., 2005. Powiat tucholski w latach 1875-1920, administracja, ludność, gospodarka,
1667 kultura. Logo, Tuchola.

1668 Taeger, S., Zang, C., Liesebach, M., Schneck, V., Menzel, A., 2013. Impact of climate and drought events
1669 on the growth of Scots pine (*Pinus sylvestris* L.) provenances. *For Ecol Manage* 307, 30–42.
1670 <https://doi.org/10.1016/j.foreco.2013.06.053>

1671 Tinner, W., Hu, F.S., 2003. Size parameters, size-class distribution and area-number relationship of
1672 microscopic charcoal: relevance for fire reconstruction. *Holocene* 13, 499–505.

1673 Tipton, J., 2023. Past anthropogenic impacts on peatland through the forest management practices in the
1674 Polish Tuchola Forest (Master Thesis). Adam Mickiewicz University, Poznań.

1675 [Tobolski, K., 2000. Przewodnik do oznaczania torfów i osadów jeziornych. PWN, Warszawa.](#)

1676 Trouet, V., Van Oldenborgh, G.J., 2013. KNMI Climate Explorer: A Web-Based Research Tool for High-

1677 Resolution Paleoclimatology. *Tree Ring Res* 69, 3–13. <https://doi.org/10.3959/1536-1098-69.1.3>

1678 Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science* (1979) 349, 814–

1679 818. <https://doi.org/10.1126/science.aac6759>

1680 Ulrich, B., 1980. Production and consumption of Hydrogen ions in the ecosphere, in: Hutchinson, T.,

1681 Havas, M. (Eds.), *Effects of Acid Precipitation on Terrestrial Ecosystems*. Plenum Press, New York,

1682 London, pp. 255–282.

1683 UNESCO, 2024. *Tuchola Forests [WWW Document]. Man and the Biosphere Programme (MAB)*.

1684 Väiliranta, M., Korhola, A., Seppä, H., Tuittila, E.S., Sarmaja-Korjonen, K., Laine, J., Alm, J., 2007. High-

1685 resolution reconstruction of wetness dynamics in a southern boreal raised bog, Finland, during the

1686 late Holocene: a quantitative approach. *Holocene* 17, 1093–1107.

1687 <https://doi.org/10.1177/0959683607082550>

1688 van der Linden, M., Heijmans, M.M., van Geel, B., 2014. Carbon accumulation in peat deposits from

1689 northern Sweden to northern Germany during the last millennium. *Holocene* 24, 1117–1125.

1690 <https://doi.org/10.1177/0959683614538071>

1691 van der Schrier, G., Allan, R.P., Ossó, A., Sousa, P.M., Van de Vyver, H., Van Schaeybroeck, B.,

1692 Coscarelli, R., Pasqua, A.A., Petrucci, O., Curley, M., Mietus, M., Filipiak, J., Štěpánek, P.,

1693 Zahradníček, P., Brázdil, R., Rezníčková, L., van den Besselaar, E.J.M., Trigo, R., Aguilar, E., 2021.

1694 The 1921 European drought: impacts, reconstruction and drivers. *Climate of the Past* 17, 2201–2221.

1695 <https://doi.org/10.5194/cp-17-2201-2021>

1696 Waller, M., 2013. Drought, disease, defoliation and death: forest pathogens as agents of past vegetation

1697 change. *J Quat Sci* 28, 336–342. <https://doi.org/10.1002/jqs.2631>

1698 Wardenaar, E.C.P., 1987. A new hand tool for cutting peat profiles. *Canadian Journal of Botany* 65, 1772–

1699 1773. <https://doi.org/10.1139/b87-243>

1700 Warner, B.G., 1993. Palaeoecology of floating bogs and landscape change in the Great Lakes drainage

1701 basin of North America, in: Chambers, F.M. (Ed.), *Climate Change and Human Impact on the*

1702 *Landscape*. Chapman & Hall, pp. 237–248.

1703 Westerling, A.L., 2016. Increasing western US forest wildfire activity: sensitivity to changes in the timing

1704 of spring. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371, 20150178.

1705 <https://doi.org/10.1098/rstb.2015.0178>

1706 Westman, C.J., Laiho, R., 2003. Nutrient dynamics of drained peatland forests. *Biogeochemistry* 63, 269–

1707 298. <https://doi.org/10.1023/A:1023348806857>

1708 Whitlock, C., Larsen, C., 2001. Charcoal as a fire proxy., in: *Tracking Environmental Change Using Lake*

1709 *Sediments. Terrestrial, Algal, and Siliceous Indicators*. J. P. Smol, H. J. B. Birks, and W. M. Last,

1710 Eds. Dordrecht: Kluwer, pp. 75–97.

1711 Wilson, J.K., 2012. *The German Forest. Nature, Identity, and the Contestation of a National Symbol,*

1712 1871–1914. University of Toronto Press.

1713 Wilson, R.M., Hopple, A.M., Tffaily, M.M., Sebestyén, S.D., Schadt, C.W., Pfeifer-Meister, L., Medvedeff,

1714 C., McFarlane, K.J., Kostka, J.E., Kolton, M., Kolka, R.K., Kluber, L.A., Keller, J.K., Guilderson,

1715 T.P., Griffiths, N.A., Chanton, J.P., Bridgman, S.D., Hanson, P.J., 2016. Stability of peatland carbon

1716 to rising temperatures. *Nat Commun* 7, 13723. <https://doi.org/10.1038/ncomms13723>

1717 <http://www.nature.com/articles/ncomms13723#supplementary-information>

1718 Woodward, C., Haines, H.A., 2020. Unprecedented long-distance transport of macroscopic charcoal from

1719 a large, intense forest fire in eastern Australia: Implications for fire history reconstruction. *Holocene*

1720 30, 947–952. <https://doi.org/10.1177/0959683620908664>

1721 Wotton, B.M., Nock, C.A., Flannigan, M.D., 2010. Forest fire occurrence and climate change in Canada.

1722 *Int J Wildland Fire* 19, 253. <https://doi.org/10.1071/WF09002>

1723 Young, D.M., Baird, A.J., Charman, D.J., Evans, C.D., Gallego-Sala, A. V., Gill, P.J., Hughes, P.D.M.,

1724 Morris, P.J., Swindles, G.T., 2019. Misinterpreting carbon accumulation rates in records from near-

1725 surface peat. *Sci Rep* 9, 17939. <https://doi.org/10.1038/s41598-019-53879-8>

Z komentarzem [MB101]: We removed it.

- 1726 Yu, Z., Loisel, J., Brosseau, D.P., Beilman, D.W., Hunt, S.J., 2010. Global peatland dynamics since the
1727 Last Glacial Maximum. *Geophys Res Lett* 37. <https://doi.org/10.1029/2010GL043584>
- 1728 Zajączkowski, J., Brzeziecki, B., Kozak I., 2013. Wpływ potencjalnych zmian klimatycznych na zdolność
1729 konkurencyjną głównych gatunków drzew w Polsce. *Sylvan* 157, 253–261.
- 1730 Zang, C., Biondi, F., 2015. treeclim: an R package for the numerical calibration of proxy-climate
1731 relationships. *Ecography* 38, 431–436. <https://doi.org/10.1111/ecog.01335>
- 1732 Zielski, A., 1996. Wpływ temperatury i opadów na szerokość słoików rocznych drewna u sosny zwyczajnej
1733 (<s>Pinus sylvestris<s> L. w rejonie Torunia). *Sylvan* 2, 71–80.
- 1734 Zielski, A., Barankiewicz, A., 2000. Dendrochronologiczna analiza przyrostów radialnych sosny
1735 zwyczajnej na terenie leśnictwa Dąbie, Nadleśnictwa Włocławek. *Sylvan* 2000, 69–74.
- 1736 Zielski, A., Błaszczkowski, A., Barankiewicz, A., 1998. Dynamika przyrostu radialnego sosny zwyczajnej
1737 (*Pinus sylvestris* L.) na obszarze leśnym eksploatowanym turystycznie nad jeziorem Wielkie
1738 Partęczyny (Nadleśnictwo Brodnica). *Sylvan* 3, 69–78.
- 1739 Zielski, A., Krąpiec, M., 2004. *Dendrochronologia*. PWN.
- 1740 Zielski, A., Krąpiec, M., Koprowski, M., 2010. Dendrochronological data, in: Przybylak, R., Majorowicz,
1741 J., Brázdil, R., Kejna, M. (Eds.), *The Polish Climate in the European Context: An Historical*
1742 *Overview*. pp. 191–217.
- 1743 Zielski, A., Sygit, W., 1998. Wpływ klimatu na przyrost radialny sosny zwyczajnej (*Pinus sylvestris* L.),
1744 badania wzdłuż równoleżnika 52°N i transekcje Śląsk-Białowieża. In: In: Breymeyer, A., Roo-
1745 Zielińska, E. (Eds.), *Reakcja borów sosnowych na zmianę klimatu wzdłuż równoleżnika 52°N (12-
1746 32°E) oraz na zmiany w depozycji zanieczyszczeń chemicznych na transekcji Śląsk-Białowieża*.
1747 *Dokumentacja Geograficzna IG PAN Warszawa* 13, 161–185.
- 1748