

1 **Assessing the impact of forest management and climate on a peatland under Scots pine monoculture**
2 **using a multidisciplinary approach**
3

4 Mariusz Bąk¹, Mariusz Lamentowicz¹, Piotr Kołaczek¹, Daria Wochal¹, Paweł Matulewski², Dominik
5 Kopec^{3,4}, Martyna Wietecha^{4,5}, Dominika Jaster², Katarzyna Marcisz¹
6

7 ¹Climate Change Ecology Research Unit, Faculty of Geographical and Geological Sciences, Adam
8 Mickiewicz University, Poznań, Poland

9 ²Anthropocene Research Unit, Faculty of Geographical and Geological Sciences, Adam Mickiewicz
10 University, Poznań, Poland

11 ³Department of Biogeography, Paleoecology and Nature Conservation, University of Lodz, Łódź, Poland

12 ⁴MGGP Aero Sp. z o. o., Tarnów, Poland

13 ⁵Doctoral School of Exact and Natural Sciences, University of Lodz, Łódź, Poland
14
15

16 *Correspondence to:* Mariusz Bąk, mariusz.bak@amu.edu.pl
17

18 **Abstract:** Assessing the scale, rate and consequences of climate change, manifested primarily by rising
19 average air temperatures and altered precipitation regimes, is a critical challenge in contemporary scientific
20 research. These changes are accompanied by various anomalies and extreme events that negatively impact
21 ecosystems worldwide. Monoculture forests, including Scots pine (*Pinus sylvestris* L.) monocultures, are
22 particularly vulnerable to these changes due to their homogeneous structure and simplified ecosystem
23 linkages compared to mixed forests, making them more sensitive to extreme events such as insect outbreaks,
24 droughts, fires and strong winds. In the context of global warming, forest fires are becoming extremely
25 dangerous, and the risk of their occurrence increases as average temperatures rise. The situation becomes
26 even more dramatic when fire enters areas of peatlands, as these ecosystems effectively withdraw carbon
27 from the rapid carbon cycle and store it for up to thousands of years. Consequently, peatlands become
28 emitters of carbon dioxide into the atmosphere.

29 In this study, we aim to trace the last 300 years of historical development of a peatland situated in a Scots
30 pine monoculture. Our focus is on the Okoniny (Jezierzba) peatland located within the Tuchola Pinewoods
31 in northern Poland, one of the country's largest forest complexes. We delved into the phase when the
32 peatland's surroundings transitioned from a mixed forest to a pine monoculture and investigated the impact
33 of changes in forest management on the peatland vegetation and hydrology. Our reconstructions are based
34 on a multi-proxy approach using: pollen, plant macrofossils, micro- and macrocharcoal and testate amoebae.
35 We combine the peatland palaeoecological record with the dendrochronology of *Pinus sylvestris* to compare
36 the response of these two archives. Our results show that a change in forest management and progressive
37 climate warming affected the development of the peatland. We note an increase in acidity over the analyzed

38 period and a decrease in the water table over the last few decades that led to the lake-peatland transition.
39 These changes progressed with the strongest agricultural activity in the area in the 19th century. However,
40 the 20th century was a period of continuous decline in agriculture and an increase in the dominance of Scots
41 pine in the landscape as the effect of afforestation. Dendroclimatic data indicate a negative effect of
42 temperature on Scots pine and pressure from summer rainfall deficiency. Additional remote sensing
43 analysis, using hyperspectral, LiDAR and thermal airborne data, provided information about the current
44 condition of the peatland vegetation. With the application of spectral indices and the analysis of land surface
45 temperature, spatial variations in peatland drying have been identified. Considering the context of forest
46 management and the protection of valuable ecosystems in monocultural forests, the conclusions are relevant
47 for peatland and forest ecology, palaeoecology and forestry.

48
49 **Keywords:** palaeoecological data, palaeoecology, dendrochronology, dendroclimatic data, climate change,
50 monoculture forests, plantation, historical data, historical maps, multi-proxy, high-resolution, airborne data,
51 thermal data, vegetation indices, remote sensing

52
53 **1. Introduction**

54 Peatlands are vulnerable to various types of change, which play an important role in the global carbon cycle
55 and whose destabilization can create positive feedback for climate warming (Gallego-Sala et al., 2018;
56 Wilson et al., 2016). Peatlands, although they only cover about 3% of the Earth's total land area (Parish et
57 al., 2008; Rydin and Jeglum, 2013), store more than 30% of the organic carbon (C) (Freeman et al., 2004;
58 Gorham, 1991; Harenda et al., 2018), which is far more carbon than the entire biomass of the world's forests
59 (Beaulne et al., 2021b). Their advantage over forests is not only due to their ability to accumulate C but also
60 to the fact that they do not emit decomposed carbon from the so-called rapid C cycle for up to thousands of
61 years (Blodau, 2002; Gorham, 1991). The estimation of C content accumulated in peatlands is challenging
62 (Sanderson et al., 2023), although some studies indicate ca. 600 Gt of C in the Northern Hemisphere alone
63 (Yu et al., 2010). It has recently been shown that even the smallest kettle-hole peatlands effectively
64 accumulate C and serve as important C hot spots (Karpínska-Kołaczek et al., 2024).

65 Insufficient awareness of the ecological importance of peatlands has led to them being treated as
66 wastelands and drained for hundreds of years to obtain land for agriculture, and forestry or exploited
67 commercially as an energy resource (Joosten et al., 2012; Łuców et al., 2022; Paavilainen and Päivänen,
68 1995). Many of these areas have also had to adapt to a changing environment resulting from the use of
69 various forest management techniques, e.g., the replacement of mixed forests with more easily managed
70 monoculture forests (plantations) (Lee et al., 2023; Łuców et al., 2021; Słowiński et al., 2019). Mixed
71 forests, through greater biodiversity, are more resilient and better able to adapt to environmental change

72 (Bauhus et al., 2017; Messier et al., 2022), providing a more comprehensive range of ecosystem services
73 (Felton et al., 2016; Huuskonen et al., 2021).

74 Despite being more straightforward to manage, forest monocultures are characterized by simplified
75 ecosystem linkages (Chapin et al., 2012). As a result, they are more susceptible to various extreme events
76 and disturbances, both natural and anthropogenic, including droughts, fires, strong winds, and pest
77 infestations (Grondin et al., 2014). This is particularly important as disturbances of these types of forests
78 are becoming more common (Seidl et al., 2014; Westerling, 2016). Natural disturbance regimes in forests
79 are mainly a response to climate change (Hanson and Weltzin, 2000; Pureswaran et al., 2015; Seidl et al.,
80 2017; Trumbore et al., 2015), therefore they are expected to increase in frequency and severity in the coming
81 years (Gregow et al., 2017; Moritz et al., 2012; Wotton et al., 2010). Moreover, the problem applies to all
82 kinds of monoculture forests regardless of the dominant species and climate zones (Booth, 2013; Guariguata
83 et al., 2008; McNulty et al., 2013; Spiecker, 2000), including pine plantations in the temperate climate zone
84 of Central and Eastern Europe (Łuców et al., 2021; Schüle et al., 2023). Thus, peatlands, which are so crucial
85 in terms of their impact on global climate change, located in the area of forest monocultures are even more
86 vulnerable to extreme phenomena and disturbance, despite the already high climatic and anthropogenic
87 pressure.

88 The history of peatlands' development can be traced using palaeoecological analyses, which allow
89 numerous reconstructions of past environmental conditions, including climate change (Lamentowicz et al.,
90 2015; Mauquoy and Yeloff, 2008). These include reconstructions of vegetation changes in the peatland and
91 its surroundings, changes in the water table, and reconstructions of past fire activity (Gałka et al., 2022;
92 Kołaczek et al., 2018; Marcisz et al., 2020b, 2017; Mroczkowska et al., 2021). Peat archive records contain
93 a wide range of preserved micro- and macrofossils for example, pollen, spores, microbial remains, and
94 charcoal are deposited in situ and brought in by wind or water (Godwin, 1981). While paleoenvironmental
95 reconstructions based on peat records have become common, few studies integrate palaeoecological data
96 with other methods. For example, studies that combine palaeoecological and dendrochronological records,
97 including dendroclimatic reconstructions based on analysis of the annual growth of tree rings, are still
98 relatively rare (Ballesteros-Cánovas et al., 2022; Beaulne et al., 2021a; Dinella et al., 2021; Edvardsson et
99 al., 2022, 2019, 2016; González de Andrés et al., 2022; Kuosmanen et al., 2020; Lamentowicz et al., 2009b).
100 Yet, combining peat records with dendrochronological data can benefit interpretations of trees and forest
101 resilience and resistance to disturbances compared to local environmental changes recorded in peat. Such a
102 view of past environmental changes through several proxies and other archive types is fundamental and will
103 be helpful for forest management and nature conservation in the future. To assess the current state of the
104 peatland, we also included remote sensing data in the analysis. Remote sensing methods have been applied
105 to study wetland conditions for over 50 years and are currently regarded as one of the most useful methods

106 in this research area (FAO, 2020; Guo et al., 2017). Remote sensing technologies enable the remote and
107 non-invasive acquisition of information about the research object using specialized sensors, typically
108 mounted aboard satellites or aircraft. In this study, data obtained from a multisensor aerial platform were
109 used to assess the extent of peatland, the identification of drainage ditches, and the current vegetation
110 condition.


111 Our study aims to assess the impact of forest management (introduction of pine monoculture) and
112 changing climate on the vegetation, as well as hydrological, and trophic conditions of a peatland in CE
113 Europe by integrating various data sources - palaeoecology, dendrochronology, remote sensing, and
114 historical information. We assumed that the introduction of pine monoculture led to changes in the species
115 composition of peatlands in favor of *Sphagnum* mosses, as well as to the stabilization of the water table. We
116 also undertook to confirm whether peatlands register and respond to extreme events, both *in situ* and in the
117 immediate environment. We assumed that the disturbances that occurred in the monoculture forest would
118 be recorded in the tree rings (annual growths) record of Scots pine (*Pinus sylvestris* L.) and would confirm
119 and complement the palaeoecological reconstruction of the peatland. Thus, we have identified peat layers
120 corresponding to the occurrence of extremes known from historical sources and compared
121 dendrochronological (dendroclimatic) data with them.

122

123 2. Materials and methods

124 2.1. Study site

125 The Okoniny (Jezierzba) peatland (53°40'52"–53°41'21"N 18°03'09"–18°03'40"E according to
126 standard WGS 84) is located in northern Poland, about 60 km north of Bydgoszcz and about 20 km northeast
127 of Tuchola (Fig. 1). The study area is located within the Tuchola Pinewoods mesoregion (Kondracki, 2001),
128 close to the Pomeranian ice margin of the Vistulian Glaciation dated to ca. 17,000–16,000 cal. BP (Marks,
129 2012). The entire area of the Tuchola Pinewoods is a young glacial landscape covered by glacial till, sand,
130 and numerous depressions and other forms originating from melting dead ice (Błaszkiwicz et al., 2015).
131 Based on the analysis of remote sensing data, it was determined that the surface area of the peatland is 27.08
132 hectares, with approximately 7.00 hectares designated as non-forested area. The direct catchment area of
133 the peatland covers a surface of 33.23 hectares. The current elevation of the peatland is around 119 m asl,
134 with the highest elevated area within the direct catchment reaching around 128 m asl. It is part of a protected
135 area (Regulation No. 64/97, 1997), included within the boundaries of the Tuchola Landscape Park (created
136 in 1985). Moreover, since 2008 the entire complex of the Tuchola Pinewoods has been included on the
137 Natura 2000 list as a Special Protection Area. Since 2010, it has been listed as a UNESCO Biosphere
138 Reserve (UNESCO, 2024).



139
 140 Figure 1. Location of the study area. (A) Location on a map of north-western Poland. (B) Location of the
 141 two study sites – dendrochronological sampling site and peat coring site. (C) Okoniny (Jezierzba) peatland
 142 sampling site with current peatland boundaries.

143
 144 The Okoniny (Jezierzba) peatland is located in a temperate latitude zone, with a transitional climate
 145 influenced by continental air masses from eastern Europe and oceanic air masses from the Atlantic Ocean
 146 (Beck et al., 2018). According to climate data obtained from the Institute of Meteorology and Water
 147 Management for the meteorological station in Chojnice (35 km west of the study area) for the period
 148 between 1991-2020, the coldest month is January with an average temperature of -1.5 °C, the warmest
 149 month is July with an average temperature of 18.0 °C. Between 1961-1990, both January and July were
 150 cooler by 1.6 °C compared to 1991-2020. The average annual temperature increased from 6.9 °C in 1951-

151 1990 to 8.1 °C in 1991-2020. In terms of precipitation, February has the least amount with an average of
 152 31.1 mm for the period 1991-2020, and July has the most with an average of 80.7 mm for the period 1991-
 153 2020. Compared to 1951-1990, the average precipitation for February increased by 7.7 mm and for July
 154 decreased by 4.1 mm. Mean annual rainfall increased from 558.1 mm for 1951-1990 to 612.4 mm for 1991-
 155 2020.

156 Samples for dendroclimatic analysis were taken from forest division no. 91 in the Woziwoda
 157 Forestry, Woziwoda Forest District, about 9.5 km west of the study site (Fig. 1). The oldest pine trees in the
 158 forest district were selected for the study according to the indications of the forest survey and taxonomic
 159 descriptions.

160

161 2.2. Peat and tree core sampling

162 A peat core was taken from the north-western part of the peatland in February 2022 using a Wardenaar
 163 corer (chamber dimension: 10 cm × 10 cm × 100 cm) (Wardenaar, 1987). The entire length of the sampled
 164 peat core – 96 cm-long monolith – was analyzed. The core was sampled continuously every 1 cm, except
 165 for the top 10 cm, which contained a living *Sphagnum* layer. The first sample covered 4 cm of the surface
 166 layer (0-4 cm), and the following three samples were taken every 2 cm (4-6, 6-8 and 8-10 cm). 90 samples
 167 were obtained and analyzed for bulk density, ash content, peat and carbon accumulation rates, plant
 168 macrofossils, testate amoebae, macroscopic and microscopic charcoal, and pollen.

169 The research tree stem material was taken in April 2023 from 23 living and healthy trees at the
 170 Woziwoda site, ca. 9.5 km west of the Okoniny (Jezierzba) peatland. From each tree, a minimum of two
 171 cores were taken (from the east and west sides) at a breast height (1.3 m) with a Pressler increment corer. In
 172 total, 50 cores were acquired from the Scots pine tree stems.

173

174 2.3. Radiocarbon dating and chronology

175 Ten samples containing *Sphagnum* stems and leaves were used for accelerator mass spectroscopy
 176 (AMS) ¹⁴C dating of the entire length of the profile. The survey was conducted at the Poznan Radiocarbon
 177 Laboratory in Poland (laboratory code marked Poz; Tab. 1). The IntCal20 (Reimer et al., 2020) and
 178 Bomb21NH1 (Hua et al., 2021) atmospheric curves were used to calibrate the dates.

179

180 Table 1. The list of radiocarbon dates from Okoniny (Jezierzba) peatland with calibration in the OxCal
 181 v4.4.4 software using the IntCal20 calibration curve for the atmospheric data and Bomb21NH1 curve for
 182 bomb series.

No	Laboratory code – number sample	Depth (cm)	¹⁴ C date (¹⁴ C BP)	Calibrated dates [cal. CE (2σ – 95.4%)	Dated material
----	------------------------------------	---------------	---	---	----------------

1	Poz-150386	10.5	100.86 ± 0.33 pMC	1952-1958 (33.9%) 2013-... (61.5%)	<i>Sphagnum</i> stems
2	Poz-150387	20.5	107.92 ± 0.34 pMC	1952-1958 (11.1%) 1996-2009 (84.4%)	<i>Sphagnum</i> stems
3	Poz-150388	30.5	132.8 ± 0.36 pMC	1958-1962 (20.8%) 1972-1984 (74.6%)	<i>Sphagnum</i> stems
4	Poz-150445	40.5	165 ± 30	1661-1706 (17.2%) 1720-1818 (44.0%) 1832-1892 (14.9%) 1906-... (19.5%)	<i>Sphagnum</i> stems
5	Poz-150446	50.5	85 ± 30	1688-1730 (26.1%) 1806-1924 (69.3%)	<i>Sphagnum</i> stems
6	Poz-150447	60.5	105 ± 30	1682-1736 (25.9%) 1802-1936 (69.5%)	<i>Sphagnum</i> stems
7	Poz-150449	70.5	135 ± 30	1674-1766 (32.8%) 1774-1776 (0.6%) 1798-1942 (62.0%)	<i>Sphagnum</i> stems
8	Poz-150450	80.5	165 ± 30	1661-1706 (17.2%) 1720-1818 (44.0%) 1832-1892 (14.9%) 1906-... (19.5%)	<i>Sphagnum</i> stems
9	Poz-150631	90.5	280 ± 30	1505-1596 (55.0%) 1616-1665 (37.8%) 1784-1794 (2.6%)	<i>Sphagnum</i> stems
10	Poz-150633	95.5	100 ± 30	1683-1735 (26.1%) 1802-1930 (69.3%)	<i>Sphagnum</i> stems

183

184 The absolute chronology of the entire core was based on a Bayesian age-depth model using OxCal
185 v4.4.4 (Bronk Ramsey, 2021). The *P_Sequence* command with a parameter *k* of 0.1 cm⁻¹ was used to
186 calculate the model, assuming $\log_{10}(k/k_0) = 2$, and interpolation = 1 cm. The most pronounced change in
187 peat composition, as manifested by changes in pollen concentration, testate amoeba species composition
188 and species composition of plant macrofossils, which may signal changes in peat accumulation rates, was
189 input using the *Boundary* command at a depth of 66 cm. For better readability of the age-depth model, mean
190 values (μ) were introduced and used to illustrate the modeled age.

191
192 **2.4. Peat properties and carbon accumulation rate**
193 Analyses of bulk density, loss on ignition, and peat carbon accumulation rate (PCAR) were carried out
194 for each of the ninety samples. Each sample's volume [cm³] was carefully measured beforehand using
195 calipers to avoid compressing the material. Each sample was then placed in a separate crucible and dried to
196 determine the percentage of water content. The weighed and dried samples were incinerated at 550 °C for
197 12 hours and reweighed according to the protocol of Heiri et al. (2001) to determine the ash mass [g]. Bulk
198 density [g/cm³] was obtained by dividing the dry sample mass by the volume of the fresh sample according
199 to Chambers et al. (2010). Loss on ignition [g] was obtained by subtracting the ash mass from the dry sample
200 mass. Accumulation rates obtained from the peat core chronologies were multiplied by measuring the bulk
201 density without ash and by 50% to obtain the PCAR, following the protocol of Loisel et al. (2014). The top
202 eleven centimeters of the core (0-11 cm) were discarded for PCAR assessment due to the unrepresentative
203 nature of the results obtained, as increased values of carbon accumulation in near-surface peat cannot be
204 used for inference (Young et al., 2019).

205
206 **2.5. Plant macrofossil analysis**
207 The analysis of plant macrofossils was carried out using the modified protocol of Mauquoy et al., 2010.
208 Each sample of approximately 5cm³ was wet sieved (mesh diameter: 200 µm). The generalized content of
209 the sample was estimated in percentage using a binocular microscope. Fruits, seeds, caryopses, achenes,
210 perigynia, bud scales, catkin scales, whole preserved leaves, whole preserved needles, cones, anthers,
211 sporangia, opercula, fungi sclerotia, and wood pieces were counted as total numbers in each sample. The
212 tissues of monocotyledon species and moss leaves (brown and *Sphagnum* mosses) were identified on slides
213 using a magnification of ×200 and ×400. The material was compared with the guides (Anderberg, 1994;
214 Berggren, 1969; Bojňanský and Fargašová, 2007; Mauquoy and van Geel, 2007). The diagram for the
215 analyzed proxy was plotted using the riojaPlot package for R (plant macrofossils) (Juggins, 2023).

216
217 **2.6. Testate amoebae analysis**
218 Samples for testate amoeba analysis (volume: ca. 5cm³) were washed under 300 µm sieves following
219 the method described by Booth et al. (2010). Testate amoebae were analyzed under a light microscope with
220 ×200 and ×400 magnifications until the sum of 100 tests per sample was reached (Payne and Mitchell,
221 2009). Several keys and taxonomic monographs (Clarke, 2003; Mazei and Tsyganov, 2006; Meisterfeld,
222 2001; Ogden and Hedley, 1980) as well as online resources (Siemensma, 2023) were used to achieve the
223 highest possible taxonomic resolution. The results of a testate amoebae analysis were used for the
224 quantitative depth-to-water table (DWT) and pH reconstructions. Both the full diagram and the

225 reconstructions were performed in C2 software (Juggins, 2007) using the European training set (Amesbury
226 et al., 2016).

227

228 **2.7. Pollen and non-pollen palynomorphs**

229 Samples for palynological analysis (volume: 2 cm³) were prepared using standard laboratory
230 procedures (Berglund and Ralska-Jasiewiczowa, 1986). To remove the carbonates, samples were treated
231 with 10% hydrochloric acid. This step was followed by digestion in hot 10% potassium hydroxide (to
232 remove humic compounds) and soaking in 40% hydrofluoric acid for 24 h (to remove the mineral fraction).
233 Next, acetolysis was carried out. Three *Lycopodium* tablets (Batch 280521291, containing 18407 spores per
234 tablet; produced by Lund University) were added to each sample during the laboratory procedures for the
235 calculation of microfossil concentration (Stockmarr, 1971). Pollen, spores, and selected non-pollen
236 palynomorphs (NPPs) were counted under an upright microscope (Zeiss Axio SCOPE A1) until the number
237 of total pollen sum (TPS) grains in each sample reached at least 500, apart from 23 samples in which pollen
238 concentrations were very low. Sporomorphs were identified with the assistance of atlases, keys (Beug, 2004;
239 Moore et al., 1991), various publications, and the image database in the case of NPPs, for which there are
240 no atlases (Miola, 2012; Shumilovskikh et al., 2022; Shumilovskikh and van Geel, 2020). The results of the
241 palynological analysis were expressed as percentages, calculations are based on the ratio of an individual
242 taxon to the TPS, i.e., the sum of AP (arboreal pollen) and NAP (non-arboreal pollen), excluding aquatic
243 and wetland plants (together with Cyperaceae and Ericaceae), cryptogams, and fungi. The diagram for the
244 analyzed proxy was plotted using Tilia/Tilia graph software (pollen) (Grimm, 1992, 1991).

245

246 **2.8. Macro- and microcharcoal analysis**

247 Microscopic charcoal particles (size: > 10 µm) were counted from the same slides as pollen until the
248 number of charcoal particles and *Lycopodium* spores counted together, exceeded 200 (Finsinger and Tinner,
249 2005; Tinner and Hu, 2003). Microscopic charcoal influx or accumulation rates (MIC) were calculated by
250 multiplying charcoal concentrations by peat accumulation rates (PAR) (Davis and Deevey, 1964; Tinner and
251 Hu, 2003).

252 For macroscopic charcoal analysis, samples (volume: 2 cm³) were prepared by bleaching to create a
253 more visible contrast between the charcoal and the remaining organic matter following the method described
254 by Whitlock and Larsen (2001). Samples were sieved through a 500-µm mesh and only large charcoal
255 fragments > 600 µm were analyzed to obtain a local fire signal (Adolf et al., 2018). Samples were analyzed
256 with a binocular under 60× magnification. Macroscopic charcoal influx or accumulation rates (MAC,
257 particles/cm²/year) were calculated using the charcoal concentrations and PAR.

258

259 **2.9. Tree core chronology construction**

260 Tree cores underwent a standardized dendrochronological procedure (Zielski and Krapiec, 2004).
261 Polished cores were scanned between 1200 - 2400 DPI using an Epson Perfection V700 Photo scanner.
262 Annual growth rings were measured on digital images with an accuracy of 0.01mm using CooRecorder.
263 This facilitated the selection of individual growth sequences, which were utilized to form a chronology for
264 each plot. Visual comparisons were made between individual sequences, and the significance of correlations
265 was assessed using Student's t-test (Baillie and Pilcher, 1973). Subsequently, cross-dating was conducted
266 using COFECHA software (Grissino-Mayer, 2001), which evaluates each data series concerning the
267 reference chronology created and compares the correlation coefficients obtained. Raw chronologies were
268 derived by employing an arithmetic mean. For climate-growth analysis standardized chronologies were
269 used, obtained by fitting a spline function (i.e., the "n-year spline" was set at 2/3 of the wavelength of n
270 years of single growth series) using the 'dplR' package (Bunn, 2008) package version 1.7.6 (2023) in the
271 software R version 4.3.0 (R Core Team, 2022). By using this standardization method, random variation in
272 the radial growth was removed (Cook et al., 1990). For the obtained chronologies i.e., raw (TRW) and
273 standardized (RWI), values for the following descriptive statistics were computed: the mean correlation
274 between series (inter-series correlation or Rbar), the GLK index (Gleichläufigkeit; Eckstein and Bauch,
275 1969), and EPS (express population signal) (McCarroll and Loader, 2004).

276

277 **2.10. Dendroclimatological and pointer years analysis**

278 The 'chron' function from 'dplr' package allowed for the making of a residual chronology, which was
279 used for climate-growth analysis. The 'dcc' function and its moving response (25-yr window) function
280 method were used to determine the effects of climate conditions on the growth of Scots pine using the
281 'treeclim' package (Zang and Biondi, 2015) version 2.0.6.0 in R (R Core Team, 2022). This package allows
282 the use of the bootstrap procedure to test the significance and stability of the coefficients of determination
283 (r^2) over a set period (Guiot, 1991). Monthly mean air temperature (TEMP) and total monthly precipitation
284 (PREC) were used to analyze climate-growth for the period 1920-2022 (Klein Tank et al., 2002). Climate
285 data were acquired via Climate Explorer (Trouet and van Oldenborgh, 2013) and calculated from the
286 monthly gridded observational dataset E OBS v. 25.0e (Haylock et al., 2008) obtained for the 17.75-18.00°E
287 and 53.50-53.75°N grid.

288 The Becker algorithm (Becker et al., 1994) was used to determine the pointer years in the Woziwoda
289 chronology. Calculations were made using the 'dplr' package in R and the 'pointer' function (Bunn, 2008).
290 Pointer years were calculated using adjustable thresholds of relative variation in radial growth set to a 10-
291 year time window and the number of series exhibiting a similar incremental growth pattern. The main
292 criterion for determining pointer years was the occurrence of unidirectional changes (i.e., a decrease or

293 increase in the number of annual rings) in a minimum of 85% of the tested sequences of annual increments
294 observed in a group of trees at the Woziwoda site.

295

296 **2.11. Acquisition and post-processing of remote sensing data**

297 The analysis of the current state of Okoniny (Jezierzba) peatland was conducted using airborne remote
298 sensing data. The data were acquired through a multisensor aerial platform by the MGGP Aero company on
299 March 25, 2022 (leaf-off collection) and July 20, 2022, one of the warmest days of the year, which was
300 particularly important for acquiring thermal data (leaf-on collection). Multispectral images (acquired with
301 the IXM-100 camera) and Airborne Laser Scanning data (ALS; acquired with the Riegl VQ780-II scanner)
302 were obtained in the leaf-off season. Subsequently, during the vegetation season, the dataset was enhanced
303 by acquiring hyperspectral data (collected using the HySpex VS-725 scanner) and thermal data (obtained
304 with the InfraTEC 9400 camera). Based on the multispectral images, an orthophotomap was generated with
305 a Ground Sampling Distance (GSD) of 10 cm. Hyperspectral data were used to create a mosaic consisting
306 of 430 bands (in the range from 400 to 2500 nm), ALS data were applied for the development of a Digital
307 Terrain Model (DTM), and thermal data were used to produce a land surface temperature (LST) mosaic.
308 Thermal and hyperspectral mosaics and DTM were prepared with GSD = 1 m.

309 Photo interpretation was carried out to assess the extent of peatlands and the course of drainage ditches
310 using orthophotos and DTM as a base map. DTM was also used to delineate the catchment area of the
311 peatland. Hydrological modelling methods based on watershed analyses were employed for this purpose. A
312 hyperspectral mosaic was used to calculate spectral indices such as the Normalized Difference Vegetation
313 Index (NDVI; Rouse et al., 1974) and Moisture Stress Index (MSI; Hunt and Rock, 1989). Spectral indices
314 are mathematical formulas that enable the simultaneous analysis of reflectance across multiple spectral
315 ranges. The NDVI is a measure of healthy, green vegetation ranging from -1 to 1. Vegetation values
316 typically range from 0.2 to 0.8, with higher values indicating healthier and denser vegetation. The MSI index
317 is sensitive to increasing leaf water content. Its values range from 0 to more than 3, but the common values
318 for vegetation are from 0.4 to 2. Higher values indicate greater water stress and less water content in this
319 case. Thermal data was used for calculating Land Surface Temperature (LST), measured in degrees Celsius.

320

321 **2.12. Historical and cartographic information**

322 Several historical cartographic studies were used to assess changes to the peatland and its surroundings.
323 The oldest of the materials used is the Schrötter-Engelhardt map of 1803. Work on creating the map began
324 in 1796 under the leadership of the Prussian government minister Friedrich Leopold von Schrötter (1743-
325 1815) and topographer Friedrich Bernhard Engelhardt (1768-1854). The manuscript was produced at a scale
326 of 1:50,000. Still, due to the concerns of the Prussian army command about the map being too detailed and

327 capable of being used by enemy armies, a generalized version was eventually published at a scale of
328 1:150,000. A larger-scale version of the map was not available until the 1920s (Jäger, 1982, 1981). In this
329 article, the generalized version of the map is interpreted.

330 The Prussian topographic map Messtischblatt of 1874 on a scale of 1:25 000, sheet No. 982, Zalesie
331 section, was also analyzed. and the Detailed Map of Poland issued by the Military Geographical Institute in
332 1933 at a scale of 1:25,000, PAN map sheet 34 - SLUP 26 - B (Linsk). In addition, a geological-agricultural
333 map compiled between 1899 and 1900 on the topographic Messtischblatt of 1874 was considered. The
334 Prussian Geological Survey produced the map (Königlich-Preußische Geologische Landesanstalt) and
335 provides information on alluvial and diluvial deposits covering the area under study. The maps show the
336 changes in the peat bog and its surroundings from the early 19th century to the 1930s. Aerial images from
337 1964, 1984, and 1997 obtained from the Central Office of Geodesy and Cartography were also used for the
338 same purpose (license no. DIO.7211.457.2023_PL_N).

339 Insect outbreak data are based on the literature (Orłowicz, 1924; Schütte, 1893; Wilson, 2012).


340

341 **3. Results and interpretation**

342 **3.1. Age-depth model and peat accumulation rate**

343 The age-depth model showed a model agreement index (A_{model}) of 60% (Fig. 2), precisely at the limit
344 of the recommended minimum for its reliability (60% according to Bronk Ramsey, 2008). The model
345 spanned the period of ca. 282 years, with a maximum uncertainty of ca. 30 years (mostly in the section of
346 ca. 1883-1783 cal. CE). Most of the core consisted of well-preserved *Sphagnum* peat, while the lower part
347 consisted of sedge peat. The peat accumulation rate averaged 3.6 mm/yr, with the highest values associated
348 with the undecomposed acrotelm zone. The upper layers located between 0 and 11 cm were excluded from
349 the analysis of peat accumulation rates. The fastest rate was 0.71 cm/yr (at 11.5 cm), and the slowest was
350 0.1 cm/yr (at 91.5 cm). The mean BD value across the core was 0.07 g/cm³. It was highest in the lower part
351 of the core with 0.10 g/cm³ between 96 and 70 cm, and lowest in the middle part - 0.05 g/cm³, between 69
352 and 30 cm. In the upper part between 29 and 0 cm, it was 0.06 g/cm³. Similarly, this upper, undecomposed
353 layer was excluded from the peat carbon accumulation rate (PCAR) analysis. For the rest of the core (11-
354 96 cm), PCAR averaged 112 gC/m²/yr. The mean water content of the wet sample was 93.8%, and the mean
355 organic matter content of the dry sample was 95.5%.

356



357
 358 Figure 2. ¹⁴C age-depth model of the Okoniny (Jezierzba) peat profile. Water content, organic matter content,
 359 bulk density, and PCAR are also marked.
 360

361 3.2. Palaeoecological analyses

362 3.2.1. Phase 1 (~1726–1838, 96–74 cm): wet conditions and low human impact

363 The plant macrofossils and pollen analyses point to the presence of a shallow water body during this
 364 time interval. Plant macrofossil analysis (Fig. 3) showed that the peatland vegetation in this phase was
 365 strongly dominated by vascular vegetation, mainly monocotyledons with *Carex* spp. Shallow waters and
 366 edges of the water body were overgrown by sedge communities (Cyperaceae pollen) (2.8-14.5%) (Fig. 5).
 367 Additionally, this was indicated by the presence of macrophytes represented by pollen of *Potamogeton*
 368 subgen. *Eupotamogeton* (0-0.9%), *Nymphaea* (0-0.4%), and *Utricularia* (0-0.3%) (Fig. 5). The high shares
 369 of aquatic non-pollen palynomorphs (NPPs) such as cyanobacteria and the algae *Tetraëdron minimum*,
 370 *Scenedesmus*, *Botryococcus*, and *Pediastrum* (Fig. 5) confirms results of plant macrofossil and pollen
 371 analyses.


372 This phase was also characterized by the brown moss *Straminergon stramineum* (max. 9% of the
 373 subsample content) (Fig. 3). This species occurs in a wide range of habitats (Hedenäs, 1993) but is most
 374 common in wet, moderately acidic habitats (Blockeel, 2010). *Straminergon stramineum* is usually found as

375 scattered stems or small patches among other mosses but occasionally forms scattered mats, sometimes
 376 partially submerged in water, next to lakes, on the edges of peat bogs or in lakeside marshes (Hill and
 377 Blockeel, 2014).

378 This phase of peatland development was characterized by a very low concentration of testate amoebae
 379 in the samples. *Centropyxis aculeata* was the most abundant species (Fig. 4). The dominance of plagiostomic
 380 species from the genus *Centropyxis* may point to the presence of mineral input into the peatland
 381 (Lamentowicz et al., 2009a; Marcisz et al., 2020a). The water level in the peatland was quite unstable and
 382 fluctuated between 4.3 and 16.5 cm below the ground and the pH value ranged between 4.5 and 5.2, but due
 383 to the low number of identified tests, these reconstructions should be viewed with caution (Fig. 4).

384 The surrounding vegetation was characterized by the dominance of forests, as evidenced by the high
 385 proportion of arboreal pollen (AP) (83.6-91.1%) in total pollen content (TP) (Fig. 5). The main species
 386 recorded were *Pinus sylvestris* (62.6-81.3% AP) and *Betula* (6.8-16.0% AP), with admixtures of *Alnus* (2.5-
 387 7.7% AP), *Quercus* (1.8-8.1% AP), *Corylus avellana* (0.6-3.8% AP), *Carpinus betulus* (0-3.4% AP) and
 388 *Fagus sylvatica* (0.4-3.3% AP). Values of Cerealia pollen sum (0-7.8% TP) with *Centaurea cyanus*, a crop
 389 weed, indicated a stable presence of cultivated fields.

390




391
 392 Figure 3. Diagram showing macrofossil percentages, macroscopic charcoal concentrations, and influx as a
 393 local fires proxy. 10 times exaggeration is marked.


394

395 This phase also had the highest influx of macroscopic charcoal (MAC) of all three distinguished
 396 phases (Fig. 3). Towards the end of the phase, at depths of 79.5 and 78.5 cm (1st half of the 1820s according

397 to calibrated dates), influx reached the highest values throughout the core and equaled 24.5 and 11.5
 398 particles/cm²/year, respectively. The highest influx of MAC in both subsamples corresponded with the influx
 399 of microscopic charcoal (MIC), reaching over 53,200 particles/cm²/year for the 79.5 cm subsample and over
 400 125,000 particles/cm²/year for the 78.5 cm subsample (Fig. 5). This distinct fire event was followed by a
 401 slight decrease in pH, an appearance of wet indicator mixotrophic testate amoeba species (*Amphitrema*
 402 *wrightianum*, *Archerella flavum*, *Hyalosphenia papilio*), and the disappearance of cyanobacteria and algae
 403 (Fig. 4).



404
 405 Figure 4. Testate amoebae and selected aquatic non-pollen palynomorphs (*Cyanobacteria*, *Tetradion*
 406 *minimum*, *Scenedesmus*, *Botryococcus*, and *Pediastrum*) diagram. Percentages are shown in black and 10
 407 times exaggeration is marked. Testate amoeba-based depth-to-water table (DWT) and pH reconstructions
 408 as well as the sum of testate amoeba shells counted in each sample (TA sum) are presented.



409
 410 Figure 5. Arboreal pollen and microscopic charcoal diagram. Percentages are shown in black and 10
 411 times exaggeration is marked. Microscopic charcoal-based depth-to-water table (DWT) and pH reconstructions
 412 as well as the sum of microscopic charcoal counted in each sample (MIC sum) are presented.

410 Figure 5. Pollen diagram with selected taxa presented (list of taxa presented in the associated open dataset).
411 Pollen percentages are shown in black, and 10 times exaggeration is marked. Microscopic charcoal
412 concentrations and influx as an extra-local fires proxy are also presented.

413
414 **3.2.2. Phase 2: (~1838–1945, 74–37 cm): stabilization of water table and increase in acidity, a**
415 **transition from mixed forest to pine monoculture and agricultural development**

416
417 The local vegetation (Fig. 3) in this phase was dominated by *Sphagnum*, first by the subgenus
418 *Subsecunda*, then for most of this period by *Sphagnum papillosum*. *S. papillosum* occupies the more
419 oligotrophic lawns with a preference for open space (Clymo and Hayward, 1982; Laine et al., 2018). Along
420 with the appearance of *Sphagnum* from the subgenus *Subsecunda*, *Drosera intermedia* was also recorded.
421 Currently, in Poland, it is a very rare species, found in dispersed peatlands (Mirek et al., 2006). Individuals
422 often stand in the water even throughout the season. *Andromeda polifolia* also appeared in this phase.
423 Initially, the presence of *Sphagnum* was accompanied by *Straminergon stramineum* (max. 10%), but later it
424 disappeared completely. By the beginning of the twentieth century, a relatively high proportion of
425 monocotyledonous plants was also observed, represented in the samples by their epidermis, averaging about
426 20% in a sample, with a much higher proportion in the early stages. All these taxa indicate an intermediate
427 environment between a shallow lake and a moss peatland.

428 After an initial decline (from 9.2 cm at 73.5 cm, 1838 cal. CE, to 13.0 cm at 66.5 cm, 1862 cal. CE),
429 the water table level increased and stabilized at a high level, reaching a maximum of 6.8 cm at 47.5 cm,
430 1907 cal. CE (Fig. 4). The abundance of individual testate amoeba species also increased. Initially, *C.*
431 *aculeata* dominated, but later *Amphitrema wrightianum* and *Hyalosphenia papilo*, mixotrophic taxa that
432 contain endosymbiotic photosynthetic algae, begin to prevail (Lamentowicz and Mitchell, 2005a; Marcisz
433 et al., 2020a) (Fig. 4). Subsequently, the proportion of *A. wrightianum* and *H. papilo* begun to decline in
434 favour of *Archerella flavum* and *Hyalosphenia elegans* (Fig. 4). All four species are associated with the
435 presence of *Sphagnum*, with *A. flavum* and *A. wrightianum* tolerating very wet or even submerged
436 *Sphagnum* habitats, which corresponds to a stably high-water table. Then, from the mid-1880s for another
437 ca. 20 years, *C. aculeata* again became dominant. After this period, species associated with *Sphagnum*–*A.*
438 *wrightianum*, *A. flavum* and *Heleopera sphagni* – began to dominate again. During this phase, further
439 acidification of the site was noted through a drop in the pH value from the initial 4.8 to 4.1 (Fig. 4).

440 The forests surrounding the peatland (55.1–92.7% TP) were still dominated by pine (64.5–92.8% AP),
441 although their percentage has decreased in comparison to phase 1, especially during the 1920s and 1930s
442 (Fig. 5). Deciduous taxa such as *Quercus*, *Corylus avellana*, *Carpinus betulus* and *Fagus sylvatica* retreated.
443 The percentage of Cerealia in the TP increased significantly, from 0–7.8% TP in the first phase to 2.8–19.8%

444 in the second phase, with a peak in the late 1910s and early 1920s, indicating the development of agriculture
445 in the vicinity of the peatland (Fig. 5). Around the same time, the proportion of *Rumex* also increases
446 significantly (0-11.5%). The low values of MAC (Fig. 3) and MIC (Fig. 5) indicate a low fire activity in the
447 studied area.

448
449 **3.2.3. Phase 3: (~1945–present, 37–0 cm): Lowering of the groundwater table, further afforestation**
450 **with *Pinus sylvestris*, a succession of *Betula***

451
452 The local vegetation (Fig. 3) underwent several changes during this phase. Although *Sphagnum*
453 dominated for the entire time, the subgenus *Sphagnum* receded in favour of first the subgenus *Cuspidata*
454 and then the subgenus *Acutifolia*. The beginning of the phase was marked by *Pohlia nutans*, which can win
455 the competition in unstable habitat conditions, such as during the dry season (Boulc'h et al., 2020). Its
456 occurrence correlated with the presence of *Phryganella acropodia* among testate amoebae (Fig. 4), which
457 is an indicator of low water levels in *Sphagnum* peatland (Diaconu et al., 2017; Lamentowicz and Mitchell,
458 2005b).

459 This was followed by *Alabasta militaris* ($\bar{x} = 25.5\%$), *Galeripora discoides* ($\bar{x} = 10.5\%$) and *Nebela*
460 *tincta* ($\bar{x} = 8.2\%$) beginning to dominate (Fig. 4). *G. discoides* is typically present in acidic sites with
461 unstable hydrological conditions (Lamentowicz and Mitchell, 2005b; Sullivan and Booth, 2011). *N. tincta*
462 tolerates dry, highly acidic conditions with mineral matter supply (Booth, 2002; Koenig et al., 2018;
463 Lamentowicz et al., 2011). *A. militaris*, dominant in recent years, is indicative of dry and markedly acidic
464 conditions (Amesbury et al., 2016; Booth, 2002; Lamentowicz et al., 2011; Marcisz et al., 2020a; Sullivan
465 and Booth, 2011). Based on testate amoebae, this phase was distinguished by a significant drop in the
466 groundwater table, from an average level of 9.6 cm below the ground surface in the second phase to 15.7
467 cm. In the last decade, the most significant decline was observed, with an average level of 21.9 cm, with a
468 maximum of 27.5 cm, 1983 cal. CE. The pH continued to decrease – from 4.4 to 4.0 (Fig. 4).

469 On a regional scale, there is an increase in the relative abundance of *Pinus* pollen in the TP, from about
470 46% at the beginning of the phase to about 85% today as an effect of afforestation (Fig. 5). *Betula* pollen
471 share has an apparent increase, from 0,7-11,3% in the second phase to 5,6-32,5%. The increased percentage
472 of *Betula* pollen, combined with macroscopic remains in the form of achenes and catkin scales, indicates
473 the intensive succession of this species on the peatland surface. The ruderal species *Urtica* and *Artemisia*
474 were also more strongly manifested. The average proportion of *Urtica* pollen in the TPS increased distinctly
475 (from 0-0.7% to 0-2.9%). The percentage of Cerealia in TP has decreased significantly, from nearly 20% in
476 the early 1920s to just over 1% today.

477 Local (Fig. 3) and regional (Fig. 5) fire activity continued to be low, although two slightly more
 478 intensive periods of regional fires were marked – ca. 1945-1963 and the early 2020s.

479

480 3.3. Dendrochronological and pointer years analysis

481 A total of 50 tree-ring series of 23 *Pinus sylvestris* L. trees from the Woziwoda site were successfully
 482 cross-dated. Based upon the TRW (Fig. 6) and RWI sites, well-synchronised tree-ring series spanning 222
 483 years (1801-2022) was developed. The statistical characteristics of the ring-width series and the statistical
 484 parameters indicating the signal strength of the regional RWI chronology are shown in Tab. 2. The mean
 485 EPS was 0.93, which is well above the threshold value (EPS = 0.85) required to produce a statistically robust
 486 RWI chronology. Mean series inter-correlation, MS, SNR, and other statistical parameters indicating the
 487 strength of chronology signals were also high, indicating the suitability of chronology for climate-growth
 488 analysis.

489 Tab. 2 Descriptive statistics of standardized *Pinus sylvestris* L. (RWI) chronology for Woziwoda site

Chronology length	1801-2022
Mean tree age [yrs]	197
Number of tree/cores	23/50
Mean ring width (mm) ± SD	1.256 ± 0.702
Series intercorrelation	0.623
Average mean sensitivity	0.265
Expressed Population Signal (EPS)	0.93
Signal-to-noise ratio (SNR)	12.97
Rbar.eff (effective chronology signal)	0.361


490

491 Across the study period (1920-2022) a significant positive relationship between growth and
 492 February mean temperature was identified (Fig. 7). The moving correlation analysis showed an increasing
 493 trend in the sensitivity of tree growth to climatic factors (Fig. 8). The positive response of tree growth to
 494 February mean temperature remained constant throughout the study period (1920-2022) (Fig. 8). However,
 495 the sensitivity of tree growth to summer temperature increased. The relationship between annual growth and
 496 summer temperature was not stable during the period 1920-2022. Nevertheless, in the last 30 years, a
 497 significant negative relationship between annual growth and June mean temperature was observed.

498 Climate-growth analysis for monthly data did not show a statistically significant relationship
 499 between growth and precipitation (Fig. 7). However, moving response analysis revealed significant short-
 500 term relationships between tree growth and precipitation. Furthermore, it was demonstrated that the
 501 influence of precipitation in the current year's months on tree growth calculated for the years 1960-2022


502 was more significant than the relationships calculated for the years 1921-1959. In recent years, a particularly
 503 positive relationship between tree growth and early-year (February-April) precipitation as well as June
 504 precipitation has become apparent.

505 For Woziwoda site 8 positive and 13 negative pointer years were identified for the period 1814-
 506 2022 (with a minimum sample depth of 10 trees) (Fig. 6). The most pronounced positive pointer years with
 507 more than 90% tree response were as follows: 1847, 1863, 1912, 1941, 1945, 1957, and 1983. The most
 508 pronounced negative pointer years were: 1839, 1868, 1869, 1911, 1925, 1940, and 1950. Figure 6 provides
 509 marks of pointer years together with meteorological and ecological characteristics.




510
 511 Figure 6. The grey lines depict the individual tree ring series of each tree, while the green line represents
 512 the average raw chronology of *Pinus sylvestris* L. at the Woziwoda site. Identified within the Scots pine

513 chronology from Woziwoda are pointer years, categorized as negative (NEG) (A) and positive (POS) (B).
 514 These pointer years are highlighted with colored asterisks: red for positive pointer years and blue for
 515 negative pointer years. The position of the asterisks refers to a scale of 0-100%. Information on extreme
 516 phenomena is based on Orłowicz, 1924; Schütte, 1893, Broda 2000, Wilson 2012.




517
 518 Figure 7. Response function coefficients between residual *Pinus sylvestris* L. chronology and climate
 519 variables: (A) mean air temperature (Temp), and (B) precipitation (Prec) for the period 1920–2022. Names
 520 of the previous year’s months start with a lowercase letter. Solid lines represent significant coefficients at p
 521 < 0.05.

(A) Temp



(B) Prec




522
 523 Figure 8. Moving response correlations (25-year window) between residual *Pinus sylvestris* L. chronology
 524 and climate variables: (A) mean air temperature (Temp), and (B) precipitation (Prec) for the period 1920–
 525 2022. The color code represents the response function coefficients. Significant correlations are indicated by
 526 white asterisks.


527

528 **3.4. The current state of the peatland based on remote sensing data analysis**

529 Presently, the non-forested part of the peatland is drained by two parallel ditches. One is located in
530 the northern, and the other is in the southern non-forested part of the peatland. The analysis of thermal data
531 obtained on a midsummer day indicates that the average LST for the non-forested part of the peatland is
532 approximately 34.29 °C, with a temperature range extending from 19.22 °C to 46.37 °C. There is a distinct
533 internal variability in LST values within the studied area. Higher values, indicative of more significant
534 dehydration, were identified in the eastern part of the peatland, while lower values were observed in the
535 western part. A repeating spatial pattern of values was observed in the analysis of vegetation indices (NDVI
536 and MSI). High NDVI values and low MSI values, indicative of good vegetation condition and low water
537 stress, were observed in the western and southwestern parts of the peatland (Fig. 9). The average NDVI
538 value in these areas is 0.71, and MSI is 0.6. Conversely, low NDVI values and high MSI values, indicative
539 of significant dehydration of the peatland and low vegetation vigor, were observed in the eastern part of the
540 object (Fig. 9), where NDVI averages 0.63, and MSI is around 0.69. The overall average NDVI for the
541 object was 0.65, and for MSI, it was 0.68.



542
543 Figure 9. Remote sensing characteristics of Okoniny (Jezierzba) peatland based on multisensorial airborne
544 data acquired in 2022.
545



547
 548 Figure 10. Changes in the peatland and its surroundings since the beginning of the 19th century based on
 549 historical maps and aerial images. (A) Schrötter-Engelhardt map 1:150 000 (1802), (B) Messtischblatt map
 550 no. 982 1:25 000 (1874), (C) Prussian geological and agricultural map no. 2374 1:25 000 (1905), (D)
 551 Detailed Map of Poland 1:25 000 (1933), (E) Aerial photograph from 1964, (F) Aerial photograph from
 552 1984, (G) Aerial photograph from 1997. Maps no. A, B, C, and D are in the public domain. Aerial
 553 photographs were obtained from © Central Office of Geodesy and Cartography in Poland, license no.
 554 DIO.7211.457.2023_PL_N.

555
 556 Analysis of historical materials (Fig. 10), including maps and airborne images, confirms the results
 557 of the palaeoecological analysis. Both the Schrötter-Engelhardt map of 1802 and the Messtischblatt of 1874
 558 indicate the existence of a small lake in the coring area. Again, however, it should be noted that the Schrötter-
 559 Engelhardt map is a highly generalized study and does not give much information about the surroundings

560 of today's peatland, other than that we are dealing with an area with the character of a dense forest complex
561 with wetlands in isolated places. Messtischblatt allows us to better interpret the surroundings of the analyzed
562 modern peatland at the time in which the map was prepared. A small lake named "Kolze See" is observed
563 in an advanced stage of development, i.e., progressive overgrowth. This lake is located in the surroundings
564 of wetlands (Bruch in German) somewhat distant heathland (Heide in German) and wasteland (Ödland in
565 German) (the original nomenclature of the map legend was adopted). This lake and two other lakes close
566 by are enclosed within a single catchment area. To the south, the area of the current peatland was adjacent
567 to an open, extensive meadow.

568 Even more information is provided by a 1905 geological-agricultural map prepared on the
569 topographic base Messtischblatt map of 1874. In addition to land use, it shows the type and thickness of
570 alluvial and diluvial deposits. According to this map, the area around the lake was covered by alluvial
571 sediments – humus with peat subsoil and shallow groundwater (org. in German: Humus (Peat) mit Torf-
572 Untergrund und nahem Grundwasser). The thickness of the peat was marked at two meters. However, it
573 should be noted that drilling surveys at that time only covered a maximum depth of two meters, so the maps
574 do not provide information on the total thickness of the sediments (Jasnowski, 1962). Places that were used
575 as heathland and wasteland on the topographic map are covered by sandy humus on a sandy substrate with
576 shallow groundwater (org. in German: Sandiger Humus mit Sand-Untergrund und nahem Grundwasser) and
577 by humic sands on a substrate of permeable sands with shallow groundwater (org. in German: Humoser
578 Sand mit durchlässigen Sand-Untergrund und nahem Grundwasser).

579 A Detailed Map of Poland from 1933 documents the change in an ecosystem from lake to land. The
580 area, which on Prussian topographic maps was a lake with a surrounding bog, is described as a meadow on
581 this map. Moreover, the meadows adjacent to the south were marked with drainage ditches, which were not
582 on the Prussian maps. The area's surroundings, as before, were dominated by coniferous forests.

583 Aerial photos document subsequent changes in the ecosystem. The 1964 photo shows the northern part
584 of today's peatland's agricultural use (regular surface layout). Lake Kały, located nearby, became completely
585 overgrown, and its area was later dug by a drainage ditch, brought to the studied peat bog. The surrounding
586 area of the peatland is dominated by dense forest with occasional open clear-cutting areas. A photo from
587 1984 documents the succession of trees in the north-central part of the peatland. In the surrounding area,
588 open forest areas have entirely disappeared. A photo from 1997 clearly shows the development of trees on
589 the peatland, which have formed a dense block in its north-central part. A distinct area of *Sphagnum*-
590 dominated peatland with a well-marked edge has also emerged. Currently, the northernmost part of the
591 peatland is overgrown by pine; it is almost impossible to identify the maximum extent of the peatland surface
592 in the field (Fig. 1).

593

594 4. Discussion

595 4.1. Exceptionally high peat accumulation rate

596 In the Okoniny (Jezierzba) peatland, a rapid rate of peat accumulation is observed, averaging 3.56
597 mm/yr, with a maximum value of 7.1 mm/yr at a depth between 11 and 12 cm. This accumulation rate is not
598 commonly observed. There are only few peatlands in Poland for which higher accumulation rates were
599 reported. In the Tuchola Pinewoods, these were Dury – 10 mm/yr (Pawlyta and Lamentowicz, 2010),
600 Mukrza – 4.6 mm/yr (Lamentowicz and Obremska, 2010), Jelenia Wyspa mire where the accumulation rates
601 reached 0.4 mm/yr for the first 3000 years but accelerated to 3 mm/yr in the last 150 years (Lamentowicz
602 et al., 2007), and the Tuchola kettle-hole bog – 1.2 mm and after ca. 1320 cal. yr BP the accumulation rate
603 dropped to 0.4 mm/yr (Lamentowicz et al., 2008b). In other pine monocultures, such as the Noteć Forest,
604 the Rzecin peatland stands out for its high accumulation rate – an average of 6.8 mm/yr in one profile and
605 7.5 mm/yr in the other one (Milecka et al., 2017). Peatlands in Tuchola Pinewoods, including Okoniny
606 (Jezierzba) peatland, generally have a faster accumulation rate than peatlands located in other parts of
607 Pomerania, especially small kettle-hole peatlands that accumulate carbon the fastest of all peatland types
608 (Karpińska-Kołodziej et al., 2024). In Pomeranian peatlands, the highest accumulation rates were reported
609 for the period between ca. 150-1230 AD and reached 2.2 mm/yr in Stążki (Lamentowicz et al., 2008a), and
610 1.38 mm between 1830 and 2006, although the highest accumulation rate was 5 mm/yr (during AD 840-
611 860) in Słowińskie Błota raised bog (Lamentowicz et al., 2009b). At the Gołębiewo sites the maximum
612 accumulation rates were 1.85 mm/yr and 0.36 mm/yr (Pędziszewska and Latałowa, 2016). For many
613 *Sphagnum*-dominated peatlands in other parts of Poland, the average PAR varied between 1.4-2.5 mm/yr
614 (Gałka et al., 2015; Lamentowicz et al., 2020; M. Lamentowicz et al., 2015; Marcisz et al., 2020b). Such
615 high accumulation rate values are also rare in other parts of the temperate climate zone of Europe. Teici bog
616 (Latvia) showed similar accumulation rates - 3.5 mm/yr - from 1835 to 1965 AD and 10 mm/yr after 2000
617 (Stivrins et al., 2018). Okoniny (Jezierzba) peatland after 2000 (between 21.5 and 11.5 cm) recorded an
618 accumulation of 5.7 mm/yr. Saxnäs mosse in Sweden showed an almost linear peat accumulation rate of 2-
619 2.5 mm/yr (van der Linden et al., 2014). The maximum accumulation was recorded at around 2310-2250
620 cal on the Estonian Hara bog. BP (31-15 cm) reaching 2.4 mm/yr (Łuców et al., 2022). A comparison with
621 other regions of Poland and Europe shows that the exceptionally high accumulation rates at the analyzed
622 site are worth highlighting.

623

624 4.2. Relationships between forest management and pollen analysis

625 4.2.1. The complex history of the Tuchola Pinewoods and its influence on the forest

626 The results of pollen analysis of the collected core enabled us to illustrate how the forest was managed
627 over the past 300 years. Due to political changes and several administrative decisions, the management

628 strategies of the Tuchola Pinewoods underwent vital changes. The consequences of the implementation of
629 forest management techniques were visible in the palaeoecological record.

630 With the first partition of Poland in 1772 by Prussia, regulations for planned forest management began
631 to be introduced. The main planting species was Scots pine, which over time began to dominate the forest,
632 replacing deciduous admixture species (Broda, 1993). The region's forest cover and forest composition were
633 also affected by later political and administrative developments. For more information on the history of
634 forest management in the late 18th and early 19th centuries, see Supplementary File 1.

635 Our data confirm an increase in the proportion of pine pollen in the forest composition and a decrease
636 in the proportion of pollen of other species. From the 1730s to the mid-1860s, the share of pine pollen in
637 the pollen of all trees increased from about 60% to about 90%. Our pollen diagram shows the rapid increase
638 in *Pinus sylvestris* pollen percentage after 1850. It can, therefore, be assumed that this resulted from *Pinus*
639 *sylyestris* introduced by mass monoculture plantings in the early 1830s reaching reproductive capacity. Pine
640 usually reaches sexual maturity between 10 and 15 years (Sullivan, 1993), although the threshold age has
641 been set at 25 years (Matthias and Giesecke, 2014). The decline in the share of deciduous species and the
642 increase in the share of Scots pine in the landscape began in Poland with the formation of the state. However,
643 at that time, it was associated with the expansion of agriculture and the harvesting of preferred species such
644 as *Carpinus betulus* (Czerwiński et al., 2021) Nevertheless, in the Prussian partition, planned forest
645 management permanently changed the composition of Poland's largest forest complexes, which were
646 dominated by easy-to-grow pine (Broda, 1993) (see Supplementary File 1). A dynamic increase in the share
647 of pine pollen until the 1860s in the Tuchola Pinewoods was also recorded at the Czechowskie Lake
648 (Słowiński et al., 2019). An increase in pine pollen percentage since the 19th century was also shown in
649 pollen diagrams of other sites from Pomerania – Stażki (Lamentowicz et al., 2008a), Słowińskie Błota
650 (Lamentowicz et al., 2009b) – and in other monoculture plantation complexes from the Prussian partitioning
651 area – Rzecin peatland in the Noteć Forest (Milecka et al., 2017).

652 Although attempts were undertaken to correct earlier mistakes, this did not stop the massive
653 deforestation (among other consequences of war events and administrative regulations on settlement, more
654 in Supplementary File 1). Until the 1870s, the feudal system was still mixed with capitalist components, but
655 from the 1870s onward, under monopoly capitalism, timber trade and processing began to reach a significant
656 size (Broda, 2000). However, it has been noted that forests regulate air temperature, store water in the soil
657 more efficiently, and reduce wind speed, preventing soil erosion, which can help local agriculture face
658 difficult environmental conditions (Wilson, 2012). For this reason, as early as the 1870s, the state
659 administration encouraged landowners to protect forest stands on their lands and establish forestry
660 cooperatives. The government also guaranteed funds for the reforestation of private and municipal lands. In
661 the mid-1870s, the Landtag set aside a budget for the purchase and reforestation of wasteland by the state.

662 However, these funds were used to a small extent, although this somewhat reduced the share of forested
663 private property (Broda, 2000; Wilson, 2012). In 1886, the Royal Settlement Commission (in German:
664 Königliche Ansiedlungskommission) was established to buy up the estates of impoverished Polish nobility
665 to acquire agricultural land for German settlers (Wilson, 2012).

666 At the end of the 19th century, Tuchola Pinewoods became the largest timber production hub in the
667 Prussian partition. The Bydgoszcz timber industry region also played a major role in wood processing. The
668 first steam sawmill in the Bydgoszcz region was built in 1873, and by 1913, there were 20 of them,
669 processing some 500,000 m³ of wood and employing more than 1,600 people (Broda, 2000). All this resulted
670 in a significant decline in the share of tree pollen in the total pollen share in our diagram, to less than 60%
671 by the late 1920s and early 1930s. At the same time, we have seen intensive agricultural development. At
672 Okoniny (Jezierzba), the proportion of Cerealia pollen doubled between ca. 1900 and 1920. This trend is
673 also confirmed by pollen data from the site in Okoniny Nadjeziorne, on the other side of Okonińskie Lake
674 (Tipton, 2023), as well as from Czechowskie Lake, about 25 km northeast of our site (Słowiński et al.,
675 2019). Despite intensive deforestation in general, further afforestation with pine was also progressing. In
676 1893, pine forests accounted for 99% of all forests in Tuchola County (Szwankowski, 2005). Intense
677 changes in forest management (pine dominance) and agricultural development (high percentage of Cerealia
678 pollen) in the 19th century are also evident in records of profiles outside large, dense forest complexes –
679 Kusowskie Bagno (Gałka et al., 2014), Linje mire (Marcisz et al., 2015).

680

681 **4.2.2. Impact of forest management on peatland vegetation**

682 As a result of changes related to forest management, the lake-to-peatland transition occurred rapidly.
683 We assume that this was primarily the result of drainage, which was undertaken in the area at the end of the
684 19th century (see drainage ditches on the southern side and a dike in the middle part of the site on maps in
685 Figure 6), and secondly, to a lesser extent, the transition from mixed forests to pine monoculture. These
686 activities contributed to an increase in the acidity of the peatland. Forest drainage is often associated with
687 the acidification of surface waters (Miller et al., 1990). The introduction of forest drainage, on or near
688 peatlands, to improve tree growth has been quite common in northern and northeastern Europe (Westman
689 and Laiho, 2003). The oxidation of organic sediments and the detachment of H⁺ ions increase acidity
690 (Ulrich, 1980). In addition, the supply of alkaline cations to the peat is impeded by drainage ditches
691 (Minkkinen et al., 2008). However, the long-term consequences of drainage are devastating to peatlands, as
692 they initiate vegetation succession, in which species typical of peatlands are replaced by forest vegetation
693 (Laine et al., 1995). In the example of our palaeoecological data, the dynamic succession of pine and birch
694 in the Okoniny (Jezierzba) peatland is evident, which is also supported by aerial imaging. As already
695 mentioned, the successive decline in pH is also the result of the impact of pine plantations growing in

696 catchments. A drop in pH in Okoniny (Jezierzba) has likely enabled the rapid growth and expansion of
697 *Sphagnum* and the peatland initiation. The crowns of forests, especially the needles, can increase the uptake
698 of atmospheric pollutants such as sulfur and nitrogen components, contributing to the acidification of surface
699 waters (Nisbet, 2001; Reynolds et al., 1994). Conifers also can capture ions of marine origin - Na and Mg
700 cations. These, in turn, displace hydrogen and aluminium cations from the soil, leading to acid runoff from
701 the forests along with surface runoff, which is known as the "sea-salt effect" (Drinan et al., 2013; Harriman
702 et al., 2003; Reynolds et al., 1994). We observed the presence of *Pinus* needles at the beginning of phase 2
703 (from 1838 cal. CE), at the transition from pond to peatland ecosystem. Moreover, *Pinus* stomata were also
704 present in palynological samples at that time, pointing to more frequent needle falls. The presence of *Pinus*
705 stomata has been suggested as a possibly important proxy for insect outbreaks in palaeoecological records
706 in previously published studies from another Pine monoculture forest in Poland, the Noteć Forest (Barabach,
707 2015), where this phenomenon has been observed (Słowiński et al., 2019). More pine trees in the Tuchola
708 Pinewoods resulted in much higher amounts of needles and other pine fragments accumulating on the forest
709 ground, leading to soil acidification. This, together with drier conditions, could quickly lead to acidification
710 around the pond, forming perfect conditions for *Sphagnum* to encroach – first as a floating mat that
711 successively overgrown the pond. We sampled the peat core close to the edge of the peatland, thus in the
712 place where moss encroachment on the open water body began; therefore, we were able to track this
713 succession in our record. This succession and disappearance of Lake Kolze are also clearly visible in
714 historical maps (Figure 10). Other examples of quick encroachment of floating mats on the surface of the
715 lake have been observed and mapped in other open water bodies in the Tuchola Pinewoods (Kowalewski,
716 2003; Kowalewski and Milecka, 2003) and other regions (Warner, 1993).

717

718 **4.3. Anomalies and extreme events**

719 **4.3.1. The impact of droughts and fires on the forest and peatland**

720 Historical sources indicate that in the 18th and 19th centuries, the Tuchola Pinewoods were relatively
721 often affected by droughts resulting in fires (Wilson, 2012). In 1781, there was a fire in Tuchola (ca. 16 km
722 SW), during which a large part of the city with the church and town hall burned down, and in 1792, Starogard
723 Gdański (ca. 42 km NE) burned almost to the ground (Orłowicz, 1924). Major fires also occurred in 1794,
724 and 1807, when more than 34,000 hectares of forest burned (Orłowicz, 1924; Schütte, 1893). Fires in 1809,
725 1810, 1812, 1813 and 1828 in the Świt forest district about 15 km from the study site were also recorded
726 (Cyzman, 2008). Palaeoecological data, especially MIC, confirm high fire activity in the first decades of the
727 19th century (a rapid increase). Słowiński et al. (2019) emphasized that data on fires before the 1830s,
728 especially regarding their area, should be treated with caution due to the lack of accurate measurement
729 techniques. In the Woziwoda Forest District, within which the Okoniny (Jezierzba) peatland is located, the

730 forests of the Biała and Barłogi forest districts also burned in 1842 (Cyzman, 2008). Intense fires also
731 appeared in the Tuchola Pinewoods between 1846 and 1848 (Orłowicz, 1924; Schütte, 1893).

732 Later, numerous fires were also reported in the Woziwoda Forest District. Between 1860 and 1889,
733 310 fires were observed, destroying 4206 hectares of the forest (Orłowicz, 1924; Schütte, 1893). The highest
734 number of fires in this period was registered in 1862-1864 and 1874-1875 when 3565 hectares of forest
735 burned; altogether, nearly 85% of the area burned in 1860-1889 (Schütte, 1893). The largest area burned in
736 1863 equaled 2333 hectares, including more than 1250 hectares in the Woziwoda forest district; altogether,
737 25% of all the forest burned in 1860-1889 (Orłowicz, 1924; Schütte, 1893). Meteorological data confirm
738 dry years in the period from 1862 to 1865. In 1862 and 1863, the annual precipitation in Bydgoszcz was
739 only a little over 450 mm (Kirschenstein, 2005), and it was then that the largest number of hectares of forest
740 in the known history of the Tuchola Forest burned (Dietze et al., 2019).

741 The number of fires can also be linked to political events (Orłowicz, 1924; Schütte, 1893; Wilson,
742 2012). In 1901, in the nearby Trzebciny and Gołębek Forest Districts, a fire consumed 663 hectares of forest
743 (there was a parallel children's strike in Września Province) (Orłowicz, 1924; Wilson, 2012). Fires could
744 also be caused by agricultural activities and land preparation for crops (Poraj-Górska et al., 2017). By the
745 1830s, charcoal production was widespread (McGrath et al., 2015), and forest burning was used to create
746 heathlands for beekeeping (Bienias, 2009).

747 Fires of the 1860s provide a regional signal at another site in the Tuchola Pinewoods – Czechowskie
748 Lake (Dietze et al., 2019). Increased fire activity in the mid-19th century was also observed at the Lake
749 Jaczno site (Poraj-Górska et al., 2017). At the Okoniny (Jezierzba) peatland, MIC and MAC values
750 decreased after 1850, but at the same time, the water level stabilized and remained high. Fire activity
751 remained low in areas where wet conditions prevailed, such as southern Finland (Väliiranta et al., 2007) and
752 eastern Estonia (Sillasoo et al., 2011).

753 In 1948, about 450 hectares of forest were burned near Osieczna, and in 1954, 80 hectares were burned
754 near Ocypel (Cherek, 2007). Palaeoecological data record an increased MIC supply during this period. The
755 first of these fires was also recorded in the sediments of Czechowskie Lake (Słowiński et al., 2019). The
756 summer drought of 1921 occurred over a larger area of Europe, from Poland and the Czech Republic to the
757 UK (van der Schrier et al., 2021). Summer droughts also affected the Tuchola Pinewoods in 1951 and 1959.
758 In 1959 Bydgoszcz received only 37 mm of precipitation from August to October (Mitosek, 1960), and from
759 1950 to 1958 Bydgoszcz received less than 500 mm of rain per year (Kirschenstein, 2005). Our
760 palaeoecological data confirm droughts in the 1950s. There is a sharp increase in the proportion of
761 *Phryganella acropodia* among the testate amoebae, an indicator of dry conditions (Diaconu et al., 2017) and
762 an expansion of brown mosses in the form of *Pohlia nutans* (up to 30% of the peat sample composition) are

763 also marked. Dendroclimatic data recorded the negative impact of climatic conditions on pine, especially
764 strongly in 1950 and 1956.

765 Studies show that particle size illustrates the distance of the fire from the site, the heavier the particles,
766 the shorter distances they travel (Clark, 1988; Peters and Higuera, 2007). However, many factors determine
767 the particles' transport—the fire's intensity, the burning areas, and the wind direction. Adolf et al. (2018) point
768 out that the charcoal source area of occurrence of both MIC and MAC can reach a radius of 40 km. However,
769 it is often assumed that MAC indicates fires that occurred up to 1-3 km (Clark, 1990; Higuera et al., 2007;
770 Oris et al., 2014). The distances to which particles move are also determined by terrain and vegetation. They
771 move longer distances on flat terrain covered with grasses (Woodward and Haines, 2020), while they move
772 shorter distances in dense forests (Kelly et al., 2013; Oris et al., 2014). In this context, it should be assessed
773 that the local fire activity in the studied peatland was low, with an average of 0.36 particles/cm³/year,
774 although from historical sources, fires are known to have occurred nearby.

775

776 **4.3.2. Insect outbreaks and their impact on pine monoculture**

777 Palaeoecological studies based on the presence of insect head capsules and/or faeces, as well as other
778 insect remains could be helpful, but these methods are rarely used (Bhiry and Fillion, 1996; Lavoie et al.,
779 2009; Simard et al., 2006; Waller, 2013). Often the main obstacle to performing this method is bad
780 preservation of insect remains in peat. In the Okoniny (Jezierzba) peatland, we found no insect remains,
781 even though quite a large sample volume has been analyzed for the plant macrofossil analysis. Therefore,
782 we can interpret the effect of insect outbreaks using other sources of evidence.

783 The earliest information on insect outbreaks from the forests of the Tuchola Pinewoods under planned
784 forest management dates back to 1836-1838. An infestation of *Panolis flammea* occurred at that time
785 (Schütte, 1893). The insects also attacked between 1866 and 1868. As a result of this infestation, 1380
786 hectares of forest were destroyed in the Woziwoda forest district alone (Schütte, 1893). The pollen diagram
787 from the Okoniny (Jezierzba) peatland documents the phenomenon in the 1860s with a decrease in *Pinus*
788 *sylvestris* pollen and an increased presence of *Pinus* stomata that may indicate the event of the insect
789 outbreak (Barabach, 2015). The needles that fell were partially decomposed and carried downwind to the
790 peatland, where they were preserved (Słowiński et al., 2019). The same effect was noted in another closely
791 located peatland in Okoniny Nadjeziorne, where the 1866-1868 infestation also corresponds with increased
792 numbers of *Pinus* stomata (Tipton, 2023). In 1855, *Lymantria monachal* appeared in large numbers but
793 damaged only some of the younger stands (Schütte, 1893).

794 A serious incidence of *Panolis flammea* gradation also occurred in 1922-1924 (Kielczewski, 1947;
795 Mokrzecki, 1928). Between 1978 and 1985, with a peak in 1982, the forests of the northern part of the
796 country were overrun by *Lymantria monacha*, and this was the largest infestation since the establishment of

797 the National Forests in 1924, with salvage treatments covering more than 6.3 million hectares of forest over
798 seven years (Broda, 2000; Jabłoński, 2015; Śliwa, 1989, 1987). Both major gradations are reflected in
799 palynological data, manifested by declines in the pollen percentage of trees, primarily *Pinus* and *Picea*. A
800 decrease in conifer pollen during the gradation period has also been shown by studies of other sites in the
801 Tuchola Pinewoods (Łuców et al., 2021; Tipton, 2023). Other pine monoculture in Poland, the Noteć Forest
802 was also affected by gradation in 1922-1924, and this event manifested itself in palaeoecological data
803 (Barabach, 2015; Lamentowicz et al., 2015; Milecka et al., 2017). Among other things, Barabach (2015)
804 noted an increase in *Glomeromycota* fungal spores, which according to this author may indicate intense soil
805 erosion caused by the felling of dead trees, and a marked increase in *Calluna* and *Poaceae* indicating an
806 increase in the openness of the landscape. Lamentowicz et al. (2015) noted an increase in mineral content
807 in the sediment as indicated by *Centropyxis platystoma*, which was confirmed by XMT analysis of the peat.
808 Milecka et al. (2017) described higher ash and charcoal content in the sediments. Although the Tuchola
809 Pinewoods and the Noteć Forest are in the region of highest risk of outbreaks, other areas of Poland were
810 also affected, such as the Kampinos Forest in 1972 (Śliwa, 1974), or over the last decade, the Białowieża
811 Primeval Forest (Grodzki, 2016; Kamińska et al., 2021).

812 It's difficult to assess unequivocally whether the gradations affected the immediate vicinity of the
813 peatland, or whether this is a regional signal. Historic maps could be helpful, but these usually do not show
814 the difference between old and new plantings (Barabach, 2012). However, dendrochronological data
815 obtained from pine trees could help to reconstruct the extent of the outbreak. The main problem in
816 monoculture forests though is that the forest is successively cut and new trees are planted regularly.
817 However, for our dendrochronological record, we were able to obtain samples from the oldest pine trees in
818 the area. The oldest trees in the region analyzed in this study were planted over 200 years ago in the close
819 vicinity of the Woziwoda Forest District, after the introduction of the Prussian forest management strategies,
820 and have been kept there by foresters for obtaining tree saplings and for monitoring. The influence of insect
821 outbreaks has been recorded in these pine trees and we were able to track all the outbreak events in the
822 wood. The first years after the gradations - 1839, 1869, 1925, and 1982 - manifested very strongly in the
823 dendrochronological data as negative indicator years.

824

825 **4.4. Current condition of the peatland vs. remote sensing and dendroclimatic data**

826 The assessed growth reactions of pine trees to climate factors at the Woziwoda site may be considered
827 typical. The effect of February air temperatures on Scots pine growth in northern Poland was previously
828 noted (Cedro, 2001; Cedro and Lamentowicz, 2011; Feliksik and Wilczyński, 2009; Koprowski et al., 2012,
829 2011; Matulewski et al., 2019; Zielski, 1996; Zielski et al., 2010; Zielski and Sygit, 1998). Although the
830 studied pines from Woziwoda showed a similar growth response to climate as other pines from northern

831 Poland, their climate sensitivity was greater. The highest negative correlation for pine radial growth from
832 the Woziwoda site was found with July's mean air temperature.

833 Another factor commonly affecting the radial growth of Scots pine, according to the literature, is pluvial
834 conditions in February. This linkage was identified by Cedro (2001), Feliksik and Wilczyński (2009),
835 Koprowski et al. (2011) in the Pomeranian region (Northern Poland). The present study confirmed a short-
836 term relationship between pine radial growth and precipitation sums in February (Fig. 7). Late February and
837 early March are when additional water is required due to the initiation of biochemical processes in trees
838 (Przybylski, 1993). Additionally, in our study, a stronger dependence of pine radial growth on precipitation
839 was demonstrated in June. A similar result for pine from northern Poland was obtained by Matulewski et al.
840 (2019), Zielski and Barankiewicz (2000), where pine growth was threatened by a water deficit in the summer
841 season. Increased pine demand for water occurs in June and July, the months of the most intense growth
842 (Obmiński, 1970). At the same time, these are the months when droughts have become more frequent in
843 recent years (Łabędzki, 2004; Spinoni et al., 2018). Our results confirm that within the temperature and
844 monthly precipitation values typically observed in Central Europe, the primary environmental factor
845 influencing the diversity of species growth in the near future will be the availability of water (Boczoń et al.,
846 2017; Taeger et al., 2013). This availability is determined by the precipitation level and losses caused by
847 evapotranspiration (Boczoń and Wróbel, 2015; Zajączkowski et al., 2013).

848 The higher climatic sensitivity of pines at the Woziwoda site was manifested also by a higher number
849 of pointer years. The pointer years identified in this study are confirmed by earlier studies performed on
850 pine trees in northern Poland for 1910-2014 (Matulewski et al., 2019; Zielski et al., 1998; Zielski and
851 Barankiewicz, 2000). The years 1911, 1940, 1950, and 1982 attract particular attention. These are years in
852 which dry and hot summers were recorded (Matulewski et al., 2019; Zielski, 1996).

853 Our data show that *Pinus sylvestris* has been under critical climatic pressure and is responding
854 negatively to a warming climate and changing precipitation regime. Models predict a severe decline in
855 coniferous species in the next 50 years, including *Pinus sylvestris* in the temperate zone of Europe (Dyderski
856 et al., 2018; Hanewinkel et al., 2013; Schueler et al., 2014). The disappearance of species currently dominant
857 in the forests of central and eastern Europe will result in the profound disruption or disappearance of
858 ecosystems functionally related to them, such as peatlands (Dyderski et al., 2018).

859 Peatlands are also affected by accelerating climate change and on top of that they are at risk of losing
860 their favourable environment, especially in *Pinus sylvestris* monoculture forests, which are particularly
861 vulnerable to increasing extreme events. Studies conducted by various researchers confirm that remote
862 sensing data, provide a valuable source of information about peatlands and help in monitoring their
863 condition (Czapiewski and Szumińska, 2021; Kaplan et al., 2019; Lees et al., 2021; Rapinel et al., 2023).
864 The analyses conducted in this study have demonstrated that multisensor airborne data can be successfully

865 utilized to assess the current state of peatland vegetation. The application of simple remote sensing indices
866 enabled the detection of spatial differences in the condition and water stress of vegetation in the Okoniny
867 (Jezierzba) peatland. According to Rastogi et al. (2019), NDVI values for peatland vegetation may decrease
868 in areas affected by stress factors such as warming and reduced precipitation. Moreover, NDVI values for
869 healthy *Sphagnum* moss in peatland usually range from 0.8 to 0.9 during the summer, but they are also
870 species-dependent (Harris, 2008; Letendre et al., 2008; Péli et al., 2015). Consequently, the values of NDVI
871 observed in this study (averaging 0.65) may indicate a prevailing drought situation in certain areas of the
872 Okoniny (Jezierzba) peatland. Comparable findings can be drawn from the spatial variation of MSI values
873 presented in this study. Harris et al. (2006, 2005) demonstrated that MSI is significantly correlated with
874 near-surface moisture condition of *Sphagnum* moss. Despite the wide application of optical data and spectral
875 indices in assessing peatland conditions, Gerhards (2018) found that spectral indices may only be useful
876 under conditions of severe or prolonged water stress. For the pre-visual detection of initial vegetation water
877 stress symptoms, temperature-based indices are most suitable, exemplified by the LST index used in this
878 study. Although aerial thermal data has been previously applied in peatland research (Kopeć et al., 2016),
879 further research into the potential use of airborne thermal data in assessing peatland vegetation conditions
880 is recommended. To date, there have been few works in Poland using spectral data in peatland monitoring
881 (Bandopadhyay et al., 2021, 2019). However, none has attempted to collate palaeoecological,
882 dendrochronological, and remote sensing data.

883

884 5. Conclusions

885 Our data show that peatlands are highly sensitive to the progressive rise in Earth's temperatures and
886 changing precipitation regimes. Groundwater levels have dropped dramatically in recent years, causing
887 intense heating of the peatland surface in summer and stressing peat-forming vegetation to water scarcity.
888 The pine monocultures surrounding the peatlands are also sensitive to climate change. They are currently
889 responding very strongly to summer precipitation deficiency, and these data fit into dendrological predictive
890 models. Planned forest management has permanently changed the composition of the forest. Deciduous tree
891 species such as *Quercus*, *Fagus*, *Carpinus*, and *Corylus avellana* have almost disappeared. Forest
892 management has also contributed to increased acidity in the peatland, and thus the rapid development of
893 *Sphagnum* specialized for life in acidic conditions. After the expansion of *Sphagnum*, the water level in the
894 peatland stabilized. Peatlands are also valuable archives of past climatic anomalies and catastrophic events.
895 Pest gradations are recorded, among other things, by the presence of *Pinus* stomata, and periods of drought
896 by an increase in the values of coprophilous fungi. These events correspond with dendrochronological
897 records. There is a strong correlation between the first years after hailstorms and smaller increments of tree
898 rings. Our study shows that the combining of different data (palaeoecological, dendrochronological, remote

899 sensing and historical) can complement each other and create a more complete picture of past environmental
900 changes and expand knowledge of best practices for local (Konczal et al., 2024) and global (Joosten, 2021)
901 recommendations for peatland conservation in forests. Healthy wetlands could be key to protecting forests
902 and slowing the transformation of forests caused by climate change (Marcisz et al., 2024). The results are
903 essential for peatland conservation in planned forest management.

904

905 **Competing interests**

906 The contact author has declared that none of the authors has any competing interests.

907

908 **Acknowledgments**

909 The study was funded by the National Science Centre, Poland, grant 2020/39/D/ST10/00641. Remote
910 sensing data collection and visualization were done within the scope of the project "Protection of Valuable
911 Ecosystems of Tuchola Forest" funded by the European Economic Area Financial Mechanism 2014-2021
912 within the framework of the Environment, Energy and Climate Change Programme MF EEA 2014-2021
913 "Implementation of Ecosystem Management Plans".

914 We want to thank Stefan Konczal and other foresters from the Woziwoda Forestry Unit for their cooperation
915 and help in the field, for providing us with historical maps, and for sharing knowledge of the forest's history
916 and management. We thank Małgorzata Suchorska (Adam Mickiewicz University, Poznań) for her help in
917 the field.

918

919 **Data availability**

920 All data associated with this article are openly available on Mendeley Data repository under the DOI:
921 10.17632/prdgmjcg69.3

922

923 **Authors contribution**

924 MB – fieldwork, laboratory analyses (bulk density, carbon accumulation, plant macrofossils, selection of
925 plant macrofossils for AMS radiocarbon dating), age-depth modelling, data interpretation, visualization,
926 writing (original draft)

927 ML – fieldwork, support in plant macrofossil analysis, data interpretation, writing (commenting and editing)

928 PK – fieldwork, laboratory analyses (pollen and spores), age-depth modelling, data interpretation,
929 visualization, writing (commenting and editing)

930 DW – laboratory analyses (testate amoebae), testate amoeba-based reconstructions, data interpretation

931 PM – fieldwork, laboratory analyses (dendrochronology), data interpretation, visualization, writing
932 (commenting and editing)

933 DK, MW – fieldwork, remote sensing analyses and interpretation, writing (commenting and editing)
934 DJ – laboratory analyses (dendrochronology), data interpretation
935 KM – funding acquisition, conceptualization, fieldwork, laboratory analyses (charcoal), testate amoeba-
936 based reconstructions, data interpretation, visualization, writing (commenting and editing)

937

938 **References**

939

- 940 Adolf, C., Wunderle, S., Colombaroli, D., Weber, H., Gobet, E., Heiri, O., van Leeuwen, J.F.N., Bigler, C.,
941 Connor, S.E., Gałka, M., La Mantia, T., Makhortykh, S., Svitavská-Svobodová, H., Vannière, B.,
942 Tinner, W., 2018. The sedimentary and remote-sensing reflection of biomass burning in Europe. *Global*
943 *Ecology and Biogeography* 27, 199–212. <https://doi.org/10.1111/geb.12682>
944 Amesbury, M.J., Swindles, G.T., Bobrov, A., Charman, D.J., Lamentowicz, M., Mallon, G., Mazei, Y.,
945 Mitchell, E.A.D., Payne, R.J., Roland, T.P., Turner, E.T., Warner, B.G., 2016. Development of a new
946 pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quat Sci*
947 *Rev* 152, 132–151. <https://doi.org/10.1016/j.quascirev.2016.09.024>
948 Anderberg, A.-L., 1994. Atlas of seeds and small fruits of Northwest-European plant species with
949 morphological descriptions. Part 4: Resedaceae - Umbelliferae. Risbergs Tryckeri AB, Uddevalla.
950 Baillie, M.G.L., Pilcher, J., 1973. A simple cross-dating program for tree-ring research. *Tree-Ring Bull.* 33,
951 7–14.
952 Ballesteros-Cánovas, J.A., Edvardsson, J., Corona, C., Mažeika, J., Stoffel, M., 2022. Estimation of recent
953 peat accumulation with tree saplings. *Progress in Physical Geography: Earth and Environment* 46,
954 515–529. <https://doi.org/10.1177/03091333211073786>
955 Bandopadhyay, S., Rastogi, A., Cogliati, S., Rascher, U., Gąbka, M., Juszczak, R., 2021. Can Vegetation
956 Indices Serve as Proxies for Potential Sun-Induced Fluorescence (SIF)? A Fuzzy Simulation Approach
957 on Airborne Imaging Spectroscopy Data. *Remote Sens (Basel)* 13, 2545.
958 <https://doi.org/10.3390/rs13132545>
959 Bandopadhyay, S., Rastogi, A., Rascher, U., Rademske, P., Schickling, A., Cogliati, S., Julitta, T., Mac
960 Arthur, A., Hueni, A., Tomelleri, E., Celesti, M., Burkart, A., Stróżecki, M., Sakowska, K., Gąbka, M.,
961 Rosadziński, S., Sojka, M., Iordache, M.-D., Reusen, I., Van Der Tol, C., Damm, A., Schuettemeyer,
962 D., Juszczak, R., 2019. Hyplant-Derived Sun-Induced Fluorescence—A New Opportunity to
963 Disentangle Complex Vegetation Signals from Diverse Vegetation Types. *Remote Sens (Basel)* 11,
964 1691. <https://doi.org/10.3390/rs11141691>
965 Barabach, J., 2015. Zapis zdarzeń katastrofalnych na obszarze Puszczy Noteckiej w osadach Torfowiska
966 Rzecin. *Wydział Nauk Geograficznych i Geologicznych*.
967 Barabach, J., 2012. The history of Lake Rzecin and its surroundings drawn on maps as a background to
968 palaeoecological reconstruction. *Limnological Review* 12, 103–114. [https://doi.org/10.2478/v10194-](https://doi.org/10.2478/v10194-011-0050-0)
969 [011-0050-0](https://doi.org/10.2478/v10194-011-0050-0)
970 Bauhus, J., Forrester, D.I., Gardiner, B., Jactel, H., Vallejo, R., Pretzsch, H., 2017. Ecological Stability of
971 Mixed-Species Forests, in: *Mixed-Species Forests*. Springer, Berlin, Heidelberg, pp. 337–382.
972 https://doi.org/10.1007/978-3-662-54553-9_7
973 Beaulne, J., Boucher, É., Garneau, M., Magnan, G., 2021a. Paludification reduces black spruce growth rate
974 but does not alter tree water use efficiency in Canadian boreal forested peatlands. *For Ecosyst* 8, 28.
975 <https://doi.org/10.1186/s40663-021-00307-x>
976 Beaulne, J., Garneau, M., Magnan, G., Boucher, É., 2021b. Peat deposits store more carbon than trees in
977 forested peatlands of the boreal biome. *Sci Rep* 11, 2657. [https://doi.org/10.1038/s41598-021-82004-](https://doi.org/10.1038/s41598-021-82004-x)
978 x

- 979 Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. Present and
980 future Köppen-Geiger climate classification maps at 1-km resolution. *Sci Data* 5, 180214.
981 <https://doi.org/10.1038/sdata.2018.214>
- 982 Becker, M., Nieminen, T., Géréma, F., 1994. Short-term variations and long-term changes in oak
983 productivity in northeastern France. The role of climate and atmospheric CO₂. *Annales des Sciences*
984 *Forestières* 51, 477–492. <https://doi.org/10.1051/forest:19940504>
- 985 Berggren, G., 1969. Atlas of seeds and small fruits of Northwest-European plant species (Sweden, Norway,
986 Denmark, East Fennoscandia and Iceland) with morphological descriptions. Part 2: Cyperaceae.
987 Berlingska Boktryckeriet, Lund.
- 988 Berglund, B.E., Ralska-Jasiewiczowa, M., 1986. Pollen analysis and pollen diagrams, in: Berglund, B.E.
989 (Ed.), *Handbook of Holocene Palaeoecology and Palaeohydrology*. John Wiley & Sons, Chichester,
990 pp. 455–484.
- 991 Beug, H.-J., 2004. *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete*. Verlag Dr.
992 Friedrich Pfeil, München.
- 993 Bhiry, N., Filion, L., 1996. Mid-Holocene Hemlock Decline in Eastern North America Linked with
994 Phytophagous Insect Activity. *Quat Res* 45, 312–320. <https://doi.org/10.1006/qres.1996.0032>
- 995 Bienias, D., 2009. Las i człowiek w Borach Tucholskich (uwagi o bartnictwie i smolarstwie w Borach
996 Tucholskich, in: Woźny, J. (Ed.), *Dziedzictwo Techniczne Boro' w Tucholskich*. Przedsiębiorstwo
997 Marketingowe LOGO, Bydgoszcz, pp. 43–51.
- 998 Błaszkiwicz, M., Piotrowski, J.A., Brauer, A., Gierszewski, P., Kordowski, J., Kramkowski, M.,
999 Lamparski, P., Lorenz, S., Noryskiewicz, A.M., Ott, F., Słowiński, M., Tyszkowski, S., 2015. Climatic
1000 and morphological controls on diachronous postglacial lake and river valley evolution in the area of
1001 Last Glaciation, northern Poland. *Quat Sci Rev* 109, 13–27.
1002 <https://doi.org/10.1016/j.quascirev.2014.11.023>
- 1003 Blockeel, T., 2010. *Straminergon stramineum*, in: Atherton, I., Bosanquet, S., Lawley, M. (Eds.), *Mosses*
1004 *and Liverworts of Britain and Ireland a Field Guide*. British Bryological Society, Plymouth, p. 720.
- 1005 Blodau, C., 2002. Carbon cycling in peatlands – A review of processes and controls. *Environmental Reviews*
1006 10, 111–134. <https://doi.org/10.1139/a02-004>
- 1007 Boczoń, A., Kowalska, A., Gawryś, R., 2017. Glebowo-wodne uwarunkowania prowadzenia gospodarki
1008 leśnej w perspektywie zmian klimatu. *Sylwan* 161, 763–771.
- 1009 Boczoń, A., Wróbel, M., 2015. Wpływ suszy na pobór wody przez sosnę zwyczajną (*Pinus sylvestris* L.) o
1010 różnej pozycji w drzewostanie. *Leśne Prace Badawcze* 76, 370–376.
- 1011 Bojňanský, V., Fargašová, A., 2007. *Atlas of seeds and fruits of central and east-european flora. The*
1012 *Carpathian Mountains Region*. Springer, Dordrecht.
- 1013 Booth, R.K., 2002. Testate amoebae as paleoindicators of surface-moisture changes on Michigan peatlands:
1014 modern ecology and hydrological calibration. *J Paleolimnol* 28, 329–348.
- 1015 Booth, R.K., Lamentowicz, M., Charman, D.J., 2010. Preparation and analysis of testate amoebae in
1016 peatland paleoenvironmental studies. *Mires and Peat* 7 (2010/11), 1–7.
- 1017 Booth, T.H., 2013. Eucalypt plantations and climate change. *For Ecol Manage* 301, 28–34.
1018 <https://doi.org/10.1016/j.foreco.2012.04.004>
- 1019 Boulc'h, P.-N., Caullireau, E., Faucher, E., Gouerou, M., Guérin, A., Miray, R., Couée, I., 2020. Abiotic
1020 stress signalling in extremophile land plants. *J Exp Bot* 71, 5771–5785.
1021 <https://doi.org/10.1093/jxb/eraa336>
- 1022 Broda, J., 2000. *Historia leśnictwa w Polsce*. Wydawnictwo Akademii Rolniczej im. Augusta
1023 Cieszkowskiego w Poznaniu, Poznań.
- 1024 Broda, J., 1993. Sosna w czasach historycznych, in: Białobok, S., Boratyński, A., Bugała, W. (Eds.),
1025 *Biologia Sosny Zwyczajnej*. Instytut Dendrologii PAN, Poznań-Kórnik, pp. 17–31.
- 1026 Bronk Ramsey, C., 2021. OxCal v4.4.4 [WWW Document]. URL <https://c14.arch.ox.ac.uk/oxcal.html>
1027 (accessed 11.21.23).
- 1028 Bronk Ramsey, C., 2008. Deposition models for chronological records. *Quat Sci Rev* 27, 42–60.

- 1029 Bunn, A.G., 2008. A dendrochronology program library in R (dplR). *Dendrochronologia* (Verona) 26, 115–
1030 124. <https://doi.org/10.1016/j.dendro.2008.01.002>
- 1031 Cedro, A., 2001. Próba oceny oddziaływania temperatury powietrza i opadów atmosferycznych na przyrost
1032 radialny sosny zwyczajnej (<s>Pinus sylvestris</s>) na Pomorzu Zachodnim. *Annales Universitatis*
1033 *Mariae Curie-Skłodowska. Sectio B, Geographia, Geologia, Mineralogia et Petrographia* 55/56, 105–
1034 112.
- 1035 Cedro, A., Lamentowicz, M., 2011. Contrasting responses to environmental changes by pine (*Pinus*
1036 *syvestris* L.) growing on peat and mineral soil: An example from a Polish Baltic bog. *Dendrochronologia* (Verona) 29, 211–217. <https://doi.org/10.1016/j.dendro.2010.12.004>
- 1037 Chambers, F.M., Beilman, D.W., Yu, Z., 2010. Methods for determining peat humification and for
1038 quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate and
1039 peatland carbon dynamics. *Mires and Peat* 7, 1–10.
- 1041 Chapin, F.S., Matson, P.A., Vitousek, P., 2012. Managing and Sustaining Ecosystems, in: Chapin, F.S. (Ed.),
1042 *Principles of Terrestrial Ecosystem Ecology*. Springer, p. 447.
- 1043 Cherek, E., 2007. Ochotnicza Straż Pożarna w Kasparusie 1932-2007. *Kasparus*.
- 1044 Clark, J.S., 1990. Fire and climate change during the last 750 yr in northwestern Minnesota. *Ecol. Monogr.*
1045 60, 135–159. <https://doi.org/10.2307/1943042>
- 1046 Clark, J.S., 1988. Particle Motion and the Theory of Charcoal Analysis: Source Area, Transport, Deposition,
1047 and Sampling. *Quat Res* 30, 67–80. [https://doi.org/10.1016/0033-5894\(88\)90088-9](https://doi.org/10.1016/0033-5894(88)90088-9)
- 1048 Clarke, K.J., 2003. Guide to Identification of Soil Protozoa - Testate Amoebae, Soil Biodiversity Programme
1049 Research Report No. 4. Freshwater Biological Association, Ambleside, U.K.
- 1050 Clymo, R.S., Hayward, P.M., 1982. The Ecology of Sphagnum, in: Smith, A.J.E. (Ed.), *Bryophyte Ecology*.
1051 Chapman & Hall, London, New York, pp. 229–289.
- 1052 Cook, E.R., Briffa, K., Shiyatov, S., Mazepa, A., Jones, P.D., 1990. Data analysis, in: Cook, E.R.,
1053 Kairiukstis, L.A. (Eds.), *Methods of Dendrochronology: Applications in the Environmental Sciences*.
1054 Kluwer Academic Publ., Dordrecht, pp. 97–162.
- 1055 Cyzman, W., 2008. Jednolity Program Gospodarczo-Ochronny dla Leśnego Kompleksu Promocyjnego
1056 „Bory Tucholskie”.
- 1057 Czapiewski, S., Szumińska, D., 2021. An Overview of Remote Sensing Data Applications in Peatland
1058 Research Based on Works from the Period 2010–2021. *Land* (Basel) 11, 24.
1059 <https://doi.org/10.3390/land11010024>
- 1060 Czerwiński, S., Guzowski, P., Lamentowicz, M., Gałka, M., Karpińska-Kołaczek, M., Poniak, R., Łokas, E.,
1061 Diaconu, A.-C., Schwarzer, J., Miecznik, M., Kołaczek, P., 2021. Environmental implications of past
1062 socioeconomic events in Greater Poland during the last 1200 years. Synthesis of paleoecological and
1063 historical data. *Quat Sci Rev* 259, 106902.
1064 <https://doi.org/https://doi.org/10.1016/j.quascirev.2021.106902>
- 1065 Davis, M.B., Deevey, E.S., 1964. Pollen Accumulation Rates: Estimates from Late-Glacial Sediment of
1066 Rogers Lake. *Science* (1979) 145, 1293–1295. <https://doi.org/10.1126/science.145.3638.1293>
- 1067 Diaconu, A.-C., Tóth, M., Lamentowicz, M., Heiri, O., Kuske, E., Tanțău, I., Panait, A.-M., Braun, M.,
1068 Feurdean, A., 2017. How warm? How wet? Hydroclimate reconstruction of the past 7500 years in
1069 northern Carpathians, Romania. *Palaeogeogr Palaeoclimatol Palaeoecol* 482, 1–12.
1070 <https://doi.org/10.1016/j.palaeo.2017.05.007>
- 1071 Dietze, E., Brykała, D., Schreuder, L.T., Jażdżewski, K., Blarquez, O., Brauer, A., Dietze, M., Obremska,
1072 M., Ott, F., Pieńczewska, A., Schouten, S., Hopmans, E.C., Słowiński, M., 2019. Human-induced fire
1073 regime shifts during 19th century industrialization: A robust fire regime reconstruction using northern
1074 Polish lake sediments. *PLoS One* 14, e0222011. <https://doi.org/10.1371/journal.pone.0222011>
- 1075 Dinella, A., Giammarchi, F., Prendin, A.L., Carrer, M., Tonon, G., 2021. Xylem traits of peatland Scots
1076 pines reveal a complex climatic signal: A study in the Eastern Italian Alps. *Dendrochronologia*
1077 (Verona) 67, 125824. <https://doi.org/10.1016/j.dendro.2021.125824>

1078 Drinan, T.J., Graham, C.T., O'Halloran, J., Harrison, S.S.C., 2013. The impact of catchment conifer
1079 plantation forestry on the hydrochemistry of peatland lakes. *Science of The Total Environment* 443,
1080 608–620. <https://doi.org/10.1016/j.scitotenv.2012.10.112>

1081 Dyderski, M.K., Paż, S., Frelich, L.E., Jagodziński, A.M., 2018. How much does climate change threaten
1082 European forest tree species distributions? *Glob Chang Biol* 24, 1150–1163.
1083 <https://doi.org/10.1111/gcb.13925>

1084 Eckstein, D., Bauch, J., 1969. Beitrag zur Rationalisierung eines dendrochronologischen Verfahrens und zur
1085 Analyse seiner Aussagesicherheit. *Forstwissenschaftliches Centralblatt* 88, 230–250.
1086 <https://doi.org/10.1007/BF02741777>

1087 Edvardsson, J., Baužienė, I., Lamentowicz, M., Šimanauskienė, R., Tamkevičiūtė, M., Taminskas, J.,
1088 Linkevičienė, R., Skuratovič, Ž., Corona, C., Stoffel, M., 2019. A multi-proxy reconstruction of
1089 moisture dynamics in a peatland ecosystem: A case study from Čepkeliai, Lithuania. *Ecol Indic* 106,
1090 105484. <https://doi.org/https://doi.org/10.1016/j.ecolind.2019.105484>

1091 Edvardsson, J., Corona, C., Mažeika, J., Pukienė, R., Stoffel, M., 2016. Recent advances in long-term
1092 climate and moisture reconstructions from the Baltic region: Exploring the potential for a new multi-
1093 millennial tree-ring chronology. *Quat Sci Rev* 131, Part A, 118–126.
1094 <https://doi.org/https://doi.org/10.1016/j.quascirev.2015.11.005>

1095 Edvardsson, J., Helama, S., Rundgren, M., Nielsen, A.B., 2022. The Integrated Use of Dendrochronological
1096 Data and Paleoecological Records From Northwest European Peatlands and Lakes for Understanding
1097 Long-Term Ecological and Climatic Changes—A Review. *Front Ecol Evol* 10.
1098 <https://doi.org/10.3389/fevo.2022.781882>

1099 FAO, 2020. Peatlands mapping and monitoring. Recommendations and technical overview. Rome.
1100 <https://doi.org/10.4060/ca8200en>

1101 Feliksik, E., Wilczyński, S., 2009. The Effect of Climate on Tree-Ring Chronologies of Native and
1102 Nonnative Tree Species Growing Under Homogenous Site Conditions. *Geochronometria* 33, 49–57.
1103 <https://doi.org/10.2478/v10003-009-0006-4>

1104 Felton, A., Gustafsson, L., Roberge, J.-M., Ranius, T., Hjältén, J., Rudolphi, J., Lindbladh, M., Weslien, J.,
1105 Rist, L., Brunet, J., Felton, A.M., 2016. How climate change adaptation and mitigation strategies can
1106 threaten or enhance the biodiversity of production forests: Insights from Sweden. *Biol Conserv* 194,
1107 11–20. <https://doi.org/10.1016/j.biocon.2015.11.030>

1108 Finsinger, W., Tinner, W., 2005. Minimum count sums for charcoal-concentration estimates in pollen slides:
1109 accuracy and potential errors. *Holocene* 15, 293–297.

1110 Freeman, C., Fenner, N., Ostle, N., Kang, H., Dorwick, D.J., Reynolds, B., Lock, M.A., Sleep, D., Hughes,
1111 S., Hudson, J., 2004. Export of dissolved organic carbon from peatlands under elevated carbon dioxide
1112 levels. *Nature* 430, 195–198.

1113 Gałka, M., Knorr, K.-H., Tobolski, K., Gallego-Sala, A., Kołaczek, P., Lamentowicz, M., Kajukał-
1114 Drygalska, K., Marcisz, K., 2022. How far from a pristine state are the peatlands in the Białowieża
1115 Primeval Forest (CE Europe) – Palaeoecological insights on peatland and forest development from
1116 multi-proxy studies. *Ecol Indic* 143, 109421. <https://doi.org/10.1016/j.ecolind.2022.109421>

1117 Gałka, M., Miotk-Szpiganiowicz, G., Marczevska, M., Barabach, J., van der Knaap, W.O., Lamentowicz,
1118 M., 2015. Palaeoenvironmental changes in Central Europe (NE Poland) during the last 6200 years
1119 reconstructed from a high-resolution multi-proxy peat archive. *Holocene* 25, 421–434.
1120 <https://doi.org/10.1177/0959683614561887>

1121 Gałka, M., Tobolski, K., Górska, A., Milecka, K., Fiałkiewicz-Kozieł, B., Lamentowicz, M., 2014.
1122 Disentangling the drivers for the development of a Baltic bog during the Little Ice Age in northern
1123 Poland. *Quaternary International* 328–329, 323–337.
1124 <https://doi.org/http://dx.doi.org/10.1016/j.quaint.2013.02.026>

1125 Gallego-Sala, A. V., Charman, D.J., Brewer, S., Page, S.E., Prentice, I.C., Friedlingstein, P., Moreton, S.,
1126 Amesbury, M.J., Beilman, D.W., Björck, S., Blyakharchuk, T., Bochicchio, C., Booth, R.K., Bunbury,
1127 J., Camill, P., Carless, D., Chimner, R.A., Clifford, M., Cressey, E., Courtney-Mustaphi, C., De
1128 Vleeschouwer, F., de Jong, R., Fiałkiewicz-Kozieł, B., Finkelstein, S.A., Garneau, M., Githumbi, E.,

1129 Hribljan, J., Holmquist, J., Hughes, P.D.M., Jones, C., Jones, M.C., Karofeld, E., Klein, E.S., Kokfelt,
1130 U., Korhola, A., Lacourse, T., Le Roux, G., Lamentowicz, M., Large, D., Lavoie, M., Loisel, J.,
1131 Mackay, H., MacDonald, G.M., Makila, M., Magnan, G., Marchant, R., Marcisz, K., Martínez
1132 Cortizas, A., Massa, C., Mathijssen, P., Mauquoy, D., Mighall, T., Mitchell, F.J.G., Moss, P., Nichols,
1133 J., Oksanen, P.O., Orme, L., Packalen, M.S., Robinson, S., Roland, T.P., Sanderson, N.K., Sannel,
1134 A.B.K., Silva-Sánchez, N., Steinberg, N., Swindles, G.T., Turner, T.E., Uglow, J., Välranta, M., van
1135 Bellen, S., van der Linden, M., van Geel, B., Wang, G., Yu, Z., Zaragoza-Castells, J., Zhao, Y., 2018.
1136 Latitudinal limits to the predicted increase of the peatland carbon sink with warming. *Nat Clim Chang*
1137 8, 907–913. <https://doi.org/https://www.nature.com/articles/s41558-018-0271-1>

1138 Gerhards, M., 2018. Advanced Thermal Remote Sensing for Water Stress Detection of Agricultural Crops.
1139 Godwin, H., 1981. *Archives of the Peat Bogs*. Cambridge University Press, Cambridge.

1140 González de Andrés, E., Shestakova, T.A., Scholten, R.C., Delcourt, C.J.F., Gorina, N. V, Camarero, J.J.,
1141 2022. Changes in tree growth synchrony and resilience in Siberian *Pinus sylvestris* forests are
1142 modulated by fire dynamics and ecohydrological conditions. *Agric For Meteorol* 312, 108712.
1143 <https://doi.org/https://doi.org/10.1016/j.agrformet.2021.108712>

1144 Gorham, E., 1991. Northern Peatlands: Role in the Carbon Cycle and Probable Responses to Climatic
1145 Warming. *Ecological Applications* 1, 182–195. <https://doi.org/10.2307/1941811>

1146 Gregow, H., Laaksonen, A., Alper, M.E., 2017. Increasing large scale windstorm damage in Western, Central
1147 and Northern European forests, 1951–2010. *Sci Rep* 7, 46397. <https://doi.org/10.1038/srep46397>

1148 Grimm, E.C., 1992. *Tilia and Tilia-Graph. Pollen Spreadsheet and Graphics Programs*. 8th International
1149 Palynological Congress (President: A. Pons) Aix-en-Provence, September 6-12, 1992. Program and
1150 Abstracts, p. 56.

1151 Grimm, E.C., 1991. *Tilia and Tilia Graph*. Illinois State Museum.

1152 Grissino-Mayer, H.D., 2001. Evaluating crossdating accuracy: A manual and tutorial for the computer
1153 program COFECHA. *Tree Ring Res* 57, 205–221.

1154 Grodzki, W., 2016. Mass outbreaks of the spruce bark beetle *Ips typographus* in the context of the
1155 controversies around the Białowieża Primeval Forest. *Forest Research Papers* 77, 324–331.
1156 <https://doi.org/10.1515/frp-2016-0033>

1157 Grondin, P., Gauthier, S., Borcard, D., Bergeron, Y., Noël, J., 2014. A new approach to ecological land
1158 classification for the Canadian boreal forest that integrates disturbances. *Landsc Ecol* 29, 1–16.
1159 <https://doi.org/10.1007/s10980-013-9961-2>

1160 Guariguata, M.R., Cornelius, J.P., Locatelli, B., Forner, C., Sánchez-Azofeifa, G.A., 2008. Mitigation needs
1161 adaptation: Tropical forestry and climate change. *Mitig Adapt Strateg Glob Chang* 13, 793–808.
1162 <https://doi.org/10.1007/s11027-007-9141-2>

1163 Guiot, J., 1991. The bootstrapped response function. *Tree-Ring Bulletin* 51, 39–41.

1164 Guo, M., Li, J., Sheng, C., Xu, J., Wu, L., 2017. A Review of Wetland Remote Sensing. *Sensors* 17, 777.
1165 <https://doi.org/10.3390/s17040777>

1166 Hanewinkel, M., Cullmann, D.A., Schelhaas, M.-J., Nabuurs, G.-J., Zimmermann, N.E., 2013. Climate
1167 change may cause severe loss in the economic value of European forest land. *Nat Clim Chang* 3, 203–
1168 207. <https://doi.org/10.1038/nclimate1687>

1169 Hanson, P.J., Weltzin, J.F., 2000. Drought disturbance from climate change: response of United States
1170 forests. *Science of The Total Environment* 262, 205–220. [https://doi.org/10.1016/S0048-9697\(00\)00523-4](https://doi.org/10.1016/S0048-9697(00)00523-4)

1171 Harenda, K.M., Lamentowicz, M., Samson, M., Chojnicki, B.H., 2018. The Role of Peatlands and Their
1172 Carbon Storage Function in the Context of Climate Change. pp. 169–187. https://doi.org/10.1007/978-3-319-71788-3_12

1173 Harriman, R., Watt, A.W., Christie, A.E.G., Moore, D.W., McCartney, A.G., Taylor, E.M., 2003. Quantifying
1174 the effects of forestry practices on the recovery of upland streams and lochs from acidification. *Science
1175 of The Total Environment* 310, 101–111. [https://doi.org/10.1016/S0048-9697\(02\)00626-5](https://doi.org/10.1016/S0048-9697(02)00626-5)

1176 Harris, A., 2008. Spectral reflectance and photosynthetic properties of *Sphagnum* mosses exposed to
1177 progressive drought. *Ecohydrology* 1, 35–42. <https://doi.org/10.1002/eco.5>

1180 Harris, A., Bryant, R., Baird, A., 2005. Detecting near-surface moisture stress in spp. *Remote Sens Environ*
1181 97, 371–381. <https://doi.org/10.1016/j.rse.2005.05.001>

1182 Harris, A., Bryant, R.G., Baird, A.J., 2006. Mapping the effects of water stress on Sphagnum: Preliminary
1183 observations using airborne remote sensing. *Remote Sens Environ* 100, 363–378.
1184 <https://doi.org/10.1016/j.rse.2005.10.024>

1185 Haylock, M.R., Hofstra, N., Klein Tank, A.M.G., Klok, E.J., Jones, P.D., New, M., 2008. A European daily
1186 high-resolution gridded data set of surface temperature and precipitation for 1950–2006. *Journal of*
1187 *Geophysical Research: Atmospheres* 113. <https://doi.org/10.1029/2008JD010201>

1188 Hedenäs, L., 1993. A generic revision of the *Warnstorfia-Calliergon* group. *J Bryol* 17, 447–479.
1189 <https://doi.org/10.1179/jbr.1993.17.3.447>

1190 Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and carbonate
1191 content in sediments: Reproducibility and comparability of results. *J Paleolimnol* 25, 101–110.
1192 <https://doi.org/10.1023/A:1008119611481>

1193 Higuera, P., Peters, M., Brubaker, L., Gavin, D., 2007. Understanding the origin and analysis of sediment-
1194 charcoal records with a simulation model. *Quat Sci Rev* 26, 1790–1809.
1195 <https://doi.org/10.1016/j.quascirev.2007.03.010>

1196 Hill, M.O., Blockeel, T.L., 2014. *Straminergon stramineum*, in: Blockeel, T.L., Bosanquet, S.D.S., Hill,
1197 M.O., Preston, C.D. (Eds.), *Atlas of British and Irish Bryophytes*. British Bryological Society,
1198 Newbury, Berkshire, p. 464.

1199 Hua, Q., Turnbull, J.C., Santos, G.M., Rakowski, A.Z., Ancapichún, S., De Pol-Holz, R., Hammer, S.,
1200 Lehman, S.J., Levin, I., Miller, J.B., Palmer, J.G., Turney, C.S.M., 2021. ATMOSPHERIC
1201 RADIOCARBON FOR THE PERIOD 1950–2019. *Radiocarbon* 1–23.
1202 <https://doi.org/10.1017/RDC.2021.95>

1203 Hunt, E., Rock, B., 1989. Detection of changes in leaf water content using Near- and Middle-Infrared
1204 reflectances☆. *Remote Sens Environ* 30, 43–54. [https://doi.org/10.1016/0034-4257\(89\)90046-1](https://doi.org/10.1016/0034-4257(89)90046-1)

1205 Huuskonen, S., Domisch, T., Finér, L., Hantula, J., Hynynen, J., Matala, J., Miina, J., Neuvonen, S.,
1206 Nevalainen, S., Niemistö, P., Nikula, A., Piri, T., Siitonen, J., Smolander, A., Tonteri, T., Uotila, K.,
1207 Viiri, H., 2021. What is the potential for replacing monocultures with mixed-species stands to enhance
1208 ecosystem services in boreal forests in Fennoscandia? *For Ecol Manage* 479, 118558.
1209 <https://doi.org/10.1016/j.foreco.2020.118558>

1210 Jabłoński, T., 2015. Występowanie i zwalczanie leśnych foliofagów - trendy i prognozy. *Postępy Techniki*
1211 *w Leśnictwie* 132, 13–19.

1212 Jäger, E., 1982. Prussia-Karten 1542-1810: Geschichte der kartographischen Darstellung Ostpreussens vom
1213 16. bis zum 19. Jahrhundert. Entstehung der Karten - Kosten - Vertrieb: bibliographischer Katalog.
1214 Anton H. Konrad Verlag, Weißenhorn.

1215 Jäger, E., 1981. Die Schroettersche Landesaufnahme von Ost- und Westpreußen (1796-1802).
1216 Entstehungsgeschichte, Herstellung und Vertrieb der Karte. *Z Ostforsch* 30, 359–389.

1217 Jasnowski, M., 1962. Budowa i roślinność torfowisk Pomorza Szczecińskiego. *Societas Scientiarum*
1218 *Stetinensis, Szczecin*.

1219 Joosten, H., 2021. Global guidelines for peatland rewetting and restoration. Ramsar Technical Report No.
1220 11. Gland, Switzerland.

1221 Joosten, H., Tapio-Biström, M.-L., Tol, S., 2012. Peatlands - guidance for climate change mitigation through
1222 conservation, rehabilitation and sustainable use, 2nd ed. Food and Agriculture Organization of the
1223 United Nations, Rome.

1224 Juggins, S., 2023. Rioja: Analysis of Quaternary Science Data [WWW Document]. URL [https://cran.r-](https://cran.r-project.org/web/packages/rioja/index.html)
1225 [project.org/web/packages/rioja/index.html](https://cran.r-project.org/web/packages/rioja/index.html) (accessed 12.4.23).

1226 Juggins, S., 2007. C2 Version 1.5 User guide. Software for ecological and palaeoecological data analysis
1227 and visualisation. Newcastle University, Newcastle upon Tyne, UK, p. 73.

1228 Kamińska, A., Lisiewicz, M., Kraszewski, B., Stereńczak, K., 2021. Mass outbreaks and factors related to
1229 the spatial dynamics of spruce bark beetle (*Ips typographus*) dieback considering diverse management

1230 regimes in the Białowieża forest. *For Ecol Manage* 498, 119530.
1231 <https://doi.org/10.1016/j.foreco.2021.119530>

1232 Kaplan, G., Yigit Avdan, Z., Avdan, U., 2019. Mapping and Monitoring Wetland Dynamics Using Thermal,
1233 Optical, and SAR Remote Sensing Data, in: *Wetlands Management - Assessing Risk and Sustainable*
1234 *Solutions*. IntechOpen. <https://doi.org/10.5772/intechopen.80264>

1235 Karpińska-Kołaczek, M., Kołaczek, P., Marcisz, K., Gałka, M., Kajukało-Drygalska, K., Mauquoy, D.,
1236 Lamentowicz, M., 2024. Kettle-hole peatlands as carbon hot spots: Unveiling controls of carbon
1237 accumulation rates during the last two millennia. *Catena (Amst)* 237, 107764.
1238 <https://doi.org/10.1016/j.catena.2023.107764>

1239 Kelly, R., Chipman, M.L., Higuera, P.E., Stefanova, I., Brubaker, L.B., Hu, F.S., 2013. Recent burning of
1240 boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy*
1241 *of Sciences* 110, 13055–13060. <https://doi.org/10.1073/pnas.1305069110>

1242 Kiełczewski, B., 1947. Klęska sówki chojnowki jako zagadnienie biocenotyczne. *Prace Komisji*
1243 *Matematyczno-Przyrodniczej* 10, 167–171.

1244 Kirschenstein, M., 2005. Wieloletnie zmiany sum opadów atmosferycznych na wybranych stacjach
1245 północno-zachodniej Polski. *Śląskie Prace Geograficzne* 2, 199–214.

1246 Klein Tank, A.M.G., Wijngaard, J.B., Können, G.P., Böhm, R., Demarée, G., Gocheva, A., Mileta, M.,
1247 Pashiardis, S., Hejkrlik, L., Kern-Hansen, C., Heino, R., Bessemoulin, P., Müller-Westermeier, G.,
1248 Tzanakou, M., Szalai, S., Pálsdóttir, T., Fitzgerald, D., Rubin, S., Capaldo, M., Maugeri, M., Leitass,
1249 A., Bukantis, A., Aberfeld, R., van Engelen, A.F. V., Forland, E., Miletus, M., Coelho, F., Mares, C.,
1250 Razuvaev, V., Nieplova, E., Cegnar, T., Antonio López, J., Dahlström, B., Moberg, A., Kirchhofer, W.,
1251 Ceylan, A., Pachaliuk, O., Alexander, L. V., Petrovic, P., 2002. Daily dataset of 20th-century surface
1252 air temperature and precipitation series for the European Climate Assessment. *International Journal of*
1253 *Climatology* 22, 1441–1453. <https://doi.org/10.1002/joc.773>

1254 Koenig, I., Mulot, M., Mitchell, E.A.D., 2018. Taxonomic and functional traits responses of *Sphagnum*
1255 peatland testate amoebae to experimentally manipulated water table. *Ecol Indic* 85, 342–351.
1256 <https://doi.org/https://doi.org/10.1016/j.ecolind.2017.10.017>

1257 Kołaczek, P., Karpińska-Kołaczek, M., Marcisz, K., Gałka, M., Lamentowicz, M., 2018. Palaeohydrology
1258 and the human impact on one of the largest raised bogs complex in the Western Carpathians (Central
1259 Europe) during the last two millennia. *Holocene* 28, 595–608.
1260 <https://doi.org/10.1177/0959683617735587>

1261 Konczal, S., Lamentowicz M., Bąk, M., Czerwiński, S., Kołaczek, P., Wochal, D., Marcisz, M., Chojnicki,
1262 B., Harenda, K., Poczta, P., Gąbka, M., Jaster, D., Matulewski, P., Jedliński, J., Niedzielko, J.,
1263 Wylazłowska, J., Żmuda, M., Żmuda, D., Kopeć, D., Rosadziński, S., Wietecha, M., Landowska, J.,
1264 Landowski, J., 2024. Rekomendacje dla ochrony mokradeł w lasach, in: Lamentowicz, M., Konczal,
1265 S. (Eds.), *Jak Chronić Torfowiska w Lasach?* ArchaeGraph, Łódź, pp. 161–165.

1266 Kondracki, J., 2001. *Geografia regionalna Polski*. Wydawnictwo Naukowe PWN, Warszawa.

1267 Kopeć, D., Michalska-Hejduk, D., Sławik, Ł., Berezowski, T., Borowski, M., Rosadziński, S., Chormański,
1268 J., 2016. Application of multisensoral remote sensing data in the mapping of alkaline fens Natura 2000
1269 habitat. *Ecol Indic* 70, 196–208. <https://doi.org/10.1016/j.ecolind.2016.06.001>

1270 Koprowski, M., Przybylak, R., Zielski, A., Pospieszynska, A., 2012. Tree rings of Scots pine (*Pinus*
1271 *sylvestris* L.) as a source of information about past climate in northern Poland. *Int J Biometeorol* 56,
1272 1–10. <https://doi.org/10.1007/s00484-010-0390-5>

1273 Koprowski, M., Zielski, A., Skowronek, T., 2011. Analiza przyrostów rocznych dwóch sosen (*Pinus*
1274 *sylvestris*) o nietypowej budowie strzały na terenie Nadleśnictwa Borne Sulino. *Sylwan* 155, 555–
1275 562.

1276 Kowalewski, G., 2003. Shoreline changes of basins in the mire-lake reserves in s Tuchola Pinewoods.
1277 *Limnological Review* 3, 119–126.

1278 Kowalewski, G., Milecka, K., 2003. Palaeoecology of basins of organic sediment accumulation in the
1279 Reserve Dury. *Studia Quaternaria* 20, 73–82.

1280 Kuosmanen, N., Čada, V., Halsall, K., Chiverrell, R.C., Schafstall, N., Kuneš, P., Boyle, J.F., Knížek, M.,
 1281 Appleby, P.G., Svoboda, M., Clear, J.L., 2020. Integration of dendrochronological and
 1282 palaeoecological disturbance reconstructions in temperate mountain forests. *For Ecol Manage* 475,
 1283 118413. <https://doi.org/10.1016/j.foreco.2020.118413>
 1284 Łabędzki, L., 2004. Problematyka susz w Polsce. *Woda-Środowisko-Obszary Wiejskie* 4, 47–66.
 1285 Laine, J., Flatberg, K.I., Harju, P., Timonen, T., Minkinen, K., Laine, A., Tuittila, E.-S., Vasander, H., 2018.
 1286 Sphagnum mosses. The Stars of European Mires . Department of Forest Sciences, University of
 1287 Helsinki, Sphagna Ky, Helsinki.
 1288 Laine, J., Vasander, H., Laiho, R., 1995. Long-Term Effects of Water Level Drawdown on the Vegetation of
 1289 Drained Pine Mires in Southern Finland. *J Appl Ecol* 32, 785–802. <https://doi.org/10.2307/2404818>
 1290 Lamentowicz, Ł., Gałka, M., Rusińska, A., Sobczyński, T., Owsiany, P.M., Lamentowicz, M., 2011.
 1291 Testate amoeba (Arcellinida, Euglyphida) ecology along a poor-rich gradient in fens of western
 1292 Poland. *Int Rev Hydrobiol* 96, 256–380.
 1293 Lamentowicz, M., Balwierz, Z., Forysiak, J., Płóciennik, M., Kittel, P., Kloss, M., Twardy, J., Żurek, S.,
 1294 Pawlyta, J., 2009a. Multiproxy study of anthropogenic and climatic changes in the last two millennia
 1295 from a small mire in central Poland. *Hydrobiologia* 631, 213–230. [https://doi.org/10.1007/s10750-](https://doi.org/10.1007/s10750-009-9812-y)
 1296 009-9812-y
 1297 Lamentowicz, M., Cedro, A., Gałka, M., Goslar, T., Miotk-Szpiganowicz, G., Mitchell, E.A.D., Pawlyta, J.,
 1298 2008a. Last millennium palaeoenvironmental changes from a Baltic bog (Poland) inferred from stable
 1299 isotopes, pollen, plant macrofossils and testate amoebae. *Palaeogeogr Palaeoclimatol Palaeoecol* 265,
 1300 93–106.
 1301 Lamentowicz, M., Gałka, M., Lamentowicz, Ł., Obremska, M., Kühl, N., Lücke, A., Jassey, V.E.J., 2015.
 1302 Reconstructing climate change and ombrotrophic bog development during the last 4000years in
 1303 northern Poland using biotic proxies, stable isotopes and trait-based approach. *Palaeogeogr*
 1304 *Palaeoclimatol Palaeoecol* 418, 261–277. <https://doi.org/10.1016/j.palaeo.2014.11.015>
 1305 Lamentowicz, M., Marcisz, K., Guzowski, P., Gałka, M., Diaconu, A.-C., Kołaczek, P., 2020. How
 1306 Joannites' economy eradicated primeval forest and created anthroecosystems in medieval Central
 1307 Europe. *Sci Rep* 10, 18775. <https://doi.org/10.1038/s41598-020-75692-4>
 1308 Lamentowicz, M., Milecka, K., Gałka, M., Cedro, A., Pawlyta, J., Piotrowska, N., Lamentowicz, Ł., van
 1309 der Knaap, W.O., 2009b. Climate and human induced hydrological change since AD 800 in an
 1310 ombrotrophic mire in Pomerania (N Poland) tracked by testate amoebae, macro-fossils, pollen tree-
 1311 rings of pine. *Boreas, An international journal of Quaternary research* 38, 214–229.
 1312 Lamentowicz, M., Mitchell, E.A.D., 2005a. Testate amoebae (Protists) as palaeoenvironmental indicators
 1313 in peatlands. *Polish Geological Institute Special Papers* 16, 58–64.
 1314 Lamentowicz, M., Mitchell, E.A.D., 2005b. The ecology of testate amoebae (Protists) in Sphagnum in north-
 1315 western Poland in relation to peatland ecology. *Microb Ecol* 50, 48–63.
 1316 Lamentowicz, M., Mueller, M., Gałka, M., Barabach, J., Milecka, K., Goslar, T., Binkowski, M., 2015.
 1317 Reconstructing human impact on peatland development during the past 200 years in CE
 1318 Europe through biotic proxies and X-ray tomography. *Quaternary International* 357, 282–294.
 1319 <https://doi.org/http://dx.doi.org/10.1016/j.quaint.2014.07.045>
 1320 Lamentowicz, M., Obremska, M., 2010. A rapid response of testate amoebae and vegetation to inundation
 1321 of a kettle hole mire. *J Paleolimnol* 43, 499–511. <https://doi.org/10.1007/s10933-009-9347-2>
 1322 Lamentowicz, M., Obremska, M., Mitchell, E.A.D., 2008b. Autogenic succession, land-use change, and
 1323 climatic influences on the Holocene development of a kettle hole mire in Northern Poland. *Rev*
 1324 *Palaeobot Palynol* 151, 21–40. <https://doi.org/10.1016/j.revpalbo.2008.01.009>
 1325 Lamentowicz, M., Tobolski, K., Mitchell, E.A.D., 2007. Palaeoecological evidence for anthropogenic
 1326 acidification of a kettle-hole peatland in northern Poland. *Holocene* 17, 1185–1196.
 1327 Lavoie, M., Filion, L., Robert, É.C., 2009. Boreal peatland margins as repository sites of long-term natural
 1328 disturbances of balsam fir/spruce forests. *Quat Res* 71, 295–306.
 1329 <https://doi.org/10.1016/j.yqres.2009.01.005>

- 1330 Lee, D., Holmström, E., Hynynen, J., Nilsson, U., Korhonen, K.T., Westerlund, B., Bianchi, S., Aldea, J.,
 1331 Huuskonen, S., 2023. Current state of mixed forests available for wood supply in Finland and Sweden.
 1332 Scand J For Res 1–11. <https://doi.org/10.1080/02827581.2023.2259797>
- 1333 Lees, K.J., Artz, R.R.E., Chandler, D., Aspinall, T., Boulton, C.A., Buxton, J., Cowie, N.R., Lenton, T.M.,
 1334 2021. Using remote sensing to assess peatland resilience by estimating soil surface moisture and
 1335 drought recovery. Science of The Total Environment 761, 143312.
 1336 <https://doi.org/10.1016/j.scitotenv.2020.143312>
- 1337 Letendre, J., Poulin, M., Rochefort, L., 2008. Sensitivity of spectral indices to CO₂ fluxes for several plant
 1338 communities in a *Sphagnum*-dominated peatland. Canadian Journal of Remote Sensing 34, S414–
 1339 S425. <https://doi.org/10.5589/m08-053>
- 1340 Loisel, J., Yu, Z., Beilman, D.W., Camill, P., Alm, J., Amesbury, M.J., Anderson, D., Andersson, S.,
 1341 Bochicchio, C., Barber, K., Belyea, L.R., Bunbury, J., Chambers, F.M., Charman, D.J., De
 1342 Vleeschouwer, F., Fiałkiewicz-Kozieł, B., Finkelstein, S.A., Gałka, M., Garneau, M., Hammarlund,
 1343 D., Hinchcliffe, W., Holmquist, J., Hughes, P., Jones, M.C., Klein, E.S., Kokfelt, U., Korhola, A.,
 1344 Kuhry, P., Lamarre, A., Lamentowicz, M., Large, D., Lavoie, M., MacDonald, G., Magnan, G., Mäkilä,
 1345 M., Mallon, G., Mathijssen, P., Mauquoy, D., McCarroll, J., Moore, T.R., Nichols, J., O'Reilly, B.,
 1346 Oksanen, P., Packalen, M., Peteet, D., Richard, P.J.H., Robinson, S., Ronkainen, T., Rundgren, M.,
 1347 Sannel, A.B.K., Tarnocai, C., Thom, T., Tuittila, E.-S., Turetsky, M., Väiliranta, M., van der Linden,
 1348 M., van Geel, B., van Bellen, S., Vitt, D., Zhao, Y., Zhou, W., 2014. A database and synthesis of
 1349 northern peatland soil properties and Holocene carbon and nitrogen accumulation. Holocene 24, 1028–
 1350 1042. <https://doi.org/10.1177/0959683614538073>
- 1351 Łuców, D., Küttim, M., Słowiński, M., Kołaczek, P., Karpińska-Kołaczek, M., Küttim, L., Salme, M.,
 1352 Lamentowicz, M., 2022. Searching for an ecological baseline: Long-term ecology of a post-extraction
 1353 restored bog in Northern Estonia. Quaternary International 607, 65–78.
 1354 <https://doi.org/10.1016/j.quaint.2021.08.017>
- 1355 Łuców, D., Lamentowicz, M., Kołaczek, P., Łokas, E., Marcisz, K., Obremska, M., Theuerkauf, M.,
 1356 Tyszkowski, S., Słowiński, M., 2021. Pine Forest Management and Disturbance in Northern Poland:
 1357 Combining High-Resolution 100-Year-Old Paleoecological and Remote Sensing Data. Front Ecol
 1358 Evol 9. <https://doi.org/10.3389/fevo.2021.747976>
- 1359 Marcisz, K., Bąk, M., Kołaczek, P., Lamentowicz, M., Wochal, D., 2024. Historia lasu i mokradeł zapisana
 1360 w torfowiskach, in: Lamentowicz, M., Konczal, S. (Eds.), Jak Chronić Torfowiska w Lasach?
 1361 ArchaeGraph, Łódź, pp. 29–45.
- 1362 Marcisz, K., Gałka, M., Pietrala, P., Miotk-Szpiganiowicz, G., Obremska, M., Tobolski, K., Lamentowicz,
 1363 M., 2017. Fire activity and hydrological dynamics in the past 5700 years reconstructed from *Sphagnum*
 1364 peatlands along the oceanic–continental climatic gradient in northern Poland. Quat Sci Rev 177, 145–
 1365 157. <https://doi.org/10.1016/j.quascirev.2017.10.018>
- 1366 Marcisz, K., Jassey, V.E.J., Kosakyan, A., Krashevskaya, V., Lahr, D.J.G., Lara, E., Lamentowicz, Ł.,
 1367 Lamentowicz, M., Macumber, A., Mazei, Y., Mitchell, E.A.D., Nasser, N.A., Patterson, R.T., Roe,
 1368 H.M., Singer, D., Tsyganov, A.N., Fournier, B., 2020a. Testate Amoeba Functional Traits and Their
 1369 Use in Paleoecology. Front Ecol Evol 8, 340. <https://doi.org/10.3389/fevo.2020.575966>
- 1370 Marcisz, K., Kołaczek, P., Gałka, M., Diaconu, A.-C., Lamentowicz, M., 2020b. Exceptional hydrological
 1371 stability of a *Sphagnum*-dominated peatland over the late Holocene. Quat Sci Rev 231, 106180.
 1372 <https://doi.org/https://doi.org/10.1016/j.quascirev.2020.106180>
- 1373 Marcisz, K., Tinner, W., Colombaroli, D., Kołaczek, P., Słowiński, M., Fiałkiewicz-Kozieł, B., Łokas, E.,
 1374 Lamentowicz, M., 2015. Long-term hydrological dynamics and fire history over the last 2000 years in
 1375 CE Europe reconstructed from a high-resolution peat archive. Quat Sci Rev 112, 138–152.
 1376 <https://doi.org/10.1016/j.quascirev.2015.01.019>
- 1377 Marks, L., 2012. Timing of the Late Vistulian (Weichselian) glacial phases in Poland. Quat Sci Rev 44, 81–
 1378 88. <https://doi.org/https://doi.org/10.1016/j.quascirev.2010.08.008>

- 1379 Matthias, I., Giesecke, T., 2014. Insights into pollen source area, transport and deposition from modern
1380 pollen accumulation rates in lake sediments. *Quat Sci Rev* 87, 12–23.
1381 <https://doi.org/10.1016/j.quascirev.2013.12.015>
- 1382 Matulewski, P., Buchwal, A., Makohonienko, M., 2019. Higher climatic sensitivity of Scots pine (*Pinus*
1383 *sylvestris* L.) subjected to tourist pressure on a hiking trail in the Brodnica Lakeland, NE Poland.
1384 *Dendrochronologia* (Verona) 54, 78–86. <https://doi.org/10.1016/j.dendro.2019.02.008>
- 1385 Mauquoy, D., Hughes, P.D.M., van Geel, B., 2010. A protocol for plant macrofossil analysis of peat deposits.
1386 *Mires and Peat* 7, 1–5.
- 1387 Mauquoy, D., van Geel, B., 2007. Mire and peat macros, in: *Encyclopedia of Quaternary Science*. Elsevier,
1388 Heidelberg, pp. 2315–2336.
- 1389 Mauquoy, D., Yeloff, D., 2008. Raised peat bog development and possible responses to environmental
1390 changes during the mid- to late-Holocene. Can the palaeoecological record be used to predict the nature
1391 and response of raised peat bogs to future climate change? *Biodivers Conserv* 17, 2139–2151.
1392 <https://doi.org/10.1007/s10531-007-9222-2>
- 1393 Mazei, Y., Tsyganov, A.N., 2006. Freshwater testate amoebae. KMK, Moscow.
- 1394 McCarroll, D., Loader, N.J., 2004. Stable isotopes in tree rings. *Quat Sci Rev* 23, 771–801.
1395 <https://doi.org/10.1016/j.quascirev.2003.06.017>
- 1396 McGrath, M.J., Luyssaert, S., Meyfroidt, P., Kaplan, J.O., Bürgi, M., Chen, Y., Erb, K., Gimmi, U.,
1397 McInerney, D., Naudts, K., Otto, J., Pasztor, F., Ryder, J., Schelhaas, M.-J., Valade, A., 2015.
1398 Reconstructing European forest management from 1600 to 2010. *Biogeosciences* 12, 4291–4316.
1399 <https://doi.org/10.5194/bg-12-4291-2015>
- 1400 McNulty, S., Caldwell, P., Doyle, T.W., Johnsen, K., Liu, Y., Mohan, J., Prestemon, J., Sun, G., 2013. Forests
1401 and Climate Change in the Southeast USA, in: Ingram, K.T., Dow, K., Carter, L., Anderson, J. (Eds.),
1402 *Climate of the Southeast United States*. Island Press, Washington, pp. 165–189.
- 1403 Meisterfeld, R., 2001. Testate amoebae, in: Costello, M.J., Emblow, C.S., White, R. (Eds.), *Patrimoines*
1404 *Naturels*. Muséum National d’Histoire Naturelle - Institut d’Ecologie et de Gestion de la Biodiversité
1405 (I.E.G.B.) - Service du Patrimoine Naturel (S.P.N.), Paris, pp. 54–57.
- 1406 Messier, C., Bauhus, J., Sousa-Silva, R., Auge, H., Baeten, L., Barsoum, N., Bruelheide, H., Caldwell, B.,
1407 Cavender-Bares, J., Dhiedt, E., Eisenhauer, N., Ganade, G., Gravel, D., Guillemot, J., Hall, J.S.,
1408 Hector, A., Hérault, B., Jactel, H., Koricheva, J., Kreft, H., Mereu, S., Muys, B., Nock, C.A., Paquette,
1409 A., Parker, J.D., Perring, M.P., Ponette, Q., Potvin, C., Reich, P.B., Scherer-Lorenzen, M., Schnabel,
1410 F., Verheyen, K., Weih, M., Wollni, M., Zemp, D.C., 2022. For the sake of resilience and
1411 multifunctionality, let’s diversify planted forests! *Conserv Lett* 15. <https://doi.org/10.1111/conl.12829>
- 1412 Milecka, K., Kowalewski, G., Fiałkiewicz-Kozieł, B., Gałka, M., Lamentowicz, M., Chojnicki, B.H.,
1413 Goslar, T., Barabach, J., 2017. Hydrological changes in the Rzecin peatland (Puszcza Notecka, Poland)
1414 induced by anthropogenic factors: Implications for mire development and carbon sequestration.
1415 *Holocene* 27, 651–664. <https://doi.org/10.1177/0959683616670468>
- 1416 Miller, J.D., Anderson, H.A., Ferrier, R.C., Walker, T.A.B., 1990. Hydrochemical Fluxes and their Effects
1417 on Stream Acidity in Two Forested Catchments in Central Scotland. *Forestry* 63, 311–331.
1418 <https://doi.org/10.1093/forestry/63.4.311>
- 1419 Minkinen, K., Byrne, K.A., Trettin, C., 2008. Climate impacts of peatland forestry, in: Strack, M. (Ed.),
1420 *Peatlands and Climate Change*. International Peat Society, Saarijärvi, pp. 98–122.
- 1421 Miola, A., 2012. Tools for Non-Pollen Palynomorphs (NPPs) analysis: A list of Quaternary NPP types and
1422 reference literature in English language (1972–2011). *Rev Palaeobot Palynol* 186, 142–161.
1423 <https://doi.org/10.1016/j.revpalbo.2012.06.010>
- 1424 Mirek, Z., Zarzycki, K., Wojewoda, W., Szelaż, Z., 2006. Red list of plants and fungi in Poland. *Czerwona*
1425 *lista roślin i grzybów Polski*. Instytut Botaniki im. W. Szafera, Polska Akademia Nauk, Kraków.
- 1426 Mitosek, H., 1960. Letnio-jesienna susza 1959 r. *Postępy Nauk Rolniczych* 7, 53–64.
- 1427 Mokrzecki, Z., 1928. *Strzygonia choinówka*. Związek Zawodowy Leśników w Rzeczypospolitej Polskiej,
1428 Warszawa.
- 1429 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen Analysis*. Blackwell Scientific Publications, Oxford.

- 1430 Moritz, M.A., Parisien, M.-A., Batllori, E., Krawchuk, M.A., Van Dorn, J., Ganz, D.J., Hayhoe, K., 2012.
 1431 Climate change and disruptions to global fire activity. *Ecosphere* 3, 1–22.
 1432 <https://doi.org/10.1890/ES11-00345.1>
- 1433 Mroczkowska, A., Kittel, P., Marcisz, K., Dolbunova, E., Gauthier, E., Lamentowicz, M., Mazurkevich, A.,
 1434 Obremaska, M., Płóciennik, M., Kramkowski, M., Łuców, D., Kublitskiy, Y., Słowiński, M., 2021.
 1435 Small peatland with a big story: 600-year paleoecological and historical data from a kettle-hole
 1436 peatland in Western Russia. *Holocene* 31, 1761–1776. <https://doi.org/10.1177/09596836211033224>
- 1437 Nisbet, T.R., 2001. The role of forest management in controlling diffuse pollution in UK forestry. *For Ecol*
 1438 *Manage* 143, 215–226. [https://doi.org/10.1016/S0378-1127\(00\)00519-3](https://doi.org/10.1016/S0378-1127(00)00519-3)
- 1439 Obmiński, Z., 1970. Zarys ekologii, in: Białobok, S. (Ed.), *Sosna Zwyczajna. Nasze Drzewa Leśne*.
 1440 Warszawa-Poznań, pp. 152–231.
- 1441 Ogden, C.G., Hedley, R.H., 1980. *An Atlas of Freshwater Testate Amoebae*. Oxford University Press,
 1442 London.
- 1443 Oris, F., Ali, A.A., Asselin, H., Paradis, L., Bergeron, Y., Finsinger, W., 2014. Charcoal dispersion and
 1444 deposition in boreal lakes from 3 years of monitoring: Differences between local and regional fires.
 1445 *Geophys Res Lett* 41, 6743–6752. <https://doi.org/10.1002/2014GL060984>
- 1446 Orłowicz, M., 1924. *Ilustrowany przewodnik po województwie pomorskiem*. Książnica Polska, Lwów.
- 1447 Paavilainen, E., Päivänen, J., 1995. *Peatland Forestry: Ecology and Principles*, Ecological Studies. Springer,
 1448 Berlin.
- 1449 Parish, F., Sirin, A., Charman, D.J., Joosten, H., Minayeva, T., Silviu, M., Stringer, L., 2008. Assessment
 1450 on peatlands, biodiversity and climate change: main report, Global Environment Centre, Kuala
 1451 Lumpur and Wetlands International, Wageningen.
- 1452 Pawlyta, J., Lamentowicz, M., 2010. Age-depth model for modern peat core. Methodological approach,
 1453 methods of absolute chronology., in: 10th International Conference. GADAM Centre of Excellent.
 1454 Department of Radioisotopes, Institute of Physics, Silesian University of Technology, Gliwice, p.
 1455 109.
- 1456 Payne, R.J., Mitchell, E.A.D., 2009. How many is enough? Determining optimal count totals for ecological
 1457 and palaeoecological studies of testate amoebae. *J. Paleolimnol.* 42, 483–495.
 1458 <https://doi.org/10.1007/s10933-008-9299-y>
- 1459 Pędziszewska, A., Latałowa, M., 2016. Stand-scale reconstruction of late Holocene forest succession on the
 1460 Gdańsk Upland (N. Poland) based on integrated palynological and macrofossil data from paired sites.
 1461 *Veg Hist Archaeobot* 25, 239–254. <https://doi.org/10.1007/s00334-015-0546-7>
- 1462 Péli, E.R., Nagy, J.G., Cserhalmi, D., 2015. In situ measurements of seasonal productivity dynamics in two
 1463 sphagnum dominated mires in Hungary . *Carpathian Journal of Earth and Environmental Sciences* 10,
 1464 231–240.
- 1465 Peters, M.E., Higuera, P.E., 2007. Quantifying the source area of macroscopic charcoal with a particle
 1466 dispersal model. *Quat Res* 67, 304–310. <https://doi.org/10.1016/j.yqres.2006.10.004>
- 1467 Poraj-Górska, A.I., Żarczyński, M.J., Ahrens, A., Enters, D., Weisbrodt, D., Tylmann, W., 2017. Impact of
 1468 historical land use changes on lacustrine sedimentation recorded in varved sediments of Lake Jaczno,
 1469 northeastern Poland. *Catena* (Amst) 153, 182–193.
 1470 <https://doi.org/http://dx.doi.org/10.1016/j.catena.2017.02.007>
- 1471 Przybylski, T., 1993. Ekologia, in: Białobok, S., Boratyński, A., Bugała, W. (Eds.), *Biologia Sosny*
 1472 *Zwyczajnej*. Sorus, Poznań-Kórnik, pp. 255–300.
- 1473 Pureswaran, D.S., De Grandpré, L., Paré, D., Taylor, A., Barrette, M., Morin, H., Régnière, J., Kneeshaw,
 1474 D.D., 2015. Climate-induced changes in host tree–insect phenology may drive ecological state-shift in
 1475 boreal forests. *Ecology* 96, 1480–1491. <https://doi.org/10.1890/13-2366.1>
- 1476 R Core Team, 2022. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical
 1477 Computing [WWW Document]. URL <https://www.R-project.org> (accessed 12.4.23).
- 1478 Rapinel, S., Panhelleux, L., Gayet, G., Vanacker, R., Lemercier, B., Laroche, B., Chambaud, F., Guelmami,
 1479 A., Hubert-Moy, L., 2023. National wetland mapping using remote-sensing-derived environmental

1480 variables, archive field data, and artificial intelligence. *Heliyon* 9, e13482.
1481 <https://doi.org/10.1016/j.heliyon.2023.e13482>

1482 Rastogi, A., Stróżecki, M., Kalaji, H.M., Łuców, D., Lamentowicz, M., Juszczak, R., 2019. Impact of
1483 warming and reduced precipitation on photosynthetic and remote sensing properties of peatland
1484 vegetation. *Environ Exp Bot* 160, 71–80.
1485 <https://doi.org/https://doi.org/10.1016/j.envexpbot.2019.01.005>

1486 Regulation No. 64/97 of the Bydgoszcz Voivode of October 30, 1997, on the recognition as ecological sites
1487 of natural objects in the Bydgoszcz Voivodeship, 1997.

1488 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng,
1489 H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G.,
1490 Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht,
1491 J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F.,
1492 Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S.,
1493 Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S., 2020. The IntCal20 Northern
1494 Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62, 725–757.
1495 <https://doi.org/10.1017/RDC.2020.41>

1496 Reynolds, B., Ormerod, S.J., Gee, A.S., 1994. Spatial patterns concentrations in upland Wales in relation to
1497 catchment forest cover and forest age. *Environmental Pollution* 84, 27–33.
1498 [https://doi.org/10.1016/0269-7491\(94\)90067-1](https://doi.org/10.1016/0269-7491(94)90067-1)

1499 Rouse, J.W., Haas, R.H., Schell, J.A., Deering, D.W., 1974. Monitoring vegetation systems in the Great
1500 Plains with ERTS, Third ERTS Symposium. ed. NASA.

1501 Rydin, H., Jeglum, J.K., 2013. The biology of peatlands (Second Edition). Oxford University Press.

1502 Sanderson, N., Loisel, J., Gallego-Sala, A., Anshari, G., Novita, N., Marcisz, K., Lamentowicz, M., Bąk,
1503 M., Wochal, D., 2023. Setting a new research agenda for tropical peatlands, recent carbon
1504 accumulation and ecosystem services. *Past Global Changes Magazine* 31, 121–121.
1505 <https://doi.org/10.22498/pages.31.2.121>

1506 Schueler, S., Falk, W., Koskela, J., Lefèvre, F., Bozzano, M., Hubert, J., Kraigher, H., Longauer, R., Olrik,
1507 D.C., 2014. Vulnerability of dynamic genetic conservation units of forest trees in Europe to climate
1508 change. *Glob Chang Biol* 20, 1498–1511. <https://doi.org/10.1111/gcb.12476>

1509 Schüle, M., Domes, G., Schwanitz, C., Heinken, T., 2023. Early natural tree regeneration after wildfire in a
1510 Central European Scots pine forest: Forest management, fire severity and distance matters. *For Ecol*
1511 *Manage* 539, 120999. <https://doi.org/10.1016/j.foreco.2023.120999>

1512 Schütte, R., 1893. Die Tucheler Haide vornehmlich in forstlicher Beziehung. Bertling, Danzig.

1513 Seidl, R., Schelhaas, M.-J., Rammer, W., Verkerk, P.J., 2014. Increasing forest disturbances in Europe and
1514 their impact on carbon storage. *Nat Clim Chang* 4, 806. <https://doi.org/10.1038/nclimate2318>
1515 <https://www.nature.com/articles/nclimate2318#supplementary-information>

1516 Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D., Petr,
1517 M., Honkaniemi, J., Lexer, M.J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel, T.A.,
1518 Reyer, C.P.O., 2017. Forest disturbances under climate change. *Nat Clim Chang* 7, 395.
1519 <https://doi.org/10.1038/nclimate3303> [https://www.nature.com/articles/nclimate3303#supplementary-](https://www.nature.com/articles/nclimate3303#supplementary-information)
1520 [information](https://www.nature.com/articles/nclimate3303#supplementary-information)

1521 Shumilovskikh, L.S., Shumilovskikh, E.S., Schlütz, F., van Geel, B., 2022. NPP-ID: Non-Pollen
1522 Palynomorph Image Database as a research and educational platform. *Veg Hist Archaeobot* 31, 323–
1523 328. <https://doi.org/10.1007/s00334-021-00849-8>

1524 Shumilovskikh, L.S., van Geel, B., 2020. Non-Pollen Palynomorphs, in: Henry, E.G. (Ed.), Handbook for
1525 the Analysis of Micro-Particles in Archaeological Samples. Springer Cham, pp. 65–94.

1526 Siemensma, F., 2023. Microworld, world of amoeboid organisms [WWW Document]. URL
1527 <https://arcella.nl/> (accessed 11.23.23).

1528 Sillasoo, U., Väiliranta, M., Tuittila, E.S., 2011. Fire history and vegetation recovery in two raised bogs at
1529 the Baltic Sea. *Journal of Vegetation Science* 22, 1084–1093.

- 1530 Simard, I., Morin, H., Lavoie, C., 2006. A millennial-scale reconstruction of spruce budworm abundance in
1531 Saguenay, Québec, Canada. *Holocene* 16, 31–37. <https://doi.org/10.1191/0959683606hl904rp>
- 1532 Śliwa, E., 1989. Przebieg masowego pojawu brudnicy mniszki (*Lymantria monacha* L.) i jej zwalczania w
1533 Polsce w latach 1978-1985 oraz regeneracja aparatu asymilacyjnego w uszkodzonych drzewostanach.
1534 *Prace Instytutu Badawczego Leśnictwa* 710, 3–120.
- 1535 Śliwa, E., 1987. Występowanie i zwalczanie ważniejszych folio fagów w drzewostanach sosnowych w
1536 latach 1946-1985. *Sylvan* 11–12.
- 1537 Śliwa, E., 1974. Gradacja strzygoni choinówki (*Panolis flammea* Schiff.) w Kampinowskim Parku
1538 Narodowym. *Sylvan* 11, 61–66.
- 1539 Słowiński, M., Lamentowicz, M., Łuców, D., Barabach, J., Brykała, D., Tyszkowski, S., Pieńczewska, A.,
1540 Śnieszko, Z., Dietze, E., Jażdżewski, K., Obremska, M., Ott, F., Brauer, A., Marcisz, K., 2019.
1541 Paleocological and historical data as an important tool in ecosystem management. *J Environ Manage*
1542 236, 755–768. <https://doi.org/10.1016/j.jenvman.2019.02.002>
- 1543 Spiecker, H., 2000. Growth of Norway Spruce (*Picea abies* [L.] Karst.) under Changing Environmental
1544 Conditions in Europe, in: Klimo, E., Hager, H., Kulhavý, J. (Eds.), *Spruce Monocultures in Central*
1545 *Europe – Problems and Prospects*. European Forest Institute, Joensuu, pp. 11–26.
- 1546 Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., Dosio, A., 2018. Will drought events become more
1547 frequent and severe in Europe? *International Journal of Climatology* 38, 1718–1736.
1548 <https://doi.org/10.1002/joc.5291>
- 1549 Stivrins, N., Liiv, M., Ozola, I., Reitalu, T., 2018. Carbon accumulation rate in a raised bog in Latvia, NE
1550 Europe, in relation to climate warming. *Estonian Journal of Earth Sciences* 67, 247.
1551 <https://doi.org/10.3176/earth.2018.20>
- 1552 Stockmarr, J., 1971. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13, 615–621.
- 1553 Sullivan, J., 1993. <https://www.fs.usda.gov/database/feis/plants/tree/pinsyl/all.html> [WWW Document].
1554 *Pinus sylvestris*. In: *Fire Effects Information System*.
- 1555 Sullivan, M.E., Booth, R.K., 2011. The Potential Influence of Short-term Environmental Variability on the
1556 Composition of Testate Amoeba Communities in Sphagnum Peatlands. *Microb Ecol* 62, 80–93.
1557 <https://doi.org/10.1007/s00248-011-9875-y>
- 1558 Szwankowski, J., 2005. Powiat tucholski w latach 1875-1920, administracja, ludność, gospodarka, kultura.
1559 Logo, Tuchola.
- 1560 Taeger, S., Zang, C., Liesebach, M., Schneck, V., Menzel, A., 2013. Impact of climate and drought events
1561 on the growth of Scots pine (*Pinus sylvestris* L.) provenances. *For Ecol Manage* 307, 30–42.
1562 <https://doi.org/10.1016/j.foreco.2013.06.053>
- 1563 Tinner, W., Hu, F.S., 2003. Size parameters, size-class distribution and area-number relationship of
1564 microscopic charcoal: relevance for fire reconstruction. *Holocene* 13, 499–505.
- 1565 Tipton, J., 2023. Past anthropogenic impacts on peatland through the forest management practices in the
1566 Polish Tuchola Forest (Master Thesis). Adam Mickiewicz University, Poznań.
- 1567 Trouet, V., Van Oldenborgh, G.J., 2013. KNMI Climate Explorer: A Web-Based Research Tool for High-
1568 Resolution Paleoclimatology. *Tree Ring Res* 69, 3–13. <https://doi.org/10.3959/1536-1098-69.1.3>
- 1569 Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science* (1979) 349, 814–
1570 818. <https://doi.org/10.1126/science.aac6759>
- 1571 Ulrich, B., 1980. Production and consumption of Hydrogen ions in the ecosphere, in: Hutchinson, T., Havas,
1572 M. (Eds.), *Effects of Acid Precipitation on Terrestrial Ecosystems*. Plenum Press, New York, London,
1573 pp. 255–282.
- 1574 UNESCO, 2024. Tuchola Forests [WWW Document]. *Man and the Biosphere Programme (MAB)*.
- 1575 Väiliranta, M., Korhola, A., Seppä, H., Tuittila, E.S., Sarmaja-Korjonen, K., Laine, J., Alm, J., 2007. High-
1576 resolution reconstruction of wetness dynamics in a southern boreal raised bog, Finland, during the late
1577 Holocene: a quantitative approach. *Holocene* 17, 1093–1107.
1578 <https://doi.org/10.1177/0959683607082550>

- 1579 van der Linden, M., Heijmans, M.M., van Geel, B., 2014. Carbon accumulation in peat deposits from
1580 northern Sweden to northern Germany during the last millennium. *Holocene* 24, 1117–1125.
1581 <https://doi.org/10.1177/0959683614538071>
- 1582 van der Schrier, G., Allan, R.P., Ossó, A., Sousa, P.M., Van de Vyver, H., Van Schaeybroeck, B., Coscarelli,
1583 R., Pasqua, A.A., Petrucci, O., Curley, M., Mietus, M., Filipiak, J., Štěpánek, P., Zahradníček, P.,
1584 Brázdil, R., Řezníčková, L., van den Besselaar, E.J.M., Trigo, R., Aguilar, E., 2021. The 1921
1585 European drought: impacts, reconstruction and drivers. *Climate of the Past* 17, 2201–2221.
1586 <https://doi.org/10.5194/cp-17-2201-2021>
- 1587 Waller, M., 2013. Drought, disease, defoliation and death: forest pathogens as agents of past vegetation
1588 change. *J Quat Sci* 28, 336–342. <https://doi.org/10.1002/jqs.2631>
- 1589 Wardenaar, E.C.P., 1987. A new hand tool for cutting peat profiles. *Canadian Journal of Botany* 65, 1772–
1590 1773. <https://doi.org/10.1139/b87-243>
- 1591 Warner, B.G., 1993. Palaeoecology of floating bogs and landscape change in the Great Lakes drainage basin
1592 of North America, in: Chambers, F.M. (Ed.), *Climate Change and Human Impact on the Landscape*.
1593 Chapman & Hall, pp. 237–248.
- 1594 Westerling, A.L., 2016. Increasing western US forest wildfire activity: sensitivity to changes in the timing
1595 of spring. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371, 20150178.
1596 <https://doi.org/10.1098/rstb.2015.0178>
- 1597 Westman, C.J., Laiho, R., 2003. Nutrient dynamics of drained peatland forests. *Biogeochemistry* 63, 269–
1598 298. <https://doi.org/10.1023/A:1023348806857>
- 1599 Whitlock, C., Larsen, C., 2001. Charcoal as a fire proxy., in: *Tracking Environmental Change Using Lake*
1600 *Sediments. Terrestrial, Algal, and Siliceous Indicators*. J. P. Smol, H. J. B. Birks, and W. M. Last, Eds.
1601 Dordrecht: Kluwer, pp. 75–97.
- 1602 Wilson, J.K., 2012. *The German Forest. Nature, Identity, and the Contestation of a National Symbol, 1871–*
1603 *1914*. University of Toronto Press.
- 1604 Wilson, R.M., Hopple, A.M., Tfaily, M.M., Sebestyen, S.D., Schadt, C.W., Pfeifer-Meister, L., Medvedeff,
1605 C., McFarlane, K.J., Kostka, J.E., Koltun, M., Kolka, R.K., Kluber, L.A., Keller, J.K., Guilderson, T.P.,
1606 Griffiths, N.A., Chanton, J.P., Bridgham, S.D., Hanson, P.J., 2016. Stability of peatland carbon to rising
1607 temperatures. *Nat Commun* 7, 13723. <https://doi.org/10.1038/ncomms13723>
1608 <http://www.nature.com/articles/ncomms13723#supplementary-information>
- 1609 Woodward, C., Haines, H.A., 2020. Unprecedented long-distance transport of macroscopic charcoal from a
1610 large, intense forest fire in eastern Australia: Implications for fire history reconstruction. *Holocene* 30,
1611 947–952. <https://doi.org/10.1177/0959683620908664>
- 1612 Wotton, B.M., Nock, C.A., Flannigan, M.D., 2010. Forest fire occurrence and climate change in Canada.
1613 *Int J Wildland Fire* 19, 253. <https://doi.org/10.1071/WF09002>
- 1614 Young, D.M., Baird, A.J., Charman, D.J., Evans, C.D., Gallego-Sala, A. V., Gill, P.J., Hughes, P.D.M.,
1615 Morris, P.J., Swindles, G.T., 2019. Misinterpreting carbon accumulation rates in records from near-
1616 surface peat. *Sci Rep* 9, 17939. <https://doi.org/10.1038/s41598-019-53879-8>
- 1617 Yu, Z., Loisel, J., Brosseau, D.P., Beilman, D.W., Hunt, S.J., 2010. Global peatland dynamics since the Last
1618 Glacial Maximum. *Geophys Res Lett* 37. <https://doi.org/10.1029/2010GL043584>
- 1619 Zajączkowski, J., Brzeziecki, B., Kozak I., 2013. Wpływ potencjalnych zmian klimatycznych na zdolność
1620 konkurencyjną głównych gatunków drzew w Polsce. *Sylwan* 157, 253–261.
- 1621 Zang, C., Biondi, F., 2015. treeclim: an R package for the numerical calibration of proxy-climate
1622 relationships. *Ecography* 38, 431–436. <https://doi.org/10.1111/ecog.01335>
- 1623 Zielski, A., 1996. Wpływ temperatury i opadów na szerokość słoju rocznych drewna u sosny zwyczajnej
1624 (<s>Pinus sylvestris<s> L. w rejonie Torunia). *Sylwan* 2, 71–80.
- 1625 Zielski, A., Barankiewicz, A., 2000. Dendrochronologiczna analiza przyrostów radialnych sosny zwyczajnej
1626 na terenie leśnictwa Dębie, Nadleśnictwa Włocławek. *Sylwan* 2000, 69–74.
- 1627 Zielski, A., Błaszowski, A., Barankiewicz, A., 1998. Dynamika przyrostu radialnego sosny zwyczajnej
1628 (*Pinus sylvestris* L.) na obszarze leśnym eksploatowanym turystycznie nad jeziorem Wielkie
1629 Partęczyny (Nadleśnictwo Brodnica). *Sylwan* 3, 69–78.

- 1630 Zielski, A., Krapiec, M., 2004. Dendrochronologia. PWN.
- 1631 Zielski, A., Krapiec, M., Koprowski, M., 2010. Dendrochronological data, in: Przybylak, R., Majorowicz,
1632 J., Brázdil, R., Kejna, M. (Eds.), The Polish Climate in the European Context: An Historical Overview.
1633 pp. 191–217.
- 1634 Zielski, A., Sygit, W., 1998. Wpływ klimatu na przyrost radialny sosny zwyczajnej (*Pinus sylvestris* L.),
1635 badania wzdłuż równoleżnika 52°N i transekcje Śląsk-Białowieża. In: In: Breymeyer, A., Roo-
1636 Zielińska, E. (Eds.), Reakcja borów sosnowych na zmianę klimatu wzdłuż równoleżnika 52°N (12-
1637 32°E) oraz na zmiany w depozycji zanieczyszczeń chemicznych na transekcje Śląsk-Białowieża.
1638 Dokumentacja Geograficzna IG PAN Warszawa 13, 161–185.
- 1639