

# Burning conditions and transportation pathways determine biomass-burning aerosol properties in the Ascension Island marine boundary layer

Amie Dobracki<sup>1</sup>, Ernie R. Lewis<sup>1</sup>, Arthur J. Sedlacek III<sup>1</sup>, Tyler Tatro<sup>2</sup>, Maria A. Zawadowicz<sup>1</sup>, Paquita Zuidema<sup>2</sup>

<sup>1</sup>Brookhaven National Laboratory, Upton, NY, USA

<sup>2</sup>Department of Atmospheric Sciences, Rosenstiel School, University of Miami, Miami, Florida, USA

*Correspondence to:* Amie Dobracki (adobracki@bnl.gov) and Paquita Zuidema (pzuidema@miami.edu)

## Abstract:

10 African biomass-burning aerosol (BBA) in the southeast Atlantic Ocean (SEA) marine boundary layer (MBL) is an important contributor to Earth's radiation budget, yet its representation remains poorly constrained in regional and global climate models. Data from the Layered Atlantic Smoke Interactions with Clouds (LASIC) field campaign on Ascension Island (7.95°S, 14.36°W) provide insight into how burning conditions, fuel type, transport pathways, and atmospheric processing affect the chemical, microphysical, and optical properties of BBA between June and September 2017. Ten individual plume  
15 events characterize the seasonal evolution of the BBA properties. Early-season inefficient fires, determined by low refractory black carbon to above-background carbon monoxide mixing ratios ( $r_{BC:\Delta CO}$ ), led to enhanced concentrations of organic- and sulfate-rich aerosols. Mid-season efficient fires, determined by higher  $r_{BC:\Delta CO}$  values, led to  $r_{BC}$ -enriched BBA. A mix of efficient and inefficient fires later in the season resulted in conflicting BBA properties. Prolonged transport (~10 days) through the MBL and lower free troposphere (FT), facilitated chemical and aqueous-phase processing, which led to a reduction in  
20 organic aerosol mass concentrations. This resulted in lower OA to  $r_{BC}$  (OA: $r_{BC}$ ) mass ratios (2–5) in the MBL compared to higher values (5–15) in the nearby FT. These atmospheric and cloud oxidation processes yield more light-absorbing BBA and explain the notably low single scattering albedo at 530 nm ( $SSA_{530}$ ) values (< 0.80) observed in the MBL. This study establishes a robust correlation between  $SSA_{530}$  and OA: $r_{BC}$  across the MBL and FT, underscoring the dependency of optical properties on chemical composition.

25

## 1. Introduction

Biomass burning injects 16-34 Tg of particulate matter into Earth's atmosphere each year, with Africa contributing over 50% of that amount (Akagi et al., 2011; Bond et al., 2013). BBA and the copious amounts of co-emitted gases such as carbon dioxide, carbon monoxide, and ozone, have major effects on air quality, climate change, human health, and Earth's geochemical cycles (Bowman et al., 2009, Bond et al., 2013). Biomass burning is a major global source of black carbon (BC), the second largest contributor to climate warming after carbon dioxide (CO<sub>2</sub>). However, BC is a minor component of the total biomass burning aerosol (BBA), which also includes organic aerosol (OA), sulfate (SO<sub>4</sub>), nitrate (NO<sub>3</sub>), and ammonium (NH<sub>4</sub>). The large impacts of BBA on direct aerosol radiative forcing and aerosol-cloud interactions, both of which contribute substantial uncertainty to Earth's radiative budget (IPCC, 2021), underlie the importance of detailed characterization of its chemical, microphysical, and optical properties.

A deck of persistent marine stratocumulus clouds offshore of southwestern Africa coexists with biomass burning in southern Africa from June to October of each year, creating a natural laboratory to investigate how long-range transported BBA interacts with low clouds far to the west of the continent (Zuidema et al., 2016; Adebisi and Zuidema, 2018). Recent field campaigns have determined that the BBA is commonly transported across the southeast Atlantic Ocean (SEA) in both the free troposphere (FT) and the marine boundary layer (MBL) during these five months (Zuidema et al., 2018; Haywood et al., 2021; Redemann et al., 2021). The BBA in the FT is highly absorbing of shortwave radiation (Pistone et al., 2019; Denjean et al., 2020a; Taylor et al., 2020; Wu et al., 2020; Dobracki et al., 2023) with single scattering albedo (the ratio of the aerosol scattering coefficient to the sum of the aerosol scattering and absorption coefficients) at wavelength 530 nm (SSA<sub>530</sub>) values increasing from 0.80 to 0.95 between June and October in the continental boundary layer across southern Africa (Eck et al., 2013). Measurements in the FT over the SEA show August-September mean SSA<sub>530</sub> values near 0.84 (Pistone et al., 2019; Wu et al., 2020, Dobracki et al., 2023) and measurements in the MBL at Ascension Island (7.95 °S, 14.36 °W), a remote location in the tropical Atlantic, yield even lower SSA<sub>530</sub> values, with June-August monthly-mean values near 0.80 (Zuidema et al., 2018; Che et al., 2022a). The lower SSA<sub>530</sub> values at Ascension Island in the MBL compared to those in the FT above the island have not been previously explained (Barrett et al., 2022; Sedlacek et al., 2022).

Multiple field campaigns have collected spatially comprehensive data sets on aerosol properties across the SEA, yet substantial observation-model discrepancies for this region persist (Chylek et al., 2019; Hodzic et al., 2020; Mallet et al., 2020; Shinozuka et al., 2020; Brown et al., 2021; Doherty et al., 2022; Howes et al., 2023). Measured SSA<sub>530</sub> values in the FT and MBL across the SEA are much lower than those used in many global and regional climate models (Shinozuka et al., 2020; Mallet et al., 2020; Doherty et al., 2022). The aerosol over the SEA is highly oxidized (Hodzic et al., 2020; Dobracki et al., 2023), with measured molecular ratios of organic matter to organic carbon (OM:OC) near 2.25 in the FT in September, exceeding the value of 1.4 that is used in many climate models (Hodzic et al., 2020). Aerosol removal through chemical and photolytic processing is typically not represented in models (Hodzic et al., 2020), which contributes to the underestimation of OM:OC. Solar absorption in the FT over the SEA is also not well captured in climate models because of unrealistic aerosol

chemical composition (Mallet et al., 2021), inaccurate representation of mixing states (Brown et al., 2021), and overestimation  
60 of the rate at which absorbing aerosol descends in the atmosphere, which also leads to underestimation in the aerosol loading  
in the FT (Das et al., 2017). These studies demonstrate that more realistic representations of aerosol properties and processes  
in the SEA are required to yield accurate results.

Much of the research over the SEA has focused on BBA properties in the FT, with fewer studies characterizing and  
constraining BBA properties in the MBL. Here, data from the Layered Atlantic Smoke Interactions with Clouds (LASIC) field  
65 campaign on Ascension Island are used to examine how source regions and conditions (i.e., burning conditions and fuel type),  
chemical processes, and transport pathways affect the chemical, microphysical, and optical properties of the BBA in the MBL  
between June and September of 2017. Results from LASIC are compared to those of Wu et al. (2020) and of Taylor et al.  
(2020), both of whom analyzed BBA properties during the Cloud-Aerosol Radiation Interaction and Forcing Year 2017  
(CLARIFY) aircraft campaign in August-September 2017 (Haywood et al., 2021). The CLARIFY analysis, while based on  
70 only three weeks of data, covered times when BBA was present in only the FT, only the MBL, and both the FT and the MBL.  
We also compare our results to those of Dobracki et al. (2023), who analyzed BBA properties in the FT near the African coast  
for September, 2016 during the ObserVations of Aerosols above CLouds and their IntEractionS (ORACLES 2016-2018)  
campaign (Redemann et al., 2021). Moreover, we investigate whether the strong relationship between  $SSA_{530}$  and the mass  
ratio of OA to BC (OA:BC) observed during ORACLES, documented in Dobracki et al. (2023), also applies to the BBA within  
75 the remote MBL.

We distinguish BBA characteristics that are primarily determined at the source, such as BC core diameter and above-  
background CO ( $\Delta$ CO) mixing ratio, from those that can change during long-range transport, such as chemical composition  
and size distribution, by examining back trajectories for selected time periods when BC mass concentrations in the MBL were  
sufficiently high. We then combine the back-trajectory estimates of BBA source locations with maps of fire density, surface  
80 relative humidity (RH), and land use to help infer the burning conditions and fuel types at the fire sources. We address the  
following questions:

1. *How do the chemical, microphysical, and optical properties of African BBA in the MBL at Ascension Island change  
between June and September?*
2. *How do transport pathways from Africa to Ascension Island impact the chemical, microphysical, and optical properties  
85 of African BBA in the MBL there?*
3. *How do differences in burning conditions and fuel type affect the chemical, microphysical, and optical properties of  
African BBA in the MBL at Ascension Island?*
4. *How do the chemical, microphysical, and optical properties of African BBA in the MBL at Ascension Island differ from  
those in the FT?*
- 90 5. *Is there a clear relationship between  $SSA_{530}$  and OA:BC for the MBL at Ascension Island similar to that reported by  
Dobracki et al. (2023) for the FT closer to the coast?*

In Sect. 2, we introduce the LASIC field campaign, the sampling and instrumentation methods, the data products, and  
the data analysis techniques used in this study. In Sect. 3.1, we identify 10 events with high BC mass concentrations, aggregated  
95 into three temporal regimes distinguished by burning conditions, fuel type, and the conserved tracers BC and  $\Delta$ CO. In Sect. 3.2,

we present the chemical, microphysical, and optical properties of the BBA in the Ascension Island MBL for the 10 events. Next, in Sect. 3.3, we discuss the aerosol transport pathways of the BBA to Ascension Island. In Sect. 4, we discuss the evolution of BC properties (Sects. 4.1-4.2), oxidation processes, and transport pathways (Sects. 4.3-4.4 and the dependence of SSA<sub>530</sub> on BBA chemical properties (Sect. 4.5). Sect. 5 highlights a change in aerosol transport in early September (Sect. 5.1) and discusses remaining questions on enhanced aerosol absorption (Sect. 5.2). Lastly, in Sect. 6, we conclude with a summary of our major findings.

## 2. Methods

### 2.1 Instrumentation and data products

The LASIC field campaign deployed the Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) First Mobile Facility (AMF1) (Zuidema et al., 2018) on Ascension Island between June 2016 and October 2017 to measure aerosol and cloud properties with the goal of constraining uncertainties in BBA aging processes and aerosol-cloud interactions. Here we focus on surface measurements collected from June to September of 2017, as aerosol composition was available only for those times. The utilized data products, quantities measured, and instrumentation suite are listed in Table 1. While the LASIC data are constrained to surface based measurements, they are predominantly representative of the boundary layer surrounding Ascension Island. It is crucial to consider that the clouds were intermittently coupled with the surface during the early morning and evening. The observed intermittent coupling of clouds with the surface during early morning and evening hours highlights a more complex interaction than the constant decoupling observed by aircraft measurements (Zhang and Zuidema., 2019). Despite these complexities, our findings provide a reliable representation of the prevailing boundary layer conditions (Zhang and Zuidema, 2019; Abel et al., 2020; Wu et al., 2020).

Mass concentrations of OA, SO<sub>4</sub>, NO<sub>3</sub>, and NH<sub>4</sub> were measured with the quadrupole Aerosol Chemical Speciation Monitor (ACSM; Ng et al., 2011), which separates the ions of different elemental compositions at each mass-to-charge ratio (m/z) to provide unit-mass resolution. The sampling efficiency of the LASIC ACSM was previously discussed in Barrett et al. (2022); however, a comparison (Appendix A) of the aerosol volume concentration calculated using the ACSM with that calculated from the size distribution measurements from the scanning mobility particle sizer (SMPS) indicates the ACSM sampled effectively. Masses of the refractory BC (rBC) in individual particles within the rBC-diameter range of 80 to 500 nm were measured with the Single Particle Soot Photometer (SP2, DMT), allowing the determination of size distributions of number concentration of rBC-containing particles and mass concentrations of rBC (Sedlacek et al., 2022). CO mixing ratios were measured with the CO/NO<sub>2</sub>/H<sub>2</sub>O integrated cavity output spectroscopy (ICOS) analyzer. (Table 1).

During the ORACLES 2016 campaign non-refractory aerosols were sampled with a High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS, Aerodyne Inc.)(Dobracki et al., 2023). During the CLARIFY campaign, the same type of aerosol was sampled with a Compact Time-of-Flight Aerosol Mass Spectrometer (C-ToF-AMS) (Taylor et al., 2020; Barrett et al., 2022). During the ORACLES 2016 and CLARIFY campaigns, rBC was measured with a four-channel Single

Particle Soot Photometer (SP2, DMT) (Barrett et al., 2022; Sedlacek et al., 2022). Despite these slight differences in instrumentation and sampling strategies, we utilize the data from both campaigns in this study to ensure a comprehensive analysis.

Size distributions of aerosol number concentration for mobility diameters between 10 and 500 nm were measured with the SMPS, and total number concentrations of particles with diameters between 10 and 3000 nm were measured with a Condensation Particle Counter (CPC). Light-scattering coefficients ( $\sigma_s$ ) at wavelengths 450, 550, and 700 nm were measured with the nephelometer (TSI 3563), and light-absorption coefficients ( $\sigma_a$ ) at 465, 530, and 650 nm were measured with the Particle Soot Absorption Photometer (PSAP, Radiance Research). These quantities were taken from the ARM Aerosol Optical Properties Value Added Product (AOP VAP; Flynn et al., 2020), as were values of  $SSA_{530}$  that were calculated from them. The mass absorption cross-section at 530 nm ( $MAC_{530}$ ), calculated as the ratio of the absorption coefficient at 530 nm determined by the PSAP to the rBC mass concentration determined by the SP2, is used below to quantify aerosol absorption enhancement. A mass absorption cross-section of  $7.5 \text{ m}^2 \text{ g}^{-1}$  at 550 nm, which corresponds to  $\sim 7.8 \text{ m}^2 \text{ g}^{-1}$  at 530 nm under the assumption of an inverse relationship between absorption coefficient and wavelength (i.e., absorption Ångström exponent (AAE) equal to unity), is universally used for fresh, uncoated rBC (Bond and Bergstrom, 2006). Therefore, a value of  $MAC_{530}$  substantially greater than this indicates enhanced light absorption (Bond et al., 2013; Zanetta et al., 2016). The absorption Ångström exponent between wavelengths 470 and 660 nm ( $AAE_{470-660}$ ) is the negative of the ratio of the logarithm of the absorption coefficient at 470 nm divided by that at 660 nm to the logarithm of 470 divided by 660. These values were also taken from the ARM AOP VAP.

The aerosol optical properties measured in this study were also compared to those from the ORACLES and CLARIFY campaigns. During ORACLES, the aerosol optical properties were measured with similar instruments as those in the LASIC campaign. Specifically, scattering coefficients at wavelengths of 450, 550, and 700 nm were measured using the TSI 3563 nephelometer, while absorption coefficients at 465, 530, and 650 nm were measured with the PSAP (Radiance Research) (Dobracki et al., 2023). Corrections for the scattering coefficients were applied using both Virkkula and Bond & Ogren methods (Pistone et al., 2019; Dobracki et al., 2023). During the CLARIFY, absorption coefficients at 405, 515, and 660 nm, along with aerosol dry extinction at 405 and 658 nm, were measured using the EXtinction SCattering and Absorption of Light for Airborne Aerosol Research (EXSCALABAR) (Wu et al., 2020; Barrett et al., 2022). EXSCALABAR sampled downstream of a  $1.3 \text{ }\mu\text{m}$  aerodynamic diameter impactor (Taylor et al., 2020), whereas a  $1.0 \text{ }\mu\text{m}$  aerodynamic impactor was positioned upstream of the instruments deployed in LASIC. Variations in impactor cutoff sizes could introduce biases in the measured aerosol optical properties, particularly the scattering coefficients. Nevertheless, data from both campaigns are utilized in this study to facilitate a comparative analysis.

160

Table 1. Instrument and Analysis Techniques Table.

Measurement	Instrument / Technique	Notes	Uncertainty
Mass concentrations of Organic, Nitrate, Sulfate, Ammonium, Chloride particulate matter	Aerosol Chemical Speciation Monitor (ACSM) (Aerodyne Inc.)	Data only available in 2017	30 %
Mass and number concentrations and coating-to-core mass ratio of black carbon-containing particles	Single Particle Soot Photometer (SP2) (Droplet Measurement Technologies)	SP2 disconnected from the sampling line after September 21, 2017. Sampled particles between 80 and 500nm	20 %
Size distribution and total number concentration	Scanning Mobility Particle Sizer (SMPS) (TSI Inc 3936)	Sampled particles between 10 to 500nm	10 %
Total particle number concentration	Condensation Particle Counter (CPC) (TSI Inc.)	Sampled particles between 10 and 3000nm	10 %
Aerosol light-scattering coefficient at 470, 530, and 660 nm	Nephelometer (TSI Inc.)	Data was corrected using both Virkkula and Bond & Ogren corrections.	10 %
Aerosol light-absorption coefficient at 450, 550, and 700 nm	Particle Soot Absorption Photometer (PSAP) (Radiance Research)		
CO mole mixing ratio	CO/NO <sub>2</sub> /H <sub>2</sub> O Analyzer ICOS Los Gatos Research		<1 %
Surface wind speed and direction	Radiosonde	Launched 4x daily	
Single Scattering Albedo at 530 nm (SSA <sub>530</sub> )	$SSA = \frac{cs \text{ at } \lambda_{530}}{(cs+ca)}$	Data products available as an ARM Value Added Product (VAP) file	~15%
Mass Absorption Cross-section at 530 nm (MAC <sub>530</sub> ) <i>f44, f60</i>	$MAC_{530} = \frac{ca \text{ at } \lambda_{530} \text{ Mm}^{-1}}{\text{black carbon } (\mu\text{gm}^{-3})}$ Extracted using positive matrix factorization and k means clustering techniques with ACSM organic aerosol data		~20%
Fraction of black carbon (FrBC)	$FrBC = \frac{\text{rBC \# concentration}}{\text{cpc \# concentration}}$	Analysis performed at Brookhaven National Lab with ME-2 engine and SoFi software	30 %
Fire count / distribution	SUOMI Visible Infrared Imaging Radiometer Suite (VIIRS)	Data available on NASA FIRMS Archive. Data binned in 2°x2° boxes 4x Daily Average at 850mb	5-10%
Meridional and Zonal winds reanalysis data	National Centers for Environmental Prediction (NCEP)		
CO (ppbv) reanalysis data	Copernicus Atmosphere Monitoring Service (CAMS)	Data available on Copernicus Atmosphere Monitoring Service archive	

## 2.2 Data Analysis Techniques

$\Delta$ CO mixing ratios at Ascension Island were calculated by subtracting the 5<sup>th</sup> percentile value of the measured CO mixing ratio from the average value for each month, following Che et al. (2022a), and then converted to mass concentrations. These mass concentrations, together with rBC mass concentrations measured by the SP2, were used to determine mass ratios of rBC: $\Delta$ CO. These mass ratios can be used as an indicator of burning conditions at the source because rBC is chemically inert (Wang 2004; Cape et al., 2012; Lund et al., 2018) and the average lifetime of CO is approximately two months (Khalil et al., 1990), much longer than typical transport times. rBC: $\Delta$ CO was used to classify fires as either inefficient or efficient, with inefficient fires having values of rBC: $\Delta$ CO less than 0.01 and efficient fires having values of rBC: $\Delta$ CO greater than 0.01, consistent with previous classifications (Vakkari et al., 2018; Che et al., 2022a). Surface RH fields provided by the NOAA National Center for Environmental Prediction (NCEP) reanalysis are robustly used to assess the burning condition classification. RH values >50% are generally associated with efficient fires and <50% are generally associated with inefficient fires. The surface RH analysis serves as supplementary information for interpreting the rBC: $\Delta$ CO ratios, rather than as a primary classification criterion, similar to the approach used in Che et al. (2022a). Inefficient fires typically produce relatively more OA and SO<sub>4</sub> and relatively less rBC than efficient fires (Collier et al., 2016; Rickly et al., 2022). The rBC: $\Delta$ CO classification is used as a general framework in this study, with supplementary context provided by additional factors such as surface RH, land use maps, fire locations, and the OA:rBC and SO<sub>4</sub>:rBC ratios. This approach is consistent with methodologies commonly used in other studies investigating biomass burning emissions and plume characteristics in this region (Vakkari et al., 2018; Che et al., 2022a). Although modified combustion efficiency (MCE) may be a better determinant of burning conditions (Collier et al., 2016; Dobracki et al., 2023), this quantity could not be calculated because CO<sub>2</sub> was not sampled at Ascension Island during the LASIC campaign.

The locations of the fires between 12.0 °W-52.0 °E, and 0 °N-34.5 °S, encompass the sources of most of the BBA measured at Ascension Island. Fire locations were determined from fire distributions and counts obtained from NASA's Fire Information for Resource Management System (FIRMS) archive, which uses the Visible Infrared Imaging Radiometer Suite (VIIRS) aboard the S-NPP and NOAA 20 satellites. The vegetation types contributing to the fuel for the fires were identified using annual land use maps from the Moderate Resolution Imaging Spectroradiometer (MODIS), combined with the NOAA Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) back trajectories initialized with Global Data Assimilation System (GDAS) meteorology inputs. Transport pathways were determined using a combination of back trajectories from HYSPLIT and CO mixing ratios from the Copernicus Atmosphere Monitoring Service (CAMS) combined with European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis (ERA5) 950, 800, and 700 hPa winds (Figs. S1a-d).

Positive matrix factorization (PMF) was applied to the ACSM-derived compositions during BBA-laden time periods to apportion the organic mass spectra into various factors, following Aiken et al. (2008), Lanz et al. (2010), and Zhang et al. (2011). This PMF analysis yielded only two factors: oxygenated organic aerosol (OOA) and low-volatility oxygenated organic

aerosol (LV-OOA). This result is not surprising, as BBA dominated the composition during the time periods during which  
 230 PMF was applied. This analysis also provided robust calculations of the ion fractions  $f_{44}$  and  $f_{60}$ , with  $f_{44}$  indicating the  
 presence of the  $\text{CO}_2^+$  ion, a product of oxidation (Canagaratna et al., 2015), and  $f_{60}$  indicating the presence of  $\text{C}_2\text{H}_4\text{O}_2$ , a  
 fragment of levoglucosan, which is a known tracer for BBA (Cubison et al., 2011). Further details on the PMF analysis can be  
 found in Appendix A.

Monthly-mean above-background  $\Delta\text{SO}_4$  mass concentrations were calculated by subtracting  $\text{SO}_4$  mass concentrations  
 235 averaged over times each month when rBC mass concentrations were less than  $20 \text{ ng m}^{-3}$  (Table 2), from the total  $\text{SO}_4$  mass  
 concentration during BBA-laden times (Table 2). This quantity excludes background  $\text{SO}_4$  produced from marine or local  
 sources in the MBL. Background  $\text{SO}_4$  concentrations were typically higher later in the biomass burning season and did not  
 correlate with windspeed, suggesting some remaining BBA residual  $\text{SO}_4$ . Changes in the mass ratios of  $\Delta\text{SO}_4$  to rBC  
 ( $\Delta\text{SO}_4:\text{rBC}$ ),  $\text{OA}:\text{rBC}$ , and  $f_{44}$  can indicate changes in BBA properties from aqueous-phase chemistry and chemical processing  
 240 during transport, respectively. Oxalate, an organic acid that is a well-known tracer of aqueous-phase oxidation of OA that  
 contributes to  $f_{44}$  and can indicate that the aerosol has interacted with cloud (Sorooshian et al., 2010; Ervens et al., 2011).  
 However, the ACSM cannot specifically distinguish oxalate from other species that contribute to  $f_{44}$ .

Table 2.  $\Delta\text{SO}_4$ , rBC, and  $\Delta\text{CO}$  from clean periods

Start time	End time	rBC ( $\mu\text{g m}^{-3}$ ) $\times 10^{-4}$	$\Delta\text{CO}$ (ppb)	$\Delta\text{SO}_4$ ( $\mu\text{g m}^{-3}$ ) $\times 10^{-1}$
6/4/17 3:01	6/6/17 0:26	2.46	1.13	2.40
6/11/17 7:19	6/12/17 20:24	5.36	3.91	2.89
6/20/17 19:18	6/20/17 22:27	6.06	3.96	4.93
7/11/17 17:12	7/12/17 19:32	3.40	0.44	2.18
7/20/17 16:21	7/21/17 13:17	9.79	6.17	1.86
7/25/17 18:34	7/26/17 20:52	6.14	6.24	7.97
7/29/17 20:57	7/31/17 6:03	5.18	6.55	1.01
8/21/17 8:32	8/24/17 2:10	1.41	3.65	8.89
9/13/17 3:18	9/13/17 15:27	3.8	6.53	9.71
9/15/17 2:57	9/16/17 3:16	8.3	1.67	1.36

245

The number fraction of rBC-containing particles (FrBC) in the BBA was calculated by dividing the number  
 concentrations of rBC-containing particles measured by the SP2 by the total particle number concentration measured by the  
 CPC. This value is a lower limit because the SP2 detects only particles with rBC-core diameters between 80 and 500 nm,  
 whereas the CPC detects particles with total diameters between 10 and 3000 nm. However, there were likely very few rBC  
 250 particles with diameters less than 80 nm or greater than 500 nm (Taylor et al., 2020; Dobracki et al., 2023). The FrBC values  
 are not likely influenced by the background aerosol emissions because aerosol number concentrations in the accumulation

mode during these clean periods are only a fraction of the BBA aerosol number concentrations during plume events. We further highlight the differences in aerosol number concentration between the clean and polluted periods in Sect. 3.2.3). The rBC geometric peak diameter ( $rBC_{gpd}$ ) is defined as the diameter at the maximum value of the size distribution of number concentration of rBC-containing particles in the representation  $dN/d\log D$ , and the rBC core mass-equivalent peak diameter ( $rBC_{mpd}$ ) is defined as the diameter at the maximum value of the size distribution of mass concentration in the representation  $dM/d\log D$ . The coating-to-core mass ratio of rBC-containing particles is defined as the daily average of the ratio of the coating mass to the rBC core mass and has an uncertainty of  $\sim 20\%$  (Sedlacek et al., 2022).

### 3 Results

260 A time series of rBC and  $\Delta CO$  spanning from June through September 2017 indicates significant synoptic variability, with times when rBC mass concentrations were near  $1 \mu\text{g m}^{-3}$  interspersed with times with rBC less than  $0.02 \mu\text{g m}^{-3}$  (Fig. 1, see also Pennypacker et al., 2020). Previous studies from LASIC have often classified BBA using monthly means (Zuidema et al., 2018; Carter et al., 2021; Che et al., 2022). However, due to the variability in aerosol loading across the four months, monthly means may not adequately capture the BBA properties in the Ascension Island MBL. Here we explore ten plume  
265 events, denoted P1 to P10, corresponding to rBC mass concentrations exceeding  $0.15 \mu\text{g m}^{-3}$ , the 70<sup>th</sup> percentile of all rBC mass concentrations. These synoptically defined events last from two days to two weeks and may contain BBA from multiple sources, each with its own combustion history. Temporal trends in the plume rBC: $\Delta CO$  mass ratios (Fig. 2) together with maps of fire density, surface RH, land use, and fuel type (Table 3) support a classification of the 10 events into three temporal regimes, reflecting different conditions at the beginning, middle, and end of the biomass-burning period as perceived within  
270 the remote MBL.

275

280




Figure 1. Time series of black carbon mass concentration ( $\mu\text{g m}^{-3}$ ) (black) and  $\Delta\text{CO}$  mixing ratio (ppb) (blue), calculated by removing the bottom 5th percentile of the monthly  $\Delta\text{CO}$  distribution from the total, from June 1, 2017 through September 15, 2017. Pink boxes indicate selected plume events and purple boxes indicate clean events.




Figure 2.  $r\text{BC}:\Delta\text{CO}$  box-whisker distributions for major plume events, with lowest whisker at 10<sup>th</sup> percentile, lowest bar at 25<sup>th</sup> percentile, middle bar at 50<sup>th</sup> percentile, upper bar at 75<sup>th</sup> percentile, and highest whisker at 90<sup>th</sup> percentile. Filled circles represent the mean. Red dashed lines distinguish the three regimes discussed in the text.

Table 3. Mean  $\pm$  standard deviation of rBC,  $\Delta$ CO, rBC: $\Delta$ CO, fire location, fuel type, and burning conditions for each regime.

	rBC ( $\mu\text{g m}^{-3}$ )	$\Delta$ CO (ppb)	rBC: $\Delta$ CO mass ratio ( $10^{-3}$ )	Fire location	Fuel type	Burning conditions
Regime 1 06/05-07/20	0.28 $\pm$ 0.10	32 $\pm$ 11	8.3 $\pm$ 1.1	Angola, DRC	Wood savannas and broadleaf forests	Inefficient
Regime 2 07/21-08/17	0.43 $\pm$ 0.11	39 $\pm$ 13	11 $\pm$ 0.2	Zambia, Mozambique, DRC	Grasslands and savannas	Efficient
Regime 3 08/24-09/11	0.34 $\pm$ 0.11	50 $\pm$ 20	7.1 $\pm$ 0.4	Tanzania, Mozambique, and Zimbabwe	Grasslands and savannas	Inefficient and efficient

*The data presented here encompasses only the plume events in each regime.*

### 330 3.1 Regime-based analysis determined from rBC: $\Delta$ CO

The first five plume events, spanning June 8 to July 20, have relatively low rBC: $\Delta$ CO mass ratios of 0.0083 $\pm$ 0.0011 (mean $\pm$ standard deviation; Fig. 2), indicating inefficient combustion. Four of the events have mean rBC: $\Delta$ CO values near 0.009, with P2 significantly lower. Despite the anomalously low rBC: $\Delta$ CO value of 0.006 $\pm$ 0.001 in P2, the rBC: $\Delta$ CO ratios of the other four plumes suggest that burning conditions remained mostly homogeneous over the six weeks. Most fires were  
335 located near 10°S and west of 30°E (Fig. 3a; near the coast in northern Angola and in western Congo), coinciding with surface RH values greater than 50 % (Fig. 3b). Woodier landscapes (e.g. woody savannas and broadleaf forests) dominated in the burning region (Fig. 3c). These five events, based primarily on their similar rBC: $\Delta$ CO values, are grouped into Regime 1.

Plume events 5-8, extending from July 21 to August 17, extend into the heart of the burning season. The mean rBC: $\Delta$ CO is higher, at 0.0112 $\pm$ 0.0002 (Fig. 2), indicating efficient combustion. Most fires occurred south of 10°S (Fig. 4a; in  
340 Zambia, the Democratic Republic of the Congo (DRC), and western Mozambique), coinciding with surface RH values less than 50 % (Fig. 4b). The burned regions are more grassy and less woody (Fig. 4c) than those burned in Regime 1. These plume events are grouped into Regime 2.

The final two plume events, spanning August 24 to September 11, have the lowest overall rBC: $\Delta$ CO, with a mean of 0.0071 $\pm$ 0.0004. These low rBC: $\Delta$ CO values indicate that the fires were inefficient. Most of the fires occurred east of 30°E and  
345 south of 10°S (Fig. 5a, in northeast Zambia, southwest Tanzania, Mozambique, and Zimbabwe), with surface RH ranging between 30 and 60 % (Fig. 5b), over vegetation types varying from grasslands to woody savannas (Fig. 5c). Some of the fires occurred over dry central Africa and on the eastern African coast where precipitation was greater in September (Ryoo et al., 2021). The variation in surface RH, and vegetation insinuate that the two plumes in this regime may have originated from fires that are both efficient and inefficient. We further describe these conflicting conditions in Section 4. A notable feature of this

350 regime is that the strong free-tropospheric winds known as the African Easterly Jet-South became active around August 20, at  
approximately 700 hPa (Ryoo et al., 2022).

355


360

365

370

375

380



415 Figure 3. a) Regime 1 fire density maps showing  $2^\circ \times 2^\circ$  bins of the number of fires detected by the NASA SUOMI NPP satellite between May 31, 2017 and July 13, 2017. b) NCEP reanalysis Regime 1 mean surface relative humidity, with contours indicating fire number density. c) MODIS annual (2017) mean land use map with Regime 1 fire number density contours. Fire number density is shown for total fires beginning 10 days prior to the regime initiation date.





Figure 4. a) Regime 2 fire density maps showing  $2^{\circ} \times 2^{\circ}$  bins of the number of fires detected by the NASA SUOMI NPP satellite between July 14, 2017 and August 10, 2017. b) NCEP reanalysis Regime 2 mean surface relative humidity, with contours indicating fire number density. c) MODIS annual (2017) mean land use map with Regime 2 fire number density contours. Fire number density is shown for total fires beginning 10 days prior to the regime initiation date.

455


### Regime 3

460




465

470



475

480



485

490

Figure 5. a) Regime 3 fire density maps showing  $2^\circ \times 2^\circ$  bins of number of fires detected by the NASA SUOMI NPP satellite between August 19, 2017 and September 4, 2017. b) NCEP reanalysis Regime 3 mean surface relative humidity, with contours indicating fire number density. c) MODIS annual (2017) mean land use map with Regime 3 fire number density contours. Fire number density is shown for total fires beginning 10 days prior to the regime initiation date.

### 3.2 BBA properties measured at Ascension Island

Here, we present the chemical, microphysical, and optical properties of the African BBA sampled in the Ascension Island MBL. By examining rBC size distributions, rBC coating-to-core mass ratios, and number fraction of rBC-containing particles, we can infer how burning conditions and fuel type varied across the three regimes (Table 4). Additionally, we  
 495 examine changes in non-refractory mass concentrations and values of  $f_{44}$  across the four months to understand how long-range transport affected the chemical composition of African BBA. BBA size distributions indicate whether the BBA had undergone cloud processing or mixed with clean marine air in the MBL. Light-absorption coefficients and light-scattering coefficients are used to assess how the optical properties changed as a result of changing chemical and microphysical properties during long-range transport. Lastly, HYSPLIT back trajectories facilitate an understanding of the BBA transport pathways to  
 500 Ascension Island in June, July, August, and September.


Table 4. Mean  $\pm$  standard deviation of aerosol properties across the three regimes. For plume events only.

	Regime 1 06/05-07/20	Regime 2 07/21-08/17	Regime 3 08/24-09/11
rBC: $\Delta$ CO $\times 10^{-3}$	8.3 $\pm$ 1.1	11 $\pm$ 0.20	7.1 $\pm$ 0.40
rBC <sub>gmd</sub> (nm)	125 $\pm$ 3	134 $\pm$ 5	135 $\pm$ 1
FrBC	0.21 $\pm$ 0.04	0.29 $\pm$ 0.04	0.30 $\pm$ 0.02
rBC coating-to-core mass ratio	2.3 $\pm$ 0.1	2.0 $\pm$ 0.2	2.5 $\pm$ 0.01
Aerosol number concentration (accumulation mode) (cm <sup>-3</sup> )	626 $\pm$ 200	730 $\pm$ 257	590 $\pm$ 240
OA:rBC	3.3 $\pm$ 0.73	2.04 $\pm$ 0.32	3.5 $\pm$ 1.8
OA: $\Delta$ CO	0.026 $\pm$ 0.003	0.021 $\pm$ 0.003	0.023 $\pm$ 0.011
$\Delta$ SO <sub>4</sub> :rBC	2.0 $\pm$ 1.1	1.3 $\pm$ 0.17	1.4 $\pm$ 0.96
$f_{44}$	0.27 $\pm$ 0.01	0.26 $\pm$ 0.01	0.26 $\pm$ 0.03
Acc. Dia. (nm)	165 $\pm$ 8	175 $\pm$ 6	209 $\pm$ 11
MAC <sub>530</sub> m <sup>2</sup> g <sup>-1</sup>	16 $\pm$ 1	15 $\pm$ 1	19 $\pm$ 3
SSA <sub>530</sub>	0.81 $\pm$ 0.01	0.78 $\pm$ 0.02	0.80 $\pm$ 0.01

#### 3.2.1 Black carbon aerosol properties

rBC size distributions, rBC coating-to-core mass ratios, and number fraction of rBC-containing particles provide  
 505 additional information on fuel type variations across the three regimes. The rBC geometric and mass-equivalent peak diameters rBC<sub>gpd</sub> and rBC<sub>mpd</sub>, increase monotonically if undramatically from P1 to P10, with rBC<sub>gpd</sub> increasing from 122 to 136 nm and rBC<sub>mpd</sub> increasing from 193 to 203 nm (Fig. 6, Table 4). Mean FrBC also increases from 0.21 $\pm$ 0.04 during Regime 1 to 0.29 $\pm$ 0.03 for the latter two regimes (Fig. 7a), although internal variability is high. Interestingly, the coating-to-core mass ratio trends opposite to rBC: $\Delta$ CO, decreasing from 2.3 $\pm$ 0.1 in Regime 1 to 2.0 $\pm$ 0.2 in Regime 2 before increasing to 2.50 $\pm$ 0.01 in

510 Regime 3 (Fig. 7b). These variations suggest particles with larger rBC cores also contain a higher fraction of rBC-containing  
515 particles, with coating thicknesses that vary less with rBC core size.



520  
525  
530 Figure 6. Mean  $rBC_{mpd}$  (circles) and  $rBC_{gpd}$  (triangles) for multiple field campaigns in the SEA. Filled colored circles and triangles represent the daily mean geometric peak diameters from the LASIC MBL (black, red), ORACLES FT (blue, yellow) (Dobrcki et al., 2023), CLARIFY MBL (pink, green) (Wu et al., 2020), and the Amazonian Tall Tower Observatory lower FT (cyan) (Holanda et al., 2022). Error bars are the standard deviation of each plume event.

535

540

545

550




Figure 7. Box-whisker distributions of a) FrBC and b) rBC coating-to-core mass ratio with lowest whisker at 10<sup>th</sup> percentile, lowest bar at 25<sup>th</sup> percentile, middle bar at 50<sup>th</sup> percentile, upper bar at 75<sup>th</sup> percentile, and highest whisker at 90<sup>th</sup> percentile. Filled circles represent the mean. Red dashed lines distinguish the three regimes discussed in the text.

### 3.2.2 Organic and sulfate aerosol properties

OA, SO<sub>4</sub>, NO<sub>3</sub>, and NH<sub>4</sub> were also commonly present in the MBL at Ascension Island between June and September of 2017 (Fig. 8). Organics provided 45 % of total aerosol mass over the 10 plume events, with above-background sulfate contributing 33 %, rBC 13 %, and nitrate and ammonium the remaining 9 % (not shown). OA:rBC mass ratios generally decreased from P1 to P9, from 3.3±1.3 (P1) to 2.2±0.8 (P9), before increasing to a maximum of 4.8±1.9, during P10 (Fig. 9a, Table 4). The trend in ΔSO<sub>4</sub>:rBC approximately follows that in OA:rBC across the 10 plumes (Fig. 9b; Table 4) but is more pronounced, decreasing from 3.2±1.0 in P2 to 0.82±0.30 in P9 and then increasing to 2.2±0.9 in P10.




Figure 8. Time series of the mass concentrations of bulk chemical species OA, NO<sub>3</sub>, ΔSO<sub>4</sub>, and NH<sub>4</sub> from the ACSM, and rBC from the SP2. Pink boxes indicate selected plume events and purple boxes indicate clean events. A color vision deficiency friendly version of this figure can be found in the supplement.

The two factors (OOA and LV-OOA) generally had similar peak intensities at  $m/z$  values of 18, 28, 43, and 44, and only slight differences in minor peaks (Figs. S2 and S3). Most of the plumes had similar  $f_{44}$  values (Fig. S3), with the exception of P5 and P10 (Fig. 10), for which Factor 1  $f_{44}$  values were  $0.23 \pm 0.01$  and  $0.26 \pm 0.01$ , respectively, and Factor 2  $f_{44}$  values were  $0.29 \pm 0.01$  and  $0.313 \pm 0.002$ , respectively. These differences possibly result from regime transitions and/or more complex fuel source mixtures.  $f_{44}$  values from both Factor 1 and Factor 2 for the other 8 plumes were not substantially different; therefore, averages of the two factors are used to describe each of the 10 events. These values ranged between 0.24 and 0.28 (Fig. 10), indicating highly-oxidized BBA. Mean  $f_{44}$  values decrease only slightly from  $0.284 \pm 0.001$  in Regime 1 to  $0.26 \pm 0.03$  in Regime 3 (Table 4) but may still provide information about transport and oxidation processes in the MBL.

615

620




Figure 9. Box-whisker distributions of a) OA:rBC, b)  $\Delta$ SO<sub>4</sub>:rBC with lowest whisker at 10<sup>th</sup> percentile, lowest bar at 25<sup>th</sup> percentile, middle bar at 50<sup>th</sup> percentile, upper bar at 75<sup>th</sup> percentile, and highest whisker at 90<sup>th</sup> percentile. Filled circles represent the mean. Red dashed lines distinguish the three regimes discussed in the text.




Figure 10. Box-whisker distributions of *f44* factor 1 (red) and *f44* factor 2 (blue). Filled white circles represent the mean of each factor with the lowest whisker at the 10<sup>th</sup> percentile, the lowest bar at the 25<sup>th</sup> percentile, the middle bar at the 50<sup>th</sup> percentile, the upper bar at the 75<sup>th</sup> percentile, and the highest whisker at the 90<sup>th</sup> percentile. Red dashed lines distinguish the three regimes discussed in the text. Red-filled circles represent the mean of factor 1 and factor 2.

### 3.2.3 Size distributions

665 The shape of the aerosol size distribution provides information on whether or not the aerosol has undergone cloud processing. A bimodal size distribution has an Aitken mode at small diameters and an accumulation mode at larger diameters, with the minimum between these modes referred to as the Hoppel minimum (Hoppel et al., 1986). The bimodality results from the activation of larger particles into cloud drops, which then either precipitate or evaporate to form even larger aerosol particles due to the uptake of other substances and coagulation after collisions with other cloud drops. The smaller particles are typically  
670 formed from gas-to-particle conversion of organic and sulfate vapors from the ocean or the biomass burning plume, and have not activated. Therefore, bimodal aerosol size distributions in the MBL suggest aerosol that has undergone cloud processing, be it clean marine aerosol or BBA (Figs. 11, S4).

Daily-mean number size distributions during clean time periods in Regime 1 were always bimodal, with noticeable Aitken and accumulation modes (Fig. 11a). The average of all clean size distributions (in the representation  $dN/d\log D$ ) had an  
675 Aitken mode with a peak near  $200\text{ cm}^{-3}$  at a diameter near 35 nm and an accumulation mode with a peak also near  $200\text{ cm}^{-3}$  at a diameter near 135 nm. The Hoppel minimum diameter was near 60 nm. Daily-mean size distributions during BBA-laden times in Regime 1 were bimodal on 25 of the 28 days (Fig. S4a). The average of the bimodal size distributions during BBA-laden times had an Aitken mode with a peak  $200\text{ cm}^{-3}$  at a larger diameter than that during the pristine period, near 45 nm, and an accumulation mode with a much larger peak than that during the pristine period ranging from  $400\text{--}700\text{ cm}^{-3}$  at a diameter  
680 near 165 nm (Fig. 11a), with a Hoppel minimum diameter near 70 nm.

The pristine size distributions in Regime 2 were also bimodal (Fig. 11b) but with a more pronounced Aitken mode than during Regime 1, with number concentrations greater than  $500\text{ cm}^{-3}$ , at a diameter near 35 nm. The accumulation mode peaks at slightly less than  $250\text{ cm}^{-3}$  at a slightly larger diameter, near 145 nm, but the Hoppel minimum diameter remained near 60 nm. In contrast to Regime 1, only slightly more than half (12 out of 20) of the BBA-laden days in Regime 2 had  
685 bimodal size distributions (Fig. S4b). When BBA was present, the Hoppel minimum was less pronounced than that in Regime 1, spanning a broader range of diameters between 45 and 85 nm. The Aitken mode for the average of these bimodal size distributions was also less pronounced, with peak number concentrations less than  $150\text{ cm}^{-3}$  occurring at diameters between 30 and 50 nm. The accumulation mode peak number concentration ranged from 400 and  $900\text{ cm}^{-3}$  and occurred at slightly larger diameters than for Regime 1, near 175 nm (Fig. 11b).

690 Lastly, in Regime 3, the number size distributions from the clean time periods were more weakly bimodal than in Regimes 1 and 2 (Fig. 11c), primarily because the accumulation mode was less well defined. The Hoppel minimum diameter remained between 60 and 80 nm, similar to Regimes 1 and 2. The Aitken mode for the average of these size distributions had a maximum number concentration near  $250\text{ cm}^{-3}$  at a smaller diameter, near 20 nm, than in the two other regimes, and the accumulation mode was less pronounced, with a maximum number concentration near  $100\text{ cm}^{-3}$  at diameters around 150 nm.  
695 Only 4 out of the 12 polluted days in Regime 3 exhibited a bimodal number size distribution (Fig. S4c), which is fewer than

in both Regimes 1 and 2. The monomodal number size distributions only contained the accumulation mode which had number concentrations between 400 and 800  $\text{cm}^{-3}$  and had a modal diameter near 200 nm, the largest of the three regimes.





Figure 11. Plume mean SMPS aerosol size distributions ( $dN/d\log D \text{ cm}^{-3}$ ) for each selected plume event in a) Regime 1, b) Regime 2, c) Regime 3. Monthly mean distributions from clean periods are denoted as black dashed lines.

### 730 3.2.4 Optical Properties

There was no obvious trend in  $MAC_{530}$  (Fig. 12a),  $SSA_{530}$  (Fig. 12b), or  $AAE_{470-660}$  (not shown) over the 10 plume events but  $MAC_{530}$  and  $SSA_{530}$  do correlate positively over the first 9 plume events (Fig. S5). Plume-mean  $MAC_{530}$  values remained between 15 and 18  $m^2 g^{-1}$  during P1 to P9 and increased only to  $20 \pm 4 m^2 g^{-1}$  in P10 (Fig. 12a). The lowest  $MAC_{530}$  value,  $15 \pm 2 m^2 g^{-1}$ , was observed during P1 (Fig. 12a), but this is statistically indistinguishable from the campaign mean of  $16 \pm 2 m^2 g^{-1}$ .  $SSA_{530}$  varied between  $0.75 \pm 0.01$  and  $0.83 \pm 0.02$  for individual plume events, with a campaign mean of  $0.80 \pm 0.01$  (Fig. 12b). Lower  $SSA_{530}$  values were typically associated with higher FrBC and lower OA:rBC and  $\Delta SO_4$ :rBC. For example, the lowest  $SSA_{530}$ ,  $0.75 \pm 0.01$ , in P6, also had the second highest FrBC ( $0.32 \pm 0.04$ ) and low OA:rBC and  $\Delta SO_4$ :rBC ( $2.3 \pm 0.8$  and  $1.1 \pm 0.4$ , respectively), whereas the highest  $SSA_{530}$ ,  $0.83 \pm 0.02$ , in P2, corresponded to the lowest FrBC ( $0.17 \pm 0.04$ ) and high OA:rBC and  $\Delta SO_4$ :rBC ( $4.5 \pm 1.3$  and  $3.2 \pm 1.0$ , respectively).  $AAE_{470-660}$  had a campaign mean of  $1.03 \pm 0.04$  and varied little between plume events.




745  
750  
755 Figure 12. Box-whisker distributions of a)  $MAC_{530}$  and b)  $SSA_{530}$  with lowest whisker at 10<sup>th</sup> percentile, lowest bar at 25<sup>th</sup> percentile, middle bar at 50<sup>th</sup> percentile, upper bar at 75<sup>th</sup> percentile, and highest whisker at 90<sup>th</sup> percentile. Filled circles represent the mean. Red dashed lines distinguish the three regimes discussed in the text.

### 3.3 BBA transport to Ascension Island


Differences in the transport patterns between clean and BBA-laden days highlight the conditions that allow BBA to reach the remote island. HYSPLIT back trajectories, initialized at 800, 1100, and 1400 m, are shown with NCEP reanalysis 850 hPa wind speeds for representative days with and without BBA at Ascension Island (Figs. 13a-d, Figs. 14a-d). The 8 selected days are representative of the overall transport processes occurring during June through September.

A common feature on clean days (Figs. 14a-d) was the presence of a sea-level pressure high between 0° and 20°W, which promoted the advection of pristine air from the southern oceans (also noted by Pennypacker et al., 2020) and forced the BBA to north of Ascension Island. On polluted days, the lack of a sea-level pressure high allowed easterly winds to transport  
 765 BBA to Ascension Island in the upper levels of the MBL or lower levels of the FT.



790 Figure 13. HYSPLIT back trajectories at 800 m (red), 1100 m (blue), and 1400 m (green) above sea level for polluted days  
 791 initiated at 1200 UTC for a) June 24, 2017, b) July 23, 2017, c) August 06, 2017, and d) September 11, 2017. Each filled  
 792 colored circle represents the trajectory location at 00 UTC after initiation. Grey solid lines represented NCEP reanalysis  
 793 geopotential height at 850 hPa. Wind direction and magnitude are represented by arrows colored by windspeed (0-25m s<sup>-1</sup>).  
 794 Fire density is shown in red contours for total fires two weeks prior to the initiation date.

795



835 Figure 14. HYSPLIT back trajectories at 800 m (red), 1100 m (blue), and 1400 m (green) above sea level for clean days initiated at 1200 UTC for clean days in e) June 5, 2017, f) July 26, 2017, g) August 22, 2017, and h) September 14, 2017. Each filled colored circle represents the trajectory location at 00 UTC after initiation. Grey solid lines represented NCEP reanalysis geopotential height at 850 hPa. Wind direction and magnitude are represented by arrows colored by windspeed (0-25m s<sup>-1</sup>). Fire density is shown in red contours for total fires two weeks prior to the initiation date.

840 In June, the BBA transport to Ascension Island occurred mostly near 850 hPa, in the upper levels of the MBL, with a transit time of approximately 10 days. The lack of a southeast Atlantic sea-level pressure high near the African continent allows a direct if slow westward movement of BBA. Back trajectories from June 24 (Fig. 13a) demonstrate the BBA originated from northern Angola and the southern region of the DRC, in agreement with the conclusion above based on fire density maps

(Fig. 3a). Additional HYSPLIT back trajectories (not shown) highlight that P2 with the anomalously low rBC: $\Delta$ CO ratio also originated from northern Angola, which suggests that the source of this plume was likely similar to those from the other four plume events in this regime.

The BBA transport in late July also occurred near 850 hPa, but because the MBL was shallower than in June (Zhang and Zuidema 2021), more of the transport may have occurred in the lower FT, with the BBA later becoming entrained into the MBL further away from the continent. The transport took less time ( $\sim$ 8 days) because of stronger winds further south near 20°S (Fig. 13b), despite most of the fires being located more inland, in central Africa near southern Angola and from the central region of the DRC, similar to the monthly-mean fire density maps (Fig. 4a). The BBA transport in early August was similar, also occurring near 850 hPa and requiring  $\sim$ 8 days. However, the fire locations shifted further south and east to southern Angola and western Zambia (Fig. 13c), also seen on the fire density maps (Fig. 5a).

The African Easterly Jet South became active after Aug. 20 causing a dramatic switch in the BBA transport to much higher in the FT, near 700 hPa (Ryoo et al., 2022). This would imply that little BBA was transported to the MBL at Ascension Island, despite more African fires occurring in September than any other month (Fig. 5a). Nevertheless, one last BBA-laden episode was observed in early September (Fig. 12d). The back trajectories on September 11 suggest that the BBA may have arrived after  $\sim$ 8 days from fires originating in Mozambique and the DRC. This last plume event is further investigated in Sect. 5.1.

#### 860 4. Discussion

The continuous sampling of BBA in the MBL during the LASIC campaign provides a unique opportunity to characterize the evolution of the African BBA properties across the biomass-burning season. We first describe how burning conditions and fuel type, as inferred from rBC: $\Delta$ CO (Sect. 3.1), affect the microphysical and chemical properties of BBA. We then discuss how heterogeneous oxidation and evaporation, along with transport pathways and aqueous-phase processes, alter the composition and masses of the BBA particles by the time they reach the Ascension Island MBL. Lastly, we examine the dependence of SSA<sub>530</sub> on the chemical properties of the BBA and explain why the BBA in the MBL has a lower SSA<sub>530</sub> than that in the FT. Throughout this discussion, we also compare our results to those from the ORACLES and CLARIFY campaigns, which sampled African BBA in the FT and MBL between the African coast and Ascension Island, and those from the Amazonian Tall Tower Observatory (ATTO) site (Holanda et al., 2020, 2023), which sampled African BBA in Brazil, beyond Ascension Island, in September 2014. These comparisons are summarized in Table 5 and are discussed further below.

875

Table 5. Field campaign comparison table (mean  $\pm$  standard deviation)

	LASIC (June 2017)	LASIC (July 2017)	LASIC (August 2017)	LASIC (September 2017)	CLARIFY (August 2017 MBL)	ATTO (LPL)
Age (days)	10+	8-10	~8	~8-10	--	10+
<i>f44</i>	0.27 $\pm$ 0.01	0.26 $\pm$ 0.02	0.25 $\pm$ 0.01	0.29 $\pm$ 0.005	0.2-0.25	~0.30
OA (mass frac. %)	43	39	45	50	52	51
SO <sub>4</sub> (mass frac. %)	37	38	32	30	26	23
NO <sub>3</sub> (mass frac. %)	3	3	2	4	2	1
NH <sub>4</sub> (mass frac. %)	7	6	6	5	9	13
rBC (mass frac. %)	10	15	14	11	11	11
OA:rBC	3.34 $\pm$ 0.7	2.27 $\pm$ 0.4	2.32 $\pm$ 0.2	4.82 $\pm$ 1.85	3.8-6.25	4.7 $\pm$ 0.7
rBC: $\Delta$ CO (x10 <sup>-3</sup> )	8.7 $\pm$ 2	9.7 $\pm$ 0.7	8.7 $\pm$ 3	7 $\pm$ 2	6-11	3 $\pm$ 2
SSA <sub>530</sub>	0.81 $\pm$ 0.02	0.79 $\pm$ 0.03	0.79 $\pm$ 0.02	0.80 $\pm$ 0.03	0.8-0.86 (658nm)	--
MAC <sub>530</sub>	16 $\pm$ 1.7	15 $\pm$ 0.5	16.6 $\pm$ 1.8	20.45 $\pm$ 4.3	--	--
Median dia. (nm)	167 $\pm$ 9	167 $\pm$ 7	191 $\pm$ 15	217 $\pm$ 15	197-213	105 (optical)
FrBC (%)	23 $\pm$ 5	26 $\pm$ 6	29 $\pm$ 6	26 $\pm$ 5	33 $\pm$ 10	9 $\pm$ 2
rBC <sub>gmd</sub> (mm)	124 $\pm$ 2	132 $\pm$ 6	133 $\pm$ 2	136 $\pm$ 3	117	--

#### 910 4.1 Dependence of BBA properties on burning conditions and fuel type

In this section, we discuss how burning conditions and fuel types change across the three temporal regimes, and how these changes may affect BBA properties such as FrBC, rBC coating-to-core mass ratios, and OA:rBC and  $\Delta\text{SO}_4$ :rBC mass ratios. These analyses offer further BBA characterization than what has typically been presented for biomass-burning events. However, given the limited data set, we cannot definitively describe conditions over land and instead rely on supporting data  
915 from previous studies in this region to interpret the observed patterns.

The mean rBC: $\Delta\text{CO}$  values in Regime 1 were less than 0.01, indicating that the fires across the woodlands and savannas of Angola and the southern DRC were inefficient, as concluded above (Figs. 3a, 13a). The high OA:rBC and  $\Delta\text{SO}_4$ :rBC ratios sampled in this regime had elevated *f44* values and extended transport times which support the interpretation that inefficient fires at the source may have emitted greater amounts of OA relative to rBC. The BBA in this regime also had  
920 large rBC coating-to-core mass ratios and low FrBC (Sects. 3.2-3.3; Figs. 6-7). These results are consistent with those from the Southern African Regional Science Initiative 2000 campaign (SAFARI 2000; Sinha et al., 2004), which demonstrated that woodland fires across central Africa are inefficient, emitting approximately three times as much OA mass as rBC mass.

The mean rBC: $\Delta\text{CO}$  values in Regime 2 were greater than 0.01, indicating that fires across the grasslands of Angola, the DRC, and Zambia were efficient, as concluded above (Figs. 4a, 13b). The low OA:rBC and  $\Delta\text{SO}_4$ :rBC values sampled in  
925 this regime had lower *f44* values and faster transport times which support the interpretation that efficient fires at the source may have emitted less OA relative to rBC. In contrast to BBA properties from Regime 1, the BBA from efficient fires in this regime had small coating-to-core mass ratios, high FrBC, and low  $\Delta\text{SO}_4$ :rBC values (Sects. 3.2-3.3; Figs. 6-7). These results are also consistent with those from the SAFARI 2000 campaign, which showed that burning conditions became more efficient as the soil moisture content across Zambia decreased by 80 % between June and August and rBC emissions increased relative  
930 to those of OA (Hoffa et al., 1999; Korontzi et al., 2003).

The mean rBC: $\Delta\text{CO}$  values in Regime 3 were less than 0.01, indicating that the fires across the DRC and Mozambique were inefficient, as concluded above (Figs. 5a, 13c-d). However, the low OA:rBC values, low  $\Delta\text{SO}_4$ :rBC values, and high FrBC observed during P9 (late August) suggest that BBA during this plume event originated from efficient fires. A similar discrepancy is seen in P10 (early September), where high OA:rBC, large rBC coating-to-core mass ratios, and high  $\Delta\text{SO}_4$ :rBC  
935 would suggest that the fires were inefficient, while the high FrBC would indicate that the fires were efficient. These conflicting results imply that despite the overall low rBC: $\Delta\text{CO}$  values in Regime 3, the BBA from earlier in this regime likely resulted from efficient fires across central African grasslands, whereas the BBA from later in this regime likely resulted from both efficient fires across the grasslands and inefficient fires near the eastern coast (Jiang et al., 2020). These intriguing BBA properties observed in early September are further investigated in Sect. 5.1. Overall, the efficient fires in late August and the  
940 combination of efficient and inefficient fires in September are consistent with observations reported by Che et al. (2022a), who concluded that burning conditions in this region become less efficient as cloud cover increases, precipitation increases, surface wind speed decreases, and soil moisture increases from August to October.

## 4.2 Black carbon core size properties

945 Average  $rBC_{mpd}$  values from the current study generally compare well with those from other studies in this region  
(Fig. 6). The average  $rBC_{mpd}$  value of 180 nm observed in the MBL in August during the CLARIFY campaign is slightly  
smaller (although still within instrument uncertainty) than the corresponding LASIC value of 200 nm, while the average  $rBC_{mpd}$   
value of 200 nm observed in the FT closer to the African coast in September during the ORACLES campaign (Dobracki et al.,  
2023) is the same as that sampled during LASIC at this time. The average  $rBC_{mpd}$  value of 238 nm sampled at the ATTO site,  
950 further west, in September of 2014, was considerably larger than that sampled between the African coast and Ascension Island  
during LASIC, CLARIFY, and ORACLES (Fig. 6). Explanations include differences in burning conditions and fuel types,  
which seems less likely than a combination of burning and meteorological conditions that select for larger rBC (Adebiyi and  
Zuidema, 2016), possibly enhanced by opportunities for the coagulation of rBC cores over the  $\sim 10$  day transport time.

Both the  $rBC_{gpd}$  and  $rBC_{mpd}$  increased between June and September, with the largest values of 140 and 210 nm,  
955 respectively, occurring when the rBC: $\Delta$ CO mass ratio was the highest (0.011) in P7 (late July). This suggests that efficient  
fires produce either larger rBC core diameters or more and smaller rBC particles that coagulate to form larger rBC particles as  
the BBA undergoes long-range transport (Pan et al., 2017). However, large  $rBC_{gpd}$  and  $rBC_{mpd}$  values of 135 and 205 nm,  
respectively, were also observed during Regime 3, in late August and early September when burning conditions were less  
efficient (rBC: $\Delta$ CO = 0.007). These results are consistent with Holder et al. (2016), who conclude woodier, inefficient fires  
960 across the southeast United States emit larger rBC particles than those from efficient grassland fires. Yet, smaller  $rBC_{gpd}$  and  
 $rBC_{mpd}$  values in Regime 1 (125 and 195 nm) occurred when fires were also less efficient (rBC: $\Delta$ CO=0.008), which is  
contradictory to the higher values of  $rBC_{gpd}$  and  $rBC_{mpd}$  in Regime 3 that support Holder et al., (2016). These contrasting results  
suggest that further research is needed on the dependence of rBC core diameters on burning conditions and fuel types.

## 4.3 Heterogeneous oxidation as an aerosol mass removal mechanism in the MBL

965 The evolutions of  $f_{44}$  and  $f_{60}$  help quantify changes in aerosol chemical composition from heterogeneous and  
aqueous-phase oxidation, and thus to determine if heterogeneous oxidation is a dominant chemical process that can explain  
the low mass ratios of OA:rBC (Fig. 9a). In fresh BBA,  $f_{44}$  values are typically near 0.05 and those of  $f_{60}$  are typically near  
0.04 (Cubison et al., 2011; Garofalo et al., 2019). Heterogeneous oxidation can increase  $f_{44}$  through the formation of carboxylic  
acid groups, and decrease  $f_{60}$  through the fragmentation of levoglucosan ( $C_6H_{10}O_5$ ) (Ng et al., 2010; Canonaco et al., 2015).  
970 Heterogeneous oxidation can also fragment carbon-carbon bonds in the OA, allowing the smaller fragments to subsequently  
evaporate, because of the higher vapor pressure of compounds with lower molecular weight (Kroll et al., 2009; 2015).

ORACLES data show  $f_{44}$  increasing from 0.18 to 0.22 and  $f_{60}$  decreasing from 0.006 to 0.004 after 4 to 6 days of  
transport in the free troposphere (Fig. 15; Dobracki et al., 2023). This study also shows  $f_{44}$  increasing and  $f_{60}$  decreasing with

increasing transport time. Eight-day-old plumes at Ascension Island had  $f_{44}$  values near 0.24 and  $f_{60}$  values near 0.005, whereas  $f_{44}$  values for ten-day-old plumes, occurring in June, were near 0.29, and  $f_{60}$  values were lower, near 0.003 (Fig. 15). The coupled temporal increase in  $f_{44}$  and decrease in  $f_{60}$  with increasing transport time observed in both the ORACLES and the LASIC campaigns demonstrate that heterogeneous oxidation may still be continuing for up to ten days, with contributions from aqueous-phase oxidation discussed in Section 4.4. This result is particularly noteworthy because to date, few studies have observed evolution in  $f_{44}$  for times greater than a few hours (see also Che et al., 2022b; Dang et al., 2022).




Figure 15.  $f_{44}$  versus  $f_{60}$  with LASIC plume event data (filled squares), colored by month, ORACLES 2016 level flight average data where  $OA > 20 \mu\text{g m}^{-3}$  (black squares) (Dobracki et al., 2023). Error bars represent the standard deviation of the data set.

Changes in  $f_{44}$  and  $f_{60}$  indicate that oxidation has occurred, but do not capture simultaneous loss of OA with transport time and oxidation. Fragmentation has previously been invoked to explain why  $OA:rBC$  mass ratios decrease by 30% in the FT between the African coast and the ORACLES sampling locations (Dobracki et al., 2023). Here, the relationship between changes in  $f_{44}$  and in the  $OA:rBC$  mass ratio during the ORACLES, CLARIFY, and LASIC campaigns and at the ATTO site is used to determine the extent of OA mass loss caused by heterogeneous oxidation across the SEA (Fig. 16).  $f_{44}$  values were typically near 0.18 and  $OA:rBC$  values were near 8 for BBA sampled near the African coast during the ORACLES campaign (Dobracki et al., 2023).  $f_{44}$  values were greater than 0.22 and  $OA:rBC$  values were less than 5 for BBA sampled in the FT and MBL near Ascension Island (Wu et al., 2020; Dang et al., 2022).  $f_{44}$  values increase further to near 0.30 while  $OA:rBC$  values decrease to less than 2.5 for BBA sampled at the ATTO site in Brazil (Holanda et al., 2020). The BBA sampled at ATTO was

transported primarily within the FT, facilitated by the southern African Easterly Jet, resulting in rapid transport at higher altitudes. As a result, the BBA reached Brazil in a similar time frame to Ascension Island, which likely explains the similarities in  $f_{44}$  and OA:rBC values between the two studies. Notably, the lower pollution layer (LPL) in their study corresponds to the lower FT and not the MBL. As removal through precipitation does not occur in the southeast Atlantic free troposphere during the June-September time period (Adebiyi et al., 2015), these results suggest that heterogeneous oxidation is likely the primary process causing loss of OA across the SEA and that it continues to occur at time scales longer than four days. Loss of OA mass, also attributed to heterogeneous oxidation, is also evident in decreasing rBC coating-to-core mass ratios with increasing transport time (Sedlacek et al., 2022).




Figure 16. OA:rBC versus  $f_{44}$  for the FT during ORACLES (2016-2018) (red), CLARIFY FT (2017) (blue) (Wu et al., 2020), ATTO (2014) (black) (Holanda et al., 2020), and for the LASIC MBL (2017) (green). Error bars represent the standard deviation of the data set.

The rBC coating-to-core mass ratios between 2 and 4 observed during LASIC were also much lower than the values between 5 and 10 observed in the FT for African BBA aged between 7 and 10 days (Sedlacek et al., 2022). This indicates that most of the coating had been removed by the time the BBA reaches Ascension Island.

One other aerosol removal mechanism we consider in the MBL is through thermodynamic evaporation. Results from the ORACLES campaign found OA:rBC decreases from only  $14.5 \pm 2.6$  to  $11.7 \pm 1.9$  as the air temperature increased from  $5^\circ\text{C}$  to  $35^\circ\text{C}$  (Fig. S6), suggesting some thermodynamic evaporation may contribute to OA mass loss. Given MBL air temperatures remain between  $20$  and  $25^\circ\text{C}$ , evaporation is likely not an important loss mechanism, however. Another reason why the OA:rBC ratio is substantially lower in the MBL than in the FT may be higher relative humidities in the MBL encourage faster heterogeneous oxidation (fragmentation) (Wong et al., 2015; Li et al., 2018). Ascension Island RH regularly exceeds 65 %

(Zhang and Zuidema., 2021). During ORACLES, however, OA:rBC mass ratios are 10 % greater when RH is greater than 60 % (Dobracki et al., 2023), again suggesting humidity-induced oxidation may not be dominating. We next consider aqueous-phase processing.

#### 1050 **4.4 Aqueous-phase processes and aerosol transport pathways are powerful determinants of aerosol size distributions, *f44*, and $\Delta\text{SO}_4$ variability**

BBA in the MBL have opportunity to directly interact with the semi-permanent deck of marine stratocumulus clouds between the African coast and Ascension Island during their transit. These aerosol-cloud interactions could encourage a net loss of BBA through wet removal (the formation of cloud droplets subsequently removed by precipitation). A previous comparison of the BBA size distributions in the FT and in the MBL near Ascension Island does not indicate a preferential removal of larger particles, as expected during precipitation events (Wu et al., 2020). Instead, non-precipitating aerosol-cloud interactions can alter both the chemical and the physical properties of the BBA. For example, oxidants such as OH, commonly found in high concentrations in cloud drops, can increase *f44* (McNeill et al., 2015) and alter aerosol composition. Additionally, OA and  $\Delta\text{SO}_4$  and their gaseous precursors can interact with  $\text{H}_2\text{O}_2$  within cloud drops to form additional aerosol mass (Yang et al., 2011; Bianco et al., 2020). Here, we use the aerosol size distributions, transport pathways, and values of *f44* to explore the extent to which aqueous-phase processes may have occurred and contributed to additional OA loss, also helping to explain the low OA:rBC values.

The BBA size distributions observed in the MBL were monomodal, weakly bimodal, or distinctly bimodal, whereas the BBA size distributions observed in the FT were uniformly monomodal and broad, with most of the particles having diameters greater than 100 nm but with a small fraction of particles having diameters less than 60 nm (Kacarab et al., 2019; Howell et al., 2021; Dobracki et al., 2023). The monomodal BBA size distributions in the MBL indicate that this BBA likely had not interacted with marine clouds. The weakly bimodal BBA size distributions with indistinct Hoppel minima in the MBL (Fig. 11) indicate that this BBA had undergone insufficient processing to fully alter the shape of the distinctly bimodal background aerosol size distributions (Atwood et al., 2020). Lastly, the distinctly bimodal BBA size distributions with pronounced Hoppel minima (Figs. 11, S4) suggest that this BBA had undergone interactions with clouds.

Regime 1 was dominated by bimodal aerosol size distributions, which, along with inferred transport pathways, high *f44* values, and high  $\Delta\text{SO}_4$ :rBC mass ratios, suggest BBA that had undergone appreciable cloud processing. The BBA transport was the slowest of the three regimes, and because upper-level FT winds were still weak, likely confined the BBA to the deeper MBL (Zhang and Zuidema 2021). These conditions allow more time for interaction with clouds. The highest fraction of activated aerosol particles occurred during this time (Zuidema et al., 2018), consistent with larger accumulation-mode aerosol (Dedrick et al. 2024). Relatively high *f44* values (Fig. 10), suggest some oxidation also occurred through aqueous-phase processing (Giorio et al., 2017; Che et al., 2022b). The  $\Delta\text{SO}_4$ :rBC mass ratios were also the highest during Regime 1, consistent with more conversion of  $\text{SO}_2$  gas (co-emitted with the BBA) into  $\text{SO}_4$  aerosol within cloud drops (Yang et al., 2011; Bianco

et al., 2020). These observations support the conclusion that the BBA in this regime likely experienced cloud oxidation and that additional aqueous-phase aerosol production.

1080           Thereafter, fewer and less distinct bimodal aerosol size distributions, faster transport pathways, lower  $f_{44}$  values, and lower  $\Delta\text{SO}_4:\text{rBC}$  mass ratios in Regime 2 indicate that the BBA has undergone less cloud processing than in Regime 1. The MBL is also slightly shallower (Zhang and Zuidema, 2021), allowing the BBA to remain in the FT for longer. These conditions discourage opportunities for aerosol-cloud microphysical interactions, despite a cloud fraction near Ascension Island (Zhang and Zuidema., 2021). Values of  $f_{44}$  and  $\Delta\text{SO}_4:\text{rBC}$  in Regime 2 were lower than those in Regime 1, also indicating less  
1085 additional oxidation through aqueous-phase processing.


          During Regime 3, mostly monomodal aerosol size distributions, faster transport pathways in the FT, lower  $f_{44}$  values, and lower  $\Delta\text{SO}_4:\text{rBC}$  mass ratios suggest that the BBA has not undergone extensive cloud processing. BBA is more likely to be transported above the MBL, in part by stronger FT easterly winds after August 20, and thereby less likely entrained into the MBL until near Ascension Island (Ryoo et al., 2022). Delayed entrainment into the MBL would result in the BBA replacing  
1090 the marine aerosol initially, with little mixing.

          Cloud processing, which is most active in Regime 1, can also produce small amounts of additional aerosol mass, increasing the mass fractions of OA and  $\Delta\text{SO}_4$  (Ervens et al., 2011; Gilardoni et al., 2016). This may have occurred during Regime 1, for which rBC particles also possess a slightly greater coating-to-core mass ratios than in the latter regimes, and for which OA:rBC and  $f_{44}$  correlate positively ( $r^2 = 0.61$ ), as do  $\Delta\text{SO}_4:\text{rBC}$  and  $f_{44}$  ( $r^2 = 0.58$ ; Fig. S7). However, the correlations  
1095 between OA:rBC and  $f_{44}$  and between  $\Delta\text{SO}_4:\text{rBC}$  and  $f_{44}$  could also have been influenced by differences in aerosol composition at the fire sources and differences in transport pathways. Nevertheless, our conclusions that aqueous-phase processing in non-precipitating clouds oxidizes the OA but does not substantially remove it are consistent with the results of Che et al. (2022b) and numerous laboratory studies (Ervens et al., 2011; Gilardoni et al., 2016). The smaller values of OA:rBC in the MBL compared to the FT, in August, are most likely explained by a combination of longer transport paths allowing for  
1100 more heterogeneous oxidation, including within clouds.

#### **4.5 SSA<sub>530</sub> values in the MBL are best explained by OA:rBC**

          The SSA<sub>530</sub> values in the Ascension Island MBL, between 0.75 and 0.83 (Section 3.2.4), are among the lowest in the world from BBA (Eck et al., 2013; Zuidema et al., 2018) and will strengthen the boundary layer semi-direct effect (Zhang and Zuidema, 2019) as a result. These values are even lower (by about 0.07) than those sampled in the FT between the African  
1105 plateau and Ascension Island during the September 2016 ORACLES campaign. This result is surprising, as the mean background SSA<sub>530</sub> value was  $0.98 \pm 0.01$ , and mixing with clean marine air should increase SSA<sub>530</sub> in the MBL, especially near the surface (Wu et al., 2020). However, the low SSA<sub>530</sub> values of  $\sim 0.79$  across P8–P9 compare relatively well with the polluted-MBL SSA<sub>550</sub> value of  $\sim 0.81$  measured during CLARIFY in the polluted MBL for the same time period. This minor difference in SSA can likely be explained by the variations in impactor cutoff sizes between LASIC and CLARIFY (Section

1110 2.2; Barrett et al., 2022), where larger particles near the surface may contribute to higher SSA values. The slight reduction in SSA<sub>530</sub> from P8 to P9 (Fig. 12) is consistent with the observed increase in rBC fraction from Period 1 to Period 3 during CLARIFY (Wu et al., 2020). Our results further solidify that SSA<sub>530</sub> is negatively correlated with FrBC (Fig. 17a) and positively correlated with OA:rBC (Fig. 17b) in both the MBL and in the FT between the African plateau and Ascension Island. Moreover, SSA<sub>530</sub> is not correlated with the geometric peak diameter of the accumulation mode (Fig. S8) in either the MBL or FT, consistent with Denjean et al. (2020b) using in-situ data over the Gulf of Guinea and Dobracki et al. (2023). FrBC values for both the MBL and the FT are almost identical between ORACLES and LASIC, ranging between 0.15 and 0.40, so variations in FrBC cannot account for the reduced MBL SSA<sub>530</sub> values either (Fig. 18a). Instead, one difference is that the OA:rBC mass ratios in the MBL (ranging from 2 to 5) are markedly lower than those in the FT (between 5-15). Of further note, a robust correlation exists between SSA<sub>530</sub> and OA:rBC that applies to both the MBL and FT: SSA<sub>530</sub> = 0.78 + 0.0077×OA:rBC (Fig. 18b). This relation is similar to that proposed by Dobracki et al. (2023) for the ORACLES 2016 campaign. The SSA<sub>530</sub> dependence on OA:rBC shows that accurate estimates of SSA<sub>530</sub> can be obtained from the BBA chemical properties alone.



1140 Figure. 17 a) SSA<sub>530</sub> versus FrBC b) SSA<sub>530</sub> versus OA:rBC. Markers are colored by month and error bars represent the standard deviation of the data set. The Black dashed line is the best fit line calculated by linear regression analysis. The best-fit line is weighted by  $\mu\text{g m}^{-3} \text{min}^{-1}$ .

1145

1150

1155

1160

1165




Figure 18. a)  $SSA_{530}$  versus FrBC for ORACLES level leg data (black) (Dobracki et al., 2022) and LASIC 2017 major plume event data (red). b)  $SSA_{530}$  versus OA:rBC for ORACLES level leg data (black) and LASIC 2017 major plume event data (red). Error bars represent the standard deviation of the data set and the black dashed line is the best-fit line calculated by linear regression analysis.


## 1170 5. Remaining Questions

### 5.1 Case Study: 6-11 September 2017

The last plume event to reach the MBL at Ascension Island between September 6-11 (P10), is unique in that the BBA originated from both efficient fires across central Africa and inefficient fires along the eastern coast of Africa (Section 4.1). This plume is notable for possessing large rBC core diameters (Fig. 6), and the largest OA:rBC mass ratio (Fig. 9),  $f_{44}$  values (Fig. 10; factor 2 only), and  $MAC_{530}$  of the 10 plumes-

Additional HYSPLIT back trajectories (Fig. 19) indicate that the BBA sampled before September 10 may have experienced more time in the FT than that from 1-2 days later, by first circulating anticyclonically  $\sim 10^\circ$  to the southeast of Ascension Island in the lower FT (1-2.5 km) before entering the MBL (Fig. 19a), similar to the case discussed in Diamond et al. (2022). The African Easterly Jet South would have advected the BBA aloft prior to entrainment into the MBL (Ryoo et al., 2022). In contrast, the BBA sampled on September 10 was transported directly westward from the continent (Fig. 19b), potentially from as far away as eastern Africa. This direct westward transport, meteorologically unusual for September (Ryoo et al., 2022), is more likely to occur at lower altitudes (Fig. 13d), and the different source region may explain why the aerosol properties of this plume differed from those from previous times.

1185



1200 Figure 19. 10-day HYSPLIT back trajectories at 1000 m (red), 1500 m (blue), and 2500 m (green) on a) September 8, 2017 at  
1205 1200 UTC and b) September 10, 2017 at 1200 UTC.

The high values of  $rBC:\Delta CO$  (Fig. S9a) and  $FrBC$  and the low values of  $OA:rBC$  (Fig. S9b) and  $\Delta SO_4:rBC$  (Fig. S9c) indicate that the BBA sampled before September 10 originated from efficient fires. Lower values of  $rBC:\Delta CO$  and  $FrBC$  paired with the higher values of  $OA:rBC$  and  $\Delta SO_4:rBC$  on September 10 indicate BBA during the last two days from inefficient combustion. Although these chemical properties are consistent with inefficient combustion, the optical properties are not. Intuitively  $MAC_{530}$  would increase with  $OA:rBC$  because of greater absorption by particles with larger OA coatings (so-called lensing) while  $SSA_{530}$  would increase with  $OA:rBC$  because of relatively more scattering than absorption, for either internally or externally mixed aerosol. Instead, we find the opposite relationship. Although the  $MAC_{530}$  did increase from 17.5 to 29.5  $m^2 g^{-1}$  between September 7 and September 10, it decreased to 25  $m^2 g^{-1}$  on September 11 (Fig. S9d). Meanwhile, the  $SSA_{530}$  decreased from 0.835 to 0.785 across the five days (Fig. S9e). The highest  $MAC_{530}$  and lowest  $SSA_{530}$  occurred on the day when  $\Delta CO$  was the highest (Fig. S9f). Overall  $MAC_{530}$  correlates positively and  $SSA_{530}$  correlates negatively with  $\Delta CO$  (Fig. S10), indicating greater absorption and absorptivity by particles more deeply within the smoke plume. We cannot fully explain the underlying processes.

1215

## 5.2 Aerosol absorption enhancement

The larger analysis of LASIC also brings the typical explanation for the high  $MAC_{530}$  values ( $> 15 m^2 g^{-1}$ ) observed in the Ascension Island MBL, of lensing-enhanced absorption from thick coatings, into question.  $MAC_{530}$  does not correlate with the geometric mean diameter of the accumulation mode determined from the BBA size distributions (Fig. S11), in the presence of more constant  $rBC$  core sizes, also concluded by Denjean et al. (2020b) and Dobracki et al. (2023) for FT aerosol.

1220

The high  $MAC_{530}$  values in the FT have been attributed to thick coatings (Taylor et al., 2020; Sedlacek et al., 2022), but rBC-containing particles in the MBL are more thinly coated (Sedlacek et al., 2022).

$MAC_{530}$  and  $SSA_{530}$  are positively correlated (Fig. S5), indicating that both scattering and absorption are enhanced when soot particles are coated with transparent OA and  $\Delta SO_4$  (Khalizov et al., 2009). In addition, both  $MAC_{530}$  and  $AAE_{470-660}$  in the MBL increase with increasing  $f_{44}$  (Figs. 20a-b), suggesting that chemical processing, either through heterogeneous oxidation or aqueous-phase processing, can act to enhance aerosol light absorption, also shown in Zhang et al. (2018). The increased light absorption resulting from chemical changes must have been small, however, as average  $AAE_{470-660}$  values are near 1 in both the MBL and the FT near Ascension Island, consistent with BrC contributing less than 10 % to the aerosol absorption (Taylor et al. 2020; Zhang et al. 2022). We conclude the processes contributing to aerosol absorption enhancement in the MBL are different than those in the FT and require additional study.




Figure 20. a)  $AAE_{470-660nm}$  versus average  $f_{44}$  for the 10 selected plume events. b) Average  $f_{44}$  versus  $MAC_{530}$  ( $m^2 g^{-1}$ ) for the 10 selected plume events. Markers are colored by month and error bars represent the standard deviation of the data set. The black dashed line is the best-fit line calculated by linear regression analysis. The best-fit line is weighted by  $\mu g m^{-3} min^{-1}$ .

## 6. Conclusions and summary

The current analysis focuses solely on 10 specific BBA plumes within the remote MBL of the southeast Atlantic, in contrast to BBA plumes in the FT. The synoptically-modulated BBA plumes in the MBL occur from June until early September, before further aerosol transport moves to the free troposphere. BBA in June and early July (Regime 1) contained more OA and  $\Delta SO_4$  relative to rBC, within primarily bimodal BBA size distributions. These BBA properties resulted from inefficient burning conditions (characterized by rBC: $\Delta CO$  mass ratios less than 0.01) combined with a slow westward transport of BBA in the MBL that allowed more time for cloud processing, arguably most notable in the enhanced  $\Delta SO_4$  mass. The BBA

1255 in late July and early August (Regime 2) had lower contributions of OA and  $\Delta\text{SO}_4$  relative to rBC, and fewer, broader bimodal  
size distributions, than those earlier in the season. These BBA properties resulted from efficient burning conditions  
(characterized by rBC: $\Delta\text{CO}$  mass ratios greater than 0.01) and from slightly faster transport pathways that were less confined  
to the MBL, reducing time for aqueous-phase processing. The BBA in late August and early September (Regime 3) had the  
highest contribution from OA relative to rBC. Mostly monomodal BBA size distributions indicate little mixing with marine  
1260 air. BBA properties reflect both efficient and inefficient burning conditions. These BBA properties resulted from both efficient  
and inefficient burning conditions and aerosol transport pathways being mostly in the FT due to the strong easterly winds  
during this time that did not allow for many aerosol-cloud interactions.

The BBA transport pathways to Ascension Island in the MBL and lower FT between June and September are slow,  
overall, and allow not only for heterogeneous oxidation processes but also for aqueous-phase processes. Continuous  
1265 heterogeneous oxidation processes for up to 10 days contribute to a net loss of OA mass, with the contribution from aqueous-  
phase processes (in June and July) which are difficult to distinguish from fuel type variations. Although  $f_{44}$  values and BBA  
size distributions suggest that aqueous-phase processing occurred, further effort is required to better quantify the relative  
contributions of heterogeneous oxidation and aqueous-phase processing on the evolution of BBA.

SSA values measured in the Ascension Island MBL, less than 0.80, are among the lowest in the world from BBA.  
1270 The low OA:rBC ratios in the MBL, a consequence of oxidation processes, likely explains these low  $\text{SSA}_{530}$  values and is  
arguably the most striking result of this study. Current global and regional aerosol models typically do not include  
heterogeneous and aqueous-phase oxidation processes as major aerosol loss mechanisms, but the resulting changes in chemical  
composition can have substantial effects on aerosol optical properties, demonstrating the need for further effort placed on  
constraining the loss of OA in the MBL. A remaining puzzle is a strong positive correlation between  $\text{MAC}_{530}$  and  $\text{SSA}_{530}$ ,  
1275 which is not easily explained by lensing effects.

## Appendix A

### Experimental design and Instrumentation

#### 1.1 Aerosol Sampling Inlet

The inlet used in each Aerosol Observing System (AOS) was based on the design from the Environmental  
1280 Measurements Laboratory (Leifer et al., 1994; Uin et al., 2019). The inlet was positioned 10 m above the ground and drew a  
flow rate of  $1000 \text{ L min}^{-1}$ , with  $150 \text{ L min}^{-1}$  distributed evenly among five sample lines (Uin et al., 2019). The flow through  
the sample lines was maintained by bypass flow with support from a vacuum source and from individual instrument flows on  
that line (Uin et al., 2019). The bypass flows were measured and monitored by rotameters. Trace gas measurements were  
sampled through a separate  $\frac{1}{2}$ -inch tube that ran parallel to the main inlet (Uin et al., 2019, Figure 4). A comprehensive  
1285 description of the AOS set up and operation within the ARM program is given by Uin et al. (2019).

## 1.2 Aerosol Chemical Speciation Monitor (ACSM)

The aerosol chemical speciation monitor (ACSM, Aerodyne Inc.) measures non-refractory particle mass concentrations of organic, nitrate, sulfate, ammonium, and chloride species in real time for particles with aerodynamic diameters between 75 and 650 nm at 30 min time resolution (ACSM handbook, Watson, 2014). The ACSM has a detection limit of  $0.3 \mu\text{g m}^{-3}$  (30-minute signal average) for organic aerosol,  $0.2 \mu\text{g m}^{-3}$  for nitrate,  $0.4 \mu\text{g m}^{-3}$  for sulfate,  $0.5 \mu\text{g m}^{-3}$  for ammonium, and  $0.2 \mu\text{g m}^{-3}$  for chloride. The aerosol was drawn in through a sampling pump at  $3 \text{ L min}^{-1}$ , focused through an aerodynamic lens, and impacted on a collector plate kept at  $600^\circ\text{C}$  to vaporize the particles, and then ionized by electron ionization at 70 eV. The ions were analyzed by a quadrupole mass spectrometer which provides unit mass resolution. The ACSM has an uncertainty of 30-40 % in mass concentration.

1295 The ACSM was calibrated twice in the field using ammonium nitrate. A misaligned laser filament was replaced in January 2017, and the instrument was re-calibrated. A time- and composition-dependent collection efficiency (CDCE) corrects for the incomplete vaporization of mixed-phase particles (Middlebrook et al., 2012). The CDCE was calculated at each sampling interval using methods from Shilling and Levin (2021). The average CDCE between June and September was 0.54.

## 1.3 Single Particle Soot Photometer (SP2)

1300 The single-particle soot photometer (SP2, Droplet Measurement Technology) measured time-dependent scattering and incandescence signals from refractory black carbon-containing particles with aerodynamic diameter between 80 and 500 nm. The SP2 used an 8-channel neodymium-doped yttrium-aluminum-garnet (Nd:YAG) laser (1064 nm) that induced incandescence of the particles to determine rBC mass and number concentrations (Stephens et al., 2003, Sedlacek 2017, ARM Handbook). The SP2 was calibrated twice in the field using fullerene soot with effective density from Gysel et al. (2011). The SP2 has a detection limit of  $0.1 \mu\text{g m}^{-3}$  with 75 % certainty (May et al., 2014) and a mass uncertainty of ~20 %.

## 1.4 Scanning Mobility Particle Sizer (SMPS)

1310 The scanning mobility particle sizer (SMPS, TSI 3936) determines size distributions of number concentration of particles with mobility diameters between 10 and 1000 nm. The SMPS uses a bipolar aerosol charger to establish an equilibrium charge distribution on the aerosol. A long-column differential mobility analyzer (DMA) classifies charged particles according to their electrical mobility. The particle concentration was measured with a condensation particle counter (CPC). The SMPS and CPC were calibrated with certified polystyrene latex (PSLs), silver chloride, and sodium chloride particles (Hermann et al., 2007). The SMPS has an uncertainty of 10 %.

## 1.5 Particle Soot Absorption Photometer

The particle soot absorption photometer (PSAP, Radiance Research) measured optical transmittance through a cellulose filter on which particles were deposited to determine the light absorption at 470, 530, and 660 nm. PSAP data were corrected for loading and scattering effects following the ARM protocol ARM handbook (Virkkula 2010; Ogren 2010; Bond et al. 1999).

## 1.6 Nephelometer

The nephelometer (TSI, Inc) measured the total light scattering coefficient and backscatter fraction by aerosol particles at 450, 550, and 700 nm (Anderson and Ogren 1998). The aerosol optical properties and their derived products incorporate correction algorithms from Bond/Ogren (Bond 1999, extended by Ogren 2010) and Virkkula 2010, as done previously in Zuidema et al. (2018) and Pistone et al. (2019). Further information can be found in Flynn et al. (2020). The nephelometer deployed at ASI was installed in series with a drying/humidification system to measure the change in scattering due to water uptake or loss by the particles and has a systematic uncertainty of 10 % (Uin 2016b).

## 1.7 CAPS-SSA

The cavity attenuated phase shift single-scattering albedo monitor (CAPS-SSA, Aerodyne Inc) was deployed to Ascension Island between August 4, 2017, and September 22, 2017. The CAPS-SSA collected 1 s time response measurements of aerosol extinction and scattering. The CAPS-SSA measurement of ambient SSA at 530 nm wavelength (0.78) compared well with  $SSA_{530}$  (0.79) calculated from the PSAP (absorption) and nephelometers (scattering) (Fig. A1) (Onasch et al., 2015).




Figure A1.  $SSA_{530}$  calculated from the PSAP and nephelometer absorption and scattering coefficients (black) and CAPS-SSA (green). rBC mass concentration (grey dashed line) indicates this comparison was done in the plume.

## 1345 **1.8 CO/N<sub>2</sub>O/H<sub>2</sub>O Analyzer**

Carbon monoxide, nitrous oxide, and water vapor mixing ratios were measured with a CO/N<sub>2</sub>O/H<sub>2</sub>O Analyzer (Los Gatos Research) at 1 s resolution. The measurement precision is 0.05 ppb under quiet ambient conditions (Springston, 2015).  $\Delta$ CO was calculated for each month by removing (subtracting) the 5<sup>th</sup> percentile of the measured CO from the total measured CO. Background values of CO were 50.2 ppb for June, 54.4 ppb for July, 58.8 ppb for August, and 60.2 ppb for September.

1350 Methods for calculating  $\Delta$ CO were adopted from Che et al. (2022).

## **2. Data analysis**

### **2.1 Positive Matrix Factorization (PMF)**

A PMF apportions the unit mass resolution data using a bilinear model through a multilinear engine (ME-2). A graphical user interface, Source Finder (SoFi), developed at the Paul Scherrer Institute, was run using IGOR Pro (Wavemetrics v8) to  
1355 facilitate the testing of different rotational techniques within the ME-2 (Canonaco et al., 2013; Crippa et al., 2014; Zhu et al., 2018; Canonaco et al., 2021). The PMF results in this paper were obtained by running 400-1000 clusters of the unit-mass resolution data from the 10 major plume events with SoFi through IGOR Pro (v8). The clusters were then analyzed using a k-means statistical function to find clusters of comparable spatial extent. The average mass spectra result of the clusters presented two factors of source and process-related apportionments.

### **2.2 SSA<sub>530</sub> and MAC<sub>530</sub> Calculations**


The single-scattering albedo (SSA) is defined as the ratio of the (light) scattering coefficient to the sum of the absorption and scattering coefficients. The absorption Ångström exponent (AAE) describes the spectral dependence of light absorption by aerosols (Helin et al., 2021) and is calculated from the linear fit of  $\log(\sigma_a)$  to  $\log(\lambda)$ . The MAC<sub>530</sub> was calculated by dividing the absorption coefficient at 530 nm by the rBC mass concentration determined by the SP2.

1365 The Cavity Attenuated Phase Shift (CAPS) extinction monitor (Aerodyne Inc.) was deployed in August, 2017 to measure aerosol extinction (Sedlacek 2017). The calculated SSA<sub>530</sub> from the ARM VAP file agrees well with the SSA<sub>530</sub> from CAPS during smokey periods (Figure A1), indicating that the assumed corrections to the PSAP and nephelometers are appropriate in the MBL where there could be a large presence of sea-salt particles.

### **2.3 ACSM volume comparison**

1370 The total volume concentration calculated using the size distribution from the SMPS was compared to the sum of the volume concentrations from the ACSM and SP2, calculated for each substance as the mass concentration divided by the density the bulk species based on the mass fraction of each species (OA, SO<sub>4</sub>, NO<sub>3</sub>, NH<sub>4</sub>, rBC). The values of the densities of most of the primary substances measured by the ACSM are well established (nitrate:  $\rho=1.72$  g cm<sup>-3</sup>, sulfate and ammonium:  $\rho=1.77$  g cm<sup>-3</sup>, and chloride:  $\rho=1.35$  g cm<sup>-3</sup>; Nakao et al., 2013); however, values for OA between 0.8 and 1.8 g cm<sup>-3</sup> have

1375 been reported. Some of this range is due to differences in the chemical composition of the individual organic species and some  
from the age of the OA, with values of aged OA being typically less than those of freshly emitted OA (Rudich et al., 2006;  
Denjean et al., 2020b). Here a density of  $0.9 \text{ g cm}^{-3}$  for OA was assumed, as aging metrics suggest that the aerosol at Ascension  
Island is highly aged (greater than seven days old; Section 3.2.3). The volume concentration calculated as the sum of those  
determined by the SMPS agrees with that determined from the ACSM and SP2 to within 30 % (Fig. A2), which is roughly the  
1380 sampling error expected for the latter two instruments.



1385

1390

1395

1400 Figure A2. ACSM and SMPS volume comparison for the 10 selected plume events. Solid lines represent the best-fit linear  
regression analysis for each plume event. Lines and markers are colored by plume number. Black dashed line represents the  
1:1 relationship.

1405

1410

## Appendix A References

- Anderson, T. L. and Ogren, J. A.: Determining Aerosol Radiative Properties Using the TSI 3563 Integrating Nephelometer, *Aer. Sci. Tech.*, 29, 57–69, <https://doi.org/10.1080/02786829808965551>, 1998.
- 1415 Bond, T. C., Anderson, T. L., and Campbell, D.: Calibration and Intercomparison of Filter-Based Measurements of Visible Light Absorption by Aerosols, *Aerosol Science and Technology*, 30, 582–600, <https://doi.org/10.1080/027868299304435>, 1999.
- 1420 Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: SoFi, an IGOR-based interface for the efficient use of the generalized multiline engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos Meas Tech*, 6, 3649–3661, <https://doi.org/10.5194/amt-6-3649-2013>, 2013.
- 1425 Canonaco, F., Tobler, A., Chen, G., Sosedova, Y., Slowik, J. G., Bozzetti, C., Daellenbach, K. R., El Haddad, I., Crippa, M., Huang, R.-J., Furger, M., Baltensperger, U., and Prévôt, A. S. H.: A new method for long-term source apportionment with time-dependent factor profiles and uncertainty assessment using SoFi Pro: application to 1 year of organic aerosol data, *Atmos Meas Tech*, 14, 923–943, <https://doi.org/10.5194/amt-14-923-2021>, 2021.
- 1430 Che, H., Segal-Rozenhaimer, M., Zhang, L., Dang, C., Zuidema, P., Sedlacek III, A. J., Zhang, X., and Flynn, C.: Seasonal variations in fire conditions are important drivers in the trend of aerosol optical properties over the south-eastern Atlantic, *Atmos Chem Phys*, 22, 8767–8785, <https://doi.org/10.5194/acp-22-8767-2022>, 2022.
- 1435 Crippa, M., Canonaco, F., Lanz, V. A., Äijälä, M., Allan, J. D., Carbone, S., Capes, G., Ceburnis, D., Dall’Osto, M., Day, D. A., DeCarlo, P. F., Ehn, M., Eriksson, A., Freney, E., Hildebrandt Ruiz, L., Hillamo, R., Jimenez, J. L., Junninen, H., Kiendler-Scharr, A., Kortelainen, A.-M., Kulmala, M., Laaksonen, A., Mensah, A. A., Mohr, C., Nemitz, E., O’Dowd, C., Ovadnevaite, J., Pandis, S. N., Petäjä, T., Poulain, L., Saarikoski, S., Sellegri, K., Swietlicki, E., Tiitta, P., Worsnop, D. R., Baltensperger, U., and Prévôt, A. S. H.: Organic aerosol components derived from 25 AMS data sets across Europe using a consistent ME-2 based source apportionment approach, *Atmos Chem Phys*, 14, 6159–6176, <https://doi.org/10.5194/acp-14-6159-2014>, 2014.
- 1440 Flynn, C., Chand, D., Ermold, B., and Koontz, A.: The ARM Aerosol Optical Properties (AOP) The ARM Aerosol Optical Properties (AOP) Value-Added Product, November 2020.
- 1445 Gysel, M., Laborde, M., Olfert, J. S., Subramanian, R., and Gröhn, A.: Effective density of Aquadag and fullerene soot black carbon reference materials used for SP2 calibration, *Atmos. Meas. Tech. Disc.*, 4, 4937–4955, <https://doi.org/10.5194/amt-d-4-4937-2011>, 2011.
- Hermann, M., Wehner, B., Bischof, O., Han, H.-S., Krinke, T., Liu, W., Zerrath, A., and Wiedensohler, A.: Particle counting efficiencies of new TSI condensation particle counters, *J. Aerosol Sci.*, 38, 674–682, <https://doi.org/https://doi.org/10.1016/j.jaerosci.2007.05.001>, 2007.
- 1450 Keabian, P. L., Robinson, W. A., and Freedman, A.: Optical extinction monitor using cw cavity enhanced detection, *Rev. Sci. Instrum.*, 78, <https://doi.org/https://doi.org/10.1063/1.2744223>, 2007.
- 1455 Onasch, T. B., Massoli, P., Keabian, P. L., Hills, F. B., Bacon, F. W., and Freedman, A.: Single Scattering Albedo Monitor for Airborne Particulates, *Aerosol Science and Technology*, 49, 267–279, <https://doi.org/10.1080/02786826.2015.1022248>, 2015.
- Ogren, J. A.: Comment on ‘‘Calibration and Intercomparison of Filter-Based Measurements of Visible Light Absorption by Aerosols’’, *Aerosol Science and Technology*, 44, 589–591, <https://doi.org/10.1080/02786826.2010.482111>, 2010.

- 1460 Leifer, R., R. H. Knuth, and H. N. Lee, 1994: Surface aerosol measurements at Lamont, Oklahoma. *Proc. Third Atmospheric Radiation Measurement Science Team Meeting*, Washington, DC, U.S. Department of Energy, 349–351.
- Pistone, K., Redemann, J., Doherty, S., Zuidema, P., Burton, S., Cairns, B., Cochrane, S., Ferrare, R., Flynn, C., Freitag, S., Howell, S., Kacenenbogen, M., LeBlanc, S., Liu, X., Schmidt, K. S., Sedlacek III, A. J., Segal-Rosenhaimer, M., Shinozuka, Y., Starnes, S., van Dierenhoven, B., Van Harten, G., and Xu, F.: Intercomparison of biomass burning aerosol optical properties from in-situ and remote-sensing instruments in ORACLES-2016, *Atmos. Chem. Phys.*, 19, 9181–9208, <https://doi.org/10.5194/acp-19-9181-2019>, 2019.
- 1465 Shilling, J. E. and Levin, M. S.: Aerosol Chemical Speciation Monitor (ACSM) Composition-Dependent Collection Efficiency (CDCE) Value-Added Product Report, August 2021.
- 1470 Sedlacek, A. J., 2016: Cavity attenuated phase shift (CAPS) monitor instrument handbook. ARM Tech. Rep. DOE/SC-ARM-TR-155, 23 pp., <https://doi.org/10.2172/1251390>.
- 1475 Sedlacek, A. J., 2017: Single-Particle Soot Photometer (SP2) instrument handbook. ARM Tech. Rep. DOE/SC-ARM-TR-169, 24 pp., <https://doi.org/10.2172/1344179>.
- Springston, S. R., 2015: Carbon monoxide analyzer (CO-ANALYZER) instrument handbook. ARM Tech. Rep. DOE/SC-ARM-TR-159, 30 pp., <https://doi.org/10.2172/1495422>.
- 1480 Stephens, M., Turner, N., and Sandberg, J.: Particle identification by laser-induced incandescence in a solid-state laser cavity, *Appl. Opt.*, 42, 3726–3736, <https://doi.org/https://doi.org/10.1364/AO.42.003726>, 2003.
- Uin, J., 2016: Integrating nephelometer instrument handbook. ARM Tech. Rep. DOE/SC-ARM-TR-165, 16 pp., <https://doi.org/10.2172/1246075>.
- 1485 Uin, J., Aiken, A. C., Dubey, M. K., Kuang, C., Pekour, M., Salwen, C., Sedlacek, A. J., Senum, G., Smith, S., Wang, J., Watson, T. B., and Springston, S. R.: Atmospheric Radiation Measurement (ARM) Aerosol Observing Systems (AOS) for Surface-Based In Situ Atmospheric Aerosol and Trace Gas Measurements, *J Atmos Ocean Technol*, 36, 2429–2447, <https://doi.org/10.1175/JTECH-D-19-0077.1>, 2019.
- Virkkula, A.: Correction of the Calibration of the 3-wavelength Particle Soot Absorption Photometer (3 PSAP), *Aerosol Sci. Tech.*, 44, 706–712, 2010.
- 1495 Watson, T. B., 2016b: Proton transfer time-of-flight mass spectrometer. *U ARM Tech. Rep.* DOE/SC-ARM-TR-160, 32 pp., <https://doi.org/10.2172/1251396>.
- 1500 Zhu, Q., Huang, X.-F., Cao, L.-M., Wei, L.-T., Zhang, B., He, L.-Y., Elser, M., Canonaco, F., Slowik, J. G., Bozzetti, C., El-Haddad, I., and Prévôt, A. S. H.: Improved source apportionment of organic aerosols in complex urban air pollution using the multilinear engine (ME-2), *Atmos Meas Tech*, 11, 1049–1060, <https://doi.org/10.5194/amt-11-1049-2018>, 2018.
- Zuidema, P., Sedlacek III, A. J., Flynn, C., Springston, S., Delgadillo, R., Zhang, J., Aiken, A. C., Koontz, A., and Muradyan, P.: The Ascension Island Boundary Layer in the Remote Southeast Atlantic is Often Smoky, *Geophys. Res. Lett.*, 45, 4456–4465, <https://doi.org/10.1002/2017gl076926>, 2018.
- 1505

## **Data availability**

1510 The following data are publicly available through the Atmospheric Radiation Measurement (ARM) Data Archive (for definitions see Table 1).

- ACSM: <https://doi.org/10.5439/1762267> (Watson 2017; Shilling and Levin, 2021).

- SP2: <http://www.arm.gov/campaigns/amf2016lasic/> (Sedlacek 2017)

- SMPS: <https://doi.org/10.5439/1476898> (Kuang 2016)

1515 - CPC: <http://doi.org/10.5439/1352536> (Kuang 2016)

- Nephelometer: <https://doi.org/10.5439/1369240> (Uin 2016)

- PSAP: <https://doi.org/10.5439/1369240> (Springston 2018)

- Surface winds: <https://doi.org/10.5439/1595321> (Holdridge 2020)

- CO analyzer: <http://www.arm.gov/campaigns/amf2016lasic/> (Springston 2015)

1520 - Optical properties VAP: <https://doi.org/10.5439/1369240> (Flynn 2020)

- Fire data: <https://firms.modaps.eosdis.nasa.gov/download/>

- NCEP winds: <https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.html>

- CAMS CO: <https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-reanalysis-eac4?tab=overview>

## **Supplement**

1525

### **Author contributions**

PZ designed the research. AD and PZ conceived this study. AJS and MZ prepared datasets of ACSM and SP2. AD, MZ, AJS analyzed datasets. AD, PZ, and EL wrote the paper with edits from TT, MZ, and AJS.

### **1530 Competing interests**

The contact author has declared that none of the authors has any competing interests.

### **Disclaimer**

### **Special issue statement**

1535

### **Acknowledgments**

We are grateful to the LASIC instrument mentors, AOS technicians, and logistics staff, who made this analysis possible through their efforts in deploying and maintaining the instruments at this remote location and to those who processed and calibrated the campaign data. We thank the editor, Dr. Rebecca Garland and reviewers for their insightful feedback and  
1540 suggestions, which have enhanced the clarity and impact of our manuscript.

**Financial Support** A. D., T. T., and P. Z. were supported by DOE Atmospheric System Research (ASR) award DE-SC0021250 and by NASA award NASA-80NSSC21K1344: “Constraining Fire-Cloud Interactions in the South Atlantic.” E. R. L., A. J. S., and M. Z. were supported by the US Department of Energy’s Office of Biological & Environmental Sciences (OBER) Program under contract DE-SC0012704 (the Process-level Advancements of Climate through Cloud and Aerosol Lifecycle Studies, PASCALS)."

## Review statement

## 1550 References

- Adebisi, A. A. and Zuidema, P.: The role of the southern African easterly jet in modifying the southeast Atlantic aerosol and cloud environments, *Q. J. Roy. Meteor. Soc.*, 142, 1574–1589, <https://doi.org/10.1002/qj.2765>, 2016.
- Aiken, A. C., DeCarlo, P. F., Kröll, J. H., Worsnop, D., Huffman, J. A., Docherty, K. S., Ulbrich, I., and et al., C. M.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ Sci Technol.*, 42, 4478–4485, 2008.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- 1565 Atwood, S. A., Reid, J. S., Kreidenweis, S. M., Blake, D. R., Jonsson, H. H., Lagrosas, N. D., Xian, P., Reid, E. A., Sessions, W. R., and Simpas, J. B.: Size-resolved aerosol and cloud condensation nuclei (CCN) properties in the remote marine South China Sea – Part 1: Observations and source classification, *Atmos Chem Phys*, 17, 1105–1123, <https://doi.org/10.5194/acp-17-1105-2017>, 2017.
- 1570 Barrett, P. A., Abel, S. J., Coe, H., Crawford, I., Dobracki, A., Haywood, J. M., Howell, S., Jones, A., Langridge, J., McFarquhar, G., Nott, G., Price, H., Redemann, J., Shinzuka, Y., Szpek, K., Taylor, J., Wood, R., Wu, H., Zuidema, P., Bauguitte, S., Bennett, R., Bower, K., Chen, H., Cochrane, S. P., Cotterell, M., Davies, N., Delene, D., Flynn, C., Freedman, A., Freitag, S., Gupta, S., Noone, D., Onasch, T. B., Podolske, J., Poellot, M. R., Schmidt, S. K., Springston, S., III, A. J. S., Trembath, J., Vance, A., Zawadowicz, M., and Zhang, J.: Intercomparison of airborne and surface-based measurements during the CLARIFY, ORACLES and LASIC field experiments, *Atmos. Meas. Tech.*, 15, 6329–6371, <https://doi.org/10.5194/amt-15-6329-2022>, 2022.
- 1580 Bianco, A., Passananti, M., Brigante, M., and Mailhot, G.: Photochemistry of the Cloud Aqueous Phase: A Review, *Molecules*, 25, <https://doi.org/10.3390/molecules25020423>, 2020.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aer. Sci. Tech.*, 40, 27–67, <https://doi.org/10.1080/02786820500421521>, 2006.
- 1585 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., and et al., Y. K.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.

- 1590 Bowman, D. M., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., and et al., R. S. D.: Fire in the Earth System, *Science* (1979), 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- 1595 Brown, H., Liu, X., Pokhrel, R., Murphy, S., Lu, Z., Saleh, R., Mielonen, T., Kokkola, H., Bergman, T., Myhre, G., Skeie, R. B., Watson-Paris, D., Stier, P., Johnson, B., Bellouin, N., Schulz, M., Vakkari, V., Beukes, J. P., vanZyl, P. G., Liu, S., and Chand, D.: Biomass burning aerosols in most climate models are too absorbing, *Nat. Comm.*, 12, 277, <https://doi.org/10.1038/s41467-020-20482-9>, 2021.
- 1600 Canagaratna, M. R., Jimenez, J. L., and et al., J. H. K.: Elemental ratio measurements of organic compounds using aerosol mass spectrometry: characterization, improved calibration, and implications, *Atmos. Chem. Phys.*, 15, 253–272, <https://doi.org/10.5194/acp-15-253-2015>, 2015.
- 1605 Canonaco, F., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: Seasonal differences in oxygenated organic aerosol composition: implications for emissions sources and factor analysis, *Atmos Chem Phys*, 15, 6993–7002, <https://doi.org/10.5194/acp-15-6993-2015>, 2015.
- 1610 Cantrell Will Hart, I.: Relationship between the aerosol number distribution and the cloud condensation nuclei supersaturation spectrum, <http://hdl.handle.net/11122/9516>, 1999.
- 1615 Cape, J. N., Coyle, M., and Dumitrean, P.: The atmospheric lifetime of black carbon, *Atmos. Env.*, 59, 256–263, <https://doi.org/10.1016/j.atmosenv.2012.05.030>, 2012.
- 1620 Cappa, C. D. and Jimenez, J. L.: Quantitative estimates of the volatility of ambient organic aerosol, *Atmos Chem Phys*, 10, 5409–5424, <https://doi.org/10.5194/acp-10-5409-2010>, 2010.
- 1625 Che, H., Segal-Rozenhaimer, M., Zhang, L., Dang, C., Zuidema, P., Sedlacek III, A. J., Zhang, X., and Flynn, C.: Seasonal variations in fire conditions are important drivers in the trend of aerosol optical properties over the south-eastern Atlantic, *Atmos Chem Phys*, 22, 8767–8785, <https://doi.org/10.5194/acp-22-8767-2022>, 2022a.
- 1630 Che, H., Segal-Rozenhaimer, M., Zhang, L., Dang, C., Zuidema, P., Dobracki, A., Sedlacek, A. J., Coe, H., Wu, H., Taylor, J., Zhang, X., Redemann, J., and Haywood, J.: Cloud processing and weeklong ageing affect biomass burning aerosol properties over the south-eastern Atlantic, *Commun Earth Environ*, 3, 182, <https://doi.org/10.1038/s43247-022-00517-3>, 2022b.
- 1635 Chýlek, P., Lee, J. E., Romonosky, D. E., Gallo, F., Lou, S., and et al., M. S.: Mie Scattering Captures Observed Optical Properties of Ambient Biomass Burning Plumes Assuming Uniform Black, Brown, and Organic Carbon Mixtures, *J. Geophys. Res.*, 124, 11406–11427, <https://doi.org/10.1029/2019jd031224>, 2019.
- 1640 Clarke, A. D., Freitag, S., Simpson, R. M. C., Hudson, J. G., Howell, S. G., Brekhovskikh, V. L., Campos, T., Kapustin, V. N., and Zhou, J.: Free troposphere as a major source of CCN for the equatorial pacific boundary layer: long-range transport and teleconnections, *Atmos Chem Phys*, 13, 7511–7529, <https://doi.org/10.5194/acp-13-7511-2013>, 2013.
- 1645 Collier, S., Zhou, S., Onasch, T. B., Jaffe, D. A., Kleinman, L., A. J. Sedlacek, I. I. I., Briggs, N. L., Hee, J., Fortner, E., and et al., J. E. S.: Regional Influence of Aerosol Emissions from Wildfires Driven by Combustion Efficiency: Insights from the BBOP Campaign, *Environ Sci Technol.*, 50, 8613–8622, <https://doi.org/10.1021/acs.est.6b01617>, 2016.
- 1650 Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., and et al., D.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049–12064, <https://doi.org/10.5194/acp-11-12049-2011>, 2011.

- 1640 Dang, C., Segal-Rozenhaimer, M., Che, H., Zhang, L., Formenti, P., Taylor, J., Dobracki, A., Purdue, S., Wong, P.-S., Nenes, A., Sedlacek, A., Coe, H., Redemann, J., Zuidema, P., and Haywood, J.: Biomass burning and marine aerosol processing over the southeast Atlantic Ocean: A TEM single particle analysis, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-2021-724>, 2022.
- 1645 Das, S., Harshvardhan, H., Bian, H., Chin, M., Curci, G., Protonotariou, A. P., Mielonen, T., Zhang, K., Wang, H., and Liu, X.: Biomass burning aerosol transport and vertical distribution over the South African-Atlantic region, *J. Geophys. Res.*, 122, 6391–6415, <https://doi.org/10.1002/2016JD026421>, 2017.
- 1650 Dedrick, J. L., Russell, L. M., Sedlacek, A. J., III, Kuang, C., Zawadowicz, M. A., & Lubin, D., Aerosol-correlated cloud activation for clean conditions in the tropical Atlantic boundary layer during LASIC. *J. Geophys. Res.*, 51, e2023GL105798. <https://doi.org/10.1029/2023GL105798>, 2024
- 1655 Denjean, C., Bourriane, T., Burnet, F., Mallet, M., Maury, N., Colomb, A., Dominutti, P., Brito, J., Dupuy, R., Sellegri, K., Schwarzenboeck, A., Flamant, C., and Knippertz, P.: Overview of aerosol optical properties over southern West Africa from DACCWA aircraft measurements, *Atmos. Chem. Phys.*, 20, 4735–4756, <https://doi.org/10.5194/acp-20-4735-2020>, 2020a.
- 1655 Denjean, C., Brito, J., Libois, Q., Mallet, M., Bourriane, T., and et al., F. B.: Unexpected Biomass Burning Aerosol Absorption Enhancement Explained by Black Carbon Mixing State, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2020gl089055>, 2020b.
- 1660 Diamond, M. S., Saide, P. E., Zuidema, P., Ackerman, A. S., Doherty, S. J., Fridlind, A. M., Gordon, H., Howes, C., Kazil, J., Yamaguchi, T., Zhang, J., Feingold, G., and Wood, R.: Cloud adjustments from large-scale smoke–circulation interactions strongly modulate the southeastern Atlantic stratocumulus-to-cumulus transition, *Atmos Chem Phys*, 22, 12113–12151, <https://doi.org/10.5194/acp-22-12113-2022>, 2022.
- 1665 Donahue, N. M., Henry, K. M., Mentel, T. F., Kiendler-Scharr, A., Spindler, C., Bohn, B., Brauers, T., Dorn, H. P., Fuchs, H., Tillmann, R., Wahner, A., Saathoff, H., Naumann, K.-H., Möhler, O., Leisner, T., Müller, L., Reinnig, M.-C., Hoffmann, T., Salo, K., Hallquist, M., Frosch, M., Bilde, M., Tritscher, T., Barmet, P., Praplan, A. P., DeCarlo, P. F., Dommen, J., Prévôt, A. S. H., and Baltensperger, U.: Aging of biogenic secondary organic aerosol via gas-phase OH radical reactions, *Proceedings of the National Academy of Sciences*, 109, 13503–13508, <https://doi.org/10.1073/pnas.1115186109>, 2012.
- 1670 Dobracki, A., Zuidema, P., Howell, S. G., Saide, P., Freitag, S., Aiken, A. C., Burton, S. P., Sedlacek III, A. J., Redemann, J., and Wood, R.: An attribution of the low single-scattering albedo of biomass burning aerosol over the southeastern Atlantic, *Atmos Chem Phys*, 23, 4775–4799, <https://doi.org/10.5194/acp-23-4775-2023>, 2023.
- 1675 Doherty, S. J., Saide, P., Zuidema, P., Shinozuka, Y., Ferrada, G., and et al., H. G.: Modeled and observed properties related to the direct aerosol radiative effect of biomass burning aerosol over the Southeast Atlantic, *Atmos. Chem. Phys.*, 22, 1, 46, <https://doi.org/10.5194/acp-22-1-2022>, 2022.
- 1680 Eck, T. F.: Correction to “Variability of biomass burning aerosol optical characteristics in southern Africa during the SAFARI 2000 dry season campaign and a comparison of single scattering albedo estimates from radiometric measurements” by T. F. Eck et al., *Journal of Geophysical Research: Atmospheres*, 108, n/a-n/a, <https://doi.org/10.1029/2003JD001606>, 2003.
- 1685 Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos Chem Phys*, 11, 11069–11102, <https://doi.org/10.5194/acp-11-11069-2011>, 2011.
- 1685 Flynn, C., Chand, D., Ermold, B., and Koontz, A.: The ARM Aerosol Optical Properties (AOP) The ARM Aerosol Optical Properties (AOP) Value-Added Product, November 2020.

- George, I. J. and Abbatt, J. P. D.: Chemical evolution of secondary organic aerosol from OH-initiated heterogeneous oxidation, *Atmos. Chem. Phys.*, 10, 5551–5563, <https://doi.org/10.5194/acp-10-5551-2010>, 2010.
- 1690
- Gilardoni, S., Massoli, P., Paglione, M., Giulianelli, L., Carbone, C., Rinaldi, M., Decesari, S., Sandrini, S., Costabile, F., Gobbi, G. P., Pietrogrande, M. C., Visentin, M., Scotto, F., Fuzzi, S., and Facchini, M. C.: Direct observation of aqueous secondary organic aerosol from biomass-burning emissions, *Proceedings of the National Academy of Sciences*, 113, 10013–10018, <https://doi.org/10.1073/pnas.1602212113>, 2016.
- 1695
- Giorio, C., Monod, A., Brégonzio-Rozier, L., DeWitt, H. L., Cazaunau, M., Temime-Roussel, B., Gratien, A., Michoud, V., Pangui, E., Ravier, S., Zielinski, A. T., Tapparo, A., Vermeylen, R., Claeys, M., Voisin, D., Kalberer, M., and Doussin, J.-F.: Cloud Processing of Secondary Organic Aerosol from Isoprene and Methacrolein Photooxidation, *J Phys Chem A*, 121, 7641–7654, <https://doi.org/10.1021/acs.jpca.7b05933>, 2017.
- 1700
- Haywood, J. M., Abel, S. J., Barrett, P. A., Bellouin, N., Blyth, A., Bower, K. N., Brooks, M., Carslaw, K., Che, H., Coe, H., Cotterell, M. I., Crawford, I., Cui, Z., Davies, N., Dingley, B., Field, P., Formenti, P., Gordon, H., de Graaf, M., Herbert, R., Johnson, B., Jones, A. C., Langridge, J. M., Malavelle, F., Partridge, D. G., Peers, F., Redemann, J., Stier, P., Szpek, K., Taylor, J. W., Watson-Parris, D., Wood, R., Wu, H., and Zuidema, P.: Overview: The CLOUD-Aerosol-Radiation Interaction and Forcing: Year-2017 (CLARIFY-2017) measurement campaign, *Atmos. Chem. Phys.*, 21, 1049–1084, <https://doi.org/10.5194/acp-21-1049-2021>, 2021.
- 1705
- Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J., Madronich, S., and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmos. Chem. Phys.*, 16, 7917–7941, <https://doi.org/10.5194/acp-16-7917-2016>, 2016.
- 1710
- Hodzic, A., Campuzano-Jost, P., Bian, H., Chin, M., Colarco, P. R., Day, D. A., Froyd, K. D., Heinold, B., and et al., D. S. J.: Characterization of organic aerosol across the global remote troposphere: a comparison of ATom measurements and global chemistry models, *Atmos. Chem. Phys.*, 20, 4607–4635, <https://doi.org/10.5194/acp-20-4607-2020>, 2020.
- 1715
- Hoffa, E. A., Ward, D. E., Hao, W. M., Susott, R. A., and Wakimoto, R. H.: Seasonality of carbon emissions from biomass burning in a Zambian savanna, *Journal of Geophysical Research: Atmospheres*, 104, 13841–13853, <https://doi.org/https://doi.org/10.1029/1999JD900091>, 1999.
- 1720
- Holanda, B. A., Pöhlker, M. L., Walter, D., Saturno, J., Sörgel, M., Ditas, J., Ditas, F., Schulz, C., Franco, M. A., Wang, Q., Donth, T., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Braga, R., Brito, J., Cheng, Y., Dollner, M., Kaiser, J. W., Klimach, T., Knote, C., Krüger, O. O., Fütterer, D., Lavrič, J. V., Ma, N., Machado, L. A. T., Ming, J., Morais, F. G., Paulsen, H., Sauer, D., Schlager, H., Schneider, J., Su, H., Weinzierl, B., Walser, A., Wendisch, M., Ziereis, H., Zöger, M., Pöschl, U., Andreae, M. O., and Pöhlker, C.: Influx of African biomass burning aerosol during the Amazonian dry season through layered transatlantic transport of black carbon-rich smoke, *Atmos. Chem. Phys.*, 20, 4757–4785, <https://doi.org/10.5194/acp-20-4757-2020>, 2020.
- 1725
- Holanda, B. A., Franco, M. A., Walter, D., Artaxo, P., Carbone, S., Cheng, Y., Chowdhury, S., Ditas, F., Gysel-Ber, M., Klimach, T., Kremper, L. A., Krüger, O. O., Lavric, J. V., Lelieveld, J., Ma, C., Machado, L. A. T., Modini, R. L., Morais, F. G., Pozzer, A., Saturno, J., Su, H., Wendisch, M., Wolff, S., Pöhlker, M. L., Andreae, M. O., Pöschl, U., and Pöhlker, C.: African biomass burning affects aerosol cycling over the Amazon, *Commun Earth Environ*, 4, 154, <https://doi.org/10.1038/s43247-023-00795-5>, 2023.
- 1730
- Holder, A. L., Hagler, G. S., Aurell, J., Hays, M. D., and Gullett, B. K.: Particulate matter and black carbon optical properties and emission factors from prescribed fires in the southeastern United States, *J. Geophys. Res.*, 121, 3465–3483, <https://doi.org/10.1002/2015JD024321>, 2016.
- 1735

- Hoppel, W., Frick, G., and Larson, R.: Effect of non-precipitating clouds on the aerosol size distribution, *Geophysical Research Letters - GEOPHYS RES LETT*, 13, <https://doi.org/10.1029/GL013i002p00125>, 1986.
- 1740 Hoppel, W. A. and Frick, G. M.: Submicron aerosol size distributions measured over the tropical and South Pacific, *Atmospheric Environment. Part A. General Topics*, 24, 645–659, [https://doi.org/10.1016/0960-1686\(90\)90020-N](https://doi.org/10.1016/0960-1686(90)90020-N), 1990.
- 1745 Howell, S. G., Freitag, S., Dobracki, A., Smirnow, N., and III, A. J. S.: Undersizing of aged African biomass burning aerosol by an ultra-high-sensitivity aerosol spectrometer, *Atmos. Meas. Tech.*, 14, 7381–7404, <https://doi.org/10.5194/amt-14-7381-2021>, 2021.
- 1750 Howes, C., Saide, P. E., Coe, H., Dobracki, A. N., Freitag, S., Haywood, J. M., Howell, S. G., Gupta, S., Uin, J., Kacarab, M., Kuang, C., Leung, L. R., Nenes, A., McFarquhar, G., Redemann, J., Sedlacek, A. J., Thornhill, K. L., Wong, J. P. S., Wood, R., Wu, H., Zhang, Y., Zhang, J., and Zuidema, P.: Biomass-burning smoke properties and its interactions with marine stratocumulus clouds in WRF-CAM5 and southeastern Atlantic field campaigns, *EGU*sphere, 2023, 1–47, <https://doi.org/10.5194/egusphere-2023-886>, 2023.
- 1755 Huffman, J. A., Docherty, K. S., Mohr, C., Cubison, M. J., Ulbrich, I. M., Ziemann, P. J., Onasch, T. B., and Jimenez, J. L.: Chemically-resolved volatility measurements of organic aerosol from different sources, *Environ. Sci. Technol.*, 43, 5351–5357, <https://doi.org/10.1021/es803539d>, 2009.
- 1760 IPCC, 2022: *Climate Change 2022: Impacts, Adaptation, and Vulnerability*. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp., doi:10.1017/9781009325844.
- 1765 Jiang, Y., Zhou, L., and Raghavendra, A.: Observed changes in fire patterns and possible drivers over Central Africa, *Environmental Research Letters*, 15, 0940b8, <https://doi.org/10.1088/1748-9326/ab9db2>, 2020.
- 1770 Jolleys, M. D., Coe, H., McFiggans, G., Capes, G., Allan, J. D., Crosier, J., Williams, P. I., Allen, G., Bower, K. N., Jimenez, J. L., Russell, L. M., Grutter, M., and Baumgardner, D.: Characterizing the aging of biomass burning organic aerosol by use of mixing ratios: a meta-analysis of four regions, *Environ. Sci. Technol.*, 46, 13093–13102, 2012.
- 1775 Kacarab, M., Thornhill, K. L., Dobracki, A., Howell, S. G., O'Brien, J. R., Freitag, S., Poellot, M. R., Wood, R., Zuidema, P., Redemann, J., and Nenes, A.: Biomass burning aerosol as a modulator of the droplet number in the southeast Atlantic region, *Atmos Chem Phys*, 20, 3029–3040, <https://doi.org/10.5194/acp-20-3029-2020>, 2020.
- 1780 Khalil, M. A. K. and Rasmussen, R. A.: The global cycle of carbon monoxide: Trends and mass balance, *Chemosphere*, 20, 227–242, [https://doi.org/10.1016/0045-6535\(90\)90098-E](https://doi.org/10.1016/0045-6535(90)90098-E), 1990.
- Khalizov, A. F., Xue, H., Wang, L., Zheng, J., and Zhang, R.: Enhanced Light Absorption and Scattering by Carbon Soot Aerosol Internally Mixed with Sulfuric Acid, *J Phys Chem A*, 113, 1066–1074, <https://doi.org/10.1021/jp807531n>, 2009.
- 1785 Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Lioussé, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCWA project: Dynamics-Aerosol-Chemistry-Cloud Interactions in West Africa, *B. Am. Meteorol. Soc.*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015.

- Korontzi, S., Justice, C. O., and Scholes, R. J.: Influence of timing and spatial extent of savanna fires in southern Africa on atmospheric emissions, *J Arid Environ*, 54, 395–404, <https://doi.org/https://doi.org/10.1006/jare.2002.1098>, 2003.
- 1790 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Env.*, 42, 3593–3624, 2008.
- Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Env.*, 42, 3593–3624, 2008.
- 1795 Kroll, J.H., Smith, J. D., Che, D. L., Kessler, S. H., Worsnop, D. R., and Wilson, K. R.: Measurement of fragmentation and functionalization pathways in the heterogeneous oxidation of oxidized organic aerosol, *Phys. Chem. Chem. Phys.*, 11, 8005–8014, <https://doi.org/10.1039/B905289E>, 2009.
- 1800 Kroll, J. H., Lim, C. Y., Kessler, S. H., and Wilson, K. R.: Heterogeneous Oxidation of Atmospheric Organic Aerosol: Kinetics of Changes to the Amount and Oxidation State of Particle-Phase Organic Carbon, *J Phys Chem A*, 119, 10767–10783, <https://doi.org/10.1021/acs.jpca.5b06946>, 2015.
- Lack, D. A., Langridge, J., Bahreni, R., Cappa, C. D., Middlebrook, A., and Schwarz, J. P.: Brown Carbon and Internal Mixing in Biomass Burning Particles, *P. Natl. Acad. Sci. USA*, 10, 14802–14807, <https://doi.org/10.1073/pnas.1206575109>, 2012.
- 1805 Lanz, V. A., Prévôt, A. S. H., Alfarra, M. R., Weimer, S., Mohr, C., DeCarlo, P. F., Gianini, M. F. D., Hueglin, C., Schneider, J., Favez, O., D’Anna, B., George, C., and Baltensperger, U.: Characterization of aerosol chemical composition with aerosol mass spectrometry in Central Europe: an overview, *Atmos Chem Phys*, 10, 10453–10471, <https://doi.org/10.5194/acp-10-10453-2010>, 2010.
- 1810 Lee, J. E., Gorkowski, K., Meyer, A., Benedict, K., Aiken, A. C., and Dubey, M. K.: Wildfire smoke demonstrates significant and predictable black carbon light absorption enhancements, *Geophys. Res. Lett.*, 49, 2022.
- Li, H. and May, A. A.: Estimating mass-absorption cross-section of ambient black carbon aerosols: Theoretical, empirical, and machine learning models, *Aerosol Science and Technology*, 56, 980–997, <https://doi.org/10.1080/02786826.2022.2114311>, 2022.
- 1815 Li, J., Pósfai, M., Hobbs, P. V., and Buseck, P. R.: Individual aerosol particles from biomass burning in southern Africa: 2, Compositions and aging of inorganic particles, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/https://doi.org/10.1029/2002JD002310>, 2003.
- 1820 Li, Z., Smith, K. A., and Cappa, C. D.: Influence of relative humidity on the heterogeneous oxidation of secondary organic aerosol, *Atmos Chem Phys*, 18, 14585–14608, <https://doi.org/10.5194/acp-18-14585-2018>, 2018.
- 1825 Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S., Williams, P. I., Ting, Y.-C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, *Nat Geosci*, 10, 184–188, <https://doi.org/10.1038/ngeo2901>, 2017.
- 1830 Lund, M. T., Samset, B. H., Skeie, R. B., Watson-Parris, D., Katich, J. M., Schwarz, J. P., and Weinzierl, B.: Short Black Carbon lifetime inferred from a global set of aircraft observations, *npj Clim. Atmos. Sci.*, 1, <https://doi.org/10.1038/s41612-018-0040-x>, 2018.
- 1835 Mallet, M., Solmon, F., Nabat, P., Elguindi, N., Waquet, F., and et al., D. B.: Direct and semi-direct radiative forcing of biomass-burning aerosols over the southeast Atlantic (SEA) and its sensitivity to absorbing properties: a regional climate modeling study, *Atmos. Chem. Phys.*, 20, 13191–13216, <https://doi.org/10.5194/acp-20-13191-2020>, 2020.

- 1840 Mallet, M., Nabat, P., Johnson, B., Michou, M., Haywood, J. M., Chen, C., and Dubovik, O.: Climate models generally underrepresent the warming by Central Africa biomass-burning aerosols over the Southeast Atlantic, *Sci. Adv.*, 7, eabg9998, <https://doi.org/10.1126/sciadv.abg9998>, 2021.
- 1845 McNeill, V. F., Woo, J. L., Kim, D. D., Schwier, A. N., Wannell, N. J., Sumner, A. J., and Barakat, J. M.: Aqueous-Phase Secondary Organic Aerosol and Organosulfate Formation in Atmospheric Aerosols: A Modeling Study, *Environ Sci Technol*, 46, 8075–8081, <https://doi.org/10.1021/es3002986>, 2012.
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Chhabra, P. S., Seinfeld, J. H., and Worsnop, D. R.: Changes in organic aerosol composition with aging inferred from aerosol mass spectra, *Atmos. Chem. Phys.*, 11, 6465–6474, <https://doi.org/10.5194/acp-11-6465-2011>, 2011.
- 1850 Nguyen, T. B., Lee, P. B., Updyke, K. M., Bones, D. L., Laskin, J., Laskin, A., and Nizkorodov, S. A.: Formation of nitrogen- and sulfur-containing light-absorbing compounds accelerated by evaporation of water from secondary organic aerosols, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/https://doi.org/10.1029/2011JD016944>, 2012.
- 1855 O'Brien, R. E. and Kroll, J. H.: Photolytic Aging of Secondary Organic Aerosol: Evidence for a Substantial Photo-Recalcitrant Fraction, *J. Phys. Chem. Lett.*, 10, 4003–4009, <https://doi.org/10.1021/acs.jpcclett.9b01417>, 2019.
- Pan, X., Kanaya, Y., Taketani, F., Miyakawa, T., Inomata, S., Komazaki, Y., Tanimoto, H., Wang, Z., Uno, I., and Wang, Z.: Emission characteristics of refractory black carbon aerosols from fresh biomass burning: a perspective from laboratory experiments, *Atmos Chem Phys*, 17, 13001–13016, <https://doi.org/10.5194/acp-17-13001-2017>, 2017.
- 1860 Pennypacker, S., Diamond, M., and Wood, R.: Ultra-clean and smoky marine boundary layers frequently occur in the same season over the southeast Atlantic, *Atmos Chem Phys*, 20, 2341–2351, <https://doi.org/10.5194/acp-20-2341-2020>, 2020.
- 1865 Pistone, K., Redemann, J., Doherty, S., Zuidema, P., Burton, S., Cairns, B., Cochrane, S., Ferrare, R., Flynn, C., Freitag, S., Howell, S., Kacenelenbogen, M., LeBlanc, S., Liu, X., Schmidt, K. S., Sedlacek III, A. J., Segal-Rosenhaimer, M., Shinozuka, Y., Stamnes, S., van Diedenhoven, B., Van Harten, G., and Xu, F.: Intercomparison of biomass burning aerosol optical properties from in-situ and remote-sensing instruments in ORACLES-2016, *Atmos. Chem. Phys.*, 19, 9181–9208, <https://doi.org/10.5194/acp-19-9181-2019>, 2019.
- 1870 Redemann, J., Wood, R., Zuidema, P., Doherty, S. J., Luna, B., LeBlanc, S. E., Diamond, M. S., Shinozuka, Y., Chang, I. Y., Ueyama, R., Pfister, L., Ryoo, J.-M., Dobracki, A. N., da Silva, A. M., Longo, K. M., Kacenelenbogen, M. S., Flynn, C. J., Pistone, K., Knox, N. M., Piketh, S. J., Haywood, J. M., Formenti, P., Mallet, M., Stier, P., Ackerman, A. S., Bauer, S. E., Fridlind, A. M., Carmichael, G. R., Saide, P. E., Ferrada, G. A., Howell, S. G., Freitag, S., Cairns, B., Holben, B. N., Knobelspiesse, K. D., Tanelli, S., L'Ecuyer, T. S., Dzambo, A. M., Sy, O. O., McFarquhar, G. M., Poellot, M. R., Gupta, S., O'Brien, J. R., Nenes, A., Kacarab, M., Wong, J. P. S., Small-Griswold, J. D., Thornhill, K. L., Noone, D., Podolske, J. R., Schmidt, K. S., Pilewskie, P., Chen, H., Cochrane, S. P., Sedlacek, A. J., Lang, T. J., Stith, E., Segal-Rozenhaimer, M., Ferrare, R. A., Burton, S. P., Hostetler, C. A., Diner, D. J., Seidel, F. C., Platnick, S. E., Myers, J. S., Meyer, K. G., Spangenberg, D. A., Maring, H., and Gao, L.: An overview of the ORACLES (ObseRvations of Aerosols above CLouds and their intERactionS) project: aerosol–cloud–radiation interactions in the southeast Atlantic basin, *Atmos. Chem. Phys.*, 21, 1507–1563, <https://doi.org/10.5194/acp-21-1507-2021>, 2021.
- 1880 Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: and Physics A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics Discussions*, 4, 799–825, <https://doi.org/10.5194/acpd-4-5135-2004>, 2005.

- 1885 Rickly, P. S., Guo, H., Campuzano-Jost, P., Jimenez, J. L., Wolfe, G. M., Bennett, R., Bourgeois, I., Crouse, J. D., Dibb, J. E., DiGangi, J. P., Diskin, G. S., Dollner, M., Gargulinski, E. M., Hall, S. R., Halliday, H. S., Hanisco, T. F., Hannun, R. A., Liao, J., Moore, R., Nault, B. A., Nowak, J. B., Peischl, J., Robinson, C. E., Ryerson, T., Sanchez, K. J., Schöberl, M., Soja, A. J., St. Clair, J. M., Thornhill, K. L., Ullmann, K., Wennberg, P. O., Weinzierl, B., Wiggins, E. B., Winstead, E. L., and Rollins, A. W.: Emission factors and evolution of SO<sub>2</sub> measured from biomass burning in wildfires and agricultural fires, *Atmos Chem Phys*, **22**, 15603–15620, <https://doi.org/10.5194/acp-22-15603-2022>, 2022.
- Ryoo, J.-M., Pfister, L., Ueyama, R., Zuidema, P., Wood, R., Chang, I., and Redemann, J.: A meteorological overview of the ORACLES (ObseRvations of Aerosols above CLouds and their intEractionS) campaign over the southeast Atlantic during 2016–2018: Part 1 - Climatology, *Atmos. Chem. Phys.*, **21**, 16689–16707, <https://doi.org/10.5194/acp-21-16689-2021>, 2021.
- 1895 Ryoo J.-M., L. Pfister, R. Ueyama, P. Zuidema, R. Wood, I. Chang, J. Redemann, 2022: A meteorological overview of the ORACLES (ObseRvations of Aerosols above CLouds and their intEractionS) campaign over the southeast Atlantic during 2016–2018: Part 2 – daily and synoptic characteristics. *Atmos. Chem. Phys.*, **22**, p. 14209–14241, doi:10.5194/acp-22-14209-2022
- 1900 Schnaiter, M., Linke, C., Möhler, O., Naumann, K.-H., Saathoff, H., Wagner, R., Schurath, U., and Wehner, B.: Absorption amplification of black carbon internally mixed with secondary organic aerosol, *Journal of Geophysical Research: Atmospheres*, **110**, <https://doi.org/https://doi.org/10.1029/2005JD006046>, 2005.
- Sedlacek, A., Lewis, E., Onasch, T., Zuidema, P., Redemann, J., Jaffee, D., and Kleinman, L.: Using the black carbon particle mixing state to characterize the lifecycle of biomass burn aerosols, *Env. Sci. Tech.*, **56**, 14315–14325, <https://doi.org/10.1021/acs.est.2c30851>, 2022.
- 1910 Shinozuka, Y., Saide, P. E., Ferrada, G. A., Burton, S. P., Ferrare, R., Doherty, S. J., Gordon, H., Longo, K., Mallet, M., Feng, Y., Wang, Q., Cheng, Y., Dobracki, A., Freitag, S., Howell, S. G., LeBlanc, S., Flynn, C., Segal-Rosenhaimer, M., Pistone, K., Podolske, J. R., Stith, E. J., Bennett, J. R., Carmichael, G. R., da Silva, A., Govindaraju, R., Leung, R., Zhang, Y., Pfister, L., Ryoo, J.-M., Redemann, J., Wood, R., and Zuidema, P.: Modeling the smoky troposphere of the southeast Atlantic: a comparison to ORACLES airborne observations from September of 2016, *Atmos. Chem. Phys.*, **20**, 11, 411–491, 526, <https://doi.org/10.5194/acp-20-11491-2020>, 2020.
- 1915 Sinha, P., Hobbs, P. V., Yokelson, R. J., Blake, D. R., Gao, S., and Kirchstetter, T. W.: Emissions from miombo woodland and dambo grassland savanna fires, *Journal of Geophysical Research: Atmospheres*, **109**, <https://doi.org/https://doi.org/10.1029/2004JD004521>, 2004.
- 1920 Sorooshian, A., Murphy, S. M., Hersey, S., Bahreini, R., Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Constraining the contribution of organic acids and AMS m/z 44 to the organic aerosol budget: On the importance of meteorology, aerosol hygroscopicity, and region, *Geophys Res Lett*, **37**, <https://doi.org/https://doi.org/10.1029/2010GL044951>, 2010.
- Taylor, J. W., Wu, H., Szpek, K., Bower, K., Crawford, I., Flynn, M. J., Williams, P. I., Dorsey, J., Langridge, J. M., and et al.: Absorption closure in highly aged biomass burning smoke, *Atmos. Chem. Phys.*, **20**, 11201–11221, <https://doi.org/10.5194/acp-20-11201-2020>, 2020.
- 1925 Uin, J., Aiken, A. C., Dubey, M. K., Kuang, C., Pekour, M., Salwen, C., Sedlacek, A. J., Senum, G., Smith, S., Wang, J., Watson, T. B., and Springston, S. R.: Atmospheric Radiation Measurement (ARM) Aerosol Observing Systems (AOS) for Surface-Based In Situ Atmospheric Aerosol and Trace Gas Measurements, *J Atmos Ocean Technol*, **36**, 2429–2447, <https://doi.org/10.1175/JTECH-D-19-0077.1>, 2019.
- 1930 Vakkari, V., Beukes, J. P., Maso, M. D., Aurela, M., Josipovic, M., and van Zyl, P. G.: Major secondary aerosol formation in southern African open biomass burning plumes, *Nature Geo.*, **11**, 580–583, <https://doi.org/10.1038/s41561-018-0170-0>, 2018.

- 1935 Wang, C.: A modeling study on the climate impacts of black carbon aerosols, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004084>, 2004.
- Wong, J. P. S., Zhou, S., and Abbatt, J. P. D.: Changes in Secondary Organic Aerosol Composition and Mass due to Photolysis: Relative Humidity Dependence, *J Phys Chem A*, 119, 4309–4316, <https://doi.org/10.1021/jp506898c>, 2015.
- 1940 Wu, H., Taylor, J. W., Szpek, K., Langridge, J. M., Williams, P. I., Flynn, M., Allan, J. D., Abel, S. J., Pitt, J., Cotterell, M. I., Fox, C., Davies, N. W., Haywood, J., and Coe, H.: Vertical variability of the properties of highly aged biomass burning aerosol transported over the southeast Atlantic during CLARIFY-2017, *Atmos. Chem. Phys.*, 20, 12697–12719, <https://doi.org/10.5194/acp-20-12697-2020>, 2020.
- 1945 Yang, M., Huebert, B. J., Blomquist, B. W., Howell, S. G., Shank, L. M., McNaughton, C. S., Clarke, A. D., Hawkins, L. N., and et al., L. M. R.: Atmospheric sulfur cycling in the southeastern Pacific – longitudinal distribution, vertical profile, and diel variability observed during VOCALS-REx, *Atmos. Chem. Phys.*, 11, 5079–5097, <https://doi.org/10.5194/acp-11-5079-2011>, 2011.
- 1950 Zanatta, M., Gysel, M., Bukowiecki, N., Müller, T., Weingartner, E., Areskou, H., Fiebig, M., Yttri, K. E., Mihalopoulos, N., Kouvarakis, G., Beddows, D., Harrison, R. M., Cavalli, F., Putaud, J. P., Spindler, G., Wiedensohler, A., Alastuey, A., Pandolfi, M., Sellegri, K., Swietlicki, E., Jaffrezo, J. L., Baltensperger, U., and Laj, P.: A European aerosol phenomenology-5: Climatology of black carbon optical properties at 9 regional background sites across Europe, *Atmos Environ*, 145, 346–364, <https://doi.org/10.1016/j.atmosenv.2016.09.035>, 2016.
- 1955 Zhang, J. and Zuidema, P.: The diurnal cycle of the smoky marine boundary layer observed during August in the remote southeast Atlantic, *Atmos Chem Phys*, 19, 14493–14516, <https://doi.org/10.5194/acp-19-14493-2019>, 2019.
- 1960 Zhang, J. and Zuidema, P.: Sunlight-absorbing aerosol amplifies the seasonal cycle in low-cloud fraction over the southeast Atlantic, *Atmos. Chem. Phys.*, 21, 11179–11199, <https://doi.org/10.5194/acp-21-11179-2021>, 2021.
- 1965 Zhang, L., Segal-Rozenhaimer, M., Che, H., Dang, C., Sedlacek III, A. J., Lewis, E. R., Dobracki, A., Wong, J. P. S., Formenti, P., Howell, S. G., and Nenes, A.: Light absorption by brown carbon over the South-East Atlantic Ocean, *Atmos Chem Phys*, 22, 9199–9213, <https://doi.org/10.5194/acp-22-9199-2022>, 2022.
- Zhang, Q., Jimenez, J. L., and Canagaratna, M. R.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry : a review, 3045–3067, <https://doi.org/10.1007/s00216-011-5355-y>, 2011.
- 1970 Zhang, Y., Favez, O., and et al., F. C.: Evidence of major secondary organic aerosol contribution to lensing effect black carbon absorption enhancement, *npj Clim. Atmos. Sci.*, 1, <https://doi.org/10.1038/s41612-018-0056-2>, 2018.
- 1975 Zuidema, P., Redemann, J., Haywood, J., Wood, R., Piketh, S., Hipondoka, M., and Formenti, P.: Smoke and Clouds above the Southeast Atlantic: Upcoming Field Campaigns Probe Absorbing Aerosol’s Impact on Climate, *Bull. Am. Meteor. Soc.*, 97, 1131–1135, <https://doi.org/10.1175/bams-d-15-00082.1>, 2016.
- Zuidema, P., Sedlacek III, A. J., Flynn, C., Springston, S., Delgadillo, R., Zhang, J., Aiken, A. C., Koontz, A., and Muradyan, P.: The Ascension Island Boundary Layer in the Remote Southeast Atlantic is Often Smoky, *Geophys. Res. Lett.*, 45, 4456–4465, <https://doi.org/10.1002/2017gl076926>, 2018.