

# 1 Water masses in the Atlantic Ocean: water mass ages and ventilation

2 Mian Liu<sup>1,2</sup> and Toste Tanhua<sup>2</sup>

3 <sup>1</sup>School of Environmental Science and Engineering, Xiamen University of Technology, Xiamen, 361021, China

4 <sup>2</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Marine Biogeochemistry, Chemical Oceanography, Wischhofstraße  
5 1-3, 24148 Kiel, Germany

6 Correspondence to: Mian Liu (2023000069@xmut.edu.cn)

7 **Abstract:** The distribution of water masses and their characteristics, including ventilation, provides fundamental insights into  
8 large-scale oceanographic processes such as thermohaline circulation and marine biogeochemical cycles. The characteristics  
9 of main water masses in the Atlantic Ocean have been comprehensively documented in a companion study (Liu and Tanhua,  
10 2021), this study presents quantitative assessments of water mass age characteristics and ventilation time scales through an  
11 analysis using the transient tracers chlorofluorocarbon-12 (CFC-12), sulfur hexafluoride (SF<sub>6</sub>), and argon-39 (<sup>39</sup>Ar). We use  
12 two distinct age concepts: mean-age as an integrative metric of water mass chronology, and mode-age as a proxy for advective  
13 time scales. Empirical results demonstrate systematic age progression with increasing depth and along water mass trajectories.  
14 Surface layer central waters exhibit mean ages up to ~100 years and mode ages reaching ~30 years. In the intermediate layer,  
15 meridional age gradients characterize the Antarctic Intermediate Water (AAIW) reaching maximum mean-age (~300 years)  
16 and mode-age (~80 years) at 30 °N, whereas zonal variations manifest in Mediterranean Water (MW) with peak values (~400  
17 years in mean-age, ~100 years in mode-age) observed in equatorial regions. As the dominant deep water component, North  
18 Atlantic Deep Water (NADW) exhibits extreme ages in the Antarctic Circumpolar Current (ACC) region at 50°S, achieving  
19 mean age ~600 years and mode age ~100 years. Bottom layer water masses display their oldest signatures: Antarctic Bottom  
20 Water (AABW) from the Weddell Sea reaches ~600 years mean-age and ~100 years mode-age at equatorial latitudes, while its  
21 extension, Northeast Atlantic Bottom Water (NEABW), attains exceptional values of ~800 years (mean) and ~120 years (mode)  
22 at 50°N. The age analysis reveals significant basin-scale asymmetries, with western basins exhibiting younger ages compared  
23 to western counterparts. Ventilation efficiency modulates these age distributions, as evidenced by lower mode-ages and reduced  
24 apparent oxygen utilization (AOU) in better-ventilated western basins. The calculated oxygen utilization rate (OUR)  
25 demonstrates spatial concordance with dissolved oxygen (DO) concentrations, corroborating enhanced oxidative processes in  
26 high-oxygen regimes. This integrated age framework provides novel insights into water mass ventilation dynamics and their  
27 biogeochemical implications through quantitative characterization of temporal-spatial age distributions across multiple  
28 oceanographic provinces.

29 **Key words:** Water Mass, Atlantic Ocean, Transient Tracer, Mean- and Mode-age, Ventilation, GLODAPv2 data product

## 30 1 Introduction

31 The Atlantic Ocean is composed by a complex vertical stratification of water masses characterized by differences in properties  
32 such as salinity, temperature and oxygen. Liu and Tanhua (2021) have systematically characterized 16 main water masses that  
33 have been extensively considered and studied due to their significant impacts on ocean circulation patterns and global climate  
34 regulation (e.g., Hall & Bryden, 1982; Bryden et al., 2005; Kuhlbrodt et al., 2007; Clark et al., 2012). Ocean ventilation  
35 constitutes a fundamental thermohaline process responsible for the vertical displacement of surface waters into the deep ocean,  
36 effectively redistributing surface heat and salinity anomalies through density-driven stratification. This process inherently  
37 obeys the principle of mass conservation, simultaneously enabling the upwelling of deep-sea waters to the surface ocean. The  
38 resultant vertical exchange plays a critical role in maintaining global heat distribution patterns and sustaining the  
39 thermodynamic equilibrium of the ocean system (Talley, 2013; Armour et al., 2016; Talley et al., 2016). Beyond its  
40 thermodynamic significance, ocean ventilation emerges as a pivotal mechanism governing marine biogeochemical cycles. By  
41 mediating the downward transport of atmospheric oxygen and carbon dioxide to abyssal depths, this process fundamentally  
42 regulates dissolved oxygen concentrations and carbon sequestration rates in the deep ocean (Tanhua et al., 2006; Ziska et al.,  
43 2013; Skinner et al., 2017). Related studies have further elucidated ventilation's deterministic role in controlling abyssal redox  
44 conditions and organic carbon remineralization processes (van Heuven et al., 2011; Ito et al., 2016; Schmidtke et al., 2017).

45 Early studies of the water mass ventilation are proposed by Sandström already in the early 20th century (Sandström, 1908;  
46 1916). Sverdrup and Bjerknes also reported the impacts of ventilation and currents on the climate through the air-sea  
47 interactions (Sverdrup, 1940; Bjerknes, 1964). Hereafter many studies have been focusing on the ventilation. For instance,  
48 Dickson et al., (1994) estimated the formation and pathway of the North Atlantic Deep Water (NADW) based on the  
49 hydrodynamics, Orsi et al., (1999) investigated the circulation of the Antarctic Bottom Water (AABW) by using  
50 Chlorofluorocarbons (CFCs). Recent studies suggest that the intensity of ocean ventilation varies with environmental factors,  
51 such as intensity of the westerly winds (e.g. Rahmstorf, 2010; Purkey et al., 2010; Morrison et al., 2015). Several studies  
52 investigate the regional scale or specific water masses, for instance, the upwelling of Circumpolar Deep Water (CDW) in the  
53 Southern Ocean (Tamsitt et al., 2017), or the ventilation in the South China Sea (Wang et al., 2021). In parallel, advances in  
54 ocean modeling have provided new insights into water mass transformation and tracer transport on basin to global scales,  
55 offering valuable perspectives on the climatic drivers of ventilation variability (e.g., Li et al., 2023). However, few studies  
56 look at the ventilation over ocean basin scales, which is helpful to understand the effect of ocean-climate interactions, and in  
57 addition can provide a basis for the biogeochemical studies.

58 The water mass age is an important parameter to evaluate the ventilation and refers to as the elapsed time on the pathway. The  
59 CFCs and Sulfur hexafluoride ( $\text{SF}_6$ ) are recognized as effective transient tracers for water masses and to estimate their ages,  
60 i.e. the time since the water left the surface ocean (Fine, 2011). The concentrations of CFC-12 in surface seawater increased  
61 continuously following the increase of the atmospheric concentration since their introduction in the early 20<sup>th</sup> century  
62 (Gammon et al., 1982; Warner and Weiss, 1985) up to about the turn of the century. The use of CFCs as a tracer in the

63 oceanography is well established (e.g. Bullister and Weiss, 1983; Doney and Bullister, 1992; Fine, 2010; 2011), and transient  
64 tracers is a core variable in the GO-SHIP repeat hydrography program. Similarly, the transient tracer is an Essential Ocean  
65 Variable (EOV) in the Global Ocean Observing System (GOOS) framework. SF<sub>6</sub> has been used as a new tracer since mid-  
66 1990s complementing the CFCs due to the reduction of CFCs in the atmosphere (Maiss et al., 1996; Bullister et al., 2002). The  
67 application of SF<sub>6</sub> is generally appropriate for recently formed water masses in the upper ocean (Tanhua et al., 2008), i.e. for  
68 well ventilated, or young, waters. The CFCs are used to trace the relative deeper water masses (before mid-1990s or partial  
69 pressure of CFC-12 lower than about 450 parts per trillion, ppt), while the SF<sub>6</sub> is the better choice for recently formed shallow  
70 water masses (after mid-1990s or partial pressure of CFC-12 higher than about 450 ppt, Tanhua et al., 2008). The  
71 complementary nature of CFCs and SF<sub>6</sub> allows for more accurate tracing of water masses (Law and Watson, 2001; Vollmer  
72 and Weiss, 2002; Watanabe et al., 2003; Tanhua et al., 2004, 2005; Bullister et al., 2006). In addition, the application of  
73 transient tracers also provides support to the hydrographic and biogeochemical field in calculating the upwelling velocity  
74 (Tanhua and Liu, 2015) or estimating the ventilation and anthropogenic carbon cycle (van Heuven et al., 2011; Tanhua et al.,  
75 2013; Patara et al., 2021).

76 As two typical representatives of transient tracers, the CFC-12 and SF<sub>6</sub> are commonly employed for determining water mass  
77 ages, verifying hydrological models, and estimating rates of marine biogeochemical processes, such as anthropogenic carbon  
78 storage and oxygen consumption. They exhibit advantages in tracing upper and middle water masses with relatively younger  
79 ages. However, challenges arise when attempting to obtain data from deep and bottom layers with old waters. The chemically  
80 inert isotope argon-39 (<sup>39</sup>Ar), which has a half-life of 269 years, addresses this gap. <sup>39</sup>Ar is particularly advantageous for tracing  
81 deep and bottom water masses with ages exceeding 500 years (Broecker & Peng, 2000; Holzer & Primeau, 2010; Lu et al.,  
82 2014). Initially, its application was constrained by the requirement for large sampling volumes—typically around 1000 liters—  
83 and the limitations of low-level counting (LLC) detection methods (Loosli et al., 1983; Schlosser et al., 1994; Schlitzer et al.,  
84 1985). Consequently, <sup>39</sup>Ar was not widely utilized in oceanographic tracer studies. With advancements in Atom Trap Trace  
85 Analysis (ATTA) technology (Chen et al., 1999; Jiang et al., 2011), <sup>39</sup>Ar has gradually gained prominence as an effective tracer  
86 for investigating deep and bottom waters (Lu et al., 2014; Ebser et al., 2018), the accuracy and stability have also been  
87 significantly improved (Jia et al., 2025). Therefore, the multi-parameter combination of CFCs, SF<sub>6</sub>, and <sup>39</sup>Ar can transcend the  
88 temporal limitations associated with single-tracer dating methodologies while providing more comprehensive insights into  
89 biogeochemical processes. In this study, we introduce <sup>39</sup>Ar to investigate water mass ages within the Atlantic Ocean, in  
90 particular for the older waters.

91 The apparent oxygen utilization (AOU,  $\mu\text{mol kg}^{-1}$ ) is an important indicator in characterizing the respiration of organic matter  
92 consuming oxygen, and balancing the oxygen supply through ventilation and photosynthesis. The oxygen utilization rate (OUR,  
93  $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ ), which can be calculated from AOU and water mass age, is a significant parameter in estimating the  
94 biogeochemical processes such as primary productivity or remineralization (e.g. Jenkins, 1982; Bender, 1990). The OUR has  
95 been recognized to be a function of pressure for a long time (e.g. Tseitlin, 1992). However, the OUR is the integrated oxygen

consumption rate along the pathway of a water mass and represents a regional view of export flux (e.g. Stanley et al., 2012; Koeve and Kähler, 2016), and thus an imperfect measure of local OUR. In this study, the OUR is calculated from water mass ages estimated from the CFC-12 and SF<sub>6</sub>. The change of OUR over time in different water masses are presented and the impacts from the currents and topography are discussed.

In the study by Liu and Tanhua, (2021), the characteristics of main water masses in the Atlantic Ocean are defined by six key properties, using the Global Ocean Data Analysis Project version 2 the annual update 2023 (GLODAPv2.2023, Lauvset et al., 2016; Olsen et al., 2016; Lauvset et al., 2024) data product as a source of unbiased data. The static distribution of the water masses is estimated with the Optimal Multi-Parameter analysis (OMP analysis, Karstensen and Tomczak, 1997; 1998). As a continuation of that work, here we estimate the ventilation of water masses in the Atlantic Ocean with the combination of the above mentioned three transient tracers (CFC-12, SF<sub>6</sub> and <sup>39</sup>Ar). The water mass ages are estimated and the OURs are calculated. The goal of this study is to further improve the report of water masses in the Atlantic Ocean by adding the age as a measure of ventilation, and to provide a hydrographic assistant for the biogeochemical researches.

In this article, we start in Section 2 with an overview of the data sources and methodologies employed. Section 3 presents the core results, illustrating the spatial distributions of mean and mode ages for the principal Atlantic Ocean water masses, stratified from surface to bottom. In Section 4, the derived water mass ages are applied to estimate Oxygen Utilization Rates (OUR) and to examine associated biogeochemical implications. Section 5 offers a comparative analysis of water mass ages obtained from traditional transient tracers (CFC-12 and SF<sub>6</sub>) and the radioisotope <sup>39</sup>Ar. Finally, Section 6 provides a comprehensive summary of conclusions, discusses the broader relevance of the findings, and acknowledges the limitations of the present study.

## 2 Data and methods

### 2.1 Transit Time Distribution (TTD) and the Inverse Gaussian Model

To quantify the ventilation timescales of water masses, we apply the Transit Time Distribution (TTD) framework. The TTD, denoted as  $G(\tau)$ , represents the distribution of transit times  $\tau$  that water parcels take to travel from the surface formation region to an interior observation point (Hall and Plumb, 1994; Waugh et al., 2003). The concentration of a transient tracer in the ocean interior,  $c(t)$ , can be modeled as the convolution of its time-evolving atmospheric source function,  $c_0(t)$ , with the TTD:

$$c(t) = \int_0^{\infty} c_0(t - \tau) G(\tau) d\tau$$

A common and practical parameterization of the TTD is the one-parameter Inverse Gaussian (IG) distribution (Waugh et al., 2003, 2004), which represents a solution to the one-dimensional advection-diffusion equation for a steady-state flow. The IG-TTD is defined as:

$$G(\tau) = \sqrt{\frac{\Gamma^3}{4\pi\Delta^2\tau^3}} \exp\left(-\frac{\Gamma(\tau-\Gamma)^2}{4\Delta^2\tau}\right);$$

125 Where:

- 126 •  $\tau$  is the transit time (in years);
- 127 •  $\Gamma$  is the mean-age (in years), representing the first moment of the TTD:  $\Gamma = \int_0^\infty \tau G(\tau) d\tau$ , which signifies the average  
128 time elapsed since the water was last in contact with the atmosphere;
- 129 •  $\Delta$  is the width (in years) of the TTD, which quantifies the spread of transit times around the mean age due to mixing and  
130 dispersion processes.

131 The mixing ratio, defined as  $(\Delta/\Gamma)$ , unitless), indicates the relative importance of mixing versus advection in the transport. A  
132 low ratio ( $\ll 1$ ) suggests advection-dominated transport with a narrow range of transit times, while a high ratio ( $\gg 1$ ) indicates  
133 strong mixing and a wide range of transit times.

134 While other forms of TTD, for instance, a two-IG distribution to characterize a mix of two water masses with different TTSs  
135 was used by Stöven and Tanhua (2014) for the Mediterranean Sea, or non-steady-state circulations may better represent reality  
136 in certain regions, the one-parameter IG-TTD remains a widely adopted and practical model. This is particularly true given  
137 the typical scarcity of multiple, independent tracer observations needed to constrain more complex distributions across large  
138 ocean basins. In this study, we apply the one-parameter IG-TTD framework as a robust approach for basin-scale analysis.


## 139 2.2 Definitions of Water Mass Age: Tracer-age, Mean-age, and Mode-age

140 The fundamental concept of water mass age is the time elapsed since a water parcel was last in equilibrium with the atmosphere  
141 at the surface (Thiele and Sarmiento, 1990). From the TTD framework, we derive and utilize three specific age definitions,  
142 each with its own physical interpretation and application (see conceptual diagram in Fig. 1).

143 **Tracer-age** is the simplest age estimate, obtained by matching the observed tracer partial pressure in a water sample directly  
144 to the historical atmospheric record (Fig. 1 a). It assumes the transport along a purely advective pathway (i.e., no mixing),  
145 corresponding to a TTD that approaches a delta function as  $\Delta/\Gamma \rightarrow 0$ . This concept systematically underestimates the “true”  
146 age in the presence of mixing, as it ignores the distribution of transit times/pathways (Sonnerup et al., 2001).

147 **Mean-age ( $\Gamma$ )** is the first moment of the TTD. The mean-age represents the average value of the transport times of all water  
148 parcel and represents an integrative timescale (Fig. 1 b), particularly applicable to the study of biogeochemical accumulation  
149 processes, such as calculating integrated oxygen consumption rates (e.g., Jenkins, 1987; Sonnerup et al., 2013).

150 **Mode-age ( $\tau_{\text{mode}}$ )** is the transit time at which the TTD,  $G(t)$ , reaches its maximum (Fig. 1 b). For the IG-TTD, it is calculated  
151 as  $\tau_{\text{mode}} = \Gamma(\sqrt{1 + (3\Delta^2/\Gamma^2)} - 1)$ . The mode-age represents the most probable transit time of the dominant water fraction in  
152 the sample, making it a better proxy for the advective timescale along the core pathway of a water mass.



**Figure 1.** Schematic diagram illustrating three concepts of water mass ages.

- (a) The tracer age: The transient tracer (such as CFC-12) in the water sample in equilibrium with the atmospheric at a certain year. This approach assumes that the ocean is a purely advective process (without mixing);
- (b) The concepts of mean- and mode-ages: Based on the inverse Gaussian transport time distribution model, assuming the mixing ratio ( $\Delta/\Gamma = 1$ ) and observed tracer partial pressure CFC-12 = 300 ppt, as an example. Shown are the positions of the mean-age ( $\Gamma$ ), tracer age, and the mode-age ( $\tau_{\text{mode}}$ ).

153 To calculate tracer partial pressures from measured concentrations, the saturation of transient tracers (CFC-12 and SF<sub>6</sub>) in  
 154 surface seawater must be determined. This saturation is a function of potential temperature ( $\theta$ ) and practical salinity (S),  
 155 following established solubility relationships (Warner and Weiss, 1985; Bullister et al., 2002).


### 156 2.3 Determination of the Mixing Ratio and age calculation procedure

157 A critical step in applying the IG-TTD is specifying the shape parameter, the mixing ratio ( $\Delta/\Gamma$ ). The atmospheric  
 158 concentrations of these tracers provide the essential input for age calculations: the CFC-12 increased monotonically until mid-  
 159 1990s, while the SF<sub>6</sub> is still increasing (Fig. 2a). Ideally, the mixing ratio is determined by fitting the TTD model to observations  
 160 of two or more transient tracers with different input histories. However, the similar atmospheric growth curves of CFC-12 and  
 161 SF<sub>6</sub> often preclude a robust, independent determination of  $\Delta/\Gamma$  across large ocean basins (Waugh et al., 2004), and the short  
 162 atmospheric history of SF<sub>6</sub> reduce this method to recently ventilated waters. Furthermore, the limited number of collocated  
 163 transient tracer observations restricts the application of more complex models. In this context, understanding how the mixing  
 164 ratio influences age estimates is crucial. For instance, under given tracer partial pressure (e.g., CFC-12 = 300 ppt) and sampling  
 165 year (e.g., 1990), an increase in the  $\Delta/\Gamma$  ratio of the IG-TTD leads to a corresponding increase in the calculated mean-age,  
 166 while the mode-age decreases. This behavior arises from the shape of the Inverse Gaussian distribution: a higher mixing ratio  
 167 extends the distribution's tail, elevating the mean-age, while simultaneously shifting its peak earlier, thereby reducing the  
 168 mode-age (Fig. 2b). Observational evidence from hydrographic sections further supports this theoretical relationship,  
 169 demonstrating how the resulting age fields vary systematically under different assumed mixing ratios (Fig. 3)

170 To ensure a consistent and widely comparable analysis across the whole Atlantic Ocean, we adopt the standard mixing  
 171 ratio  $\Delta/\Gamma = 1$ . This value represents a balance between advective and diffusive processes and is commonly used in large-scale  
 172 oceanographic studies (Waugh et al., 2004; Stanley et al., 2012; Stöven and Tanhua, 2014; Thomas et al., 2020). We


173 acknowledge this as a limitation but note it is a necessary and widely accepted approach for basin -scale analyses where tracer  
 174 data are limited.

175 The mean-age ( $\Gamma$ ) for each sample was calculated by finding the value that minimizes the difference between the observed  
 176 tracer partial pressures and those predicted by convolving the IG-TTD (with  $\Delta/\Gamma = 1$ ) with the atmospheric histories of CFC-  
 177 12 and SF<sub>6</sub>. The atmospheric histories for the Northern Hemisphere were taken from Bullister et al., (2002). Where both CFC-  
 178 12 and SF<sub>6</sub> data were available, they were used jointly in the inversion. Following Tanhua et al. (2008), in waters where the  
 179 CFC-12 partial pressure was above ~450 ppt (typically younger, shallower waters), the solution relied more heavily on SF<sub>6</sub>,  
 180 whereas CFC-12 was the primary tracer used at lower partial pressures (typically older, deeper waters). The corresponding  
 181 mode-age was then calculated from the derived mean-age and the fixed mixing ratio. For the radioactive tracer <sup>39</sup>Ar, the  
 182 convolution integral incorporates an additional exponential decay term,  $e^{-\lambda t}$ , where  $\lambda$  is the decay constant corresponding to  
 183 its 269-year half-life.



**Figure 2.** The relationships between mean-age, mode-age and mixing ratio ( $\Delta/\Gamma$ ).







- (a) Historical atmospheric partial pressure data of transient tracers over time (CFC-12 and SF<sub>6</sub>) in the Northern Hemisphere (data source: Bullister, 2002);
- (b) Relationship between mean-ages and mode-ages under same partial pressure (CFC-12 = 300 ppt) but different mixing ratios ( $\Delta/\Gamma$ ) assuming the sampling year is 1990.



**Figure 3.** (a) Map of the A16 section under the framework of the WOCE/GO-SHIP project in 2013. (b) Distributions of mean-ages and (c) mode-ages under varying mixing ratios ( $\Delta\Gamma = 0.8, 1.0, 1.2$ , and  $1.4$ ) derived from GLODAPv2 observational data. Green contour lines correspond to mean-ages of 50, 300, and 700 years and mode-ages of 50, 100, and 150 years, respectively.

### 185 3 Results: Ages of water masses in the Atlantic Ocean

186 In this study, we follow the division of vertical layers and distributions of water masses presented by Liu and Tanhua (2021),  
 187 but now focus on the transient tracers and water mass ages. Three hydrographic sections are selected from the WOCE/GO-  
 188 SHIP sections in this work to represent the mean-ages of the main water masses (Table 1). The A16 sections in 2013 (Expo-  
 189 code 33RO20130803 and 33RO2013 1223) show the meridional distribution across the whole Atlantic Ocean, while the A05  
 190 section in 2010 (Expo-code 74DI20100106) and A10 section in 2011 (Expo-code 33RO20110926) show the zonal distributions  
 191 in the North and South Atlantic Ocean respectively (Fig. 4). The partial pressures of CFC-12 and SF<sub>6</sub> along the above sections  
 192 are shown in Fig. 4 (Only CFC-12 data are displayed along A05 section due to the lack of SF<sub>6</sub> data in this section). Both tracers  
 193 show similar distributions with high values in the shallow layers, and a general decrease with increasing pressure.



**Figure 4.** Map of three selected WOCE/GO-SHIP sections to represent the ages of the main water masses in the Atlantic Ocean (middle left) and partial pressures of CFC-12 and SF<sub>6</sub> along the three selected sections. Note the different colorbars between upper and lower parts divided at pressure of 1000 dbar. The green contour lines indicate the partial pressures of CFC-12 at 10, 50 and 300 ppt and SF<sub>6</sub> at 0.2, 1, 2 and 5 ppt, respectively.


195 **Table 1:** Summary of hydrographic cruises in the Atlantic Ocean selected for the oceanographic sections in this study

| WOCE/GO-SHIP<br>section | Year | Expo-code    | Ship            | Chief Scientist                    | PI of transient<br>tracers |
|-------------------------|------|--------------|-----------------|------------------------------------|----------------------------|
| A16N                    | 2013 | 33RO20130803 | Ronald H. Brown | John Bullister &<br>Molly Baringer | John Bullister             |
| A16S                    | 2013 | 33RO20131223 | Ronald H. Brown | Rik Wanninkhof                     | John Bullister             |
| A05                     | 2010 | 74DI20100106 | Discovery       | Brian King                         | Marie-José Messias         |
| A10                     | 2011 | 33RO20110926 | Ronald H. Brown | Molly Baringer                     | John Bullister             |

196 The upper water layers have the lowest mean-ages within ~100 years and mode-ages within ~30 years (Fig. 5), the surface  
197 water has, by definition, zero age. The distributions of mean-ages and mode-ages also show spatial differences in the horizontal  
198 direction. Relatively low ages are found in the high latitude regions (Fig. 5, a and b). This indicates that the deep water masses  
199 are newly formed here, and thus have low ages. In the region between 20 and 40 °N, the mean-age reaches the peak value up  
200 to ~1000 years and the mode-age up to ~150 years, suggesting the sluggish water exchange and long residence time of old  
201 water masses. In the zonal direction, both mean-ages and mode-ages are significantly lower in the west in the general region  
202 below 1500m dbar, suggesting that these regions are better ventilated due to the western boundary current (Fig. 5 c to f).

203 In the analysis of water mass ages in each layer, maps with mean-ages and mode-ages in each station are firstly presented as a  
204 qualitative overview. The mean values of the ages at the core pressure of each water mass are plotted at certain station. In  
205 addition, the quantitative calculations are made along the three selected sections to represent the distributions of water mass  
206 ages in detail and estimate the impacts from currents and mixing.

207



**Figure 5.** Mean-ages and Mode-ages along the three selected WOCE/GO-SHIP sections


### 3.1 The upper layer

The spreading of central waters can be traced by the distributions of ages. After leaving their formation areas in the mid-latitude regions, the West North Atlantic Central Water (WNACW) and West South Atlantic Central Water (WSACW) spread generally in zonal direction towards the formation areas of the East North Atlantic Central Water (ENACW) and East South Atlantic Central Water (ESACW) with Azores Current and South Atlantic Current, respectively (Fig. 6, a). The mean -ages show that the “mixed” age of all the compositions in the western central waters is ~50 years (Fig. 6, b and Fig. 7). When focusing on the main body of WNACW and WSACW, the mode-ages show that the advective time-scale is ~15 years for both water masses (i.e. Both WNACW and WSACW take ~15 years from the formation area to the equator, Fig. 6, c and Fig. 7).

216 At greater depths, the ENACW and ESACW are transported from their formation areas and spread with the main currents  
 217 towards the equator (Fig. 6, a). In the northern hemisphere, the ENACW spreads southward with the Canary Current, while  
 218 the ESACW in the southern hemisphere is transported northward with the Benguela Current and the South Equatorial Current.  
 219 The difference in mean-age is  $\sim 100$  years between the formation areas and the equatorial region (Fig. 6, b and Fig. 7). The  
 220 mode-ages indicate that the main body of ENACW and ESACW takes approximately 30 years to be transported from formation  
 221 areas to the equator (Fig. 6, c and Fig. 7).


222 The central waters occupy the upper layer above the neutral density isoline of  $27.10 \text{ kg m}^{-3}$ . The core pressures of western and  
 223 eastern central waters are around the neutral density ( $\gamma$ ) of 26.5 and  $26.9 \text{ kg m}^{-3}$  respectively (Liu and Tanhua, 2021). The ages  
 224 of central waters at their core neutral densities are shown in Fig. 6. The eastern central waters have mean-ages of up to  
 225 approximately 100 years and mode-ages up to approximately 30 years, while the western central waters are younger with ages  
 226 of  $\sim 50$  years and  $\sim 15$  years respectively (Fig. 6). The relatively low ages in the upper layer indicate that these water masses  
 227 are newly formed.

228 In the meridional direction along the A16 section, both mean-ages and mode-ages show low values of around 20 years in the  
 229 mid-latitude regions that are close to the formation areas of central waters (Fig. 6 and Fig. 7). In contrast, the ages are higher  
 230 ( $\sim 100$  years for mean-age and  $\sim 30$  years for mode-ages) in the tropical region. In the zonal direction, the ages in the eastern  
 231 basin are slightly higher than in the western basin for the central waters, but the difference is only within  $\sim 10$  years for both  
 232 mean-ages and mode-ages, indicating that the influence from western boundary current is not significant in the upper layer  
 233 (Fig 7).



**Figure 6.** Ages of central water masses in the Atlantic Ocean.

- (a) The main currents in the upper layer with warm (red) and cold (blue) currents and the approximate formation areas (rectangular shadows) of central water masses.
- (b) The mean- and (c) mode-ages of central water masses at their core neutral densities ( $\text{kg m}^{-3}$ ). The colored dots show the ages in each station. The grey dots show all the GLODAPv2 stations that have less than 20% contribution of the water mass in question or lack of transient-tracer data.



**Figure 7.** The mean- and mode-ages of central waters along the three sections for the upper 1500 dbar. The solid green isolines show the 50% fractions of water masses and the dashed green lines show the 20 % fractions. The grey color indicate areas with less than 20% contribution from central waters. White vertical lines show cross overs with other sections.

### 234 3.2 The Intermediate Layer

235 Three main water masses belong to the intermediate layer ( $\gamma$  between 27.10 and 27.90 kg m<sup>-3</sup>), including the Antarctic  
236 Intermediate Water (AAIW), the Subarctic Intermediate Water (SAIW) and the Mediterranean Water (MW). The ages of SAIW  
237 are not displayed in this study due to the lack of tracer data with a complete section through the distribution region of this  
238 water mass.

239 **Antarctic Intermediate Water (AAIW):** The AAIW spreads from the formation area between 40 and 50 °S northward to  
240 approximately 30 °N along the Western Boundary Current (WBC) (Fig. 8, a). The northward flow of this water mass is part  
241 of the Atlantic western boundary current system (Talley, 1996) and the upper limb of AMOC (Kirchner et al., 2009).

242 Compared to the central waters, AAIW has significantly higher ages. In principle, the AAIW is supposed to get higher ages  
243 towards the north, as the transport time increases with the distance from the formation area in the south. In the meridional  
244 direction, the mean- and mode-ages increase from 0 to ~300 years and ~80 years from formation area to the equator, and  
245 further increase up to ~400 years and ~100 years respectively to 20 °N (Fig. 8 and 9). However, both ages decrease instead  
246 and approach even towards 0 in the further north region between 20 °N and 30 °N (Fig 8, b and c), giving the impression of  
247 the AAIW being younger in the north Atlantic Ocean. The observed decrease in ages likely results from mixing with younger  
248 surrounding water masses, although other factors such as limitations of the steady-state IG-TTD assumption (e.g., Waugh et  
249 al., 2004) or regional variations in the mixing ratio (e.g., Thomas et al., 2020) may also contribute. The maximum distance of  
250 AAIW to the north can reach 30 °N. However, the mixing is speculated between 20 °N and 30 °N, since the ENACW and  
251 upper NADW comes into contact with AAIW from the upper and lower (Liu and Tanhua, 2021) (Fig 7, a, Fig. 9, a and Fig.  
252 11, a). Both ENACW and upper NADW are newly formed in this region, so the “mixed” AAIW obtains younger ages. Age  
253 differences are also found in the zonal direction. In better ventilated western basins, the AAIW transports northward with the  
254 western boundary current (WBC), so the mean- and mode-ages are lower. By contrast, the ages are significantly higher in the  
255 eastern basins with poor ventilation. The mean-ages reach ~400 years and the mode-ages are up to ~100 years in the east  
256 between 0 and 20 °S (Fig. 8, b and c). The mean- and mode-ages along the A05 section also show the zonal difference (Fig. 9,  
257 b). In the eastern basin (east of the Mid-Atlantic-Ridge, MAR), the mean-age and mode-age appears up to ~400 and ~80 years  
258 respectively, in contrast only ~200 years and ~50 years in the west. This distribution confirms that the AAIW is transported  
259 by the WBC and takes a longer time to cross the MAR. Generally lower ages are found along the A10 Section in contrast to  
260 the A05 Section due to closer distance to the formation area of AAIW (Fig. 9, c). Similar zonal difference exists with mean-  
261 ages of ~50 years in the west ~100 years in the east and mode-age of ~10 years in the west and ~20 years in the east.


262 Both mean- and mode-ages of AAIW also vary in the vertical direction. These variations are consistent with the influence of  
263 mixing with adjacent water masses along its boundaries. Near the upper boundary, the AAIW is characterized by relatively  
264 lower ages, which is consistent with mixing with the generally younger eastern central waters above. Conversely, at the lower  
265 boundary with the deep and overflow layer, AAIW encounters the southward upper NADW. Along the A10 section, the upper

266 NADW exhibits high mean-ages ( $\sim 600$  years) and mode-ages ( $\sim 80$  years) due to the long southward transport. The resulting  
 267 composite water mass at this interface shows intermediate ages ( $\sim 300$  years mean-age,  $\sim 60$  years mode-age), higher than those  
 268 in the AAIW core ( $\sim 100$  years mean-age,  $\sim 20$  years mode-age) (Fig. 9 c).

269 A contrasting pattern is observed along the A05 section. In the northern Atlantic sampled by A05, the AAIW obtains higher  
 270 water mass ages, while the locally encountered upper NADW is relatively younger. Consequently, the ages at the interface  
 271 ( $\sim 200$  years mean-age,  $\sim 50$  years mode-age) are lower than those in the AAIW core at this location ( $\sim 300$  years mean-age,  
 272  $\sim 60$  years mode-age) (Fig. 9 b). These observed patterns at the boundaries are interpreted as reflecting the mixing of AAIW  
 273 with water masses of contrasting ages, and the resulting composite age signals are consistent with the independent age estimates  
 274 for NADW presented in the following section.

275 **Mediterranean Water (MW):** The MW is referred to as the product of the Mediterranean Overflow Water (MOW) that flows  
 276 across the Strait of Gibraltar and mixes with the ENACW (Liu and Tanhua, 2021). This water mass is formed in the Gulf of  
 277 Cadiz where the MOW exits the Strait of Gibraltar as a deep current and turns into two branches (Fig 8, a). The northward  
 278 branch spreads into the West European Basin until  $50^\circ\text{N}$ , while the westward branch spreads across the MAR into the west  
 279 basin of the North Atlantic Ocean (Price et al., 1993; Carracedo et al., 2016).


280 The spreading of MW can also be traced by the transient tracers (Fig 9, a), in addition to the water mass variables that goes  
 281 into the OMP analysis (Liu and Tanhua, 2021). The ages of MW have an increasing trend towards the south (Fig. 8, b and c).  
 282 The northward flow spreads faster so the mean-age is  $\sim 100$  years and the mode-age is  $\sim 20$  years at  $50^\circ\text{N}$ . In contrast, the  
 283 southward flow shows a higher mean-age of  $\sim 400$  years and mode-age of  $\sim 100$  years at  $25^\circ\text{N}$  along the A05 Section (Fig 9,  
 284 b). In the zonal direction, the ages decrease to the west (Fig 9, b), since the fraction of MW is only between 20% and 30%  
 285 along this section (green dash contour lines in Fig 9, b), and is therefore influenced by the lower ages from the NADW.



**Figure 8.** Ages of intermediate water masses in the Atlantic Ocean.

(a) Main currents in the intermediate layer. The currents (arrows) and the formation areas (rectangular shadows) of water masses in the intermediate layer.

(b) The mean- and (c) mode-ages of intermediate water masses at their core neutral densities ( $\text{kg m}^{-3}$ ). The colored dots show the ages in each station. The grey dots show all the GLODAPv2 stations that have less than 20% contribution of the water mass in question or lack of transient-tracer data.




**Figure 9.** Mean- and mode-ages of intermediate water masses along the three sections for the upper 4000 dbar. The solid green isolines show the 50% fractions of water masses and the dashed green lines show the 20 % fractions. The grey blank indicates less than 20% contribution from intermediate water mass. White vertical lines show cross overs with other sections.

### 286 3.3 Deep and Overflow Layer


287 The North Atlantic Deep Water (NADW), including its upper and lower portions, dominates the deep and overflow layer ( $\gamma$   
288 between 27.90 and 28.10  $\text{kg m}^{-3}$ ) of the Atlantic Ocean. This water mass is formed in the high latitude region in the North  
289 Atlantic and spreads southward at pressures between 2000 and 4000 dbar until it meets AAIW and AABW in the Antarctic  
290 Circum-polar Current (ACC) region. The southward spreading pathway is along the Deep Western Boundary Current (DWBC,  
291 Fig. 10, a). Meanwhile, the NADW also extends eastward with the eddies and finally covers the deep and overflow layer  
292 (Dickson and Brown, 1994).

293 **Upper and lower North Atlantic Deep Water (NADW):** The NADW is a water mass that is transported far southward from  
294 its formation area, with a mean-age of  $\sim 600$  years (Fig. 10, b) and mode-age of  $\sim 100$  years (Fig. 10, c) at the southern limb in  
295 the Antarctic Circum-polar Current (ACC) region. The increasing ages during the pathway to the south is shown along the A16  
296 section (Fig. 11, a). After leaving the formation area, the mean- and mode-ages increase  $\sim 200$  and  $\sim 20$  years respectively from  
297  $60^\circ\text{N}$  to  $40^\circ\text{N}$ . Afterwards, a sharply increase in both ages appear in the region between  $30^\circ\text{N}$  and  $10^\circ\text{N}$ . In this interval, the  
298 highest value of mean-age reaches  $\sim 1000$  years and the mode-age is up to  $\sim 150$  years. This is because the A16 section at this  
299 latitude range passed through the east part of the Atlantic Ocean (east of MAR) and the distance from the Deep Western  
300 Boundary Current (DWBC, Fig. 4 and Fig. 10, a) was high. High ages in this region are due to the sluggish water exchange  
301 over the ridge. This result can also be confirmed by the observation along the A05 Section (Fig. 11, b): The NADW near the  
302 DWBC region has a low mean-age of  $\sim 200$  years and mode-age of  $\sim 40$  years, while the ages are up to  $\sim 600$  years and  $\sim 100$   
303 years east of the MAR. After passing the equator, when the A16 section returns to the west of the MAR, the mean-ages of  
304 NADW are  $\sim 400$  years and the mode-ages are  $\sim 80$  years in the latitude range between  $0$  and  $20^\circ\text{S}$ , which further increase to  
305  $\sim 600$  (mean-ages) and  $\sim 100$  (mode-ages) years along the pathway southward until  $40^\circ\text{S}$ . In the ACC region between  $40^\circ\text{S}$   
306 and  $60^\circ\text{S}$ , the mean- and mode-ages of upper and lower NADW decrease to  $\sim 400$  and  $\sim 80$  years respectively. Similar to the  
307 situation of AAIW in the north Atlantic, this is also the result of mixing with newly formed AAIW and AABW. The spreading  
308 of NADW in the zonal direction is slower eastward, so the ages are generally higher in the eastern basin of the Atlantic Ocean  
309 (Fig. 10, b and c). The tracer data along the A05 and A10 sections also reflect the above result. In the A05 section, the mean-  
310 ages are  $\sim 200$  years in the west and  $\sim 600$  or even  $\sim 800$  years in the east, and the mode-ages show  $\sim 50$  years in the west and  
311  $\sim 100$  to  $\sim 120$  years in the east (Fig. 11, b). In the A10 section, the mean-ages are  $\sim 400$  years in the west and  $\sim 800$  years in the  
312 east and the mode-ages are  $\sim 80$  years in the west and  $\sim 120$  years in the east (Fig. 11, c).


313 In the vertical direction, the upper NADW mixes with the AAIW from above and the lower NADW mixes with the AABW  
314 from below. As the NADW is transported in the opposite direction to these two water masses, the “mixed” ages in the north,  
315 which is closer to the formation area of NADW, are higher than the age at the core density of NADW, while the opposite  
316 situation exists when spreading to south. After entering the ACC region, both mean-age and mode-age even decrease since the  
317 fractions of newly formed AAIW and AABW are high.



(a)



(b)




(c)

**Figure 10.** Ages of upper and lower NADW in the Atlantic Ocean.

(a) Main currents in the deep and overflow layer. The currents (arrows) and the formation areas (rectangular shadows) of water masses in the deep and overflow layer.

(b) The mean- and (c) mode-ages of upper and lower NADW at their core neutral densities ( $\text{kg m}^{-3}$ ). The colored dots show the ages in each station. The grey dots show all the GLODAPv2 stations that have less than 20% contribution of the water mass in question or lack of transient-tracer data.




**Figure 11.** Mean- and mode ages of upper and lower NADW along the A16 (a), A05 (b) and A10 (c) section lines. The green contour lines show fractions of water masses in 50 % and the dashed green lines show the 20 % fractions. The grey blank indicates less than 20% contribution from NADW. The white vertical lines show cross overs with other section lines.

### 320 3.4 Bottom Layer

321 The bottom layer is noticed already by Wüst, (1933) as it is filled with water from the south. Sverdrup, (1940) further pointed  
322 out the role of ACC plays in the formation and spreading of Antarctic Bottom Water (AABW). On this basis, Mantyla and  
323 Reid, (1983) elaborated that the most original dense bottom water is restricted near the Antarctic region by topography and the  
324 northward flow of bottom water is the mixture of original dense water and the overlying warm water. Gordon and Huber (1990)  
325 and Orsi et al., (1999) illustrated that the pathway of AABW to the north is through the Drake Passage sill into the Argentina  
326 and Brazilian Basin (Fig. 12, a). In the Atlantic Ocean, the AABW dominates the bottom layer ( $\gamma > 28.10 \text{ kg m}^{-3}$ ), which  
327 originates from the Weddell Sea and spreads to the north. After passing the equator, the AABW is redefined as Northeast Atlantic  
328 Bottom Water (NEABW, Liu and Tanhua, 2021).

329 **Antarctic Bottom Water (AABW) & Northeast Atlantic Bottom Water (NEABW):** The transport of bottom water is a long  
330 process going from the south towards the north. In the meridional direction, the AABW is transported towards the equator with  
331 a mean-age of ~600 years and mode-age of ~100 years at the equator, and further northward as NEABW to 40 °N with a mean-  
332 age of ~1000 years and mode-age of ~150 years (Fig. 12, b and c). Combined with the A16 Section, more specific  
333 segments/details can be seen. In the early stage from the surface of Weddell Sea to the bottom (below the pressure 4000 dbar)  
334 at 40 °S, the mean-age of AABW is ~300 years and mode-age is ~50 years. In the range between 10 and 30 °S, high values of  
335 ~800 years (mean-age) and ~120 years (mode-age) appear due to the mixing with the lower NADW with high ages (Fig. 13,  
336 a). After passing the equator, the redefined NEABW contains a mean-age up to ~1000 years and mode-age up to ~150 years  
337 at 40 °N, indicating that the bottom water, including the AABW and NEABW, takes about 150 years (mode-age) for the  
338 transport from the formation area (Weddell Sea) to 40 °N. North of 40 °N, the ages decrease due to the mixing with newly  
339 formed lower NADW in this region (Fig. 12, b and c, Fig. 13, a). The distribution of mean- and mode-ages also show a  
340 difference in the zonal direction. As shown in Fig. 12 b and c, the general age distribution presents a trend with lower ages in  
341 the west, and higher in the east. Along the A10 section at 30 °S, the AABW has a mean-age of ~400 in the west and ~600 years  
342 in the east, whereas the mode-age is ~50 in the west and ~100 in the east. A similar situation is recorded along the A05 section;  
343 the mean-age of NEABW is ~600 years in the west and ~800 years in the east, whereas the mode-age is ~80 years in the west  
344 and between ~120 years in the east (Fig. 13, b). An additional factor that affects the age distributions is the mixing between  
345 AABW and NADW. In the south hemisphere, both ages are relatively low (~400 years for mean-age and ~80 years for mode-  
346 age) at the pressures below 4000 dbar, where the AABW has a fraction higher than 50%, while the ages are significantly higher  
347 when the AABW mixes with lower NADW at pressure of ~3000 dbar, especially in the region between 0 and 10 °W, the mean-  
348 age reaches up to ~800 years and the mode-age is ~120 years. This is because the “old” NADW takes up a fraction for more  
349 than 50% to 80%. The situation in the north hemisphere is the opposite, the ages of NEABW are higher in the bottom layer  
350 below 4000 dbar but lower in the deep layer at 3000 dbar due to the mixing with the young NADW. In the western basin, the  
351 mean-age is ~600 years and mode-age is ~80 years in the bottom, while ~400 years and ~50 years when mixes with NADW.  
352 Similar in the east basin, the mean-age is ~800 years and mode-age is ~120 years below 4000 dbar, while ~600 years and ~100


353 years at 3000 dbar (Fig. 13, b).



**Figure 12.** Ages of bottom water masses in the Atlantic Ocean.

(a) Main currents in the bottom layer. The currents (arrows) and the formation areas (rectangular shadows) of water masses in the deep and overflow layer.

(b) The mean- and (c) mode-ages of bottom water masses at their core neutral densities ( $\text{kg m}^{-3}$ ). The colored dots show the ages in each station. The grey dots show all the GLODAPv2 stations that have less than 20% contribution of the water mass in question or lack of transient-tracer data.



**Figure 13.** Mean-ages of AABW and NEABW along the A16 (a), A05 and A10 (b) section lines.

The green contour lines show fractions of water masses in 50 and the dashed green lines show the 20 % fractions. The grey blank indicates less than 20% contribution from bottom water masses. The white vertical lines show cross overs with other section lines.

#### 354 4 The application of water mass ages in estimating the oxygen utilization rate (OUR)

355 The transit time distribution (TTD) provides a framework for using transient tracers to estimate the time scales governing

356 ventilation, which in turn can be applied to quantify biogeochemical rates. For instance, based on the TTD determined from

357 the tracer observations, Wang et al., (2021) estimated the OUR in the northern South China Sea.

358 The oxygen concentration in seawater under steady state is maintained by the balance of supply from oxygen -rich surface

359 water and consumption by the respiration of organic matter. The Apparent Oxygen Utilization (AOU) shows the accumulated

360 oxygen consumption in a water mass since isolated from the atmosphere, and is defined as the difference between saturated  
361 and measured oxygen concentration (i.e.  $\text{AOU} = \text{Oxy}_{\text{saturated}} - \text{Oxy}_{\text{measured}}$ , Redfield, 1942; Redfield et al., 1963; Pytkowicz,  
362 1971). The Oxygen Utilization Rate OUR ( $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ ) can be calculated from AOU and water mass age and shows the  
363 integrated oxygen utilization rate (oxidation rate) during the pathway of a water mass from surface of the formation area to the  
364 sampling location (Jenkins, 1987; Doney and Bullister, 1992; Sonnerup et al., 2013). In addition, the OUR is also an important  
365 indicator in characterizing the combination of ventilation and respiration of the interior ocean (Koeve and Kähler, 2016;  
366 Thomas et al., 2020).

367 The true, or local OUR is difficult to estimate, and the main difficulty focuses on the determination of the “true” water mass  
368 age since the seawater comes from different pathways due to the mixing in the ocean interior (Tomczak, 1999; Koeve and  
369 Kähler, 2016). Researchers have tried a variety of ways to define the age of a water mass. For instance, the tracer-age (CFC  
370 age) is used in Karstensen et al., (2008), while the mean-age (partial pressure ages) is used in Sonnerup et al., (2015) in  
371 estimating the “integrated” water mass age that effects the oxygen consumption. As described in Section 2.2, the mean-age  
372 represents the average of all the parts in one water sample, so the OUR calculated from mean-age is a “mixed” value of all the  
373 contributing water masses in the sample. In the steady state, the mean-age can reflect the “mixed” or “integrated” OUR  
374 objectively. The OUR is not constant over the life-time of a water mass, but is strongly depth and temperature dependent.  
375 Neither using the mode-age or the mean-age will be a reflection of the local OUR. The mean-age ( $\Gamma$ ) represents the average  
376 time elapsed since all water parcels in the sample were last in contact with the atmosphere, integrating over all pathways  
377 including long tails of older water, that most often has seen very low OUR in the interior ocean. In contrast, the mode -age,  
378 representing the most probable transit time, better captures the advective timescale of the dominant water mass along its  
379 dominant pathway from its formation region. The OUR calculated from mode-ages thus more closely approximates the time  
380 available for respiration to occur along the core of the water mass's trajectory. Using the mean-age, which includes much older  
381 water fractions will yield unrealistically low rates. We choose to use mode-age since it provides values in approximately the  
382 right magnitude, noting again that this value is not an absolute measure of local OUR.


383 Water masses with mode-ages larger than ~50 years in some parts of their distribution, including the AAIW, NADW, AABW  
384 and NEABW in the Atlantic Ocean, are defined here as wide-spread water masses. The transports of these wide-spread water  
385 masses are significantly influenced by the currents and topography. Therefore, their mode-age shows regional differences.  
386 Such differences also exist in DO and AOU, and lead to the spatial differences in OUR, especially between the eastern and  
387 western basins (Fig. 14). In the meridional direction, the mode-age increases with the distance from formation area. In the  
388 zonal direction, significant differences exist between the east and west of MAR. The wide-spread water masses generally have  
389 lower mode-ages in the western basin, which is better ventilated by the WBC and the DWBC (Fig. 14, left). In addition, surface  
390 oxygen-rich water is transported by these water masses to the deeper layers and are gradually consumed during the path-way.  
391 As a result, the DO is highest in the formation area and decreases with the distance. Meanwhile, the western basin displays  
392 higher DO due to the better ventilation (Fig. 14, middle left). The OUR shows a similar distribution to DO indicating that

393 higher oxygen consumption rate, or oxidation rate is co-located with the regions with high oxygen concentration. At the same  
394 time, Fig. 14 also shows that the OUR approaches 0 when the mode-age is larger than ~50 years.

395 In the intermediate layer, the AAIW show low mode-ages, high DO and low AOU in the formation area near the Antarctic  
396 region. During the northward transport between 40 °S and 20 °N, the mode-ages increase and the DOs decrease with distance  
397 from the formation area (Fig. 14, upper panel). In general, the mode-ages are lower in the west, while the zonal difference for  
398 DOs are not significant except for some regions in the eastern basins. The mode-ages are mostly between ~60 and ~90 years  
399 and reaches up to ~120 years in the poor ventilated eastern basin. The DOs are between 160 and 220  $\mu\text{mol kg}^{-1}$  in the south  
400 hemisphere and mostly between 100 and 160  $\mu\text{mol kg}^{-1}$  in the north hemisphere. The OURs in the Antarctic region obtain the  
401 highest value in the latitude region between 20 and 40 °S and decrease on the pathway to the north until 20 °N.

402 In the deep and overflow layer, the NADW is newly formed in the high north latitude. In most regions covered by the NADW  
403 between 30 °N and 40 °S, the differences between west and east exist with obvious low mode-age and AOU (high DO) and in  
404 the west, where is well ventilated by the DWBC (Fig. 14). The mode-age is between ~30 and ~60 years in the west and between  
405 ~90 and ~150 years in the east. The AOU is lower in the west and higher in the east. The OUR, which shows the opposite  
406 distribution as the AOU in the zonal direction, obtains the highest value in the formation area of NADW, maintains relative  
407 higher in the west along the DWBC region and shows a lower value in the eastern basin.

408 The bottom layer is occupied by the AABW and NEABW in the south and north hemisphere respectively. The AABW is  
409 formed in the Antarctic region with an original mode-age lower than ~30 years and increases on the way to the north. Similar  
410 to the AAIW and NADW, the western basin is better ventilated, therefore the mode-ages are lower in the west at the same  
411 latitude (Fig. 14, lower panel). In contrast to the AAIW and NADW, the AABW obtains relative lower DOs and higher AOU  
412 when formed (see also Fig. 15, b). In the western basin of the South Atlantic, the DOs decrease to 220  $\mu\text{mol kg}^{-1}$  and the AOU  
413 increase to 130  $\mu\text{mol kg}^{-1}$  on the pathway, while in the eastern basin, the DOs are 240  $\mu\text{mol kg}^{-1}$  and the AOU are 110  $\mu\text{mol}$   
414  $\text{kg}^{-1}$ . This is because the eastern basin is more influenced by the relative oxygen-rich NADW from the north. Similarly, the  
415 DOs and AOU of NEABW in the North Atlantic also show similar states to the NADW. In west regions ventilated by the  
416 DWBC, the DOs reach the values of higher than 260  $\mu\text{mol kg}^{-1}$  and AOU is lower than 70  $\mu\text{mol kg}^{-1}$ . This result can also be  
417 confirmed in Figure 15. Although the distributions of DOs and AOU are more complex in the bottom layer, the OURs still  
418 match the distribution of mode-ages. The highest OURs appears in the formation area near the Antarctic region with values  
419 higher than 4  $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ . In the western Atlantic region with low mode-ages, the OURs show relative higher values (>  
420 between 1 and 3  $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ ) and low value (below 1  $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ ) in the east (Fig. 14, lower panel).




**Figure 14.** Mode-age, DO, AOU and OUR of wide-spread water masses in the Atlantic Ocean.

The colored dots show the values in each station at core neutral densities of each water mass (AAIW at 27.20, upper NADW at 27.95, lower NADW at 28.05, AABW at 28.15, NEABW at 28.10  $\text{kg m}^{-3}$ , the upper and lower NADW, AABW and NEABW are shown in one plot. The grey dots show all the GLODAPv2 stations that have less than 20% contribution of the water mass in question. Note: Negative AOU values indicate oxygen supersaturation, which is typically observed in well-ventilated surface or near-surface waters.

421 The above four properties are also represented along the A16 section (Fig. 15). The AAIW has the lowest mode-age among  
 422 the three water masses, but also the largest variations in the DO and AOU. The mode-age indicates that in total of ~40 to 60  
 423 years is needed for the main body of AAIW (fractions > 80%) to be transported to the equator (Fig 15, a). The DO decreases  
 424 rapidly from 260 to below 180  $\mu\text{mol kg}^{-1}$  at the oxygen minimum zone at 20 °S, while the AOU increases from 60 to 150  $\mu\text{mol}$   
 425  $\text{kg}^{-1}$  during the northward pathway (Fig 15, b and c). The OUR is higher than 8  $\mu\text{mol kg}^{-1} \text{yr}^{-1}$  at the beginning and decreases  
 426 to 3  $\mu\text{mol kg}^{-1} \text{yr}^{-1}$  close to the equator (Fig 15, d). The mode-age of NADW is ~140 years at 40 °S and reaches a peak of ~160  
 427 years in the region of 20 °N due to the sluggish water exchange caused by the topography. The NADW contains the highest  
 428 oxygen concentration and the lowest AOU and OUR in the abyssal Atlantic Ocean. The value of DO is 260  $\mu\text{mol kg}^{-1}$  at the  
 429 starting point at 60 °N and still remains 220  $\mu\text{mol kg}^{-1}$  at 40 °S where it meets the AABW and AAIW. The AOU of NADW  
 430 remains below 100  $\mu\text{mol kg}^{-1}$  during the entire pathway. The OUR of NADW is below 5  $\mu\text{mol kg}^{-1} \text{yr}^{-1}$  close to the formation  
 431 area, and below 1  $\mu\text{mol kg}^{-1} \text{yr}^{-1}$  south of 60 °N. The transport of the main body of AABW takes in total of ~100 years from  
 432 the Weddell Sea to the equator. The DO and AOU remain unchanged at 220 and 130  $\mu\text{mol kg}^{-1}$  respectively on the pathway to  
 433 the equator, indicating very low OUR. Although derived from the AABW, the characteristics of the NEABW are more similar

434 to the NADW in all the above properties.




**Figure 15.** Distributions of (a) mode-age, (b) dissolved oxygen, (c) AOU and (d) OUR along the A16 section line. The color solid lines show the fractions of 80% in each water mass and the dashed lines show the fractions of 50%.

435

## 436 5 Estimation of water mass ages based on $^{39}\text{Ar}$

437 As a representative of noble gas radionuclides,  $^{39}\text{Ar}$  is considered to exhibit nearly ideal chemical and physical properties for  
 438 studying earth and environmental processes (Lu et al., 2014). In this study, the mean- and mode- ages of Atlantic wide-spread  
 439 water masses, including AAIW, NADW, AABW, and NEABW, are also estimated based on observations of  $^{39}\text{Ar}$ . For this  
 440 purpose, we make a distinction between the western (green dots) and eastern (red dots) Atlantic, divided by the Mid-Atlantic-  
 441 Ridge (Fig. 16a). The distribution of  $^{39}\text{Ar}$  is shown in Fig. 16 b. Near the formation area of each wide-spread water mass, the  
 442 concentration of  $^{39}\text{Ar}$  approaches approximately 100%, indicating that the surface waters are in equilibrium with the  
 443 atmosphere. As these water masses are transported over long distances and extended periods, the continuous decay of  $^{39}\text{Ar}$   
 444 leads to concentrations decreasing to roughly 30% at their furthest locations. By integrating this information with its half-life  
 445 of 269 years, the water mass ages can be estimated. For  $^{39}\text{Ar}$  we focus our attention to the older water masses, as the younger  
 446 ones are better characterized by CFC12 and  $\text{SF}_6$ .



**Figure 16.** (a) Map of the  $^{39}\text{Ar}$  sampling stations. The green circle dots show the stations in the west Atlantic Ocean (west of M-A-R), while the red stars show the stations in the east Atlantic Ocean. (b) Distributions of  $^{39}\text{Ar}$  in the Atlantic Ocean (% Modern). The circle dots show the data in the west, while the stars show the data in the east Atlantic Ocean. The solid isolines show the 80% fractions of water masses and the dashed lines show the 50% fractions.

447 Similar to the findings from CFC-12 and  $\text{SF}_6$  observations, the mean- and mode-ages derived from measurements using  $^{39}\text{Ar}$   
 448 increase with transport from the source of formation (Fig. 17). At a latitude of approximately 40°S, the AAIW exhibits mean-  
 449 ages ranging between ~100 to ~200 years (with mode-age around ~30 years). When transported northward to about 20°S, both  
 450 mean- and mode-ages rise to roughly ~300 years and ~60 years respectively. Upon reaching approximately 10°N during further  
 451 northward transport, these values escalate to around ~700 years for mean-age and ~100 years for mode-age. However, near  
 452 the region at about 20°N, the terminus for northward flow, the mean-age decreases back down to roughly ~400 years while  
 453 maintaining a mode-age close to ~60 years due to mixing with younger upper NADW. In the deep and overflow layer, the  
 454 newly formed NADW initiates at high northern latitudes, exhibiting both mean- and mode-ages approaching zero. During

455 southward transport at around 30 °S, the concentration data indicates corresponding mean- and mode-ages of approximately  
 456 ~400 years and ~60 years respectively. Similarly, the bottom water masses (AABW and NEABW) display mean- and mode-  
 457 ages of around ~500 years and ~100 years respectively at the farthest end of the transport distance approximately 50 °N.




Figure 17. Mean- (a) and mode-ages (b) of the wide-spread water masses estimated by  $^{39}\text{Ar}$  in the Atlantic Ocean. The circle dots show the data in the west, while the stars show the data in the east Atlantic Ocean. The solid isolines show the 80% fractions of water masses and the dashed lines show the 50 % fractions.

458 The contrast between the western and eastern Atlantic Ocean is distinctly evident (Fig. 16, b). Generally, higher concentrations  
 459 of  $^{39}\text{Ar}$  are observed in the west compared to the east at equivalent latitudes. For instance, in the vicinity of the AAIW formation  
 460 area at 40 °S, the  $^{39}\text{Ar}$  concentration is approximately 100% in the west and around 70% in the east. In the intermediate layer  
 461 near the equator, these concentrations decrease to about 70% (west) and 50% (east). These findings suggest that both mean-  
 462 and mode-ages of AAIW are close to zero at 40 °S. The mean-age is estimated to be ~300 years in the west and ~500 years in  
 463 the east north of 40°S. Similar results are also observed in the deep water and overflow layers. Based on elevated levels of  
 464  $^{39}\text{Ar}$  near its formation area, it can be inferred that both mean- and mode-ages of NADW are close to zero at 60 °N across both  
 465 West and East Atlantic Oceans. Between 20 °N and the equator, the mean-age is estimated to be ~300 years in the west and  
 466 ~500 years in the east; while the mode-age is around ~60 years (west) and ~90 years (east), respectively. For bottom water  
 467 masses below a depth of 4000 m, concentrations of  $^{39}\text{Ar}$  generally fall below 50%, with higher levels observed in the west  
 468 (50%-60%) compared to lower levels seen in the east (40%-50%) between 20 °S and 40 °N, indicating the water mass ages  
 469 range from ~400–500 years (mean-age) and ~60–90 years (mode-age) in the west, and from ~500–600 years (mean-age) and  
 470 ~90–120 years (mode-age) in the east.

471

| Water Masses | Age Type (years) | Tracer                   | 40°S / 40°N                   | 20°S / 20°N     | Destination                   |
|--------------|------------------|--------------------------|-------------------------------|-----------------|-------------------------------|
| AAIW         | Mean-age         | CFC-12 & SF <sub>6</sub> | ~70 (West) / ~200(East)       | ~300            | ~400 (maximun) / ~100 (mixed) |
|              |                  | <sup>39</sup> Ar         | ~100-200 (West) / ~150 (East) | ~300            | ~700 (East) / ~400 (mixed)    |
|              | Mode-age         | CFC-12 & SF <sub>6</sub> | ~10 (West) / ~30 (East)       | ~50             | ~100 (maximun) / ~10 (mixed)  |
|              |                  | <sup>39</sup> Ar         | ~30 (West) / ~35 (East)       | ~60             | ~100 (East) / ~60 (mixed)     |
| NADW         | Mean-age         | CFC-12 & SF <sub>6</sub> | ~200 (A05)                    | ~400 (A10)      | ~600 (ACC, 50 °S)             |
|              |                  | <sup>39</sup> Ar         | ~130                          | ~200            | ~230                          |
|              | Mode-age         | CFC-12 & SF <sub>6</sub> | ~40 (A05)                     | ~80 (A10)       | ~100 (ACC, 50 °S)             |
|              |                  | <sup>39</sup> Ar         | ~30-50                        | ~60-80          | ~60–90                        |
| AABW & NEABW | Mean-age         | CFC-12 & SF <sub>6</sub> | ~400 (A10)                    | ~600 (A05)      | ~800 (East, A05)              |
|              |                  | <sup>39</sup> Ar         | ~500–600 (East)               | ~500–600 (East) | ~500–600 (East)               |
|              | Mode-age         | CFC-12 & SF <sub>6</sub> | ~50 (A10)                     | ~80 (A05)       | ~120 (East, A05)              |
|              |                  | <sup>39</sup> Ar         | ~90–120 (East)                | ~90–120 (East)  | ~90–120 (East)                |

473

474

475

As expected, the inferred ages of water masses differ depending on the tracer employed. These discrepancies arise not only from potential limitations in the assumed IG-TTD, but also from the distinct temporal validity ranges and detection limits of each tracer. Our comparative analysis (as summarized in Table 2) reveals a systematic pattern: mean- and mode-ages derived

476 from  $^{39}\text{Ar}$  are generally lower than those estimated from CFC-12 and  $\text{SF}_6$  for deeper/older water masses. This is exemplified  
477 by the NADW in the eastern basin near  $20^\circ\text{N}$ , where CFC-12/ $\text{SF}_6$  yield a mean-age of  $\sim 1000$  years and a mode-age of  $\sim 150$   
478 years, whereas  $^{39}\text{Ar}$  suggests younger ages of  $\sim 600$  and  $\sim 90$  years, respectively. Similarly, for AABW/NEABW near  $50^\circ\text{N}$ ,  
479 CFC-12 and  $\text{SF}_6$  based estimates ( $\sim 800$  years mean,  $\sim 120$  years mode) exceed those from  $^{39}\text{Ar}$  ( $\sim 500$  years mean,  $\sim 100$  years  
480 mode). A likely explanation for this systematic tendency towards higher age estimates from CFC-12 and  $\text{SF}_6$  in deep and  
481 bottom layers might be attributed to their extremely low concentrations at great depths and for older waters. These  
482 concentrations often approach or fall below analytical detection limits, which can amplify uncertainties and appear to skew  
483 age calculations towards higher values. In contrast, the  $^{39}\text{Ar}$  with 269-year half-life appears to be better suited for dating waters  
484 with transit times on the order of centuries to a millennium, and thus could provide relative more robust age estimates for these  
485 older water masses, as its slower decay rate remains measurable over such longer timescales.

## 486 **6 Conclusions and discussion**

487 The geographic distributions of main water masses in the Atlantic Ocean are reported in Liu and Tanhua (2021). In the present  
488 work, the transports of water masses are traced by transient tracers (CFC-12,  $\text{SF}_6$  and  $^{39}\text{Ar}$ ) and their ventilation timescales  
489 (water mass ages) are quantified. Three distinct concepts of water mass age are discussed within the analytical framework.  
490 While the tracer-age concept (assuming purely advective transport) is introduced for context, the quantitative analysis and  
491 results presented herein are based on the mean-age and mode-age derived from the Transit Time Distribution (TTD)  
492 model. The application of a standard mixing ratio ( $\Delta/\Gamma = 1$ ) and 100% surface saturation is followed. The mean-age,  
493 representing the average transit time of all water parcels in a sample and providing an integrative timescale, is used to  
494 characterize the overall ventilation status. The mode-age, representing the most probable transit time and serving as a proxy  
495 for advective timescales, is applied to trace the dominant transport pathways and to estimate process rates like the oxygen  
496 utilization rate (OUR).

497 In general, the water mass age increases with the depth and distance from the formation area, so the central waters in the upper  
498 layer obtain the lowest ages of within  $\sim 100$  years. In the intermediate layer, the AAIW and MW are two conspicuous water  
499 masses and both take  $\sim 300$  years to spread from south to north (AAIW) and east to west (MW) respectively. As the dominant  
500 water mass in the deep and overflow layer, the NADW takes  $\sim 500$  years to spread along the DWBC in the west Atlantic Ocean,  
501 and takes  $\sim 900$  years eastward to cover the east part. The bottom waters origin from south of Antarctic Circumpolar Current  
502 region take  $\sim 600$  years to spread northward and cover the bottom layer. Similar as the NADW, the mean-age of both AABW  
503 and NEABW show spatial difference between the west (low mean-age) and the east (high mean-age). This is because the west  
504 Atlantic Ocean is well ventilated by the DWBC while the water exchange is sluggish in the east.

505 The transient tracers (CFC-12,  $\text{SF}_6$  and  $^{39}\text{Ar}$ ), serve as effective tools for identifying and quantifying oceanographic ventilation,  
506 each tracer possessing distinct advantages and limitations in determining and tracing the depths and mean ages of water masses.  
507 CFC-12 and  $\text{SF}_6$  exhibit optimal applicability for water masses range within 2000 dbar pressure (or mean-ages  $< \sim 500$  years).

508 However, their utility becomes constrained in deeper aqueous regimes (below 2000 dbar or mean-age > ~500 years) due to  
509 inherent limitations in tracer concentration sensitivity, which may induce systematic overestimations of water mass ages when  
510 applied to deep or bottom layers. The chemically inert radioisotope  $^{39}\text{Ar}$ , characterized by a 269-year half-life, effectively  
511 addresses this observational gap through its unique temporal resolution. Consequently, the synergistic application of  $\text{SF}_6$ , CFCs,  
512 and  $^{39}\text{Ar}$  in multi-tracer frameworks overcomes temporal constraints inherent to single-tracer approaches, enabling  
513 comprehensive investigation of coupled biogeochemical-climate interactions. This integrated methodology facilitates novel  
514 insights into climate-driven water mass reorganization, carbon cycle dynamics, and other critical marine processes essential  
515 for understanding oceanic responses to global change.

516 The ventilations and respiration rates (oxidation rates, OUR) of wide-spread water masses are estimated by the mode-ages.  
517 The currents and topography have significant impacts on the mode-ages and furthermore the the apparent oxygen utilisation  
518 rate (OUR). The western basins of the Atlantic Ocean are better ventilated with lower the mode-ages, while the OURs show  
519 the opposite distribution, suggesting that the mode-age is an important determinant of OUR in seawater than AOU and higher  
520 respiration rates exist when the region is better ventilated.

521 The quantitative age framework and ventilation rates established in this study provide a crucial baseline dataset. This baseline  
522 is essential for detecting future changes in ocean circulation and biogeochemical cycles, and will serve to validate the next  
523 generation of climate and ocean models.

524 This study provides a quantitative, basin-scale assessment of Atlantic water mass ages using a multi-tracer approach. However,  
525 several methodological assumptions should be considered when interpreting the results. First, the calculated ages rely on the  
526 assumption of a steady-state circulation and an Inverse Gaussian Transit Time Distribution (IG-TTD) with a fixed mixing ratio  
527 ( $\Delta/\Gamma = 1$ ). While this is a necessary and common simplification for basin-scale analysis, regional deviations from this  
528 idealized transport are likely. Second, the complementary use of tracers (CFC-12,  $\text{SF}_6$ ,  $^{39}\text{Ar}$ ) mitigates but does not fully  
529 eliminate the inherent uncertainties associated with each method, particularly in deep layers with very low tracer concentrations.  
530 Finally, the apparent oxygen utilization rates (OUR) derived here represent integrated rates along flow paths and are subject  
531 to the same age estimation uncertainties.

532 Notwithstanding these limitations, the comprehensive age and ventilation maps presented here establish a baseline of the  
533 contemporary Atlantic Ocean ventilation structure. This dataset can serve to validate and constrain the next generation of ocean  
534 circulation and biogeochemical models. Furthermore, by quantifying current ventilation timescales and their spatial patterns,  
535 this work provides a reference frame against which future observational studies can detect changes in ocean circulation and  
536 ventilation in response to climatic forcing.

537 **Author contributions.** ML designed the research study and conducted the analysis. ML and TT interpreted the results and  
538 wrote the paper.

539 **Competing interests.** The contact author has declared that none of the authors has any competing interests.

## 540 Acknowledgements

541 This work is based on the comprehensive and detailed data from the GLODAP data product throughout the past few decades.  
542 In particular, we are grateful to the efforts from all the scientists and crews on cruises, who generated funding and dedicated  
543 time on committing the collection of data. We also would like to thank the working groups of GLODAP for their support and  
544 information of the collation, quality control and publishing of data. Their contributions and selfless sharing are prerequisites  
545 for the completion of this work. Thank all the colleagues from the School of Environmental Science and Engineering, Xiamen  
546 University of Technology for their support in my research work. In addition, we are grateful to Dr. Tim Stöven for his support  
547 and advices in running calculating programs of water mass ages. Thanks also goes to Prof. Minggang Cai and his research  
548 group at the College of Ocean and Earth Sciences, Xiamen University for the help and support during Mian Liu's postdoctoral  
549 work.

550 **Financial support.** The authors gratefully thank the Research Projects for Overseas Scholars in Xiamen City (No.  
551 0108/50103250009); Research Start-up Project for the Introduction (Cultivation) of High-level Talents of Xiamen University  
552 of Technology (No. YKJ23021R); National Natural Science Foundation of China (No. 22106128) and the Xiamen University  
553 of Technology National Natural Science Foundation of China Training Programme for Excellence (No. XPYM2409). Mian  
554 Liu received support from the China Scholarship Council (CSC) to support the PhD study at GEOMAR Helmholtz Centre for  
555 Ocean Research Kiel, Germany.

## 556 References

- 557 Armour, K. C., J. Marshall, J. R. Scott, A. Donohoe, and E. R. Newsom: Southern Ocean warming delayed by circumpolar  
558 upwelling and equatorward transport, *Nature Geoscience*, 9(7), 549-554, doi:10.1038/ngeo2731, 2016.
- 559 Bender, M. L.: The  $\delta^{18}\text{O}$  of dissolved  $\text{O}_2$  in seawater: A unique tracer of circulation and respiration in the deep sea, *Journal of*  
560 *Geophysical Research: Oceans*, 95(C12), 22243-22252, doi:https://doi.org/10.1029/JC095iC12p22243, 1990.
- 561 Bjerknes, J.: Atlantic Air-Sea Interaction, in *Advances in Geophysics*, edited by H. E. Landsberg and J. Van Mieghem, pp. 1-  
562 82, Elsevier, doi:https://doi.org/10.1016/S0065-2687(08)60005-9, 1964.
- 563 Bryden, H. L., H. R. Longworth, and S. A. Cunningham: Slowing of the Atlantic meridional overturning circulation at 25  
564 degrees N, *Nature*, 438(7068), 655-657, doi:10.1038/nature04385, 2005.
- 565 Bullister, J. L., and R. F. Weiss: Anthropogenic Chlorofluoromethanes in the Greenland and Norwegian Seas, *Science*,  
566 221(4607), 265-268, doi:10.1126/science.221.4607.265, 1983.
- 567 Bullister, J. L., D. P. Wisegarver, and F. A. Menzia: The solubility of sulfur hexafluoride in water and seawater, *Deep Sea*  
568 *Research Part I: Oceanographic Research Papers*, 49(1), 175-187, doi:10.1016/s0967-0637(01)00051-6, 2002.
- 569 Clark, P. U., J. D. Shakun, P. A. Baker, P. J. Bartlein, S. Brewer, E. Brook, A. E. Carlson, H. Cheng, D. S. Kaufman, and Z.  
570 Liu: Global climate evolution during the last deglaciation, *Proceedings of the National Academy of Sciences*, 109(19), E1134-  
571 E1142, doi:10.1073/pnas.1116619109, 2012.
- 572 Dickson, R. R., and J. Brown: The production of North Atlantic Deep Water: Sources, rates, and pathways, *Journal of*  
573 *Geophysical Research: Oceans*, 99(C6), 12319-12341, doi:10.1029/94jc00530, 1994.
- 574 Doney, S. C., and J. L. Bullister: A chlorofluorocarbon section in the eastern North Atlantic, *Deep-Sea Res*, 39(11-12A), 1857-  
575 1883, doi:10.1016/0198-0149(92)90003-c, 1992.
- 576 Fine, R. A.: Observations of CFCs and  $\text{SF}_6$  as Ocean Tracers, in *Annual Review of Marine Science*, Vol 3, edited by C. A.  
577 Carlson and S. J. Giovannoni, pp. 173-195, doi:10.1146/annurev.marine.010908.163933, 2011.
- 578 Gammon, R. H., J. Cline, and D. Wisegarver: Chlorofluoromethanes in the northeast Pacific Ocean: Measured vertical  
579 distributions and application as transient tracers of upper ocean mixing, *Journal of Geophysical Research-Oceans*, 87(NC12),  
580 9441-9454, doi:10.1029/JC087iC12p09441, 1982.
- 581 Hall, M. M., and H. L. Bryden: Direct Estimates and Mechanisms of Ocean Heat-Transport, *Deep-Sea Res*, 29(3), 339-359,  
582 doi:10.1016/0198-0149(82)90099-1, 1982.
- 583 Hall, T. M., and R. A. Plumb: Age as a diagnostic of stratospheric transport, *Journal of Geophysical Research-Atmospheres*,  
584 99(D1), 1059-1070, doi:10.1029/93jd03192, 1994.

585 Huhn, O., M. Rhein, M. Hoppema, and S. van Heuven: Decline of deep and bottom water ventilation and slowing down of  
 586 anthropogenic carbon storage in the Weddell Sea, 1984-2011, *Deep Sea Research Part I: Oceanographic Research Papers*, 76,  
 587 66-84, doi:10.1016/j.dsr.2013.01.005, 2013.

588 Ito, T., A. Nenes, M. S. Johnson, N. Meskhidze, and C. Deutsch: Acceleration of oxygen decline in the tropical Pacific over  
 589 the past decades by aerosol pollutants, *Nature Geoscience*, 9(6), 443-447, doi:10.1038/ngeo2717, 2016.

590 Jenkins, W. J.: Oxygen utilization rates in North Atlantic subtropical gyre and primary production in oligotrophic systems,  
 591 *Nature*, 300(5889), 246-248, doi:10.1038/300246a0, 1982.

592 Jenkins, W. J.:  $^3\text{H}$  and  $^3\text{He}$  in the Beta Triangle: Observations of Gyre Ventilation and Oxygen Utilization Rates, *Journal of*  
 593 *Physical Oceanography*, 17(6), 763-783, doi:10.1175/1520-0485(1987)017<0763:Aitbto>2.0.Co;2, 1987.

594 Jia, Z.-H., P. Zhang, L.-B. Li, Q.-W. Chen, J.-L. Liu, Y.-G. Liu, Q. Wu, Y. Yang, L.-T. Sun G.-M. Yang, W. Jiang, Z.-T. Lu:  
 595 Stability and reliability study of an  $^{39}\text{Ar}$  enrichment system for accurate  $^{39}\text{Ar}$  dating, *Nuclear Instruments and Methods in*  
 596 *Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 1075, 170438,  
 597 doi:https://doi.org/10.1016/j.nima.2025.170438, 2025.

598 Karstensen, J., and M. Tomczak, Ventilation processes and water mass ages in the thermocline of the southeast Indian Ocean,  
 599 *Geophysical Research Letters*, 24(22), 2777-2780, doi:10.1029/97gl02708, 1997.

600 Karstensen, J., and M. Tomczak: Agedetermination of mixed water masses using CFC and oxygen data, *Journal of Geophysical*  
 601 *Research-Oceans*, 103(C9), 18599-18609, doi:10.1029/98jc00889, 1998.

602 Karstensen, J., L. Stramma, and M. Visbeck: Oxygen minimum zones in the eastern tropical Atlantic and Pacific oceans,  
 603 *Progress in Oceanography*, 77(4), 331-350, doi: 10.1016/j.pocean.2007.05.009, 2008.

604 Kirchner, K., M. Rhein, S. Huttel-Kabus, and C. W. Böning: On the spreading of South Atlantic Water into the Northern  
 605 Hemisphere, *Journal of Geophysical Research-Oceans*, 114, doi:10.1029/2008jc005165, 2009.

606 Koeve, W., and P. Kähler: Oxygen utilization rate (OUR) underestimates ocean respiration: A model study, *Global*  
 607 *Biogeochemical Cycles*, 30(8), 1166-1182, doi:10.1002/2015gb005354, 2016.

608 Kuhlbrodt, T., A. Griesel, M. Montoya, A. Levermann, M. Hofmann, and S. Rahmstorf: On the driving processes of the Atlantic  
 609 meridional overturning circulation, *Reviews of Geophysics*, 45(1), doi:10.1029/2004rg000166, 2007.

610 Lauvset, S. K., R. M. Key, A. Olsen, S. van Heuven, A. Velo, X. Lin, C. Schirnack, A. Kozyr, T. Tanhua, M. Hoppema, S.  
 611 Jutterström, R. Steinfeldt, E. Jeansson, M. Ishii, F. F. Pérez, T. Suzuki and S. Watelet. A new global interior ocean mapped  
 612 climatology: the 1°x1° GLODAP version 2, *Earth Syst. Sci. Data*, 8, 325–340, https://doi:10.5194/essd-8-325-2016, 2016.

613 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Brown, P. J., Carter,  
 614 B. R., Cotrim da Cunha, L., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson, E., Murata, A., Müller, J. D., Pérez, F. F.,

615 Schirnick, C., Steinfeldt, R., Suzuki, T., Ulfsbo, A., Velo, A., Woosley, R. J., and Key, R. M.: The annual update  
616 GLODAPv2.2023: the global interior ocean biogeochemical data product, *Earth Syst. Sci. Data*, 16, 2047–2072,  
617 <https://doi.org/10.5194/essd-16-2047-2024>, 2024.

618 Li, Q., England, M. H., Hogg, A. M., Rintoul, S. R., & Morrison, A. K., Abyssal ocean overturning slowdown and warming  
619 driven by Antarctic meltwater. *Nature*, 615, (7954), 841–847, 2023.

620 Liu, M., and T. Tanhua: Water masses in the Atlantic Ocean: characteristics and distributions, *Ocean Science*, 17(2), 463–486,  
621 [doi:10.5194/os-17-463-2021](https://doi.org/10.5194/os-17-463-2021), 2021.

622 Maiss, M., L. P. Steele, R. J. Francey, P. J. Fraser, R. L. Langenfelds, N. B. A. Trivett, and I. Levin: Sulfur hexafluoride - A  
623 powerful new atmospheric tracer, *Atmospheric Environment*, 30(10–11), 1621–1629, [doi:10.1016/1352-2310\(95\)00425-4](https://doi.org/10.1016/1352-2310(95)00425-4),  
624 1996.

625 Morrison, A. K., T. L. Frolicher, and J. L. Sarmiento: Upwelling in the Southern Ocean, *Physics Today*, 68(1), 27–32,  
626 [doi:10.1063/pt.3.2654](https://doi.org/10.1063/pt.3.2654), 2015.

627 Olsen, A., R. M. Key, S. van Heuven, S. K. Lauvset, A. Velo, X. Lin, C. Schirnick, A. Kozyr, T. Tanhua, M. Hoppema, S.  
628 Jutterström, R. Steinfeldt, E. Jeansson, M. Ishii, F. F. Pérez and T. Suzuki. The Global Ocean Data Analysis Project version 2  
629 (GLODAPv2) – an internally consistent data product for the world ocean, *Earth Syst. Sci. Data*, 8, 297–  
630 323, <https://doi.org/10.5194/essd-8-297-2016>, 2016.

631 Orsi, A. H., G. C. Johnson, and J. L. Bullister: Circulation, mixing, and production of Antarctic Bottom Water, *Progress in*  
632 *Oceanography*, 43(1), 55–109, [doi:10.1016/s0079-6611\(99\)00004-X](https://doi.org/10.1016/s0079-6611(99)00004-X), 1999.

633 Patara, L., C. W. Böning, and T. Tanhua: Multidecadal Changes in Southern Ocean Ventilation since the 1960s Driven by Wind  
634 and Buoyancy Forcing, *Journal of Climate*, 34(4), 1485–1502, [doi:10.1175/Jcli-D-19-0947.1](https://doi.org/10.1175/Jcli-D-19-0947.1), 2021.

635 Purkey, S. G., and G. C. Johnson: Warming of Global Abyssal and Deep Southern Ocean Waters between the 1990s and 2000s:  
636 Contributions to Global Heat and Sea Level Rise Budgets, *Journal of Climate*, 23(23), 6336–6351, [doi:10.1175/2010jcli3682.1](https://doi.org/10.1175/2010jcli3682.1),  
637 2010.

638 Pytkowicz, R. M.: On the apparent oxygen utilization and the preformed phosphate in the oceans<sup>1</sup>, *Limnology and*  
639 *Oceanography*, 16(1), 39–42, <https://doi.org/10.4319/lo.1971.16.1.0039>, 1971.

640 Rahmstorf, S.: The Great Ocean Conveyor: Discovering the Trigger for Abrupt Climate Change, *Nature*, 464(7289), 681–681,  
641 [doi:10.1038/464681a](https://doi.org/10.1038/464681a), 2010.

642 Redfield, A. C.: The processes determining the concentration of oxygen, phosphate and other organic derivatives within the  
643 depths of the Atlantic Ocean, *Pap. Phys. Oceanogr. Meteorol. Mass. Inst. Technol. Woods Hole Oceanogr. Inst.*, 9, 1–22,  
644 <https://doi.org/10.1575/1912/1053>, 1942.

645 Redfield, A. C., B. H. Ketchum, F. A. Richards, The influence of organisms on the composition of seawater, *The Sea. Ideas*  
646 *and Observations on Progress in the Study of the Seas* M. N. Hill, 26–77, John Wiley, New York, 1963.

647 Sandström, J. W.: Dynamische versuche mit meerwasser. *Annalen der Hydrographie und Maritimen Meteorologie*, 36, 6–23,  
648 1908.

649 Sandström, J. W.: Meteorologische studien im Schwedischen Hochgebirge. Göteborgs Kungl. Vetenskaps- och  
650 Vitterhetssamhälles Handlingar, 17, 1–48, 1916.

651 Schmidt, S., L. Stramma, and M. Visbeck: Decline in global oceanic oxygen content during the past five decades, *Nature*,  
652 542(7641), 335–339, doi:10.1038/nature21399, 2017.

653 Schneider, A., T. Tanhua, A. Kortzinger, and D. W. R. Wallace: High anthropogenic carbon content in the eastern Mediterranean,  
654 *Journal of Geophysical Research-Oceans*, 115, doi:10.1029/2010jc006171, 2010.

655 Skinner, L. C., F. Primeau, E. Freeman, M. de la Fuente, P. A. Goodwin, J. Gottschalk, E. Huang, I. N. McCave, T. L. Noble,  
656 and A. E. Scrivner: Radiocarbon constraints on the glacial ocean circulation and its impact on atmospheric CO<sub>2</sub>, *Nature*  
657 *Communications*, 8, doi:10.1038/ncomms16010, 2017.

658 Sonnerup, R. E., Quay, P. D., and Bullister, J. L.: Thermocline ventilation and oxygen utilization rates in the subtropical North  
659 Pacific based on CFC-11 and CFC-12 measurements from 1991 to 2000, *J. Geophys. Res.*, 106, 9259–  
660 9287, <https://doi.org/10.1029/1999JC000086>, 2001.

661 Sonnerup, R. E., S. Mecking, and J. L. Bullister: Transit time distributions and oxygen utilization rates in the Northeast Pacific  
662 Ocean from chlorofluorocarbons and sulfur hexafluoride, *Deep-Sea Res Pt I*, 72, 61–71, doi:10.1016/j.dsr.2012.10.013, 2013.

663 Sonnerup, R. E., S. Mecking, J. L. Bullister, and M. J. Warner: Transit time distributions and oxygen utilization rates from  
664 chlorofluorocarbons and sulfur hexafluoride in the Southeast Pacific Ocean, *Journal of Geophysical Research-Oceans*, 120(5),  
665 3761–3776, doi:10.1002/2015jc010781, 2015.

666 Stanley, R. H. R., S. C. Doney, W. J. Jenkins, and D. E. Lott: Apparent oxygen utilization rates calculated from tritium and  
667 helium-3 profiles at the Bermuda Atlantic Time-series Study site, *Biogeosciences*, 9(6), 1969–1983, doi:10.5194/bg-9-1969-  
668 2012, 2012.

669 Stöven, T., and T. Tanhua: Ventilation of the Mediterranean Sea constrained by multiple transient tracer measurements, *Ocean*  
670 *Science*, 10(3), 439–457, doi:10.5194/os-10-439-2014, 2014.

671 Sverdrup, H. U.: The unity of the sciences of the sea, *Sigma Xi Quarterly*, 28(3), 105–115,  
672 <https://www.jstor.org/stable/23049117>, 1940.

673 Talley, L.: Antarctic Intermediate Water in the South Atlantic, in *The South Atlantic: Present and Past Circulation*, edited by  
674 G. Wefer, W. H. Berger, G. Siedler and D. J. Webb, pp. 219–238, Springer Berlin Heidelberg, Berlin, Heidelberg,

doi:10.1007/978-3-642-80353-6\_11, 1996.

Talley, L.: Closure of the Global Overturning Circulation Through the Indian, Pacific, and Southern Oceans: Schematics and Transports, *Oceanography*, 26(1), 80-97, doi:10.5670/oceanog.2013.07, 2013.

Talley, L., R. Feely, B. Sloyan, R. Wanninkhof, M. Baringer, J. Bullister, C. Carlson, S. Doney, R. Fine, and E. Firing, N. Gruber, D.A. Hansell, M. Ishii, G.C. Johnson, K. Katsumata, R.M. Key, M. Kramp, C. Langdon, A.M. Macdonald, J.T. Mathis, E.L. McDonagh, S. Mecking, F.J. Millero, C.W. Mordy, T. Nakano, C.L. Sabine, W.M. Smethie, J.H. Swift, T. Tanhua, A.M. Thurnherr, M.J. Warner, and J.-Z. Zhang: Changes in ocean heat, carbon content, and ventilation: A review of the first decade of GO-SHIP global repeat hydrography, *Annual review of marine science*, 8, 185-215, <https://doi.org/10.1146/annurev-marine-052915-100829>, 2016.

Tamsitt, V., H. F. Drake, A. K. Morrison, L. D. Talley, C. O. Dufour, A. R. Gray, S. M. Griffies, M. R. Mazloff, J. L. Sarmiento, J. Wang, W. Weijer: Spiraling pathways of global deep waters to the surface of the Southern Ocean, *Nature Communications*, 8(1), 172, doi:10.1038/s41467-017-00197-0, 2017.

Tanhua, T., A. Biastoch, A. Kortzinger, H. Luger, C. Boning, and D. W. R. Wallace: Changes of anthropogenic CO<sub>2</sub> and CFCs in the North Atlantic between 1981 and 2004, *Global Biogeochemical Cycles*, 20(4), doi:10.1029/2006gb002695, 2006.

Tanhua, T., D. W. Waugh, and D. W. R. Wallace: Use of SF<sub>6</sub> to estimate anthropogenic CO<sub>2</sub> in the upper ocean, *Journal of Geophysical Research-Oceans*, 113(C4), doi:10.1029/2007jc004416, 2008.

Tanhua, T., D. W. Waugh, and J. L. Bullister: Estimating changes in ocean ventilation from early 1990s CFC-12 and late 2000s SF<sub>6</sub> measurements, *Geophysical Research Letters*, 40(5), 927-932, doi:10.1002/grl.50251, 2013.

Tanhua, T., and M. Liu: Upwelling velocity and ventilation in the Mauritanian upwelling system estimated by CFC-12 and SF<sub>6</sub> observations, *Journal of Marine Systems*, 151, 57-70, doi:10.1016/j.jmarsys.2015.07.002, 2015.

Thiele, G., and J. L. Sarmiento: Tracer Dating and Ocean Ventilation, *Journal of Geophysical Research-Oceans*, 95(C6), 9377-9391, doi:10.1029/JC095iC06p09377, 1990.

Thomas, J. L., D. W. Waugh, and A. Gnanadesikan: Relationship between Age and Oxygen along Line W in the Northwest Atlantic Ocean, *Ocean Science Journal*, 55(2), 203-217, doi:10.1007/s12601-020-0019-5, 2020.

Tomczak, M.: Some historical, theoretical and applied aspects of quantitative water mass analysis, *Journal of Marine Research*, 57(2), 275-303, doi:10.1357/002224099321618227, 1999.

Tseitlin, V.: Depth dependence of oxygen utilization rate, *OKEANOLOGIYA*, 32(2), 264-269, 1992.

van Heuven, S., M. Hoppema, O. Huhn, H. A. Slagter, and H. J. W. de Baar: Direct observation of increasing CO<sub>2</sub> in the Weddell Gyre along the Prime Meridian during 1973-2008, *Deep-Sea Research Part II-Topical Studies in Oceanography*,

704 58(25-26), 2613-2635, doi:10.1016/j.dsr2.2011.08.007, 2011.

705 Wang, W. M., M. G. Cai, P. Huang, H. W. Ke, M. Liu, L. H. Liu, H. X. Deng, B. J. Luo, C. H. Wang, X. H. Zheng, W. Q. Li:  
 706 Transit Time Distributions and Apparent Oxygen Utilization Rates in Northern South China Sea Using Chlorofluorocarbons  
 707 and Sulfur Hexafluoride Data, *Journal of Geophysical Research-Oceans*, 126(8), doi:10.1029/2021jc017535, 2021.

708 Warner, M. J., and R. F. Weiss: Solubility of Chlorofluorocarbon-11 and Chlorofluorocarbon-12 in water and seawater, *Deep-*  
 709 *Sea Res*, 32(12), 1485-1497, doi:10.1016/0198-0149(85)90099-8, 1985.

710 Waugh, D. W., T. M. Hall, and T. W. N. Haine: Relationships among tracer ages, *Journal of Geophysical Research: Oceans*,  
 711 108(C5), doi:10.1029/2002jc001325, 2003.

712 Waugh, D. W., T. W. N. Haine, and T. M. Hall: Transport times and anthropogenic carbon in the subpolar North Atlantic Ocean,  
 713 *Deep Sea Research Part I: Oceanographic Research Papers*, 51(11), 1475-1491, doi:10.1016/s0967-0637(04)00145-1, 2004.

714 Ziska, F., B. Quack, K. Abrahamsson, S. D. Archer, E. Atlas, T. Bell, J. H. Butler, L. J. Carpenter, C. E. Jones, N. R. P. Harris,  
 715 H. Hepach, K. G. Heumann, C. Hughes, J. Kuss, K. Krüger, P. Liss, R. M. Moore, A. Orlikowska, S. Raimund, C. E. Reeves,  
 716 W. Reifenhäuser, A. D. Robinson, C. Schall, T. Tanhua, S. Tegtmeier, S. Turner, L. Wang, D. Wallace, J. Williams, H.  
 717 Yamamoto, S. Yvon-Lewis, and Y. Yokouchi: Global sea-to-air flux climatology for bromoform, dibromomethane and methyl  
 718 iodide, *Atmospheric Chemistry and Physics*, 13(17), 8915-8934, doi:10.5194/acp-13-8915-2013, 2013.