

Supporting Information for

Technical note: Quantified organic aerosol subsaturated hygroscopicity by a simple optical scatter monitor system through field measurements

Jie Zhang¹, Tianyu Zhu^{1,2}, Alexandra Catena¹, Yaowei Li³, Margaret J. Schwab¹, Pengfei Liu⁴, Akua A. Asa-Awuku⁵, James Schwab¹

¹Atmospheric Sciences Research Center, University at Albany, State University of New York, Albany, NY, 12226, United States

²Department of Atmospheric and Environmental Sciences, University at Albany, State University of New York, Albany, NY, 12226, United States

³School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, 02138, United States

⁴School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, 30332, United States

⁵Department of Chemical and Biomolecular Engineering, A. James Clark School of Engineering, University of Maryland, College Park, MD, 20742, United States

Correspondence to: Jie Zhang (jzhang35@albany.edu)

Contents of this file

Figures S1 to S6

Introduction

Supporting Information includes additional figures that show the self-calibration of two pDRs under low RH conditions (Fig. S1), the representative back trajectories for each subperiod in Fig. 4 of the three groups with different aerosol sources (Fig. S2-S7), and the relationship between κ_{OA} and O:C of each subperiod of Group1(urban) and Group2(rural) (Fig. S8).




Figure S1. The head-to-head-correlation scatterplot between the two pDRs under the lowest relative humidity conditions reported by PDR_{wet} (RH < 45%)




Figure S2. The representative back trajectories of each subperiod (P1, P3, P5, and P9) of Group1(urban) with aerosol having urban sources




Figure S3. The representative back trajectories of each subperiod (P2, P6, P10, P11) of Group2(rural) with aerosol having rural sources




Figure S4. The representative back trajectory on July 19 of P4 of Group3(wildfire) when aerosol affected by the wildfire plumes, with the PM_{2.5} AQI distribution map on July 19 verifying the wildfire plumes influence and on the previous date (July 16) showing the place of original wildfire and its transport.




Figure S5. The representative back trajectory on July 26 of P7 of Group3(wildfire) when aerosol affected by the wildfire plumes, with the PM_{2.5} AQI distribution map on July 26 verifying the wildfire plumes influence and on the previous date (July 24) showing the place of original wildfire and its transport.




Figure S6. The representative back trajectory on Aug. 05 of P8 of Group3(wildfire) with aerosol affected by the wildfire plumes, with the PM_{2.5} AQI distribution map on Aug. 05 verifying the wildfire plumes influence and on the previous date (Aug. 03) showing the place of original wildfire and its transport.




Figure S7. The representative back trajectory on Sep. 04 of P12 of Group3(wildfire) group with aerosol affected by the wildfire plumes, with the PM_{2.5} AQI distribution map on Sep. 04 verifying the wildfire plumes influence and on the previous date (Sep. 04) showing the place of original wildfire and its transport.




Figure S8. (a) The relationship between bin-averaged κ_{OA} and O:C for Group1(urban) with the urban aerosol sources (in grey squares with their standard deviation as error bar), and the relationship between κ_{OA} and O:C of each subperiod of Group1(urban) with a time resolution of 1-hr. (b) The relationship between bin-averaged κ_{OA} and O:C for Group2(rural) with the rural aerosol sources (in green squares with their standard deviation as error bar), and the relationship between κ_{OA} and O:C of each subperiod of Group2(rural) with a time resolution of 1-hr.