

1 **Seasonal and interannual variability on the chemical composition of the**
2 **Svalbard surface snowpack**
3 Azzurra Spagnesi^{a,b}, Elena Barbaro^{a,b} *, Matteo Feltracco^{a,b}, Federico Scoto^{c,b}, Marco Vecchiato^b,
4 Massimiliano Vardè^a, Mauro Mazzola^a, François Burgay^{d,e}, Federica Bruschi^f, Clara Jule Marie Hoppe^g,
5 Allison Bailey^g, Andrea Gambaro^{a,b}, Carlo Barbante^{a,b}, Andrea Spolaor^{a,b}

6 ^a Institute of Polar Sciences - National Research Council of Italy (ISP-CNR), Via Torino 155, 30172, Venice, Italy
7 ^b Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Via Torino 155,
8 30172, Venice, Italy
9 ^c Institute of Atmospheric Sciences and Climate - National Research Council of Italy (ISAC-CNR), Campus Ecotekne,
10 Lecce, 73100, Italy
11 ^d Laboratory of Environmental Chemistry (LUC), Paul Scherrer Institut (PSI), Villigen, 5232, Switzerland
12 ^e Oeschger Centre for Climate Change Research, University of Bern, Bern, 3012, Switzerland
13 ^f Department of Chemistry, Biology and Biotechnology, University of Perugia, Via dell'Elce di Sotto 8, 06123, Perugia,
14 Italy
15 ^g Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, 27570 Bremerhaven, Germany

18 Corresponding: Elena Barbaro (elena.barbaro@cnr.it)

Definizione stile: Normale: Tipo di carattere: (Predefinito)
Calibri, Inglese (Regno Unito)

Definizione stile: Titolo 1: Tipo di carattere: (Predefinito)
Calibri, Colore carattere: Colore
personalizzato(RGB(47;84;150)), Inglese (Regno Unito)

Definizione stile: Titolo 4: Tipo di carattere: (Predefinito)
Calibri, Colore carattere: Colore
personalizzato(RGB(47;84;150)), Inglese (Regno Unito)

Definizione stile: Paragrafo elenco: Tipo di carattere:
(Predefinito) Calibri, Inglese (Regno Unito)

Definizione stile: Piè di pagina: Tipo di carattere:
(Predefinito) Calibri, Inglese (Regno Unito)

Definizione stile: Testo commento: Tipo di carattere:
(Predefinito) Calibri, Inglese (Regno Unito)

Definizione stile: Normale (Web)

Definizione stile: Default: Tipo di carattere: Legature:
Nessuna

Definizione stile: Revisione: Tipo di carattere: (Predefinito)
Calibri

Definizione stile: Intestazione: Tipo di carattere:
(Predefinito) Calibri, Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

Formattato: SpazioPrima: 0 pt, Dopo: 14 pt

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: elena.barbaro@cnr.it

ha eliminato: 1 ¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo),
Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A
destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni:
8,5 cm, Allineato al centro + 17 cm, Allineato a destra

32 **Abstract**

33 Arctic Amplification (AA) is driving long-term changes in the Arctic climate system, including
34 glacier ice melt, rapid sea ice decline, and alterations in atmospheric and geochemical processes, with
35 consequences on the formation, transport, and chemical composition of aerosols and seasonal
36 snowpack. Svalbard, particularly vulnerable to AA, provides a valuable site to assess changes in local
37 environmental processes contributing to the seasonal snow chemical composition. From 2018 to
38 2021, sampling campaigns at the Gruebadet Snow Research Site in Ny-Ålesund, in the North-West
39 of the Svalbard Archipelago, captured the interannual variability in ionic and elemental impurities
40 within surface snow, reflecting seasonal differences in atmospheric and oceanic conditions. Notably,
41 warmer conditions prevailed in 2018-19 and 2020-21, contrasting with the relatively colder season
42 of 2019-20. Our findings suggest that impurity concentrations in the 2019-20 colder season are
43 impacted by enhanced sea spray aerosol production, likely driven by a larger extent of sea ice, and
44 drier, windy conditions. This phenomenon was particularly evident in March 2020, when extensive
45 sea ice was present in Kongsfjorden and around Spitsbergen due to an exceptionally strong, cold
46 stratospheric polar vortex and unusual Arctic Oscillation (AO) index positive phase. By comparing
47 the snow chemical composition of the 2019-20 season with 2018-19 and 2020-21, we provide insights
48 into the interplay between short-term meteorological variability and the long-term climatic impacts
49 of AA in Svalbard, as well as associated shifts in aerosol production process.

50 **1. Introduction**

51 Chemical analysis of surface Arctic snow and ice can provide valuable comprehension of the
52 composition of Arctic aerosols, its deposition, and exchange processes (Lai et al., 2017). These
53 dynamics are influenced by a range of factors, including Arctic Amplification (AA) – a pronounced,
54 long-term increase in near-surface air temperature observed since 1975 (Chylek et al., 2022). AA is
55 recognized as an inherent characteristic of the global climate system, with multiple intertwined causes
56 operating on a spectrum of spatial and temporal scales. These include, but are not limited to, changes
57 in sea ice extent that impact heat fluxes between the ocean and the atmosphere, and water vapour that
58 alters longwave radiation (Serreze and Barry, 2011). The Svalbard Archipelago is particularly
59 sensitive to these effects due to the relatively low altitude of its main ice fields and its geographical
60 location in the higher North Atlantic, where the impact of AA is especially pronounced (Spolaor et
61 al., 2024). Therefore, in the 21st century, predicting and characterising climate change in Svalbard is
62 particularly crucial, as changes in near-surface air temperature, precipitation, and sea ice extent occur
63 at an extremely high pace (Gjermundsen et al., 2020; Rantanen et al., 2022). The Svalbard region,

ha formattato: Tipo di carattere: Times New Roman, 12 pt, Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: .

ha formattato: Inglese (Regno Unito)

ha eliminato: processes

ha formattato: Inglese (Regno Unito)

ha eliminato: vapor

ha formattato: Inglese (Regno Unito)

ha eliminato: characterizing

ha formattato: Inglese (Regno Unito)

ha eliminato: 2

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

68 located at the southern edge of the seasonal Arctic sea-ice zone, is ~~characterised~~ by a maritime climate
69 with strong temperature variations during winter (Hansen et al., 2014; Barbaro et al., 2021). In the
70 Arctic winter, the stratospheric polar jet fosters a high-atmospheric vorticity zone. This winter vortex
71 typically acts as a strong barrier for the long-range transport of pollutants from mid-latitudes
72 (Lawrence et al., 2020). However, it occasionally allows warm southern air to penetrate the region
73 (Schoeberl and Newman, 2015). Additionally, Svalbard frequently experiences intense cyclonic
74 storms in autumn and winter, which bring both heat and moisture from lower latitudes (Rinke et al.,
75 2017). These intense meteorological variations, generally linked with a weaker polar vortex (Sobota
76 et al., 2020; Salzano et al., 2023), ~~favour~~ long-range transport of aerosols to the archipelago, including
77 pollutants from continental sources (Stohl et al., 2006b; Yttri et al., 2014a; Vecchiato et al., 2024;
78 D'Amico et al., 2024).

79 Arctic snow captures dry and wet deposition and forms an archive that includes a range of seasonal
80 chemical species such as major ions and trace elements, as well as human-made pollutants emitted
81 into the Arctic atmosphere (Koziol et al., 2021). Ny-Ålesund is a well-monitored area and a natural
82 laboratory for complex system observations, ideal for exploring both long-range contaminants from
83 mid- to high-latitude regions of Eurasia and Canada (Nawrot et al., 2016; Song et al., 2020; Vecchiato
84 et al., 2024; D'Amico et al., 2024), and local inputs from both natural processes and human settlement
85 (Vecchiato et al., 2018). Previous research has extensively investigated the chemistry of Arctic snow
86 and the exchange of inorganic species between the cryosphere and the atmosphere across multiple
87 sites, including Barrow, Summit Greenland, Alert, Sodankylä, and over the Arctic Ocean during the
88 MOSAiC expedition (e.g., Beine et al., 2003; Björkman et al., 2013; Jacobi et al., 2019). Specific
89 studies in Ny-Ålesund and surrounding areas have explored the temporal and compositional aspects
90 of the lower atmosphere (Stohl et al., 2006a; Eleftheriadis et al., 2009; Geng et al., 2010; Zhan et al.,
91 2014; Feltracco et al., 2020, 2021; Turetta et al., 2021), though relatively few have addressed the
92 detailed seasonal dynamics of snow-atmosphere interactions in this region. Building on this existing
93 research, our study aims to enhance the understanding of these interactions, particularly in the context
94 of recent climatic changes.

95 In this study, we evaluate the surface snow concentration of ionic (Cl^- , Br^- , NO_3^- , SO_4^{2-} , MSA, Na^+ ,
96 NH_4^+ , K^+ , Ca^{2+}) and elemental impurities (Li, Be, Mg, Al, Ca, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Se,
97 Rb, Sr, Ag, Cd, Sb, Cs, Ba, Tl, Pb, Bi, U) for the snow seasons between 2018-2021, at the Gruevbadet
98 Snow Research Site (GSRS) location, 1 km south of Ny-Ålesund, where clean and undisturbed snow
99 conditions are guaranteed throughout the whole sampling season. The differences in average
100 meteorological and climatological conditions across the studied seasons are analysed to assess how

ha eliminato: characterized

ha formattato: Inglese (Regno Unito)

ha eliminato:).

ha formattato: Inglese (Regno Unito)

ha eliminato:).

ha formattato: Inglese (Regno Unito)

ha eliminato: favor

ha formattato: Inglese (Regno Unito)

ha eliminato:).

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: ↵

ha eliminato: 3¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

107 sea ice extent, polar vortex, and Arctic Oscillation (AO) conditions influence the composition of
108 surface snow in connection with the aerosol-producing and deposition processes in Kongsfjorden.
109 Additionally, a Kruskal-Wallis's test, followed by Dunn's post-hoc analysis, has been employed to
110 identify species most affected by inter-annual and seasonal variations, with the goal of uncovering
111 potential correlations between meteorological-climatic conditions and chemical concentrations.

112 2. Methodology

113 2.1 Sampling and processing

114 Three sampling campaigns were conducted in Svalbard between 2018 and 2021, covering the period
115 from October to May according to the onset of the snowpack formation and melting. Sample
116 collection followed the protocol presented in Spolaor et al. (2019), further adopted by Bertò et al.
117 (2021), where consecutive and adjacent sampling was carried in a 3x3 meters snow sampling area,
118 within a clean sampling snowfield of 100 m wide. Each sample was collected 10 cm apart from the
119 previous one, along a precise path. This method was designed to minimise the temporal variability
120 between consecutive samples and reduce the impact of potential spatial variability (within the 5-15%
121 range, according to Spolaor et al., 2019).

122 During the first sampling campaign, carried out from October 4th, 2018 to May 10th, 2019, 133 snow
123 samples were collected at the Gruvebadet Snow Research Site (GSRS), a clean-area located about 1
124 km south of Ny-Ålesund, nearby the Gruvebadet Atmospheric Laboratory (GAL), dedicated to the
125 chemical and physical monitoring of the seasonal snowpack (Scoto et al., 2023; Fig. 1).

126 The surface snow was sampled within the upper 3 cm, as this is the snow layer most impacted by
127 aerosol deposition and exchange processes at the snow-atmosphere interface (Spolaor et al., 2018,
128 2021b). This choice also minimised the effect of different physical snow conditions (density, crystal
129 shape, and size).

130 Throughout the season, the sampling resolution varied based on light conditions. During the polar
131 night (from October to early March), snow sampling was carried out every 3-5 days. With the
132 beginning of the polar day, daily sampling was conducted until the end of the snow season.

133 The second campaign was carried out from October 26th, 2019 to May 25th, 2020, with a total of 107
134 samples collected. Consecutive samples represent the same snow layer in the absence of snowfall or
135 wind drift/erosion. However, factors such as snow aging, potential element re-emission,
136 transformation, and dry deposition can introduce variability, making continuous monitoring essential.
137 Finally, during the third snow sampling campaign, lasting from October 27th, 2020 to June 15th, 2021,
138 a weekly sampling was conducted at GSRS, with a total of 32 samples collected.

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 0 pt, Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:2 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Inglese (Regno Unito)

ha eliminato: 114 surface

ha spostato (inserimento) [1]

ha eliminato: in a delimited snow field located ~ 100 m south of

ha spostato (inserimento) [2]

ha eliminato: "Dirigibile Italia Station" in Ny-Ålesund (78.92° N 11.93° E, Ny-Ålesund, Svalbard).

ha formattato: Inglese (Regno Unito)

ha spostato in alto [1]: a clean-area located about 1 km south of Ny-Ålesund, nearby the Gruvebadet Atmospheric Laboratory (GAL), dedicated to the chemical and physical monitoring of the seasonal snowpack (Scoto et al., 2023; Fig. 1).

ha spostato in alto [2]: 1).

ha eliminato: Concurrently, additional 133 snow samples were collected at the Gruvebadet Snow Research Site (GSRS) to evaluate the spatial variability with respect to the snow samples collected in Ny-Ålesund. The GSRS is

ha eliminato: daily at Ny-Ålesund, and

ha eliminato: at the GSRS

ha eliminato: both in Ny-Ålesund and at the GSRS in March, and then continued only at the GSRS

ha eliminato: in June due to the lower contamination of the site, more distant from the fervent local activities. This sampling resolution overlap during March ensured a good comparison of results in both snow fields (Fig. S1).

ha eliminato: Starting from the

ha eliminato: , snow sampling activities were conducted only at the GSRS site, since clean conditions of the field in Ny-Ålesund could not be guaranteed due to construction works. The snow sampling

ha eliminato: The surface snow layer was sampled every 3-5 days during the polar night (until February 24th, 2020), and daily from the beginning of the polar day until the end of the snow season. ...

ha eliminato: 4

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

171 During snow sampling, the temperature and density of surface snow were measured, and the density
172 of snow was calculated based on weighting a 100 cc cylinder. After collection, snow samples were
173 melted, and two different aliquots were obtained and stored in separate vials. In a 1.5 mL
174 polypropylene (PP) vial, 1 mL of sample was stored for ionic species, while another aliquot was
175 stored in a 5 mL LDPE vials for trace elements analysis. PP vials designated to ionic species analysis
176 were previously sonicated for 30 min in UltraPure Water (UPW) (18 MΩ cm⁻¹ at 25 °C) for
177 decontamination. LDPE vial used for trace elements analysis were instead conditioned with HNO₃
178 2% and sonicated for 30 min. All sample aliquots were stored at -20°C in dark conditions and
179 transported to the Venice ISP-CNR laboratories.

180 To assess potential contamination during sampling, handling, and transport, field blanks were
181 collected during each campaign. Metal-free vials (Avantor, VWR Centrifuge Tubes, CHN) were
182 opened to ambient air at the sampling sites for a few minutes without collecting snow, then sealed
183 and transported to the Ny-Ålesund laboratory. There, they were filled with 2% HNO₃ and stored under
184 the same conditions as the snow samples. In parallel, analytical blanks were prepared by opening
185 vials to air, sealing them, and transporting them directly to Venice, where they were filled with 2%
186 HNO₃ and ultrapure water from the laboratory. Both field and analytical blanks were analysed
187 alongside the snow samples, confirming that background contamination levels were consistently
188 below LODs or one order of magnitude lower than the lowest concentration detected for all analytes.

189

190 Furthermore, seawater temperatures and salinity at 10 m depth were also monitored in Kongsfjorden
191 (Kb3; 78°57.228'N, 11°57.192'E) during 2019-2021 spring seasons, with data collected every 3-6
192 days (Assmy et al. 2023). Data was derived from Conductivity Temperature Depth (CTD) casts with
193 either a MiniSTD model SD-204 (SAIV A/S, Bergen, Norway) or a XR-620 CTD (RBR Ltd, Ottawa,
194 Canada). Combined casts of both instruments conducted in May 2020 and 2021 did not reveal
195 differences in temperature or salinity in the reported accuracy (two post comma digits).

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: analyzed

ha eliminato: below detection limits

ha eliminato: target

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: 5¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra




Fig. 1. Gruvebadet Snow Research Site (GSRS) location with respect to Ny-Ålesund Research Station, in the Brogger peninsula (Svalbard Islands).

2.2 Analysis of ionic species

The analysis of anionic species (Cl^- , Br^- , NO_3^- , SO_4^{2-} , MSA) was carried out using an ion chromatograph (IC, Thermo Scientific Dionex™ ICS-5000, Waltham, MA, USA) coupled with a single quadrupole mass spectrometer (MS, MSQ Plus™, Thermo Scientific, Bremen, Germany). The separation was performed using an anionic exchange column (Dionex Ion Pac AS 19 2 mm ID $\times$ 250 mm length) equipped with a guard column (Dionex Ion Pac AG19 2 mm ID $\times$ 50 mm length). Sodium hydroxide (NaOH), used as mobile phase, was produced by an eluent generator (Dionex ICS 5000EG, Thermo Scientific). The NaOH gradient with a 0.25 mL min⁻¹ flow rate was: 0–6 min at 15 mM; 6–15 min gradient from 15 to 45 mM; 15–23 min column cleaning with 45 mM; 23–33 min equilibration at 15 mM. The injection volume was 100 μL . A suppressor (ASRS 500, 2 mm, Thermo Scientific) removed NaOH before entering the MS source. The IC-MS operated with a negative electrospray source (ESI) with a temperature of 500°C and a needle voltage of 3 kV. The other MS parameters are reported by Barbaro et al. (2017). The same IC system was simultaneously used to determine cationic species (Na^+ , K^+ , Ca^{2+} and NH_4^+). However, Ca^{2+} was not measured within the samples collected during the second campaign due to instrumental limitations.

The separation occurred with a capillary cation-exchange column (Dionex Ion Pac CS19–4 mm 0.4 mm ID $\times$ 250 mm length), equipped with a guard column (Dionex Ion Pac CG19–4, 0.4 mm ID $\times$ 50 mm length), and the species were determined using a conductivity detector. Analytical blanks of ultrapure water ($> 18 \text{ M}\Omega \text{ cm}$) were included in the analysis, and the Method Detection Limit (MDL) was set to 3 times the standard deviation of the blank values. Checks for accuracy were made using

ha eliminato:

ha eliminato: Sampling sites of

ha eliminato: and Gruvebadet, located

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: 6

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

225 certified multi-element standard solutions for anions (Cl^- , Br^- , NO_3^- , SO_4^{2-} , no. 89886-50ML-F, Sigma
226 Aldrich) and cations (Na^+ , K^+ , Ca^{2+} , no. 89316-50ML-F, Sigma Aldrich) at a concentration of 10 mg
227 $\text{L}^{-1} \pm 0.2\%$. The analytical precision was quantified as the relative standard deviation (RSD) for
228 replicates ($n > 3$) of standard solutions and was always $< 10\%$ for each ion.

229 *2.3 Trace Elements analysis*

230 Twenty-six elements (Li, Be, Mg, Al, Ca, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Se, Rb, Sr, Ag, Cd, Sb,
231 Cs, Ba, Tl, Pb, Bi and U) were analysed on samples previously melted and acidified to 2% v/v with
232 HNO_3 (UpA grade, Romil, UK) for 24 hours before analysis (Spolaor et al., 2018; Spolaor et al.,
233 2021a).

234 The analysis was performed using Inductively Coupled Plasma Mass Spectrometry (ICP-MS, iCAP
235 RQ, Thermo Scientific, US). The ICP-MS was equipped with an ASX-560 autosampler (Teledyne
236 Cetac Technologies), PolyPro PFE nebulizer, PFE cyclonic spray chamber thermostated at 2.7°C ,
237 sapphire injector, quartz torch and Ni cones. The acquisition was performed at 1550 W of plasma RF
238 power in Kinetic Energy Discrimination (KED) – high matrix mode, using He as collision gas (4.3
239 mL min^{-1}). Instrument parameters were optimised for best sensitivity in the whole mass range,
240 minimum oxides ($< 1\%$) and double charges ($< 3\%$). Quantification was obtained by external
241 calibration with multi-elemental standards prepared in ultrapure water ($18 \text{ M}\Omega \text{ cm}^{-1}$ at 25°C) with
242 2% v/v ultrapure grade HNO_3 (UpA grade, Romil, UK), with a combination of certified level multi-
243 elemental solutions IMS-102 and IMS-104 from UltraScientific. Analytical quality control was
244 performed by memory test blank (repeated analysis of ultrapure grade HNO_3 2% v/v blank solution)
245 after each sample and calibration verification (repeated analysis of reference standards) every 11
246 samples. More details are found in Spolaor et al., 2021a.

248 *2.4 Transport modelling, sea ice, Kongsfjorden condition, and polar vortex*

250 The Lagrangian particle dispersion model HYSPLIT (Draxler, 1998; Stein et al., 2015) was used to
251 determine the source region of air masses over Ny-Ålesund. This model has previously been shown
252 to be an effective tool for the prediction of transport pathways into and within the Arctic and Antarctic
253 regions (Barbaro et al., 2015; Feltracco et al., 2021). The simulations were driven using
254 meteorological data from the Global Data Assimilation System (GDAS) one-degree archive, set the
255 top of the model at 10000 m and the height source equal to the GSRS altitude. Back-trajectories were
256 calculated every 6 h, with a propagation time of 120 h for each sampling period. The choice of a 6-
257 hour interval for the calculation of back-trajectories allows for the capture of temporal variability in

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Inglese (Regno Unito)

ha eliminato: analyzed

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato:).

ha formattato: Inglese (Regno Unito)

ha eliminato: optimized

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Nero

Formattato: Normale, Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato:).

ha eliminato: 7

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

262 air mass origins over the day, which is particularly important in polar regions where atmospheric
263 circulation patterns can change rapidly. This temporal resolution strikes a balance between
264 computational efficiency and the need for sufficient detail to characterise the variability in source
265 regions during each sampling period. The propagation time of 120 hours was selected to provide an
266 adequate temporal window to trace long-range transport pathways that influence air mass
267 composition at Ny-Ålesund. This configuration is consistent with previous studies on atmospheric
268 circulation in the same site (Feltracco et al., 2021).

269 This approach was used to ensure an envelope working for all investigated tracers. The resulting
270 multiple trajectories were based on the screen-plot analyses of total spatial variance.

271 The Ice Service provided by the Norwegian Meteorological Institute (NIS) was employed to analyse
272 the weather conditions via remotely sensed data and to generate ice charts of Svalbard, ice-edge
273 information, and sea surface temperatures trends. Sea ice extent variability in Kongsfjorden was
274 evaluated based on dataset made available by Gerland et al. (2022).

275 Differences between the sampling campaigns were evaluated through the National Centers for
276 Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) Reanalysis
277 data from NOAA Physical Sciences Lab's daily composites tool, used to calculate the near-surface
278 air temperatures across the Northern Hemisphere from October to May.

279

280 2.5 *Statistical procedures and Enrichment Factors (EFs) Analysis*

281

282 Results below the limit of detection were assumed to be equal to ½ of Method Determination Limit
283 (MDL) prior to perform statistical analysis, to approximate their likely level based on the data
284 distribution curve (best approximated as log-normal for most of the studied variables) (George et al.,
285 2021).

286 Enrichment Factors (EFs) using Ba as a crustal element of reference (Widepohl, 1995; Spolaor et al.,
287 2021a; Ruppel et al., 2023), were calculated to explore the mixed sources of the investigated elements.

288 To complement the EFs analysis, a Hierarchical Cluster Analysis (HCA) was performed using Ward's
289 algorithm and Euclidean distances as clustering criteria, to determine the presence of some clusters
290 and simplify the interpretation of the dataset.

291 Additionally, to examine the inter-annual and seasonal variations in chemical's concentrations and to
292 identify the species most affected by these changes, a Kruskal-Wallis' test was employed, followed
293 by Dunn's post-hoc test for pairwise comparisons. The Kruskal-Wallis' test, a nonparametric
294 alternative to Anova, which is typically applied when samples do not follow a normal distribution
295 (Van Hecke, 2013), was chosen to assess differences in chemical concentrations across the years of

ha formattato: Colore carattere: Nero

Formattato: Normale, Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: The Wilcoxon rank-sum test was applied on data from the 2018–19 sampling campaign to assess whether there was a statistically significant difference in the surface snow contamination levels (ionic loads) between the two sites: Ny-Ålesund and Gruvebadet. This model assumes that the data is sampled from two matched or dependent populations, and data is assumed to be continuous. Because it is a nonparametric test, it does not require a particular probability distribution of the dependent variable in the analysis. Furthermore,

ha eliminato: 8¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

sampling campaigns and multiple seasons. This allowed for an objective evaluation of temporal differences in concentration levels, while Dunn’s post-hoc analysis, which uses the same shared rankings and pooled variance assumed under the Kruskal-Wallis null hypothesis, helped identify which specific seasons showed statistically significant differences from each other. The Bonferroni correction was used for the Dunn’s test to control for false positives when performing multiple pairwise comparisons (Dunn, 1961). The ultimate goal of this analysis was to uncover potential correlations between meteorological-climatic conditions and chemical concentrations in surface snow, helping to better understand how environmental factors contribute to contamination trends. These combined statistical approaches provided a comprehensive framework for analysing the complex dataset.

3. Results

3.1 Interannual trends of chemical species on the surface snow

Three consecutive snow seasons were evaluated to define the chemical composition of the surface snow in the Arctic site of GSRs. The sea salt ions Cl⁻ (50 %), and Na⁺ (23%) represent the most abundant species, followed by SO₄²⁻ (11 %), Mg (7 %), Ca (2%), Fe (1%) and Al (1%). Similar relative concentration abundances were also found in previous studies on the snow of the Svalbard Archipelago (Beaudon and Moore, 2009; Vega et al., 2015; Barbaro et al., 2021; Spolaor et al., 2021b). Table 1 reports the average ionic loads of the most abundant (> 1%) species in the surface snow, considering three different seasons: autumn is defined until December 21st, winter until March 21st, and spring from then to the melt onset, identified by 5-6 consecutive days of negative snow accumulation. The average ionic loads of the less abundant (< 1%) species are reported instead in Table S1.

Table 1. Average ionic loads of the most abundant (>1%) ionic and elemental species in the surface snow during each season of the three consecutive sampling campaigns. The standard deviation is shown in brackets, while in the case of nss-SO₄²⁻ the brackets represent the percentage of nss-SO₄²⁻ compared to the total SO₄²⁻. “n” indicates the number of samples considered for the calculation of the average. Annual averages are reported at the end of each sampling year.

mg m ⁻²	Total	Cl ⁻	Na ⁺	SO ₄ ²⁻	nss-SO ₄ ²⁻	Mg	Fe	Ca	NO ₃ ⁻	K ⁺	NH ₄ ⁺
--------------------	-------	-----------------	-----------------	-------------------------------	-----------------------------------	----	----	----	------------------------------	----------------	------------------------------

ha eliminato: ¶

ha formattato: Colore carattere: Nero

ha eliminato: 3.1 Comparison between concentration trends at Gruvebadet and Ny-Ålesund ¶
The concentration variations between an undisturbed area in Ny-Ålesund village and GSRs sites were compared during the 2018-19 sampling campaign to better understand the effect of spatial variability between the two sampling sites. The concentration trends of Na⁺, as sea salt tracer, Pb as anthropogenic species, and Ca²⁺ as crustal tracer, are reported in Fig. S1, for both sampling sites. Although the difference in time resolution between sites is apparent in Fig. S1, the difference in concentration trends appears not statistically significant (p values < 0.05, Wilcoxon test) for all the analysed species, except for sporadic peaks in sea salt and crustal tracers present in the Ny-Ålesund record from November to February. These peaks do not consistently correlate with positive temperature anomalies and precipitation events, as some occurred under low temperatures and without significant precipitation (e.g., January and early February), suggesting that other processes, such as long-range transport, wind-driven deposition, or dry deposition may also play a role (Fig. S1). Concordant Pb trends emerge at Ny-Ålesund and Gruvebadet, with the highest concentrations observed from February to May. ¶
To evaluate the differences in concentration range and spatial distribution of surface snow impurity content, we applied the Wilcoxon test for the 2018-19 sampling period by comparing the distributions for positive and negative differences of the ranks of their absolute values. This comparison is crucial for assessing spatial variability in the measured parameters, providing context and serving as a reference for local-scale differences. However, at a significance level of 0.01, the two distributions from GSRs and Ny-Ålesund sites appeared not statistically different for all the trace elements and most of the inspected ions, except for Na⁺, K⁺, NO₃⁻, Br⁻, and MSA – which are typical sea salt and biological species (MSA). ¶
For this reason, only the GSRs temporal trend has been considered throughout the manuscript, referring to ionic loads (mg m⁻²) instead of concentrations (ng g⁻¹), to highlight the seasonal trends of specific tracers. The ionic load is calculated as ionic concentrations multiplied by the density and the thickness of sampled strata. ¶

ha formattato: Colore carattere: Nero

ha eliminato:).

ha eliminato: ¶

ha eliminato: ¶

ha formattato: Inglese (Regno Unito)

Tabella formattata

ha eliminato: 9¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

autumn 2018 (n=22)	32 (25)	15 (21)	7 (11)	3 (4)	1 (36%)	3 (3)	2 (5)	0.3 (0.3)	0.5 (0.5)	0.3 (0.4)	0.04 (0.03)
winter 2018-19 (n=41)	115 (80)	55 (68)	31 (39)	16 (16)	8 (51%)	8 (8)	0.4 (0.3)	0.4 (0.3)	2 (2)	2 (2)	0.3 (0.3)
spring 2019 (n=51)	75 (50)	36 (43)	19 (24)	9 (9)	4 (48%)	6 (5)	2 (3)	0.6 (0.4)	1 (1)	1 (1)	0.5 (0.4)
Year 2018-19	222 (25)	106 (20)	58 (12)	28 (7)	13 (48%)	17 (3)	4 (1)	1 (0.1)	4 (0.8)	3 (0.6)	0.8 (0.2)
autumn 2019 (n=15)	212 (98)	101 (71)	40 (53)	26 (20)	16 (61%)	10 (6)	1 (1)	9 (12)	5 (4)	2 (3)	4 (5)
winter 2019-20 (n=43)	335 (120)	159 (88)	79 (73)	41 (20)	21 (52%)	16 (8)	2 (5)	9 (10)	3 (2)	3 (4)	7 (8)
spring 2020 (n=49)	264 (131)	110 (91)	52 (77)	28 (25)	15 (53%)	21 (21)	6 (9)	13 (16)	4 (2)	2 (4)	5 (8)
Year 2019-20	811 (39)	370 (31)	171 (20)	95 (8)	52 (55%)	47 (6)	9 (3)	31 (6)	12 (1)	7 (1)	16 (2)
autumn 2020 (n=6)	796 (542)	435 (466)	191 (205)	66 (83)	18 (27%)	84 (165)	1 (2)	2 (5)	6 (11)	9 (10)	2 (2)
winter 2020-21 (n=13)	327 (203)	207 (194)	64 (48)	41 (36)	24 (60%)	6 (5)	0.2 (0.4)	0.2 (0.1)	5 (3)	3 (2)	1 (1)
spring 2021 (n=13)	178 (92)	107 (86)	36 (26)	16 (12)	7 (43%)	9 (10)	4 (7)	1 (2)	3 (2)	2 (2)	1 (1)
Year 2020-21	1300 (194)	749 (168)	291 (83)	123 (25)	49 (40%)	99 (44)	5 (2)	3 (1)	14 (2)	14 (4)	4 (0.5)

ha formattato: Inglese (Regno Unito)

ha eliminato: 116

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha eliminato: 76

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha eliminato: 214

ha formattato: Inglese (Regno Unito)

Tabella formattata

ha formattato: Inglese (Regno Unito)

ha eliminato: 339

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: 273 (132)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha formattato: Colore carattere: Rosso scuro, Inglese (Regno Unito)

ha eliminato: 803

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

Tabella formattata

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha eliminato:

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Colore carattere: Automatico, Inglese (Regno Unito)

ha eliminato: 181

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato

ha eliminato: S2

ha eliminato:),

ha eliminato: S2

ha eliminato: 10¶

ha formattato: Colore carattere: Nero

Formattato

422 ($\chi^2 = 16.9$), MSA ($\chi^2 = 81.3$), SO_4^{2-} ($\chi^2 = 21.3$), nss-SO_4^{2-} ($\chi^2 = 23.4$), Bi ($\chi^2 = 19.2$), Cd ($\chi^2 = 31.9$),
423 Cu ($\chi^2 = 44.5$), Fe ($\chi^2 = 21.9$), Pb ($\chi^2 = 45.1$), Ni ($\chi^2 = 20.6$), Ag ($\chi^2 = 24.5$), U ($\chi^2 = 22.1$), V ($\chi^2 =$
424 40.9), Zn ($\chi^2 = 29.9$). To identify which seasons accounted for these differences, a Dunn's test with
425 Bonferroni correction was applied as a post-hoc test, revealing significant seasonal differences for
426 multiple elements (Fig. S2), with pronounced differences particularly between autumn and winter,
427 and between spring and winter ($P_{\text{adj}} < 0.05$; Table S5).

428 Finally, correlations between meteo-climatic variables (air temperature, wind speed, wind direction,
429 precipitation, AO index) and the concentrations of the selected species were computed to assess
430 potential meteorological influences. The results indicated generally weak positive correlations ($\rho <$
431 0.5, **p-values < 0.05**) for the variables considered. Notably, MSA showed a correlation with air
432 temperature ($\rho = 0.2$, **p-values < 0.05**), as did Bi ($\rho = 0.2$, **p-values < 0.05**), Cu ($\rho = 0.2$, **p-values <**
433 **0.05**), Fe ($\rho = 0.2$, **p-values < 0.05**), Pb (0.1, **p-values < 0.05**), Ni, ($\rho = 0.2$, **p-values < 0.05**), Ag ($\rho =$
434 0.2, **p-values < 0.05**), U ($\rho = 0.3$, **p-values < 0.05**), V ($\rho = 0.3$, **p-values < 0.05**), and Zn ($\rho = 0.3$, **p-**
435 **values < 0.05**).

436 3.2 Polar vortex and Arctic Sea ice extent in 2019-20

437 According to the 2023 survey conducted by the National Snow and Ice Data Center (NSIDC), the
438 maximum extent of Arctic Sea ice since 2014 has been recorded in March 2020, with 14.73 million
439 square kilometres of the Arctic Ocean surface, in a decadal trend **characterised** by a -2.53% of decline,
440 due to the Arctic Amplification. Considering the Kongsfjorden area, the total sea ice extent varied
441 from 63.94 km² in March 2019 to 129.81 km² in March 2020, and was with 46.26 km² lowest in
442 March 2021 (Gerland et al., 2022). Specifications on Drift Ice (DI), Fast Ice (FI), and Open Water
443 (OW) extent are reported in Table S2. The 2020 maximum sea ice extent followed the exceptionally
444 strong and cold stratospheric polar vortex that took place in the Northern Hemisphere (NH) during
445 the 2019-20 polar winter, together with low wave activity from the troposphere, which allowed the
446 polar vortex to remain relatively undisturbed (Lawrence et al., 2020). Notably, the 2020 Arctic Sea
447 ice extent is 16% and 9% higher than previous (2018-19) and following (2020-21) records (dataset
448 NSIDC, NOAA). Lower surface mean air temperatures (-14.7°C, compared to the -11.4°C recorded
449 in 2018-19, and the -8.0°C recorded in the 2020-21), average reduced precipitations (2.1 mm
450 compared to 2.6 mm in 2018-19, and 3.2 mm in 2020-21), higher maximum mean wind speed (17.7
451 ms⁻¹ for 6 days, compared to 15.4 ms⁻¹ for 3 days in 2018-19, and 13.8 ms⁻¹ for 5 days in 2020-21),
452 and higher mean snow height (62.63 cm compared to 58.06 cm in 2018-19, and 57.77 cm in 2020-
453 21) were observed during the 2019-20 winter season, attributed to the influence of a strong polar
454 vortex triggered by a net positive Arctic Oscillation (AO) phase. The 2020 anomalous AO index is

ha eliminato: S3

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:2 + Allineamento: A sinistra + Allinea a: 1,9 cm + Imposta un rientro di: 2,54 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: characterized

ha eliminato: 11

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

457 displayed in Fig. S3. Seasonal values of mean air temperatures (°C), mean precipitation (mm),
458 maximum mean wind speed (m sec⁻¹) and mean snow depth (cm) during the three consecutive
459 sampling campaigns are reported in Table S3. Temperature data were provided by the Norwegian
460 Centre for Climate Services (NCCS), while sea ice extent data were supplied by National Snow and
461 Ice Data Center (NSIDC). Seawater temperature data collected at 10 m depth at a mid-fjord station
462 near Ny-Ålesund (Kb3) was found to be colder during 2020 compared to 2019 and 2021 spring
463 seasons (Table S4). Although these temperatures were still above the freezing point for the observed
464 salinity levels, they contributed to colder overall conditions in Kongsfjorden, supporting the
465 formation and prolonged presence of sea ice when combined with cold atmospheric conditions.

466 **4. Discussion**

467 *4.1 Ny-Ålesund seasonal and interannual trends variability in surface snow*

468 The three consecutive sampling campaigns conducted from 2018 to 2021 confirmed the dominance
469 of sea salt input in the surface snow of Svalbard, due to the proximity of the sampling site to the
470 coastline (Barbaro et al., 2021). The dominant ions are Na⁺, Cl⁻, and SO₄²⁻, associated with the
471 scavenging precipitation of marine aerosol (Hodgkins and Tranter, 1998). The observed mean
472 seasonal trends (Fig. S1) display the highest concentrations of marine species in autumn 2020,
473 followed by 2020-21 and 2019-20 winter seasons. However, wintry concentrations are presumably
474 linked to weakened (2019-20) or destroyed (2020-21) polar vortex (Fig. 2) and intense cyclonic
475 storms, associated with anomalous warming events capable of transporting both heat and moisture
476 from lower latitudes to Svalbard (Rinke et al., 2017). In addition to these factors, it is important to
477 account for snow ablation or erosion by wind. The change in snow height (Δh) between consecutive
478 sampling events provides valuable insights into such processes. A decrease in snow height signals
479 ablation can indeed result from snow melting (under positive temperatures), snow aging, sublimation,
480 or snow drift. In particular, when wind speeds exceed 5 m s⁻¹, the threshold commonly accepted for
481 initiating snow drift and ablation episodes (Pomeroy, 1989), we can infer the snow erosion due to
482 wind drift likely occurred (Fig. 2). This effect may explain some of the variability in snow ion
483 concentrations, especially during intense storms or strong winds.

484 Autumn 2020 represents most likely an outlier, due to scarce precipitations (Fig. 3) that led to more
485 concentrated impurities in the surface snow. In contrast, late spring 2020 saw a notable increase in
486 typical marine ions (Na⁺, Cl⁻, Br⁻, MSA, SO₄²⁻) and geogenic elements (Al, Ca, Mn, Fe, Sr), compared
487 to spring 2019 (Fig. 3). This increase may be attributed to the very close drift Arctic Sea ice presence
488 in Kongsfjorden (Table S2), which reached its maximum extent in March 2020 due to low-

ha eliminato: S4

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 0 pt, Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:4 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: sites to the coastline (Barbaro et al., 2021).


ha eliminato: (Hodgkins and Tranter, 1998). The observed mean seasonal trends (Fig. S2

ha eliminato: 12

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

493 temperature anomalies and intensified atmospherically driven sea ice transport and deformation,
494 caused by higher than normal winter wind speeds (Fig. S4). These conditions likely enhanced the
495 production of sea spray aerosols, which, when carried by winds, may have increased the deposition
496 of marine species onto the snowpack. The increased deposition of geogenic elements might also have
497 been influenced by low temperature anomalies (as seen with Fe), and/or by stronger wind speeds,
498 although significant correlations have not emerged in this preliminary statistical analysis.
499 Additionally, an outstanding positive phase of the Arctic Oscillation (AO) in the troposphere (Fig. 2)
500 was recorded in January-March 2020 (Lawrence et al., 2020; Dethloff et al., 2022). Although the
501 entire period was remarkable, January and February featured as clear outliers in the historical
502 timeseries (1950–2023) reported by the NOAA service, while March 2020 exhibited significantly
503 elevated values but did not exceed the statistical threshold for outliers (Fig. S3).



504 **Fig. 2.** AO Index, air temperature (°C), precipitation (mm), snow height (cm), Δh snow height (cm), wind speed (m sec⁻¹), and wind direction (°) from the NCEP/NCAR Reanalysis data. The green horizontal line above the wind speed graph


505

506

ha eliminato: S5

ha eliminato: S4).

ha formattato: Tipo di carattere: Calibri, 11 pt, Non Evidenziato



ha eliminato:

ha eliminato: 13

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

510 indicates the 5 m sec⁻¹ threshold, above which wind drift may occur on surface snow layers. The colour of the line refers
511 to the Δh color scale, which indicates negative values of Δh, NOAA Physical Sciences Lab's daily composites tool was
512 used to calculate the near-surface air temperatures across the Northern Hemisphere from October to May. Grey bands
513 indicate the winter periods.

514 A 2021 spring peak of marine species was also recorded, although more attenuated than in spring
515 2020 (Fig. 3, Fig. [S1](#)). This variation is likely attributable to different extents of sea ice in the fjord.
516 Nonetheless, seawater temperatures in 2021, similar to those in 2020 and 2.3°C colder than in 2019
517 (Table S4), along with comparable wind speed conditions (Fig. [S4](#)), may also have contributed to the
518 observed trends in marine species concentrations. Similarly, the spring peak of Mg, Sr, Mn, Fe, Al,
519 and V in 2021 seems to reflect the high wind speed and positive AO index recorded from March to
520 April 2021. Positive anomalies for air temperatures (A) and wind speed (W) conditions, together with
521 negative anomalies in seawater (O) conditions were observed during the 2020-21 campaign, while
522 negative A and O conditions were accompanied to positive W during 2019-20. On the contrary, 2018-
523 19 diverges from the other campaigns for positive O condition associated to negative W condition
524 anomalies. These observations provide valuable insights into how shifts in atmospheric and oceanic
525 conditions impact the concentrations of ionic and elemental species in surface snow, enhancing our
526 understanding of the underlying mechanisms that govern these changes in the context of AA
527 conditions.

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: S2

ha eliminato: S5

ha eliminato: 14 ¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra





Fig. 3. Wind direction ($^{\circ}$), wind speed (m sec^{-1}), and ionic loads (mg m^{-2}) of Na^+ , Cl^- , Mg , SO_4^{2-} , nss-SO_4^{2-} , MSA , Br^- , Ca , Sr , Mn , Fe , Al , Pb , V , Ni in the surface snow for the three sampling campaigns: 2018-19, 2019-20, 2020-21. Seasonal trends are here presented for specific elements to provide a detailed view of how concentrations vary across distinct sampling periods.

A singular case is represented by Pb , which showed a 12.5-fold increase during winter-spring 2019 period compared to autumn 2018. The EF for Pb during these seasons exceeded 100, indicating a strong anthropogenic contribution. In contrast, the peak observed in 2020 no longer shows such elevated EF values (i.e., $\text{EF} = 26$), suggesting a mixed source (Fig. 4). This implies that the April-May 2020 peak is largely of crustal origin, as the overlap with V ($\text{EF} < 10$) also suggests, possibly due to local dust events driven by strong winds exceeding the 5 m sec^{-1} threshold (Fig. 3; Table S3).





ha eliminato:

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: 15

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra: (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra



ha eliminato:

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha formattato: Inglese (Regno Unito)

ha eliminato: 16

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

556 The 2019 springtime Pb concentration maxima are typically consistent with a mixture of eastern and
557 western European sources (Sherrell et al., 2000; Bazzano et al., 2015, 2021). In this study, cluster
558 mean trajectories obtained for winter 2018-2019 highlighted a 25% of air mass provenance from
559 Russian Arctic and a 13% from eastern Siberia (Fig. S5), possibly explaining the higher
560 concentrations of Pb revealed in spring 2019, following a reduced precipitation regime that occurred
561 in January 2019. A local anthropogenic origin can be excluded though, since no activities (ordinary-
562 extraordinary maintenance or particular events) were recorded in the vicinity of the sampling site in
563 the 2019 sampling season. However, at present, the long-range transport of Pb remains a hypothesis,
564 likely supported by the breakdown of the wintry polar vortex (Fig. 2).

565 Backward trajectories (Fig. S5) for Ny-Ålesund area (78.92° N, 11.89° E) appear mostly in line with
566 literature findings (Platt et al., 2022; Vecchiato et al., 2024), showing three main seasonal characters:
567 a prevalent mass movement from ice-covered Central Arctic Ocean, Kara Sea, and Greenland Sea
568 during autumn, a main provenance from Central Arctic Ocean and Kara Sea during winter, and a
569 predominant trajectory from Northern Canada in addition to air masses arriving from Arctic Ocean
570 and Kara seas during spring.

571 4.2 *The main ion sources in the seasonal snow of Ny-Ålesund*

572 Examining the dominant ions associated with marine aerosol, we found Cl⁻/Na⁺ median ratios ranging
573 from 1.3 to 1.5 w⁻¹, slightly lower than the expected value of 1.8 w⁻¹ in the pure seawater (Zhuang
574 et al., 1999), pointing the occurrence of a minimum Cl⁻ depletion in aerosol, quantified as 14% for
575 the 2018-19 and 2019-20 campaigns, and as just 2% for the 2020-21 campaign. A possible
576 explanation for this phenomenon could be the de-chlorination of sea-spray aerosol during transport.
577 This reaction occurs when sea-salt particles containing NaCl interact with acids such as HNO₃,
578 H₂SO₄, or organic acids, leading to the release of gaseous HCl (Zhuang et al., 1999, and reference
579 therein). Although less likely, this process could also occur at the snow-atmosphere interface. On the
580 contrary, a possible influence of biomass burning on Cl⁻ depletion process has been excluded by the
581 very low correlation (0.18, p-value < 0.05) found between Cl⁻ depletion values and nss-K⁺/K⁺ ratios,
582 which is a tracer of relative contribution of biomass burning (Song et al., 2018).

583 Positive correlations between Mg, Ca, and K⁺ with Na⁺ and Cl⁻ (Fig. 5, p-value < 0.05) suggest a
584 common sea-spray source. However, the concentrations of Mg are also positively correlated with nss-
585 Ca (ρ_{load} = 0.55, p-value < 0.05), calculated according to Morales et al. (1998), indicating a shared
586 non-marine source. This suggests that in addition to their marine origin, these ions (Mg, Ca) also have

ha eliminato: S6

ha eliminato: In addition, both GSRS and Ny-Ålesund (Fig. 1), located at 1 km of distance from each other, recorded comparable high concentrations of Pb, thus ruling out a possible contamination.

ha eliminato: S6

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello: 2 + Stile numerazione: 1, 2, 3, ... + Comincia da: 2 + Allineamento: A sinistra + Allinea a: 1,26 cm + Imposta un rientro di: 1,89 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra: (Nessun bordo)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato:

ha eliminato:

ha eliminato: 17

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra: (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

595 contributions from a non-marine, crustal source. Further evidence comes from the Ca/Mg ratios in
596 surface snow samples collected during the three campaigns, which were higher than those found in
597 seawater (0.32, Millero et al., 2008). This excess indicates the likely presence of mineral particles
598 (i.e., calcite and dolomite), potentially originating from local rock or soil dust (e.g., limestone,
599 dolostone, and marble, which are abundant in Svalbard), as previously observed by Barbaro et al.
600 (2021). To further explore the mixed sources of these elements, we calculated the Enrichment Factors
601 (EFs) of Mg and Ca. For Mg, the EF values were consistently above 10 in all the seasons analysed,
602 indicating a significant non-crustal source (e.g., marine). In contrast, Ca displayed EF values greater
603 than 10 only during autumn 2019, suggesting that its non-crustal contribution (likely from sea spray)
604 was more pronounced during that specific season. Based on these findings, we conclude that Mg and
605 Ca effectively share a common sea spray origin, but the sea-salt contribution of Ca was mainly
606 significant in autumn 2019, while the excess of Ca in other seasons likely reflects inputs from crustal
607 source, such as local mineral dust.

608 Similarly, sulphate (SO_4^{2-}) is highly and significantly correlated ($p\text{-values} < 0.05$) with both Na^+ ($\rho_{\text{load}} = 0.80$) and Cl^- ($\rho_{\text{load}} = 0.91$), indicating that sea-spray is its main source. Nonetheless, $\text{Na}^+/\text{SO}_4^{2-}$ and
609 $\text{Cl}^-/\text{SO}_4^{2-}$ ratios are significantly lower than typical seawater values (3.97 and 7.13, respectively,
610 according to Millero et al., 2008) for the former two campaigns (2018-19, 2019-20). The elevated
611 sulphate concentrations compared to sodium, also in winter snow, suggest the absence of a strong
612 frost flower signature. Additionally, the lack of significant depletion in nss-SO_4^{2-} further supports the
613 minimal role of frost flowers in contributing to the snow composition. Therefore, while frost flowers
614 are known to impact snow chemistry in Svalbard (Rankin et al., 2002; Beaudon and Moore, 2009;
615 Roscoe et al., 2011), our analysis indicates that their contribution to the observed sea salt peaks in
616 Kongsfjorden during the 2018-19, 2019-20 campaigns was likely limited. This analysis highlights
617 that, while sea ice extent supports higher sea-salt concentrations in snow, the specific sulphate and
618 ion ratios observed point to sea spray as the main source, with only a minor role for frost flower-
619 related inputs.

621 Instead, the presence of nss-SO_4^{2-} suggests potential inputs from other sources, such as crustal
622 material, anthropogenic emissions (e.g., fossil fuel combustion), or the oxidation of dimethylsulfide
623 (DMS) released from marine biological activities.

624 To estimate the crustal fraction of sulphate (cr-SO_4^{2-}), the nss-Ca (as crustal marker) content was
625 multiplied by 0.59 ($\text{SO}_4^{2-}/\text{Ca}$ w/w ratio in the uppermost Earth crust - Wagenbach et al. 1996),
626 obtaining variable contributions for the three sampling campaigns, ranging from 2.45% up to 12.94%.

ha eliminato: 18

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

Conversely, the anthropogenic contribution to nss-SO_4^{2-} concentrations was investigated by the application of the $[\text{ex-SO}_4^{2-}]$ concentration formula (Schwikowski et al., 1999), considering the average concentration of $[\text{Ca}]$ instead of the average ionic concentration $[\text{Ca}^{2+}]$ because Ca^{2+} concentrations were not measured in the samples collected during the second campaign due to instrumental limitations:

$$[\text{ex-SO}_4^{2-}] = [\text{SO}_4^{2-}] - (0.12 [\text{Na}^+]) - (0.175 [\text{Ca}^{2+}])$$

The obtained results showed a 50 up to 60% of anthropogenic contribution for the nss-SO_4^{2-} input. This finding corroborates previous results from Amore et al. (2022), who noted that anthropogenic sulphate was the most abundant apportioned component at Gruvebadet, accounting for at least 50% all over the year during the 2010-2019 investigated period. The plausible source of the anthropogenic fraction is the atmospheric transport of secondary aerosols containing SO_4^{2-} , and ammonium sulphate. This sulphate can be formed by SO_x emitted from coal combustion throughout the winter and biomass burning in the spring (Barbaro et al., 2021 and reference therein). The nss-SO_4^{2-} does not correlate significantly with other ionic species (except for Mg), thus suggesting a separate origin (Fig. 5).




Fig. 5. Spearman correlation plots. A) Correlation plot for the three sampling seasons; B) Correlation plot for the 2018-2019 season; C) Correlation plot for the 2019-2020 season; D) Correlation plot for the 2020-2021 season.

ha eliminato: 19

ha eliminato: 19

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

653 To quantify the biogenic nss-SO₄²⁻ contribution, the methanesulfonic acid (MSA) loads - the final
654 product of DMS oxidation - were multiplied by 3.0 (Udisti et al., 2016), revealing biogenic SO₄²⁻
655 contributions ranging from 0.15% (2018-19, 2020-21) up to 0.38% (2019-20). Furthermore, the
656 MSA/nss-SO₄²⁻ ratio was inspected, revealing a mean value of 0.02 ± 0.03 during the first (2018-19)
657 and the third (2020-21) sampling campaigns, and a maximum ratio equal to 0.06 ± 0.18 reached
658 during the second campaign (2019-20), similar to the subarctic western North Pacific ratio found by
659 Jung et al. (2014). However, several factors can influence MSA formation, a univocal marker of
660 biogenic emissions, including higher biological productivity related to higher nutrient input; the
661 concentrations of NO₃ radicals as key oxidants for DMS decomposition (higher NO₃ gives higher
662 MSA); and lower air temperatures, which tend to yield higher MSA levels (Andreae et al., 1985;
663 Udisti et al., 2020).

664 For the 2019-20 campaign, it seems likely that a combination of these three factors, together with the
665 positive expansion of sea ice and the very close drift ice presence in March 2020, as revealed from
666 satellite reconstructions (Fig. S6), contributed to the increased release of MSA in aerosol, and its
667 consistent deposition in surface snow (Fig. 3). Indeed, DMS was likely accumulated under the sea
668 ice cover in the fjord and surrounding areas, and then being released and oxidised in the atmosphere
669 when the ice broke off and melted (April-May). Furthermore, lower temperatures (-17.3°C in March
670 2020, compared to -16.1°C in March 2019, and -9.8°C in March 2021), positive correlation between
671 MSA and NO₃⁻ ($p_{\text{load}} = 0.49$, $p\text{-value} < 0.05$), and high-speed short-range transport (wind directions
672 between 0°-60° and speeds > 5 m sec⁻¹) from the source to the near-coast sink site (GSRS) would
673 have aided elevated concentrations of MSA in atmospheric depositions. However, the dominant south
674 and southwest winds (180°-240°) during the major MSA peak in April 2020 (Fig. 3) likely transported
675 marine aerosols and DMS from open ocean regions, further facilitating the increased MSA
676 concentrations observed.

677 Contrarily, in the 2018-19 season, sea ice lasted only until April and was restricted to the inner,
678 shallower parts of Kongsfjorden (Assmy et al., 2023), possibly not allowing enough time with
679 adequate sunlight for substantial biological activity to accumulate beneath or within it. This occurred
680 despite the dominance in 2019, unlike the following year, of *Phaeocystis pouchetii*, a phytoplankton
681 species known for its capacity to generate DMS in significant quantities, according to Assmy et al.
682 (2023).

683 Finally, the ammonium (NH₄⁺) load showed positive correlations (Fig. 5) with Na⁺ ($p_{\text{load}} = 0.71$, $p\text{-value} < 0.05$),
684 Cl⁻ ($p_{\text{load}} = 0.52$, $p\text{-value} < 0.05$) and K⁺ ($p_{\text{load}} = 0.72$, $p\text{-value} < 0.05$), as well as with

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: S7

ha eliminato: 20¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

686 SO_4^{2-} ($\rho_{\text{load}} = 0.54$, **p-value < 0.05**), NO_3^- ($\rho_{\text{load}} = 0.45$, **p-value < 0.05**), MSA ($\rho_{\text{load}} = 0.37$, **p-value <**
687 **0.05**) and Br^- ($\rho_{\text{load}} = 0.53$, **p-value < 0.05**), suggesting a close link with sea-salt ions and biogenic
688 emissions. However, some contribution from biomass burning events and potential influence from
689 anthropogenic activities cannot be excluded.

690 4.3 Bromine enrichment

691 The bromine enrichment factor (Br_{enr}) is well known to reflect specific processes (i.e., sea ice gas
692 phase Br emission) that affect the Br concentration and load in the snowpack (Spolaor et al., 2014).
693 Therefore, calculating the relative enrichment over the Br/Na ratio in sea water can offer crucial
694 insights on sea ice variability for the investigated Arctic areas (Barbaro et al., 2021). As reported in
695 previous studies (Maffezzoli et al., 2017; Barbaro et al., 2021), the Br enrichment factor (Br_{enr}) can
696 be calculated as $\text{Br}_{\text{enr}} = \text{Br}^- / (0.0065 \text{ Na}^+)$, where 0.0065 represents the Br/Na^+ seawater mass ratio.
697 Contrarily to what observed in a former study (Barbaro et al., 2021) for the Hornsund area and north-
698 western Spitsbergen, where the Br_{enr} mean values were often < 1 , indicating some Br^- depletion, in
699 this study we observed Br_{enr} mean values ranging from 1.5 up to 17.7, with the highest value
700 associated to the second sampling campaign conducted in 2019-20, which showed the most extensive
701 sea ice coverage. These results support the impact of the sea ice expansion and the close drift ice in
702 the Kongsfjorden on the snow chemical composition. Indeed, newly formed sea ice releases gas-
703 phase Br into the polar atmosphere, thus supplying an extra Br^- source in addition to sea spray
704 (Spolaor et al., 2016).

705 4.4 Anthropogenic and natural sources of ions and particulate trace elements

706 To complement the EFs analysis and further distinguish possible anthropogenic contributions from
707 natural ones (marine and geogenic) for ions and particulate trace elements, a Hierarchical Cluster
708 Analysis (HCA) method was carried out. Results of clustering (Fig. 6) clearly disentangle marine
709 (Na^+ , Cl^- , K^+ , NH_4^+ , SO_4^{2-} , NO_3^- , Br^- , Mg, Sr, bio- SO_4^{2-} , MSA), anthropogenic (As, Co, Ag, Ba, Cd,
710 Zn, Pb, Bi, Cr, Cu, Ni), and geogenic (nss-Ca, Tl, Li, Al, Cs, Rb, Fe, Mn, U, Be, Se, V) sources of
711 ionic and elemental species, considering the whole sampling campaign period (2018-2021).
712 Interestingly, nss- SO_4^{2-} is brought together with the marine cluster, suggesting that its presence is
713 largely influenced by marine biogenic sources, alongside contributions from secondary sulphate
714 formation in the atmosphere. This indicates that nss- SO_4^{2-} , despite having a variety of sources such
715 as human contribution or dust, is closely linked to the marine environment. One important reason for
716 this is the emission of DMS by phytoplankton. Additionally, secondary sulphate formation may have

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello: 2 + Stile numerazione: 1, 2, 3, ... + Comincia da: 2 + Allineamento: A sinistra + Allinea a: 1,26 cm + Imposta un rientro di: 1,89 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello: 2 + Stile numerazione: 1, 2, 3, ... + Comincia da: 2 + Allineamento: A sinistra + Allinea a: 1,26 cm + Imposta un rientro di: 1,89 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Inglese (Regno Unito)

Formattato: SpazioPrima: 14 pt, Dopo: 14 pt

ha eliminato: 21 ¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

717 further contributed to the nss-SO₄²⁻. **Cobalt** was grouped within the anthropogenic cluster in HCA, in
718 contrast with EFs that suggested a crustal origin. This apparent discrepancy may reflect the relatively
719 larger uncertainties typically associated with EF calculations, which can inherit errors from the choice
720 of reference element, assumptions about crustal composition, and variability in background
721 concentrations, compared to the more integrative approach of HCA (e.g., Reimann and de Caritat,
722 2000). Moreover, the trend for **Co** closely aligns with anthropogenic ones (e.g., As), while distinct
723 trends are evident for crustal elements. Therefore, while the HCA results offer a reliable perspective
724 on source differentiation and clustering patterns, they are best interpreted as complementary to the
725 EF calculations rather than directly integrated with them. Incorporating EFs directly into the HCA
726 could introduce significant inaccuracies, as EFs rely on reference element ratios that may vary and
727 thus add complexity to the clustering process, potentially skewing the results. Consequently, HCA
728 independently provides robust insights in this context, enhancing our understanding without the
729 additional uncertainties that EF-based clustering might introduce.

ha eliminato: Co
ha formattato: Inglese (Regno Unito)

ha eliminato: cobalt
ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)





Fig. 6. Hierarchical cluster analysis applied to further disentangle the particulate trace element non-crustal sources.

5. Summary and Conclusion



ha eliminato:
ha formattato: Colore carattere: Nero

Formattato: Normale, SpazioPrima: 14 pt, Dopo: 14 pt, Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:4 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: 22

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

In this study, trace elements and major ions were investigated in surface snow samples collected in Ny-Ålesund between October 2018 to June 2021. Seasonal and interannual variations of impurities have been observed, with general higher concentrations of marine species revealed in late spring 2020, associated to more extensive sea ice in Kongsfjorden in March 2020, promoted by negative temperature anomalies in both atmosphere and ocean and likely related to higher air mass recycle within the Arctic. These results provide direct evidence of how sea ice extent modulates the storage, release, and transport of marine-derived impurities, thereby influencing snow-atmosphere chemical exchange processes under varying climatic conditions. Higher concentrations in spring 2020 for geogenic and anthropogenic species are consistent with the influence of higher wind speeds, low atmospheric temperature anomalies, and generally drier conditions resulting from the exceptional occurrence of a strong and cold stratospheric polar vortex, accompanied by an unprecedentedly positive phase of the Arctic Oscillation in the troposphere during January-March 2020. While correlations between meteorological variables and concentrations were generally weak, the overall meteorological context suggests that these unusual conditions likely contributed to the observed patterns. This is particularly evident especially in cold years, when the production of sea spray related aerosol likely derives by a larger extension of sea ice and stronger local Arctic circulation. The identification of geogenic, marine, and anthropogenic sources in the snowpack was allowed by a chemometric approach (HCA), which clarified the EFs results. The use of chemometric techniques (HCA) and back-trajectory analysis enabled a clearer attribution of sources and transport pathways, improving the interpretation of snow composition in relation to meteorological drivers. Specifically, the distinct seasonal air mass patterns revealed, characterised by dominant Arctic-origin air masses in fall and winter and a lack of expected mid-latitude inputs in spring, underscore the changing dynamics of snow-atmosphere interactions in a warming Arctic. Overall, these insights advance our understanding of how recent climatic anomalies, such as altered sea ice extent, shifts in Arctic Oscillation phases, and stronger polar vortices, modulate the chemical composition of the snowpack in Svalbard. Our findings highlight the sensitivity of snow-atmosphere exchanges to both local and large-scale climatic processes, offering important context for interpreting snow chemistry trends in a rapidly changing Arctic environment. Further statistical investigations, including multivariate and longer-term analyses, would be valuable for better quantifying these correlations.

Data availability

The data supporting the findings of this study are available within the article and its supplementary materials. Other data that support the findings of this study are available from the corresponding author upon request.

ha formattato: Inglese (Regno Unito)
Formattato: SpazioPrima: 14 pt, Dopo: 14 pt
ha formattato: Inglese (Regno Unito)
ha formattato: Inglese (Regno Unito)

ha eliminato: Such findings illustrate how large-scale atmospheric circulation anomalies associated with Arctic Amplification may significantly alter the deposition patterns of both natural and anthropogenic species in the snowpack.
ha formattato: Inglese (Regno Unito)

ha eliminato: 23¶
ha formattato: Colore carattere: Nero
Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

774 **Author contribution**

775 AS: Conceptualization, Data curation, Investigation, Writing-original draft, Writing-review and
776 editing. EB: Conceptualization, Field work, Data curation, Formal Analysis, Writing-original draft,
777 Writing-review and editing, Funding acquisition. MF: Formal Analysis, Field work, Data curation,
778 Writing-review and editing. FS: Field work, Formal analysis, Investigation, Writing-review and
779 editing. MV: Writing-review and editing. MV: Field work, Writing-review and editing. MM:
780 Investigation. FB: Investigation, Writing-review and editing. FB: Investigation. Field work. CJMH:
781 Investigation, Data curation, Writing-review and editing. AB: Field work, Data curation, Writing-
782 review and editing. AG: Resources, Supervision, Validation, Writing-review and editing, Funding
783 acquisition. CB: Resources, Supervision, Validation, Writing-review and editing, Funding
784 acquisition. AS: Funding acquisition, Supervision, Validation, Writing-review and editing.

785 **Competing interests**

786 The authors declare that they have no conflict of interest.

787 **Acknowledgments**

788 This project has received funding from the European Union's Horizon 2020 research and innovation
789 programme under grant agreement no. 689443 via ERA_PLANET Strand 4 project iCUPE
790 (Integrative and Comprehensive Understanding on Polar Environments). We are grateful to the Arctic
791 Station Dirigibile Italia from the Italian National Research Council – Institute of Polar Science (CNR-
792 ISP) for logistical support. The authors gratefully acknowledge Claudio Artoni, Maria Papale, Ivan
793 Sartorato, Federico Dallo, Alice Callegaro, Marco Casula, Mariasilvia Giamberini, Fabio Giardi, and
794 all the station leaders of “Dirigibile Italia” who participated and offered valuable help and logistic
795 support during the 2018-2021 sampling campaigns. Furthermore, we acknowledge Klara Wolf, Linda
796 Rehder, and Ane Kvernvik for help with CTD sampling. We acknowledge the help of ELGA
797 LabWater in providing the PURELAB Pulse and PURELAB Flex, which produced the ultrapure
798 water used in these experiments.

799 **References**

800 Amore, A., Giardi, F., Becagli, S., Caiazzo, L., Mazzola, M., Severi, M., Traversi, R.: Source
801 apportionment of sulphate in the High Arctic by a 10 yr-long record from Gruebadet Observatory
802 (Ny-Ålesund, Svalbard Islands). Atmos. Environ., 270, 118890, 1-10,
803 <https://doi.org/10.1016/j.atmosenv.2021.118890>, 2022
804

ha formattato: Tipo di carattere: 12 pt, Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: <https://doi.org/10.1016/j.atmosenv.2021.118890>

ha formattato: Tipo di carattere: 12 pt, Colore carattere: Nero

ha eliminato: 24

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

807 Andreae, M.O., Ferek, R.J., Bermond, F., Byrd, K.P., Engstrom, R.T., Hardin, S., Houmere,
808 P.D., LeMarrec, F., Raemdonck, H., Chatfield, R.B.: Dimethyl sulfide in the marine atmosphere,
809 J. Geophys. Res., 90, D7, 12891-12900, <https://doi.org/10.1029/JD090iD07p12891>, 1985

810 Assmy, P., Kvernvik, A.C., Hop, H., Hoppe, C.J.M., Chierici, M., David, D., Duarte, P.,
811 Fransson, A., Garcia, L.M., Patuła, W., Kwaśniewski, S., Maturilli, M., Pavlova, O., Tatarek, A.,
812 Wiktor, J.M., Wold, A., Wolf, K.K.E., Bailey, A.: Seasonal plankton dynamics in Kongsfjorden
813 during two years of contrasting environmental conditions, Prog. Oceanogr., 213, 102996,
814 <https://doi.org/10.1016/j.pocan.2023.102996>, 2023

815 Barbaro, E., Koziol, K., Björkman, M.P., Vega, C.P., Zdanowicz, C., Martma, T., Gallet, J.C.,
816 Kępski, D., Larose, C., Luks, B., Tolle, F., Schuler, T. V., Uszczyk, A., Spolaor, A.: Measurement
817 report: Spatial variations in ionic chemistry and water-stable isotopes in the snowpack on glaciers
818 across Svalbard during the 2015-2016 snow accumulation season, Atmos. Chem. Phys., 21, 3163–
819 3180, <https://doi.org/10.5194/acp-21-3163-2021>, 2021

820 Barbaro, E., Padoan, S., Kirchgeorg, T., Zangrando, R., Toscano, G., Barbante, C., Gambaro,
821 A.: Particle size distribution of inorganic and organic ions in coastal and inland Antarctic aerosol,
822 Environ. Sci. Pollut. Res., 24, 2724-2733, <https://doi.org/10.1007/s11356-016-8042-x>, 2017

823 Barbaro, E., Zangrando, R., Vecchiato, M., Piazza, R., Cairns, W.R.L., Capodaglio, G.,
824 Barbante, C., Gambaro, A.: Free amino acids in Antarctic aerosol: Potential markers for the evolution
825 and fate of marine aerosol, Atmos. Chem. Phys., 15, 5457–5469, <https://doi.org/10.5194/acp-15-5457-2015>, 2015

827 Bazzano, A., Ardini, F., Becagli, S., Traversi, R., Udisti, R., Cappelletti, D., Grotti, M.: Source
828 assessment of atmospheric lead measured at Ny-Ålesund, Svalbard, Atmos. Environ., 113, 20-26,
829 <https://doi.org/10.1016/j.atmosenv.2015.04.053>, 2015

830 Bazzano, A., Bertinetti, S., Ardini, F., Cappelletti, D., Grotti, M.: Potential Source Areas for
831 atmospheric Lead reaching Ny-Ålesund from 2010 to 2018, Atmos., 12, 388, 1-17,
832 <https://doi.org/10.3390/atmos12030388>, 2021

833 Beaudon, E., Moore, J.: Frost flower chemical signature in winter snow on Vestfonna ice cap,
834 Nordaustlandet, Svalbard, Cryosphere, 3, 147–154, <https://doi.org/10.5194/tc-3-147-2009>, 2009

835 Beine, H.J., Dominé, F., Ianniello, A., Nardino, M., Allegrini, I., Teinilä, K., Hillamo, R.:
836 Fluxes of nitrates between snow surfaces and the atmosphere in the European high Arctic, Atmos.
837 Chem. Phys., 3, 335-346, <https://doi.org/10.5194/acp-3-335-2003>, 2003

838 Bertò, M., Cappelletti, D., Barbaro, E., Varin, C., Gallet, J.-C., Markowicz, K.,
839 Rozwadowska, A., Mazzola, M., Crocchianti, S., Poto, L., Laj, P., Barbante, C., Spolaor, A.:
840 Variability in black carbon mass concentration in surface snow at Svalbard, Atmos. Chem. Phys., 21,
841 12479-12493, <https://doi.org/10.5194/acp-21-12479-2021>, 2021

842 Björkman, M., Kühnel, R., Partridge, D.G., Roberts, T.J., Aas, W., Mazzola, M., Viola, A.,
843 Hodson, A., Ström, J., Isaksson, E.: Nitrate dry deposition in Svalbard, Tellus B, 65(1),
844 <https://doi.org/10.3402/tellusb.v65i0.19071>, 2013

845 Chylek, P., Folland, C., Klett, J.D., Wang, M., Hengartner, N., Lesins, G., Dubey, M.K.:
846 Annual Mean Arctic Amplification 1970-2020. Observed and Simulated by CMIP6 Climate Models,
847 Geophys. Res. Lett., 49 (13), e2022GL099371, <https://doi.org/10.1029/2022GL099371>, 2022

Formattato: Regola lo spazio tra testo asiatico e in alfabeto latino, Regola lo spazio tra caratteri asiatici e numeri

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1029/JD090iD07p12891>, 1985

ha eliminato: <https://doi.org/10.1016/j.pocan.2023.102996>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-21-3163-2021>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1007/s11356-016-8042-x>, 2017

ha eliminato: <https://doi.org/10.5194/acp-15-5457-2015>

ha eliminato: <https://doi.org/10.1016/j.atmosenv.2015.04.053>, 2015

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3390/atmos12030388>

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-3-335-2003>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-21-12479-2021>, 2021...

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3402/tellusb.v65i0.19071>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1029/2022GL099371>

ha eliminato: 25¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

863 D'Amico, M., Kallenborn, R., Scotto, F., Gambaro, A., Gallet, J.-C., Spolaor, A., Vecchiato,
864 M.: Chemicals of Emerging Arctic Concern in north-western Spitsbergen snow: Distribution and
865 sources, *Sci. Total Environ.*, 908, 168401, <https://doi.org/10.1016/j.scitotenv.2023.168401>, 2024

866 Dethloff, K., Maslowski, W., Hendricks, S., Lee, Y.J., Goessling, H.F., Krumpfen, T., Haas,
867 C., Handorf, D., Ricker, R., Bessnov, V., Cassano, J.J., Kinney, J.C., Osinski, R., Rex, M., Rinke, A.,
868 Sokolova, J., Sommerfeld, A.: Arctic sea ice anomalies during the MOSAiC winter 2019/20,
869 *Cryosphere*, 16, 981-1005, <https://doi.org/10.5194/tc-16-981-2022>, 2022

870 Draxler, R.R.: An overview of the HYSPLIT 4 modelling system for trajectories, dispersion
871 and deposition, *Aust. Meteorol. Mag.*, 47, 295–308, 1998

872 Dunn, O.J.: Multiple comparisons among means, *J. Am. Stat. Assoc.*, 56, 293,
873 <https://doi.org/10.1080/01621459.1961.10482090>, 1961

874 Eleftheriadis, K., Vratolis, S., Nyeki, S.: Aerosol black carbon in the European Arctic:
875 Measurements at Zeppelin station, Ny-Ålesund, Svalbard from 1998-2007, *Geophys. Res. Lett.*, 36,
876 1–5, <https://doi.org/10.1029/2008GL035741>, 2009

877 Feltracco, M., Barbaro, E., Hoppe, C.J.M., Wolf, K.K.E., Spolaor, A., Layton, R., Keuschig,
878 C., Barbante, C., Gambaro, A., Larose, C.: Airborne bacteria and particulate chemistry capture
879 Phytoplankton bloom dynamics in an Arctic fjord, *Atmos. Environ.*, 256, 118458,
880 <https://doi.org/10.1016/j.atmosenv.2021.118458>, 2021

881 Feltracco, M., Barbaro, E., Tedeschi, S., Spolaor, A., Turetta, C., Vecchiato, M., Morabito,
882 E., Zangrando, R., Barbante, C., Gambaro, A.: Interannual variability of sugars in Arctic aerosol:
883 Biomass burning and biogenic inputs, *Sci. Total Environ.*, 706, 136089,
884 <https://doi.org/10.1016/j.scitotenv.2019.136089>, 2020

885 Geng, H., Ryu, J., Jung, H.J., Chung, H., Ahn, K.H.O., Ro, C.U.N.: Single-particle
886 characterization of summertime arctic aerosols collected at Ny-Ålesund, Svalbard, *Environ. Sci.*
887 *Technol.*, 44, 2348–2353, <https://doi.org/10.1021/es903268j>, 2010

888 George, B.J., Gains-Germain, L., Broms, K., Black, K., Furman, M., Hays, M.D., Thomas,
889 K.W., Simmons, J.E.: Censoring trace-level environmental data: statistical analysis considerations to
890 limit bias, *Environ. Sci. Technol.*, 55(6), 3786-3795, <https://doi.org/10.1021/acs.est.0c02256>, 2021

891 Gerland, S., Pavlova, O., Marnela, M., Divine, D.V., Kohler, J., Renner, A., Skoglund, A.:
892 Sea ice extent variability in Kongsfjorden, Svalbard during 2003-2021, based on visual observations
893 from the mountain Zeppelinfjellet, Norwegian Polar Institute,
894 <https://doi.org/10.21334/npolar.2022.d6d31f5b>, 2022

895 Gjermundsen, A., Seland Graff, L., Bentsen, M., Anders Breivik, L., Boldingh Debernard, J.,
896 Makkonen, R., Oliivié, D.J.L., Seland, Ø., Zieger, P., Schulz, M.: How representative is Svalbard for
897 future Arctic climate evolution? An Earth system modelling perspective (SvalCLIM), SESS Report
898 2020 – The State of Environmental Science in Svalbard, <https://doi.org/10.5281/zenodo.4034104>,
899 2020

900 Hansen, B.B., Isaksen, K., Benestad, R.E., Kohler, J., Pedersen, Å., Loe, L.E., Coulson, S.J.,
901 Larsen, J.O., Varpe, Ø.: Warmer and wetter winters: Characteristics and implications of an extreme
902 weather event in the High Arctic, *Environ. Res. Lett.*, 9, <https://doi.org/10.1088/1748-9326/9/11/114021>, 2014

ha eliminato: <https://doi.org/10.1016/j.scitotenv.2023.168401>
1...

ha eliminato: <https://doi.org/10.5194/tc-16-981-2022>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1080/01621459.1961.10482090>
90...

ha formattato: Colore carattere: Testo 1

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1029/2008GL035741>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.atmosenv.2021.118458>
8...

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.scitotenv.2019.136089>
9...

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1021/es903268j>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1021/acs.est.0c02256>

ha eliminato: 10.21334/npolar.2022.d6d31f5b

ha eliminato: <https://doi.org/10.5281/zenodo.4034104>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1088/1748-9326/9/11/114021>...

ha formattato: Inglese (Regno Unito)

ha eliminato: 26¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo),
Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A
destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni:
8,5 cm, Allineato al centro + 17 cm, Allineato a destra

920 Hodgkins, R., Tranter, M.: Solute in High Arctic glacier snow cover and its impact on runoff
 921 chemistry, *Ann. Glaciol.*, 26, 156-160, <https://doi.org/10.3189/1998AoG26-1-156-160>, 1998

922 Jacobi, H.-W., Obleitner, F., Da Costa, S., Ginot, P., Eleftheriadis, K., Aas, W., Zanatta, M.:
 923 Deposition of ionic species and black carbon to the Arctic snowpack: combining snow pit
 924 observations with modeling, *Atmos. Chem. Phys.*, 19, 10361-10377,
 925 <https://doi.org/10.3402/tellusb.v65i0.19071>, 2019

926 Jung, J., Furutani, H., Uematsu, M., Park, J.: Distributions of atmospheric non-sea-salt
 927 sulphate and methanesulfonic acid over the Pacific Ocean between 48°N and 55°S during summer,
 928 *Atmos. Environ.*, 99, 374-384, <https://doi.org/10.1016/j.atmosenv.2014.10.009>, 2014

929 Koziol, K., Uszczyk, A., Pawlak, F., Frankowski, M., Polkowska, Ż.: Seasonal and Spatial
 930 Differences in Metal and Metalloid Concentrations in the Snow Cover of Hansbreen, Svalbard, *Front.*
 931 *Earth Sci.*, 8, 1-8, <https://doi.org/10.3389/feart.2020.538762>, 2021

932 Lai, A.M., Shafer, M.M., Dibb, J.E., Polashenski, C.M., Schauer, J.J.: Elements and inorganic
 933 ions as source tracers in recent Greenland snow, *Atmos. Environ.*, 164, 205-215,
 934 <https://doi.org/10.1016/j.atmosenv.2017.05.048>, 2017

935 Lawrence, Z.D., Perlwitz, J., Butler, A.H., Manney, G.L., Newman, P.A., Lee, S.H., Nash,
 936 E.R.: The Remarkably Strong Arctic Stratospheric Polar Vortex of Winter 2020: Links to Record-
 937 Breaking Arctic Oscillation and Ozone Loss, *J. Geophys. Res. Atmos.*, 125, 1-21,
 938 <https://doi.org/10.1029/2020JD033271>, 2020

939 Maffezzoli, N., Spolaor, A., Barbante, C., Bertò, M., Frezzotti, M., Vallelonga, P.: Bromine,
 940 iodine and sodium in surface snow along the 2013 Talos Dome-GV7 traverse (northern Victoria Land,
 941 East Antarctica), *Cryosphere*, 11, 693-705, <https://doi.org/10.5194/tc-11-693-2017>, 2017

942 Millero, F.J., Feistel, R., Wright, D.G., McDougall, T.J.: The composition of Standard
 943 Seawater and the definition of the Reference-Composition Salinity Scale, *Deep-Sea Res.*, 55, 50-72,
 944 2008

945 Morales, J.A., Pirela, D., de Nava, M.G., de Borrego, B.S., Velásquez, H., Durán, J.: Inorganic
 946 water soluble ions in atmospheric particles over Maracaibo Lake Basin in the western region of
 947 Venezuela, *Atmos. Res.*, 46, [https://doi.org/10.1016/S0169-8095\(97\)00071-9](https://doi.org/10.1016/S0169-8095(97)00071-9), 1998

948 Nawrot, A.P., Migala, K., Luks, B., Pakszys, P., Głowacki, P.: Chemistry of snow cover and
 949 acidic snowfall during a season with a high level of air pollution on the Hans Glacier, Spitsbergen,
 950 *Polar Sci.*, 10, 249-261, <https://doi.org/10.1016/j.polar.2016.06.003>, 2016

951 Platt, S.M., Hov, Ø., Berg, T., Breivik, K., Eckhardt, S., Eleftheriadis, K., Evangeliou, N.,
 952 Fiebig, M., Fisher, R., Hansen, G., Hansson, H.-C., Heintzenberg, J., Hermansen, O., Heslin-Rees,
 953 D., Holmén, K., Hudson, S., Kallenborn, R., Krejci, R., Krognes, T., Larssen, S., Lowry, D., Lund
 954 Myhre, C., Lunder, C., Nisbet, E., Nizzetto, P.B., Park, K.-T., Pedersen, C.A., Pfaffhuber, K.A.,
 955 Röckmann, T., Schmidbauer, N., Solberg, S., Stohl, A., Ström, J., Svendby, T., Tunved, P., Tørnkvist,
 956 K., van der Veen, C., Vratolis, S., Yoon, Y.L., Yttri, K.E., Zieger, P., Aas, W., Tørseth, K.:
 957 Atmospheric composition in the European Arctic and 30 years of the Zeppelin Observatory, Ny-
 958 Ålesund, *Atmos. Chem. Phys.*, 22, 3321-3369, <https://doi.org/10.5194/acp-22-3321-2022>, 2022

959 Pomeroy, J.W.: A process-based model of snow drifting, *Ann. Glaciol.*, 13, 237-240,
 960 <https://doi.org/10.3189/S0260305500007965>, 1989

ha eliminato: <https://doi.org/10.3189/1998AoG26-1-156-160>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3402/tellusb.v65i0.19071>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.atmosenv.2014.10.009>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3389/feart.2020.538762>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.atmosenv.2017.05.048>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1029/2020JD033271>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/tc-11-693-2017>

ha formattato: Inglese (Regno Unito)

ha eliminato: [https://doi.org/10.1016/S0169-8095\(97\)00071-9](https://doi.org/10.1016/S0169-8095(97)00071-9)

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.polar.2016.06.003>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-22-3321-2022>

ha eliminato: <https://doi.org/10.3189/S0260305500007965>

ha eliminato: 27¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

Rankin, A.M., Wolff, E.W.: Frost flowers: Implications for tropospheric chemistry and ice core interpretation, *J. Geophys. Res.*, 107, D23, 4683, <https://doi.org/10.1029/2002JD002492>, 2002

Rantanen, M., Karpechko, A.Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022

Reimann, C., De Caritat, P.: Intrinsic flaws of element Enrichment Factors (EFs) in environmental geochemistry, *Environ. Sci. Technol.*, 34, <https://doi.org/10.1021/es001339o>, 2000

Rinke, A., Maturilli, M., Graham, R.M., Matthes, H., Handorf, D., Cohen, L., Hudson, S.R., Moore, J.C.: Extreme cyclone events in the Arctic: Wintertime variability and trends, *Environ. Res. Lett.*, 12, <https://doi.org/10.1088/1748-9326/aa7def>, 2017

Roscoe, H.K., Brooks, B., Jackson, A.V., Smith, M.H., Walker, S.J., Obbard, R.W., Wolff, E.W.: Frost flowers in the laboratory: Growth, characteristics, aerosol, and the underlying sea ice, *J. Geophys. Res.*, 116, D12301, <https://doi.org/10.1029/2010JD015144>, 2011

Ruppel, M.M., Khedr, M., Liu, X., Beaudon, E., Szidat, S., Tunved, P., Ström, J., Koponen, H., Sippula, O., Isaksson, E., Gallet, J.-C., Hermanson, M., Manninen, S., Schnelle-Kreis, J.: Organic compounds, radiocarbon, trace elements and atmospheric transport illuminating sources of elemental carbon in a 300-year Svalbard ice core, *J. Geophys. Res.: Atmos.*, 128, <https://doi.org/10.1029/2022JD038378>, 2023

Salzano, R., Cerrato, R., Scoto, F., Spolaor, A., Valentini, E., Salvatore, M., Esposito, G., Sapio, S., Taramelli, A., Salvatori, R.: Detection of winter heat wave impact on surface runoff in a periglacial environment (Ny-Ålesund, Svalbard), *Remote Sens.*, 15(18), 4435, <https://doi.org/10.3390/rs15184435>, 2023

Schoeberl, M.R., Newman, P.A.: *Middle Atmosphere: Polar Vortex*, Encyclopedia of Atmospheric Sciences, Second Edition, Elsevier, 12-17, <https://doi.org/10.1016/B978-0-12-382225-3.00228-0>, 2015

Schwikowski, M., Döschner, A., Gäggeler, H.W., Schotterer, U.: Anthropogenic versus natural sources of atmospheric sulphate from an Alpine ice core. *Tellus B: Chemical and Physical Meteorology*, 51:5, 938-951, <https://doi.org/10.3402/tellusb.v51i5.16506>, 1999

Scoto, F., Pappaccogli, G., Mazzola, M., Donato, A., Salzano, R., Monzali, M., de Blasi, F., Larose, C., Gallet, J.-C., Decesari, S., Spolaor, A.: Automated observation of physical snowpack properties in Ny-Ålesund, *Front. Earth Sci.*, 11:1123981, 1-9, <https://doi.org/10.3389/feart.2023.1123981>, 2023

Serreze, M.C., Barry, R.G.: Processes and impacts of Arctic amplification: A research synthesis, *Glob. Planet. Change*, 77, 85-96, <http://dx.doi.org/10.1016/j.gloplacha.2011.03.004>, 2011

Sherrell, R.M., Boyle, E.A., Harris, N.R., Falkner, K.K.: Temporal variability of Cd, Pb, and Pb isotope deposition in central Greenland snow, *Geochem Geophys*, 1 (1), 1-22, <https://doi.org/10.1029/1999GC000007>, 2000

Sobota, I., Weckwerth, P., Grajewski, T.: Rain-On-Snow (ROS) events and their relations to snowpack and ice layer changes on small glaciers in Svalbard, the high Arctic, *J. Hydrol*, 590, 125279, <https://doi.org/10.1016/j.jhydrol.2020.125279>, 2020

ha eliminato: <https://doi.org/10.1029/2002JD002492>

ha eliminato: <https://doi.org/10.1038/s43247-022-00498-3>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1021/es001339o>

ha formattato: Colore carattere: Testo 1

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1088/1748-9326/aa7def>

ha eliminato: <https://doi.org/10.1029/2010JD015144>

ha eliminato: <https://doi.org/10.1029/2022JD038378>

ha eliminato: <https://doi.org/10.3390/rs15184435>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/B978-0-12-382225-3.00228-0>

ha eliminato: <https://doi.org/10.3402/tellusb.v51i5.16506>

ha formattato: Legature: Nessuna

ha formattato: Inglese (Regno Unito)

ha eliminato: <http://dx.doi.org/10.1016/j.gloplacha.2011.03.004>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1029/1999GC000007>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.jhydrol.2020.125279>

ha eliminato: 28¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

1029 Song, C., Becagli, S., Beddows, D.C.S., Brean, J., Browse, J., Dai, Q., Dall'Osto, M., Ferracci,
1030 V., Harrison, R.M., Harris, N., Li, W., Jones, A.E., Kirchg   ner, A., Kramawijaya, A.G., Kurganskiy,
1031 A., Lupi, A., Mazzola, M., Severi, M., Traversi, R., Shi, Z.: Understanding sources and drivers of
1032 size-resolved aerosol in the High Arctic islands of Svalbard using a receptor model coupled with
1033 machine learning, *Environ. Sci. Technol.*, 56, 11189-11198, <https://doi.org/10.1021/acs.est.1c07796>,
1034 2022.

ha formattato: Inglese (Regno Unito)

1035 Song, J., Zhao, Y., Zhang, Y., Fu, P., Zheng, L., Yuan, Q., Wang, S., Huang, X., Xu, W., Cao,
1036 Z., Gromov, S., Lai, S.: Investigation of biomass burning on atmospheric aerosols over the western
1037 South China Sea: Insights from ions, carbonaceous fractions and stable carbon isotope ratios,
1038 Environ. Pollut., 242, 1800-1809, <https://doi.org/10.1016/j.envpol.2018.07.088>, 2018

ha eliminato: <https://doi.org/10.1021/acs.est.1c07796>, 2022

ha formattato: Inglese (Regno Unito)

1039 Spolaor, A., Scoto, F., Larose, C., Barbaro, E., Burgay, F., Bjorkman, M., Cappelletti, D.,
1040 Dallo, F., de Blasi, F., Divine, D., Dreossi, G., Gabrieli, J., Isaksson, E., Kohler, J., Martma, T.,
1041 Schmidt, L.S., Schuler, T.V., Stenni, B., Turetta, C., Luks, B., Casado, M., Gallet, J.-C.: Climate
1042 change is rapidly deteriorating the climatic signal in Svalbard glaciers, *Cryosphere*, 18, 307-320,
1043 <https://doi.org/10.5194/tc-18-307-2024>

ha eliminato: <https://doi.org/10.1016/j.envpol.2018.07.088>

1044 Spolaor, A., Varin, C., Pedeli, X., Christille, J.M., Kirchgeorg, T., Giardi, F., Cappelletti, D.,
1045 Turetta, C., Cairns, W.R.L., Gambaro, A., Bernagozzi, A., Gallet, J.C., Björkman, M.P., Barbaro, E.:
1046 Source, timing and dynamics of ionic species mobility in the Svalbard annual snowpack, *Sci. Total*
1047 *Environ.*, 751, 141640, <https://doi.org/10.1016/j.scitotenv.2020.141640>, 2021b

ha eliminato: <https://doi.org/10.5194/tc-18-307-2024>

Spolaor, A., Moroni, B., Luks, B., Nawrot, A., Roman, M., Larose, C., Stachnik, Ł., Bruschi, F., Koziol, K., Pawlak, F., Turetta, C., Barbaro, E., Gallet, J.-C., Cappelletti, D.: Investigation on the Sources and Impact of Trace Elements in the Annual Snowpack and the Firn in the Hansbreen (Southwest Spitsbergen). *Front. Earth Sci.*, 8, 1–10, <https://doi.org/10.3389/feart.2020.536036>, 2021a

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.scitotenv.2020.141640>

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

Spolaor, A., Barbaro, E., Cappelletti, D., Turetta, C., Mazzola, M., Giardi, F., Björkman, M.P.,
Lucchetta, F., Dallo, F., Aspö, P., Pfaffhuber, K., Angot, H., Dommergue, A., Maturilli, M., Saiz-
Lopez, A., Barbante, C., Cairns, W.R.L.: Diurnal cycle of iodine, bromine, and mercury
concentrations in Svalbard surface snow, *Atmos. Chem. Phys.*, 19, 13325–13339,
<https://doi.org/10.5194/acp-19-13325-2019>, 2019

ha eliminato: <https://doi.org/10.3389/feart.2020.536036>

ha formattato: Inglese (Regno Unito)

1057 Spolaor, A., Angot, H., Roman, M., Dommergue, A., Scarchilli, C., Vardè, M., Del Guasta,
1058 M., Pedeli, X., Varin, C., Sprovieri, F., Magand, O., Legrand, M., Barbante, C., Cairns, W.R.L:
1059 Feedback mechanisms between snow and atmospheric mercury: Results and observations from field
1060 campaigns on the Antarctic plateau, *Chemosphere*, 197, 306–317,
1061 <https://doi.org/10.1016/j.chemosphere.2017.12.180>, 2018

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.1016/j.chemosphere.2017.12.180>...

ha formattato: Inglese (Regno Unito)

Spolaor, A., Opel, T., McConnell, J.R., Maselli, O.J., Spreen, G., Varin, C., Kirchgeorg, T.,
Fritzsche, D., Saiz-Lopez, A., Vallenga, P.: Halogen-based reconstruction of Russian Arctic sea ice
area from the Akademii Nauk ice core (Severnaya Zemlya), Cryosphere, 10, 245-256,
<https://doi.org/10.5194/tc-10-245-2016>, 2016

ha eliminato: <https://doi.org/10.5194/tc-10-245-2016>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-14-9613-2014>

1066 Spolaor, A., Vallelonga, P., Gabrieli, J., Martma, T., Björkman, M.P., Isaksson, E., Cozzi, G.,
1067 Turetta, C., Kjær, H.A., Curran, M.A.J., Moy, A.D., Schönhardt, A., Blechschmidt, A.-M., Burrows,
1068 J.P., Plane, J.M.C., Barbante, C.: Seasonality of halogen deposition in polar snow and ice, *Atmos.*
1069 *Chem. Phys.*, 14, 18, 9613-9622, <https://doi.org/10.5194/acp-14-9613-2014>, 2014.

ha formattato: Inglese (Regno Unito)

ha eliminato: 29

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8.5 cm. Allineato al centro + 17 cm. Allineato a destra

1081 Stein, A.F., Draxler, R.R., Rolph, G.D., Stunder, B.J.B., Cohen, M.D., Ngan, F.: Noaa's
 1082 hysplit atmospheric transport and dispersion modeling system, Bull. Am. Meteorol. Soc., 96, 2059–
 1083 2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015

1084 Stohl, A., Berg, T., Burkhardt, J.F., Fjæraa, a. M., Forster, C., Herber, A., Hov, Ø., Lunder, C.,
 1085 McMillan, W.W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth,
 1086 K., Treffeisen, R., Virkkunen, K., Yttri, K.E.: Arctic smoke – record high air pollution levels in the
 1087 European Arctic due to agricultural fires in Eastern Europe, Atmos. Chem. Phys. Discuss., 6, 9655–
 1088 9722, <https://doi.org/10.5194/acpd-6-9655-2006>, 2006b

1089 Stohl, A., Berg, T., Burkhardt, J.F., Fjæraa, a. M., Forster, C., Herber, A., Hov, Ø., Lunder, C.,
 1090 McMillan, W.W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth,
 1091 K., Treffeisen, R., Virkkunen, K., Yttri, K.E.: Arctic smoke – record high air pollution levels in the
 1092 European Arctic due to agricultural fires in Eastern Europe, Atmos. Chem. Phys. Discuss., 6, 9655–
 1093 9722, <https://doi.org/10.5194/acpd-6-9655-2006>, 2006a

1094 Turetta, C., Feltracco, M., Barbaro, E., Spolaor, A., Barbante, C., Gambaro, A.: A year-round
 1095 measurement of water-soluble trace and rare earth elements in arctic aerosol: Possible inorganic
 1096 tracers of specific events, Atmosphere, 12, <https://doi.org/10.3390/atmos12060694>, 2021

1097 Udisti, R., Traversi, R., Becagli, S., Tomasi, C., Mazzola, M., Lupi, A., Quinn, P.K.: Arctic
 1098 Aerosols, in: Physics and Chemistry of the Arctic Atmosphere, Springer Polar Sciences, edited by:
 1099 Kokhanovsky, A., Tomasi, C., 209-329, https://doi.org/10.1007/978-3-030-33566-3_4, 2020

1100 Udisti, R., Bazzano, A., Becagli, S., Bolzacchini, E., Caiazzo, L., Cappelletti, D., Ferrero, L.,
 1101 Frosini, D., Giardi, F., Grotti, M., Lupi, A., Malandrino, M., Mazzola, M., Moroni, B., Severi, M.,
 1102 Traversi, R., Viola, A., Vitale, V.: Sulphate source apportionment in the Ny-Ålesund (Svalbard
 1103 Islands) Arctic aerosol, Rend. Fis. Acc. Lincei, 27, Suppl 1: S85-S94, <https://doi.org/10.1007/s12210-016-0517-7>, 2016

1105 Van Hecke, T.: Power study of anova versus Kruskal-Wallis test, J. Stat. Manag. Syst., 15, 2-
 1106 3, <https://doi.org/10.1080/09720510.2012.10701623>, 2013

1107 Vecchiato, M., Barbante, C., Barbaro, E., Burgay, F., Cairns, W.R.L., Callegaro, A.,
 1108 Cappelletti, D., Dallo, F., D'Amico, M., Feltracco, M., Gallet J.-C., Gambaro, A., Larose, C.,
 1109 Maffezzoli, N., Mazzola, M., Sartorato, I., Scoto, F., Turetta, C., Vardè, M., Xie, Z., Spolaor, A.: The
 1110 seasonal change of PAHs in Svalbard surface snow, Environ. Pollut., 340, 122864,
 1111 <https://doi.org/10.1016/j.envpol.2023.122864>, 2024

1112 Vecchiato, M., Barbaro, E., Spolaor, A., Burgay, F., Barbante, C., Piazza, R., Gambaro, A.:
 1113 Fragrances and PAHs in snow and seawater of Ny-Ålesund (Svalbard): Local and long-range
 1114 contamination, Environ. Pollut., 242, 1740-1747, <https://doi.org/10.1016/j.envpol.2018.07.095>, 2018

1115 Vega, C.P., Björkman, M.P., Pohjola, V.A., Isaksson, E., Pettersson, R., Martma, T., Marca,
 1116 A., Kaiser, J., Vega, C.P., Björkman, M.P., Pohjola, V.A., Isaksson, E., Pettersson, R., Martma, T.,
 1117 Marca, A., Kaiser, J., Pohjola, V.A., Isaksson, E., Pettersson, R., Vega, C.P., Bjo, M.P.: Nitrate stable
 1118 isotopes and major ions in snow and ice samples from four Svalbard sites Nitrate stable isotopes and
 1119 major ions in snow and ice samples from four Svalbard sites, Polar Res., 2015, 34, 23246,
 1120 <https://doi.org/10.3402/polar.v34.23246>, 2015

ha eliminato: <https://doi.org/10.1175/BAMS-D-14-00110.1>

ha formattato: Inglese (Regno Unito)

ha spostato (inserimento) [3]

ha formattato: Inglese (Regno Unito)

Formattato: Regola lo spazio tra testo asiatico e in alfabeto latino, Regola lo spazio tra caratteri asiatici e numeri

ha eliminato: <https://doi.org/10.5194/acpd-6-9655-2006>, 2006b

ha eliminato: <https://doi.org/10.5194/acpd-6-9655-2006>

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3390/atmos12060694>

ha eliminato: https://doi.org/10.1007/978-3-030-33566-3_4

ha eliminato: <https://doi.org/10.1007/s12210-016-0517-7>

ha eliminato: <https://doi.org/10.1080/09720510.2012.10701623>, 2013...

ha eliminato: <https://doi.org/10.1016/j.envpol.2023.122864>

ha eliminato: <https://doi.org/10.1016/j.envpol.2018.07.095>

ha formattato: Inglese (Regno Unito)

ha formattato: Motivo: Trasparente, Evidenziato

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.3402/polar.v34.23246>

ha eliminato: 30

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

Wagenbach, D., Preunkert, S., Schäfer, J., Jung, W., Tomadin, L.: Northward transport of Saharan dust recorded in a deep alpine ice core, in: The impact of desert dust across the mediterranean, Springer, The Netherlands, edited by: Guerzoni, S., Chester, R., 291-300, 1996

Wedepohl, K.H.: The composition of the chemical crust. Geoch. Cosm. Act., 59(7), 1217-1232, [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2), 1995

Yttri, K.E., Lund Myhre, C., Eckhardt, S., Fiebig, M., Dye, C., Hirdman, D., Ström, J., Klimont, Z., Stohl, A.: Quantifying black carbon from biomass burning by means of levoglucosan - A one-year time series at the Arctic observatory Zeppelin, Atmos. Chem. Phys., 14, 6427–6442, <https://doi.org/10.5194/acp-14-6427-2014>, 2014a

Zhan, J., Gao, Y., Li, W., Chen, L., Lin, H., Lin, Q.: Effects of ship emissions on summertime aerosols at Ny-Alesund in the Arctic, Atmos. Pollut. Res., 5, 500–510, <https://doi.org/10.5094/APR.2014.059>, 2014

Zhuang, H., Chan, C.K., Fang, M., Wexler, A.S.: Size distributions of particulate sulphate, nitrate, and ammonium at a coastal site in Hong Kong, Atmos. Environ., 33, 843-853, 1999

ha eliminato: [https://doi.org/10.1016/0016-7037\(95\)00038-2](https://doi.org/10.1016/0016-7037(95)00038-2)

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5194/acp-14-6427-2014>

ha formattato: Inglese (Regno Unito)

ha eliminato: <https://doi.org/10.5094/APR.2014.059>

ha formattato: Inglese (Regno Unito)

ha eliminato: 31

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

Pagina 9: [1] ha eliminato	azzurra spagnesi	27/08/25 15:52:00
----------------------------	------------------	-------------------

Pagina 1: [2] Formattato	azzurra spagnesi	27/08/25 15:52:00
--------------------------	------------------	-------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 8,5 cm, Allineato al centro + 17 cm, Allineato a destra

Pagina 10: [3] ha formattato	azzurra spagnesi	27/08/25 15:52:00
------------------------------	------------------	-------------------

Colore carattere: Automatico, Inglese (Regno Unito)

Pagina 19: [4] ha eliminato	azzurra spagnesi	27/08/25 15:52:00
-----------------------------	------------------	-------------------