

<https://doi.org/10.5194/egusphere-2024-1422>

Preprint. Discussion started: 3 June 2024

© Author(s) 2024. CC BY 4.0 License.



Impact of methane and other precursor emission reductions on surface ozone in Europe: Scenario analysis using the EMEP MSC-W model

Willem E. van Caspel¹, Zbigniew Klimont², Chris Heyes², and Hilde Fagerli¹

¹EMEP MSC-W, Norwegian Meteorological Institute, Oslo, Norway

²International Institute for Applied Systems Analysis, Laxenburg, Austria

Correspondence: W.E. van Caspel (willemvc@met.no)



Abstract.

The impacts of future methane (CH_4) and other precursor emission changes are investigated for surface ozone (O_3) in the United Nations Economic Commission for Europe (UNECE) region excluding North America and Israel (the “EMEP region”, for European Monitoring and Evaluation Programme) for the year 2050. The analysis includes a Current Legislation (CLE) and Maximum Feasible Technical (MFR) reduction scenario, and a scenario that combines MFR reductions with an additional dietary shift that also meets the Paris Agreement objectives with respect to greenhouse gas emissions (LOW). For each scenario, background CH_4 concentrations are calculated using a probabilistic Earth System model emulator, and combined with other precursor emissions in a three-dimensional Eulerian chemistry-transport model. While focus is placed on peak season maximum daily 8-hour average (MDA8) O_3 concentrations, a range of other indicators for health and vegetation impacts are also discussed. Our analysis show that roughly one-thirds of the total peak season MDA8 reduction achieved between the 2050 CLE and MFR scenarios is attributable to CH_4 reductions, resulting predominantly from CH_4 emission reductions outside of the EMEP region. The impact of other precursor emission reductions is split nearly evenly between the reductions inside and outside of the EMEP region. However, the relative importance of CH_4 and other precursor emission reductions is shown to depend on the choice of O_3 indicator, though indicators sensitive to peak O_3 show generally consistent results. The analysis also highlights the synergistic impacts of CH_4 mitigation as reducing solely CH_4 achieves, beyond air quality improvement, nearly two-thirds of the total global warming reduction calculated for the LOW scenario compared to the CLE case.



Plain language summary

Methane in the atmosphere contributes to the production of ozone gas, which is an air pollutant as well as a greenhouse gas.

- 20 In this study, the impact of reducing methane emissions on surface ozone is investigated for the United Nations Economic Commission for Europe (UNECE) region excluding North America and Israel (the “EMEP region”), in particular in terms of its importance in reaching the ozone exposure guideline limits set by the World Health Organization. The relative importance of reducing emissions of other pollutants that lead to the formation of ozone, such as nitrogen oxides, is also investigated. To this end, our study employs emission scenarios up to the year 2050, each having different assumptions about future human-
- 25 caused emissions. Relative to a scenario where only the already agreed emission reductions are implemented, one-third of surface ozone reductions in the most ambitious emission reduction scenario are attributable to methane reductions. The other two-thirds are attributable to emission reductions of other ozone forming pollutants, with reductions inside and outside of the EMEP region contributing roughly equally.



1 Introduction

30 Surface ozone (O_3) is an important source of air pollution, impacting both human and ecosystem health (Lefohn et al., 2018; Monks et al., 2015). In the lower troposphere, the majority of O_3 is produced by the photochemical reaction of nitrogen oxides ($NO_x = NO + NO_2$) in carbonaceous-rich environments (Crutzen et al., 1999). The most abundant carbonaceous O_3 precursor species is methane (CH_4), having a present-day volume mixing ratio of around 1915 parts-per-billion (ppb). Moreover, CH_4 mixing ratios are likely to increase further, as anthropogenic CH_4 emissions are anticipated to increase in the coming decade
35 (UNEP, 2021; Saunio et al., 2020; Höglund-Isaksson et al., 2020). In addition to being a source of air pollution, CH_4 is also the second most important anthropogenic greenhouse gas (GHG), with its importance as both an air pollutant and global warming agent having received considerable attention in recent years (Mar et al., 2022; Abernethy et al., 2021; Fiore et al., 2008; Dentener et al., 2005).

In this study, the impact of CH_4 and other precursor emissions is investigated for the European Monitoring and Evaluation
40 Programme (EMEP) region, which includes the member countries of the United Nations Economic Commission for Europe (UNECE) region excluding North America and Israel. Focus is placed on the population-weighted exposure to peak season (April-September) average maximum daily 8-hour mean (MDA8) O_3 concentrations, being the health indicator employed by the new World Health Organization (WHO) guidelines (WHO, 2021). The latter recommends a peak season MDA8 exposure limit of $60 \mu g m^{-3}$ based on the association between long-term O_3 exposure and all-cause mortality, with an interim target of
45 $70 \mu g m^{-3}$ for areas where initial exposure is high. To our knowledge, neither the guideline nor interim target values are met in any of the countries within the EMEP region at present. The focus on peak season MDA8 is also motivated by the broader association between the exposure to peak O_3 and all-cause mortality (Huangfu and Atkinson, 2020).

In the current work, the impacts on O_3 are investigated for a Current Legislation (CLE), Maximum technical Feasible Reduction (MFR), and MFR with an additional dietary shift and Paris Agreement policy scenario (LOW) up to the year 2050.
50 The CLE scenario includes the currently agreed upon policies for the abatement of air pollutant and GHG emissions, while the MFR scenario combines the economic activity pathway of the CLE scenario with the full implementation of the best available emission reduction technologies defined in the GAINS (Greenhouse gas – Air pollution Interactions and Synergies) model (Amann et al., 2011). The LOW scenario extends the MFR by including climate policies compatible with the Paris Agreement objectives and an additional shift in agricultural practices, bringing further CH_4 and precursor emission reductions. Relative
55 to the year 2015, global anthropogenic CH_4 emissions decline by 35 % and 50 % in the LOW scenario by 2030 and 2050, respectively, making the reductions comparable to those of the Methane Pledge (30 % by 2030, Malley et al., 2023) and Global Methane Assessment (45 % abatement target for 2050, UNEP, 2021).

The emission scenarios are combined with the Model for the Assessment of Greenhouse-gas Induced Climate Change v7.5.3 (MAGICC7) (Meinshausen et al., 2020, 2011, 2009) to calculate their respective background CH_4 concentrations up to the
60 year 2050. To calculate their impacts on surface O_3 , the CH_4 projections are specified in the three-dimensional Eulerian Chemistry-Transport Model (CTM) developed at the EMEP Meteorological Synthesising Centre – West (hereafter “EMEP model”), where they are combined with the other precursor scenario emissions. The EMEP model has a long history of policy



support and research development (e.g., Jonson et al., 2018; Simpson, 2013; Simpson et al., 2012), with one of its main tasks being the modeling of transboundary fluxes of air pollutants as part of the UNECE Convention on Long-range Transboundary Air Pollution (CLRTAP) (Fagerli et al., 2023). In this capacity, the EMEP model has previously been used in support of the review of the UNECE Gothenburg Protocol (Protocol to Abate Acidification, Eutrophication and Ground-level Ozone).

The emission scenarios and their implementation are described in more detail Sect. 2. The MAGICC7 model is described in Sect. 3, where it is used to calculate background CH_4 concentrations up to the year 2050. Sect. 4 describes the EMEP model configuration, while also evaluating the baseline configuration against five years of observations across Europe. For the scenario calculations presented in Sect. 5, the default modeling configuration involves averaging all results over five meteorological years, while a linear latitudinal CH_4 gradient is imposed to capture the effects of inter-hemispheric variations in emissions. Sect. 5 further combines regional EMEP model simulations with global simulations to quantify the separate impacts of emission changes inside and outside of the EMEP region. While focus is placed on the peak season MDA8 indicator, scenario results for a range of other O_3 health and vegetation indicators are also presented. The results are discussed and compared against earlier studies in Sect. 6, followed by a conclusion in Sect. 7.



Table 1. Global emission totals for the CLE, MFR, and LOW emission scenarios in units of Tg yr⁻¹. Emission totals within the EMEP region, as defined in Sect. 1, are listed in brackets for the years 2015, 2030, and 2050.

Species	Scenario	2015	2020	2025	2030	2035	2040	2045	2050
NO _x	CLE	119 (16)	111	106	103 (10)	102	103	104	106 (9)
NO _x	MFR				65 (6)	52	42	40	38 (4)
NO _x	LOW				62 (6)	46	33	29	25 (3)
NMVOC	CLE	121 (15)	120	120	118 (14)	119	119	120	121 (14)
NMVOC	MFR				68 (10)	63	59	59	59 (8)
NMVOC	LOW				63 (9)	57	52	50	48 (8)
CO	CLE	517 (50)	474	449	427 (37)	418	411	408	405 (43)
CO	MFR				160 (22)	139	123	123	123 (18)
CO	LOW				149 (22)	124	102	96	91 (17)
CH ₄	CLE	334 (62)	345	360	371 (60)	385	401	416	428 (60)
CH ₄	MFR				229 (28)	224	226	220	210 (22)
CH ₄	LOW				219 (27)	208	202	195	168 (14)

2 Emissions

The emission scenarios were developed using the global version of the GAINS model (Winiwarter et al., 2018; Klimont et al., 2017; Höglund-Isaksson, 2012; Amann et al., 2011), and provided by the EMEP Center for Integrated Assessment Modelling (CIAM) hosted by the Institute for Applied Systems Analysis (IIASA). The scenarios include annual anthropogenic emission
80 totals of CH₄, NO_x, Non-Methane Volatile Organic Compounds (NMVOC), carbon monoxide (CO), sulphur oxides (SO_x), ammonia (NH₃), primary fine Particulate Matter (PM_{2.5}), and primary coarse PM (PM_{co}), as well as carbonaceous fraction of primary PM represented by black carbon (BC) and organic carbon (OC). For the current work, the key emission species are CH₄, NO_x, CO, and NMVOC, where the latter three affect the lifetime of CH₄ by acting as either net sources (NO_x) or
85 sinks (CO and NMVOC) of hydroxyl (OH). The latter in turn affects the lifetime of CH₄ by loss against oxidation. The global emission totals for the key species are shown in Table 1, along with their respective emissions within the EMEP region for the years 2015, 2030, and 2050.

The emission scenarios span the period from the baseline year 2015 up to 2050 in 5-year intervals, with the MFR and LOW scenarios diverging from the CLE scenario from 2025 onwards. In the EMEP model, natural emissions of soil NO_x are included based on monthly climatological values from the CAMS-GLOB-SOIL v2.4 inventory (Simpson and Darras, 2021), noting that
90 soil NO_x emissions from the application of manures and mineral nitrogen fertilizers on agricultural land are calculated in the GAINS model. Forest fire emissions are included based on the daily Fire INventory from NCAR version 2.5 (FINNV2.5, Wiedinmyer et al., 2023) dataset, derived from fire detections from both the Moderate Resolution Imaging Spectroradiometer (MODIS) and Visible Infrared Imaging Radiometer Suite (VIIRS) satellite instruments.



2.1 Emission scenarios

95 2.1.1 CLE scenario

The CLE scenario assumes the implementation and effective enforcement of all currently committed energy and environmental policies affecting emissions of air pollutants and greenhouse gases. CIAM has undertaken a review and update of historical data (up to 2020) driving emissions of all species in the GAINS model, drawing on information from the statistical office of the European Union (EUROSTAT), International Energy Agency (IEA), UN Food and Agriculture Organization (FAO), in
100 addition to data and emissions reported to the Center on Emission Inventories and Projections (CEIP). For the EU27 countries, the energy and agriculture projections are consistent with the objectives of the European Green Deal and ‘Fit for 55’ package to make the EU carbon neutral by 2050, while also being consistent with the projections used in the EU 3rd Clean Air Outlook (https://environment.ec.europa.eu/topics/air/clean-air-outlook_en, last access: April 2024). For West Balkan, Republic of
105 Moldova, Georgia, and Ukraine, a similar set of modelling tools was used as for the EU, developing a new consistent set of projections. For other world regions, the GAINS model down-scales projections from IEA and FAO (Alexandratos and Bruinsma, 2012; IEA, 2018), considering updated air pollution legislation from national and international sources (e.g., He et al., 2021; Zhang, 2018), including EU legislation and their implementation in consultation with the EU Member States. For the
110 CLE scenario, the socio-economic activity assumptions are similar to that of the Shared Socioeconomic Pathway 2 with an end-of-century radiative forcing of 4.5 W m^{-2} (SSP2-4.5). The SSP2-4.5 scenario describes the ‘middle of the road’ for future societal development, as described in Meinshausen et al. (2020), O’Neill et al. (2017), and Riahi et al. (2017) for a range of SSP scenarios. For the background CH_4 calculations described in Sect. 3, the CLE scenario emissions are therefore combined with GHG emissions (e.g., CO_2 and hydrofluorocarbons) from the SSP2-4.5 scenario.

2.1.2 MFR mitigation scenario

The MFR mitigation scenario assumes the full implementation of the proven technical mitigation potential as included in the
115 GAINS model for precursor emissions (Amann et al., 2020, 2013; Rafaj et al., 2018) and CH_4 (Gomez Sanabria et al., 2022; Höglund-Isaksson et al., 2020; Höglund-Isaksson, 2012). Technologies to abate air pollution precursor emissions include, for example, end of pipe technologies applied in the power, industry, and transport sector, technology change in industry and residential combustion, as well as measures in agriculture addressing emissions from manures and mineral fertilizer application
120 by, for example, improved manure management techniques and construction low emission housing including covered manure stores. The fossil fuel and solvent sector emissions include improved flaring, maintenance, leakage, and distribution control measures, as well as low-solvent product substitutions. Global emissions of NO_x , NMVOC, and CO decline by nearly 80 % by 2050 relative to the 2015 baseline, while CH_4 emissions fall by 37 %. These reductions are driven by rapid introduction of stringent emission limit values for stationary and mobile sources, strong decline in fossil fuel use, and access to clean energy for cooking. The MFR energy and agricultural activity projections are the same as those of the CLE scenario, with the MFR
125 scenario also being combined with other GHG emissions from the SSP2-4.5 scenario.



2.1.3 LOW mitigation scenario

The LOW mitigation scenario extends the MFR by including several additional policies targeting significant transformations in the agricultural sector. This transformation leads to strong reductions of livestock numbers, especially cattle and pigs. The scenario is based in part on the ‘Growing Better report 2019’ (The Food and Land Use Coalition, 2019) and other studies
130 addressing healthy dietary requirements (Kanter et al., 2020; Willett et al., 2019), as used in earlier scenarios for global air pollution studies (Amann et al., 2020). While the LOW scenario has the same energy projections as for the CLE for EU27 countries, the rest of the world now includes climate policies compatible with Paris Agreement goals, making the GHG emissions consistent with those of the ‘taking the green road’ SSP1-2.6 scenario (Riahi et al., 2017; O’Neill et al., 2017). In the LOW scenario, global CH₄ emission decline by 34 % and 50 % by 2030 and 2050 relative to the 2015 baseline, respectively.

135 2.2 Model implementation

The annual mean national and sector (e.g., road traffic and agriculture) emission totals are distributed in time using a set of monthly, weekly, daily, and hourly time-factors based on the global and European CAMS-TEMPO datasets described in Guevara et al. (2021, 2020a, b). For the regional EMEP domain discussed in Sect. 4, the native 0.5° × 0.5° scenario emissions are redistributed to the 0.1° × 0.1° spatial distribution of the most recent EMEP reported emissions (2021) for countries within
140 the EMEP region (EMEP/CEIP, 2023). However, following the approach used for the Gothenburg Protocol review, native 0.1° × 0.1° gridded emissions from CIAM are used for countries located within the West-Balkan and Economic Co-operation and Development, Eastern Europe, Caucasus and Central Asia (EECCA) regions, and for Türkiye. Countries that lie (partially) within the regional modelling domain but that are not part of the EMEP region, such as North African countries, follow the global 0.5° × 0.5° gridded emissions. International shipping emissions also follow the global 0.5° × 0.5° spatial distribution
145 provided by CIAM for all simulations. We further note that direct emissions of CH₄ are not included in the EMEP model, with concentrations instead being specified on an annual mean basis, as discussed in Sect. 4.1.



3 Background CH₄

Earth System emulators, sometimes known as Reduced Complexity Models (RCMs), have a long history of development as low-cost alternatives to full complexity climate models. RCMs include simplified parameterizations of, for example, ocean heat uptake, GHG effective radiative forcing, and climate feedbacks, to efficiently estimate future change in climate variables such as GHG concentrations and global-mean surface air temperature (GSAT) (Nicholls et al., 2021, 2020). To this end, the MAGICC7 v7.5.3 RCM has been used in the Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report (AR6) (Forster et al., 2021), being calibrated to capture the relationship between emissions and GSAT for the AR6 historical temperature assessment (Nicholls et al., 2022). In the current work, the MAGGIC7 model is run using the 5-yearly annual totals from Table 1 linearly interpolated to annual values and combined with their respective SSP GHG scenario emissions.

In the MAGICC7 model, CH₄ sinks are represented by loss against OH in the troposphere, loss to the stratosphere, and soil uptake (Meinshausen et al., 2011). Climate sensitivities for these mechanisms arise from, for example, temperature-driven changes in atmospheric composition, changes in the Brewer-Dobson circulation strength, and changing soil properties. CH₄ sources are controlled by the separate contributions arising from anthropogenic, natural, and permafrost (Schneider von Deimling et al., 2012) emissions. Natural emissions are estimated by closing the CH₄ budget between the years 2015-2023, for which the IIASA emissions are the same for all scenarios, using observed global mean background CH₄ concentrations up to the most recent year as reference (1923 ppb by 2023, Lan et al., 2022). With this approach, natural emissions are estimated at 214.9 Tg yr⁻¹, falling within the top-down range of 194–267 Tg yr⁻¹ reported by Saunio et al. (2020) for the year 2017. The natural emissions are kept constant throughout the simulation period.

A key feature of the MAGICC7 model is that it can be run in a probabilistic mode, where the results of its 600-member ensemble reflect the uncertainties in the parameters controlling future climate change (Nicholls et al., 2022). The initial parameter values controlling the CH₄-cycle are the same for each ensemble run, however, with parameters such as the initial lifetime of CH₄ (9.95 yr⁻¹) and temperature-sensitivity of the loss against OH (0.07 K⁻¹) calibrated to match the projections by Holmes et al. (2013) across the range of Representative Concentration Pathway (RCP) scenarios (Meinshausen et al., 2020). As a result, the inter-ensemble variations for the calculated CH₄ projections represent the sensitivity of the different CH₄ source and sink terms to temperature change. We note that the net land-to-atmosphere CH₄ flux from permafrost is found to have a minimal contribution to the simulation results, with the 600-ensemble emissions falling below 4 Tg yr⁻¹ by 2050 for all scenarios.

3.1 CH₄ projections

Fig. 1 shows the CH₄ projections calculated for the CLE, MFR, and LOW scenarios, with the shaded regions indicating the 5th to 95th percentile (5-95 %) range of the 600-ensemble output. Here the CH₄ projections for the SSP1-2.6, SSP2-4.5, and SSP5-8.5 scenarios are included for reference, noting that the IIASA scenario projections fall within the range of the optimistic (SSP1-2.6) and pessimistic (SSP5-8.5) scenarios. For the CLE, MFR, and LOW scenarios, the 2050 global mean CH₄ concentrations and their 5-95 % range are calculated as 2236 [2166-2299], 1651 [1597-1700], and 1574 [1512-1627] ppb, respectively. For other years, ensemble mean CH₄ concentrations are shown in supplementary Table S1.




Figure 1. Projected background CH₄ concentrations up to 2050 for the CLE, MFR, and LOW scenarios described in the text. Projections for the SSP5-8.5, SSP2-4.5, and SSP1-2.6 scenarios are included for reference. The shaded area represents the 600-ensemble 5-95% range.

180 Fig. 1 shows that the temperature-driven MFR and LOW scenarios uncertainties partly overlap. However, the inter-scenario difference between the CLE and the MFR (and LOW) scenarios far exceeds the temperature-driven uncertainties, with the 2050 ensemble mean difference amounting to 585 ppb. In the current work, the difference between the 2050 CLE and the MFR scenarios represents an important measure of the impact of CH₄ emission changes, as this represents the largest inter-scenario concentration change. While both scenarios have a 5-95 % range of approximately 100 ppb by 2050, the 5-95 % interval of
 185 the difference between the 2050 CLE and MFR scenarios amounts to 571-598 ppb. Thus illustrating that ensemble members with a comparatively high CH₄ concentration in the CLE scenario also have a comparatively high concentration in the MFR scenario.

Diagnostic simulations for a scenario where CH₄ emissions follow the LOW scenario while all other emissions follow those of the CLE scenario (LOW-CH₄) are also performed. This hypothetical scenario thereby reflects a situation where CH₄ emis-
 190 sions are reduced strongly, while no further abatement policies are implemented for the other emissions. In reality, however, CH₄ reductions might also lead to a reduction in other co-emitted species. The resulting 2050 LOW-CH₄ concentration of 1440 [1392-1484] ppb is comparable to that of the LOW scenario, although lower by 134 ppb (-8.5 %) due to higher emissions of other lifetime-affecting precursor species. The LOW-CH₄ scenario thereby illustrates that the difference in CH₄ concentrations between the 2050 CLE and LOW scenarios (and corollary MFR) is primarily driven by the difference in the direct emissions
 195 of CH₄, and to a lesser extent by the difference in other precursor and GHG emissions. We further note that continuing the



CH₄ projections into 2055 with the emissions fixed to that of 2050 leads to an additional change in the ensemble-mean concentrations of 38 ppb (1.7 %), -45 ppb (-2.7 %), and -70 ppb (-4.45 %) for the CLE, MFR, and LOW scenarios, respectively. The latter illustrates that the CH₄ source and sink terms have not yet reached equilibrium by 2050, owing to the relatively long lifetime of CH₄.



200 4 EMEP model description

The current work uses EMEP model version rv5.0, as described in more detail by EMEP MSC-W (2023) and others (e.g., Ge et al., 2024; van Caspel et al., 2023; Stadtler et al., 2018; Simpson et al., 2012). The model employs 20 vertical hybrid pressure- σ levels for the regional $0.1^\circ \times 0.1^\circ$ EMEP modelling domain (30°N - 82°N , 30°W - 90°E), and 19 vertical levels for the global $0.5^\circ \times 0.5^\circ$ modelling domain. Both the regional and global grids use 3-hourly meteorological data derived from the ECMWF
205 Integrated Forecasting System (IFS) cycle 40r1 model (ECMWF, 2014). 3-hourly IFS O_3 concentrations are specified at the model top (100 hPa) boundary condition, while output surface concentrations are adjusted to an equivalent altitude of 3 meters. Hydrogen gas (H_2) is specified with a fixed global concentration of 500 ppb. For the lateral boundary conditions (LBCs) in the regional simulations, 6-hourly output fields from global simulations are used, with each of the global simulations employing a spin-up period of six months. Diagnostic simulations find that the choice of LBC time-resolution has a negligibly small impact
210 on the simulation results, while choosing 6-hourly over 3-hourly LBCs saves considerable computation time. Output fields from the global model are also used as initial conditions for the regional runs. The geographical region spanned by the regional EMEP modelling domain contains the EMEP region, but also parts of North Africa and Asia, whose emissions are consistently treated as Rest-Of-World (ROW) between the global and regional simulations. For reference, the EMEP region as represented in the regional EMEP modelling domain is shown in supplementary Fig. S1.

215 4.1 CH_4 implementation

In the EMEP model, global mean CH_4 concentrations are specified at the start of each run and remain unchanged over the course of the simulation. However, observed CH_4 concentrations display a marked latitudinal gradient, primarily due to the presence of large natural and anthropogenic emission sources in the Northern Hemisphere. The latitudinal gradient can be described by its two leading Empirical Orthogonal Functions (EOFs), or principal components (Meinshausen et al., 2017). The
220 first EOF (EOF1) represents a nearly linear North-South gradient, while the second EOF (EOF2) represents a local northern mid-latitude maximum of ~ 10 ppb. EOF1 has a pre-industrial North to South pole gradient of around 40-50 ppb, and of around 90 ppb for the year 2014 (Meinshausen et al., 2017). To capture the main characteristics of the latitudinal gradient, the contribution of EOF1 is included in the EMEP model by specifying

$$\text{CH}_4'(\phi, \text{CH}_4) = \text{CH}_4 \left(1 + \frac{0.025\phi}{90} \right), \quad (1)$$

225 where CH_4 represents the global mean background concentration and ϕ is latitude in degrees. For pre-industrial (808 ppb) and the year 2015 (1834 ppb) global mean CH_4 concentrations, Eq. 1 yields latitudinal gradients of 40 ppb and 92 ppb, respectively, consistent with those described in Meinshausen et al. (2017). By applying Eq. 1 also for the projected CH_4 concentrations, an approach similar to that of Meinshausen et al. (2020) is followed, by effectively using EOF1 to extrapolate the latitudinal gradient into the future based on anthropogenic CH_4 emissions.



Table 2. EMEP model configurations for the scenario analysis discussed in Sect. 5. The CH₄ concentrations refer to the 2050 global mean values calculated in Sect. 3. Each of the scenarios is simulated for the five meteorological years between 2013-2017.

Experiment long-name	Short-name	ROW emis	EMEP region emis	CH ₄ (ppb)
Baseline 2015	bs15_bs15ch4	2015 baseline	2015 baseline	1834
Baseline 2015 to ROW 2050 CLE emis	rowcle50_bs15ch4	2050 CLE	2015 baseline	1834
Baseline 2015 to global 2050 CLE emis	cle50_bs15ch4	2050 CLE	2050 CLE	1834
Global 2050 CLE	cle50_cle50ch4	2050 CLE	2050 CLE	2236
2050 CLE to ROW 2050 MFR emis	rowmfr50_cle50ch4	2050 MFR	2050 CLE	2236
2050 CLE to global 2050 MFR emis	mfr50_cle50ch4	2050 MFR	2050 MFR	2236
Global 2050 MFR	mfr50_mfr50ch4	2050 MFR	2050 MFR	1651
2050 MFR to ROW 2050 LOW emis	rowlow50_mfr50ch4	2050 LOW	2050 MFR	1651
2050 MFR to global 2050 LOW emis	low50_mfr50ch4	2050 LOW	2050 LOW	1651
Global 2050 LOW	low50_low50ch4	2050 LOW	2050 LOW	1574

230 4.2 Scenario configurations

To simulate the effects of precursor emission changes inside and outside of the EMEP region, regional simulations are combined with LBCs from the global model configuration. Simulations where only background CH₄ concentrations are changed serve to isolate the impact of global CH₄ change. Since CH₄ is a globally well-mixed gas, and since the concentration changes are the result of anthropogenic CH₄ emission changes, the impact of the total global mean CH₄ change is split into its EMEP region and ROW contributions based on the CH₄ emission changes within these respective regions. This approach is supported by the surface O₃ response being effectively linear in the range of CH₄ concentrations relevant to the current work, as discussed in Sect. 6.1. An overview of the scenario simulations is shown in Table 2, noting that each of the configurations is simulated for each of the five meteorological years between 2013-2017 for both the regional and global setups, as discussed below.

4.3 Baseline evaluation against observation

240 The efficacy of the EMEP model to simulate peak season MDA8 is evaluated by comparing the baseline configuration to surface observations. To this end, the baseline 2015 configuration is used to perform simulations for the 2013-2017 meteorological years, and compared against surface observations from the EBAS database (Tørseth et al., 2012). While the emissions are fixed to that of the year 2015, inter-annual variability in the emissions is generally small. The 56 EBAS stations are located within the European part of the EMEP region (as shown in supplementary Fig. S2), and are selected from all available stations based on the requirement that they each measure peak season MDA8 for each of the five meteorological years. For MDA8, data availability guidelines stipulate that for each 8-hourly mean 75% of the hourly values must be present, while at least 75% of the eight hour averages must be present in a day to assign a maximum daily 8-hour mean (EU, 2008). Data availability guidelines



similar to those for annual mean O₃ are also adopted, requiring that at least 90 % of the days between April-September have MDA8 measurements available to assign a peak season average.

250 Fig. 2a compares the five-year average modelled and observed peak season MDA8 values at each of the 56 stations. A clear relationship between the modelled and observed values is present, having a Pearson correlation coefficient (r) of 0.87. The normalized mean bias (NMB) amounts to 4.2 %, indicating that the model has a slight tendency to overestimate. Fig. 2b shows the annual averages across all 56 stations, illustrating that the total inter-annual variability for both model and measurements corresponds to around 4-5 $\mu\text{g m}^{-3}$. The difference between the total annual average modelled and observed concentrations is
255 greatest for the year 2014, amounting to 6.9 $\mu\text{g m}^{-3}$ (8.2 %), while being as low as 0.6 $\mu\text{g m}^{-3}$ (0.7 %) for the year 2013. The difference in the five-year average measured (84.9 $\mu\text{g m}^{-3}$) and modelled (88.5 $\mu\text{g m}^{-3}$) concentrations follows that of the NMB (3.6 $\mu\text{g m}^{-3}$, or 4.2 %). In Fig. 2c, annual averages across all stations within Sweden, Germany, Spain, the United Kingdom, and Poland are shown, illustrating that the model generally captures the observed variability between high and low O₃ years also at regional scales. Observed concentrations were the lowest in 2017, except for in Spain, as also reproduced
260 by the model. The observed differences between the highest (2015) and lowest year (2017) can be as large as 12.7 $\mu\text{g m}^{-3}$ (16.80 %), for example for Poland. The modelled inter-annual variability in the different regions is approximately equal to, or sometimes smaller than (e.g., Poland, Spain), the observed variability. For Poland, the difference between the highest and lowest modelled year amounts to 7.6 $\mu\text{g m}^{-3}$ (8.51 %), being lower by 5.1 $\mu\text{g m}^{-3}$ than the observed maximum variability.

Overall, the EMEP model displays good agreement with observations across the five meteorological years, while highlight-
265 ing that inter-annual peak season MDA8 variability can be on the order of 10-15 % on regional scales, and around 5 % across Europe. To reduce the effects of meteorological variability, each of the scenarios listed in Table 2 is therefore simulated for the years 2013-2017, with the results presented in the following representing five-year averages.




Figure 2. Modelled versus observed peak season MDA8 across Europe. Panel (a) shows five-year averaged values at each of the 56 stations, while panel (b) compares the annual values averaged over all stations. Panel (c) shows the yearly averages for Sweden, Germany, Spain, the United Kingdom, and Poland, with the number in brackets indicating the number of stations in each of the countries.



5 Results

While the focus in this section lies on peak season MDA8, results for other O₃ indicators are included in the supplementary material, as referred to in the text. In addition, the following discusses a number of weighted averaging approaches for both health and vegetation O₃ indicators, with the different population and crop-area maps shown for reference in supplementary Fig. S3.

5.1 EMEP region peak season MDA8

Fig. 3 shows a so-called cascade-plot of the EMEP region population weighted peak season MDA8 changes between the 2015 baseline and the 2050 CLE, MFR, and LOW scenarios. Here the population weighting is calculated using the Global Human Settlement Layer (GHSL) population distribution for the year 2015 (Schiavina et al., 2023), aggregated from its native 3 arc-second resolution to the regional EMEP grid, remaining unchanged for all scenarios. In Fig. 3, the impacts arising from NO_x, CO, and NMVOC precursor emissions changes and from CH₄ are shown as separate cascade-steps. In the cascades, the separate contributions arising from EMEP region and ROW emission changes are also highlighted, calculated using the model configurations described in Table 2. For example, the difference between the ‘bs15_bs15ch4’ and ‘rowcle50_bs15ch4’ simulations yields the change due to 2050 CLE precursor emission changes in the ROW region relative to the 2015 baseline, while the difference between the ‘cle50_bs15ch4’ and ‘cle50_cle50ch4’ simulations yields the change due to background CH₄ changes. As noted in Sect 4, the impact of global CH₄ emission changes is split into its EMEP and ROW region contributions based on the emission changes within these respective regions. In effect, the cascade-plot summarizes the impact of each of successive precursor and CH₄ change from the 2015 baseline down to the 2050 LOW scenario.

Fig. 3 shows that average peak season MDA8 concentrations are reduced from 93.3 to 90.3 µg m⁻³ between the 2015 baseline and 2050 CLE scenarios, resulting largely from a decrease in precursor emissions in the EMEP region (-4.8 µg m⁻³) and to a lesser extent in the ROW region (-1.4 µg m⁻³). However, these reductions are partially offset by an increase of 3.2 µg m⁻³ arising from increased background CH₄ concentrations, being almost entirely the result of increased CH₄ emissions in the ROW region. Going from the 2050 CLE to 2050 MFR scenario, the net reduction from 90.3 to 76.8 µg m⁻³ (-15 %) is split into three nearly equal parts arising from EMEP region precursor reductions, ROW precursor reductions, and background CH₄ reductions. Further, the 2050 LOW scenario differs relatively little from the MFR, with roughly half of the change from 76.8 to 73.4 µg m⁻³ arising from further precursor emission reductions within the EMEP region. Cascade-plots for the annual O₃ mean, SOMO35, and POD₃IAM_{WH} indicators discussed in Sect. 5.2, are shown in supplementary Fig. S4-S6.

5.1.1 Geographical distribution

To illustrate the impact of geographical location on the O₃ changes resulting from precursor and CH₄ emission changes, the difference in peak season MDA8 between the 2050 CLE and LOW scenarios is shown across the regional EMEP modelling domain in Fig. 4. Here the 2050 CLE to LOW impacts are calculated by combining the results from the 2050 CLE to 2050 MFR simulations with the 2050 MFR to 2050 LOW simulations described in Table 2. In Fig. 4a, the change in peak season MDA8




Figure 3. Cascade plot of the population weighted EMEP region average peak season MDA8 scenario changes arising from NO_x, CO, and NMVOC emission changes within the EMEP (black) and ROW (blue) regions, and from background CH₄ changes arising from EMEP (red) and ROW (green) region emission changes. The black and grey dashed horizontal line denote guideline and interim WHO target values, respectively. Note that the y-axis starts at 50 µg m⁻³.

300 resulting from the change to ROW LOW emissions is shown. As expected, the ROW LOW impacts are most pronounced in the ROW countries within the regional modelling domain (e.g., North African countries). Nevertheless, countries along the Southern border of the EMEP region as well as along the Western coast of Europe also see reductions ranging from 5-15 µg m⁻³. The reductions along the Western coast of Europe are likely the result of emission reductions in North America, with the associated O₃ perturbations carried over the Atlantic ocean by the prevailing Westerlies. Fig. 4b shows that the impact of regional LOW emissions is largely centered on the EMEP region, ranging from approximately 5 µg m⁻³ in Western Europe to 15 µg m⁻³ in the West-Balkan and EECCA countries. While local in nature, the impact of emission reductions in both the EMEP and ROW regions can lead to increases of as much as 30 µg m⁻³ in large urban areas (as highlighted in Fig. 4b for Almaty, Kazakhstan), due to reductions in the titration effect of NO_x. The impact of background CH₄ reductions from 2236 to 1574 ppb is shown in Fig. 4c, with the latitudinal gradient arising from the latitudinal variations in insolation. The resulting peak season MDA8 reductions amount to around 5 µg m⁻³ across the EMEP region.

Fig. 4d shows the results for the full 2050 LOW scenario, illustrating that peak season MDA8 concentrations fall below 60 µg m⁻³ over parts of Scandinavia, while ranging from 80 to 90 µg m⁻³ over Northern Italy and Kazakhstan. In Central Europe, concentrations typically range from 60-70 µg m⁻³, highlighting that the WHO exposure guideline of 60 µg m⁻³ is not met anywhere in the majority of EMEP countries. However, the interim target of 70 µg m⁻³ is reached in a number of




Figure 4. Reductions in peak season MDA8 achieved by 2050 ROW LOW (panel a) and EMEP region (panel b) precursor emission changes relative to the 2050 CLE scenario. Panel (b) also highlights the simulation results for Almaty, Kazakhstan. Panel (c) shows the reductions arising from the background CH₄ change from 2236 to 1574 ppb, while panel (d) shows the peak season MDA8 as simulated for the full 2050 LOW scenario. Note the difference in color-scale for panel (d).

315 Western European countries, such as the Netherlands, France, Germany, and the United Kingdom. The population weighted
 LOW scenario concentrations for each of the individual countries in the EMEP region are shown in supplementary Fig. S7,
 along with their 2015 baseline and 2050 CLE and MFR concentrations. In addition, supplementary Fig. S8 follows that of
 Fig. 4, but instead compares the impacts of the LOW scenario against the 2015 baseline. For the latter, the impact of regional
 emission reductions is comparatively higher, while that of CH₄ changes is comparatively lower, consistent with the results
 320 shown in Fig. 3.

5.2 Other O₃ indicators

This section serves to provide reference to earlier studies by showing the scenario results for a range of other health and
 vegetation O₃ indicators. For example, earlier works have investigated the impact of precursor and CH₄ emission changes on
 (area weighted) annual mean surface O₃ (“O₃ mean”) concentrations (Turnock et al., 2018; Jonson et al., 2018), while the
 325 Sum of Ozone Means Over 35 ppb (SOMO35), 4th highest annual MDA8, and summertime (JJA) average daily maximum O₃



concentrations have been used for health impact studies (Fleming et al., 2018). Furthermore, JJA average O_3 concentrations were used for the study of the climate-impact on surface O_3 by Colette et al. (2015), as will be discussed in more detail in Sect. 6. For the impacts on vegetation, the growing-season accumulated Phyto-toxic Ozone Dose (POD_Y) uptake over a certain threshold value Y ($\text{nmole } O_3 \text{ m}^{-2} \text{ s}^{-1}$) can induce reductions in crop and semi-natural biomass (Emberson, 2020; Mills et al., 2018). To this end, the Integrated Assessment Modelling (IAM) vegetation-type specific POD_Y indicators (POD_{Y-IAM}) serve as simplified risk assessment indicators for use in CTMs such as the EMEP model (Simpson et al., 2012, 2007), as also described in the UNECE ‘Mapping Manual’ (UNECE, 2017). The $POD_{3-IAM_{WH}}$ indicator represents the cumulative growing-season (~ 90 days) stomatal O_3 uptake for a generic temperate or boreal crop, being largely based on wheat (WH), and is used as an indicator for wheat yield loss (Pandey et al., 2023; Mills et al., 2018). In addition, the $POD_{1-IAM_{DF}}$ indicator is used in the risk assessment of reductions in annual living deciduous forest (DF) biomass growth (UNECE, 2017), having a ~ 180 day growing season at 50°N .

Table 3 shows the absolute and percentage change scenario results across the range of O_3 indicators, for an extended range of (constructed) scenarios. For example, the ‘2015 Base to 2050 MFR’ scenario is constructed using the differences between the ‘2015 Base to 2050 CLE’ and the ‘2050 CLE to 2050 MFR’ scenarios described in Table 2, while the ‘2050 CLE to 2050 LOW’ scenario is constructed using the differences between the ‘2050 CLE to 2050 MFR’ and ‘2050 MFR to 2050 LOW’ scenarios (as described in Sect. 5.1.1). Likewise, the ‘2015 Base to 2050 LOW’ scenario is constructed using the differences between the ‘2015 Base to 2050 CLE’, ‘2050 CLE to 2050 MFR’, and ‘2050 MFR to 2050 LOW’ scenarios. Note that for peak season MDA8, the absolute numbers shown for the ‘2015 Base to 2050 CLE’, ‘2050 CLE to 2050 MFR’, and ‘2050 MFR to 2050 LOW’ scenarios correspond to those shown in Fig. 3. Furthermore, since the relative importance of CH_4 emission changes inside the EMEP region is small, Table 3 only includes the impact of global CH_4 changes. While health-related O_3 indicators are shown as population weighted averages, the POD_Y indicators are shown as their respective vegetation-area weighted averages (i.e., average values per square meter of vegetation, as illustrated in supplementary Fig. S3).

Table 3 illustrates that different uses of threshold values and time and length of averaging or accumulation periods leads to differences in the relative importance of precursor and CH_4 emission changes. For example, indicators most sensitive to O_3 concentrations during its peak photochemical production period (peak season MDA8, JJA O_3 max, JJA O_3 mean, and 4th highest MDA8) are most strongly impacted by regional precursor emission reductions, especially when compared against the 2015 baseline scenario. In contrast, regional emission reductions are much less important for annual O_3 , due to the competing effects of local wintertime NO_x titration. The importance of ROW emissions is, broadly speaking, proportional to the length of the averaging or accumulation period, while also being most relevant to the 2050 scenarios (i.e., 2050 CLE and MFR). For indicators employing a threshold value, the percentage-change impacts are proportional to the height of the threshold relative to the baseline (or background) value, which effectively determines the degree of which the natural background is filtered out. For example, the total percentage-change reduction from the 2015 baseline to 2050 LOW scenarios for the SOMO35, $POD_{3-IAM_{WH}}$, and $POD_{1-IAM_{DF}}$ indicators amounts to 63.5 %, 36.9 %, and 32.1 %, respectively. While already implied in Fig. 3, Table 3 also shows that the impact of CH_4 emission reductions is most important relative to the 2050 CLE scenario and less so when compared against the 2015 baseline. However, O_3 mean is an exception to the latter, with CH_4 having the largest



Table 3. Absolute and percentage change (in brackets) scenario impacts across the EMEP region. Changes resulting from precursor emission changes in the EMEP (reg.) and ROW regions, and from global CH₄ changes, are shown relative to the scenario starting points. End values correspond to the weighted averages at each of the scenario end-points.

Scenario		2015 Base to 2050 CLE	2015 Base to 2050 MFR	2015 Base to 2050 LOW	2050 CLE to 2050 MFR	2050 CLE to 2050 LOW	2050 MFR to 2050 LOW
PS MDA8 ^a	Reg. emis	-4.8 (-5.1 %)	-9.3 (-10.0 %)	-11.2 (-12.0 %)	-4.5 (-5.0 %)	-6.4 (-7.1 %)	-1.9 (-2.4 %)
	ROW emis	-1.4 (-1.5 %)	-6.1 (-6.5 %)	-7.0 (-7.5 %)	-4.7 (-5.2 %)	-5.6 (-6.3 %)	-1.0 (-1.2 %)
	CH ₄	3.1 (3.4 %)	-1.2 (-1.2 %)	-1.7 (-1.9 %)	-4.3 (-4.8 %)	-4.9 (-5.4 %)	-0.6 (-0.8 %)
	End value	90.3	76.8	73.4	76.8	73.4	73.4
O ₃ mean ^a	Reg. emis	1.3 (2.0 %)	-0.2 (-0.3 %)	-0.9 (-1.4 %)	-1.5 (-2.3 %)	-2.1 (-3.3 %)	-0.7 (-1.2 %)
	ROW emis	-1.1 (-1.8 %)	-5.5 (-8.8 %)	-6.6 (-10.6 %)	-4.4 (-6.7 %)	-5.5 (-8.5 %)	-1.1 (-2.0 %)
	CH ₄	2.2 (3.4 %)	-0.8 (-1.3 %)	-1.2 (-2.0 %)	-3.0 (-4.6 %)	-3.4 (-5.2 %)	-0.4 (-0.7 %)
	End value	65.0	56.2	54.0	56.2	54.0	54.0
4th MDA8 ^a	Reg. emis	-9.3 (-6.4 %)	-15.2 (-10.4 %)	-18.0 (-12.3 %)	-5.8 (-4.2 %)	-8.7 (-6.2 %)	-2.8 (-2.3 %)
	ROW emis	-1.0 (-0.7 %)	-4.7 (-3.2 %)	-5.6 (-3.8 %)	-3.7 (-2.7 %)	-4.6 (-3.3 %)	-0.9 (-0.7 %)
	CH ₄	3.1 (2.1 %)	-1.2 (-0.8 %)	-1.7 (-1.2 %)	-4.3 (-3.1 %)	-4.9 (-3.5 %)	-0.6 (-0.5 %)
	End value	139.1	125.2	120.9	125.2	120.9	120.9
JJA O ₃ max ^d	Reg. emis	-7.4 (-7.1 %)	-13.3 (-12.7 %)	-15.7 (-14.9 %)	-5.9 (-5.9 %)	-8.2 (-8.3 %)	-2.3 (-2.7 %)
	ROW emis	-1.2 (-1.1 %)	-5.2 (-5.0 %)	-5.9 (-5.6 %)	-4.0 (-4.0 %)	-4.7 (-4.7 %)	-0.7 (-0.8 %)
	CH ₄	3.7 (3.6 %)	-1.3 (-1.3 %)	-2.0 (-1.9 %)	-5.1 (-5.1 %)	-5.8 (-5.8 %)	-0.7 (-0.8 %)
	End value	99.9	84.9	81.2	84.9	81.2	81.2
JJA O ₃ mean ^a	Reg. emis	-3.6 (-4.9 %)	-6.9 (-9.3 %)	-8.3 (-11.2 %)	-3.2 (-4.5 %)	-4.7 (-6.5 %)	-1.4 (-2.3 %)
	ROW emis	-0.9 (-1.3 %)	-4.4 (-6.0 %)	-5.0 (-6.7 %)	-3.5 (-4.8 %)	-4.0 (-5.6 %)	-0.6 (-0.9 %)
	CH ₄	2.9 (3.9 %)	-1.0 (-1.4 %)	-1.6 (-2.2 %)	-3.9 (-5.5 %)	-4.5 (-6.2 %)	-0.5 (-0.9 %)
	End value	72.0	61.4	58.8	61.4	58.8	58.8
SOMO35 ^b	Reg. emis	-427 (-13.2 %)	-891 (-27.5 %)	-1049 (-32.4 %)	-464 (-15.4 %)	-622 (-20.7 %)	-158 (-10.5 %)
	ROW emis	-162 (-5.0 %)	-809 (-25.0 %)	-919 (-28.3 %)	-647 (-21.5 %)	-756 (-25.2 %)	-109 (-7.3 %)
	CH ₄	355 (11.0 %)	-44 (-1.4 %)	-89 (-2.8 %)	-399 (-13.3 %)	-445 (-14.8 %)	-45 (-3.0 %)
	End value	3009	1498	1185	1498	1185	1185
POD ₃ IAM _{WH} ^c	Reg. emis	-1.7 (-10.1 %)	-3.0 (-18.2 %)	-3.5 (-21.2 %)	-1.3 (-8.7 %)	-1.8 (-11.9 %)	-0.5 (-4.4 %)
	ROW emis	-0.4 (-2.5 %)	-1.9 (-11.3 %)	-2.1 (-12.8 %)	-1.4 (-9.4 %)	-1.7 (-11.0 %)	-0.2 (-2.2 %)
	CH ₄	1.0 (5.8 %)	-0.3 (-1.9 %)	-0.5 (-3.0 %)	-1.3 (-8.3 %)	-1.4 (-9.4 %)	-0.2 (-1.5 %)
	End value	15.4	11.3	10.4	11.3	10.4	10.4
POD ₁ IAM _{DF} ^c	Reg. emis	-2.4 (-10.1 %)	-4.3 (-18.2 %)	-4.9 (-20.9 %)	-1.9 (-8.8 %)	-2.5 (-11.7 %)	-0.6 (-3.7 %)
	ROW emis	-0.5 (-2.1 %)	-1.8 (-7.8 %)	-2.1 (-8.9 %)	-1.3 (-6.2 %)	-1.6 (-7.4 %)	-0.3 (-1.5 %)
	CH ₄	1.0 (4.2 %)	-0.4 (-1.5 %)	-0.5 (-2.3 %)	-1.3 (-6.2 %)	-1.5 (-7.0 %)	-0.2 (-1.1 %)
	End value	21.5	16.9	15.9	16.9	15.9	15.9

^aPopulation weighted EMEP region average in µg m⁻³. ^bPopulation weighted EMEP region average in ppb day⁻¹. ^cCrop-area weighted EMEP region average in mmol m⁻².

^dPopulation weighted average converted from ppb to µg m⁻³ using the standard-atmosphere O₃ conversion factor of 1.96.



percentage-change impact from the 2015 baseline to 2050 CLE scenario. Furthermore, CH₄ reductions contribute roughly one thirds of the total reductions for each of the peak O₃ indicators for the 2050 CLE to 2050 MFR scenario, although this is closer to one-fourths for SOMO35 (26.4 %).

365 For the population weighted O₃ indicators (i.e., all except those for vegetation), the corresponding area weighted averages are shown in supplementary Table S2. While the results are generally consistent between the two weighted averaging approaches, indicators sensitive to peak O₃ concentrations are comparatively less impacted by regional precursor emission changes when calculated as area weighted averages. However, the area weighted impacts of regional precursor emission changes are considerably larger for annual O₃ mean, since NO_x titration effects in urban areas is weighted less heavily. For example, reducing regional emissions between the 2015 baseline to 2050 MFR scenarios sees a population weighted O₃ mean reduction of 0.2
370 µg m⁻³ (0.3 %), while the corresponding area weighted reduction amounts to 4.5 µg m⁻³ (6.7 %).



6 Discussion

In the current setup, the EMEP model is unable to capture the effects of future climate change on surface O_3 concentrations. This effect, often described as the O_3 climate penalty (e.g., Fu and Tian, 2019; Rasmussen et al., 2013), can affect surface O_3 for example through climate-change induced changes in water vapour concentrations and biogenic VOC emissions. For European land surfaces, Colette et al. (2015) estimated the 95 % confidence interval of the mid-century (2041-2070) surface JJA O_3 mean climate penalty to range from 0.44-0.64 ppb, based on an ensemble of 25 chemistry-climate model simulations. Compared to the JJA O_3 mean changes between the 2050 CLE and MFR scenarios shown in Table 3, amounting to $10.6 \mu\text{g m}^{-3}$ (or 5.3 ppb using the standard-atmosphere O_3 conversion factor of 1.96), the impact of the climate penalty on the results of the current work is expected to be small. Other climate-uncertainties relate to the calculated CH_4 projections, with natural soil emissions estimated to increase by $22.8 \pm 3.6 \text{ Tg CH}_4 \text{ yr}^{-1}$ by the year 2100 in the SSP5-8.5 scenario (Guo et al., 2023). Such changes are nevertheless small relative to the total natural emissions, estimated at 210 Tg yr^{-1} in Sect. 3, also considering that the scenario analysis of the current work goes up to the year 2050.

While constructing emission data sets based on a wide variety of information is by itself challenging (e.g., de Meij et al., 2024; Thunis et al., 2022), the emission scenarios employed in the current work are also inherently based on a number of socio-economic activity projections. In practice, the reliable quantification of the uncertainty on the input parameters to the GAINS model is itself considered the most uncertain element of the analysis (Amann et al., 2011). In this light, the emission scenarios arguably represent the largest source of uncertainty for the current work, which is unavoidable and not directly quantifiable. Nevertheless, the GAINS model by design attempts to minimize the impact of uncertainties on policy-relevant model output, to increase the robustness (i.e., the priorities and control needs between countries, sectors and pollutants do not significantly change due to uncertainties in the model elements) of the emission control strategies (Amann et al., 2011).

6.1 O_3 production efficiency of CH_4

The CH_4 oxidation reaction that leads the production of O_3 depends on the availability of NO_x and OH (Crutzen et al., 1999). OH is produced through the photolysis of O_3 and subsequent reaction of $\text{O}(1\text{D})$ with water vapour (H_2O), with the majority of surface O_3 being produced by the photolysis of NO_x in carbonaceous-rich environments. In addition, CO and VOCs (including CH_4) are net sinks of OH, creating a non-linear relationship between their atmospheric abundance and the O_3 production efficiency (OPE) of CH_4 (Isaksen et al., 2014). In the current work, the OPE is taken as the capacity of CH_4 to produce surface O_3 in the EMEP region. To investigate the impact of OPE on the calculated O_3 response, diagnostic simulations are performed where background CH_4 concentrations are varied between 850 to 2600 ppb in 250 ppb steps, using both the 2015 baseline and the 2050 CLE and LOW emission scenarios as the source of background precursor emissions. The resulting CH_4 impacts on peak season MDA8 are shown in Fig. 5, noting that the starting point of 850 ppb corresponds roughly to pre-industrial CH_4 concentrations. For simplicity, the simulations shown here are only calculated for the 2015 meteorological year.




Figure 5. Change in EMEP region population weighted peak season MDA8 for background CH₄ concentrations ranging from 850 to 2600 ppb in 250 ppb intervals, relative to peak season MDA8 concentrations at 850 ppb CH₄. The impacts are calculated for the baseline 2015 and the 2050 CLE and LOW emission scenarios defined in Table 2, with the dashed vertical lines marking their calculated 2050 background CH₄ concentrations (1574, 1834, and 2236 ppb, respectively).

Fig. 5 illustrates that the OPE is highest in the 2015 baseline scenario, when EMEP region NO_x emissions are also highest (Table 1). Regional NO_x emissions are reduced strongly in the 2050 CLE scenario, while other emissions change relatively little. As a result, the OPE is a factor of 0.88 smaller relative to the 2015 baseline across the range of CH₄ concentrations. Similarly, the OPE in the 2050 LOW scenario is a factor of 0.88 lower than that of the 2050 CLE scenario, and by a factor of 0.78 relative to the 2015 baseline scenario. In the analysis of Sect. 5, the largest CH₄ differences occur between the 2050 CLE and LOW scenarios, ranging from 2236 to 1574 ppb. For this concentration interval, the decrease in peak season MDA8 amounts to 5.4 and 4.7 µg m⁻³ when calculated from the 2050 CLE and LOW precursor emission scenarios, respectively. Since this represents a comparatively small difference, the CH₄ impacts described in Sect. 5 are robust with respect to the choice of emission scenario in which the CH₄ concentrations are reduced. Fig. 5 furthermore illustrates that the peak season MDA8 response is approximately linear in the range of CH₄ concentrations relevant to the current work, supporting the approach of splitting the O₃ impacts based on the separate emission changes within the EMEP and ROW regions. Another corollary is that the contribution of anthropogenic background CH₄ to total peak season MDA8 amounts to approximately 10.7 (11.5 %), 12.7 (14.0 %), and 6.4 (8.7 %) µg m⁻³ in the 2015 baseline, 2050 CLE, and 2050 LOW scenarios, respectively. Here the percentage



contributions are based on the scenario totals shown in Fig. 3 and Table 3. Recognizing that the MFR and LOW precursor emission scenarios are nearly identical except for CH₄ emissions, the anthropogenic CH₄ contribution calculated for the 2050 MFR scenario (1651 ppb) amounts to 7.0 µg m⁻³ (9.1 %).

6.2 Comparison to previous studies

420 Comparing to the results of the current work to earlier studies can be challenging, for example due to differences in source-receptor area definitions, model configuration, weighted averaging approach, and emission scenarios. Nevertheless, the EMEP region total peak season MDA8 exposure reduction by 15 % between the 2050 CLE and MFR scenarios is consistent with the 16 % reduction found by Belis and Van Dingenen (2023) across the entire UNECE region. However, in our calculations the total 2050 MFR anthropogenic CH₄ contribution amounts to 7.0 µg m⁻³ (or 3.5 ppb using the standard-atmosphere O₃ conversion factor of 1.96), which is lower than their estimate of ~ 5 ppb (based on their Fig. S4). This can largely be reconciled considering that our estimate was calculated with the 2050 LOW scenario as the source of background precursor emissions, while theirs is based on O₃ sensitivities calculated from a 2010 baseline emission scenario. When using the 2015 baseline emission scenario as the source of background precursor emissions in our calculations, the total 2050 MFR anthropogenic CH₄ contribution amount to 9.0 µg m⁻³, or 4.5 ppb, which is more comparable.

430 In the work of Turnock et al. (2018), the box-model described in Holmes et al. (2013) is used to estimate the 2050 CLE and MFR CH₄ concentrations to amount to 2361 and 1420 ppb, respectively. They further estimate the 2050 CLE increase in CH₄ to contribute 1.6 ppb to annual mean area weighted O₃ across Europe relative to a 2010 baseline concentration of 1798 ppb, based on the parameterized response of 14 models. While the latter is higher than our estimate of 1.1 ppb for the EMEP region (Table S3, using the standard-atmosphere O₃ conversion factor of 1.96), our results find a more comparable contribution of 1.4 ppb when the response is calculated as the European area weighted average following the land-area definition of Turnock et al. (2018). However, in our results the 2050 CLE and MFR ensemble mean CH₄ concentrations amount to 2236 and 1651 ppb, respectively, with the total difference between the CLE and MFR scenarios therefore being 403 ppb (or 43 %) less than that of Turnock et al. (2018). While this may in part be due to their MFR scenario diverging from the CLE from 2020 rather than 2025 onwards, it nevertheless highlights the importance of the methodology used to estimate CH₄ concentrations, as the cumulative difference between scenarios can quickly diverge. The difference in CH₄ estimates also has implications for the impact of the 2050 MFR emissions relative to the baseline, which in our analysis (-183 ppb) is around half that determined by Turnock et al. (2018) (-378 ppb).

6.3 Air-pollution and global warming co-benefits

445 While a detailed discussion is beyond the scope of the current work, the global mean temperature change relative to the reference period of 1986-2005, as calculated for the 600-ensemble mean and 5-95 % range using the MAGICC7 model in Sect. 3, amounts to 2.21 [1.61-2.94], 2.02 [1.45-2.74], and 1.92 [1.33-2.67] degrees K for the 2050 CLE, MFR, and LOW scenarios, respectively. In the LOW-CH₄ scenario, where CH₄ emissions follow the LOW scenario while all other emissions follow that of the CLE, this change amounts to 2.03 [1.47-2.74] degrees K. Thus illustrating that around two-thirds of the

<https://doi.org/10.5194/egusphere-2024-1422>

Preprint. Discussion started: 3 June 2024

© Author(s) 2024. CC BY 4.0 License.



global warming reduction between the 2050 CLE (SSP2-4.5 GHGs) and LOW (SSP1-2.6 GHGs) scenarios can be achieved by
450 solely reducing CH₄ emissions. Note that all emissions follow those of the corresponding SSP scenarios before 2015, as the
IIASA scenarios only start from 2015 onwards.



7 Conclusion

This work investigates the impact of CH₄ and other precursor emissions on surface O₃ concentrations in the EMEP region for the CLE, MFR, and LOW emission scenarios up to the year 2050. In the CLE scenario, background CH₄ concentrations are projected to increase by 402 ppb (22 %) relative to 2015 baseline concentrations, while they are reduced by 183 ppb (-10 %) in the MFR scenario. The difference between the 2050 MFR and CLE scenario therefore amounts to 585 ppb (or 26.1 % less in the MFR compared to the CLE scenario), while the LOW scenario achieves a modest further 77 ppb reduction. The MFR CH₄ reductions lead to a peak season MDA8 exposure reduction of 4.3 µg m⁻³ (4.8 %) relative to the 2050 CLE case, contributing around one-thirds of the total peak season MDA8 reduction (13.5 µg m⁻³, or 15 %). The other two-thirds are split almost equally between the impact of other precursor emission (NO_x, CO, NMVOC) reductions in the EMEP and ROW regions, respectively. As for CH₄, the impact of further abatement policies for the other precursor emissions is comparatively small in the LOW scenario. While the impacts of background CH₄ changes are relatively small when measured against the 2015 baseline, the increase in CH₄ in the CLE scenario nevertheless offsets the peak season MDA8 reductions achieved by precursor emissions reductions in the EMEP region almost entirely.

In terms of the total reductions, the 2050 MFR scenario brings the EMEP region average peak season MDA8 exposure down from 90.3 to 76.8 µg m⁻³ relative to the CLE, against a 2015 baseline exposure of 93.3 µg m⁻³. Nevertheless, in the MFR scenario the majority of countries in the EMEP region (38 out of 49) are projected to stay above the interim WHO exposure target of 70 µg m⁻³. While the more stringent emission policies of the LOW scenario reduce the number of countries to 31, it still highlights the difficulties in reaching WHO guideline values, given that the MFR and LOW scenarios already include optimistic global and regional emission reductions. For some mostly Northern European countries, the LOW scenario brings surface peak season MDA8 concentrations close to the 60 µg m⁻³ WHO guideline value, even though none of their population weighted averages reach below this limit.

While the current work focuses on the peak season MDA8 indicator, the scenario results are also discussed for a range of other health and vegetation O₃ indicators. These results find that the relative importance of CH₄ and other precursor emission reductions depends on the choice of indicator, and to some extent on the spatial averaging approach between area and population weighted. The percentage change scenario impacts can vary greatly, being mostly dependent on the degree to which a threshold value applies. For example, the total reduction between the CLE and MFR scenarios for the SOMO35 health indicator and the POD₃IAM_{WH} vegetation indicator amounts to 50 % and 26 %, respectively, compared to a 15 % total reduction for peak season MDA8. Nevertheless, O₃ indicators emphasizing peak O₃ concentrations yield results largely consistent with those for peak season MDA8 in terms of the relative importance of the different emission changes. For these indicators, reducing precursor emissions other than CH₄ within the EMEP region, or Europe, further has the largest potential to reduce the impact of surface O₃ exposure relative to the 2015 baseline.

The EMEP modelling configuration described in the current work also serves to define the setup for future scenario work performed at MSC-W, both for the EMEP region as well as other source-receptor regions. Future work can also go out to quantifying the risks for mortality and vegetation yield loss based on their associations with the range of O₃ indicators. In addition,



the results of the current work contribute to the discussion surrounding the second revision of the Gothenburg Protocol, for which the impact of CH₄ on surface O₃ plays a prominent role. The current work also highlights that reducing CH₄ emissions achieves considerable global warming reductions, with solely reducing CH₄ emissions achieving roughly two-thirds of the global warming reduction potential between the full 2050 CLE (SSP2-4.5 GHGs) and LOW (SSP1-2.6 GHGs) scenarios.

490 However, the global warming and surface air quality reductions are almost entirely the result of, and can only be achieved by, CH₄ emission reductions outside of the EMEP region.



495 *Code and data availability.* The EMEP MSC-W CTM version rv5.0 is available from <https://zenodo.org/record/8431553> (EMEP MSC-W, 2023) (last access February 2024). The EMEP input files and output data fields specific to the current work, in addition to the Python scripts used for the data analysis and figure creation, are available from van Caspel et al. (2024). The latter data repository also contains the Python scripts used to create the MAGICC7 input and run files. The MAGICC7 model, 600-ensemble probabilistic distribution, and SSP emission scenarios can be downloaded after registration from <https://magicc.org/download/magicc7> (last access February 2024). The EBAS data are available from <https://ebas.nilu.no/> (last access February 2024).

500 *Author contributions.* HF and WEvC conceptualized the work, while WEvC performed the simulations, did the analysis, and wrote the manuscript. ZK contributed to the text of Sect. 2. CH created the emission scenario files. All authors reviewed the manuscript before submission.

Competing interests. The authors declare that no competing interests are present.

505 *Acknowledgements.* This work has been funded by the EMEP Trust Fund. IT infrastructure in general was available through the Norwegian Meteorological Institute (MET Norway). Some computations were performed on resources provided by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway (grant NN2890k and NS9005k). The CPU time made available by ECMWF has been critical for both for generation of meteorology used as input for the EMEP MSC-W model as well as the calculations presented in the current work.

510 The EBAS database has largely been funded by the UN-ECE CLRTAP (EMEP), AMAP and through NILU internal resources. Specific developments have been possible due to projects like EUSAAR (EU-FP5)(EBAS web interface), EBAS-Online (Norwegian Research Council INFRA) (upgrading of database platform) and HTAP (European Commission DG-ENV) (import and export routines to build a secondary repository in support of www.htap.org, last access April 2024). A large number of specific projects have supported development of data and meta data reporting schemes in dialog with data providers (EU, CREATE, ACTRIS and others). For a complete list of programmes and projects for which EBAS serves as a database, please consult the information box in the Framework filter of the web interface. These are all highly acknowledged for their support.

515 We acknowledge the use of Pyaerocom (Gliss et al., 2020, <https://pyaerocom.met.no/>) (last access February 2024) for creating the co-located model and measurement data files used in the comparison of EBAS data against the simulations. We also acknowledge the use of the MAGICC7 Python wrapper Pymagicc version 2.1.4, available from <https://github.com/openscm/pymagicc> (last access February 2024).



References

- Abernethy, S., O'Connor, F. M., Jones, C. D., and Jackson, R. B.: Methane removal and the proportional reductions in surface temperature and ozone, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 379, 20210104, <https://doi.org/10.1098/rsta.2021.0104>, 2021.
- Alexandratos, N. and Bruinsma, J.: World agriculture towards 2030/2050: the 2012 revision, ESA Working paper No. 12-03. Rome, FAO., <https://www.fao.org/3/ap106e/ap106e.pdf>, [Last access: April 2024], 2012.
- Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Hoeglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schoepp, W., Wagner, F., and Winiwarter, W.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications, *Environmental Modelling & Software*, 26, 1489–1501, <https://doi.org/10.1016/j.envsoft.2011.07.012>, 2011.
- Amann, M., Klimont, Z., and Wagner, F.: Regional and Global Emissions of Air Pollutants: Recent Trends and Future Scenarios, *Annual Review of Environment and Resources*, 38, 31–55, <https://doi.org/10.1146/annurev-environ-052912-173303>, 2013.
- Amann, M., Kiesewetter, G., Schöpp, W., Klimont, Z., Winiwarter, W., Cofala, J., Rafaj, P., Höglund-Isaksson, L., Gomez-Sabriana, A., Heyes, C., Purohit, P., Borken-Kleefeld, J., Wagner, F., Sander, R., Fagerli, H., Nyiri, A., Cozzi, L., and Pavarini, C.: Reducing global air pollution: the scope for further policy interventions, *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences*, 378, 20190331, <https://doi.org/10.1098/rsta.2019.0331>, 2020.
- Belis, C. A. and Van Dingenen, R.: Air quality and related health impact in the UNECE region: source attribution and scenario analysis, *Atmospheric Chemistry and Physics*, 23, 8225–8240, <https://doi.org/10.5194/acp-23-8225-2023>, 2023.
- Colette, A., Andersson, C., Baklanov, A., Bessagnet, B., Brandt, J., Christensen, J. H., Doherty, R., Engardt, M., Geels, C., Giannakopoulos, C., Hedegaard, G. B., Katragkou, E., Langner, J., Lei, H., Manders, A., Melas, D., Meleux, F., Rouil, L., Sofiev, M., Soares, J., Stevenson, D. S., Tombrou-Tzella, M., Varotsos, K. V., and Young, P.: Is the ozone climate penalty robust in Europe?, *Environmental Research Letters*, 10, 084015, <https://doi.org/10.1088/1748-9326/10/8/084015>, 2015.
- Crutzen, P. J., Lawrence, M. G., and Pöschl, U.: On the background photochemistry of tropospheric ozone, *Tellus B: Chemical and Physical Meteorology*, 51, 123–146, <https://doi.org/10.3402/tellusb.v51i1.16264>, 1999.
- de Meij, A., Cuvelier, C., Thunis, P., Pisoni, E., and Bessagnet, B.: Sensitivity of air quality model responses to emission changes: comparison of results based on four EU inventories through FAIRMODE benchmarking methodology, *Geoscientific Model Development*, 17, 587–606, <https://doi.org/10.5194/gmd-17-587-2024>, 2024.
- Dentener, F., Stevenson, D., Cofala, J., Mechler, R., Amann, M., Bergamaschi, P., Raes, F., and Derwent, R.: The impact of air pollutant and methane emission controls on tropospheric ozone and radiative forcing: CTM calculations for the period 1990-2030, *Atmospheric Chemistry and Physics*, 5, 1731–1755, <https://doi.org/10.5194/acp-5-1731-2005>, 2005.
- ECMWF: IFS Documentation CY40R1 - Part IV: Physical Processes, Tech. Rep. 4, ECMWF, <https://doi.org/10.21957/f56vvey1x>, 2014.
- Emberson, L.: Effects of ozone on agriculture, forests and grasslands, *Philosophical Transactions of the Royal Society A*, 378, 20190327, <https://doi.org/10.1098/rsta.2019.0327>, 2020.
- EMEP MSC-W: OpenSource v5.0 (202310), <https://doi.org/10.5281/zenodo.8431553>, 2023.
- EMEP/CEIP: Present state of emission data, <https://www.ceip.at/webdab-emission-database/reported-emissiondata>, [last access February 2024], 2023.



- EU: Consolidated text: Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe, Tech. rep., Publications Office, <https://eur-lex.europa.eu/EN/legal-content/summary/cleaner-air-for-europe.html>, [Last Access: February 2024], 2008.
- 555 Fagerli, H., Tsyro, S., Jonson, J. E., and al et: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components. EMEP Status Report 1/2023, Tech. rep., The Norwegian Meteorological Institute, Oslo, Norway, available from www.emep.int [last access: March 2024], 2023.
- Fiore, A. M., West, J. J., Horowitz, L. W., Naik, V., and Schwarzkopf, M. D.: Characterizing the tropospheric ozone response to methane emission controls and the benefits to climate and air quality, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- 560 Fleming, Z. L., Doherty, R. M., von Schneidmesser, E., Malley, C. S., Cooper, O. R., Pinto, J. P., Colette, A., Xu, X., Simpson, D., Schultz, M. G., Lefohn, A. S., Hamad, S., Moolla, R., Solberg, S., and Feng, Z.: Tropospheric Ozone Assessment Report: Present-day ozone distribution and trends relevant to human health, *Elementa: Science of the Anthropocene*, 6, 12, <https://doi.org/10.1525/elementa.273>, 2018.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, p. 923–1054, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021.
- 565 Fu, T.-M. and Tian, H.: Climate change penalty to ozone air quality: review of current understandings and knowledge gaps, *Current Pollution Reports*, 5, 159–171, 2019.
- 570 Ge, Y., Solberg, S., Heal, M., Reimann, S., van Caspel, W., Hellack, B., Salameh, T., and Simpson, D.: Evaluation of modelled versus observed NMVOC compounds at EMEP sites in Europe, *EGUsphere*, 2024, 1–73, <https://doi.org/10.5194/egusphere-2023-3102>, 2024.
- Gliss, J., Griesfeller, J., and Svennevik, H.: metno/pyaerocom: Release version 0.8.0, <https://doi.org/10.5281/zenodo.4159570>, 2020.
- Gomez Sanabria, A., Kiesewetter, G., Klimont, Z., Schoepp, W., and Haberl, H.: Potential for future reductions of global GHG and air pollutants from circular waste management systems, *Nature Communication*, 13, <https://doi.org/10.1038/s41467-021-27624-7>, 2022.
- 575 Guevara, M., Jorba, O., Tena, C., Denier van der Gon, H., Kuenen, J., Elguindi, N., Darras, S., Granier, C., and Pérez García-Pando, C.: Copernicus Atmosphere Monitoring Service TEMPORal profiles for the Global domain version 2.1 (CAMSGLOB-TEMPOv2.1), Copernicus Atmosphere Monitoring Service, ECCAD, <https://doi.org/10.24380/ks45-9147>, 2020a.
- Guevara, M., Jorba, O., Tena, C., Denier van der Gon, H., Kuenen, J., Elguindi, N., Darras, S., Granier, C., and Pérez García-Pando, C.: Copernicus Atmosphere Monitoring Service TEMPORal profiles for the regional European domain version 2.1 (CAMSGLOB-TEMPOv2.1), Copernicus Atmosphere Monitoring Service, ECCAD, <https://doi.org/10.24380/1cx4-zy68>, 2020b.
- 580 Guevara, M., Jorba, O., Tena, C., Denier van der Gon, H., Kuenen, J., Elguindi, N., Darras, S., Granier, C., and Pérez García-Pando, C.: Copernicus Atmosphere Monitoring Service TEMPORal profiles (CAMSGLOB-TEMPO): global and European emission temporal profile maps for atmospheric chemistry modelling, *Earth System Science Data*, 13, 367–404, <https://doi.org/10.5194/essd-13-367-2021>, 2021.
- Guo, J., Feng, H., Peng, C., Chen, H., Xu, X., Ma, X., Li, L., Kneeshaw, D., Ruan, H., Yang, H., and Wang, W.: Global Climate Change Increases Terrestrial Soil CH₄ Emissions, *Global Biogeochemical Cycles*, 37, e2021GB007255, <https://doi.org/10.1029/2021GB007255>, 2023.
- He, X., Shen, W., Wallington, T. J., Zhang, S., Wu, X., Bao, Z., and Wu, Y.: Asia Pacific road transportation emissions, 1900–2050, *Faraday Discuss.*, 226, 53–73, <https://doi.org/10.1039/D0FD00096E>, 2021.



- Holmes, C. D., Prather, M. J., Søvde, O. A., and Myhre, G.: Future methane, hydroxyl, and their uncertainties: key climate and emission parameters for future predictions, *Atmospheric Chemistry and Physics*, 13, 285–302, <https://doi.org/10.5194/acp-13-285-2013>, 2013.
- Huangfu, P. and Atkinson, R.: Long-term exposure to NO₂ and O₃ and all-cause and respiratory mortality: A systematic review and meta-analysis, *Environment international*, 144, 105998, 2020.
- Höglund-Isaksson, L.: Global anthropogenic methane emissions 2005-2030: technical mitigation potentials and costs, *acpd*, p. 11275–11315, 2012.
- 595 Höglund-Isaksson, L., Gómez-Sanabria, A., Klimont, Z., Rafaj, P., and Schöpp, W.: Technical potentials and costs for reducing global anthropogenic methane emissions in the 2050 timeframe –results from the GAINS model, *Environmental Research Communications*, 2, 025004, <https://doi.org/10.1088/2515-7620/ab7457>, 2020.
- IEA: World Energy Outlook 2018, <https://www.iea.org/reports/world-energy-outlook-2018>, [Last access: April 2024], 2018.
- Isaksen, I. S. A., Berntsen, T. K., Dalsøren, S. B., Eleftheratos, K., Orsolini, Y., Rognerud, B., Stordal, F., Søvde, O. A., Zerefos, C., and
600 Holmes, C. D.: Atmospheric Ozone and Methane in a Changing Climate, *Atmosphere*, 5, 518–535, <https://doi.org/10.3390/atmos5030518>, 2014.
- Jonson, J. E., Schulz, M., Emmons, L., Flemming, J., Henze, D., Sudo, K., Lund, M. T., Lin, M., Benedictow, A., Koffi, B., Dentener, F., Keating, T., Kivi, R., and Davila, Y.: The effects of intercontinental emission sources on European air pollution levels, *acp*, 18, 13655–13672, <https://doi.org/10.5194/acp-18-13655-2018>, 2018.
- 605 Kanter, D. R., Winiwarter, W., Bodirsky, B. L., Bouwman, L., Boyer, E., Buckle, S., Compton, J. E., Dalgaard, T., de Vries, W., Leclère, D., Leip, A., Müller, C., Popp, A., Raghuram, N., Rao, S., Sutton, M. A., Tian, H., Westhoek, H., Zhang, X., and Zurek, M.: A framework for nitrogen futures in the shared socioeconomic pathways, *Global Environmental Change*, 61, 102029, <https://doi.org/10.1016/j.gloenvcha.2019.102029>, 2020.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions
610 of particulate matter including black carbon, *Atmospheric Chemistry and Physics*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.
- Lan, X., Thoning, K., and Dlugokencky, E.: Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements. Version 2024-04, <https://doi.org/10.15138/P8XG-AA10>, 2022.
- Lefohn, A. S., Malley, C. S., Smith, L., Wells, B., Hazucha, M., Simon, H., Naik, V., Mills, G., Schultz, M. G., Paoletti, E., De Marco, A., Xu,
615 X., Zhang, L., Wang, T., Neufeld, H. S., Musselman, R. C., Tarasick, D., Brauer, M., Feng, Z., Tang, H., Kobayashi, K., Sicard, P., Solberg, S., and Gerosa, G.: Tropospheric ozone assessment report: Global ozone metrics for climate change, human health, and crop/ecosystem research, *Elementa: Science of the Anthropocene*, 6, 27, <https://doi.org/10.1525/elementa.279>, 2018.
- Malley, C. S., Borgford-Parnell, N., Haeussling, S., Howard, I. C., Lefèvre, E. N., and Kuylenstierna, J. C. I.: A roadmap to achieve the global methane pledge, *Environmental Research: Climate*, 2, 011003, <https://doi.org/10.1088/2752-5295/acb4b4>, 2023.
- 620 Mar, K. A., Unger, C., Walderdorff, L., and Butler, T.: Beyond CO₂ equivalence: The impacts of methane on climate, ecosystems, and health, *Environmental Science & Policy*, 134, 127–136, <https://doi.org/10.1016/j.envsci.2022.03.027>, 2022.
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C., Frieler, K., Knutti, R., Frame, D. J., and Allen, M. R.: Greenhouse-gas emission targets for limiting global warming to 2 C, *Nature*, 458, 1158–1162, <https://doi.org/10.1038/nature08017>, 2009.
- Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model,
625 MAGICC6 – Part 1: Model description and calibration, *Atmospheric Chemistry and Physics*, 11, 1417–1456, <https://doi.org/10.5194/acp-11-1417-2011>, 2011.



- 630 Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), *Geoscientific Model Development*, 10, 2057–2116, <https://doi.org/10.5194/gmd-10-2057-2017>, 2017.
- 635 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geoscientific Model Development*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.
- Mills, G., Sharps, K., Simpson, D., Pleijel, H., Broberg, M., Uddling, J., Jaramillo, F., Davies, W. J., Dentener, F., Van den Berg, M., et al.: Ozone pollution will compromise efforts to increase global wheat production, *Global change biology*, 24, 3560–3574, <https://doi.org/10.1111/gcb.14157>, 2018.
- 640 Monks, P. S., Archibald, A., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G., et al.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric Chemistry and Physics*, 15, 8889–8973, <https://doi.org/10.5194/acp-15-8889-2015>, 2015.
- Nicholls, Z., Meinshausen, M., Lewis, J., Corradi, M. R., Dorheim, K., Gasser, T., Gieseke, R., Hope, A. P., Leach, N. J., McBride, L. A., Quilcaille, Y., Rogelj, J., Salawitch, R. J., Samset, B. H., Sandstad, M., Shiklomanov, A., Skeie, R. B., Smith, C. J., Smith, S. J., Su, X., Tsutsui, J., Vega-Westhoff, B., and Woodard, D. L.: Reduced Complexity Model Intercomparison Project Phase 2: Synthesizing Earth System Knowledge for Probabilistic Climate Projections, *Earth's Future*, 9, e2020EF001900, <https://doi.org/10.1029/2020EF001900>, 2021.
- 645 Nicholls, Z., Meinshausen, M., Lewis, J., Smith, C. J., Forster, P. M., Fuglestedt, J. S., Rogelj, J., Kikstra, J. S., Riahi, K., and Byers, E.: Changes in IPCC Scenario Assessment Emulators Between SR1.5 and AR6 Unraveled, *Geophysical Research Letters*, 49, e2022GL099788, <https://doi.org/10.1029/2022GL099788>, e2022GL099788 2022GL099788, 2022.
- 650 Nicholls, Z. R. J., Meinshausen, M., Lewis, J., Gieseke, R., Dommenges, D., Dorheim, K., Fan, C.-S., Fuglestedt, J. S., Gasser, T., Golüke, U., Goodwin, P., Hartin, C., Hope, A. P., Kriegler, E., Leach, N. J., Marchegiani, D., McBride, L. A., Quilcaille, Y., Rogelj, J., Salawitch, R. J., Samset, B. H., Sandstad, M., Shiklomanov, A. N., Skeie, R. B., Smith, C. J., Smith, S., Tanaka, K., Tsutsui, J., and Xie, Z.: Reduced Complexity Model Intercomparison Project Phase 1: introduction and evaluation of global-mean temperature response, *Geoscientific Model Development*, 13, 5175–5190, <https://doi.org/10.5194/gmd-13-5175-2020>, 2020.
- 655 O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century, *Global Environmental Change*, 42, 169–180, <https://doi.org/10.1016/j.gloenvcha.2015.01.004>, 2017.
- Pandey, D., Sharps, K., Simpson, D., Ramaswami, B., Cremades, R., Booth, N., Jamir, C., Büker, P., Sinha, V., Sinha, B., et al.: Assessing the costs of ozone pollution in India for wheat producers, consumers, and government food welfare policies, *Proceedings of the National Academy of Sciences*, 120, e2207081120, <https://doi.org/10.1073/pnas.2207081120>, 2023.
- 660 Rafaj, P., Kiesewetter, G., Gül, T., Schöpp, W., Cofala, J., Klimont, Z., Purohit, P., Heyes, C., Amann, M., B.-K., and J., Cozzi, L.: Outlook for clean air in the context of sustainable development goals, *Glob. Environ. Change*, 53, 1–11, <https://doi.org/10.1016/j.gloenvcha.2018.08.008>, 2018.



- Rasmussen, D., Hu, J., Mahmud, A., and Kleeman, M. J.: The ozone–climate penalty: past, present, and future, *Environmental science & technology*, 47, 14 258–14 266, 2013.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O’Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentretter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- Schiavina, M., Melchiorri, M., Pesaresi, M., Politis, P., Carneiro Freire, S., Maffeni, L., Florio, P., Ehrlich, D., Goch, K., Carioli, A., Uhl, J., Tommasi, P., and Kemper, T.: GHSL Data Package 2023, Publications Office of the European Union, <https://doi.org/10.2760/098587>, 2023.
- Schneider von Deimling, T., Meinshausen, M., Levermann, A., Huber, V., Frieler, K., Lawrence, D. M., and Brovkin, V.: Estimating the near-surface permafrost-carbon feedback on global warming, *Biogeosciences*, 9, 649–665, <https://doi.org/10.5194/bg-9-649-2012>, 2012.
- Simpson, D.: The EMEP MSC-W modelling programme: Its relationship to policy support, current challenges and future perspectives, in: *Air Pollution Modelling and its Application XXII*, edited by Steyn, D., Builtjes, P., and Timmermans, R., NATO Science for Peace and Security Series C. Environmental Security, pp. 265–268, Springer, Dordrecht, 32nd NATO/SPS International Technical Meeting, 2013.
- Simpson, D. and Darras, S.: Global soil NO emissions for Atmospheric Chemical Transport Modelling: CAMS-GLOB-SOIL v2.2, *Earth System Science Data Discussions*, 2021, 1–35, <https://doi.org/10.5194/essd-2021-221>, 2021.
- Simpson, D., Ashmore, M., Emberson, L., and Tuovinen, J.-P.: A comparison of two different approaches for mapping potential ozone damage to vegetation. A model study, *Environmental Pollution*, 146, 715–725, <https://doi.org/10.1016/j.envpol.2007.05.011>, 2007.
- Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, A., and Wind, P.: The EMEP MSC-W chemical transport model — technical description, *Atmospheric Chemistry and Physics*, 12, 7825–7865, <https://doi.org/10.5194/acp-12-7825-2012>, 2012.
- Stadtler, S., Simpson, D., Schröder, S., Taraborrelli, D., Bott, A., and Schultz, M.: Ozone impacts of gas–aerosol uptake in global chemistry-transport models, *acp*, 18, 3147–3171, <https://doi.org/10.5194/acp-18-3147-2018>, 2018.

The Food and Land Use Coalition: Growing Better: Ten Critical Transitions to Transform Food and Land Use, <https://www.foodandlandusecoalition.org/>, 2019.

705 Thunis, P., Clappier, A., Pisoni, E., Bessagnet, B., Kuenen, J., Guevara, M., and Lopez-Aparicio, S.: A multi-pollutant and multi-sectorial approach to screening the consistency of emission inventories, *Geoscientific Model Development*, 15, 5271–5286, <https://doi.org/10.5194/gmd-15-5271-2022>, 2022.

Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmospheric Chemistry and Physics*, 12, 5447–5481, <https://doi.org/10.5194/acp-12-5447-2012>, 2012.

710 Turnock, S. T., Wild, O., Dentener, F. J., Davila, Y., Emmons, L. K., Flemming, J., Folberth, G. A., Henze, D. K., Jonson, J. E., Keating, T. J., Kengo, S., Lin, M., Lund, M., Tilmes, S., and O'Connor, F. M.: The impact of future emission policies on tropospheric ozone using a parameterised approach, *Atmospheric Chemistry and Physics*, 18, 8953–8978, <https://doi.org/10.5194/acp-18-8953-2018>, 2018.

715 UNECE: Mapping critical levels for vegetations. Revised Chapter 3 of the Manual on Methodologies and Criteria for Modelling and Mapping Critical Loads and Levels and Air Pollution Effects, Risks and Trends., ICP vegetation, CEH, UK, https://unece.org/fileadmin/DAM/env/documents/2017/AIR/EMEP/Final__new_Chapter_3_v2__August_2017_.pdf, [last access: March 2024], 2017.

UNEP: United Nations Environment Programme and Climate and Clean Air Coalition. Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions., Nairobi: United Nations Environment Programme, ISBN: 978-92-807-3854-4, Job No: DTI/2352/PA, 2021.

720 van Caspel, W. E., Simpson, D., Jonson, J. E., Benedictow, A. M. K., Ge, Y., di Sarra, A., Pace, G., Vieno, M., Walker, H. L., and Heal, M. R.: Implementation and evaluation of updated photolysis rates in the EMEP MSC-W chemistry-transport model using Cloud-J v7.3e, *Geoscientific Model Development*, 16, 7433–7459, <https://doi.org/10.5194/gmd-16-7433-2023>, 2023.

van Caspel, W. E., Klimont, Z., Heyes, C., and Fagerli, H.: Dataset supporting the article: Impact of methane and other precursor emission reductions on surface ozone in Europe: Scenario analysis using the EMEP MSC-W model, <https://doi.org/10.5281/zenodo.11184508>, 2024.

725 WHO: WHO global air quality guidelines. Particulate Matter (PM_{2.5} and PM₁₀), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide, World Health Organization Geneva, Switzerland, Licence: CC BY-NC-SA 3.0 IGO., 2021.

Wiedinmyer, C., Kimura, Y., McDonald-Buller, E. C., Emmons, L. K., Buchholz, R. R., Tang, W., Seto, K., Joseph, M. B., Barsanti, K. C., Carlton, A. G., and Yokelson, R.: The Fire Inventory from NCAR version 2.5: an updated global fire emissions model for climate and chemistry applications, *EGUsphere*, 2023, 1–45, <https://doi.org/10.5194/egusphere-2023-124>, 2023.

730 Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., et al.: Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems, *The lancet*, 393, 447–492, [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4), 2019.

Winiwarter, W., Höglund-Isaksson, L., Klimont, Z., Schöpp, W., and Amann, M.: Technical opportunities to reduce global anthropogenic emissions of nitrous oxide, *Environmental Research Letters*, 13, 014011, <https://doi.org/10.1088/1748-9326/aa9ec9>, 2018.

735 Zhang, X.: Emission standards and control of PM_{2.5} from coal power plants (ISBN: 978–92–9029–590–7), IEA Clean Coal Centre, London, 2018.