



# The optical properties of stratospheric aerosol layer perturbation of the Hunga volcano eruption of January 15th, 2022

Pasquale Sellitto<sup>1,2</sup>, Redha Belhadji<sup>1</sup>, Bernard Legras<sup>3</sup>, Aurélien Podglajen<sup>3</sup>, Clair Duchamp<sup>3</sup>

<sup>1</sup>Univ Paris Est Creteil and Université de Paris, CNRS, LISA, F-94010 Créteil, France

5 <sup>2</sup>Istituto Nazionale di Geofisica e Vulcanologia, Osservatorio Etno, Catania, Italy

<sup>3</sup>Laboratoire de Météorologie Dynamique (LMD-IPSL), CNRS, Sorbonne Université, ENS-PSL, École Polytechnique, Paris, France

Correspondence to: Pasquale Sellitto ([pasquale.sellitto@lisa.ipsl.fr](mailto:pasquale.sellitto@lisa.ipsl.fr))

**Abstract.** The Hunga volcano violently erupted on January 15th, 2022, and produced the largest stratospheric aerosol layer  
10 perturbation of the last 30 years. One notable effect of the Hunga eruption was the significant modification of the size  
distribution (SD) of the stratospheric aerosol layer with respect to background conditions and other recent moderate  
stratospheric eruptions, with larger mean particles size and smaller SD spread for Hunga. Starting from satellite-based SD  
retrievals, and the assumption of pure sulphate aerosol layers, in this work we calculate the optical properties of both  
15 background and Hunga-perturbed stratospheric aerosol scenarios using a Mie code. We found that the intensive optical  
properties of the stratospheric aerosol layer (i.e., single scattering albedo, asymmetry parameter, aerosol extinction per unit  
mass and the broad-band average Ångström exponent) were not significantly perturbed by the Hunga eruption, with respect to  
background conditions. The calculated Ångström exponent was found consistent with multi-instrument satellite observations  
of the same parameter. Thus, the basic impact of the Hunga eruption on the optical properties of the stratospheric aerosol layer  
was an increase of the stratospheric aerosol extinction (or optical depth), without any modification of the shortwave and  
20 longwave relative absorption, angular scattering and broad-band spectral trend of the extinction, with respect to background.  
This highlights a marked difference of the Hunga perturbation of the stratospheric aerosol layer and those from other larger  
stratospheric eruptions, like Pinatubo 1991 and El Chichon 1982. With simplified radiative forcing estimations, we show that  
the Hunga eruption produced an aerosol layer likely 3-10 times more effective in producing a net cooling of the climate system  
with respect to Pinatubo and El Chichon eruptions, due to more effective shortwave scattering. As intensive optical properties  
25 are seldom directly measured, e.g. from satellite, our calculations can support the estimation of radiative effects for the Hunga  
eruption with climate or offline radiative models.

## 1 Introduction

The Hunga volcano (Kingdom of Tonga, 20.54° S, 175.38° W) violently erupted on January 15th, 2022. Due to its very specific  
shallow submarine volcanological setting, and the subsequent interaction of seawater with the volcanic magma chamber, this  
30 eruption was characterised by a large explosivity (e.g. Poli and Shapiro, 2022) and injected volcanic material at altitudes as



large as 58 km, well into the deep stratosphere with parts up to the lower mesosphere (Carr et al., 2020). Basing on a wide array of satellite, ground-based and in situ measurements, it was demonstrated that the Hunga eruption produced the largest perturbation of the global stratospheric aerosol layer since the eruption of Mount Pinatubo in 1991 and the largest perturbation of stratospheric water vapour ever observed (Millàn et al., 2022, Sellitto et al., 2022a). An uncommonly fast conversion of volcanic sulphur dioxide emissions to sulphate aerosols (SA) was observed (e.g. Sellitto et al., 2023) and explained by the gas-to-particle kinetics acceleration due to very large water vapour concentrations, through modelling studies (Zhu et al., 2022). Thus, these very specific perturbations are likely due to the large amount of the injected water vapour, which contribute to generate an exotic environment for the perturbed stratosphere and drove peculiar chemical and microphysical evolution within the Hunga plume. The persistent Hunga stratospheric aerosol perturbations, at the southern-hemispheric scale (Legras et al., 2022, Taha et al., 2022), did not show any optical signature of ash, which were likely removed from the stratosphere over the very few days after the eruption (Legras et al., 2022, Sellitto et al., 2022a). The Hunga stratospheric aerosol perturbation can then solely be associated to SA. These effects proved long-lasting, with significantly perturbed aerosol extinctions extending well into the years 2022 and 2023 (Duchamp et al., 2023, Sellitto et al., 2023), and probably also in 2024. Using limb satellite observations, Duchamp et al. (2023) observed aerosol size distributions (SD) in the stratosphere significantly different than usual volcanic perturbations for moderate stratospheric eruptions and the background stratospheric aerosol layer, with significantly larger mean radii and smaller widths of the aerosol size distribution. The liquid-droplets nature and the relatively large mean size of the Hunga aerosol particles was also confirmed with direct measurements with sondes balloons (Kloss et al., 2022). In the absence of volcanic or other perturbations, a local maximum of the vertical aerosol distribution is found in the lower stratosphere, due to troposphere-to-stratosphere exchanges flux of sulphur-containing aerosol precursors at low latitudes and a very limited aerosol sink at these altitudes (Kremser et al., 2016). Thus, a background stratospheric aerosol layer can be defined in the absence of episodic perturbation of this layer, i.e. stratospheric volcanic eruptions and pyroconvective fires. Secondary SA largely dominate the composition of the background aerosol layer (Kremser et al., 2016). The stratospheric aerosol layer perturbations induced by the Hunga eruption are expected to have an impact on the optical properties of the stratospheric aerosol layer and on the Earth radiative balance, producing radiative and climatic impacts (Sellitto et al., 2022a).


In this paper, we use the size distributions (in the following, SD) determined by Duchamp et al. (2023), for both the Hunga-perturbed and the background stratospheric aerosol layer, and we derive their optical properties. These optical properties and their possible effects on the radiative balance are then compared with those from the major recent past eruptions, El Chichon in 1982 and Pinatubo in 1991. Our results are expected to contribute to the new estimations of the radiative forcing of the Hunga eruption, which are presently ongoing to characterise the long-term radiative impacts of this event. This paper is structured as follows: in Sect. 2 the data and methods used in the paper are described; in Section 3 results are shown and discussed; conclusions are drawn in Section 4.



## 2 Data and Methods

### 2.1 Chemical composition and refractive index

65 As discussed in the introduction, both the background stratospheric aerosol layer and its perturbation brought by the Hunga eruption can be solely characterised, in terms of composition, as secondary SA particles. Thus, we model both the background and Hunga-perturbed stratospheric aerosol layers as composed of SA. These particles are usually represented as spherical liquid droplets of binary aqueous solution of sulphuric acid ( $\text{H}_2\text{SO}_4$ ) as done, e.g., by Sellitto and Legras (2016). The link between the chemical composition and the optical properties of a specific composition is provided by the complex refractive index (CRI). Among the available laboratory measurements of SA CRI, we have selected, for this work, the one from Hummel et al. (1988). This is one of the few datasets that extend the CRI spectra in both the shortwave (SW) and the longwave (LW) spectral ranges, from the ultraviolet (UV) to part of the far infrared (FIR), for representative stratospheric conditions in terms of the  $\text{H}_2\text{SO}_4$  mass mixing ratio and temperature. For these latter, a 75%  $\text{H}_2\text{SO}_4$  mixing ratio and a 215 K temperature are selected in this work. Both the background stratospheric SA (e.g. Kremser et al., 2016) and the Hunga perturbations (Duchamp et al., 2023) are characterised by very acidic particles with typically 70-80%  $\text{H}_2\text{SO}_4$  mass mixing ratio. Among the available particles temperatures in the Hummels et al. (1988) database, a temperature of 215 K is selected as the more suitable to represent mid-stratospheric conditions. Real and imaginary parts of CRI used in this work are shown in Fig. 1.



80 **Figure 1: Real (a) and imaginary part (b) of the complex refractive index of a SA layer with 75%  $\text{H}_2\text{SO}_4$  mass mixing ratio and 215 K temperature, from Hummel et al., 1988.**

### 2.2 Size distribution


The number density SD  $n(r)$  is defined so that  $n(r)dr$  is the number of particles per unit volume, in the aerosol layer, with radius between  $r$  and  $r+dr$ . All SD, in this work, are modelled as mono-modal log-normal distributions (Eq. 1). In Eq. 1,  $N_0$  is the



total number concentration (in particles per  $\text{cm}^{-3}$ ),  $r_m$  is the median radius and  $S=\ln\sigma$  is the SD spread i.e. the unitless standard  
85 deviation of  $\ln(r/r_m)$ .

$$n(r) = \frac{N_0}{r \ln \sigma \sqrt{2\pi}} e^{-\frac{1}{2} \left( \frac{\ln\left(\frac{r}{r_m}\right)}{\ln \sigma} \right)^2} \quad (\text{Eq. 1})$$

Both background and Hunga-perturbed typical SD are derived from the results of Duchamp et al. (2023). In this work, SD  
parameters of a mono-modal log-normal SD, i.e.  $N_0$ ,  $r_m$  and  $S$  in Eq. 1, are obtained using SAGE III/ISS (Stratospheric Aerosol  
and Gas Experiment on the International Space Station) satellite observations with the method initially developed by Wrana  
90 et al. (2021). Duchamp et al. (2023) applied that method to the Hunga plume and extended back these retrievals to immediately  
previous unperturbed periods, so to derive SD parameters also for a representative background stratospheric aerosol layer. It  
is usually convenient, in remote sensing applications, to define an effective radius  $r_e$  (the cube particle radius divided by the  
square particle radius, averaged over the SD);  $r_e$  is directly linked to the extinction of the aerosol layer. For a mono-modal log-  
normal SD, an effective radius can be defined as  $r_e = r_m e^{2.5 \ln^2 \sigma}$ . For the background stratospheric aerosol layer a typical  
95 combination of  $r_m = 0.20 \mu\text{m}$  and  $\sigma = 1.50$  is considered (Fig. 2a and Tab. 1). The Hunga eruption produced an increase of the  
mean particle size and decrease of the spread of the particle SD with respect to background conditions (Duchamp et al., 2023).  
Typical values of Hunga-perturbed stratospheric aerosol layers are used here, with  $r_e$  varying between 0.35 and 0.45  $\mu\text{m}$  and  
 $\sigma$  varying between 1.20 and 1.30 (Fig. 2a and Tab. 1). In Fig. 2,  $N_0$  is fixed to 1 particle/ $\text{cm}^3$  for the background and all Hunga  
100 also considered (Fig. 2b and Tab. 1).



105 **Figure 2:** Typical background (black curve) and Hunga-perturbed SDs (yellow, red and dark red curves), modelled as mono-modal log-normal SDs with  $r_e$  and  $\sigma$  estimated using SAGE III/ISS by Duchamp et al. (2023) (a). Background (black curve), Hunga-perturbed (red curve, case for  $r_e = 0.40 \mu\text{m}$  and  $\sigma = 1.25$ ), and Mount Pinatubo- and El Chichon-perturbed SD (blue and sky blue curves, respectively). Log-normal parameters for Pinatubo and El Chichon SD are taken from Asano, 1993 and Hoffmann and Rosen, 1983, respectively.

For a SA layer of given SD, the effective SA mass  $M_e$  can be calculated as done in Eq. 2, where  $\rho$  is the mass density of SA (here taken as  $1.75 \text{ g/cm}^3$ ) and  $N_e = N_0 e^{-3 \ln^2 \sigma}$  is the mono-modal log-normal effective number density. The  $M_e$  values for the SDs used in the present work are also indicated in Tab. 1.

110 
$$M_e = \frac{4}{3} \pi r_e^3 \rho N_e \quad (\text{Eq. 2})$$

**Table 1:** Parameters  $r_m$ ,  $r_e$ ,  $\sigma$  and  $M_e$  for the SD in Fig. 2a for the background and Hunga-perturbed stratospheric aerosol layers used in this manuscript. All SD in Fig. 2a are for a value of  $N_0 = 1 \text{ particles/cm}^3$ . In the table,  $N_0$  values for each SD are also reported, in the case of a fixed  $M_e = 1 \mu\text{g/m}^3$  (last column).

$r_m$ ( $\mu\text{m}$ )	$r_e$ ( $\mu\text{m}$ )	$\sigma$	$M_e$ ( $\mu\text{g/m}^3$ )	$N_0$ ( $\#/ \text{cm}^3$ )
-------------------------	-------------------------	----------	-----------------------------	-----------------------------



<b>Background</b>				
0.20	0.30	1.50	0.12	8.27
<b>Hunga-perturbed</b>				
0.32	0.35	1.20	0.28	3.52
0.31	0.35	1.25	0.27	3.69
0.28	0.35	1.30	0.26	3.91
0.37	0.40	1.20	0.42	2.36
0.35	0.40	1.25	0.40	2.47
0.34	0.40	1.30	0.38	2.62
0.41	0.45	1.20	0.60	1.65
0.40	0.45	1.25	0.55	1.74
0.38	0.45	1.30	0.54	1.84
<b>Pinatubo-perturbed (Asano, 1993)</b>				
0.60	0.90	1.50	-	0.31
<b>El Chichon-perturbed (Hoffmann and Rosen, 1983)</b>				
0.72	1.71	1.80	-	0.08

### 2.3 Optical properties calculation

115 The optical properties of background and Hunga-perturbed (as well as for El Chichon and Pinatubo eruptions) stratospheric aerosol layers are calculated with the scheme shown in Fig. 3. Starting from CRI and SD input data described in Sects. 2.1-2, a Mie code estimates the extinction coefficient ( $\beta_{\text{ext}}$ ), single scattering albedo (SSA) and asymmetry parameter ( $g$ ) spectra for both cases, and for Mount Pinatubo and El Chichon as a further comparison. As Mie code, the IDL (Interactive Data Language, <http://www.exelisvis.com/ProductsServices/IDL.aspx>) Mie scattering routines of the Earth Observation Data Group of the Department of Physics of Oxford University are used (<https://eodg.atm.ox.ac.uk/MIE/>). The outputs of the Mie calculations are intended to represent the overall extinction of typical stratospheric background and Hunga-perturbed layers, as well as their absorption/scattering properties (through SSA) and angular distribution of the scattered radiation (through  $g$ ). The SSA is the

120



ratio of the scattering to extinction (i.e. scattering plus absorption) efficiency factors  $Q_{\text{sca}}$  and  $Q_{\text{ext}}$  obtained with Mie calculations (van de Hulst, 1957). Values of the SSA approaching 1.0 (0.0) point at pure scattering (absorbing) particles. The  
125  $g$  parameter is the mean value of the cosine of the scattering angle weighted through the scattering phase function obtained with Mie calculations (van de Hulst, 1957). Values of the  $g$  parameter approaching 1.0 (0.0) point at pure forward (isotropic) scattering. The overall extinction of the layers, measured by  $\beta_{\text{ext}}$  has been calculate with two different assumptions: a fixed  $N_0$  and a fixed  $M_e$ . With this latter, extinction spectra per unit mass concentration ( $\beta_{\text{ext}}/M_e$ ) have also been derived. The SSA,  $g$  and  $\beta_{\text{ext}}/M_e$  are intensive optical properties of the aerosol layer, i.e. they do not depend on the injected aerosol mass. It was  
130 discussed previously that these optical parameters, together with surface reflectivity information, fed to a radiative transfer model (RTM), are sufficient to describe the radiative properties and impacts of an aerosol layer, like the instantaneous radiative forcing and the vertical profiles of radiative/cooling heating rates (e.g. Sellitto et al., 2022b, 2023). In this work, optical properties in both the solar shortwave (SW) and the terrestrial longwave (LW) spectral ranges are estimated, so to produce results usable in RTM when interested in the whole radiant energy spectrum in Earth's atmosphere. The output spectral range  
135 of the optical properties is somewhat limited by the available spectral range of the input CRI. In our case, we are limited at 25  $\mu\text{m}$  on the upper end, due to available SA CRIs in the literature, so a part of the far infrared range is not represented in this work and this latter impact is probably underestimated.




Figure 3: Scheme of the optical properties calculations used in this work.



## 140 3 Results and discussion

### 3.1 Impact of size distribution on the optical properties

Figure 4 shows  $\beta_{ext}/N_0$  and  $\beta_{ext}/M_e$  calculation, for the background and Hunga-perturbed scenarios, as described in Sect. 2. The extinction is an extensive optical property, which depends on the aerosol layer mass (the SA mass, in the present case). Thus, large differences in  $\beta_{ext}/N_0$  among background and Hunga-perturbed scenarios can be observed (Fig. 4a). This is principally  
145 due to the larger SA mass, for a fixed number density  $N_0=1.0 \text{ \#/cm}^3$ , of the larger Hunga particles with respect to the smaller background particles. The effective mass concentration varies from  $0.12 \text{ \mu g/cm}^3$ , for the background layer, to up to  $0.60 \text{ \mu g/cm}^3$ , for the Hunga-perturbed layers (see Tab. 1). On the contrary,  $\beta_{ext}/M_e$  is much less variable depending on the scenario, except at very short wavelengths in the SW (Fig. 4b). In general, a clear exponential decrease of  $\beta_{ext}$  can be observed in the UV/Vis and part of the NIR (thus, in most on the SW spectral range), following the empirical Ångström law in Eq. 3,  
150 where  $\lambda$  is the wavelength ( $\lambda_{ref}$  is a reference wavelength, in some cases  $1 \text{ \mu m}$ ) and AE is the Ångström exponent.

$$\beta_{ext}(\lambda) = \beta_{ext}(\lambda_{ref})\lambda^{-AE} \quad (\text{Eq. 3})$$

At longer wavelengths, absorption features of SA particles appear, including the peculiar MIR signatures at  $8.0$  to  $11.0 \text{ \mu m}$ . These absorption features can be associated with the roto-vibrational absorption bands of the undissociated  $\text{H}_2\text{SO}_4$  in the concentrated solution droplets discussed e.g. by Sellitto and Legras (2016). More absorption features are visible in the NIR,  
155 from about  $3.0$  to  $6.0 \text{ \mu m}$ , and in the FIR, from  $15.0$  to  $18.0 \text{ \mu m}$ . The two regimes, dominated by scattering in the SW, for wavelengths shorter than about  $3.0 \text{ \mu m}$ , and by absorption in the LW, for longer wavelengths, is discussed further in the following (associated with SSA), and is linked to a large variability of  $\beta_{ext}/M_e$ . This latter ranges between values up to  $4.5 \cdot 10^{-3} \text{ km}^{-1}$  per unit effective mass, below  $3.0 \text{ \mu m}$ , to less than  $1.0 \cdot 10^{-3} \text{ km}^{-1}$  per unit effective mass, above  $3.0 \text{ \mu m}$ .


Figure 5 shows the SSA (Fig. 5a) and  $g$  (Fig. 5b) calculation, for the background and Hunga-perturbed scenarios. One first  
160 take-away message from these results is that both SSA and  $g$  are not perturbed significantly by Hunga eruption, and then their absolute values and variability are similar for background and Hunga-perturbed scenarios. The two scattering- and absorption-dominated regimes, discussed for the aerosol extinction, can be seen here with SSA approximately  $1.0$  (pure scattering aerosol layers), for wavelength shorter than about  $3.0 \text{ \mu m}$ , and sharply decreasing to values lower than about  $0.2$ , for wavelength longer than about  $3.0 \text{ \mu m}$ . The SSA approaches values of  $0.0$  (pure absorbing particles), for wavelengths longer than about  $6.0 \text{ \mu m}$ .  
165 The  $g$  parameter, starting from values of about  $0.6$  to  $0.8$ , steeply decreases to values lower than  $0.1$ , for wavelengths longer than  $6.0 \text{ \mu m}$ . This can be associated with a markedly dominating forward scattering in the SW and a quasi-isotropic scattering in the LW.

These results are summarized in Fig. 6, where spectral-band average values of the intensive optical properties  $\beta_{ext}/M_e$ , SSA and  $g$ , for background and Hunga-perturbed layers, are shown for the UV/Vis, NIR, MIR and FIR broad-band ranges. Large  
170 values for the three intensive parameters are found in the UV/Vis, thus pointing at very extinction-effective layers, dominated






by a largely forward scattering. While this characteristic behaviour of the aerosol layers stands for the whole SW, the extinction-efficiency markedly decreases in the NIR. In the LW, the optical characterisation of the layers is more homogeneous, with relatively small extinction dominated by absorption and a small quasi-isotropic scattering, with small differences between the MIR and the FIR.



175

**Figure 4:** Extinction coefficient spectra  $\beta_{\text{ext}}$  at fixed number concentration  $N_0$  (a) and effective mass  $M_e$  (b), for background (black curve) and Hunga-perturbed aerosol layers (from yellow, red and dark red dotted, dashed and solid curves – see corresponding values of the  $r_e$  and  $\sigma$  assumption in the figures captions). Grey vertical dotted lines in panel a indicate the two wavelength bands used to average the extinction to calculate the UV/Vis-to-MIR AE of Fig. 7.



180

**Figure 5: Same as Fig. 4 but for single scattering albedo SSA (a) and asymmetry parameter g (b) spectra.**


More in general, the most apparent effect, visible in Figs. 4-6, is that all intensive optical properties of Hunga-perturbed layer are very similar to those in background conditions. It can be concluded that, despite the significant perturbation in stratospheric aerosol SD and the large SA injected mass, the Hunga eruption had minimal effects on the intensive optical properties of the stratospheric aerosol layer. Thus, the only impact of Hunga eruption on the optical properties of the stratospheric aerosol layer is the marked increase in the overall extinction and the aerosol optical depth (AOD). For this latter property, Hunga produced the largest global perturbation since Mount Pinatubo (Philippines) eruption in 1991 (Sellitto et al., 2022a). The specific absorption and scattering properties of the stratospheric aerosol layer were not significantly perturbed in the SW and LW by the Hunga eruption, as shown by SSA and g calculations of Fig. 5 and 6b,c. It has to be noted that Hunga perturbations on the stratospheric aerosol size distributions, though significant, are much smaller than for other stronger recent eruptions like El

185

190



Chichon in 1982 (Hofmann and Rosen, 1983) or Pinatubo in 1991 (e.g., Asano 1993), see respective SDs in Fig. 2b. In these cases, a fraction of relatively large particles (larger than  $1.0 \mu\text{m}$ ) was present, which is not the case for the Hunga eruption (see Fig. 2b). We performed additional Mie calculations for Pinatubo and El Chichon SD, using the SD parameters of Asano (1993) and Hoffmann and Rosen (1983), respectively, summarised in Tab. 1. In both these cases, the UV/Vis  $\beta_{\text{ext}}/M_e$  is significantly smaller, and the overall LW SSA and  $g$  are significantly larger, than for Hunga and stratospheric background (Fig. 6). Such a change of optical regime implies a stronger LW climate warming effect, that can counterbalance the SW cooling. More in general, Lacis et al. (1992) have shown that a marked change in optical and radiative regime occurs for volcanic aerosol layers with mean size exceeding  $1.0 \mu\text{m}$ , which is the case for Pinatubo and El Chichon but not for the Hunga perturbations. This change in the SW/LW optical properties can result in less effective cooling at top of the atmosphere or even in aerosol-related warming, which could have been initially the case for Pinatubo and El Chichon eruptions (before a marked and relatively long-term cooling effect, due to the removal of larger aerosol particles) but was likely not the case for Hunga, even in the first phases after the eruption (e.g. Sellitto et al., 2022a). The mean particle size for Hunga was larger than any post-Pinatubo stratospheric eruption (Wrana et al., 2023). Most post-Pinatubo eruptions perturbed the stratospheric aerosol layer with a decrease of mean particle size, rather than an increase. The radiative effects of the Hunga-related aerosols SD perturbations are discussed more in Sect. 3.2.




**Figure 6: Band average (UV/Vis, NIR, MIR and FIR) extinction coefficient per unit effective mass (a), single scattering albedo (b) and asymmetry parameter (c), for background (black circles with error bars) and Hunga-perturbed (averaged over all scenarios in e.g. Fig. 2, red circles with error bars) stratospheric aerosol layer. Band-average values of these intensive aerosol optical properties are also shown for Pinatubo 1991 (blue circles with error bars) and El Chichon (sky blue circles with error bars). UV/Vis (ultra-violet and visible): 0.30-0.85  $\mu\text{m}$ ; NIR (near infrared): 0.85-3.0  $\mu\text{m}$ ; MIR (mid infrared): 3.0-15.0  $\mu\text{m}$ ; FIR (far infrared): 15.0-25.0  $\mu\text{m}$ .**

210

Another intensive aerosol optical parameter is the Ångström exponent AE, see also Eq. 3. An average AE can be obtained by combining aerosol extinction or AOD information at two different wavelengths. In Sellitto et al. (2023) the average UV/Vis-  
 215 to-MIR AE was calculated using combination of OMPS-LP (Ozone Mapping and Profiler Suite – Limb Profiler) and IASI (Infrared Atmospheric Sounding Interferometer) observations, at 0.7 and 8.5  $\mu\text{m}$ , respectively. Figure 7 shows this value compared with our calculated background and Hunga-perturbed scenarios. As for other intensive optical properties shown above, the stratospheric aerosol layer UV/Vis-to-MIR AE is not significantly perturbed by the Hunga eruption, and background and Hunga-perturbed values stay around a value of 1.0. These results are not dissimilar from the observed value ( $1.13 \pm 0.23$ ),  
 220 even if slightly smaller. Thus, this shows the consistency of our calculated and the observed optical properties for the Hunga plume. It has to be noted that Taha et al. (2023) have shown that the observed Vis AE, i.e. estimated with two different Vis



bands of the OMPS-LP instrument (0.52 and 1.02  $\mu\text{m}$ ), is significantly perturbed by the Hunga eruption. The AE estimation from observations is very sensitive to the selection of the two wavelengths used to derive it. Using the same wavelengths as in Taha et al. (2023), we obtain Vis AE of 1.35 and 0.80 for the background and Hunga-perturbed scenarios. This is very consistent with past theoretical studies, i.e. the one of Schuster et al. (2006) (see their figure 4a, with effective radii 0.30  $\mu\text{m}$  and 0.40-0.45  $\mu\text{m}$ , respectively). The variability of the UV/Vis AE, in terms of the selected background or Hunga-perturbed scenarios, can also be seen Fig. 4, reflected by the different slopes of the spectral variability of the extinction coefficient.



**Figure 7: Average UV/Vis-to-MIR Ångström exponent for Hunga and the three  $r_e$  assumptions ( $r_e=0.35$ , 0.40 and 0.45  $\mu\text{m}$ ) and their average (HTHH) and background (BG) conditions. The Ångström exponents are calculated using Eq. 3 and averaging around the wavelengths 0.7 and 8.5  $\mu\text{m}$ , see Fig. 4a. The value observed with the OMPS-LP/IASI coupling, discussed by Sellitto et al. (2024), is also shown (Obs). A 20% error bar is associated with this latter estimation.**

### 3.2 Impact of optical properties on the radiative forcing


A simple parameterisation of the top-of-atmosphere (TOA) radiative forcing is used to estimate the radiative impact, through optical properties perturbations, of changing SD of stratospheric aerosol by Hunga eruption. For partly absorbing aerosols in a layer over a given surface of spectral reflectivity  $R_s$  and placed in a stratified atmosphere of spectral transmissivity  $T_{\text{atm}}$ , a broad-band TOA radiative forcing  $\Delta F$  per unit SA mass can be defined as in Eq. 4. This quantity is estimated at the four broad-band spectral ranges, as defined, e.g., in Fig. 6 (UV/Vis, NIR, MIR and TIR) and then these broad-band estimations are added up to obtain a total  $\Delta F$ . In Eq. 4, the optical properties  $\beta_{\text{ext}}/M_e$  and SSA are the same as in Fig. 6. The optical parameter  $b$  is the hemispheric backscatter ratio, which can be derived using our asymmetry parameter  $g$  estimations, using the method described by Marshall et al. (1995). The aerosols have been put in a 1-km deep layer. In the parameterisation,  $S$  is the radiation source, taken as pure Planck functions at temperatures 5770 K, in the SW range (UV/Vis and NIR broad bands) and at the



temperature 300 K, in the LW range (MIR and FIR broad bands), to simulate the solar and the terrestrial radiation sources. The solar S component is then scaled at the mean sun-Earth distance. The diurnal cycle of S was not considered in this example. 245 The underlying surface have been considered as marine ( $R_s$  0.05 in the UV/Vis and FIR and 0.00 in the NIR and MIR) and average values of  $T_{atm}$  have also been considered (0.6, 0.5, 0.5 and 0.2 in the UV/Vis, NIR, MIR and FIR).

$$\Delta F = -S(\lambda)T_{atm}(\lambda) \frac{\beta_{ext}(\lambda)}{M_e} SSA(\lambda)b(\lambda) \left( (1 - R_s(\lambda))^2 + 2R_s(\lambda) \left( \frac{1-SSA(\lambda)}{b(\lambda)SSA(\lambda)} \right) \right) \Delta\lambda \quad (\text{Eq. 4})$$

The SW (UV/Vis + NIR), LW (MIR + FIR) and total (SW+LW) radiative forcings obtained with Eq. 4 and our estimated optical parameters are shown in Fig. 8, for a background stratospheric aerosol layer, and layers perturbed by Hunga, Pinatubo and El Chichon eruptions. Two different radiative regimes can be observed in Fig. 6: one clearly dominated by the SW scattering, thus a large negative radiative forcing per unit mass, for smaller effective radii, and one less and less dominated by the decreasing SW scattering with increasing mean size and the increases in importance of the LW warming effect due to LW absorbing effect, for larger effective radii. In this latter case, the total radiative forcing per unit SA mass decreases significantly with respect to background conditions. In addition, SW/LW absolute ratio of the  $\Delta F$  is approximately 99, 98, 80 and 60% for 255 the background, Hunga, Pinatubo and El Chichon scenarios. These results point at a very different overall radiative regimes between background and Hunga, with respect to Pinatubo and El Chichon. The Hunga eruption did not modify significantly the radiative regime of the stratospheric aerosol layer with respect to background conditions, with values around  $-0.15 \text{ W/m}^2$  per  $\mu\text{g/m}^3$  SA, dominated by SW scattering. This is 3 times more effective, per unit SA mass, than Pinatubo eruption (about  $-0.05 \text{ W/m}^2$  per  $\mu\text{g/m}^3$  SA) and nearly 10 times more effective than El Chichon eruption (about  $-0.02 \text{ W/m}^2$  per  $\mu\text{g/m}^3$  SA). For 260 these latter, the SW cooling markedly decreased, as well as the SW/LW ratio, with respect to background and Hunga eruption. Our results are consistent with previous studies, in particular the one of Lacis et al. (1992) who showed that the cooling potential at TOA of stratospheric aerosols significantly decreases for effective radii larger than about  $1.0 \mu\text{m}$ , and can even switch from negative (cooling) to positive (warming) for effective radii larger than  $2.0 \mu\text{m}$  (see their Fig. 2). It is important to recall that Hunga eruption is characterised by much less injected SA mass than Pinatubo and El Chichon eruptions so, despite 265 the larger radiative effectiveness of its stratospheric aerosol perturbations, in terms of the SD, its overall radiative effect is expected to be smaller than Pinatubo and El Chichon.



270 **Figure 8: TOA RF ( $\Delta F$ ) in the SW (negative, dark blue lines), LW (dark red lines) and total SW+LW (negative, pink lines and symbols) as a function of the effective radius for: background conditions (pink dot), Hunga (pink triangle), and Pinatubo and El Chichon (pink large squares), see text for details.**

Optical properties of aerosol layers are needed, in radiative calculations with both climate models and offline radiative models, for a given source of radiative forcing to estimate its impacts (e.g. Sellitto et al., 2022b, 2023). Our results provide a ready-to-use benchmark for the radiative impacts estimations of the Hunga eruption versus stratospheric aerosol background. Thus, we provide as Supplementary datasets optical properties: (1) computed at the wavelengths at which the refractive indices of Hummel et al. are available, and (2) averaged over the 30 broad bands of the Rapid Radiative Transfer Model (RRTM, Iacono et al., 2008) in the ECMWF ECRAD implementation (Hogan, and Bozzo, 2018), for future studies on the Hunga radiative impacts. Averaged values for the UV/Vis, NIR, MIR and FIR bands are also reported in Tab. 2 (same as Fig. 6). Caution must be taken for the radiative impacts estimation of Hunga eruption because of the additional important radiative impacts of water vapour injections, that can be even larger than aerosol impacts during the first months (Sellitto et al., 2022). This additional radiative effect for the Hunga eruption was not present in recent stratospheric volcanic eruptions from subaerial volcanoes.

280 **Table 2: Summary of band-average ready-to-use optical properties inputs for radiative calculations**

Spectral interval ( $\mu\text{m}$ )	AE	SSA	g
<b>Background</b>			
0.3-0.8	0.94	0.99±0.00	0.72±0.02
0.8-3.0		0.88±0.25	0.45±0.15
3.0-15.0		0.05±0.04	0.08±0.06



15.0-25.0		0.01±0.01	0.01±0.00
<b>Hunga-perturbed</b>			
0.3-0.8	0.97±0.02	0.99±0.00	0.73±0.05
0.8-3.0		0.89±0.24	0.46±0.19
3.0-15.0		0.06±0.06	0.06±0.06
15.0-25.0		0.01±0.01	0.01±0.00

#### 4 Conclusions

285 The submarine Hunga volcano violently erupted on January 15<sup>th</sup>, 2022 and produced the largest stratospheric aerosol layer  
 perturbation, in terms of the aerosol extinction and the injected aerosol mass, since the climate-relevant eruption of Pinatubo  
 volcano in 1991. One notable feature of this perturbation is the significant modification of the stratospheric aerosol SD, with  
 larger particles (0.35-0.45  $\mu\text{m}$  effective radius) and a smaller SD width (1.20-1.30 spread in a modelled mono-modal size  
 distribution) than the background stratospheric aerosol layer and than for most other post-Pinatubo stratospheric eruptions. In  
 this paper, using a Mie code and the assumption of pure SA layers, we calculate the optical properties of the Hunga-perturbed  
 290 stratospheric aerosol layer and compare with those for a background scenario. We found that, despite the sensible impact on  
 the aerosol SD, the Hunga eruption have only a minor impact on the intensive optical properties, namely the aerosol extinction  
 per unit effective mass  $\beta_{\text{ext}}/M_e$ , the single scattering albedo SSA, the asymmetry parameter  $g$  and the broad-band UV/Vis-to-  
 MIR Ångström exponent AE. Thus, while producing an historical perturbation on the stratospheric aerosol extinction and  
 optical depth, the Hunga eruption did not modify the absorption and angular-scattering properties of the stratospheric aerosol  
 295 layer. Our UV/Vis-to-MIR AE calculations are consistent with those observed with the combination of OMPS-LP and IASI  
 satellite observations. We further calculate intensive aerosol optical properties for past eruptions of Pinatubo 1991 and El  
 Chichon 1982 and found that these latter events produced enough particles with radii  $> 1 \mu\text{m}$  to change their SW/LW optical  
 regimes with respect to stratospheric background, which was not the case for Hunga. We demonstrate here, with a simplified  
 radiative forcing parameterisation, that the Hunga eruption has likely produced stratospheric aerosol layers with a 3-10 times  
 300 larger cooling potential than the Pinatubo and El Chichon eruptions, due to SD-related different radiative regimes. Hunga-  
 related stratospheric aerosol perturbations are more effective in SW cooling than Pinatubo- and El Chichon-related  
 perturbations. Values as large as  $-0.15 \text{ W/m}^2$  per  $\mu\text{g/m}^3$  of injected SA are found for Hunga eruption ( $-0.05$  and  $-0.02 \text{ W/m}^2$   
 per  $\mu\text{g/m}^3$  of injected SA are obtained for Pinatubo and El Chichon, respectively). Our optical properties calculations are a





ready-to-use dataset for future estimations of the radiative impacts of the Hunga eruption compared with background,  
305 combined with pertinent water vapour observations, which is another important forcing agent for this submarine eruption.

### Author contribution

PS conceived and performed the study, and wrote the first version of the manuscript. All authors participated to the discussions of the results and contributed to the final version of the manuscript?

### Competing interests

310 At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics. The peer-review process will be guided by an independent editor. The authors also have no other competing interests to declare.

### References

- Asano, S. (1993), Estimation of the size distribution of Pinatubo volcanic dust from Bishop's ring simulations, *Geophys. Res. Lett.*, 20(6), 447–450, doi:10.1029/93GL00512.
- 315 Duchamp, C., Wrana, F., Legras, B., Sellitto, P., Belhadji, R., & von Savigny, C. (2023). Observation of the aerosol plume from the 2022 Hunga Tonga—Hunga Ha'apai eruption with SAGE III/ISS. *Geophysical Research Letters*, 50, e2023GL105076. <https://doi.org/10.1029/2023GL105076>
- Carr, J. L., Horvath, A., Wu, D. L., & Friberg, M. D. (2022). Stereo Plume Height and Motion Retrievals for the Record-Setting Hunga Tonga-Hunga Ha'apai Eruption of 15 January 2022. *Geophysical Research Letters*, 49, 270 e2022GL098131.  
320 doi: 10.1029/2022GL098131
- Hofmann, D. J., Rosen, J. M. (1983) Sulfuric Acid Droplet Formation and Growth in the Stratosphere After the 1982 Eruption of El Chichón. *Science* 222, 325–327(1983). DOI:10.1126/science.222.4621.325
- Hogan, R. and Bozzo, A.: ECRAD: A new radiation scheme for the IFS, <https://doi.org/10.21957/whntqkfdz>, 2016. [5] Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *Journal of Advances in Modeling Earth Systems*, 10, 1990–2008, <https://doi.org/10.1029/2018MS001364>, 2018.  
325
- Hummel, J. R., Shettle, E. P., Longtin, D. R. (1988). A new background stratospheric aerosol model for use in atmospheric radiation models}, Tech. Rep., Provided by the SAO/NASA Astrophysics Data System, url: <https://ui.adsabs.harvard.edu/abs/1988nbsa.rept.....H>



- Iacono, M. J., J. S. Delamere, E. J. Mlawer, M. W. Shephard, S. A. Clough, and W. D. Collins (2008), Radiative forcing by  
330 long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.*, 113, D13103,  
doi:10.1029/2008JD009944.
- Kloss, C., Sellitto, P., Renard, J.-B., Baron, A., Bègue, N., Legras, B., et al. (2022). Aerosol characterization of the stratospheric  
plume from the volcanic eruption at Hunga Tonga 15 January 2022. *Geophysical Research Letters*, 49, e2022GL099394.  
<https://doi.org/10.1029/2022GL099394>
- 335 Lacis, A., Hansen, J. and Sato, M. (1992) Climate forcing by stratospheric aerosols. *Geophys. Res. Lett.*, 19, 1607-1610,  
doi:10.1029/92GL01620.
- Legras, B., Duchamp, C., Sellitto, P., Podglajen, A., Carboni, E., Siddans, R., . . ., Ploeger, F. (2022). The evolution and  
dynamics of the Hunga Tonga–Hunga Ha’apai sulfate aerosol plume in the stratosphere. *Atmospheric Chemistry and Physics*,  
22 (22), 14957–14970. doi: 10.5194/acp-22-14957-2022
- 340 Marshall, S. F., D. S. Covert, and R. J. Charlson (1995), Relationship between asymmetry parameter and hemispheric  
backscatter ratio: Implications for climate forcing by aerosols, *Appl. Opt.*, 34(27), 63066311.
- Millán, L., Santee, M. L., Lambert, A., Livesey, N. J., Werner, F., Schwartz, M. J., et al. (2022). Hunga Tonga-Hunga Ha’apai  
hydration of the stratosphere. *Geophysical Research Letters*, 49, e2022GL099381.
- Poli, P., & Shapiro, N. M. (2022). Rapid Characterization of Large Volcanic Eruptions: Measuring the Impulse of the Hunga  
345 Tonga Ha’apai Explosion From Teleseismic Waves. *Geophysical Research Letters*, 49 (8), e2022GL098123.  
doi:10.1029/2022GL098123
- Schuster, G. L., O. Dubovik, and B. N. Holben (2006), Angstrom exponent and bimodal aerosol size distributions, *J. Geophys.*  
*Res.*, 111, D07207, doi:10.1029/2005JD006328.
- Sellitto, P. and Legras, B.: Sensitivity of thermal infrared nadir instruments to the chemical and microphysical properties of  
350 UTLS secondary sulfate aerosols, *Atmos. Meas. Tech.*, 9, 115–132, <https://doi.org/10.5194/amt-9-115-2016>, 2016.
- Sellitto, P., Podglajen, A., Belhadji, R., Boichu, M., Carboni, E., Cuesta, J., . . ., Legras, B. (2022a). The unexpected radiative  
impact of the Hunga Tonga eruption of 15th January 2022. *Communications Earth & Environment*, 3 (1), 288, doi:  
10.1038/s43247-022-00618-z
- Sellitto, P., Belhadji, R., Kloss, C., and Legras, B. (2022b) Radiative impacts of the Australian bushfires 2019–2020 – Part 1:  
355 Large-scale radiative forcing, *Atmos. Chem. Phys.*, 22, 9299–9311, <https://doi.org/10.5194/acp-22-9299-2022>



- Sellitto, P., Belhadji, R., Cuesta, J., Podglajen, A., and Legras, B. (2023) Radiative impacts of the Australian bushfires 2019–2020 – Part 2: Large-scale and in-vortex radiative heating, *Atmos. Chem. Phys.*, 23, 15523–15535, <https://doi.org/10.5194/acp-23-15523-2023>
- 365 Sellitto, P., Siddans, R., Belhadji, R., Carboni, E., Legras, B., Podglajen, A., Duchamp, C., Kerridge, B. (2023) Observing the SO<sub>2</sub> and Sulphate Aerosol Plumes from the 2022 Hunga Tonga-Hunga Ha'apai Eruption with IASI. [10.22541/essoar.169091894.48592907/v1](https://doi.org/10.22541/essoar.169091894.48592907/v1) [preprint].
- Taha, G., Loughman, R., Colarco, P. R., Zhu, T., Thomason, L. W., & Jaross, G. (2022). Tracking the 2022 Hunga Tonga-Hunga Ha'apai aerosol cloud in the upper and middle stratosphere using space-based observations. *Geophysical Research Letters*, 49, e2022GL100091. <https://doi.org/10.1029/2022GL100091>
- 365 Wrana, F., von Savigny, C., Zalach, J., & Thomason, L. W. (2021). Retrieval of stratospheric aerosol size distribution parameters using satellite solar occultation measurements at three wavelengths. *Atmospheric Measurement Techniques*, 14(3), 2345–2357. <https://doi.org/10.5194/amt-14-2345-2021>
- Wrana, F., Niemeier, U., Thomason, L. W., Wallis, S., and von Savigny, C. (2023) Stratospheric aerosol size reduction after volcanic eruptions, *Atmos. Chem. Phys.*, 23, 9725–9743, <https://doi.org/10.5194/acp-23-9725-2023>.