

Disentangling the drivers behind the post-2000 retreat of Sermeq Kujalleq, Greenland (Jakobshavn Isbrae)

Ziad Rashed¹, Alexander A. Robel¹, and Hélène Seroussi²

¹School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

²Thayer School of Engineering, Dartmouth College, Hanover, NH, USA

Correspondence: Ziad Rashed (zrashed3@gatech.edu)

Abstract. Ocean temperatures have warmed in the fjords surrounding the Greenland Ice Sheet, causing increased melt along their ice fronts, rapid glacier retreat, and contributing to rising global sea levels. However, there are many physical mechanisms that ~~may-can~~ mediate the glacier response to ocean warming and variability. Warm ocean waters can directly cause melt at horizontal and vertical ice interfaces, or promote iceberg calving by weakening proglacial mélange or undercutting the glacier front. Sermeq Kujalleq (also known as Jakobshavn Isbræ) is the largest and fastest glacier in Greenland and has undergone substantial retreat starting in the late 1990s. In this study, we use a large ensemble modeling approach to disentangle the dominant mechanisms ~~driving-that drive~~ the retreat of Sermeq Kujalleq. Within this ensemble, we vary the sensitivity of three different glaciological parameters to ocean ~~warmingtemperature~~: frontal melt, subshelf melt, and a calving stress threshold. Comparing results to the observed retreat behavior from 1985-2018, we select a best-fitting simulation which reproduces the observed retreat well. In this simulation, the arrival of warm water at the front of Sermeq Kujalleq in the late 1990s ~~leads-led~~ to enhanced rates of subshelf melt, ~~leading-to-triggering~~ the disintegration of the floating ice tongue over a decade. ~~Retreat-The recession of the calving front~~ into a substantially deeper bed trough around 2010 accelerates ~~the~~ retreat, which continues nearly unabated despite local ocean cooling in 2016. An extended ensemble of simulations with varying calving threshold shows evidence of hysteresis in ~~the~~ calving rate, which can only be inhibited by a substantial increase in ~~the~~ calving stress threshold beyond ~~the~~ values suggested for the historical period. Our findings indicate that accurate simulation of rapid calving-driven glacier retreats requires more sophisticated models of iceberg mélange and calving evolution coupled to ice flow models.

1 Introduction

Observations indicate that many glaciers in Greenland have undergone rapid retreat over the last few decades. Sermeq Kujalleq (hereafter SK, also known as Jakobshavn Isbræ) is a fast-flowing marine-terminating glacier in West Greenland, which has been the fastest and largest contributor ~~of-to~~ Greenland ice discharge for the past several decades (Mouginot et al., 2019). SK has experienced considerable thinning and retreat with terminus velocities almost doubling from ~6,700 m/yr in 1985 to ~12,600 m/yr in 2003 (Holland et al., 2008; Joughin et al., 2014). As a result of this mass loss, SK was singularly responsible for 4% of the total ~~rise of sea level in the 21st century-sea-level-rise century~~. Prior to this rapid acceleration, SK was considerably slower, experiencing moderate thickening ~~from 1991-1997 (Schweinsberg et al., 2017; Joughin et al., 2004). Prior to-between~~

25 1991 and 1997 (Schweinsberg et al., 2017; Joughin et al., 2004). Before 2000, SK had a floating ice tongue providing that
provided some stability by buttressing of the terminus (Echelmeyer and Harrison, 1990) and buffering of frontal melt rates :
The intrusion by acting as a heat sink for warmer fjord waters. The arrival of warmer subsurface waters into in Disko Bay and
the Illulisat Icefjord in the late 1990s preceded the collapse of the floating ice tongue (Holland et al., 2008), which is widely
thought widely believed to have initiated the retreat and acceleration of SK over the next 20 years. Gladish et al. (2015) suggest
30 that warmer waters may not have reached SK's front due to the downstream sill, however, we keep consistent in our assumption
of intrusion to contextualize our findings in the context of similar efforts.

While it is generally agreed that warm subsurface waters triggered the most recent retreat phase of SK (Holland et al., 2008;
Myers and Ribergaard, 2013), there is still debate about which physical processes are responsible for mediating the glacier
response to ocean warming. Here we are primarily concerned in answering whether the question: was SK's recent acceleration
35 and retreat caused by interannual variability in calving activity or by amplified melting of the ice front?. Though there is some
uncertainty over the extent to which warm waters penetrated the fjord beyond a submersive sill (Gladish et al., 2015) here
we test the extent to which observed retreat could have been triggered by warm ocean waters and amplified by glaciological
processes. The greatest increases in SK's terminus flow speed occurred in the summers of 2012 to 2015 (Khazendar et al.,
2019) but initial warming (1-2°C) of Disko Bay fjord waters occurred a decade prior earlier, indicating a lag in SK's delay
40 in the SK flow response to warming ocean conditions. Conversely In contrast, SK flow speeds decreased concurrently with
the cooling of Disko Bay water by 1.5 °C in 2017. Based on this relationship between glacier speed and fjord tempera-
tures, it has been argued that enhanced melting of the terminus caused greater calving, retreat, and speed-up, particularly in
summer when buoyant subglacial meltwater plumes should enhance circulation at the terminus (Khazendar et al., 2019). On
the other hand, However, observational records indicate that previous intervals of enhanced fjord heat content prior to the
45 1980s did not result in the same dramatic retreat (Slater et al., 2018), leading to the natural question: why did the most re-
cent period of warming beginning in 1985 cause such a dramatic and unprecedented retreat? While water temperatures in
Disko Bay are associated with melt and retreat at SK, it is still not clear whether this association indicates a causal rela-
tionship between enhanced terminus melt and observed thinning and retreat (Joughin et al., 2020), and whether the recent
retreat is the direct result of oceanographic or glaciological factors. Alternatively, the strength of iceberg mélange and under-
50 cutting via frontal melt have been observed to have strong control on calving frequency and style at SK and other glaciers
(Joughin et al., 2004; Amundson et al., 2010; Cassotto et al., 2015; Luckman et al., 2015; Kajanto et al., 2023). Disentangling
the (Joughin et al., 2004; Amundson et al., 2010; Cassotto et al., 2015; Luckman et al., 2015; Xie et al., 2019; Kajanto et al., 2023)
. Although this is not a universal feature among marine terminating glaciers in West Greenland (Amaral et al., 2020), the
cliff-like geometry of SK's front may make it more susceptible to retreat in the presence of a weakening mélange. Thus,
55 disentangling the drivers behind SK's response to warming ocean conditions requires distinguishing between retreat driven
through direct melting and calving ocean-induced melting due to increased local temperatures and calving due to a weakened
pro-glacial mélange.

Understanding the response of SK to warming ocean waters poses a difficult challenge ,owing due to the complex range of
processes occurring at its interface with the ocean. Many processes that play a critical role in glacier stability (i.e., subshelf

60 melting, calving, melt undercutting, mélange buttressing) remain poorly understood, despite recent advancements in high-fidelity glacier models (Benn et al., 2017; Slater et al., 2021). Furthermore, the process of acquiring the necessary observations to ~~limit the potential values of modeling constrain model~~ parameters becomes challenging due to the presence of icebergs in winter seasons. ~~Simplified~~ Observational records of environmental variables such as mélange density, detailed calving event catalogs, and calving front geometry are difficult to collect because icebergs act as physical barriers to oceanographic vessels, especially near the glacier front, where they are most dense. To avoid this, simplified parameterizations that relate calving rates to glacier stress and geometric conditions are used in many ice sheet models (Benn et al., 2007), but do not always capture the complex interactions between glacier and ocean state. Additionally, parameterizations ~~which that~~ may perform well in describing one glacier might not perform as well at other glaciers (Amaral et al., 2020). Here, we use ~~SK's the~~ rapid retreat and complex evolution ~~over the past of SK in recent~~ decades as a natural experiment to better understand ~~the~~ uncertainties and
70 shortcomings in simple parameterizations of ice-ocean interactions.

In this study, we simulate the historical evolution of SK from 1985 to 2018 using the Ice-Sheet and Sea-Level System Model (Larour et al., 2012). We perform a large ensemble of simulations of SK retreat through perturbation of three sensitivity parameters that ~~impact control three processes which directly influence~~ its retreat: subshelf melt, melt at the calving front, and calving threshold modulated by mélange rigidity. We compare model simulations to the relevant period of the observational
75 record by scoring their ability to reproduce observed calving front geometry. We investigate the trade-off between different processes in driving SK's temporal and spatial sensitivity to melt- and stress-based mechanisms of mass loss and highlight the possible mechanisms most likely responsible for observed retreat. We emphasize that our goal here is not to accurately simulate mélange, but to offer a scenario in which disentangling the causes of SK's retreat can be accomplished with the consideration of long-term changes in mélange buttressing of the terminus through modification of the calving threshold parameter.

80 In section 2, we lay out the methodology for simulating SK's evolution from 1985-2018. We describe how simulations are initialized in the Ice-Sheet and Sea-Level System Model and the specific data used to recreate the state of SK in 1985. We then explain how melt and calving processes are parameterized with respect to ocean forcing and how we design our ensemble to determine which parameter combination results in model ~~output states~~ that most closely matches observational data. In section 3, we present our model ensemble results and highlight key relationships between model parameters that best fit observations.
85 ~~Additionally~~In addition, we analyze the timing and extent of retreat within the model with the best match to observations. In section 4, we discuss implications for SK's future evolution given its current state. We also ~~discuss how our results compare to previous studies and hypotheses regarding contextualize our findings in the context of Bondzio et al. (2018)'s study and use it as a control to compare against in assessing relative contributions of melt and calving to SK's rapid retreat evolution.~~

2 Methods

90 2.1 Model Configuration

We use the Ice-Sheet and Sea-Level System Model (ISSM) to simulate the retreat of SK from 1985 to 2018. ISSM is a state-of-the-art thermomechanical ice sheet model that has been used to simulate ~~ice sheet evolution of glacier and at the~~

evolution of glaciers and ice sheets from catchment to continental scales (Larour et al., 2012). Our modeling approach draws on some aspects of the configuration of a previous SK modeling study by Bondzio et al. (2018), with some key enhancements detailed later in this section. The domain of our simulations includes the fast-flowing parts of the SK catchment and extends upstream deep into the SK catchment area. ~~We use a mesh with a horizontal resolution which ranges~~ Using ISSM's built-in bidimensional anisotropy mesh generator, the domain mesh is refined using a metric based on the product of bedrock slope and surface velocity. The resulting mesh elements range in length from 400 m ~~near the grounding line to 7 km at the most upstream parts of the catchment~~ at high-velocity locations closer to the calving front to 4 km at lower-velocity locations deeper within the catchment area. A 2D shelfy-stream approximation (SSA; Morland and Zainuddin, 1987; MacAyeal, 1989) is used to simplify the three-dimensional flow equations as vertical gradients in velocity are relatively small and basal sliding is the dominant contributor to ice velocity at SK. The SSA approximation greatly reduces the computational expense of simulating marine-terminating glacier evolution and thus enables the large ~~ensembles~~ ensemble of simulations in this study. A linear-viscous Budd sliding relation (Budd et al., 1984) is used ~~and appropriate sliding coefficients were inverted by using composite velocities collected in 1991 and~~ to relate basal shear stresses to basal speed. To obtain effective pressures, we first use a 1985 DEM (Korsgaard et al., 2016) and a 2009 ~~for the area close to the ice front (Rignot and Mouginot, 2012), which have been sealed down to give a smooth transition between both data sets. A DEM (Morlighem et al., 2017).~~ To fill in the gaps in the 1985 surface, we use the 2009 DEM and apply a height offset proportional to the rate of surface height change in 2009. We follow a similar process in initializing the velocity field but instead use a velocity offset proportional to the velocity ratio between 2009 (Joughin, 2015) and 1991 (Mouginot et al., 2019) to fill in the 1985 velocity gaps. We then deduce initial Budd sliding coefficients from the updated 1985 surface heights and velocities. Grounding line migration is modeled using a sub-element migration scheme which allows the simulated grounding line to evolve continuously through mesh elements and ~~limit~~ reduces the dependence on mesh resolution (Seroussi et al., 2014). ~~At the upstream boundary of the domain, ice velocities are set to observed ice velocities and the corresponding ice thicknesses are kept constant. The~~ We use a model time step is of approximately 5.5 hours in order to capture rapid changes in ice ~~terminus front~~ geometry while maintaining numerical stability and accuracy. ~~e-SMB over-~~

Along the domain boundary, ice velocities are set to observed 1985 ice velocities, and the corresponding ice thicknesses are kept constant. Although SMB in our region of interest changed ~~over during~~ the time period considered, the ~~magnitude of its variability relative to the time average is much smaller than that of oceanic forcings (Hanna et al., 2011).~~ Additionally ~~relative variations in SMB are small relative to the relative variations in ocean temperature (Hanna et al., 2011).~~ Furthermore, it has been shown that the SK region is weakly sensitive to ~~atmospheric forcings~~ SMB forcing within our time period of interest (Seroussi et al., 2013) and ~~we can consider atmospheric forcings to be more influential in its role modulating ocean conditions that oceanic energy fluxes to the glacier increased while atmospheric energy fluxes to the glacier remained relatively constant during our study period (Wang et al., 2020).~~ Following this, we use a spatially variable surface mass balance held constant in time, based on a multidecadal mean from the RACMO regional climate model (Ettema et al., 2009). ~~We initialize simulations with 1985 ice surface topography obtained via photogrammetry (Korsgaard et al., 2016) with gaps filled using the~~

Greenland Ice Mapping Project (Howat et al., 2014) and corrected with an offset proportional to the ice flow velocity. Bedrock topography used in these simulations is from BedMachine Greenland v3 (Morlighem et al., 2017).

130 Migration of the calving front is simulated using a level set formulation (Bondzio et al., 2016), where the migration rate of the calving front (w) is determined using the difference between ice velocity and frontal ablation rate,

$$w = v - (c + m_{fr}) n \quad (1)$$

135 where v is the ice velocity at the calving front, and $c + m_{fr}$ is the ablation rate. Ablation at the calving front is driven by two parameterized processes: iceberg calving (c is calving rate) and direct melt of the calving front by heat flux from the ocean (m_{fr} is frontal melt rate). In the level set approach, ice flow advects the calving front downstream and ablation mechanisms move the calving front upstream.

2.2 Stress-based iceberg calving threshold

Iceberg-In ISSM, iceberg calving rate is calculated with using a tensile-stress based criterion (inspired by criteria based on the von-Mises principal tensile stress)

$$c = |v| \frac{\sigma}{\sigma_{thr}}, \quad (2)$$

140 where σ is the principal tensile stress, σ_{thr} is a prescribed stress threshold, and v is the ice velocity magnitude at the ice front (Morlighem et al., 2016).

Calving-induced retreat of the glacier front is initiated once local principal tensile stresses exceed the stress threshold parameter (σ_{thr}). The stress threshold parameter can be thought to conceptually represent many physical processes that material characteristics such as fracture toughness, grain-scale deformation, and ice strength, which have the ability to modify the propensity for calving events. We parameterize a linear decrease in calving threshold, σ_{thr} , with increasing fjord ocean temperatures (T)

$$\sigma_{thr}(T) = \sigma_{max} - \frac{T - \min(T)}{\max(T) - \min(T)} \frac{T - \min(T)}{\max(T) - \min(T)} (\sigma_{max} - \sigma_{min}). \quad (3)$$

150 Bondzio et al. (2018) use a similar framework to effect seasonality in calving in which their calving stress threshold alternates between a low stress threshold in the summer seasons and a high stress threshold in winter seasons, effectively shutting off the calving in the winter seasons and allowing the calving in the summer seasons. The minimum stress threshold, σ_{min} , corresponds to the stress threshold in the presence of the warmest temperature in the fjord temperature time series, T , and conversely the maximum stress threshold, σ_{max} , corresponds to the stress threshold in the presence of the coldest temperature in the fjord temperature time series. The linear variation in calving stress threshold is a simplified realization of the effects of mélange weakening/strengthening on buttressing the calving front, which has been posited as a possible explanation for observations indicating mélange weakening and breakup in concert with ocean temperature seasonality, rather than atmospheric

temperatures (Kehrl et al., 2017; Bevan et al., 2019; Joughin et al., 2020). We use a linear sensitivity to relate ice mélange strength to ocean temperature for a simple comparison with other linear sensitivities assumed in this study. Due to the fractured nature of near-terminus ice at SK, we set ~~this the~~ minimum calving stress threshold, σ_{\min} , to 100 kPa, below the measured mechanical strength properties of laboratory samples of pristine ice, typically in the range of 0.1-1 MPa (Lee and Schulson, 1988; Petrovic, 2003). The ~~temperature dependence of the stress threshold ensures that calving activity increases when the ocean in contact with the glacier is warmer. This stress threshold is intended to parameterize the maximum stress threshold,~~ σ_{\max} , parameterizes the potential roles of rigid iceberg mélange and melt undercutting in modulating the relationship between fjord ocean temperatures and calving rate. ~~Though and we use it to set the sensitivity of calving activity to local ocean temperature. Within our ensemble, we use a σ_{\max} range of 220-350 kPa to capture a large swathe of potential calving behavior.~~ The temperature dependence of the stress threshold ensures that calving activity increases when the ocean in contact with the glacier is warmer and vice versa. Although other material properties and glacier processes may play a role in setting the propensity for calving (e.g., ~~fracture toughness of ice~~ ice fracture toughness, surface melt), they ~~are not able to~~ cannot explain the timing of seasonal and ~~multi-annual changes in calving style at~~ multiannual changes in the calving style in SK (Joughin et al., 2008). In ~~section~~ Section 4, we further discuss the shortcomings of such a simplified representation of the effect of mélange on calving.

2.3 Frontal and submarine melt

Ocean melt of the glacier calving front (referred to hereafter as frontal melt) also contributes to the glacier response to fjord temperatures. We compute the frontal melt rates using the empirical parameterization from ~~Rignot et al. (2016); Xu et al. (2013);~~ Rignot et al. (2016):

$$m_{\text{fr}} = (A h Q_{\text{sg}}^{\alpha} + B) T^{\beta}, \quad (4)$$

where Q_{sg} is the subglacial discharge ~~which that~~ is taken as the ~~decadal average of~~ total runoff of ~~SK's drainage basin as given by the SK drainage basin according to~~ RACMO 2.3 (Noël et al., 2015), T is the ~~ocean thermal forcing obtained via the depth-integrated temperature of the water column near the glacier front,~~ thermal forcing of the ocean obtained using the depth-averaged fjord water temperatures from the water temperature in the Egedesminde Dyb in the coupled ocean and sea ice simulation provided by the Estimating the Circulation and Climate of the Ocean, Phase II (Menemenlis et al., 2008), and h is the depth of the water column. A is a tuning parameter and B is a parameter that ensures that the heat flux does not vanish in the absence of melt water (Rignot et al., 2016). This empirical equation has been shown to be a good approximation of how frontal melt relates to ocean temperature and subglacial discharge at ~~some several~~ glaciers in Greenland, based on in-situ observations and numerical simulations (Rignot et al., 2016). Since Q_{sg} and α are sufficiently small, and β is sufficiently close to one, we assume that frontal melt is linearly proportional to thermal forcing to simplify the following analysis. We do not account for the effects of convective plume forcing on frontal melting because the sensitivity of frontal melting to subglacial discharge is lower than to ocean temperature (Xu et al., 2013), especially after the removal of the ice tongue.

Submarine melt on the floating portion of the glacier is the final process ~~we consider contributing to that we consider to contribute to~~ the glacier response to fjord temperatures. ~~We simulate the submarine melting process by assuming that the water column is stratified such that water at maximum depth is also the warmest water. We use this assumption to justify the warmest waters overcoming the shallow sill and infiltrating SK's fjord. Following this, we take the depth-average of the water column's temperature between the surface and maximum of bedrock and sill height (-250 m). We fill the data gap between 1986-1992 by repeating the temperature time series between 1992-1996.~~ The submarine melting rate is ~~then~~ computed using the simplified ~~two-equation~~ parameterization from Holland and Jenkins (1999):

$$m_{\text{sm}} = -\rho_M c_{pM} \gamma_T (T - T_{\text{pmp}}). \quad (5)$$

where ρ_M is the ~~mixed-layer density~~density of the mixed layer, c_{pM} is the specific heat capacity of the mixed layer, γ_T is the thermal exchange velocity ~~and~~ $(T - T_{\text{pmp}})$ is the difference between the temperature of the mixed layer and the ~~pressure~~ melting point temperature at the base of the ice.

2.4 ~~Model-Observation Mismatch Score~~

<u>Parameter</u>	<u>Value</u>	<u>Units</u>
<u>A</u>	<u>3×10^{-4}</u>	<u>Unitless</u>
<u>B</u>	<u>0.15</u>	<u>Unitless</u>
<u>α</u>	<u>0.39</u>	<u>Unitless</u>
<u>β</u>	<u>1.18</u>	<u>Unitless</u>
<u>ρ_M</u>	<u>1023</u>	<u>Kg m^{-3}</u>
<u>c_{pM}</u>	<u>3974</u>	<u>$\text{J K}^{-1} \text{Kg}^{-1}$</u>
<u>γ_T</u>	<u>1×10^{-4}</u>	<u>$\text{m}^2 \text{s}^{-1}$</u>
<u>T_{pmp}</u>	<u>-1.85</u>	<u>$^{\circ}\text{C}$</u>

Table 1. Overview of all relevant melt parameters.

We ~~then~~ choose a parameter space ~~for multipliers of forcing of multipliers for melt~~ mechanisms such that we encompass a large range of possible ~~forcing-melting~~ scenarios. For our frontal and submarine melt ~~parameters space, we choose multipliers, α_{mf} and α_{ms} respectively, we chose~~ a range of 0-4x the empirical parameterizations mentioned in equations (5) and (4). ~~Following a series of experiments within different ranges, this range was selected to capture as much variability in simulation output while producing simulated retreats at least somewhat resembling the observed evolution of SK.~~

2.4 Model-Observation Mismatch Score

We choose to plot results for a max stress threshold varying over ~~220-360kPa~~ 220-350kPa with an interval of 10 kPa to ensure ~~that~~ we capture both potentially ~~rigorous-vigorous~~ and negligible calving activity. We also conducted additional simulations




Figure 1. Observed and simulated calving fronts from our highest scoring simulation at SK from 1985-2018. The solid lines denote outputs from our best fitting ensemble member and the dashed lines indicate the observed positions of SK's calving front with the color corresponding to the year of observation.

outside the aforementioned parameter bounds, particularly with a much wider range of calving stress threshold. ~~But~~ However, simulations with prescribed stress thresholds outside the bounds plotted in the next section produced retreats (or lack of retreat) very different from those observed in reality. Thus, we opted not to plot those here. Each simulation is scored ~~based on its correspondence to~~ on the basis of its ability to match the observed position and geometry of the glacier calving front. We use a historical catalog of SK's calving front geometry (i.e., a 2-D curve) obtained from observational records which includes multiple snapshots of calving front geometry derived from Landsat 5–8, ERS-1 and 2, and TerraSAR-X satellite imagery (Figure 1). ~~We pair each~~ For each point in time in which we have an observation, we pair each simulation's observed terminus geometry with the nearest-in-time (always within ~~less than~~ 5 days) modeled terminus geometry from ISSM. For each observation-model pair, we calculate the area between the modeled and observed terminus geometry. ~~Though we~~ The resulting mismatch vector contains the difference in geometric area between modeled and observed geometries, and we take the root mean square of this vector to assign a score. This is different from the approach of Bondzio et al. (2018) which only considers the center line position of the SK calving front; considering the entire calving front geometry allows for more accurate tracking of the glacier when the front bifurcates, as it did in 2006. Additionally, while we do not weight ~~scores-area~~ differences to account for changing observational density in time, almost all of the retreat of SK occurs during a time period (post-2000) when observational density is high. Thus, the mismatch score is unlikely to be strongly dependent on observational density. Using this scoring method, we can accurately capture changes in calving front position and shape ~~and~~, and we are only limited

by the resolution of the observational records and model meshing. Convergence studies indicate that at our chosen model
 225 time step, potential errors due to the mismatch in timing between model and observation contribute negligibly to the overall
 scored metric. ~~For every ensemble member, we take the root mean square of the error vector as the overall score for the
 simulation to measure how well the model did in matching calving front geometry. This is different from the approach of
 Bondzio et al. (2018) which only considers the center line position of the SK calving front, as considering the entire calving
 front geometry allows for more accurate tracking of the glacier when the front bifurcates, as it did in 2006.~~
 230 ~~Observed and simulated calving fronts from our highest scoring simulation at SK from 1985-2018. Thicker lines denote
 outputs from our best fitting ensemble member. The dashed lines indicate the observed positions of SK's calving front with the
 color corresponding to the year of observation.~~

3 Results

The speed and timing of the simulated retreat of SK vary depending on the sensitivities of the calving threshold, submarine
 235 melt, and (to a lesser extent, discussed below in the following section) frontal melt to the local ocean temperature, but there
 are some commonalities between simulations in our perturbed-parameter ensemble. Figure 1 shows the observed calving front
 positions (~~thin~~ thick lines) and the results of one simulation with the best correspondence to observations (~~thick~~ dashed lines).
 Generally, the ~~calving front~~ simulated calving fronts of SK pauses at locations where there are geometric pinning points such
 as bedrock peaks and fjord narrowings. However, the length of time that the simulated glacier remains at such pinning points
 240 varies, due to the interplay of calving and melting. Initially, the ice tongue of SK stabilized the glacier front by buttressing inland
 ice. As in observations (Motyka et al., 2011), simulations ~~generally~~ indicate that intensified submarine melting due to warming
 of fjord waters starting around 1997 thinned the ice tongue and weakened its buttressing capacity. The observed weakening and
 subsequent disintegration of SK's ice tongue ~~result~~ resulted in dynamic thinning of the terminus and an acceleration in retreat.
~~Most~~ Similarly, most of the simulations in our perturbed parameter ensemble ~~yield~~ yielded submarine melting rates which ~~peak~~
 245 peaked between 1995 and 2000, as warm water entered Illulisat Icefjord via Disko Bay. The simulated disintegration of the ice
 tongue and subsequent front retreat leads to the bifurcation of the calving front into two branches and exposure of a much larger
 frontal area to warm ocean waters (Figure 1). The response of SK's two branches is not homogeneous owing to large differences
 in their bed topography and fjord geometry. Ice fluxes are greatest along the southern calving front where the bedrock is deeper
 and upstream topography is characterized by more extensive retrograde slopes (Figure 1). The combination of a thicker and
 250 steeper glacial terminus and deepening grounding line bed slopes facilitate ice loss via calving. Thicker ice results in a greater
 overburden pressure which, at the grounding line, is counterbalanced by hydrostatic pressure and buttressing stresses from
 floating ice and mélange. Thus, ~~following the removal of the~~ after removing the ice tongue, tensile stresses grow rapidly along
 the southern front (Figure S1) and calving quickly becomes the dominant ice loss mechanism (Figure 3). The northern branch
 of SK experiences much less retreat, similarly to what observations show (Figure 1). The shallow bed topography and an
 255 abundance of pinning points constrain the upper branch from rapidly retreating following ~~the~~ ice tongue disintegration. ~~Tensile
 stresses along the upper branch's calving front grow but do not result in rapid calving as observed in the lower or main branch.~~




Figure 2. Parameter combination sweep of relevant melt values. Darker colors indicate a smaller RMS ~~error of cumulative area mismatch~~ between modeled and observational records of SK’s calving front position. Panel a displays the RMS of cumulative area mismatch for specified maximum stress levels. Panels b-d show respective RMS slices for $\sigma_{\max} = 300\text{kPa}$, $\alpha_{mf} = 0$, $\alpha_{ms} = 2$, and $\alpha_{mf} = 1$. The red diamond in panel b indicates where our best-fit parameter combination lies in parameter space.

3.1 Perturbed-parameter ensemble of SK retreat simulations

In our large perturbed-parameter ensemble, multiple simulations were able to achieve nearly equivalent matches to the observed retreat of SK (Figure 2). There appears to be a minimum achievable match to observations (with RMSE of approximately ~~40~~ 50 km²) related to observed changes in calving style unrelated to ocean temperatures, which we discuss in more detail in section 4. We also expanded our ensemble extent within parameter space beyond the ranges plotted in Figure 2 to verify that the best fitting simulation was indeed the best fit. We do not plot simulations from this greater ensemble due to the lack of glacier retreat under a sufficiently large stress threshold regime. The best scoring simulations all occur within a ~~similar~~ region of parameter space ~~where submarine melt rates are relatively low and maximum stress thresholds relatively high, as~~ plotted in Figure 2. ~~Outside of this region of parameter space, model-observation mismatch scores are consistently much worse.~~ The cold-ocean maximum of calving stress threshold in this region of parameter space is 250-400 kPa, generally much lower than suggested by laboratory studies, and near the low end of the range of observationally derived values for fractured glacier fronts (Vaughan, 1993; Choi et al., 2018). Outside this region of parameter space, model-observation mismatch scores are consistently much worse, indicating an implicit role for rigid mélange in buttressing the calving front and preventing calving.

The simulations best matching observations generally require submarine melt to be more sensitively dependent on ocean temperatures (~~$M_s > 1$~~ $\alpha_{ms} > 1$) than suggested by the ~~two-equation-parameterization~~ parameterization (Rignot et al., 2016). However, there is a notable trade-off between the stress threshold and submarine melt, such that simulations with higher

cold-ocean stress thresholds (i.e., less calving in ~~cold-waters~~coldwaters) also need higher submarine melt rates to ~~compensate-~~
~~Diminished~~achieve reasonably low RMS (Figure 2b). Early in the simulation (1985-2000), the ~~diminished~~diminished calving of the ice
 275 tongue ~~needs-to-be-compensated-must be compensated for~~ by amplified submarine melt ~~in-order~~ to accurately simulate the
 timing of the ice tongue collapse. ~~The~~Consequently, the greatest mismatch between models occurs following the collapse of
 the ice tongue, coinciding with the onset of calving-dominated retreat (Figure 3b).

Though the absolute best-fitting simulation requires high sensitivities of front melting to ocean temperatures (~~$M_f = 4$~~ $\alpha_{mf} = 4$),
 there are several simulations with very similar RMSE requiring little to no front melt at all to fit observations (Figure S2). In
 280 these simulations, the greatest mismatch occurs following ice tongue collapse which is when calving becomes the dominant
 mode of ice loss. It is also by this point that thermal forcing from warmer waters contributes less to SK's dynamic response.
 If the stress threshold is sufficiently low, retreat following ice tongue collapse is controlled by bed topography and ~~ice-flux-is~~
~~dominated-by~~ calving in the southern trunk. Conversely, retreat is controlled by submarine melt when stress thresholds are con-
 siderably ~~larger-greater~~ than tensile stresses generated at the glacier front. Changing frontal melt rates predicted by the Rignot
 285 et al. (2016) parameterization by modifying ~~M_f did α_{mf} does~~ not significantly change ~~model behavior, nor improve our best~~
~~fit parameter combination~~the behavior of the model or improve our combination of best fit parameters. The weak dependence
 of RMSE in Figure 2 to the sensitivity of frontal melt to ocean temperature indicate that, at least for SK, calving and submarine
 melt control the speed and timing of glacier retreat.

3.2 Best-match simulation

290 ~~The simulation best-fitting observations captures the observed SK calving front geometry well, though there do remain some~~
~~periods of mismatch. In Figure 3, the fluxes contributing to ice loss at the calving front are decomposed (panel a), along with~~
~~the fit to observations (panel b), and the ocean forcing from the ECCO2 reanalysis product (panel c).~~

~~During the period from~~As observed from 1985 to 2000, SK maintained a floating ~~ice-tongue-ahead-of-its-terminus. While~~
~~the-tongue that at some locations extended more than 10 km from its grounding line. As long as a~~ floating ice tongue existed,
 295 observed surface ice flow velocities exhibited very little seasonal variability (Echelmeyer and Harrison, 1990), indicating a
 glaciological state in which ~~dynamic thinning due to ocean melt is~~ calving and melting are consistently balanced by surface
 accumulation and ice flow ~~from-upstream~~. SK's ice tongue acted as a buffer for retreat ~~through two primary mechanisms: (1)~~
~~by~~ transmitting buttressing back stress from the Illulisat Icefjord walls to the grounding line, ~~and (2) acting as a heat sink~~
~~for warm ocean waters. Local ocean.~~ Following the influx of warmer ocean water into Disko Bay in 1997, local fjord water
 300 temperatures experienced an abrupt increase ~~in-1997~~, which led to the disintegration of SK's ice tongue over several years and
 the subsequent thinning and accelerated retreat of the newly exposed terminus (Joughin et al., 2020).

In Similarly, in our best-fitting simulation, the ~~rise-increase~~ in ocean temperature beginning in 1997 (Figure 3c) causes an
 increase in submarine melt fluxes, which is then followed by the simulated ice tongue thinning and retreat. Submarine melt
 initially dominates ice loss, peaking as ocean temperatures reach their maximum in 2000. Following this peak, submarine
 305 melt fluxes slowly decrease, albeit at a rate faster than ocean temperatures decrease ~~due-to-because of~~ the decreasing area of
 the floating ice tongue. During this period, the base of the ice tongue steepens, which causes a subsequent ~~rise-increase~~ in




Figure 3. Best fitting model flux contribution and performance. (**Top-Panel,** a) Mass flux contribution of relevant ablation mechanisms to the evolution of SK's calving front. (**Middle-Panel,** b) ~~Difference between modeled and observed calving front areas~~ Observed area change time series vs modelled area time series. (**Bottom-Panel,** c) Estimated Disko Bay fjord temperatures sourced from ECCO2 (Menemenlis et al., 2008).

driving stress and extensional stresses, which promotes calving activity. ~~However, we do not see a corresponding~~ Although we expected to see a similar increase in calving fluxes ~~immediately following the temperature increase due to the decreased stress threshold under warm water conditions, we did not see an increase until 2010 (Figure 3a).~~ This is most likely due to the response timescale associated with the geometric adjustment of the floating ice tongue and corresponding stress state of the terminus, which depends more on the instantaneous glaciological state rather than the oceanic state. It should also be noted that during this period of retreat, SK's front is still far enough downstream that it has not begun ~~retreating via its two branches. This is important because this part of Ilulissat Icefjord has a slight~~ to retreat into its two separate branches (Figure 1). Throughout this period, SK's grounding line sits on a slightly prograde bed slope and also has not yet begun our rapid retreat. Thus, this period is marked by a relatively slow retreat of the floating calving front, driven by submarine melt, which sets ~~up the glacier~~ the glacier up for further retreat as persistently warm water continues to ~~foree reach~~ force reach the glacier front.

During the period from 2000-2012, in our best-fitting simulation, SK's retreat begins to accelerate. In this phase of the retreat, the change in SK's geometry begins to play an important role in setting the calving rate at the terminus. ~~In this period, we also mark a point at which the dominant mechanism of retreat transitions from melt-dominated to calving-dominated, with~~

320 ~~melt accounting for a greater ice flux at the beginning of the period and calving at the end of the period. This transition is~~
~~caused by the recession of the grounding line into a deeper bed and a final removal of the last part of the floating ice tongue~~as
the front retreats onto a much narrower and deeper bed (Figure 1). From 2000-2005, there is a decrease in submarine melt flux
caused mostly by the reduction in floating ice tongue area, and to a lesser extent ~~by~~ the relative decrease in ocean temperature
forcing. By 2005, SK's calving front ~~has had~~ retreated enough to lie across the upper and lower branches, which introduced
325 a greater variability in bed topography along the front (Figure 1). The greatest sustained discrepancy between modeled and
observed glacier geometries occurs over the 2005-2010 period, as our modeled terminus lags downstream of the observed
terminus (Figure 3). The mismatch in geometry is largely amplified by the differences in calving rate as there is a trade-off
between capturing accurate calving rates after 2010 and insuring minimal mismatch between 2005 and 2010 as the dominant
mode of ice loss changes. ~~During Throughout~~ this period, we observe the transition to a calving-dominated retreat owing to
330 geometric changes at the calving front. The combination of a retrograde bed and a steep ice cliff front geometry promotes
intensified calving activity along the lower branch of SK. By 2010, SK's ice tongue has been completely removed and the
calving front is a steep ice cliff. The time at which model ensemble members reach this transition point would be the greatest
source of variability in overall error between modeled and observed calving fronts.

~~By After~~ 2010, the best-fit model simulates a rapid acceleration in calving rate and a better fit between modeled and observed
335 calving fronts. Between 2010 and 2015, the dominant retreat mechanism transitions from submarine melt dominated to calving
dominated with melt accounting for a greater ice flux at the beginning of the period and calving at the end of the period (Figure
3a). By 2011, calving fluxes overtake submarine melt fluxes as the dominant source of mass loss. This transition is caused by the
recession of the grounding line into a deeper retrograde sloping bed. From 2010-2016, calving fluxes increase and SK's front
retreats considerably more along its lower branch than its upper branch. The transition of SK to a calving-dominated retreat
after 2010 (Figure 3), and the lack of substantial re-advance during the brief cooling between 2016-2018 ~~raise suggests~~ the
340 possibility that ~~the our simulated~~ SK has undergone a hysteretic change to irreversible calving-dominated retreat (~~as~~ discussed
further in the next section). ~~Other studies have also raised this possibility.~~ Due to the strong topographical control on retreat
rate, we expect that once the lower branch's grounding line retreats past the over-deepened trough, a rapid acceleration of retreat
is very likely regardless of climate forcing ~~Kajanto et al. (2020). This~~ (Kajanto et al., 2020). Further retreat is largely driven
345 by calving fluxes as floating ice area considerably decreases and ~~the effects of submarine melting are diminished.~~ submarine
melting becomes less influential on glaciological state. We also note that frontal melt fluxes begin to overtake submarine melt
fluxes during this period due to increased exposure area of the ice front to warm waters, but this increase is still an order of
magnitude smaller than the relative increases in calving fluxes.

3.3 Potential hysteresis effects of ice mélange

350 In our best-fitting simulations, calving rate at SK rapidly accelerates in concert with retreat into a deep trough on the south-
ern flank of the glacier catchment. The glacier bed in this trough further deepens ~~over~~ 20-30 km upstream of the 2018
calving front position, raising the ~~spectre-specter~~ of hysteresis to permanently high calving rates, regardless of future cli-
mate forcing. However, other studies have raised the possibility that increased calving will produce a thicker ice mélange

(Xie et al., 2019; Cassotto et al., 2015) which could inhibit calving. As described in section 2, our simulations do not account for this potential negative feedback, ~~assuming and assume~~ that calving rate is only sensitive to ocean temperatures and glacier front geometry.

To simulate the potential for hysteresis effects associated with calving, we continue the best-matching simulation from the large ensemble described in the previous section in a series of simulations with an increased calving stress threshold until 2100. To do this, we run an ensemble of simulations continuing the best-fit model from 2018-2100 where maximum calving stress threshold is set to a multiple (1-2.5x) of the calving stress threshold at 2018 and held constant until 2100. Similarly, we keep the temperature forcing constant at 2018 values until 2100 such that frontal and submarine melting rates remain constant. Figure 4 shows that the resulting simulations exhibit two types of behavior. When the calving stress threshold is ~~held-maintained~~ at 260 kPa, the stress threshold value given by the ~~best-fitting-best-fitting~~ simulation in 2018, calving rate continues to increase, peaking around 2050 at rates more than twice 2018 rates. This behavior is shared with a grouping of ensemble members (260-437 kPa) or (1-1.67x) and consistently results in an almost complete retreat into the ice sheet interior by 2100. Ensemble members with a stress threshold above 437 kPa (~1.67x) stabilize, but do not readvance. In these simulations, the calving front stabilized in the deepest trough along the flowline, and the glacier experienced gradual thickening. We note here that such calving stress thresholds were not attained at any point during the historical time period used to calibrate the stress threshold (i.e., they are above the ~~cold-ocean- σ_{max} -cold ocean σ_{max}~~ found for all best-fitting simulations in the parameter sweep described in the previous section. Thus, even if ocean temperatures returned to the coldest values achieved during the historical period over the next 80 years, the rapid acceleration of calving and retreat would likely continue unabated. Assuming that calving threshold is only determined by ocean temperature and the stress state at the calving front, this runaway retreat highlights that simulations which represent calving in such a simplistic way will tend to produce irreversible retreat, which may not be the case if mélange is realistically represented. Importantly, almost all current ice sheet models that use stress-based calving parameterizations hold such parameters constant in time.

4 Discussion

In this study, we have investigated the drivers behind SK's recent retreat from 1985-2018. By considering only processes related to ocean forcing, we are able to closely reproduce the observed evolution of SK and analyze the mechanisms responsible for its retreat before and after ice tongue collapse. Our best-fitting model simulations match those sparse observations ~~which do~~ exist, that do exist but also allow us to understand the physical mechanisms that drove the observed retreat in a physically self-consistent manner. Thus, we can paint a fuller picture of the dynamics responsible for SK's evolution.

We noted two distinct behaviors in glacier evolution in the simulations beyond 2018 plotted in Figure 4, ~~extending past 2018-~~. The first is ~~arrest-stabilization~~ of the calving front retreat on a prograde slope following rapid retreat. The second is characterized by a rapidly accelerating retreat which extends 10's of km into SK's interior followed by ~~arrest-a slow down~~ of retreat. The ~~location at which arrest or rapid retreat occurs is consistent across these simulations. However, the~~ timing of rapid retreat and arrest varies marginally between ensemble members. By 2021, ensemble members which experience a slow




Figure 4. Top Panel (a): Calving fluxes of 2100-1985-2100 run ensemble members from 1985 to 2100 with the onset of stress amplification at 2018. Bottom Panel (b): Along flowline profiles of ensemble members at year 2030. The initial 2018 profile is denoted with a thick black dashed outline and sea level is denoted with a blue dashed line.

down in retreat stabilize near the bedrock peak approximately 40 km along the flowline (Figure 4b). This further reinforces the notion that SK's current retreat is controlled by bed topography, with the stress threshold (and all the factors which determine it) influencing the amount of time it takes to reach the rapidly calving state. This behavior is indicative of threshold behavior, wherein a small change in a parameter (in this case calving stress threshold) will lead to either a stable and slightly advancing SK or a rapidly retreating and unstable SK.

Additionally, throughout the parameter space considered in this study, there is a bimodality in calving front positions of 2018. As seen in Figure 5, there exists two clusters of final calving front position. On one end, a subset of simulations that do not experience ice tongue loss and subsequently do not experience rapid calving. These simulations are characterized by either lower calving stress thresholds or melting multipliers (Figure S2). Although calving front locations are more sensitive to calving thresholds, insufficiently strong ocean melt is unable to generate a rapid removal of the ice tongue, which is a necessary condition to produce mechanical imbalances at the calving front and subsequent retreat. Within these simulations, the calving front is quickly prevented from retreating and remains at the same position. On the other end, we see simulations that quickly lose their ice tongue and subsequently accelerate beyond observational records. In this set of simulations, the calving stress threshold plays a greater role than melting in setting the rate of retreat as the grounding line recedes onto deeper bedrock and the glacier front experiences greater mechanical stresses. When letting these simulations continue beyond 2018, we note that rapid calving continues along the southern trunk of SK, but stops across the northern trunk, owing to the presence or lack of steepening bedrock gradients, respectively. There are no simulations with calving fronts stabilizing along a position between

the observed calving front in 2018 and a few kms inland from the initial calving front in 1985. The most likely reason for this lies in our simplified parameterization of a dynamic ice mélange. In the absence of mélange, the gross behavior of the SK calving front as modeled here can be characterized either by rapid and vigorous retreat or terminal stability. This leads to a bimodality in model response, while, in reality, ice mélange acts as a stress buffer in response to rapid calving.

In designing simulations that can be used to disentangle the drivers of SK retreat, certain simplifications were necessary, which limit the broad applicability of this study to other glacier settings. In order to assess the main drivers behind SK's retreat, certain assumptions were made in our simulations. We do not simulate higher-order use higher-order approximations for the glacier momentum balance (i.e., Full Stokes). We instead rely, instead relying on the SSA in order to reduce computational expense, enabling a large ensemble of simulations, while maintaining an accurate representation of grounding line dynamics. A direct consequence of simplifying the flow equations is the overestimation of the basal drag coefficient near the grounding line (Morlighem et al., 2010). On the other hand, Bondzio et al. (2017) showed that the basal sliding law a linear-viscous Budd sliding relation (Budd et al., 1984) and coefficients captured SK's behavior fairly-velocity well.

Calibrating multiple model parameters (in this case M_f , M_s and σ_{max} , α_{mf} , α_{ms} and σ_{max}) allows us to produce simulations which fit observed retreat of SK reasonably well. In Bondzio et al. (2018)'s study, a novel large ensemble approach was used to forecast SK's behavior under an behavior under a collection of parameter combinations of M_f , M_s and σ_{max} which were held constant in time α_{mf} , α_{ms} , and σ_{max} , which were kept constant over time with a rectified seasonal cycle in calving threshold. We expanded on this work by using a similar methodology to discern which mechanisms were most responsible for initiating and maintaining SK's retreat. We find that dynamically adjusting calving propensity via mélange's control on σ_{max} yields dynamically changing calving propensity through inter-annual change in ocean temperature. Our modified approach produces modeled calving fronts which closely match observed calving fronts without requiring melt forcing frontal melt to be multiple factors larger than observed. These historical simulations enabled us to better understand the factors which drove SK's retreat. times higher than suggested by empirical parameterizations (Rignot et al., 2016). However, ultimately such an approach is limited in its ability to explain the role of physical processes not included in our modeling system and how they may evolve outside of the historical sample of satellite observations of SK. The extended simulations described in section Section 3.3 indicate that the only way to arrest the future retreat of SK is by increasing in our modeling system is to increase the calving threshold to values never attained during the historical period and not achievable by changes in ocean temperature alone. We do (Figure 4a). We recognize that it is may be possible that SK is engaged in indeed engaged in a runaway retreat that will not be arrested by any mechanism in the next century. But, to even However, to allow for the possibility that SK retreat could slow down or arrest stop in the future, other processes which that play a role in modulating calving must be simulated. Given SK's potential to generate large and dense ice mélange fields, this (Cassotto et al., 2015) and its strong dependence on calving front conditions (Bondzio et al., 2017) suggests that a strong enough ice mélange, potentially generated through vigorous calving, may be able to suppress runaway calving behavior. Simulating such a feedback would require a dynamic model of ice mélange coupled to the ice sheet and the ocean below. No mélange model currently exists which fits these requirements, though prior efforts have produced useful parameterizations (Vaňková and Holland, 2017; Pollard et al., 2018; Amundson and Burton, 2018; Schlemm and Levermann, 2021b) and very computationally expense-expensive tools unfit for coupling to ice sheet




Figure 5. Top panel (a) depicts the average rate of surface height change for all ensemble members along a flowline (denoted in(b)) along the southern SK branch. The average rate of observed height change is obtained from ICESat and ICESat-2 data from 2003-2019 (Smith et al., 2020). Bottom panel (b) displays the 2018 calving front positions for all ensemble members. Additionally the best fit model calving front is highlighted with a dark orange and the observed 2019 winter calving front Smith et al. (2020) with black. The thin yellow line indicates the flowline on which we calculate surface height change.

models (Robel, 2017; Burton et al., 2018). However, our study indicates While our goal in this study was to provide a different perspective on how ocean-temperature-based processes could modulate calving in the context of SK, our findings indicate that a fully capable and coupled model of mélange is a pre-requisite for any attempts to accurately model SK's future evolution.

5 Conclusions

In this study, we conducted numerical simulations of Sermeq Kujalleq using ISSM to disentangle the relative importance of different mechanisms in driving the retreat of SK from 1985-2017. Using a large ensemble of parameter-perturbed simulations, we explored a wide parameter space of calving and ocean melt parameterizations and compared calving front positions to key observations to score a given simulation on how well it could match observations of retreat. We found that submarine melting and calving both played critical roles in the timing and magnitude of SK's retreat. Specifically, we note that intensified submarine melting due to anomalous deep-water a potential intrusion of anomalous deep-water temperatures was necessary to instigate rapid retreat by melting SK's ice tongue. Following the loss of the ice tongue, calving became the dominant mechanism of retreat, owing retreat mechanism, due to the exposure of a tall calving front with correspondingly high stresses

exceeding the calving threshold. Calving rates rapidly increased as SK retreated into deeper waters and therefore experienced greater tensile stresses along its calving front. ~~The best-fitting simulations display the largest discrepancies with observations during periods of rapid retreat or periods of transition between dominant ablation modes.~~

Top panel (a) depicts the average rate of surface height change for all ensemble members along a flowline (denoted in (b)) along the southern SK branch. The average rate of observed height change is obtained from ICESat and ICESat-2 data from 2003–2019 (Smith et al., 2020). Bottom panel (b) displays the 2018 calving front positions for all ensemble members. Additionally the best fit model calving front is highlighted with a dark orange and the observed 2019 winter calving front Smith et al. (2020) with black. The thin yellow line indicates the flowline on which we calculate surface height change.

Additionally, throughout our model space, a bimodality in calving front positions by 2018 occurs. As seen in Figure 5 there exists two states of final calving front positions. On one end, there exists the set of simulations which do not experience ice tongue loss and subsequently do not experience rapid calving ablation. These simulations are characterized by either lower calving stress thresholds or melting multipliers. While calving front locations are more sensitive to calving thresholds, an insufficiently powerful melting regime is unable to generate a rapid enough removal of the ice tongue which would eventually cascade into the growth of mechanical imbalances at the calving front and subsequent retreat. Within these simulations, the calving front is quickly arrested and remains quasi-stable. On the other end we see simulations which quickly lose their ice tongue and subsequently accelerate beyond observational records. In this set of simulations, the calving stress threshold plays a greater role than melting in setting the rate of retreat as the grounding line recedes onto deeper bedrock and the glacier front experiences greater mechanical stresses. When letting these simulations continue beyond 2018, we note that rapid calving continues along the southern trunk of SK, but stops across the northern trunk, owing to the presence or lack of steepening bedrock gradients, respectively. Interestingly, we do not witness simulations with calving fronts arresting along a position between the observed calving front in 2018 and a few kms inland from the initial calving front in 1985. The most likely reason for this lies in our simplified parameterization of a dynamic ice mélange. In the absence of mélange, the gross behavior of the SK calving front as modeled here can either be characterized by rapid and vigorous retreat or terminal stability. This leads to a bimodality in model response, while, in reality, ice mélange acts as a stress buffer in response to rapid calving events.

A central finding of this study is that the ability of ice mélange to buttress SK's calving front ~~is increasing~~ increases in importance as ~~its~~ SK's calving front rapidly retreats onto deeper beds. ~~By~~ We tested this by extending simulations until 2100 ~~we and~~ analyzed the potential for ~~ice mélange~~ increased calving stress threshold to arrest further retreat of SK. Our simulations reveal that a sufficiently robust ice mélange ~~is able to~~ could suppress calving activity during SK's most vigorous calving ~~phases~~ phase. The loss of SK's ice tongue subjected the calving front to greater tensile stress, which consequently amplified the importance of ice mélange importance in modulating calving rates. ~~A weakening of ice mélange is likely to have occurred in response to~~ We hypothesize that the influx of warm waters into ~~Ilulissat Icefjord, which melted integral ice bonds and the Ilulissat icefjord~~ facilitated downstream movement and fragmentation ~~of ice mélange. Further testing of this hypothesis is required, but we have shown here that calving variability, in its current state, is the dominant control on SK's evolution. Additionally, simple parameterizations of mélange such as those employed here and other studies attempting to quantify mélange buttressing effects~~ (Schlemm and Levermann, 2021a; Parsons et al., 2024) are ultimately meant as upper bounds on glacier-mélange interactions

since they do not represent the true rheology of mélange nor its two-way with the calving front. This finding emphasizes the importance of future development of dynamic models of ice mélange evolution that can be coupled to ~~ice sheet models~~ models of vigorously calving glaciers .

Code availability. Relevant run and analysis scripts required to produce results found at a persistent Zenodo repository located at
490 <https://doi.org/10.5281/zenodo.11176342>.

Author contributions. All author help conceive the study and contributed to the manuscript. ZR conducted model simulations, post-processing and analysis.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. Thanks to Blake Castleman for early work on simulations, and Johannes Bondzio for answering questions on ISSM
495 configuration. Vincent Verjans and John Christian provided useful comments during the completion of this work. This research and authors were supported by funding from a JPL Strategic University Research Partnership (SURP) SP20007. Z. Rashed and A. Robel were also supported by NSF OPP Grant 2025692. H. Seroussi was also supported by NSF NNA Grant 2127246.

References

- Amaral, T., Bartholomaus, T. C., and Enderlin, E. M.: Evaluation of iceberg calving models against observations from Greenland outlet glaciers, *Journal of Geophysical Research: Earth Surface*, 125, e2019JF005444, 2020.
- Amundson, J. M. and Burton, J.: Quasi-static granular flow of ice mélange, *Journal of Geophysical Research: Earth Surface*, 123, 2243–2257, 2018.
- Amundson, J. M., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M. P., and Motyka, R. J.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001405>,
_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009JF001405>, 2010.
- Benn, D. I., Warren, C. R., and Mottram, R. H.: Calving processes and the dynamics of calving glaciers, *Earth-Science Reviews*, 82, 143–179, <https://doi.org/10.1016/j.earscirev.2007.02.002>, 2007.
- Benn, D. I., Cowton, T., Todd, J., and Luckman, A.: Glacier Calving in Greenland, *Current Climate Change Reports*, 3, 282–290, <https://doi.org/10.1007/s40641-017-0070-1>, 2017.
- Bevan, S. L., Luckman, A. J., Benn, D. I., Cowton, T., and Todd, J.: Impact of warming shelf waters on ice mélange and terminus retreat at a large SE Greenland glacier, *The Cryosphere*, 13, 2303–2315, <https://doi.org/10.5194/tc-13-2303-2019>, 2019.
- Bondzio, J. H., Seroussi, H., Morlighem, M., Kleiner, T., Rückamp, M., Humbert, A., and Larour, E. Y.: Modelling calving front dynamics using a level-set method: application to Jakobshavn Isbræ, West Greenland, *The Cryosphere*, 10, 497–510, <https://doi.org/10.5194/tc-10-497-2016>, publisher: Copernicus GmbH, 2016.
- Bondzio, J. H., Morlighem, M., Seroussi, H., Kleiner, T., Rückamp, M., Mouginot, J., Moon, T., Larour, E. Y., and Humbert, A.: The mechanisms behind Jakobshavn Isbræ’s acceleration and mass loss: A 3-D thermomechanical model study, *Geophysical Research Letters*, 44, 6252–6260, <https://doi.org/10.1002/2017GL073309>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL073309>, 2017.
- Bondzio, J. H., Morlighem, M., Seroussi, H., Wood, M. H., and Mouginot, J.: Control of Ocean Temperature on Jakobshavn Isbræ’s Present and Future Mass Loss, *Geophysical Research Letters*, 45, 12,912–12,921, <https://doi.org/10.1029/2018GL079827>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL079827>, 2018.
- Budd, W. F., Jenssen, D., and Smith, I. N.: A Three-Dimensional Time-Dependent Model of the West Antarctic Ice Sheet, *Annals of Glaciology*, 5, 29–36, <https://doi.org/10.3189/1984AoG5-1-29-36>, 1984.
- Burton, J. C., Amundson, J. M., Cassotto, R., Kuo, C.-C., and Dennin, M.: Quantifying flow and stress in ice mélange, the world’s largest granular material, *Proceedings of the National Academy of Sciences*, 115, 5105–5110, 2018.
- Cassotto, R., Fahnestock, M., Amundson, J. M., Truffer, M., and Joughin, I.: Seasonal and interannual variations in ice mélange and its impact on terminus stability, Jakobshavn Isbræ, Greenland, *Journal of Glaciology*, 61, 76–88, <https://doi.org/10.3189/2015JoG13J235>, publisher: Cambridge University Press, 2015.
- Choi, Y., Morlighem, M., Wood, M., and Bondzio, J. H.: Comparison of four calving laws to model Greenland outlet glaciers, *The Cryosphere*, 12, 3735–3746, 2018.
- Echelmeyer, K. and Harrison, W. D.: Jakobshavns Isbræ, West Greenland: Seasonal Variations in Velocity - or Lack Thereof, *Journal of Glaciology*, 36, 82–88, <https://doi.org/10.3189/S0022143000005591>, publisher: Cambridge University Press, 1990.
- Ettema, J., van den Broeke, M. R., van Meijgaard, E., van de Berg, W. J., Bamber, J. L., Box, J. E., and Bales, R. C.: Higher surface mass balance of the Greenland ice sheet revealed by high-resolution climate modeling, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL038110>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009GL038110>, 2009.

- 535 Gladish, C. V., Holland, D. M., and Lee, C. M.: Oceanic Boundary Conditions for Jakobshavn Glacier. Part II: Provenance and Sources of Variability of Disko Bay and Ilulissat Icefjord Waters, 1990–2011, *Journal of Physical Oceanography*, 45, 33–63, <https://doi.org/https://doi.org/10.1175/JPO-D-14-0045.1>, 2015.
- Hanna, E., Huybrechts, P., Cappelen, J., Steffen, K., Bales, R. C., Burgess, E., McConnell, J. R., Peder Steffensen, J., Van den Broeke, M., Wake, L., Bigg, G., Griffiths, M., and Savas, D.: Greenland Ice Sheet surface mass balance 1870 to 2010 based
540 on Twentieth Century Reanalysis, and links with global climate forcing, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016387>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2011JD016387>, 2011.
- Holland, D. M. and Jenkins, A.: Modeling Thermodynamic Ice–Ocean Interactions at the Base of an Ice Shelf, *Journal of Physical Oceanography*, 29, 1787–1800, [https://doi.org/10.1175/1520-0485\(1999\)029<1787:MTIOIA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1787:MTIOIA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Physical Oceanography*, 1999.
- 545 Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm sub-surface ocean waters, *Nature Geoscience*, 1, 659–664, <https://doi.org/10.1038/ngeo316>, number: 10 Publisher: Nature Publishing Group, 2008.
- Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *The Cryosphere*, 8, 1509–1518, <https://doi.org/10.5194/tc-8-1509-2014>, publisher: Copernicus GmbH, 2014.
- 550 Joughin, I.: MEaSUREs Greenland ice sheet velocity map from InSAR data, version 2, Data Set, 2015.
- Joughin, I., Abdalati, W., and Fahnestock, M.: Large fluctuations in speed on Greenland’s Jakobshavn Isbræ glacier, *Nature*, 432, 608–610, <https://doi.org/10.1038/nature03130>, number: 7017 Publisher: Nature Publishing Group, 2004.
- Joughin, I., Howat, I., Alley, R. B., Ekstrom, G., Fahnestock, M., Moon, T., Nettles, M., Truffer, M., and Tsai, V. C.: Ice-front variation and tidewater behavior on Helheim and Kangerdlugssuaq Glaciers, Greenland, *Journal of Geophysical Research: Earth Surface*, 113,
555 <https://doi.org/10.1029/2007JF000837>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007JF000837>, 2008.
- Joughin, I., Smith, B. E., and Medley, B.: Marine Ice Sheet Collapse Potentially Under Way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735–738, <https://doi.org/10.1126/science.1249055>, publisher: American Association for the Advancement of Science, 2014.
- Joughin, I., Shean, D. E., Smith, B. E., and Floricioiu, D.: A decade of variability on Jakobshavn Isbræ: ocean temperatures pace speed through influence on mélange rigidity, *The Cryosphere*, 14, 211–227, <https://doi.org/10.5194/tc-14-211-2020>, publisher: Copernicus
560 GmbH, 2020.
- Kajanto, K., Seroussi, H., de Fleurian, B., and Nisancioglu, K. H.: Present day Jakobshavn Isbræ (West Greenland) close to the Holocene minimum extent, *Quaternary Science Reviews*, 246, 106492, <https://doi.org/https://doi.org/10.1016/j.quascirev.2020.106492>, 2020.
- Kajanto, K., Straneo, F., and Nisancioglu, K.: Impact of icebergs on the seasonal submarine melt of Sermeq Kujalleq, *The Cryosphere*, 17, 371–390, <https://doi.org/10.5194/tc-17-371-2023>, 2023.
- 565 Kehrl, L. M., Joughin, I., Shean, D. E., Floricioiu, D., and Krieger, L.: Seasonal and interannual variabilities in terminus position, glacier velocity, and surface elevation at Helheim and Kangerlussuaq Glaciers from 2008 to 2016, *Journal of Geophysical Research: Earth Surface*, 122, 1635–1652, <https://doi.org/https://doi.org/10.1002/2016JF004133>, 2017.
- Khazendar, A., Fenty, I. G., Carroll, D., Gardner, A., Lee, C. M., Fukumori, I., Wang, O., Zhang, H., Seroussi, H., Moller, D., Noël, B. P. Y., van den Broeke, M. R., Dinardo, S., and Willis, J.: Interruption of two decades of Jakobshavn Isbrae acceleration and thinning as regional
570 ocean cools, *Nature Geoscience*, 12, 277–283, <https://doi.org/10.1038/s41561-019-0329-3>, number: 4 Publisher: Nature Publishing Group, 2019.

- Korsgaard, N. J., Nuth, C., Khan, S. A., Kjeldsen, K. K., Bjørk, A. A., Schomacker, A., and Kjær, K. H.: Digital elevation model and orthophotographs of Greenland based on aerial photographs from 1978–1987, *Scientific Data*, 3, 160032, <https://doi.org/10.1038/sdata.2016.32>, number: 1 Publisher: Nature Publishing Group, 2016.
- 575 Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002140>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2011JF002140>, 2012.
- Lee, R. W. and Schulson, E. M.: The Strength and Ductility of Ice Under Tension, *Journal of Offshore Mechanics and Arctic Engineering*, 110, 187–191, <https://doi.org/10.1115/1.3257049>, 1988.
- 580 Luckman, A., Benn, D. I., Cottier, F., Bevan, S., Nilsen, F., and Inall, M.: Calving rates at tidewater glaciers vary strongly with ocean temperature, *Nature communications*, 6, 8566, 2015.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment: Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research: Solid Earth*, 94, 4071–4087, <https://doi.org/10.1029/JB094iB04p04071>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/JB094iB04p04071>, 1989.
- 585 Menemenlis, D., Campin, J.-M., Heimbach, P., Hill, C., Lee, T., Nguyen, A., Schodlok, M., and Zhang, H.: ECCO2: High resolution global ocean and sea ice data synthesis, *Mercator Ocean Quarterly Newsletter*, 31, 13–21, 2008.
- Morland, L. W. and Zainuddin, R.: Plane and Radial Ice-Shelf Flow with Prescribed Temperature Profile, in: *Dynamics of the West Antarctic Ice Sheet*, edited by Van der Veen, C. J. and Oerlemans, J., *Glaciology and Quaternary Geology*, pp. 117–140, Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-94-009-3745-1_7, 1987.
- 590 Morlighem, M., Rignot, E., Seroussi, H., Larour, E., Ben Dhia, H., and Aubry, D.: Spatial patterns of basal drag inferred using control methods from a full-Stokes and simpler models for Pine Island Glacier, West Antarctica, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL043853>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2010GL043853>, 2010.
- Morlighem, M., Bondzio, J., Seroussi, H., Rignot, E., Larour, E., Humbert, A., and Rebuffi, S.: Modeling of Store Gletscher’s calving dynamics, West Greenland, in response to ocean thermal forcing, *Geophysical Research Letters*, 43, 2659–2666, 2016.
- 595 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44, 11,051–11,061, <https://doi.org/10.1002/2017GL074954>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL074954>, 2017.
- 600 Motyka, R. J., Truffer, M., Fahnestock, M., Mortensen, J., Rysgaard, S., and Howat, I.: Submarine melting of the 1985 Jakobshavn Isbræ floating tongue and the triggering of the current retreat, *Journal of Geophysical Research: Earth Surface*, 116, <https://doi.org/10.1029/2009JF001632>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009JF001632>, 2011.
- Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six
605 years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, 116, 9239–9244, <https://doi.org/10.1073/pnas.1904242116>, publisher: Proceedings of the National Academy of Sciences, 2019.
- Myers, P. G. and Ribergaard, M. H.: Warming of the Polar Water Layer in Disko Bay and Potential Impact on Jakobshavn Isbrae, *Journal of Physical Oceanography*, 43, 2629–2640, <https://doi.org/10.1175/JPO-D-12-051.1>, publisher: American Meteorological Society Section: Journal of Physical Oceanography, 2013.

- 610 Noël, B., van de Berg, W. J., van Meijgaard, E., Kuipers Munneke, P., van de Wal, R. S. W., and van den Broeke, M. R.: Evaluation of the updated regional climate model RACMO2.3: summer snowfall impact on the Greenland Ice Sheet, *The Cryosphere*, 9, 1831–1844, <https://doi.org/10.5194/tc-9-1831-2015>, publisher: Copernicus GmbH, 2015.
- Parsons, R., Sun, S., Gudmundsson, G. H., Wuite, J., and Nagler, T.: Quantifying the Buttressing Contribution of Sea Ice to Crane Glacier, *EGUosphere*, 2024, 1–17, <https://doi.org/10.5194/egusphere-2024-1499>, 2024.
- 615 Petrovic, J. J.: Review Mechanical properties of ice and snow, *Journal of Materials Science*, 38, 1–6, <https://doi.org/10.1023/A:1021134128038>, 2003.
- Pollard, D., DeConto, R. M., and Alley, R. B.: A continuum model (PSUMEL1) of ice mélange and its role during retreat of the Antarctic Ice Sheet, *Geoscientific Model Development*, 11, 5149–5172, 2018.
- Rignot, E. and Mouginot, J.: Ice flow in Greenland for the International Polar Year 2008–2009, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051634>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2012GL051634>, 2012.
- 620 Rignot, E., Xu, Y., Menemenlis, D., Mouginot, J., Scheuchl, B., Li, X., Morlighem, M., Seroussi, H., den Broeke, M. v., Fenty, I., Cai, C., An, L., and Fleurian, B. d.: Modeling of ocean-induced ice melt rates of five west Greenland glaciers over the past two decades, *Geophysical Research Letters*, 43, 6374–6382, <https://doi.org/10.1002/2016GL068784>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016GL068784>, 2016.
- 625 Robel, A. A.: Thinning sea ice weakens buttressing force of iceberg mélange and promotes calving, *Nature Communications*, 8, 14 596, 2017.
- Schlemm, T. and Levermann, A.: A simple parametrization of mélange buttressing for calving glaciers, *The Cryosphere*, 15, 531–545, <https://doi.org/10.5194/tc-15-531-2021>, 2021a.
- Schlemm, T. and Levermann, A.: A simple parametrization of mélange buttressing for calving glaciers, *The Cryosphere*, 15, 531–545, 2021b.
- 630 Schweinsberg, A. D., Briner, J. P., Miller, G. H., Bennike, O., and Thomas, E. K.: Local glaciation in West Greenland linked to North Atlantic Ocean circulation during the Holocene, *Geology*, 45, 195–198, <https://doi.org/10.1130/G38114.1>, 2017.
- Seroussi, H., Morlighem, M., Rignot, E., Khazendar, A., Larour, E., and Mouginot, J.: Dependence of century-scale projections of the Greenland ice sheet on its thermal regime, *Journal of Glaciology*, 59, 1024–1034, <https://doi.org/10.3189/2013JoG13J054>, publisher: Cambridge University Press, 2013.
- 635 Seroussi, H., Morlighem, M., Larour, E., Rignot, E., and Khazendar, A.: Hydrostatic grounding line parameterization in ice sheet models, *The Cryosphere*, 8, 2075–2087, 2014.
- Slater, D. A., Straneo, F., Das, S. B., Richards, C. G., Wagner, T. J. W., and Nienow, P. W.: Localized Plumes Drive Front-Wide Ocean Melting of A Greenlandic Tidewater Glacier, *Geophysical Research Letters*, 45, 12,350–12,358, <https://doi.org/10.1029/2018GL080763>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL080763>, 2018.
- 640 Slater, D. A., Benn, D. I., Cowton, T. R., Bassis, J. N., and Todd, J. A.: Calving Multiplier Effect Controlled by Melt Under-cut Geometry, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006 191, <https://doi.org/10.1029/2021JF006191>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JF006191>, 2021.
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, 368, 1239–1242, <https://doi.org/10.1126/science.aaz5845>, 2020.
- 645 Vaňková, I. and Holland, D. M.: A model of icebergs and sea ice in a joint continuum framework, *Journal of Geophysical Research: Oceans*, 122, 9110–9125, 2017.

- Vaughan, D. G.: Relating the occurrence of crevasses to surface strain rates, *Journal of Glaciology*, 39, 255–266, 1993.
- Wang, K., Cheng, X., Chen, Z., Hui, F., Liu, Y., and Tian, Y.: Ocean contributes to the melting of the Jakobshavn Glacier front, *Science*
650 *China Earth Sciences*, 63, 405–411, <https://doi.org/10.1007/s11430-019-9394-6>, 2020.
- Xie, S., Dixon, T. H., Holland, D. M., Voytenko, D., and Vaňková, I.: Rapid iceberg calving following removal of tightly packed pro-glacial
mélange, *Nature Communications*, 10, 3250, <https://doi.org/10.1038/s41467-019-10908-4>, 2019.
- Xu, Y., Rignot, E., Fenty, I., Menemenlis, D., and Flexas, M. M.: Subaqueous melting of Store Glacier, west Greenland
655 from three-dimensional, high-resolution numerical modeling and ocean observations, *Geophys. Res. Lett.*, 40, 4648–4653,
<https://doi.org/10.1002/grl.50825>, 2013.