

Representation of iron aerosol size distributions of anthropogenic emissions is critical in evaluating atmospheric soluble iron input to the ocean

Mingxu Liu^{1,2}, Hitoshi Matsui¹, Douglas S. Hamilton³, Sagar D. Rathod⁴, Kara D. Lamb⁵, Natalie M. Mahowald⁶

¹Graduate School of Environmental Studies, Nagoya University, Nagoya, Japan

²State Key Joint Laboratory of Environmental Simulation and Pollution Control, College of Environmental Science and Engineering, Peking University, Beijing, China

³Department of Marine, Earth, and Atmospheric Sciences, North Carolina State University, Raleigh, NC, USA

⁴La Follette School of Public Affairs, University of Wisconsin-Madison, Madison, WI, USA

⁵Department of Earth and Environmental Engineering, Columbia University, New York, NY, USA

⁶Department of Earth and Atmospheric Science, Cornell University, Ithaca, NY, USA

Correspondence to: M.L. (liumingxu@pku.edu.cn) and H.M. (matsui@nagoya-u.jp)

Abstract. Atmospheric aerosol deposition acts as a major source of soluble (bioavailable) iron in open ocean regions where it limits phytoplankton growth and primary production. The aerosol size distribution of emitted iron particles, along with particle growth from mixing with other atmospheric components, is an important modulator of its long-range transport potential. There currently exists a large uncertainty in the particle size distribution of iron aerosol, and the extent to which such uncertainty shapes global soluble iron deposition remains unclear. Here, we couple a sophisticated microphysical, size-resolved aerosol model with an iron-speciated and -processing module to disentangle the impact of iron emission size distributions on soluble iron input to the ocean, with a focus on anthropogenic combustion and metal smelting sources. We evaluate our model results against a global-scale flight measurement dataset for anthropogenic iron concentration and show that the different representations of iron size distribution at emission, as adopted in previous studies, introduces a variability in modeled iron concentrations over remote oceans of a factor of 10. Shifting the iron aerosol size distribution toward finer particle sizes (<1 μm) enables longer atmospheric lifetime (a doubling), promoting atmospheric processing that enhances the soluble iron deposition to ocean basins by up to 50% on an annual basis. The monthly enhancements reach 110% and 80% over the Southern Ocean and North Pacific Ocean, respectively. Uniquely, our results highlight that compared with emission flux variability, iron emission size distribution plays an equally important role in regulating soluble iron deposition, especially to the remote oceans. Our new findings can help to interpret inter-model differences in iron deposition estimation and to better quantify the effects of atmospheric nutrients input on marine biogeochemistry, including but not limited to iron, phosphorus, and others.

30 **1. Introduction**

31 Iron is a critical micronutrient supporting marine primary production, which is closely associated with marine carbon-nitrogen
32 cycles in the Earth system (Mahowald et al., 2018; Moore et al., 2001). The atmospheric deposition of soluble iron to many
33 ocean basins has long been regarded as an important source of bioavailable iron for ocean biota uptake in iron-limited areas
34 (Jickells et al., 2005; Jickells and Moore, 2015; Tagliabue et al., 2017). Understanding the amount and past-to-future evolution
35 of atmospheric iron deposition to the ocean is critical in assessing the ocean carbon sequestration under a changing climate
36 (Bergas-Massó et al., 2023; Myriokefalitakis et al., 2020; Hamilton et al., 2020b).

37 The quantification of soluble (bioavailable) iron input to the ocean is linked to differences in iron emission source properties,
38 the degree to which iron aerosol undergoes acidic or organic chemistry, and atmospheric transport (Hamilton et al., 2022).
39 Atmospheric iron comes from three major emission sources, i.e., wind-blowing dust, wildfire and biomass burning, and
40 anthropogenic activities, such as fossil fuel combustion and iron smelting. Dust storms, which frequently occur in arid or semi-
41 arid regions of the world, such as North Africa and East Asia, provide an abundant iron source to the ocean and support primary
42 production (Mahowald et al., 2009; Westberry et al., 2023). In addition, a growing body of evidence is showing that pyrogenic
43 iron, with higher fractional solubility than dust (Ito et al., 2019), is a large source of atmospheric soluble iron deposition to
44 many ocean basins, including the Southern Ocean, Northern Pacific Ocean, and Northern Atlantic Ocean (Conway et al., 2019;
45 Liu et al., 2022; Matsui et al., 2018; Seo and Kim, 2023). Because the strength of each source could be affected by future
46 climate change and/or human activities, their contributions to bioavailable iron input to the ocean may vary regionally and
47 temporally by the end of the century (Bergas-Massó et al., 2023).

48 Atmospheric transport provides the essential pathway in which iron aerosol emitted from the land is supplied to the remote
49 ocean. Atmospheric circulation patterns dictate the main transport pathways for aerosol to follow and thereby which source
50 regions are important to consider in terms of their supply to ocean basins. Additionally, atmospheric transport enables internal
51 mixing of iron-bearing aerosols with other aerosol and gas components, like sulfates and organics; a process commonly known
52 as aging that facilitates the dissolution of iron from an insoluble state to a soluble state (Shi et al., 2020; Shi et al., 2012;
53 Solmon et al., 2009; Li et al., 2017). The atmospheric aging processes can significantly increase iron solubility and subsequent
54 soluble iron deposition, evidenced by both in-field and laboratory observations and global model simulations (Ito, 2015; Li et
55 al., 2017; Longo et al., 2016). Uncovering the underlying mechanisms of the aging processes and associated enhancement of
56 iron solubility during transport is an ongoing topic of investigation (Meskhidze et al., 2019; Shi et al., 2020).

57 To elucidate atmospheric flux of iron-containing aerosols to the ocean, global-scale aerosol models have been developed to
58 include a range of iron emission sources that currently show a large intermodal difference in flux estimates (Myriokefalitakis
59 et al., 2018). Part of the problem is that it is difficult to realistically reproduce the distribution of soluble iron concentrations
60 across all the different regions of the world, and especially over the remote polar oceans that are often characterized by low
61 iron concentrations with a high fractional solubility (Ito et al., 2019). Among those global aerosol simulations, the size
62 distribution of iron, which is an important consideration when determining aerosol lifetimes and thus its long-range transport
63 potential, is key in shaping atmospheric iron distributions (Hamilton et al., 2020a; Myriokefalitakis et al., 2018). Compared to
64 coarse-sized particles (e.g., larger than 1 μm), smaller particles generally feature lower loss rates with respect to dry deposition
65 and wet removal, resulting in longer atmospheric lifetimes; being transported a longer distance increases the potential for
66 atmospheric processing (i.e., longer period of aerosol ageing) and thus higher soluble iron deposition. Evidence from recent
67 works (Zhang et al., 2023; Zhang et al., 2022) pointed out that Fe sources and physicochemical properties largely differ
68 between coarse and fine particles, leading to different Fe solubility. Representation of iron size distribution in models could
69 be therefore important.

70 Iron aerosol characteristics depend in part on differences between source types. The iron mass size distribution associated with
71 natural dust sources commonly pertains to mineral dust aerosols, with the coarse-sized (diameter greater than 1 μm) fraction
72 dominant (Mahowald et al., 2014; Albani et al., 2014). Similarly, fire iron emissions are dominated (>80%) by coarse mode
73 particles (Hamilton et al., 2019), suggested to be due to the entrainment of local dust iron-bearing aerosol in the pyro convective
74 updrafts generated by a fire (Hamilton et al., 2022). For iron aerosol with an anthropogenic source, however, the relative
75 fractions between the fine and coarse particle size distribution at emission are more divergent among previous investigations.
76 Recent observational constraints reveal large mass concentrations of anthropogenic iron oxide in the fine mode (Moteki et al.,
77 2017), while a subset of modelling studies have treated most of this iron in the coarse mode (Wang et al., 2015). As opposed
78 to those natural sources, anthropogenic iron size distributions may be highly variable with respect to diverse fuels, combustion
79 temperatures, and industrial processes, as well as the abatement technologies applied to control air pollution (Rathod et al.,
80 2020; Hamilton et al., 2020a).

81 The extent to which iron aerosol size distributions shape the pattern of atmospheric soluble iron inputs to different ocean
82 regions is currently unknown. Herein, by leveraging a size-resolved global aerosol model configured with iron processes, we
83 focus on the representation of anthropogenic iron size distributions at emission, primarily involving its roles in altering global
84 long-range transport and deposition fluxes of iron. We further put the effect of iron size distribution in the context with iron
85 emission uncertainty to shed light on their relative importance in controlling global-scale iron deposition.

86

87 **2. Methods and Materials**

88 **2.1 Global aerosol model**

89 We conducted global aerosol simulations using the Community Atmospheric Model version 5 (CAM5.3) with the Aerosol
90 Two-dimensional bin module for foRmation and Aging Simulation version 2 (CAM-ATRAS) (Matsui, 2017; Matsui and
91 Mahowald, 2017). The model treats a series of aerosol chemical and microphysical processes in a size-resolved manner with
92 12 aerosol size bins from 1 to 10,000 nm in diameter. Our recent study suggests that this size-resolved method can well
93 represent the growth of small particles to larger ones and the evolution of particle size distributions during atmospheric
94 transport (Liu and Matsui, 2022). Both dry deposition (Zhang et al., 2001) and wet deposition (Liu et al., 2012) of aerosols are
95 treated in our model. The improvement of aerosol in-cloud wet scavenging process was included to improve the modelling of
96 aerosol long-range transport efficiency (Liu and Matsui, 2021). The CAM-ATRAS model has been adequately validated for
97 aerosol mass and number concentrations at a global scale using comprehensive measurements from the ground to the upper
98 troposphere (Matsui and Liu, 2021; Gliß et al., 2021; Kawai et al., 2021; Matsui et al., 2022; Matsui and Moteki, 2020).

99 To represent the iron cycle from emission to deposition, we explicitly treated iron constituents within the aerosol model, similar
100 to our previous study (Liu et al., 2022), while with updates for iron processing in the current work. All iron-bearing components
101 were assumed to be internally mixed with other aerosols and underwent emission, physical transport, chemical aging (e.g.,
102 solubilization), and deposition in the atmosphere. The internal mixing assumption is reasonable for anthropogenic iron aerosols,
103 which are often mixed together with other aerosol compounds in the polluted environments, e.g., East Asia, and enable the
104 growth of iron aerosols via condensation and coagulation. Our model simulated iron solubility and atmospheric processing of
105 iron-bearing aerosols through the online coupling with the Mechanism of Intermediate complexity for Modelling Iron
106 (Hamilton et al., 2019). For anthropogenic iron, we consider five different minerals, namely magnetite, hematite, illite,
107 kaolinite, and sulfate iron, following the global emission inventory by Rathod et al. (2020), which was developed by a bottom-
108 up approach at 1° spatial resolution and 1-month temporal resolution. The reference year of the inventory was 2010. We
109 account for a wide range of anthropogenic sources including iron smelting and fossil fuel combustion sources.

110 Dust iron emission was calculated by assuming a constant iron content of 3.5% in dust aerosol emission (Shi et al., 2012). The
111 model estimated total dust emission fluxes using the scheme of Zender et al. (2003), with modifications by Albani et al. (2014)
112 and the size distribution from Kok (2011). Dust mineralogy was not involved in our simulations. The global interannual mean
113 iron (insoluble + soluble) emissions from dust, biomass burning, and anthropogenic sources were 87 Tg Fe yr⁻¹, 1.1 Tg Fe yr⁻¹,
114 and 2.2 Tg Fe yr⁻¹, respectively.

115 We validated our modeled anthropogenic iron oxide concentrations against a global-scale aircraft measurement in the
116 troposphere consisting of eight campaigns for the periods of 2009–2011 and 2016–2018 (Lamb et al., 2021). These
117 observations provide mass concentrations of anthropogenic iron oxide, i.e., magnetite, with volume equivalent diameters
118 between 180–1290 nm. The model results were extracted along the flight tracks in time and space and further averaged in
119 several latitudinal bands across Pacific and Atlantic Oceans (see Fig. S1). More details can be seen in Liu et al. (2022).

120 The model was compiled with a horizontal resolution of 1.9° × 2.5° (latitude × longitude) and 30 vertical layers from the
121 surface to 40 km. We ran the model for the two periods, 2008–2011 and 2015–2018, with the first year in each period as spin-
122 up. The meteorological fields were nudged by the Modern-Era Retrospective analysis for Research and Applications Version
123 2. In addition, to provide implications for ocean biogeochemistry, we estimated the changes in marine net primary production
124 induced by iron inputs following the methods used by Rathod et al. (2022) and Okin et al. (2011), in which a cut-off (4 μmol
125 L⁻¹) of nitrate concentrations at surface water was chosen to define the geographical areas of iron-limited ocean basins.

126 **2.2 Representation of iron size distribution of anthropogenic emissions**

127 The number and mass size distribution of aerosol at emission is an essential parameter in aerosol modelling. However, due to
128 the limited knowledge about iron emission characteristics, the representation of iron size distributions varies greatly for
129 anthropogenic sources. We tested four different size distributions of anthropogenic iron at emission (Fig. 1), briefly described
130 as follows.

131 First, we adopted the same size distribution with our previous studies (Matsui et al., 2018; Liu et al., 2022) based on Moteki
132 et al. (2017), abbreviated as Moteki2017 hereafter, which was derived from a power law function to fit the observed
133 anthropogenic iron oxides concentrations within the boundary layer in the outflow of East Asian sources. Note that the
134 observed size-resolved number concentrations were confined within 170 to 2,100 nm in diameter given the detection limit. By
135 extrapolating the observation results, we obtained the mass size distribution between 1 nm and 10,000 nm with negative values
136 excluded. In this case, more than 90% of iron mass was allocated to the size range of 100 nm to 2,500 nm. In addition, we
137 varied the emissions by a factor of two (×0.5 and ×2.0, respectively) in another two parallel experiments to account for the
138 potential uncertainties in iron emission estimates.


139 The second case was derived from Rathod et al. (2020) (abbreviated as Rathod2020 hereafter). They developed a new
140 mineralogy-based iron emission inventory by introducing more details in anthropogenic sources, especially the inclusion of a
141 metal smelting source. These improvements increase the fine aerosol (less than 1 μm in diameter) fractions by a factor of 10
142 higher than most previous inventories. Consequently, this inventory was characterized with almost equal fractions between
143 fine- and coarse-sized emissions, while previous inventories always applied a much larger fraction for the coarse mode.
144 Consistent with Rathod et al., we allocated 10% and 90% of fine iron mass to the Aitken mode and the accumulation modes
145 of aerosols, respectively.

146 The third case was derived from Hamilton et al. (2019) (abbreviated as Hamilton2019 hereafter). They revised the
147 anthropogenic iron emission inventory based on Luo et al. (2008) (no metal smelting) and showed that the ratio of fine-sized
148 iron mass with that of the coarse-sized was 1:5.6, which resulted in the coarse mode dominating. A similar ratio for

149 anthropogenic iron emissions was applied by Ito (2013). Also, 10% of fine-sized emissions were allocated to the Aitken mode
150 and the remaining 90% to the accumulation mode.

151 The fourth case was derived from Wang et al. (2015) (abbreviated as Wang2015 hereafter). In their combustion-iron inventory,
152 the ratio of fine-mode mass to the coarse mode was as low as 1:24, because only 0.1–0.3 % of iron mass from coal fly ash
153 were emitted in the fine mode. Thus, we allocate 96% of iron in the coarse mode and the remaining 4% in the fine mode.

154 To enable the intercomparison among these cases, we used the global-scale anthropogenic iron emission mass inventory from
155 Rathod et al. (2020) but with different allocations between fine and coarse sizes in each case. The size distribution of
156 anthropogenic iron emission in each case was treated uniformly on a global scale. Therefore, the simulated variability in
157 atmospheric iron input to the ocean between cases should be attributable to iron size distributions rather than emission amount.
158 For the last three cases, we adopted three constant log-normal modes to distribute iron emissions, namely Aitken, accumulation,
159 and coarse modes, with their determining parameters including geometric standard deviations, number mode diameter, and
160 density reported by Hamilton et al. (2019). We then separated these three modes into 12 size bins from 1 nm to 10,000 nm
161 adapted for our size-resolved aerosol modelling. The size distributions of iron from biomass burning and dust sources were
162 consistent with Liu et al. (2022) in all cases. In the following analysis, we grouped the Moteki2017 and Rathod2020 as the
163 fine-sized group while the Hamilton2019 and Wang2015 as the coarse-sized group.



164
165 **Fig. 1.** Mass size distribution functions for anthropogenic iron emission adopted in four previous studies. The Moteki2017
166 curve was provided by fitting in-situ measurements for iron particles within the range of 170 and 2,100 nm in diameter over
167 East Asia; the Rathod2020 was based on the latest combustion iron emission inventory, with updates on iron estimate in fine-
168 mode sizes (<1 μ m); the Hamilton2019 and Wang2015 cases were modeling studies that assumed most anthropogenic
169 combustion in the coarse-mode bin.


170

171 3. Results and Discussion

172 3.1 Atmospheric iron aerosol concentrations


173 We first examined the effect of changes to iron particle size distributions for anthropogenic sources (unless otherwise stated)
174 on the atmospheric iron aerosol burden and its global distributions. Figure 2 illustrates that the global-mean anthropogenic
175 iron lifetimes differ by about a factor of 2 among the four examined cases. Both the Moteki2017 and Rathod2020 cases

176 simulated a lifetime around 3.0 days. In contrast, the Hamilton2019 and Wang2015 cases simulated a lifetime around half as
177 long (between 1.4 and 1.7 days). As only the size distribution is changed in these simulations the change in lifetime is directly
178 linked to the apportionment of mass aerosol between fine and coarse particle size modes. This result is in line with previous
179 reports (Hamilton et al., 2020a), and demonstrates that shifting emitted iron toward fine-sized diminishes atmospheric loss
180 rates of iron aerosols via dry (sedimentation) and wet (precipitation) removal pathways and extends their lifetime.
181 Consequently, given the same emission amount, the atmospheric iron burdens are enhanced, accordingly by approximately a
182 factor of 2 from ~9.0 Gg in the coarse-sized cases (Hamilton2019 and Wang2015) to ~18.0 Gg in the fine-sized cases
183 (Moteki2017 and Rathod2020). In a similar manner, the lifetimes and mass burdens of anthropogenic iron in the soluble form
184 are almost doubled in the fine-sized cases (Fig. 2b). The extended lifetimes also enhance the globally averaged iron solubility
185 (Fig. S2), by allowing more iron subject to aerosol aging and solubilization processes.



186
187 **Fig. 2.** Global anthropogenic iron concentration burdens and lifetimes varied by the emission size distributions. The scatter
188 plots are shown for (a) anthropogenic iron (labeled as AN iron) burden vs. lifetime and (b) soluble anthropogenic iron vs.
189 lifetime. Four representative cases are examined in this work: Moteki2017, Rathod2020, Hamilton2019, and Wang2015. The
190 first two cases are grouped into “FINE-sized” and the other two are grouped into “COARSE-sized”.

191



192


193 **Fig. 3.** Comparison of modelled anthropogenic iron mineral (magnetite) vertical concentration profiles in four cases with
 194 aircraft measurements across global oceans. The flight routes and model-observation sampling methods have been documented
 195 in Lamb et al. (2021) and Liu et al. (2022). The geographical location of each oceanic area is marked in Fig. S1. We also scaled
 196 up and down emission fluxes by a factor of 2 from the Moteki2017 case, respectively, to account for potential uncertainties in
 197 emission estimates (red dashed lines in the panels).

198 To evaluate each iron simulation, we compare simulated aerosol characteristics against global-scale aircraft measurements of
 199 anthropogenic magnetite within nine regions of the troposphere (Fig. 3). We extracted the modeled mass concentrations of
 200 iron aerosols with the size range similar to that of the measurements. Note that magnetite, comprised about 70% of
 201 anthropogenic iron emissions, can be used as an indicator of anthropogenic iron abundance in the atmosphere (Rathod et al.,
 202 2020; Matsui et al., 2018). Despite the same emission fluxes considered in all cases, their simulated magnetite aerosol
 203 concentrations can differ by up to a factor of 10. Specifically, the Moteki2017 and Rathod2020 cases show a much higher

204 performance in reproducing the observed profiles over all ocean basins compared to Hamilton2019 and Wang2015 cases. In
 205 particular, the cases with a more uniformly distributed particle size distribution across modes captures the high concentration
 206 ($>1 \text{ ng m}^{-3}$) in North Pacific, which can be linked to atmospheric plumes transported from East Asia with intensive emission
 207 rates (Seo and Kim, 2023; Moteki et al., 2017). Some underestimations still exist in near-surface or high altitudes. Doubling
 208 the emission fluxes based the Moteki2017 case can appreciably narrow the gaps with the observation (dashed red lines in Fig.
 209 3). In contrast, the coarse-sized dominated simulations (i.e., Hamilton2019 and Wang2015) underrepresents the magnetite
 210 concentrations over global remote oceans, particularly by up to one order of magnitude over the Pacific Ocean The shorter
 211 lifetimes in this group limit the long-range transport of iron aerosols from continental sources to the remote atmosphere.

212 These results imply that agreement between observations and model simulations can be improved by reducing uncertainties in
 213 the emission inventory and in the long-range transport efficiency associated with representation of iron size distributions.
 214 Moreover, as illustrated in Fig. 3, the variabilities (the ratio between the maximum and minimum) of simulated magnetite
 215 vertical profiles by iron size distribution changes are wider than those by the emission uncertainties, for which emission fluxes
 216 were perturbed by a factor of 2 ($\times 2.0$ and $\times 0.5$, respectively) to test the sensitivity of simulated iron concentrations. We
 217 therefore highlight that in order to observationally constrain iron emissions more realistically in global aerosol simulations, it
 218 is a prerequisite to use a realistic empirical representation of anthropogenic iron aerosol size distributions.

219




220

221 **Fig. 4.** Global map of variability in yearly averaged iron column concentrations between the fine-sized group and the coarse-
 222 sized group. The variability ratios (fine-sized/coarse-sized) are calculated for (a) anthropogenic iron, (b) total iron, and (c)
 223 soluble iron, respectively. Herein, total iron is a combination of iron from dust, biomass burning, and anthropogenic sources,
 224 the soluble form of which denotes soluble iron. The ratios represent the maximum differences between the fine-sized group
 225 (the Moteki2017 and Rathod2020 cases) and the coarse-sized group (the Hamilton2019 and Wang2015 cases) and indicate the
 226 spread of iron simulations.

227 Our simulations further demonstrate that the representation of iron size distribution shapes the iron aerosol concentrations at a
 228 global scale (Fig. 4). Anthropogenic iron in the fine-sized group shows higher column concentrations by more than a factor of
 229 2 than in the coarse-sized over oceans. The differences (ratios) are larger in those remote oceans compared to source regions
 230 like East Asia, southern Africa and South America, because iron-bearing aerosols in smaller sizes can be transported to a
 231 longer distance. The variability between the two groups is relatively minor for total iron (Fig. 4b), which includes contributions
 232 from dust and biomass burning sources. By contrast, the differences for soluble iron are more pronounced than for total iron
 233 over much of the global Ocean (Fig 4c), because of the higher solubility of anthropogenic iron than dust iron. Over East Asia
 234 and its outflow areas, the rapid aging process in the polluted environments are capable of enhancing iron solubility, particularly
 235 those of anthropogenic origin, and thus amplify the differences of soluble iron concentrations between the fine-sized and
 236 coarse-sized groups (Li et al., 2017; Zhu et al., 2022).

237




239

240 **Fig. 5.** Comparison of yearly accumulated soluble iron input to the ocean between the fine-sized and coarse-sized cases. Shown
 241 here are (a) the ratio of anthropogenic soluble iron simulated in the fine-sized case to the result of the coarse-sized, (b) the
 242 ratio of total soluble iron, (c) the fractional contribution (in percentage) of anthropogenic emission to total soluble iron input
 243 in the fine-sized case, and (d) the fractional contribution of anthropogenic emission in the coarse-sized case.

244 Next, we examine the extent to which the iron size distributions at emission can alter soluble iron input to the global ocean
 245 basins, which is vital to net primary productivity, especially in the high-nutrient, low-chlorophyll (HNLC) areas (Hamilton et
 246 al., 2022; Moore et al., 2013). As illustrated in Fig. 5, though the global emission and the resulting annual iron (insoluble +
 247 soluble) deposition amount are the same between cases, their geographical distributions vary substantially. Using the ratio of
 248 annual soluble iron deposition in the fine-sized group to that of the coarse-sized as a proxy of the variability, we find the fine-
 249 sized distributions lead to much more soluble iron input to remote ocean basins including North Pacific and Southern Ocean,
 250 by up to a factor of 4 for the anthropogenic sources and 1.5 for the total of all sources (i.e., anthropogenic + fire + dust). Similar
 251 spatial patterns emerge regarding the total iron deposition, suggesting the importance of long-range transport efficiency in
 252 regulating iron distributions (Fig. S3). The variability is negligible in the equatorial and subtropical Atlantic, where dust iron
 253 dominates soluble iron input to this area. The ratios less than 1, indicating reduced deposition fluxes in the fine-sized group,
 254 are found near the continental sources, including western U.S., Australia, and southern Africa, because of the slower deposition
 255 speed. However, in East Asia, which has most intensive anthropogenic iron emissions, the shift toward finer sizes also increase
 256 soluble iron deposition near the sources (e.g., eastern China). This is attributable to the rapid aging processes of the fine-sized
 257 iron in such polluted environment that convert more insoluble iron to its soluble form (Baldo et al., 2022; Zhang et al., 2022).


258 The source appointment of soluble iron deposition across ocean basins also varies with iron size distributions (Fig. 5c-d). The
 259 anthropogenic iron source becomes more dominant in North Pacific, North Atlantic, and parts of the Southern Ocean, with its
 260 fractional contribution reaching more than 50% in the fine-sized group compared to that of 30–40% in the coarse-sized. Such
 261 variability is attributable to the enhancement of anthropogenic soluble iron fluxes to those remote oceans by the shift of
 262 anthropogenic iron emission toward finer size bins. Globally, the soluble iron deposition from anthropogenic sources is 55.0
 263 Gg per year for the fine-sized group, larger than that of 35.3 Gg per year for the coarse-sized group. Hence, even though the
 264 same emission is applied in these simulations, the diversity of iron size distributions at emission yields a considerable
 265 variability of soluble iron deposition on a global basis. As discussed earlier, the extended iron lifetime by about a factor to 2
 266 in the fine-sized group allows more iron to be transported to a remote region and simultaneously increases the amount of
 267 atmospheric iron processing and dissolution to a soluble form. Of the total iron deposition, the soluble fraction is thus notably
 268 elevated. We also find that the chemical aging process, as the major source of soluble iron, controls the differences of soluble
 269 iron deposition over remote oceans between the fine-sized and coarse-sized groups (Fig. S4).



270
 271 **Fig. 6.** Ratios of yearly accumulated soluble iron deposition for four anthropogenic iron-containing minerals, comparing fine-
 272 sized and coarse-sized cases.


273 In this study, the explicit treatment of anthropogenic iron mineralogy enables us to identify the iron mineral-dependent
 274 variability. We find that for those coming primarily from fossil fuel combustion and iron smelting on land, namely magnetite,
 275 hematite, and clays, the shift of iron emissions toward finer size bins promotes their long-range transport and enhances
 276 corresponding soluble iron deposition to North Pacific, Equatorial Pacific, and Southern Ocean by more than a factor of 4 (Fig.
 277 6a-c). However, the size distribution effects on iron sulfates are pronounced only in the polar areas, which are subject to plumes

278 of middle latitude shipping emissions. In line with previous study (Rathod et al., 2022), the iron sulfates constitute an important
 279 contribution to soluble iron deposition away from major continental sources, predominately associated with shipping emission
 280 over local oceans rather than long-range transport from land (Fig. S5). Hence, the variability induced by iron size distributions
 281 is less remarkable for iron sulfates than for other anthropogenic minerals. These results also suggest that the relative importance
 282 of iron sulfates in total soluble iron deposition to remote oceans is altered by the size distributions of all other iron minerals
 283 that originate from continental sources. Because the size distributions of anthropogenic iron minerals may depend on different
 284 combustion processes, source- and region-specific size distribution representation is desirable in the future work.



285 **Fig. 7.** Differences of monthly total soluble iron deposition between the fine-sized and coarse-sized groups over specific ocean
 286 basins. Only anthropogenic iron emission sizes have been examined here. Histograms describe the ratio of monthly results in
 287 the fine-sized case to that of the coarse-sized for (a) Southern Ocean, (b) North Pacific, and (c) Equatorial Pacific, respectively.
 288 For comparison, black dots describe the ratio of the fine-sized results to that with global anthropogenic emission amount
 289 scaling down by a factor of 2. The ocean basins of interest are indicated at the top-left corner of each panel. The red dashed
 290 lines indicate the ratio of 1.0.
 291

292 It is critical to examine monthly soluble iron availability altered by emission size distributions, because ocean primary
 293 production can respond to iron inputs on the order of days to months (Guieu et al., 2014). From our results in Fig. 7, the
 294 importance of anthropogenic emission size distributions in shifting soluble iron deposition varies by month over potentially
 295 iron-limited ocean basins, i.e., HNLC regions, due to the episodic nature of natural iron sources (dust and wildfire) and their
 296 deposition. For the Southern Ocean, the monthly ratios of the fine-sized case to the coarse-sized span from 1.1 in September
 297 to 2.1 in December (Fig. 7a). The September peak of fire iron (shown in Fig. S6), possibly linked to low precipitation in
 298 southern winter, masks the variability in anthropogenic iron contributions by emission size distributions. Conversely, the
 299 largest difference in December is associated with the lowest contribution of natural sources (Fig. S6). By contrast, the monthly
 300 differences are less fluctuated in North Pacific, ranging between 1.3 and 1.8 (Fig. 7b). Anthropogenic emission dominates
 301 soluble iron throughout the year except in March-May, during which dust storms originating from East Asia frequently occur
 302 and regulate soluble iron inputs to North Pacific. The Equatorial Pacific has the lowest ratio amongst the three regions, because
 303 anthropogenic aerosol bearing plumes rarely arrive in this region and lots of rain out here can efficiently remove aerosols. For
 304 the three ocean basins, such differences related to the emission size treatments are even larger than those by adjusting the iron
 305 emission amount by a factor of 2 with the consideration of emission uncertainty (black dots in Fig. 7). We therefore suggest
 306 that compared with iron emission fluxes, the representation of size distributions for anthropogenic iron is equally or even more
 307 important to the estimation of total soluble iron deposition to remote oceans.



308

309 **Fig. 8.** Difference of marine net primary production sustained by atmospheric soluble iron between two iron size distribution
 310 groups. The panels display (a) absolute differences and (b) percentage differences in net primary production of the fine-sized
 311 group relative to the coarse-sized group. Here, following Rathod et al. (2022), we focus only on the iron-limited ocean basins,
 312 which are defined using a cut-off of nitrate concentrations at surface oceans.

313 We also provide an estimate of the changes in iron-sustained marine net primary production between the finer-sized and coarse-
 314 sized groups (Fig. 8). In line with the distributions of soluble iron deposition, the effects of finer-sized iron distributions can
 315 enhance primary production over remote oceans including the North Pacific Ocean and Southern Ocean as high as 50%.
 316 Considering that anthropogenic iron aerosols may contribute to >10% of the total marine productivity in the North Pacific
 317 Ocean (Rathod et al., 2022; Ito et al., 2020), the representation of their size distributions at emissions, mostly from East Asia,
 318 is particularly important in the Earth system modeling. The evolution of atmospheric iron-aerosol size characteristics and their
 319 emission fluxes can be critical players to ocean carbon sequestration from past to future. Hamilton et al. (2020) found that
 320 historical air pollution controls has cut down anthropogenic emission amounts of particles in coarse sizes, in turn elevating the
 321 mass fraction of finer-sized iron particles and thus the overall lifetime of atmospheric iron. Hence, the complex interactions of
 322 iron and the Earth system is linked to human activity effects on soluble iron availability to ocean basins.

323

324 4. Conclusion

325 This study explores the extent to which iron size distribution at emission, specifically from anthropogenic sources, alters
 326 estimates of soluble iron deposition to the open ocean. A global microphysical, size-resolved aerosol model is used to simulate
 327 the iron cycle, involving emission, atmospheric processing, and deposition on a global scale. The model treats iron mineralogy,
 328 size evolution, and chemical aging processes during atmospheric transport, which enables the investigation on the relationship
 329 between iron size distributions and iron long-range transport and subsequent deposition. We test four representative size
 330 distribution schemes for anthropogenic iron sources employed in previous studies.

331 We find that allocating a more balanced fraction of iron aerosol at emission into particle sizes less than $1 \mu\text{m}$, results in a longer
 332 atmospheric lifetime and mass burden of total iron aerosols by about a factor of 2 compared to a coarse-sized dominated case,
 333 primarily associated with the decreased loss rates via dry and wet removal processes. The evaluation of anthropogenic iron
 334 aerosols against the global-scale observation dataset reveals that despite the same emission fluxes considered in all cases, their
 335 simulated magnetite aerosol concentrations differ by up to a factor of 10, while the higher fine-sized cases agree better with
 336 the observations. It is therefore necessary to accurately represent iron size distributions in order to constrain iron emission
 337 fluxes more realistically with aerosol simulations and observations (Liu et al., 2022). Our simulations show that the resulting

338 annual soluble iron deposition differs by up to a factor of 1.5 over remote oceans including the North Pacific Ocean and
339 Southern Ocean, because the fine-sized group allows more iron to be transported to a long distance with enhanced atmospheric
340 processing. More importantly, the monthly soluble iron deposition, which is relevant to ocean primary production responses
341 over days to months, would be enhanced by 110% and 80% in the fine-sized case over the Southern Ocean and North Pacific
342 Ocean, respectively. Such differences are similar to or even larger than those with the consideration of emission uncertainty,
343 suggesting the equally important role of iron size distribution treatment.

344 This study unravels the critical role of iron size distributions in shaping atmospheric soluble iron inputs to global oceans,
345 especially to the remote regions. However, the realistic understanding of iron emission size distributions is still inadequate
346 given limited observation data. Targeted in-site measurements on iron aerosol size along with its mass and solubility at source
347 areas are highly desirable. Higher resolution models with finer grids and detailed microphysics are useful to explore iron
348 aerosol deposition and chemical aging processes at regional scales. Furthermore, our finding may be extended to other key
349 trace elements of importance to ocean biogeochemistry, like copper, manganese, and phosphorus.

350

351 **Competing interests:** The authors declare that they have no conflict of interest.

352 **Data availability:** The model data used to generate the figures can be available upon request.

353 **Code availability:** The CESM source code is publicly available from NCAR at: <https://www.cesm.ucar.edu/>.

354 **Acknowledgement**

355 This study was supported by the Ministry of Education, Culture, Sports, Science, and Technology and the Japan Society for
356 the Promotion of Science (MEXT/JSPS) KAKENHI Grants (JP19H04253, JP19H05699, JP19KK0265, JP20H00196,
357 JP20H00638, JP22H03722, JP22F22092, JP23H00515, JP23H00523, JP23K18519, JP23K24976, and JP24H02225); by the
358 MEXT Arctic Challenge for Sustainability II (ArCS II) Project (JPMXD1420318865); and by the Environment Research and
359 Technology Development Fund 2–2003 (JPMEERF20202003) and 2–2301 (JPMEERF20232001) of the Environmental
360 Restoration and Conservation Agency. Mingxu Liu was supported by special fund of State Key Joint Laboratory of
361 Environmental Simulation and Pollution Control (24Y03ESPCP) and JSPS Postdoctoral Fellowships for Research in Japan
362 (Standard). DSH was partly supported by the Scientific Committee on Oceanic Research International Working Group 167:
363 Reducing Uncertainty in Soluble Aerosol Trace Element Deposition. We acknowledged the NOAA Atmospheric Composition
364 and Climate Program for providing the aircraft observational data.

365

366 **References:**

- 367 Albani, S., Mahowald, N. M., Perry, A. T., Scanza, R. A., Zender, C. S., Heavens, N. G., Maggi, V., Kok, J. F.,
368 and Otto-Bliesner, B. L.: Improved dust representation in the Community Atmosphere Model, *J. Adv. Model.*
369 *Earth Syst.*, 6, 541-570, <https://doi.org/10.1002/2013MS000279>, 2014.
- 370 Baldo, C., Ito, A., Krom, M. D., Li, W., Jones, T., Drake, N., Ignatyev, K., Davidson, N., and Shi, Z.: Iron from
371 coal combustion particles dissolves much faster than mineral dust under simulated atmospheric acidic
372 conditions, *Atmos. Chem. Phys.*, 22, 6045-6066, 10.5194/acp-22-6045-2022, 2022.
- 373 Bergas-Massó, E., Gonçalves Ageitos, M., Myriokefalitakis, S., Miller, R. L., van Noije, T., Le Sager, P., Montan
374 é Pinto, G., and Pérez García-Pando, C.: Pre-Industrial, Present and Future Atmospheric Soluble Iron
375 Deposition and the Role of Aerosol Acidity and Oxalate Under CMIP6 Emissions, *Earth's Future*, 11,
376 10.1029/2022ef003353, 2023.

377 Conway, T. M., Hamilton, D. S., Shelley, R. U., Aguilar-Islas, A. M., Landing, W. M., Mahowald, N. M., and John,
378 S. G.: Tracing and constraining anthropogenic aerosol iron fluxes to the North Atlantic Ocean using iron
379 isotopes, *Nat Commun*, 10, 2628, 10.1038/s41467-019-10457-w, 2019.

380 Gliß, J., Mortier, A., Schulz, M., Andrews, E., Balkanski, Y., Bauer, S. E., Benedictow, A. M. K., Bian, H., Checa-
381 Garcia, R., Chin, M., Ginoux, P., Griesfeller, J. J., Heckel, A., Kipling, Z., Kirkevåg, A., Kokkola, H., Laj, P.,
382 Le Sager, P., Lund, M. T., Lund Myhre, C., Matsui, H., Myhre, G., Neubauer, D., van Noije, T., North, P.,
383 Olivié, D. J. L., Rémy, S., Sogacheva, L., Takemura, T., Tsigaridis, K., and Tsyro, S. G.: AeroCom phase III
384 multi-model evaluation of the aerosol life cycle and optical properties using ground- and space-based remote
385 sensing as well as surface in situ observations, *Atmos. Chem. Phys.*, 21, 87-128, 10.5194/acp-21-87-2021,
386 2021.

387 Guieu, C., Aumont, O., Paytan, A., Bopp, L., Law, C. S., Mahowald, N., Achterberg, E. P., Marañón, E., Salihoglu,
388 B., Crise, A., Wagener, T., Herut, B., Desboeufs, K., Kanakidou, M., Olgun, N., Peters, F., Pulido-Villena, E.,
389 Tovar-Sanchez, A., and Völker, C.: The significance of the episodic nature of atmospheric deposition to Low
390 Nutrient Low Chlorophyll regions, *Global Biogeochemical Cycles*, 28, 1179-1198,
391 <https://doi.org/10.1002/2014GB004852>, 2014.

392 Hamilton, D. S., Scanza, R. A., Rathod, S. D., Bond, T. C., Kok, J. F., Li, L., Matsui, H., and Mahowald, N. M.:
393 Recent (1980 to 2015) Trends and Variability in Daily-to-Interannual Soluble Iron Deposition from Dust,
394 Fire, and Anthropogenic Sources, *Geophysical Research Letters*, 47, e2020GL089688,
395 10.1029/2020gl089688, 2020a.

396 Hamilton, D. S., Scanza, R. A., Feng, Y., Guinness, J., Kok, J. F., Li, L., Liu, X., Rathod, S. D., Wan, J. S., Wu,
397 M., and Mahowald, N. M.: Improved methodologies for Earth system modelling of atmospheric soluble iron
398 and observation comparisons using the Mechanism of Intermediate complexity for Modelling Iron (MIMI
399 v1.0), *Geosci. Model Dev.*, 12, 3835-3862, 10.5194/gmd-12-3835-2019, 2019.

400 Hamilton, D. S., Moore, J. K., Arneeth, A., Bond, T. C., Carslaw, K. S., Hantson, S., Ito, A., Kaplan, J. O., Lindsay,
401 K., Nieradzik, L., Rathod, S. D., Scanza, R. A., and Mahowald, N. M.: Impact of Changes to the Atmospheric
402 Soluble Iron Deposition Flux on Ocean Biogeochemical Cycles in the Anthropocene, *Global Biogeochemical
403 Cycles*, 34, e2019GB006448, 10.1029/2019gb006448, 2020b.

404 Hamilton, D. S., Perron, M. M. G., Bond, T. C., Bowie, A. R., Buchholz, R. R., Guieu, C., Ito, A., Maenhaut, W.,
405 Myriokefalitakis, S., Olgun, N., Rathod, S. D., Schepanski, K., Tagliabue, A., Wagner, R., and Mahowald, N.
406 M.: Earth, Wind, Fire, and Pollution: Aerosol Nutrient Sources and Impacts on Ocean Biogeochemistry, 14,
407 303-330, 10.1146/annurev-marine-031921-013612, 2022.

408 Ito, A.: Global modeling study of potentially bioavailable iron input from shipboard aerosol sources to the ocean,
409 *Global Biogeochemical Cycles*, 27, 1-10, <https://doi.org/10.1029/2012GB004378>, 2013.

410 Ito, A.: Atmospheric Processing of Combustion Aerosols as a Source of Bioavailable Iron, *Environ. Sci. Technol.
411 Lett.*, 2, 70-75, 10.1021/acs.estlett.5b00007, 2015.

412 Ito, A., Ye, Y., Yamamoto, A., Watanabe, M., and Aita, M. N.: Responses of ocean biogeochemistry to atmospheric
413 supply of lithogenic and pyrogenic iron-containing aerosols, *Geol. Mag.*, 157, 741-756,
414 10.1017/S0016756819001080, 2020.

415 Ito, A., Myriokefalitakis, S., Kanakidou, M., Mahowald, N. M., Scanza, R. A., Hamilton, D. S., Baker, A. R.,
416 Jickells, T., Sarin, M., Bikkina, S., Gao, Y., Shelley, R. U., Buck, C. S., Landing, W. M., Bowie, A. R., Perron,
417 M. M. G., Guieu, C., Meskhidze, N., Johnson, M. S., Feng, Y., Kok, J. F., Nenes, A., and Duce, R. A.:
418 Pyrogenic iron: The missing link to high iron solubility in aerosols, *Sci. Adv.*, 5, eaau7671,
419 10.1126/sciadv.aau7671, 2019.

420 Jickells, T. and Moore, C. M.: The Importance of Atmospheric Deposition for Ocean Productivity, *Annu. Rev.
421 Ecol. Evol. Syst.*, 46, 481-501, 10.1146/annurev-ecolsys-112414-054118, 2015.

422 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce,
423 R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M.,
424 Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean
425 Biogeochemistry, and Climate, *Science*, 308, 67, 10.1126/science.1105959, 2005.

426 Kawai, K., Matsui, H., and Tobo, Y.: High Potential of Asian Dust to Act as Ice Nucleating Particles in Mixed-
427 Phase Clouds Simulated With a Global Aerosol-Climate Model, *J. Geophys. Res.-Atmos*, 126,
428 e2020JD034263, <https://doi.org/10.1029/2020JD034263>, 2021.

429 Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate
430 the size of the global dust cycle, *Proceedings of the National Academy of Sciences of the United States of
431 America*, 108, 1016, 10.1073/pnas.1014798108, 2011.

432 Lamb, K. D., Matsui, H., Katich, J. M., Perring, A. E., Spackman, J. R., Weinzierl, B., Dollner, M., and Schwarz,
433 J. P.: Global-scale constraints on light-absorbing anthropogenic iron oxide aerosols, *npj Clim. Atmos. Sci.*, 4,
434 10.1038/s41612-021-00171-0, 2021.

435 Li, W., Xu, L., Liu, X., Zhang, J., Lin, Y., Yao, X., Gao, H., Zhang, D., Chen, J., Wang, W., Harrison, R. M.,
436 Zhang, X., Shao, L., Fu, P., Nenes, A., and Shi, Z.: Air pollution–aerosol interactions produce more
437 bioavailable iron for ocean ecosystems, *Sci. Adv.*, 3, e1601749, 10.1126/sciadv.1601749, 2017.

438 Liu, M. and Matsui, H.: Improved Simulations of Global Black Carbon Distributions by Modifying Wet
439 Scavenging Processes in Convective and Mixed-Phase Clouds, *J. Geophys. Res.-Atmos*, 126, e2020JD033890,
440 <https://doi.org/10.1029/2020JD033890>, 2021.

441 Liu, M. and Matsui, H.: Secondary Organic Aerosol Formation Regulates Cloud Condensation Nuclei in the Global
442 Remote Troposphere, *Geophysical Research Letters*, 49, 10.1029/2022gl100543, 2022.

443 Liu, M., Matsui, H., Hamilton, D. S., Lamb, K. D., Rathod, S. D., Schwarz, J. P., and Mahowald, N. M.: The
444 underappreciated role of anthropogenic sources in atmospheric soluble iron flux to the Southern Ocean, *npj*
445 *Clim. Atmos. Sci.*, 5, 10.1038/s41612-022-00250-w, 2022.

446 Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J. F., Gettelman, A., Morrison, H., Vitt,
447 F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono,
448 M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in
449 climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*,
450 5, 709-739, 10.5194/gmd-5-709-2012, 2012.

451 Longo, A. F., Feng, Y., Lai, B., Landing, W. M., Shelley, R. U., Nenes, A., Mihalopoulos, N., Violaki, K., and
452 Ingall, E. D.: Influence of Atmospheric Processes on the Solubility and Composition of Iron in Saharan Dust,
453 *Environmental Science & Technology*, 50, 6912-6920, 10.1021/acs.est.6b02605, 2016.

454 Luo, C., Mahowald, N., Bond, T., Chuang, P. Y., Artaxo, P., Siefert, R., Chen, Y., and Schauer, J.: Combustion
455 iron distribution and deposition, *Global Biogeochemical Cycles*, 22, GB1012,
456 <https://doi.org/10.1029/2007GB002964>, 2008.

457 Mahowald, N., Albani, S., Kok, J. F., Engelstaeder, S., Scanza, R., Ward, D. S., and Flanner, M. G.: The size
458 distribution of desert dust aerosols and its impact on the Earth system, *Aeolian Res.*, 15, 53-71,
459 <https://doi.org/10.1016/j.aeolia.2013.09.002>, 2014.

460 Mahowald, N. M., Hamilton, D. S., Mackey, K. R. M., Moore, J. K., Baker, A. R., Scanza, R. A., and Zhang, Y.:
461 Aerosol trace metal leaching and impacts on marine microorganisms, *Nat. Commun.*, 9, 2614,
462 10.1038/s41467-018-04970-7, 2018.

463 Mahowald, N. M., Engelstaedter, S., Luo, C., Sealy, A., Artaxo, P., Benitez-Nelson, C., Bonnet, S., Chen, Y.,
464 Chuang, P. Y., Cohen, D. D., Dulac, F., Herut, B., Johansen, A. M., Kubilay, N., Losno, R., Maenhaut, W.,
465 Paytan, A., Prospero, J. M., Shank, L. M., and Siefert, R. L.: Atmospheric Iron Deposition: Global Distribution,
466 Variability, and Human Perturbations, *Annual Review of Marine Science*, 1, 245-278,
467 10.1146/annurev.marine.010908.163727, 2009.

468 Matsui, H.: Development of a global aerosol model using a two-dimensional sectional method: 1. Model design, *J.*
469 *Adv. Model. Earth Syst.*, 9, 1921-1947, 10.1002/2017ms000936, 2017.

470 Matsui, H. and Liu, M.: Importance of Supersaturation in Arctic Black Carbon Simulations, *Journal of Climate*, 34,
471 7843-7856, 10.1175/jcli-d-20-0994.1, 2021.

472 Matsui, H. and Mahowald, N.: Development of a global aerosol model using a two-dimensional sectional method:
473 2. Evaluation and sensitivity simulations, *J. Adv. Model. Earth Syst.*, 9, 1887-1920, 10.1002/2017ms000937,
474 2017.

475 Matsui, H. and Moteki, N.: High sensitivity of Arctic black carbon radiative effects to subgrid vertical velocity in
476 aerosol activation, *Geophysical Research Letters*, 47, e2020GL088978, 10.1029/2020GL088978, 2020.

477 Matsui, H., Mori, T., Ohata, S., Moteki, N., Oshima, N., Goto-Azuma, K., Koike, M., and Kondo, Y.: Contrasting
478 source contributions of Arctic black carbon to atmospheric concentrations, deposition flux, and atmospheric
479 and snow radiative effects, *Atmos. Chem. Phys. Discuss.*, 2022, 1-31, 10.5194/acp-2021-1091, 2022.

480 Matsui, H., Mahowald, N. M., Moteki, N., Hamilton, D. S., Ohata, S., Yoshida, A., Koike, M., Scanza, R. A., and
481 Flanner, M. G.: Anthropogenic combustion iron as a complex climate forcer, *Nat. Commun.*, 9, 1593,
482 10.1038/s41467-018-03997-0, 2018.

483 Meskhidze, N., Völker, C., Al-Abadleh, H. A., Barbeau, K., Bressac, M., Buck, C., Bundy, R. M., Croot, P., Feng,
484 Y., Ito, A., Johansen, A. M., Landing, W. M., Mao, J., Myriokefalitakis, S., Ohnemus, D., Pasquier, B., and
485 Ye, Y.: Perspective on identifying and characterizing the processes controlling iron speciation and residence
486 time at the atmosphere-ocean interface, *Marine Chemistry*, 217, 103704,
487 <https://doi.org/10.1016/j.marchem.2019.103704>, 2019.

488 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R.
489 J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E.,
490 Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A., and Ulloa,
491 O.: Processes and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6, 701-710, 10.1038/ngeo1765, 2013.

492 Moore, J. K., Doney, S. C., Glover, D. M., and Fung, I. Y.: Iron cycling and nutrient-limitation patterns in surface
 493 waters of the World Ocean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 49, 463-507,
 494 [https://doi.org/10.1016/S0967-0645\(01\)00109-6](https://doi.org/10.1016/S0967-0645(01)00109-6), 2001.

495 Moteki, N., Adachi, K., Ohata, S., Yoshida, A., Harigaya, T., Koike, M., and Kondo, Y.: Anthropogenic iron oxide
 496 aerosols enhance atmospheric heating, *Nat. Commun.*, 8, 15329, 10.1038/ncomms15329, 2017.

497 Myriokefalitakis, S., Gröger, M., Hieronymus, J., and Döscher, R.: An explicit estimate of the atmospheric nutrient
 498 impact on global oceanic productivity, *Ocean Sci.*, 16, 1183-1205, 10.5194/os-16-1183-2020, 2020.

499 Myriokefalitakis, S., Ito, A., Kanakidou, M., Nenes, A., Krol, M. C., Mahowald, N. M., Scanza, R. A., Hamilton,
 500 D. S., Johnson, M. S., Meskhidze, N., Kok, J. F., Guieu, C., Baker, A. R., Jickells, T. D., Sarin, M. M., Bikkina,
 501 S., Shelley, R., Bowie, A., Perron, M. M. G., and Duce, R. A.: Reviews and syntheses: the GESAMP
 502 atmospheric iron deposition model intercomparison study, *Biogeosciences*, 15, 6659-6684, 10.5194/bg-15-
 503 6659-2018, 2018.

504 Okin, G. S., Baker, A. R., Tegen, I., Mahowald, N. M., Dentener, F. J., Duce, R. A., Galloway, J. N., Hunter, K.,
 505 Kanakidou, M., Kubilay, N., Prospero, J. M., Sarin, M., Surapipith, V., Uematsu, M., and Zhu, T.: Impacts of
 506 atmospheric nutrient deposition on marine productivity: Roles of nitrogen, phosphorus, and iron, *Global
 507 Biogeochemical Cycles*, 25, <https://doi.org/10.1029/2010GB003858>, 2011.

508 Rathod, S. D., Hamilton, D. S., Mahowald, N. M., Klimont, Z., Corbett, J. J., and Bond, T. C.: A Mineralogy-
 509 Based Anthropogenic Combustion-Iron Emission Inventory, *J. Geophys. Res.-Atmos*, 125, e2019JD032114,
 510 10.1029/2019jd032114, 2020.

511 Rathod, S. D., Hamilton, D. S., Li, L., Mahowald, N. M., Matsui, H., Pierce, J. R., and Bond, T. C.: Atmospheric
 512 Radiative and Oceanic Biological Productivity Responses to Increasing Anthropogenic Combustion-Iron
 513 Emission in the 1850–2010 Period, *Geophys. Res. Lett.*, 49, 10.1029/2022gl099323, 2022.

514 Seo, H. and Kim, G.: Anthropogenic Iron Invasion into the Ocean: Results from the East Sea (Japan Sea), *Environ.
 515 Sci. Technol.*, 57, 10745-10753, 10.1021/acs.est.3c01084, 2023.

516 Shi, J., Guan, Y., Ito, A., Gao, H., Yao, X., Baker, A. R., and Zhang, D.: High Production of Soluble Iron Promoted
 517 by Aerosol Acidification in Fog, *Geophysical Research Letters*, 47, 10.1029/2019gl086124, 2020.

518 Shi, Z., Krom, M. D., Jickells, T. D., Bonneville, S., Carslaw, K. S., Mihalopoulos, N., Baker, A. R., and Benning,
 519 L. G.: Impacts on iron solubility in the mineral dust by processes in the source region and the atmosphere: A
 520 review, *Aeolian Res.*, 5, 21-42, <https://doi.org/10.1016/j.aeolia.2012.03.001>, 2012.

521 Solmon, F., Chuang, P. Y., Meskhidze, N., and Chen, Y.: Acidic processing of mineral dust iron by anthropogenic
 522 compounds over the north Pacific Ocean, *J. Geophys. Res.-Atmos*, 114,
 523 <https://doi.org/10.1029/2008JD010417>, 2009.

524 Tagliabue, A., Bowie, A. R., Boyd, P. W., Buck, K. N., Johnson, K. S., and Saito, M. A.: The integral role of iron
 525 in ocean biogeochemistry, *Nature*, 543, 51-59, 10.1038/nature21058, 2017.

526 Wang, R., Balkanski, Y., Boucher, O., Bopp, L., Chappell, A., Ciais, P., Hauglustaine, D., Peñuelas, J., and Tao,
 527 S.: Sources, transport and deposition of iron in the global atmosphere, *Atmos. Chem. Phys.*, 15, 6247-6270,
 528 10.5194/acp-15-6247-2015, 2015.

529 Westberry, T. K., Behrenfeld, M. J., Shi, Y. R., Yu, H., Remer, L. A., and Bian, H.: Atmospheric nourishment of
 530 global ocean ecosystems, 380, 515-519, doi:10.1126/science.abq5252, 2023.

531 Zender, C. S., Bian, H., and Newman, D.: Mineral Dust Entrainment and Deposition (DEAD) model: Description
 532 and 1990s dust climatology, *J. Geophys. Res.-Atmos*, 108, 4416, <https://doi.org/10.1029/2002JD002775>,
 533 2003.

534 Zhang, H., Li, R., Dong, S., Wang, F., Zhu, Y., Meng, H., Huang, C., Ren, Y., Wang, X., Hu, X., Li, T., Peng, C.,
 535 Zhang, G., Xue, L., Wang, X., and Tang, M.: Abundance and Fractional Solubility of Aerosol Iron During
 536 Winter at a Coastal City in Northern China: Similarities and Contrasts Between Fine and Coarse Particles, *J.
 537 Geophys. Res.-Atmos*, 127, e2021JD036070, <https://doi.org/10.1029/2021JD036070>, 2022.

538 Zhang, H., Li, R., Huang, C., Li, X., Dong, S., Wang, F., Li, T., Chen, Y., Zhang, G., Ren, Y., Chen, Q., Huang,
 539 R., Chen, S., Xue, T., Wang, X., and Tang, M.: Seasonal variation of aerosol iron solubility in coarse and fine
 540 particles at an inland city in northwestern China, *Atmos. Chem. Phys.*, 23, 3543-3559, 10.5194/acp-23-3543-
 541 2023, 2023.

542 Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric
 543 aerosol module, *Atmospheric Environment*, 35, 549-560, [https://doi.org/10.1016/S1352-2310\(00\)00326-5](https://doi.org/10.1016/S1352-2310(00)00326-5),
 544 2001.

545 Zhu, Y., Li, W., Wang, Y., Zhang, J., Liu, L., Xu, L., Xu, J., Shi, J., Shao, L., Fu, P., Zhang, D., and Shi, Z.: Sources
 546 and processes of iron aerosols in a megacity in Eastern China, *Atmos. Chem. Phys.*, 22, 2191-2202,
 547 10.5194/acp-22-2191-2022, 2022.

548

549