

# Bayesian reconstruction of sea-level and hydroclimates from coastal landform inversion: application to Santa Cruz (US) and Gulf of Corinth

Gino de Gelder<sup>1,2,3</sup>, Navid Hedjazian<sup>3</sup>, Laurent Husson<sup>1</sup>, Thomas Bodin<sup>4</sup>, Anne-Morwenn Pastier<sup>5</sup>, Yannick Boucharat<sup>1</sup>, Kevin Podoja<sup>6</sup>, Tubagus Solihuddin<sup>2</sup>, Sri Y. Cahyarini<sup>2</sup>

<sup>1</sup>ISTerre, IRD, CNRS, Université Grenoble-Alpes, Saint-Martin d'Hères, 38400, France

<sup>2</sup>Res. Group of Paleoclimate & Paleoenvironment, Res. Centr. for Climate and Atmosphere, Res. Org. of Earth Sciences and Maritime, National Research and Innovation Agency, Bandung, 40135, Indonesia

<sup>3</sup>ENS de Lyon, CNRS, LGL-TPE, Université Claude Bernard Lyon1, Villeurbanne, 69100, France

<sup>4</sup>Instituto de Ciencias del Mar (ICM) - CSIC, Barcelona, Spain

<sup>5</sup>GeoForschungsZentrum, Potsdam, 14473, Germany

<sup>6</sup>Université de Caen Normandie, Caen, 14000, France

*Correspondence to:* Gino de Gelder (gino.de-gelder@univ-grenoble-alpes.fr)

**Abstract.** Quantifying Quaternary sea-level changes and hydroclimatic conditions is an important challenge given their intricate relation with paleo-climate, ice-sheets and geodynamics. The world's coastlines provide an enormous geomorphologic archive, from which forward landscape evolution modelling studies have shown their potential to unravel paleo sea-levels, albeit at the cost of assumptions to the genesis of these landforms. We take a next step, by applying a Bayesian approach to jointly invert the geometries of multiple coastal terrace sequences to paleo sea- and lake level variations and extract past hydroclimatic conditions. Using a Markov chain Monte Carlo sampling method, we first test our approach on synthetic marine terrace profiles as proof of concept and benchmark our model on an observed marine terrace sequence in Santa Cruz (US). We successfully reproduce observed sequence morphologies and simultaneously obtain probabilistic estimates for past sea-level variations, as well as for other model parameters such as uplift and erosion rates. When applied to the semi-isolated Gulf of Corinth (Greece), our method allows deciphering the geomorphic Rosetta stone at an unprecedented resolution, revealing the connectivity between the Lake/Gulf of Corinth and the open sea for different hydroclimatic conditions. Eustatic sea-level and changing sill depths drive marine and transitional phases during interglacial and interstadial periods, whereas wetter and drier hydroclimates respectively over- and under-fill Lake Corinth during interstadial and glacial periods.

**Short summary.** Marine terrace sequences - staircase-shaped coastal landforms - record sea-level changes, vertical motions and erosional processes that are difficult to untangle. To do so we developed a numerical inversion approach: using the observed landscape as input, we constrain the ensemble of parameter ranges that could have created this landscape. We apply the model to marine terrace sequences in Santa Cruz (US) and Corinth (Greece) to reveal past sea/lake levels, uplift rates and hydro-climates.

## 1 Introduction

Reconstructions of Quaternary sea-level variations provide crucial constraints on thresholds and feedbacks within climatic and geodynamic systems that help understand how contemporary climate change may affect future sea level (Lambeck and Chappell, 2001; Hay et al., 2014; Dutton et al., 2015; Shakun et al., 2015; Austermann et al., 2017). A key archive of past sea-level is exposed within the geomorphology of most of the world's coastal areas in the form of paleo-shorelines (Johnson and Libbey, 1997; Podoja et al., 2011, 2014; Rovere et al., 2023; Fig. 1a), but it remains difficult to accurately translate coastal observations and measurements into paleo-sea-level estimates, and to evaluate the uncertainties inherent to these conversions. Major challenges include 1) dating of these landforms, as most paleo-


shorelines are erosive in nature (Pedoja et al., 2014) and absolute dating techniques themselves are complex and prone to large uncertainties (Strobl et al., 2014; Hibbert et al., 2016; Ott et al., 2019), 2) **observational** bias, which are mostly restricted to the most recent glacial cycle(s) and to periods where relative sea level was at similar elevations to present-day (Medina-Elizalde, 2013; Hibbert et al., 2016), 3) absence of reciprocity between paleo-shorelines and sea-level stands, as not all highstands lead to paleo-shorelines, and paleo-shorelines may have formed during one or many sea-level cycles (Guilcher, 1974; Malatesta et al., 2021; Chauveau et al., 2023), and 4) separating the tectonic from the sea-level component within relative sea-level changes (Pedoja et al., 2011).

Numerical models of landscape evolution started to overcome some of these limitations, by providing a means to quantitatively interpret undated paleo-shorelines, incorporate full sea-level curves instead of highstands only, unravelling the creation of paleo-shorelines formed over multiple glacial cycles, and considering multiple sea-level curves (e.g. Webster et al., 2007; Jara-Muñoz et al., 2019; Leclerc and Feuillet, 2019; De Gelder et al., 2020; 2023). So far, such numerical models have mainly been used for forward modelling approaches, where a number of proposed sea-level curves are used to predict shorelines, which are then compared to actual observations. However, this only provides a limited way to explore the full ensemble of possible sea-level histories and other model parameters like rock erosion rates or effective wave base depths, which are difficult to estimate. It follows that uncertainties in sea-level estimates from marine terraces remain poorly known, regardless of the method used, and in spite of uniformization attempts (Lorscheid and Rovere, 2019).

Semi-isolated marine basins, i.e. bodies of water that have been connected to the open sea in some intervals of their geologic history, and little or disconnected from the sea in other intervals, **develop in** hydrodynamic settings for which it is particularly complex to reconstruct sea and lake level. Such **basins**, like the Red Sea, Sea of Marmara (Turkey), Carioco Basin (Venezuela) and Gulf of Corinth (Greece), have a special geologic interest, given the active tectono-sedimentary processes driving their formation (e.g. Van Daele et al., 2011; McNeill et al., 2019), their sensitivity to rapid sea-level and climatic changes (e.g. Aksu et al., 1999; Siddall et al., 2004), and their role in dispersion of species (e.g. Derricourt, 2005). The main complexity in reconstructing sea-/lake-level fluctuations in such settings, is that 1) during disconnected phases these basins may have been underfilled or overfilled depending on local hydroclimate, and 2) the structural highs (sills) separating the basins from the sea can be simultaneously affected by tectonic vertical motion, sedimentation and erosion.

In this study, we intend to overcome common marine terrace analysis limitations, by using a Bayesian approach to invert the geometry of paleo-shoreline sequences. Our approach provides probabilistic estimates of paleo sea-level, erosion rates, uplift rates, wave-based depths and initial slopes. We focus on erosive marine terraces (Fig. 1b), which are both the most common type of paleo-shoreline (Pedoja et al., 2014), and are simpler to model than their depositional and bio-constructed equivalents (e.g. Pastier et al., 2019). We first apply our probabilistic inversion approach to a set of synthetic coastal profiles to test and illustrate the method, after which we invert a well-studied marine terrace sequence in Santa Cruz (US) to benchmark our model on a natural example. Finally, we use our approach on the semi-isolated Gulf of Corinth to decipher the complex combination of tectonic uplift, sea- and lake-level fluctuations, local climatic drivers and sill dynamics. **The Gulf of Corinth has been considered as a natural rift laboratory and has therefore received a lot of attention as a primary example for studying tectonic and surface processes in young rift systems (e.g. Armijo et al., 1996; Bernard et al., 2006; Nixon et al., 2016, Gallen & Fernández-Blanco, 2021). As a semi-isolated marine basin it has also been subjected to many studies on its environmental and climatic evolution (e.g. Collier et al., 2000; Watkins et al., 2019), yet, so far, previous studies could not resolve to what extent sea and lake levels have fluctuated.**

These case studies highlight how we can derive probabilistic estimates of past sea-level from marine terraces, and how the natural archive of paleo-shorelines can be further utilized to improve both paleo sea-level estimates, and unravel complex tectono-hydro-climatic interactions.



**Figure 1. Paleo-shorelines and paleo-sea level.** **a)** Global compilation of paleoshoreline sequences, adjusted from Pedoja et al., 2014, **b)** pictures of marine terrace sequences from Santa Cruz (US) and the Corinth Rift (Greece) and **c)** paleo-sea-level estimates for the past 450 ka, showing a sea-level curve (SpLi16, blue; Spratt and Lisiecki, 2016) derived from principal component analysis of 7 sea-level curves with its 2.5% and 97.5% likelihood range (grey envelope), an approximation of that curve with nodes and a cubic spline interpolation (red), global mean sea-level highstand estimates adjusted for glacio-istostatic adjustments (GMSL, brown dots; Kopp et al., 2009; Dutton et al., 2015; Pico et al., 2016; Creveling et al., 2017; Dyer et al., 2021; Tawil-Morsink et al., 2022), selected relative sea-level highstand estimates >130 ka (RSL, pink dots; Stirling et al., 2001; Murray-Wallace, 2002; Andersen et al., 2010; de Gelder et al., 2022; Marra et al., 2023), global mean sea-level lowstand estimates from ice sheet data (orange dots; Batchelor et al., 2019), and red boxes that represent the likely admissible range of relative sea-level elevations at locations far from the major ice-sheets (details in Supplementary Information and Table S1) that we consider in this study. Marine Isotope Stages (MIS) are given in bold, and based on Railsback et al., 2015.

## 2 Marine terrace sequence inversion

Marine terraces are relatively flat surfaces of coastal origin, either horizontal or gently inclined seawards (Fig 1b; Pirazzoli, 2005). They are bounded inland by a fossil sea-cliff, and can be covered by a layer of coastal sediments. Here we model erosive marine terraces, which are primarily formed by sea-cliff retreat in response to wave action. The superposition of Quaternary sea-level variations (Fig. 1c) and vertical land movement typically leads to a staircase landscape exhibiting marine terraces sequences (Fig. 1b).

The landscape evolution model we use (the code 'REEF'; Husson et al., 2018; Pastier et al., 2019) has a wave erosion module based on the wave energy dissipation model developed by Anderson et al. (1999). The model assumes that the vertical seabed erosion rate is a linear function of the rate of wave energy dissipation against the seabed (Sunamura, 1992). Horizontal erosion rates depend on the energy available at the sea-cliff after dissipation of the far-field wave energy (Anderson et al., 1999). The dissipation rate is dictated by the water depth profile, which increases landwards exponentially with decreasing water depth. The initial erosion rate  $ER$  is expressed as an effective eroded volume per unit of time and coastal length, in which a fraction of erosional residual power  $ER_r$  erodes the foundation at each

location along the profile. This fraction  $ER_r$  depends on the water level,  $h$ , the water depth for wave base erosion  $WB$ , and a coefficient for sea bed erodibility  $K$ , so that:

$$\frac{dER_r}{dt} = K \times ER_r \times \exp\left(-\frac{h}{WB}\right) \quad (1)$$

Then a residual power ( $ER - dER_r/dt$ ) carves out a 1 m high notch and all its overhanging material to form a cliff. Following previous studies (Anderson et al., 1999; Pastier et al., 2019), for  $K$  we use 0.1 as bedrock erodibility and 1 for notch carving. Finally, the 2D model we use consists of a landmass with a seaward dipping linear initial slope ( $IS$ ; Fig. 2a), an initial erosion rate ( $ER$ ; Fig. 2a) that evolves as platforms are being carved, a wave base depth ( $WB$ ; Fig. 2a) that determines the vertical range over which erosion takes place, a land uplift rate ( $UR$ ; Fig. 2a) and a sea-level history.


To invert the morphology of the marine terrace sequences, we parameterize sea-level history with a finite number of unknown parameters. We use nodes interpolated through a cubic spline scheme (Fig. 2b; light blue). This creates sea-level curves with similar characteristics to published sea-level curves (red line, Fig. 1c), in which the nodes represent sea-level minima (lowstands) and maxima (highstands) that are typically linked to even and odd-numbered marine isotope stages (MIS), respectively.

Reconstructing the sea-level history from present day observations of marine terrace sequences can be mathematically formulated as a highly non-linear inverse problem where the solution is non-unique. To embrace this non-uniqueness, the problem can be cast in a Bayesian (probabilistic) framework where the solution is a posterior probability distribution describing the probability of the model parameters (here the past sea-level variations), given the observed data (here the geometry of marine terraces). We use a Markov chain Monte Carlo algorithm to sample the posterior distribution and explore the range of models that can explain the observed topography within errors. In the Monte Carlo exploration of the model space, nodes can either be fixed at certain ages and elevations, or left free to move within a prescribed range (e.g. red boxes in Fig. 2b,d,e). The 4 main erosion model parameters ( $IS$ ,  $ER$ ,  $WB$ ,  $UR$ ) can also be fixed to chosen values, as done in the synthetic tests below, or left free within chosen ranges, as done for the Santa Cruz and Corinth examples below. The algorithm samples the parameter space as a random walk, where at each iteration a new sea level history model is proposed by perturbing the current one. The proposed model is then either accepted or rejected following an acceptance rule based on the level of data fit of the current and proposed models. The final solution is a large ensemble of paleo sea-level models that approximates the probabilistic solution. That is, the distribution of models follows the posterior probability solution. Of the 1 million model runs per figure, the first 50 000 models were discarded as burn-in models. To verify that the random walk samples the target distribution, we show a number of convergence diagnostics in Fig. S1, which includes parameter acceptance ratios and likelihood evolution for all model runs in this article. For a review of Bayesian inference and Monte Carlo methods in the geosciences, we refer the reader to Mosegaard and Sambridge (2002), and Gallagher et al. (2009).

One benefit of Bayesian inference is the ability to propagate uncertainty estimates from the observed measurements towards the unknown model parameters. For that, a likelihood probability distribution needs to be defined, based on a misfit function and on uncertainty estimates associated with observations. In this work, the data vector is defined as a set of points measured on the shoreline with a vertical step size ( $ipstep$ ; Fig. 2a). The misfit between this observed topography and the modelled paleo-shoreline sequences is calculated on the horizontal axis. The elevation axis is therefore divided in a regular grid, and the level of data fit is measured by comparing the horizontal distance between observed and simulated horizontal position at each grid point. In this way, the terrace width is used as an observed parameter. Uncertainties about the observed shorelines account for the inability of our numerical model to explain observations, and can be due to both observational or modelling errors. These errors are supposed Gaussian and described by a standard deviation ( $\sigma$ ; Fig. 2a) and the level of spatial correlation related to the spatial resolution of the data ( $corrl$ ; Fig 2a). In preliminary tests (Fig. S2) we also tested inversions with the misfit calculated along vertical axes (i.e. comparing elevations between observed and estimated profiles), but found it harder to reproduce realistic marine terrace sequences with a few m of terrace height variability, and a few hundred meter of terrace width variability (e.g. Regard et al., 2017; De Gelder et al., 2020). We note that changing the misfit calculation from horizontal to vertical misfit does not change the paleo sea-level posterior distribution (Fig. S2). The assigned ranges



for input parameters are all treated as flat uniform priors, i.e. a prior distribution that assigns equal probability to all the values within the specified ranges.



**Figure 2. Inversion of synthetic marine terrace profiles.** **a)** Synthetic topography (Obs, blue) created from a forward model with known input parameters: IS = Initial Slope, ER = Erosion Rate, UR = Uplift Rate and WB = Wave Base. The posterior range of models that fit the observed topography with the given  $\sigma$ ,  $ipstep$  and  $corr1$  (see text) is represented by the median (orange), and the 2.5 and 97.5 percentiles of the inverted models (grey envelope). **b)** Posterior probability distribution for the sea-level histories. Each individual paleo sea-level history is described with 6 sea-level nodes linked with a cubic spline interpolation, of which the nodes at 78, 6 and 0 ka are fixed in time and elevation, and the other three nodes can move within the three red boxes. The input (target) sea-level history is given in light blue, and the probabilistic solution is depicted by the median (solid orange line) and the 2.5 and 97.5 percentiles (dashed orange lines). **c)** Same as **a**, but with different uplift rates and sea-level histories. **d)** Sea-level histories for the inverted profiles in **c**, similar to **b** but with a different input sea-level history including more nodes. **e)** Similar to **d**, but inverting the two profiles simultaneously to find a common sea-level curve explaining both profiles. MIS are marked in white.

### 3 Synthetic marine terrace profiles

To test and illustrate the potential of the inversion approach, we inverted synthetic topographic profiles that were produced by forward models with known input parameters (Fig. 2). To start with a relatively short and simple sea-level range, we defined an 80 ka sea-level history consisting of 6 nodes (Fig. 2b; light blue). For the inversion, we fixed the nodes at 78, 6 and 0 ka, and the positions of the other three nodes were left as unknown model parameters to be recovered. In the Monte Carlo exploration of the model space, these three nodes were left free to move within a prescribed range (red boxes in Fig. 2b). All other erosion model parameters (IS, ER, WB, UR; Fig. 2a) were fixed

during the inversion at the values used to produce the observed topographic profile. The parameters  $\sigma$ , *ipstep* and *corrl* were set at 50, 1 and 10 m, respectively. We inverted the topographic profile between 0 and 60 m elevation by sampling the parameter space with 1 million forward simulations. The solution is a large ensemble of sea-level histories that reflect the probability of the paleo sea-level, given the synthetic coastline topography.

The resulting profiles show an MIS 3 terrace at an elevation range of ~15-30 m (Fig. 2a), whereas an MIS 1 terrace lies below the present-day sea level, and is thus not considered in the inversion. As such, the range of sea-level histories that could have created the MIS 3 terrace is narrower than for the MIS 1 terrace (Fig. 2b). This range is particularly limited for the period of sea-level rise leading up to the MIS 3 peak, suggesting that uplifted marine terraces are more likely to form during periods of relative sea-level rise. This is theoretically expected, as erosion scales with the total duration of sea-level occupation (Malatesta et al., 2021), and simultaneous sea-level rise and land uplift implies favorable conditions for the formation of marine terraces. Another notable feature is the distribution of possible sea-level histories along a diagonal line that corresponds to the uplift rate. This line would reach the maximum terrace elevation when extrapolated to  $t=0$  ka, in line with classic graphical methods (Bloom and Yonekura, 1990). Although the MIS 1 terrace is not inverted, there are some limitations to the magnitude and rate of sea-level rise between MIS 2 and MIS 1 (Fig. 2b), probably because this period determines how much of the MIS 3 terrace is eroded at its distal edge.

For the inversion of every individual profile there should be a trade-off between younger, higher sea-level peaks and older, lower sea-level peaks in line with the fixed uplift rate (as in Fig. 2b). These trade-off effects can be overcome through the joint inversion of multiple profiles with different uplift rates, reducing the uncertainty in sea-level reconstructions. To show this, we also inverted two different topographic profiles produced with different fixed uplift rates but with the same sea-level history over a 135 ka timescale (the last glacial-interglacial cycle; Fig. 2c-e). When the two profiles are inverted individually, the range of possible sea-level histories is relatively wide, and again the sea-level peaks would follow a diagonal line parallel to the uplift rate (Fig. 2c, d). However, if we jointly invert both profiles, i.e. assuming that a unique sea-level history would have created both marine terrace staircase morphologies, the probability distribution for past sea-level narrows, and the median sea-level of the inversion better approximates the input curve (Fig. 2e). The range is particularly narrow for the transgressions leading up to the MIS 5a and 5c highstands, for which the corresponding terraces are well developed in the topographic profiles (Fig. 2c). Similar to the MIS 1 terrace in Fig. 2a, the MIS 1 and 3 terraces in Fig. 2c would be located below sea level for the given parameters, and thus the possible sea-level range is wider for the transgressions leading up to MIS 1 and 3 (Fig. 2d). Also for these highstands though, the sea-level is better constrained for the joint inversion (Fig. 2e) than with the individual inversions (Fig. 2d). This suggests that jointly inverting more profiles would increase even further our ability to constrain sea-level histories. To understand what happens in scenarios in which more parameters are unknown, we repeated the same tests (from Figures 2b-d) with broad prior ranges for the uplift rate, initial slope, erosion rate and wave base height (Fig. S3). Also in this case the joint inversion provides much narrower posterior ranges for paleo sea-level compared to the two individual profile inversions, and not too different from the cases with fixed model parameters (Figs. 2b-d). The posterior ranges for all parameters are consistently smaller for the joint inversions, compared to the individual inversions (Fig. S3).

These synthetic tests imply that in natural examples, sea-level reconstruction should also benefit from the inversion of multiple marine terrace profiles if conditions change between those profiles. In this example we used two different uplift rates for the joint inversion, which lead to a range in different terrace sequence morphologies (Fig. 2c), but an approach where all parameters, including wave base, erosion rate or initial slope, are undefined a priori (or only within a given range), should lead to a more realistic range of possible sea-level histories.

To put this method to the test in real cases, we selected two well-documented yet contrasting cases, Santa Cruz and the Gulf of Corinth, each having their peculiarities that make them ideal to study the inversion of marine terraces. Santa Cruz is well documented and possible parametric windows are quite narrow, making it an ideal benchmark site for our method. On the other hand, the staircase sequence in the Gulf of Corinth, while equally well documented, is particularly relevant to take advantage of the predictive capacities of our method. There, we can decipher the complex interplay between vertical land motion and sea-/lake-level fluctuations in a semi-closed basin, and better reconstruct its hydrodynamic history where paleoclimatic data are insufficient.


#### 229 4 Santa Cruz marine terrace sequence inversion

230 The marine terraces along the Santa Cruz coastline (central California, US) formed through a combination of  
231 Quaternary sea-level oscillations and tectonic uplift by nearby active faults (e.g. Bradley, 1957; Anderson and  
232 Menking, 1994; Anderson et al., 1999; Perg et al., 2001; Matsumoto et al., 2022). We invert a topographic profile  
233 from Rosenbloom and Anderson (1994), who distinguished the original eroded bedrock surface, which we use, from  
234 its overlying colluvium for 5 marine terraces. We followed the age interpretation of Perg et al. (2001), suggesting  
235 these terraces were formed, from bottom to top, during MIS 1, 3, 5a, 5c and 5e. Unlike in the synthetic tests, here we  
236 left uplift rate, erosion rate, wave base depth and initial slope parameters free within a **prior** range of values. We use  
237 the elevation (~170 m) and age (~118-128 ka) of the upper terrace to derive a range of possible uplift rates (1.3-1.65  
238 mm/yr), and simultaneously consider ranges for initial slope (5-15%), wave base depth (1-10 m) and erosion rates  
239 (100-800 mm<sup>3</sup>/yr) in the terrace inversion. We use the same inversion parameters as for the synthetic tests, running 1  
240 million models over 450 ka with the sea-level high- and lowstands limited to the red boxes in Fig. 1 (See  
241 Supplementary Information and Table S1). We tested different inversion parameters (Fig. S4), and settled on values  
242 of 1 m *ipstep*, 100 m  $\sigma$  and 10 m *corrl*, as for these values the corresponding 95% posterior ranges for the variability  
243 in terrace height (~5 m; Fig. S4) and width (~150 m; Fig. S4), are along the same order of magnitude as the variability  
244 in a swath profile across the Santa Cruz terraces (Fig. S5). Higher values for  $\sigma$  and *corrl* result in both broader ranges  
245 for accepted terrace heights/widths, and consequently also in broader posterior ranges for paleo sea-level, although  
246 we note that the first order shape of the paleo sea-level curves remains similar irrespective of the tested *ipstep*,  $\sigma$  and  
247 *corrl* values (Fig. S4).

248 The sampled sea-level histories successfully reproduce the terrace morphology, as evidenced by the low misfit of 2 m  
249 (Fig. 3a). As with the synthetic tests (Fig. 2), periods of sea-level rise are better constrained than periods of sea-level  
250 fall, and highstands better constrained than lowstands (Fig. 3b). Also here there is a trade-off in sea-level peaks, in  
251 which younger, higher sea-level peaks could result in similar shaped marine terraces as older, lower sea-level peaks  
252 (e.g. for MIS 3c, Fig. 3e). The models limit the uplift rate to ~1.35-1.6 mm/yr, the initial slope to ~7-9.5%, the wave  
253 base depth to 4-10 m and the erosion rate to 200-800 mm/yr (Figs. 3c,d). Notably there is a positive correlation between  
254 wave base depth and erosion rate (Fig. 3d), suggesting a higher value for wave base depth would require a higher  
255 erosion rate to create the same marine terrace sequence morphology.

256 Compared to our proposed **prior** range of possible sea-level elevations for MIS 3 (-30 to -80 m; Fig. 3b), the **posterior**  
257 **distribution of the** inversion suggests paleo sea-level values on the higher end of that spectrum. This is in agreement  
258 with a growing number of studies suggesting oxygen-isotope derived sea-level curves underestimate sea-level for that  
259 period (Pico et al., 2016; Dalton et al., 2019, 2022; Gowan et al., 2021; De Gelder et al., 2022). For MIS 5a on the  
260 other hand, the **posterior distribution of the** inversion suggests a sea-level peak on the lower end of our proposed range  
261 of sea-level elevations (Fig. 3b). Although the highstand **posterior distributions** still span a broad elevation range of  
262 ~25 m, these results tend to align with studies proposing an overall decrease in sea-level between MIS 5e, 5c and 5a  
263 (e.g. Chappell and Shackleton, 1986; Schellmann and Radtke, 2004; Tawil-Morsink et al., 2022).

264 We also tested additional uplift rate scenarios (Fig. S6), given that there has been concerns on the terrace chronology  
265 that we adopted (Brown and Bourlès, 2002), and other studies have suggested the terrace at 27 m elevation might be  
266 formed during MIS 5a, 5c or 5e instead of MIS 3 (Bradley and Addicott, 1968; Lajoie et al., 1975; Kennedy et al.,  
267 1982; Weber et al., 1990). These uplift rates can fit the terrace sequence morphology equally well in terms of  
268 topographic misfit, but generally imply a larger possible range of paleo sea level. This can be explained by the  
269 increased terrace re-occupation for lower uplift rates (Malatesta et al., 2021), which also explains why the **posterior**  
270 ranges for the initial slope and wave base depth change increase for lower uplift rate scenarios (Fig. S6), and erosion  
271 rate estimates decrease. These tests suggest that locations with higher uplift rates will generally provide narrower  
272 constraints on paleo sea-level, while still providing realistic and unbiased parameter estimates.



**Figure 3. Inversion of NW-Santa Cruz marine terrace sequence.** a) Observed topography (from Rosenbloom and Anderson, 1994; Obs, blue) with the age interpretation of Perg et al. (2001) marked in bold, together with the modeled best-fit, median, 2.5% and 97.5% percentile profiles. b) Probabilistic sea-level reconstruction for the profiles in a, MIS in white c) **Posterior ranges** for uplift rates and initial slopes (histogram of the sampled models), d) **Posterior ranges** for wave base depths and erosion rates e) **Posterior ranges** for the MIS 3c peak, i.e. distribution for the position of the 50-57 ka node within the paleo sea level curve.

## 5 Gulf of Corinth marine terrace sequence inversion


The complexity of semi-isolated basins, connected to sea during some time intervals and isolated in others, make them ultimate testing grounds for our modelling approach. We focus on one of such basins, the SE Gulf of Corinth, to derive a sea-/lake-level history from terrace sequence geometries, and compare its outcomes to paleoclimate data, tectonic structures and sill dynamics.

Natural interaction between the Gulf of Corinth and the open sea is currently restricted by the Rion and Acheloos-Cape Pappas sills at its W entrance at ~45-60 m depth (Fig. 4; Beckers et al., 2016). In the past there was an additional connection at its E end along the Corinth Isthmus, currently uplifted at ~80 m elevation but consisting of Quaternary marine sediments (Fig. 4; Caterina et al., 2023). These sills have controlled the Gulf's connection with the open sea over the past few hundred thousand years and lead to an alternation of marine and (semi-)isolated lake environments within the Gulf (McNeill et al., 2019). Although we approximately know the timing of these alternations, it remains unclear whether sill depths remained stable or fluctuated throughout the Quaternary (Roberts et al., 2009; McNeill et al., 2019), and in addition, whether lake levels were stable or fluctuating during periods with no connection to the sea. Hydroclimatic conditions, i.e. the balance between inflow, outflow, precipitation and evaporation, determine the level of the lake, and state-of-the-art paleo-environmental reconstructions (e.g. Kafetzidou et al., 2023) are insufficient to infer its fluctuations over time.

Terrace sequences are well exposed in the SE of the Gulf, where the Gulf of Corinth Fault System (Fig. 4) has lead to differential coastal uplift rates (Armijo et al., 1996; De Gelder et al., 2019; Fernández-Blanco et al., 2019; Fig. S7). This peculiarity allows us to test on a natural example whether the joint inversion of multiple terrace sequence profiles with different uplift rates provides a better-constrained sea-/lake-level history (as in Fig. 2). To account for the



unknown range of possible lake-level elevations, we carried out inversions with all nodes from (semi-)isolated periods broadly constrained between -15 and -150 m elevation. We selected three topographic profiles with little river incision and ~0.4-1.4 mm/yr uplift rates (Fig. S7), and avoided modelling the broad coastal plains at the base of all profiles that appear to have been modified by human presence (Fig. S7). We used the 90% percentile of 100-m wide swath profiles to obtain representative terrace sequence morphologies (Fig. S7). For the three profiles we assigned prior ranges of possible uplift rates of 1.25-1.4, 0.7-0.9 and 0.4-0.55 mm/yr (De Gelder et al., 2019; Fig. S6), and broad prior ranges for erosion rate (100-1500 mm/yr), initial slope (1-20%) and wave base depth (1-12 m).




**Figure 4. Tectonomorphology of the Gulf of Corinth.** Map showing the main features of the Gulf of Corinth, including the active faults, marine terraces, profile locations used in the inversion and connections to the Ionian Sea (Rion Sill) and Aegean Sea (Corinth Isthmus). M0078 and M0079 indicate the IODP-381 sedimentary coring locations (McNeill et al., 2019). Psat = Psathopyrgos Fault, Pisia = Pisia Fault, Kala = Kalamaki Fault. Modified from de Gelder et al. (2019) and Fernández-Blanco et al. (2020).

The individual profile inversions mostly constrain paleo sea/lake level for profile 1 (Fig. 5a), because it has the highest uplift rate and contains most terraces. The other two profiles provide limited constraints on paleo sea/lake level when inverted individually (Fig. 5b/c), but when jointly inverted with profile 1 they provide a much narrower range in terms of posterior distribution for sea-level (Fig. 5d). The cumulative misfit for the individual inversions (28 m) is slightly better than for the joint inversion (46 m), but there are no major visible differences between the terrace sequence profiles for the two inversions, and apart from the highest terrace of profile 2 (Fig. 5b) the terrace sequences are all near perfectly reconstructed. The three profiles show variations in initial slopes that are in line with the overall morphology, i.e. present-day profile 1 is steeper than 2, which is steeper than 3, which is also what we find for their initial slopes. The three profiles do have similar posterior distributions for wave base depths and erosion rates (Fig. 5e). Although we might expect lateral differences in these rates given variability in sediment types, catchment area and coastal orientation, the broad ranges for the posterior distributions indicate we cannot quantify these lateral differences from the profile morphology alone. The posterior parameter ranges mostly remain the same between the individual and joint inversion, with exception of the uplift rates for profiles 2 and 3 that became a little lower for the joint inversion. As for the sea-/lake-level inversion, all the other posterior parameter ranges become narrower for the joint inversion (Fig. 5e).

The inverted sea-/lake-level history (Fig. 5d) shows a few notable features. To a first order, fluctuations resemble global sea-level trends, with relatively fast periods of sea-/lake-level rise prior to major sea-level highstands, followed

329 by long periods of slow sea-/lake-level fall (Fig. 1c). Yet, unlike global sea-level trends, there are several periods of  
330 prolonged stability, in particular around 180-200 ka and 275-300 ka, and possibly also around 75-95 ka and 360-370  
331 ka. In addition, glacial periods are often surprisingly poorly resolved, like during the period 20-75 ka or 160-180 ka.  
332 In the last section we discuss our interpretation of these trends, and show how they provide insightful arguments to  
333 decipher the relation between water level, fault activity, paleoclimate and tectonics.



**Figure 5. Inversion of SE Corinth Rift marine terrace sequence.** a-c) Observed topography (left) from 3 different profiles in the SE Corinth Rift (locations see Fig. S7), together with the modeled best-fit, median, 2.5% and 97.5% percentile profiles for both an individual profile inversion and a joint inversion of the three profiles (horizontally offset by an arbitrary value). Corresponding probabilistic sea/lake-level ranges for the individually inverted profiles are given on the right, d) Posterior distribution for sea/lake-level history from joint inversion, with red boxes describing prior ranges e) Prior and posterior parameter ranges from both individual and joint profile inversions

## 6 Discussion

### 6.1 Beyond sea level: tectono-hydro-climatic processes in the Gulf of Corinth

While the results for the joint inversion of the 3 profiles in the Gulf of Corinth permit to unravel sea- and lake-levels through time as expected, we also stress that such inversions yield more, when integrated within the broader environmental setting: these reconstructions allow for a more detailed look into sea- and lake-level fluctuations within a (semi-)isolated basin. Figure 6 compares our inverted sea-lake level to the stratigraphy, facies and pollen content within two sedimentary cores from the sea floor of the central basin (McNeil et al., 2019; Gawthorpe et al., 2022; Kafetzidou et al., 2023). Based on those combined datasets, we propose that the main hydroclimatic modes that have occurred in the Gulf of Corinth throughout the past 450 ka, are 1) marine Gulf of Corinth, 2) transitional Gulf of Corinth or overfilled Lake Corinth and 3) underfilled Lake Corinth. The first 2 of those have been proposed before based on sedimentary cores (McNeill et al., 2019; Gawthorpe et al., 2022), whereas we base the occurrence of intervals with an underfilled Lake Corinth on our marine/lake terrace inversion, as available paleoclimatic proxies are unable to uncover them.

The major peaks in our reconstructed sea/lake-level curve occurred during interglacial sea-level highstands, when sea level in the Gulf of Corinth was similar to eustatic sea level (marine mode M). Sedimentary cores indicate marine conditions (McNeill et al., 2019), the corresponding stratal packages are bioturbated, and associated sedimentary facies are types FA1 and FA6 (Fig. 6; see caption for facies description). From pollen records, the typical reconstructed biomes are cool mixed evergreen needleleaf and deciduous broadleaf forests, indicating relatively warm and wet conditions with low amounts of steppic taxa (Kafetzidou et al., 2023; Fig. 6).

We interpret the interstadial periods around 75-95 ka, 180-200 ka, 275-300 ka and 360-370 ka as periods with an overfilled Lake Corinth, possibly with some marine incursions indicating a transitional Gulf of Corinth (T/O mode). This would explain the prolonged sea-/lake-level stability, during interstadial periods when eustatic sea-level fluctuated by tens of meters (e.g. Spratt and Lisiecki, 2016; De Gelder et al., 2022). In that case, sea-/lake-level elevations would correspond to the paleo-sill depth of the Rion Sill and/or Corinth Isthmus (white line, Fig. 6). Within the sedimentary cores, these periods are mostly characterized by laminated stratal packages, and associated sedimentary facies are types FA2, FA3 and FA4 (Fig. 6; see caption for facies description). The occurrence of marine incursions into Lake Corinth during these interstadial periods is suggested by dated corals of ~76 ka, ~178 ka and ~201 ka (Roberts et al., 2009; Houghton et al., 2003) as well as the white, aragonite-rich laminations of FA3 and FA4. In other locations such laminations have been linked to (seasonal) mixing of marine and non-marine surface waters (Sondi and Juracic, 2010; Roeser et al., 2016).

The glacial periods are characterized by relatively low sea/lake-level elevations, possibly even down to the lower limit of -150 m we used in the inversion. We interpret these periods as underfilled Lake Corinth conditions (U mode), during which water inflow was lower than water evaporation within the lake, and lake level fell down to tens of meters below the sill depth. Sedimentary cores indicate non-marine conditions (McNeill et al., 2019), the corresponding stratal packages are mostly bedded and associated sedimentary facies are types FA5 and FA11 (Fig. 6; see caption for facies description). Reconstructed biomes from pollen suggest an increase in open vegetation such as grassland and steppe communities under colder and drier conditions (Kafetzidou et al., 2023; Fig. 6), matching reconstructed periods of lake underfilling. In general the inverted resolution of the lake-level elevation is much lower for these periods, with large probabilistic ranges. We attribute this to the occurrence of rapid lake-level fluctuations, like in other isolated E-Mediterranean water bodies such as the Dead Sea (Stein et al., 2010) and Lake Van (Turkey; Landmann et al., 1996). In such environments, the lake level is determined by the budget between runoff and evaporation, and quick variations




are expected. Alternatively, this could also be due to the fact that terraces formed during low sea/lake level get increasingly eroded during transgressions.

The interstadial periods with prolonged sea-/lake-level stability also allow for a possible reconstruction of sill depths through time (white lines, Fig. 6). The westernmost sill, the Acheloos-Cape Pappas Sill (Fig. 4), is currently at a depth of ~45-48 m. While there are no major faults there, we can't exclude slow subsidence or uplift at a few tenths of mm/yr, cumulating to a few tens of meters on the 100 kyr time scale. The Rion Sill, at the western entrance to the Gulf of Corinth, is currently at ~62 m depth (Perissoratis et al., 2000). As it is located in the hanging wall of the Psathopyrgos Fault (Fig. 5), active since at least the past ~200 ka (Houghton et al., 2003), the Rion Sill was unlikely deeper in the past. We reconstruct the Rion Sill depth assuming marine incursions around ~76 ka (Roberts et al., 2009) took place through this sill, and the Rion Sill was not lower than sea/lake level during the overfilled/transitional interval around ~200 ka. Extrapolating the trend, it would make sense for the older connections between Lake/Gulf of Corinth and the open sea to have occurred primarily through the Corinth Isthmus at the eastern end of the Gulf of Corinth.

The Corinth Isthmus is currently at an elevation of ~80 m, and has been uplifted through the Pisira Fault, Kalamaki Fault and/or a regional uplift (Armijo et al., 1996; Roberts et al., 2009; Caterina et al., 2022). We reconstruct the Corinth Isthmus depth assuming lake/sea level during overfilled/transitional intervals around ~290 and ~360 ka correspond to the Corinth Isthmus depth, and the isthmus was not lower than sea/lake level during the overfilled/transitional interval around ~200 ka. Extrapolating this trend fits with the current Corinth Isthmus elevation of ~80 m. The isthmus elevation before ~360 ka is difficult to constrain from our data, but was possibly shallower before, given the small amount of marine sediments deposited around the ~400 ka interglacial period, and the lack of deposits within the Isthmus stratigraphy older than ~350 ka (Collier and Dart, 1991; Caterina et al., 2022). Our reconstruction of both Rion Sill and Corinth Isthmus fits with the sedimentary interpretation of a tidal strait around ~300 ka at the isthmus, with marine connections on both ends of the Gulf of Corinth (Caterina et al., 2022).

Our exploration of sea-/lake-level variations in the Gulf of Corinth demonstrates the strength of using marine terrace sequence inversion. Although several questions remain – like the effects of erosion and sedimentation on sill evolution, or the effects of non-constant uplift rates of the marine terrace sequences – we are able to provide a solid framework that can explain several different tectonic and hydro-climatic processes simultaneously. We distinguish 4 different hydroclimatic modes for the Lake/Gulf of Corinth, that have probably also occurred in other (semi-)isolated basins like the Sea of Marmara or the Carioco Basin. Marine and transitional modes will most likely depend on eustatic sea-level elevations and sill depths, whereas over- or underfilled lakes likely depend on sill depths and local climatic conditions. For the Corinth Lake we show that this transition from over- to underfilled lakes occurs during changes from interstadial to glacial periods, and is accompanied by changes in vegetation that imply drier conditions.



**Figure 6. Comparison of sea-/lake-level reconstruction to other datasets.** **a)** Comparison to IODP-381 cores M0078 and M0079, with steppic taxa from Kafetzidou et al. (2023), facies from McNeill et al. (2019) and stratal packages with U/Th ages from Gawthorpe et al. (2022). FA1: homogenous mud, FA2: greenish gray mud with dark gray to black silty-to-sandy beds (cm-scale), FA3: light gray to white sub-mm laminations (cc or aragonite) alternating with mud-silt beds, FA4: laminated greenish gray to gray mud with muddy beds, FA5: greenish gray mud with homogeneous cm thick gray mud beds, FA6: green bedded partly bioturbated mud, silt and sand, FA11: interbedded mud/silt and cm thick sand beds. **b)** Inversion result from Fig. 5d with proposed sill/isthmus elevations, marking periods with marine (M), transitional or overfilled (T/O) and underfilled (U) hydroclimatic modes. **c)** Schematic illustrations of the different hydroclimatic modes in the Lake/Gulf of Corinth.

## 6.2 Inversion of marine terrace sequences

In the examples above, we showed how to assess paleo sea-level variations, and simultaneously extract quantified metrics for morphotectonics and hydrodynamics, from the geometry of marine terrace sequences. Using a probabilistic inversion methodology set in a Bayesian framework, we avoid the simplifications of bijective approaches in which a single marine terrace is always linked to a single sea-level highstand and vice-versa (e.g. Pastier et al., 2019; Malatesta et al., 2021). By considering a full sea-level curve and its possible variability, it is possible to provide quantitative constraints on highstands, lowstands, sea-level rise and fall, filling the observational gap for time periods for which field measurements are scarce. We admit that some model simplifications and approximations may alter our

interpretations. In particular, we neglect subaerial erosion, and kept uplift rate, erosion rate, initial slope and wave base depth parameters time-constant for each individual sampled paleo sea level curve. Both could be fine-tuned in future developments.

To apply this model procedure to other sites, we can make several recommendations based on our findings in this paper. Any coastal area with marine terraces will have lateral variability in terms of terrace width and terrace elevation. Given that the posterior ranges for the model parameters (paleo sea-level, uplift rate, etc.) will directly depend on the chosen inversion parameters (*ipstep*,  $\sigma$  and *corr1*), it is important to adjust those inversion parameters to the naturally observed variability in the geometry of a marine terrace sequence. In the Santa Cruz example the similarity in terrace width/height variability between the modeled posterior ranges (Fig. S4), and the observed variability within a section of the marine terrace sequence (Fig. S5) justifies our choice of inversion parameters. But, if we wanted to characterize a larger area along the Santa Cruz coastline, we would either have to increase  $\sigma$  and/or *corr1* to account for the higher degree of variability, or jointly invert multiple cross-sections in which every cross-section has a similar variability in terrace height/width.

The prior information required to obtain reliable results, will also be dependent on the context of a specific marine terrace sequence, and on the parameter cut-off choices deemed realistic. For Santa Cruz and Corinth, the posterior distributions for wave base depth suggest that, purely based on marine terrace sequence morphology, this cut-off could have been deeper than 10-12 m (Figs. 3 and 5). However, based on observations and models of cliff erosion it seems unlikely that the wave base for bedrock erosion is more than 10 m for Santa Cruz (Kline et al., 2014), and wave base depths/heights should be smaller for the calmer Gulf of Corinth. This choice in cut-off in turn affects the posterior distribution of other parameters, and should thus be chosen carefully.

In the cases of Corinth and Santa Cruz we assigned relatively narrow windows for the uplift rate, based on chronological information that was already available. As such we only obtained refinements, rather than ‘new’ information about the uplift rate. For the Santa Cruz case, we tested four uplift rate scenarios matching different possible chrono-stratigraphies (Fig. S6) that could all explain the morphology equally well, albeit with different ranges for the other parameters. We also inverted the Santa Cruz terrace sequence morphology with a broad range between 0.3 and 1.5 mm/yr, but the inversion would converge on only one of the four scenarios in Fig. S6. This suggests that at least an approximate prior idea on the uplift rate is a prerequisite for a reliable inversion of a marine terrace sequence morphology.

Many paleo sea-level studies that use geomorphic/geologic observations tend to have a confirmation bias regarding sea-level curves, and propose refinements of paleo sea-level estimates to sub-m scale (e.g. Murray-Wallace, 2002; Roberts et al., 2012) or uplift rates to precisions of  $\sim 0.01$  mm/yr (e.g. Pedoja et al., 2018; Meschis et al., 2022). In this study, we take a step back by allowing more freedom to possible paleo sea-level variations, as well as uplift rate, erosion rate, initial slope and wave base depth, to provide a more reliable way to translate morphologic observations to paleo sea-level constraints. For instance, the low uplift rate examples from the Corinth Rift (Fig. 5b, c) and Santa Cruz (Fig. S6) reveal very little about paleo sea/lake levels, even if the uplift rate is roughly known. As a marine terrace is formed over several sea-level cycles, the resulting terrace width and height will depend on all those cycles, as well as wave base depths, erosion rates and initial slopes, all of which are generally poorly constrained. Even in the hypothetical case that these parameters are known (Fig. 2), there is still a wide spectrum of sea-level histories that could have created the specific morphology of a marine terrace sequence. It suggests that estimating paleo sea-level based on the comparison of a present-day landform to a paleo-landform (Rovere et al., 2016), may be too simplistic in many cases, at least for erosive marine terraces. Although uncertainties that we provide on paleo sea-level are much larger than what calculations based on hydrodynamic ranges would suggest (Lorchsteid and Rovere, 2019), we do consider them to be reliable as they take in a large number of unknowns.

Although here we focused on erosive marine terraces to develop a proof of concept, a promising avenue is to apply this inversion method to bio-constructed (coral reef) terraces, which tend to be better dated (e.g. Pedoja et al., 2014; Hibbert et al., 2016) and for which modelling routines also exist (e.g. Toomey et al., 2013; Pastier et al., 2019). One of our key findings is that inverting multiple profiles simultaneously provides much better paleo sea-level constraints than focusing on individual profiles (Figs. 2, 5). The global archive of paleo-shorelines (Fig. 1a) presents a huge potential for such multi-profile marine terrace inversions. This massive inversion would not only lead to improved

estimates of local relative sea-level histories, but may also complement studies on glacio-isostatic adjustments that are relevant to a global sea-level perspective.

## 7 Conclusions

In this study we demonstrated the use of a probabilistic inversion approach to decipher the formation of marine terrace sequences in general, and the tectonic and hydro-climatic evolution of (semi-)isolated basins in particular. With this approach, we provide the tools (see below) to simultaneously estimate past sea-level variations, uplift rates, erosion rates, initial slopes and wave heights.

From synthetic tests, benchmarking on a terrace sequence near Santa Cruz marine terrace sequence, and application to the sequence in the SE Gulf of Corinth, our results bring a theoretical advance by showing that : 1) Paleo sea-level and other parameter ranges can be better constrained from sequences that are uplifting at higher rates compared to lower rates, and better constrained from a joint inversion of multiple profiles than from inversion of a single profile. 2) Uplift rates, sea-level variations and wave erosion parameters are intricately linked. By allowing more freedom to possible ranges of all the relevant parameters, we provide a more reliable way to translate morphologic observations to paleo sea-level constraints. Resulting uncertainties may be higher compared to ‘classic’ approaches of comparing present to past shoreline elevations, but are more realistic. 3) Probabilistic inversion of marine terrace sequences is a powerful method, applicable to a large portion of the world’s coastlines to disentangle tectonic and hydro-climatic processes.

**Beyond the methodological achievement,** by applying our method to a complex case -the semi-isolated Gulf of Corinth (Greece), we show that this method can be a powerful tool to explore subtle environmental forcings, like the balance between precipitation and evaporation, that may have had a prime importance in setting the lake level during certain periods of time. We found that eustatic sea-level and tectonically changing sill depths drive marine and transitional phases during interglacial and interstadial periods, respectively. Wetter and drier conditions drive over- and underfilling of Lake Corinth during interstadial and glacial periods, respectively. We expect such transitions to be different for each unique tectono-hydro-climatic setting, with our inversion approach providing a new way to decipher such geomorphic Rosetta stones.

## Code availability

The marine terrace inversion code used in this study can be found at <https://github.com/ginodegelder/Rosetta>.

## Supplement link

The supplementary material to this study can be found at <https://doi.org/10.31223/X5B117>.

## Author contribution

**Conceptualization** by GdG, LH and TB, **Formal analysis** by GdG, **Funding acquisition** by TB, **Investigation** by GdG, NH, LH and TB, **Methodology** by GdG, NH, AMP, TB and YB, **Visualization** by GdG, NH and YB, **Writing - original draft** by GdG, **Writing - review & editing** by all authors.

## Competing interests

The authors declare that they have no conflict of interest.



517 **Acknowledgements**

518 This study has been funded by the European Union horizon 2020 research and innovation program under grant  
519 agreement 716542. Furthermore, GdG acknowledges postdoctoral funding from the IRD and the Manajemen Talenta  
520 BRIN fellowship program, and research permit 52/SIP.EXT/IV/FR/5/2023 provided by the Indonesian government  
521 on May 11<sup>th</sup> 2023 . GdG likes to thank Katerina Kouli for sharing her pollen data, and other IODP-381 expedition  
522 members, as well as David Fernández-Blanco, Robin Lacassin and Rolando Armijo, for the many fruitful discussions  
523 on the Corinth Rift.

524 **References**

- 525 Aksu, A. E., Hiscott, R. N., & Dogan, Y. (1999). Oscillating Quaternary water levels of the Marmara Sea and vigorous  
526 outflow into the Aegean Sea from the Marmara Sea–Black Sea drainage corridor. *Marine Geology*, 153(1-4), 275-  
527 302.
- 528 Andersen, M. B., Stirling, C. H., Potter, E. K., Halliday, A. N., Blake, S. G., McCulloch, M. T., ... & O’Leary, M. J.  
529 (2010). The timing of sea-level high-stands during Marine Isotope Stages 7.5 and 9: Constraints from the uranium-  
530 series dating of fossil corals from Henderson Island. *Geochimica et Cosmochimica Acta*, 74(12), 3598-3620.
- 531 Anderson, R.S., Densmore, A.L., Ellis, M.A., 1999. The generation and degradation of marine terraces. *Basin Res.* 11  
532 (1), 7e19.
- 533 Anderson, R. S., & Menking, K. M. (1994). The Quaternary marine terraces of Santa Cruz, California: Evidence for  
534 coseismic uplift on two faults. *Geological Society of America Bulletin*, 106(5), 649-664.
- 535 Armijo, R., Meyer, B. G. C. P., King, G. C. P., Rigo, A., & Papanastassiou, D. (1996). Quaternary evolution of the  
536 Corinth Rift and its implications for the Late Cenozoic evolution of the Aegean. *Geophysical Journal International*,  
537 126(1), 11-53.
- 538 Austermann, J., Mitrovica, J. X., Huybers, P., & Rovere, A. (2017). Detection of a dynamic topography signal in last  
539 interglacial sea-level records. *Science Advances*, 3(7), e1700457.
- 540 Batchelor, C. L., Margold, M., Krapp, M., Murton, D. K., Dalton, A. S., Gibbard, P. L., ... & Manica, A. (2019). The  
541 configuration of Northern Hemisphere ice sheets through the Quaternary. *Nature communications*, 10(1), 3713.
- 542 Beckers, A., Beck, C., Hubert-Ferrari, A., Tripsanas, E., Crouzet, C., Sakellariou, D., ... & De Batist, M. (2016).  
543 Influence of bottom currents on the sedimentary processes at the western tip of the Gulf of Corinth, Greece. *Marine*  
544 *Geology*, 378, 312-332.
- 545 Bernard, P., Lyon-Caen, H., Briole, P., Deschamps, A., Boudin, F., Makropoulos, K., ... & Linde, A. (2006).  
546 Seismicity, deformation and seismic hazard in the western rift of Corinth: New insights from the Corinth Rift  
547 Laboratory (CRL). *Tectonophysics*, 426(1-2), 7-30.
- 548 Bloom, A. L., & Yonekura, N. (1990). Graphic analysis of dislocated Quaternary shorelines. *Sea-level change*, 104-  
549 115.
- 550 Bradley, W. C. (1957). Origin of marine-terrace deposits in the Santa Cruz area, California. *Geological Society of*  
551 *America Bulletin*, 68(4), 421-444.
- 552 Bradley, W. C., & Addicott, W. O. (1968). Age of first marine terrace near Santa Cruz, California. *Geological Society*  
553 *of America Bulletin*, 79(9), 1203-1210.
- 554 Brown, E. T., & Bourles, D. L. (2002). Use of a new <sup>10</sup>Be and <sup>26</sup>Al inventory method to date marine terraces, Santa  
555 Cruz, California, USA: Comment and Reply: COMMENT. *Geology*, 30(12), 1147-1148.
- 556 Caterina, B., Rubi, R., & Hubert-Ferrari, A. (2023). Stratigraphic architecture, sedimentology and structure of the  
557 Middle Pleistocene Corinth Canal (Greece). *Geological Society, London, Special Publications*, 523(1), 279-304.
- 558 Chappell, J., & Shackleton, N. (1986). Oxygen isotopes and sea level. *Nature*, 324(6093), 137-140.
- 559 Chauveau, D., Pastier, A.-M., de Gelder, G., Husson, L., Authemayou, C., Pedoja, K. et al. (2024) Unravelling the  
560 morphogenesis of coastal terraces at Cape Laundi (Sumba Island, Indonesia): Insights from numerical models. *Earth*  
561 *Surface Processes and Landforms*, 49(2), 549–566.

Collier, R. L., & Dart, C. J. (1991). Neogene to Quaternary rifting, sedimentation and uplift in the Corinth Basin, Greece. *Journal of the Geological Society*, 148(6), 1049-1065.

Collier, R. E., Leeder, M. R., Trout, M., Ferentinos, G., Lyberis, E., & Papatheodorou, G. (2000). High sediment yields and cool, wet winters: Test of last glacial paleoclimates in the northern Mediterranean. *Geology*, 28(11), 999-1002.

Creveling, J. R., Mitrovica, J. X., Clark, P. U., Waelbroeck, C., & Pico, T. (2017). Predicted bounds on peak global mean sea level during marine isotope stages 5a and 5c. *Quaternary Science Reviews*, 163, 193-208.

Dalton, A. S., Finkelstein, S. A., Forman, S. L., Barnett, P. J., Pico, T., & Mitrovica, J. X. (2019). Was the Laurentide Ice Sheet significantly reduced during marine isotope stage 3?. *Geology*, 47(2), 111-114.

Dalton, A. S., Pico, T., Gowan, E. J., Clague, J. J., Forman, S. L., McMartin, I., ... & Helmens, K. F. (2022). The marine  $\delta^{18}\text{O}$  record overestimates continental ice volume during Marine Isotope Stage 3. *Global and Planetary Change*, 212, 103814.

De Gelder, G., Fernández-Blanco, D., Melnick, D., Duclaux, G., Bell, R. E., Jara-Muñoz, J., ... & Lacassin, R. (2019). Lithospheric flexure and rheology determined by climate cycle markers in the Corinth Rift. *Scientific Reports*, 9(1), 4260.

De Gelder, G., Jara-Munoz, J., Melnick, D., Fernández-Blanco, D., Rouby, H., Pedoja, K., ... & Lacassin, R. (2020). How do sea-level curves influence modeled marine terrace sequences?. *Quaternary Science Reviews*, 229, 106132.

De Gelder, G., Husson, L., Pastier, A. M., Fernández-Blanco, D., Pico, T., Chauveau, D., ... & Pedoja, K. (2022). High interstadial sea levels over the past 420ka from the Huon Peninsula, Papua New Guinea. *Communications Earth & Environment*, 3(1), 256.

De Gelder, G., Solihuddin, T., Utami, D. A., Hendrizan, M., Rachmayani, R., Chauveau, D., ... & Cahyarini, S. Y. (2023). Geodynamic control on Pleistocene coral reef development: insights from northwest Sumba Island (Indonesia). *Earth Surface Processes and Landforms*, 48(13), 2536-2553.

Derricourt, R. (2005). Getting “Out of Africa”: sea crossings, land crossings and culture in the hominin migrations. *Journal of world prehistory*, 19, 119-132.

Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., ... & Raymo, M. E. (2015). Sea-level rise due to polar ice-sheet mass loss during past warm periods. *science*, 349(6244), aaa4019.

Dyer, B., Austermann, J., D’Andrea, W. J., Creel, R. C., Sandstrom, M. R., Cashman, M., ... & Raymo, M. E. (2021). Sea-level trends across The Bahamas constrain peak last interglacial ice melt. *Proceedings of the National Academy of Sciences*, 118(33), e2026839118.

Fernández-Blanco, D., de Gelder, G., Lacassin, R., & Armijo, R. (2019). A new crustal fault formed the modern Corinth Rift. *Earth-Science Reviews*, 199, 102919.

Fernández-Blanco, D., de Gelder, G., Lacassin, R., & Armijo, R. (2020). Geometry of flexural uplift by continental rifting in Corinth, Greece. *Tectonics*, 39(1), e2019TC005685.

Gallagher, K., Charvin, K., Nielsen, S., Sambridge, M., & Stephenson, J. (2009). Markov chain Monte Carlo (MCMC) sampling methods to determine optimal models, model resolution and model choice for Earth Science problems. *Marine and Petroleum Geology*, 26(4), 525-535.

Gallen, S. F., & Fernández-Blanco, D. (2021). A new data-driven Bayesian inversion of fluvial topography clarifies the tectonic history of the corinth rift and reveals a channel steepness threshold. *Journal of Geophysical Research: Earth Surface*, 126(3), e2020JF005651.

Gawthorpe, R. L., Fabregas, N., Pechlivanidou, S., Ford, M., Collier, R. E. L., Carter, G. D., ... & Shillington, D. J. (2022). Late Quaternary mud-dominated, basin-floor sedimentation of the Gulf of Corinth, Greece: Implications for deep-water depositional processes and controls on syn-rift sedimentation. *Basin Research*, 34(5), 1567-1600.

Gowan, E. J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L., ... & Lohmann, G. (2021). A new global ice sheet reconstruction for the past 80 000 years. *Nature communications*, 12(1), 1199.

Guilcher, A., 1974. Les «rasas»: un problème de morphologie littorale générale. *Annales de Géographie*, 83, 1–33.

Hay, C., Mitrovica, J. X., Gomez, N., Creveling, J. R., Austermann, J., & Kopp, R. E. (2014). The sea-level fingerprints of ice-sheet collapse during interglacial periods. *Quaternary Science Reviews*, 87, 60-69.

- Hibbert, F. D., Rohling, E. J., Dutton, A., Williams, F. H., Chutcharavan, P. M., Zhao, C., & Tamisiea, M. E. (2016). Coral indicators of past sea-level change: A global repository of U-series dated benchmarks. *Quaternary Science Reviews*, 145, 1-56.
- Houghton, S. L., Roberts, G. P., Papanikolaou, I. D., McArthur, J. M., & Gilmour, M. A. (2003). New <sup>234</sup>U-<sup>230</sup>Th coral dates from the western Gulf of Corinth: Implications for extensional tectonics. *Geophysical Research Letters*, 30(19).
- Husson, L., Pastier, A.-M., Pedoja, K., Elliot, M., Paillard, D., Authemayou, C., et al., 2018. Reef carbonate productivity during quaternary sea level oscillations. *Geochem. Geophys. Geosyst.* 19 (4), 1148e1164.
- Jara-Muñoz, J., Melnick, D., Pedoja, K., & Strecker, M. R. (2019). TerraceM-2: A Matlab® interface for mapping and modeling marine and lacustrine terraces. *Frontiers in Earth Science*, 255.
- Johnson, M. E., & Libbey, L. K. (1997). Global review of upper Pleistocene (substage 5e) rocky shores: tectonic segregation, substrate variation, and biological diversity. *Journal of Coastal Research*, 297-307.
- Kafetzidou, A., Fatourou, E., Panagiotopoulos, K., Marret, F., & Kouli, K. (2023). Vegetation Composition in a Typical Mediterranean Setting (Gulf of Corinth, Greece) during Successive Quaternary Climatic Cycles. *Quaternary*, 6(2), 30.
- Kennedy, G. L., Lajoie, K. R., & Wehmiller, J. F. (1982). Aminostratigraphy and faunal correlations of late Quaternary marine terraces, Pacific Coast, USA. *Nature*, 299(5883), 545-547.
- Kline, S. W., Adams, P. N., & Limber, P. W. (2014). The unsteady nature of sea cliff retreat due to mechanical abrasion, failure and comminution feedbacks. *Geomorphology*, 219, 53-67.
- Kopp, R. E., Simons, F. J., Mitrovica, J. X., Maloof, A. C., & Oppenheimer, M. (2009). Probabilistic assessment of sea level during the last interglacial stage. *Nature*, 462(7275), 863-867.
- Landmann, G., Reimer, A., & Kempe, S. (1996). Climatically induced lake level changes at Lake Van, Turkey, during the Pleistocene/Holocene transition. *Global Biogeochemical Cycles*, 10(4), 797-808.
- Lajoie, K. R., Wehmiller, J. F., Kvenvolden, K. A., Peterson, E., & White, R. H. (1975). Correlation of California marine terraces by amino acid stereochemistry. In *Geological Society of America Abstracts with Programs* (Vol. 7, No. 3, pp. 338-339).
- Lajoie, K.R., 1986. Coastal tectonics. In: Press, N.A. (Ed.), *Active Tectonics*. National Academic Press, Washington DC, pp. 95e124.
- Lambeck, K., & Chappell, J. (2001). Sea level change through the last glacial cycle. *Science*, 292(5517), 679-686.
- Leclerc, F., & Feuillet, N. (2019). Quaternary coral reef complexes as powerful markers of long-term subsidence related to deep processes at subduction zones: Insights from Les Saintes (Guadeloupe, French West Indies). *Geosphere*, 15(4), 983-1007.
- Lorscheid, T., & Rovere, A. (2019). The indicative meaning calculator—quantification of paleo sea-level relationships by using global wave and tide datasets. *Open Geospatial Data, Software and Standards*, 4, 1-8.
- Malatesta, L. C., Finnegan, N. J., Huppert, K. L., & Carreño, E. I. (2022). The influence of rock uplift rate on the formation and preservation of individual marine terraces during multiple sea-level stands. *Geology*, 50(1), 101-105.
- Marra, F., Sevink, J., Tolomei, C., Vannoli, P., Florindo, F., Jicha, B. R., & La Rosa, M. (2023). New age constraints on the MIS 9–MIS 5.3 marine terraces of the Pontine Plain (central Italy) and implications for global sea levels. *Quaternary Science Reviews*, 300, 107866.
- Matsumoto, H., Young, A. P., & Carilli, J. E. (2022). Modeling the relative influence of environmental controls on marine terrace widths. *Geomorphology*, 396, 107986.
- McNeill, L. C., Shillington, D. J., Carter, G. D., Everest, J. D., Gawthorpe, R. L., Miller, C., ... & Green, S. (2019). High-resolution record reveals climate-driven environmental and sedimentary changes in an active rift. *Scientific Reports*, 9(1), 3116.
- Medina-Elizalde, M. (2013). A global compilation of coral sea-level benchmarks: implications and new challenges. *Earth and Planetary Science Letters*, 362, 310-318.
- Meschis, M., Roberts, G. P., Robertson, J., Mildon, Z. K., Sahy, D., Goswami, R., ... & Iezzi, F. (2022). Out of phase Quaternary uplift-rate changes reveal normal fault interaction, implied by deformed marine palaeoshorelines. *Geomorphology*, 416, 108432.

Mosegaard, K., & Sambridge, M. (2002). Monte Carlo analysis of inverse problems. *Inverse problems*, 18(3), R29.

Murray-Wallace, C. V. (2002). Pleistocene coastal stratigraphy, sea-level highstands and neotectonism of the southern Australian passive continental margin—a review. *Journal of Quaternary Science: Published for the Quaternary Research Association*, 17(5-6), 469-489.

Nixon, C. W., McNeill, L. C., Bull, J. M., Bell, R. E., Gawthorpe, R. L., Henstock, T. J., ... & Kranis, H. (2016). Rapid spatiotemporal variations in rift structure during development of the Corinth Rift, central Greece. *Tectonics*, 35(5), 1225-1248.

Ott, R. F., Gallen, S. F., Wegmann, K. W., Biswas, R. H., Herman, F., & Willett, S. D. (2019). Pleistocene terrace formation, Quaternary rock uplift rates and geodynamics of the Hellenic Subduction Zone revealed from dating of paleoshorelines on Crete, Greece. *Earth and Planetary Science Letters*, 525, 115757.

Pastier, A.-M., Husson, L., Pedoja, K., Bezos, A., Authemayou, C., Arias-Ruiz, C., Cahyarini, S.Y. (2019). Genesis and architecture of sequences of quaternary coral reef terraces: Insights from numerical models. *Geochem. Geophys. Geosyst.*, 20 (8), 4248e4272.

Pedoja, K., Husson, L., Regard, V., Cobbold, P. R., Ostanciaux, E., Johnson, M. E., ... & Delcaillau, B. (2011). Relative sea-level fall since the last interglacial stage: are coasts uplifting worldwide?. *Earth-Science Reviews*, 108(1-2), 1-15.

Pedoja, K., Husson, L., Johnson, M. E., Melnick, D., Witt, C., Pochat, S., ... & Garestier, F. (2014). Coastal staircase sequences reflecting sea-level oscillations and tectonic uplift during the Quaternary and Neogene. *Earth-Science Reviews*, 132, 13-38.

Pedoja, K., Jara-Muñoz, J., De Gelder, G., Robertson, J., Meschis, M., Fernández-Blanco, D., ... & Pinel, B. (2018). Neogene-Quaternary slow coastal uplift of Western Europe through the perspective of sequences of strandlines from the Cotentin Peninsula (Normandy, France). *Geomorphology*, 303, 338-356.

Perg, L. A., Anderson, R. S., & Finkel, R. C. (2001). Use of a new  $^{10}\text{Be}$  and  $^{26}\text{Al}$  inventory method to date marine terraces, Santa Cruz, California, USA. *Geology*, 29(10), 879-882.

Perissoratis, C., Piper, D. J. W., & Lykousis, V. (2000). Alternating marine and lacustrine sedimentation during late Quaternary in the Gulf of Corinth rift basin, central Greece. *Marine Geology*, 167(3-4), 391-411.

Pico, T., Mitrovica, J. X., Ferrier, K. L., & Braun, J. (2016). Global ice volume during MIS 3 inferred from a sea-level analysis of sedimentary core records in the Yellow River Delta. *Quaternary Science Reviews*, 152, 72-79.

Pirazzoli, P.A., 2005. Marine terraces. In: Schwartz, M.L. (Ed.), *Encyclopedia of Coastal Science*. Springer Netherlands, Dordrecht, pp. 632-633.

Railsback, L. B., Gibbard, P. L., Head, M. J., Voarintsoa, N. R. G., & Toucanne, S. (2015). An optimized scheme of lettered marine isotope substages for the last 1.0 million years, and the climatostratigraphic nature of isotope stages and substages. *Quaternary Science Reviews*, 111, 94-106.

Regard, V., Pedoja, K., De La Torre, I., Saillard, M., Corte\_s-Aranda, J., Nexer, M., 2017. Geometrical trends within sequences of Pleistocene marine terraces: selected examples from California, Peru, Chile and New-Zealand. *Zeitschrift Fur Geomorphologie* 61 (1), 53e73.

Roberts, G. P., Houghton, S. L., Underwood, C., Papanikolaou, I., Cowie, P. A., van Calsteren, P., ... & McArthur, J. M. (2009). Localization of Quaternary slip rates in an active rift in 105 years: An example from central Greece constrained by  $^{234}\text{U}$ - $^{230}\text{Th}$  coral dates from uplifted paleoshorelines. *Journal of Geophysical Research: Solid Earth*, 114(B10).

Roberts, D. L., Karkanas, P., Jacobs, Z., Marean, C. W., & Roberts, R. G. (2012). Melting ice sheets 400,000 yr ago raised sea level by 13 m: Past analogue for future trends. *Earth and Planetary Science Letters*, 357, 226-237.

Roeser, P., Franz, S. O., & Litt, T. (2016). Aragonite and calcite preservation in sediments from Lake Iznik related to bottom lake oxygenation and water column depth. *Sedimentology*, 63(7), 2253-2277.

Rosenbloom, N. A., & Anderson, R. S. (1994). Hillslope and channel evolution in a marine terraced landscape, Santa Cruz, California. *Journal of Geophysical Research: Solid Earth*, 99(B7), 14013-14029.

Rovere, A., Raymo, M. E., Vacchi, M., Lorscheid, T., Stocchi, P., Gomez-Pujol, L., ... & Hearty, P. J. (2016). The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world. *Earth-Science Reviews*, 159, 404-427.



- Rovere, A., Ryan, D. D., Vacchi, M., Dutton, A., Simms, A. R., & Murray-Wallace, C. V. (2023). The World Atlas of Last Interglacial Shorelines (version 1.0). *Earth System Science Data*, 15(1), 1-23.
- Schellmann, G., & Radtke, U. (2004). A revised morpho- and chronostratigraphy of the Late and Middle Pleistocene coral reef terraces on Southern Barbados (West Indies). *Earth-Science Reviews*, 64(3-4), 157-187.
- Shakun, J. D., Lea, D. W., Lisiecki, L. E., & Raymo, M. E. (2015). An 800-kyr record of global surface ocean  $\delta^{18}\text{O}$  and implications for ice volume-temperature coupling. *Earth and Planetary Science Letters*, 426, 58-68.
- Siddall, M., Smeed, D. A., Hemleben, C., Rohling, E. J., Schmelzer, I., & Peltier, W. R. (2004). Understanding the Red Sea response to sea level. *Earth and Planetary Science Letters*, 225(3-4), 421-434.
- Sondi, I., & Juračić, M. (2010). Whiting events and the formation of aragonite in Mediterranean Karstic Marine Lakes: new evidence on its biologically induced inorganic origin. *Sedimentology*, 57(1), 85-95.
- Spratt, R. M., & Lisiecki, L. E. (2016). A Late Pleistocene sea level stack. *Climate of the Past*, 12(4), 1079-1092.
- Stein, M., Torfstein, A., Gavrieli, I., & Yechieli, Y. (2010). Abrupt aridities and salt deposition in the post-glacial Dead Sea and their North Atlantic connection. *Quaternary Science Reviews*, 29(3-4), 567-575.
- Stirling, C. H., Esat, T. M., Lambeck, K., McCulloch, M. T., Blake, S. G., Lee, D. C., & Halliday, A. N. (2001). Orbital forcing of the marine isotope stage 9 interglacial. *Science*, 291(5502), 290-293.
- Strobl, M., Hetzel, R., Fassoulas, C., & Kubik, P. W. (2014). A long-term rock uplift rate for eastern Crete and geodynamic implications for the Hellenic subduction zone. *Journal of Geodynamics*, 78, 21-31.
- Sunamura, T. (1992). *Geomorphology of Rocky Coasts*, vol. 3. John Wiley & Son Ltd.
- Tawil-Morsink, K., Austermann, J., Dyer, B., Dumitru, O. A., Precht, W. F., Cashman, M., ... & Raymo, M. E. (2022). Probabilistic investigation of global mean sea level during MIS 5a based on observations from Cave Hill, Barbados. *Quaternary Science Reviews*, 295, 107783.
- Toomey, M., Ashton, A. D., & Perron, J. T. (2013). Profiles of ocean island coral reefs controlled by sea-level history and carbonate accumulation rates. *Geology*, 41(7), 731-734.
- Van Daele, M., van Welden, A., Moernaut, J., Beck, C., Audemard, F., Sanchez, J., ... & De Batist, M. (2011). Reconstruction of Late-Quaternary sea- and lake-level changes in a tectonically active marginal basin using seismic stratigraphy: The Gulf of Cariaco, NE Venezuela. *Marine Geology*, 279(1-4), 37-51.
- Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., Mcmanus, J. F., Lambeck, K., ... & Labracherie, M. (2002). Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. *Quaternary science reviews*, 21(1-3), 295-305.
- Watkins, S. E., Whittaker, A. C., Bell, R. E., McNeill, L. C., Gawthorpe, R. L., Brooke, S. A., & Nixon, C. W. (2019). Are landscapes buffered to high-frequency climate change? A comparison of sediment fluxes and depositional volumes in the Corinth Rift, central Greece, over the past 130 ky. *GSA Bulletin*, 131(3-4), 372-388.
- Weber, G.E. (1990). Late Pleistocene slip rates on the San Gregorio fault zone at Point Ano Nuevo, San Mateo County, California, in Garrison, R.E., et al., eds., *Geology and tectonics of coastal California, San Francisco to Monterey* (volume and guidebook): Bakersfield, California, Pacific Section, American Association of Petroleum Geologists, p. 193-203.
- Webster, J. M., Wallace, L. M., Clague, D. A., & Braga, J. C. (2007). Numerical modeling of the growth and drowning of Hawaiian coral reefs during the last two glacial cycles (0-250 kyr). *Geochemistry, Geophysics, Geosystems*, 8(3).