

Influence of minor hydrocarbon seepage on sulfur cycling in marine subsurface sediments

Ellen Schnabel¹, Aurèle Vuillemin¹, Cédric C. Laczny², Benoit J. Kunath², André R.
5 Soares³, Rolando Di Primio⁴, Jens Kallmeyer^{1*} and the PROSPECTOMICS Consortium[†]

¹ GFZ German Research Centre for Geosciences, Section Geomicrobiology, Telegrafenberg, 14473
Potsdam, Germany

² Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg

³ Environmental Metagenomics, Research Center One Health Ruhr of the University Alliance Ruhr, Faculty
10 of Chemistry, University of Duisburg-Essen, Essen, Germany.

⁴ AkerBP, 1366 Lysaker, Norway

*<https://www.prospectomics.eu/>

[†]A full list of authors and their affiliations appears at the end of the paper.

**Corresponding author*

Jens Kallmeyer, GFZ German Research Centre for Geosciences, Section Geomicrobiology, Telegrafenberg,
14473 Potsdam, Germany, email: kallm@gfz-potsdam.de

20 **Abstract.**

All hydrocarbon (HC) reservoirs leak to some extent. When small quantities of HCs escape offshore
reservoirs and migrate through overlying organic-poor marine sediments towards the surface, these HCs are
often completely metabolized by microorganisms before reaching the sediment-water interface. However,
these low and often unnoticed upwards HC fluxes still influence the geochemistry of surrounding sediments
25 and potentially stimulate metabolic activity by microbial populations in shallow seafloor environments.

In this study, we examined how localized HC seepage affects microbial sulfate reduction in
organic-poor sediments from the SW Barents Sea, focusing on three sampling areas overlying known HC
deposits and two pristine seabed reference areas. The analysis of 50 gravity cores revealed significant
variability in the predicted sulfate depletion depth, ranging from 3 to 12 m below the seafloor. We observed
30 nearly linear pore water sulfate and alkalinity profiles, alongside low rates of sulfate reduction ($\mu\text{mol} \times \text{cm}^3$

× d⁻¹). Metagenomic and metatranscriptomic data indicated metabolic potential and activity towards sulfate reduction and anaerobic oxidation of methane (AOM). Expression of functional marker genes (*aprAB*, *dsrAB*, *mcrA*) revealed metabolic activities by sulfate-reducing Desulfobacterota and methanotrophic ANME-1 archaea sustained by HC traces in the sediment. Further, in concomitance to AOM processes, we found that archaea amongst the class Lokiarchaeia and Thoarchaeia were expressing genes involved in sulfur reduction (*hydB*, *hydG*).

Overall, our study demonstrates that the gradient in pore water geochemistry, the rates of sulfate reduction processes, and the genetic features of microbial populations actively involved in sulfate-driven AOM processes are all affected by low and often unnoticed upwards HC fluxes. This slight HC seepage resulted in sulfate depletion at shallower depths compared to what would be expected at a comparable, non-affected site, and produced concomitant biogeochemical signatures in the shallow subsurface that enable the inference of deeply buried reservoirs.

1. Introduction



All hydrocarbon (HC) reservoirs leak to some degree (Hunt, 1995; Yergin, 2009; Heggland, 1998), with over 80 % of all seeps occurring directly above the reservoirs (Ciotoli et al. (2020)). While large leakages result in conspicuous manifestations at the seafloor, (e.g. Cramm et al. (2021)), minor seeps, characterized by low, primarily diffusive HC fluxes usually remain invisible at the sediment-water interface (SWI), as there are no visible signs of seepage, e.g. carbonate crusts or chemosynthetic communities such as sulfur bacteria mats, clams, tubeworms. In cases where HC fluxes are so low that hydrocarbons are fully mineralized and do not reach the SWI, sediment geochemistry and microbiology at minor seepage sites will be altered (Rasheed et al., 2013; Abrams, 2020; Riedinger et al., 2014; Joye et al., 2004). Even if HC fluxes were higher and hydrocarbons do indeed reach the SWI, they are often overlooked (Abrams, 2020; Joye, 2020).

HC seepage triggers both geochemical and microbiological processes, changing physicochemical properties, such as sediment porosity, mineral and pore water composition, as well as microbial community composition and activity (Abrams, 2020; Hvoslef et al., 1996; Joye, 2020). Likewise, the presence of free gas in methane-containing pore fluids alters sediment porosity and compressibility (Jang et al., 2018), which can be detected through seismic profiling due to the resulting impedance contrast (Rovere et al., 2020). These changes may also affect the sediment's geochemical properties, such as clay texture, organic matter (OM) reactivity, and metal content (Chen et al., 2023). HC influx and subsequent HC consumption will also alter pore water geochemistry and pH, promoting ion exchange reactions, dissolution and

precipitation (Smrzka et al., 2019; Turchyn et al., 2021; Torres et al., 1996) and structural conversion of minerals (Jiang, 2012). As a consequence, specific trace metals and metalloids (e.g. Cr, Cu, As, Se, Sb) that can be metabolized by specific microorganisms (Raab and Feldmann, 2003) tend to become enriched in the sediment pore water at seepage sites (Rasheed et al., 2013; Guseva et al., 2021). Finally, most offshore sediments are composed of pelagic deposits, i.e. clays, silts or biogenic oozes, containing only low amounts of bioavailable OM that becomes increasingly recalcitrant with depth of burial (Middelburg, 2018). Thus, the influx of HCs in such sedimentary systems represents an additional source of electron donors, providing chemical energy for microorganisms and potentially fueling various metabolic processes. The metabolic potential to break down and assimilate HCs has been identified in a large number of microorganisms, e.g. reviewed in Joye (2020), whose activities may induce concomitant effects in the pore water, organic and mineral fraction of the sediment (Kim et al., 2004).

The quantitatively most important anaerobic OM degradation process in the upper layers of marine sediments, where oxygen is depleted, but sulfate is still present in relatively high concentrations is sulfate reduction (Jørgensen et al., 2024; Bowles et al., 2014; Kasten and Jørgensen, 2000; Jørgensen, 1982). Due to its abundance in seawater, sulfate diffuses into the sediment pore water where it is used as an electron acceptor by sulfate-reducing microorganisms composed of sulfate reducing bacteria and ANME (Widdel et al., 2010). Sulfate reduction (SR) can be divided into two main pathways, namely organoclastic and methanotrophic SR. The respective overall reactions are:

80



Due to the abundance of easily biodegradable OM and sulfate close to the sediment-water interface, organoclastic SR normally prevails in anaerobic OM degradation within the uppermost part of the sediment column, leading to a decrease in sulfate concentration. Methanotrophic SR becomes predominant only when biogenic or thermogenic methane diffusing from deeper sediments reaches the zone where sulfate is still available but approaches depletion (Martens and Berner, 1974). The narrow zone where both methane and sulfate are available is referred to as the sulfate-methane transition zone (SMTZ). Within the SMTZ, mainly methane, and potentially also other OM compounds (Beulig et al., 2019; Jørgensen et al., 2019b), are oxidized by microorganisms while sulfate is reduced (Equation 2).

Thus, the respective methane and sulfate fluxes exert control on the depth of the SMTZ (Borowski et al., 1996, 1999; Jørgensen and Kasten, 2006), which, under steady-state conditions, is usually a reflection of methanogenesis rates (Henrichs and Reeburgh, 1987) or fluxes of thermogenic methane (Hu et al., 95 2017). However, the SMTZ is highly dynamic and responds relatively rapidly (i.e. within decades) to shifts in pore water geochemistry (Sultan et al., 2016; Hong et al., 2016). Thus, microbially active subseafloor sediments exhibit significant variability in the depth of the SMTZ across different oceans, from 1 to 10 of meters on the continental shelf (Egger et al., 2018; Treude et al., 2020; Treude et al., 2014; Riedinger et al., 2017; Riedinger et al., 2014; März et al., 2008; Niewöhner et al., 1998). On the continental slope, the 100 SMTZ can be down to 100 m below seafloor (mbsf) in extreme cases (Nunoura et al., 2009), but is usually not deeper than 25 mbsf.

The subseafloor biosphere hosts phylogenetically diverse and metabolically active microbial communities (Parkes et al., 2014; D'hondt et al., 2004). Their metabolic activity depends mainly on biogeochemical cycling of nitrogen, phosphate and sulfur. At HC seepage sites, microbial communities are 105 well-defined in terms of taxonomy and highly specialized in terms of metabolic functions, and are referred to as “seep biomes” (Ruff et al., 2015). It has been known for many decades that methanogenic hydrocarbon degradation is carried out jointly by bacteria and archaea in underground hydrocarbon reservoirs. New studies suggest that *Candidatus methanoliparum* may be able to independently degrade long-chain alkanes and produce methane (Wegener et al., 2022; Zhou et al., 2021; Zhang et al., 2024). Seep 110 biomes show spatial heterogeneity, but are distinct from non-seep environments (Pop Ristova et al., 2015). The reported key microbial guilds include methane and oil oxidizers, sulfate reducers, and nitrogen fixers. Taxa found near HC seeps involve members of the bacterial phyla *Desulfobacterota* and *Chloroflexota* that can respectively use simple and halogenated HCs as electron acceptors (Kleindienst et al., 2014; Zhang et al., 2023), whereas archaeal groups, like the ANME or Asgardarchaeota, consume methane or make use of 115 HCs or sulfur compounds (Firrincieli et al., 2021; Macleod et al., 2019).

Our study aims to determine in what ways minor HC seepage modifies the diversity and activity of anaerobic methanotrophs and sulfate-reducing bacteria in surface sediments of the SW Barents Sea. We hypothesize that minor HC seepage influences pore water geochemical properties, even in near-surface sediment, and that resident microbial communities respond quickly in terms of composition and activity to 120 the geochemical alteration of their habitats.

For this, we collected a total of 50 gravity cores, of which 40 cores from areas affected by light HC seepage from underlying HC reservoirs (i.e., HC-affected sites) but without any visible surface

manifestations like carbonate crusts or chemosynthetic communities such as sulfur bacteria mats, clams, tubeworms, and 10 cores from seabed areas without seepage (i.e., reference sites). We combine detailed analyses of pore water sulfate and alkalinity with potential and modeled sulfate reduction rates, dissolved methane (CH₄) and carbon dioxide (CO₂) concentrations, and supplement them with metagenomic (i.e. total DNA) and metatranscriptomic (i.e. total RNA) data on taxonomic and functional marker genes in order to trace metabolic activity by microbial sulfate-reducing consortia in their geochemical context.

2. Material and methods

2.1 Geological context of the Barents Sea, sampling sites

The Barents Sea is an epicontinental shelf sea surrounded by the marginal Norwegian Sea to the southwest, the Arctic Ocean to the north and the Russian island Novaya Zemlya to the east (Fig. 1a). Since its formation during the Caledonian orogeny (i.e. 490-390 Ma), the Barents Sea Basin experienced several phases of uplift and subsidence, with subsequent tilting and erosion. These geologic events resulted in the characteristic structure of the Barents Sea in the form of a succession of structural highs and basins (Gabrielsen et al., 1990; Larssen et al., 2002).

HC exploration in the Barents Sea has a history stretching back several decades (Doré, 1995; Johansen et al., 1993). The sediments have been characterized in numerous studies, e.g. Sættem et al. (1991); Elverhøi and Solheim (1983), and are mainly composed of organic-poor silty clays (Knies and Martinez, 2009; Nickel et al., 2013) originating from eroded sedimentary and igneous rocks.

During an expedition to the southern Loppa High region of the Barents Sea (Fig. 1a) from October 29th to November 8th 2021, we collected 50 gravity cores (up to 3 m in length) sediment at about 350 m water depth at three areas with known underlying HC reservoirs (i.e. Zone 1, Zone 10, Zone 12) and at two reference areas (i.e. Ref 5, Ref 6) without underlying HC reservoirs (Fig. 1b), labelled thereafter HC-affected sites and reference sites, respectively. The cores were subsampled for sediment and pore water on board the research vessel.




Figure 1. Sampling locations. (a) Map of the Norwegian coast. The square marks the sampling area. (b) Fifty cores were retrieved from three HC-affected sites (Zone 1, Zone 10, Zone 12) and two reference sites (Ref 5, Ref 6) (bathymetric map: Emodnet (2022)) (c) bathymetric map of the 3 HC-affected sites. (d) Each site was sampled in a grid of cores, Zone 1 is enlarged here as an example. The missing samples in the middle are due to the increased occurrence of dropstones in this area, preventing the deployment of a gravity corer.

155 2.2 Pore water and sediment sampling

Pore water samples were extracted from all cores at a resolution of 10 cm, using rhizons (Rhizosphere Research Products) (Seeberg-Elverfeldt et al., 2005), collecting a minimum volume of ca. 5 mL over 24 h. The pore water was filtered through 0.2 μm pore size cellulose esters (MCE) membrane syringe filters (Merck MF-Millipore) and aliquoted as follows: 1.5 mL transferred into a plastic screw cap vial without further treatment for anion and cation measurements; 2 mL into a glass vial amended with 50 μL saturated HgCl_2 solution to inhibit microbial activity and sealed without headspace for alkalinity measurements (Edenborn et al., 1985); and 1.5 mL into a plastic screw cap vial mixed with 200 μL ZnCl_2 (20 % weight \times vol⁻¹) to precipitate dissolved hydrogen sulfide as ZnS for subsequent sulfide measurements. All samples were stored at +4 °C until analysis in the home lab.

165 Immediately after core recovery, the lowermost 20 cm of each sediment core were cut off and subsampled for DNA and RNA extraction. The sediment was pushed out of the liner and the outer 1-2 cm of sediment rims were removed with a sterile spatula. Aliquots of the remaining sediment were transferred into gas-tight aluminum foil bags, flushed with nitrogen gas, heat-sealed and stored at -80 °C for later extraction of various biomolecules.

170

2.3 Pore water sulfate, sulfide and alkalinity

Anion concentrations were determined using an Ion Chromatography (IC) System equipped with a SykroGel A \times 300 AB-A01 column (all Sykam). The eluent contained 7.3 mg \times L⁻¹ NaSCN and 636 mg \times L⁻¹ NaCO₃. The pump rate was set to 1 mL \times min⁻¹ and the injected sample volume was 50 μL . All samples were measured in triplicates and the results were averaged. Because of their high salt concentrations, all samples had to be diluted 1:50 with MilliQ water before injection. The detection limit of the IC system is 0.5 mg \times L⁻¹. The average standard deviation of replicate measurements was always better than 3 %.

Sulfide concentrations were measured according to the protocol of (Cline, 1969). Aliquots with absorption values greater than 1 were diluted and remeasured. Sulfide concentrations were calculated by comparing adsorption values measured for pore water samples to those of a Na₂S standard. The detection limit of the method is 0.1 μM. All samples were measured in triplicates and the results were averaged, with standard deviation ≤ 3.5 %.

Pore water alkalinity was determined using the Visocolor HE alkalinity AL 7 kit (Macherey-Nagel GmbH). As we only had a total of ca. 2 mL of pore water per sample, we scaled down the volume of sample to 500 μL in order to perform all measurements in triplicates. For this, 10 μL of indicator was added to 500 μL of pore water and titrated with the Visocolor solution until we observed a change in color from blue to orange. Alkalinity of the pore water samples was calculated based on the titration solution volume. Replicates differed by less than 3 % on average.

190 **2.4 Sulfate reduction rates, modeled sulfate reduction rates**

For sulfate reduction measurements, each gravity core was subsampled in triplicate every 40-50 cm and at the bottom end. Structurally intact sediment plugs (volume ca. 3.5 cm³) were recovered by inserting 5 mL glass barrels fitted with a syringe plunger into the sediment. Following the whole core injection method (Jørgensen, 1978), 15 μL of radioactive ³⁵SO₄-sulfate tracer (200 kBq) were injected into the glass barrels containing the samples retrieved on the same day (Fossing et al., 2000). The glass barrels were closed with butyl rubber stoppers, and the samples incubated at *in situ* temperature (ca. 4 °C) for 24 hours. Each sample was then transferred into a previously weighed 50 mL centrifuge tube containing 10 mL of 20 % zinc acetate solution to stop all microbial activity and to precipitate all volatile hydrogen sulfide as zinc sulfide. Each vial was then thoroughly shaken to break down all sediment aggregates and stored frozen until analysis. Blank samples were prepared by adding the radiolabeled sulfate tracer just before transferring the sample into the zinc acetate solution. The tracer blank was prepared in the same way without sediment.

Sulfate reduction rates (SRR) were quantified in the home lab, using single-step cold chromium distillation (Kallmeyer et al., 2004). The sample vials were thawed, centrifuged and the supernatant carefully removed. A small aliquot of supernatant was kept for quantification of total radioactivity, the rest was discarded. The sediment pellet was then quantitatively transferred into a distillation flask. Due to the clayey consistence of the sediments, we had to re-freeze the vials and transfer the frozen sediment pellet from the centrifuge tube to the distillation flask. SRR were calculated according to the following formula:

Sulfate reduction rate:
$$SRR = \frac{SO_4 * \varphi * a_{tris} * 1.06 * 10^6}{a_{tot} * t} \quad (\text{Equation 3})$$

SRR: sulfate reduction rate [$\text{pmol cm}^{-3} \times \text{day}^{-1}$]; SO_4 : sulfate concentration in the pore water [$\text{mmol} \times \text{L}^{-1}$];
 210 φ : porosity of the sediment, set to 0.7; a_{tris} : radioactivity of total reduced inorganic sulfur [cpm]; a_{tot} : total
 radioactivity used in the spiking [cpm]; t : incubation time [d]; 1.06: correction factor for isotopic
 fractionation; 10^6 : unit conversion factor from [$\mu\text{mol} \times \text{L}^{-1}$] to [$\text{pmol} \times \text{L}^{-1}$].

Assuming steady state conditions, the net rate of sulfate production, or consumption, can be
 modeled based on the measured sulfate concentration profiles by the diffusion-reaction modeling software
 215 PROFILE (Berg et al., 1998). This software considers three different kinds of vertical transport (i.e.
 diffusion, bioturbation, irrigation) together with the flux across the SWI. The final output corresponds to
 the minimum number of intervals, as defined by their respective sulfate production or consumption rates,
 that are necessary to reproduce the measured pore water profiles. For the modeling calculation, we used:
 coefficient of sulfate diffusion in water at 4 °C $D_{\text{Sulfate}} = 0.56 \times 10^{-5} [\text{cm}^2 \times \text{s}^{-1}]$ (Iversen and Jørgensen
 220 (1993); porosity $\varphi = 0.7$; and sulfate seawater concentration $28 \text{ mmol} \times \text{cm}^3$. Because the sediment
 appeared to be free of macrobenthos, the biodiffusivity and irrigation coefficients were set to $D_B = 0 \text{ cm}^2 \times$
 s^{-1} , and $\alpha = 0$, respectively.

2.5 Methane and carbon dioxide concentrations

225 For dissolved CH_4 and CO_2 concentrations, we subsampled the lowermost 20 cm of sediment from each
 gravity core directly after recovery. Using 5 mL cut-off syringes, we extracted ca. 4 cm^3 of sediment per
 core. The first and last 0.5 cm^3 of sediment in the syringe were discarded while the central 3 cm^3 were
 transferred into a 10 mL glass crimp vial containing a saturated NaCl solution. The vials were immediately
 sealed with thick butyl rubber stoppers, crimped, and stored upside down without headspace until analysis
 230 in the home lab.

Prior to measurement, we introduced 3 mL of ultrapure helium gas as headspace while
 withdrawing the same amount of NaCl solution from the vial. To equilibrate dissolved gases with the
 headspace, the content of the vials was mixed at 220 rpm on an orbital shaker for 18 h, and further vortexed
 to break up the remaining small clayey aggregates. We extracted ca. $350 \mu\text{L}$ of headspace from the vial
 235 using a gas-tight syringe. Prior to measurement, ca. $100 \mu\text{L}$ of the sampled gas were flushed through the
 injection needle, and $250 \mu\text{L}$ of gas sample were then injected into a 7890A Gas Chromatography System
 equipped with a flame ionization detector (FID), a thermal conductivity detector (TCD) and HP PLOT Q

column (all Agilent). Oven temperature was set to 50°C, flow rate to 17.2 mL × min⁻¹ and pressure to 13 psi. The detectors both worked at 200°C with flow rates of 40 mL × min⁻¹ (FID) and 15 mL × min⁻¹ (TCD).
240 The system was calibrated, using 250 µL of analytic pure standards, injecting CO₂ concentrations of 310 ppm and 5270 ppm, and CH₄ concentrations of 10 ppm and 5170 ppm. The initial CH₄ and CO₂ concentrations were converted from ppm to molar concentrations by applying the ideal gas law.

2.6 DNA/RNA extractions, sequencing libraries, functional marker genes

245 DNA was extracted for a total of 12 samples from 8 different cores corresponding to 4, 2 and 2 samples from HC-affected sites, i.e. Zone 1, Zone 10 and Zone 12, respectively, along with 3 and 1 samples from Ref 5 and Ref 6, respectively. Samples were subjected to DNA extraction using the DNEasy PowerMax Soil Kit (Qiagen, Germany) following manufacturer's instructions with few adaptations. For 2.5 g of starting sediment material, the final elution was performed in two steps using twice 2.5 mL of solution C6,
250 and successively incubated and centrifuged. Eluted DNA was then concentrated by adding 510 µL of a 3M sodium acetate (pH 5.2) / glycogen (0.4 µg × µL⁻¹) solution and incubated for one hour at -20°C and then centrifuged to pellet DNA. After ethanol cleaning and air-drying, DNA was resuspended in 100 µL of 10 mM Tris buffer. Finally, DNA was cleaned and concentrated using the Zymo DNA Clean & Concentrator (Zymo, UK) following the manufacturer's instructions and eluted in a final elution volume of 35 µL.

255 DNA sequencing was achieved by using 32.5 µL of purified and concentrated DNA for metagenomic library preparation with the QIAseq FX DNA library UDI A/B kit (Qiagen, Germany) irrespective of the individual sample concentrations. The genomic DNA was enzymatically fragmented for 10 min and DNA libraries were prepared with 7 PCR cycles for the PCR-less procedure, and without dedicated PCR step for the PCR-free procedure. The average library sizes were ±400 base pairs. Prepared
260 libraries were quantified using a Qubit fluorometer (ThermoFischer, USA) and quality-checked on a Bioanalyzer (Agilent, USA). Sequencing was performed on an Illumina NextSeq2000 platform using 2 × 151 bp read length at the Luxembourg Centre for Systems Biomedicine (LCSB) Sequencing Platform, aiming at an average of 10 Giga base pairs per sequencing library.

RNA was extracted from 2 × 5 g of sediment per sample, using the RNeasy PowerSoil Total RNA
265 kit (QIAGEN) according to the manufacturer's instructions, yielding 30 µL of RNA-containing extract, and quantified on a Qubit4 fluorometer, using the RNA high sensitivity kit (ThermoFisher Scientific). RNA extracts were directly subjected to library preparation in technical duplicates, using the Revelo™ RNA-Seq High Sensitivity kit (Tecan Life Sciences) with 8 µL of RNA extracts as template. This trio kit includes

first and second strand cDNA synthesis, adaptor ligation and library amplification through 10 PCR cycles.
270 Prepared libraries were quantified using a Qubit4 fluorometer (ThermoFisher Scientific) and quality-
checked on a Bioanalyzer (Agilent) to determine the molarity and overall size distribution of RNA
molecule fragments. The average library fragment size was ± 300 bps and the yields were between 4 and 12
 $\text{ng} \times \mu\text{L}^{-1}$. Metatranscriptomic libraries displaying high and low molarities were pooled into two different
batches, and both sequenced by the commercial company EMBL (Heidelberg, Germany) on a NextSeq
275 2000 with a P3 kit and an Illumina micro kit (2×150 bps), respectively.

De novo assembly of metagenomic reads was performed (Table S1) using a customized
Snakemake workflow (Mölder et al., 2021), including quality-control, binning into contigs, gene prediction
and annotation as described in (Bornemann et al., 2023). Paired-end reads were interleaved, Illumina
adapters and controls removed, reads quality-based trimmed, and de-interleaved into forward and reverse
280 read files. Bacterial and archaeal scaffolds were concatenated and mapped back to raw reads to retrieve
coverage, % GC and contig lengths. Genes were then predicted by extracting open reading frames (ORFs)
from metagenome-assembled scaffolds using Prodigal (Hyatt et al., 2010). Metatranscriptomic reads were
quality-filtered and trimmed using Trimmomatic (Bolger et al., 2014). Functional profiles were obtained by
mapping the filtered and trimmed reads to their corresponding metagenomic databases using BWA-mem
285 (Li and Durbin, 2009). ORFs were retrieved using Bakta (Schwengers et al., 2021) and combined with the
mapping using the featureCounts tool (Liao et al., 2014).

Taxonomic identifications of all ORFs were further integrated with functional annotations by
performing BLASTp searches against an aggregated database of predicted proteins (Orsi et al., 2018), using
DIAMOND protein aligner v. 0.9.24 (Buchfink et al., 2015). Cut-off values for assigning the best match to
290 specific taxa present in the aggregated database were performed at a minimum bit score of 50, minimum
amino acid similarity of 60, and an alignment length of 50 residues. We used this approach to draw
conclusions about metabolic traits derived specifically from –phylum to class taxonomic levels only
(Vuillemin et al., 2020a; Vuillemin et al., 2020b).

Here, we focus on ORFs encoding genes involved in sulfur cycling and methane-related processes
295 only. To trace metabolic potential in anaerobic respiration of sulfur compounds (i.e. sulfate, sulfite,
thiosulfate, sulfur, polysulfide), we looked for the presence of ORF- encoding genes with similarity to
adenylylsulfate reductase (*apr*), dissimilatory sulfite reductase (*dsr*), anaerobic sulfite reductase (*asr*),
thiosulfate/polysulfide reductase (*phs/psr*), and sulfhydrogenase (*hyd*) (Vuillemin et al., 2022). For methane
production and anaerobic consumption, we targeted ORFs encoding proteins of the methyl coenzyme M


300 reductase (*mcr*). To identify and characterize metabolically active taxa, we targeted the small subunit ribosomal ribonucleic acid (SSU rRNA) as a taxonomic marker and the filamenting temperature-sensitive mutant A-Z (*FtsAZ*) for cell division.

To confirm taxonomic assignments and metabolic activity of taxa involved in sulfate reduction and AOM processes, we performed a phylogenetic analysis of the RNA polymerase sigma factor (*rpoD*) and ribosomal protein S3 (*rps3*) gene proteins, *aprAB*, *dsrAB*, *mcrA* and *hydB* that we could detect in the 305 transcriptomes (Vuillemin et al., 2020a). All ORFs annotated to these genes were aligned against their top two BLASTp hits in the NCBI database, using MUSCLE (Edgar, 2004). Conserved regions of the alignments were selected using Gblocks 0.91b (Castresana, 2000) with the following settings: allowing for smaller final blocks, gap positions within the final blocks and less strict flanking positions (Dereeper et al., 310 2008). Phylogenetic analysis of the resulting amino acid alignments of the predicted proteins were conducted in SeaView version 5.0.5 (Gouy et al., 2010), using PhyLM maximum likelihood (Guindon et al., 2010), with BLOSUM62 as the evolutionary model and 100 bootstrap replicates

3. Results

315 3.1 Pore water geochemical profiles

All sulfate concentrations in the uppermost profiles roughly match the sulfate concentration of the Barents Sea bottom waters (ca. 28 mM). In all cores, sulfate concentrations decrease almost linearly with depth, linear regressions showing correlation coefficients of $R^2 > 0.85$ (Fig. 2a-2b and Supplementary Fig. S1). The slopes of the regression lines can be divided into two groups corresponding to one group encompassing 320 reference sites Ref 5 and Ref 6, and another group including the three HC-affected sites Zone 1, Zone 10 and Zone 12. Based on a t-test using the software PAST v. 3.20 (Hammer et al., 2001), the slopes of these two groups are statistically significantly different ($t = 5.574$, $p = 1.25E-6$) from each other (Fig 2). The sulfate profiles from HC-affected sites show a steeper decrease with depth than those from reference sites. On average, sulfate concentrations decrease by $10 \text{ mmol} \times \text{m}^{-1}$, which would correspond to complete sulfate 325 depletion at roughly 3 mbsf at HC-affected sites. At reference sites, the average slope is $3 \text{ mmol} \times \text{m}^{-1}$ with little variations, which implies that sulfate would be depleted at ca. 12 mbsf.



330 **Figure 2. Pore-water linear regression for sulfate concentrations, alkalinity, sulfide, along with some representative profiles.**

(a, b) Regression lines for sulfate profiles (derived from measured pore water concentrations) are linear ($R^2 > 0.85$) and exhibit decreasing trends with depth. The regression lines shown in (a) represent HC-affected

sites (Zone 1, Zone 10, Zone 12), which display a steeper slope compared to those in (b) reference sites (Ref 5, Ref 6). (c, d) Regression lines for pore-water alkalinity (calculated from measured pore water concentrations) are linear ($R^2 \geq 0.85$) and increase with depth. The regression lines for (c) HC-affected sites show a more gradual increase compared to (d) reference sites. (e) Sulfide concentrations measured across all cores from HC-affected sites (represented by dots), with their corresponding regression lines shown where sulfide concentrations were quantifiable. (f) Subset of representative profiles selected for each sampling site. The uppermost and lowermost rows depict cores with the highest and lowest R^2 values, respectively, from HC-affected and reference sites.

All alkalinity profiles display a clear linear increase with depth ($R^2 \geq 0.85$). From $5 \text{ mmol} \times \text{L}^{-1}$ at the SWI, pore water alkalinity increases differently with depth, corresponding to 8, 5 and 6 $\text{mmol} \times \text{L}^{-1} \times \text{m}^{-1}$ at HC-affected sites Zone 1, Zone 10, and Zone 12, and $\leq 1 \text{ mmol} \times \text{L}^{-1} \times \text{m}^{-1}$ at reference sites Ref 5 and Ref 6, respectively (Fig. 2c-2d and Supplementary Fig. S2).


Based on the slopes of pore water profiles, the calculated alkalinity fluxes are larger at HC-affected sites than at reference sites. The same is true for sulfate fluxes, but differences across HC-affected sites and reference sites are less pronounced (Supplementary Fig. S3). In comparison, variations observed within each sampling site are small.

Sulfide could hardly be detected in pore water samples collected from reference sites, most values being below $1 \mu\text{M}$. In contrast, sulfide was detected in most of the cores from HC-affected sites. Especially at Zone 1 and Zone 10, sulfide concentrations increase linearly with depth (Fig. 2e). The majority (ca. 70 %) of the cores from Zone 1 show an increase of $2.5 \text{ mM} \times \text{m}^{-1}$ in sulfide concentrations. One core (Z01C03) stands out, reaching sulfide concentrations of ca. 11 mM at 1 mbsf. Several cores (ca. 57 %) from Zone 10 show a binary distribution of sulfide concentrations, corresponding either to linear profiles reaching ca. 1mM sulfide at 1 mbsf, or to a steeper increase reaching ca. 3.7 mM at the same depth. At Zone 12, sulfide is detectable in only 2 of the 17 cores, with much lesser increase in concentrations compared to Zone 1 and Zone 10.

3.2 Measured and modeled sulfate reduction rates

SRR were detectable in only 29 out of the 50 cores (Fig. 3), with rates between 4 and $320 \text{ pmol} \times \text{cm}^{-3} \times \text{d}^{-1}$. In 403 of the 455 samples across the 50 cores analyzed (i.e. 89 %), SRR were below the detection limit (1

$\text{pmol} \times \text{cm}^{-1} \times \text{d}^{-1}$). The occurrence of very low but detectable SRR increased with sediment depth, which mostly stands true for HC-affected sites (Fig. 3).



365

Figure 3. Sulfate reduction rates (SRR) and percentages of samples with sulfate reduction rates above the detection limit. Low SRR became increasingly detectable with increasing sediment depth. The number of samples with measurable SRR (% grey bars) is higher in sediments from HC-affected sites (dots) compared to reference sites (triangles), *n.B.* due to a lack of samples, the two uppermost intervals were merged.

370

Diffusion-reaction modeling with PROFILE (Berg et al., 1998) suggests steady SRR in 47 of the 50 cores (Supplementary Fig. S1), implying that SRRs remain constant over the entire length of the cores. The modeled SRR profiles show SR activity (i.e. $>0 \text{ pmol} \times \text{cm}^{-3} \times \text{d}^{-1}$) in 33 cores, of which 6 display rates $\leq 0.01 \text{ pmol} \times \text{cm}^{-3} \times \text{d}^{-1}$, and 27 with supposed SRR $\geq 1 \text{ pmol} \times \text{cm}^{-3} \times \text{d}^{-1}$. For three specific cores (i.e. Z01C02, Z01C09, Z12C17), PROFILE proposes two intervals with different SRRs, suggesting that SR processes may only be active in the upper 50 cmbsf and inactive in the sediment below.

375


The modeled and measured SRR are indicative of net and gross turnover rates, respectively. The gross rate reflects microbial turnover whereas the net turnover rate includes the recycling of reduced sulfur compounds (Berg et al., 1998). The modeled (i.e. net) rates for most samples are higher than the measured (i.e. gross) rates, which is rather unusual and can be explained by very low values whose absolute changes

380

have a drastic effect on the calculated values. The modeled and measured SRR are nevertheless within the same order of magnitude, and thus comparable.

385 3.3 Methane and carbon dioxide concentrations

All samples contain measurable methane concentrations, ranging from 0.08 to 19.79 $\mu\text{mol} \times \text{L}^{-1}$. All samples from reference sites contain less than 5 $\mu\text{mol} \times \text{L}^{-1}$ of methane, detection of any higher concentrations being restricted to some samples from HC-affected sites and only below 100 cmbsf (Fig. 4). CO_2 was detectable in all samples, with concentrations ranging from 29 $\mu\text{mol} \times \text{L}^{-1}$ to 512 $\mu\text{mol} \times \text{L}^{-1}$, but without any specific trend with depth (Fig. 4). The reason is probably that any CO_2 from whichever source enters the carbonate equilibrium and the gas is removed from the pore water, both in HC-affected sites and reference sites. The low methane and carbon dioxide concentrations prevented any measurement of their stable isotopic signatures.



395 **Figure 4. Concentrations of dissolved methane and carbon dioxide. (a)** At HC-affected sites, methane concentrations are highest in samples originating from 100 cmblf or deeper, whereas methane concentrations at reference sites remain below 5 $\mu\text{mol} \times \text{L}^{-1}$. **(b)** CO_2 concentrations across samples do not display any specific trend with sediment depth.

400 3.4 Statistical correlations

To identify geochemical differences between HC-affected and reference sites, we performed a Pearson correlation analysis. Correlation coefficients calculated for each separate gravity core (Table S2) indicate

that alkalinity is strongly negatively correlated ($p \geq -0.85$) with sulfate concentrations in sediments from HC-affected sites. The same applies to quantifiable sulfate and sulfide concentrations ($p \geq -0.7$). For reference sites, alkalinity and sulfate concentrations are also strongly correlated ($p \approx 0.8$), whereas sulfide concentrations are below the detection limit and hence have been omitted for the correlation analysis.

We established ratios of sulfate to alkalinity (HCO_3^-) fluxes (Fig. 5). As about 90% of seawater alkalinity can be contributed to bicarbonate (89.8% HCO_3^- , 2.9 % CO_3^{2-}) (Kerr et al., 2021) we set alkalinity synonymous with bicarbonate concentration. We compared the measured fluxes to those expected from organoclastic (1:2 based on Eq. 1) and methanotrophic (1:1 based on Eq. 2) SRR activities. At reference sites, the average ratio of sulfate to alkalinity fluxes equals 2:1 whereas it is 1.25:1 (5:4) at HC-affected sites, imply that either HCO_3^- is being removed from the pore water or that pore water sulfate is eventually being replenished via sulfide oxidation.





Figure 5. Relationship between sulfate and alkalinity fluxes at HC-affected sites and reference sites. The majority of samples from HC-affected sites (dots) have lower alkalinity than sulfate fluxes. This is even more pronounced at reference sites (triangles). Black lines signify theoretical ratios for methanotrophic (1:1) and organoclastic (1:2) SR.

3.5 Functional marker genes specific to the SMTZ

From the 12 sequenced metagenomes, we extracted a total of 307 234 ORFs with multiple annotations that corresponded to 821 850 predicted genes. From the 5 sequenced metatranscriptomes, we extracted a total of 31 093 ORFs corresponding to 47 316 predicted genes (Supplementary Table S2).



425 **Figure 6. Open reading frames encoding functional marker genes involved in sulfate reduction, production and anaerobic consumption of methane and their taxonomic assignments at the phylum to class level.** The relative abundances of certain phyla increase at HC-affected sites compared to reference sites. Concomitant expression of *apr* and *dsr* genes by Desulfobacterota, *phs* genes by Chloroflexota, *hyd* genes by Asgardarchaeota, and *mcr* genes by ANMEs and Halobacterota are consistent with
 430 methanotrophic sulfate-reducing activities. **Abbreviations** - *apr*: adenylylsulfate reductase; *dsr*: dissimilatory sulfite reductase; *asr*: anaerobic sulfite reductase; *phs/psr*: thiosulfate and polysulfide reductase; *hyd*: sulfhydrogenase; *mcr*: methyl coenzyme M reductase; MG: Metagenomics; MT: Metatranscriptomics; BDL: below detection limit.

435 For each of the targeted marker genes, we plotted the relative percentage of total predicted genes and their corresponding taxonomic assignments at the phylum and class level (Fig. 6). Taxonomic assignments of *SSU* rRNA and *FtsAZ* proteins show that metabolically active and growing microbial populations at HC-affected sites are affiliated with the phyla Chloroflexota, Desulfobacterota and Euryarchaeota, and more specifically with anaerobic methanotrophs (i.e. ANMEs). The phyla
 440 Desulfobacterota and Chloroflexota are actively engaged in processes of sulfate reduction, as shown by the expression of ORFs encoding the *aprAB* and *dsrAB* genes (Fig. 7a-7b). Taxonomic assignments of expressed ORFs encoding *mcrA* genes show an increased abundance of methanotrophic ANMEs (Fig. 7c) over methanogenic Halobacterota. ORFs encoding *hydB-G* genes are expressed by taxa related to the class Lokiarchaeia and Thorarchaeia (Fig. 7d) among the phylum Asgardarchaeota, indicating metabolic
 445 capability to use reduced sulfur compounds (e.g. polysulfide, elemental sulfur). However, their overall expression levels appear limited and these Asgardarchaeota do not seem to actively divide (Fig. 6). At

reference sites, Chloroflexota appear to be the only phylum capable of growing and dividing in these organic-poor sediments (Fig. 6). However, concomitant expression of functional marker genes involved in sulfate reduction remains below the limit of detection. We acknowledge that the non-detection of certain genes may also result from limited sequencing depths.

Phylogenetic trees plotted for transcript ORFs assigned to the *rps3* and *rpoD* taxonomic marker genes (Supplementary Figs. S4-S5) provided concomitant evidence for metabolic activity by the phyla Desulfobacterota, Chloroflexota (class Dehalococcoidia), Acidobacterota (class Aminicenentia), Asgardarchaeota (class Thorarchaeia), and Halobacterota (class ANME-1).

455




Figure 7. Phylogenetic trees of *aprAB*, *dsrAB*, *mcrA* and *hydB-G* protein-encoding genes. PhyML maximum likelihood trees of transcript ORFs encoding: **(a)** the adenylylsulfate reductase (714 amino acids) subunit alpha (*aprA*) and beta (*aprB*); **(b)** the dissimilatory sulfite reductase (493 amino acids) subunit alpha (*dsrA*) and beta (*dsrB*); **(c)** the methyl coenzyme M reductase reductase (698 amino acids) subunit alpha (*mcrA*); and **(d)** the sulfhydrogenase, or sulfur reductase (286 amino acids), subunit beta (*hydB*) and gamma (*hydG*). Phylogenetic trees are based on 100 bootstrap replicates with BLOSUM62 as evolutionary model. Boldface types signify sequence accession numbers to the NCBI database.

465 4. Discussion

4.1 Pore water geochemical profiles correlate to meta-omics features

Pore water profiles reflect the spatial distribution of net turnover processes. The shape of the profile allows to distinguish intervals in which transport is purely diffusive (i.e. linear profile) from those in which there is either net production (i.e. convex profile) or consumption (i.e. concave profile) of chemical species
470 (Schulz, 2006; Hesse and Schacht, 2011).

Based on the largely linear profiles for pore water alkalinity, sulfate and sulfide (Figs. 2-3, Figs. S1-S2), we infer the absence or near-absence of net sulfate consumption in the depth interval covered by the cores, and a sink for sulfate as well as a source of sulfide and alkalinity at greater depth. Microbially-mediated AOM with sulfate as electron acceptor yields hydrogen sulfide, bicarbonate and water (Eq. 2),
475 providing an explanation for both sink and source (Martens and Berner, 1974; Iversen and Jorgensen, 1985), assuming sufficient supply of methane at greater depth. The occurrence of AOM at deeper depth is further supported by the relationship between sulfate and alkalinity fluxes (Fig. 5) and negligibly small methane concentrations ($< 10 \mu\text{M L}^{-1}$) at the bottom end of all cores from HC-affected sites and reference sites (Fig. 4), indicating almost complete consumption of methane originating from biogenic or
480 thermogenic sources below our sampling interval.

Linear concentration profiles are only obtained in regions with no net consumption or production of the respective compound (Schulz, 2006). Despite the apparent linearity of sulfate profiles (Fig. 2), we observed very low SRR ($\leq 300 \text{ pmol} \times \text{cm}^{-3} \times \text{d}^{-1}$), confirming previous reports of sporadic SRR in Barents Sea sediments with similar rates above the SMTZ (Nickel et al., 2012). In our case, sulfate reduction is
485 mainly detectable at HC-affected sites (Fig. 3), with a greater prevalence in deeper sediments (Fig. 5). With seepage supplying electron donors, like methane and/or other HCs, from deeper reservoirs to near-surface sediment, sulfate reduction is fuelled by those HCs rather than from the buried and more recalcitrant OM. Inconspicuous supplies of methane and/or other HCs to the overlying sediment can stimulate SRR by providing alternative electron donors. Such sporadic energy supply promotes geochemical conditions that
490 stimulate benthic HC-degrading microbial communities. Although these communities can be physiologically and phylogenetically diverse, they are mainly composed of anaerobic guilds of methanogens, methanotrophs, sulfate reducers and fermenters (Teske, 2019).

The analysis of functional marker genes supports the notion of sporadic HC supply, providing complementary insights into the metabolic potential and activity of specific microbial populations in the

495 sediment. The occurrence and abundance of *aprAB*, *dsrAB*, and *mcrA* genes supports interpretations of
geochemical data in terms of sulfate-driven AOM processes and metabolic activity (Fig. 6). Taxonomic
profiling of populations expressing ORFs encoding genes involved in sulfate reduction (i.e. *aprAB*, *dsrAB*)
reveals Desulfobacterota (family Desulfobacteraceae) and Chloroflexota (class Dehalococcoidia) as the
prevalent bacterial phyla constituting the consortium of sulfate-reducing bacteria at HC-affected sites (Fig.
500 7, Supplementary Figs. S4-S5). The *aprAB* and *dsrAB* phylogenetic trees clearly show that none of these
genes correspond to their reverse function and the ORFs annotated as *aprAB* and *dsrAB* are predicted to
function in dissimilatory sulfate reduction. Moreover, the sediment being fully anoxic, any detection of
sulfur oxidation via reversible *aprAB* or reverse *dsrAB* was unexpected as such activities are mostly
reported for aerobic Gammaproteobacteria (Vuillemin et al., 2022).

505 Further, ORFs encoding genes involved in anaerobic production and consumption of methane (i.e.
mcrA) clearly show that the clade ANME-I exhibits concomitant, although low metabolic activity towards
anaerobic oxidation of methane (Fig. 7c), namely they express ORFs encoding the reversible *mcrA* gene.
The higher expression levels of methanotrophic clade ANME-I over methanogenic Halobacterota (Fig. 6)
also suggest that metabolic activity by ANMEs could be sustained by an alternative source of methane
510 (Dong et al., 2020), although cryptic production of biogenic methane by the ANME-1 clade has also been
observed in experiments replicating SMTZ conditions (Beulig et al., 2019). Because methane
concentrations measured at both HC-affected sites and reference sites were similarly low (Fig. 4), it appears
unlikely that AOM processes are sustained by in situ production of biogenic methane only (Dong et al.,
2020). Finally, the class Lokiarchaeia and Thorarchaeia in the phylum Asgardarchaeota expressed ORFs
515 encoding beta and gamma subunits of *hydB-G* genes (Fig. 7d), and thereby appear metabolically active,
using various sulfur and sulfide species, potentially produced as by-products of sulfate reduction.

In contrast, Chloroflexota (class Dehalococcoidia) was the only phylum at reference sites showing
sufficient metabolic activity was adequate to achieve cell division (i.e. FstAZ). In the absence of an univocal
signal towards active dissimilatory sulfate reduction by Dehalococcoidia at HC-affected sites (Fig. 7b), we
520 infer that these Chloroflexota could persist and gain energy through fermentation processes in organic-lean
marine sediments instead (Vuillemin et al., 2020a). Due to this metabolic versatility, Chloroflexota were
found to be active at both HC-affected and reference sites, which rank them as poor microbial indicators for
minor HC seepage (Iasakov et al., 2022).

Together, these microbial patterns extracted from meta-omics corroborate the geochemical data,
525 arguing in favor of discrete AOM processes concomitant to sulfate reduction at HC-affected sites, and for

limited in situ production of biogenic methane at both HC-affected sites and reference sites. Alongside an active AOM consortium composed of Desulfobacterota and ANME-1, concomitant production of reduced sulfur compounds could apparently stimulate metabolic activity by Lokiarchaeia and Thorarchaeia (Figs. 6 and 7).

530 **4.2 Geochemical indicators for minor HC seepage**

The combined use of meta-omics microbial patterns with the determination of SR and related geochemical features and fluxes in upper sediments, enables to a certain extent to distinguish between HC-affected sites and reference sites. Sediments of the SW Barents Sea are organic-poor (TOC <0.5 %), and thus characterized by a lack of electron donors (Knies and Martinez, 2009) and low microbial activity (Nickel et al., 2012; Nickel et al., 2013) with consequent sulfate penetration to greater sediment depths. In organic-rich sediments, any signal inherent to minor HC seepage fueling SR would be difficult to detect as it would be covered by organoclastic SR processes. The organic-lean nature of Barents Sea sediments should enable the detection of discrete HC seeps, because the slightest supply of electron donors leads to a direct increase in the metabolic activity of HC-degrading microbial populations. This slight increase in metabolic activity
540 can then be discriminated from the background signal observed at reference sites. Diffusion of HCs was thus expected to promote sulfate consumption, resulting in its depletion in the pore water at shallower sediment depths. This, combined with variable fluxes of thermogenic and biogenic methane, can account for variations in the depth of the SMTZ.

At different locations of the SW Barents Sea, the depth of the SMTZ, or rather the sulfate depletion depth, was postulated, for instance, at 37 mbsf in the Hammerfest/Loppa High and the Tromsø Basin/Ingøydjupet area (Nickel et al., 2012), and between 3.5 and 29.2 mbsf in the Ingøydjupet area (Argentino et al., 2021a). These values are consistent with the global average depth of the SMTZ (12.8 ± 12.1 mbsf) on the continental slope (Egger et al., 2018). For comparison, at the Vestnesa Ridge which is under the influence of gas hydrates, the depth of the SMTZ fluctuates massively between seepage (ca. 1
550 mbsf) and non-seepage (ca. 3 to 75 mbsf) sites (Hong et al., 2016). Similarly, Barents Sea sediments with and without methane seeps show an SMTZ located between a few cm and about 2 mbsf (Argentino et al., 2021b).

Linear extrapolation of sulfate profiles (Fig. 2) indicates complete sulfate depletion, indicative of an SMTZ, at 12 and 3 mbsf for reference and HC-affected sites, respectively. We suggest that the
555 difference in sulfate penetration depth is caused by upwards migration of HCs. The sulfate and alkalinity fluxes inherent to organoclastic (Eq. 1) and methanotrophic (Eq. 2) SR have respective theoretical ratios of

1:2 and 1:1 while mixed SR processes have intermediate values. Current ratios of sulfate to alkalinity fluxes (Fig. 5) at reference sites imply that either HCO_3^- is being removed from the pore water, or that pore water sulfate is eventually being replenished via sulfide oxidation. From other fieldwork in the area, we know that the sediment is fully anoxic below the uppermost few cm. Phylogenetic analysis of *aprAB* and *dsrAB* genes clearly shows that none of the identified amino acid sequences correspond to their reverse function. Further, we did not identify any *dsrC* sequences, which potentially work in cellular sulfur oxidation (Ruiz-Blas et al., 2024). While dissolved inorganic carbon (DIC) can be assimilated via acetogenesis, methanogenesis, or methanotrophy (Kellermann et al., 2012), authigenic carbonate minerals are known to precipitate from the pore water as by-products of AOM processes (Turchyn et al., 2021), sometimes even forming carbonate hardground at the seabed. However, from the low level of detection of AOM-related transcripts, we inferred only a minor imprint on the geochemistry and therefore assume that bicarbonate ions originating from AOM processes at greater depths are diffusing upward and eventually precipitate as calcium carbonate. Up to 80 % of the sulfide produced through SR can also be converted back into sulfate through reoxidation of reduced sulfur compounds (Jørgensen et al., 2004) and disproportionation (Pellerin et al., 2015; Jørgensen et al., 2019a). Finally, the dissolution of sulfate minerals, such as barite (BaSO_4), may also act as an additional source of sulfate (Griffith and Paytan, 2012). Altogether, this suggests either the reoxidation of sulfide, or the removal of bicarbonate from the pore water, e.g. by mineral precipitation, or both. The former is further supported by concomitant sulfate reduction, detected by radiotracer incubations (Fig. 3) and microbial gene expression patterns (Fig. 6), the latter by the ratios of sulfate to alkalinity flux (Fig. 5). Thus, we could identify several biogeochemical features that argue for ongoing sulfate reduction (most likely AOM-related) at HC-affected sites below our sampled depth range, which allow their discrimination from reference sites. Given the sulfate concentration profiles and the small amounts of methane detected in the samples, it becomes clear that AOM is not a quantitatively important process in the cored depth interval, producing an erratic signal in HC-affected sediments.

5. Conclusions

Our study, combining modeled and measured SRR as well as pore water geochemistry with functional marker genes, revealed remarkable differences between samples originating from a pristine seafloor and

those from areas affected by small and inconspicuous HC seepage. The linear shape of pore water profiles was indicative of purely diffusive transport under steady-state conditions and allowed estimation of the depth of sulfate depletion, which appeared markedly different between HC-affected and reference sites. At the HC-affected sites, we found higher pore water fluxes of sulfate, sulfide and alkalinity, and could also demonstrate active microbial SR by turnover measurements and modeling. Furthermore, meta-omics provided additional evidence for enhanced metabolic activity related to microbial sulfate reduction coupled with minor AOM processes at HC-affected sites, which could be clearly discriminated from the background signal inherent to fermentative activities and minor *in situ* production of methane by the subseafloor biosphere. These observations demonstrate that minor HC seepage, which is not even visible at the seafloor, significantly influences sedimentary biogeochemical cycles driven by microbial populations resident in organic-lean sediments.

6. Data availability

Geochemical data, ORFs sequences and their BlastP outputs are available as Supplementary Information. All scripts, code and instructions regarding how to conduct the BLASTp workflow for taxonomic and functional annotations are posted on GitHub with a link to the MetaProt page (github.com/williamorsi/MetaProt-database).

7. Author contributions

JK and RDP organized the sampling cruise. ES and JK designed the study and sampled the cores. ES performed sulfate, alkalinity, SRR and gas analyses. AV, BK, CL and AS performed DNA and RNA analyses. ES and AV wrote the manuscript and all authors contributed to writing and revision.

8. Competing interests

The authors declare that they have no conflict of interest.

9. Acknowledgments

We thank all crew members of the SV *Sverdrup*, as well as Steffen Okolski, Jan Axel Kitte and Edgar
610 Kutschera for their great help during the sampling cruise. The help of Steffen Okolski in conducting the ion
chromatography, alkalinity and sulfide analyses and Simone Bernsee during radioisotope experiments is
acknowledged.

10. Financial support

This research has received funding from the European Union's Horizon 2020 research and innovation
615 programme under grant agreement no. 899667.

PROSPECTOMICS Consortium

The principal investigators of the PROSPECTOMICS project are Jens Kallmeyer¹, Paul Wilmes²,
Alexander J. Probst³, Dörte Becher⁴, Thomas Rattei⁵, and Rolando di Primio⁶. The project managers are
620 Aurèle Vuillemin¹, Cédric C. Laczny², André R. Soares³, and Anke Trautwein-Schult⁴. Scientists and
technicians include Ellen Schnabel¹, Kai Mangelsdorf⁷, Steffen Okolski¹, J. Axel Kitte¹, Benoit J. Kunath²,
Oskar Hickl², Tuesday Lowndes², Zainab Zafar², Sarah Esser³, Anne Ostrzinski⁴, Sebastian Grund⁴, and
Alexander Pfundner⁵.

¹ GFZ German Research Centre for Geosciences, Section Geomicrobiology, Telegrafenberg, 14473
625 Potsdam, Germany; ²Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-
Alzette; ³Environmental Metagenomics, Research Center One Health Ruhr of the University Alliance Ruhr,
Faculty of Chemistry, University of Duisburg-Essen, Essen, Germany; ⁴Department of Microbial
Proteomics, University of Greifswald, Greifswald, Germany; ⁵Computational Systems Biology, Centre for
Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria; ⁶Aker BP ASA,
630 Sandvika, Viken, Norway; ⁷GFZ German Research Centre for Geosciences, Section Organic Geochemistry,
Telegrafenberg, 14473 Potsdam, Germany.

11. References

Abrams, M. A.: Marine seepage variability and its impact on evaluating the surface migrated hydrocarbon
seep signal, *Marine and Petroleum Geology*, 121, 10.1016/j.marpetgeo.2020.104600, 2020.

- 635 Argentino, C., Waghorn, K. A., Bünz, S., and Panieri, G.: Sulfate reduction and anaerobic oxidation of methane in sediments of the South-Western Barents Sea, *Biogeoscience*, 10.5194/bg-2021-58, 2021a.
 Argentino, C., Waghorn, K. A., Vadakkepuliyambatta, S., Polteau, S., Bunz, S., and Panieri, G.: Dynamic and history of methane seepage in the SW Barents Sea: new insights from Leirdjupet Fault Complex, *Sci Rep*, 11, 4373, 10.1038/s41598-021-83542-0, 2021b.
- 640 Berg, P., Risgaard-Petersen, N., and Rysgaard, S.: Interpretation of measured concentration profiles in sediment pore water, *Limnol Oceanogr*, 43, 1500-1510, 10.4319/lo.1998.43.7.1500, 1998.
 Beulig, F., Roy, H., McGlynn, S. E., and Jorgensen, B. B.: Cryptic CH₄ cycling in the sulfate-methane transition of marine sediments apparently mediated by ANME-1 archaea, *ISME J*, 13, 250-262, 10.1038/s41396-018-0273-z, 2019.
- 645 Bolger, A. M., Lohse, M., and Usadel, B.: Trimmomatic: a flexible trimmer for Illumina sequence data, *Bioinformatics*, 30, 2114-2120, 10.1093/bioinformatics/btu170, 2014.
 Bornemann, T. L. V., Esser, S. P., Stach, T. L., Burg, T., and Probst, A. J.: uBin: A manual refining tool for genomes from metagenomes, *Environ Microbiol*, 25, 1077-1083, 10.1111/1462-2920.16351, 2023.
 Borowski, W. S., Paull, C. K., and Ussler, W.: Marine pore-water sulfate profiles indicate in situ methane flux from underlying gas hydrate, *Geology*, 24, 655-658, 10.1130/0091-7613(1996)024<0655:Mpwspi>2.3.Co;2, 1996.
- 650 Borowski, W. S., Paull, C. K., and Ussler, W.: Global and local variations of interstitial sulfate gradients in deep-water, continental margin sediments: Sensitivity to underlying methane and gas hydrates, *Marine Geology*, 159, 131-154, 10.1016/S0025-3227(99)00004-3, 1999.
- 655 Bowles, M. W., Mogollon, J. M., Kasten, S., Zabel, M., and Hinrichs, K. U.: Global rates of marine sulfate reduction and implications for sub-sea-floor metabolic activities, *Science*, 344, 889-891, 10.1126/science.1249213, 2014.
 Buchfink, B., Xie, C., and Huson, D. H.: Fast and sensitive protein alignment using DIAMOND, *Nat Methods*, 12, 59-60, 10.1038/nmeth.3176, 2015.
- 660 Castresana, J.: Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis, *Mol Biol Evol*, 17, 540-552, 10.1093/oxfordjournals.molbev.a026334, 2000.
 Chen, C., Wu, X., Wan, Z., Shang, J., Huang, W., Zhang, W., Liang, J., Xiao, Z., Zhou, W., and Zhong, L.: Geochemical characteristics of sediment and pore water affected by cold seeps in southern South China Sea, *Frontiers in Marine Science*, 10, 10.3389/fmars.2023.1167578, 2023.
- 665 Ciotoli, G., Procesi, M., Etiope, G., Fracassi, U., and Ventura, G.: Influence of tectonics on global scale distribution of geological methane emissions, *Nat Commun*, 11, 2305, 10.1038/s41467-020-16229-1, 2020.
 Cline, J. D.: Spectrophotometric Determination of Hydrogen Sulfide in Natural Waters¹, *Limnol Oceanogr*, 14, 454-458, 10.4319/lo.1969.14.3.0454, 1969.
- 670 Cramm, M. A., Neves, B. M., Manning, C. C. M., Oldenburg, T. B. P., Archambault, P., Chakraborty, A., Cyr-Parent, A., Edinger, E. N., Jaggi, A., Mort, A., Tortell, P., and Hubert, C. R. J.: Characterization of marine microbial communities around an Arctic seabed hydrocarbon seep at Scott Inlet, Baffin Bay, *Sci Total Environ*, 762, 143961, 10.1016/j.scitotenv.2020.143961, 2021.
 D'Hondt, S., Jorgensen, B. B., Miller, D. J., Batzke, A., Blake, R., Cragg, B. A., Cypionka, H., Dickens, G. R., Ferdelman, T., Hinrichs, K. U., Holm, N. G., Mitterer, R., Spivack, A., Wang, G., Bekins, B., Engelen, B., Ford, K., Gettemy, G., Rutherford, S. D., Sass, H., Skilbeck, C. G., Aiello, I. W., Guerin, G., House, C. H., Inagaki, F., Meister, P., Naehr, T., Niitsuma, S., Parkes, R. J., Schippers, A., Smith, D. C., Teske, A., Wiegand, J., Padilla, C. N., and Acosta, J. L.: Distributions of microbial activities in deep seafloor sediments, *Science*, 306, 2216-2221, 10.1126/science.1101155, 2004.

- 680 Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J. F., Guindon, S., Lefort, V., Lescot, M., Claverie, J. M., and Gascuel, O.: Phylogeny.fr: robust phylogenetic analysis for the non-specialist, *Nucleic Acids Res*, 36, W465-469, 10.1093/nar/gkn180, 2008.
- Dong, X., Rattray, J. E., Campbell, D. C., Webb, J., Chakraborty, A., Adebayo, O., Matthews, S., Li, C., Fowler, M., Morrison, N. M., MacDonald, A., Groves, R. A., Lewis, I. A., Wang, S. H., Mayumi, D., Greening, C., and Hubert, C. R. J.: Thermogenic hydrocarbon biodegradation by diverse depth-stratified microbial populations at a Scotian Basin cold seep, *Nat Commun*, 11, 5825, 10.1038/s41467-020-19648-2, 2020.
- Doré, A. G.: Barents Sea Geology, Petroleum Resources and Commercial Potencial, *ARCTIC*, 48, 1995.
- Edenborn, H. M., Paquin, Y., and Chateauneuf, G.: Bacterial contribution to manganese oxidation in a deep coastal sediment, *Estuarine, Coastal and Shelf Science*, 21, 801-815, 10.1016/0272-7714(85)90074-5, 1985.
- 690 Edgar, R. C.: MUSCLE: multiple sequence alignment with high accuracy and high throughput, *Nucleic Acids Res*, 32, 1792-1797, 10.1093/nar/gkh340, 2004.
- Egger, M., Riedinger, N., Mogollón, J. M., and Jørgensen, B. B.: Global diffusive fluxes of methane in marine sediments, *Nature Geoscience*, 11, 421-425, 10.1038/s41561-018-0122-8, 2018.
- 695 Elverhøi, A. and Solheim, A.: The Barents Sea ice sheet - a sedimentological discussion, *Polar Research*, 1, 23-42, 10.1111/j.1751-8369.1983.tb00729.x, 1983.
- EMODnet project (European Commission): https://erddap.emodnet.eu/erddap/files/bathymetry_2022/, last access: 27-nov-2024.
- Firringioli, A., Negroni, A., Zanaroli, G., and Cappelletti, M.: Unraveling the Metabolic Potential of Asgardarchaeota in a Sediment from the Mediterranean Hydrocarbon-Contaminated Water Basin Mar Piccolo (Taranto, Italy), *Microorganisms*, 9, 10.3390/microorganisms9040859, 2021.
- Fossing, H., Ferdelman, T. G., and Berg, P.: Sulfate reduction and methane oxidation in continental margin sediments influenced by irrigation (South-East Atlantic off Namibia), *Geochimica Et Cosmochimica Acta*, 64, 897-910, Doi 10.1016/S0016-7037(99)00349-X, 2000.
- 705 Gabrielsen, R. H., Faerseth, R. B., Jensen, L. N., Kalheim, J. E., and Riis, F.: Structural elements of the Norwegian continental shelf, Part 1: The Barents Sea Region, *NPD Bulletin*, 6, 1-33, 1990.
- Gouy, M., Guindon, S., and Gascuel, O.: SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building, *Mol Biol Evol*, 27, 221-224, 10.1093/molbev/msp259, 2010.
- 710 Griffith, E. M. and Paytan, A.: Barite in the ocean – occurrence, geochemistry and palaeoceanographic applications, *Sedimentology*, 59, 1817-1835, 10.1111/j.1365-3091.2012.01327.x, 2012.
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O.: New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0, *Syst Biol*, 59, 307-321, 10.1093/sysbio/syq010, 2010.
- 715 Guseva, N., Moiseeva, Y., Purgina, D., Gershelis, E., Yakushev, E., and Semiletov, I.: The Impact of Methane Seepage on the Pore-Water Geochemistry across the East Siberian Arctic Shelf, *Water*, 13, 10.3390/w13040397, 2021.
- Heggland, R.: Gas seepage as an indicator of deeper prospective reservoirs. A study based on exploration 3D seismic data, *Marine and Petroleum Geology*, 15, 1-9, 10.1016/S0264-8172(97)00060-3, 1998.
- 720 Henrichs, S. M. and Reeburgh, W. S.: Anaerobic mineralization of marine sediment organic matter: Rates and the role of anaerobic processes in the oceanic carbon economy, *Geomicrobiology Journal*, 5, 191-237, 10.1080/01490458709385971, 1987.

- Hesse, R. and Schacht, U.: Early Diagenesis of Deep-Sea Sediments, in: Deep-Sea Sediments, edited by: Hüneke, H., and Mulder, T., Developments in Sedimentology, Elsevier, 557-713, 10.1016/b978-0-444-53000-4.00009-3, 2011.
- 725 Hong, W.-L., Sauer, S., Panieri, G., Ambrose, W. G., James, R. H., Plaza-Faverola, A., and Schneider, A.: Removal of methane through hydrological, microbial, and geochemical processes in the shallow sediments of pockmarks along eastern Vestnesa Ridge (Svalbard), *Limnol Oceanogr*, 61, S324-S343, 10.1002/lno.10299, 2016.
- 730 Hu, C.-Y., Frank Yang, T., Burr, G. S., Chuang, P.-C., Chen, H.-W., Walia, M., Chen, N.-C., Huang, Y.-C., Lin, S., Wang, Y., Chung, S.-H., Huang, C.-D., and Chen, C.-H.: Biogeochemical cycles at the sulfate-methane transition zone (SMTZ) and geochemical characteristics of the pore fluids offshore southwestern Taiwan, *Journal of Asian Earth Sciences*, 149, 172-183, 10.1016/j.jseae.2017.07.002, 2017.
- Hunt, J. M.: *Petroleum Geochemistry and Geology*, Freeman & co, New York 1995.
- 735 Hvoslef, S., Christie, O. H. J., Sassen, R., Kennicutt, M. C., Requejo, A. G., and Brooks, J. M.: Test of a new surface geochemistry tool for resource prediction in frontier areas, *Marine and Petroleum Geology*, 13, 107-124, Doi 10.1016/0264-8172(95)00032-1, 1996.
- Hyatt, D., Chen, G. L., Locascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J.: Prodigal: prokaryotic gene recognition and translation initiation site identification, *BMC Bioinformatics*, 11, 119, 10.1186/1471-2105-11-119, 2010.
- 740 Iasakov, T. R., Kanapatskiy, T. A., Toshchakov, S. V., Korzhenkov, A. A., Ulyanova, M. O., and Pimenov, N. V.: The Baltic Sea methane pockmark microbiome: The new insights into the patterns of relative abundance and ANME niche separation, *Mar Environ Res*, 173, 105533, 10.1016/j.marenvres.2021.105533, 2022.
- 745 Iversen, N. and Jørgensen, B. B.: Anaerobic methane oxidation rates at the sulfate-methane transition in marine sediments from Kattegat and Skagerrak (Denmark)1, *Limnol Oceanogr*, 30, 944-955, 10.4319/lno.1985.30.5.0944, 1985.
- Iversen, N. and Jørgensen, B. B.: Diffusion coefficients of sulfate and methane in marine sediments: Influence of porosity, *Geochemica et Cosmochemica Acta*, 57, 571-778, 10.1016/0016-7037(93)90368-7, 1993.
- 750 Jang, J., Cao, S. C., Stern, L. A., Jung, J., and Waite, W. F.: Impact of Pore Fluid Chemistry on Fine-Grained Sediment Fabric and Compressibility, *Journal of Geophysical Research: Solid Earth*, 123, 5495-5514, 10.1029/2018jb015872, 2018.
- Jiang, S.: Clay Minerals from the Perspective of Oil and Gas Exploration, in: *Clay Minerals in Nature - Their Characterization, Modification and Application*, 10.5772/47790, 2012.
- 755 Johansen, S. E., Ostisty, B. K., Birkeland, Ø., Fedorovsky, Y. F., Martirosjan, V. N., Bruun Christensen, O., Cheredeev, S. I., Ignatenko, E. A., and Margulis, L. S.: Hydrocarbon potential in the Barents Sea region: play distribution and potential, *Norwegian Petroleum Society Special Publications*, 2, 273-320, 10.1016/B978-0-444-88943-0.50024-1, 1993.
- 760 Jørgensen, B. B.: A comparison of methods for the quantification of bacterial sulfate reduction in coastal marine sediments, *Geomicrobiology Journal*, 1, 11-27, 10.1080/01490457809377721, 1978.
- Jørgensen, B. B.: Mineralization of organic matter in the sea bed—the role of sulphate reduction, *Nature*, 296, 643-645, 10.1038/296643a0, 1982.
- Jørgensen, B. B. and Kastan, S.: Sulfur Cycling and Methane Oxidation, in: *Marine Geochemistry*, 271-309, 10.1007/3-540-32144-6_8, 2006.
- 765 Jørgensen, B. B., Egger, M., and Canfield, D. E.: Sulfate distribution and sulfate reduction in global marine sediments, *Geochimica et Cosmochimica Acta*, 364, 79-88, 10.1016/j.gca.2023.11.015, 2024.

- Jørgensen, B. B., Findlay, A. J., and Pellerin, A.: The Biogeochemical Sulfur Cycle of Marine Sediments, *Front Microbiol*, 10, 849, 10.3389/fmicb.2019.00849, 2019a.
- 770 Jørgensen, B. B., Nelson, D. C., Amend, J. P., Edwards, K. J., and Lyons, T. W.: Sulfide oxidation in marine sediments: Geochemistry meets microbiology, in: *Sulfur Biogeochemistry - Past and Present*, Geological Society of America, 0, 10.1130/0-8137-2379-5.63, 2004.
- Jørgensen, B. B., Beulig, F., Egger, M., Petro, C., Scholze, C., and Røy, H.: Organoclastic sulfate reduction in the sulfate-methane transition of marine sediments, *Geochimica et Cosmochimica Acta*, 254, 231-245, 10.1016/j.gca.2019.03.016, 2019b.
- 775 Joye, S. B.: The Geology and Biogeochemistry of Hydrocarbon Seeps, *Annual Review of Earth and Planetary Sciences*, 48, 205-231, 10.1146/annurev-earth-063016-020052, 2020.
- Joye, S. B., Boetius, A., Orcutt, B. N., Montoya, J. P., Schulz, H. N., Erickson, M. J., and Lugo, S. K.: The anaerobic oxidation of methane and sulfate reduction in sediments from Gulf of Mexico cold seeps, *780 Chemical Geology*, 205, 219-238, 10.1016/j.chemgeo.2003.12.019, 2004.
- Kallmeyer, J., Ferdelman, T. G., Weber, A., Fossing, H., and Jørgensen, B. B.: A cold chromium distillation procedure for radiolabeled sulfide applied to sulfate reduction measurements, *Limnol Oceanogr-Meth*, 2, 171-180, DOI 10.4319/lom.2004.2.171, 2004.
- 785 Kasten, S. and Jørgensen, B. B.: Sulfate Reduction in Marine Sediments, in: *Marine Geochemistry*, Springer, Heidelberg, 263-281, 10.1007/978-3-662-04242-7_8, 2000.
- Kellermann, M. Y., Wegener, G., Elvert, M., Yoshinaga, M. Y., Lin, Y. S., Holler, T., Mollar, X. P., Knittel, K., and Hinrichs, K. U.: Autotrophy as a predominant mode of carbon fixation in anaerobic methane-oxidizing microbial communities, *Proc Natl Acad Sci U S A*, 109, 19321-19326, 10.1073/pnas.1208795109, 2012.
- 790 Kerr, D. E., Brown, P. J., Grey, A., and Kelleher, B. P.: The influence of organic alkalinity on the carbonate system in coastal waters, *Marine Chemistry*, 237, 10.1016/j.marchem.2021.104050, 2021.
- Kim, J., Dong, H. L., Seabaugh, J., Newell, S. W., and Eberl, D. D.: Role of microbes in the smectite-to-illite reaction, *Science*, 303, 830-832, DOI 10.1126/science.1093245, 2004.
- 795 Kleindienst, S., Herbst, F. A., Stagars, M., von Netzer, F., von Bergen, M., Seifert, J., Peplies, J., Amann, R., Musat, F., Lueders, T., and Knittel, K.: Diverse sulfate-reducing bacteria of the *Desulfosarcina/Desulfococcus* clade are the key alkane degraders at marine seeps, *ISME J*, 8, 2029-2044, 10.1038/ismej.2014.51, 2014.
- Knies, J. and Martinez, P.: Organic matter sedimentation in the western Barents Sea region: Terrestrial and marine contribution based on isotopic composition and organic nitrogen content, *Norwegian Journal of 800 Geology*, 89, 79-89, 2009.
- Larssen, G. B., Elvebakk, G., Henriksen, L. B., Kristensen, S.-E., Nilsson, I., Samuelsen, T. J., Svana, T. A., Stemmerik, L., and Worsley, D.: Upper Palaeozoic lithostratigraphy of the Southern Norwegian Barents Sea, in, 2002.
- Li, H. and Durbin, R.: Fast and accurate short read alignment with Burrows-Wheeler transform, *805 Bioinformatics*, 25, 1754-1760, 10.1093/bioinformatics/btp324, 2009.
- Liao, Y., Smyth, G. K., and Shi, W.: featureCounts: an efficient general purpose program for assigning sequence reads to genomic features, *Bioinformatics*, 30, 923-930, 10.1093/bioinformatics/btt656, 2014.
- MacLeod, F., Kindler, G. S., Wong, H. L., Chen, R., and Burns, B. P.: Asgard archaea: Diversity, function, and evolutionary implications in a range of microbiomes, *AIMS Microbiol*, 5, 48-61, 10.3934/microbiol.2019.1.48, 2019.
- 810 Martens, C. S. and Berner, R. A.: Methane production in the interstitial waters of sulfate-depleted marine sediments, *Science*, 185, 1167-1169, 10.1126/science.185.4157.1167, 1974.

- März, C., Hoffmann, J., Bleil, U., de Lange, G. J., and Kasten, S.: Diagenetic changes of magnetic and geochemical signals by anaerobic methane oxidation in sediments of the Zambezi deep-sea fan (SW Indian Ocean), *Marine Geology*, 255, 118-130, 10.1016/j.margeo.2008.05.013, 2008.
- 815 Middelburg, J. J.: Reviews and syntheses: to the bottom of carbon processing at the seafloor, *Biogeosciences*, 15, 413-427, 10.5194/bg-15-413-2018, 2018.
- Mölder, F., Jablonski, K. P., Letcher, B., Hall, M. B., Tomkins-Tinch, C. H., Sochat, V., Forster, J., Lee, S., Twardziok, S. O., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen, S., and Koster, J.: Sustainable data analysis with Snakemake, *F1000Res*, 10, 33, 10.12688/f1000research.29032.2, 2021.
- 820 Nickel, J. C., di Primio, R., Mangelsdorf, K., Stoddart, D., and Kallmeyer, J.: Characterization of microbial activity in pockmark fields of the SW-Barents Sea, *Marine Geology*, 332-334, 152-162, 10.1016/j.margeo.2012.02.002, 2012.
- Nickel, J. C., di Primio, R., Kallmeyer, J., Hammer, Ø., Horsfield, B., Stoddart, D., Brunstad, H., and Mangelsdorf, K.: Tracing the origin of thermogenic hydrocarbon signals in pockmarks from the southwestern Barents Sea, *Organic Geochemistry*, 63, 73-84, 10.1016/j.orggeochem.2013.08.008, 2013.
- Niewöhner, C., Hensen, C., Kasten, S., Zabel, M., and Schulz, H. D.: Deep Sulfate Reduction Completely Mediated by Anaerobic Methane Oxidation in Sediments of the Upwelling Area off Namibia, *Geochimica et Cosmochimica Acta*, 62, 455-464, 10.1016/s0016-7037(98)00055-6, 1998.
- 830 Nunoura, T., Soffientino, B., Blazejak, A., Kakuta, J., Oida, H., Schippers, A., and Takai, K.: Subseafloor microbial communities associated with rapid turbidite deposition in the Gulf of Mexico continental slope (IODP Expedition 308), *FEMS Microbiol Ecol*, 69, 410-424, 10.1111/j.1574-6941.2009.00718.x, 2009.
- Orsi, W. D., Richards, T. A., and Francis, W. R.: Predicted microbial secretomes and their target substrates in marine sediment, *Nat Microbiol*, 3, 32-37, 10.1038/s41564-017-0047-9, 2018.
- 835 Parkes, R. J., Cragg, B., Roussel, E., Webster, G., Weightman, A., and Sass, H.: A review of prokaryotic populations and processes in sub-seafloor sediments, including biosphere:geosphere interactions, *Marine Geology*, 352, 409-425, 10.1016/j.margeo.2014.02.009, 2014.
- Pellerin, A., Bui, T. H., Rough, M., Mucci, A., Canfield, D. E., and Wing, B. A.: Mass-dependent sulfur isotope fractionation during reoxidative sulfur cycling: A case study from Mangrove Lake, Bermuda, *Geochimica et Cosmochimica Acta*, 149, 152-164, 10.1016/j.gca.2014.11.007, 2015.
- 840 Pop Ristova, P., Wenzhofer, F., Ramette, A., Felden, J., and Boetius, A.: Spatial scales of bacterial community diversity at cold seeps (Eastern Mediterranean Sea), *ISME J*, 9, 1306-1318, 10.1038/ismej.2014.217, 2015.
- Raab, A. and Feldmann, J.: Microbial transformation of metals and metalloids, *Sci Prog*, 86, 179-202, 10.3184/003685003783238671, 2003.
- 845 Rasheed, M. A., Lakshmi, M., Rao, P. L. S., Kalpana, M. S., Dayal, A. M., and Patil, D. J.: Geochemical evidences of trace metal anomalies for finding hydrocarbon microseepage in the petroliferous regions of the Tatipaka and Pasarlapudi areas of Krishna Godavari Basin, India, *Petroleum Science*, 10, 19-29, 10.1007/s12182-013-0245-x, 2013.
- 850 Riedinger, N., Formolo, M. J., Lyons, T. W., Henkel, S., Beck, A., and Kasten, S.: An inorganic geochemical argument for coupled anaerobic oxidation of methane and iron reduction in marine sediments, *Geobiology*, 12, 172-181, 10.1111/gbi.12077, 2014.
- Riedinger, N., Brunner, B., Krastel, S., Arnold, G. L., Wehrmann, L. M., Formolo, M. J., Beck, A., Bates, S. M., Henkel, S., Kasten, S., and Lyons, T. W.: Sulfur Cycling in an Iron Oxide-Dominated, Dynamic Marine Depositional System: The Argentine Continental Margin, *Frontiers in Earth Science*, 5, 10.3389/feart.2017.00033, 2017.
- 855 Rovere, M., Mercorella, A., Frapiccini, E., Funari, V., Spagnoli, F., Pellegrini, C., Bonetti, A. S., Veneruso, T., Tassetti, A. N., Dell'Orso, M., Mastroianni, M., Giuliani, G., De Marco, R., Fabi, G., Ciccone, F., and

- Antoncecchi, I.: Geochemical and Geophysical Monitoring of Hydrocarbon Seepage in the Adriatic Sea, *Sensors* (Basel), 20, 10.3390/s20051504, 2020.
- 860 Ruff, S. E., Biddle, J. F., Teske, A. P., Knittel, K., Boetius, A., and Ramette, A.: Global dispersion and local diversification of the methane seep microbiome, *Proc Natl Acad Sci U S A*, 112, 4015-4020, 10.1073/pnas.1421865112, 2015.
- Sættem, J., Rise, L., and Westgaard, D. A.: Composition and properties of glacial sediments in the southwestern Barents Sea, *Marine Geotechnology*, 10, 229-255, 10.1080/10641199109379893, 1991.
- 865 Schulz, H. D.: Quantification of Early Diagenesis: Dissolved Constituents in Pore Water and Signals in the Solid Phase, in: *Marine Geochemistry*, 73-124, 10.1007/3-540-32144-6_3, 2006.
- Schwengers, O., Jelonek, L., Dieckmann, M. A., Beyvers, S., Blom, J., and Goesmann, A.: Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification, *Microb Genom*, 7, 10.1099/mgen.0.000685, 2021.
- 870 Seeberg-Elverfeldt, J., Schluter, M., Feseker, T., and Kolling, M.: Rhizon sampling of porewaters near the sediment-water interface of aquatic systems, *Limnol Oceanogr-Meth*, 3, 361-371, 10.4319/lom.2005.3.361, 2005.
- Smrzka, D., Zwicker, J., Bach, W., Feng, D., Himmler, T., Chen, D., and Peckmann, J.: The behavior of trace elements in seawater, sedimentary pore water, and their incorporation into carbonate minerals: a review, *Facies*, 65, 10.1007/s10347-019-0581-4, 2019.
- 875 Sultan, N., Garziglia, S., and Ruffine, L.: New insights into the transport processes controlling the sulfate-methane-transition-zone near methane vents, *Sci Rep*, 6, 26701, 10.1038/srep26701, 2016.
- Teske, A.: Hydrocarbon-Degrading Microbial Communities in Natural Oil Seeps, in: *Microbial Communities Utilizing Hydrocarbons and Lipids: Members, Metagenomics and Ecophysiology*, 1-31, 10.1007/978-3-319-60063-5_3-2, 2019.
- 880 Torres, M. E., Brumsack, H. J., Bohrmann, G., and Emeis, K. C.: Barite fronts in continental margin sediments: a new look at barium remobilization in the zone of sulfate reduction and formation of heavy barites in diagenetic fronts, *Chemical Geology*, 127, 125-139, 10.1016/0009-2541(95)00090-9, 1996.
- 885 Treude, T., Krause, S., Maltby, J., Dale, A. W., Coffin, R., and Hamdan, L. J.: Sulfate reduction and methane oxidation activity below the sulfate-methane transition zone in Alaskan Beaufort Sea continental margin sediments: Implications for deep sulfur cycling, *Geochimica et Cosmochimica Acta*, 144, 217-237, 10.1016/j.gca.2014.08.018, 2014.
- Treude, T., Krause, S., Steinle, L., Burwicz, E., Hamdan, L. J., Niemann, H., Feseker, T., Liebetrau, V., Krastel, S., and Berndt, C.: Biogeochemical Consequences of Nonvertical Methane Transport in Sediment Offshore Northwestern Svalbard, *Journal of Geophysical Research: Biogeosciences*, 125, 10.1029/2019jg005371, 2020.
- 890 Turchyn, A. V., Bradbury, H. J., Walker, K., and Sun, X.: Controls on the Precipitation of Carbonate Minerals Within Marine Sediments, *Frontiers in Earth Science*, 9, 10.3389/feart.2021.618311, 2021.
- 895 Vuillemin, A., Coskun, O. K., and Orsi, W. D.: Microbial Activities and Selection from Surface Ocean to Subseafloor on the Namibian Continental Shelf, *Appl Environ Microbiol*, 88, e0021622, 10.1128/aem.00216-22, 2022.
- Vuillemin, A., Kerrigan, Z., D'Hondt, S., and Orsi, W. D.: Exploring the abundance, metabolic potential and gene expression of subseafloor Chloroflexi in million-year-old oxic and anoxic abyssal clay, *FEMS Microbiol Ecol*, 96, 10.1093/femsec/fiaa223, 2020a.
- 900 Vuillemin, A., Vargas, S., Coskun, O. K., Pockalny, R., Murray, R. W., Smith, D. C., D'Hondt, S., and Orsi, W. D.: Atribacteria Reproducing over Millions of Years in the Atlantic Abyssal Subseafloor, *mBio*, 11, 10.1128/mBio.01937-20, 2020b.

- Widdel, F., Knittel, K., and Galushko, A.: Anaerobic Hydrocarbon- Degrading Microorganisms: An
905 Overview, in: Handbook of Hydrocarbon and Lipid Microbiology, Springer, Berlin, Heidelberg, 1997-
2021, <https://doi.org/10.1007/978-3-540-77587-4>, 2010.
- Yergin, D.: The Prize: The Epic Quest for Oil, Money & Power, Free Press 2009.
- Zhang, C., Fang, Y. X., Yin, X., Lai, H., Kuang, Z., Zhang, T., Xu, X. P., Wegener, G., Wang, J. H., and
910 Dong, X.: The majority of microorganisms in gas hydrate-bearing seafloor sediments ferment
macromolecules, *Microbiome*, 11, 37, 10.1186/s40168-023-01482-5, 2023.