

1 **HESS Opinion paper:**  
2 **Towards a common vision for the future of hydrological**  
3 **observatories**

4 Paolo Nasta<sup>1</sup>, Günter Blöschl<sup>2</sup>, Heye R. Boga<sup>3</sup>, Steffen Zacharias<sup>4</sup>, Roland Baatz<sup>5</sup>, Gabriëlle De  
5 Lannoy<sup>6</sup>, Karsten H. Jensen<sup>7</sup>, Salvatore Manfreda<sup>8</sup>, Laurent Pfister<sup>9,10</sup>, Ana M. Tarquis<sup>11</sup>, Ilja van  
6 Meerveld<sup>12</sup>, Marc Voltz<sup>13</sup>, Yijian Zeng<sup>14</sup>, William Kustas<sup>15</sup>, Xin Li<sup>16</sup>, Harry Vereecken<sup>3</sup>, Nunzio  
7 Romano<sup>1,\*</sup>

8 <sup>1</sup> Department of Agricultural Sciences, Division of Agricultural, Forest and Biosystems Engineering, University of  
9 Naples Federico II, Portici (Naples), Italy.

10 <sup>2</sup> Institute of Hydraulic Engineering and Water Resources Management, TU Wien, Austria.

11 <sup>3</sup> Agrosphere Institute, Forschungszentrum Jülich GmbH, Jülich, Germany.

12 <sup>4</sup> Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany.

13 <sup>5</sup> Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany.

14 <sup>6</sup> Department of Earth and Environmental Sciences, Division Soil and Water Management, KULeuven, Belgium.

15 <sup>7</sup> Department of Geosciences and Natural Resource Management, Geology section, University of Copenhagen,  
16 Copenhagen, Denmark.

17 <sup>8</sup> Department of Civil, Building and Environmental Engineering (DICEA), University of Naples Federico II, Naples,  
18 Italy.

19 <sup>9</sup> Luxembourg Institute of Science & Technology (LIST), Esch-sur-Alzette, Luxembourg.

20 <sup>10</sup> University of Luxembourg, Faculty of Science, Technology and Medicine, Luxembourg.

21 <sup>11</sup> CEIGRAM, Department of Applied Mathematics, Universidad Politécnica de Madrid (UPM), Madrid, Spain.

22 <sup>12</sup> Department of Geography, University of Zurich, Zurich, Switzerland.

23 <sup>13</sup> Laboratoire sur les Interactions Sol-Agrosystème-Hydrosystème UMR INRAE-IRD-Institut Agro 2, Montpellier  
24 Cedex, France.

25 <sup>14</sup> Department of Water Resources, Faculty of Geo-Information Science and Earth Observation (ITC), University of  
26 Twente in Enschede, the Netherlands.

27 <sup>15</sup> USDA-ARS Hydrology & Remote Sensing Lab, Beltsville, USA.

28 <sup>16</sup> National Tibetan Plateau Data Center, State Key Laboratory of Tibetan Plateau Earth System, Environment and  
29 Resources, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China.

30  
31 *Correspondence to:* Nunzio Romano (nunzio.romano@unina.it)

32  
33

34 **Abstract.** The Unsolved Problems in Hydrology (UPH) initiative has emphasized the need to establish networks of  
35 multi-decadal hydrological observatories to gain a deep understanding of the complex hydrologic processes occurring  
36 within diverse environmental conditions. The already existing monitoring infrastructures have provided an enormous  
37 amount of hydrometeorological data, facilitating detailed insights into the causal mechanisms of hydrological  
38 processes, the testing of scientific theories and hypotheses, and the revelation of the physical laws governing  
39 catchment behavior. Yet, hydrological monitoring programs have often produced limited outcomes due to the  
40 intermittent availability of financial resources and the substantial efforts required to operate observatories and conduct  
41 comparative studies to advance previous findings. Recently, some initiatives have emerged aiming at coordinating  
42 data acquisition and hypothesis testing to facilitate an efficient cross-site synthesis of findings. To this end, a common  
43 vision and practical data management solutions need to be developed. This opinion paper provocatively discusses two  
44 potential end members of a future hydrological observatory (HO) network, based on a given hypothesized community  
45 budget: a comprehensive set of moderately instrumented observatories or, alternatively, a small number of highly  
46 instrumented super-sites.

47 A network of moderately instrumented monitoring sites would provide a broad spatial coverage across the major  
48 pedoclimatic regions by supporting cross-site synthesis of the lumped hydrological response (e.g., rainfall-runoff  
49 relationship, Budyko analysis) across diverse continental landscapes. However, the moderate instrumentation at each  
50 site may hamper an in-depth understanding of complex hydrological processes. In contrast, a small number of  
51 extensively instrumented research sites would enable community-based experiments in an unprecedented manner,  
52 thereby facilitating a deeper understanding of complex, non-linear processes modulated by scale-dependent feedback  
53 and multiscale spatiotemporal heterogeneity. Lumping resources has proven to be an effective strategy in other  
54 geosciences, e.g. research vessels in oceanography and drilling programs in geology. On the downside, a potential  
55 limitation of this approach is that a few catchments will not be representative of all pedoclimatic regions, necessitating  
56 the consideration of generalization issues.

57 A discussion on the relative merits and limitations of these two visions on HOs is presented to build consensus on the  
58 optimal path for the hydrological community to address the UPH in the coming decades. A final synthesis proposes  
59 the potential for integrating the two end members into a flexible management strategy.


60  
61 **Keywords:** hydrological observatory network, experimental catchments, cross-site synthesis, hypothesis testing vs.  
62 exploratory science, unsolved problems in hydrology, societal needs, technology advancements.

- 63  
64 **Highlights**
- 65 • The historical situation of HOs has led to fragmented knowledge and sub-optimal research progress.
  - 66 • Some initiatives emerged to coordinate and standardize data and models resulting in efficient cross-site synthesis.
  - 67 • It is important to stimulate discussion within the hydrological community to arrive at a consensus view on HOs.

68  
69  
70  
71  
72  
73  
74  
75

76  
77  
78  
79  
80  
81  
82

### Graphical Abstract



83

## 84 **1. How do we advance scientific understanding of hydrological processes?**

85 Water is under increasing threat due to human activities. Rapid changes in land use, such as the adoption of more  
86 intensive farming practices, the expansion of urbanization, and the abandonment of land in rural areas, have a  
87 significant impact on the hydrological cycle and water quality, whereas unsustainable water withdrawals lead to the  
88 depletion of resources. Global warming is expected to exacerbate hydrological extremes, resulting in more disastrous  
89 floods and severe droughts that will further threaten water security. In light of these challenges, the mission of  
90 hydrologists and water managers is to sustainably meet human needs while preserving biodiversity and ecosystem  
91 services, based on the most accurate and up-to-date information. However, the extent to which anthropogenic stressors  
92 influence the hydrologic cycle is not yet fully understood and the effectiveness of adaptation actions to guide the  
93 management of water resources has yet to be fully evaluated. Hydrology is a data-hungry discipline but the limited  
94 observations on all components of the terrestrial hydrosphere, from bedrock to the lower atmosphere, represent a  
95 significant obstacle to progress in the understanding of hydrologic process dynamics (Siebert et al., 2024).

96 To grasp the daunting complexity of the hydrological cycle, particularly in relation to the impact of human activities  
97 on the critical zone and catchment functionality, and to address the Unsolved Problems in Hydrology (UPH), several  
98 hydrological observatories (HOs) have been established around the world with the specific purpose of monitoring  
99 hydrological states and flows (Blöschl et al., 2019; Arora et al., 2023).

100 The concept of hydrological observatories (HOs) can be traced back to the early 1900s when scientists began to  
101 recognize the significance of long-term data collection for understanding hydrological processes (McDonnell et al.,  
102 2007). In 1903, runoff and other hydrological variables were initially collected in the Sperbelgraben and  
103 Rappengraben experimental catchments in the Emmental region of Switzerland. These catchments remain operational  
104 and hold one of the longest continuous discharge records in the world (Stähli et al., 2011). In the United States, the  
105 first HOs were the Wagon Wheel Gap Experiment in Colorado (Bates and Henry, 1928), the Coweeta Hydrological  
106 Laboratory in North Carolina (Neary et al., 2012), and a catchment network across the continental U.S. established by  
107 the USDA-Agricultural Research Service (Goodrich et al., 2021). These sites were designed to study the influence of  
108 human activities on hydrological systems, with a particular focus on deforestation and afforestation, land-use changes,  
109 and agricultural practices (Whitehead and Robinson, 1993). Since the 1950s, there has been a notable increase in the  
110 number of HOs established across the globe. The HO sites have provided invaluable information for the effective  
111 management of water resources. Currently, a diverse range of entities, including government agencies (e.g., the  
112 Hydrologic Benchmark Network of the US Geological Survey), universities and research institutions, international  
113 organizations, and non-governmental organizations, provide funding and support for these sites.

114

## 115 **2. Building integrated observation platforms**

116 A considerable number of rivers worldwide have been equipped with gauges to monitor precipitation and streamflow  
117 by governmental agencies for the purpose of water management. The data collected have been primarily utilized at  
118 the national level, although there are several transnational initiatives, including the Global Runoff Data Centre in  
119 Koblenz, Germany, and the Camels datasets, such as those for the US, Chile, and Brazil, (Addor et al.; Alvarez-  
120 Garreton et al. 2018; Chagas et al. 2020). HOs extend beyond these conventional networks, striving to gain a more  
121 comprehensive understanding of hydrological processes, typically in smaller catchments.

122 A hydrological observatory is defined as a cyber-physical infrastructure established within a catchment area to monitor  
123 the hydrological variables and fluxes, as well as to characterize the hydrological behavior of the three-dimensional

124 spatial domain. The catchment is assumed to be the fundamental hydrological unit with well-defined system  
125 boundaries. It is from this unit that the impact of anthropogenic disturbances (global warming, land use change, aquifer  
126 contamination, etc.) on water resources can be evaluated through a long-term data analysis. Given the impracticality  
127 of full catchment coverage, the hydrological observatory focuses on a selected cluster of sub-catchments (spatial  
128 resolution of hectares) which are representative of land use, geomorphology, topography, and pedology similarities  
129 (Bogena et al., 2006). Consequently, the selected sub-catchments are equipped with wireless sensor networks for  
130 continuous data collection and subjected to disparate field campaigns, contingent upon budgetary constraints.

131 The selection of sensors is crucial for the effective collection of hydrometeorological data within a hydrological  
132 observatory. Weather station networks (also called synoptic stations) ensure the collection of meteorological data and  
133 have been integrated in many countries with weather radar networks for the purpose of detailed precipitation  
134 estimation (Sokol et al., 2021). Snow water equivalent is already measured on a routine basis with snow pillows (e.g.  
135 by the SNOTEL network in the United States) or, more experimentally, by airborne LiDAR snow depth surveys  
136 (Painter et al., 2016). Groundwater levels are monitored on a routine basis, whereas distributed temperature sensing  
137 technology is a more novel approach for estimating infiltration rates and potentially catchment-scale groundwater  
138 recharge (Medina et al., 2020). The measurements of soil water content and matric potential, soil temperature, and  
139 soil bulk electrical conductivity are conducted across soil profiles at the point scale (Hoffmann et al., 2015; Peng et  
140 al., 2019; Bogena et al., 2022). Cosmic-ray neutron sensors, meanwhile, are capable of extending the footprint of soil  
141 moisture to approximately 150-200 meters in radius (Romano, 2014; Köhli et al., 2015; Baatz et al., 2017). At  
142 experimental sites, surface and subsurface runoff from hillslopes are measured using flowmeters in runoff plots (Fu  
143 et al., 2024). In addition, the mapping of saturation areas on hillslopes (Silasari et al. 2017) and channel-network  
144 dynamics (Jenssen et al., 2019; Strelnikova et al., 2023; Noto et al., 2024) provide insight into the spatial patterns of  
145 catchment-scale processes that extend beyond point measurements. Topographic surveys assist in determining surface  
146 flow paths within a catchment, thus enabling the extension of point measurements to the catchment scale (e.g.,  
147 Rinderer et al., 2019; Fan et al., 2019; Refsgaard et al., 2021). The rates of soil erosion and deposition are quantified  
148 through the use of sediment fences, soil profile surveys, and cosmogenic nuclide analysis, in addition to repeated high-  
149 precision topographic surveys. The measurement of soil physical, chemical, and hydraulic properties is typically  
150 conducted in field campaigns and laboratory experiments, with remote sensing serving as a complementary technique.  
151 Geophysical tools, such as electromagnetic (EM) surveys, have the potential to provide valuable insights into the  
152 imaging of aquifer systems and the characterization of subsurface heterogeneity (Nasta et al., 2019; Dewar and Knight,  
153 2020). To elucidate the interactions of the water cycle with the biochemical, energy, and carbon cycles, numerous  
154 other variables are monitored (Valdes-Abellan et al., 2017). The key vegetation characteristics that are monitored  
155 include canopy height, leaf area index (LAI), leaf water potential, sap flow, rooting depth and distribution, plant water  
156 stress, canopy/vegetation water content, and temperature (Poyatos et al., 2021; Loritz et al., 2022; Zeng and Su, 2024).  
157 Eddy covariance measurements, some of which are connected through networks, such as Fluxnet, are used to obtain  
158 evapotranspiration and carbon fluxes at the local level. Sapflow sensors, some of which are organized in the Sapflux  
159 network (Poyatos et al., 2021), can be used to quantify transpiration rates. The use of tracer measurements, such as  
160 isotope and dye studies, enables the tracking and differentiation of water fluxes (Klaus and McDonnell, 2013; Penna  
161 et al., 2018). Lysimeters are used to determine groundwater recharge and the associated concentrations of, e.g., nitrate,  
162 at the point scale.

163 The use of unmanned aerial systems (UAS; e.g. Dugdale et al., 2022; Romano et al. 2023) and satellite platforms (e.g.  
164 Durand et al., 2021, De Lannoy et al., 2022) for remote sensing has emerged as a valuable supplementary method to

165 ground-based observation in HOs for gathering information over large heterogeneous areas as well as for upscaling  
166 or downscaling hydrological variables (e.g., McCabe et al., 2017; Manfreda et al., 2018, 2024; Su et al., 2020).  
167 Recently, higher-resolution observations of various hydrological variables have become available, including soil  
168 moisture (Burdun et al., 2023; Han et al., 2023), snow depth (Lievens et al., 2021), and irrigation rate (Dari et al.,  
169 2023). These observations can be used together with coarser-scale products, including total water storage data from  
170 the Gravity Recovery and Climate Experiment (GRACE) mission, and discharge data from the Surface Water and  
171 Ocean Topography (SWOT) mission. The deployment of multiple sensors, as seen in the various Sentinel and Landsat  
172 missions, can enhance the accuracy and resolution of the data. The European Space Agency (ESA) and the United  
173 States National Aeronautics and Space Administration (NASA) are engaged in collaborative efforts with public and  
174 private organizations to develop relevant new missions and to disseminate a range of products, including  
175 evapotranspiration estimates through the SEN-ET (Guzinski et al., 2019; 2020) and OpenET (Melton et al., 2021)  
176 initiatives.

177 It is evident that the key to progress in hydrological understanding will be contingent upon the integration of these  
178 observation platforms. These platforms should integrate technologies such as remote sensing, high-performance  
179 computing resources, artificial intelligence, and the Internet of Things, while acknowledging the influence of  
180 geochemical and biotic heterogeneity, as well as socio-economic processes, on water and energy fluxes.

181 While observations are the cornerstone of progress in hydrological understanding (Sivapalan and Blöschl, 2017),  
182 models are equally essential for hypothesis testing and making predictions that are practically relevant (Brooks et al.,  
183 2015; Baatz et al., 2018; Bogena et al., 2018; Bechtold et al., 2019; Nearing et al., 2024). However, hydrological  
184 models, particularly those of a complex nature, frequently rely on lumped parameter calibration. This means that  
185 model parameters are adjusted based on aggregated (or lumped) fluxes, such as those observed in streamflow  
186 measurements at the outlet of the catchment. Although this approach can be effective, it can also result in limitations.  
187 A significant challenge is the assumption that the model's behavior is uniform across the entire catchment. This  
188 assumption might not hold true, especially in heterogeneous catchments with diverse topography, land use, and soil  
189 types. In such cases, relying exclusively on lumped fluxes may result in suboptimal model performance. An integrated  
190 observation approach enables the calibration based on insightful analysis of process complexity through systematic  
191 learning from distributed hydrometeorological data given that catchments are complex systems with structured  
192 heterogeneities, which give rise to non-linear interactions and feedback between the component processes (Vereecken  
193 et al., 2015; Li et al., 2022). One aspect of integration is the assimilation of observations into hydrological models  
194 (Mwangi et al., 2020; Kumar et al., 2022; De Lannoy et al., 2022) to estimate unobserved variables, improve  
195 predictions, and calibrate and validate satellite retrieval (Colliander et al., 2021). Paleo-reconstructions represent  
196 another example of integration and are instrumental in developing a more comprehensive understanding of how  
197 dynamic, abiotic, and biotic catchment characteristics have co-evolved well before the advent of instrumental records  
198 (Troch et al., 2013). Climate shifts leave a multitude of signatures in the natural world, influencing processes such as  
199 tree growth and the distribution of plant species. The advent of increasingly sophisticated analytical techniques has  
200 facilitated a rapid growth in knowledge regarding past climate and river ecosystem variability. Of particular benefit  
201 are reconstructions of river flow and erosion derived from natural archives (Chaussé et al., 2008; Torbenson et al.,  
202 2021; Büntgen et al., 2021; Schöne et al., 2020; Strelnikova et al., 2023).

203  
204  
205

### 206 **3. HO networks and hydrological synthesis**

207 The sustainability of HO is a matter of concern. Financial and logistical constraints have posed challenges to the long-  
208 term operation of HOs, jeopardizing essential maintenance, equipment upgrades, and personnel training. This  
209 ultimately compromises the quality and continuity of hydrological data collection and analysis. Data gaps and the lack  
210 of continuity in the data collection process hamper the identification and understanding of hydrological change, which  
211 represents a significant societal need for hydrology in the present and the future (Montanari et al., 2013). In light of  
212 the frequently constrained budgetary resources available for each site, many studies have focused on measuring  
213 lumped hydrological fluxes (e.g., the streamflow at the catchment outlet), while observatories that prioritize the  
214 analysis of spatial details remain relatively scarce (e.g., Blöschl et al., 2016). Site-specific methods, tailored to site-  
215 specific UPH, have frequently resulted in advancements in the understanding of a specific hydrological process, but  
216 have not fully exploited the potential for synergies with other HOs. Consequently, the outcome has frequently been  
217 an increase in fragmented knowledge, rather than progress in understanding the interactions of hydrological processes,  
218 which is so urgently needed.

219 To address these issues, scientists have proposed initiatives to sustain long-term operation, harmonize, and standardize  
220 both hydrometeorological data and eco-hydrological models in HO networks (Zoback 2001; Reid et al., 2010;  
221 Kulmala, 2018). In numerous instances, hydrological observations are now integrated into interdisciplinary research  
222 programs in terrestrial observatories which are scientific facilities designed to observe and study various aspects of  
223 the Earth's surface, atmosphere, and interior. Terrestrial observatories collect data on a range of phenomena, including  
224 earthquakes, volcanic activity, weather patterns, climate change, and the movement of tectonic plates. Hydrological  
225 observations play a crucial role in the context of terrestrial observatories. Notable initiatives that have integrated  
226 existing environmental research infrastructures include the pan-European ENVRI initiative (<https://envri.eu>) and the  
227 global GERI initiative (<https://global-ecosystem-ri.org/>, Loescher et al., 2022). Networks such as Fluxnet  
228 (<https://fluxnet.org>) and the Integrated Carbon Observation System (<https://www.icos-cp.eu>) collect standardized data  
229 on the soil surface energy balance and evapotranspiration. The network of Critical Zone Observatories aims at  
230 understanding critical zone processes, with a particular focus on hydrologic monitoring (Brantley et al., 2017;  
231 Anderson et al., 2018; Gaillardet et al., 2018). The integrated European Long-Term Ecosystem, Critical Zone, and  
232 socio-ecological Research Infrastructure (<https://elter-ri.eu>) is establishing a network of approximately 200 integrated  
233 terrestrial observatories across Europe, with hydrological monitoring forming a component of this initiative. In the  
234 field of agriculture, the United States Department of Agriculture (USDA) is providing support for the Long-Term  
235 Agroecosystem Research (LTAR) initiative (<https://ltar.ars.usda.gov/>), which combines strategic research projects  
236 with common measurements across multiple agroecosystems, including croplands, rangelands, and pasturelands.

237 The advent of digital technology and data exchange platforms has enabled scientists to aggregate and jointly analyze  
238 data streams from disparate locations in a manner that was previously unfeasible. This is contingent upon the  
239 standardization and harmonization of existing protocols and methods for hydrological observation. The extant  
240 research infrastructures have already established standards for the environmental variables they collect. The  
241 harmonization of such standards across disciplinary infrastructures represents a crucial building block toward  
242 enhanced integration and should be reflected in future strategies for designing international environmental research.


243 The cross-site synthesis of hydrological processes serves to fill the gap between site-specific studies and broader, more  
244 generalizable knowledge (Zacharias et al., 2024). The objective is to integrate information from multiple sites and  
245 sources to identify patterns, trends, and relationships that can lead to the development of a more robust and transferable

246 body of knowledge for model development and, ultimately, more effective decision-making. The implementation of  
247 cross-site synthesis typically entails the following steps, as illustrated in Figure 1:

- 248 – Formulating the UPH;
- 249 – Data collection by using standardized protocols;
- 250 – Use of community-shared hydrological models;
- 251 – Comparative hydrology;
- 252 – Meta-analyses to consolidate results.

253  
254 The initial step is to formulate scientifically interesting questions that address existing knowledge gaps and contribute  
255 to a broader understanding and societal benefits in the field of hydrology (see Appendix). To ensure consistency in  
256 data quality, it is essential to harmonize the measurement techniques and quality control protocols employed.  
257 Community networks and centralized data repositories can facilitate this process and provide access to standardized  
258 and curated datasets. Community-shared hydrological models can be employed to represent the complex interactions  
259 between hydrological processes, ecosystems, and human activities. The models are calibrated and tested using the  
260 harmonized data from multiple sites, thereby enhancing their predictive capabilities and generalizability. Notable  
261 initiatives are already operational, including the Unified Forecast System (UFS) which is a community-based, coupled,  
262 comprehensive Earth modeling system used for weather forecast applications  
263 (<https://www.ufscommunity.org/articles/hierarchical-system-development-for-the-ufs/>). Comparative studies have  
264 been instrumental in identifying the key drivers of hydrological variability and in establishing generalizable principles.  
265 This is accomplished through a comprehensive and systematic comparison of hydrological processes and responses  
266 across a range of sites, while accounting for several factors, including climate, topography, land use, and management  
267 practices. In addition, meta-analyses can synthesize and compare findings from multiple studies, identify recurrent  
268 patterns and trends in integrated measurements and model simulations, and present consolidated results in a coherent  
269 manner.

270 By following these steps, hydrologists can effectively implement cross-site synthesis, thereby advancing the field of  
271 hydrology toward a more generalizable and transferable body of knowledge. This can inform more effective decision-  
272 making with regard to the management of water resources and the adaptation to climate change in a variety of contexts.



273  
274 **Figure 1. Schematic of the proposed cross-site synthesis. Obs, Db, M, and UPH indicate Observatories, Database, Models,**  
275 **and Unsolved Problems in Hydrology, respectively.**



277 Cross-site synthesis helps unveil hidden assumptions that may be embedded in site-specific studies, thereby enabling  
278 researchers to critically assess the validity of these assumptions and explore alternative perspectives. The identification  
279 of common principles and practices allows researchers to develop transferable knowledge that can be applied to other  
280 settings, thereby accelerating progress in research and practice. Examples of cross-site synthesis already exist in the  
281 literature. For example, Wlostowski et al. (2021) conducted a meta-analysis of hydrologic signatures from 15  
282 catchments in the U.S. Critical Zone Observatory (CZO) network, which revealed consistent relationships between  
283 critical zone structure and hydrologic response across sites. Similarly, Addor et al. (2018) examined the predictability  
284 of hydrologic signatures for the catchments included in the Camels dataset but found that the relationship between  
285 these signatures and catchment attributes other than climate characteristics was weak.

286 Comparative analyses have yielded a range of interesting results, although there is not yet complete concordance. For  
287 instance, some studies have indicated that afforestation may result in a decrease in water yield, whereas others have  
288 identified an increase. Two distinct theoretical frameworks have been put forth to explain the aforementioned  
289 conflicting results (Ellison et al., 2012). The *demand-side* perspective places emphasis on the increase in transpiration  
290 and the subsequent reduction in streamflow, particularly in catchments smaller than a few square kilometers (Schilling  
291 et al., 2008; Kim et al., 2013; Nasta et al., 2017). In contrast, the *supply-side* perspective posits that afforestation will  
292 intensify precipitation, thereby increasing streamflow, in downwind catchments (Ellison et al., 2012). Similarly, the  
293 impact of reforestation on dry season flows is contingent upon the relative importance of increased infiltration and  
294 evapotranspiration rates (Bruijnzeel, 1989).

295 As demonstrated by the preceding case studies, cross-site synthesis provides a valuable approach for quantifying the  
296 spatial variability of hydrological processes and identifying consistent patterns of phenomena, such as droughts and  
297 floods. These examples will ultimately inform water management practices around the world while maintaining track  
298 and awareness of local hydrological particularities.


299

#### 300 **4. How to manage a network of hydrological observatories?**

301 For the sake of the argument, we assume that a fixed community budget has been allocated for the establishment and  
302 operation of a hypothetical network of HOs in the European Union (EU). Two potential scenarios can be envisaged.  
303 In the first scenario (*Scenario 1*), the available financial resources are distributed among a multitude of moderately  
304 instrumented HOs situated throughout the EU, with the objective of addressing challenges in hydrology with extensive  
305 geographical coverage. Figure 2 illustrates an example of a moderately instrumented site belonging to a hydrological  
306 observatory network in *Scenario 1*. This plan reflects the current status of the majority of HO networks around the  
307 world. The principal benefit of this approach is that the HOs are widely and effectively distributed, enabling the  
308 identification of cause-and-effect relationships and supporting the cross-site synthesis of the lumped hydrological  
309 responses (e.g., rainfall-runoff relationship, Budyko analysis) across diverse continental landscapes (Wagner et al.,  
310 2007; Ehret et al., 2014; Jones et al., 2012; Kuentz et al., 2017; Templer et al., 2022).

311 In *Scenario 1*, a combination of centralized and distributed components is utilized. Distributed components provide  
312 observed data that is managed by different entities (e.g., universities, research institutions, government agencies, etc.)  
313 across geographically spread sites. To guarantee the comparability of data, it is essential to implement standardized  
314 protocols for data collection, storage, quality assurance, and analysis. This will alleviate the burden associated with

315 the cross-site synthesis. Centralized data management facilitates the accessibility of data across multiple sites.  
 316 Furthermore, additional central thematic elements may be provided, such as those pertaining to communication and  
 317 knowledge transfer, or those relevant to modeling applications. The organizational structure may be based on other  
 318 successfully established or planned distributed continental infrastructures. Notable examples include ICOS (Integrated  
 319 Carbon Observation System) and eLTER (Integrated European Long-Term Ecosystem, Critical Zone and Socio-  
 320 ecological Research Infrastructure). Free data availability and accessibility of the sites should be a fundamental aspect  
 321 of the scenario designs.  
 322



323  
 324 **Figure 2. Graphical illustration of a hydrological observatory (HO) network in the European Union (EU) in Scenario 1.**  
 325 **Each sub-catchment is equipped with basic instrumentation: a weather station, a runoff gauging station, a Cosmic-Ray**  
 326 **Neutron Sensor (CRNS) with a wireless sensor network controlling soil profile sensors, a streamflow sensor at the**  
 327 **catchment's outlet. Satellite products are available anywhere in the world. The soil profile cross-section illustrates the soil**  
 328 **profile sensor unit and the stationary CRNS.**

329  
 330 In *Scenario 1*, collaboration and partnership among different stakeholders are crucial. Such collaboration may  
 331 facilitate broader opportunities for citizen and stakeholder participation, particularly given the distributed nature of  
 332 the scenario and the encouragement of local initiatives.  
 333 In the second scenario (*Scenario 2*), research efforts and financial resources are pooled into a limited number of pilot  
 334 HOs, each equipped with massive instrumentation. Similar initiatives can be found in related fields of study. In  
 335 oceanography, a limited number of costly research vessels are made available, primarily through the financial support  
 336 of national governments. This approach enables numerous researchers to engage in collaborative community  
 337 experiments, thereby facilitating a deeper understanding of specific oceanic regions. One illustrative example is the

338 Multidisciplinary Drifting Observatory for the Study of the Arctic Climate (MOSAiC), which undertook a drift with  
339 the Arctic Sea ice from October 2019 to September 2020 aboard the Polarstern research vessel (Rabe et al., 2022). In  
340 the field of geology, the cost of drilling into the Earth is almost equally expensive. The consolidation of resources  
341 permits geologists from diverse geographical locations to engage in collaborative drilling programs, such as the  
342 International Continental Scientific Drilling Program (ICDP) (Harms et al., 2007). The scientific drilling programs  
343 were conducted at locations of global geological significance, which have been designated as World Geological Sites.  
344 Of course, in both cases, the research questions or aims are explicitly delineated. In the case of MOSAiC, the aim was  
345 to gain a deeper understanding of the influence of the Arctic on the global climate, given that the Arctic has  
346 experienced a more pronounced warming trend than any other region of the world. Concerning ICDP, the objective  
347 was to gain a deeper understanding of the Earth's processes and structure at the most interesting locations. In both  
348 instances, participation is contingent upon the successful completion of an application and review process overseen  
349 by an international committee.


350 Similarly, a small number of HOs equipped with comprehensive instrumentation and managed by an international  
351 team of experts from various disciplines could represent the pinnacle of hydrological field research. Figure 3 shows a  
352 hypothetical super-site established along an ideal transect within the European Union (*Scenario 2*). A high-density  
353 network of sampling and monitoring units for soil hydrology research is designed and planned for each super-site.  
354 This infrastructure, as yet unrealized, would facilitate a comprehensive understanding of water dynamics in the  
355 groundwater-soil-plant-atmosphere continuum and water circulation in the surface and subsurface domains within a  
356 few sites on each continent. In this case, cross-site synthesis would support the application and refinement of complex  
357 hydrological models based on fundamental insights into complex, non-linear processes that are modulated by scale-  
358 dependent feedback and multiscale spatiotemporal heterogeneity.

359 A wealth of data would enable an unprecedented unraveling of hydrological processes at the hillslope/catchment scale,  
360 based on observations of water and energy fluxes in the groundwater-soil-vegetation-atmosphere continuum at high  
361 spatial and temporal resolutions.

362 In *Scenario 2*, the research questions should be presented boldly (see discussion in Davis, 1926; Beven and Germann,  
363 2013; McDonnell, 2014; Burt and McDonnell, 2015; Kirchner, 2016; Blöschl et al., 2019; Gao et al., 2023). Some  
364 interesting examples of *Scenario 2* have already been documented in controlled environments. Biosphere-2 (B2) in  
365 Tucson, Arizona, (Evaristo et al., 2019) is a research facility comprising a tropical rainforest biome, a mesocosm  
366 enclosed in a pyramidal glass structure. Additionally, the Landscape Evolution Observatory (LEO) comprises three  
367 artificial hillslopes equipped with a dense network of soil sensors (Pangle et al., 2015). The observatory is focused on  
368 understanding the interaction between water and weathering processes (Van Den Heuvel, 2018; Bauser et al., 2022).  
369 Another example is the artificial Chicken Creek catchment in Germany, which has served as the fulcrum of  
370 comparative community research on runoff generation (Holländer et al., 2009).

371 Once more, as with the sister disciplines, the choice of location is of the utmost importance. The selected locations  
372 should represent hydrological situations that are particularly conducive to addressing the primary research question.  
373 The Austrian Hydrological Open Air Laboratory (HOAL) (Blöschl et al., 2016) was designed with the specific  
374 objective of facilitating a more comprehensive understanding of rainfall-runoff processes. It is ideally suited for this  
375 purpose, featuring a range of different runoff generation processes, including surface runoff, springs, tile drains, and  
376 wetlands. Another example is provided by the Alento hydrological observatory, which aims at elucidating the effects  
377 of the typical Mediterranean seasonality of climate, as well as land-use/land-cover changes on water flow in the critical  
378 zone of a representative southern European catchment (Nasta et al., 2017; Romano et al., 2018). To explore land-

379 atmosphere feedback, it is recommended that a catchment of considerable size be selected (Späth et al., 2023). In  
 380 addition, when selecting a location for a new, high-budget HO, it may be beneficial to consider the existence of so-  
 381 called environmental archives in the area of interest. The ease of accessibility and the availability of infrastructure  
 382 may be another factor to consider, but this could result in a geographic and climatic bias of the research sites.  
 383



384

385 **Figure 3. Graphical illustration of a hydrological observatory (HO) network in the European Union (EU) in Scenario 2.**  
 386 **Each sub-catchment established along an ideal transect is equipped with a high-density network of sampling and**  
 387 **monitoring units for soil hydrology research. Frequent unmanned aerial system (UAS) and aircraft surveys are organized**  
 388 **over the experimental area. Satellite products are available anywhere in the world. Frequent campaigns of geophysical**  
 389 **(electromagnetic induction, EMI technique) and tracing (stable isotopes in water such as  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ ) measurements are**  
 390 **carried out across the HO. Flow monitoring and water sampling are carried out along the stream. The soil profile cross-**  
 391 **section shows the monitoring and sampling activities in the groundwater-soil-plant-atmosphere continuum in a position of**  
 392 **the dense point grid (purple circles).**


393

394

395 Few super-sites would require a central governing body that would likely be responsible for overseeing all aspects of  
 396 the super-sites, including instrument deployment and maintenance, as well as data collection and analysis. Such an  
 397 entity could be a dedicated government agency with a specific mandate or a research consortium with significant  
 398 resources. The establishment of a single entity in charge, operating as a central authority, would facilitate the decision-  
 399 making process with regard to instrument upgrades, research focus, and site and data access.

400 Super-sites with advanced instrumentation might attract highly specialized researchers, leading to a concentration of  
 401 expertise in specific areas. The implementation of standardized sensors would result in cost savings and enhanced  
 402 efficiency in the collection and processing of data. Conversely, the specific hydrological environment may require the

403 use of specialized instrumentation or measurement techniques. A lack of flexibility in standardization can limit the  
 404 ability to adapt to new research questions or emerging challenges. Notable examples of standardization efforts include  
 405 the Global Network of River Observatories (GLORIA) and the World Meteorological Organization (WMO)  
 406 guidelines for hydrological stations. By taking these factors into careful consideration and adopting a balanced  
 407 approach, hydrological observatories can harness the power of standardization while maintaining flexibility and  
 408 adaptability. To ensure equity and stimulate greater involvement in *Scenario 2*, it is essential to establish a  
 409 collaborative governance structure that incorporates a diverse range of stakeholders in decision-making processes  
 410 pertaining to super site operations and data utilization. The governance and site access aspects are well presented in  
 411 initiatives such as the International Continental Scientific Drilling Program (ICDP), which addresses geodynamic  
 412 processes, solid Earth geohazards, sustainable geo-resources, and environmental change (<https://www.icdp-online.org/about-icdp/entities/>). Another noteworthy example is the Alfred Wegener Institute (AWI), which aims to  
 414 understand the complex processes in the Earth system and the impact of global warming on the oceans and polar  
 415 regions (<https://www.awi.de/en/>). The AWI maintains a network of well-instrumented long-term observatories,  
 416 comprising both stationary devices and mobile components that are used for studies pertaining to oceanography,  
 417 meteorology, and geophysics (<https://www.awi.de/en/expedition/observatories.html>).  
 418 The selection of the optimal scenario is contingent upon the research questions deemed most pertinent, the capacity  
 419 to secure funding, and the extent to which the hydrological community is willing and able to collaborate. In the context  
 420 of financial constraints, no single alternative can be considered inherently superior. A distributed network of numerous  
 421 HOs is well-suited to broad-scale inquiries, whereas a network of a few super-sites is particularly adept at facilitating  
 422 in-depth process understanding.  
 423



424  
 425 **Figure 4. Possible configurations of Hydrological Observatory (HO) networks are illustrated, spanning a range from a few**  
 426 **(color-coded in red) to numerous (color-coded in blue) HOs. The thickness of the arrows indicates the quantity of**  
 427 **instrumentation present in each HO. The data obtained from the HO network and remote sensing platforms are used to**  
 428 **inform hydrological models of different complexities, enabling the addressing of specific scientific questions across**  
 429 **disparate spatial scales.**  
 430

431  
 432 A potential way forward to the aforementioned issues could be to merge the two scenarios into a dynamic or adaptive  
 433 hybrid approach (see Figure 4). The establishment of a network of geographically distributed observatories would  
 434 facilitate the achievement of a high level of representativeness regarding existing gradients of geology, climate, and  
 435 land use. This would assist in the identification of priority areas requiring further investigation, and alleviate some of  
 436 the bias in current hydrological studies (Burt and McDonnell, 2015; Tarasova et al., 2024). The development of some  
 437 of these observatories into hydrological super-sites, contingent on the availability of resources, would allow for the

438 investigation of specific questions, such as karst hydrology, water scarcity, floodplains, forest hydrology, precision  
439 agriculture, and different runoff generation mechanisms. This approach enables targeted investigations at specific  
440 locations with high-resolution data, which can then be exploited to support the development of high-fidelity models.  
441 It is similarly feasible to reverse this scenario. Should a super-site situated in a particular bioclimatic zone yield  
442 scientific breakthroughs, it may be possible to establish a network of HOs in regions exhibiting analogous hydrological  
443 behavior. The key factor is to leverage the strengths of each approach while operating within the confines of the  
444 allocated budgetary resources. By integrating the strengths of the two approaches, one can attain a balance between  
445 representativeness (a distributed network) and a detailed understanding (super-sites). This approaches ensures the  
446 optimal exploitation of resources while maximizing the scientific output.

447

## 448 **5. Concluding remarks**

449 To address water-related issues at the catchment scale across diverse global contexts, it is imperative to develop  
450 adaptation and mitigation strategies that are grounded in evidence gathered in HOs. The previous situation, which was  
451 characterized by a myriad of relatively unconnected, moderately and differently instrumented HOs that were supported  
452 by grant-to-grant funding, has resulted in significant but fragmented knowledge. This has impeded comparative  
453 studies and has hindered scientific progress. New initiatives are being proposed with the objective of enhancing the  
454 coordination of HO networks, thereby enabling efficient cross-site synthesis. In light of financial constraints, we need  
455 to identify a common vision for the optimal allocation of resources.

456 A network of numerous HOs provides broad spatial coverage, enabling the capture of variations in environmental  
457 conditions across diverse regions, ecosystems, and land uses. Environmental change can manifest itself differently  
458 across regions due to the influence of local climate, geography, and human activities. A network of numerous  
459 observatories offers the opportunity to monitor these interactions, capture feedbacks, teleconnections, and cross-scale  
460 dynamics that may not be observable at individual observatories. In contrast, a limited number of intensively  
461 instrumented observatories permit the collection of high-resolution data, the testing of novel hypotheses, and  
462 informing comprehensive process-oriented hydrological models. This choice can capture variations at smaller spatial  
463 and finer temporal scales, thereby providing a more nuanced understanding of environmental and hydrological  
464 processes. Nevertheless, the strategy of pooling all financial efforts into a limited number of intensely instrumented  
465 hydrological observatories will exacerbate the issue of knowledge transferability and geographic bias in hydrological  
466 data and understanding. It is therefore necessary to devise strategies for generalization, potentially drawing inspiration  
467 from the Prediction in Ungauged Basins initiative.

468 In the context of accelerated global transformation, there is a pressing need to establish a network of HOs. The question  
469 of how to organize and manage such a global network, including the number of observatories, remains a topic of  
470 discussion. Both distributed networks and super-sites offer valuable contributions to the advancement of hydrological  
471 understanding. We envision a dynamic hybrid approach that combines the two aforementioned visions in a manner  
472 that does not exclude either of them from consideration. It is important to raise public awareness about the significance  
473 of hydrological research and its linkages with a multitude of other disciplines, including atmospheric science, soil  
474 science, biochemistry, pedology, ecology, microbiology, geology, plant physiology, and remote sensing. Such an  
475 approach can garner support and increase funding opportunities. It is our hope that all hydrologists will engage in a  
476 discussion process with the aim of refining and building upon the ideas presented in this paper.

477

478 **Acknowledgments**

479 This paper is a result of the presentations and discussions held at the 8<sup>th</sup> Galileo Conference, entitled “A European Vision for  
480 Hydrological Observations and Experimentation”, which took place in Naples, Italy, from June 12th to 15th, 2023. In his keynote  
481 lecture, entitled “The Future of Hydrology: Nature or Nurture”, on June 14th, 2023, Günter Blöschl discussed the concept of  
482 focusing research efforts on a limited number of catchments worldwide. This prompted a lively exchange of opinions among the  
483 attendees. The USDA is an equal opportunity provider and employer. Salvatore Manfreda acknowledges the RETURN Extended  
484 Partnership which received funding from the Next-Generation EU (National Recovery and Resilience Plan - NRRP, Mission 4,  
485 Component 2, Investment 1.3 - D.D. 1243 2/8/2022, PE0000005). Paolo Nasta and Nunzio Romano acknowledge the partial  
486 support provided by the the Ministry of University and Research (MiUR) through the MiUR-PRIN-PNRR-2022 project entitled  
487 “ASAP: Assessing and mapping novel agroecosystem vulnerability and resilience indicators in southern Italy” (grant  
488 P20228L3KX). The authors would also like to express their gratitude to the executive editor in charge, Thom Bogaard, and the two  
489 reviewers (A. Guswa and an anonymous) for their constructive comments, which have contributed to enhance the quality of this  
490 paper.

491  
492 **Appendix**

493 The 8<sup>th</sup> Galileo Conference “A European Vision for Hydrological Observations and Experimentation” was held in  
494 Naples (Italy) on 12-15 June 2023. Upon presentations and discussions, we report the most intriguing questions in  
495 hydrology that emerged from the conference. Additionally, we conducted a literature review and identified several  
496 key points that warrant further investigation:  
497

---

**Research questions in Scenario 1**

---

- 1 How can we improve the quantity and quality of measurements in data-poor regions?
- 2 Where and how can we deploy the sensors to get the most information without wasting too much effort?
- 3 Are measurements taken in the past still valid in the future? How will accuracy/precision change with as technological advances? Do we need to remove all “inaccurate” historical data and keep only “currently accurate” data? Is the assumption of a steady hydrological system valid? Can we simplify the system by linearizing a nonlinear system behavior?
- 4 What role(s) do continuous and ephemeral water bodies, including ponds, lakes, rivers, streams, marshes, swamps, etc., play in influencing water quantity and quality in the catchment?

---

**Research questions in Scenario 2**

---

- 1 What are the hydrologic laws at the catchment scale, and how do they change with scale?
- 2 How can we use innovative technologies to measure surface and subsurface properties, states, and fluxes at a range of spatial and temporal scales?
- 3 How can different multi-scale observations be assimilated into a hydrological model to improve model predictability?
- 4 How do we obtain large-scale flux measurements and feedbacks to analyze the water dynamics within and between the compartments of the groundwater-soil-plant-atmosphere continuum?
- 5 How is the water cycle influenced by the other cycles (carbon, nitrogen, etc.)?
- 6 How can the dynamics and feedbacks at groundwater-soil, groundwater-surface water, soil-plant, soil-atmosphere, and plant-atmosphere interfaces be assessed?
- 7 How do we incorporate plant physiological aspects into hydrological models?

---

**Research questions in both scenarios**

---

- 1 What is the impact of preferential flow on catchment-scale water flow dynamics?
  - 2 How can remote sensing provide more reliable information on soil moisture, changes in water storage, surface energy balance, and evapotranspiration at appropriate spatial and temporal scales (Lettenmaier et al., 2015)?
  - 3 What causes spatial heterogeneity and homogeneity in runoff, evapotranspiration, subsurface water and material fluxes (carbon and other nutrients, sediments), and in their sensitivity to their controls (e.g., snowfall regime, aridity, response coefficients)?
  - 4 How can hydrological models be adapted to be able to extrapolate changing conditions, including changing vegetation dynamics?
-

- 5 How can we disentangle and reduce model structure/parameter/input uncertainty in hydrological prediction?

---

- 6 Is it better to emphasize uncertainty or causality?

---

- 7 How do vegetation types, distribution, and dynamics shape hydrological processes, particularly in terms of water quality, quantity, and energy fluxes at the catchment scale?

---

- 8 How can we integrate the different spatial and temporal scales of observations, processes, and models?

---

- 9 How can we develop socio-hydrological models by allowing for anthropogenic disturbances in the ecosystem?

---



- 499 **References**
- 500 Addor, N., Newman, A. J., Mizukami, N., and Clark, M. P.: The CAMELS data set: catchment attributes and  
501 meteorology for large-sample studies, *Hydrol. Earth Syst. Sci.*, 21, 5293–5313, [https://doi.org/10.5194/hess-21-](https://doi.org/10.5194/hess-21-5293-2017)  
502 5293-2017, 2017.
- 503 Addor, N., Nearing, G., Prieto, C., Newman, A. J., Le Vine, N., and Clark, M. P.: A ranking of hydrological  
504 signatures based on their predictability in space, *Water Resour. Res.*, 54, 8792–8812,  
505 <https://doi.org/10.1029/2018WR022606>, 2018.
- 506 Alvarez-Garreton, C., Mendoza, P. A., Boisier, J. P., Addor, N., Galleguillos, M., Zambrano-Bigiarini, M., ... and  
507 Ayala, A.: The CAMELS-CL dataset: catchment attributes and meteorology for large sample studies—Chile  
508 dataset, *Hydrol. Earth Syst. Sci.*, 22(11), 5817–5846, 2018.
- 509 Anderson, S. P., Bales, R. C., and Duffy, C. J.: Critical Zone Observatories: Building a network to advance  
510 interdisciplinary study of Earth surface processes, *Mineral. Mag.*, 72(1), 7–10, 2008.
- 511 Arora, B., Kuppel, S., Wellen, C., Oswald, C., Groh, J., Payandi-Rolland, D., Stegen, J., Coffinet, S.: Building Cross-  
512 Site and Cross-Network collaborations in critical zone science, *J. Hydrol.*, 618, 129248  
513 <https://doi.org/10.1016/j.jhydrol.2023.129248>, 2023
- 514 Baatz, R., Franssen, H.J.H., Han, X., Hoar, T., Bogena, H.R., and Vereecken, H.: Evaluation of a cosmic-ray neutron  
515 sensor network for improved land surface model prediction, *Hydrol. Earth Syst. Sci.* 21 (5), 2509–2530.  
516 <https://doi.org/10.5194/hess-21-2509-2017>, 2017.
- 517 Baatz, R., Sullivan, P. L., Li, L., Weintraub, S. R., Loescher, H. W., Mirtl, M., Groffman, P. M., Wall, D. H., Young,  
518 M., White, T., Wen, H., Zacharias, S., Kühn, I., Tang, J., Gaillardet, J., Braud, I., Flores, A. N., Kumar, P., Lin,  
519 H., Ghezzehei, T., Jones, J., Gholz, H. L., Vereecken, H., and Van Looy, K.: Steering operational synergies in  
520 terrestrial observation networks: opportunity for advancing Earth system dynamics modelling, *Earth Syst. Dynam.*,  
521 9, 593–609, <https://doi.org/10.5194/esd-9-593-2018>, 2018.
- 522 Bates, C. G., and Henry, A. J.: Monthly weather, *Monthly Weather Review*, 56(3), 79–85, 1928.
- 523 Bauser, H. H., Kim, M., Ng, W.-R., Bugaj, A., and Troch, P. A.: Richards equation at the hillslope scale: Can we  
524 resolve the heterogeneity of soil hydraulic material properties?, *Water Resour. Res.*, 58, e2022WR032294.  
525 <https://doi.org/10.1029/2022WR032294>, 2022.
- 526 Bechtold, M., De Lannoy, G. J. M., Koster, R. D., Reichle, R. H., Mahanama, S. P., Bleuten, W., et al.: PEAT-CLSM:  
527 A specific treatment of peatland hydrology in the NASA Catchment Land Surface Model, *J. Adv. Model. Earth*  
528 *Syst.*, 11, 2130–2162. <https://doi.org/10.1029/2018MS001574>. 2019.
- 529 Benettin, P., Rodriguez, N. B., Sprenger, M., Kim, M., Klaus, J., Harman, C. J., et al.: Transit time estimation in  
530 catchments: Recent developments and future directions, *Water Resour. Res.*, 58, e2022WR033096,  
531 <https://doi.org/10.1029/2022WR033096>, 2022.
- 532 Beven, K., and Germann, P.: Macropores and water flow in soils revisited, *Water Resour. Res.*, 49, 3071 – 3092.  
533 <https://doi.org/10.1002/wrcr.20156>, 2013.
- 534 Beven, K. J.: On hypothesis testing in hydrology: Why falsification of models is still a really good idea, *WIREs*  
535 *Water*, 5, e1278. <https://doi.org/10.1002/wat2.1278>, 2018.
- 536 Blöschl, G., Hydrologic synthesis: Across processes, places, and scales, *Water Resour. Res.*, 42,  
537 W03S02, [doi:10.1029/2005WR004319](https://doi.org/10.1029/2005WR004319), 2006.
- 538 Blöschl, G., Blaschke, A. P., Broer, M., Bucher, C., Carr, G., Chen, X., Eder, A., Exner-Kittridge, M., Farnleitner, A.,  
539 Flores-Orozco, A., Haas, P., Hogan, P., Kazemi Amiri, A., Oismüller, M., Parajka, J., Silasari, R., Stadler, P.,  
540 Strauß, P., Vreugdenhil, M., Wagner, W., and Zessner, M.: The Hydrological Open Air Laboratory (HOAL) in  
541 Petzenkirchen: a hypothesis-driven observatory, *Hydrol. Earth Syst. Sci.*, 20, 227–255, [doi:10.5194/hess-20-227-](https://doi.org/10.5194/hess-20-227-2016)  
542 2016, 2016.
- 543 Blöschl, G., Bierkens, M. F. P., Chambel, A., Cudennec, C., Destouni, G., Fiori, A., et al.: Twenty-three unsolved  
544 problems in hydrology (UPH)—A community perspective, *Hydrolog. Sci. J.*, 64(10), 1141–1158, 2019.
- 545 Bogena, H., Schulz, K., and Vereecken, H.: Towards a network of observatories in terrestrial environmental research,  
546 *Adv. Geosci.*, 9, 109–114, <https://doi.org/10.5194/adgeo-9-109-2006>, 2006.
- 547 Bogena, H.R., White, T., Bour, O., Li, X., and Jensen, K.H.: Toward better understanding of terrestrial processes  
548 through long-term hydrological observatories, *Vadose Zone J.*, 17, 180194. [doi:10.2136/vzj2018.10.0194](https://doi.org/10.2136/vzj2018.10.0194), 2018.
- 549 Bogena, H. R., Schrön, M., Jakobi, J., Ney, P., Zacharias, S., Andreasen, M., et al.: COSMOS-Europe: A European  
550 network of cosmic-ray neutron soil moisture sensors. *Earth Syst. Sci. Data*, 14(3), 1125–1151.  
551 <https://doi.org/10.5194/essd-14-1125-2022>, 2022.
- 552 Brantley, S. L., McDowell, W. H., Dietrich, W. E., White, T. S., Kumar, P., Anderson, S. P., Chorover, J., Lohse, K.  
553 A., Bales, R. C., Richter, D. D., Grant, G., and Gaillardet, J.: Designing a network of critical zone observatories  
554 to explore the living skin of the terrestrial Earth, *Earth Surf. Dynam.*, 5, 841–860, [https://doi.org/10.5194/esurf-5-](https://doi.org/10.5194/esurf-5-841-2017)  
555 841-2017, 2017
- 556 Brooks, P. D., Chorover, J., Fan, Y., Godsey, S. E., Maxwell, R. M., McNamara, J. P., and Tague, C.: Hydrological  
557 partitioning in the critical zone: Recent advances and opportunities for developing transferable understanding of  
558 water cycle dynamics, *Water Resour. Res.*, 51, 6973–6987, [doi:10.1002/2015WR017039](https://doi.org/10.1002/2015WR017039), 2015.
- 559 Bruijnzeel, L. A.: (De)Forestation and dry season flow in the tropics: A closer look. *J. Trop. For. Sci.*, 1(3), 229–243,  
560 1989.

561 Bunting, U., Urban, O., Krusic, P.J. et al.: Recent European drought extremes beyond Common Era background  
562 variability, *Nature Geoscience*, 14, 190–196, doi:10.1038/s41561-021-00698-0, 2021.

563 Burdun, I., Bechtold, M., Aurela, M., De Lannoy, G., Desai, A. R., Humphreys, E., et al.: Hidden becomes clear:  
564 Optical remote sensing of vegetation reveals water table dynamics in northern peatlands, *Remote Sens. Environ.*,  
565 296, 113736, <https://doi.org/10.1016/j.rse.2023.113736>, 2023.

566 Burt, T. P., and McDonnell, J. J.: Whither field hydrology? The need for discovery science and outrageous  
567 hydrological hypotheses, *Water Resour. Res.*, 51(8), 5919–5928, 2015.

568 Chagas, V. B., Chaffe, P. L., Addor, N., Fan, F. M., Fleischmann, A. S., Paiva, R. C., and Siqueira, V. A.: CAMELS-  
569 BR: hydrometeorological time series and landscape attributes for 897 catchments in Brazil, *Earth Syst. Sci. Data*,  
570 12(3), 2075–2096, 2020.

571 Chaussé C., Leroyer, C., Girardclos, O. et al.: Holocene history of the River Seine, Paris, France: bio-  
572 chronostratigraphic and geomorphological evidence from the Quai-Branly, *The Holocene*, 18, 967–980,  
573 doi:10.1177/0959683608093, 2008.

574 Colliander, A., Reichle, R., Crow, W., Cosh, M., Chen, F., Chan, S., Das, N., Bindlish, R., Chaubell, J., Kim, S., Liu,  
575 Q., O’Neill, P., Dunbar, S., Dang, L., Kimball, J., Jackson, T., AlJassar, H., Asanuma, J., Bhattacharya, B., Berg,  
576 A., Bosch, D., Bourgeau-Chavez, L., Caldwell, T., Calvet, J.-C., Collins, C.H., Jensen, K., Livingston, S., López-  
577 Baeza, E., Martínez-Fernández, J., McNairn, H., Moghaddam, M., Montzka, C., Notarnicola, C., Pellarin, T., Pfeil,  
578 I., Pulliainen, J., Ramos, J., Seyfried, M., Starks, P., Su, Z., Thibeault, M., van der Velde, R., Vreugdenhil, M.,  
579 Walker, J., Zeng, Y., Zribi, M., Entekhabi, D., Yueh, S.: Validation of soil moisture data products from the NASA  
580 SMAP mission, *IEEE J. Sel. Top. Appl. Earth Observ. Remote Sens.*, 15, 364–392, <https://doi.org/10.1109/JSTARS.2021.3124743>, 2021.

582 Dari, J., Brocca, L., Modanesi, S., Massari, C., Tarpanelli, A., Barbetta, S., Quast, R., Vreugdenhil, M., Freeman, V.,  
583 Barella-Ortiz, A., Quintana-Seguí, P., Bretreger, D., and Volden, E.: Regional data sets of high-resolution (1 and  
584 6 km) irrigation estimates from space, *Earth Syst. Sci. Data*, 15, 1555–1575, [https://doi.org/10.5194/essd-15-1555-](https://doi.org/10.5194/essd-15-1555-2023)  
585 2023, 2023.

586 Davis, W. M.: The value of outrageous geological hypotheses. *Science*, 63(1636), 463–468, 1926.

587 De Lannoy, G.J.M., Bechtold, M., Albergel, C., Brocca, L., Calvet, J.-C., Carrasi, A., Crow, W.T., de Rosnay, P.,  
588 Durand, M., Forman, B., Geppert, G., Girotto, M., Hendricks Franssen, H.-J., Jonas, T., Kumar, S., Lievens, H.,  
589 Lu, Y., Massari, C., Pauwels, V.R.N., Reichle, R.H., and Steele-Dunne, S.: Perspective on satellite-based land data  
590 assimilation to estimate water cycle components in an era of advanced data availability and model sophistication,  
591 *Front. Water*, 4, 981745, doi: 10.3389/frwa.2022.981745, 2022.

592 Dewar, N., and Knight, R., Estimation of the top of the saturation zone from airborne electromagnetic data,  
593 *Geophysics*, 85(8), EN63–EN76. <https://doi.org/10.1190/geo2019-0539.1>, 2020.

594 Dugdale, S. J., Klaus, J., and Hannah, D. M.: Looking to the skies: realising the combined potential of drones and  
595 thermal infrared imagery to advance hydrological process understanding in headwaters. *Water Resour. Res.*, 58(2),  
596 e2021WR031168, 2022.

597 Durand, M., Barros, A., Dozier, J., Adler, R., Cooley, S., Entekhabi, D., et al.: Achieving breakthroughs in global  
598 hydrologic science by unlocking the power of multisensor, multidisciplinary Earth observations, *AGU Advances*,  
599 2, e2021AV000455, <https://doi.org/10.1029/2021AV000455>, 2021

600 Ellison, D., Futter, M.N., Bishop, K.: On the forest cover-water yield debate: From demand- to supply-side thinking,  
601 *Global Change Biol.*, 18, 806–820, 2012.

602 Evaristo, J., Kim, M., van Haren, J., Pangle, L. A., Harman, C. J., Troch, P. A., and McDonnell, J. J.: Characterizing  
603 the fluxes and age distribution of soil water, plant water, and deep percolation in a model tropical ecosystem,  
604 *Water Resour. Res.*, 55, 3307–3327, <https://doi.org/10.1029/2018WR023265>, 2019.

605 Ehret, U., Gupta, H. V., Sivapalan, M., Weijs, S. V., Schymanski, S. J., Blöschl, G., ... and Winsemius, H. C.:  
606 Advancing catchment hydrology to deal with predictions under change, *Hydrol. Earth Syst. Sci.*, 18(2), 649–671,  
607 2014.

608 Evaristo, J., Kim, M., van Haren, J., Pangle, L. A., Harman, C. J., Troch, P. A., and McDonnell, J. J.: Characterizing  
609 the fluxes and age distribution of soil water, plant water, and deep percolation in a model tropical ecosystem, *Water*  
610 *Resour. Res.*, 55, 3307 – 3327, <https://doi.org/10.1029/2018WR023265>, 2019.

611 Fan, Y., Clark, M., Lawrence, D. M., Swenson, S., Band, L. E., Brantley, S. L., et al.: Hillslope hydrology in global  
612 change research and Earth system modeling, *Water Resour. Res.*, 55, 1737–1772,  
613 <https://doi.org/10.1029/2018WR023903>, 2019.

614 Fu, T., Liu, J., Gao, H., Qi, F., Wang, F., and Zhang, M.: Surface and subsurface runoff generation processes and  
615 their influencing factors on a hillslope in northern China, *Sci. Total Environ.*, 906, 167372,  
616 <https://doi.org/10.1016/j.scitotenv.2023.167372>, 2024.

617 Gaillardet, J., Braud, I., Hankard, F., Anquetin, S., Bour, O., Dorfliger, N., de Dreuzy, J.R., Galle, et al.: OZCAR:  
618 The French Network of Critical Zone Observatories, *Vadose Zone J.*, 17, 1–24 180067.  
619 <https://doi.org/10.2136/vzj2018.04.0067>, 2018.

620 Gao, H., Fenicia, F., and Savenije, H. H. G.: HESS Opinions: Are soils overrated in hydrology?, *Hydrol. Earth Syst.*  
621 *Sci.*, 27, 2607–2620, <https://doi.org/10.5194/hess-27-2607-2023>, 2023.

622 Goodrich, D. C., Heilman, P., Anderson, M., Baffaut, C., Bonta, J., Bosch, D., et al.: The USDA-ARS Experimental  
623 Watershed Network: Evolution, lessons learned, societal benefits, and moving forward, *Water Resour. Res.*,  
624 57,e2019WR026473, <https://doi.org/10.1029/2019WR026473>, 2021.

625 Guzinski, R., and Nieto, H.: Evaluating the feasibility of using Sentinel-2 and Sentinel-3 satellites for high-resolution  
626 evapotranspiration estimations, *Remote Sens. Environ.*, 221, 157–172. <https://doi.org/10.1016/j.rse.2018.11.019>,  
627 2019.

628 Guzinski, R., Nieto, H., Sandholt, I., and Karamitilios, G.: Modelling high-resolution actual evapotranspiration  
629 through Sentinel-2 and Sentinel-3 data fusion, *Remote Sensing*, 12(9), 1433, <https://doi.org/10.3390/rs12091433>,  
630 2020.

631 Han, Q., Zeng, Y., Zhang, L. et al.: Global long term daily 1 km surface soil moisture dataset with physics informed  
632 machine learning, *Sci Data*, 10, 101, <https://doi.org/10.1038/s41597-023-02011-7>, 2023.

633 Harms, U., Koeberl, C., and Zoback, M. D.: Continental scientific drilling: a decade of progress, and challenges for  
634 the future, Springer, Berlin/Heidelberg, p. 366, 2007.

635 Hoffmann, M., Jurisch, N., Albiac Borraz, E., Hagemann, U., Drösler, M., Sommer, M., and Augustin, J.: Automated  
636 modeling of ecosystem CO<sub>2</sub> fluxes based on periodic closed chamber measurements: A standardized conceptual  
637 and practical approach, *Agric. For. Meteorol.*, 200, 30–45, 2015.

638 Holländer, H.M., Blume, T., Bormann, H., Buytaert, W., Chirico, G.B., Exbrayat, J.-F., Gustafsson, D., Hölzel, H.,  
639 Kraft, P., Stamm, C., Stoll, S., Blöschl, G., Flühler, H.: Comparative predictions of discharge from an artificial  
640 catchment (Chicken Creek) using sparse data, *Hydrol. Earth Syst. Sci.*, 13, 2069 - 2094, 2009.

641 Hrachowitz, M., Savenije, H. H. G., Blöschl, G., McDonnell, J. J., Sivapalan, M., Pomeroy, J. W., ... and Cudennec,  
642 C.: A decade of Predictions in Ungauged Basins (PUB)—a review, *Hydrol. Sci. J.*, 58(6), 1198-1255,  
643 10.1080/02626667.2013.803183, 2013.

644 Jensen, C.K., McGuire, K.J., McLaughlin, D.L. et al.: Quantifying spatiotemporal variation in headwater stream  
645 length using flow intermittency sensors, *Environ. Monit. Assess.*, 191, 226, <https://doi.org/10.1007/s10661-019-7373-8>, 2019.

647 Jones, J.A., Creed, I. F., Hatcher, K. L., Warren, R. J., Adams, M. B., Benson, M. H., Boose, E., Brown, W. A.,  
648 Campbell, J. L., Covich, A., Clow, D. W., Dahm, C. N., Elder, K. , Ford, C. R., Grimm, N. B., Henshaw, D. L.,  
649 Larson, K. L., Miles, E. S., Miles, K. M., Sebestyen, S. D., Spargo, A. T., Stone, A. B., Vose, J. M., Williams, M.  
650 W.: Ecosystem Processes and Human Influences Regulate Streamflow Response to Climate Change at Long-Term  
651 Ecological Research Sites, *BioScience*, 62, 390–404, <https://doi.org/10.1525/bio.2012.62.4.10>, 2012.

652 Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P., and Zacharias, S.: Footprint characteristics revised for  
653 field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resour. Res.*, 51(7), 5772-5790, 2015.

654 Kim, Y., Band, L.E., Song, C.: The influence of forest regrowth on the stream discharge in the North Carolina  
655 Piedmont Watersheds, *J. Am. Water Resour. Assoc.*, 50, 57-73, 2013.

656 Kirchner, J. W.: Aggregation in environmental systems – Part 1: Seasonal tracer cycles quantify young water  
657 fractions, but not mean transit times, in spatially heterogeneous catchments, *Hydrol. Earth Syst. Sci.*, 20, 279 –  
658 297. <https://doi.org/10.5194/hess-20-279-2016>, 2016.

659 Klaus, J., and McDonnell, J. J.: Hydrograph separation using stable isotopes: Review and evaluation, *J. Hydrol.*, 505,  
660 47–64, <https://doi.org/10.1016/j.jhydrol.2013.09.006>, 2013.

661 Kuentz, A., Arheimer, B., Hundecha, Y., and Wagener, T.: Understanding hydrologic variability across Europe  
662 through catchment classification, *Hydrol. Earth Syst. Sci.*, 21(6), 2863-2879, 2017.

663 Kulmala, M.: Build a global Earth observatory, *Nature*, 553(7686), 21-23, 2018.

664 Kumar, S., Kolassa, J., Reichle, R., Crow, W., de Lannoy, G., de Rosnay, P., MacBean, N., Girotto, M., Fox, A.,  
665 Quaipe, T., Draper, C., Forman, B., Balsamo, G., Steele-Dunne, S., Albergel, C., Bonan, B., Calvet, J., Dong, J.,  
666 Liddy, H. and Ruston, B.: An agenda for land data assimilation priorities: realizing the promise of terrestrial water,  
667 energy, and vegetation observations from space, *J. Adv. Model. Earth Syst.*, 14, e2022MS003259.  
668 <https://doi.org/10.1029/2022ms003259>, 2022.

669 Lettenmaier, D. P., Alsdorf, D., Dozier, J., Huffman, G. J., Pan, M., and Wood, E. F.: Inroads of remote sensing into  
670 hydrologic science during the WRR era, *Water Resour. Res.*, 51, 7309-7342, 2015.

671 Li, Z., Li, X., Zhou, S., Yang, X., Fu, Y., Miao, C., Wang, S., Zhang, G., Wu, X., Yang, C., Deng, Y.: A comprehensive  
672 review on coupled processes and mechanisms of soil-vegetation-hydrology, and recent research advances, *Science*  
673 *China Earth Sci.*, 65(11), 2083–2114, <https://doi.org/10.1007/s11430-021-9990-5>, 2022.

674 Lievens, H., Brangers, I., Marshall, H.-P., Jonas, T., Olefs, M., and De Lannoy, G.: Sentinel-1 snow depth retrieval at  
675 sub-kilometer resolution over the European Alps, *The Cryosphere*, 16, 159–177, <https://doi.org/10.5194/tc-16-159-2022>, 2022.

677 Loescher, H. W., Vargas, R., Mirtl, M., Morris, B., Pauw, J., Yu, X., et al.: Building a global ecosystem research  
678 infrastructure to address global grand challenges for macrosystem ecology, *Earth's Future*, 10(5), e2020EF001696,  
679 <https://doi.org/10.1029/2020ef001696>, 2022.

680 Loritz, R., Bassiouni, M., Hildebrandt, A., Hassler, S.K., and Zehe, E.: Leveraging sap flow data in a catchment-scale  
681 hybrid model to improve soil moisture and transpiration estimates, *Hydrol. Earth Syst. Sci.*, 26 (18), 4757–4771,  
682 <https://doi.org/10.5194/hess-26-4757-2022>, 2022.

683 Manfreda, S., Miglino, D., Saddi, K. C., Jomaa, S., Etner, A., Perks, M., Strelnikova, D., Peña-Haro, S., Maddock, I.,  
684 Tauro, F., Grimaldi, S., Zeng, Y., Gonçalves, G., Bogaard, T., van Emmerik, T., Bussettini, M., Mariani, S.,  
685 Marchetti, G., Lastoria, B., Su, B., and Rode, M.: Advancing river monitoring using image-based techniques:  
686 challenges and opportunities, *Hydrol. Sci. J.*, 69, 657–677, <https://doi.org/10.1080/02626667.2024.2333846>, 2024.

687 Manfreda, S., McCabe, M. F., Miller, P. E., Lucas, R., Pajuelo Madrigal, V., Mallinis, G., ... and Toth, B.: On the use  
688 of unmanned aerial systems for environmental monitoring. *Remote sens.*, 10(4), 641, 2018.

689 McCabe, M.F., et al.: The future of Earth observation in hydrology, *Hydrol. Earth Syst. Sci.*, 21(7), 3879–3914.  
690 doi:10.5194/hess-21-3879-2017, 2017.

691 McDonnell, J. J., et al.: Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology,  
692 *Water Resour. Res.*, 43, W07301, doi:10.1029/2006WR005467, 2007.

693 McDonnell, J. J., McGuire, K., Aggarwal, P., Beven, K., Biondi, D., Destouni, G., et al.: How old is stream water?  
694 Open questions in catchment transit time conceptualization, modeling, and analysis, *Hydrol. Process.*, 24(12),  
695 1745–1754. <https://doi.org/10.1002/hyp.7796>, 2010.

696 McDonnell, J. J.: The two water worlds hypothesis: Ecohydrological separation of water between streams and  
697 trees? *WIREs Water*, 1, 323 – 329. <https://doi.org/10.1002/wat2.1027>, 2014.

698 Medina, R., Pham, C., Plumlee, M.H., Hutchinson, A., Becker, M.W., and O’Connell, P.J.: Distributed temperature  
699 sensing to measure infiltration rates across a groundwater recharge basin, *Groundwater*, 58(6), 913–923,  
700 <https://doi.org/10.1111/gwat.13007>, 2020.

701 Melton, F.S., Huntington, J., Grimm, R., Herring, J., Hall, M., Rollison, D., Erickson T., et al.: OpenET: Filling a  
702 Critical DataGap in Water Management for the Western United States, *J. Am. Water Resour. As.*, 58 (6), 971–  
703 994, <https://doi.org/10.1111/1752-1688.12956>, 2022.

704 Montanari, A., Young, G., Savenije, H. H. G., Hughes, D., Wagener, T., Ren, L. L., ... and Belyaev, V.: Panta Rhei—  
705 everything flows: Change in hydrology and society—the IAHS scientific decade 2013–2022, *Hydrol. Sci. J.*, 58(6),  
706 1256–1275, 2013.

707 Mwangi, S., Zeng, Y., Montzka, C., Yu, L., and Su, Z.: Assimilation of cosmic-ray neutron counts for the estimation  
708 of soil ice content on the eastern Tibetan Plateau, *J. Geophys. Res.: Atmospheres*, 125, e2019JD031529,  
709 <https://doi.org/10.1029/2019JD031529>, 2020.

710 Nasta, P., Boaga, J., Deiana, R., Cassiani, G., and Romano, N.: Comparing ERT- and scaling-based approaches to  
711 parameterize soil hydraulic properties for spatially distributed model applications, *Adv. Water Resour.*, 126, 155–  
712 167, 2019.

713 Nasta, P., Palladino, M., Ursino, N., Saracino, A., Sommella, A., and Romano, N.: Assessing long-term impact of  
714 land-use change on hydrological ecosystem functions in a Mediterranean upland agro-forestry catchment, *Sci.*  
715 *Total Environ.*, 605–606, 1070–1082, 2017.

716 Nearing, G., Cohen, D., Dube, V., Gauch, M., Gilon, O., Harrigan, S., ... and Matias, Y.: Global prediction of extreme  
717 floods in ungauged watersheds, *Nature*, 627(8004), 559–563, 2024.

718 Neary, D., Hayes, D., Rustad, L., Vose, J., Gottfried, G., Sebestyen, S., Johnson, S., Swanson, F. and Adams, M.: US  
719 Forest Service Experimental Forests and Ranges Network: a continental research platform for catchment-scale  
720 research, In: *Revisiting Experimental Catchment Studies in Forest Hydrology* (ed. by A. E. Webb), 49–57  
721 (Proceedings of a Workshop held during the XXV IUGG General Assembly in Melbourne, June–July 2011) IAHS  
722 Publ. 353. IAHS Press, Wallingford, UK (this volume), 2012.

723 Noto, S., Durigetto, N., Tauro, F., Grimaldi, S., and Botter, G.: Characterizing space-time channel network dynamics  
724 in a Mediterranean intermittent catchment of central Italy combining visual surveys and cameras, *Water Resour.*  
725 *Res.*, 60, e2023WR034682, <https://doi.org/10.1029/2023WR034682>, 2024.

726 Painter, T.H., Berisford, D.F., Boardman, J.W., Bormann, K.J., Deems, J.S., Gehrke, F., Hedrick, A., Joyce, M.,  
727 Laidlaw, R., Marks, D., Mattmann, C., McGurk, B., Ramirez, P., Richardson, M., Skiles, S.K., Seidel, F.C., and  
728 Winstal, A.: The airborne snow observatory: fusion of scanning lidar, imaging spectrometer, and physically-based  
729 modeling for mapping snow water equivalent and snow albedo, *Remote Sens. Environ.*, 184, 139 – 152,  
730 <https://doi.org/10.1016/j.rse.2016.06.018>, 2016.

731 Pangle, L. A., DeLong, S. B., Abramson, N., Adams, J., Barron-Gafford, G. A., Breshears, D. D., ... and Zeng, X.:  
732 The Landscape Evolution Observatory: A large-scale controllable infrastructure to study coupled Earth-surface  
733 processes, *Geomorphology*, 244, 190–203, 2015.

734 Peng, W., Lu, Y., Xie, X., Ren, T., and Horton, R.: An improved thermo-TDR technique for monitoring soil thermal  
735 properties, water content, bulk density, and porosity, *Vadose Zone J.*, 18, 190026, doi:10.2136/vzj2019.03.0026,  
736 2019.

737 Penna, D., et al.: Ideas and perspectives: Tracing ecosystem water fluxes using hydrogen and oxygen stable isotopes—  
738 Challenges and opportunities from an interdisciplinary perspective, *Biogeosciences*, 15(21), 6399–6415,  
739 <https://doi.org/10.5194/bg-15-6399-2018>, 2018.

740 Peters-Lidard, C. D., Hossain, F., Leung, L. R., McDowell, N., Rodell, M., Tapiador, F. J., et al.: 100 years of progress  
741 in hydrology, In Chapter 25 in *AMS Meteorological Monographs* (Vol. 59, pp. 50–25.51). Boston, MA: American,  
742 <https://doi.org/10.1175/amsmonographs-d-18-0019.1>, 2018.

743 Poyatos, R., Granda, V., Flo, V., Adams, M.A., Adorján, B., Aguadé, D., Aidar, M.P.M., Allen, S., Alvarado-  
744 Barrientos, M.S., Anderson-Teixeira, K.J., et al.: Global transpiration data from sap flow measurements: the  
745 SAPFLUXNET database, *Earth Syst. Sci. Data*, 13, 2607-2649, doi.org/10.5194/essd-13-2607-2021, 2021.

746 Rabe, B., Heuzé, C., Regnery, J., Aksenov, Y., Allerholt, J., Athanase, M., ... and Zhu, J.: Overview of the MOSAiC  
747 expedition: Physical oceanography. *Elementa: Science of the Anthropocene*, 10(1), 00062.  
748 https://doi.org/10.1525/elementa.2021.00062, 2022.

749 Refsgaard, J.C., Stisen, S., and Koch, J.: Hydrological process knowledge in catchment modelling – Lessons and  
750 perspectives from 60 years development, *Hydrol. Process.*, 36, e14463, https://doi.org/10.1002/hyp.1446, 2021.

751 Reid, W.V., Chen, D., Goldfarb, L., Hackmann, H., Lee, Y.T., Mokhele, K., Ostrom, E., Raivio, K., Rockström, J.,  
752 Schellnhuber, H.J., and Whyte, A.: Earth system science for global sustainability: Grand Challenges, *Science* 330,  
753 916-917, 2010.

754 Rinderer, M., Van Meerveld, H. J., and McGlynn, B. L.: From points to patterns: using groundwater time series  
755 clustering to investigate subsurface hydrological connectivity and runoff source area dynamics, *Water Resour.*  
756 *Res.*, 55(7), 5784-5806, 2019.

757 Romano, N.: Soil moisture at local scale: Measurements and simulations, *J. Hydrol.*, 516, 6-20, 2014.

758 Romano, N., Nasta, P., Bogena, H., De Vita, P., Stellato, L., and Vereecken, H.: Monitoring hydrological processes  
759 for land and water resources management in a Mediterranean ecosystem: The Alento River catchment observatory,  
760 *Vadose Zone J.*, 17, 0042, 2018.

761 Romano, N., Szabò, B., Belmonte, A., Castrignanò, A., Ben Dor, E., Francos, N., and Nasta, P.: Mapping soil  
762 properties for unmanned aerial system-based environmental monitoring. In “Unmanned Aerial Systems for  
763 Monitoring Soil, Vegetation, and Riverine Environments” (S. Manfreda, E. Ben Dor, eds.), pp. 155-178, Elsevier,  
764 Amsterdam, Netherlands, ISBN: 978-0-323-85283-8, 2023.

765 Schilling, K.E., Jha, M.K., Zhang, Y.K., Gassman, P.W., and Wolter, C.F.: Impact of land use and land cover change  
766 on the water balance of a large agricultural watershed: Historical effects and future directions, *Water Resour. Res.*,  
767 44, doi:10.1029/2007WR006644, 2008.

768 Schöne B.R., Meret, A.E., Baier, S.M., (...), and Pfister, L.: Freshwater pearl mussels from northern Sweden serve as  
769 long-term, high-resolution stream water isotope recorders. *Hydrol. Earth Syst. Sci.*, 24, 673–696,  
770 doi:10.5194/hess-24-673-2020, 2020.

771 Seibert, J., Clerc-Schwarzenbach, F. M., and van Meerveld, H. J.: Getting your money's worth: Testing the value  
772 of data for hydrological model calibration. *Hydrol. Process.*, 38, e15094, https://doi.org/10.1002/hyp.15094, 2024.

773 Silasari, R., Parajka, J., Ressler, C., Strauss, P., and Blöschl, G.: Potential of time-lapse photography for identifying  
774 saturation area dynamics on agricultural hillslopes. *Hydrol. Process.*, 31, 3610–3627, doi: 10.1002/hyp.11272,  
775 2017.

776 Sivapalan, M.: Prediction in ungauged basins: a grand challenge for theoretical hydrology. *Hydrol. Process.*, 17(15),  
777 3163–3170, 2003.

778 Sivapalan, M., and Blöschl, G.: The growth of hydrological understanding: Technologies, ideas, and societal needs  
779 shape the field, *Water Resour. Res.*, 53, 8137–8146, 2017.

780 Sokol, Z., Szturc, J., Orellana-Alvear, J., Popová, J., Jurczyk, A., and Célleri, R.: The Role of Weather Radar in  
781 Rainfall Estimation and Its Application in Meteorological and Hydrological Modelling—A Review, *Remote Sens.*,  
782 13, 351. https://doi.org/10.3390/rs13030351, 2021.

783 Späth, F., Rajtschan, V., Weber, T.K.D., Morandage, S., Lange, D., Abbas, S.S., Behrendt, A., Ingwersen, J., Streck,  
784 T., and Wulfmeyer, V.: The land–atmosphere feedback observatory: a new observational approach for  
785 characterizing land–atmosphere feedback, *Geosci. Instrum. Method. Data Syst.*, 12, 25–44,  
786 https://doi.org/10.5194/gi-12-25-2023, 2023.

787 Stähli, M., Badoux, A., Ludwig, A., Steiner, K., Zappa, M., and Hegg, C.: One century of hydrological monitoring in  
788 two small catchments with different forest coverage. *Environ. Monit. and Assess.*, 174(1 – 4), 91 – 106,  
789 doi:10.1007/s10661-010-1757-0, 2011.

790 Strelnikova, D., Perks, M. T., Dal Sasso, S. F., and Pizarro, A.: River flow monitoring with unmanned aerial system.  
791 In *Unmanned aerial systems for monitoring soil, vegetation, and riverine environments* (pp. 231-269), Elsevier,  
792 2023.

793 Su, Z., Zeng, Y., Romano, N., Manfreda, S., Francés, F., Ben Dor, E., Szabó, B., Vico, G., Nasta, P., Zhuang, R., et  
794 al.: An Integrative Information Aqueduct to Close the Gaps between Satellite Observation of Water Cycle and  
795 Local Sustainable Management of Water Resources, *Water*, 12, 1495. https://doi.org/10.3390/w12051495, 2020.

796 Tarasova, L., Gnann, S., Yang, S., Hartmann, A., Wagener, T.: Catchment characterization: Current descriptors,  
797 knowledge gaps and future opportunities, *Earth-Science Reviews*, 252, 104739,  
798 https://doi.org/10.1016/j.earscirev.2024.104739, 2024.

799 Templer, P.H., Harrison, J.L., Pilotto, F. et al.: Atmospheric deposition and precipitation are important predictors of  
800 inorganic nitrogen export to streams from forest and grassland watersheds: a large-scale data synthesis,  
801 *Biogeochemistry*, 160, 219–241, https://doi.org/10.1007/s10533-022-00951-7, 2022.

802 Torbenson, M.C.A., and Stagge, J.H.: Informing seasonal proxy-based flow reconstructions using baseflow  
803 separation: An example from the Potomac River, United States, *Water Resour. Res.*, 57, e2020WR027706,  
804 doi:10.1029/2020WR027706, 2021.

805 Troch, P.A., Carrillo, G., Sivapalan, M., et al.: Climate-vegetation-soil interactions and long-term hydrologic  
806 partitioning: signatures of catchment co-evolution, *Hydrol. Earth Syst. Sci.*, 17, 2209–2217, doi:10.5194/hess-17-  
807 2209-2013, 2013.

808 Valdes-Abellan, J., Jiménez-Martínez, J., Candela, L., Jacques, D., Kohfahl, C., Tamoh, K., Reactive transport  
809 modelling to infer changes in soil hydraulic properties induced by non-conventional water irrigation, *J. Hydrol.*,  
810 549, 114–124, 2017.

811 Van Den Heuvel, D. B., Troch, P. A., Booij, M. J., Niu, G. Y., Volkmann, T. H., and Pangle, L. A.: Effects of  
812 differential hillslope-scale water retention characteristics on rainfall–runoff response at the Landscape Evolution  
813 Observatory, *Hydrol. Process.*, 32(13), 2118-2127, 2018.

814 Vereecken, H., Huisman, J. A., Hendricks Franssen, H. J., Brüggemann, N., Bogena, H. R., Kollet, S., Javaux, M.,  
815 van der Kruk, J., and Vanderborght, J.: Soil hydrology: Recent methodological advances, challenges, and  
816 perspectives, *Water Resour. Res.*, 51, 2616–2633, doi:10.1002/2014WR016852, 2015.

817 Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment classification and hydrologic similarity, *Geography*  
818 *compass*, 1(4), 901-931, 2007.

819 Whitehead, P. G., and Robinson, M.: Experimental basin studies-an international and historical perspective of forest  
820 impacts, *J. Hydrol.*, 145(3–4), 217–230. [https://doi.org/10.1016/0022-1694\(93\)90055-E](https://doi.org/10.1016/0022-1694(93)90055-E), 1993.

821 Wlostowski, A. N., Molotch, N., Anderson, S. P., Brantley, S. L., Chorover, J., Dralle, D., et al.: Signatures of  
822 hydrologic function across the Critical Zone Observatory network, *Water Resour. Res.*, 57, e2019WR026635.  
823 <https://doi.org/10.1029/2019WR026635>, 2021.

824 Zacharias, S., Loescher, H., Bogena, H.R., Kiese, R., Schrön, M., ... and Vereecken, H.: Fifteen years of Integrated  
825 Terrestrial Environmental Observatories (TERENO) in Germany: Functions, services and lessons Learned, *Earth’s*  
826 *Future*, 12(6), e2024EF004510. DOI: 10.22541/essoar.170808463.36288013/v1, 2024.

827 Zeng, Y. and Su, Z.: Digital twin approach for the soil-plant-atmosphere continuum: think big, model small, *Front.*  
828 *Sci.*, 2, 1376950, doi: 10.3389/fsci.2024.1376950, 2024.

829 Zoback, M.L.: Grand challenges in earth and environmental sciences: sciences, stewardship, and service for the  
830 twenty-first century, *GSA Today*, 12, 41-47, 2001.

831

832