



# Variability and long-term changes of tropical cold point temperature and water vapor

Mona Zolghadrshojaee<sup>1</sup>, Susann Tegtmeier<sup>1</sup>, Sean M. Davis<sup>2</sup>, Robin Pilch Kedzierski<sup>3</sup>

<sup>1</sup>Institute of Space and Atmospheric Studies, University of Saskatchewan, Saskatoon, Canada

5 <sup>2</sup>National Oceanic and Atmospheric Administration Chemical Sciences Laboratory, Boulder, CO 80305, USA

<sup>3</sup>Meteorological Institute, Universität Hamburg, Hamburg, Germany

*Correspondence to:* Susann Tegtmeier (susann.tegtmeier@usask.ca)

**Abstract.** The tropical tropopause layer (TTL) is the main gateway for air transiting from the troposphere to the stratosphere and therefore impacts the chemical composition of the stratosphere. In particular, the cold point tropopause, where air parcels  
10 encounter their final dehydration, effectively controls the water vapor content of the lower stratosphere. Given the important role of stratospheric water vapor for the global energy budget, it is crucial to understand the long-term changes in cold point temperature and their impact on water vapor trends.

Our study uses GNSS-RO data to show that, there has been no overall cooling trend of the TTL over the past two decades, in contrast to observations prior to 2000. Instead, the cold point is warming, with the strongest trends of up to 0.7 K/decade during  
15 boreal winter and spring. The cold point warming shows longitudinal asymmetries with the smallest warming over the central Pacific and the largest warming over the Atlantic. These asymmetries are anti-correlated with patterns of tropospheric warming, and regions of strongest cold point warming are found to show slight cooling trends in the upper troposphere. Overall, the here identified warming of the cold point is consistent with model prediction under global climate change, which attributes the warming trends to radiative effects. The seasonal signals and zonal asymmetries of the cold point temperature and height  
20 trends, on the other hand, seemed to be related to dynamical responses to enhanced upper tropospheric heating, changing convection, or trends of the stratospheric circulation.

Water vapor observations in the TTL show mostly positive trends for 2004–2021 consistent with cold point warming. We find a decrease of the amplitude of the cold point temperature seasonal cycle by ~7% driving a reduction of the seasonal cycle in  
25 100 hPa water vapor by 5–6%. Our analysis shows that this reduction in the seasonal cycle is transported upwards together with the seasonal anomalies and has reduced the amplitude of the well-known tape recorder over the last two decades.

## 1 Introduction

The tropical tropopause layer (TTL) is the transition region between the turbulent, moist troposphere and the stable, dry stratosphere (Fueglistaler et al., 2009). The TTL acts as the main gateway for air entering the stratosphere and extends from the region of strong convective outflow near 12–14 km to the highest altitudes reached by convective overshooting events,  
30 around 18 km (e.g., Highwood and Hoskins, 1998; Folkins et al., 1999; Fueglistaler et al., 2009). In consequence, air masses

in the TTL are controlled by numerous processes on a wide range of lengths and timescales and show dynamical and chemical properties of both the troposphere and the stratosphere. Monitoring long-term changes in the TTL is crucial as the interactions among circulation, convection, trace gases, clouds, and radiation make this region a key player in radiative forcing and chemistry–climate coupling (e.g., Mote et al., 1996; Holton and Gettelman, 2001; Fueglistaler et al., 2011; Randel and Jensen, 2013). Atmospheric radiative fluxes in the TTL are particularly sensitive to temperatures, water vapor concentrations as well as the presence of thin cirrus clouds and convective anvils (Wang et al., 1996; Sassen et al., 2009). Theoretical arguments suggest that the TTL radiative balances can in turn impact altitudes of deep convective outflow (Hartmann and Larson, 2002; Harrop and Hartmann, 2012).

An important characteristic of the TTL is the change of the negative vertical temperature gradient of the troposphere into the positive temperature gradient of the stratosphere. The vertical location of the temperature minimum is termed the Cold Point Tropopause (CPT; Highwood and Hoskins, 1998). Air parcels transitioning from the troposphere to the stratosphere encounter final dehydration at these minimum temperatures and in consequence, the cold point tropopause effectively controls the water vapor content of the lower stratosphere (Randel et al., 2004; Fueglistaler et al., 2009; Randel and Park, 2019). Stratospheric water vapor plays an important role in the global energy budget with increasing stratospheric water vapor cooling the stratosphere and heating the troposphere (De F. Forster and Shine, 1999; Smith et al., 2001) and can impact stratospheric ozone loss (Vogel et al., 2011).

The temperature structure of the TTL is determined by the interplay of tropospheric and stratospheric processes. Ongoing strong convection across continents and the western Pacific Ocean contributes significantly to the convective equilibrium that governs the temperatures in the lower TTL. In particular, the connection between El Niño Southern Oscillation (ENSO) and TTL temperatures via convective activities can cause large interannual variations in the latter (Yulaeva and Wallace, 1994). The enhancement of deep convection over the tropical West Pacific is caused by ENSO's negative phase (i.e., La Niña) which is accompanied by TTL cold anomalies right above (Gettelman et al., 2001). Similar TTL cold anomalies occur over the central Pacific during the positive phase (El Niño).

Conversely, the Brewer-Dobson circulation (BDC) with large-scale tropical upwelling, which keeps temperatures well below radiative equilibrium, affects lower stratospheric temperatures. The BDC has a distinct annual cycle due to the seasonality of extratropical planetary wave driving, which imprints a pronounced seasonal cycle in upper TTL temperature. Significant interannual variations in upper TTL temperature are driven by the Quasi-Biennial Oscillation (QBO; Baldwin et al., 2001), a descending pattern of alternating zonal winds. The QBO temperature variations oscillate between the easterly and westerly shear phases following thermal wind balance. Easterly wind shear results in increased upwelling and cold anomalies, while westerly wind shear results in decreased upwelling and warm anomalies (Plumb and Bell, 1982; Baldwin et al., 2001). If the QBO easterly phase occurs during boreal winter, seasonal mean deep convection over the western Pacific will be strengthened (Collimore et al., 2003; Liess and Geller, 2012) further amplifying the QBO cold anomalies in this region (Tegtmeier et al., 2020b). When compared to the equivalent stratospheric changes ( $\pm 4$  K at 25 km), the QBO temperature signal in the TTL is relatively small ( $\pm 0.5$  K at the CPT) and peaks between  $10^{\circ}$ S and  $10^{\circ}$ N (Randel and Wu, 2015).



65 Volcanic eruptions have long been recognized as major natural events capable of influencing global climate including TTL  
temperature. For the three major volcanic eruptions that have occurred since the late 1950s, analysis of observational and  
reanalysis data has revealed warming signals in the tropical lower stratosphere (Free and Lanzante, 2009; Fujiwara et al., 2015)  
and upper TTL (e.g., Krüger et al., 2008; Fujiwara et al., 2015; Tegtmeier et al., 2020a)

Given the important role that the TTL plays in the climate system and the chemical composition of the stratosphere, it is crucial  
70 to understand long-term changes in TTL temperature. In particular, trends of the cold point temperature are of interest since  
they can lead to changes in stratospheric water vapor. Long-term changes of the TTL temperatures are regulated by complex  
coupling of chemical, dynamic, and radiative processes (Fueglistaler et al., 2009; Grise and Thompson, 2013). Changes in  
Greenhouse Gas (GHG) concentrations (e.g., Xie et al., 2008), the strength of the BDC (e.g., Wang et al., 2013; Fu et al.,  
2015), convection (e.g., Nishimoto and Shiotani, 2012), and Sea Surface Temperatures (SSTs; e.g., Hu et al., 2014) all have  
75 an impact on TTL temperatures. Among these factors, the SST and GHG changes as well as convective activities play a key  
role in the long-term cold point temperature changes (e.g., Rosenlof and Reid, 2008; Xie et al., 2014).

Different data sources exist for deriving trends of TTL temperature. An extended satellite temperature record based on the  
Stratospheric Sounding Unit (SSU), the Advanced Microwave Sounding Unit (AMSU-A), the Microwave Limb Sounder  
(MLS), and the Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) offers very good global  
80 coverage, but relatively low vertical resolution (Maycock et al., 2018). The Global Navigation Satellite System – Radio  
Occultation (GNSS-RO) data also have very good global coverage and, in addition, offer a very high vertical resolution. GNSS-  
RO data are available for the period starting in 2002 thus restricting trend estimates to the last 20 years. TTL temperature  
measurements from radiosonde stations extend back in time much longer; however, climate records of radiosonde temperature  
often suffer from time-varying biases or inhomogeneities caused by changes in measurement practices (Seidel and Randel,  
85 2006; Wang et al., 2012). Reanalysis models offer TTL temperature records with different vertical resolutions (Tegtmeier et  
al., 2020a). While meteorological reanalysis data are often utilized as “stand-ins” for observations, trends in reanalysis fields  
can be impacted by spurious changes in the quality and quantity of the observations used as input data (e.g., Long et al., 2017).  
Over the past several decades the global mean stratosphere has cooled due to increasing GHGs and decreases in stratospheric  
ozone (e.g., Maycock et al., 2018). The onset of ozone recovery in the late 1990s has led to a reduction in ozone-induced  
90 cooling with stratospheric cooling trends over 1998–2016 compared to 1979–1997. The troposphere, on the other hand, is  
warming due to the increasing GHGs. Amplified warming of the upper tropical troposphere compared to the surface (Santer  
et al., 2005), the so-called tropical amplification is driven by convective process and water vapor feedback as well as oceanic  
dynamical process and heat storage (e.g., Song et al., 2014; Li et al., 2019). Trends in TTL temperature vary between cooling  
and warming depending on the time period considered, season, and region. Reanalysis data for 1979–2014 suggest zonally  
95 asymmetric cold point temperature trends with positive trends of 0.22 K/decade over the tropical central and eastern Pacific  
and negative trends of -0.08 K/decade throughout the remaining tropical regions (Hu et al., 2016). Zonal mean temperature  
trends from reanalysis over 2002–2020 suggest small but substantial cooling of -0.3 to -0.6 K/decade at 100 hPa and the cold  
point, which are statistically compatible with trends based on the adjusted radiosonde data sets (Tegtmeier et al., 2020a). Based



on MSU and AMSU satellite records, Garfinkel et al. (2013) showed that there has been a strong warming trend in the tropical  
100 upper troposphere near the Indo-Pacific warm pool, while the trends in the western and central Pacific are much weaker. They  
found these patterns to be opposite to the lower stratosphere trends, where the cooling trend is strongest over the Indo-Pacific  
warm pool and weaker over western and central Pacific waters. GNSS-RO data available for the last two decades have revealed  
a continued strong upper tropospheric warming extending into the lower tropical stratosphere (Ladstädter et al., 2023).

Water vapor measurements in the upper troposphere and lower stratosphere are available from the Microwave Limb Sounder  
105 (MLS) at near-global coverage starting in 2004. The Stratospheric Water and OzOne Satellite Homogenized (SWOOSH)  
dataset combines water vapor data from multiple instruments (Davis et al., 2016). Based on these two data sets a significant  
moistening of the stratosphere was found starting in 2002 and with relatively high values persisting over the last decade  
(Konopka et al., 2022). This moistening is prominent in late boreal winter and spring and shows a clear hemispheric  
asymmetry, with the largest positive values in the Northern Hemisphere and negative trends in the polar Southern Hemisphere.  
110 The eruption of the Hunga Tonga–Hunga Ha’apai volcano (Tonga) in January 2022 resulted in a substantial injection of water  
vapor directly into the upper atmosphere (Millán et al., 2022; Vömel et al., 2022) which will impact estimates of stratospheric  
water vapor trends that go beyond the end of 2021.

Here, we update available long-term trends of cold point tropopause temperatures from GNSS-RO data and of stratospheric  
water vapor from MLS and SWOOSH data. We investigate spatial and seasonal patterns of the cold point temperature trends  
115 and their relation to tropospheric and stratospheric changes in Sect. 3. The impact of the seasonality of the cold point  
temperature trends on the seasonality of water vapor trends and specifically on the tape recorder signal are presented in Sect.  
4. Data sets and methodology are introduced in Sect. 2, and a discussion and summary of the results are provided in Sect. 5.

## 2 Data and Methods

### 2.1 GNSS-RO temperature data

120 The Radio Occultation (RO) data derived from the Global Navigation Satellite System (GNSS) exhibit widespread coverage,  
remain unaffected by clouds and precipitation, and boast exceptional vertical resolution and accuracy, as highlighted by  
Kursinski et al. (1997). In this study, we leverage GNSS-RO measurements collected from various satellite missions, including  
CHAMP (Wickert et al., 2001), COSMIC (Anthes et al., 2008), GRACE (Beyerle et al., 2005), Metop-A (Von Engeln et al.,  
2011), successive Metop-B, METOP-C, SAC-C (Hajj et al., 2004), TerraSAR-X (Beyerle et al., 2011), and KOMPSAT-5  
125 (Bowler, 2018).

The dataset employed in this analysis represents an updated version of the data used in Tegtmeier et al. (2020a). Specifically,  
we utilize a dataset encompassing monthly mean CPT temperature, height, and pressure data, along with temperature values  
at various pressure levels (30, 50, 60, 70, 100, 150, 200, 300, and 400 hPa). These data are sourced from all GNSS-RO missions  
and are spatially gridded on a  $30^\circ \times 10^\circ$  grid within the latitudinal range of  $30^\circ$  N to  $30^\circ$  S, covering the period from 2002 until  
130 October 2022.



## 2.2 MLS water vapor data

The Microwave Limb Sounder (MLS) instrument onboard the Aura spacecraft is a part of NASA's Earth Observing System (EOS) platform launched in 2004. MLS uses the microwave limb sounding technique to measure vertical profiles of many atmospheric constituents from the upper troposphere through the mesosphere (Waters et al., 2006). In this study, we use  
135 MLS/Aura Level 3 monthly binned water vapor ( $\text{H}_2\text{O}$ ) mixing ratios, data version 5.1 (Lambert et al., 2020) during 2004-2021. The spatial coverage is near-global ( $82^\circ\text{S}$  to  $82^\circ\text{N}$ ) at a resolution of  $4^\circ$  latitude by  $5^\circ$  longitude. A recently identified drift in the MLS v4  $\text{H}_2\text{O}$  has been corrected in version 5 and as a result, the MLS v5  $\text{H}_2\text{O}$  record shows no statistically significant drifts compared to ACE-FTS and reduced drifts when compared to balloon-borne frost point hygrometer measurements (Livesey et al., 2021). We omit data after the end of 2021 to avoid having the long-term changes impacted by  
140 the anomalous entrainment of water vapor with the Hunga Tonga–Hunga Ha’apai eruption.

## 2.3 SWOOSH water vapor data

The Stratospheric Water and OzOne Satellite Homogenized (SWOOSH) v2.6 data set is a merged record of stratospheric ozone and water vapor measurements taken by a number of limb sounding and solar occultation satellites over the previous ~30 years (Davis et al., 2016). The SWOOSH record spans 1984 to the present and is comprised of data from the SAGE-II/III,  
145 UARS HALOE, UARS MLS, and Aura MLS instruments (v4.2). The measurements are homogenized by applying corrections that are calculated from data taken during time periods of instrument overlap. The SWOOSH data product consists of monthly mean zonal-mean values on a pressure grid at a resolution of  $5^\circ$  latitude by  $20^\circ$  longitude.

## 2.4 Regression Method


In order to isolate tropospheric and stratospheric drivers of TTL temperature variability from long-term changes, a multiple  
150 linear regression analysis was used. The regression was applied to the CPT, 30, 50, 60, 70, 100, 150, 200, 300, and 400 hPa time series of deseasonalized monthly mean temperature anomalies for each  $30^\circ \times 10^\circ$  grid cell and tropical mean values. The analysis isolates the linear long-term changes from temperature variability driven by QBO, ENSO, and SAOD based on the regression described in Eq. (1).

$$T(t) = A + B \cdot t + C_1 \cdot QBO_1(t) + C_2 \cdot QBO_2(t) + D \cdot ENSO(t) + E \cdot SAOD(t) + \epsilon(t), \quad (1)$$

155 where  $T(t)$  are the temperature anomalies and  $\epsilon(t)$  are the residuals. The output of the regression provides the linear trend in the form of the regression coefficient  $B$ . The  $QBO_1(t)$  and  $QBO_2(t)$  terms are orthogonal time series representing QBO variations constructed as the first two empirical orthogonal functions of the Freie Universität Berlin (FUB) radiosonde stratospheric winds (Naujokat, 1986; Wallace et al., 1993).  $ENSO(t)$  is the multivariate ENSO index (<https://www.esrl.noaa.gov/psd/enso/mei/>), and  $SAOD(t)$  is the detrended stratospheric aerosol optical depth from the Global  
160 Space-based Stratospheric Aerosol Climatology (Thomason et al., 2018). Significance is tested based on a two-tailed test with a 95% confidence interval.

### 3 Cold point temperature trends

The long-term change in tropical mean CPT temperature has been calculated for GNSS-RO 2002-2022 data (Fig. 1) by applying the multiple linear regression described in section 2.4. The deseasonalized temperature anomalies (blue line), used as input data to the regression analysis, exhibit strong interannual variations with a range of approximately  $\pm 2\text{K}$ . The R-squared value of 0.3 indicates that approximately 30% of the variability in CPT temperature variations can be explained by the long-term trend, QBO, ENSO, and SAOD. The QBO results in periodic oscillations in CPT temperature, albeit with a relatively modest magnitude (reddish line). ENSO variations in TTL temperature are slightly larger (purple line), showing often, but not always, warm anomalies coinciding with positive ENSO phases (e.g., 2011 and 2021-2022). The fluctuations in SAOD are relatively minor. After taking into account these factors, the regression suggests a positive linear trend (black line) of  $0.36 \pm 0.06 \text{ K/decade}$ . It should be emphasized that substantial residuals (not shown here) indicate a significant portion of variability that cannot be accounted for by the prescribed regressors. This variability may arise from non-linear processes or processes not captured by the proxies, such as fluctuations in tropical upwelling with the BDC due to extratropical wave driving and changes in convective activity not associated with ENSO.




**Figure 1. Tropical mean (20°S-20°N) deseasonalized cold point tropopause (CPT) temperature anomalies (Ano CPT), regression terms (QBO, ENSO, SAOD), and linear trend based on GNSS-RO data.**

In the second step, spatial and seasonal patterns of CPT temperature trends have been analyzed based on the regression analysis applied to seasonal and monthly temperature time series at each grid point. The findings, depicted in Fig. 2, indicate that over the past two decades, there has been no overall cooling trend of the CPT, in contrast to observations prior to 2000 (e.g., Randel et al., 2006; Rosenlof and Reid, 2008). For the full monthly mean time series, clear warming of the CPT with the strongest and significant trends over the tropical regions of South America, the Atlantic, Africa, the Indian Ocean, and Australia of up to  $0.71 \text{ K/decade}$  can be observed (Fig. 2a). Temperature trends minimize to around zero over the central Pacific region.





185 Seasonal mean CPT temperature trends (Fig. 2b-e) are similar to trends based on the full-time series. Temperature changes in the Northern Hemisphere (NH) winter (December, January, and February, DJF), spring (March, April, and May, MAM), summer (June, July, and August, JJA), and fall (September, October, and November, SON) indicate prevalent CPT warming across the tropics. The warming is especially pronounced over the tropical Atlantic and South America in NH winter and spring (up to 1.13 K/decade) as well as Central Africa and the Maritime Continent in NH spring (up to 0.91 K/decade). During  
190 NH summer and fall, the cold point tropopause shows weak and non-significant warming trends except for Australia, which shows significant cold point warming (up to 1.25 K/decade). Similar results are found for NH fall with mostly non-significant cold point trends except for warming over the Atlantic (up to 0.94 K/decade).



195 **Figure 2. Long-term (2002-2022) temperature trends [K/decade] of the cold point tropopause (CPT) temperature based on the GNSS-RO dataset for the monthly mean (a) and seasonal mean time series (b-e). Trends not significant at the 95% confidence level are marked with grey crosses.**


Trends of cold point height and pressure can be expected to be linked to long-term changes in CPT temperature. A comparison of the seasonal trends in CPT temperature (Fig. 2b-e), height (Fig. 3a-d), and pressure (Fig. 3e-h) reveals a noteworthy consistency of the seasonality of the signals. Seasons with the strongest warming (DJF and MAM) show a decrease in CPT height (up to -0.12 km/decade) and an increase in CPT pressure (up to 2.4 hPa/decade). Regional patterns between temperature and height trends are similar for MAM, but less so for DJF. However, these trends in cold point pressure and specifically in  
200



cold point height are significant only over a few smaller regions. During NH summer and fall, most regions experience a rise of the CPT (positive trend in height and negative trend in pressure) while temperature showing weak warming trends.


205 The investigation of the seasonal cycle and the vertical structure of TTL temperature trends highlights discernible patterns (Fig. 4). Overall, the well-known upper tropospheric warming extends into the lower stratosphere up to 80 hPa during the NH spring and summer, 50 hPa during NH winter and all the way up to 30 hPa during October. Temperature trends at the cold point show a clear seasonal cycle with a pronounced warming during boreal winter and spring and weaker, non-significant warming for the rest of the year. For the upper troposphere, the seasonal cycle of the temperature trends is nearly opposite to

210 that of the CP, with largest warming during NH late summer and winter.




215 **Figure 3.** Cold point tropopause (CPT) height [km/decade] (left column) and pressure [hPa/decade] (right column) trends for each season during 2002-2022 based on GNSS-RO data. Trends not significant at the 95% confidence level are marked with gray crosses.





**Figure 4. The seasonal cycle of temperature trends [K/decade] for 2002-2022 at the cold point (CP) and different pressure levels based on GNSS-RO data. Trends not significant at the 95% confidence level are marked with crosses.**

220 The connections between temperature trends at the cold point trends and in the lower stratosphere and upper troposphere (at  
fixed pressure levels at 30, 50, 70, 100, 200, 300, and 400 hPa) are analysed by comparing their spatial patterns (Fig. 5).  
Focusing on boreal winter trends at various pressure levels, the findings highlight a notable pattern: in regions where the cold  
point exhibits the strongest warming (specifically, the tropical Atlantic and South America), temperatures in the upper  
troposphere (400 and 300 hPa) experience a slight cooling trend, although this lower-level cooling trend is not statistically  
225 significant. Conversely, regions characterized by weaker and nonsignificant cold point cooling such as parts of East Asia  
demonstrate substantial warming in the upper troposphere. Such reversed dipole patterns are also observed during the boreal  
fall season when the cold point shows some local cooling over the tropical East Pacific and East Asia (Fig 5, approximately  
90–140°E and 10°N). This cooling is also evident at 70 and 100 hPa, while in the upper troposphere at 300 and 400 hPa the  
same regions show the strongest warming trends. During MAM and JJA, the observed trends do not exhibit significant dipole  
230 patterns or clear trend reversals between the CP and the upper troposphere. Instead, these seasons predominantly display  
warming trends across most tropical regions and all levels between CP and 400 hPa (not shown here).




**Figure 5.** Temperature trends [K/decade] during DJF and SON 2002-2022 at different pressure levels and the cold point (CP) based on GNSS-RO dataset. Trends not significant at the 95% confidence level are marked with grey crosses.



235 The connections between temperature trends at the cold point and in the upper troposphere as well as lower stratosphere are further analyzed by calculating the Pearson correlation coefficients between the gridded trends at the CP and the trends at all other levels, respectively (Fig. 6). Notably, the cold point temperature trend patterns exhibit striking similarities with the temperature trend patterns at levels slightly below (100 hPa) and above (70 hPa) the CPT. These similarities extend upwards into the lower stratosphere (60 to 50 hPa) during NH winter. The connection between temperature trends at the cold point and

240 in the upper troposphere manifests itself in the form of clear anti-correlation between the trend patterns, particularly during DJF and SON seasons as discussed above. Anticorrelations maximize at 300 hPa for DJF and 400 hPa for SON. This suggests that the intensity of cold point warming is connected to patterns of enhanced and reduced upper troposphere warming via some regional dynamical or radiative processes.



245 **Figure 6.** Correlation coefficients between gridded temperature trends at the CPT and at different pressure levels as given on the y-axis based on GNSS-RO data for 2002-2022. Correlation coefficients in grey boxes are not significant at the 95% confidence level.


#### 4 Stratospheric water vapor trends

The temperature of the cold point plays a crucial role in governing the water vapor content of the lower stratosphere. In order to examine the influence of the spatial variations in cold point temperature trends, the patterns of CPT temperature and water

250 vapor trends are compared in Fig. 7 for the 2004-2021 time period. The analyzed data sets, SWOOSH and MLS, agree on the overall trend patterns, with some differences for individual regions and with respect to the trend significance. These differences are largely driven by the fact that SWOOSH uses MLS v4.2, which is compared here to MLS v5. While the stratosphere shows



overall moistening, these positive trends are only significant over South America for DJF and MAM and over the Maritime  
Continent for MAM for SWOOSH. Spatial variations in temperature trends at the CPT (and 70 hPa, not shown here) are found  
255 to coincide with spatial variations of water vapor trends at 100 hPa (and 82 hPa, respectively, not shown here). The strongest  
moistening is observed over the regions that show the largest warming trends (e.g., South America in DJF). Regions with small  
cold point cooling, on the other hand, show weak and not significant drying (e.g., Southeast Asia in DJF).




260 **Figure 7. Temperature trends [K/decade] at the cold point tropopause based on GNSS-RO data (a-b), and water vapor trends [ppm/decade] by MSL data and SWOOSH data at 100 hPa (c-f) for DJF and MAM 2004-2021. Trends not significant at the 95% confidence level are marked with grey crosses.**

Given the strong seasonal cycle in cold point temperature trends, one can expect water vapor trends to have similar seasonal  
variations. Fig. 8 shows the seasonal cycles of cold point temperature and 100 hPa water vapor trends for 10°S-10°N. The cold  
point displays the strongest warming during the boreal winter and spring months with significant trends from February to June  
265 (see also Fig. 4). Most of the boreal summer and fall months, on the other hand, experience weak and non-significant warming.  
This seasonality of the CPT temperature trend extracted from GNSS-RO aligns closely with the seasonality found for the water  
vapor trends sourced from MLS and SWOOSH data. Consistent with peak warming in boreal winter and spring, water vapor  
at 100 hPa shows the strongest positive trends during the same time of the year and small, non-significant trends during boreal  
autumn. It is worth noting that both water vapor datasets indicate a maximum trend slightly later than the maximum cold point  
270 warming. The reasons for this discrepancy are currently unclear and may be related to limitations of the vertical resolution of  
the measurements or involve other processes potentially influencing water vapor trends during boreal spring.

MLS water vapor trends are smaller than SWOOSH trends but agree with their uncertainties. MLS water vapor trends also  
show weaker seasonal fluctuations (amplitude of 0.1 ppm/decade) compared to SWOOSH (0.2 ppm/decade). The seasonal

cycle of the cold point warming shows an amplitude of 0.36 K/decade. The ratios of the water vapor and temperature trend amplitudes of 0.3 ppm/K and 0.6 ppm/K for MLS and SWOOSH, respectively, are consistent with the Clausius-Clapeyron equation as first shown in Konopka et al. (2022).




**Figure 8.** Seasonal cycle of cold point tropopause (CPT) temperature climatology (brown line) and trend (blue line) based on GNSS-RO data (a), and water vapor climatology (brown line) and trends (blue line) at 100 hPa (b-c), and 21 hPa (d-e) based on MLS and SWOOSH data for 2004-2021. Error bars show  $\pm 2$ -sigma confidence intervals of the trend.

The stronger moistening during boreal winter and spring when TTL water vapor shows minimum values (Fig. 8b-c), along with the weaker moistening during NH fall when water vapor shows maximum values, suggest a weakening of the water vapor seasonal cycle. The same principle applies to the seasonal cycle in cold point temperatures where the strongest warming





coincides with the season of lowest temperatures (Fig. 8a) thus leading to a decrease of the seasonal cycle amplitude over time.  
285 As the cold point and water vapor seasonal cycles set the entry values of the tape recorder (Mote et al., 1996), their weakening can lead to a weakening of the tape recorder signal. Indeed, the seasonal cycle of water vapor trends in the middle stratosphere (38 hPa, Fig. 8 d-e) illustrates how the signal propagates upward in time with the strongest moistening now appearing during boreal late fall and winter. As the climatological seasonal cycle propagates upward in the same way, the largest moistening still coincides with the timing of the lowest water vapor values while maximum water vapor experiences weaker moistening.  
290 The upward transport of the deseasonalized water vapor anomalies is shown in Fig. 9a and c for MLS and SWOOSH data over 10°N-10°S. All time-series are boxcar smoothed (width = 5 months) to remove sub-seasonal variability. The mean seasonal cycle of the tape recorder shows the expected minimum during NH winter and spring followed by an increase during summer and fall at 100 hPa and the slow upward propagation of this pattern to higher levels. Water vapor trends have been calculated based on the multilinear regression for each level and month of the year for MLS and SWOOSH (Fig. 9b and d). Noticeably,  
295 the findings illustrate an upward transport of the trends in water vapor at similar time scales as the upward transport of the seasonal cycle in water vapor. The black dotted lines illustrate the timing of the upward transport of water vapor seasonal maxima and minima. The same black lines are overlaid on top of the water vapor trend demonstrating that the minimum (maximum) in water vapor coincides with the stronger (weaker) moistening on most levels.



300 **Figure 9. Deviation in water vapor [ppm] from the time mean averaged profile averaged over 10°N-10°S for MLS (a), and SWOOSH (c) data for 2004-2021. Upward transport of seasonal maxima and minima are shown as black dotted lines. Water vapor trend [ppm/decade] from MLS (b), and SWOOSH (d) data with black dotted lines from panel a and c overlaid on top.**

The percentage change in amplitude of the water vapor seasonal cycle based on the long-term linear trend has been calculated by assuming that the climatological seasonal cycle changes only according to the linear trend for the respective time period  
305 (Table 1). At 100 hPa, the tape recorder signal is estimated to decrease by as 5 to 6% per decade. Such decreasing amplitude



is also found at most other levels consistent with our hypothesis of upward propagation of the weakening. An exception is the levels 56 hPa and 46 hPa which show a slightly increasing amplitude, inconsistent with trends at the other levels. The weakening of amplitude relative to the background seasonal cycle changes with increasing altitude, in particular above 30 hPa, possibly due to the influence of other seasonal signals in the middle stratosphere.

310 **Table 1. Change in amplitude of the tropical water vapor tape recorder [% per decade] averaged over 10°N-10°S for MLS and SWOOSH data for 2004-2021.**

Pressure levels (hPa)	SWOOSH data	MLS data
<b>21</b>	-23.8	-24.4
<b>26</b>	-27.8	-25.6
<b>31</b>	-15.5	-15.9
<b>38</b>	-13.6	-16
<b>46</b>	1.9	-4.2
<b>56</b>	4.8	1.8
<b>68</b>	-2.7	-6.2
<b>82</b>	-0.05	-4.4
<b>100</b>	-5.7	-5.2

## 5 Discussion and summary

High vertical resolution GNSS-RO data demonstrates that over the past two decades, there has been no overall cooling trend in the TTL region consistent with previous studies (e.g., Randel et al., 2017; Steiner et al., 2020; Ladstädter et al., 2023).

330 Focusing on the cold point, which sets the entry values of stratospheric water vapor, our study shows clear warming of this level with strongest trends over the tropical regions of South America, the Atlantic, Africa, the Indian Ocean, and Australia of up to 0.7 K/decade. This cold point warming over 2002-2022 is in contrast to observations prior to 2000, which show a clear cold point cooling (Randel et al., 2006; Wang et al., 2012). In addition, we find an upward shift of the cold point during boreal summer and autumn and a downward shift during boreal winter and spring in good agreement with results derived from the  
335 ozonesonde network SHADOZ (Southern Hemisphere Additional Ozonesondes; Thompson et al., 2021).

Modelling studies have suggested warmer and higher tropopause levels under future global warming due to local radiative effects from increased GHGs and due to a warmer troposphere (Kim et al., 2013; Lin et al., 2017). This temperature signal agrees with the here identified cold point warming, suggesting that radiative effects drive the trends observed over the last 20 years. However, our tropopause height trends with a downward shift during the seasons of maximum warming (DJF and  
340 MAM) are not consistent with the model results. This implies that dynamical processes might also play a role for the observed cold point tropopause trends as discussed below.



Upper tropospheric warming is known to be an amplification of the surface warming signal maximizing in the tropics around 200 hPa and is predicted to decrease to zero near the tropical tropopause (e.g., Santer et al., 2005; Keil et al., 2023). The strong upper tropospheric warming is evident from the GNSS-RO observations (Fig. 4 and 5) and agrees well with estimates from other observational data sets (Vergados et al., 2021; Steiner et al., 2020). Previous studies have suggested that observed temperature trends in the tropical upper troposphere and in the lower stratosphere are anti-correlated (Garfinkel et al., 2013; Hu et al., 2016) in agreement with our results. While 1979–2014 reanalysis datasets suggest strongest warming over the tropical central and eastern Pacific (Hu et al., 2016), our analysis of 2002–2022 GNSS-RO data finds the warming signal to minimize over the central Pacific and to maximize over the Atlantic (Fig. 2). The regions of strongest tropopause warming reveal slight cooling trends in the upper troposphere while regions characterized by weaker and nonsignificant tropopause cooling demonstrate substantial warming in the upper troposphere (Figs. 5 and 6). The identified anticorrelations suggest that dynamical responses to the enhanced moist heating in the upper troposphere or SST driven enhanced convection can be expected to result in negative cold point trends potentially cancelling some of the radiatively driven warming.

Global stratospheric cooling has weakened over the last two decades due to the reduction in ozone-induced cooling after the onset of ozone recovery (Randel et al., 2017; Maycock et al., 2018). However, the tropical lower stratosphere continues to experience negative ozone trends (e.g., Godin-Beekmann et al., 2022), which leads to ongoing cooling and is therefore not consistent with the warming of the tropical lower stratosphere extending up to 30–70 hPa (Fig. 4). Dynamical changes related to upwelling with the stratospheric Brewer-Dobson circulation (BDC), or local impacts of wave activity can also impact tropopause trends. Tropical TTL warming could result from decreased upwelling due to a weakening of the BDC. While this weakening is not expected from model projections under increasing GHGs, some observational data sets suggest positive, but non-significant age-of-air trends in the TTL for 2002–2012 (Haenel et al., 2015) indicating weakened BDC transport in this region. Global lower-stratospheric temperature observations from MSU/AMSU also suggest that the annual mean BDC accelerated from 1980–1999 and decelerated from 2000–2018 (Fu et al., 2019) consistent with changes in TTL temperatures from GNSS-RO data identified here. We find the strongest cold point warming during boreal winter which would be consistent with a more pronounced weakening of the BDC during this time of year. In addition to boreal winter, the cold point and lower stratosphere show a strong warming signal in October, which has also been suggested to be related to the weakening of the BDC during this month as it coincides with a strong negative temperature trend at the southern hemisphere polar regions (Ladstädter et al., 2023).

The cold point temperatures have a clear seasonal cycle with the lowest temperatures during boreal winter and spring due to increased upwelling with BDC. The cold point temperature trends also show a seasonal cycle with the strongest warming during boreal winter and spring. Weaker warming trends found for the rest of the year were not significant due to the strong variability of the temperature time series. We find that the amplitude of the temperature seasonal cycle decreased by 7% as a result of the seasonal alignment of the coldest temperatures with the strongest warming.

Water vapor observations at 100 hPa, a level very close to the cold point, show positive trends for 2004–2021 throughout most of the tropics consistent with increasing cold point temperatures. This is consistent with multi-decadal variation in water vapor



identified for a longer time series extending over the past 40 years (Tao et al., 2023). The positive trend occurred after a sharp drop in water vapor in 2001 (Konopka et al., 2022). The authors showed that the tropical moistening after 2000 occurred mainly during boreal winter and spring, consistent with our results and the seasonal cycle of the cold point warming (Fig. 8). We find that the seasonal cycle in water vapor at 100 hPa is reduced by 5-6% (Table 1) as a result of strongest moistening and maximum warming during the season of lowest water vapor and coldest temperatures. Our analysis shows that this reduction in the seasonal cycle is transported upwards together with the seasonal anomalies and has reduced the amplitude of the well-known tape recorder over the last two decades.

Overall, the here identified warming of the cold point is consistent with model simulations which attribute the warming trends to radiative effects. Seasonal signals and zonal asymmetries in cold point temperature and height trends, on the other hand, seemed to be related to dynamical responses to the enhanced moist heating in the upper troposphere, changing convection or trends of the stratospheric BDC. These seasonal variations have important implications for cross-tropopause transport of water vapor. We find a weakened seasonal cycle in cold point temperature that leads to a weakened seasonal cycle in stratospheric moistening and thus a reduction of the tape recorder signal.

*Data Availability:* The GNSS-RO data are available from <https://cdaac-www.cosmic.ucar.edu/cdaac/products.html>. MLS data can be accessed at [https://disc.gsfc.nasa.gov/datasets/ML3MBH2O\\_005/summary](https://disc.gsfc.nasa.gov/datasets/ML3MBH2O_005/summary), and SWOOSH data can be obtained from <https://csl.noaa.gov/groups/csl8/swoosh/>.

*Author contribution:* MZ performed the analysis and wrote the manuscript. ST developed the research question and guided the research process. RPK provided the GNSS-RO data. SMD provided the SWOOSH data. All authors took part in the process of the manuscript preparation.

*Competing interests:* The contact author has declared that none of the authors has any competing interests.

*Financial support:* This research has been supported by the Natural Sciences and Engineering Research Council of Canada (NSERC, grant no. RGPIN-2020-06292).

## References

Anthes, R. A., Bernhardt, P. A., Chen, Y., Cucurull, L., Dymond, K. F., Ector, D., Healy, S. B., Ho, S.-P., Hunt, D. C., Kuo, Y.-H., Liu, H., Manning, K., McCormick, C., Meehan, T. K., Randel, W. J., Rocken, C., Schreiner, W. S., Sokolovskiy, S. V., Syndergaard, S., Thompson, D. C., Trenberth, K. E., Wee, T.-K., Yen, N. L., and Zeng, Z.: The COSMIC/FORMOSAT-3 Mission: Early Results, Bull. Amer. Meteor. Soc., 89, 313–334, <https://doi.org/10.1175/BAMS-89-3-313>, 2008.



- Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnerson, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The quasi-biennial oscillation, *Reviews of Geophysics*, 39, 179–229, <https://doi.org/10.1029/1999RG000073>, 2001.
- 410 Beyerle, G., Schmidt, T., Michalak, G., Heise, S., Wickert, J., and Reigber, C.: GPS radio occultation with GRACE: Atmospheric profiling utilizing the zero difference technique, *Geophysical Research Letters*, 32, 2005GL023109, <https://doi.org/10.1029/2005GL023109>, 2005.
- Beyerle, G., Grunwaldt, L., Heise, S., Köhler, W., König, R., Michalak, G., Rothacher, M., Schmidt, T., Wickert, J., Tapley, B. D., and Giesinger, B.: First results from the GPS atmosphere sounding experiment TOR aboard the TerraSAR-X satellite, 415 *Atmos. Chem. Phys.*, 11, 6687–6699, <https://doi.org/10.5194/acp-11-6687-2011>, 2011.
- Bowler, N. E.: An initial assessment of the quality of RO data from KOMPSAT-5, *GRAS SAF Rep*, 32, 18, 2018.
- Collimore, C. C., Martin, D. W., Hitchman, M. H., Huesmann, A., and Waliser, D. E.: On the relationship between the QBO and tropical deep convection, *Journal of climate*, 16, 2552–2568, 2003.
- Davis, S. M., Rosenlof, K. H., Hassler, B., Hurst, D. F., Read, W. G., Vömel, H., Selkirk, H., Fujiwara, M., and Damadeo, R.: 420 The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) database: a long-term database for climate studies, *Earth Syst. Sci. Data*, 8, 461–490, <https://doi.org/10.5194/essd-8-461-2016>, 2016.
- De F. Forster, P. M. and Shine, K. P.: Stratospheric water vapour changes as a possible contributor to observed stratospheric cooling, *Geophysical Research Letters*, 26, 3309–3312, <https://doi.org/10.1029/1999GL010487>, 1999.
- Folkens, I., Loewenstein, M., Podolske, J., Oltmans, S. J., and Proffitt, M.: A barrier to vertical mixing at 14 km in the tropics: 425 Evidence from ozonesondes and aircraft measurements, *Journal of Geophysical Research: Atmospheres*, 104, 22095–22102, 1999.
- Free, M. and Lanzante, J.: Effect of Volcanic Eruptions on the Vertical Temperature Profile in Radiosonde Data and Climate Models, *Journal of Climate*, 22, 2925–2939, <https://doi.org/10.1175/2008JCLI2562.1>, 2009.
- Fu, Q., Lin, P., Solomon, S., and Hartmann, D. L.: Observational evidence of strengthening of the Brewer–Dobson circulation 430 since 1980, *JGR Atmospheres*, 120, <https://doi.org/10.1002/2015JD023657>, 2015.
- Fu, Q., Solomon, S., Pahlavan, H. A., and Lin, P.: Observed changes in Brewer–Dobson circulation for 1980–2018, *Environ. Res. Lett.*, 14, 114026, <https://doi.org/10.1088/1748-9326/ab4de7>, 2019.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkens, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Reviews of Geophysics*, 47, 2008RG000267, <https://doi.org/10.1029/2008RG000267>, 2009.
- 435 Fueglistaler, S., Haynes, P. H., and Forster, P. M.: The annual cycle in lower stratospheric temperatures revisited, *Atmos. Chem. Phys.*, 11, 3701–3711, <https://doi.org/10.5194/acp-11-3701-2011>, 2011.
- Fujiwara, M., Hibino, T., Mehta, S. K., Gray, L., Mitchell, D., and Anstey, J.: Global temperature response to the major volcanic eruptions in multiple reanalysis data sets, *Atmos. Chem. Phys.*, 15, 13507–13518, <https://doi.org/10.5194/acp-15-13507-2015>, 2015.
- 440 Garfinkel, C. I., Waugh, D. W., Oman, L. D., Wang, L., and Hurwitz, M. M.: Temperature trends in the tropical upper troposphere and lower stratosphere: Connections with sea surface temperatures and implications for water vapor and ozone, *JGR Atmospheres*, 118, 9658–9672, <https://doi.org/10.1002/jgrd.50772>, 2013.





- Gottelman, A., Randel, W. J., Massie, S., Wu, F., Read, W. G., and Russell, J. M.: El Nino as a natural experiment for studying the tropical tropopause region, *Journal of climate*, 14, 3375–3392, 2001.
- 445 Godin-Beekmann, S., Azouz, N., Sofieva, V. F., Hubert, D., Petropavlovskikh, I., Effertz, P., Ancellet, G., Degenstein, D. A., Zawada, D., Froidevaux, L., Frith, S., Wild, J., Davis, S., Steinbrecht, W., Leblanc, T., Querel, R., Tourpali, K., Damadeo, R., Maillard Barras, E., Stübi, R., Vigouroux, C., Arosio, C., Nedoluha, G., Boyd, I., Van Malderen, R., Mahieu, E., Smale, D., and Sussmann, R.: Updated trends of the stratospheric ozone vertical distribution in the 60° S–60° N latitude range based on the LOTUS regression model, *Atmos. Chem. Phys.*, 22, 11657–11673, <https://doi.org/10.5194/acp-22-11657-2022>, 2022.
- 450 Grise, K. M. and Thompson, D. W. J.: On the Signatures of Equatorial and Extratropical Wave Forcing in Tropical Tropopause Layer Temperatures, *Journal of the Atmospheric Sciences*, 70, 1084–1102, <https://doi.org/10.1175/JAS-D-12-0163.1>, 2013.
- Haenel, F. J., Stiller, G. P., Von Clarmann, T., Funke, B., Eckert, E., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., and Reddmann, T.: Reassessment of MIPAS age of air trends and variability, *Atmos. Chem. Phys.*, 15, 13161–13176, <https://doi.org/10.5194/acp-15-13161-2015>, 2015.
- 455 Hajj, G. A., Ao, C. O., Iijima, B. A., Kuang, D., Kursinski, E. R., Mannucci, A. J., Meehan, T. K., Romans, L. J., De La Torre Juarez, M., and Yunck, T. P.: CHAMP and SAC-C atmospheric occultation results and intercomparisons, *J. Geophys. Res.*, 109, 2003JD003909, <https://doi.org/10.1029/2003JD003909>, 2004.
- Harrop, B. E. and Hartmann, D. L.: Testing the Role of Radiation in Determining Tropical Cloud-Top Temperature, *Journal of Climate*, 25, 5731–5747, <https://doi.org/10.1175/JCLI-D-11-00445.1>, 2012.
- 460 Hartmann, D. L. and Larson, K.: An important constraint on tropical cloud - climate feedback, *Geophysical Research Letters*, 29, <https://doi.org/10.1029/2002GL015835>, 2002.
- Highwood, E. J. and Hoskins, B. J.: The tropical tropopause, *Quart J Royal Meteorol Soc*, 124, 1579–1604, <https://doi.org/10.1002/qj.49712454911>, 1998.
- Holton, J. R. and Gottelman, A.: Horizontal transport and the dehydration of the stratosphere, *Geophysical Research Letters*, 465 28, 2799–2802, <https://doi.org/10.1029/2001GL013148>, 2001.
- Hu, D., Tian, W., Xie, F., Shu, J., and Dhomse, S.: Effects of meridional sea surface temperature changes on stratospheric temperature and circulation, *Adv. Atmos. Sci.*, 31, 888–900, <https://doi.org/10.1007/s00376-013-3152-6>, 2014.
- Hu, D., Tian, W., Guan, Z., Guo, Y., and Dhomse, S.: Longitudinal Asymmetric Trends of Tropical Cold-Point Tropopause Temperature and Their Link to Strengthened Walker Circulation, *Journal of Climate*, 29, 7755–7771, <https://doi.org/10.1175/JCLI-D-15-0851.1>, 2016.
- 470 Keil, P., Schmidt, H., Stevens, B., Byrne, M. P., Segura, H., and Putrasahan, D.: Tropical tropospheric warming pattern explained by shifts in convective heating in the Matsuno–Gill model, *Quart J Royal Meteorol Soc*, 149, 2678–2695, <https://doi.org/10.1002/qj.4526>, 2023.
- Kim, J., Grise, K. M., and Son, S.-W.: Thermal characteristics of the cold-point tropopause region in CMIP5 models, *Journal of Geophysical Research: Atmospheres*, 118, 8827–8841, 2013.
- 475 Konopka, P., Tao, M., Ploeger, F., Hurst, D. F., Santee, M. L., Wright, J. S., and Riese, M.: Stratospheric Moistening After 2000, *Geophysical Research Letters*, 49, e2021GL097609, <https://doi.org/10.1029/2021GL097609>, 2022.



- Krüger, K., Tegtmeier, S., and Rex, M.: Long-term climatology of air mass transport through the Tropical Tropopause Layer (TTL) during NH winter, *Atmos. Chem. Phys.*, 8, 813–823, <https://doi.org/10.5194/acp-8-813-2008>, 2008.
- 480 Kursinski, E. R., Hajj, G. A., Schofield, J. T., Linfield, R. P., and Hardy, K. R.: Observing Earth's atmosphere with radio occultation measurements using the Global Positioning System, *J. Geophys. Res.*, 102, 23429–23465, <https://doi.org/10.1029/97JD01569>, 1997.
- Ladstädter, F., Steiner, A. K., and Gleisner, H.: Resolving the 21st century temperature trends of the upper troposphere–lower stratosphere with satellite observations, *Sci Rep*, 13, 1306, <https://doi.org/10.1038/s41598-023-28222-x>, 2023.
- 485 Lambert, A., Livesey, N., Read, W., and Fuller, R.: MLS/Aura Level 3 Monthly Binned Water Vapor (H<sub>2</sub>O) Mixing Ratio on Assorted Grids V004, <https://doi.org/10.5067/AURA/MLS/DATA/3209>, 2020.
- Li, Y., Yang, S., Deng, Y., Hu, X., Cai, M., and Zhou, W.: Detection and attribution of upper-tropospheric warming over the tropical western Pacific, *Clim Dyn*, 53, 3057–3068, <https://doi.org/10.1007/s00382-019-04681-9>, 2019.
- Liess, S. and Geller, M. A.: On the relationship between QBO and distribution of tropical deep convection, *J. Geophys. Res.*, 490 117, 2011JD016317, <https://doi.org/10.1029/2011JD016317>, 2012.
- Lin, P., Paynter, D., Ming, Y., and Ramaswamy, V.: Changes of the Tropical Tropopause Layer under Global Warming, *Journal of Climate*, 30, 1245–1258, <https://doi.org/10.1175/JCLI-D-16-0457.1>, 2017.
- Livesey, N. J., Read, W. G., Froidevaux, L., Lambert, A., Santee, M. L., Schwartz, M. J., Millán, L. F., Jarnot, R. F., Wagner, P. A., Hurst, D. F., Walker, K. A., Sheese, P. E., and Nedoluha, G. E.: Investigation and amelioration of long-term instrumental drifts in water vapor and nitrous oxide measurements from the Aura Microwave Limb Sounder (MLS) and their implications for studies of variability and trends, *Atmos. Chem. Phys.*, 21, 15409–15430, <https://doi.org/10.5194/acp-21-15409-2021>, 2021.
- 495 Long, C. S., Fujiwara, M., Davis, S., Mitchell, D. M., and Wright, C. J.: Climatology and interannual variability of dynamic variables in multiple reanalyses evaluated by the SPARC Reanalysis Intercomparison Project (S-RIP), *Atmos. Chem. Phys.*, 17, 14593–14629, <https://doi.org/10.5194/acp-17-14593-2017>, 2017.
- 500 Maycock, A. C., Randel, W. J., Steiner, A. K., Karpechko, A. Y., Christy, J., Saunders, R., Thompson, D. W. J., Zou, C., Chrysanthou, A., Luke Abraham, N., Akiyoshi, H., Archibald, A. T., Butchart, N., Chipperfield, M., Dameris, M., Deushi, M., Dhomse, S., Di Genova, G., Jöckel, P., Kinnison, D. E., Kirner, O., Ladstädter, F., Michou, M., Morgenstern, O., O'Connor, F., Oman, L., Pitari, G., Plummer, D. A., Revell, L. E., Rozanov, E., Stenke, A., Visionsi, D., Yamashita, Y., and Zeng, G.: Revisiting the Mystery of Recent Stratospheric Temperature Trends, *Geophysical Research Letters*, 45, 9919–9933, 505 <https://doi.org/10.1029/2018GL078035>, 2018.
- Millán, L., Santee, M. L., Lambert, A., Livesey, N. J., Werner, F., Schwartz, M. J., Pumphrey, H. C., Manney, G. L., Wang, Y., Su, H., Wu, L., Read, W. G., and Froidevaux, L.: The Hunga Tonga-Hunga Ha'apai Hydration of the Stratosphere, *Geophysical Research Letters*, 49, e2022GL099381, <https://doi.org/10.1029/2022GL099381>, 2022.
- Mote, P. W., Rosenlof, K. H., McIntyre, M. E., Carr, E. S., Gille, J. C., Holton, J. R., Kinnarsley, J. S., Pumphrey, H. C., 510 Russell, J. M., and Waters, J. W.: An atmospheric tape recorder: The imprint of tropical tropopause temperatures on stratospheric water vapor, *J. Geophys. Res.*, 101, 3989–4006, <https://doi.org/10.1029/95JD03422>, 1996.
- Naujokat, B.: An update of the observed quasi-biennial oscillation of the stratospheric winds over the tropics, *Journal of Atmospheric Sciences*, 43, 1873–1877, 1986.



- 515 Nishimoto, E. and Shiotani, M.: Seasonal and interannual variability in the temperature structure around the tropical tropopause and its relationship with convective activities, *J. Geophys. Res.*, 117, 2011JD016936, <https://doi.org/10.1029/2011JD016936>, 2012.
- Plumb, R. A. and Bell, R. C.: Equatorial waves in steady zonal shear flow, *Quart J Royal Meteor Soc*, 108, 313–334, <https://doi.org/10.1002/qj.49710845603>, 1982.
- 520 Randel, W. and Park, M.: Diagnosing Observed Stratospheric Water Vapor Relationships to the Cold Point Tropical Tropopause, *JGR Atmospheres*, 124, 7018–7033, <https://doi.org/10.1029/2019JD030648>, 2019.
- Randel, W. J. and Jensen, E. J.: Physical processes in the tropical tropopause layer and their roles in a changing climate, *Nature Geosci*, 6, 169–176, <https://doi.org/10.1038/ngeo1733>, 2013.
- Randel, W. J. and Wu, F.: Variability of Zonal Mean Tropical Temperatures Derived from a Decade of GPS Radio Occultation Data, *Journal of the Atmospheric Sciences*, 72, 1261–1275, <https://doi.org/10.1175/JAS-D-14-0216.1>, 2015.
- 525 Randel, W. J., Wu, F., Oltmans, S. J., Rosenlof, K., and Nedoluha, G. E.: Interannual changes of stratospheric water vapor and correlations with tropical tropopause temperatures, *Journal of the Atmospheric Sciences*, 61, 2133–2148, 2004.
- Randel, W. J., Wu, F., Vömel, H., Nedoluha, G. E., and Forster, P.: Decreases in stratospheric water vapor after 2001: Links to changes in the tropical tropopause and the Brewer-Dobson circulation, *J. Geophys. Res.*, 111, 2005JD006744, <https://doi.org/10.1029/2005JD006744>, 2006.
- 530 Randel, W. J., Polvani, L., Wu, F., Kinnison, D. E., Zou, C., and Mears, C.: Troposphere-Stratosphere Temperature Trends Derived From Satellite Data Compared With Ensemble Simulations From WACCM, *JGR Atmospheres*, 122, 9651–9667, <https://doi.org/10.1002/2017JD027158>, 2017.
- Rosenlof, K. H. and Reid, G. C.: Trends in the temperature and water vapor content of the tropical lower stratosphere: Sea surface connection, *J. Geophys. Res.*, 113, 2007JD009109, <https://doi.org/10.1029/2007JD009109>, 2008.
- 535 Santer, B. D., Wigley, T. M. L., Mears, C., Wentz, F. J., Klein, S. A., Seidel, D. J., Taylor, K. E., Thorne, P. W., Wehner, M. F., Gleckler, P. J., Boyle, J. S., Collins, W. D., Dixon, K. W., Doutriaux, C., Free, M., Fu, Q., Hansen, J. E., Jones, G. S., Ruedy, R., Karl, T. R., Lanzante, J. R., Meehl, G. A., Ramaswamy, V., Russell, G., and Schmidt, G. A.: Amplification of Surface Temperature Trends and Variability in the Tropical Atmosphere, *Science*, 309, 1551–1556, <https://doi.org/10.1126/science.1114867>, 2005.
- 540 Sassen, K., Wang, Z., and Liu, D.: Cirrus clouds and deep convection in the tropics: Insights from CALIPSO and CloudSat, *J. Geophys. Res.*, 114, 2009JD011916, <https://doi.org/10.1029/2009JD011916>, 2009.
- Seidel, D. J. and Randel, W. J.: Variability and trends in the global tropopause estimated from radiosonde data, *J. Geophys. Res.*, 111, 2006JD007363, <https://doi.org/10.1029/2006JD007363>, 2006.
- 545 Smith, C. A., Haigh, J. D., and Toumi, R.: Radiative forcing due to trends in stratospheric water vapour, *Geophysical Research Letters*, 28, 179–182, <https://doi.org/10.1029/2000GL011846>, 2001.
- Song, X., Zhang, G. J., and Cai, M.: Quantifying contributions of climate feedbacks to tropospheric warming in the NCAR CCSM3.0, *Clim Dyn*, 42, 901–917, <https://doi.org/10.1007/s00382-013-1805-x>, 2014.
- Steiner, A. K., Ladstädter, F., Randel, W. J., Maycock, A. C., Fu, Q., Claud, C., Gleisner, H., Haimberger, L., Ho, S.-P., Keckhut, P., Leblanc, T., Mears, C., Polvani, L. M., Santer, B. D., Schmidt, T., Sofieva, V., Wing, R., and Zou, C.-Z.: Observed



- 550 Temperature Changes in the Troposphere and Stratosphere from 1979 to 2018, *Journal of Climate*, 33, 8165–8194, <https://doi.org/10.1175/JCLI-D-19-0998.1>, 2020.
- Tao, M., Konopka, P., Wright, J. S., Liu, Y., Bian, J., Davis, S. M., Jia, Y., and Ploeger, F.: Multi-decadal variability controls short-term stratospheric water vapor trends, *Commun Earth Environ*, 4, 441, <https://doi.org/10.1038/s43247-023-01094-9>, 2023.
- 555 Tegtmeier, S., Anstey, J., Davis, S., Dragani, R., Harada, Y., Ivanciu, I., Pilch Kedzierski, R., Krüger, K., Legras, B., Long, C., Wang, J. S., Wargan, K., and Wright, J. S.: Temperature and tropopause characteristics from reanalyses data in the tropical tropopause layer, *Atmos. Chem. Phys.*, 20, 753–770, <https://doi.org/10.5194/acp-20-753-2020>, 2020a.
- Tegtmeier, S., Anstey, J., Davis, S., Ivanciu, I., Jia, Y., McPhee, D., and Pilch Kedzierski, R.: Zonal Asymmetry of the QBO Temperature Signal in the Tropical Tropopause Region, *Geophysical Research Letters*, 47, e2020GL089533, 560 <https://doi.org/10.1029/2020GL089533>, 2020b.
- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, *Earth Syst. Sci. Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.
- Thompson, A. M., Stauffer, R. M., Wargan, K., Witte, J. C., Kollonige, D. E., and Ziemke, J. R.: Regional and Seasonal Trends in Tropical Ozone From SHADOZ Profiles: Reference for Models and Satellite Products, *JGR Atmospheres*, 126, e2021JD034691, <https://doi.org/10.1029/2021JD034691>, 2021.
- Vergados, P., Ao, C. O., Mannucci, A. J., and Kursinski, E. R.: Quantifying the Tropical Upper Tropospheric Warming Amplification Using Radio Occultation Measurements, *Earth and Space Science*, 8, e2020EA001597, <https://doi.org/10.1029/2020EA001597>, 2021.
- 570 Vogel, B., Feck, T., and Grooß, J.-U.: Impact of stratospheric water vapor enhancements caused by CH<sub>4</sub> and H<sub>2</sub>O increase on polar ozone loss, *J. Geophys. Res.*, 116, D05301, <https://doi.org/10.1029/2010JD014234>, 2011.
- Vömel, H., Evan, S., and Tully, M.: Water vapor injection into the stratosphere by Hunga Tonga-Hunga Ha’apai, *Science*, 377, 1444–1447, <https://doi.org/10.1126/science.abq2299>, 2022.
- Von Engel, A., Andres, Y., Marquardt, C., and Sancho, F.: GRAS radio occultation on-board of Metop, *Advances in Space Research*, 47, 336–347, <https://doi.org/10.1016/j.asr.2010.07.028>, 2011.
- Wallace, J. M., Panetta, R. L., and Estberg, J.: Representation of the equatorial stratospheric quasi-biennial oscillation in EOF phase space, *Journal of the atmospheric sciences*, 50, 1751–1762, 1993.
- Wang, J. S., Seidel, D. J., and Free, M.: How well do we know recent climate trends at the tropical tropopause?, *J. Geophys. Res.*, 117, 2012JD017444, <https://doi.org/10.1029/2012JD017444>, 2012.
- 580 Wang, P., Minnis, P., McCormick, M. P., Kent, G. S., and Skeens, K. M.: A 6-year climatology of cloud occurrence frequency from Stratospheric Aerosol and Gas Experiment II observations (1985–1990), *J. Geophys. Res.*, 101, 29407–29429, <https://doi.org/10.1029/96JD01780>, 1996.
- Wang, W., Matthes, K., Schmidt, T., and Neef, L.: Recent variability of the tropical tropopause inversion layer, *Geophysical Research Letters*, 40, 6308–6313, <https://doi.org/10.1002/2013GL058350>, 2013.



- 585 Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. E., Filipiak, M. J., Flower, D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo, M. S., Perun, V. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, K. M., Chavez, M. C., Gun-Shing Chen, Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M. A., Jiang, J. H., Yibo Jiang, Knosp, B. W., LaBelle, R. C., Lam, J. C., Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, 590 D. M., Van Snyder, W., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth observing system microwave limb sounder (EOS MLS) on the aura Satellite, *IEEE Trans. Geosci. Remote Sensing*, 44, 1075–1092, <https://doi.org/10.1109/TGRS.2006.873771>, 2006.
- Wickert, J., Reigber, C., Beyerle, G., König, R., Marquardt, C., Schmidt, T., Grunwaldt, L., Galas, R., Meehan, T. K., Melbourne, W. G., and Hocke, K.: Atmosphere sounding by GPS radio occultation: First results from CHAMP, *Geophysical Research Letters*, 28, 3263–3266, <https://doi.org/10.1029/2001GL013117>, 2001. 595
- Xie, F., Tian, W., and Chipperfield, M. P.: Radiative effect of ozone change on stratosphere-troposphere exchange, *J. Geophys. Res.*, 113, 2008JD009829, <https://doi.org/10.1029/2008JD009829>, 2008.
- Xie, F., Li, J., Tian, W., Li, Y., and Feng, J.: Indo-Pacific Warm Pool Area Expansion, Modoki Activity and Tropical Cold-Point Tropopause Temperature Variations, *Sci Rep*, 4, 4552, <https://doi.org/10.1038/srep04552>, 2014.
- 600 Yulaeva, E. and Wallace, J. M.: The signature of ENSO in global temperature and precipitation fields derived from the microwave sounding unit, *Journal of climate*, 7, 1719–1736, 1994.