

Measurement Report: The effects of SECA regulations on the atmospheric SO₂ concentrations in the Baltic Sea, based on long-term observations at the Finnish Utö Island.

Androniki Maragkidou¹, Tiia Grönholm¹, Laura Rautiainen¹, Juha Nikmo¹, Jukka-Pekka Jalkanen¹, Timo Mäkelä¹, Timo Anttila¹, Lauri Laakso^{1,2} and Jaakko Kukkonen^{1,3}

¹Finnish Meteorological Institute, Helsinki, FI-00560, Finland

²Atmospheric Chemistry Research Group, Chemical Resource Beneficiation, North-West University, Potchefstroom 2531, South Africa

³Centre for Climate Change Research (C3R), University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK

10 *Correspondence to:* Androniki Maragkidou (androniki.maragkidou@fmi.fi)

Abstract. The designation of the Baltic Sea as a Sulfur Emission Control Area (SECA) in May 2006, with subsequent tightening of regulations in 2010 and 2015 has reduced the sulfuric emission from shipping traffic. This study assesses the impacts of SECA on observed SO₂ concentrations by providing a long-term analysis of 1-minute time resolution air quality data from 2006 to 2020 at Utö island (Baltic Sea) supported by the predictions from the Ship Traffic Emission Assessment Model (STEAM). Additionally, 1-hour resolution data from 2003 to 2005 is utilized to investigate changes due to the SECA limits set in 2006. The observed SO₂ concentrations at Utö have continuously decreased since 2003 due to an overall decrease in SO₂ emissions in Northern Europe, combined with reduced emissions from shipping traffic due to SECA regulations. Three-year average SO₂ concentrations dropped from pre-SECA (2003–2005) to post-SECA periods (2007–2009, 2011–2013, 2016–2018) by 38 %, 39 %, and 67 %, respectively. No clear trends were observed in the concentrations of other measured pollutants. Furthermore, we investigated wind direction resolved SO₂ concentrations for two selected years (2014 and 2019) and the results showed a significant decrease in high SO₂-concentration shipping plumes due to the implementation of SECA in 2015. Our study brings out the importance of long-term, high time-resolution air quality observations at remote marine research station in providing means for both quantitative and qualitative analyses of the impacts of regulatory environmental legislation.

25 1 Introduction

In recent years, international maritime trade has experienced sustained growth, primarily due to its recognized economic efficiency, accounting for over 90 % of global trade (IMO, 2016). This growth has been projected to continue, with the United Nations Conference on Trade and Development (UNCTAD, 2019) indicating a prospective annual growth rate of 2.6 % to 3.4 % until 2024. In 2019, 11.08 billion tonnes of goods were shipped, and the global commercial fleet, consisting of 98 140 vessels exceeding 100 gross tonnes, had a total capacity of 2.06 billion deadweight tonnes (UNCTAD, 2020). In the Baltic

Sea, the shipping activity has increased fairly steadily during 2006–2020 (HELCOM, 2021). For example, in 2020 the total number of vessel fleet was increased by 271 %, 140 % and 46 % in comparison to 2006, 2010 and 2015, respectively.

35 However, maritime activities have led to significant environmental challenges. In particular, shipping emissions are a major source of air pollution, contributing annually to approximately 20.88×10^6 tonnes NO_x , 9.7×10^6 tonnes SO_2 and 1.5×10^6 tonnes $\text{PM}_{2.5}$ globally (Johansson et al., 2017). Smith et al. (2015) evaluated that global shipping is responsible for approximately 13 % and 12 % of total global anthropogenic emissions of NO_x and SO_x , respectively. On European scale, the European Environment Agency (EEA, 2017) evaluated that 90 % of SO_2 emissions from transport were attributed to maritime transport, while $\text{PM}_{2.5}$, PM_{10} and NO_x transport-associated emissions accounted for 45 %, 28 %, and 35 %, respectively.

40 In the Baltic Sea region, emissions of SO_2 , NO_x , and NO_2 from international shipping were estimated to account for as much as 80 % of the overall concentrations of these pollutants in the near-surface environment (Haglund et al., 2016), deteriorating the air quality in coastal areas (Jonson et al., 2015). Apart from their air quality impacts, these emissions had also the potential to cause acidification and eutrophication of marine waters and surrounding terrestrial ecosystems, with serious implications in the Baltic Sea environment (HELCOM, 2009; Hunter et al., 2011; Raudsepp et al., 2013).

45 Notably, the impacts of shipping emissions are not limited to sea areas. Around 70 % of ship emissions occur within 400 km of land (Corbett et al., 1999), while more than one-third of the world's population lives within 100 km from the coast (UNEP, 2024). Pollution from shipping can reach the inhabited land areas and have a severe impact on human health. Shipping emissions have been shown to negatively impact the air quality, especially on coastal regions and portside (Chen et al., 2019; Donato et al., 2014; Liu et al., 2017), the climate (Contini et al., 2015; Merico et al., 2016; Sofiev et al., 2018) and the economy (Jalkanen et al., 2014). Andersson et al. (2009) and Brandt et al. (2013) evaluated that international shipping was responsible for approximately 50 000 premature deaths annually in Europe. Furthermore, Barregard et al. (2019) reported that shipping in the Baltic Sea and the North Sea alone may have resulted in approximately 14 000 premature deaths in Europe in 55 2011.

60 While sulfur reductions have improved air quality and subsequently human health (Jonson et al. 2015; Sofiev et al., 2018; Barregard et al., 2019), it should be kept in mind that reducing sulfuric emissions from shipping may lead to unintended adverse impacts on climate and marine ecosystems. The impact on climate, due to decrease in the cooling effect of ship-emitted aerosols, is likely leading to an increase in global sea surface and atmospheric temperatures (Lauer et al., 2009; Partanen et al., 2013; Hausfather and Forster, 2023). The impacts of sulfur reduction on marine ecosystems includes concentrated aquatic phase pollutant emissions from the exhaust gas cleaning systems as discussed in Picone et al. (2023) and Hermansson et al. (2024). Thus, the potential success of sulfur reduction regulations discussed in this study may require further investigations in

a wider context not focusing only on direct human health benefits.

65

As international maritime trade and shipping emissions are increasing, measures to limit and monitor their adverse impacts are necessary. To mitigate shipping emissions, the International Maritime Organization (IMO) has implemented a series of regulations aimed at reducing emissions from ships. These regulations include the establishment of Sulfur Emission Control Areas (SECAs), which require ships to use low sulfur fuel in designated areas, or, alternatively, reduce the sulfuric emission to air with exhaust gas cleaning system. In May 2006, the Baltic Sea was designated as a SECA region along with the North Sea and the English Channel. SECA regions have also been set up along the North American East and West Coast and the U.S Caribbean. The maximum allowed sulfur content of fuel in the SECA areas was decreased in 2006 from 2.7 to 1.5 %. Since 1 of July 2010, the International Maritime Organization (IMO) implemented a rule that required ships sailing in SECA regions to use fuel with a sulfur content no higher than 1.0 % (Van Aardenne et al., 2013). This rule was further tightened in 2015, reducing the sulfur limit to 0.1 %.

75

Given these regulations, it is essential to assess their effectiveness through compliance studies across various regions. For instance, Beecken et al. (2015) conducted two campaigns in the Gulf of Finland and Neva Bay during the summers of 2011 and 2012, sampling 466 plumes from 311 individual vessels using both ground-based and helicopter-borne measurements. Their analysis revealed that 90 % of the observed plumes complied with the 1 % sulfur content limit set for SECAs in 2011, with this compliance increasing to 97 % in 2012. Similarly, Yang et al. (2016) observed a significant decrease in ship-emitted SO₂ following the implementation of SECA regulations in January 2015 in the English Channel, underscoring the substantial impact of these policies on air quality. Mellqvist et al. (2017) reported compliance levels of 92–94% around Denmark during 2015–2016, and Jonson et al. (2019) revealed strong compliance with SECA regulations in the Baltic Sea based on emission modelling. Additionally, Repka et al. (2019) conducted a comprehensive study and found compliance rates exceeding 98 % in Göteborg and Gdynia, 95–97 % in St. Petersburg, and 94% in the central Baltic Sea. However, compliance rates were slightly lower at SECA borders, around 87 %.

85

In this study, the impacts of SECA regulations on SO₂ concentrations in a small island in the Baltic Sea are studied. More specifically, this research will present unique fine temporal resolution data of SO₂, NO_x, NO, PM_{2.5} and O₃ concentrations measured at Utö island at the outer edge of the Archipelago Sea in the Baltic Sea, providing further insights into the impacts and the effectiveness of SECA regulations. The observations were carried out during the period 2006-2020, covering the period with three separate changes in SECA limits. Additionally, sparser hourly data from 2003 to 2005 has also been utilized.

90

95

Moreover, the observations of air quality parameters are supported with local meteorological data and emission estimates utilizing AIS data. To further contextualize these measurements, we also computed the emissions attributed to shipping in the

Baltic Sea during the target period using the STEAM model.

It is also noteworthy that the dataset used in this study, which includes wind direction analysis of SO₂ concentrations relative to a major shipping lane, has not been previously published or analyzed.

2 Measurement location and site characteristics

The observations utilized in this study have been carried out at Utö (59° 46' 50 N, 21° 22' 23 E), a small island in the Baltic Sea (Fig. 1a) with an area less than 1 km². It is located 70 km from the coast of the mainland of Finland and surrounded only by the open sea and a few smaller islands. The vegetation between the bare rocky areas consists mainly of different shrubs and a few trees. The island is flat, the highest point is less than 15 m above the sea surface. Less than 40 inhabitants live at Utö during the winter. Population, including tourists during the summer months, varies between 100 and 200. Ships that pass Utö are pilot, passenger, military, fishing, cargo, commercial and tanker vessels. An important shipping lane to or from Finland is located at a distance of approximately 1 km west of the island (Fig. 1b).

Meteorological observations at Utö started in 1881. Since the beginning of observations, there has been a gradual increase in measured atmospheric, air quality, marine and electromagnetic parameters (Ahl \ddot{u} s, 1961; Laurila and Hakola, 1996; Komppula et al., 2007; Kyll \ddot{o} nen et al., 2009; Kilkki et al., 2015; Laapas and Ven \ddot{a} l \ddot{a} inen, 2017; Laakso et al. 2018; Honkanen et al., 2018, 2021; Gr \ddot{o} nholm et al., 2021; Kraft et al., 2021; Sepp \ddot{a} l \ddot{a} et al., 2021; Rautiainen et al., 2023; Honkanen et al., 2024; Hell \ddot{e} n et al., 2024). As the island is sparsely populated and there is very little land traffic, there are no local sources of NO_x and SO₂. Consequently, Ut \ddot{o} is an ideal location for plume tracking.

The long-term marine and meteorological observations and other characteristics of Ut \ddot{o} Atmospheric and Marine Research Station are discussed in Laakso et al. (2018) and the references listed above, and thus only few results relevant for this study are summarized here. The prevailing meteorological conditions in the area are characterized by high wind speeds, with monthly averages ranging from 5.6 m s⁻¹ in July to 8.9 m s⁻¹ in December, resulting in an annual average of 7.1 m s⁻¹. The prevailing wind is from the southwest, while the sector from south (clockwise) to northwest dominates, accounting for approximately 60 % of the observed wind patterns (Honkanen et al., 2018). An examination of the wind time series from 1959 to 2016 (Laapas and Ven \ddot{a} l \ddot{a} inen, 2017) indicated no substantial alterations in wind direction or wind speed, aligning with a recent investigation covering the period of 1979 to 2008.

125




Figure 1: a) The location of the Utö island in the Baltic Sea and b) Utö air quality station (orange diamond) and the shipping lanes adjacent to the island (black dashed lines). Major ships operate on the north-south shipping lane at a distance of approximately 500 m west of Utö (or 1 km from the quality station). The wind sector representing the directions from the main shipping lane (185°–315°), is indicated by the orange lines. The ship traffic to and from the harbour of the island is presented by the shaded orange sector.

3 Measurements and methods

3.1 Air quality and wind observations

To study the effects of SECA regulations on air quality, concentrations of SO₂, PM_{2.5}, NO, NO_x, and O₃ were measured at 1-minute time resolution and analysed. The period of the study spanned 15 years, from 2006 to 2020 (supporting hourly data from 2003 to 2005). During this time, some of the instruments were changed (Table 1). The data has gone through normal quality control procedures as defined by EMEP and World Meteorological Organization (WMO) standards (see e.g. Anttila and Tuovinen, 2010 and references therein). In addition to the air quality observations, wind data on site was used. The temporal resolution of the wind data was 10 minutes and therefore it was interpolated linearly for intervals of 1 minute to match the air quality data.

Table 1. Air quality observations.

Observation	Device	Start	End	Notes	Uncertainty *
-------------	--------	-------	-----	-------	---------------

Formatted Table

SO₂	Thermo 43i-TLE	09/09/2011		<u>1 % of reading or 1 ppb (whichever is greater)</u>
	Thermo 43s	01/01/1996	08/09/2011	<u>1 % of reading or 1 ppb (whichever is greater)</u>
PM_{2.5}	Fidas 200E	11/06/2021		<u>9.7 %</u>
	5030 Sharp	22/11/2017		<u>± 2.0 µg m⁻³ <80 µg/m⁻³; ± 5 µg m⁻³ >80 µg m⁻³ (24-hour)</u>
	Thermo FH62 I-R	01/06/2003	22/11/2017	<u>~ 17 % **</u>
NO_x	Thermo 42i-TL	07/02/2014		<u>± 0.4 ppb in the range of 0–500 ppb</u>
	Horiba APNA-370	18/08/2007	07/02/2014	<u>0.5 ppb</u>
	TEI 42CTL		18/08/2007	Data gap 2006-2007 <u>± 0.4 ppb in the range of 0–500 ppb</u>
O₃	Thermo 49i	12/02/2015		<u>1 ppb</u>
	Horiba Ltd APOA-360	03/10/2003	11/02/2015	<u>± 1.0 %</u>

* Based on instruments' or equivalent models' datasheets, manuals, or manufacturer-provided specifications available online

** Based on Lagler et al. (2011)

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Font: 9 pt

145

Calibration of the instruments is carried out with precision to ensure accuracy and reliability. Initially, instruments are calibrated at the National Reference Laboratory for Air Quality at the Finnish Meteorological Institute (FMI). Following this, instruments are calibrated in the field using reference values from the laboratory, involving zero level and three span points to

150

confirm linearity.

SO₂ instrument is calibrated with the Gasetm Ansyco system, which utilizes a permeation chamber to produce SO₂ test gas and can also generate zero air. For PM_{2.5/10}, the Fidas instrument is calibrated according to the manufacturer's procedure using a Monodust (reference particles) provided with the instrument to define the number of particles entering the device. Zero calibration is achieved with a HEPA filter, eliminating the need for reference values from the National Reference Laboratory.

NO_x is calibrated using the Sonimix 3012/3022 multigas calibrator, which includes a [gas phase titration \(GPT\)](#), ozone, and a self-regenerating zero air generator. Calibration for NO specifically is conducted with externally supplied diluted NO in N₂. O₃ is calibrated with the 49i-PS, a UV photometric primary standard designed for the calibration of ozone analyzers. These procedures ensure that measurements are both accurate and reliable.

3.2 Automatic Identification System (AIS) data

The Automatic Identification System (AIS) is a mechanism for the automated generation and transmission of vessel-related information to both other vessels and coastal authorities (IMO, 2020). The AIS data encompassed main ship attributes, including Maritime Mobile Service Identity (MMSI), latitude, longitude, true heading, course over ground, and speed over ground. Based on the HELCOM AIS data, vessels were categorized into six distinct types that best characterized their specific roles: cargo ships, large passenger ships, medium-sized passenger vessels, large work vessels, small vessels, and others. In the vicinity of Utö, cargo and large passenger ships constituted the majority of vessels.

In our study, AIS data was utilized to calculate emissions using the STEAM model and to investigate potential changes in the shipping traffic density close to the Utö measurement site.

3.3. The STEAM model

The STEAM model integrates AIS-derived data alongside technical insights into the global fleet and fundamental naval architecture principles (Jalkanen et al., 2009, 2012, 2014; Johansson et al., 2017). The model is subsequently used to predict vessel water resistance and instantaneous engine power of the main and auxiliary engines. The STEAM model facilitates the prediction of instantaneous fuel consumption and the emissions of specific pollutants (Jalkanen et al., 2016).

The input data for the STEAM model concerning ship properties, encompasses, among other factors, measured emission factors (when available), shaft generators, specific fuel oil consumption, and fuel type and sulfur content for main and auxiliary engines, along with installed abatement techniques (Jalkanen et al., 2009). The computed emissions for NO_x used in this study encompass the International Maritime Organization (IMO) registered traffic. The assessment includes all vessel traffic equipped with AIS transceivers while specifically excluding those navigating in inland waterways from the dataset. In this work, STEAM was used to run the Baltic Sea ship emission timeseries between years 2006 and 2020, including the regulatory

changes to sulfur content of marine fuels. Further, STEAM was used to study the vessel traffic statistics near the Utö island, and determine hourly vessel counts. STEAM was also used as a supporting tool in interpretation of air quality measurement results to determine the emission intensity of local plumes

4 Results and discussion

The results are divided in subsections addressing the changes in long-range transport and shipping, investigating (i) the air quality time series and (ii) the wind-direction dependence of observations. For convenience, we will use the following abbreviations throughout the text: SECA2006, SECA2010, and SECA2015 to refer to the SECA regulations established on 19 May 2006, 1 July 2010, and 1 January 2015, respectively.

4.1 Changes of long-range transport and shipping emissions

The SO₂ emissions in the Northern Europe have shown a decreasing trend in most of the emission sectors. Figure 2 illustrates the combined emissions in each emission sector from countries that have a major impact on the northern Baltic Proper / southern border of Archipelago Sea during the period 2003–2020. The dominant emissions originate from the energy (indicated as “public power” in Fig. 2) and industry sectors, exceeding the emissions from shipping by more than an order of magnitude. The average lifetime of SO₂ in the lower atmosphere is approximately 1–3 days (Lee *et al.*, 2011; Beirle *et al.*, 2014). Part of the SO₂ observed at Utö therefore originates from long-range transport of regional background pollution, while another portion is attributed to shipping traffic in the vicinity of Utö (e.g., Lee *et al.*, 2011). Long-range transported background pollution is well-diluted, whereas local emissions tend to manifest as short-term, high-concentration peaks.




Figure 2: The combined annual SO_x emissions of Estonia, Finland, Lithuania, Latvia, Sweden, Russia and Poland divided in source sectors during the period 2003-2020. The implementation of subsequent SECA regulations is indicated with vertical dashed lines. Note that the figure is presented on a logarithmic scale (y-axis). The source of SO_x emissions data for the different sectors is EMEP (European Monitoring and Evaluation Programme).

205


Figure 3 displays the monthly shipping emissions in the Baltic Sea calculated using the STEAM model. The changes in SO₂ emissions are a combination of changes in emissions due to the implementation of SECA regulations, and changes in shipping traffic density:

210

From 2006 onward, shipping activity in the Baltic Sea has generally increased, with some exceptions. In particular, notable downturns in the total number of ships and the transport work were observed during the economic recession, which started in 2008 and continued until 2009 affecting most areas of the neighbouring countries of the Baltic Sea (Jalkanen et al., 2014), and again during the COVID-19 pandemic in 2019 (HELCOM, 2021). An exception to this upward trend occurred in 2013, with

215 approximately 350 000 ship crossings in the used passage lines compared to 2012, likely due to the reductions in cargo ships attributed to reduced shipping activity resulting from the economic recession during these years (HELCOM, 2014). Finland also experienced a unique gross domestic product (GDP) decline from 2011 to 2012, largely due to weak export performance (Bank of Finland, 2018).

220 Emission computations from STEAM (Fig. 3) indicate a gradual decline in SO_x and PM_{2.5} emissions attributed to maritime traffic in the Baltic Sea starting in 2006. This decline in SO_x and PM_{2.5} emissions continued gradually until 2014, at which point Baltic economies began to recover, reaching pre-crisis production levels (Bank of Finland, 2018). Following the introduction of SECA2015, a sharp decrease in SO_x and PM_{2.5} emissions is observed, highlighting the impact of stricter emissions controls. The magnitude of the relative reduction of PM_{2.5} emissions attributed to shipping is smaller than that of
225 the SO_x emissions. The predicted annual NO_x emissions during the period between 2006 and 2020 have remained relatively stable, except for between 6 November 2017 and 31 December 2017, when a 95 % decrease in AIS data was observed due to an issue with AIS data reception.



230 **Figure 3: Predicted emissions of NO_x, SO_x, and PM_{2.5} attributed to maritime traffic in the Baltic Sea from January 2006 to December 2020, computed using the STEAM model. The SECA regulations introduced in 2006, 2010 and 2015 are represented by vertical dashed lines.**

4.2 Observed changes in ship traffic and concentrations of SO₂ and other air quality parameters at Utö

235 At Utö, local marine traffic typically consists of a few larger vessels per day. We analysed the AIS data to quantify the number of ships passing within 5 km of the island of Utö on a daily and yearly basis (Fig. 4). As expected, results showed a declining trend in the yearly number of ships up to 2010, which coincided with the 2008–2009 economic recession affecting the majority of the Baltic Sea region. The decline continued through 2013, with a slight increase observed in 2014, as production levels across the region began to recover (Bank of Finland, 2018).

240

From 2015 to 2020, however, no clear increasing or decreasing trend in the number of ships was detected.




Figure 4: The total number of ships per year (left y-axis) that have passed the island of Utö at distances closer than 5 km (blue vertical bars). The total number of ships per day (orange line) is shown on the right y-axis.


The observed 30-day moving averages and moving percentiles of the concentration data for SO₂, PM_{2.5}, NO, NO_x and O₃ from 2006 to the end of 2020 are presented in Fig. 5. The dashed vertical lines represent the introduction of SECA2006, SECA2010 and SECA2015, corresponding to 1.5 %, 1.0 % and 0.1 % sulfur fuel content limits, respectively. The effects of the sulfur cap in 2006 and 2015 are clearly visible, while the impact of 2010 is negligible. The change in 2015 is apparent in the shifts of the 1st and 99th percentiles corresponding to short-duration local ship plumes. Additionally, the mean and median values after 2015 are closer to each other, indicating that the higher peaks in the data occur less frequently (Fig. 5a).


Table 2: Annual average and median concentrations of SO₂ at Utö. Values for the period 2003–2005 are based on hourly averages, while for the period 2006–2020, 1-minute time resolution data has been used. STD is the standard deviation. N (%) represents the fraction of the year for which high-quality data is available, expressed as a percentage. High-quality data, used in our analysis, are defined as valid data recorded, excluding those compromised by factors such as instrument malfunctions, environmental interference, or calibration issues.


	Mean [µg m ⁻³]	Median [µg m ⁻³]	STD [µg m ⁻³]	N (%)
2003	1.38	-	2.15	98.3
2004	1.17	-	1.13	91.5
2005	1.32	-	1.45	97.1
2006	0.93	0.55	1.51	55.7

2007	0.97	0.56	1.58	99.8
2008	0.70	0.40	1.19	98.9
2009	0.73	0.42	1.07	98.4
2010	1.12	0.58	1.80	89.0
2011	1.00	0.57	1.81	93.0
2012	0.78	0.45	1.18	95.8
2013	0.58	0.38	0.81	84.5
2014	0.75	0.46	1.08	99.5
2015	0.37	0.29	0.47	99.4
2016	0.45	0.32	0.68	99.4
2017	0.34	0.27	0.54	98.3
2018	0.50	0.36	0.60	99.3
2019	0.41	0.36	0.50	99.1
2020	0.34	0.32	0.35	99.1


260 In general, the reduction in the release of SO₂ emissions attributed to shipping could also result in a decline in the production of particulate matter originating from precursor compounds from shipping. However, as the results (Fig. 5 and Appendix A) demonstrate, there were no step changes in the PM_{2.5} concentration during the considered period. This does not mean that the emissions of particulate matter from ships were at the same level as before the introduction of SECA regulations, but rather that the measurement of PM_{2.5} is not sensitive enough to show the changes in the ultrafine particle size range. PM_{2.5} concentrations also had some negative values, which were removed, leading to gaps in the time series (Fig. 5b). Seppälä et al. (2021) studied the impact of SECA on particle number size distribution (size range 7–537 nm) at Utö and found that the main changes due to SECA in particle number occurred in sizes between 33 and 144 nm. Thus, the impact on particle mass, resulting mainly from larger particles, is minor. Similarly to PM_{2.5}, no decreases were observed in the NO_x, NO and O₃ concentrations. However, for NO and NO_x, a period of data from 22 May 2010 to 15 June 2011 was removed (Figs. 5c and 5d) due to abnormally low values, likely caused by overly strict data processing. The year-to-year temporal variation of PM_{2.5}, NO, NO_x and O₃ (yearly mean and median values and standard deviation) are presented in Appendix A.







275







Figure 5: Time series of measured concentrations for (a) SO₂, (b) PM_{2.5}, (c) NO, (d) NO_x, and (e) O₃ from 2003 to the end of 2020 at the air quality station in Utö. The concentrations are presented as moving percentiles based on 1-minute resolution data within a 30-day window, with 30-day moving mean, median, and 1st–99th percentiles. For the years 2003–2005, only the moving mean is shown, based on 1-hour resolution data. The implementation of SECA regulations in 2006, 2010, and 2015 is indicated by three vertical dashed lines.

Since the years of 2006, 2010 and 2015 marked significant milestones, as they represented the introduction of the new SECA regulations, we decided to focus on the periods that occurred before and after the implementation of SECAs: 2003–2005, 2007–2009, 2011–2013 and 2016–2018. Specifically, we compared the three-year average of SO₂ concentrations before the SECA introduction (2003–2005) to those following each SECA enactment (2007–2009, 2011–2013, and 2016–2018) (Table 2). Our findings revealed that three-year average SO₂ concentrations from the pre-SECA period (2003–2005) decreased by 38 %, 39 %, and 67 % in comparison to the post-SECA periods (2007–2009, 2011–2013, and 2016–2018), respectively.

As indicated in the previous section, SO₂ emissions have been widely reduced across various industrial sectors. To estimate the local effect of shipping, we utilized NO, which is highly reactive and has a short lifetime in the air. Since it is one of the main gases in ship exhaust, it can serve as a marker for ship plumes originating near Utö. Therefore, we selected periods when

295 SO₂ and NO both showed clear peaks simultaneously in the data, and when the wind was blowing from the western sector (180–360 degrees). The prominence of the peak indicating the peak height relative to other data was chosen to be 2 μg m⁻³ for both NO and SO₂. Figure 6 depicts the SO₂ concentration normalized by NO concentration during these plumes. Normalization using CO₂ concentrations would have allowed further analysis of fuel sulfur content, unfortunately such data has not been measured at Utö in a location suitable for ship SO₂ plume research prior to implementation of SECA in 2015. There is a clear difference before and after January 2015. Prior to the implementation of the strictest SECA regulation (i.e. SECA2015), the
300 SO₂ concentrations during local pollution plumes were of the same magnitude as NO. However, after the SECA2015 regulation was enacted, the SO₂ concentrations decreased significantly within these near-source plumes.






305 **Figure 6:** Time series of the observed SO₂/NO concentration ratio, when the wind was blowing from the westerly directions (wind direction 180-360°) and the plume from the passing ships was transported over the Utö island. Only the periods with simultaneous peaks in both SO₂ and NO signals were included into the analysis.

4.3. Dependence of concentrations on local wind direction

To study the impacts of ships passing by Utö, we selected SO₂ concentration data based on wind direction. First, we separated
310 the datapoints measured when the wind was blowing from the shipping lane (covering wind directions from 185° to 315°) towards the measurement site from the rest of the data.

All data, along with data from when the wind was blowing from the direction of the shipping lane and data from the background sector (wind directions excluding the shipping lane sector) are shown in Figs. 7a–c. Similar to the previously presented results,
315 there is an evident decrease in SO₂ concentrations after the SECA2015, and a slight decrease after the SECA2010. This decreasing trend is visible in all three plots, but the most pronounced decrease occurs after 2015 when only wind directions from shipping lane were considered.




320

325 **Figure 7: The time series of measured SO₂ concentrations (1-min. temporal resolution; grey solid lines) and their moving means (blue solid lines) and medians (red solid lines) in 30 days window with a) all the data points included (unstratified), b) data only from 130° wide wind sector from direction of shipping lane, extending from 185° to 315° and c) data only from background wind sectors excluding shipping lane, less than 185° and over 315°. The temporal resolution of the SO₂ concentration data was 1 minute on average.**

330 To analyse the effect of wind direction in more detail, we generated concentration rose plots in finer 10-degree intervals (Fig. 8). To assess the effects of the implementation of SECA regulation in 2015, we chose two years: one immediately preceding the new regulation, 2014, and the other representing the 5-year period after, 2019. The years 2016, 2017, 2018 and 2019 were not substantially different in terms of SO_x emissions from shipping (cf. Fig. 2) or the number of ships (Fig. 4). In this regard, any of these years could have been selected as an example year for the post-SECA 2015 analysis. The selected year 2019 was prior to the COVID-19 pandemic and selected for comparison. The pandemic did not affect the emissions in Europe in 2019; its effects were felt only during the subsequent years.

335 The measured annual distribution of wind directions is determined by the local climate at the site and potentially by local factors affecting wind measurements. The most common wind directions in this region are southwesterly, with relatively high contributions also from northwesterly directions at angles from 320° to 350° (Fig. 8a and Fig. 8c). This northwesterly sector coincidentally corresponds to the routes of shipping to and from the harbor and wharf area of the island. The shipping traffic to the island typically consists of smaller vessels on average compared to the average marine transport in the neighboring ship lane. The upper panels in Figs. 8a–b include all measured concentrations. The observed SO₂ concentrations were systematically and substantially higher in 2014 compared to the values in 2019. Furthermore, if only peak concentrations during the plumes are considered (lower panels, Figs. 8c–d), it is evident that the numbers of the highest concentrations were drastically higher in 2014 compared to the corresponding values in 2019 and originated from the direction of the adjacent shipping lane (i.e.,
340 from the westerly directions).
345



350 Figure 8: The measured concentrations of SO₂ at the island of Utö as a function of the wind direction, as polar histograms with 1-minute resolution. The panels a) and c) correspond to the data in 2014 and the panels b) and d) to those in 2019. These two years were selected to represent the situation before and after the SECA regulation in 2015. The upper panels (a and b) include all datapoints, whereas the lower panels (c and d) include only the data, for which the concentrations were higher than a selected threshold value ($5 \mu\text{g m}^{-3}$). In panels a) and b), the red colour in the legend represents all data points with values larger than $0 \mu\text{g m}^{-3}$. Overlaid on the red, the yellow colour shows data points with values below $1.5 \mu\text{g m}^{-3}$, and the violet colour, which overlays the yellow, indicates data points with values below $0.5 \mu\text{g m}^{-3}$. The radial axis represents the number of measured cases for each wind direction sector.

355

4.4. Uncertainties

Given the nearly two decade long observation period, several factors could introduce uncertainties in the results. Instrumentation for all variables changed over time, and quality assurance methods improved. Furthermore, there were changes in the personnel responsible for maintaining and ensuring data quality. Although standard protocols were followed and measurement diaries were kept, these changes may have affected the results.

360

5 Conclusions

In this study, we have presented and analysed a high temporal resolution air quality dataset of SO₂, PM_{2.5}, NO, NO_x, and O₃ concentrations and of wind data for one and a half decades at the island of Utö in the Baltic Sea. This dataset is particularly unique, as there are no other similar long-term air quality observations from remote locations in the Baltic Sea, and the data set has not been previously published or analyzed. Utö's central location next to northern Baltic Proper, with minimal influence from other pollution sources, makes it an ideal site to study long-term pollution trends from shipping. The period of the study includes all three consecutive revisions of the SECA regulations.

The findings derived from the analysis of the air quality and meteorological datasets and STEAM modelling indicated that the SECA regulations, introduced in 2006, tightened in 2010 and further strengthened in 2015, have been successful in reducing the observed SO₂ concentrations at Utö during the studied period 2003–2020. Comparing three-year average SO₂ concentrations pre-SECA (2003–2005) to post-SECA periods (2007–2009, 2011–2013, 2016–2018) revealed reductions of 38 % (SECA 2006), 39 % (SECA 2010), and 67 % (SECA 2015), respectively. During the target period, there were no clear decreasing trends for the concentrations of PM_{2.5}, NO_x, NO and O₃. The year-to-year variations of the concentrations were substantial for all pollutants; these were attributed partly to the variations in regional meteorology, partly to the variations of emissions.

We also analysed polar histograms of SO₂-concentration during two representative years, 2014 and 2019, i.e., before and after the implementation of the SECA2015. This analysis was done both for the whole dataset and for selecting only the highest measured concentrations, above a specified threshold. The numbers of the cases with the highest SO₂ concentrations were drastically higher in 2014 compared to the corresponding values in 2019. The highest concentrations were mostly originating from the direction of the adjacent shipping lane. This analysis therefore indicates that the highest measured concentrations attributed to shipping have substantially decreased after the SECA2015.

By filling an important gap in the current literature, this study provides a thorough view of the air quality trends in the Baltic Sea and highlights the importance of good quality, high temporal resolution, long-term air quality data at remote marine research stations. Such observations are crucial for both quantitative and qualitative analyses of the impacts of regulatory environmental legislation. The findings of this work will provide valuable insights into the effectiveness of SECA regulations and serve as a benchmark for local and regional dispersion modelling efforts for future research utilizing the data of this study.

Appendix A

Table A1: Annual average and median concentrations of PM_{2.5} at Utö. Values for the period 2003–2005 are based on hourly averages, while for the period 2006–2020 on 1-minute time resolution data has been used. STD is the standard deviation. N (%) represents the fraction of the year for which high-quality data is available, expressed as a percentage. High-quality data, used in our analysis, are defined as valid data recorded, excluding those compromised by factors such as instrument malfunctions, environmental interference, or calibration issues.

PM _{2.5}	Mean [µg m ⁻³]	Median [µg m ⁻³]	STD [µg m ⁻³]	N (%)
2003	5.97	-	5.57	20.0
2004	6.82	-	6.07	84.8
2005	6.07	-	6.13	93.8
2006	8.53	5.67	9.44	68.9
2007	5.28	3.93	5.49	99.9
2008	5.25	3.87	5.99	98.8
2009	5.30	4.37	4.84	98.0
2010	7.73	6.02	7.07	58.2
2011	7.60	6.02	7.23	79.3
2012	6.14	4.70	5.54	72.0
2013	5.92	4.54	5.32	58.5
2014	8.97	7.24	7.34	82.4
2015	7.06	5.66	6.15	71.3
2016	6.38	5.28	5.15	62.1
2017	5.78	4.27	6.31	63.1
2018	5.61	3.85	4.93	97.3
2019	5.56	4.14	4.53	89.5
2020	5.51	3.99	5.00	86.2

Table A2: Annual average and median concentrations of NO at Utö. Values for the period 2003–2005 are based on hourly averages, while for the period 2006–2020 1-minute time resolution data has been used. STD is the standard deviation. N (%) represents the fraction of the year for which high-quality data is available, expressed as a percentage. High-quality data, used in our analysis, are defined as valid data recorded, excluding those compromised by factors such as instrument malfunctions, environmental interference, or calibration issues.

NO	Mean m ⁻³]	[µg	Median m ⁻³]	[µg	STD m ⁻³]	[µg	N (%)
2003	0.40		-		0.76		98.3
2004	0.39		-		0.60		91.1
2005	0.35		-		0.57		70.4
2006	0.46		0.14		1.85		71.3
2007	0.40		0.16		1.62		86.9
2008	0.29		0.15		1.30		99.1
2009	0.33		0.22		1.04		97.9
2010	0.64		0.48		1.27		35.9 *
2011	0.33		0.25		0.98		50.2 *
2012	0.33		0.24		1.11		98.1
2013	0.30		0.21		1.14		98.0
2014	0.29		0.09		1.09		99.4
2015	0.22		0.08		1.08		98.6
2016	0.23		0.09		1.25		99.5
2017	0.29		0.09		1.86		86.0
2018	0.34		0.11		1.76		99.3
2019	0.31		0.12		1.73		99.1
2020	0.22		0.10		1.41		95.9

*Low N (%) is attributed to the removal of data from 22 May 2010 to 15 June 2011, due to abnormally low values, likely caused by overly strict data processing.

Table A3: Annual average and median concentrations of NO_x at Utö. Values for the period 2003–2005 are based on hourly averages, while for the period 2006–2020 1-minute time resolution data has been used. STD is the standard deviation. N (%) represents the fraction of the year for which high-quality data is available, expressed as a percentage. High-quality data, used in our analysis, are defined as valid data recorded, excluding those compromised by factors such as instrument malfunctions, environmental interference, or calibration issues.

NO _x	Mean m ⁻³]	[µg	Median m ⁻³]	[µg	STD [µg m ⁻³]	N (%)
-----------------	---------------------------	-----	-----------------------------	-----	---------------------------	-------

2003	-	-	-	-
2004	-	-	-	-
2005	-	-	-	-
2006	-	-	-	-
2007	-	-	-	-
2008	2.57	1.51	4.43	47.1
2009	3.32	2.12	4.46	98.1
2010	4.20	2.72	5.66	35.9 *
2011	3.41	2.39	4.09	50.2 *
2012	3.87	2.59	4.93	98.1
2013	3.69	2.46	4.75	98.0
2014	4.32	2.98	4.97	99.4
2015	3.31	2.13	4.35	98.6
2016	3.47	2.25	4.91	99.5
2017	3.50	2.13	5.97	86.0
2018	4.49	3.01	5.92	99.3
2019	4.06	2.59	5.91	99.1
2020	3.27	2.31	4.78	95.9

*Low N (%) is attributed to the removal of data from 22 May 2010 to 15 June 2011, due to abnormally low values, likely caused by overly strict data processing.

- 415 Table A4: Annual average and median concentrations of O₃ at Utö. Values for the period 2003–2005 are based on hourly averages, while for the period 2006–2020 on 1-minute time resolution data has been used. STD is the standard deviation. N (%) represents the fraction of the year for which high-quality data is available, expressed as a percentage. High-quality data, used in our analysis, are defined as valid data recorded, excluding those compromised by factors such as instrument malfunctions, environmental interference, or calibration issues.

O ₃	Mean [µg m ⁻³]	Median [µg m ⁻³]	STD [µg m ⁻³]	N (%)
2003	67.04	-	19.97	92.3

2004	69.73	-	16.47	90.0
2005	68.71	-	18.20	97.1
2006	74.94	74.08	22.23	77.6
2007	65.66	65.87	16.46	99.8
2008	64.99	63.46	20.38	98.8
2009	60.09	59.91	16.90	98.3
2010	64.43	64.49	16.35	99.1
2011	66.17	66.10	16.42	95.1
2012	65.64	66.28	16.00	97.5
2013	70.28	68.93	18.05	98.4
2014	62.36	62.43	15.98	82.0
2015	67.58	68.13	13.65	99.3
2016	66.68	66.51	15.86	99.5
2017	65.61	66.83	14.53	98.7
2018	67.64	67.59	17.67	99.3
2019	67.25	67.72	17.92	99.1
2020	64.55	65.35	15.55	99.1

420 Code availability

Codes for the analysis of long-term pollutant concentrations are available upon request from Androniki Maragkidou (androniki.maragkidou@fmi.fi). Codes for the concentration analyses are available from Tiia Grönholm (Tiia.Gronholm@fmi.fi).

Data availability

425 The 1-minute air quality data from 2006-2020 is available in Zenodo [DOI to be added]. The data set also includes 10-minute resolution meteorological data (wind speed, wind direction, air temperature, air pressure, relative humidity and precipitation) and 1-hour air quality data from 2003-2005. Meteorological data is also available from FMI Open Data (<https://en.ilmatieteenlaitos.fi/open-data>).

Author contributions

430 AM conducted part of the analyses, wrote the first draft of the manuscript, and lead the processing of the manuscript to its final
format. TG performed most of the data analysis and part of the writing. LR processed the AQ data and computed some of the
figures. LR, LL, and JK contributed to the planning of the study and the writing of the article. TM and TA were responsible
for the measurements at the Utö island and contributed to the analysis of the measurements. J-PJ has done the computations
on the emissions of shipping using the STEAM model. JN is responsible of the computer analyses underlying Fig. 7. All
435 authors commented on the manuscript and provided feedback.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

[The authors would like to thank Marko Juola for supporting the measurements.](#)

440 Financial support

This work was partly funded by the European Union's Horizon 2020 research and innovation programme Evaluation, control
and Mitigation of the EnviRonmental impacts of shippinG Emissions (EMERGE) [grant agreement No 874990]. This work
reflects only the authors' view and CINEA is not responsible for any use that may be made of the information it contains. Part
of the research has been funded through H2020 project Joint European Research Infrastructure of Coastal Observatories:
445 Science, Service, Sustainability (JERICO-S3) [grant agreement No 871153]. Observations carried out at Utö have utilized
ICOS (Integrated Carbon Observing System), ACTRIS (Aerosol, Clouds and Trace Gases Research Infrastructure) and
FINMARI (Finnish Marine Research Infrastructure) research facilities and resources.

References

Ahlnäs, K.: Variations in Salinity at Utö 1911–1961, *Geophysica*, 8, 135–149, 1961.

450 Andersson, C., Bergström, R., and Johansson, C.: Population exposure and mortality due to regional background PM in Europe
– Long-term simulations of source region and shipping contributions, *Atmospheric Environment*, 43, 3614–3620,
<https://doi.org/10.1016/j.atmosenv.2009.03.040>, 2009.

455 Anenberg, S., Miller, J., Henze, D., and Minjares, R.: A global snapshot of the air pollution-related health impacts of transportation sector emissions in 2010 and 2015, The International Council on Clean Transportation (ICCT), Washington, DC, USA, 2019.

Anttila, P., Tuovinen, J.-P., Trends of primary and secondary pollutant concentrations in Finland in 1994–2007. *Atmos. Environ.*, 44, 38–49, 2010.

Bank of Finland: https://www.bofbulletin.fi/en/2018/1/euro-countries-recovered-from-crisis-at-different-paces/?fbclid=IwY2xjawGO5nhleHRuA2FlbQIxMAABHY1m3KbNYmUoxS2MUxncUyxp0qms9F38Pw43a7hd9V2WPg6IpiOoctbgyg_aem_cHbm-fQgSJrD6g0gAiOAZQ (last access: 30 October 2024), 2018.

465 Barregard, L., Molnàr, P., Jonson, J. E., and Stockfelt, L.: Impact on Population Health of Baltic Shipping Emissions, *IJERPH*, 16, 1954, <https://doi.org/10.3390/ijerph16111954>, 2019.

Beecken, J., Mellqvist, J., Salo, K., Ekholm, J., Jalkanen, J.-P., Johansson, L., Litvinenko, V., Volodin, K., and Frank-Kamenetsky, D. A.: Emission factors of SO₂, NO_x and particles from ships in Neva Bay from ground-based and helicopter-borne measurements and AIS-based modeling, *Atmos. Chem. Phys.*, 15, 5229–5241, <https://doi.org/10.5194/acp-15-5229-2015>, 2015.

475 Beirle, S., Hormann, C., Penning de Vries, M., Dörner, S., Kern, C., and Wagner, T.: Estimating the volcanic emission rate and atmospheric lifetime of SO₂ from space: A case study for Kilauea Volcano, Hawaii, *Atmos. Chem. Phys.*, vol. 14, pp. 8309–8322, <https://doi.org/10.5194/acp-14-8309-2014>, 2014.

Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E., and Frohn, L. M.: Assessment of past, present and future health-cost externalities of air pollution in Europe and the contribution from international ship traffic using the EVA model system, *Atmos. Chem. Phys.*, 13, 7747–7764, <https://doi.org/10.5194/acp-13-7747-2013>, 2013.

485 Chen, C., Saikawa, E., Comer, B., Mao, X., and Rutherford, D.: Ship Emission Impacts on Air Quality and Human Health in the Pearl River Delta (PRD) Region, China, in 2015, With Projections to 2030, *GeoHealth*, 3, 284–306, <https://doi.org/10.1029/2019GH000183>, 2019.

Contini, D., Donato, A., Gambaro, A., Argiriou, A., Melas, D., Cesari, D., Poupkou, A., Karagiannidis, A., Tsakis, A., Merico, E., Cesari, R., and Dinoi, A.: Impact of Ship Traffic to PM2.5 and Particle Number Concentrations in Three Port-Cities of the Adriatic/Ionian Area, 9, 2015.

490

Corbett, J. J., Fischbeck, P. S., and Pandis, S. N.: Global nitrogen and sulfur inventories for oceangoing ships, *J. Geophys. Res.*, 104, 3457–3470, <https://doi.org/10.1029/1998JD100040>, 1999.

Corbett, J. J., Winebrake, J. J., Green, E. H., Kasibhatla, P., Eyring, V., and Lauer, A.: Mortality from Ship Emissions: A
495 Global Assessment, *Environ. Sci. Technol.*, 41, 8512–8518, <https://doi.org/10.1021/es071686z>, 2007.

Donato, A., Gregoris, E., Gambaro, A., Merico, E., Giua, R., Nocioni, A., and Contini, D.: Contribution of harbour activities and ship traffic to PM2.5, particle number concentrations and PAHs in a port city of the Mediterranean Sea (Italy), *Environ Sci Pollut Res*, 21, 9415–9429, <https://doi.org/10.1007/s11356-014-2849-0>, 2014.

500

EEA (European Environment Agency): Energy efficiency and specific CO2 emissions, <https://www.eea.europa.eu/data-and-maps/indicators/energy-efficiency-and-specific-co2-emissions/energy-efficiency-and-specific-co2-9> (last access: 20 March 2017), 2017 (not up-to-date content, to be replaced with more recent reference)

Eyring, V., Isaksen, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H., and Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping, *Atmospheric Environment*, 44, 4735–4771, <https://doi.org/10.1016/j.atmosenv.2009.04.059>, 2010.

Grönholm, T., Mäkelä, T., Hatakka, J., Jalkanen, J.-P., Kuula, J., Laurila, T., Laakso, L., and Kukkonen, J.: Evaluation of
510 Methane Emissions Originating from LNG Ships Based on the Measurements at a Remote Marine Station, *Environ. Sci. Technol.*, 55, 13677–13686, <https://doi.org/10.1021/acs.est.1c03293>, 2021.

Haglund, K., Claremar, B., and Rutgeresson, A.: Temporal and Spatial variation of Contribution from Ship Emissions to the concentration and deposition of air pollutants in the Baltic Sea, *Management of the Earth system: sustainability science*,
515 <https://doi.org/10.5194/esd-2016-46>, 2016.

Hausfather, Z., and Forster, P.: Analysis: How low-sulfur shipping rules are affecting global warming, <https://www.carbonbrief.org/analysis-how-low-sulfur-shiping-rules-are-affecting-global-warming/> (last access: 19 November 2023), 2023.

520

HELCOM: Eutrophication in the Baltic Sea – An integrated thematic assessment of the effects of nutrient enrichment and eutrophication in the Baltic Sea region., *Balt. Sea Environ. Proc. No. 115B*, 2009.

HELCOM: Annual report on shipping accidents in the Baltic Sea in 2013, 2014.

525

HELCOM: Emissions from Baltic Sea shipping in 2006 - 2020, *Maritime 21-4.2. INF*, 2021.

Hellén, H., Kouznetsov, R., Kraft, K., Seppälä, J., Vestenius, M., Jalkanen, J.-P., Laakso, L., and Hakola, H.: Shipping and algae emissions have a major impact on ambient air mixing ratios of NMHCs and methanethiol on Utö island in the Baltic Sea, *EGUsphere*, <https://doi.org/10.5194/egusphere-2023-2323>, accepted, 2024.

530

Hermansson, A.L., Hassellöv, I.-M., Grönholm, T., Jalkanen, J.-P., Fridell, E., Parsmo, R., Hassellöv, J., and Ytreberg, E., "Monetary return on scrubber installations at the expense of environmental damage", Accepted to *Nature Sustainability* 5.3.2024, preprint <https://www.researchsquare.com/article/rs-3534127/v1>, 2024.

535

Honkanen, M., Tuovinen, J.-P., Laurila, T., Mäkelä, T., Hatakka, J., Kielosto, S., and Laakso, L.: Measuring turbulent CO₂ fluxes with a closed-path gas analyzer in a marine environment, *Atmos. Meas. Tech.*, 11, 5335–5350, <https://doi.org/10.5194/amt-11-5335-2018>, 2018.

540

Honkanen, M., Müller, J. D., Seppälä, J., Rehder, G., Kielosto, S., Ylöstalo, P., Mäkelä, T., Hatakka, J., and Laakso, L.: The diurnal cycle of pCO₂ in the coastal region of the Baltic Sea, *Ocean Sci.*, 17, 1657–1675, <https://doi.org/10.5194/os-17-1657-2021>, 2021.

545

Honkanen, M., Aurela, M., Hatakka, J., Haraguchi, L., Kielosto, S., Mäkelä, T., Seppälä, J., Siiriä, S.-M., Stenbäck, K., Tuovinen, J.-P., Ylöstalo, P., and Laakso, L.: Interannual and seasonal variability of the air-sea CO₂ exchange at Utö in the coastal region of the Baltic Sea, *EGUsphere*, [preprint], <https://doi.org/10.5194/egusphere-2024-628>, 2024.

550

Hunter, K. A., Liss, P. S., Surapipith, V., Dentener, F., Duce, R., Kanakidou, M., Kubilay, N., Mahowald, N., Okin, G., Sarin, M., Uematsu, M., and Zhu, T.: Impacts of anthropogenic SO_x, NO_x and NH₃ on acidification of coastal waters and shipping lanes, *Geophys. Res. Lett.*, 38, L13602, <https://doi.org/10.1029/2011GL047720>, 2011.

IMO (International Maritime Organization), <https://business.un.org/en/entities/13> (last access: 6 October 2017), 2016.

IMO (International Maritime Organization), <https://www.imo.org/en/OurWork/Safety/Pages/AIS.aspx> (last access: 28
555 November 2022), 2020.

Jalkanen, J.-P., Brink, A., Kalli, J., Kukkonen, J., and Stipa, T.: A modelling system for the exhaust emissions of marine traffic
and its application in the Baltic Sea area, *Atmos. Chem. Phys.*, 9, 9209–9223, <https://doi.org/10.5194/acp-9-9209-2009>, 2009.

560 Jalkanen, J.-P., Johansson, L., Kukkonen, J., Brink, A., Kalli, J., and Stipa, T.: Extension of an assessment model of ship traffic
exhaust emissions for particulate matter and carbon monoxide, *Atmos. Chem. Phys.*, 12, 2641–2659,
<https://doi.org/10.5194/acp-12-2641-2012>, 2012.

Jalkanen, J.-P., Johansson, L., and Kukkonen, J.: A Comprehensive Inventory of the Ship Traffic Exhaust Emissions in the
565 Baltic Sea from 2006 to 2009, *AMBIO*, 43, 311–324, <https://doi.org/10.1007/s13280-013-0389-3>, 2014.

Jalkanen, J.-P., Johansson, L., and Kukkonen, J.: A comprehensive inventory of ship traffic exhaust emissions in the European
sea areas in 2011, *Atmos. Chem. Phys.*, 16, 71–84, <https://doi.org/10.5194/acp-16-71-2016>, 2016.

570 Johansson, L., Jalkanen, J.-P., and Kukkonen, J.: Global assessment of shipping emissions in 2015 on a high spatial and
temporal resolution, *Atmospheric Environment*, 167, 403–415, <https://doi.org/10.1016/j.atmosenv.2017.08.042>, 2017.

Jonson, J. E., Gauss, M., Jalkanen, J.-P., and Johansson, L.: Effects of strengthening the Baltic Sea ECA regulations, *Atmos.
Chem. Phys.*, 19, 13469–13487, <https://doi.org/10.5194/acp-19-13469-2019>, 2019.

575

Jonson, J. E., Jalkanen, J. P., Johansson, L., Gauss, M., and Denier Van Der Gon, H. A. C.: Model calculations of the effects
of present and future emissions of air pollutants from shipping in the Baltic Sea and the North Sea, *Atmos. Chem. Phys.*, 15,
783–798, <https://doi.org/10.5194/acp-15-783-2015>, 2015.

580 Kilkki, J., Aalto, T., Hatakka, J., Portin, H., and Laurila, T.: Atmospheric CO₂ observations at Finnish urban and rural sites,
Boreal Environ. Res., 20, 227–242, available at: <http://hdl.handle.net/10138/228120> (last access: 9 July 2018), 2015.

Komppula, M., Vana, M., Kerminen, V.-M., Lihavainen, H., Viisanen, Y., Hörrak, U., Komsaare, K., Tamm, E., Hirsikko, A.,
Laakso, L., and Kulmala, M.: Size distributions of atmospheric ions in Baltic Sea region, *Boreal Env. Res.* 12: 323–336, 2007.

585

Kraft, K., Seppälä, J., Hällfors, H., Suikkanen, S., Ylöstalo, P., Anglès, S., Kielosto, S., Kuosa, H., Laakso, L., Honkanen, M., Lehtinen, S., Oja, J., and Tamminen, T.: Application of IFCB High-Frequency Imaging-in-Flow Cytometry to Investigate Bloom-Forming Filamentous Cyanobacteria in the Baltic Sea, <https://www.frontiersin.org/article/10.3389/fmars.2021.594144>, *Frontiers in Marine Science*, 2021.

590

Kyllönen K., Karlsson V., and Ruoho-Airola T.: Trace element deposition and trends during a ten year period in Finland, *Sci. Total Environ.*, 407, 2260-2269, 2009.

595

Laakso, L., Mikkonen, S., Drebs, A., Karjalainen, A., Pirinen, P., and Alenius, P.: 100 years of atmospheric and marine observations at the Finnish Utö Island in the Baltic Sea, *Ocean Sci.*, 14, 617–632, <https://doi.org/10.5194/os-14-617-2018>, 2018.

600

Laapas, M. and Venäläinen, A.: Homogenization and trend analysis of monthly mean and maximum wind speed time series in Finland, 1959–2015, *Intl Journal of Climatology*, 37, 4803–4813, <https://doi.org/10.1002/joc.5124>, 2017.

Lagler, F., Belis, C., and Borowiak, A.: A Quality Assurance and Control Program for PM2.5 and PM10 Measurements in European Air Quality Monitoring Networks, EUR Scientific and Technical Research Series, Publications Office of the European Union, Luxembourg, 118 pp., ISBN 978-92-79-20480-7, <https://doi.org/10.2788/31381>, 2011.

605

Lauer, A., Eyring, V., Corbett, J. J., Wang, C., and Winebrake, J. J.: Assessment of Near-Future Policy Instruments for Ocean-going Shipping: Impact on Atmospheric Aerosol Burdens and the Earth's Radiation Budget, *Environ. Sci. Technol.*, 43, 5592–5598, <https://doi.org/10.1021/es900922h>, 2009.

610

Laurila, T. and Hakola, H.: Seasonal Cycle of C2–C5 hydrocarbons over the Baltic Sea and Northern Finland, *Atmos. Environ.*, 30, 1597-1607, 1996.

615

Lee, C., Martin, R. V., Van Donkelaar, A., Lee, H., Dickerson, R. R., Hains, J. C., Krotkov, N., Richter, A., Vinnikov, K., and Schwab, J. J.: SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.*, 116, D06304, <https://doi.org/10.1029/2010JD014758>, 2011.

Liu, Z., Lu, X., Feng, J., Fan, Q., Zhang, Y., and Yang, X.: Influence of Ship Emissions on Urban Air Quality: A Comprehensive Study Using Highly Time-Resolved Online Measurements and Numerical Simulation in Shanghai, *Environ. Sci. Technol.*, 51, 202–211, <https://doi.org/10.1021/acs.est.6b03834>, 2017.

620 Matthias, V., Bewersdorff, I., Aulinger, A., and Quante, M.: The contribution of ship emissions to air pollution in the North Sea regions, *Environ. Pollut.*, 158, 2241–2250, <https://doi.org/10.1016/j.envpol.2010.02.013>, 2010.

Mellqvist, J., Beecken, J., Conde, V., and Ekholm, J.: Surveillance of Sulfur Emissions from Ships in Danish Waters, Chalmers University of Technology, Sweden, Report, <https://doi.org/10.17196/DEPA.001>, 2017.

625

Merico, E., Donato, A., Gambaro, A., Cesari, D., Gregoris, E., Barbaro, E., Dinoi, A., Giovanelli, G., Masieri, S., and Contini, D.: Contribution of in-port shipping emissions to gaseous pollutants and particulate matter of different sizes in an Adriatic port-city, *Proscience*, 78–83, <https://doi.org/10.14644/dust.2016.013>, 2016.

630 Partanen, A. I., Laakso, A., Schmidt, A., Kokkola, H., Kuokkanen, T., Pietikäinen, J.-P., Kerminen, V.-M., Lehtinen, K. E. J., Laakso, L., and Korhonen, H.: Climate and air quality trade-offs in altering ship fuel sulfur content, *Atmos. Chem. Phys.*, 13, 12059–12071, <https://doi.org/10.5194/acp-13-12059-2013>, 2013.

635 Picone, M., Russo, M., Distefano, G., Baccichet, M., Marchetto, D., Volpi Ghirardini, A., Hermansson, A., Petrovic, M., Gros, M., Garcia, E., Giubilato, E., Calgareo, L., Magnusson, K., Granberg, M. and Marcomini, A.: Impacts of exhaust gas cleaning systems (EGCS) discharge waters on planktonic biological indicators, *Mar. Pollut. Bull.*, 190, 114846, <https://doi.org/10.1016/j.marpolbul.2023.114846>, 2023.

640 Raudsepp, U., Laanemets, J., Maljutenko, I., Hongisto, M., and Jalkanen, J.-P.: Impact of ship-borne nitrogen deposition on the Gulf of Finland ecosystem: an evaluation, *Oceanologia*, 55, 837–857, <https://doi.org/10.5697/oc.55-4.837>, 2013.

Rautiainen, L., Tyynelä, J., Lensu, M., Siiriä, S., Vakkari, V., O’Connor, E., Hämäläinen, K., Lonka, H., Stenbäck, K., Koistinen, J., and Laakso, L.: Utö Observatory for Analysing Atmospheric Ducting Events over Baltic Coastal and Marine Waters, *Remote Sensing*, 15, 2989, <https://doi.org/10.3390/rs15122989>, 2023.

645

Repka, S., Mellqvist, J., Borkowski, T., Jalkanen, J.-P., Jonson, J. E., Barregard, L., Olaniyi, E., Prause, G. K., Gauss, M., Walden, J., Svensson, E., Genikhovich, E., Rumyantsev, D., Boyesen, J., Wenske, C., de Carvalho, T. P., Lähteenmäki-Uutela, A., Johansson, L., Conde, V., ... & Atari, S. Final report - Clean Shipping: Exploring the impact of emission regulation (EnviSuM), 2019.

650

Seppälä, S. D., Kuula, J., Hyvärinen, A.-P., Saarikoski, S., Rönkkö, T., Keskinen, J., Jalkanen, J.-P., and Timonen, H.: Effects of marine fuel sulfur restrictions on particle number concentrations and size distributions in ship plumes in the Baltic Sea, *Atmos. Chem. Phys.*, 21, 3215–3234, <https://doi.org/10.5194/acp-21-3215-2021>, 2021.

655 Smith, T. W. P., Jalkanen, J.-P., Anderson, B. A., Corbett, J. J., Faber, J., Hanayama, S., O’Keeffe, E., Parker, S., Johansson, L., Aldous, L., Raucci, C., Traut, M., Ettinger, S., Nelissen, D., Lee, D. S., Ng, S., Agrawal, A., Winebrake, J. J., Hoen, M., Chesworth, S., and Pandey, A.: Third IMO Greenhouse Gas Study 2014, International Maritime Organization, London, UK, 2015.

660 Sofiev, M., Winebrake, J. J., Johansson, L., Carr, E. W., Prank, M., Soares, J., Vira, J., Kouznetsov, R., Jalkanen, J.-P., and Corbett, J. J.: Cleaner fuels for ships provide public health benefits with climate tradeoffs, *Nat Commun*, 9, 406, <https://doi.org/10.1038/s41467-017-02774-9>, 2018.

665 Tian, L., Ho, K., Louie, P. K. K., Qiu, H., Pun, V. C., Kan, H., Yu, I. T. S., and Wong, T. W.: Shipping emissions associated with increased cardiovascular hospitalizations, *Atmos. Environ.*, 74, 320–325, <https://doi.org/10.1016/j.atmosenv.2013.04.014>, 2013.

UNCTAD (United Nations Conference on Trade and Development): Review of Maritime Transport 2019, United Nations Conference on Trade and Development, 132 pp., 2019.

670 UNEP, Coastal zone management, <https://www.unep.org/topics/ocean-seas-and-coasts/regional-seas-programme/coastal-zone-management>, cited 4.4.2024.

675 Van Aardenne, J., Colette, A., Degraeuwe, B., Hammingh, P., Viana, M., and Vlieger, I.: The impact of international shipping on European air quality and climate forcing, Technical Report No. 4/2013, Technical Report No. 4/2013., EEA (European Environment Agency), Copenhagen, Denmark, 1–84 pp., 2013.

680 Yang, M., Bell, T. G., Hopkins, F. E., and Smyth, T. J.: Attribution of atmospheric sulfur dioxide over the English Channel to dimethyl sulfide and changing ship emissions, *Atmos. Chem. Phys.*, 16, 4771–4783, <https://doi.org/10.5194/acp-16-4771-2016>, 2016.