

Future Changes of Compound Explosive Cyclones and Atmospheric Rivers in the North Atlantic

Ferran Lopez-Marti^{1,2,*}, Mireia Ginesta^{3,*}, Davide Faranda³, Anna Rutgersson^{1,2}, Pascal Yiou³, Lichuan Wu¹, and Gabriele Messori^{1,4,5}

¹Department of Earth Sciences, Uppsala University, Uppsala, Sweden

²Centre of Natural Hazards and Disaster Science (CNDS), Uppsala, Sweden

³Laboratoire des Sciences du Climat et de l'Environnement, UMR8212 CEA-CNRS-UVSQ, IPSL & U Paris-Saclay, Gif-sur-Yvette, France

⁴Department of Meteorology and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

⁵Swedish Centre for Impacts of Climate Extremes (climes), Uppsala University, Uppsala, Sweden

*These authors contributed equally to this work.

Correspondence: Ferran Lopez-Marti (ferran.lopez-marti@geo.uu.se)

Abstract. The explosive development of extratropical cyclones and atmospheric rivers play a crucial role in driving extreme weather in the mid-latitudes, such as compound windstorm-flood events. Although both explosive cyclones and atmospheric rivers are well-understood and their relationship has been studied previously, there is still a gap in our understanding of how a warmer climate may affect their concurrence. Here, we focus on evaluating the current climatology and assessing changes in the future concurrence between atmospheric rivers and explosive cyclones in the North Atlantic. To accomplish this, we independently detect and track atmospheric rivers and extratropical cyclones and study their concurrence in both ERA5 reanalysis and CMIP6 historical and future climate simulations. In agreement with the literature, atmospheric rivers are more often detected in the vicinity of explosive cyclones than non-explosive cyclones in all datasets, and the atmospheric river intensity increases in all the future scenarios analysed. ~~Moreover~~Furthermore, we find that explosive cyclones associated with atmospheric rivers are tend to be longer-lasting and deeper than ~~other explosive cyclones~~those without. Notably, we identify a significant and systematic future increase in the ~~eylones-cyclones~~ – atmospheric river concurrences. Finally, under the ~~worst-case~~ high-emission scenario, the explosive cyclone – atmospheric river concurrences show an increase and model agreement over western Europe. As such, our work provides a novel statistical relation between explosive cyclones and atmospheric rivers in CMIP6 climate projections and a characterization of their joint changes in intensity and location.

15 1 Introduction

Atmospheric Rivers (ARs) are narrow and elongated corridors of horizontal moisture transport usually associated with the cold front of an extratropical cyclone (Bao et al., 2006; Ralph et al., 2004, 2017). They play an essential role in the atmospheric water vapour cycle in the mid-latitudes, accounting for 90% of the poleward moisture transport (Zhu and Newell, 1998; Guan and Waliser, 2015). Moreover, ARs drive wet and windy extreme weather events, particularly in western continental coasts, such as Western Europe (Zhu and Newell, 1998; Guan and Waliser, 2015; Lavers and Villarini, 2015, 2013; Gimeno et al.,

2016). In some coastal areas, intense ARs are associated up to 95% of the time with extreme precipitation and up to 75% with wind extremes (Waliser and Guan, 2017). As a result, ARs have widespread socioeconomic impacts; for example, the majority of European storms causing insurance losses of a billion US dollars or more are linked to ARs (De Vries, 2020).

Another meteorological feature that can lead to wet and windy extremes are explosive cyclones (ECs), also known as weather bombs (Roebber, 1984; Reale et al., 2019). ECs are rapidly intensifying mid-latitude cyclones. Historically, ECs have been identified as those with a deepening rate of more than 24 hPa in 24 hours, and scaled by the latitude (Sanders and Gyakum, 1980), although this definition has been repeatedly challenged, especially for the Southern Hemisphere (Allen et al., 2010). Events such as the Presidents' Day Snowstorm of 1979 (Schultz, 2022), characterized by poor forecast accuracy, sparked extensive research into this phenomenon. Upon landfall during their intensification phase, ECs can produce widespread damage and impacts associated with strong winds, heavy precipitation, and storm surges (Fink et al., 2012; Liberato et al., 2013; Ludwig et al., 2015; Seiler and Zwiers, 2016b; Reale et al., 2019; Ginesta et al., 2023).

ECs and the associated ARs thus play a crucial role in driving extreme weather ~~events~~ hazards in the mid-latitudes (Liberato et al., 2013; Davolio et al., 2020). An illustrative example is Storm Alex, an EC associated with an AR that at first produced extreme winds in France and the UK and then led to record-breaking precipitation in Italy in October 2020 (Davolio et al., 2023; Ginesta et al., 2023). The climatological relationship between ECs and ARs has been previously studied and the literature evidences that ARs are more often found in the surroundings of EC than non-ECs (Eiras-Barca et al., 2018; Zhang et al., 2019; Guo et al., 2020). ~~The physical reason behind this is that ARs incorporate moisture into~~ ARs are important sources of moisture for cyclonic systems, leading to a faster and stronger deepening of the cyclone (Zhu and Newell, 1994; Ferreira et al., 2016) ~~-In this context, an important process for explosive cyclogenesis is the latent heat release from the moisture brought by the AR. Cyclones and it has been suggested that they can enhance cyclone deepening through moist diabatic processes (Zhu and Newell, 1994; Ferreira et al., 2016), such as cloud condensation (Pinto et al., 2009). In addition, ECs with ARs show larger moisture inflow and deepen more rapidly than ECs without an AR but do not show significant differences in low-level baroclinicity nor upper-level potential vorticity, suggesting that diabatic~~ processes are important contributors to their intensification (Zhang and Ralph, 2021) ~~processes are the main contributors to explosive intensification (Pinto et al., 2009; Zhang and Ralph, 2021).~~ However, the extent to which these moist diabatic processes, compared to other factors such as upper-level forcing, influence cyclone intensification can vary from case to case (Pfahl and Sprenger, 2016; Ginesta et al., 2024).

A range of studies have investigated the impact of anthropogenic climate change on ~~Extratropical Cyclones~~ extratropical cyclones and ARs individually (Lavers et al., 2015; Zappa et al., 2013). The thermodynamic response of ARs to climate change is characterized by an increase in Integrated Water Vapor Transport (IVT). This increase is driven by the Clausius-Clapeyron relation, which implies a rise in moisture content in a warmer atmosphere. However, ~~the vertically integrated water vapor content undergoes further amplification compared to surface~~ integrated water vapor vapour (IWV) is expected to experience a larger increase than surface water vapour under climate change (Payne et al., 2020). This thermodynamic signal would act to increase the number of ARs detected in a warmer climate (Thandlam et al., 2022; Espinoza et al., 2018; Zhang et al., 2024; O'Brien et al., 2022; Wang et al., 2023). Similarly, the thermodynamic response acts to increase the precipitation within extratropical cyclones (Yettella and Kay, 2017). The dynamical response to climate change, such as changes in atmospheric

circulation patterns, is more uncertain (Shepherd, 2014). In the North Atlantic, dynamic changes are mostly driven by changes in the eddy-driven jet, which serves as a guide for extratropical cyclones. The tug of war between the upper tropospheric warming and the Arctic amplification leads to a high uncertainty regarding the changes in the jet over the North Atlantic and western Europe (Shaw et al., 2016). However, climate models indicate a decline in the number of extreme extratropical cyclones in the North Atlantic, and a local increase over the North Sea in EC frequency (Priestley and Catto, 2022; Zappa et al., 2013; Seiler and Zwiers, 2016a). Regarding ARs, studies also point to an increase in frequency, intensity, and size in western Europe and a northward shift of the AR location and landfall (Lavers et al., 2013; Ramos et al., 2016; Gao et al., 2016; Zhang et al., 2024).

The hazards associated with the joint occurrence of explosive cyclones and atmospheric rivers, especially along the western coast of Europe, underscores the importance of evaluating the projected changes in their concurrence in future climates. In this study, we assess future projections of the interplay between ECs and ARs using state-of-the-art [models from the Coupled Model Intercomparison Project phase 6 \(CMIP6 data, Eyring et al. \(2016b\)\)](#). We specifically evaluate the frequency of EC and AR concurrence in the present climate for the ERA5 reanalysis and CMIP6 models, and in three end-of-century scenarios for the CMIP6 models. Moreover, we also assess the future changes in the intensity and location of such compound events.

This manuscript is structured as follows: Section 2 describes the datasets employed in this study, while Section 3 explains the methodologies for tracking ARs and cyclones, and the calculation of their concurrences. In Section 4 we evaluate and discuss the performance of the CMIP6 models with ERA5 reanalysis. Section 5 shows the results and discussion of the future changes in frequency, intensity, and location of this compound event. Finally, Section 6 summarizes the key findings and provides the conclusions of this study.

2 Data

We use the ECMWF reanalysis ERA5 (Hersbach et al., 2020) with a horizontal resolution of $0.25^\circ \times 0.25^\circ$ as an ~~observationally-constrained~~ [observationally constrained](#) reference for the current climate and validation of the Global Climate Models (GCMs). For the AR detection, the variables used are specific humidity, meridional and zonal wind components at 1000, 925, 850, 700, 600, 500, 400 and 300 hPa (Section [??3.2](#)). For cyclone detection, the variable used is the sea level pressure (SLP). In both cases, the variables are at 6 hourly resolution during the extended winter period (October to March) between 1980 and 2009 for the North Atlantic region [[25-65°N; 8015-75°N; 90°W-10W-20°E](#)] ~~-(hereafter referred to as the Data Domain)~~. [To mitigate issues caused by the cyclone tracking algorithm, which tends to generate stationary 'artefact' cyclones along the western boundary, we apply a buffer zone of 10° on this data domain. This adjustment ensures that artefacts at the western boundaries are excluded. As a result, the domain used for all analyses presented in the results section is \[25-65°N; 80°W-10°E\], referred to simply as domain from here on.](#)

Further, we use one member each of six different GCMs from the CMIP6 dataset (Eyring et al., 2016a): MPI-ESM1-2-LR, MPI-ESM1-2-HR, NorESM2-MM, EC-Earth3, CMCC-ESM2, MIROC6. More detailed information about the GCMs used can be found in the Appendix A1. We evaluate the listed GCMs for two periods: the current climate (1980-2009) using historical

simulations and the future climate at the end of the 21st century (2070-2099), where we used simulations following the SSP1-
 90 2.6; SSP2-4.5 and SSP5-8.5 forcing scenarios (Riahi et al., 2017). These 30-year datasets consist of 29 full winters and two
 partial winters (January to March for 1980 and 2070, and October to December for 2009 and 2099). Variables used for tracking
 ARs and cyclones in the GCMs are the same as used for ERA5. The current list of GCMs used in this study is limited by the
 availability of 6-hourly instantaneous variables for the historical and the three scenarios experiments in hybrid-sigma pressure
 model levels, which are required to interpolate to the necessary pressure levels for the IVT calculation when detecting ARs.
 95 This limitation in the analyzed data prevents a complete assessment of model performance, as only a single member from each
 model is used, and a multi-member ensemble for each model would be necessary for a more robust evaluation of the individual
 model uncertainty. For this reason, we evaluate the results using the multi-model mean of the ensemble.

3 Methods

3.1 Extratropical cyclone tracking

100 There are several detection and tracking algorithms for extratropical cyclones, most of them using either SLP or lower-
 tropospheric vorticity (Neu et al., 2013). Here, we detect and track cyclones based on SLP using the TempestExtreme algorithm
 developed by Ullrich et al. (2021). This command-line software facilitates adaptable and rapid feature detection and tracking
 for extratropical cyclones ~~but also and~~ for ARs.

To identify extratropical cyclones, we recognize candidate "nodes" corresponding to local minima in the SLP field with
 105 the same set-up as in Ullrich et al. (2021). Nodes within a 6° great circle distance (GCD) of each other are merged. Next,
 to connect these candidate nodes into tracks the distance between consecutive detections should not exceed 6 GCD degrees.
 The tracks must persist for a minimum of 24 hours, the maximum duration between two detections is set at 6 hours, and the
 cyclones must have moved at least 12 GCD degrees to filter out stationary lows, such as the Icelandic Low. The relevant codes
and the detailed set-up are available in the Appendix A2.

110 In addition, extratropical cyclones are classified as ECs if their Normalized Deepening Rate (NDR), as defined by Sanders
 and Gyakum (1980), is equal to or higher than 1:

$$NDR_c = \frac{DR_{24h} DR_{24h} \sin(60^\circ)}{24h \ 24hPa \ \sin(\varphi)} \quad (1)$$

where DR_{24h} is the pressure difference over 24 hours measured at the storm centre and φ is the latitude at its second-time
 step. Cyclones that do not fulfil this condition are classified as non-ECs. Table A2 in Appendix A4 lists the number of cyclone
 115 tracks of ECs and non-ECs ~~tracked~~ in each model and scenario and in ERA5 within the North Atlantic domain. The number of
~~eyelones~~ cyclone tracks detected fits inside the range number of cyclones detected in Neu et al. (2013) using several cyclone
 tracking algorithms.

Figure 1 shows the ~~storm track density~~ number of cyclone timesteps detected within a 3° spherical cap per month, hereafter
referred to as cyclone track density, for ECs and non-ECs in both ERA5 and CMIP6 models. These results agree with Priestley

120 and Catto (2022) and Zappa et al. (2013) track density for CMIP6 and Zappa et al. (2013) for CMIP5 track densities despite using different tracking algorithms. Our tracking method shows some differences between ERA5 and CMIP6 models (Figure 1c,d). In the models, there are fewer ECs and non-ECs around the south tip of Greenland, Iceland, west Scandinavia and the Gulf of Genoa in the Mediterranean Sea. In particular, for ECs the and in the North Sea. For non-ECs, CMIP6 models underestimate track density in the northern part of the domain, whereas for the non-EC CMIP6 show a northward shift of the storm track with lower cyclone track density in the south and higher track density in the north of the domain.




Figure 1. EC track density climatology during the extended winter (ONDJFM) [1980-2009] for ERA5 (a), historical simulations of CMIP6 (b), and difference between ERA5 and CMIP6 (c). In panels (d,e,f) same as (a,b,c) but for non-ECs. Units are the number of eyelones-cyclone timesteps per 1.53° spherical cap per month.

3.2 AR detection and tracking

There are two primary approaches to detecting ARs: one involves using the Integrated Water Vapor, commonly applied to satellite data, while the other, more broadly used, consists in computing the IVT (Gimeno et al., 2014; Shields et al., 2018). In this study, we calculate the IVT for both reanalysis and model data. The IVT is defined for each grid point as:

$$IVT = \left[\left(\frac{1}{g} \int_{1000hPa}^{300hPa} qu dp \right)^2 + \left(\frac{1}{g} \int_{1000hPa}^{300hPa} qv dp \right)^2 \right]^{1/2} \quad (2)$$

where q is the specific humidity, u is the zonal wind component, v is the meridional wind component, and g is the gravitational acceleration. Moreover, we separately compute the eastward (IVT_E) and northward (IVT_N) components of IVT, which correspond to the two terms inside of the brackets in Eq. (2) respectively.

135 For the detection of ARs, we also use the TempestExtreme algorithm developed by Ullrich et al. (2021). We find candidates for atmospheric rivers by detecting ridges in the IVT field. Ridges are defined as points where the Laplacian of the IVT is below $-4 \times 10^4 \text{ kg m}^{-2} \text{ s}^{-1} \text{ rad}^{-2}$, as this operator identifies elongated areas and regions of local minima. In addition, the IVT should be higher than $250 \text{ kg m}^{-1} \text{ s}^{-1}$. Each candidate should have an area larger than $4 \times 10^5 \text{ km}^{-2}$. These thresholds were defined and tested by Ullrich et al. (2021), and show strong agreement with other tracking algorithms (Collow et al., 2022). The detected candidates are concatenated if at least one grid point is ~~detected as AR in sequential timesteps~~ identified as an AR in consecutive timesteps, meaning that the AR area at a consecutive timestep spatially overlaps with the previous AR. In addition, they should last 60 hours. The successful ARs that fulfil all these requirements are labelled as AR grid-points which will be used later to determine if cyclones are concurrent to an AR (Section 3.3). This AR tracking methodology is less sensitive to a generalized increase of IVT in the future climate due to the Clausius-Capeyron relationship because we detect AR candidates using the Laplacian of the IVT instead of the IVT field. Thus, AR candidates are detected when having a pronounced gradient of IVT (not dependent on the background IVT), as discussed further in Section 5, ~~this may affect the detected changes in.~~ This detection algorithm may influence the observed changes in the number of AR tracks in future climates. The relevant codes are available in Appendix A3 and the detailed number ARs tracks detected are reported in Table A3 in Appendix A4.

150 Figure 2a shows the ~~frequency of ARs as the percentage of time-steps detected with an percentage of timesteps with a detected~~ AR for ERA5, hereafter referred to as AR frequency. Our tracking methodology ~~reproduces well the AR climatology accurately reproduces the AR frequency~~ when compared with Guan and Waliser (2015). The AR ~~climatological frequency for frequency in~~ the historical simulations of the CMIP6 models is ~~larger when higher~~ compared to ERA5 and ~~shows a shift south. In other words, more ARs are detected in CMIP6 and those tracks are chiefly located in~~ exhibits a southward shift, with more AR timesteps detected primarily in the lower midlatitudes (Figure 2b,c).




Figure 2. AR frequency climatology during the extended winter (ONDJFM) [1980-2009] for ERA5 (a), historical simulations of CMIP6 CMIP6 models (b), and difference between ERA5 and CMIP6 (c). Units are the percentage of ~~time-steps timesteps~~ detected as AR.

155 3.3 Concurrences

For each extratropical cyclone (both ECs and non-ECs) we compute the Maximum Deepening Point (MDP), which is the maximum difference in SLP between two consecutive 6-hourly timesteps. This metric allows us to evaluate the influence of

ARs on the development of the cyclone before and after its maximum intensification, which is when the potential impact of ARs on cyclone deepening is expected, following the approach used by Eiras-Barca et al. (2018).

160 Subsequently, we determine whether a specific timestep of an extratropical cyclone (EC or non-EC) is linked to an AR **by detecting the presence of an AR within** if at least one grid point within 1500 km from the centre of the cyclone is part of an AR (detected and tracked independently, see Section 3.2). This AR search radius is consistently applied across all time steps of the EC and non-EC tracks. By selecting a 1500 km **of the cyclone radius**, our methods align with those of (Eiras-Barca et al., 2018) , with the primary difference between the two methods being the AR and cyclone tracking algorithms used. Almost 50% of the
165 identified ARs are located in the southeastern quadrant of the cyclone (Supplementary Figures S1 and S2). The rationale for using a 1500 km radius is based on the consideration that an AR can influence the cyclone by delivering moisture to the warm conveyor belt or feeder airstream (Dacre et al., 2019), which are typically located in the southeastern quadrant and within this distance from the cyclone center.

Figure 3 shows an example of our detection methodology applied to storm Xynthia. EC Xynthia underwent rapid intensifica-
170 tion before making landfall on February 27, 2010. It caused widespread damage across Western European countries, specially France and Spain. In Figure 3, the black dots represent the cyclone's path, while the red crosses indicate the time steps concurrent with the presence of an AR. The shaded areas depict the regions identified as ARs at each concurrence time step during the cyclone trajectory. Xynthia was associated with an AR during its intensification phase until landfall, suggesting that the AR may have contributed to its intensification (Liberato et al., 2013).




Figure 3. Example of detection and tracking of an EC ([Xynthia, 26/02/2010 18 UTC – 01/03/2010 00 UTC](#)) concurrent to an AR in ERA5. Red crosses mark the location of the cyclone at times when it concurred with an AR. Black dots indicate no concurrence. Shaded areas depict the regions identified as ARs at each concurrence time step during the cyclone trajectory.

175 [We further evaluate the internal variability of the CMIP6 concurrences by analyzing the spread within the multi-model ensemble. Additionally, we assess the inter-annual variability of the concurrences by calculating the standard deviation at each time step of the rate of coincidence between cyclones and ARs over the 29 full extended winter seasons \(ONDJFM\).](#)

4 Concurrences in Present Climate

In this section, we evaluate the concurrence of ECs and ARs in the ERA5 reanalyses and we compare it with those obtained
 180 in climate models. Figure 4a shows the rate of coincidence between ECs and ARs, indicating the fraction of ECs that are associated with an AR (see Section 3.3), as a function of time from the MDP point. ERA5 shows a maximum rate of coincidence at around 6 hours after the MDP point the MDP. This maximum is about 0.650.72, meaning that 6572% of the ECs are associated with ARs. The rate of coincidence is minimum at the initial stages of cyclone formation, with 4536 hours before and after the MDP, with 55% of the ECs associated with an AR. During the dissipation stages of the cyclones, around 55% of ECs are
 185 associated with an AR. In other words, the rate of coincidence is higher after the MDP, when the ECs are deeper (see Section 5.3). This suggests that a mature cyclone, when its SLP is lower, favours the detection of an AR in its surroundings. These

results agree with the ones obtained by Eiras-Barca et al. (2018) ~~despite a shift of 6–12 hours in the concurrence peak, which comes from how the MDP is calculated (See Section 6), despite changes of up to 0.05 in the rate of coincidence probably due to differences in tracking method and data used.~~ The curves of the CMIP6 models show a similar shape to that of ERA5, with a maximum rate of coincidence centred ~~6/12 hours after at~~ the MDP, ~~a minimum at the formation of the cyclone, and a secondary minimum at the dissipation stage and two minimums~~ 36 hours before and after the MDP. Some of the models, specifically CMCC, MPI-HR, ~~and MPI-LR, overestimate the rate and MIROC6, have higher rates~~ of coincidence for almost the whole lifetime of the ECs. CMCC and MPI-LR show a maximum of ~~75~~ around 80% of ECs associated with an AR, which is 10% more than ERA5. On the contrary, ~~EC-EARTH3, MIROC6, and NorESM underestimate the rate of coincidence before the MDP, while they overestimate it, especially MIROC6, after the MDP. EC-EARTH3 and NorESM are the models that are more similar to the reanalysis, with maximum biases of around 0.05 along the lifetime of the ECs~~ EC-Earth3 and NorESM2-MM show very similar rates of coincidence along the ECs lifetime, with differences up to 0.02 with respect to ERA5. Figure 4c shows the ~~inter-seasonal~~ evolution of inter-annual variability as the standard deviation of the rate of coincidence between ECs and ARs over the ~~30 winter seasons~~ 29 winter seasons at each time from -36 h to +36 h from the MDP. CMIP6 models and ERA5 show an ~~inter-seasonal variability between 0.07 and 0.1, being a bit~~ inter-annual variability between 0.06 and 0.18, being higher on the first and last time steps from the MDP. CMIP6 models reproduce similar concurrence variability as ERA5 and differences in concurrence rate between them and ERA5 are within the internal variability, ~~computed as the spread of the multi-model ensemble.~~ However, MPI-LR has a higher rate of coincidence with respect to ERA5 and the rest of the models 36 hours before the MDP.

Figure 4b shows the rate of coincidence of non-ECs and ARs. ERA5 ~~shows a much flatter curve than for of~~ has a much smaller variation of the rate of coincidence for non-ECs than for the ECs. There is a maximum in the coincidence rate for ERA5 at the MDP, around 50%, with two minimums at ~~the formation and dissipation stages around 0.45%~~ 36 hours before and after the MDP of the cyclones, ~~approximately 45%.~~ Thus, there is a relatively small variation in the coincidence rate throughout the lifetime of the non-ECs. These results for non-ECs also agree with the ones obtained by Eiras-Barca et al. (2018). All models resemble ERA5 regarding the shape of the curve. However, CMCC ~~overestimates the coincidence rate~~ has a higher rate of coincidence throughout the lifetime of the cyclones ~~by around 0.05~~ of up to 0.02 with respect to ERA5. On the contrary, ~~EC-EARTH3 underestimates this rate by approximately 0.05~~ EC-Earth3 and MIROC6 have lower rates of coincidence, of up to almost 0.1 with respect to ERA5. The models that ~~perform better, especially around the MDP, are~~ have the most similar rates with respect to ERA5 are MPI-HR, MPI-LR, MIROC6, and NorESM. ~~They nonetheless underestimate or overestimate the rate by around 0.03 at the formation and dissipation stages, and NorESM2-MM.~~ We emphasize that caution is needed when interpreting these model performances, as they may be influenced by internal variability due to the use of only a single member. Additional sensitivity tests of the impact of resolution on ERA5 are available in the supplementary material. Figure 4d shows the ~~inter-seasonal~~ inter-annual variability for the rate of coincidence between non-ECs and ARs. In this case, CMIP6 models and ERA5 show an ~~inter-seasonal variability between 0.09~~ inter-annual variability between 0.08 and 0.12, showing larger values on the first and last time steps for some models, reaching 0.18 for MPI-LR. CMIP6 models

reproduce similar concurrence variability as ERA5 and the differences in concurrence rate between them and ERA5 are within their ~~internal~~ inter-annual variability.

In summary, for ECs, ~~maximum model biases occur~~ the coincidence rate maximum model difference with respect to ERA5 occurs during the MDP, reaching around ~~0.080~~ 1. Even though this represents almost 15% of the ERA5 value, the ~~internal~~ inter-annual variability of the datasets is ~~slightly larger~~ comparable. For non-ECs, ~~models have biases particularly during the formation and dissipation stages of cyclones, peaking at around~~ the different models show a more constant difference with ERA5 with a maximum of 0.05, or around 11% of the ERA5 value, but ~~again~~ smaller than the ~~internal~~ inter-annual variability. Overall, the models reproduce the qualitative features of the life cycle of the rate of coincidence between both ECs and non-ECs and ARs. The ~~quantitative~~ quantitative differences with ERA5 are generally smaller than the datasets' internal variability.




Figure 4. Rate of coincidence between **explosive-cyclones (ECs)** and **atmospheric rivers (ARs)** (a) and **non-explosive-cyclones (non-EC)** and ARs (b) for ERA5 and the historical runs of CMIP6 models [1980–2009] during the extended winter (ONDJFM). **Inter-seasonal Inter-annual** variability as the standard deviation of the rate of coincidence over the 30 winter seasons between ECs and ARs (c) and non-ECs and ARs (d).

230 5 Future projections

In this section, we analyze changes in the concurrence of cyclones and ARs in CMIP6 models between the historical simulations (1980–2009) and the three scenario simulations (2070–2099).

5.1 Changes in concurrence frequency

Figure 5a,b show a clear increase in the rate of coincidence of ARs with ECs and non-EC events across all warming scenarios. 235 In the SSP5-8.5 scenario, the multi-model mean rate of coincidence increases by approximately **+2%** 0.14 throughout the

lifetime of ~~both classes of cyclones~~ ECs and 0.12 for non-ECs. For the SSP1-2.6 and SSP2-4.5 scenarios, the increases are around ~~5% and 9%~~ 0.07 and 0.08 for ECs, respectively, and ~~4% and 8% for non-EC events~~ 0.05 and 0.07 for non-ECs.

Table A2 in Appendix A4 shows the number of ECs and non-ECs tracks and table A3 shows the number of AR tracks detected for ERA5 and the CMIP6 models in each scenario. Hence, these tables also show the changes ~~in~~ (in brackets) in the occurrence of the features individually in future scenarios compared to historical ones. The number of ECs decreases in all future scenarios with respect to the historical and for all models, ~~usually~~ with the largest decrease for the ~~worst-case high-emission~~ scenario (SSP5-8.5). ~~There is little change~~ The decrease in the number of non-ECs ~~, although most of the models also project a small decrease in future scenarios is smaller when compared with the EC, although for SSP1-2.6 not all the models show a decrease of non-ECs but for SSP2-4.5 and SSP5-8.5 all the models project a decrease.~~ Regarding the number of ARs, ~~for almost all the models there is a small increase in the number in the worst-case scenario~~ AR tracks, in four models (EC-Earth3, NorESM2-MM, MIROC6, CMCC-ESM2), there is an increase in all emission scenarios. However, the changes are not linear, that is, some of the models (MPI-LR, MPI-HR, ~~NorESM~~ NorESM2-MM, MIROC6) depict a larger number of ARs in the SSP1-2.6 scenario than in the SSP5-8.5. The ~~lower-than-expected non-proportional~~ increase in the number of AR tracks ~~in the with~~ warmer scenarios at the end of the century could be explained by the nature of the AR tracking, which detects ARs in the Laplacian of the IVT (Section ~~??3.2~~). However, Table A3 only shows the number of AR tracks and does not account for the AR duration or extension, which might result in an increase of AR activity under a warmer climate (Zhang et al., 2024; O'Brien et al., 2022). Despite not finding a strong increase ~~of in~~ ARs' tracks, the total number of times an AR is detected in the surroundings of both types of cyclones nonetheless shows a clear increase in all scenarios ~~, (Fig. 5a,b)~~, with the increase becoming larger with higher warming. ~~In addition,~~

The increase in the rate of coincidence between cyclones and ARs is partly the result of a decrease in the total number of cyclone time-steps and a direct increase in the number of concurrent cyclone/AR time-steps. To clarify this, we calculated the absolute number of compound events of ECs or non-ECs together with ARs also increases in all scenarios (not shown) (including all CMIP6 models) of cyclones concurrent with ARs at the MDP and it increases across all scenarios: 4.6%, 7.1% and 6.3% for SSP1-2.6, SSP2-4.5 and SSP5-8.5 respectively. In addition, the absolute number of cyclone time-steps also at the MDP decreases: -3.6%, -5.2% and -12.3% for SSP1-2.6, SSP2-4.5 and SSP5-8.5 respectively. Hence, the ~~combination of a small combined effect of an~~ increase in AR ~~frequency and a small occurrence and a~~ decrease in the number of ~~ECs and non-ECs results in an increase in the ratio of coincidence with cyclones, the last one especially relevant for SSP5-8.5, explains the rise in the~~ rate of coincidence as the level of warming increases (Figure 5a,b). ~~This points to~~ These findings suggest that changes in the characteristics of ~~the ARs or ARs~~, cyclones, or ~~of their interactions,~~ as drivers of the identified ~~their interactions~~ may be driving the observed changes, rather than ~~these being a simple statistical effect of more individual~~ the result being merely a statistical artefact of more cyclones and ARs occurring individually.

The spread across CMIP6 models is the primary source of uncertainty when evaluating changes in the rate of coincidence, and exhibits overlap between different scenarios (Fig. 5a, b). The choice of model thus has a significant impact on the results when quantifying the increase in cyclone-AR compound events. Nevertheless, the inter-model spread of the SSP5-8.5 scenario ~~'s inter-model spread~~ is well-separated from that of the historical period for both ECs and non-ECs (Fig. 5a, b). The inter-model

spread further exceeds ~~inter-seasonal~~inter-annual variability for the models we analyse (cf. Fig. 5a, b and 5c, d). Indeed, the ~~inter-seasonal~~inter-annual variability of the model ensemble is consistently smaller than the multi-model mean differences between any of the three warming scenarios and historical for ECs (Fig. 5c), and systematically smaller than the multi-model mean differences for non-ECs in the SSP5-8.5 scenario and, at most lead times, for the SSP2-4.5 scenario (Fig. 5d). Our results
275 thus point to anthropogenic climate change exerting a strong influence on the increase in coincidence rates of cyclones and ARs, which is most evident for higher warming levels and, for the latter, cannot be attributed to natural variability or inter-model spread.




Figure 5. Rate of coincidence [between ECs and ARs \(a\), and non-ECs and ARs \(b\)](#) for the historical runs and ERA5 [1980–2009] and as well as the future [scenarios-scenario runs \[2070–2099\]](#) [between ECs and ARs during the extended winter \(a\) and non-EC and ARs \(b\)](#). Solid lines show the multi-model mean, shape-points individual models and shades the inter-model spread. The difference in the multi-model mean rate of coincidence between the forcing scenarios and historical runs for ECs and ARs (c) and non-EC and ARs (d). Solid lines show the multi-model mean difference, shades show the [inter-seasonal-inter-annual](#) variability of the multi-model ensembles in future scenarios and historical, and the dashed line the [inter-seasonal-inter-annual](#) variability for ERA5.

5.2 Changes in AR intensity

The intensity of ARs can be quantified in different ways, here we use the maximum value of IVT (IVT-max) within the detected AR. We use this metric because it represents a good proxy for the total transport of moisture of an AR and is not dependent on the boundaries of the detected AR (Ralph et al., 2017). Moreover, the IVT-max is usually located at the core of the AR, this makes this variable also a good proxy for AR extension and duration as ARs reaching higher IVT values at their core tend to

be larger and last longer (Guan et al., 2023). Our results indicate that the IVT-max of ARs associated with ECs and non-ECs increases proportionally to the warming in future scenarios (Figure 6). When comparing the IVT-max between scenarios, it is particularly relevant that under the SSP5-8.5 the multi-model mean remains above $1250 \text{ kg}\cdot\text{m}\cdot\text{s}^{-1}$ for more than 48h, ~~this means that on average ARs under such conditions will be primarily hazardous according to the AR scale by~~. This extended duration suggests that, on average, ARs under these conditions will primarily be hazardous, as classified by the AR scale from Eiras-Barca et al. (2021). For ARs with non-EC events, the IVT-max is constant throughout the entire cyclone lifetime, suggesting a uniform inflow of atmospheric moisture transport. On the other hand, for ARs associated with ECs, the IVT-max reaches its peak around the MDP of the cyclone's lifecycle, showing the maximum moisture transport during the cyclone's most active phase (Figure 6a,b). The AR intensity for ERA5 is larger than any model for the historical period because ERA5's resolution is almost 4 times higher than the ~~CMIP6~~ CMIP6 models, and attains larger values of IVT-max. Additional sensitivity tests of the impact of resolution on the ERA5 results are available in the supplementary material. Despite this, the ERA5 and CMIP6 curves show the same qualitative behaviour during the cyclones' lifetime and the ~~inter-seasonal~~ inter-annual variability in both datasets is of the same order of magnitude (Figure 6).

When evaluating uncertainties for changes in the AR intensity, ~~inter-seasonal variability is slightly~~ inter-annual variability is smaller than the inter-model spread. Both are dependent on model resolution, as the IVT-max magnitude is sensitive to resolution and all datasets are treated on their native grids. For both ARs with ECs and non-ECs, the inter-model spread of both the SSP2-4.5 and the SSP5-8.5 scenarios is well-separated from that of the historical period (Fig. 6a, b). Moreover, the difference in the multi-model mean between the future scenarios and the historical is more pronounced than the ~~inter-seasonal~~ inter-annual variability for all three scenarios and for both ECs and non-ECs (Figure 6c,d). Thus, the IVT-max increases in all scenarios and our results point to this being larger than both inter-model spread and internal model variability in most scenarios, and thus ascribable to anthropogenic warming.




Figure 6. Mean IVT-max for the historical runs and ERA5 [1980–2009] and the future scenarios runs [2070–2099] of ARs associated with ECs (a) and ARs associated with non-EC (b) during the extended winter (ONDJFM). Solid lines show the multi-model mean, shape-points individual models and shades the inter-model spread. The difference in the multi-model mean IVT-max between the forcing scenarios and historical run for ECs and ARs (c) and non-EC and ARs (d). Solid lines show the multi-model mean difference, shades show the ~~inter-seasonal~~ inter-annual variability of the multi-model ensembles in future scenarios and historical, and the dashed line the ~~inter-seasonal~~ inter-annual variability for ERA5.

5.3 Changes in **explosive** cyclone intensity

305 In this section, we examine the cyclone core SLP to assess changes in the intensity of ECs and non-ECs with the presence or not of ARs under climate change. The presence of ARs influences the evolution of SLP through the life-cycle of the two types of cyclones differently, but in both cases when an AR is present the cyclone is deeper (Fig. 7e, f). The concurrence of ECs with ARs makes the ECs substantially deeper, especially before the MDP, where ECs are between ~~7.5 hPa and 15~~ 2.5

hPa and almost 10 hPa deeper. After the MDP the influence of the AR on the cyclone intensity decreases. Thus, ARs play an important role before the EC MDP, making these storms deeper from an earlier stage and resulting in longer-lasting and more intense cyclones. The particular influence in the first half of the cyclone life span suggests that moisture brought by the AR plays a key role in the EC intensification. On the other hand, the non-ECs also get deeper when occurring together with an AR, with a deepening between 2 and 7 hPa, having its peak also just before the MDP. The results from ERA5 show ~~the same-similar~~ behaviour for both types of cyclones ~~but with lower intensity when compared to the models~~ (Fig. 7b,d). Before the MDP, the models tend to simulate lower SLP for ECs with ARs and higher SLP for ECs without ARs. After the MDP, the models generally simulate higher SLP for both ECs with and without ARs. For non-ECs, the models have higher SLP values after the MDP compared to ERA5. However, the ERA5 values fall within the ensemble spread of historical values, indicating that they are within the uncertainty range of the models.

The ~~climate change influence~~ influence of climate change on cyclone intensity for any of the four types of compound event events is very limited, as the historical and scenario lines, along with their respective spreads in Fig. 7, are close to each other or overlap. All three forcing scenarios show the same life-cycle behaviour and intensity changes as the historical ensemble. Figure A1 shows that the difference of the multi-model mean between the forcing scenarios and the historical is negligible (less than 1 hPa). It is slightly larger only for the ECs without ARs, probably due to the reduced number of these events (but as a result their inter-seasonal-inter-annual variability is also larger). The <1 hPa shift can be compared to the models' inter-seasonal inter-annual variability of between 1 and 2 hPa (Fig. A1). Similarly, the multi-model spreads of all scenarios are about ± 4 hPa (Fig. 7). Thus, the models do not show a robust change in ~~the cyclone intensity~~ cyclone intensity when averaged across the North Atlantic basin under climate change scenarios.




Figure 7. Mean SLP for the historical runs and ERA5 [1980–2009] and the future scenarios runs [2070–2099] of ECs with ARs (a), non-EC with ARs (b), ECs without ARs (c) and non-ECs without ARs (c) during the extended winter (ONDJFM). Solid lines show the multi-model mean, shape-points individual models and shades the inter-model spread. The difference in the multi-model mean SLP between the ECs with AR and without AR (e) and non-EC with AR and without AR (f).

5.4 Changes in location of concurrent ECs and ARs

To assess the spatial distribution of changes observed across the North Atlantic, we show the rate of EC-AR coincidence over a 3-degree spherical cap. In the historical period, the rate is highest along the climatological North Atlantic storm track, extending from the western to the northeastern part of the basin (Fig. 8a,b). CMIP6 models reproduce closely this spatial pattern. Differences between future projections and the historical period show an increase in the rate of coincidence, all the stronger for higher warming levels. The **worst-case scenario (SSP5-8.5) high-emission scenario** shows the largest increase and a high model agreement across almost the entire domain. The southern part of the domain shows a noisier pattern and weaker model agreement due to the reduced number of events in the area. Only the **worst-case high-emission** scenario shows a clear increase in the rate of coincidence over Europe, showing model agreement over the British Isles, northern France and the Iberian peninsula. In other words, a larger proportion of landfalling ECs together with ARs at their maximum deepening point is expected under the SSP5-8.5 scenario. This highlights the possibility of an increase in wet and windy extremes in Western Europe in a high-emissions future.




Figure 8. Rate of coincidence between ECs and ARs for ERA5 (a), and the multimodel-mean of the CMIP6 historical runs (b). Differences between future projections and historical periods of the three scenarios: SSP1-2.6 (c), SSP2-4.5 (d), and SSP5-8.5 (e). Dots denote where all the CMIP6 models agree on the sign of change.

340 6 Conclusions

We have used six Global Climate Models (GCMs) participating in CMIP6 to evaluate the change in the concurrences of cyclones and Atmospheric Rivers (ARs) in three different future scenarios under climate change in the North Atlantic. We have ~~evaluated~~ compared the performance of the models using the ERA5 reanalysis. Our main findings are summarized as follows:

- 345 – For the present period, in ERA5 nearly ~~65~~72% of the Explosive Cyclones (ECs) are associated with an AR ~~within the 6 hours following the~~ at the Maximum Deepening Point (MDP). ~~The fact that AR concurrences are larger after the MDP suggests that mature ECs (when they are deeper) can facilitate the formation of ARs in their surroundings~~ This higher rate of coincidence around the MDP indicates that ARs are more likely to occur when the EC is at its peak deepening stage. Despite some biases in the magnitude of the coincidence rate, CMIP6 models exhibit qualitatively similar concurrence
- 350 rates to ERA5, ~~peaking 6 to 12 hours after the MDP. Conversely, the evolution of the~~ In contrast, the concurrence rate over the ~~eyclone's lifespan is less pronounced for~~ lifespan of non-Explosive Cyclones (non-ECs) shows much smaller variation compared to ECs.
- In future scenarios, there is an increase in the rate of coincidence between ECs and ARs, with the magnitude of the increase proportional to the level of warming. All models agree on the sign of the change. In the ~~worst-case~~ high-emission scenario (SSP5-8.5), there is a maximum increase in the rate of coincidence from ~~0.68 to 0.80~~ 0.72 to 0.87. Even in the best-case scenario (SSP1-2.6), there is an increase of around ~~6%~~ 0.07. For non-ECs, there is an increase in the rate of coincidence, ranging from ~~5% to 12%~~ 0.05 to 0.13 across scenarios. The increase in SSP5-8.5 and SSP2-4.5 scenarios is larger than the ~~inter-seasonal~~ inter-annual model variability for both ECs and non-ECs, but the inter-model spread of the historical period is only well-separated from that of the ~~SSP5-RCP8~~ SSP5-8.5 scenario.
- 360 – In all warming scenarios there is an increase in AR intensity. This is larger than the ~~inter-seasonal~~ inter-annual variability for ARs associated with both ECs and non-ECs. Moreover, the inter-model spread of both the SSP2-4.5 and the SSP5-8.5 scenarios is well-separated from that of the historical period. Under the SSP5-8.5 scenario, the maximum Integrated Vapour Transport (IVT) of ARs associated with ECs is projected to exceed on average $1250 \text{ kgm}^{-1}\text{s}^{-1}$ for more than 48 hours, indicating exceptional and hazardous AR conditions.
- 365 – ECs are deeper when ~~they are~~ associated with an AR, ~~especially before the MDP. This indicates that ECs with an AR get deeper.~~ The presence of an AR is associated with ECs deepening earlier compared to ~~ECs without ARs, making these eyclones those without, making them~~ deeper for a longer time period and potentially more hazardous. Non-ECs are also deeper when having an AR in their surroundings but their intensification is lesser compared to the ECs with ARs.
- The concurrence of ARs with ECs will be more frequent in the North Atlantic basin in future climates. There is an
- 370 increase in both agreement among models and magnitude of the change with the degree of warming, with the SSP5-8.5 scenario showing the largest change. Under this most severe scenario, Europe is exposed to this increase, particularly the southern Iberian peninsula, the British Isles, France and Scandinavia.

Our results for concurrence rates of cyclones and ARs in the present climate are broadly consistent with Eiras-Barca et al. (2018), ~~despite the latter study using the lower resolution reanalysis ERA-Interim (Dee et al., 2011), different tracking and selection algorithms and a slightly different historical analysis period. The main difference in our results is in the peak of concurrence for ECs, which in Eiras-Barca et al. (2018) is at the MDP. The latter authors. One key difference is that they calculated the MDP using a 24-hour time difference, whereas we calculated it with window, while we used a 6-hour difference. Thus, our methodology can provide a more accurate MDP, which may explain the above discrepancy. window. These differences in time intervals, as well as detection and tracking configurations for cyclones and ARs, may account for the minor differences in concurrence rates. Furthermore, our study uses the latest ECMWF reanalysis, ERA5 (Hersbach et al., 2020), whereas their study used the earlier ERA-Interim dataset.~~

Previous studies found an increase in the IVT of ARs under climate change (Payne et al., 2020), as well as an increase in their frequency (Espinoza et al., 2018; Wang et al., 2023; Ramos et al., 2016). These results align with the increase of IVT-max detected for ARs associated with ECs and non-ECs and the increase of concurrences between cyclones and ARs, partly driven by an enhanced AR frequency. Moreover, other studies have found an increase in the frequency and severity of extratropical cyclones under climate change, mainly over the British Isles, and an eastward extension of the storm track activity over Europe (Priestley and Catto, 2022; Zappa et al., 2013; Seiler and Zwiers, 2016a). Our results show a generalized increase in compound events of ECs with ARs in most of the North Atlantic basin. A robust increase in concurrence over the British Isles, Iberia and north France is only observed under the most severe climate change scenario. However, we did not detect a clear upward trend in the individual frequency of ECs or ARs tracks across the entire North Atlantic basin. This apparent contradiction suggests that changes in the characteristics or dynamics of ECs and ARs, rather than their frequency, may be driving the observed increase in concurrence. This is a significant finding that needs further investigation, as the underlying physical mechanisms for this increase remain unclear.

Our analysis has limitations that should be acknowledged. The main constraint is the reduced number of CMIP6 models and members used. The number of models or ensemble members used is limited by the availability of data on multiple vertical levels at the 6-hourly resolution, necessary to compute IVT. We acknowledge that using only one member per model does not facilitate a comprehensive model intercomparison; more members for each model would be needed to adequately assess model uncertainty, specifically the biases of the models relative to reanalysis data. We chose all models and members from CMIP6 where these variables were available for the historical, SSP1-2.6, SSP2-4.5 and SSP5-8.5 experiments. In particular, we deemed it important to go beyond the ~~worst-case high-emission~~ scenario (SSP5-8.5), and also look at the implications of lower warming levels. While most of the results presented indicate a stronger signal for the highest emission scenario (SSP5-8.5), this scenario has been deemed unrealistic (Hausfather and Peters, 2020). Therefore, we emphasize that our results should be interpreted with consideration of various scenarios. A further caveat is that we used a single tracking algorithm, namely the TempestExtreme software (Ullrich et al., 2021). While we have compared our results for the present period with a previous study that uses different tracking algorithms (Eiras-Barca et al., 2018), this does not detract from the fact that our results depend on the detection and tracking method (Neu et al., 2013). Furthermore, future studies should aim to delve deeper into

isolating the dynamic signal from the thermodynamics of the climate change response. Finally, future work should explore future changes in the wet and windy extremes associated with the compound meteorological events investigated here.

Code availability. The scripts are available upon reasonable request.

410 *Data availability.* ERA5 data are available on the C3S Climate Data Store at <https://cds.climate.copernicus.eu/#/home>. CMIP6 data are available on the ESGF Metagrid web application at <https://aims2.llnl.gov/search>.

Appendix A

A1 CMIP6 models information

Table A1. Description of the CMIP6 models and member used for the historical, SSP1-2.6, SSP2-4.5 and SSP5-8.5 scenarios.

Model Name	Member	Resolution	Reference
MPI-ESM1-2-LR	r1i1p1f1	T63 spectral truncation (~200 km): 192 x 96 longitude/latitude; 47 vertical levels (top level 0.01 hPa)	Mauritsen et al. (2019)
MPI-ESM1-2-HR	r1i1p1f1	T127 spectral truncation (~100 km): 384 x 192 longitude/latitude; 95 vertical levels (top level 0.01 hPa)	Müller et al. (2018)
NorESM2-MM	r1i1p1f1	1.25° × 0.9375° regular grid: 288 x 192 longitude/latitude; 32 vertical levels (top level 3.6 hPa)	Seland et al. (2020)
EC-Earth3	r1i1p1f1	T255 spectral truncation (~80km): 512 x 256 longitude/latitude; 91 vertical levels (top level 0.01 hPa)	Döscher et al. (2022)
CMCC-ESM2	r1i1p1f1	Regular grid 0.9° × 1.25°: 288 x 192 longitude/latitude; 30 vertical levels (top level 2 hPa)	Cherchi et al. (2019)
MIROC6	r1i1p1f1	T85 spectral truncation (~160 km): 256 x 128 longitude/latitude; 81 vertical levels (top level 0.004 hPa)	Tatebe et al. (2019)

A2 TempestExtremes Code for Detecting and Tracking Extratropical Cyclones

415 To identify extratropical cyclones, we use the executable *DetectNodes* [from the TempestExtremes tracking algorithm \(Ullrich et al., 2021\)](#), which recognizes candidate "nodes" corresponding to local minima in the SLP field. Subsequently, we employ *StitchNodes* [also from the TempestExtremes tracking algorithm \(Ullrich et al., 2021\)](#) to connect these candidate nodes into tracks. The tracking codes [set-up-set up](#) with the parameters used are as follows:

<pre> DetectNodes -in_data "data_slp" -out "detect_nodes_output" -searchbymin slp -mergedist 6.0 -minlon -80.0 <u>-90.0</u> -maxlon 10.0 <u>20.0</u> -minlat 25.0 <u>15.0</u> -maxlat 65.0 <u>75.0</u> -regional -outputcmd "slp,min,0" </pre>	<pre> StitchNodes -in_data "detect_nodes_output" -out "cyclone_tracks" -in_fmt "lon,lat,PSL" -range 6.0 -mintime 24h -maxgap 6h -min_endpoint_dist 12.0 -out_file_format "csv" </pre>
--	---

420 A3 TempestExtremes Code for Detecting and Tracking ARs

For the detection ~~of ARs and tracking of ARs~~, we use the ~~executable~~ executables from the TempestExtremes tracking algorithm (Ullrich et al., 2021) *DetectBlobs to detect ARs* and to connect ~~candidates ARs~~ or "blobs" we use the executable *StitchBlobs* with the following parameters:

<pre> ./DetectBlobs -in_data "data_IVT" -out "detect_blobs_output" -latname LAT -lonname LON -thresholdcmd "_LAPLACIAN{8,10} (_VECMAG(UQ_FLUX,VQ_FLUX)),<=,-40000,0; _VECMAG(UQ_FLUX,VQ_FLUX),>=,250,0" -geofiltercmd 'area,>=,4e5km2' -minlat 25 -minabslat 15 -minlon -80 -maxlon 10 -maxlat 65 -regional </pre>	<pre> ./StitchBlobs -in "detect_blobs_output" -out "ar_tracks" -latname LAT -lonname LON -var "binary_tag" -mintime 10 -regional </pre>
---	---

425 A4 Number of ~~Cyclones~~ cyclone tracks and ARs tracks detected in ERA5 and CMIP6

Section 3.1 describes how cyclones are detected, tracked and classified as ECs or non-ECs. The result of this process is the track of each cyclone. In the following table we summarize the number of individual ECs and non-ECs tracks for each dataset and the percentual difference from the historical period:

Table A2. Number of EC and non-EC tracks detected in each dataset. In brackets the percentual difference of the future scenario with respect to the historical for each model.

	ECs tracks				non-ECs tracks		
	Historical	SSP1-2.6	SSP2-4.5	SSP5-8.5	Historical	SSP1-2.6	SSP2-4.5
	<u>1980-2009</u>	2070-2099			<u>1980-2009</u>	2070-2099	
ERA5	1591 <u>1372</u>	-	-	-	3100 <u>3210</u>	-	-
MPI-LR	1072 <u>870</u>	1007 <u>822 (-5.5)</u>	958 <u>785 (-9.8)</u>	939 <u>743 (-14.6)</u>	2273 <u>2387</u>	2255 <u>2435 (2.0)</u>	2071 <u>2231 (-6.0)</u>
MPI-HR	1342 <u>1168</u>	1304 <u>1147 (-1.8)</u>	1229 <u>1104 (-5.5)</u>	1186 <u>1042 (-10.8)</u>	2772 <u>3046</u>	2664 <u>2840 (-6.8)</u>	2703 <u>2885 (-5.0)</u>
EC-Earth3	1471 <u>1283</u>	1269 <u>1107 (-13.7)</u>	1282 <u>1082 (-15.7)</u>	1306 <u>1079 (-15.9)</u>	2600 <u>2928</u>	2728 <u>2991 (2.2)</u>	2635 <u>2905 (-0.5)</u>
NorESM2-MM	1427 <u>1193</u>	1323 <u>1115 (-6.5)</u>	1356 <u>1114 (-6.6)</u>	1360 <u>741 (-37.9)</u>	3487 <u>3619</u>	3601 <u>3731 (3.1)</u>	3410 <u>3572 (-1.0)</u>
MIROC6	1038 <u>879</u>	989 <u>841 (-4.3)</u>	984 <u>825 (-6.1)</u>	966 <u>818 (-6.9)</u>	3314 <u>3424</u>	3273 <u>3284 (-4.1)</u>	3252 <u>3360 (-1.0)</u>
CMCC-ESM2	1185 <u>1076</u>	1039 <u>916 (-14.9)</u>	1046 <u>1030 (-4.3)</u>	1130 <u>1013 (-5.9)</u>	3441 <u>3537</u>	3307 <u>3312 (-6.4)</u>	3348 <u>3384 (-4.0)</u>
<u>Total CMIP6</u>	<u>6469</u>	<u>5948 (-8.1)</u>	<u>5940 (-8.2)</u>	<u>5436 (-16.0)</u>	<u>18941</u>	<u>18593 (-1.8)</u>	<u>18337 (-3.2)</u>

430

Section 3.2 describes how ARs are detected and tracked. The result of this process is the track of each AR. In the following table we summarize the number of individual AR tracks for each dataset and the percentual difference from the historical period:

Table A3. Number of AR tracks detected in each dataset. In brackets the percentual difference of the future scenario with respect to the historical for each model.

	ARs tracks			
	Historical	SSP1-2.6	SSP2-4.5	SSP5-8.5
	<u>1980-2009</u>	2070-2099		
ERA5	1224	-	-	-
MPI-LR	1283	1282 <u>(-0.2)</u>	1223 <u>(-4.8)</u>	1246 <u>(-3.0)</u>
MPI-HR	1218	1251 <u>(2.7)</u>	1209 <u>(-0.7)</u>	1198 <u>(-1.6)</u>
EC-Earth3	1186	1213 <u>(2.3)</u>	1222 <u>(3.0)</u>	1249 <u>(5.3)</u>
NorESM2-MM	1234	1261 <u>(2.2)</u>	1270 <u>(2.9)</u>	1247 <u>(1.1)</u>
MIROC6	1151	1234 <u>(7.2)</u>	1200 <u>(4.3)</u>	1221 <u>(6.1)</u>
CMCC-ESM2	1185	1203 <u>(1.5)</u>	1234 <u>(4.1)</u>	1207 <u>(1.9)</u>
<u>Total CMIP6</u>	<u>7259</u>	<u>7444 (2.5)</u>	<u>7358 (1.4)</u>	<u>7368 (1.5)</u>

A5 Cyclone intensity differences between future scenarios and historical simulations




Figure A1. Difference in the multi-model mean SLP between the forcing scenarios and historical runs in solid lines, the [inter-seasonal inter-annual](#) variability of the multi-model ensembles of mean SLP in shades and the [inter-seasonal inter-annual](#) variability for ERA5 in the dashed line, for ECs with ARs (a), non-EC with ARs (b), ECs without ARs (c), and non-ECs without ARs (d).

Author contributions. FLM and MG developed the concept of the paper, performed the data analysis, prepared the figures and wrote the first manuscript draft. All authors contributed with ideas, interpretation of the results, and manuscript revisions.

435 *Competing interests.* Author GM is a member of the editorial board of Earth System Dynamics journal.

Acknowledgements. The authors wish to thank Lionel Guez, Atef Ben Nasser, Leonardo Olivetti, and Anastasiya Shyrokaya for useful discussions and technical support that shaped this article. This project has received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant No. 956396 (EDIPI project). GM further acknowledges the European Union's H2020 research and innovation programme under ERC grant no. 948309 (CENÆ Project). DF and MG further acknowledge the support of the COST Action FutureMed CA22162 supported by COST (European Cooperation in Science and Technology), an INSU-CNRS-LEFE-MANU grant (project CROIRE), the state aid managed by the National Research Agency under France 2030 bearing the references ANR-22-EXTR-0005 (TRACCS-PC4-EXTENDING project), the European Union's Horizon 2020 research and innovation programme under grant agreement No. 101003469 (XAIDA).

References

- 445 Allen, J. T., Pezza, A. B., and Black, M. T.: Explosive cyclogenesis: A global climatology comparing multiple reanalyses, *Journal of Climate*, 23, 6468–6484, 2010.
- Bao, J., Michelson, S., Neiman, P., Ralph, F., and Wilczak, J.: Interpretation of enhanced integrated water vapor bands associated with extratropical cyclones: Their formation and connection to tropical moisture, *Monthly weather review*, 134, 1063–1080, 2006.
- Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., Bellucci, A., et al.: Global
450 mean climate and main patterns of variability in the CMCC-CM2 coupled model, *Journal of Advances in Modeling Earth Systems*, 11, 185–209, 2019.
- Collow, A. B. M., Shields, C. A., Guan, B., Kim, S., Lora, J. M., McClenny, E. E., Nardi, K., Payne, A., Reid, K., Shearer, E. J., Tomé, R., Wille, J. D., Ramos, A. M., Gorodetskaya, I. V., Leung, L. R., O'Brien, T. A., Ralph, F. M., Rutz, J., Ullrich, P. A., and Wehner, M.: An
455 Overview of ARTMIP's Tier 2 Reanalysis Intercomparison: Uncertainty in the Detection of Atmospheric Rivers and Their Associated Precipitation, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036155, <https://doi.org/https://doi.org/10.1029/2021JD036155>, e2021JD036155 2021JD036155, 2022.
- Dacre, H. F., Martínez-Alvarado, O., and Mbengue, C. O.: Linking Atmospheric Rivers and Warm Conveyor Belt Airflows, *Journal of Hydrometeorology*, 20, 1183 – 1196, <https://doi.org/10.1175/JHM-D-18-0175.1>, 2019.
- Davolio, S., Della Fera, S., Laviola, S., Miglietta, M., and Levizzani, V.: Heavy precipitation over Italy from the Mediterranean storm “Vaia”
460 in October 2018: Assessing the role of an atmospheric river, *Monthly Weather Review*, 148, 3571–3588, 2020.
- Davolio, S., Vercellino, M., Miglietta, M. M., Pitura, L. D., Laviola, S., and Levizzani, V.: The influence of an atmospheric river on a heavy precipitation event over the western Alps, *Weather and Climate Extremes*, 39, 100542, 2023.
- De Vries, A. J.: A global climatological perspective on the importance of Rossby wave breaking and intense moisture transport for extreme precipitation events, *Weather and Climate Dynamics Discussions*, 2020, 1–56, 2020.
- 465 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597,
470 <https://doi.org/https://doi.org/10.1002/qj.828>, 2011.
- Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblus-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonnet, F., Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzic, L., van Noije, T., Nolan, P., O'Donnell, D., Ollinaho, P., van den Oord,
475 G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geoscientific Model Development*, 15, 2973–3020, <https://doi.org/10.5194/gmd-15-2973-2022>, 2022.
- Eiras-Barca, J., Ramos, A. M., Pinto, J. G., Trigo, R. M., Liberato, M. L., and Miguez-Macho, G.: The concurrence of atmospheric rivers
480 and explosive cyclogenesis in the North Atlantic and North Pacific basins, *Earth System Dynamics*, 9, 91–102, 2018.

- Eiras-Barca, J., Ramos, A. M., Algarra, I., Vázquez, M., Dominguez, F., Miguez-Macho, G., Nieto, R., Gimeno, L., Taboada, J., and Ralph, F. M.: European West Coast atmospheric rivers: A scale to characterize strength and impacts, *Weather and Climate Extremes*, 31, 100305, 2021.
- 485 Espinoza, V., Waliser, D. E., Guan, B., Lavers, D. A., and Ralph, F. M.: Global analysis of climate change projection effects on atmospheric rivers, *Geophysical Research Letters*, 45, 4299–4308, 2018.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016a.
- 490 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016b.
- Ferreira, J. A., Liberato, M. L., and Ramos, A. M.: On the relationship between atmospheric water vapour transport and extra-tropical cyclones development, *Physics and Chemistry of the Earth, Parts A/B/C*, 94, 56–65, 2016.
- Fink, A. H., Pohle, S., Pinto, J. G., and Knippertz, P.: Diagnosing the influence of diabatic processes on the explosive deepening of extratropical cyclones, *Geophysical Research Letters*, 39, 2012.
- 495 Gao, Y., Lu, J., and Leung, L. R.: Uncertainties in Projecting Future Changes in Atmospheric Rivers and Their Impacts on Heavy Precipitation over Europe, *Journal of Climate*, 29, 6711 – 6726, <https://doi.org/10.1175/JCLI-D-16-0088.1>, 2016.
- Gimeno, L., Nieto, R., Vázquez, M., and Lavers, D. A.: Atmospheric rivers: A mini-review, *Frontiers in Earth Science*, 2, 2, 2014.
- Gimeno, L., Dominguez, F., Nieto, R., Trigo, R., Drumond, A., Reason, C. J., Taschetto, A. S., Ramos, A. M., Kumar, R., and Marengo, J.: 500 Major mechanisms of atmospheric moisture transport and their role in extreme precipitation events, *Annual Review of Environment and Resources*, 41, 117–141, 2016.
- Ginesta, M., Yiou, P., Messori, G., and Faranda, D.: A methodology for attributing severe extratropical cyclones to climate change based on reanalysis data: the case study of storm Alex 2020, *Climate Dynamics*, 61, 229–253, 2023.
- Ginesta, M., Flaounas, E., Yiou, P., and Faranda, D.: Anthropogenic climate change will intensify European explosive storms analogous to 505 Alex, Eunice, and Xynthia, *Journal of Climate*, 37, 5427–5452, 2024.
- Guan, B. and Waliser, D. E.: Detection of atmospheric rivers: Evaluation and application of an algorithm for global studies, *Journal of Geophysical Research: Atmospheres*, 120, 12 514–12 535, 2015.
- Guan, B., Waliser, D. E., and Ralph, F. M.: Global application of the atmospheric river scale, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 180, 2023.
- 510 Guo, Y., Shinoda, T., Guan, B., Waliser, D. E., and Chang, E. K.: Statistical relationship between atmospheric rivers and extratropical cyclones and anticyclones, *Journal of Climate*, 33, 7817–7834, 2020.
- Hausfather, Z. and Peters, G. P.: Emissions—the ‘business as usual’ story is misleading, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- 515 Lavers, D. A. and Villarini, G.: The nexus between atmospheric rivers and extreme precipitation across Europe, *Geophysical Research Letters*, 40, 3259–3264, 2013.
- Lavers, D. A. and Villarini, G.: The contribution of atmospheric rivers to precipitation in Europe and the United States, *Journal of Hydrology*, 522, 382–390, 2015.

- Lavers, D. A., Allan, R. P., Villarini, G., Lloyd-Hughes, B., Brayshaw, D. J., and Wade, A. J.: Future changes in atmospheric rivers and their implications for winter flooding in Britain, *Environmental Research Letters*, 8, 034 010, 2013.
- Lavers, D. A., Ralph, F. M., Waliser, D. E., Gershunov, A., and Dettinger, M. D.: Climate change intensification of horizontal water vapor transport in CMIP5, *Geophysical Research Letters*, 42, 5617–5625, 2015.
- Liberato, M., Pinto, J., Trigo, R., Ludwig, P., Ordóñez, P., Yuen, D., and Trigo, I.: Explosive development of winter storm Xynthia over the subtropical North Atlantic Ocean, *Natural Hazards and Earth System Sciences*, 13, 2239–2251, 2013.
- Ludwig, P., Pinto, J. G., Hoeppe, S. A., Fink, A. H., and Gray, S. L.: Secondary cyclogenesis along an occluded front leading to damaging wind gusts: Windstorm Kyrill, January 2007, *Monthly Weather Review*, 143, 1417–1437, 2015.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., et al.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and its response to increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, 2019.
- Müller, W. A., Jungclaus, J. H., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., et al.: A higher-resolution version of the max planck institute earth system model (MPI-ESM1. 2-HR), *Journal of Advances in Modeling Earth Systems*, 10, 1383–1413, 2018.
- Neu, U., Akperov, M. G., Bellenbaum, N., Benestad, R., Blender, R., Caballero, R., Coccozza, A., Dacre, H. F., Feng, Y., Fraedrich, K., et al.: IMILAST: A community effort to intercompare extratropical cyclone detection and tracking algorithms, *Bulletin of the American Meteorological Society*, 94, 529–547, 2013.
- O'Brien, T. A., Wehner, M. F., Payne, A. E., Shields, C. A., Rutz, J. J., Leung, L.-R., Ralph, F. M., Collow, A., Gorodetskaya, I., Guan, B., et al.: Increases in future AR count and size: Overview of the ARTMIP Tier 2 CMIP5/6 experiment, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036 013, 2022.
- Payne, A. E., Demory, M.-E., Leung, L. R., Ramos, A. M., Shields, C. A., Rutz, J. J., Siler, N., Villarini, G., Hall, A., and Ralph, F. M.: Responses and impacts of atmospheric rivers to climate change, *Nature Reviews Earth & Environment*, 1, 143–157, 2020.
- Pfahl, S. and Sprenger, M.: On the relationship between extratropical cyclone precipitation and intensity, *Geophysical Research Letters*, 43, 1752–1758, 2016.
- Pinto, J. G., Zacharias, S., Fink, A. H., Leckebusch, G. C., and Ulbrich, U.: Factors contributing to the development of extreme North Atlantic cyclones and their relationship with the NAO, *Climate dynamics*, 32, 711–737, 2009.
- Priestley, M. D. K. and Catto, J. L.: Future changes in the extratropical storm tracks and cyclone intensity, wind speed, and structure, *Weather and Climate Dynamics*, 3, 337–360, <https://doi.org/10.5194/wcd-3-337-2022>, 2022.
- Ralph, F. M., Neiman, P. J., and Wick, G. A.: Satellite and CALJET aircraft observations of atmospheric rivers over the eastern North Pacific Ocean during the winter of 1997/98, *Monthly weather review*, 132, 1721–1745, 2004.
- Ralph, F. M., Iacobellis, S., Neiman, P., Cordeira, J., Spackman, J., Waliser, D., Wick, G., White, A., and Fairall, C.: Dropsonde observations of total integrated water vapor transport within North Pacific atmospheric rivers, *Journal of Hydrometeorology*, 18, 2577–2596, 2017.
- Ramos, A. M., Tomé, R., Trigo, R. M., Liberato, M. L., and Pinto, J. G.: Projected changes in atmospheric rivers affecting Europe in CMIP5 models, *Geophysical Research Letters*, 43, 9315–9323, 2016.
- Reale, M., Liberato, M. L., Lionello, P., Pinto, J. G., Salon, S., and Ulbrich, S.: A global climatology of explosive cyclones using a multi-tracking approach, *Tellus A: Dynamic Meteorology and Oceanography*, 71, 1611 340, 2019.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T.,

- Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Stre-
560 fler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont,
Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and
their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168,
<https://doi.org/https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Roebber, P. J.: Statistical analysis and updated climatology of explosive cyclones, *Monthly Weather Review*, 112, 1577–1589, 1984.
- Sanders, F. and Gyakum, J. R.: Synoptic-dynamic climatology of the “bomb”, *Monthly Weather Review*, 108, 1589–1606, 1980.
- Schultz, D. M.: The Presidents’ Day Snowstorm of 18–19 February 1979: Its History and Significance at *Monthly Weather Review*, *Monthly*
565 *Weather Review*, 150, 305 – 307, <https://doi.org/10.1175/MWR-D-22-0012.1>, 2022.
- Seiler, C. and Zwiers, F.: How will climate change affect explosive cyclones in the extratropics of the Northern Hemisphere?, *Climate*
Dynamics, 46, 3633–3644, <https://doi.org/https://doi.org/10.1007/s00382-015-2791-y>, 2016a.
- Seiler, C. and Zwiers, F. W.: How will climate change affect explosive cyclones in the extratropics of the Northern Hemisphere?, *Climate*
Dynamics, 46, 3633–3644, 2016b.
- 570 Seland, Ø., Bentsen, M., Olivié, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A.,
et al.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario
simulations, *Geoscientific Model Development*, 13, 6165–6200, 2020.
- Shaw, T., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C., Hwang, Y.-T., Li, C., O’gorman, P., Rivière, G., Simpson, I., et al.: Storm
track processes and the opposing influences of climate change, *Nature Geoscience*, 9, 656–664, 2016.
- 575 Shepherd, T. G.: Atmospheric circulation as a source of uncertainty in climate change projections, *Nature Geoscience*, 7, 703–708, 2014.
- Shields, C. A., Rutz, J. J., Leung, L.-Y., Ralph, F. M., Wehner, M., Kawzenuk, B., Lora, J. M., McClenny, E., Osborne, T., Payne, A. E.,
et al.: Atmospheric river tracking method intercomparison project (ARTMIP): project goals and experimental design, *Geoscientific Model*
Development, 11, 2455–2474, 2018.
- Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., et al.: Description and
580 basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geoscientific Model Development*, 12,
2727–2765, 2019.
- Thandlam, V., Rutgersson, A., and Sahlee, E.: Spatio-temporal variability of atmospheric rivers and associated atmospheric parameters in
the Euro-Atlantic region, *Theoretical and Applied Climatology*, pp. 1–21, 2022.
- Ullrich, P. A., Zarzycki, C. M., McClenny, E. E., Pinheiro, M. C., Stansfield, A. M., and Reed, K. A.: TempestExtremes v2. 1: A community
585 framework for feature detection, tracking, and analysis in large datasets, *Geoscientific Model Development*, 14, 5023–5048, 2021.
- Waliser, D. and Guan, B.: Extreme winds and precipitation during landfall of atmospheric rivers, *Nature Geoscience*, 10, 179–183, 2017.
- Wang, S., Ma, X., Zhou, S., Wu, L., Wang, H., Tang, Z., Xu, G., Jing, Z., Chen, Z., and Gan, B.: Extreme atmospheric rivers in a warming
climate, *Nature Communications*, 14, 3219, 2023.
- Yettella, V. and Kay, J. E.: How will precipitation change in extratropical cyclones as the planet warms? Insights from a large initial condition
590 climate model ensemble, *Climate Dynamics*, 49, 1765–1781, 2017.
- Zappa, G., Shaffrey, L. C., Hodges, K. I., Sansom, P. G., and Stephenson, D. B.: A Multimodel Assessment of Future Projec-
tions of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models, *Journal of Climate*, 26, 5846–5862,
<https://doi.org/http://dx.doi.org/10.1175/JCLI-D-12-00573.1>, 2013.

- 595 Zhang, L., Zhao, Y., Cheng, T. F., and Lu, M.: Future changes in global atmospheric rivers projected by CMIP6 models, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD039359, 2024.
- Zhang, Z. and Ralph, F. M.: The influence of antecedent atmospheric river conditions on extratropical cyclogenesis, *Monthly Weather Review*, 149, 1337–1357, 2021.
- Zhang, Z., Ralph, F. M., and Zheng, M.: The relationship between extratropical cyclone strength and atmospheric river intensity and position, *Geophysical Research Letters*, 46, 1814–1823, 2019.
- 600 Zhu, Y. and Newell, R. E.: Atmospheric rivers and bombs, *Geophysical Research Letters*, 21, 1999–2002, 1994.
- Zhu, Y. and Newell, R. E.: A proposed algorithm for moisture fluxes from atmospheric rivers, *Monthly weather review*, 126, 725–735, 1998.