

Combined CO₂ measurement record indicates Amazon forest carbon uptake is offset by Savanna carbon release

Santiago Botía¹, Saqr Munassar¹, Thomas Koch¹, Danilo Custodio¹, Luana S. Basso¹, Shujiro Komiya², Jost V. Lavric³, David Walter⁴, Manuel Gloor⁵, Giordane Martins⁶, Stijn Naus¹⁰, Gebrand Koren⁷, Ingrid T. Luijkx⁸, Stijn Hantson¹¹, John B. Miller⁶, Wouter Peters^{8,9}, Christian Rödenbeck¹, and Christoph Gerbig¹

¹Department of Biogeochemical Signals, Max Planck Institute for Biogeochemistry, Jena, Germany

²Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena, Germany

³Acoem GmbH, Hallbergmoos, Germany

⁴Multiphase Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

⁵School of Geography, University of Leeds, Leeds, LS2 9JT, UK

⁶NOAA - Global Monitoring Laboratory, Boulder, Colorado, USA

⁷Copernicus Institute of Sustainable Development, Utrecht University, Utrecht, The Netherlands

⁸Environmental Sciences Group, Dept of Meteorology and Air Quality, Wageningen University and Research, Wageningen, The Netherlands

⁹University of Groningen, Centre for Isotope Research, Groningen, The Netherlands

¹⁰National Institute of Water and Atmospheric Research, Auckland, New Zealand

¹¹Facultad de Ciencias Naturales, Universidad del Rosario, Bogotá, Colombia

Correspondence: Santiago Botía (sbotia@bgc-jena.mpg.de)

Abstract. In tropical South America there has been substantial progress on atmospheric monitoring capacity, but the region still has a limited number of continental atmospheric stations relative to its large area, hindering net carbon flux estimates using atmospheric inversions. In this study, we use dry air CO₂ mole fractions measured at the Amazon Tall Tower Observatory (ATTO) and airborne vertical CO₂ profiles in an atmospheric inversion system to estimate net carbon exchange in tropical South America from 2010 to 2018. We focus on the biogeographic Amazon, and its neighboring 'Cerrado & Caatinga' biomes. Considering all prior ensemble members, we estimate that the biogeographic Amazon was a net carbon sink with the sum of vegetation uptake, river outgassing and carbon release from fires at a median of -0.33 ± 0.33 PgC year⁻¹ (1-sigma posterior uncertainty). Using only process-based models as input in the inversion system the uptake is reduced to -0.24 ± 0.33 PgC year⁻¹. The 'Cerrado & Caatinga' biomes together represent a median carbon source of 0.31 ± 0.24 PgC year⁻¹, with contributions from both vegetation carbon release and fires. Therefore, we estimate a close-to-neutral net carbon exchange for tropical South America, but we note that the uncertainties straddle zero net exchange. In addition, we calculate the effect of systematic uncertainties in the inverse estimates by proposing a water-vapor correction to measured airborne CO₂ profiles. Assuming that the correction brings the observational data closer to the truth, implies that the Amazon is a weaker sink of carbon and that the 'Cerrado & Caatinga' is a larger source. We do not find a strong spatial shift of fluxes within the biogeographic Amazon due to the correction, nor do we find a strong impact on the interannual variations. Finally, to further reduce the uncertainty in

regional carbon balance estimates in tropical South America, we call for an expansion of the atmospheric monitoring network on the continent, mainly in the Amazon-Andes foothills.

1 Introduction

Land ecosystems constitute approximately half of the atmospheric CO₂ sink (Friedlingstein et al., 2022). However, they remain the most uncertain (Ballantyne et al., 2012) and variable (Le Quéré et al., 2018) component of the global carbon cycle. Numerous independent studies confirm that tropical land ecosystems drive most of the interannual variability (IAV) in the net land carbon flux (Keeling and Revelle, 1985; Keeling et al., 1995; Bousquet et al., 2000; Jung et al., 2011; Peylin et al., 2013; Jung et al., 2017; Rödenbeck et al., 2018a; Bastos et al., 2020), which is linked to the atmospheric CO₂ growth rate (Cox et al., 2013; Wang et al., 2013; Piao et al., 2020). Furthermore, tropical ecosystems store substantial carbon reserves in aboveground ground living biomass (Brando et al., 2019), that can be released rapidly further amplifying the CO₂ growth rate. In other words, the accumulation of CO₂ in the atmosphere heavily relies on the carbon uptake and release dynamics occurring in tropical ecosystems. In particular, South America plays a pivotal role in both aspects, uptake and release, as it hosts the Amazon rainforest, which contains 49% of tropical biomass carbon (Saatchi et al., 2011) and encompasses a third of the continent landmass (Goulding et al., 2003).

The emissions of deforestation fires (van der Werf et al., 2010) and degradation (Assis et al., 2020) carbon sources are particularly important for the Amazon region (Aragão et al., 2018; Matricardi et al., 2020; Qin et al., 2021; Kruid et al., 2021; Lapola et al., 2023). In the years after 2017 to 2022, clear-cut deforestation increased significantly in Brazil (Alencar et al., 2019; Silva Junior et al., 2021), not only releasing massive amounts of carbon (Assis et al., 2020) but also exposing larger areas of forest fragments to degradation (Matricardi et al., 2020) and inducing indirect carbon losses by edge effects (Silva Junior et al., 2020). In 2023, while deforestation decreased by 22% relative to 2022, there was a disproportionate rise of wildfires in old-growth forests (Mataveli et al., 2024). Such threats to standing forests pose a risk of gradually releasing the carbon stock of the Amazon, which amounts to 150-200 PgC (Saatchi et al., 2007; Malhi et al., 2009; Marques et al., 2017; Baccini et al., 2017). Despite the attention to deforestation and degradation in the Amazon, vegetation loss has largely been overlooked in the Cerrado biome (da Conceição Bispo et al., 2024), where agricultural expansion is more widespread (Rodrigues et al., 2022). By 2019, 46% of the original land cover in the Cerrado was converted to pastures and crops (MapBiomas, 2020). Fires in the Cerrado occur naturally and are crucial for ecosystem functioning, but agricultural expansion has brought more frequent and intense fires threatening the aboveground biomass and creating the need for near-real-time monitoring (Pletsch et al., 2022).

Top-down atmospheric inversions exploit measured CO₂ gradients of an observational network to constrain the $F_{NetLand}$. The balance between total uptake and release of carbon results in the $F_{NetLand}$, which consists of the Net Biome Exchange (NBE) and the release of carbon from fossil fuel combustion. The NBE is composed of vegetation-related fluxes, Gross Primary Productivity (GPP) and Terrestrial Ecosystem Respiration (TER), disturbance-related emission from biomass burning and degradation (F_{fire}), and river CO₂ outgassing (F_{river}). Estimates of $F_{NetLand}$ for tropical regions (Gurney et al., 2002; Rödenbeck et al., 2003; Peylin et al., 2013; Liu et al., 2017; Gloor et al., 2018; Palmer et al., 2019; Crowell et al., 2019; Peiro

et al., 2022) and specifically for the Amazon region (Molina et al., 2015; Alden et al., 2016; Gloor et al., 2018; Wang et al., 2023) have been conducted previously using global inversions, typically operating under the assumption that sources from fires and fossil fuels are separately constrained or well known. Top-down studies have shed light on how fire influences the net carbon balance in specific years (2010-2011) (Gatti et al., 2014; van der Laan-Luijkx et al., 2015), the spatial differences of NBE in areas within the Amazon for 2010 to 2012 (Alden et al., 2016), the response of the Amazon carbon cycle relative to other tropical regions in the 2015/2016 El Niño event (Liu et al., 2017; Gloor et al., 2018; Crowell et al., 2019), and the main drivers of the NBE anomaly in 2015/2016 in the Amazon using satellite-constrained inverse estimates (Wang et al., 2023). At the time of these studies, the available data allowed only analyses of the response to a drought year and normal/wet years. The longer measurement records of in-situ CO₂ measurements at the Amazon Tall Tower Observatory (ATTO) (Botía et al., 2022) and the vertical CO₂ profiles in the Amazon region (Gatti et al., 2021) now enables the community to further examine interannual and seasonal variations using atmospheric inversions (Koren, 2020; Basso et al., 2023) or column budget techniques (Gatti et al., 2021, 2023).

Previous results, using a column budget technique suggested that the $F_{NetLand}$ of the eastern Amazon was on average a carbon source from 2010 to 2018, mainly explained by fire emissions, but also by a vegetation-related source in the southeastern part of the Amazon forest (Gatti et al., 2021). The authors attribute the carbon source to the combined impact of temperature and precipitation anomalies on vegetation, hindering its carbon uptake capacity. Basso et al. (2023) reported a smaller carbon source compared to Gatti et al. (2021) for the same period (2010-2018), using similar CO₂ observational constraints but using an atmospheric transport inversion and integrating additional CO observational constraints on fire emissions. Both studies agree on the calculation of a net carbon source (positive $F_{NetLand}$) and a small vegetation-related carbon sink after subtracting fires and fossil fuels (or assuming them negligible). In recent years, a potential bias in CO₂ observations collected from the aircraft network using nondried air samples has been recognized (Baier et al., 2020; Gatti et al., 2023; Basso et al., 2023). The presence of water vapor in the collected samples would lead to a loss of CO₂ in condensed water in the pressurized flasks. This can lead to an underestimation of fluxes when using this dataset as a constraint on the amount of CO₂ that the Amazon exchanges, with a possibility for seasonal biases due to the higher water vapor present during the wet season. However, the effect of water vapor in the aircraft samples and how that propagates to optimized fluxes has not been quantified, adding additional uncertainty to the Amazon-wide carbon budget.

Other studies using local forest plot data that is up-scaled to the Amazon found that old-growth forests are a small carbon sink with a decreasing trend over the last 30 years (Brienen et al., 2015; Hubau et al., 2020). Although these studies are broadly consistent with the existence of an Amazon-wide vegetation carbon sink and net biomass growth in old-growth plots, substantial uncertainties persist regarding their magnitudes. This agrees with a recent synthesis by Rosan et al. (2024) who reported an Amazon-wide carbon sink of 342 ± 192 TgC year⁻¹ from vegetation uptake over the period 2010-2018, albeit with considerable inter-annual variability. Although all of these studies have investigated the carbon exchange of the Amazon, they each had different definitions in area, be it the biogeographic or a selection of sub-regions defined by Eva et al. (2005). Moreover, a perspective including the neighboring biomes (i.e., 'Cerrado & Caatinga') is lacking.

In this study, we use the CarboScope Global and Regional inversion system to assimilate the 2010-2018 airborne CO₂ profile record and the continuous and long-term CO₂ record at the Amazon Tall Tower Observatory (ATTO). We build on previous studies using the CarboScope Regional system in Europe (Kountouris et al., 2018b, a; Munassar et al., 2021), to explore its ability to constrain the $F_{NetLand}$ at the continental scale over a larger domain, but with a sparser observational network. The study is structured as follows. First, we aim to quantify where the atmospheric inversion using this set of atmospheric data can provide a constraint based on uncertainty reduction. Second, a sub-continental analysis of the carbon budget, with a strong focus on the biogeographic Amazon, but not limited to it, is performed to shed light on spatial gradients. Last, we present a detailed quantification of how systematic uncertainties in measured mole fractions in the aircraft network affect the estimated fluxes in an atmospheric inversion. With this study, we provide a broad perspective on carbon exchange in tropical South America, going beyond the Amazon biome and highlighting where we need to expand our observational efforts to reduce the uncertainty in carbon exchange estimates in the region.

2 Methods and Data

2.1 CarboScope Regional Inversion System

2.1.1 Two-step scheme description

The version of the CarboScope Global inversion system used here is described in detail by Rödenbeck et al. (2018b). To refine the resolution of fluxes and atmospheric transport within our study area, we use the two-step scheme described in several publications previously (Rödenbeck et al., 2009; Trusilova et al., 2010; Kountouris et al., 2018b; Munassar et al., 2021). Our inversion set-up follows largely that of Kountouris et al. (2018a), but we use an isotropic exponential decay for the spatial error structure, mainly because in our domain, unlike in mid-latitudes, the climatic gradients are similar in both latitude and longitude.

In the two-step scheme (see Figure 1), two atmospheric transport models with different spatial resolutions are used. In step 1, a global inversion is performed using the CarboScope Global inversion system (Rödenbeck et al., 2018b) to obtain an optimized NBE flux field having Ocean and Fossil Fuel fluxes prescribed. This global inversion is performed on a coarse global scale using the TM3 atmospheric transport model (Heimann and Körner, 2003) at 4 x 5 deg resolution driven by the NCEP reanalysis meteorological fields (Kalnay et al., 1996). Using that optimized NBE flux field and the same atmospheric transport set-up, simulated mole fractions increments for all sites are obtained, except the site left for validation (i.e. the s10 station set plus the South American stations, see Sect 2.1.4). These "forward" runs represent an intermediate step and are done twice, see equation 1 and Figure 1 (adapted from Rödenbeck et al. (2009)). The first one is performed using TM3 at the coarse global resolution and for the entire time-period of the global inversion. The second forward run is performed using NBE fluxes at coarse spatial resolution, only for the regional domain (i.e. with zeros outside the regional domain) tropical South America and the desired period of interest (i.e., 2010-2018). Both forward runs result in simulated mole fraction increments (Δc_{mod1} and Δc_{mod2}) and their difference corresponds to the far-field contribution from fluxes outside of the regional domain. An initial

condition (c_{ini}) that corresponds to a well-mixed atmosphere with a given initial tracer mole fraction is then added to the far-field contribution. All this together is subtracted from the measured mole fractions at the sites within the domain of interest. This difference represents a "remaining mole fraction" (Δc_{remain}), corresponding to signals from fluxes within the regional domain, as defined by Rödenbeck et al. (2009).

$$\Delta c_{remain} = c_{meas} - (\Delta c_{mod1} - \Delta c_{mod2} + c_{ini}) \quad (1)$$

In step 2, the Δc_{remain} is assimilated in a regional inversion at high-resolution ($0.25 \times 0.25^\circ$) using the model STILT (Lin et al., 2003). STILT is driven by the ECMWF Integrated Forecasting System (IFS) (following the contemporary IFS cycle development; for more information, see <https://www.ecmwf.int/en/publications/ifs-documentation>). At each measurement location and time (x, y, z, t) in the regional domain, we released an ensemble of 100 particles back in time (10 days) to calculate the surface influence on the observations. The surface influence, with units of $ppm/(\mu mol m^{-2} s^{-1})$, provides the link between measured mole fractions and the prior surface fluxes. A set of different prior fluxes (representing a prior ensemble) is used (see Section 2.1.6), and the regional inversion is performed for each prior ensemble member individually. The domain in this study extends from $28.875^\circ S$ to $13.875^\circ N$ and from $83.875^\circ W$ to $34.125^\circ W$, see Figure 2. We have limited our domain to $28.875^\circ S$ due to the lack of observational records further south.

2.1.2 Definition of CO₂ flux components

The total CO₂ exchange with the atmosphere is denoted as Net Land Flux ($F_{NetLand}$), which consists of Net Biome Exchange (NBE) and fossil fuel CO₂ emissions (F_{ff}),

$$F_{NetLand} = NBE + F_{ff} \quad (2)$$

NBE, in turn, is composed of Net Ecosystem Exchange (NEE), the carbon exchange between atmosphere and rivers (F_{river}), and emissions from fires (F_{fire}),

$$NBE = NEE + F_{river} + F_{fire} \quad (3)$$

For all fluxes we adopt the atmospheric sign convention in which positive (+) denotes a source to the atmosphere and negative (-) denotes a sink.

The atmospheric signals reflect the total CO₂ flux, $F_{NetLand}$. However, as F_{ff} is prescribed as a prior the inversion optimizes NBE. In the figures in which the posterior $F_{NetLand}$ is reported, we have added the F_{ff} in post-processing. When subtracting F_{fire} from the NBE we obtain a flux component composed of NEE and F_{river} , reported in Section 3.2.




Figure 1. CarboScope Regional Inversion Two-step scheme (Rödenbeck et al., 2009; Trusilova et al., 2010) flow diagram showing the inputs (purple polygons) to the processes (blue squares) and their specific output (green circles).

2.1.3 Analysis regions in postprocessing

To analyze the spatial distribution of the estimated posterior fluxes and the coverage of the atmospheric network, we have used the definition of sub-regions shown in Figure 2a. One of the criteria for the choice of these regional areas is that they should be independent of our observational network and associated with a biogeographic gradient. Thus, the selection of these areas follows, to some extent, a biome-like distribution, but it does not represent individual biomes strictly. The division within the biogeographic Amazon serves to provide individual sub-regions dividing east-west but also north-south. In addition, we have also kept a separate sub-region for the main branch of the Amazon river as we are interested in the atmospheric constraint over these areas. The biogeographic limit of the Amazon we use here is the same as defined by Albert et al. (2021) as Amazon biome. When reference to the Amazon *sensu-stricto*+, the area corresponds to four subregions defined in Eva et al. (2005): Amazon *sensu stricto*, Andes, Guiana and Gurupi. The Amazon *sensu-stricto*+ area is only used for comparative reasons in the

Discussion. As we are using a regional inversion with 0.25×0.25 degree of spatial resolution with a correlation length of about 200 km, the posterior budgets as well as posterior uncertainties for these subregions within the Amazon can be individually quantified.

2.1.4 Observational network

155 The location of the measurement sites is shown in Figure 2b, together with their aggregated annual mean surface influence (sensitivities of atmospheric concentrations to surface fluxes, in the following called "footprint") from 2010 to 2018. The coverage of the observational network in our regional domain concentrates on the areas within the Amazon but also in the northeast of Brazil. For the seasonal footprints for each site see Figure A1.

In the global inversion (step 1) we have used the set of stations in the 2022 release version of the CarboScope global
 160 system with nearly continuous coverage from 2010 onward (i.e., s10oc_v2022, see http://www.bgc-jena.mpg.de/CarboScope/?ID=s10oc_v2022, for details of stations and data providers). To this default 2022 station set, we added the ATTO CO₂ record (available at: <https://attodata.org/>), five sites within the Amazon region where airborne profiles (available at: <https://doi.org/10.1594/PANGAEA.926834>) are collected (Gatti et al., 2021), and the weekly flask sampling record in Natal (NAT), a station in the northeast of Brazil (Dlugokencky et al., 2021). While the global inversion (s10, step 1) is augmented with all of
 165 the above sites (s10 + ATTO + NAT + aircraft), only the sites within our domain are used for the regional inversion (step 2). The monthly time series at each site is shown in Figure 2c, indicating the data gaps and the evolution of CO₂ over the last decade. Note that ATTO has provided continuous data since 2013 from an 80-m tower, ≈ 50 m above the canopy, and there were major gaps in the aircraft network during 2015 and 2016. For the continuous data (ATTO), we use only daytime measurements (i.e., from 13:00 to 17:00 local time), to ensure we have measurements representative for the well-mixed boundary layer when the
 170 transport model errors are smallest (see Fig S4, Supplementary material in Botía et al. (2022)). The average number of aircraft profiles per month is two, see Figure A2. For each measurement in an aircraft profile (full profile goes up to ≈ 4500 m a.s.l.), for the weekly flask measurements at NAT, and for every single data point at ATTO, we have simulated the surface influence using the STILT model. Therefore, each measurement has an individual footprint linking the observations with surface fluxes in the regional inversion. The STILT set-up follows that of Botía et al. (2022), but the spatial resolution used here is 0.25×0.25 deg. For validating the estimated posterior fluxes we use airborne data from the Manaus site (MAN) (Figure 2), which was not assimilated and is left as an independent site. The information gained from using the South American stations (Aircraft profile + ATTO) is tested using CarboScope global with (s10sam) and without (s10) these stations.

The model-data mismatch uncertainty (including the representation error of the measurements within the transport model) for the three types of sites (in-situ tower, aircraft, and weekly flasks) is chosen to be 1.5 ppm for weekly time scales, following
 180 common practice in CarboScope global (Rödenbeck, 2005; Rödenbeck et al., 2018a) which assimilates a large set of weekly flask samples. To assimilate multiple data streams, we apply a data density weighting Rödenbeck (2005): For the hourly ATTO data, the error will be inflated by $\sqrt{N_{hours/week}}$ (details see Kountouris et al. (2018a)), while for aircraft profiles (composed of several flasks) the error is scaled with $\sqrt{N_{flasks/profile}}$. The data-density weighting practically ensures that one week of

hourly ATTO observations, one aircraft profile, or one weekly flask sample have the same weight in the inversion, reflecting
185 the assumption that they provide the same amount of information due to roughly weekly error correlations.




Figure 2. South American domain (from 13.875°N to 28.875°S and 83.875°W to 34.125°W) with a sub regional division adapted from EPA (2011) (a). The percentages in the legend (adding to 67%) denote the area of the region relative to the total land area in the domain, the remaining 33% corresponds to the white areas. The annual mean footprint, associated with the period where data are available is shown in b. Note that MAN is not assimilated and is used as a validation site. The time series showing the monthly mean CO_2 mole fraction at each station or aircraft profile site is shown in c. The gaps are shown in grey and are due to logistical problems and/or instrument malfunction. The Amazon contour corresponds to the biogeographic limit of the Amazon.

2.1.5 Addressing systematic uncertainties in vertical profiles

Gatti et al. (2023) (Methods Section) reported a source of uncertainty in the aircraft profiles given by moisture in undried flask samples at time of collection. Such moisture can lead to biases in the measured CO₂ mole fractions. The bias manifests itself as an underestimate of the true mole fraction of atmospheric CO₂ in the measurements, due to the removal of gaseous CO₂ into water between the time of sampling at the aircraft inlet and analysis in the lab. Both liquid water and condensation of water vapor after pressurizing the flasks form the source for this CO₂ absorption and thus a relation with water vapor mole fraction exists (hereafter xH₂O), as described specifically for this sampling method by Baier et al. (2020) and Paul et al. (2020). Moreover, systematic errors in assimilated data can bias inverse results in various ways (Masarie et al., 2011). To quantify the effect of such systematic uncertainty on the estimated fluxes, we apply a water-vapor-dependent bias correction to the data, before it is assimilated.

To derive the water vapor bias correction, we used CO₂ vertical profiles collected approximately 80 km northeast of the city of Manaus in the central Amazon basin (site code MAN; Miller et al. (2023)) aboard a Cessna Grand Caravan. Collocated portable flask packages (PFPs; which have a water-related CO₂ bias) and in-situ, on-board measurements (Picarro Inc. CRDS analyzer model G-2401m) were performed (see Figure A4). The in-situ analyzer also provided measurements of CH₄, CO, and H₂O roughly every 3 seconds. The MAN time series of vertical profiles, extending from approximately 150 m above ground level (agl) to approximately 5 km above sea level (asl) began in April 2017 and continues to this day at a frequency of approximately twice per month. The in-situ gas stream was undried (but CO₂ mole fractions have been converted from wet to dry using the simultaneously measured water vapor), and calibrations occurred on the ground. Previous results have shown this kind of implementation has minimal ($\approx < 0.1$ ppm) impact on CO₂ values (Rella et al., 2013). Critically, the fact that the air stream was undried allows us to use the analyzer's water vapor measurement. The PFPs used were version 2 models and CO₂ measurements on PFP air samples were performed at INPE LaGEE; the sampling and analysis methodologies have been described previously (Gatti et al., 2014).

Between April 2017 and June 2019, when air sample collection via PFP stopped at MAN, there were 12 vertical profiles available that had collocated PFP discrete samples and in-situ measurements that could be compared. These 12 profiles span a range of water vapor mole fraction from 0.2 to 3.5 % that decrease with altitude (see Figure A5) and allow us to define a relationship between CO₂ bias ($\Delta\text{CO}_2 = \text{CO}_{2_{in-situ}} - \text{CO}_{2_{PFP}}$) vs. ambient water vapor. Matching between the independent PFP and on-board analyzer systems could be performed accurately by matching each system's GPS time and position signals. Because the PFP sample is collected over $\approx 15 - 40$ s (depending on altitude), we tested a wide range of time averaging windows for the in-situ analyzer, ranging from 10 s to 40 s to best match the single value of CO₂ measured in the laboratory from the PFP air sample. Fortunately, this averaging window range had a negligible impact on the final relationship between ΔCO_2 and water vapor (see Figure A3); the results presented here use a window of 30 s. We also filtered Picarro data for high variability by excluding any 30 s period where the CO₂ standard deviation was greater than 0.5 ppm. Finally, we binned all ΔCO_2 and xH₂O values in 10,000 ppm (or 1%) increments to define a bias relationship. This relationship (Figure A3) shows little bias between 0 and 1% (above 3 km agl), but an increasing, approximately linear bias with increasing xH₂O (below 2 km agl).

220 We used the STILT model to extract the water vapor mole fraction from the ECMWF-IFS short-term forecasts (as also used to drive STILT) at the same measurement and sampling locations in the Manaus flights in which PFPs and the CRDS analyzer were collocated. These flights span altitudes from 280 m agl to 5200 m agl and vertical profiles are taken at $\approx 17:00$ UTC. The correlation between the measured water vapor (CRDS analyzer) and that extracted from ECMWF-IFS was 0.94 ($p < 0.01$, $N=158$) (see Figure A5), with a mean bias of -0.2% (ECMWF-IFS biased low relative to the continuous measurements). The observational record at MAN includes both dry and wet seasons, and in combination with a large altitude range covers the range of water vapor variability at the other vertical profile sites (see Figure A6). The latter gives us confidence in ECMWF-IFS as a proxy for the water vapor mole fraction at the time and location (x, y, z) of each flask sample at ALF, RBA, TEF, TAB, SAN. Therefore, the difference between the dry air CO₂ mole fraction from the continuous measurements (using the wet-dry correction as proposed by Rella et al. (2013) and the PFP on the Manaus flights was fitted to the water vapor mole fraction from ECMWF-IFS. The linear fit is $y = 0.594x - 0.168$, where y is the resulting CO₂ increase [ppm] to be applied to the PFP data at a given water vapor mole fraction x [%], the slope (0.594) was significantly ($p < 0.01$, $N=158$) different from zero (See Figure A7). The standard error of the slope and the intercept is ± 0.11 and ± 0.20 , respectively. Note that the negative values in Figure A7 are due to the variable nature of the atmosphere causing uncertainty in both measurement strategies. We discard the possibility of being a mismatch in different air parcels sampled because of the collocated sampling lines and the tests we have done with different averaging times with the Picarro data (Figure A3). Note that these negative values are more frequent at low water vapor concentrations, which occur more often in the free troposphere (Figures A4 and A5). We have further quantified the expected error in the PFP mole fractions by the calculating the $\sqrt{\frac{SSR}{N-2}}$. Introducing the correction we obtain 0.93 ppm and without the correction 1.30 ppm. The sum of squared residuals (SSR) with the correction is obtained using the linear fit reported previously, and N is the number of measurement points ($N=158$). In the case without the correction the SSR is calculated by obtaining the residuals with respect to a line with no slope and intercept at zero in Figure A7. As introducing the correction leads to a lower $\sqrt{\frac{SSR}{N-2}}$ than without it, it indicates that by applying the water vapor correction we are decreasing the measurement error.

Using this function and the ECMWF-IFS-water-vapor at ALF, RBA, TEF, TAB, SAN we applied the water-vapor correction to each individual PFP sample (see Figure A8). Note that the uncertainty added by the accuracy of the water vapor mole fraction given by the ECMWF-IFS forecast is dealt with using the fit of the ΔCO_2 to the ECMWF-IFS water vapor. At RBA there were 6 flights (dates: 16/02/2018, 09/03/2018, 06/04/2018, 08/05/2018, 15/07/2018, 19/09/2018) that had a drier installed, so for these flights we did not apply the correction. With this, we re-ran the inversion system (Global and Regional) assimilating the data including the water vapor correction, and quantified the potential effect of such systematic uncertainty on the posterior fluxes (hereafter s10samwvc, for CarboScope global). We note that this approach was implemented to diagnostically quantify the effect of water vapor in the PFPs and how that propagates to the estimated fluxes in an inversion system. The effects of water vapor on not-dried flask samples has been established and documented previously (Baier et al., 2020; Paul et al., 2020). Here, we establish the offsets on this specific set of flask samples collected over the Amazon. The offsets used in this study are provided as a public dataset (<https://edmond.mpg.de/privateurl.xhtml?token=a6fba176-8a6b-4b59-a371-3acc804adaf9>), such that the community can use them in their inversion systems and compare their magnitude to other correction methods.

We present an ensemble of inversions based on a set of prior fluxes that differ in several aspects (see Table 1). The first aspect is their conceptual nature, having (a) the Vegetation Photosynthesis and Respiration Model (VPRM) (Mahadevan et al., 2008), a simple diagnostic model using MODIS imagery and fitted to eddy covariance data (Saleska et al. (2013), see Table A2 for the site descriptions) within the domain which provides NEE; (b) the FLUXCOM product (Bodesheim et al., 2018) and its latest
 260 version X-BASE_{NEE} (Nelson et al., 2024), which up-scales site-level eddy covariance data to the globe using a random forest regression; and (c) two process-based models, the SiB4 (Haynes et al., 2019) and SiBCASA (Schaefer et al., 2008) models, both having served as biospheric flux priors in earlier published studies focusing on the Amazon (van der Laan-Luijkx et al., 2015; van Schaik et al., 2018). By having this set of priors we have a wide representation of the potential spatio-temporal dynamics of NEE over our domain. The other variable aspect of this prior selection is their flux magnitudes and seasonal patterns (see
 265 Figure A9 and A11). Originally the eddy-covariance-based products, like the two FLUXCOM versions and VPRM, have a large sink magnitude for the Amazon. Note that the total land flux in the Amazon is highly uncertain, spanning from -0.34 to 0.29 PgC year⁻¹ (Gatti et al., 2021; Rosan et al., 2024), but this range gets larger than 1 PgC considering the uncertainties associated with each estimate, thus we decided to keep the eddy-flux based prior NEE products (VPRM and FLUXCOM), as they can be considered as a plausible first guess in an inversion. Furthermore and regardless of how they compare to current
 270 independent estimates we proceeded to make an experiment scaling two of our priors (i.e. VPRM and SiB4) such that NEE = 0.5 and 1 PgC, and thus we can test an opposing (in sign) prior scenario. To achieve this, we scaled ecosystem respiration in VPRM and SiB4 such that the total NEE integral for the biogeographic Amazon equals 0.5 PgC year⁻¹ and 1 PgC year⁻¹ (namely VPRM-0.5Pg, VPRM-1Pg, SiB4-0.5Pg, and SiB4-1Pg). An example for VPRM-0.5Pg is shown in Figure A9. Two additional sensitivity tests were performed using the original VPRM. In one, we removed the long-term mean, seasonality, and
 275 interannual variability (IAV) from VPRM (called VPRM_{flat}) and run the inversion only with a diurnal cycle in the prior. In the second one, we used VPRM as prior but left the ATTO data out from the assimilated station set (called VPRMnoATT).

Note that the CarboScope global v2022 has no diurnal cycle in the prior fluxes. The effect of this and how it propagates to the regional inversion is discussed in Munassar et al. (2024). Here, we apply a region-specific correction based on this work. The correction is based on the response of CarboScope global to having a diurnal cycle in the prior fluxes, which was
 280 derived by inverting the diurnal anomalies (hourly - dailyMean) of two forward runs (one with and the other without diurnal cycle) using CarboScope Global. The prior flux diurnal cycle was based on FLUXCOM. The posterior fluxes of such inversion correspond to the per-grid-cell correction that should be added to the posterior fluxes of the normal inversion, assuming that the FLUXCOM diurnal cycles are correct. Then, we propagate such a correction using the two-step scheme described above via the lateral boundary conditions.

285 The fire emissions used in this study (GFAS-opt CO₂) are based on the original GFAS product (Kaiser et al., 2012), with an adjustment over the Northern part of South America based on CO inversions by Naus et al. (2022). These inversions were performed using the TM5-4DVAR system (Krol et al., 2005; Meirink et al., 2008) using CO data from MOPITT (Deeter et al., 2019) inside the study area (Northern part of South America) and NOAA station data outside of this region. We calculated

Table 1. Inversion runs indicating the type of inversion (Global or regional), the selection of prior NEE fluxes used in the regional inversion and the label identifying each individual run. All these individual runs except VPRMnoATT, represent the prior ensemble. IAV stands for interannual variability and R_{eco} for ecosystem respiration. The s10 station set has nearly continuous coverage from 2010 onward.

Identifier	Station Set	Type	Comment
s10	s10	Global Inv.	
s10sam	s10+Aircraft+NAT+ATTO	Global Inv.	
VPRM	s10+Aircraft+NAT+ATTO	Regional Inv.	
VPRM _{flat}	s10+Aircraft+NAT+ATTO	Regional Inv.	only with diurnal cycle
VPRM-0.5Pg	s10+Aircraft+NAT+ATTO	Regional Inv.	R_{eco} scaled to get 0.5 PgC NEE
VPRM-1Pg	s10+Aircraft+NAT+ATTO	Regional Inv.	R_{eco} scaled to get 1 PgC NEE
VPRMnoATT	s10+Aircraft+NAT	Regional Inv.	
FLUXCOM	s10+Aircraft+NAT+ATTO	Regional Inv.	
X-BASE _{NEE}	s10+Aircraft+NAT+ATTO	Regional Inv.	
SiBCASA	s10+Aircraft+NAT+ATTO	Regional Inv.	
SiB4	s10+Aircraft+NAT+ATTO	Regional Inv.	
SIB4-0.5Pg	s10+Aircraft+NAT+ATTO	Regional Inv.	R_{eco} scaled to get 0.5 PgC NEE
SIB4-1Pg	s10+Aircraft+NAT+ATTO	Regional Inv.	R_{eco} scaled to get 1 PgC NEE

the increment ratios from prior (GFAS) to posterior CO fluxes for each grid cell and applied this factor to the GFAS CO₂ fire emissions as previously done in Koren (2020). For regions outside the domain of Naus et al. (2022) we did not scale the GFAS emissions. Thus, our approach assumes that the adjustment of the MOPITT-Inversion in CO is also applicable to CO₂. We acknowledge that this is an approximation, as the emission ratio between CO and CO₂ could also be off in GFAS. However, here we assume they are constant and interpret the underestimation in CO as an underestimation in fire emissions. This is in line with recent studies of undetected African fire emissions (Ramo et al., 2021).

Furthermore, a non-optimized set of fluxes is used to account for important CO₂ sources that contribute to the integrated signal of CO₂ in the atmosphere within our domain. Ocean fluxes are based on surface-ocean pCO₂ data (Rödenbeck et al., 2013) but specifically processed at higher (1x1 deg) resolution (Run ID: oc_1x1_v2022). Following Steinbach et al. (2011), the EDGAR 4.3 inventory, sector and fuel-type specific and scaled at the national level for each year based on the British Petroleum statistical review (BP Annual report 2020), is used to account for emissions related to the burning of fossil fuels.

The assumed regional prior uncertainty for the domain-wide and annually integrated flux is chosen to be 0.9 PgC year⁻¹, which is based on the contribution of the regional domain to the assumed prior uncertainty (2.8 PgC) in the CarboScope global inversion system. Therefore, when spatially aggregating the spatially and temporally correlated prior error, regardless of the correlation length scale, it scales to the assumed prior uncertainty for the domain.

2.2 Statistical metrics

To report our results, we have adopted the following metrics. The uncertainty associated with the posterior covariance matrix in the inversion system, what is often referred to as "Bayesian" uncertainty, is used in the context of reporting the ensemble median for particular regions. The posterior flux uncertainty can be calculated from the prior uncertainties given by the prior flux and the measurement covariance matrices as described by Rödenbeck (2005). The posterior uncertainty is calculated for each year of interest (2010-2018) and the regions in Figure 2a as the square root of the covariance matrix multiplied by a regional operator. This uncertainty primarily depends on the observation availability and the assumed uncertainties for model-data-mismatch and prior fluxes, which is independent of the biosphere or diagnostic model used as a prior. The uncertainty reduction (UR, $\Delta\sigma$) is then calculated with equation 4,

$$\Delta\sigma = \frac{\sigma_{prior} - \sigma_{post}}{\sigma_{prior}} \quad (4)$$

For the complete prior and posterior ensemble in CarboScope regional an ensemble mean is calculated for each single year (Equation 5). Then the ensemble IAV is calculated as the standard deviation of the ensemble mean across all years (Equation 6). The equation 6 refer to $F_{NetLand}$, but one can replace it with a different flux component, like NBE, to obtain the same quantities.

$$F_{NetLand_{ens.mean_k}} = \left(\frac{1}{n} \sum_{i=1}^n EnsMember_i \right)_{j,j=2010-2018,k=Region} \quad (5)$$

$$F_{NetLand_{IAV_k}} = std(F_{NetLand_{ens.mean_k}}) \quad (6)$$

3 Results

3.1 Understanding atmospheric and prior constraints

We found that the eastern part of the South American domain has a better observational constraint compared to the west. We obtained a mean (averaged over 2010-2018) Uncertainty Reduction (UR) of 44% for the Amazon region and for all the regions within it a reduction equal to or above 18% (Figure 3). Despite the uncertainty reduction of 44%, the absolute uncertainty of the posterior is $0.33 \text{ PgC year}^{-1}$. For the 'Amazon River Flat Plains' and the 'Brazilian Shield Moist Forests' the mean reduction is 53% and 54%, followed by the 'Guianan Shield Moist Forests' with 25% and the 'Amazon and Andes Piedmont' with the lowest mean reduction of 18% within the Amazon region. Note that the 'Cerrado & Caatinga' has a mean UR of 30%, higher than the 'Amazon and Andes Piedmont', indicating that on average there is better observational coverage over this eastern part of the domain.




Figure 3. Areas for spatial integration of fluxes (a) and prior / posterior uncertainty for each of these areas (b). The percentages represent the mean uncertainty reduction over the period between 2010 and 2018, the values in brackets indicate the min-max range. For a complete time series for each region see Figure A12. Region 6 (biogeographic Amazon) is the sum of 1, 2, 4 and 5.

330 The UR not only varies spatially but also from year to year, depending on the continuity of the measurement records. As expected, years with data gaps have a low uncertainty reduction (see Figure A12). For example, due to the gaps in 2015 at the sites of TEF, SAN, and partly ALF, the uncertainty reduction was affected largely in the 'Brazilian Shield Moist Forests' (decreasing by 21% from 2014 to 2015), the 'Amazon River Flat Plains' (17%) and the 'Cerrado & Caatinga' (18%). This impact on UR was also observed in the biogeographic Amazon with an effect of 10% (Figure A12). Furthermore, in the

335 'Amazon and Andes Piedmont' we observe a slight decrease in the UR throughout the years. The highest UR is observed from 2010 to 2012, when the TAB site was active; after that, the UR never gets back to the 2010-2012 values. The latter highlights the effect of the location and the continuity of the measurement record on the UR for a particular region, and specifically the low information content in the west of the domain.

The largest UR caused by adding the ATTO data is for the flux estimate from 'Guianan Shield Moist Forest'. Sensitivity

340 tests (not shown) excluding ATTO from the assimilated data show that on average the UR is 7% lower in this region, but some years reaching 12% (i.e. 2016). The constraint added by ATTO is smaller but also relevant in the 'Amazon River Flat Plains', increasing the UR by 6% also in 2016. At the biogeographic Amazon scale, the mean impact on the UR is small (2%), but in individual years it can amount to more than 5%. These changes in UR are somewhat conservative as in this study we have

345 of observations per week or per vertical profile, as described in Section 2.1.4. It is important to mention that the mean bias error of simulated mole fractions at each site assimilated, shows a better agreement between ensemble members (individual priors) relative to the simulations using prior fluxes, and the magnitude is reduced considerably from prior to posterior (see

Figure A13). For the MAN site, used as validation (i.e. not assimilated), we observed a reduction in the mean bias from -0.4 ppm to 0.1 ppm.




Figure 4. Relationship between prior and posterior mean net land flux ($F_{NetLand}$). On the y and x axes of each panel the density distribution is shown.

350 For the biogeographic Amazon and the 'Cerrado & Caatinga' we find a strong linear dependence of the posterior estimates on the prior (Figure 4a,b). Even though the spread in the marginal distribution is reduced largely from prior to posterior, the models with a large uptake in the prior (e.g. VPRM, FLUXCOM, X-BASE_{NEE}) do not converge with the main cluster of posterior estimates. We further evaluated such dependence with the VPRM_{flat} experiment, confirming that after removing the long-term mean, the VPRM_{flat} posterior $F_{NetLand}$ falls closer to the main group of estimates in both regions. Interestingly,

355 the regions in the eastern part of the Amazon ('Amazon River Flat Plains', 'Brazilian Shield Moist Forests', and the 'Guianan Shield Moist Forests') exhibit superior constraint by atmospheric data, as illustrated in Figure 4c-f. The spread in the posterior

marginal distribution and the slopes of the linear regression in these four regions are inversely proportional to their respective reduction in uncertainty (Figure 3). This inverse relationship indicates that the posterior estimates are more effectively adjusted, irrespective of the prior magnitude, in regions with a higher reduction in uncertainty. Therefore, the 'Amazon and Andes
 360 Piedmont' in the west stands out as an area where a bias in the prior fluxes would exert a more substantial impact on posterior estimates.

3.2 Carbon balance for tropical South America

The atmospheric inversion allocates a net carbon source ($F_{NetLand}$) to the 'Cerrado & Caatinga' region (except for VPRM, VPRM-0.5Pg and VPRM-1Pg) and consistently identifies a net carbon sink in the biogeographic Amazon (Figure 5 and see
 365 Figures A9 and A10 for the spatial patterns). Interestingly, the addition of South American stations amplify this pattern and reduces the posterior uncertainty (compare s10 with s10sam). Despite the considerable variability in magnitude for the biogeographic Amazon, the atmospheric constraint tends to adjust priors with a positive sign, shifting them towards a smaller source (e.g., SIB4-1Pg) or even turning them into a sink with a negative $F_{NetLand}$ (e.g., VPRM-1Pg, VPRM-0.5Pg, and SIB4-0.5Pg). Furthermore, in the global inversion when adding South American data, the resulting fluxes closely follow the sign
 370 and magnitude of those obtained in the regional inversion. Therefore, we contend that information suggesting a sink-source gradient between the Amazon and the 'Cerrado & Caatinga' is embedded in the atmospheric measurement record, even with the inherent limitations in adjusting certain individual priors.




Figure 5. Carbon balance for the biogeographic Amazon (a) and the Cerrado & Caatinga (b) regions. The posterior flux components shown with a vertical bar ($NEE + F_{river}$), result from subtracting the F_{fire} and F_{ff} from the posterior $F_{NetLand}$ (shown with the markers).

The estimates obtained using VPRM, VPRM-1Pg, and VPRM-0.5Pg for the 'Cerrado & Caatinga' undergo substantial adjustments, with some cases exceeding +1 PgC year⁻¹ (VPRM). However, this diagnostic model (VPRM) exhibits a bias
 375 towards a large sink in the prior (<-1 PgC year⁻¹) that even after the inversion, the posterior estimate is far from the other

ensemble members and beyond the posterior uncertainty. Similarly, in the biogeographic Amazon a comparable pattern is observed for VPRM, FLUXCOM, and X-BASE_{NEE}. Nevertheless, the removal of the long-term mean of the prior flux, as implemented in VPRM_{flat}, results in a $F_{NetLand}$ that aligns more closely with the main cluster of posterior estimates at ≈ -0.2 PgC year⁻¹ for the Biogeographic Amazon and at ≈ 0.3 PgC year⁻¹ for the 'Cerrado & Caatinga'. In other words, a prior flux
 380 having zero-mean, but a diurnal cycle would be closer to the main cluster of posterior estimates. To provide further insights on the main drivers of $F_{NetLand}$ and the spatial gradient reported here, we explore the individual components of the $F_{NetLand}$.

The mean fire emission estimates for the biogeographic Amazon and the 'Cerrado & Caatinga' are 0.10 PgC year⁻¹ and 0.19 PgC year⁻¹, respectively (differences between marker and vertical bar in Figure 5 and shown in Figure 6). Our findings indicate that, for most estimates, the carbon source in the 'Cerrado & Caatinga' does not solely originate from fires (Figure 5b). Within
 385 the cluster that exhibits positive posterior estimates (excluding s10), the remaining flux components ($NEE + F_{river}$) remain positive even after subtracting the fires and fossil fuels from the mean $F_{NetLand}$. Values range from neutral for s10sam with 0.00 PgC year⁻¹ to 0.35 PgC year⁻¹ for SIBCASA. Despite fires contributing 0.19 PgC year⁻¹ in this region, the persistence of positive $NEE + F_{river}$ in most inversions suggests a non-fire-related carbon source. Conversely, for the biogeographic Amazon we observe a consistently negative $NEE + F_{river}$ across all posterior estimates, with a smaller fire component (0.10
 390 PgC year⁻¹) compared to the adjacent 'Cerrado & Caatinga.' Therefore, given these results the majority of fire-related carbon sources lie outside the biogeographic Amazon, emphasizing the critical role of fire locations in determining remaining flux components in the NBE.




Figure 6. Cumulative burned area for the domain of the regional inversion using GFED5 (a) (Chen et al., 2023) and the prior and posterior fire emission estimates from GFAS and optimized with using satellite retrievals from MOPITT (Deeter et al., 2019) (b) for the biogeographic Amazon and the 'Cerrado & Caatinga' regions, the error bars denote the IAV.

The largest contribution (63%) to the mean fire emission in the biogeographic Amazon ($0.10 \text{ PgC year}^{-1}$) is from the 'Brazilian Shield Moist Forests' (see Figure A14). This fire estimate is lower than the estimates reported in other studies (Gatti et al., 2021; Basso et al., 2023), but our analysis shows that fire emissions are concentrated on the border of the Amazon and the 'Cerrado & Caatinga' region. Note that when adding the fire emissions from the 'Brazilian Shield Moist Forests' and the 'Cerrado & Caatinga', we obtain $0.25 \text{ PgC year}^{-1}$, which is close to the fire estimate of Basso et al. (2023). Furthermore, the cumulative burned area in GFED5 (Chen et al., 2023), a fire emission proxy independent of the optimized fires used here, has most of the burning in the 'Cerrado & Caatinga' (Figure 6a), which is also the case when using an alternative Amazon boundary, such as that defined by Eva et al. (2005). Burned area is indicative of fire activity but it does not scale 1:1 with fire carbon emissions, as they depend on factors like fuel load and combustion efficiency (van der Werf et al., 2010; van Wees et al., 2022). However, fire emissions in the 'Cerrado & Caatinga' region increase from prior ($0.08 \text{ PgC year}^{-1}$) to posterior estimates ($0.19 \text{ PgC year}^{-1}$) (Figure 6b), consistent with an increase in burned area from GFED4 (Giglio et al., 2013) to GFED5 Chen et al. (2023). It is worth noting that the fire emissions used here results from a top-down optimized version (Naus et al., 2022) based on CO using satellite retrievals from MOPITT (Deeter et al., 2019). Therefore, the difference in fire emissions relative to other studies is likely associated with the distinct methodologies used to attribute the fluxes as well as the spatial resolution used in the inversion, both of which can lead to accounting for fire emissions outside of the regional boundaries in the Amazon region.

3.3 Effect of systematic measurement uncertainty on posterior fluxes

The flask-specific CO_2 bias-correction (see Section 2.1.5) results in a consistent shift towards a source on the posterior F_{NetLand} for the biogeographic Amazon (Figure 7a). No significant effect is observed on interannual variability or spatial gradients (Figure A15). Applying the correction results in a mean posterior F_{NetLand} with a weaker sink or a larger source of carbon (Figure 7a) for particular years, consistent with the need to simulate higher CO_2 levels to match the data. Using the global inversion, we find a mean effect of $0.21 \text{ PgC year}^{-1}$, while in the regional inversion the effect is $0.31 \text{ PgC year}^{-1}$ (see Table A1). For the 'Cerrado & Caatinga' the mean effect is $0.10 \text{ PgC year}^{-1}$ (CarboScope Global) and $0.14 \text{ PgC year}^{-1}$ (CarboScope Regional), and for the regions within the Amazon equal or less than $0.10 \text{ PgC year}^{-1}$ for both global and regional inversions (see Table A1). Note that the correction is allocated to the regions that were selected previously having an uncertainty reduction larger than 15% consistent with the observational coverage (compare Figure 7b with Figure 2). In other words, the correction only affects the areas covered by the aircraft network.

Assuming that the correction brings the observational data closer to the truth implies that the Amazon is a weaker sink of carbon and that the 'Cerrado & Caatinga' is a larger source. We do not find a strong spatial shift of fluxes within the biogeographic Amazon due to the correction, nor do we find a strong impact on the interannual variations. The response to drought in 2010, 2015 and 2016 is affected in the absolute flux magnitude, but the year-to-year variability remains the same (Figure 7). Therefore, the findings of our sensitivity tests support the hypothesis in Gatti et al. (2023) that the water vapor bias mainly affects the absolute annual flux magnitudes. In both cases, with and without correction, our estimates for the total carbon loss to the atmosphere in the Amazon during 2015 and 2016, are lower (from 0.15 to 0.3 PgC) than other studies (Liu

et al., 2017; Gloor et al., 2018). Our total net flux is closer to the 0.5 ± 0.3 PgC from Gloor et al. (2018), but note that they used a time period from September 2015 to June 2016 and the area they refer to as Amazonia is not clearly defined. Compared to the 1.6 ± 0.29 PgC in Liu et al. (2017), our estimates are much lower, but the difference in area is large as they refer to tropical South America, including parts of the 'Cerrado & Caatinga' biomes and central America. Furthermore, they focus their study on the anomaly relative to 2011 at the peak of the El Niño phase.




Figure 7. Posterior flux estimates using the global inversion assimilating vertical profile data without a bias correction (s10sam) and with the correction (s10samwvc) (a). The integrated flux corresponds to the Biogeographic Amazon. The error bars correspond to the posterior uncertainty for the Biogeographic Amazon. In (b), the difference per grid cell of the two resulting posterior estimates averaged over 2010-2018.

4 Discussion

4.1 The Amazon carbon exchange in context

Significant progress advancing the monitoring capacity over the continent, particularly in Brazil, has allowed for an increasing number of studies (Gatti et al., 2014; van der Laan-Luijkx et al., 2015; Alden et al., 2016; Koren, 2020; Gatti et al., 2021; Botía et al., 2022; Basso et al., 2023; Gatti et al., 2023) using atmospheric mole fraction measurements to understand ecosystem carbon exchange. Nevertheless, using atmospheric inversions to estimate net carbon exchange remains a challenging task given the remaining limitations in data constraint, as shown in Figure 3. We have shown that posterior estimates have a linear dependence on the prior magnitude that varies according to the uncertainty reduction and absolute posterior uncertainty. On the biogeographic Amazon scale, such linear dependence together with the large posterior uncertainty (± 0.33 PgC year⁻¹) makes it difficult to attribute a mean carbon exchange based solely on one prior, as the magnitude and sign could be greatly influenced by that of the prior. To address this issue, our study offers a meticulous evaluation of uncertainty for specific regions, recommending areas where the inversion benefits from superior atmospheric data constraint. In light of this assessment, we

identify potential sites for new measurement stations to reduced the uncertainty in Amazon-wide top-down estimates, like the
445 'Amazon-Andes foothills'.

In this study, using an ensemble of prior fluxes in a regional inversion and a global inversion with and without South American data, we can infer that the biogeographic Amazon is a net carbon sink (i.e. negative $F_{NetLand}$) given by the direction of the adjustment introduced by the inversion. When subtracting fires and fossil fuels we obtain a negative 'NEE + River' flux component in all of the individual inversion runs, suggesting a consistent ecosystem carbon sink over 2010-2018. Taking
450 into account all posterior estimates, the median $F_{NetLand}$ is $-0.33 \text{ PgC year}^{-1}$. The posterior median without considering the inversion runs using priors with a large sink (FLUXCOM, X-BASE_{NEE}, and VPRM) is $-0.24 \text{ PgC year}^{-1}$, both with a posterior uncertainty of $\pm 0.33 \text{ PgC year}^{-1}$. We avoid reporting an ensemble mean as the large variability in the magnitude given by the setup of our experiments could be biased, and thus we rather interpret the individual posterior estimates in relative terms (as done in Section 3.1).

We acknowledge that a shortcoming of our regional inversion, common to other regional systems (Munassar et al., 2023), is the use of a single global inversion for the far-field contribution (e.g. the CO_2 mole fraction advected into the regional domain). For example, if the global inversion has a regional bias in the CO_2 mole fractions, it can propagate to the regional inversion via the far-field contribution. Yet, this influence should be smaller in well-constrained areas (those having high uncertainty reduction, Figure 3). In this regional domain, the predominant atmospheric flow changes seasonally from easterly
460 to northeasterly, both directions constrained by two background stations: Ragged Point Barbados (RPB, 13.1650°N , 59.432°W) and Ascension Island (ASC, 7.9667°S , 14.4°W). CarboScope global assimilates data from these stations and the mean bias error between posterior CO_2 mole fractions and local measurements at these sites is lower than 0.1 ppm (Botía et al. (2022), Supplementary Figures). In addition, Schuh et al. (2019) showed that atmospheric transport models (e.g. TM5) having a slow vertical and meridional transport in northern mid-latitudes can have a weaker sink from 45°N to 90°N , but a larger one from 0 to
465 45°N and thus result in a stronger global integrated carbon sink compared to the fast vertical mixing models (e.g. GEOS-Chem). We speculate that as a precursor of TM5, the model we used in the global inversion (i.e. TM3) could have a stronger global carbon sink due to a slower vertical and meridional transport, so the contribution of our South American domain to that carbon sink will propagate to the regional inversion via Step 1 of the two-step scheme. The quantification of this potential effect is a source of uncertainty in our regional inversions that should be further quantified. Finally, we have assumed weekly error
470 correlations for the model-data mismatch uncertainty to be able to assimilate multiple data streams and for consistency with the CarboScope Global inversion system. Such assumption should be revisited and evaluated with further sensitivity tests in future studies in the tropics, but maintaining consistency between the Global (which provides the boundary conditions) and Regional inversion systems.

Our estimates for the biogeographic Amazon are consistent with bottom-up approaches in the negative sign of net land
475 flux ($F_{NetLand}$) (Table 2), yet a large uncertainty remains in both approaches. For the same area, Basso et al. (2023) reports a small net source of carbon, but considering the uncertainty range, all approaches overlap. Integrating our posterior fluxes for the same Amazon boundaries as in Gatti et al. (2021) and Basso et al. (2023) results in contrasting signs, but again with an overlapping uncertainty range. Therefore, to answer the sink/source question for the Amazon region -regardless of its

boundaries-, reducing uncertainty in both bottom-up and top-down approaches should be the priority in upcoming studies.
480 From the top-down perspective, such uncertainty reduction can be achieved by expanding the observational network to the west of the Amazon, towards the Andes foothills in Peru, Colombia, and Ecuador, as this region had the smallest uncertainty reduction in our study. Thus, mean estimates over these regions should be interpreted with caution, as the selection of prior fluxes can largely influence the sign and the magnitude of such a mean, as shown in this study.

To further reduce the uncertainty in this domain, top-down estimates could combine in-situ data with satellite retrievals. The
485 inversions assimilating data from the Orbiting Carbon Observatory 2 (OCO2) (Liu et al., 2017; Crowell et al., 2019; Peiro et al., 2022; Wang et al., 2023) have shown that remotely-sensed CO₂ columns can provide a valuable constraint of net carbon exchange in tropical regions. However, the OCO2-inversions are still limited by cloud coverage during the wet season (Massie et al., 2017; Peiro et al., 2022) and the adjustment of the prior can be biased to dry season retrievals (Crowell et al., 2019). Nevertheless, our results for the response to the El Niño 2015/2016 coincide with the OCO2 inversions (Liu et al., 2017;
490 Crowell et al., 2019; Peiro et al., 2022) in a carbon source in 2015 and 2016. Yet, a direct comparison of the magnitude in those studies to our results is difficult, as the area for South America in Liu et al. (2017) includes parts of the 'Cerrado & Caatinga' regions and in Crowell et al. (2019); Peiro et al. (2022) they divide South America in three parts: 1. Northern South America: including only the north (north of the equator) of the Amazon basin, the Orinoco basin, a part of central America and the Caribbean islands. 2. Southern Tropical South America, which includes a large part of the Cerrado and the southern part
495 of the Amazon and 3. South America Temperate, which includes the Cerrado and Caatinga, extending until the southernmost point on the continent. None of these regions coincide with our regional distribution, therefore, using our domain definition on the OCO2-MIP results should be part of a next study, as it is out of the scope of this one.

The effect of fires is fundamental to understanding the NBE and quantifying this signal remains challenging. Studies like van der Laan-Luijkx et al. (2015); Koren (2020); Naus (2021); Basso et al. (2023), show that further optimization of fires in
500 a Bayesian setup can have an important impact on the magnitude of the derived fire-CO signal, which can be translated to a fire-CO₂ signal and thus impact the net carbon flux on the regional scale. This is why we selected an optimized fire emission for our domain, as did Basso et al. (2023), which results in fire emissions that are generally larger than the prior used, in both cases GFAS. The higher estimates than the original GFAS product are in line with recent findings that the global burned area has been strongly underestimated, resulting in underestimated fire emissions (e.g. Ramo et al. (2021)). Further advances in
505 global burned area mapping (Chen et al., 2023) and fire emission estimates (van Wees et al., 2022; Wiedinmyer et al., 2023) will help to reduce the estimated fire flux and help to further constrain NBE in the future. The fire flux for the Amazon area used in Gatti et al. (2021) is 0.41 ± 0.05 , which is relatively high compared to the $0.09 \text{ PgC year}^{-1}$ (Rosan et al., 2024), the $0.26 \text{ PgC year}^{-1}$ (Basso et al., 2023), and the $0.10 \text{ PgC year}^{-1}$ from this study. The Gatti et al. (2021) approach with the column budget technique solves for the total flux concerning a background signal linked to each of the aircraft profiles. Using
510 an observation-based CO:CO₂ relationship, they obtained the contribution of fires and subtract that from the $F_{NetLand}$, but the area for attributing fluxes corresponds to the regional influence of each aircraft site limited to the Amazon boundaries. As the CO:CO₂ is also influenced by areas outside of the Amazon (see Figures 5 and 6 in Cassol et al. (2020) and Figure 1 in Gatti et al. (2014)), such an approach could assign fire sources from the Cerrado to the Amazon.

In addition to the challenges posed by a limited observational constraint in this region, other sources of uncertainty could be associated with water vapor in the samples of the aircraft vertical profiles, as was mentioned in Gatti et al. (2023). Previous studies (Baier et al., 2020) focusing on the effect of water vapor in flasks, have shown that the linear dependency of the ΔCO_2 on humidity in the flasks is more evident above $\approx 1.5\%$ xH_2O . We have fitted the ΔCO_2 to the entire humidity range in the measurements and thus the range simulated by the STILT-ECMWF-IFS model (see Figure A7), so we recognize that the bias estimates at low water vapor levels could be slightly overestimated. Nevertheless, we hypothesized that the effect of that overestimation on the posterior fluxes should be minor, as water vapor mole fractions are predominantly above 1% at all sites. Thus, our approach provides a reference for future studies focusing on characterizing the water dependence in undried PFP samples and the effect of such a bias in an atmospheric inversion. We have quantified the effect of this systematic uncertainty on the posterior estimates, resulting in a weaker sink or a larger carbon source of 0.31 (based on the regional inversion) to 0.21 PgC year^{-1} (based on the global inversion). As we have shown, assimilating a bias-corrected dataset does not affect the IAV nor the spatial patterns of the retrieved fluxes and rather results in an upward shift (towards a source of carbon) of the posterior flux.

The implications of our suggested correction depend on the prior used, but factoring in the correction (see Table 2) in the median F_{NetLand} of $-0.33 \text{ PgC year}^{-1} \pm 0.33 \text{ PgC year}^{-1}$, we obtain a close-to-neutral carbon balance ($-0.02 \text{ PgC year}^{-1} \pm 0.33 \text{ PgC year}^{-1}$ if the correction from the regional inversion is used) or a small sink ($-0.13 \text{ PgC year}^{-1} \pm 0.33 \text{ PgC year}^{-1}$, if the correction from the global inversion is used). Note that in both cases after subtracting fires and fossil fuels the resulting flux will still represent an ecosystem carbon sink (a negative $\text{NEE} + F_{\text{rivers}}$). In addition, considering the posterior uncertainty of $\pm 0.33 \text{ PgC year}^{-1}$ together with the potential effect of water vapor in the flasks ($0.31 \text{ PgC year}^{-1}$, using the regional adjustment), the positive shift amounts to $+0.64 \text{ PgC year}^{-1}$ suggesting a large change in the carbon balance at the biogeographic Amazon scale. In any case, this change is within the large spread in F_{NetLand} estimates by different approaches (Table 2). Now, placing this in the context of the global carbon budget, an increase of $+0.64 \text{ PgC year}^{-1}$ is within the uncertainty of the global net land carbon fluxes ($1.1 \text{ PgC year}^{-1}$) for 2013-2022 reported in Friedlingstein et al. (2023), and would represent a third of the constraint given by atmospheric inversions with a range of $0.5\text{-}2.3 \text{ PgC year}^{-1}$ (Friedlingstein et al., 2023). Therefore, given the large uncertainties reported for the global carbon budget, it is possible to accommodate such a magnitude shift in the net land carbon flux of the biogeographic Amazon.

4.1.1 Rivers

For the carbon balance of the main branch of the Amazon River, we estimated a F_{NetLand} close to neutral. The median F_{NetLand} for the 'Amazon River Flat Plains' region was $-0.04 \pm 0.09 \text{ PgC year}^{-1}$. Note that this region is relatively well constrained by the atmospheric monitoring network with an mean UR of 53%, thus the posterior estimates have the smallest spread of all regions (see Figure 4d). After removing fires and fossil fuel emissions (0.01 and $0.007 \text{ PgC year}^{-1}$) the resulting flux, $\text{NEE} + F_{\text{river}}$, is slightly negative. In this work we have considered river CO_2 evasion (F_{river}) as an explicit component of the NBE because we believe is a land flux that should be characterized independently of the NEE, in particular for the Amazon lowland area considering that at least 31% (Fleischmann et al., 2022) can be seasonally flooded.

Table 2. Comparison of $F_{NetLand}$ for the Amazon region based on top-down and bottom-up approaches averaged over 2010-2018. We present estimates for two different definitions of Amazon boundaries, the Biogeographic Amazon and the Amazon *sensu-stricto+*. The latter corresponds to four subregions defined in Eva et al. (2005): Amazon *sensu stricto*, Andes, Guiana and Gurupi. We present our estimates for the Amazon *sensu-stricto+* so the area is comparable to previous top-downs studies. The acronym WVC, stands for water vapor correction. Units in PgC year⁻¹.

	Area	Fires	$F_{NetLand}$	Uncertainty
This study (all ensemble members)	Biogeographic Amazon	0.10	-0.33	± 0.33
This study (only process-based models)	Biogeographic Amazon	0.10	-0.24	± 0.33
Rosan et al. (2024)-Bottom-up	Biogeographic Amazon	0.09	-0.15	± 0.19
Rosan et al. (2024)-Hybrid	Biogeographic Amazon	-	-0.25	± 0.19
Rosan et al. (2024)-CARDAMOM	Biogeographic Amazon	-	-0.34	CI=[-2.94,2.45]
Basso et al. (2023) in Rosan et al. (2024)	Biogeographic Amazon	-	0.02	± 0.13
This study	Amazon <i>Sensu stricto+</i>	0.11	-0.18	± 0.34
Basso et al. (2023)	Amazon <i>Sensu stricto+</i>	0.26	0.13	± 0.17
Gatti et al. (2021)	Amazon <i>Sensu stricto+</i>	0.41	0.29	± 0.40
This study (all ensemble members) + WVC-global	Biogeographic Amazon	0.10	-0.33+0.21 = -0.12	± 0.33
This study (all ensemble members) + WVC-regional	Biogeographic Amazon	0.10	-0.33+0.31 = -0.02	± 0.33

Using output from a process-based model (ORCHILEAK, Hastie et al. (2019)), Botía et al. (2022) suggested that river outgassing could play an important role in representing the seasonal pattern of CO₂ mole fractions at ATTO, yet there are important processes in that model that are not accounted for (i.e. aquatic plants), which is why in this study we did not used ORCHILEAK as a prior. However, if we consider F_{river} as the CO₂ evasion based on the ORCHILEAK model (Hastie et al., 2019) for the 'Amazon River Flat Plains' (0.09 PgC year⁻¹), the resulting NEE for this area would be even more negative, suggesting that plant productivity in this region is larger than the respired CO₂ from the decomposition of organic matter. Nevertheless, the growth of aquatic plants in rivers could play an important role in the net balance of riverine CO₂ fluxes (Science Panel for the Amazon, 2021). This uptake of carbon, which is not taken into account in ORCHILEAK, could potentially balance out the CO₂ outgassing due to decomposition of submerged organic carbon and respiration of roots, which are the main sources in ORCHILEAK for the CO₂ evasion. Our results for the 'Amazon River Flat Plains' suggest the region is close-to-neutral, but to partition the components NEE + F_{river} using an atmospheric inversion, bottom-up estimates for rivers should consider aquatic plant productivity, which is a crucial process to determine net river evasion or uptake.

4.2 Sink-to-source gradient between the biogeographic Amazon and the Cerrado & Caatinga

Our results suggest that the Amazon is a net carbon sink and that the 'Cerrado & Caatinga' biomes are possibly net sources of carbon, even after removing fires in some individual inversions we obtain a positive NBE. This is in contrast to Gatti et al.

(2021) and Basso et al. (2023). Gatti et al. (2021) suggested that vegetation of the southeast of the Amazon was losing the capacity to capture carbon, reporting on average a positive NBE for the years 2010 to 2018, thus locating a net source of carbon within the Amazon in the southeast regions. Basso et al. (2023) report a posterior carbon source in the eastern part of their domain (see Figure A11 in Basso et al. (2023)). In both studies, the analyses were limited to the Amazon *Sensu stricto* + (as defined in Table 2) but used the same aircraft data as we used in this study. Therefore, we believe that the main reasons explaining the differences reported here are: 1. limiting their region of interest to the Amazon boundaries, while using atmospheric data that is influenced by a footprint that goes beyond the Amazon and 2. having a coarse global inversion in which the boundaries between the Amazon biome and the Cerrado are not well-defined leading to attributing sources from the Cerrado to the Amazon.

The spatial attribution of fluxes in Gatti et al. (2021) is performed using a column budget technique. In this method, the residence time of air parcels over each region of influence for each aircraft site is used to account for the time and the area that contribute to an enhancement or depletion of CO₂ mole fractions at the site relative to the background. In our study, a similar source is needed to match the atmospheric profile data, but it is further east outside the Amazon bounds. This solution is possible in our system with individual (though correlated) grid points that can be assigned extra flux, but it is not possible in the setup of Gatti et al. (2021), where a mean flux rate is assigned to the areas of influence of the airborne measurements, which is limited to the Amazon boundary. Thus, the degrees of freedom to place sources/sinks in specific biomes further upwind are not present. Sensitivity tests in Botía (2022) indicate that our result is robust against changing the spatial error structure in the inversion settings. While recognizing such spatial differences between the studies, we nevertheless conclude that the east-to-west gradient within Amazonia that Gatti et al. (2021) report is not seen in our posterior ensemble median estimates. Interestingly, an individual ensemble member (i.e. s10sam) does result in a source (0.03 gC m⁻² d⁻¹) in the 'Brazilian Shield Moist Forest', but still having a larger one (0.12 gC m⁻² d⁻¹) in the 'Cerrado & Caatinga'. Therefore, it is likely that the atmospheric signal of a carbon source is unmistakably in the data, but is attributed to different spatial regions by the different methodologies. Understanding this discrepancy, and determining the location of the eastern Brazilian CO₂ source should have the highest priority in further work. Next, despite of these discrepancies, we speculate on what could be driving a source of carbon in the 'Cerrado & Caatinga' region.

The 'Cerrado & Caatinga' biomes cover approximately 35% of the Brazilian land mass (Beuchle et al., 2015) and are characterized by a Savanna-type ecosystem (Cerrado) (Sano et al., 2007) and seasonally dry tropical forest (Caatinga) (Prado, 2003). Both biomes have a marked seasonality in precipitation, with mean annual precipitation of less than 750 mm year⁻¹ in the Caatinga (Prado, 2003; Leal et al., 2005), and from 800 to 2000 mm year⁻¹ in the Cerrado (Ratter et al., 1997; Rodrigues et al., 2022). Studies focusing on ecosystem functioning conducted in both biomes using eddy covariance measurements have shown that these ecosystems have similar seasonal patterns in net ecosystem exchange. In general, with the onset of the rainy season a higher carbon uptake is observed over converted pastures and over natural vegetation (Miranda et al., 1997; Varella et al., 2004; Santos et al., 2004; Silva et al., 2017; Mendes et al., 2020; Alves et al., 2021). Studies focusing on soil CO₂ emissions comparing converted pasture and both natural ecosystems, Caatinga (Ribeiro et al., 2016) and Cerrado (Varella et al., 2004), found no significant differences in magnitude between the pasture and the natural ecosystem. Both coincided with

higher CO₂ emissions at the beginning of the rainy season (Varella et al., 2004; Ribeiro et al., 2016). The findings of Ribeiro et al. (2016) are in line with Mendes et al. (2020), where they observed an increase in ecosystem respiration with the onset of the rainy season, but offset by GPP. Integrated over time, these ecosystems in their natural form or converted to pasture seemed to be carbon sinks when no disturbance is taken into account (Santos et al., 2004; Bustamante et al., 2012; Silva et al., 2017; Mendes et al., 2020; Alves et al., 2021).

However, these biomes have suffered considerable loss of natural vegetation due to the expansion of the agricultural frontier (Beuchle et al., 2015; Alencar et al., 2019). Beuchle et al. (2015), found that over the period between 1990 and 2010, both biomes had a continued net loss of natural vegetation. More recently, most of the agricultural expansion in the Cerrado has been concentrated in a region called MATOPIBA, which refers to portions of the Maranhão, Tocantins, Piauí and Bahia states (Spera et al., 2016; da Conceição Bispo et al., 2024). Drought periods in the Caatinga can extend over years (Leal et al., 2005), making agricultural activities more difficult to sustain, yet considerable pasture conversion for extensive livestock has changed the Caatinga landscape (Leal et al., 2005). Fire is used for the conversion of forest and shrublands to pasture or croplands, thus that conversion leads to fire CO₂ emissions (van der Werf et al., 2010; Pivello, 2011; van der Werf et al., 2017). The Cerrado biome was found to have higher fire CO₂ emissions than the Caatinga, with an increasing trend in recent years (da Silva Junior et al., 2020). Moreover, the annual fire regime fluctuates between naturally occurring low-intensity fires at the end of the wet season ignited by lightning, and anthropogenic high-intensity fires at the end of the dry season (Ramos-Neto and Pivello, 2000; Pletsch et al., 2022). Frequent fires can lead to aboveground biomass reduction, changing the ecosystem from a sink to a source of carbon (de Azevedo et al., 2020). Moreover, Bustamante et al. (2012) found that a large portion of the CO₂ emissions from pasture management (i.e. burning practices) in Brazil originated in the Cerrado. Throughout 2003 to 2013, changes in vegetation stocks due to cropland conversion in the Cerrado and specifically in the MATOPIBA region, contributed to 33% of the forest carbon emissions (Noojipady et al., 2017).

Given these studies and our findings, carbon emissions in these two biomes but primarily in the Cerrado, can be grouped into two categories. The first one is associated directly with fires (either from deforestation or pasture management), and the second arises from degradation after the conversion of natural vegetation to pastures or croplands. For the first category, we showed that fire emissions increased from prior to posterior in the 'Cerrado & Caatinga' region by a factor of 2.3, which is broadly consistent with an increase in burned area from GFED4 to GFED5, by a factor of 1.7. Emissions in the second category are associated with changes in carbon stocks, decomposition, and sink-to-source shifts due to climate change (Bustamante et al., 2012; Marengo et al., 2022). Having this in mind, we hypothesize that the carbon source in the semi-arid ecosystems of the 'Cerrado & Caatinga' given by the atmospheric inversion is likely due to agricultural expansion in the Cerrado, mainly from the second category mentioned above. On top, their capacity for secondary forest regrowth is compromised. This contrasts with the Amazon biome, which is highly resilient (Sakschewski et al., 2016; Poorter et al., 2021), and has shown a relatively rapid recovery of aboveground biomass by secondary forest growth (Poorter et al., 2016) with considerable potential to capture carbon (Heinrich et al., 2021, 2023). It is important to note here that Amazon forest resilience is heavily affected by anthropogenic disturbances (Fawcett et al., 2023; Wang et al., 2024) and it seems to be decreasing since 2000 (Boulton et al., 2022), although this finding has been questioned (Tao et al., 2023). This would make the carbon source we find in this study not only

large but likely also influence the regional carbon balance for decades to come. Therefore, the hypothesis of the 'Cerrado & Caatinga' source, driven by changes in carbon stocks and fires, should be part of a future study in which additional ways of testing the robustness of the inversion results are explored. For example, comparing several inversion systems using same data constraint could shed light on the spatial gradients reported here. Finally, an assessment of having different or even multiple global inversions for constraining the far-field contribution in the regional inversion should also be quantified and studied.

5 Conclusions

In this study, we have integrated in-situ CO₂ measurements from the Amazon Tall Tower Observatory together with a network of airborne vertical profiles in the CarboScope regional inversion system to estimate carbon fluxes in tropical South America. Our analysis is limited to regions with uncertainty reduction above 15%, which amounts to 67% of the land mass in our domain. Among these regions, the 'Amazon River Flat Plains', the 'Brazilian Shield Moist Forests' and to a lesser extent the 'Guianan Shield Moist Forests' are better constrained than others.

Furthermore, our results suggest a sink-source gradient between the Amazon (sink) and the 'Cerrado & Caatinga' (source). Across all ensemble members, this net absorption amounted to a median of -0.33 PgC year⁻¹. When relying solely on process-based models as priors, this value slightly decreased to -0.24 PgC year⁻¹, both estimates with a posterior uncertainty of ± 0.33 PgC year⁻¹. The Cerrado & Caatinga biomes together, acted as a median carbon source of 0.31 ± 0.24 PgC year⁻¹. This finding, which is in contrast to other top-down studies (Gatti et al., 2021; Basso et al., 2023), is explained by two reasons. The first one is that we do not limit our analysis to the Amazon boundaries, instead, we take into account the extended surface influence of the vertical profiles, which goes beyond the Amazon. The second reason is associated with a higher spatial resolution in our regional inversion system (0.25x0.25 degree) than in Basso et al. (2023). The latter allows us to integrate fluxes over the analysis regions with improved precision. This is particularly relevant considering that fires occur on -but are not limited to- the border of the Amazon and the Cerrado, so when having coarser grids one can attribute Cerrado fires to the Amazon.

We have quantified and reported important uncertainties associated with the data and methodology used in this work. Part of this was the assessment of how systematic uncertainties due to water vapor in the aircraft vertical profiles affect the estimated fluxes in our inversion system. In principle, we recommend to dry the air during sampling time to avoid systematic uncertainties and their propagation to flux estimates in an inversion system. Our analysis suggests that the proposed correction for water-vapor leads to an upward adjustment ranging from 0.21 to 0.31 PgC year⁻¹ in the biogeographic Amazon carbon flux. Note that other than the shift in magnitude, the interannual variability or the spatial gradients of the posterior fluxes do not change. Considering this correction and the remaining uncertainty (0.33 PgC year⁻¹) for the biogeographic Amazon, we conclude that such a change in the carbon budget is within the uncertainty of the global net land carbon flux, but represent further challenges in constraining the recent Amazon carbon balance.

Code and data availability. The data that support the findings of this study are, the ATTO CO₂ measurements, openly available in: <https://attodata.org/> and the aircraft vertical profiles, which are openly available in PANGAEA: <https://doi.pangaea.de/10.1594/PANGAEA.926834>. The data generated in this paper, the simulated water vapor at each aircraft site with its corresponding bias correction (in ppm) will be made public at the time of publication. For the review process, the reviewers can access the data under this private link <https://edmond.mpg.de/privateurl.xhtml?token=36e3c67d-d564-4c15-beaf-c0d46193a5b6>. The analysis regions in postprocessing can be found here: <https://edmond.mpg.de/privateurl.xhtml?token=a4af161f-1b77-4611-aa01-7de9203639b8>. At the time of publication, both links will be made public. The posterior fluxes for each individual inversion run and the fire-CO₂ emissions based on the MOPITT-CO inversions can be made available upon request to Santiago Botía (sbotia@bgc-jena.mpg.de). The GFED5 burned area was retrieved from the open repository Zenodo (<https://doi.org/10.5281/zenodo.7668423>) (Chen et al., 2023).

Author contributions. SB wrote the initial manuscript and run the atmospheric inversions. SB together with CG, WP and CR designed the methodology. CR developed the global and the regional inversion systems and he assisted in running the global inversions. SM and TK assisted with the regional inversion runs, DC ran the forward runs simulations for the validation site MAN. LSB helped with the data curation of the aircraft sites and further interpretation of the results. SK, JVL and DW provided the ATTO data and run the ATTO measurement system until 2021. GM is co-PI of the aircraft data and assisted with the interpretation of the results. SN, GK, IL and WP provided the fire-CO₂ emission based on the MOPITT-CO inversions. SH contributed with the analysis of burned area. JBM and GM ran the Manaus aircraft measurements and provided the data for the water vapor correction. All authors contributed to the analysis and text editing.

Competing interests. Some authors are members of the editorial board of the Atmospheric Chemistry and Physics (ACP) journal.

Acknowledgements. This work and the ATTO project was funded by the German Federal Ministry of Education and Research (BMBF, contracts 01LB1001A and 01LK1602A) and supported by the International Max Planck Research School for Global Biogeochemical Cycles (IMPRS-gBGC). The ATTO project is furthermore funded by the Brazilian Ministério da Ciência, Tecnologia e Inovação (MCTI/FINEP contract 01.11.01248.00) and the Max Planck Society. We acknowledge the Instituto Nacional de Pesquisas da Amazonia as well as the Amazon State University (UEA), FAPEAM, LBA/INPA and SDS/CEUC/RDS-Uatuma for continuous support and logistical management. Many thanks to the people coordinating the scientific support at ATTO, in particular Susan Trumbore, Carlos Alberto Quesada, Bruno Takeshi, and Reiner Ditz. We express our gratitude to the data providers, Luciana Gatti (Aircraft vertical profiles), and Jacob Nelson, Sophia Walter and Martin Jung for the FLUXCOM and X-BASE NEE fluxes. W.P. and G.K. were funded by an ERC-Consolidator grant (649087) as part of the ASICA (Airborne Stable Isotopes of Carbon from the Amazon) project. Finally, we appreciate the comments received from the two reviewers during the revision phase, they contributed with improving the text and main message of the manuscript.

Appendix A




Figure A1. Seasonal surface influence for each station used in the regional inversion. The averaging period for each station corresponds to the period of data availability, which is site-specific, see Figure 2.




Figure A2. Number of aircraft profiles per month over the period of interest in the inversion. An aircraft profile goes up until 4500 m.a.s.l and on average collects samples at 14 heights.




Figure A3. Bias between PFPs and CRDS analyzer (Picarro Inc. model G2401-m) CO_2 mole fractions at MAN as a function of water vapor mole fraction measured by the CRDS analyzer.

Table A1. Mean effect on the posterior estimates of assimilating bias corrected vertical profiles using CarboScope global and regional. The difference was calculated $\text{postFluxWVC} - \text{postFluxNonWVC}$, therefore positive numbers indicate that the WVC posterior flux is larger. Units in PgC year^{-1} .

	CarboScope Global	CarboScope Regional
Amazon and Andes Piedmont	0.02	0.04
Brazilian Shield Moist Forest	0.08	0.10
Cerrado & Caatinga	0.10	0.14
Guianan Shield Moist Forest	0.04	0.07
Amazon River Flat Plains	0.05	0.09
Biogeographic Amazon	0.21	0.31
All LAND	0.28	0.50

Table A2. Eddy flux sites used to calibrate the VPRM parameters.

Site code	Lat	Lon	Site Veg. Description	Veg. Class VPRM
STM-K67	-2.85700	-54.95900	Primary Tropical Moist Forest	Evergreen Forest
STM-K77	-3.02020	-54.88850	Pasture, then Agriculture	Cropland
STM-K83	-3.01700	-54.97070	Primary Tropical Moist Forest, sel. logging Aug/Sept 2001	Evergreen Forest
MAN-k34	-2.50000	-60.20910	Tropical Rainforest	Evergreen Forest
PA-CAX	-1.74830	-51.45360	Tropical Forest, dense lowland tropical forest	Evergreen Forest
RON-FNS	-10.76180	-62.35720	Pasture	Grassland
RON-RJA	-10.07800	-61.93310	Tropical Dry Forest	Evergreen Forest
TOC-BAN	-9.824416667	-50.15911	Seasonally flooded Forest-Savanna Ecotone	Evergreen Forest
SP-PDG	-21.61947222	-47.64989	Savanna	Savannas




Figure A4. Individual vertical profiles with the in-situ (black dots) and the discrete PFPs samples in Manaus (MAN). Each in-situ point is a 1 Hz interpolated value from the calibrated native CRDS signal at 0.3 - 0.4 Hz.




Figure A5. Correlation between STILT ECMWF-IFS water vapor and the measured water mole fraction at the Manaus flights with a Picarro (model G2401-m). The grey line corresponds to the predicted y using a linear regression.




Figure A6. Distribution of water vapor mole fractions at all sites extracted from ECMWF-IFS and the measurements at Manaus.




Figure A7. Bias between Picarro (model G2401-m) and PFPs CO₂ mole fractions at Manaus as a function of water vapor mole fraction measured by the Picarro (model G2401-m) and also extracted from ECMWF-IFS.




Figure A8. Bias (Picarro - PFP) estimated at each site using the fit to ECMWF-IFS water vapor.




Figure A9. Prior mean NEE over 2010-2018 for several of the models used. Note the different range in the colorbar.




Figure A10. Posterior mean NEE over 2010-2018 for several of the models used. Note the different range in the colorbar.




Figure A11. Prior (upper panel) and posterior (lower panel) seasonal cycle of NBE for each ensemble member aggregated for each region of interest.




Figure A12. Prior to posterior uncertainty reduction throughout the complete inversion period (2010-2018).




Figure A13. Atmospheric forward runs using the prior (a) and posterior fluxes (b). Model-data comparison is done at the sites assimilated in the inversion (NAT, ATT, ALF, TAB, TEF, SAN, RBA) and at one site that was not assimilated (MAN*, v2 (Miller et al., 2023)) as a validation site. Note that the global inversions s10 and s10sam have a zero-prior, so no forward run is available for those. In addition, note that s10 global inversion did not assimilate any of the sites shown here. At the MAN site we use aircraft profiles using a Picarro (model G2401-m) over the 2017 to 2018. For the MAN measurements the surface influence (or footprints) over were calculated for a 10s CO₂ mole fraction average. We used the same settings described for STILT-IFS in the methods.




Figure A14. Prior and posterior GFAS fire emissions for the regions within the Biogeographic Amazon (a-d), the Cerrado & Caatinga (e) and the Biogeographic Amazon (f).




Figure A15. Time series of posterior fluxes for each region using the global inversion assimilating data with the water vapor correction and without it.

References

- Albert, J., Hoorn, C., Malhi, Y., Phillips, O., Encalada, A. C., Hecht, S., Varese, M., Peña-Claros, M., and Roca, F. A.: The multiple viewpoints for the Amazon: geographic limits and meanings, Tech. rep., https://www.theamazonwewant.org/wp-content/uploads/2021/09/220105_The-multiple-viewpoints-for-the-Amazon-formatted-and-reviewed-050122.pdf, 2021.
- Alden, C. B., Miller, J. B., Gatti, L. V., Gloor, M. M., Guan, K., Michalak, A. M., van der Laan-Luijkx, I. T., Touma, D., Andrews, A., Basso, L. S., Correia, C. S. C., Domingues, L. G., Joiner, J., Krol, M. C., Lyapustin, A. I., Peters, W., Shiga, Y. P., Thoning, K., van der Velde, I. R., van Leeuwen, T. T., Yadav, V., and Diffenbaugh, N. S.: Regional atmospheric CO₂ inversion reveals seasonal and geographic differences in Amazon net biome exchange, *Global Change Biology*, 22, 3427–3443, <https://doi.org/10.1111/gcb.13305>, 2016.
- Alencar, A., Moutinho, P., Arruda, V., Balzani, C., and Ribeiro, J.: Amazon Burning: Locating the fires, Tech. rep., available at: https://ipam.org.br/wp-content/uploads/2019/09/AmazonBurning_LocatingTheFires.pdf, 2019.
- Alves, J. D. N., Ribeiro, A., Rody, Y. P., Loos, R. A., and Hall, K. B.: Carbon uptake and water vapor exchange in a pasture site in the Brazilian Cerrado, *Journal of Hydrology*, 594, 125 943, <https://doi.org/10.1016/j.jhydrol.2020.125943>, 2021.
- Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., Silva, C. V. J., Junior, C. H. L. S., Arai, E., Aguiar, A. P., Barlow, J., Berenguer, E., Deeter, M. N., Domingues, L. G., Gatti, L., Gloor, M., Malhi, Y., Marengo, J. A., Miller, J. B., Phillips, O. L., and Saatchi, S.: 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions, *Nature Communications*, 9, 1–12, <https://doi.org/10.1038/s41467-017-02771-y>, 2018.
- Assis, T. O., Aguiar, A. P. D. d., Randow, C. v., Gomes, D. M. d. P., Kury, J. N., Ometto, J. P. H. B., and Nobre, C. A.: CO₂ emissions from forest degradation in Brazilian Amazon, *Environmental Research Letters*, 15, 104 035, <https://doi.org/10.1088/1748-9326/ab9cfc>, 2020.
- Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., and Houghton, R. A.: Tropical forests are a net carbon source based on aboveground measurements of gain and loss, *Science*, 358, 230–234, <https://doi.org/10.1126/science.aam5962>, 2017.
- Baier, B. C., Sweeney, C., Choi, Y., Davis, K. J., DiGangi, J. P., Feng, S., Fried, A., Halliday, H., Higgs, J., Lauvaux, T., Miller, B. R., Montzka, S. A., Newberger, T., Nowak, J. B., Patra, P., Richter, D., Walega, J., and Weibring, P.: Multispecies Assessment of Factors Influencing Regional CO₂ and CH₄ Enhancements During the Winter 2017 ACT-America Campaign, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2019JD031339>, 2020.
- Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon dioxide uptake by land and oceans during the past 50 years, *Nature*, 488, 70–72, <https://doi.org/10.1038/nature11299>, 2012.
- Basso, L. S., Wilson, C., Chipperfield, M. P., Tejada, G., Cassol, H. L. G., Arai, E., Williams, M., Smallman, T. L., Peters, W., Naus, S., Miller, J. B., and Gloor, M.: Atmospheric CO₂ inversion reveals the Amazon as a minor carbon source caused by fire emissions, with forest uptake offsetting about half of these emissions, *Atmospheric Chemistry and Physics*, 23, 9685–9723, <https://doi.org/10.5194/acp-23-9685-2023>, 2023.
- Bastos, A., O’Sullivan, M., Ciais, P., Makowski, D., Sitch, S., Friedlingstein, P., Chevallier, F., Rödenbeck, C., Pongratz, J., Lujkx, I. T., Patra, P. K., Peylin, P., Canadell, J. G., Lauerwald, R., Li, W., Smith, N. E., Peters, W., Goll, D. S., Jain, A., Kato, E., Lienert, S., Lombardozzi, D. L., Haverd, V., Nabel, J. E. M. S., Poulter, B., Tian, H., Walker, A. P., and Zaehle, S.: Sources of Uncertainty in Regional and Global Terrestrial CO₂ Exchange Estimates, *Global Biogeochemical Cycles*, 34, <https://doi.org/10.1029/2019GB006393>, 2020.
- Beuchle, R., Grecchi, R. C., Shimabukuro, Y. E., Seliger, R., Eva, H. D., Sano, E., and Achard, F.: Land cover changes in the Brazilian Cerrado and Caatinga biomes from 1990 to 2010 based on a systematic remote sensing sampling approach, *Applied Geography*, 58, 116–127, <https://doi.org/10.1016/j.apgeog.2015.01.017>, 2015.

- Bodesheim, P., Jung, M., Gans, F., Mahecha, M. D., and Reichstein, M.: Upscaled diurnal cycles of land-atmosphere fluxes: a new global half-hourly data product, *Earth System Science Data*, 10, 1327–1365, <https://doi.org/https://doi.org/10.5194/essd-10-1327-2018>, publisher: Copernicus GmbH, 2018.
- Botía, S.: Greenhouse gas exchange in the Amazon region : carbon dioxide and methane insights from the Amazon Tall Tower Observatory (ATTO), Ph.D. thesis, Wageningen University, <https://doi.org/10.18174/573967>, 2022.
- Botía, S., Komiya, S., Marshall, J., Koch, T., Gałkowski, M., Lavric, J., Gomes-Alves, E., Walter, D., Fisch, G., Pinho, D. M., Nelson, B. W., Martins, G., Luijkx, I. T., Koren, G., Florentie, L., Carioca de Araújo, A., Sá, M., Andreae, M. O., Heimann, M., Peters, W., and Gerbig, C.: The CO₂ record at the Amazon Tall Tower Observatory: A new opportunity to study processes on seasonal and inter-annual scales, *Global Change Biology*, 28, 588–611, <https://doi.org/10.1111/gcb.15905>, 2022.
- Boulton, C. A., Lenton, T. M., and Boers, N.: Pronounced loss of Amazon rainforest resilience since the early 2000s, *Nature Climate Change*, 12, 271–278, <https://doi.org/10.1038/s41558-022-01287-8>, 2022.
- Bousquet, P., Peylin, P., Ciais, P., Le Quééré, C., Friedlingstein, P., and Tans, P. P.: Regional Changes in Carbon Dioxide Fluxes of Land and Oceans Since 1980, *Science*, 290, 1342–1346, <https://doi.org/10.1126/science.290.5495.1342>, publisher: American Association for the Advancement of Science, 2000.
- Brando, P. M., Paolucci, L., Ummenhofer, C. C., Ordway, E. M., Hartmann, H., Cattau, M. E., Rattis, L., Medjibe, V., Coe, M. T., and Balch, J.: Droughts, Wildfires, and Forest Carbon Cycling: A Pantropical Synthesis, *Annual Review of Earth and Planetary Sciences*, 47, 555–581, <https://doi.org/10.1146/annurev-earth-082517-010235>, 2019.
- Brienen, R. J. W., Phillips, O. L., Feldpausch, T. R., Gloor, E., Baker, T. R., Lloyd, J., Lopez-Gonzalez, G., Monteagudo-Mendoza, A., Malhi, Y., Lewis, S. L., Vásquez Martínez, R., Alexiades, M., Álvarez Dávila, E., Alvarez-Loayza, P., Andrade, A., Aragão, L. E. O. C., Araujo-Murakami, A., Arets, E. J. M. M., Arroyo, L., Aymard C., G. A., Bánki, O. S., Baraloto, C., Barroso, J., Bonal, D., Boot, R. G. A., Camargo, J. L. C., Castilho, C. V., Chama, V., Chao, K. J., Chave, J., Comiskey, J. A., Cornejo Valverde, F., da Costa, L., de Oliveira, E. A., Di Fiore, A., Erwin, T. L., Fauset, S., Forsthofer, M., Galbraith, D. R., Grahame, E. S., Groot, N., Hérault, B., Higuchi, N., Honorio Coronado, E. N., Keeling, H., Killeen, T. J., Laurance, W. F., Laurance, S., Licona, J., Magnussen, W. E., Marimon, B. S., Marimon-Junior, B. H., Mendoza, C., Neill, D. A., Nogueira, E. M., Núñez, P., Pallqui Camacho, N. C., Parada, A., Pardo-Molina, G., Peacock, J., Peña-Claros, M., Pickavance, G. C., Pitman, N. C. A., Poorter, L., Prieto, A., Quesada, C. A., Ramírez, F., Ramírez-Angulo, H., Restrepo, Z., Roopsind, A., Rudas, A., Salomão, R. P., Schwarz, M., Silva, N., Silva-Espejo, J. E., Silveira, M., Stropp, J., Talbot, J., ter Steege, H., Teran-Aguilar, J., Terborgh, J., Thomas-Caesar, R., Toledo, M., Torello-Raventos, M., Umetsu, R. K., van der Heijden, G. M. F., van der Hout, P., Guimarães Vieira, I. C., Vieira, S. A., Vilanova, E., Vos, V. A., and Zagt, R. J.: Long-term decline of the Amazon carbon sink, *Nature*, 519, 344–348, <https://doi.org/10.1038/nature14283>, 2015.
- Bustamante, M. M. C., Nardoto, G. B., Pinto, A. S., Resende, J. C. F., Takahashi, F. S. C., and Vieira, L. C. G.: Potential impacts of climate change on biogeochemical functioning of Cerrado ecosystems, *Brazilian Journal of Biology*, 72, 655–671, <https://doi.org/10.1590/S1519-69842012000400005>, 2012.
- Cassol, H. L. G., Domingues, L. G., Sanchez, A. H., Basso, L. S., Marani, L., Tejada, G., Arai, E., Correia, C., Alden, C. B., Miller, J. B., Gloor, M., Anderson, L. O., Aragão, L. E. O. C., and Gatti, L. V.: Determination of Region of Influence Obtained by Aircraft Vertical Profiles Using the Density of Trajectories from the HYSPLIT Model, *Atmosphere*, 11, 1073, <https://doi.org/10.3390/atmos11101073>, 2020.

- 765 Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J. T.: Multi-decadal trends and variability in burned area from the 5th version of the Global Fire Emissions Database (GFED5), *Earth System Science Data Discussions*, pp. 1–52, <https://doi.org/10.5194/essd-2023-182>, publisher: Copernicus GmbH, 2023.
- Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494, 341–344, <https://doi.org/10.1038/nature11882>, publisher: Nature Publishing Group, 2013.
- 770 Crowell, S., Baker, D., Schuh, A., Basu, S., Jacobson, A. R., Chevallier, F., Liu, J., Deng, F., Feng, L., McKain, K., Chatterjee, A., Miller, J. B., Stephens, B. B., Eldering, A., Crisp, D., Schimel, D., Nassar, R., O'Dell, C. W., Oda, T., Sweeney, C., Palmer, P. I., and Jones, D. B. A.: The 2015–2016 carbon cycle as seen from OCO-2 and the global in situ network, *Atmospheric Chemistry and Physics*, 19, 9797–9831, <https://doi.org/https://doi.org/10.5194/acp-19-9797-2019>, publisher: Copernicus GmbH, 2019.
- 775 da Conceição Bispo, P., Picoli, M. C. A., Marimon, B. S., Marimon Junior, B. H., Peres, C. A., Menor, I. O., Silva, D. E., de Figueiredo Machado, F., Alencar, A. A. C., de Almeida, C. A., Anderson, L. O., Aragão, L. E. O. C., Breunig, F. M., Bustamante, M., Dalagnol, R., Diniz-Filho, J. A. F., Ferreira, L. G., Ferreira, M. E., Fisch, G., Galvão, L. S., Giarolla, A., Gomes, A. R., de Marco Junior, P., Kuck, T. N., Lehmann, C. E. R., Lemes, M. R., Liesenberg, V., Loyola, R., Macedo, M. N., de Souza Mendes, F., do Couto de Miranda, S., Morton, D. C., Moura, Y. M., Oldekop, J. A., Ramos-Neto, M. B., Rosan, T. M., Saatchi, S., Sano, E. E., Segura-Garcia, C., Shimbo, J. Z., Silva, T. S. F., Trevisan, D. P., Zimbres, B., Wiederkehr, N. C., and Silva-Junior, C. H. L.: Overlooking vegetation loss outside forests imperils the Brazilian Cerrado and other non-forest biomes, *Nature Ecology & Evolution*, 8, 12–13, <https://doi.org/10.1038/s41559-023-02256-w>, 2024.
- 780 da Silva Junior, C. A., Teodoro, P. E., Delgado, R. C., Teodoro, L. P. R., Lima, M., de Andréa Pantaleão, A., Baio, F. H. R., de Azevedo, G. B., de Oliveira Sousa Azevedo, G. T., Capristo-Silva, G. F., Arvor, D., and Facco, C. U.: Persistent fire foci in all biomes undermine the Paris Agreement in Brazil, *Scientific Reports*, 10, 16 246, <https://doi.org/10.1038/s41598-020-72571-w>, 2020.
- 785 de Azevedo, G. B., Rezende, A. V., de Oliveira Sousa Azevedo, G. T., Miguel, E. P., de Gois Aquino, F., Bruzinga, J. S. C., de Oliveira, L. S. C., Pereira, R. S., and Teodoro, P. E.: Woody biomass accumulation in a Cerrado of Central Brazil monitored for 27 years after the implementation of silvicultural systems, *Forest Ecology and Management*, 455, 117 718, <https://doi.org/10.1016/j.foreco.2019.117718>, 2020.
- 790 Deeter, M. N., Edwards, D. P., Francis, G. L., Gille, J. C., Mao, D., Martínez-Alonso, S., Worden, H. M., Ziskin, D., and Andreae, M. O.: Radiance-based retrieval bias mitigation for the MOPITT instrument: the version 8 product, *Atmospheric Measurement Techniques*, 12, 4561–4580, <https://doi.org/10.5194/amt-12-4561-2019>, 2019.
- Dlugokencky, E., Mund, J., Crotwell, A., Crotwell, M., and Thoning, K.: Atmospheric Carbon Dioxide Dry Air Mole Fractions from the NOAA GML Carbon Cycle Cooperative Global Air Sampling Network (1968–2020)., <https://doi.org/10.15138/wkgj-f215>, version: 2021-07-30, 2021.
- 795 EPA: Level III Ecoregions of Central and South America, ftp://ftp.epa.gov/wed/ecoregions/sa/sa_eco.zip, uS Environmental Protection Agency, 2011.
- Eva, H., Huber, O., Frédéric, A., Balslev, H., Beck, S., Behling, H., Belward, A., Beuchle, R., Cleef, A. M., Colchester, M., Duivenvoorden, J., Hoogmoed, M., Junk, W., Kabat, P., Kruijt, B., Malhi, Y., Müller, J., Pereira, J., Peres, C., and Salo, J.: A proposal for defining the geographical boundaries of Amazonia; synthesis of the results from an expert consultation workshop organized by the European Commission in collaboration with the Amazon Cooperation Treaty Organization - JRC Ispra, 7-8 June 2005, 2005.
- 800

- Fawcett, D., Sitch, S., Ciais, P., Wigneron, J. P., Silva-Junior, C. H. L., Heinrich, V., Vancutsem, C., Achard, F., Bastos, A., Yang, H., Li, X., Albergel, C., Friedlingstein, P., and Arag  o, L. E. O. C.: Declining Amazon biomass due to deforestation and subsequent degradation losses exceeding gains, *Global Change Biology*, 29, 1106–1118, <https://doi.org/10.1111/gcb.16513>, 2023.
- 805 Fleischmann, A. S., Papa, F., Fassoni-Andrade, A., Melack, J. M., Wongchuig, S., Paiva, R. C. D., Hamilton, S. K., Fluet-Chouinard, E., Barbedo, R., Aires, F., Al Bitar, A., Bonnet, M.-P., Coe, M., Ferreira-Ferreira, J., Hess, L., Jensen, K., McDonald, K., Ovando, A., Park, E., Parrens, M., Pinel, S., Prigent, C., Resende, A. F., Revel, M., Rosenqvist, A., Rosenqvist, J., Rudorff, C., Silva, T. S. F., Yamazaki, D., and Collischonn, W.: How much inundation occurs in the Amazon River basin?, *Remote Sensing of Environment*, 278, 113 099, <https://doi.org/10.1016/j.rse.2022.113099>, 2022.
- 810 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Qu  r  , C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., G  rses,   , Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., K  rtzinger, A., Landsch  tzer, P., Lauvset, S. K., Lef  vre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., R  denbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., S  f  rian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth System Science Data*, 14, 1917–2005, [https://doi.org/10.5194/essd-14-1917-](https://doi.org/10.5194/essd-14-1917-2022)
- 820 2022, publisher: Copernicus GmbH, 2022.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landsch  tzer, P., Le Qu  r  , C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., G  rses,   , Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarn  kov  , T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., K  rtzinger, A., Lan, X., Lef  vre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., R  denbeck, C., Rosan, T. M., Schwinger, J., S  f  rian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth System Science Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, publisher: Copernicus GmbH, 2023.
- 835 Gatti, L. V., Gloor, M., Miller, J. B., Doughty, C. E., Malhi, Y., Domingues, L. G., Basso, L. S., Martinewski, A., Correia, C. S. C., Borges, V. F., Freitas, S., Braz, R., Anderson, L. O., Rocha, H., Grace, J., Phillips, O. L., and Lloyd, J.: Drought sensitivity of Amazonian carbon balance revealed by atmospheric measurements, *Nature*, 506, 76–80, <https://doi.org/10.1038/nature12957>, 2014.
- Gatti, L. V., Basso, L. S., Miller, J. B., Gloor, M., Gatti Domingues, L., Cassol, H. L. G., Tejada, G., Arag  o, L. E. O. C., Nobre, C., Peters, W., Marani, L., Arai, E., Sanches, A. H., Corr   la, S. M., Anderson, L., Von Randow, C., Correia, C. S. C., Crispim, S. P., and Neves, R.

840 A. L.: Amazonia as a carbon source linked to deforestation and climate change, *Nature*, 595, 388–393, <https://doi.org/10.1038/s41586-021-03629-6>, 2021.

Gatti, L. V., Cunha, C. L., Marani, L., Cassol, H. L. G., Messias, C. G., Arai, E., Denning, A. S., Soler, L. S., Almeida, C., Setzer, A., Domingues, L. G., Basso, L. S., Miller, J. B., Gloor, M., Correia, C. S. C., Tejada, G., Neves, R. A. L., Rajao, R., Nunes, F., Filho, B. S. S., Schmitt, J., Nobre, C., Corrêa, S. M., Sanches, A. H., Aragão, L. E. O. C., Anderson, L., Von Randow, C., Crispim, S. P., Silva, F. M., and Machado, G. B. M.: Increased Amazon carbon emissions mainly from decline in law enforcement, *Nature*, 621, 318–323, 845 <https://doi.org/10.1038/s41586-023-06390-0>, 2023.

Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *Journal of Geophysical Research: Biogeosciences*, 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.

Gloor, E., Wilson, C., Chipperfield, M. P., Chevallier, F., Buermann, W., Boesch, H., Parker, R., Somkuti, P., Gatti, L. V., Correia, C., 850 Domingues, L. G., Peters, W., Miller, J., Deeter, M. N., and Sullivan, M. J. P.: Tropical land carbon cycle responses to 2015/16 El Niño as recorded by atmospheric greenhouse gas and remote sensing data, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20170302, <https://doi.org/10.1098/rstb.2017.0302>, publisher: Royal Society, 2018.

Goulding, M., Barthelmie, R., and Ferreira, E.: *The Smithsonian atlas of the Amazon*, 2003.

Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I. Y., 855 Gloor, M., Heimann, M., Higuchi, K., John, J., Maki, T., Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B. C., Randerson, J., Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Towards robust regional estimates of CO₂ sources and sinks using atmospheric transport models, *Nature*, 415, 626–630, <https://doi.org/10.1038/415626a>, 2002.

Hastie, A., Lauerwald, R., Ciais, P., and Regnier, P.: Aquatic carbon fluxes dampen the overall variation of net ecosystem productivity in the Amazon basin: An analysis of the interannual variability in the boundless carbon cycle, *Global Change Biology*, 25, 2094–2111, 860 <https://doi.org/https://doi.org/10.1111/gcb.14620>, 2019.

Haynes, K. D., Baker, I. T., Denning, A. S., Stöckli, R., Schaefer, K., Lokupitiya, E. Y., and Haynes, J. M.: Representing Grasslands Using Dynamic Prognostic Phenology Based on Biological Growth Stages: 1. Implementation in the Simple Biosphere Model (SiB4), *Journal of Advances in Modeling Earth Systems*, 11, 4423–4439, <https://doi.org/10.1029/2018MS001540>, 2019.

Heimann, M. and Körner, S.: *The global atmospheric tracer model TM3*, Tech. Rep. Tech. Rep. 5., MPI-BGC, Jena (Germany), 2003.

865 Heinrich, V. H. A., Dalagnol, R., Cassol, H. L. G., Rosan, T. M., de Almeida, C. T., Silva Junior, C. H. L., Campanharo, W. A., House, J. I., Sitch, S., Hales, T. C., Adami, M., Anderson, L. O., and Aragão, L. E. O. C.: Large carbon sink potential of secondary forests in the Brazilian Amazon to mitigate climate change, *Nature Communications*, 12, 1785, <https://doi.org/10.1038/s41467-021-22050-1>, 2021.

Heinrich, V. H. A., Vancutsem, C., Dalagnol, R., Rosan, T. M., Fawcett, D., Silva-Junior, C. H. L., Cassol, H. L. G., Achard, F., Jucker, T., Silva, C. A., House, J., Sitch, S., Hales, T. C., and Aragão, L. E. O. C.: The carbon sink of secondary and degraded humid tropical forests, 870 *Nature*, 615, 436–442, <https://doi.org/10.1038/s41586-022-05679-w>, 2023.

Hubau, W., Lewis, S. L., Phillips, O. L., Affum-Baffoe, K., Breeckman, H., Cuní-Sánchez, A., Daniels, A. K., Ewango, C. E. N., Fauset, S., Mukinzi, J. M., Sheil, D., Sonké, B., Sullivan, M. J. P., Sunderland, T. C. H., Taedoumg, H., Thomas, S. C., White, L. J. T., Abernethy, K. A., Adu-Bredu, S., Amani, C. A., Baker, T. R., Banin, L. F., Baya, F., Begne, S. K., Bennett, A. C., Benedet, F., Bitariho, R., Bocko, Y. E., Boeckx, P., Boundja, P., Brienen, R. J. W., Brncic, T., Chezeaux, E., Chuyong, G. B., Clark, C. J., Collins, M., Comiskey, J. A., 875 Coomes, D. A., Dargie, G. C., de Haulleville, T., Kamdem, M. N. D., Doucet, J.-L., Esquivel-Muelbert, A., Feldpausch, T. R., Fofanah, A., Foli, E. G., Gilpin, M., Gloor, E., Gonmadje, C., Gourlet-Fleury, S., Hall, J. S., Hamilton, A. C., Harris, D. J., Hart, T. B., Hockemba,

- M. B. N., Hladik, A., Ifo, S. A., Jeffery, K. J., Jucker, T., Yakusu, E. K., Kearsley, E., Kenfack, D., Koch, A., Leal, M. E., Levesley, A., Lindsell, J. A., Lisingo, J., Lopez-Gonzalez, G., Lovett, J. C., Makana, J.-R., Malhi, Y., Marshall, A. R., Martin, J., Martin, E. H., Mbayu, F. M., Medjibe, V. P., Mihindou, V., Mitchard, E. T. A., Moore, S., Munishi, P. K. T., Bengone, N. N., Ojo, L., Ondo, F. E., Peh, K. S.-H., Pickavance, G. C., Poulsen, A. D., Poulsen, J. R., Qie, L., Reitsma, J., Rovero, F., Swaine, M. D., Talbot, J., Taplin, J., Taylor, D. M., Thomas, D. W., Toirambe, B., Mukendi, J. T., Tuagben, D., Umunay, P. M., van der Heijden, G. M. F., Verbeeck, H., Vleminckx, J., Willcock, S., Wöll, H., Woods, J. T., and Zemagho, L.: Asynchronous carbon sink saturation in African and Amazonian tropical forests, *Nature*, 579, 80–87, <https://doi.org/10.1038/s41586-020-2035-0>, 2020.
- Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *Journal of Geophysical Research: Biogeosciences*, 116, <https://doi.org/10.1029/2010JG001566>, publisher: John Wiley & Sons, Ltd, 2011.
- Jung, M., Reichstein, M., Schwalm, C. R., Huntingford, C., Sitch, S., Ahlström, A., Arneth, A., Camps-Valls, G., Ciais, P., Friedlingstein, P., Gans, F., Ichii, K., Jain, A. K., Kato, E., Papale, D., Poulter, B., Raduly, B., Rödenbeck, C., Tramontana, G., Viovy, N., Wang, Y.-P., Weber, U., Zaehle, S., and Zeng, N.: Compensatory water effects link yearly global land CO₂ sink changes to temperature, *Nature*, 541, 516–520, <https://doi.org/10.1038/nature20780>, 2017.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, B., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project., *Bulletin of the American Meteorological Society*, 77, 437–472, 1996.
- Keeling, C. D. and Revelle, R.: Effects of EL Nino/Southern Oscillation on the Atmospheric Content of Carbon Dioxide, *Meteoritics*, 20, 437, aDS Bibcode: 1985Metic...20..437K, 1985.
- Keeling, C. D., Whorf, T. P., Wahlen, M., and van der Plichtt, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, 375, 666–670, <https://doi.org/10.1038/375666a0>, publisher: Nature Publishing Group, 1995.
- Koren, G.: Constraining the exchange of carbon dioxide over the Amazon: New insights from stable isotopes, remote sensing and inverse modeling., Ph.D. thesis, Wageningen Univeristy, <https://doi.org/10.18174/524771>, 2020.
- Kountouris, P., Gerbig, C., Rödenbeck, C., Karstens, U., Koch, T. F., and Heimann, M.: Atmospheric CO₂ inversions on the mesoscale using data-driven prior uncertainties: quantification of the European terrestrial CO₂ fluxes, *Atmospheric Chemistry and Physics*, 18, 3047–3064, <https://doi.org/https://doi.org/10.5194/acp-18-3047-2018>, 2018a.
- Kountouris, P., Gerbig, C., Rödenbeck, C., Karstens, U., Koch, T. F., and Heimann, M.: Technical Note: Atmospheric CO₂ inversions on the mesoscale using data-driven prior uncertainties: methodology and system evaluation, *Atmospheric Chemistry and Physics*, 18, 3027–3045, <https://doi.org/10.5194/acp-18-3027-2018>, 2018b.
- Krol, M., Houweling, S., Bregman, B., and Bergamaschi, P.: The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmos. Chem. Phys.*, p. 16, 2005.

- Kruid, S., Macedo, M. N., Gorelik, S. R., Walker, W., Moutinho, P., Brando, P. M., Castanho, A., Alencar, A., Baccini, A., and Coe, M. T.: Beyond Deforestation: Carbon Emissions From Land Grabbing and Forest Degradation in the Brazilian Amazon, *Frontiers in Forests and Global Change*, 4, 2021.
- Lapola, D. M., Pinho, P., Barlow, J., Aragão, L. E. O. C., Berenguer, E., Carmenta, R., Liddy, H. M., Seixas, H., Silva, C. V. J., Silva-Junior, C. H. L., Alencar, A. A. C., Anderson, L. O., Armenteras, D., Brovkin, V., Calters, K., Chambers, J., Chini, L., Costa, M. H., Faria, B. L., Fearnside, P. M., Ferreira, J., Gatti, L., Gutierrez-Velez, V. H., Han, Z., Hibbard, K., Koven, C., Lawrence, P., Pongratz, J., Portela, B. T. T., Rounsevell, M., Ruane, A. C., Schaldach, R., da Silva, S. S., von Randow, C., and Walker, W. S.: The drivers and impacts of Amazon forest degradation, *Science*, 379, eabp8622, <https://doi.org/10.1126/science.abp8622>, 2023.
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-i., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth System Science Data*, 10, 2141–2194, <https://doi.org/10.5194/essd-10-2141-2018>, 2018.
- Leal, I. R., Da Silva, J. M. C., Tabarelli, M., and Lacher Jr., T. E.: Changing the Course of Biodiversity Conservation in the Caatinga of Northeastern Brazil, *Conservation Biology*, 19, 701–706, <https://doi.org/10.1111/j.1523-1739.2005.00703.x>, 2005.
- Lin, J. C., Gerbig, C., Wofsy, S. C., Andrews, A. E., Daube, B., Davis, K., and Grainger, C. A.: A near-field tool for simulating the upstream influence of atmospheric observations: The Stochastic Time-Inverted Lagrangian Transport (STILT) model, *Journal of Geophysical Research*, 108, ACH 2–1–ACH 2–17, <https://doi.org/10.1029/2002JD003161>, 2003.
- Liu, J., Bowman, K. W., Schimel, D. S., Parazoo, N. C., Jiang, Z., Lee, M., Bloom, A. A., Wunch, D., Frankenberg, C., Sun, Y., O'Dell, C. W., Gurney, K. R., Menemenlis, D., Gierach, M., Crisp, D., and Eldering, A.: Contrasting carbon cycle responses of the tropical continents to the 2015–2016 El Niño, *Science*, 358, <https://doi.org/10.1126/science.aam5690>, 2017.
- Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., Gerbig, C., Munger, J. W., Chow, V. Y., and Gottlieb, E. W.: A satellite-based biosphere parameterization for net ecosystem CO₂ exchange: Vegetation Photosynthesis and Respiration Model (VPRM), *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2006GB002735>, 2008.
- Malhi, Y., Saatchi, S., Girardin, C., and Aragão, L. E. O. C.: The Production, Storage, and Flow of Carbon in Amazonian Forests, in: *Amazonia and Global Change*, pp. 355–372, American Geophysical Union (AGU), ISBN 978-1-118-67034-7, 2009.
- MapBiomass: Collection 5.0 of the annual series of land use and land cover maps of Brazil., <https://brasil.mapbiomas.org/>, 2020.
- Marengo, J. A., Jimenez, J. C., Espinoza, J.-C., Cunha, A. P., and Aragão, L. E. O.: Increased climate pressure on the agricultural frontier in the Eastern Amazonia-Cerrado transition zone, *Scientific Reports*, 12, 457, <https://doi.org/10.1038/s41598-021-04241-4>, 2022.
- Marques, J. D. d. O., Luizão, F. J., Teixeira, W. G., Nogueira, E. M., Fearnside, P. M., and Sarrazin, M.: Soil Carbon Stocks under Amazonian Forest: Distribution in the Soil Fractions and Vulnerability to Emission, *Open Journal of Forestry*, 7, 121–142, <https://doi.org/10.4236/ojf.2017.72008>, 2017.

- Masarie, K. A., Pétron, G., Andrews, A., Bruhwiler, L., Conway, T. J., Jacobson, A. R., Miller, J. B., Tans, P. P., Worthy, D. E., and Peters, W.: Impact of CO₂ measurement bias on CarbonTracker surface flux estimates, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016270>, 2011.
- Massie, S. T., Sebastian Schmidt, K., Eldering, A., and Crisp, D.: Observational evidence of 3-D cloud effects in OCO-2 CO₂ retrievals, *Journal of Geophysical Research: Atmospheres*, 122, 7064–7085, <https://doi.org/10.1002/2016JD026111>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016JD026111>, 2017.
- Mataveli, G., Jones, M. W., Carmenta, R., Sanchez, A., Dutra, D. J., Chaves, M., de Oliveira, G., Anderson, L. O., and Aragão, L. E. O. C.: Deforestation falls but rise of wildfires continues degrading Brazilian Amazon forests, *Global Change Biology*, 30, e17202, <https://doi.org/10.1111/gcb.17202>, 2024.
- Matricardi, E. A. T., Skole, D. L., Costa, O. B., Pedlowski, M. A., Samek, J. H., and Miguel, E. P.: Long-term forest degradation surpasses deforestation in the Brazilian Amazon, *Science*, 369, 1378–1382, <https://doi.org/10.1126/science.abb3021>, 2020.
- Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmospheric Chemistry and Physics*, 8, 6341–6353, <https://doi.org/10.5194/acp-8-6341-2008>, 2008.
- Mendes, K. R., Campos, S., da Silva, L. L., Mutti, P. R., Ferreira, R. R., Medeiros, S. S., Perez-Marin, A. M., Marques, T. V., Ramos, T. M., de Lima Vieira, M. M., Oliveira, C. P., Gonçalves, W. A., Costa, G. B., Antonino, A. C. D., Menezes, R. S. C., Bezerra, B. G., and Santos e Silva, C. M.: Seasonal variation in net ecosystem CO₂ exchange of a Brazilian seasonally dry tropical forest, *Scientific Reports*, 10, 9454, <https://doi.org/10.1038/s41598-020-66415-w>, 2020.
- Miller, J. B., Martins, G. A., de Souza, R. A. F., and Schuldt, K. N.: Manaus aircraft profile data for the period 2017-2023; obspack_multi-species_1_manauas_profiles_v2.0_2023-09-26; NOAA Global Monitoring Laboratory, <https://doi.org/http://dx.doi.org/10.25925/20230922>, 2023.
- Miranda, A. C., Miranda, H. S., Lloyd, J., Grace, J., Francey, R. J., McIntyre, J. A., Meir, P., Riggan, P., Lockwood, R., and Brass, J.: Fluxes of carbon, water and energy over Brazilian cerrado: an analysis using eddy covariance and stable isotopes, *Plant, Cell & Environment*, 20, 315–328, <https://doi.org/10.1046/j.1365-3040.1997.d01-80.x>, 1997.
- Molina, L., Broquet, G., Imbach, P., Chevallier, F., Poulter, B., Bonal, D., Burban, B., Ramonet, M., Gatti, L. V., Wofsy, S. C., Munger, J. W., Dlugokencky, E., and Ciais, P.: On the ability of a global atmospheric inversion to constrain variations of CO₂ fluxes over Amazonia, *Atmospheric Chemistry and Physics*, 15, 8423–8438, <https://doi.org/10.5194/acp-15-8423-2015>, 2015.
- Munassar, S., Rödenbeck, C., Koch, T., Totsche, K. U., Galkowski, M., Walther, S., and Gerbig, C.: NEE estimates 2006-2019 over Europe from a pre-operational ensemble-inversion system, *Atmospheric Chemistry and Physics Discussions*, pp. 1–28, <https://doi.org/10.5194/acp-2021-873>, 2021.
- Munassar, S., Monteil, G., Scholze, M., Karstens, U., Rödenbeck, C., Koch, F.-T., Totsche, K. U., and Gerbig, C.: Why do inverse models disagree? A case study with two European CO₂ inversions, *Atmospheric Chemistry and Physics*, 23, 2813–2828, <https://doi.org/10.5194/acp-23-2813-2023>, 2023.
- Munassar, S., Roedenbeck, C., Gańkowski, M., Koch, F.-T., Totsche, K. U., Botía, S., and Gerbig, C.: To what extent does CO₂ diurnal cycle impact carbon flux estimates in CarboScope?, *EGUsphere*, pp. 1–24, <https://doi.org/10.5194/egusphere-2024-291>, publisher: Copernicus GmbH, 2024.
- Naus, S.: Improving estimates of the atmospheric oxidative capacity and Amazon fire emissions, Ph.D. thesis, Wageningen Univeristy, <https://doi.org/10.18174/536720>, 2021.

- Naus, S., Domingues, L. G., Krol, M., Luijkx, I. T., Gatti, L. V., Miller, J. B., Gloor, E., Basu, S., Correia, C., Koren, G., Worden, H. M., Fleming, J., Pétron, G., and Peters, W.: Sixteen years of MOPITT satellite data strongly constrain Amazon CO fire emissions, *Atmospheric Chemistry and Physics*, 22, 14 735–14 750, <https://doi.org/10.5194/acp-22-14735-2022>, 2022.
- 990 Nelson, J. A., Walther, S., Gans, F., Kraft, B., Weber, U., Novick, K., Buchmann, N., Migliavacca, M., Wohlfahrt, G., Åälgut, L., Ibrom, A., Papale, D., Göckede, M., Duveiller, G., Knohl, A., Hörtnagl, L., Scott, R. L., Zhang, W., Hamdi, Z. M., Reichstein, M., Aranda-Barranco, S., Ardö, J., Op de Beeck, M., Billdesbach, D., Bowling, D., Bracho, R., Brümmer, C., Camps-Valls, G., Chen, S., Cleverly, J. R., Desai, A., Dong, G., El-Madany, T. S., Euskirchen, E. S., Feigenwinter, I., Galvagno, M., Gerosa, G., Gielen, B., Goded, I., Goslee, S., Gough, C. M., Heinesch, B., Ichii, K., Jackowicz-Korczynski, M. A., Klosterhalfen, A., Knox, S., Kobayashi, H., Kohonen, K.-M., Korkiakoski,
- 995 M., Mammarella, I., Mana, G., Marzuoli, R., Matamala, R., Metzger, S., Montagnani, L., Nicolini, G., O'Halloran, T., Ourcival, J.-M., Peichl, M., Pendall, E., Ruiz Reverter, B., Roland, M., Sabbatini, S., Sachs, T., Schmidt, M., Schwalm, C. R., Shekhar, A., Silberstein, R., Silveira, M. L., Spano, D., Tagesson, T., Tramontana, G., Trotta, C., Turco, F., Vesala, T., Vincke, C., Vitale, D., Vivoni, E. R., Wang, Y., Woodgate, W., Yopez, E. A., Zhang, J., Zona, D., and Jung, M.: X-BASE: the first terrestrial carbon and water flux products from an extended data-driven scaling framework, *FLUXCOM-X*, *EGUsphere*, pp. 1–51, <https://doi.org/10.5194/egusphere-2024-165>, publisher: Copernicus GmbH, 2024.
- 1000 Noojipady, P., Morton, C. D., Macedo, N. M., Victoria, C. D., Huang, C., Gibbs, K. H., and Bolfe, L. E.: Forest carbon emissions from cropland expansion in the Brazilian Cerrado biome, *Environmental Research Letters*, 12, 025 004, <https://doi.org/10.1088/1748-9326/aa5986>, publisher: IOP Publishing, 2017.
- Palmer, P. I., Feng, L., Baker, D., Chevallier, F., Bösch, H., and Somkuti, P.: Net carbon emissions from African biosphere dominate pan-tropical atmospheric CO₂ signal, *Nature Communications*, 10, 3344, <https://doi.org/10.1038/s41467-019-11097-w>, 2019.
- 1005 Paul, D., Scheeren, H. A., Jansen, H. G., Kers, B. A. M., Miller, J. B., Crotwell, A. M., Michel, S. E., Gatti, L. V., Domingues, L. G., Correia, C. S. C., Neves, R. A. L., Meijer, H. A. J., and Peters, W.: Evaluation of a field-deployable Nafion-based air-drying system for collecting whole air samples and its application to stable isotope measurements of CO₂, *Atmospheric Measurement Techniques*, 13, 4051–4064, <https://doi.org/10.5194/amt-13-4051-2020>, publisher: Copernicus GmbH, 2020.
- 1010 Peiro, H., Crowell, S., Schuh, A., Baker, D. F., O'Dell, C., Jacobson, A. R., Chevallier, F., Liu, J., Eldering, A., Crisp, D., Deng, F., Weir, B., Basu, S., Johnson, M. S., Philip, S., and Baker, I.: Four years of global carbon cycle observed from the Orbiting Carbon Observatory 2 (OCO-2) version 9 and in situ data and comparison to OCO-2 version 7, *Atmospheric Chemistry and Physics*, 22, 1097–1130, <https://doi.org/10.5194/acp-22-1097-2022>, publisher: Copernicus GmbH, 2022.
- Peylin, P., Law, R. M., Gurney, K. R., Chevallier, F., Jacobson, A. R., Maki, T., Niwa, Y., Patra, P. K., Peters, W., Rayner, P. J., Rödenbeck, C.,
- 1015 van der Laan-Luijkx, I. T., and Zhang, X.: Global atmospheric carbon budget: results from an ensemble of atmospheric CO₂ inversions, *Biogeosciences*, 10, 6699–6720, <https://doi.org/https://doi.org/10.5194/bg-10-6699-2013>, publisher: Copernicus GmbH, 2013.
- Piao, S., Wang, X., Wang, K., Li, X., Bastos, A., Canadell, J. G., Ciais, P., Friedlingstein, P., and Sitch, S.: Interannual variation of terrestrial carbon cycle: Issues and perspectives, *Global Change Biology*, 26, 300–318, <https://doi.org/10.1111/gcb.14884>, 2020.
- Pivello, V. R.: The Use of Fire in the Cerrado and Amazonian Rainforests of Brazil: Past and Present, *Fire Ecology*, 7, 24–39, <https://doi.org/10.4996/fireecology.0701024>, 2011.
- 1020 Pletsch, M. A. J. S., Körting, T. S., Morita, F. C., Silva-Junior, C. H. L., Anderson, L. O., and Aragão, L. E. O. C.: Near Real-Time Fire Detection and Monitoring in the MATOPIBA Region, Brazil, *Remote Sensing*, 14, 3141, <https://doi.org/10.3390/rs14133141>, 2022.
- Poorter, L., Bongers, F., Aide, T. M., Almeyda Zambrano, A. M., Balvanera, P., Becknell, J. M., Boukili, V., Brancalion, P. H. S., Broadbent, E. N., Chazdon, R. L., Craven, D., de Almeida-Cortez, J. S., Cabral, G. A. L., de Jong, B. H. J., Denslow, J. S., Dent, D. H., DeWalt, S. J.,

- 1025 Dupuy, J. M., Durán, S. M., Espírito-Santo, M. M., Fandino, M. C., César, R. G., Hall, J. S., Hernandez-Stefanoni, J. L., Jakovac, C. C., Junqueira, A. B., Kennard, D., Letcher, S. G., Licona, J.-C., Lohbeck, M., Marín-Spiotta, E., Martínez-Ramos, M., Massoca, P., Meave, J. A., Mesquita, R., Mora, F., Muñoz, R., Muscarella, R., Nunes, Y. R. F., Ochoa-Gaona, S., de Oliveira, A. A., Orihuela-Belmonte, E., Peña-Claros, M., Pérez-García, E. A., Piotto, D., Powers, J. S., Rodríguez-Velázquez, J., Romero-Pérez, I. E., Ruíz, J., Saldarriaga, J. G., Sanchez-Azofeifa, A., Schwartz, N. B., Steininger, M. K., Swenson, N. G., Toledo, M., Uriarte, M., van Breugel, M., van der Wal, H.,
- 1030 Veloso, M. D. M., Vester, H. F. M., Vicentini, A., Vieira, I. C. G., Bentos, T. V., Williamson, G. B., and Rozendaal, D. M. A.: Biomass resilience of Neotropical secondary forests, *Nature*, 530, 211–214, <https://doi.org/10.1038/nature16512>, 2016.
- Poorter, L., Craven, D., Jakovac, C. C., van der Sande, M. T., Amissah, L., Bongers, F., Chazdon, R. L., Farrior, C. E., Kambach, S., Meave, J. A., Muñoz, R., Norden, N., Rüger, N., van Breugel, M., Almeyda Zambrano, A. M., Amani, B., Andrade, J. L., Brancalion, P. H. S., Broadbent, E. N., de Foresta, H., Dent, D. H., Derroire, G., DeWalt, S. J., Dupuy, J. M., Durán, S. M., Fantini, A. C., Finegan, B., Hernández-Jaramillo, A., Hernández-Stefanoni, J. L., Hietz, P., Junqueira, A. B., N'dja, J. K., Letcher, S. G., Lohbeck, M., López-Camacho, R., Martínez-Ramos, M., Melo, F. P. L., Mora, F., Müller, S. C., N'Guessan, A. E., Oberleitner, F., Ortiz-Malavassi, E., Pérez-García, E. A., Pinho, B. X., Piotto, D., Powers, J. S., Rodríguez-Buriticá, S., Rozendaal, D. M. A., Ruíz, J., Tabarelli, M., Teixeira, H. M., Valadares de Sá Barretto Sampaio, E., van der Wal, H., Villa, P. M., Fernandes, G. W., Santos, B. A., Aguilar-Cano, J., de Almeida-Cortez, J. S., Alvarez-Davila, E., Arreola-Villa, F., Balvanera, P., Becknell, J. M., Cabral, G. A. L., Castellanos-Castro, C., de Jong, B. H. J., Nieto, J. E., Espírito-Santo, M. M., Fandino, M. C., García, H., García-Villalobos, D., Hall, J. S., Idárraga, A., Jiménez-Montoya, J., Kennard, D., Marín-Spiotta, E., Mesquita, R., Nunes, Y. R. F., Ochoa-Gaona, S., Peña-Claros, M., Pérez-Cárdenas, N., Rodríguez-Velázquez, J., Villanueva, L. S., Schwartz, N. B., Steininger, M. K., Veloso, M. D. M., Vester, H. F. M., Vieira, I. C. G., Williamson, G. B., Zanini, K., and Hérault, B.: Multidimensional tropical forest recovery, *Science*, 374, 1370–1376, <https://doi.org/10.1126/science.abh3629>, 2021.
- 1040 Prado, D.: As caatingas da América do Sul, in: *Ecologia e Biogeografia da Caatinga*, pp. 3–73, Universidade Federal de Pernambuco, inara leal, marcelo tabarelli, j.m.c. silva edn., 2003.
- 1045 Qin, Y., Xiao, X., Wigneron, J.-P., Ciais, P., Brandt, M., Fan, L., Li, X., Crowell, S., Wu, X., Dougherty, R., Zhang, Y., Liu, F., Sitch, S., and Moore, B.: Carbon loss from forest degradation exceeds that from deforestation in the Brazilian Amazon, *Nature Climate Change*, 11, 442–448, <https://doi.org/10.1038/s41558-021-01026-5>, 2021.
- Ramo, R., Roteta, E., Bistinas, I., van Wees, D., Bastarrika, A., Chuvieco, E., and van der Werf, G. R.: African burned area and fire carbon emissions are strongly impacted by small fires undetected by coarse resolution satellite data, *Proceedings of the National Academy of Sciences*, 118, e2011160118, <https://doi.org/10.1073/pnas.2011160118>, 2021.
- 1050 Ramos-Neto, M. B. and Pivello, V. R.: Lightning Fires in a Brazilian Savanna National Park: Rethinking Management Strategies, *Environmental Management*, 26, 675–684, <https://doi.org/10.1007/s002670010124>, 2000.
- Ratter, J. A., Ribeiro, J. F., and Bridgewater, S.: The Brazilian cerrado vegetation and threats to its biodiversity, *Annals of Botany (United Kingdom)*, 1997.
- 1055 Rella, C. W., Chen, H., Andrews, A. E., Filges, A., Gerbig, C., Hatakka, J., Karion, A., Miles, N. L., Richardson, S. J., Steinbacher, M., Sweeney, C., Wastine, B., and Zellweger, C.: High accuracy measurements of dry mole fractions of carbon dioxide and methane in humid air, *Atmospheric Measurement Techniques*, 6, 837–860, <https://doi.org/10.5194/amt-6-837-2013>, publisher: Copernicus GmbH, 2013.
- Ribeiro, K., Sousa-Neto, E. R. d., Carvalho, J. A. d., Sousa Lima, J. R. d., Menezes, R. S. C., Duarte-Neto, P. J., da Silva Guerra, G., and Ometto, J. P. H. B.: Land cover changes and greenhouse gas emissions in two different soil covers in the Brazilian Caatinga, *Science of The Total Environment*, 571, 1048–1057, <https://doi.org/10.1016/j.scitotenv.2016.07.095>, 2016.
- 1060

- Rödenbeck, C.: Estimating CO₂ sources and sinks from atmospheric mixing ratio measurements using a global inversion of atmospheric transport, Tech. rep., Max Planck Institute for Biogeochemistry, <https://www.bgc-jena.mpg.de/~christian.roedenbeck/download/2005-Roedenbeck-TechReport6.pdf>, 2005.
- 1065 Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: CO₂ flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport, *Atmos. Chem. Phys.*, p. 46, 2003.
- Rödenbeck, C., Gerbig, C., Trusilova, K., and Heimann, M.: A two-step scheme for high-resolution regional atmospheric trace gas inversions based on independent models, *Atmos. Chem. Phys.*, p. 12, 2009.
- Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean p^{CO_2} and sea-air
 1070 CO₂ flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Science*, 9, 193–216, <https://doi.org/10.5194/os-9-193-2013>, publisher: Copernicus GmbH, 2013.
- Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: History of El Niño impacts on the global carbon cycle 1957–2017: a quantification from atmospheric CO₂ data, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20170303, <https://doi.org/10.1098/rstb.2017.0303>, 2018a.
- 1075 Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO₂ data, *Biogeosciences*, 15, 2481–2498, <https://doi.org/10.5194/bg-15-2481-2018>, 2018b.
- Rodrigues, A. A., Macedo, M. N., Silvério, D. V., Maracahipes, L., Coe, M. T., Brando, P. M., Shimbo, J. Z., Rajão, R., Soares-Filho, B., and Bustamante, M. M. C.: Cerrado deforestation threatens regional climate and water availability for agriculture and ecosystems, *Global Change Biology*, 28, 6807–6822, <https://doi.org/10.1111/gcb.16386>, 2022.
- 1080 Rosan, T. M., Sitch, S., O’Sullivan, M., Basso, L. S., Wilson, C., Silva, C., Gloor, E., Fawcett, D., Heinrich, V., Souza, J. G., Bezerra, F. G. S., von Randow, C., Mercado, L. M., Gatti, L., Wiltshire, A., Friedlingstein, P., Pongratz, J., Schwingshackl, C., Williams, M., Smallman, L., Knauer, J., Arora, V., Kennedy, D., Tian, H., Yuan, W., Jain, A. K., Falk, S., Poulter, B., Arneth, A., Sun, Q., Zaehle, S., Walker, A. P., Kato, E., Yue, X., Bastos, A., Ciais, P., Wigneron, J.-P., Albergel, C., and Aragão, L. E. O. C.: Synthesis of the land carbon fluxes of the Amazon region between 2010 and 2020, *Communications Earth & Environment*, 5, 1–15, <https://doi.org/10.1038/s43247-024-01205-0>,
 1085 2024.
- Saatchi, S. S., Houghton, R. A., Alvalá, R. C. D. S., Soares, J. V., and Yu, Y.: Distribution of aboveground live biomass in the Amazon basin, *Global Change Biology*, 13, 816–837, <https://doi.org/https://doi.org/10.1111/j.1365-2486.2007.01323.x>, 2007.
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents,
 1090 *Proceedings of the National Academy of Sciences*, 108, 9899–9904, <https://doi.org/10.1073/pnas.1019576108>, publisher: Proceedings of the National Academy of Sciences, 2011.
- Sakschewski, B., von Bloh, W., Boit, A., Poorter, L., Peña-Claros, M., Heinke, J., Joshi, J., and Thonicke, K.: Resilience of Amazon forests emerges from plant trait diversity, *Nature Climate Change*, 6, 1032–1036, <https://doi.org/10.1038/nclimate3109>, 2016.
- Saleska, S. R., da Rocha, H., Huete, A., Nobre, A. D., Artaxo, P., and Shimabukuro, Y.: LBA-ECO CD-32 Flux Tower Network Data
 1095 Compilation, Brazilian Amazon: 1999–2006., <https://doi.org/http://dx.doi.org/10.3334/ORNLDAAC/1174>, 2013.
- Sano, E. E., Rosa, R., Brito, J. L. S., and Ferreira, L. G.: Mapeamento de Cobertura Vegetal do Bioma Cerrado: estratégias e resultados, Tech. rep., EMBRAPA, 2007.

- Santos, A. J., Quesada, C. A., Da Silva, G. T., Maia, J. F., Miranda, H. S., Carlos Miranda, A., and Lloyd, J.: High rates of net ecosystem carbon assimilation by Brachiara pasture in the Brazilian Cerrado, *Global Change Biology*, 10, 877–885, <https://doi.org/10.1111/j.1529-8817.2003.00777.x>, 2004.
- Schaefer, K., Collatz, G. J., Tans, P., Denning, A. S., Baker, I., Berry, J., Prihodko, L., Suits, N., and Philpott, A.: Combined Simple Biosphere/Carnegie-Ames-Stanford Approach terrestrial carbon cycle model, *Journal of Geophysical Research*, 113, <https://doi.org/10.1029/2007JG000603>, 2008.
- Schuh, A. E., Jacobson, A. R., Basu, S., Weir, B., Baker, D., Bowman, K., Chevallier, F., Crowell, S., Davis, K. J., Deng, F., Denning, S., Feng, L., Jones, D., Liu, J., and Palmer, P. I.: Quantifying the Impact of Atmospheric Transport Uncertainty on CO₂ Surface Flux Estimates, *Global Biogeochemical Cycles*, 33, 484–500, <https://doi.org/10.1029/2018GB006086>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GB006086>, 2019.
- Science Panel for the Amazon: Executive Summary of the Amazon Assessment Report 2021, p. 48, UN Sustainable Development Solutions Network (SDSN), New York, USA, 2021.
- Silva, P. F. d., Lima, J. R. d. S., Antonino, A. C. D., Souza, R., Souza, E. S. d., Silva, J. R. I., and Alves, E. M.: Seasonal patterns of carbon dioxide, water and energy fluxes over the Caatinga and grassland in the semi-arid region of Brazil, *Journal of Arid Environments*, 147, 71–82, <https://doi.org/10.1016/j.jaridenv.2017.09.003>, 2017.
- Silva Junior, C. H. L., Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Shimabukuro, Y. E., Vancutsem, C., Achard, F., Beuchle, R., Numata, I., Silva, C. A., Maeda, E. E., Longo, M., and Saatchi, S. S.: Persistent collapse of biomass in Amazonian forest edges following deforestation leads to unaccounted carbon losses, *Science Advances*, 6, eaaz8360, <https://doi.org/10.1126/sciadv.aaz8360>, 2020.
- Silva Junior, C. H. L., Pessôa, A. C. M., Carvalho, N. S., Reis, J. B. C., Anderson, L. O., and Aragão, L. E. O. C.: The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade, *Nature Ecology & Evolution*, 5, 144–145, <https://doi.org/10.1038/s41559-020-01368-x>, 2021.
- Spera, S. A., Galford, G. L., Coe, M. T., Macedo, M. N., and Mustard, J. F.: Land-use change affects water recycling in Brazil’s last agricultural frontier, *Global Change Biology*, 22, 3405–3413, <https://doi.org/10.1111/gcb.13298>, 2016.
- Steinbach, J., Gerbig, C., Rödenbeck, C., Karstens, U., Minejima, C., and Mukai, H.: The CO₂ release and Oxygen uptake from Fossil Fuel Emission Estimate (COFFEE) dataset: effects from varying oxidative ratios, *Atmospheric Chemistry and Physics*, 11, 6855–6870, <https://doi.org/10.5194/acp-11-6855-2011>, publisher: Copernicus GmbH, 2011.
- Tao, S., Wigneron, J.-P., Chave, J., Tang, Z., Wang, Z., Zhu, J., Guo, Q., Liu, Y. Y., and Ciais, P.: Little evidence that Amazonian rainforests are approaching a tipping point, *Nature Climate Change*, 13, 1317–1320, <https://doi.org/10.1038/s41558-023-01853-8>, 2023.
- Trusilova, K., Rödenbeck, C., Gerbig, C., and Heimann, M.: Technical Note: A new coupled system for global-to-regional downscaling of CO₂ concentration estimation, *Atmos. Chem. Phys.*, p. 9, 2010.
- van der Laan-Luijkx, I. T., van der Velde, I. R., Krol, M. C., Gatti, L. V., Domingues, L. G., Correia, C. S. C., Miller, J. B., Gloor, M., van Leeuwen, T. T., Kaiser, J. W., Wiedinmyer, C., Basu, S., Clerbaux, C., and Peters, W.: Response of the Amazon carbon balance to the 2010 drought derived with CarbonTracker South America, *Global Biogeochemical Cycles*, 29, 1092–1108, <https://doi.org/10.1002/2014GB005082>, 2015.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997-2009), *Atmos. Chem. Phys.*, 10, 11 707–11 735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.

- 1135 van der Werf, G. R., Randerson, J. T., Giglio, L., Leeuwen, T. T. v., Chen, Y., Rogers, B. M., Mu, M., Marle, M. J. E. v., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997-2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/https://doi.org/10.5194/essd-9-697-2017>, publisher: Copernicus GmbH, 2017.
- van Schaik, E., Killaars, L., Smith, N. E., Koren, G., van Beek, L. P. H., Peters, W., and van der Laan-Luijkx, I. T.: Changes in surface hydrology, soil moisture and gross primary production in the Amazon during the 2015/2016 El Niño, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373, 20180084, <https://doi.org/10.1098/rstb.2018.0084>, 2018.
- 1140 van Wees, D., van der Werf, G. R., Randerson, J. T., Rogers, B. M., Chen, Y., Veraverbeke, S., Giglio, L., and Morton, D. C.: Global biomass burning fuel consumption and emissions at 500m spatial resolution based on the Global Fire Emissions Database (GFED), *Geoscientific Model Development*, 15, 8411–8437, <https://doi.org/10.5194/gmd-15-8411-2022>, publisher: Copernicus GmbH, 2022.
- Varela, R. F., Bustamante, M. M. C., Pinto, A. S., Kisselle, K. W., Santos, R. V., Burke, R. A., Zepp, R. G., and Viana, L. T.: Soil 1145 Fluxes of Co₂, Co, No, and N₂o from an Old Pasture and from Native Savanna in Brazil, *Ecological Applications*, 14, 221–231, <https://doi.org/10.1890/01-6014>, 2004.
- Wang, H., Ciais, P., Sitch, S., Green, J. K., Tao, S., Fu, Z., Albergel, C., Bastos, A., Wang, M., Fawcett, D., Frappart, F., Li, X., Liu, X., Li, S., and Wigneron, J.-P.: Anthropogenic disturbance exacerbates resilience loss in the Amazon rainforests, *Global Change Biology*, 30, e17006, <https://doi.org/10.1111/gcb.17006>, 2024.
- 1150 Wang, J., Zeng, N., Wang, M., Jiang, F., Chevallier, F., Crowell, S., He, W., Johnson, M. S., Liu, J., Liu, Z., Miller, S. M., Philip, S., Wang, H., Wu, M., Ju, W., Feng, S., and Jia, M.: Anomalous Net Biome Exchange Over Amazonian Rainforests Induced by the 2015/16 El Niño: Soil Dryness-Shaped Spatial Pattern but Temperature-dominated Total Flux, *Geophysical Research Letters*, 50, e2023GL103379, <https://doi.org/10.1029/2023GL103379>, 2023.
- Wang, W., Ciais, P., Nemani, R. R., Canadell, J. G., Piao, S., Sitch, S., White, M. A., Hashimoto, H., Milesi, C., and Myneni, R. B.: 1155 Variations in atmospheric CO₂ growth rates coupled with tropical temperature, *Proceedings of the National Academy of Sciences*, 110, 13061–13066, <https://doi.org/10.1073/pnas.1219683110>, 2013.
- Wiedinmyer, C., Kimura, Y., McDonald-Buller, E. C., Emmons, L. K., Buchholz, R. R., Tang, W., Seto, K., Joseph, M. B., Barsanti, K. C., Carlton, A. G., and Yokelson, R.: The Fire Inventory from NCAR version 2.5: an updated global fire emissions model for climate and chemistry applications, *Geoscientific Model Development*, 16, 3873–3891, <https://doi.org/10.5194/gmd-16-3873-2023>, publisher: Copernicus GmbH, 2023.
- 1160