



Towards deep learning solutions for classification of automated snow height measurements (CleanSnow v1.0.0)

Jan Svoboda¹, Marc Ruesch¹, David Liechti¹, Corinne Jones², Michele Volpi², Michael Zehnder^{1,3}, and Jürg Schweizer¹

¹WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland

²Swiss Data Science Center, ETH Zürich and EPFL, Zürich, Switzerland

³Institute of Integrative Biology, ETH Zürich, Zürich, Switzerland

Correspondence: Jan Svoboda (jan.svoboda@slf.ch)

Abstract. Snow height measurements are still the backbone of any snow cover monitoring whether based on modeling or remote sensing. These ground-based measurements are often realized with the use of ultrasonic or laser technologies. In challenging environments, such as high alpine regions, the quality of sensor measurements deteriorates quickly, especially in the presence of extreme weather conditions or ephemeral snow conditions. Moreover, the sensors by their nature measure the height of an underlying object and are therefore prone to return other information, such as the height of vegetation, in snow-free periods. Quality assessment and real-time classification of automated snow height measurements is therefore desirable in order to provide high-quality data for research and operational applications. To this end, we propose CleanSnow, a machine learning approach to automated classification of snow height measurements into a snow cover class and a class corresponding to everything else, which takes into account both the temporal context and the dependencies between snow height and other sensor measurements. We created a new dataset of manually annotated snow height measurements, which allowed us to train our models in a supervised manner as well as quantitatively evaluate our results. Through a series of experiments and ablation studies to evaluate feature importance and compare several different models, we validated our design choices and demonstrate the importance of using temporal information together with information from auxiliary sensors. CleanSnow achieved a high accuracy and represents a new baseline for further research in the field. The presented approach to snow height classification finds its use in various tasks, ranging from snow modeling to climate science.

1 Introduction

Snow height measurements are key in many fields, such as water resources management, avalanche forecasting, climate science, or even tourism. A variety of complex models simulating and calculating snowpack properties therefore exist. For example, estimating snow water equivalent (SWE) (e.g. Jonas et al., 2009) to assess water resources. In addition, snow height is an important parameter for snow hydrological (e.g. Mott et al., 2023) and snow cover modeling (Lehning et al., 1999) used in operational avalanche forecasting (Morin et al., 2020; Pérez-Guillén et al., 2022; Herla et al., 2023). In climate science, snow cover is one of the key variables that strongly affect the global energy balance and the atmospheric circulation, due to its high albedo, high emissivity and low thermal conductivity (e.g. Flanner et al., 2011). Snow height signals have also been used to

Summary of Comments on egusphere-2024-1752.pdf

Page: 1

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 11:41:15

How much is this accuracy? This should be in the abstract.

 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:37:29

I believe it's 97.7%



determine vegetation growth and plant phenology (e.g. Jonas et al., 2008; Fontana et al., 2008; Vitasse et al., 2017; Zehnder et al., in prep.) and to monitor climate change (e.g. Matiu et al., 2021). Finally, ¹the snow cover ²situation directly influences tourism, transportation and recreational activities (e.g. Willibald et al., 2021).

Snow height data are nowadays available, sometimes in almost real-time, from airborne or satellite remote sensing and ground-based automated weather stations (AWS). One of the sensors often mounted at meteorological stations in high alpine regions is an ultrasonic snow height sensor (Ryan et al., 2008). Due to the measurement method, snow height data come with a variety of errors that arise from the harsh mountain conditions the sensor is not originally designed to operate in. In addition, ultrasonic sensors only measure the distance to the underlying object, be it snow or anything else. It is therefore important to validate whether the information coming from the snow height sensor really corresponds to snow or not.

Arguably the most precise way of assessing the ³quality (QA) of snow height measurements is via visual inspection of the data ⁴by a human expert (Robinson, 1989). Even though it is believed the most reliable, manual quality assessment of data is a tedious procedure heavily relying on expert knowledge, which is not easily transferable and does not scale well (Fiebrich et al., 2010). ⁵A common practice in snow height ⁶QA is to distinguish between snow and grass based on static climatological or minimum snow height thresholds. Random errors, instead, are typically detected using a maximum snow height threshold or snow height variance (Avanzi et al., 2014).

There are other sensors usually mounted at an AWS, which can provide information on whether the measured snow height relates to snow or not, as well as give some indications on the precision of snow height measurement. The first attempt to leverage other sensor information was the MeteIO library developed by Bavay and Egger (2014), which contains an algorithm for grass detection based on snow surface temperature, ground surface temperature, and solar radiation. The algorithm is based on a series of thresholding rules, an approach that is known to be rather cumbersome to modify and does not generally transfer well to other station data. ⁹Observing the recent advances in machine learning, Blandini et al. (2023) ⁸have decided to deal with the ¹¹high dimensionality of the data ¹⁰by proposing a random forest approach to snow height ¹²QA, solving both snow height classification and anomaly detection at the same time. Random Forest ¹⁴(RF) models (Breiman, 2001) ¹³are possibly the most popular choice of machine learning algorithms used among data scientists worldwide. Apart from an attempt by Goehry et al. (2023), random forests, however, cannot easily and explicitly model the temporal structure of the data ¹⁵that we argue is crucial to be able to reliably say whether the snow height measurement is erroneous and whether the signal coming from the sensor shows snow or vegetation.

¹⁶Therefore, we ¹⁷aim to develop a ¹⁸machine learning model for the automated classification of snow height signals into a snow and a no-snow class, which we call CleanSnow. To approach this binary classification problem, we employed a Temporal Convolutional Network (TCN) (Lea et al., 2016) that explicitly accounts for the temporal relationships between different points in snow height time series data. To train our TCN, we created a new manually annotated snow height dataset composed of 20 measurement stations with around 20 years of data per station. This dataset also allows us to validate our design choices and evaluate the model in several different scenarios including challenging cases such as snow cover melt or plant growth periods.

Page: 2

	Number: 1	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 11:43:01
	Number: 2	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 11:43:05
	Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:48:52
	is this really an important acronym? First it's not straightforward that the acronym stands for "quality assessment", at first I thought it was for "quality" alone, and it's only used two times in another place in the text so I'd remove it in order not to overload the text with acronyms.			
	Number: 4	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:47:10
	this can be merged with the last sentence into one so that information is not repeated.			
	Number: 5	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:50:13
	Not straightforward if we're still talking about manual assessment from an expert here?			
	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:51:57	
	And in the second sentence, the "instead" is wrongly used -> Instead of what? Maybe remove it?			
	Number: 6	Author: Reviewer	Subject: Highlight	Date: 19.09.2024, 11:49:07
	Number: 7	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 12:02:09
	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:02:31	
	This sounds strange, use "addressed" instead.			
	Number: 8	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 11:56:24
	tense			
	Number: 9	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:01:48
	I think ML deserves a paragraph on its own, it feels out of place here with the other thresholding algorithm.			
	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:04:02	
	Maybe instead of having just Blandini's model, you could also refer to other studys in the field that use ML (even if not exactly the same problem)? To showcase the recent advances in ML and why you chose to use ML.			
	Number: 10	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 12:02:40
	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:02:45	
	"with a"			
	Number: 11	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:56:58
	first time this is introduced and not sure what it refers to. Why is the data high-dimensional?			
	Number: 12	Author: Reviewer	Subject: Highlight	Date: 19.09.2024, 11:49:11
	Number: 13	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:58:46
	This is a very broad statement and not entirely true anymore, only for tabular data and would require a reference.			
	Number: 14	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 11:57:40
	This acronym should come above when the word is first introduced.			
	Number: 15	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:00:30
	Why do you think temporal information is important? Would be interesting to have.			
	Number: 16	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 12:04:14
	Number: 17	Author: Reviewer	Subject: Cross-Out	Date: 19.09.2024, 12:05:20
	Number: 18	Author: Reviewer	Subject: Inserted Text	Date: 19.09.2024, 12:05:39
	developped CleanSnow,			
	Author: Reviewer	Subject: Comment on Text	Date: 19.09.2024, 12:05:59	
	Nicer to have your model at the fore-front :)			

Comments from page 2 continued on next page



determine vegetation growth and plant phenology (e.g. Jonas et al., 2008; Fontana et al., 2008; Vitasse et al., 2017; Zehnder et al., in prep.) and to monitor climate change (e.g. Matiu et al., 2021). Finally, ~~the snow cover situation~~ directly influences tourism, transportation and recreational activities (e.g. Willibald et al., 2021).

Snow height data are nowadays available, sometimes in almost real-time, from airborne or satellite remote sensing and ground-based automated weather stations (AWS). One of the sensors often mounted at meteorological stations in high alpine regions is an ultrasonic snow height sensor (Ryan et al., 2008). Due to the measurement method, snow height data come with a variety of errors that arise from the harsh mountain conditions the sensor is not originally designed to operate in. In addition, ultrasonic sensors only measure the distance to the underlying object, be it snow or anything else. It is therefore important to validate whether the information coming from the snow height sensor really corresponds to snow or not.

Arguably the most precise way of assessing the **quality (QA)** of snow height measurements is via visual inspection of the data by a human expert (Robinson, 1989). Even though it is believed the most reliable, manual quality assessment of data is a tedious procedure heavily relying on expert knowledge, which is not easily transferable and does not scale well (Fiebrich et al., 2010). A common practice in snow height QA is to distinguish between snow and grass based on static climatological or minimum snow height thresholds. Random errors, instead, are typically detected using a maximum snow height threshold or snow height variance (Avanzi et al., 2014).

There are other sensors usually mounted at an AWS, which can provide information on whether the measured snow height relates to snow or not, as well as give some indications on the precision of snow height measurement. The first attempt to leverage other sensor information was the MeteIO library developed by Bavay and Egger (2014), which contains an algorithm for grass detection based on snow surface temperature, ground surface temperature, and solar radiation. The algorithm is based on a series of thresholding rules, an approach that is known to be rather cumbersome to modify and does not generally transfer well to other station data. Observing the recent advances in machine learning, Blandini et al. (2023) ~~have decided to deal with the high dimensionality of the data by proposing~~ a random forest approach to snow height QA, solving both snow height classification and anomaly detection at the same time. Random Forest (RF) models (Breiman, 2001) are possibly the most popular choice of machine learning algorithms used among data scientists worldwide. Apart from an attempt by Goehry et al. (2023), random forests, however, cannot easily and explicitly model the temporal structure of the data that we argue is crucial to be able to reliably say whether the snow height measurement is erroneous and whether the signal coming from the sensor shows snow or vegetation.

~~Therefore, we aim to develop a~~ machine learning model for the automated classification of snow height signals into a snow and a no-snow class, ¹⁹~~which we call CleanSnow~~. To approach this binary classification problem, we employed a Temporal Convolutional Network (TCN) (Lea et al., 2016) that explicitly accounts for the temporal relationships between different points in snow height time series data. To train our TCN, we created a new manually annotated snow height dataset composed of 20 measurement stations with around 20 years of data per station. This dataset also allows us to validate our design choices and evaluate the model in several different scenarios including challenging cases such as snow cover melt or plant growth periods.



2 Data

We used snow height data from the Swiss Intercantonal Measurement and Information System (IMIS) (Lehning et al., 1999), a network of 131 AWS (as of May 2024) focused on snow measurements that are distributed throughout the Swiss Alps and Jura
 60 **1** **Region** (see Figure 1), mostly located above 2000 m a.s.l. The stations acquire data regularly in 30-minute intervals and provide, in addition to snow height, also meteorological data. To analyze snow height (HS), we also leverage measurements such as air temperature (TA), snow surface temperature (TSS), wind speed (WV), relative humidity (RH), and reflected shortwave radiation (RSWR).




Figure 1. Map of IMIS stations in Switzerland. Stations marked as full gray circles were not part of the new annotated dataset. Yellow squares are the stations that have been used for training (14 stations) and red triangles indicate stations used for testing (6 stations). Colours indicate elevation in m a.s.l.

2.1 Quality assessment of snow height measurements

65 Raw snow height measurements coming from the IMIS network contain many errors and anomalies. Due to how the sensor works, it measures the height of an underlying object, independently of whether the object is snow or not. This yields spurious measurements (e.g., vegetation growth) in summer or generally during snow-free periods.

 Number: 1 Author: Reviewer Subject: Cross-Out Date: 19.09.2024, 12:08:23

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 12:20:01

This section repeats a lot of things from the introduction but also introduce a new study that should have been in the introduction too? Maybe separate the paragraph in the intro into "classical threshold methods" and "ML methods" and then you can move most of the information here into the introduction.

 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 12:21:13

This section needs some rethinking



There have been efforts to mitigate this effect and eliminate the vegetation measurements by using signals from other sensors, mainly snow surface temperature (TSS) and ground temperature (TG), which seem to be good indicators of whether there is snow on the ground or not (Tilg et al., 2015). In particular, with the presence of snow, TSS is expected to be $\leq 0^{\circ}\text{C}$. TG is expected to be constantly around 0°C , as snow insulates the ground from atmospheric temperature variations (Domine, 2011). When no snow is present instead, both TSS and TG typically show diurnal variations, in line with the air temperature (TA). For completeness, we also analyze wind speed (WV) since it has a direct influence on snow distribution and was considered in a recent classification approach (Blandini et al., 2023).

Techniques employing thresholding rules based on the above assumptions (Tilg et al., 2015; Bavay and Egger, 2014) generally work well and allow for, in some applications, satisfactory detection of the snow disappearance date at the end of the season and the timing of the first snow in the fall. Their main drawback lies in the definition of fixed threshold values which are used together with multiple conditional statements in order to determine the presence or absence of snow. These thresholds are often sensitive to anomalies and outliers in the data and do not transfer always well from one station to another. Moreover, manual adjustment of these thresholds is rather tedious and impractical with a large number of stations.

Careful manual exploration showed that the following sensor measurements are key factors in disentangling snow from soil and vegetation measurements: snow height (HS), air temperature (TA), snow surface temperature (TSS), ground temperature (TG) and reflected short-wave solar radiation (RSWR). The latter is useful since snow has a much higher albedo than soil or vegetation.

2.2 Data preparation

For model development and validation, we prepared a dataset with reliable ground truth information. Manually annotating snow height data is a tedious process, and doing so for the whole IMIS network is intractable. Therefore, we identified a subset of IMIS stations that we then manually annotated.

It should be mentioned that annotating historical data is problematic, as there is no way of checking whether there really was snow at the station or not. This means that assessing the presence of snow with the help of information from other sensors should be considered a best effort approach.

2.2.1 Snow/no-snow dataset

A subset of 20 stations (see Appendix A) which span different locations and elevations and vary in underlying surface (e.g., vegetation, bare ground, glacier, etc.) were selected and manually annotated with binary two-class ground truth information regarding snow height data:

- Class 0 - Snow - the surface is covered by snow
- Class 1 - No Snow - the surface is snow-free (e.g., vegetation, soil, rocks, etc.)

The stations annotated with ground-truth information are depicted in yellow and red in Figure 1. An example of data annotation is shown in Figure 2, with two detailed views that emphasize the differences in behavior of TSS and RSWR in the presence

Page: 4

 Number: 1 Author: Reviewer Subject: Pencil Date: 19.09.2024, 13:48:53

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 13:47:37

This sounds very vague. Does this refer to the work done in 2.2? If yes then it should be somewhere else, because right now it feels like you're stating something without backing it up with results.

 Number: 3 Author: Reviewer Subject: Cross-Out Date: 19.09.2024, 13:38:54

 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 13:39:03
difficult

 Number: 4 Author: Reviewer Subject: Inserted Text Date: 19.09.2024, 13:48:43

the other sensors that you used are these the ones mentionned above (temp, radiation, etc)? If yes maybe here would be a good place to put them.




Figure 2. Examples of manually annotated data for the calendar year 2010 at the station SLF2. (a) shows the snow cover flag and snow height; green rectangles mark periods with snow cover. (b) focuses on the end of winter season 2009/2010 illustrating the behavior of TSS and RSWR dependent on whether there is snow or not. (c) is the same as in (b) for the beginning of the winter season 2010/2011.

100 and absence of a snow cover. The selected stations mostly contain data between 2000 and 2023, at a 30-minute frequency, with a few exceptions for stations that have been built later (BOR2, FLU2, LAG3, RNZ2 and SHE2; see Appendix A).

2.2.2 Evaluation subset

We leave part of the annotated data out during model development, which we later use as an independent test set to evaluate the generalization ability of our final approach on stations not seen at training time. We select 6 stations (SLF2, WFJ2, KLO2, TRU2, STN2, SHE2) that contain challenging scenarios and are therefore suitable test cases. In particular, these stations are located at elevations where summer snowfalls occur, the snow season duration is very different, or where grass grows during the summer periods.

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 19.09.2024, 13:44:44

I don't understand this, is this necessary if the green values are always 1 (I'm guessing the snow cover flag refers to the green rectangles)?

 Number: 2 Author: Reviewer Subject: Inserted Text Date: 19.09.2024, 13:43:40

daily?

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:03:59

might want to define at some point (maybe earlier when it first arises) what you mean by generalization for non ML experts as it's very important



3 Machine learning based snow cover classification

To distinguish whether snow or other ground cover is under the sensor, other sensor measurements can be used. Based on the domain expert analysis discussed in Section 2.1 and empirical experimentation (see Section 4.2.3), we selected four sensor measurements as input features to our models, namely HS, TA, TSS and RSWR. We omitted TG, which was used during manual annotation, as it is not available at all IMIS stations and the sensor is also prone to defects.

Having temporal information further helps in determining whether there is snow or not at a particular time step. It is often important to look at a data point in the context of its temporal neighborhood. In an operational setting, one would, however, like to be able to make a prediction for each incoming data point in real-time. This means we cannot access data points in the future and the context for each data point has to be composed of itself and preceding data points (history). To reduce computational demands while still allowing for large enough context, we chose to work with a history window of 48 time steps (corresponding to 1 day), which has shown to provide the best results, as described later in Section 4.2.4.

However, this approach leads to a multivariate temporal input signal with high dimensionality. Therefore, it would be difficult to capture correlations between different feature points manually by defining, e.g., thresholding rules. Machine learning, instead, is an appropriate choice in such cases, and has already shown its power in other tasks concerning weather and climate data (e.g. Vaughan et al., 2022; Luković et al., 2022; Lam et al., 2023). The multivariate time-series signal contains both temporal dependencies between different data points from the same sensor, as well as inter-sensor correlations between measurements from multiple different sensors. Simple models such as random forests (Breiman, 2001) or multilayer perceptron (MLP) neural networks (Rosenblatt, 1958; Hornik et al., 1989; Cybenko, 1989) cannot explicitly account for the temporal nature of the data without engineering complex and artificial features, and are therefore a rather poor design choice. To correctly capture temporal patterns in the data, we instead chose to work with neural network models specifically designed to operate on time-series data, e.g., recurrent neural networks (McCulloch and Pitts, 1943; Kleene, 1951), Temporal Convolutional Networks (Lea et al., 2016), TimesNet (Wu et al., 2023) or Transformers (Vaswani et al., 2017).

We opted to use Temporal Convolutional Networks (TCN), which have proven useful in many applications concerning time-series data (Wan et al., 2019; Pelletier et al., 2019; He and Zhao, 2019; Hewage et al., 2020). Later, Section 4.3 provides a comparison of our choice to other popular models, such as Random Forests, MLPs, LSTMs (Hochreiter and Schmidhuber, 1997), Transformers and a recently released model for time-series processing called TimesNet, which yields state-of-the-art results on standard benchmarks in several different applications, including long and short term forecasting, anomaly detection, and other time-series based tasks.

3.1 Temporal Convolutional Network (TCN)

Based on well-known convolutional neural networks (CNNs) (Fukushima, 1988; Waibel et al., 1989; Weng et al., 1993; Lecun et al., 1998), TCNs are variations that consist of dilated, causal 1D convolutional layers that have the same input and output lengths. Dilation ensures that a specific entry in the output depends on all previous entries in the input, while causal convolution

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:07:32

 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:10:28
I find a bit confusing that this is based on results that come later. I would advise you do this differently. For example you could create here in section 3 a subsection called "experiments with the model's setup" where you explain different variations that are possible with this model, i.e., input variables, size of time frame, etc. And then in results you can keep your analysis to find the optimal setup.

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:05:58
merge this into one sentence.

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:10:36
same comment as above

 Number: 4 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:11:45
This belongs in the introduction not here.

 Number: 5 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:11:11
how high is this dimension?

 Number: 6 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:12:26
Same as above, this should be in the introduction in the proposed ML paragraph. Here just focus on your model.

 Number: 7 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 14:12:32

 Number: 8 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:16:17
These two sentences are very repetitive. Just focus on your model, the TCN, and if you use the others as a baseline then say that (although from the next line it looks like you compare to other models?)

 Number: 9 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 14:15:20
CleanSnow

 Number: 10 Author: Reviewer Subject: Inserted Text Date: 23.09.2024, 14:24:07
add here that it's a RNN

 Number: 11 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 14:17:42
is there a reference for this model?

 Number: 12 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 14:17:57




Figure 3. Structure of the input data and architecture of the modified TCN employed. A time window of 48 input signals of length 4 is fed to the TCN, which causally aggregates information from all time steps into a 128-dimensional latent vector. This information is subsequently fed into the classification network, which applies a sequence of MLPs to classify the input signal into two classes - *Snow* or *No Snow*. Each dilated 1D conv block has filters described in the format (*in_feats* × *out_feats* @ *kernel_size*). The composition of each MLP is described as (*in_feats*, *out_feats*, ..., *out_feats*).

means that the i -th element of the output sequence may only depend on input elements that come before it (elements with indices $\{0, \dots, i\}$).

As shown by Lea et al. (2016), with dilations and causal convolutions, TCNs can recover the behavior of RNNs (e.g. LSTMs or GRUs (Cho et al., 2014)) and achieve state-of-the-art results compared to RNNs on many tasks. Moreover, TCNs do not suffer from typical drawbacks of RNNs, such as the vanishing gradient problem (Pascanu et al., 2013), and are therefore easier to train. The use of convolutions instead of a recurrent mechanism also potentially leads to further performance improvements due to the possibility of parallelization of the convolution operation.

We chose a 4-layer TCN architecture as shown in Figure 3, which has 4-dimensional time series with 48 time steps as the input. The number of layers and filter sizes were selected so that the output representation of the last point in the input time series is an aggregation of all previous time steps. In other words, the TCN produces an output representation of the last point in the input time series by aggregating information from the whole history available at the input. This representation is fed to an MLP classifier, which first produces a series representation and then uses this representation to produce output class probabilities.

3.2 Training

Snow height classification is a binary problem. Binary classification problems are typically optimized using the cross-entropy objective function (Good, 1952). The simple cross-entropy loss will unfortunately not yield good results in our case. At places of interest that are available in the dataset, the snow cover usually prevails, hence creating significant imbalance. Moreover, as mentioned in Section 2.2.2, in many cases the classification task is simple, and we would like our model to perform well on the challenging edge cases. We therefore chose to drive the optimization by the so-called focal loss (Lin et al., 2017), which allows

	Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:19:32
this part is pretty small in the figure and a bit hard to read, can you make this bigger? As it's pretty important.				
	Number: 2	Author: Reviewer	Subject: Pencil	Date: 23.09.2024, 14:20:39
	Number: 3	Author: Reviewer	Subject: Pencil	Date: 23.09.2024, 14:19:02
	Number: 4	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:20:27
Where do these arrows point to (last and first downward arrow?), same for Class. Network. They seem to point from/to nothing.				
	Number: 5	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:20:59
of				
	Number: 6	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:21:33
add that it's one day				
	Number: 7	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:22:59
while we might guess what "feats" refer to "hid" might be more difficult for someone not an expert in ML. I'd advise against using abbreviations in figures that are not explained.				
	Number: 8	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:23:52
	Number: 9	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:25:14
do you mean that it's the same result as with an RNN? If yes that just means the same as "can recover the behavior of RNNs".				
	Number: 10	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:26:36
very well explained :) I like this paragraph.				
	Number: 11	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:26:51
loss				
	Number: 12	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:28:39
", but this loss will not yield good results in our case." And btw, have you tested this? Meaning have you tried to use the CE loss?				
	Number: 13	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:29:57
This is a complicated sentence, do you mean that in your dataset is imbalanced because there is more snow? Can you reformulate this simpler.				
	Number: 14	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:31:23
mmh actually you said that the test cases were particularly challenging but not that the rest is simple. What makes it simple there? I'd remove this and just keep " <i>we would like our model to perform well on the challenging edge cases.</i> "				
	Number: 15	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:31:40
	Number: 16	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:31:33



the model to focus and ¹train preferentially on hard examples, while down-weighting the simple cases throughout the training process.

The focal cross-entropy loss is defined as

$$FL = - \sum_{i=0}^{N-1} \alpha_i (i - p_i)^\gamma \log_b(p_i), \quad (1)$$

where α_i is the so-called balancing factor for class i , further contributing to class balancing, γ is the focus parameter which controls the down-weighting of the easy examples, p_i is the probability of the sample belonging to the i -th class, $N = 2$ is the number of classes in the classification problem, and b is the logarithm base; typically $b = 10$.

We run training for a maximum of 300 epochs, ²feeding the model with a batch of 64 samples in each iteration. We allow for the possibility of early stopping, if the validation loss has not improved for more than 50 epochs. The optimization process was governed by the AdamW (Loshchilov and Hutter, 2019) optimizer with an initial learning rate of 10^{-3} . The learning rate was subject to step decay with factor 0.1, three times, after 50, 100 and 150 epochs.

4 ³Experiments

In this section, we summarize experiments performed to evaluate CleanSnow. We start by describing the dataset used throughout the experiments. With a series of ablation studies, we clarify various design choices and then compare our TCN, the model of choice, to other available options. We continue with a thorough evaluation of the TCN in different periods of the year, pointing out its strengths and weaknesses. Experiments are concluded with a case study that demonstrates the use of CleanSnow in vegetation science.

4.1 Dataset

In all experiments, we used the snow/no-snow dataset described in Section 2.2.1. This dataset was split into train and evaluation subsets (see Section 2.2.2). ⁴For model training, we further split the training subset into the part on which we trained CleanSnow and a validation part that was used to validate CleanSnow during training and allowed for early stopping to avoid over-fitting of the model on the training data (Ying, 2019). ⁵The available validation dataset was also used for model hyperparameter tuning.

The whole training dataset contained a ⁷huge amount of data which would be rather impractical for experimentation as it would yield ⁸extremely long training times and high compute demands, which might not always be available. To make our experiments more tractable, we selected roughly 30% of the data from every station in the training set using filtering by year. ⁹Table B1 shows which years were used from each station.

¹¹We split our training dataset randomly using a 90/10 split, meaning we used 90% of the training subset for model training and the remaining 10% for validation. We fix the random seed in all our experiments to ensure the training/validation split remains the same across different runs and also to support reproducibility of the results. Random splitting inherently takes care of having samples from different stations and different time periods throughout the whole training subset.

Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:32:45
mmh but how does it know which sample is easy or not? Is that somehow defined by you? Would be interesting to have.			
Number: 2	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:33:01
Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:33:56
Call this results to avoid confusion with model & method before, and as you're showing resultls.			
Number: 4	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 14:59:01
This sentence is a bit convoluted and hard to follow.			
	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:01:41
a possible alternative: For model training, we further (randomly) divided the training subset into two parts using a 90/10 split: 90% used for training CleanSnow and 10% for validation. The validation set was used to monitor CleanSnow's performance during training, hyperparameter tuning, and enabled early stopping to prevent overfitting on the training data (Ying, 2019).			
Number: 5	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 14:59:07
Number: 6	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:13:35
merged into previous sentence as suggested above. But would be nice to know which hyperparameters were tuned and did you do cross-validation to tune these hyperparameters?			
Number: 7	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:00:07
how much? quantify			
Number: 8	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:00:15
Number: 9	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:00:48
(Table B1)			
Number: 10	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:00:51
Number: 11	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:01:56
merged into sentence above as suggested			



4.2 Ablation studies

In the following sections, different ablation studies and model comparisons are shown to explain our design choices and their contribution to obtaining the best results. Results presented in this section may serve as guidelines for designing machine learning solutions for snow height classification. All ablation studies were performed with a version of the TCN developed before feature elimination, which took seven input features, namely HS, TSS, TA, RSWR, RH (relative humidity), WV (wind speed) and solar altitude (which encodes information about the date and time of the day).

Models were compared using the Receiver Operating Characteristic (ROC) curve (Egan, 1975), which is a plot showing the performance in terms of the true positive rate (TPR) and the false positive rate (FPR).

4.2.1 Synthetic ground-truth experiments

To demonstrate the need for annotated data, we trained a model using synthetic ground truth based on empirical rules developed according to human expert knowledge. In order for a sample to correspond to snow cover, the following condition had to be met:

$$\left(\left(\frac{1}{N} \sum_{n=0}^{N-1} \text{TSS}_n \right) \leq 0.0 \right) \wedge \left(\left(\frac{1}{N} \sum_{n=0}^{N-1} \text{RSWR}_n \right) \geq 300.0 \right), \quad (2)$$

where N is the length of the time window.

We compared the model trained with the synthetic ground truth information to the model trained with the manually annotated data. The results in Figure 4(a) demonstrate the inability of thresholding rules to generate reliable ground-truth information that could be leveraged for training. This resulted in the *TCN Synth* model not learning the correct relationships between different input variables and therefore having a much worse performance than *TCN Annot*, which was trained with our manually annotated dataset.

4.2.2 Class balancing

Our training dataset included roughly twice as many snow-covered samples as snow-free samples. We applied class balancing by adjusting the class weights of the focal cross entropy loss and observed how that affected the performance of CleanSnow. Figure 4(b) shows that class balancing improved the performance and was therefore a valid design choice in our pipeline.

4.2.3 Feature importance

We performed an ablation study training the model with a leave-one-out strategy for the input features to validate their importance for the model decision-making. We picked the TCN architecture as it is our choice for the final solution. A comparison of TCN models with different input features missing is shown in Figure 4(c).

The HS, TSS, TA and RSWR signals proved to be important, in line with what was discussed above for manual data annotation. On the other hand, WV and RH had no beneficial effect and even slightly deteriorated the overall performance.

-
-  Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:28:56
I would call this differently as you're not only removing features and the only ablation study is 4.2.3. Maybe "Experiments with CleanSnow setup".
-
-  Number: 2 Author: Reviewer Subject: Highlight Date: 23.09.2024, 15:03:15
-
-  Number: 3 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:03:58
something eventually for discussion, not results.
-
-  Number: 4 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:06:37
this is confusing as I mentioned in 3.1. I would remove this and just present it as the default, that you're going to try to optimize.
-
-  Number: 5 Author: Reviewer Subject: Highlight Date: 23.09.2024, 15:04:04
-
-  Number: 6 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:06:46
"with"
-
-  Number: 7 Author: Reviewer Subject: Highlight Date: 23.09.2024, 15:07:19
-
-  Number: 8 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:07:16
these two acronyms were already defined.
-
-  Number: 9 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:07:40
and also the F1 score right?
-
-  Number: 10 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:11:00
This feels like something for the Appendix that you can mention when you first bring up the annotated dataset, not really for results :)
-
-  Number: 11 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:16:55
This statement, and all below in other result sections, need values (for example F1 scores from the figures) to underscore them. Otherwise the reader always needs to go double check the figures. It's easier if you provide values in the text too.
-
-  Number: 12 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:12:02
what weights were given to each class?
-
-  Number: 13 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:13:03
I think it's very strange that Fig 4b also shows a comparison against TimesNet and that it's not mentioned in the text. Seems like you should remove this from the figure and keep the comparison to other models for Fig5
-
-  Number: 14 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:14:08
CleanSnow
-
-  Number: 15 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:14:34
that's obvious.
-
-  Number: 16 Author: Reviewer Subject: Inserted Text Date: 23.09.2024, 15:14:23
(Figure 4c)
-
-  Number: 17 Author: Reviewer Subject: Inserted Text Date: 23.09.2024, 15:24:02
(i.e., removing them strongly decreases the model's performance)
-
-  Number: 18 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:23:31
euh this is a bit confusing. To make it clear I would reformualte it: "On the other hand, removing WV and RH from the input features slightly improved the model's performance, hinting that they have no beneficial effect."




Figure 4. ROC curves for various ablation studies. Every plot additionally shows the macro-F1 score for the threshold where TPR = FPR (the point on each curve). (a) Importance of manually annotated ground truth data. (b) Effect of class balancing. (c) Importance of input features. (d) Influence of sequence length on model performance.

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:18:25

Acronyms from figure missing in legend. Maybe not necessary to redefine climate variables but would define TCN Synth, TCN Annot.

 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:36:48

And can we change the legend so that the order of F1 scores goes from highest to lowest? This would make reading the figure and comparing setups a lot easier as there are a lot of colors. Same for Fig 5.

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:28:12

here it would be interesting to add (maybe in ---- line) the performance of the model with the final four features for comparison and to see where it positions itself in terms of performance, otherwise one has to go look at Fig. 5)



Hence neither feature provided any additional information useful for classification. Interestingly, solar altitude, which encodes information about date and time in continuous way, deteriorated the performance of the model considerably.

Accordingly, we chose our final model to have four input features, namely HS, TSS, TA and RSWR.

4.2.4 Sequence length selection

One of the key model architectural hyperparameters is the length of the history the models can use to predict the current time step. Figure 4(d) shows the relationship between history length and model performance. The best results were obtained with a history length of 48 time steps (24 hours). Very similar results were obtained with a history of length 32 (18 hours). A history length shorter than 24 time steps deteriorated the performance. Likewise, the performance decreased for history lengths larger than 96 time steps. Accordingly, we selected the history length to be 48 time steps as a compromise between sufficient but not too much context for the model.

4.3 Model selection

To choose the right architecture for the task at hand, we experimented with several state-of-the-art machine learning models for single time-step and time-series processing, compared their performance and finally selected the one that performed the best overall. Our model of choice was TCN, which was explained in Section 3.1. A short description of the other models we evaluated is provided in Appendix C.

To have a balanced model which does not favor one of the classes, we selected the decision threshold as the point where $TPR = FPR$. We evaluated the model for two scenarios: one with all seven input features and one with only the four relevant features.

Figure 5 shows the overall best performance of the TCN. Removing RH, WV and solar altitude, which were identified as irrelevant features resulted in a significant improvement of the LSTM model performance. Nevertheless, we opted for the TCN as it was on par with the LSTM, and the results in Figure 5(a) suggest that the TCN is more resilient to unimportant features in the input. In addition, the TCN showed advantages for training over RNNs. Interestingly, for RF the performance improved when using all features, which suggests it may learn undesired and spurious (see Section 4.2.3) relationships between inputs to distinguish snow from snow-free ground based on WV, RH and solar altitude.

4.4 Performance analysis per station

To better understand the generalization capabilities of the model, we evaluated its performance for each test station separately. The results in terms of confusion matrices are presented in Figure 6 and suggest good generalization capability of the model for most stations, with the exception of SLF2 and STN2. These two stations lie in very particular locations and are therefore out of distribution samples, which are described in detail below in Section 4.5. In addition, from Figures 6 and 7 one can further conclude that the model generally performs slightly better in correctly classifying presence of snow, compared to classification of snow-free ground.

	Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:27:16
Again this is not entirely true and leads to confusion. The only thing you can conclude is that removing it leads to better model performance, but having it did not deteriorate the model's performance considerably, seeing how the model with all features is pretty ok (in pink). Reformulate this.				
	Number: 2	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:30:05
this is confusing as it's not a hyperparameter that you tune but choose yourself. I'd call this something else.				
	Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:30:18
typo				
	Number: 4	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:30:37
	Number: 5	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:31:23
(F1...)				
	Number: 6	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:32:27
	Number: 7	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:31:30
(F1...)				
	Number: 8	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:34:26
	Number: 9	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:34:57
each				
	Number: 10	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:38:38
It actually gives similar results to the TCN, I'd add this, then it explains the next sentence.				
	Number: 11	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:40:27
Euh which advantages? And here you refer probably to the LSTM as RNN but be careful, just use one name to avoid confusion.				
	Number: 12	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:40:38
	Number: 13	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:41:54
mmh why would the relationships be wrong? First of all, RFs perform their own feature importance inside and maybe it learns just other realtionships than the other models? I'd remove this as it sounds far-fetched.				
	Number: 14	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:43:48
Merge 4.4 and 4.5 as it's basically the same, also that would explain why you refer to Fig 7 in 4.4. without mentioning it before like for Fig6.				
	Number: 15	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:42:40
	Number: 16	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:42:48
(Section 4.5)				




Figure 5. Model comparison shown as ROC curves for two different versions of the six models: (a) model performance with all seven input features - HS, TSS, TA, RSWR, RH, WV and solar altitude, and (b) with the four relevant input features - HS, TSS, TA and RSWR. Every plot additionally shows the macro-F1 score for the threshold where TPR = FPR (the point on the curve).

The seemingly good performance of the model should however be taken with a grain of salt. There are periods for which it is rather easy to correctly classify snow as *snow* and snow-free ground as *no snow* and other times of the year, when the problem becomes much harder. This is discussed in detail later in Section 4.6.

4.5 Influence of station location

It is important to understand whether CleanSnow generalizes to stations at different locations with different elevations. Results presented in Figure 7 suggest that the model performance was very stable for stations at elevations between roughly 2100 and 2700 m a.s.l., while it dropped for stations located either below or above this range. This corresponds to the fact that 80% of stations in our training set were in this range and only two stations were below 2000 m and one station was at 2800 m.

The two stations where model performance was lowest, SLF2 (1563 m) and STN2 (2914 m) were considerably outside the elevation range that was available during training. Moreover, these two stations are rather special cases compared to most of the other stations. SLF2 is located on a meadow in the village of Davos which seems to have a positive effect on the classification into the class *no snow*, as it was the only station with a F1 score for class *no snow* higher than for class *snow*. STN2, instead, stands on a glacier, which results in very different ground properties compared to any other station in the dataset. This is reflected by a rather low F1 score for the class *no snow*.

	Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:44:33
Same as for figure 4, missing acronyms. Models should at least be mentioned in legend.				
	Number: 2	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:44:11
here add the models..				




Figure 6 Confusion matrices for each test station separately ordered by elevation.




Figure 7. Model performance for the six stations of the test subset as a function of elevation. The F1 score is shown separately for the classification of snow (red line) and no snow (green line). The blue columns indicate the elevation distribution in the training subset (14 stations).

	Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:45:39
Acronyms				
	Number: 2	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:45:55
: SLF2, SHE2, ...				
	Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:46:12
it's yellow				



4.6 Performance for different times of the year

Classification of snow height measurements into snow and snow-free ground can be both a simple and rather challenging task depending on the location and time of the year. We provide a per-month performance analysis in Figure 8, which shows that the model mostly had trouble predicting snow-free ground in winter months. This is because very little training data for that class were available during December, January, February and March, ¹and it was not well represented in the training set. ²The lack of data for snow-free ground in these months is further emphasized by the fact that we had no samples from this class in the test set for February and March. In summer instead, the results suggest CleanSnow was able to ³detect most of the summer snowfalls while retaining very good performance on predicting snow-free ground. At the end of winter, in May and June, the model performance was also very good, suggesting that CleanSnow ⁴can accurately predict the snow disappearance date.




Figure 8. Performance of the model for each month of the year separately. The F1 score is shown separately for the classification of snow (red line) and no snow (green line). The blue columns indicate the distribution of snow samples, while the yellow columns indicate the distribution of the no-snow samples.

In addition, we analyzed the model performance for each season. To this end, we split the test dataset into four different seasonal clusters:

- **Winter season** was defined as the period with mostly continuous snow cover (December, January, February, March and April)
- **Summer season** was the part of the year typically without snow (July, August and September)
- **End of winter season** defined the snowmelt period resulting in snow-free ground (May, June and July)
- **Start of winter season** included the months when it starts snowing more often and at some point a continuous snow cover forms on the ground (September, October and November)

In the following sections we describe the model performance for each of the four seasonal clusters in detail and point out some season-specific challenges.

 Number: 1 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 15:47:15
repeats beginning of sentence.

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:48:47
Complicated sentence to say this: Furthermore, we had no snow-free samples in the test set for February and March.

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:49:32
I'd still mention the drop for July - Sep though

 Number: 4 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:50:05
mmh how can it predict this date?

 Number: 5 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 15:47:43
yellow




Figure 9. Confusion matrices for each of the four seasonal clusters.

4.6.1 Winter season

For snow classification, the middle of winter is presumably the easiest time of the year ¹ to deal with. Besides some low-elevation stations and some exceptional seasons with a very late onset of winter or very early snowmelt, the task should be rather trivial, as the snow cover is continuous in time. Figure 9(a) demonstrates that the model confidently classified snow (TPR = 99.4%) in contrast to the classification of snow-free ground ² with ³ **TPR = 88.4%**.

	Number: 1	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:51:10
	Number: 2	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:51:51
	Number: 3	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:51:41

in parenthesis too



4.6.2 Summer season

In contrast to full winter, classification of snow in the summer ¹was more challenging. Besides snow-free ground, there were many stations where vegetation grew. ³This ²results in non-zero snow height sensor measurements, which do not correspond to snow. ⁴Exceptions were stations at high elevations (e.g., on a glacier) and winters when the snow did not melt until the beginning of summer.

The snow height signal for snow-free ground typically oscillates with high frequency and either stays around zero or grows in the presence of vegetation under the sensor. ⁵The surface temperature and air temperature will most of the time oscillate high above 0°C showing a diurnal cycle. During overcast periods or in the presence of precipitation, TA and TSS will show the same value. Due to the lower albedo of snow-free ground, smaller amounts of ⁶reflected solar radiation (RSWR) are measured. Based on the above assumptions, summer snowfalls can be detected when TA equals TSS ⁷which is followed by larger values of RSWR with a simultaneous decrease in TSS. If there is vegetation growing under the station, the HS signal counter-intuitively decreases as the plants get pressed down by the snow. In the case of snow-free ground under the sensor, the HS signal will increase as expected during a snowfall.

⁸Despite the challenging setting, Figure 9(b) demonstrates that the model accurately detected snow-free ground with 99.2% accuracy. The effect of summer vegetation is shown in Figure 10(a). On the other hand, detecting a snowfall in the summer proved to be difficult, and even more so when vegetation was present. In this very difficult setting CleanSnow achieved a performance of 81.1%. A partial detection of a summer snowfall is shown in Figure 10(c). CleanSnow succeeded in detecting the main event but failed to correctly classify a few hours both at the start and the end of the summer snowfall.


4.6.3 Start and end of winter season

The transition periods between winter and summer and vice versa are key periods for the detection of the first snow and its disappearance, which are both dates of interest in climate science. These two seasonal clusters contain both data with rather continuous snow cover and with bare ground or vegetation growth. Such data are therefore a perfect test case for ⁹the approach we developed.


In our experiments, the end-of-winter season was the easier case to classify, achieving a very competitive performance of 98% for snow and 99% for snow-free ground (Figure 9(c)). ¹⁰We attribute this high accuracy to the fact that the transition from snow-covered to snow-free ground was often rather smooth, and once the snowpack had melted, there were not many periods with snow persisting on the ground. The beginning of summer was typically represented by high air temperatures, which caused TSS to oscillate with the daily cycle indicating snow-free ground; simultaneously RSWR noticeably decreased once the snow had completely melted. ¹¹Examples for end-of-winter season detection are shown in Figure 10.

On the other hand, classification during the start-of-winter season was more challenging: the model achieved an accuracy of 95.1% for snow and 93.2% for snow-free ground (Figure 9(d)). ¹²There were multiple snowfalls at the beginning of the season after which the snow melted again completely. In addition, in late autumn and the beginning of winter, temperatures occasionally dropped and the ground froze overnight. This resulted in TSS being constantly less than or equal to 0°C even


	Number: 1	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:52:08
is				
	Number: 2	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:52:43
resulted				
	Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:56:45
But these were manually removed during your manual annotation?				
	Number: 4	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:53:20
Do you know what percentage this corresponds to approximately?				
	Number: 5	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:53:47
This sentence is missing some commas				
	Number: 6	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 15:54:05
use the acronym				
	Number: 7	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 15:54:24
,				
	Number: 8	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 16:00:12
I guess that here the "Despite the challenging setting" should go with the performance of snow detection not snow-free, as that is not difficult in summer, I guess high air temperatures dominate.				
	Number: 9	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 15:59:19
CleanSnow				
	Number: 10	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 16:01:19
and Figure 10				
	Number: 11	Author: Reviewer	Subject: Cross-Out	Date: 23.09.2024, 16:01:23
	Number: 12	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 16:02:27
and Figure 10				




(a) SLF2 (1563 m), Year 2020



(b) SHE2 (1852 m), Year 2022



(c) TRU2 (2459 m), Year 2005



(d) STN2 (2914 m), Year 2013

Figure 10. Examples of classification results. The snow height signal is depicted in blue. The model predictions in terms of probability (0 - 1) are shown in green. The dashed horizontal line denotes the decision threshold for binary classification. The red-shaded areas show classification errors. (a) shows a correct classification of summer vegetation growth. (b) is an example of early October snowfall that has been classified partially correctly. (c) demonstrates the model's capability to detect summer snowfalls as well as scattered snowfalls at the beginning of winter. (d) is evidence that the model does not always perform well, here making mistakes at the beginning of the next winter season.

	Number: 1	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 16:09:29
Really cool figure :)				
	Number: 2	Author: Reviewer	Subject: Inserted Text	Date: 23.09.2024, 16:05:54
by CleanSnow				
	Number: 3	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 16:07:30
is the height indicative of the error?				
		Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 16:09:16
Maybe specify somewhere how they are calculated.				
	Number: 4	Author: Reviewer	Subject: Comment on Text	Date: 23.09.2024, 16:08:22
you mean because it didn't classify the snow height bumps as snow? Maybe specify.				



without snow which might force the model to focus more on RSWR and HS during decision-making, potentially decreasing its decision power. ~~The tricky nature of snow height classification at the start of winter season is shown in Figure 10.~~




Figure 11. Confusion matrices for daily aggregated values on our annotations (left) vs. human observations (right). Results for station WFJ2 are in (a) and (b), followed by results for SLF2 in (c) and (d).

4.7 Comparison to manual observations

A perfect test case ~~are~~ stations with concurrent manual observations, i.e., measurements manually performed by human observers. Such measurements were available for the two stations WFJ2 and SLF2 located in the region of Davos.

 Number: 1 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 16:02:16

This is just repetitive and unnecessary if you refer to it above

 Number: 2 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 16:10:24

is



325 Since the manual measurements were done only once per day, we resampled our predictions from 30-minute intervals into 24-hour intervals. We averaged probability scores over the 24 hours (48 automatic measurements) to obtain the per-day probability score.

The performance comparison on annotated automatic measurements versus manual observations in Figure 11 confirms that we had produced high-quality annotations for the historical data. Some days with snow were erroneously annotated as snow-free ground. This can be related both to short snowfalls which disappear in daily aggregation and also to the fact that manual observations were performed around 08:00 CET in the morning, while our data were daily averaged values. Such misalignment might produce additional disagreements between manual observations and our annotations.

330 The results also show that CleanSnow achieved very good performance when evaluated against daily manual observations. The differences in performance between the two ground-truth sources were attributed to the inconsistencies between the manual annotations of automatic measurements and manual observations.

4.8 Comparison to other approaches

To further demonstrate the added value of our machine learning approach, we compared it to other state-of-the-art methods such as filtering used in the physics-based snow cover model SNOWPACK (Lehning et al., 1999). In particular, we considered the snow water equivalent (SWE) provided by SNOWPACK since the HS signal is filtered to calculate SWE. Therefore, SWE should be a good indicator of whether the HS signal relates to snow or not. If the HS signal does not represent snow, one would expect SWE to be 0. In addition, we also compared CleanSnow to thresholding-based filters implemented in the MeteIO library, which were mainly designed to filter vegetation growth measurements in summer.




Figure 12. Performance (F1 score) per station for the ¹TCN (blue), the filter based on the SWE from SNOWPACK (red) and the thresholding filter from MeteIO (²green).

Figure 12 shows the comparison of the snow height classification by our TCN model to classification based on SWE calculated by SNOWPACK and the MeteIO filter. ³The results clearly show the machine learning approach to be superior. This might be attributed to the fact that both SNOWPACK and MeteIO use thresholding-based rules based on TSS and TG to filter HS similarly to the approach described by Tilg et al. (2015). The optimal threshold values vary across different stations, which

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:17:02

Change this to CleanSnow as it's the name of your model..

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:13:16

again it's yellow in the pdf sorry

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:16:49

Mmh I don't know about "clearly", I'd be just a little bit more modest. For more than half of the stations CleanSnow and Snowpack are competitive.



requires per-station calibration of the thresholds. Moreover, TG-based filtering is problematic since, as already mentioned, the TG sensor is prone to failures and the signal is therefore often missing at some stations.

4.9 Case study: Vegetation science

Besides obvious applications in snow science, a reliable separation of snowfall from plant growth also has benefits for biological research. Removing HS measurements classified as snow allows the extraction of a clean vegetation signal and pinning down reoccurring events in the life cycle of alpine vegetation – referred to as vegetation phenology. Given the long running time of continuous snow/plant height data collection, it is possible to relate the timing of green-up (i.e. the start of vegetation growth) or other phenological phases to snow climate parameters, and study phenological shifts over time – an excellent indicator of climate change (e.g. Inouye, 2022). We extracted 25 years of vegetation growth data from HS measurement data at TUJ2 (Culmatsch, 2262 m a.s.l.), an IMIS station characterized by tall plant growth. Within the 20 years of data, the algorithm flagged all snow days during the vegetation period which were then removed. Snow disappearance and snowmelt dates were defined as the first, respectively the last, day of the continuous winter snow cover. We fitted a logistic growth curve (Kong et al., 2022) to the clean plant growth measurements and defined the start of growth by a 10% threshold of maximum plant height (Figure 13). Vegetation green-up was directly linked to the timing of snowmelt, consistent with other studies (Jerome et al., 2021; Jonas et al., 2008), while late snowfall events shifted the start of growth towards later calendar days. Linear regression analysis revealed an earlier occurrence of green-up over the study period coinciding with an increase in spring temperatures measured at the station (Zehnder et al., in prep.). Despite insignificant changes in snowmelt timing, the shorter lag between snowmelt and initiation of plant growth indicated a warming-driven advancement in phenology at the study site. This case study highlights the importance of long-term monitoring and automated machine learning approaches in understanding climate-induced phenological shifts, with implications for ecosystem dynamics in remote alpine regions.




Figure 13. An example of a logistic growth curve (in dark green) fitted to height measurements data from TUJ2, in the vegetation season of year 2019. Snow height data corresponding to snow are shown with blue stars, while plant signal is shown with green diamonds. The red cross marks the snowmelt date, while the orange diamond marks the start of plant growth.

This page contains no comments



5 Discussion

We proposed a deep learning-based approach to snow height signal classification to automate the ¹quality-checking process. In addition to selecting an appropriate model, we provided some good practices to develop machine learning models for automated snow height classification. ²In the following paragraphs, we critically review our main findings.

5.1 ³Disentangling snow height from vegetation

⁴To add labels to historical snow height measurements, we needed to understand which sensor measurements were informative to separate snow height from snow-free ground measurements. We initially selected seven signals: HS, TA, TSS, RH, RSWR, WV, and solar altitude.

375 In Section 4.2.3 we showed that only HS, TA, TSS and RSWR were important for the classification of the snow height signal into snow and snow-free ground, which is in line with domain expert knowledge. ⁵The behavior of these four variables was explained earlier in Section 2.1. In contrast to domain expertise, we did not employ TG, as it was not available at all stations and, moreover, sensors measuring TG are prone to failures. Nevertheless, TG is expected to potentially further improve the results if used.

380 The remaining sensor measurements, namely RH, WV and solar altitude, were identified as uninformative for the disentanglement of snow and snow-free ground measurements. However, for other tasks such as, e.g., snow height anomaly detection, WV might very well be an important signal carrying information about snow transport by wind and related phenomena. ⁶Interestingly, solar altitude, which carries information about date and time, led to a deterioration of model performance. We attribute this to the fact that solar altitude information potentially makes the model take decisions based on the date and time of the year, ⁷which is rather undesirable. As much as date and time information are generally valid indicators of the season and therefore have a strong influence on the presence of snow, they might hamper decision-making, especially at the beginning and end of the snow season and in the case of summer snowfalls, whose occurrences vary from year to year.

5.2 Deep learning models for snow height classification

Second, the suitability of state-of-the-art deep learning models for the snow height classification task has been studied. Several cutting-edge deep learning architectures have been evaluated against each other, resulting in the superiority of a ⁸Temporal Convolutional Network over the other compared methods. The TCN reached an accuracy of 97.7%. ⁹When we used a decision threshold that balanced the model performance on predictions for both classes - *snow* and *no-snow*, ¹⁰no data from the test stations were used during training. Hence, the results indicate that the approach generalizes well to unseen stations. A detailed performance evaluation for each station in the test set showed that the model performed very well except on SLF2 and STN2, ¹¹which are two particular cases. The station SLF2 is located low in a valley and STN2 on a glacier. Such special environments, compared to those of most other stations in the dataset, ¹²might cause slightly different behavior of the auxiliary variables used during HS analysis and result in a performance decrease.

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:18:49
check the quality of what?

 Number: 2 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 16:19:16

 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:20:15
unnecessary

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:26:00
Somehow this title does not refer to anything that is directly discussed in this section. Maybe again something about feature importance instead.

 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:26:31
And this section just repeats things from the results so I don't find it necessary. I suggest you remove it.

 Number: 4 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:23:03
This makes it sound as though you're referring to your manual labeling though I'm assuming you're taking about cleansnow, change this.

 Number: 5 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 16:21:48

 Number: 6 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:24:54
I'd put this directly when this first comes up in the results so that the reader does not have to look it up later.

 Number: 7 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:26:51
TCN

 Number: 8 Author: Reviewer Subject: Inserted Text Date: 23.09.2024, 16:27:24
on the independant test set

 Number: 9 Author: Reviewer Subject: Cross-Out Date: 23.09.2024, 16:27:28

 Number: 10 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:34:59
Somehow something that seems to be missing here from your discussion is that it generalizes well to stations that are within the distribution of the training set (not like SLF2 and STN2) and that it struggles a bit when it's outside of that. This is not surprising but a very well known challenge in ML. It would be good to mention this and be transparent.



1.3 Best practices for snow height classification using machine learning

In our analysis, we aimed to establish good practices for further development of machine learning methods for snow height classification and quality assessment. We showed that learning from synthetic ground-truth data generated using thresholding rules proposed in the past did not work well, as the predefined thresholds did not generalize to all stations without modifications. This emphasizes the need for well-annotated data for training. Next, we pointed out the importance of addressing the class imbalance problem to achieve the best possible performance. Furthermore, we demonstrated the superiority of sequence-based models (TCN, LSTM, TimesNet and Transformer) over single time-step-based models (RF and MLP), which confirms the need for temporal context to achieve a high classification performance. We acknowledge the existence of techniques that allow one to feed RF and MLP models with sequences of data, e.g., lagged features (i.e., adding data from previous time steps as extra input features). Nevertheless, we argue that such techniques do not treat sequential data as a causal sequence, which is conceptually non-ideal and might potentially lead to the resulting model becoming less explainable in how it treats temporal information. Another important aspect to consider is the sequence length. We performed an analysis of the performance for the length of the time window (i.e., the size of the temporal context), which revealed that the ideal length was around 48 time steps, as shorter and longer time windows resulted in a deterioration of the model performance. Subsequently, we showed that it was important to evaluate the model performance during the critical times of the year (the start and the end of the winter season) to reveal their true performance.

5.4 Processing of raw sensor data

One of the known limitations of CleanSnow is the fact that it operates on raw data meaning the inputs may contain both anomalies (e.g. spikes) and missing values. Even though CleanSnow seem to be resilient to anomalies, it would be good practice to perform anomaly detection and filtering before running the proposed snow height classification models. We argue that filtering obvious spikes in snow height signal is a rather trivial procedure and can be solved by employing statistical methods such as Hampel filtering (Pearson, 1999) or an exponential moving average filter (Kendall and Stuart, 1966). However, other more subtle variations are very challenging to detect by both the human eye and automated methods.

Dealing with missing data is more complicated. At the moment, in the case of missing samples in the 48-time step context, the samples were discarded without being run through the model. Therefore, CleanSnow can only be applied in cases where the full history needed to make a prediction is available. A simple solution for periods of up to several time steps would be linear interpolation. However, as the size of the interpolated interval increases, this fails to produce an accurate reconstruction of the missing data. To impute larger periods of missing data, methods that take into consideration both spatial and temporal context should be employed. This is, however, out of the scope of this work, and we therefore leave it as a possible future research direction.

 Number: 1 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:30:29

I'd start with this section in the discussion as it's most conceptual

 Number: 2 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:36:15

In this section (or in a new one) it would be great also to know how you advise people to use your model.

 Number: 3 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:32:20

This also seems a wrong title. How about "Handling of anomalies and missing data".

 Number: 4 Author: Reviewer Subject: Comment on Text Date: 23.09.2024, 16:30:57

you mean its input? Seeing how the target is not raw but annotated



6 Conclusions

Automated snow height measurements are key input data for many modeling approaches in climate sciences, snow hydrology, and avalanche forecasting. Erroneous snow height measurement deteriorate the performance of these models. We demonstrated how to mitigate the aforementioned issues by the use of deep-learning methods for automated snow height classification. Our contributions can be summarized as three-fold. First, we adapted a novel machine learning approach to snow height signal classification that operates directly on time-series data. Second, we provided an in-depth comparison of several machine learning models applied to snow height classification. Third, we introduced a new benchmark dataset with annotated snow height data, which sets a baseline and can be used for further research in the field. The proposed approach achieved a high accuracy of 97.7% and generalized well to previously unseen stations. CleanSnow can be implemented as a component of an arbitrary snow height quality assessment pipeline without the need for any special hardware.

Code availability. The exact version of the software used to produce the results in this manuscript are available at

<https://doi.org/10.5281/zenodo.12698071>, while current and future version of can be found at

<https://gitlabext.wsl.ch/jan.svoboda/snow-height-classification>.

Data availability. The manually annotated dataset that we used to both train and evaluate CleanSnow is publicly available for research under CC BY-NC¹ license at <https://www.doi.org/10.16904/envidat.512>

Appendix A: List of stations in the snow/no-snow dataset

This section provides the list of IMIS stations used in our snow/no-snow dataset (see Section 2.2.1) together with their metadata.

Table A1 shows the stations ordered by increasing elevation. The column *Subset* indicates whether a station was used for training or testing.

Appendix B: Subsampling of the training data

To run experiments in a reasonable time and make sure they were computationally tractable, we sub-sampled the training dataset to reduce the amount of training samples. In Table B1 we list which years were selected for each station for the training set.

¹<https://creativecommons.org/licenses/by-nc/4.0/>



Station ID	Latitude [°N]	Longitude [°E]	Elevation [m]	Available since	Subset
SLF2	46.8127	9.8482	1563	November 1997	test
AMD2	47.1708	9.1468	1610	October 1997	train
GLA2	46.9966	9.0375	1632	November 2000	train
SHE2	46.7488	7.8124	1852	October 2001	test
ILI2	46.1913	6.8277	2022	March 2000	train
GUT2	46.6793	8.2896	2115	November 1999	train
KLO2	46.9091	9.8738	2147	November 1996	test
TUM2	46.7810	9.0214	2191	October 2002	train
FNH2	46.1007	6.9641	2252	September 1997	train
KLO3	46.8412	9.9316	2299	November 1996	train
LAG3	46.4245	9.6977	2300	November 2009	train
FLU2	46.7527	9.9464	2394	October 2003	train
RNZ2	46.6855	8.6267	2400	December 2008	train
TRU2	46.3709	7.5855	2459	November 1996	test
BOR2	46.2905	8.1093	2517	September 2001	train
WFJ2	46.8296	9.8092	2536	January 1996	test
ARO3	46.0874	7.5620	2602	September 1996	train
SPN2	46.2294	8.1176	2620	November 1996	train
FOU2	45.9717	7.0672	2800	October 1999	train
STN2	46.1678	7.7505	2914	October 1998	test

Table A1. List of stations that are part of the snow/no-snow dataset, together with their auxiliary information, ordered by elevation.

Appendix C: Machine learning models

For completeness, we provide a short description of every machine learning model that was used in our performance comparison.

C1 Random Forest (RF)

455 Implemented in many data science libraries and easy to use, Random Forests (RFs) are a popular choice of machine learning algorithm that can provide satisfactory predictions in both classification and regression tasks. In practice, RF is an ensemble approach, which produces a final prediction as a combination of outputs of many decision trees. It often works well on tabular data, but there are no mechanisms that would allow for a more principled representation of temporal, spatial or graph structures.

This page contains no comments



Station ID	Selected years
AMD2	1998, 2001, 2004, 2007, 2010, 2013, 2016, 2019, 2022
GLA2	2001, 2004, 2007, 2010, 2013, 2016, 2019, 2022
ILI2	2002, 2005, 2008, 2011, 2014, 2017, 2020, 2023
GUT2	2000, 2003, 2006, 2009, 2012, 2015, 2018, 2021
TUM2	2004, 2007, 2010, 2013, 2016, 2019, 2022
FNH2	2000, 2003, 2006, 2009, 2012, 2015, 2018, 2021
KLO3	1999, 2002, 2005, 2008, 2011, 2014, 2017, 2020, 2023
LAG3	2011, 2014, 2017, 2020, 2023
FLU2	2005, 2008, 2011, 2014, 2017, 2020, 2023
RNZ2	2010, 2013, 2016, 2019, 2022
BOR2	2002, 2005, 2008, 2011, 2014, 2017, 2020, 2023
ARO3	1998, 2000, 2003, 2006, 2009, 2012, 2015, 2018, 2021
SPN2	1999, 2002, 2005, 2008, 2011, 2014, 2017, 2020, 2023
FOU2	2001, 2004, 2007, 2010, 2013, 2016, 2019, 2022

Table B1. List of years for each station that were selected as part of the sub-sampled training dataset.

In our experiments we used the RF classifier implementation from the Scikit-Learn library (Pedregosa et al., 2011), setting the number of decision trees to 1000 and maximum depth of each tree to 50. We left the other parameters at their default settings and trained the RFs using the Gini criterion (Gini, 1936).

C2 Multilayer Perceptron (MLP)

Being one of the first neural network models that can learn non-linear functions, MLPs have shown their power in natural language processing (NLP) and serve as a foundational component for many other neural network models nowadays. Finding their applications in both regression and classification tasks, MLPs can serve as an alternative to the RFs presented above. Putting them in comparison with RFs, MLPs can be generally more difficult to train for a given task and often exhibit lower performance, especially with tabular data. This is due to their nature of learning smooth (sometimes overly smooth) solutions, thereby causing them to not perform well on problems with non-smooth decision boundary. Grinsztajn et al. (2022) argue this is due to the gradient descent approach to MLP optimization. They also show that MLPs are more affected by, e.g., uninformative features compared to RFs.

We designed an MLP composed of an input layer with 7 input dimensions and 32 output features, followed by 3 hidden layers with 64, 128 and 256 output features, respectively. Each hidden layer had batch normalization (Ioffe and Szegedy, 2015) and Rectified Linear Unit (ReLU) activation functions (Fukushima, 1969; Nair and Hinton, 2010) appended to it. The MLP was concluded with an output layer which takes a 256-feature representation and produces the final class probability score.

This page contains no comments



475 **C3 Long short-term memory (LSTM)**

Belonging to the family of recurrent neural networks (RNNs), the original models developed for time series processing, GRU (Cho et al., 2014) and long short-term memory (LSTM) (Hochreiter and Schmidhuber, 1997) are variations that allow the model to better capture long-term dependencies compared to RNNs, which tend to forget inputs that came much earlier in history. We chose to use an LSTM in our experiments, as it is one of the gold standards in deep learning for time-series processing.

The LSTM model we used in our experiments took an input with 7 dimensions and was composed of 3 recurrent layers with hidden dimensions of 64, 128 and 256, followed by an output MLP classifier that produced the final probability scores.

C4 TimesNet

Recently released and setting the new state-of-the-art performance on many standard benchmarks, TimesNet (Wu et al., 2023) has become one of the models of choice for time series processing in general. Its main characteristic is the transformation of a 1-dimensional time series signal into a 2-dimensional one, which allows it to capture complex temporal variations in the signal. The conversion of a time series into a 2-dimensional signal is based on detecting signal periods using amplitude information from a Fast Fourier Transform (FFT) and ordering the signal chunks into a 2-dimensional array. Applying 2-dimensional convolutions to this array allows it to capture both inter- and intra-period variations in the signal.

In our experiments we used a modification where the definition of signal periods is fixed and not determined by the FFT. We used 5 periods to split the signal, namely 48, 32, 24, 16 and 8. The model was then composed of 3 layers with each layer having 2 blocks and 128 hidden features.

C5 Transformer

Since it has been brought to the public's attention in 2017, transformers have revolutionized many areas of deep learning, achieving new state-of-the-art results mostly in natural language processing and computer vision. Transformers are model based on an attention mechanism (Vaswani et al., 2017) that were originally proposed for sequence-to-sequence tasks.

Here we employed a modification of the traditional transformer. In particular, we took the classical transformer encoder in order to produce a latent representation for the input sequence, where each point is conditioned on the past context. The encoder was composed of 2 layers with hidden dimensions of 128 and 4 attention heads. Both the input positional encoding and encoder have a dropout of 0.1 applied. The latent representation produced by the transformer encoder was average pooled and passed to an MLP readout network, which produced the classification probability scores.

Author contributions. JSc, MR and DL initiated the study and together with MV prepared the research idea and main goals. DL, MR and JSv prepared the data and carried out the manual data annotation. JSv and CJ analyzed the data, developed the models, prepared the experiments,

This page contains no comments



analyzed the results and drafted the original manuscript. MZ contributed the vegetation experiment. All co-authors provided critical reviews
505 and contributed to the final paper. JSc acquired the funding to support the study.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Acknowledgements. The authors gratefully acknowledge funding from Swiss Data Science Center grant C21-15L.

This page contains no comments



References

- Avanzi, F., De Michele, C., Ghezzi, A., Jommi, C., and Pepe, M.: A processing-modeling routine to use SNOTEL hourly data in snowpack
510 dynamic models, *Adv. Water Resour.*, 73, 16–29, <https://doi.org/10.1016/j.advwatres.2014.06.011>, 2014.
- Bavay, M. and Egger, T.: MeteoIO 2.4.2: a preprocessing library for meteorological data, *Geoscientific Model Development*, 7, 3135–3151,
<https://doi.org/10.5194/gmd-7-3135-2014>, 2014.
- Blandini, G., Avanzi, F., Gabellani, S., Ponziani, D., Stevenin, H., Ratto, S., Ferraris, L., and Viglione, A.: A random forest approach to
quality-checking automatic snow-depth sensor measurements, *The Cryosphere*, 17, 5317–5333, <https://doi.org/10.5194/tc-17-5317-2023>,
515 2023.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010950718922>, 2001.
- Cho, K., van Merriënboer, B., Bahdanau, D., and Bengio, Y.: On the Properties of Neural Machine Translation: Encoder-Decoder Approaches,
in: *Proceedings of the Eighth Workshop on Syntax, Semantics and Structure in Statistical Translation (SSST@EMNLP)*, pp. 103–111,
Association for Computational Linguistics, <https://doi.org/10.48550/arXiv.1409.1259>, 2014.
- 520 Cybenko, G. V.: Approximation by superpositions of a sigmoidal function, *Mathematics of Control, Signals and Systems*, 2, 303–314,
<https://doi.org/10.1007/BF02551274>, 1989.
- Domine, F.: Physical Properties of Snow, in: *Encyclopedia of Snow, Ice and Glaciers*, edited by Singh, V. P., Singh, P., and Haritashya, U. K.,
pp. 859–863, U. K., Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-90-481-2642-2_422, 2011.
- Egan, J. P.: Signal detection theory and ROC analysis, *Series in Cognition and Perception*, Academic Press, New York, NY, ISBN
525 9780122328503, 1975.
- Fiebrich, C., Morgan, C., McCombs, A., Hall, P., and Mcpherson, R.: Quality Assurance Procedures for Mesoscale Meteorological Data,
Journal of Atmospheric and Oceanic Technology, 27, 1565–1582, <https://doi.org/10.1175/2010JTECHA1433.1>, 2010.
- Flanner, M., Shell, K., Barlage, M., Perovich, D., and Tschudi, M.: Radiative forcing and albedo feedback from the Northern Hemisphere
cryosphere between 1979 and 2008, *Nature Geoscience*, 4, 151–155, <https://doi.org/10.1038/NGEO1062>, 2011.
- 530 Fontana, F., Rixen, C., Jonas, T., Aberegg, G., and Wunderle, S.: Alpine Grassland Phenology as Seen in AVHRR, VEGETATION, and
MODIS NDVI Time Series - a Comparison with In Situ Measurements, *Sensors*, 8, 2833–2853, <https://doi.org/10.3390/s8042833>, 2008.
- Fukushima, K.: Visual Feature Extraction by a Multilayered Network of Analog Threshold Elements, *IEEE Transactions on Systems Science
and Cybernetics*, 5, 322–333, <https://doi.org/10.1109/TSSC.1969.300225>, 1969.
- Fukushima, K.: Neocognitron: A hierarchical neural network capable of visual pattern recognition, *Neural Networks*, 1, 119–130,
535 [https://doi.org/10.1016/0893-6080\(88\)90014-7](https://doi.org/10.1016/0893-6080(88)90014-7), 1988.
- Gini, C.: On the Measure of Concentration with Special Reference to Income and Statistics, *Colorado College Publication*, 208, 73–79, 1936.
- Goehry, B., Yan, H., Goude, Y., Massart, P., and Poggi, J.-M.: Random Forests for Time Series, *REVSTAT-Statistical Journal*, 21, 283–302,
<https://doi.org/10.57805/revstat.v21i2.400>, 2023.
- Good, I. J.: Rational Decisions, *Journal of the Royal Statistical Society. Series B (Methodological)*, 14, 107–114,
540 <https://doi.org/10.1111/j.2517-6161.1952.tb00104.x>, 1952.
- Grinsztajn, L., Oyallon, E., and Varoquaux, G.: Why do tree-based models still outperform deep learning on typical tabular data?, in:
Advances in Neural Information Processing Systems, <https://doi.org/10.48550/arXiv.2207.08815>, 2022.
- He, Y. and Zhao, J.: Temporal convolutional networks for anomaly detection in time series, in: *Journal of Physics: Conference Series*, vol.
1213, p. 042050, IOP Publishing, <https://doi.org/10.1088/1742-6596/1213/4/042050>, 2019.

This page contains no comments



- 545 Herla, F., Haegeli, P., Horton, S., and Mair, P.: A Large-scale Validation of Snowpack Simulations in Support of Avalanche Forecasting Focusing on Critical Layers, *EGUsphere*, 2023, 1–38, <https://doi.org/10.5194/egusphere-2023-420>, 2023.
- Hewage, P., Behera, A., Trovati, M., Pereira, E., Ghahremani, M., Palmieri, F., and Liu, Y.: Temporal convolutional neural (TCN) network for an effective weather forecasting using time-series data from the local weather station, *Soft Computing*, 24, 16453–16482, <https://doi.org/10.1007/s00500-020-04954-0>, 2020.
- 550 Hochreiter, S. and Schmidhuber, J.: Long Short-term Memory, *Neural Computation*, 9, 1735–80, <https://doi.org/10.1162/neco.1997.9.8.1735>, 1997.
- Hornik, K., Stinchcombe, M., and White, H.: Multilayer feedforward networks are universal approximators, *Neural Networks*, 2, 359–366, [https://doi.org/10.1016/0893-6080\(89\)90020-8](https://doi.org/10.1016/0893-6080(89)90020-8), 1989.
- Inouye, D. W.: Climate change and phenology, *WIREs Climate Change*, 13, e764, <https://doi.org/10.1002/wcc.764>, 2022.
- 555 Ioffe, S. and Szegedy, C.: Batch Normalization: Accelerating Deep Network Training by Reducing Internal Covariate Shift, in: *Proceedings of the 32nd International Conference on Machine Learning*, vol. 37, pp. 448–456, <https://doi.org/10.48550/arXiv.1502.03167>, 2015.
- Jerome, D., Petry, W., Mooney, K., and Iler, A.: Snowmelt timing acts independently and in conjunction with temperature accumulation to drive subalpine plant phenology, *Global Change Biology*, 27, 5054–5069, <https://doi.org/10.1111/gcb.15803>, 2021.
- Jonas, T., Rixen, C., Sturm, M., and Stoeckli, V.: How alpine plant growth is linked to snow cover and climate variability, *J. Geophys. Res.-Biogeosci.*, 113, G03 013, <https://doi.org/10.1029/2007JG000680>, 2008.
- 560 Jonas, T., Marty, C., and Magnusson, J.: Estimating the snow water equivalent from snow depth measurements in the Swiss Alps, *Journal of Hydrology*, 378, 161–167, <https://doi.org/10.1016/j.jhydrol.2009.09.021>, 2009.
- Kendall, M. G. and Stuart, A.: *The Advanced Theory of Statistics. Volume 3: Design and Analysis, and Time-Series*, vol. 3 of *Griffin's statistical monographs and courses*, Charles Griffin & Company, London, ISBN 9780852642689, 1966.
- 565 Kleene, S. C.: *Representation of Events in Nerve Nets and Finite Automata*, RAND Corporation, Santa Monica, CA, <https://doi.org/10.1515/9781400882618-002>, 1951.
- Kong, D., McVicar, T., Mingzhong, X., Zhang, Y., Peña-Arancibia, J., Filippa, G., Xie, Y., and Xihui, G.: phenofit: An R package for extracting vegetation phenology from time series remote sensing, *Methods in Ecology and Evolution*, <https://doi.org/10.1111/2041-210X.13870>, 2022.
- 570 Lam, R., Sanchez-Gonzalez, A., Willson, M., Wirnsberger, P., Fortunato, M., Alet, F., Ravuri, S., Ewalds, T., Eaton-Rosen, Z., Hu, W., Merose, A., Hoyer, S., Holland, G., Vinyals, O., Stott, J., Pritzel, A., Mohamed, S., and Battaglia, P.: Learning skillful medium-range global weather forecasting, *Science*, 382, 1416–1421, <https://doi.org/10.1126/science.adi2336>, 2023.
- Lea, C., Vidal, R., Reiter, A., and Hager, G. D.: Temporal Convolutional Networks: A Unified Approach to Action Segmentation, *CoRR*, <https://doi.org/10.48550/arXiv.1608.08242>, 2016.
- 575 Lecun, Y., Bottou, L., Bengio, Y., and Haffner, P.: Gradient-based learning applied to document recognition, *Proceedings of the IEEE*, 86, 2278–2324, <https://doi.org/10.1109/5.726791>, 1998.
- Lehning, M., Bartelt, P., Brown, B., Russi, T., Stöckli, U., and Zimmerli, M.: SNOWPACK model calculations for avalanche warning based upon a network of weather and snow stations, *Cold Regions Science and Technology*, 30, 145–157, [https://doi.org/10.1016/S0165-232X\(99\)00022-1](https://doi.org/10.1016/S0165-232X(99)00022-1), 1999.
- 580 Lin, T., Goyal, P., Girshick, R. B., He, K., and Dollár, P.: Focal Loss for Dense Object Detection, in: *Proceedings of the IEEE International Conference on Computer Vision*, pp. 2999–3007, IEEE Computer Society, <https://doi.org/10.48550/arXiv.1708.02002>, 2017.

This page contains no comments



- Loshchilov, I. and Hutter, F.: Decoupled Weight Decay Regularization, in: Proceedings of the International Conference on Learning Representations, <https://doi.org/10.48550/arXiv.1711.05101>, 2019.
- Luković, M., Zweifel, R., Thiry, G., Zhang, C., and Schubert, M.: Reconstructing radial stem size changes of trees with machine learning, *Journal of the Royal Society Interface*, 19, 20220349, <https://doi.org/10.1098/rsif.2022.0349>, 2022.
- 585 Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C. M., Marty, C., Morin, S., Schöner, W., Cat Berro, D., Chiogna, G., De Gregorio, L., Kotlarski, S., Majone, B., Resch, G., Terzagio, S., Valt, M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G., Notarnicola, C., Petitta, M., Scherrer, S. C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Faletto, M., Gaddo, M., Giovannini, L., Mercalli, L., Soubeyroux, J.-M., Sušnik, A., Trenti, A., Urbani, S., and Weilguni, V.: Observed snow depth trends in the European Alps: 1971 to 2019, *The Cryosphere*, 15, 1343–1382, <https://doi.org/10.5194/tc-15-1343-2021>, 2021.
- 590 McCulloch, W. S. and Pitts, W.: A Logical Calculus of the Ideas Immanent in Nervous Activity, *The Bulletin of Mathematical Biophysics*, 5, 115–133, <https://doi.org/10.1007/bf02478259>, 1943.
- Morin, S., Horton, S., Techel, F., Bavay, M., Coléou, C., Fierz, C., Gobiet, A., Hagenmuller, P., Lafaysse, M., Ližar, M., Mitterer, C., Monti, F., Müller, K., Olefs, M., Snook, J. S., van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting: A status report on current implementations and prospects for the future, *Cold Regions Science and Technology*, 170, 102910, <https://doi.org/10.1016/j.coldregions.2019.102910>, 2020.
- 595 Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Quéno, L., Schirmer, M., Webster, C., and Jonas, T.: Operational snow-hydrological modeling for Switzerland, *Frontiers in Earth Science*, 11, <https://doi.org/10.3389/feart.2023.1228158>, 2023.
- Nair, V. and Hinton, G.: Rectified Linear Units Improve Restricted Boltzmann Machines Vinod Nair, vol. 27, pp. 807–814, <https://doi.org/10.5555/3104322.3104425>, 2010.
- 600 Pascanu, R., Mikolov, T., and Bengio, Y.: On the difficulty of training recurrent neural networks, in: Proceedings of the International Conference on Machine Learning, vol. 28 of *JMLR Workshop and Conference Proceedings*, pp. 1310–1318, <https://doi.org/10.48550/arXiv.1211.5063>, 2013.
- Pearson, R. K.: Data cleaning for dynamic modeling and control, 1999 European Control Conference (ECC), pp. 2584–2589, <https://doi.org/10.23919/ECC.1999.7099714>, 1999.
- 605 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine Learning in Python, *Journal of Machine Learning Research*, 12, 2825–2830, <http://jmlr.org/papers/v12/pedregosa11a.html>, 2011.
- Pelletier, C., Webb, G. I., and Petitjean, F.: Temporal convolutional neural network for the classification of satellite image time series, *Remote Sensing*, 11, 523, <https://doi.org/10.3390/rs11050523>, 2019.
- 610 Pérez-Guillén, C., Techel, F., Hendrick, M., Volpi, M., van Herwijnen, A., Olevski, T., Obozinski, G., Pérez-Cruz, F., and Schweizer, J.: Data-driven automated predictions of the avalanche danger level for dry-snow conditions in Switzerland, *Natural Hazards and Earth System Sciences*, 22, 2031–2056, <https://doi.org/10.5194/nhess-22-2031-2022>, 2022.
- Robinson, D.: Evaluation of the collection, archiving and publication of daily snow data in the United States, *Physical Geography*, 10, 120–130, <https://doi.org/10.1080/02723646.1989.10642372>, 1989.
- 615 Rosenblatt, F.: The perceptron: A probabilistic model for information storage and organization in the brain., *Psychological Review*, 65 6, 386–408, <https://doi.org/10.1037/h0042519>, 1958.
- Ryan, W. A., Doesken, N. J., and Fassnacht, S. R.: Evaluation of Ultrasonic Snow Depth Sensors for U.S. Snow Measurements, *Journal of Atmospheric and Oceanic Technology*, 25, 667 – 684, <https://doi.org/10.1175/2007JTECHA947.1>, 2008.

This page contains no comments



- 620 Tilg, A.-M., Ch., M., and G., K.: An automatic algorithm for validating snow depth measurements of IMIS stations (Abstract), 13th Swiss Geoscience Meeting, Basel, Switzerland, 20-21 November 2015, 339, 2015.
- Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A. N., Kaiser, L., and Polosukhin, I.: Attention is All you Need, in: Advances in Neural Information Processing Systems, pp. 5998–6008, <https://doi.org/10.48550/arXiv.1706.03762>, 2017.
- Vaughan, A., Tebbutt, W., Hosking, J. S., and Turner, R. E.: Convolutional conditional neural processes for local climate downscaling, 625 Geoscientific Model Development, 15, 251–268, <https://doi.org/10.5194/gmd-15-251-2022>, 2022.
- Vitasse, Y., Rebetez, M., Filippa, G., Cremonese, E., Klein, G., and Rixen, C.: ‘Hearing’ alpine plants growing after snowmelt: ultrasonic snow sensors provide long-term series of alpine plant phenology, International Journal of Biometeorology, 61, 349–361, <https://doi.org/10.1007/s00484-016-1216-x>, 2017.
- Waibel, A., Hanazawa, T., Hinton, G., Shikano, K., and Lang, K.: Phoneme recognition using time-delay neural networks, IEEE Transactions 630 on Acoustics, Speech and Signal Processing, 37, 328 – 339, <https://doi.org/10.1109/29.21701>, 1989.
- Wan, R., Mei, S., Wang, J., Liu, M., and Yang, F.: Multivariate temporal convolutional network: A deep neural networks approach for multivariate time series forecasting, Electronics, 8, 876, <https://doi.org/10.3390/electronics8080876>, 2019.
- Weng, J., Ahuja, N., and Huang, T.: Learning recognition and segmentation of 3-D objects from 2-D images, in: 1993 (4th) International Conference on Computer Vision, pp. 121–128, <https://doi.org/10.1109/ICCV.1993.378228>, 1993.
- 635 Willibald, F., Kotlarski, S., Ebner, P. P., Bavay, M., Marty, C., Trentini, F. V., Ludwig, R., and Grêt-Regamey, A.: Vulnerability of ski tourism towards internal climate variability and climate change in the Swiss Alps, Science of The Total Environment, 784, 147 054, <https://doi.org/10.1016/j.scitotenv.2021.147054>, 2021.
- Wu, H., Hu, T., Liu, Y., Zhou, H., Wang, J., and Long, M.: TimesNet: Temporal 2D-Variation Modeling for General Time Series Analysis, <https://doi.org/10.48550/arXiv.2210.02186>, 2023.
- 640 Ying, X.: An Overview of Overfitting and its Solutions, Journal of Physics: Conference Series, 1168, 022 022, <https://doi.org/10.1088/1742-6596/1168/2/022022>, 2019.
- Zehnder, M., Pfund, B., Svoboda, J., Marty, C., Alexander, J., Hille Ris Lambers, J., and Rixen, C.: Snow height sensors reveal phenological advance in alpine grasslands, in prep.

This page contains no comments