

# Effects of Arctic sea-ice concentration on surface radiative fluxes in four atmospheric reanalyses

Tereza Uhlíková<sup>1,2</sup>, Timo Vihma<sup>2</sup>, Alexey Yu Karpechko<sup>2</sup>, and Petteri Uotila<sup>1</sup>

<sup>1</sup>Institute for Atmospheric and Earth System Research, Faculty of Science, University of Helsinki, 00014, Helsinki, Finland

<sup>2</sup>Finnish Meteorological Institute, Helsinki, Finland

**Correspondence:** Tereza Uhlíková (tereza.uhlikova@helsinki.fi)

**Abstract.** Spatio-temporal variations and climatological trends in the sea-ice concentration (SIC) are highly important for the energy budget of the lower atmosphere and the upper ocean in the Arctic. To better understand the local, regional, and global impacts of the recent rapid sea-ice decline, one of the key issues is to quantify the interactions of SIC and the surface radiative fluxes. We analyse these effects utilising four global atmospheric reanalyses, ERA5, JRA-55, MERRA-2, and NCEP/CFSR and evaluate the uncertainties arising from inter-reanalysis differences in the sensitivity of the surface radiative fluxes to SIC. Using daily data over the period 1980–2021, the linear orthogonal-distance regression indicates similar sensitivity of surface upward longwave radiation to SIC in all reanalyses with the greatest sensitivity in the cold season November–April (over 150 W m<sup>-2</sup> per -0.1 change in SIC) and up to 80 W m<sup>-2</sup> per -0.1 change in SIC in May–October. We find that the effect of SIC on both surface upward longwave and shortwave radiation has mostly weakened in all seasons between the study periods of 1980–2000 and 2001–2021. The decrease in the sensitivity of upward longwave radiation to SIC can be attributed to the increasing surface temperature of sea ice, which dominated in the inner ice pack, and to the sea-ice decline, which dominated in the marginal ice zone. Approximately 80 % of the decadal decrease in upward shortwave radiation in May–July was caused by a decrease in surface albedo, controlled by SIC decrease, and the rest was caused by a decrease in downward shortwave radiation due to increase in cloudiness, mostly close to sea ice margins.

## 1 Introduction

Sea ice in the Arctic Ocean both affects and is affected by thermal longwave radiation and solar shortwave radiation. The former dominates the surface net radiation over most of the year and triggers the spring onset of snowmelt on top of sea ice (Mortin et al., 2016), whereas the latter is the key driver of summertime surface melt of snow and ice (Perovich et al., 2007). In winter over the Arctic Ocean, the snow surface temperature occasionally drops below -40 °C, which strongly reduces the emitted longwave radiation (Persson et al., 2002). Simultaneously, open leads with a surface temperature close to -1.8 °C emit almost double the amount of longwave radiation, and refrozen leads have intermediate values for surface temperature and longwave radiation emission.

In summer, the surface conditions are close to isothermal, and the longwave radiation emitted is much less sensitive to the presence of sea ice, whereas the effects of sea ice and snow on reflected solar radiation are strong. New dry snow has a

25 surface albedo of approximately 0.85, and even melting ice has a surface albedo of approximately 0.4 (Light et al., 2022),  
which is much higher than that of the open sea (less than 0.1). Hence, during spring and summer, the strong reflection from  
the snow or ice surface strongly reduces the surface net shortwave radiation. Throughout the year, both open water and sea ice  
surfaces generally emit more longwave radiation than they receive from clouds and the atmosphere (Persson, 2012). This is  
30 due to the high emissivity of snow and ice, 0.97-0.98 (Liang et al., 2014), which far exceeds the typical emissivity of the Arctic  
atmosphere, even under cloudy conditions (Garrett and Zhao, 2006). An exception occurs in the presence of thick water clouds  
in summer, which emit almost like a black body and have base temperatures close to or even higher than that of the snow/ice  
surface (Persson, 2012).

The above-mentioned findings are based on data from rare field campaigns in the Arctic sea ice zone. To understand the  
processes on a regional scale as well as their seasonal, inter-annual, and decadal variations and past trends, atmospheric and  
35 ocean reanalyses, as well as satellite remote sensing products, must be applied. Comparison of different reanalyses against  
each other and observations is vital to evaluate their uncertainty. Reanalysis products for surface radiative fluxes over sea ice  
have been compared and evaluated in several studies (Walsh et al. (2009); Graham et al. (2019); Jonassen et al. (2019); Yeo  
et al. (2022)). The ERA5 (Hersbach et al., 2020) and NCEP/CFRS (Saha et al. (2010); Saha et al. (2014)) reanalyses generally  
perform better than others (Jonassen et al. (2019); Di Biagio et al. (2021)), but challenges remain, especially for clouds and  
40 downward longwave radiation in winter (Graham et al., 2019). Additionally, reanalysis products for sea-ice concentration (SIC)  
have been compared (Graham et al., 2019). However, we are not aware of any study addressing inter-reanalysis differences in  
the relationship between SIC and radiative surface fluxes. This is a key question, as SIC plays a crucial role in the radiative  
surface fluxes and the surface energy balance over the Arctic Ocean.

Relevant research questions include the spatial patterns of the relationships between SIC and radiative surface fluxes over  
45 the Arctic Ocean, and the seasonal evolution of these relationships during the spring and autumn transitions. Considering the  
threshold value of SIC for sea ice to dominate the sign of the regional surface fluxes, it is known that for turbulent surface  
fluxes in winter, the threshold typically exceeds 0.9 (Vihma (1995); Andreas et al. (2010)), but for radiative fluxes, the thresh-  
old has not received as much attention. Regarding climatological trends, according to satellite passive-microwave data from  
1979—2021, the average yearly sea-ice extent in the Arctic has declined by more than 50 000 km<sup>2</sup> per year (Parkinson, 2022).  
50 To understand at the process level how the major sea ice decline has affected the ocean and atmosphere locally, regionally, and  
globally, the necessary first step is to quantify the effects of SIC on the surface energy balance of the Arctic Ocean. Further-  
more, the range of uncertainty in these effects and their changes over recent decades deserves attention.

To meet the above-mentioned challenges, we analyze the effects of SIC on surface upward shortwave and longwave radi-  
ation and clouds based on products of four atmospheric reanalyses. This is a follow-up study to Uhlíková et al. (2024), in which  
55 we addressed the effects of SIC on the turbulent surface fluxes of sensible and latent heat over the Arctic Ocean.

## 2 Material and Methods

To investigate the relationship between SIC and radiative surface fluxes, we utilised data from four atmospheric reanalyses. Because this paper is a companion paper to Uhlíková et al. (2024) (hereafter referred to as 'the companion paper'), we use data from (1) the same reanalyses (ERA5 (Hersbach et al., 2023), JRA-55 (JMA, 2013), MERRA-2 (GMAO (2015a); GMAO (2015b); GMAO (2015c)), NCEP/CFSR (Saha et al. (2010), Saha et al. (2011))), (2) the same study periods (1980–2000 and 2001–2021), (3) the same seasons (November–December–January, February–March–April, May–June–July, August–September–October), and the same temporal resolution (daily means of data), to make the two studies comparable. The term 'NCEP/CFSR' refers to data from both NCEP Climate Forecast System Reanalysis (CFSR; covering the period 1980–2010, spatial resolution  $0.312^\circ \text{ lat} \times 0.313^\circ \text{ lon}$ ) and NCEP Climate Forecast System Version 2 (CFSv2; covering the period 2011–2021, spatial resolution  $0.204^\circ \times 0.205^\circ$ ). We unified the spatial resolution for the whole 'NCEP/CFSR' data set to  $0.4^\circ \times 0.4^\circ$  using bilinear interpolation. Besides this adjustment, we worked with the original horizontal spatial resolution of the remaining reanalyses:  $0.25^\circ \times 0.25^\circ$  (ERA5),  $0.561^\circ \times 0.563^\circ$  (JRA-55), and  $0.5^\circ \times 0.625^\circ$  (MERRA-2).

From each reanalysis, we have used the following variables: sea-ice concentration (SIC), surface upward longwave radiation (ULW), surface temperature ( $T_s$ ), surface upward shortwave radiation (USW), surface downward shortwave radiation (DSW), and cloud water (vertically integrated cloud liquid water + cloud ice; hereafter referred to 'cloud condensate content', CCC). We chose CCC as a metric for cloud conditions, as it provides better available estimate of cloud radiative properties compared to total cloud cover (Senkova et al., 2007). [All surface radiative fluxes were defined as positive.](#)

Using these data, we studied bilateral relationships between SIC and surface upward radiative fluxes (ULW, USW) utilizing linear bilateral orthogonal-distance-regression model (ODR; Boggs et al. (1988)). Because all variables in reanalyses include uncertainties, ODR model is more optimal for this data than ordinary-least-squares-regression model (OLSR), which assumes no uncertainty in the independent variable (in our case SIC). Additionally, we performed a comparison study of bilateral ODR and OLSR outputs using data from the above-mention reanalyses and noted, that while the coefficients of determination ( $R^2$ ) were 'nearly identical' (at least to five decimal points identical) for both methods, the values of slopes of the regression line varied considerably. Based on these findings, we additionally decided to utilize OLSR analyses when only studying  $R^2$ , as this regression method requires less computing resources to perform. We used linear model for both ODR and OLSR as we evaluated it as the most applicable for our purposes primarily following from the finding that typically the first order i.e. linear term dominates over higher order ones when describing the relationship between two variables with the Taylor series.

The statistical-significance testing of the results was performed using Student's t-test (95 % confidence interval) with adjusted degrees of freedom ( $DF_{\text{adj}}$ ) according to Eq. (31) from Bretherton et al. (1999) to account for autocorrelation of the time series:

$$DF_{\text{adj}} = T \frac{1 - R_1 R_2}{1 + R_1 R_2} \quad (1)$$

where T stands for number of days in one sample (in our case days in seasons in the periods of 1980–2000 or 2001–2021) and  $R_1$  respectively,  $R_1$  for correlation coefficient of lag 1 auto-correlation of SIC, and  $R_2$  for correlation coefficient for of lag 1 auto-correlation of surface radiative flux (ULW, USW) and its explanatory variable (SIC) or USW). To test the field statistical

90 significance of the coefficients of determination (OLSR) and differences in mean decadal seasonal values between the two study periods, we have used p-value  $< 0.05$  adjusted by  $\alpha_{\text{FDR}} = 0.10$  (false discovery rate, according to Wilks (2016)) to reject the null-hypothesis that the time series are independent.

As we concluded in the companion paper, the largest differences in the effects of Arctic SIC on surface turbulent fluxes in reanalyses come from the representation of the sea ice, which is modelled in NCEP/CFSR and prescribed in ERA5, JRA-55, and MERRA-2. In Table 1, we reiterate the most important differences in representation of the sea ice in reanalyses and furthermore present differences in parameterisation of the sea-ice albedo.


### 3 Results

#### 3.1 Effects of sea-ice concentration on the surface upward longwave radiative flux

Utilizing linear bilateral ODR analysis, we assessed the effects of SIC on ULW. These two variables were negatively correlated in all seasons and both study periods (Figs. 1 and S1, S3, S4), meaning less SIC—more ULW or more SIC—less ULW. The sign of the correlation was in agreement with the theoretical expectations as the open ocean surface in the Arctic is usually warmer than the sea-ice surface (and much warmer in the cold season, November–April), and accordingly emits more longwave radiation. As depicted in the above-mentioned Figures, the sensitivity of ULW to SIC (slope of the regression line) did not vary considerably among reanalyses, with the highest values over  $150 \text{ W m}^{-2}$  ULW per  $-0.1$  change in SIC in November–April in the Central Arctic (north of  $81.5^\circ \text{ N}$ ). The dark grey areas [in these Figures](#) indicate a failure of the linear bilateral ODR model to converge. For JRA-55 ([Fig. 1b panels b, f, j in Figs. 1 and S1, S3, S4](#)), this was caused by the binary representation of SIC in ~~this-the~~ reanalysis, which assigns value 1 to  $\text{SIC} > 0.55$ , and value 0 to  $\text{SIC} \leq 0.55$ . Then, because the SIC in these dark grey areas was never less than 0.55 during the 21-year periods, every grid cell was assigned a value of 1. Hence, no dependence with ULW or any other variable could be found. In other reanalyses, the ODR model failure also occurred either because of very low variability in SIC or due to high uncertainty in the slope of regression between the two variables (as shown in Figs. S1 and S2). [In the warm season May–October, the effect of SIC on ULW was generally weaker, up to  \$80 \text{ W m}^{-2}\$  ULW per  \$-0.1\$  change in SIC \(Figs. S3 and S4\).](#)

The sensitivity of ULW to SIC mostly decreased in all seasons between 1980–2000 and 2001–2021 (shades of red in panels i–l in Figs. 1 and S1, S3, S4), but strengthened in the Central Arctic (shades of blue panels i–l in Figs. 1 and S1, S3, S4). To explain these changes, in Fig. 2, we show the daily [values of](#) SIC and ULW in grid cells from ERA5, MERRA-2, and NCEP/CFSR data, where the sensitivity changed considerably between 1980–2000 and 2001–2021 in November–December–January. While in Point 1 (see Fig. 1) from the border of Chukchi and East Siberian seas, the slope of the regression line became less steep in 2001–2021 compared to 1980–2000, in Point 2 from the Central Arctic, the slope became steeper in the second (more recent) study period.

As shown in Uhlíková et al. (2024, Fig. 5), the surface temperature of the Arctic sea ice ([bare or snow-covered,  \$T\_{\text{ice}}\$](#) ) generally increased between the two study periods, ~~therefore hence,~~ the difference between ~~the surface temperature of the ice ( $T_{\text{ice}}$ ) and the ocean  $T_{\text{ice}}$  and the sea-surface temperature~~ decreased causing lower sensitivity of ULW to SIC in the majority



**Figure 1.** Change in upward longwave radiative flux ( $W\ m^{-2}$ ) per change of 0.1 in sea-ice concentration (slope of regression line) in the marine Arctic in November–December–January in four reanalyses (columns), based on the linear orthogonal-distance-regression (ODR) model. Dark grey indicates areas where the ODR model did not converge; in panels (i)–(l), dark grey shows these areas in 1980–2000 and/or 2001–2021. Only grid cells with a mean of SIC  $> 0.5$  were considered, and only the slopes whose 95 % confidence intervals do not overlap zero are shown (others masked in white). Points 1 and 2 (in black) from panels (i), (k), (l) are further analysed in Figure 2.


of the Arctic in all seasons in the second study period. Also in this study, we show in Fig. 2: Point 1, that ULW (and therefore the surface temperature) is generally higher in 2001–2021 (lower panels) than 1980–2000 (upper panels) in days with SIC = 1. Another cause of decreasing sensitivity of ULW to SIC is the fact that in areas where the SIC declined or disappeared completely between the two study periods, there is naturally smaller or no effect of SIC on ULW in the second study period. ULW is also generally not so sensitive to SIC in regions where SIC is low, because in such regions,  $T_{ice}$  is typically higher, closer to sea-surface temperature. This is illustrated in the lower panels of Fig. 2: Point 1, where all the values of ULW in the grid cells with SIC lower than approximately 0.5 fluctuate close to  $300 \text{ W m}^{-2}$ .

130 The increased sensitivity of ULW to SIC in smaller areas in the Central Arctic may be due to increased SIC in reanalyses in these areas in 2001–2021 compared to 1980–2000. As shown in Fig. 2: Point 2, there are indeed both higher SIC as well as steeper slopes of the regression lines in the second study period (lower panels) than in the first one (upper panels). We discuss the possible mechanisms of the increased SIC in Section 4.1.

**Table 1.** Parameterisation of surface albedo Forecast model and representation of the sea ice in reanalyses.

	ERA5	JRA-55	MERRA-2	NCEP/CFSR
<u>Sea-ice albedo Forecast model</u>	Prescribed—seasonal cycle, —based—on Ebert and Curry (1993)- Parameterised, —function—of—hourly $\theta_s^a$ —and— $T_s^b$ Prescribed—seasonal cycle, —based—on Duynkerke and de Roode (2001)- Parameterised <b>Sea-ice thickness</b> 1.5—m; fixed 2—m; fixed n/a <sup>c</sup> Modelled—(coupled) <b>Snow on sea ice</b> None <u>IFS CY41R2</u>	None <u>JMA GSM</u>	None <u>GEOS 5.12.4</u>	Modelled—(coupled) <u>GFS (Atmospheric model)</u> <u>MOM4 (Ocean model)</u>
<b>Sea-ice concentration</b>	Fractional, external data set (OSI SAF <sup>ea</sup> (409a) (1979/Aug 2007, OSI SAF <sup>ea</sup> oper Sep 2007-)	Binary <sup>db</sup> , external data set (COBE-SST <sup>dc</sup> )	Fractional, external data set (OISST <sup>gd</sup> 1982/Mar 2006, OSTIA <sup>he</sup> Apr 2006-)	Fractional, modelled (coupled)
<u>Sea-ice thickness</u>	<u>1.5 m, fixed</u>	<u>2 m, fixed</u>	<u>n/a<sup>f</sup></u>	<u>Modelled (coupled)</u>
<u>Snow on sea ice</u>	<u>None</u>	<u>None</u>	<u>None</u>	<u>Modelled (coupled)</u>
<u>Sea-ice albedo</u>	<u>Prescribed seasonal cycle<sup>g</sup>, based on Ebert and Curry (1993) as in ECMWF (2016)</u>	<u>Parameterised, function of hourly <math>\theta_s^h</math> and <math>T_s^i</math></u>	<u>Prescribed seasonal cycle, based on Duynkerke and de Roode (2001)</u>	<u>Parameterised (output of model SIS-1<sup>j</sup> by GFDL<sup>k</sup>)</u>

<sup>a</sup> SIC > 0.55 = 1, SIC ≤ 0.55 = 0. <sup>b</sup> Ocean and Sea Ice Satellite Application Facility. <sup>c</sup> Centennial In Situ Observation-based Estimates of the Variability of Sea Surface Temperatures and Marine Meteorological Variables. <sup>d</sup> Optimum Interpolation Sea Surface Temperature. <sup>e</sup> Operational Sea Surface Temperature and Ice Analysis. <sup>f</sup> A 7-cm ice layer for computing a prognostic ice surface temperature, which is then relaxed towards 273.15 K as a representation of the upward oceanic heat flux; n/a: not applicable. <sup>g</sup> Considering albedo of fresh snow on top of sea ice (0.85) and its simplified metamorphosis (0.85–0.5). <sup>h</sup> Solar zenith angle. <sup>i</sup> Surface temperature. <sup>j</sup> Sea Ice Simulator. <sup>k</sup> Geophysical Fluid Dynamics Laboratory.



**Figure 2.** Daily sea-ice concentration (SIC) and upward longwave radiative flux (ULW) in selected grid cells, indicated in Figure 1 in panels (i), (k), (l), where the sensitivity of ULW to SIC between 1980–2000 and 2001–2021 decreased (Point 1, grid cell nearest to 73° N, 180° W) and increased (Point 2, grid cell nearest to 83° N, 0° W). ERA5, MERRA-2, and NCEP/CFSR data, days in November–December–January (1932 days). Black solid lines depict (a part of) the regression line and illustrate their slope.

To further explore the effect of the surface type in the marine Arctic on ULW, we investigated whether the main driver of ULW is the SIC or ~~the surface temperature of sea ice ( $T_{ice}$ )~~ $T_{ice}$ . To answer this question, we compared  $R^2$  (coefficient of determination) using SIC and  $T_{ice}$  as explanatory variables for ULW. To calculate  $T_{ice}$  from the grid-averaged surface temperature ( $T_s$ ), we utilized the following equation:

$$T_{ice} = \frac{T_s - (1 - SIC)T_{ocean}}{SIC} \quad (2)$$

where we assumed temperature of the ocean ( $T_{ocean}$ ) at  $-1.8^\circ\text{C}$  (271.35 K). ~~Naturally, this~~This assumption cannot be applied in ~~all grid cells, where SIC is present, especially in the Atlantic sector (Greenland and Barents seas), and in the~~ warm season (May–October) in the majority of adjacent seas outside the Central Arctic. ~~Therefore, because the surface temperature of the ocean is likely often higher than  $-1.8^\circ\text{C}$ . Hence,~~ we focused on the cold season (November–April) in these analyses. We are also aware, that in the Greenland and Barents seas, even cold-season ocean temperature may be warmer than  $-1.8^\circ\text{C}$  due to the North Atlantic Current carrying warm Atlantic water to this area. We utilized data from only ERA5, MERRA-2, and NCEP/CFSR because JRA-55 comes with binary representation of SIC, hence, Eq. (2) is not applicable for this data set. As shown in Figs. 3 and S5, in November–April,  $T_{ice}$  ~~explains~~explained over 90 % of the variance of ULW (~~vULW~~) in areas, where SIC is very high, whereas SIC ~~explains~~explained only around 30 % of ~~vULW~~the variance in ULW in these areas. However, in the marginal ice zone, the coefficient of determination was higher for SIC (around 60 %) compared to  $T_{ice}$  (< 30 %). These results were quantitatively very similar in both study periods and we found very good agreement between the three reanalyses.


### 3.2 Effects of sea-ice thickness on the surface upward radiative flux

In addition to SIC, also sea-ice thickness and snow depth on top of sea ice affect the surface temperature and, hence, the upward longwave radiation. Due to the limited amount and accuracy of data on sea-ice thickness and snow depth in the Arctic Ocean, we estimate their effect on ULW via analytic calculations, analogous to those in Uhlřková et al. (2024). We focus on the cold season when the insulating effects of ice and snow are largest. As a first approximation, we assume that the temperature profile through ice and snow is piecewise linear, resulting in the following expression for the conductive heat flux  $C$  (Makshtas, 1991)

$$C = k_i (T_s - T_b) / [h_i + (k_i - k_s) / h_s] \quad (3)$$

where  $k_i$  stands for the heat conductivity of ice,  $T_s$  for ice surface temperature,  $T_b$  for the ice bottom temperature,  $h_i$  for the ice thickness,  $k_s$  for heat conductivity of snow, and  $h_s$  for snow thickness. We used  $-1.8^\circ\text{C}$  for  $T_b$ ,  $2.1 \text{ W m}^{-1} \text{ K}^{-1}$  for  $k_i$ , and  $0.3 \text{ W m}^{-1} \text{ K}^{-1}$  for  $k_s$ . The turbulent fluxes of latent and sensible heat were calculated applying the standard bulk formulae:

$$\text{LHF} = \rho L_E C_{HE} (Q_a - Q_s) V \quad (4)$$



**Figure 3.** Proportion of variance in the upward longwave radiation (ULW) explained by sea-ice concentration (SIC) and surface temperature of the ice ( $T_{ice}$ ) in November–December–January, 1980–2000 and 2001–2021 (columns), as represented in three reanalyses (rows), based on linear ordinary-least-square regression model (coefficient of determination,  $R^2$ ) using daily means of data from ERA5, MERRA-2, and NCEP/CFSR. Only grid cells with a mean of SIC > 0.5 were considered and only statistically significant results at the 5 % level of significance are shown (insignificant masked in white).

165 
$$\underline{\underline{SHF}} = \rho c_p C_{HE} (T_a - T_s) V$$


(5)

where  $\rho$  stands for the air density,  $L_E$  for the latent heat of sublimation,  $c_p$  for the specific heat of the air, and  $C_{HE}$  for the turbulent exchange coefficient;  $(Q_a - Q_s)$  and  $(T_a - T_s)$  are the differences in specific humidity and temperature between the lowest atmospheric level and the surface, and  $V$  stands for the wind speed at the lowest atmospheric level of the model applied in each reanalysis. The upward long-wave radiation (ULW) was calculated as:

$$170 \quad \text{ULW} = \sigma T_s^4 \quad (6)$$

where  $\sigma$  stands for the Stefan-Boltzmann constant ( $5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ ). As in Uhlíková et al. (2024), the downward longwave radiation (DLW) and the input for Eqs. (3) to (6) were taken from observations from the SHEBA campaign in the Central Arctic in February 1998 (Persson et al., 2002), when the mean values were as follows:  $155 \text{ W m}^{-2}$  for DLW,  $5.0 \text{ m s}^{-1}$  for  $V$ ,  $-32 \text{ }^\circ\text{C}$  for  $T_a$ , and 0.9 for the relative humidity, yielding  $0.17 \text{ g kg}^{-1}$  for  $Q_a$ . Then Equations (3) to (6) were solved applying the following values of  $h_i$ : 0.2, 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 m, with  $h_s$  systematically set as  $0.1 \times h_i$ . As  $T_s$  is unknown and all the fluxes except DLW depend on it, a set of calculations with various  $T_s$  values was carried out for each combination of  $h_i$  and  $h_s$  until the  $T_s$  yielded zero net heat flux (sum of radiative, turbulent and conductive fluxes) at the snow surface, representing equilibrium conditions.

The sensitivity of  $T_s$ ,  $C$ , and ULW to snow and ice thickness is presented in Fig. 4. In the case of thin ice, the snow surface temperature is highly sensitive to ice thickness, but the sensitivity decreases when ice gets thicker (Fig. 4, left panel). This is reflected in ULW. For 0.2 m ice thickness (0.02 m snow depth), ULW is  $227 \text{ W m}^{-2}$ , whereas for 3 m ice thickness (0.3 m snow depth) ULW is  $183 \text{ W m}^{-2}$ , representing a difference of  $-44 \text{ W m}^{-2}$ . The difference in ULW between ice thicknesses of 2 and 3 m is minor ( $-2 \text{ W m}^{-2}$ ), as the conductive heat flux through ice and snow is small already for 2-m thick ice (covered by 0.2 m snow pack). Comparison of Figs. 4 and S1 shows that in winter in the Bering Sea, Sea of Okhotsk, and Barents Sea, ULW is approximately equally sensitive to a decrease of ice thickness from 3 to 0.2 m and to a decrease of SIC by 0.1. However, closer to the central Arctic Ocean and in the Canadian Arctic archipelago, the sensitivity is higher for a decrease of SIC by 0.1. These high statistical sensitivities to SIC may be partly due to co-occurrence of low SIC and high ice surface temperatures.



**Figure 4.** Sensitivity of snow surface temperature (left panel), conductive heat flux through snow and ice (right panel, dashed line), and upward longwave radiation (right panel, solid line) to sea-ice thickness and snow depth (set as 10 % of the ice thickness). The numbers are representative for February in the central Arctic Ocean.

### 3.3 Effects of sea-ice concentration on surface upward shortwave radiative flux

The sea-ice ~~surface~~ (bare or snow-covered) has much higher ~~shortwave reflectivity~~ surface albedo (the proportion of incident shortwave radiation that is reflected back to space by the surface) than the open sea. Hence, as expected, we found a positive correlation between SIC and USW meaning more SIC–more USW or less SIC–less USW in all seasons with solar radiation present in the Arctic (Figs. 4.5, S6, S8). USW was the most sensitive to SIC in May–June–July in the Central Arctic – over 100 W m<sup>-2</sup> USW per 0.1 change in SIC. The ODR model did not converge in large areas of the marine Arctic in February–March–April and August–September–October due to lack of variability in both incoming solar radiation, which was mostly very low during these months, and in SIC, which was very high. This is illustrated for representative grid cells in Figs. S6 and S7.

The effect of SIC on USW weakened between 1980–2000 and 2001–2021 in nearly all of the Arctic (shades of blue in panels i–l in Figs. 4.5 and S6). As discussed in ~~the previous subsection~~ Section 3.1, the sea-ice decline in adjacent Arctic seas naturally contributes to decreased effect of SIC on ULW; the same applies also to SIC effect on USW. However, because USW is a result of both, the downward shortwave radiation (DSW) and the reflectivity of the surface (surface albedo), the decrease in USW sensitivity to SIC between the study periods could have been ~~additionally~~ caused by changes in either or both of its above-mentioned drivers. To address this issue, we created Figs. 5, 6, ~~S9–S12~~ S9–S12, S10–S13, which show changes in seasonal means of shortwave radiative fluxes between the periods ( $\Delta$ DSW,  $\Delta$ USW),  $\Delta$ USW explained by change in DSW ( $\Delta$ USW<sub>DSW</sub>), and  $\Delta$ USW explained by change in surface albedo (b,  $\Delta$ USW<sub>b</sub>). The above-mentioned variables were calculated for each grid cell using daily data according to the following equations:

$$\Delta$$
DSW = DSW<sub>2001–2021 mean</sub> – DSW<sub>1980–2000 mean</sub> (7)


$$\Delta$$
USW = USW<sub>2001–2021 mean</sub> – USW<sub>1980–2000 mean</sub> (8)

$$b = \frac{\text{USW}_{1980-2000 \text{ mean}}}{\text{DSW}_{1980-2000 \text{ mean}}} \quad (9)$$

$$\Delta$$
USW<sub>DSW</sub> = b ×  $\Delta$ DSW (10)

$$\Delta$$
USW<sub>b</sub> =  $\Delta$ USW –  $\Delta$ USW<sub>DSW</sub> (11)

210 ~~Changes in decadal means (calculated from daily means) between 1980–2000 and 2001–2021, May–June–July. Panels (a)–(h) show changes in surface downward and upward shortwave radiative fluxes ( $\Delta$ DSW,  $\Delta$ USW), panels (i)–(l) show~~



**Figure 5.** Change in upward shortwave radiative flux ( $\text{W m}^{-2}$ ) per 0.1 change in sea-ice concentration (slope of regression line) in four reanalyses (columns), marine Arctic, November–December–January, based on the linear orthogonal-distance-regression (ODR) model. Dark grey indicates areas where the ODR model did not converge; in panels (i)–(l), dark grey shows these areas in 1980–2000 and/or 2001–2021. Only grid cells with a mean of SIC  $> 0.5$  were considered, and only the slopes whose 95 % confidence intervals do not overlap zero are shown (others masked in white).

changes in USW explained by changes in DSW ( $\Delta \text{USW}_{\text{DSW}}$ ), and panels (m)–(p) changes in USW explained by changes

in albedo ( $\Delta USW_b$ ). Only statistically significant results at the 5% level of significance are shown (insignificant masked in white); statistically significant grid cells for  $\Delta USW$ ,  $\Delta USW_{DSW}$ , and  $\Delta USW_{DSW}$  are identical. Values within an interval (-0.1,0.1)  $W m^{-2}$  are also masked in white.

Changes in decadal means (calculated from daily means) 2001–2021 minus 1980–2000, May–June–July. Row (i) shows cloud condensate content (CCC, vertically integrated cloud liquid water + ice), row (ii) shows surface albedo, and row (iii) sea ice concentration (SIC). Only statistically significant results at the 5% level of significance are shown (insignificant masked in white). In rows (ii) and (iii), values within an interval (-0.01,0.01) are also masked in white.

220 For May–June–July, Fig. 5a–d–6a–d shows that reanalyses agreed on the strongest decline (around  $-15 \text{ W m}^{-2}$ ) in the mean DSW between 1980–2000 and 2001–2021 in northern Barents Sea between Svalbard and Novaja Zemlya and some smaller degree of decline in this variable in other adjacent Arctic seas. All reanalyses also agreed on an increase around  $10 \text{ W m}^{-2}$  in the mean DSW between 1980–2000 and 2001–2021 in the Central Arctic, north of Greenland and Canadian archipelago. However, the areal extent of this increase varied considerably between the data sets, with NCEP/CFSR showing the largest one, 225 followed by MERRA-2. According to Fig. 6–7 row i, the areas of increased DSW correspond with those where CCC (vertically integrated cloud water + ice) diminished between the two study periods. Vice versa, the area of strongest decadal seasonal reduction of DSW in northern Barents Sea between Svalbard and Novaja Zemlya can be connected with the one where CCC increased.

Mean USW between 1980–2000 and 2001–2021 (Fig. 5e–h–6e–h) declined in most of adjacent Arctic seas by more than  $-25$  230  $\text{W m}^{-2}$  in all reanalyses. In agreement with theoretical expectations, most of the decadal seasonal reduction in USW outside the Central Arctic (around 80 %) was attributed to decrease in surface albedo (shades of blue in Figs. 5m–p and 6–6m–p and 7 row ii) which to a large part coincided with SIC decline (shades of blue in Fig. 6–7 row iii). However, also reduction of DSW (around  $-5 \text{ W m}^{-2}$ ) played a role (Fig. 5i–l)–6i–l). Furthermore, ERA5 and NCEP/CFSR indicated an increase in mean USW (around  $+10 \text{ W m}^{-2}$ ) in 2001–2021 in the Central Arctic, north of Greenland and Canadian archipelago (shades of red in Fig. 235 5e6e, h) which spread about equally between an increase in albedo and DSW in this area (shades of red in Fig. 5i6i, l, m, p). As-

To offer a comparison of absolute values of the sea-ice albedo in ERA5 has a prescribed seasonal cycle, the same for both study periods between reanalyses, we calculated its daily and monthly means at the North Pole in six Junes in the middle of the two study periods (1989, 1990, 1991, 2009, 2010, 2011). To obtain the sea-ice albedo ( $b_{\text{ice}}$ ) from grid-averaged surface albedo ( $b_{\text{s}}$ ), we utilized the following equation: 240

$$b_{\text{ice}} = \frac{b_{\text{s}} - (1 - \text{SIC})b_{\text{ocean}}}{\text{SIC}} \quad (12)$$

where we assumed the albedo of the ocean  $b_{\text{ocean}}$  at 0.06. Monthly means of  $b_{\text{ice}}$  are shown in Table 2 and daily means are depicted in Fig. S9. In all selected peak-summer months, the decadal seasonal increase in albedo can only be explained by increased SIC in the area (as shown in Fig. 6 row iii). In NCEP sea-ice albedo in MERRA-2, which has prescribed seasonal 245 cycle, was the highest among reanalyses and the albedo parameterized in JRA-55 was the lowest in both monthly and nearly all daily means. These two datasets varied by up to around 0.2. June monthly means of the sea-ice albedo in ERA5 and NCEP/CFSR however, albedo is parameterized, therefore the increase in this variable indicates decadal seasonal changes in surface albedo in this data set (Fig. 6 row ii). At the same time, we also detected decadal seasonal growth in SIC in the Central Arctic CFSR were very similar, even though the variable is modelled in NCEP/CFSR (Fig. 6 row iii) and prescribed in ERA5. 250 The daily means of surface albedo between these two data sets varied by up to 0.1.


In February–March–April, we found very little statistically significant decadal differences in DSW, however, reanalyses generally agreed that there was an increase in CCC over the Barents Sea, between Svalbard and Novaja Zemlya, and decline along the east coast of Greenland (Fig. S10–S11 row i). We found mostly decadal reduction in USW (around  $-15 \text{ W m}^{-2}$ ) in the

marginal ice zone (shades of blue, Fig. [S9e-h](#)[S10e-h](#)). This reduction, similarly to May–June–July, was mostly attributed to  
255 decline in surface albedo (Fig. [S9m-p](#)[S10m-p](#)), but partly also to reduction in DSW (Fig. [S9i-l](#)[S10i-l](#)).


**Table 2.** Monthly mean sea-ice albedo in the grid cell nearest to the North Pole (90° N, 0° W) in three Junes in the middle of the first study period (1989, 1990, 1991) and three Junes in the middle of the second study period (2009, 2010, 2011).

	<u>1989</u>	<u>1990</u>	<u>1991</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>
<u>ERA5</u>	<u>0.69</u>	<u>0.70</u>	<u>0.71</u>	<u>0.71</u>	<u>0.71</u>	<u>0.71</u>
<u>JRA-55</u>	<u>0.59</u>	<u>0.59</u>	<u>0.59</u>	<u>0.61</u>	<u>0.59</u>	<u>0.59</u>
<u>MERRA-2</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>	<u>0.78</u>
<u>NCEP/CFSR</u>	<u>0.67</u>	<u>0.68</u>	<u>0.67</u>	<u>0.68</u>	<u>0.69</u>	<u>0.68</u>

In August–September–October, we noted decadal reduction in mean DSW around  $-10 \text{ W m}^{-2}$  in adjacent Arctic seas. All reanalyses also agreed on decadal reduction in the mean USW though disagreed on the magnitude over the Beaufort, Chukchi, East Siberian, and Laptev seas. In these areas, the decrease in USW ranged between around  $-20 \text{ W m}^{-2}$  in JRA-55 and around  $-10 \text{ W m}^{-2}$  in MERRA-2 (Fig. [S11](#)[S12e-h](#)). As in the two previously-mentioned seasons, more of the mean USW reduction between 1980–2000 and 2001–2021 was attributed to decline in surface albedo than decline in DSW. Regarding decadal changes in mean CCC, we found a strong increase across the Arctic, though reanalyses showed a large scatter on the magnitude and spatial pattern of this change (Fig. [S12](#)[S13](#) row i).



**Figure 6.** Changes in decadal means (calculated from daily means) between 1980–2000 and 2001–2021, May–June–July. Panels (a)–(h) show changes in surface downward and upward shortwave radiative fluxes ( $\Delta$ DSW,  $\Delta$ USW), panels (i)–(l) show changes in USW explained by changes in DSW ( $\Delta$  USW<sub>DSW</sub>), and panels (m)–(p) changes in USW explained by changes in albedo ( $\Delta$  USW<sub>b</sub>). Only statistically significant results at the 5% level of significance are shown (insignificant masked in white); statistically significant grid cells for  $\Delta$ USW,  $\Delta$  USW<sub>DSW</sub>, and  $\Delta$  USW<sub>b</sub> are identical. Values within an interval (-0.1,0.1) W m<sup>-2</sup> are also masked in white.



**Figure 7.** Changes in decadal means (calculated from daily means) 2001–2021 minus 1980–2000, May–June–July. Row (i) shows cloud condensate content (CCC, vertically integrated cloud liquid water + ice), row (ii) shows surface albedo, and row (iii) sea-ice concentration (SIC). Only statistically significant results at the 5% level of significance are shown (insignificant masked in white). In rows (ii) and (iii), values within an interval (-0.01, 0.01) are also masked in white.

## 4 Discussion

### 4.1 Differences between reanalyses in the effects of sea-ice concentration on surface upward longwave radiation

265 ~~Our results indicated~~ We found negative correlation between SIC and ULW in all seasons and generally the highest sensitivity of ULW to SIC in the cold season November–April in the Central Arctic (Figs. 1 and S1) ~~—over  $150 \text{ W m}^{-2}$  ULW per  $0.1$  change in SIC. Utilizing JRA-55 data, the linear bilateral ODR model did not converge in large areas due to the data set's binary representation of the sea ice, however, in the areas of adjacent Arctic seas, where the model returned values of the slope between the two variables, the results were in agreement with other reanalyses. In the warm season May–October, the sensitivity~~  
270 ~~of ULW to SIC was generally lower, up to  $80 \text{ W m}^{-2}$  ULW per  $0.1$  change in SIC (Figs. S3 and S4).~~ The magnitude of the highest sensitivities of ULW to SIC was similar ~~amongst~~ among reanalyses, though their spatial extent somewhat differed.

The effect of Arctic SIC on ULW mostly decreased in all seasons between 1980–2000 and 2001–2021 due to SIC decline and warming of the sea-ice surface, however, we ~~noted also~~ also noted an increased sensitivity of ULW to SIC in the Central Arctic, north and northeast of Greenland. As shown in Fig. 2 (Point 2) the daily SIC in November–December–January increased  
275 in the Atlantic sector of the Central Arctic between 1980–2000 and 2001–2021 in ERA5, MERRA-2, and NCEP/CFSR. Fig. ~~S10–S11~~ row iii, also indicates statistically significant decadal increase in SIC in this area in February–March–April. Greater daily SIC is then directly connected to increased sensitivity of ULW to SIC. The SIC increase may be related to thinning of the Arctic sea ice, which reduces the internal resistance of the ice field, allowing certain atmospheric and oceanic forcing to generate faster ice drift (Leppäranta, 2011). Higher drift speeds along the Transpolar Drift Stream (TDS) favour increased  
280 accumulation of ice north of Greenland (Kwok, 2015), resulting in increased SIC. Another potential factor favouring faster ice drift is increased wind speeds along TDS (Smedsrud et al., 2017). However, trends in the wind speeds are sensitive to the region and period addressed (Spreen et al. (2011); Vihma et al. (2012)). ~~—Comparing the variance in ULW explained by SIC and  $T_{\text{ice}}$  in November–April in ERA5, MERRA-2, and NCEP/CFSR, we found very good agreement between these three data sets in that variance in  $T_{\text{ice}}$  was mostly responsible for variance in ULW ( $R^2$  over 90 %) in areas, where SIC was very high. In~~  
285 ~~the marginal ice zone, however, SIC explained around 60 % of variance in ULW while  $T_{\text{ice}}$  only up to 30 % (Figs. 3, S5).~~

### 4.2 Differences between reanalyses in the effects of sea-ice concentration and clouds on surface upward shortwave radiation

~~We found~~ Our results indicated positive correlation between SIC and USW in all seasons ~~and both study periods as expected. Similarly to the effect of SIC on ULW, with the highest sensitivity of USW to SIC in May–June–July in the Central Arctic (Fig.~~  
290 ~~5). The magnitude of~~ the effect of SIC on USW ~~also was similar in all reanalyses and~~ mostly weakened between 1980–2000 and 2001–2021. While the sea-ice and its surface albedo decline plays an undeniable role in the weakening of this effect, ~~also~~ decadal changes in DSW must also be considered when assessing decadal changes in USW.

Considering May–June–July, we found the magnitude of decadal change in mean USW (~~more than  $-25 \text{ W m}^{-2}$~~ ) and its spatial pattern similar among reanalyses in adjacent Arctic seas, however, this variable somewhat differed in the Central Arctic  
295 (Fig. ~~5e–h~~ 6e–h). Namely ERA5 and NCEP/CFSR showed decadal increase in the mean USW (~~around  $+10 \text{ W m}^{-2}$~~ ) north of

Greenland and Canadian archipelago, whereas JRA-55 indicated decadal reduction in the mean USW (around  $-10 \text{ W m}^{-2}$ ) where other reanalyses did not show significant changes. ~~As the magnitude of changes in USW explained by changes in DSW ( $\Delta \text{USW}_{\text{DSW}}$ ) was very similar among all reanalyses, the differences between these data sets in decadal seasonal changes in USW were mostly attributable to decadal changes in the surface albedo (Figs. 5, S9, S11). These Such results are similar~~  
300 to those of Cao et al. (2016) who considered the surface albedo product from the Satellite Application Facility on Climate Monitoring clouds, albedo, and radiation data set (CLARA-SAL) additionally to reanalyses data from 1982–2009. According to their findings, JRA-55 data agreed the best with the satellite observations, which did not show any increase in annual surface albedo ~~in the Central Arctic~~, north of Greenland and Canadian archipelago that we saw in ERA5 and NCEP/CFRS data. ~~—In our analyses of decadal seasonal changes, we found the largest differences between reanalyses in CCC (row i in Figs. 6, S10, S12). However, in all seasons, the magnitude of  $\Delta \text{USW}_{\text{DSW}}$  was very similar among all reanalyses, so from the point of view of solar radiation, clouds did not seem to be a key factor for the inter-reanalysis differences in decadal seasonal changes—~~  
305

### 4.3 The role of clouds on surface radiative fluxes and their differences between reanalyses

The clouds in the Arctic have typically positive net radiative effect on the surface for most of the year, as they have more impact by emitting longwave radiation towards the surface (DLW) and warming it than cooling it by reflecting the shortwave  
310 radiation back to space (Wendish et al. (2019); Morrison et al. (2019)). In May–June–July, however, incoming solar radiation in the Arctic is very high and clouds regulate the melting of sea ice and partly offset the strength of the sea ice–albedo feedback (Choi et al., 2020). The sign and strength of the radiative effect of clouds mostly depend on the cloud fraction, longevity, opacity (liquid/ice phase partitioning), and temperature of the cloud layer. The presence and properties of clouds have potential to considerably affect the surface and near-surface temperature and humidity. As we showed in Figs. 3 and S5, in areas with  
315 high SIC, changes in  $T_{\text{ice}}$  are important for explaining the variance in ULW in November–April, and these may be to a large part driven by changes in clouds. At the same time, SIC also affects the formation of clouds, via turbulent surface fluxes of sensible and latent heat. As shown in observational studies by Palm et al. (2010) and Liu et al. (2012), and in the study of Schweiger et al. (2008) who used reanalysis data from ERA40 (predecessor of ERA5), cloud cover variability near the sea ice margins is strongly linked to sea-ice variability and areas with increased mid-level cloudiness coincide with those of recent  
320 sea-ice decline. Also in our results, throughout the seasons, we saw the decadal increase in CCC in areas of strong SIC decline, although, reanalyses did not always agree on the magnitude or spatial extent of this increase. The increase in CCC is in line with Sledd and L’Ecuyer (2021).

Despite their importance for the Arctic surface energy budget, the clouds appear to be one of the largest sources of uncertainty as a variable in reanalyses and as a component of the Arctic climate system. This is mostly because the retrieval of cloud  
325 fraction and cloud properties (such as optical depth, top pressure, or cloud condensate content) from satellite measurements includes considerable uncertainties when using different sensors or even different approaches to derive the data from measured radiances (Devasthale et al., 2020). Also the insufficiency of supporting ground-based observational network in the Arctic contributes to the uncertainties. In our study, we only calculated decadal seasonal differences in mean CCC, ~~however, but~~ even by using this simple calculation and just one cloud parameter, we noted ~~the a large~~ spread in values between the reanalyses (row i

330 in Figs. 7, S11, S13). However, in all seasons, the magnitude of changes in USW explained by changes in DSW ( $\Delta USW_{DSW}$ ) was very similar among reanalyses (panels i–l in Figs. 6, S10, S12), so from the point of view of solar radiation, clouds did not seem to be a key factor for the inter-reanalysis differences in decadal seasonal changes.

In reality, aerosols affect the radiative properties of Arctic clouds (Garrett and Zhao, 2006). These effects have under-  
gone notable changes due to shifts in aerosol sources and regional atmospheric conditions (Warneke et al. (2010); Stohl et al. (2013)). Among the reanalyses applied in this study, MERRA-2 is based on daily assimilation of aerosol data, whereas ERA5, JRA-55, and NCEP/CFSR apply climatological aerosol concentrations. In principle, it should be possible to distinguish the contribution of aerosols to the radiative transfer and its seasonal and decadal changes, however, the output available from the reanalyses is not sufficient for such analyses.

#### 340 4.4 The role of surface albedo and its differences between reanalyses

Surface albedo (the proportion of incident shortwave radiation that is reflected back to space by the surface) is a key component of Arctic climate system. This property of the surface is the most important in May–June–July when the incoming shortwave radiation peaks and low albedo allows a much larger part of it to penetrate into (and warm) the surface. While the snow and sea ice and their properties control the surface albedo, at the same time, surface albedo controls the mass balance of snow and sea  
ice. This effect has a seasonal cycle, when (1) the bare sea ice with large amount of melt ponds and lower albedo during the  
melt season accelerates further ice melt by allowing more shortwave radiation to be absorbed, while (2) the dry snow on top  
of the sea ice generates greater surface albedo before and after the melt season, protecting the sea ice from shortwave radiative  
warming. Pistone et al. (2014) showed the close relationship of SIC and surface albedo in satellite data from The Clouds and  
Earth’s Radiant Energy System (CERES) and the Special Sensor Microwave Imager (SSM/I) and our results demonstrated that  
the patterns of diminishing SIC coincided with the patterns of the surface albedo decrease (rows i–ii and iii in Figs. 6, S10,  
S127, S11, S13).

The albedo of the sea ice is parameterised in JRA-55 and NCEP/CFSR, considering summer melt ponds and surface temperature, whereas in ERA5 and MERRA-2, it has a prescribed seasonal cycle that is the same for the whole study period of our analyses. Pistone et al. (2014) observed pan-Arctic darkening with clear-sky albedo decreasing from 0.39 to 0.33 and all-  
sky albedo decreasing from 0.54 to 0.48 during 1979–2011. These findings and their consequences for the prescribed surface  
albedo in reanalyses are demonstrated in the the comparison study by Pohl et al. (2020), who utilized satellite data from Medium  
Resolution Imaging Spectrometer (MERIS) to derive the albedo of Arctic sea ice. In their analyses, utilizing data from May to  
September 2003–2011, ERA5 was found to generally overestimate the albedo of first-year ice and underestimate the albedo of  
multiyear ice. Overestimation of the albedo likely happens due to not accounting (1) for the warming of the sea ice and (2) for  
the increasing amount of melt ponds on top of the sea ice during the melt seasons in recent decades. Underestimation of the  
albedo may be connected to sea-ice thickness prescribed in—In our analyses, we observed differences up to around 0.2 in June  
albedo at the North Pole in both daily and monthly means between MERRA-2 and JRA-55 and around 0.1 between MERRA-2  
and ERA5(1.5 m) that, in reality, is likely higher in the multiyear ice in the Central Arctic, north of Greenland generating

365 higher real albedo measured by MERIS, and NCEP/CFSR (Table 2, Fig. S9). These findings indicate a large uncertainty in the representation of the Arctic surface energy budget in these data sets during summer.

## 5 Conclusions

In the present study, we quantified the uncertainties in the effects of Arctic sea-ice concentration on surface radiative fluxes as represented in four atmospheric reanalyses, a complement to Uhlíková et al. (2024), where we addressed turbulent surface fluxes of sensible and latent heat. Our results showed the greatest sensitivity of surface upward longwave radiation to SIC in the cold season November–April (over  $150 \text{ W m}^{-2}$  per  $-0.1$  change in SIC) and greatest sensitivity of surface upward shortwave radiation to SIC in May–July (over  $100 \text{ W m}^{-2}$  USW per  $0.1$  change in SIC). We found that the effect of SIC on both surface upward longwave and shortwave radiation has mostly weakened in all seasons between the study periods of 1980–2000 and 2001–2021. Unlike in the case of the effects of SIC on turbulent surface fluxes, we did not find ~~particularly~~ generally higher sensitivity of surface upward radiative fluxes to SIC in NCEP/CFSR (which includes both modelled sea-ice thickness and snow depth on the sea ice and accounts for their insulating effects) compared to other reanalyses (which assume a constant sea-ice thickness and do not account for the snow on sea ice).

Furthermore, we analysed decadal changes in surface downward and upward shortwave radiation and quantified differences among reanalyses in these variables and additionally in the surface albedo, sea-ice concentration, and cloud condensate content. These analyses indicated that approximately 80 % of the decadal decrease in upward shortwave radiation in May–July was caused by a decrease in surface albedo, controlled by SIC decrease, and the rest was caused by a decrease in downward shortwave radiation due to increase in cloudiness, mostly close to sea ice margins. CCC showed the largest uncertainty among reanalyses in all seasons, however, the magnitude of decadal changes in surface upward shortwave radiation explained by changes in surface downward shortwave radiation was very similar among reanalyses. Accordingly, from the point of view of solar radiation, clouds did not seem to be a key factor for the inter-reanalysis differences in decadal seasonal changes.

385 Expanding quantitative knowledge on differences in the representation of the Arctic surface energy budget in atmospheric reanalyses is needed, because the Arctic amplification of climate warming is primarily surface-based (Serreze et al. (2009); Taylor et al. (2022)) and reanalyses are broadly utilized and relied upon in studies on past climate and related processes in the Arctic.

*Code and data availability.* <https://doi.org/10.5281/zenodo.11565044> (Uotila, Uhlíková, 2024), and [https://a3s.fi/uhlitere-2000789-pub/\\*](https://a3s.fi/uhlitere-2000789-pub/*)  
390 (last access: 11 June 2024) (Hersbach et al., 2023; Japan Meteorological Agency, 2013; GMAO, 2015a, b, c; Saha et al., 2010b, 2011).  
(To download a desired file, the name of it must be entered after the last forward slash, instead of \*. Names of files can be found in codes.  
Data description can be found at [https://a3s.fi/uhlitere-2000789-pub/README2\\_data2.odt](https://a3s.fi/uhlitere-2000789-pub/README2_data2.odt), last access: 13 May 2024.

*Author contributions.* TU prepared the manuscript with contributions of TV, PU, and AYK. TV, PU, and AYK designed the concept of the  
study with contributions of TU. PU developed the code with the contribution of TU. TU collected and processed data and performed analyses.

395 *Competing interests.* The authors declare that they have no conflict of interest.

*Disclaimer.* Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or  
data it contains.

*Acknowledgements.* During most of the work, TU was a university-funded doctoral researcher at the University of Helsinki. The work of  
AYK, PU, TV and, in the final stage, TU was supported by the European Commission's Horizon 2020 Framework Programme (PolarRES;  
400 grant no. 101003590).  
Hersbach et al. (2023) was downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store (2024). The results contain  
modified Copernicus Climate Change Service information 2023. Furthermore, we acknowledge the providers of the data of the other three  
reanalyses used in our study: Japan Meteorological Agency, the National Center for Atmospheric Research (JRA-55, NCEP/CFSR, CFSv2),  
and the Global Modeling and Assimilation Office (MERRA-2).



## References

- Andreas, E. L., Persson, P. O. G., Grachev, A. A., Jordan, R. E., Horst, T., Guest, P. S., and Fairall, C.: Parameterizing Turbulent Exchange over Sea Ice in Winter, *J. Hydrometeorol.*, 11, 87–104, <https://doi.org/10.1175/2009JHM1102.1>, 2010.
- 410 Boggs, P. T., Donaldson, J. T., Schnabel, R. B., and Spiegelman, C. H.: A Computational Examination of Orthogonal Distance Regression, *Journal of Econometrics*, 38, 169–201, <https://www.sciencedirect.com/science/article/pii/0304407688900322>, 1988.
- Bretherton, C. S., Widmann, M., Dymnikov, V. P., Wallace, J. M., and Bladé, I.: The Effective Number of Spatial Degrees of Freedom of a Time-Varying Field, *Journal of Climate*, 12, 1990–2009, [https://journals.ametsoc.org/view/journals/clim/12/7/1520-0442\\_1999\\_012\\_1990\\_tenosd\\_2.0.co\\_2.xml](https://journals.ametsoc.org/view/journals/clim/12/7/1520-0442_1999_012_1990_tenosd_2.0.co_2.xml), 1999.
- Cao, Y., Liang, S., He, T., and Chen, X.: Evaluation of Four Reanalysis Surface Albedo Data Sets in Arctic Using a Satellite Product, *IEEE Geoscience and Remote Sensing Letters*, 13, <https://doi.org/10.1109/LGRS.2016.2515159>, 2016.
- 415 Choi, Y.-S., Hwang, J., Ok, J., Park, D.-S. R., Su, H., Jiang, J. H., Huang, L., and Limpasuvan, V.: Effect of Arctic clouds on the ice-albedo feedback in midsummer, *Int. J. Climatol.*, 40, 4707–4714, <https://doi.org/10.1002/joc.6469>, 2020.
- Devasthale, A., Sedlar, J., Tjernström, M., and Kokhanovsky, A.: A Climatological Overview of Arctic Clouds, in: *Physics and Chemistry of the Arctic Atmosphere*, edited by Alexander Kokhanovsky, C. T., chap. 5, pp. 331–360, Springer International Publishing, 2020.
- 420 Di Biagio, C., Pelon, J., Blanchard, Y., Loyer, S., Hudson, S. R., Walden, V. P., et al.: Toward a better surface radiation budget analysis over sea ice in the high Arctic Ocean: a comparative study between satellite, reanalysis, and local-scale observations, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD032555>, 2021.
- Duynkerke, P. and de Roode, S.: Surface energy balance and turbulence characteristics observed at the SHEBA Ice Camp during FIRE III, *Journal of Geophysical Research: Oceans*, 106, 15 313–15 322, <https://doi.org/10.1029/2000JD900537>, 2001.
- 425 Ebert, E. E. and Curry, J. A.: An intermediate one-dimensional thermodynamic sea ice model for investigating ice-atmosphere interactions, *Journal of Geophysical Research: Oceans*, 98, 10 085–10 109, <https://doi.org/10.1029/93JC00656>, 1993.
- ECMWF: IFS Documentation CY41R2 - Part IV: Physical Processes, 4, ECMWF, <https://doi.org/10.21957/tr5rv27xu>, 2016.
- Garrett, T. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes, *Nature*, 440, 787–789, <https://doi.org/10.1038/nature04636>, 2006.
- 430 GMAO, Global Modeling, A. O.: Tavgl\_2d\_flux\_Nx: MERRA-2 2D, 1-Hourly, Time-Averaged, Single-Level Assimilation, Single-Level Diagnostics., <https://doi.org/10.5067/7MCPBJ41Y0K6>, 2015a.
- GMAO, Global Modeling, A. O.: Tavgl\_2d\_slv\_Nx: MERRA-2 2D, 1-Hourly, Time-Averaged, Single-Level Assimilation, Single-Level Diagnostics., <https://doi.org/10.5067/VJAFPLI1CSIV>, 2015b.
- GMAO, Global Modeling, A. O.: Tavgl\_2d\_rad\_Nx: MERRA-2 2D, 1-hourly, Time-Averaged, Single-Level Assimilation, Surface Flux Diagnostics V5.12.4, <https://doi.org/10.5067/Q9QMY5PBNV1T>, 2015c.
- 435 Graham, R. M., Cohen, L., Ritzhaupt, N., Segger, B., Graversen, R. G., Rinke, A., Walden, V. P., Granskog, M. A., and Hudson, S. R.: Evaluation of Six Atmospheric Reanalyses over Arctic Sea Ice from Winter to Early Summer., *Journal of Climate*, 32, 4121–4143, <https://doi.org/10.1175/JCLI-D-18-0643.1>, 2019.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D.,  
440 Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, <https://doi.org/10.1002/qj.3803>, 2020.

- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., J., M. S., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present., <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- 445 JMA, J. M. A.: JRA-55: Japanese 55-year Reanalysis, Daily 3-Hourly and 6-Hourly Data., <https://doi.org/10.5065/D6HH6H41>, 2013.
- Jonassen, M. O., Välisuo, I., Vihma, T., Uotila, P., Makshtas, A. P., and Launiainen, J.: Assessment of atmospheric reanalyses with independent observations in the Weddell Sea, the Antarctic, *Journal of Geophysical Research: Atmospheres*, 124, 12 468–12 484, <https://doi.org/10.1029/2019JD030897>, 2019.
- Kwok, R.: Sea ice convergence along the Arctic coasts of Greenland and the Canadian Arctic Archipelago: Variability and extremes (1992–2014), *Geophysical Research Letters*, 42, 7598–7605, <https://doi.org/10.1002/2015GL065462>, 2015.
- 450 Leppäranta, M.: *The Drift of Sea Ice*, Springer, Berlin Heidelberg, <https://doi.org/10.1007/978-3-642-04683-4>, 2011.
- Liang, S., Zhang, X., Xiao, Z., Cheng, J., Liu, Q., and Zhao, X.: Longwave Emissivity, in: *Global LAnd Surface Satellite (GLASS) Products*, Springer Briefs in Earth Sciences, pp. 63–77, Springer, Cham, [https://doi.org/10.1007/978-3-319-02588-9\\_4](https://doi.org/10.1007/978-3-319-02588-9_4), 2014.
- Light, B., Smith, M. M., Perovich, D. K., Webster, M. A., Holland, M. M., Linhardt, F., Raphael, I. A., Clemens-Sewall, D., Macfarlane, A. R., Anhaus, P., and Bailey, D. A.: Arctic sea ice albedo: Spectral composition, spatial heterogeneity, and temporal evolution observed during the MOSAiC drift, *Elementa: Science of the Anthropocene*, 10, 000 103, <https://doi.org/10.1525/elementa.2021.000103>, 2022.
- 455 Liu, Y., Key, J. R., Liu, Z., Wang, X., and Vavrus, S. J.: A cloudier Arctic expected with diminishing sea ice, *Geophysical Research Letters*, 39, L05 705, <https://doi.org/10.1029/2012GL051251>, 2012.
- Makshtas, A. P.: *The heat budget of the Arctic ice in the winter.*, Cambridge, International Glaciological Society, 1991.
- 460 Morrison, A. L., Kay, J. E., Frey, W. R., Chepfer, H., and Guzman, R.: Cloud response to Arctic Sea ice loss and implications for future feedback in the CESM1 climate model., *Journal of Geophysical Research: Atmospheres*, 124, 1003–1020, <https://doi.org/10.1029/2018JD029142>, 2019.
- Mortin, J., Svensson, G., Graverson, R. G., Kapsch, M.-L., Stroeve, J. C., and Boisvert, L. N.: Melt onset over Arctic sea ice controlled by atmospheric moisture transport, *Geophysical Research Letters*, 43, 6636–6642, <https://doi.org/10.1002/2016GL069330>, 2016.
- 465 Palm, S. P., Strey, S. T., Spinhirne, J., and Markus, T.: Influence of Arctic sea ice extent on polar cloud fraction and vertical structure and implications for regional climate, *Journal of Geophysical Research*, 115, D21 209, <https://doi.org/10.1029/2010JD013900>, 2010.
- Parkinson, C. L.: Arctic sea ice coverage from 43 years of satellite passive-microwave observations, *Frontiers in Remote Sensing*, 3, 1021 781, <https://doi.org/10.3389/frsen.2022.1021781>, 2022.
- Perovich, D. K., Nghiem, S. V., Markus, T., and Schweiger, A.: Seasonal evolution and interannual variability of the local solar energy absorbed by the Arctic sea ice–ocean system, *Journal of Geophysical Research*, 112, C03 005, <https://doi.org/10.1029/2006jc003558>, 2007.
- 470 Persson, P. O. G.: Onset and end of the summer melt season over sea ice: Thermal structure and surface energy perspective from SHEBA, *Climate Dynamics*, 39, 1349–1371, <https://doi.org/10.1007/s00382-011-1196-9>, 2012.
- Persson, P. O. G., Fairall, C. W., Andreas, E. L., Guest, P. S., and Perovich, D. K.: Measurements near the Atmospheric Surface Flux Group tower at SHEBA: Near-surface conditions and surface energy budget, *Journal of Geophysical Research*, 107, 8045, <https://doi.org/10.1029/2000JC000705>, 2002.
- 475 Pistone, K., Eisenman, I., and Ramanathan, V.: Observational determination of albedo decrease caused by vanishing Arctic sea ice, *Proceedings of the National Academy of Sciences*, 111, 3322–3326, <https://doi.org/10.1073/pnas.1318201111>, 2014.

- 480 Pohl, C., Istomina, L., Tietsche, S., Jäkel, E., Stapf, J., Spreen, G., and Heygster, G.: Broadband albedo of Arctic sea ice from MERIS optical data, *The Cryosphere*, 14, 165–182, <https://doi.org/10.5194/tc-14-165-2020>, 2020.
- Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M., Sela, J., and Goldberg, M.: The NCEP climate forecast system reanalysis, *Bulletin of The American Meteorological Society*, 91, <https://doi.org/10.1175/2010BAMS3001.1>, 2010.
- 485 Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., van den Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: NCEP Climate Forecast System Reanalysis (CFSR) 6-hourly Products, January 1979 to December 2010., <https://doi.org/10.5065/D69K487J>, 2010.
- 490 Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H.-Y., Juang, H.-M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., van den Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: NCEP Climate Forecast System Version 2 (CFSv2) 6-hourly Products., <https://doi.org/10.5065/D61C1TXF>, 2011.
- 495 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., ya Chuang, H., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W., Chen, M., and Becker, E.: The NCEP Climate Forecast System Version 2, *Journal of Climate*, 27, 2185 – 2208, <https://journals.ametsoc.org/view/journals/clim/27/6/jcli-d-12-00823.1.xml>, 2014.
- Schweiger, A. J., Lindsay, R. W., Vavrus, S., and Francis, J. A.: Relationships between Arctic Sea Ice and Clouds during Autumn, *J. Climate*, 21, 4799–4810, <https://doi.org/10.1175/2008JCLI2156.1>, 2008.
- 500 Senkova, A., Rontu, L., and Savijarvi, H.: Parameterization of orographic effects on surface radiation in HIRLAM, *Tellus A: Dynamic Meteorology and Oceanography*, 59, 279–291, <https://doi.org/10.1111/j.1600-0870.2007.00235.x>, 2007.
- Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N., and Holland, M. M.: The emergence of surface-based Arctic amplification, *The Cryosphere*, 3, 11–19, <https://tc.copernicus.org/articles/3/11/2009/tc-3-11-2009.pdf>, 2009.
- 505 Sledd, A. and L'Ecuyer, T. S.: Emerging trends in Arctic solar absorption, *Geophysical Research Letters*, 48, e2021GL095813, <https://doi.org/10.1029/2021GL095813>, 2021.
- Smedsrud, L. H., Halvorsen, M. H., Stroeve, J. C., Zhang, R., and Kloster, K.: Fram Strait sea ice export variability and September Arctic sea ice extent over the last 80 years, *The Cryosphere*, 11, 65–79, <https://doi.org/10.5194/tc-11-65-2017>, 2017.
- Spreen, G., Kwok, R., and Menemenlis, D.: Trends in Arctic sea ice drift and role of wind forcing: 1992–2009, *Geophysical Research Letters*, 38, L19 501, <https://doi.org/10.1029/2011GL048970>, 2011.
- 510 Stohl, A. et al.: Arctic aerosols from mixed-phase clouds: impacts and variability, *Atmospheric Chemistry and Physics*, 13, 10 959–10 978, <https://doi.org/10.5194/acp-13-10959-2013>, 2013.
- Taylor, P. C., Boeke, R. C., Boisvert, L. N., Feldl, N., Henry, M., Huang, Y., Langen, P. L., Liu, W., Pithan, F., Sejas, S. A., and Tan, I.: Process Drivers, Inter-Model Spread, and the Path Forward: A Review of Amplified Arctic Warming, *Frontiers in Earth Science*, 9, <https://doi.org/10.3389/feart.2021.758361>, 2022.
- 515

- Uhlíková, T., Vihma, T., Karpechko, A. Y., and Uotila, P.: Effects of Arctic sea-ice concentration on turbulent surface fluxes in four atmospheric reanalyses, *The Cryosphere*, 18, 957–976, <https://doi.org/10.5194/tc-18-957-2024>, 2024.
- Vihma, T.: Subgrid parameterization of surface heat and momentum fluxes over polar oceans, *J. Geophys. Res.*, 100, 1995.
- 520 Vihma, T., Tisler, P., and Uotila, P.: Atmospheric forcing on the drift of Arctic sea ice in 1989–2009, *Geophysical Research Letters*, 39, L02 501, <https://doi.org/10.1029/2011GL050118>, 2012.
- Walsh, J. E., Chapman, W. L., and Portis, D. H.: Arctic Cloud Fraction and Radiative Fluxes in Atmospheric Reanalyses, *Journal of Climate*, 22, 2316–2334, <https://doi.org/10.1175/2008JCLI2213.1>, 2009.
- Warneke, C., Bahreini, R., Brioude, J., Brock, C. A., de Gouw, J. A., Fahey, D. W., Froyd, K. D., Holloway, J. S., Middlebrook, A. M., Miller, L., Montzka, S. A., Schwarz, J. P., Spackman, J. R., and Veres, P. R.: Biomass burning in Siberia and Kazakhstan as an  
525 important source for haze over the Alaskan Arctic in April 2008, *Geophysical Research Letters*, 37, L02 813, <https://doi.org/10.1029/2009GL041816>, 2010.
- Wendish, M., Macke, A., and Ehrlich, A. e. a.: The Arctic Cloud Puzzle: Using ACLOUD/PASCAL Multiplatform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification., *Bulletin of American Meteorological Society*, 100, 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.
- 530 Wilks, D. S.: “The Stippling Shows Statistically Significant Grid Points”: How Research Results are Routinely Overstated and Overinterpreted, and What to Do about It, *Bull. Amer. Meteor. Soc.*, 97, 2263–2273, <https://doi.org/10.1175/BAMS-D-15-00267.1>, 2016.
- Yeo, H., Kim, M.-H., Son, S.-W., Jeong, J.-H., Yoon, J.-H., Kim, B.-M., and Kim, S.-W.: Arctic cloud properties and associated radiative effects in the three newer reanalysis datasets (ERA5, MERRA-2, JRA-55): Discrepancies and possible causes, *Atmospheric Research*, 270, 106 080, <https://doi.org/10.1016/j.atmosres.2022.106080>, 2022.