

# Inter-model differences in 21st Century Glacier Runoff for the World's Major River Basins

Finn Wimberly<sup>1</sup>, Lizz Ultee<sup>1</sup>, Lilian Schuster<sup>2</sup>, Matthias Huss<sup>3,4,5</sup>, David R. Rounce<sup>6</sup>, Fabien Maussion<sup>7,2</sup>, Sloan Coats<sup>8</sup>, Jonathan Mackay<sup>9,10</sup>, and Erik Holmgren<sup>11</sup>

<sup>1</sup>Department of Earth & Climate Sciences, Middlebury College, Middlebury, VT, USA

<sup>2</sup>Department of Atmospheric and Cryospheric Sciences (ACINN), Universität Innsbruck, Innsbruck, Austria

<sup>3</sup>Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Zurich, Switzerland

<sup>4</sup>Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland

<sup>5</sup>Department of Geosciences, University of Fribourg, Fribourg, Switzerland

<sup>6</sup>Department of Civil Engineering, Carnegie Mellon University, Pittsburgh, PA, USA

<sup>7</sup>School of Geography, University of Bristol, Bristol, UK

<sup>8</sup>Department of Earth Sciences, University of Hawaii at Manoa, Honolulu, HI, USA

<sup>9</sup>British Geological Survey, Environmental Science Centre, Keyworth, UK

<sup>10</sup>School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, UK

<sup>11</sup>Department of Space, Earth, and Environment, Chalmers University of Technology, Gothenburg, Sweden

**Correspondence:** Finn Wimberly (fwimberly@middlebury.edu)

**Abstract.** Projected glacier change has important downstream consequences, including sea level rise, changing freshwater supply, and loss of important cultural sites. While the glacier contribution to global sea level rise and associated uncertainties have been quantified in model intercomparison studies, comparatively less focus has been directed towards the inter-annual changes in runoff caused by glacier recession. The observed effect of glacier runoff on basin-level water availability makes simulated future runoff a particularly consequential target for analysis. In this study, we compare century-scale runoff simulated by three global glacier evolution models. Aggregating annual glacier runoff contributions to 75 globally-distributed major river basins, we find that the three models agree closely in some basins but differ dramatically (up to a factor of 3.8) in others. However, when we analyze century-scale runoff changes relative to a glacier model's historical runoff baseline, annual runoff projections are much more consistent across glacier models. Glacier models project broadly consistent relative changes in seasonal runoff supply, with some differences across climatic regions. Estimates of the year of peak water are more consistent across glacier models (when driven by a climate model ensemble) than across individual climate forcings within a single glacier model. We identify the glacier models' different approaches to modifying precipitation forcing as the dominant source of inter-model differences in projected runoff. Our findings highlight the comparative roles of glacier evolution model, global climate model forcing, and emissions scenario as important sources of uncertainty across different metrics of projected glacier runoff. We recommend steps to account for glacier model uncertainties. For example, inter-glacier-model uncertainty in absolute annual runoff is large, but the year of projected peak water has much greater inter-climate-model uncertainty. We recommend that users pay particular attention to how a selected glacier model parametrizes and calibrates the glacier climatic mass balance in glacio-hydrological modelling efforts.

## 1 Introduction

20 Twenty first century glacier change has substantial downstream consequences, including sea-level rise (e.g. Marzeion et al., 2012; Gardner et al., 2013), changing freshwater supply (e.g. Kaser et al., 2010; Immerzeel et al., 2020), altering ecosystems (e.g. Bosson et al., 2023), (e.g. Bosson et al., 2023; Jacobsen et al., 2012) elevating natural hazard risk (e.g. Taylor et al., 2023), and loss of important cultural sites (e.g. Bosson et al., 2019). In recent decades the effects of these changes have already become visible. Cities such as La Paz (Kinouchi et al., 2019) and Santiago (McCarthy et al., 2022) have had to adopt severe  
25 water management strategies while numerous Himalayan communities have had to enact new preventative measures against glacial lake outburst floods (Ahmed et al., 2021). The diverse downstream consequences of glacier recession highlight the importance of refining projections of glacier change to support adaptation and mitigation strategies.

Though the relative importance of glacier runoff varies by basin (Immerzeel et al., 2020), glaciers are projected to provide critical drought buffering, especially in arid basins (Ultee et al., 2022), despite already surpassing ~~“peak” water~~ the year of  
30 greatest annual runoff (“peak water”) in many regions (Huss and Hock, 2018). Our ability to predict glacier mass and runoff changes is thus important for water planning and management. Unfortunately, global climate models (GCMs) do not simulate glacier change, and global hydrological models can only crudely represent it due to a lack of input data and parameterization constraints (van Tiel et al., 2020a). While global hydrological models can explicitly represent glaciers through coupling to a glacier evolution model, computational expense generally limits such efforts to a single hydrological model coupled to a single  
35 glacier model (Wiersma et al., 2022; Hanus et al., 2024). There is thus an urgent need to quantify uncertainties associated with projected glacier runoff from different global glacier evolution models (hereafter “glacier models”) to properly inform glacio-hydrological modeling studies.

Uncertainties in glacier runoff projections come from differences in glacier models as well as uncertainty in climate forcing. GCM-projected precipitation and land-surface runoff vary widely across GCMs and climate scenarios; in some regions, models  
40 disagree even on the sign of the expected changes (Lee et al., 2021; Wang et al., 2022). Runoff simulated by glacier models may either amplify or temper inter-GCM variability in projected hydrologic changes because glaciers respond nonlinearly to climate forcing (Oerlemans, 1989; Christian et al., 2022). The inherent nonlinearity of the glacier response is further obscured because glacier models use different methods to modify precipitation forcing and account for un-simulated processes. For example, “glacier-centric” models like the ones we study here, which simulate each of the world’s glaciers individually, must downscale  
45 and bias-correct coarse-gridded climate forcing data to the individual glacier scale. Each glacier model applies slightly different schemes to do so, calibrating parameters such as temperature lapse rates, temperature biases and multiplicative precipitation factors to match the observed glacier mass balance. The calibrated parameters often make climate biases or missing mass-balance processes implicit (Rounce et al., 2020b). Further, because only one observation per glacier is globally available for calibration (Hugonnet et al., 2021), all glacier models are over-parameterized. Thus, multiple combinations of parameters may  
50 match the observations equally well, but produce different runoff projections (Schuster et al., 2023a).

While previous glacier model intercomparison projects have revealed considerable differences in glacier mass projections due to differences in models (e.g., model parameterizations, initializations, calibration data and techniques, reanalysis data, and


bias corrections) (Hock et al., 2019; Marzeion et al., 2020), uncertainties in projected glacier runoff from a multi-glacier model ensemble have yet to be quantified. Here, we analyze the first estimate of glacier runoff projections for the three most advanced, globally capable glacier models, forced by an ensemble of CMIP6 GCMs and Shared Socioeconomic Pathways (SSPs), for 75 glaciated major river basins. For each basin, we analyze annual glacier runoff, the year of peak water, and seasonal runoff cycle projected by each glacier evolution model until the year 2100. We quantify the consistency of projections across glacier models by comparing the inter-glacier-model range with inter-GCM range. We also examine inter-SSP ranges to determine the effect of the future emission scenario on year of peak water. Ultimately, the study aims to summarize inter-model similarities and differences in runoff projections, discern general reasons for observed differences, and provide guidance for practitioners and hydrologists seeking to incorporate readily available glacier model projections into future planning.

## 2 Methods

We evaluate projected glacier runoff at annual and seasonal time scales for all 75 of the world's large river basins (> 3000 km<sup>2</sup>) with considerable glacier cover (> 30 km<sup>2</sup>). Glacier outlines are provided by the Randolph Glacier Inventory ("RGI" version 6; RGI Consortium, 2017), and we include all glaciers falling within Global Runoff Data Centre (2020) Major River Basins, including those in Europe, Asia, the Americas, and New Zealand (Figures 1 and A1). Here, we define glacier runoff as the sum of ice and snow melt and liquid precipitation minus refreezing (when modeled). We use a "fixed-gauge" station approach (e.g. Huss and Hock, 2018) where each model estimates runoff for a constant area such that it accounts for ice and snow melt as well as rainfall over the initially glaciated areas as the glacier retreats.

### 2.1 Glacier Evolution Models

The three glacier models included in our study are the Global Glacier Evolution Model ("GloGEM"; Huss and Hock, 2015), Open Global Glacier Model ("OGGM"; Maussion et al., 2019), and Python Glacier Evolution Model ("PyGEM"; Rounce et al., 2023). All three are "glacier-centric", simulating each of the world's glaciers individually. Each simulated glacier is initialized with a Randolph Glacier Inventory outline (RGI Consortium, 2017), calibrated using different reanalysis data and calibration strategies against glacier-wide mass balance from 2000–2019 for each glacier (Hugonnet et al., 2021), and forced in the future with an ensemble of GCMs and SSPs from CMIP6 (Eyring et al., 2016). Melt is estimated using a temperature-index approach and accumulation using a temperature threshold to distinguish rain from snow. GloGEM and PyGEM also include a parameterization to estimate refreezing, while OGGM does not. All three glacier models have previously been validated against seasonal mass balances from in-situ measurements of a few hundred glaciers worldwide (WGMS, 2020) showing good agreement at regional scales, even though deviations for individual glaciers can be considerable. All three models also bias correct the GCMs to align closely with the reanalysis dataset utilized during calibration by matching temperature (mean and standard deviation) and precipitation (mean) over a historical period (see e.g. Zekollari et al., 2019). Below we describe the key features of each glacier model; for further details, we refer readers to the associated description papers cited above and to [Zekollari et al. \(2024\)](#).



**Figure 1.** All 75 basins analyzed in the study. The color fill corresponds to the basin’s average annual glacier runoff over year of peak water as estimated by the entire period. Average runoff values are derived from the multi-glacier-model median of each glacier-model’s multi-GCM-median under the SSP2-4.5 scenario. This map was generated by our online interactive mapping tool (Wimberly, 2024b), available at link when published [https://fwimberly.shinyapps.io/Global\\_Glacier\\_Runoff/](https://fwimberly.shinyapps.io/Global_Glacier_Runoff/)

### 85 2.1.1 Global Glacier Evolution Model (GloGEM)

GloGEM estimates monthly snow accumulation, snow- and ice-melt, and refreezing for 10 m elevation bands on each glacier. Glacier dynamics are modeled using an empirical relation that describes ice thickness change as a function of the normalized elevation range (Huss and Hock, 2015). The model is forced with ERA5 reanalysis data (Hersbach et al., 2020) and is calibrated in a three-step calibration procedure. The main calibration parameters are the degree-day factors for relation of positive air temperature to melt, and a precipitation correction factor that accounts for differences between the reanalysis-cell forcing and the actual precipitation on the glacier. All model parameters are calibrated for every glacier individually to match observed glacier-wide mass balances (Hugonnet et al., 2021). The multiplicative precipitation correction is constrained to remain within bounds of approximately 1.0 to 2.5. These bounds vary between RGI regions and are chosen such that the misfit of modelled and measured winter balance as according to sparse observations obtained from the World Glacier Monitoring Service (Zemp et al., 2023; WGMS, 2024) is minimized. GloGEM employs separate degree-day factors for snow, firn, and ice, scaled by a multiplicative factor to account for their differing melt rates. The reanalysis climate forcing is used until 2020, after which GCM output forces the future evolution. A bias correction between the reanalysis data and future GCMs is applied based on the years 1980 to 2019. The glacier area (geometry) is assumed constant until the date associated with the RGI outline, after

which the glacier geometry evolves. The constant catchment area for the runoff computation is the RGI v6.0 glacier area in  
100 each basin.

### 2.1.2 Open Global Glacier Model (OGGM) - v1.6.1

OGGM is a modular, open source, community-based glacier evolution model framework that estimates the monthly mass  
balance using a temperature index model (Marzeion et al., 2012; Maussion et al., 2019). Glacier dynamics are modeled using  
the shallow ice approximation along elevation-band flowlines. The model is calibrated for every glacier individually to match  
105 observed glacier-wide mass balances (Hugonnet et al., 2021) when forced with W5E5v2.0 climate dataset (Lange et al., 2021).  
The glacier specific multiplicative precipitation factor is computed from an empirical function of total winter precipitation,  
leading to smaller factors for glaciers located in wetter grid points and larger factors for drier grid points. This relationship was  
derived by calibrating the model on 114 glaciers (Schuster et al., 2023a) with in-situ winter mass balance data (WGMS, 2024).  
The local precipitation factor can range from 0.1 to 10, but 90% of the global values lie between 1.6 and 6.7 (median: 3.6).  
110 High precipitation factor values may be partly explained by the fact that W5E5 is drier than ERA5 and that OGGM does not  
apply a precipitation gradient with elevation like PyGEM or GloGEM. [We analyse OGGM calibrates a single degree-day factor  
for temperature-index melt estimation, with no surface-type distinction. We analyze](#) the standard OGGM projection option of  
OGGMv1.6.1 (Maussion et al., 2023), which performs a bias correction between the reanalysis data and each GCM/scenario  
based on 2000–2019. OGGMv1.6.1 uses a dynamic spinup and calibration routine to initialise glacier states in the year 2000  
115 and ensure that glacier mass balance during the 2000–2019 historical simulation still matches observations taking elevation  
feedback into account [??\(Aguayo et al., 2023; Zekollari et al., 2024\)](#). The spin-up ensures that glacier areas at the inventory  
date are matched within 1%. The constant catchment area for the runoff computation is the glacier area in each basin in the  
year 2000 according to the dynamic spinup routine and may differ from the RGI. In this study, the GCM forcing is used from  
2000 onwards.

### 120 2.1.3 Python Glacier Evolution Model (PyGEM)

PyGEM is a modular, open source, glacier evolution model that estimates the monthly mass balance at  $\sim 10$  m elevation bins  
(Rounce et al., 2023). Sub-debris melt enhancement factors are used to account for the enhanced or reduced melting associated  
with debris thickness for debris-covered glaciers (Rounce et al., 2021). Glacier dynamics are modeled using OGGM's shallow  
ice approximation along elevation-band flowlines (OGGMv1.3). The model is calibrated using Bayesian inference (Rounce  
125 et al., 2020b, 2023) on observed glacier-wide mass balances (Hugonnet et al., 2021) forced by ERA5 reanalysis data (Lange  
et al., 2021). Prior distributions for each model parameter are determined at a subregional level based on a three-step calibration  
procedure for each glacier that constrains multiplicative precipitation factor between 0.5 and 5. During the Bayesian inference,  
the multiplicative precipitation factor is not constrained, but the choice of prior distribution ensures positivity. [PyGEM uses  
two degree-day factors to capture snow and ice melt, along with spatially dependent sub-debris melt enhancement factors,  
enhancing its ability to simulate debris-covered glaciers.](#) The GCM bias correction is performed for the period 2000–2019 and  
130 the GCM forcing is used from 2000 onwards. The initial glacier area from the RGI outline is assumed to be at 2000, i.e., unlike

the prior two models which used the specific RGI date for each glacier, after which the glacier evolves. The constant catchment area for the runoff computation is the RGI v6.0 glacier area in each basin.

### 2.1.4 Future Climatic Forcing

135 Each glacier model simulated monthly glacier runoff for all RGI glaciers from 2000–2100, forced by an ensemble of single realizations from 12 GCMs and four SSPs of the Coupled Model Intercomparison Project Phase 6 (CMIP6, Eyring et al., 2016). The SSPs include SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. The 12 GCMs were chosen to be consistent with prior studies (Compagno et al., 2022; Rounce et al., 2023) and included BCC-CSM2-MR, CESM2, CESM2-WACCM, EC-Earth3, EC-Earth3-Veg, FGOALS-f3-L, GFDL-ESM4, INM-CM4-8, INM-CM5-0, MPI-ESM1-2-HR, MRI-ESM2-0, and NorESM2-  
140 MM. For consistency, we used the “r1i1p1f1”-tagged ensemble member for each GCM from the CMIP6 archive (see Taylor et al., 2022, for description of *ripf* tags).

## 2.2 Metrics analyzed

We report annual runoff as the multi-GCM median and quartile values for each glacier evolution model, in each basin, for each SSP. Our results thus highlight the range and central tendency of projected annual runoff totals for each glacier model for the  
145 ensemble of GCMs. We also compare single-glacier-model percent-change runoff projections (i.e., for every glacier model, we divide the annual runoff by the average historical runoff):

$$\Delta_{\%}Q(y) = \frac{Q(y) - \hat{Q}}{\hat{Q}} \times 100\%, \quad (1)$$

where  $Q(y)$  is the single-glacier-model, single-GCM annual runoff in a given year  $y$ ; and  $\hat{Q}$  is the average annual runoff for the same glacier model (and GCM, if applicable) over the historical period (2000–2019). This eliminates baseline differences  
150 and allows for a more direct comparison of differences between the models’ inter-annual tendencies.

For both the annual and percent-change runoff projections, we calculated an explicit “multi-GCM range” by computing the spread (maximum - minimum) in single-GCM runoff projections for each year and averaging across the century and then across glacier models, for each basin. Similarly, our metric of “multi-glacier-model range” is the spread (maximum - minimum) of the multi-GCM median projections for the three glacier models, averaged across the century for each basin. We note that our  
155 12 forcing GCM realizations are a small sample of more than 250 realizations from the CMIP6 archive, while the three glacier models are the entire population available for global glacier runoff simulation; the expected distribution of glacier runoff for all CMIP6 realizations and all possible glacier model parameter sets is not known. As such, the multi-GCM and multi-glacier-model ranges we present do not give a complete characterization of uncertainties. Rather, we use them to contextualize whether the projections from different glacier models should be interpreted as “similar” in light of typical GCM-ensemble uncertainty.

160 Peak water, i.e., the year of maximum glacier runoff after which runoff declines, is calculated using a [20-year rolling mean 11-year rolling mean \(Huss and Hock, 2018\)](#). We computed the year of peak water from single-GCM time series and then found the multi-GCM median year for each glacier model and scenario. We also provide GCM, Glacier Model, and SSP

Ranges for year of peak water projections (see Fig. 4). The GCM Range (for year of peak water) is calculated by finding the spread (maximum - minimum) of single-GCM realizations for each glacier model and then taking the median value. We  
 165 calculated the glacier model range by finding the multi-GCM median year of peak water for each glacier model and then took the range of these three values. The SSP Range refers to the range across all glacier models' multi-GCM median peak water projections for all SSPs.

We aggregated the annual on- and off- glacier solid and liquid precipitation totals for each glacier model and basin over the historical period, 2000–2019. We report the mean annual precipitation over that period for each glacier model to highlight  
 170 differences in how the models process the raw precipitation from the climate forcing. We also calculate inter-glacier-model “relative” runoff and precipitation values. These values take the annual runoff (across the entire period) or annual precipitation (across the historical period) and divide by the mean of all three glacier models' projected annual runoff or historical precipitation. We then calculated the mean by summing the annual values and dividing by the number of years:

$$\text{Relative Q} = \frac{1}{100} \sum_{y=2000}^{2099} \frac{Q_y}{\bar{Q}_y}, \quad (2)$$

175 where  $Q_y$  indicates the single-GCM runoff for a single glacier model in a given year and  $\bar{Q}_y$  indicates the multi-glacier-model mean of single-GCM runoff for the same year. Note that we compute these values per GCM and per glacier model, and then take a multi-GCM median to arrive at a single value per glacier model for the projected period. Similarly,

$$\text{Relative P} = \frac{1}{20} \sum_{y=2000}^{2019} \frac{P_y}{\bar{P}_y}, \quad (3)$$


180 where  $P_y$  indicates multi-GCM median total precipitation as downscaled and bias-corrected by a single glacier model, summed over the initially glaciated area of the basin, for a year between 2000 and 2019, and  $\bar{P}_y$  indicates the multi-glacier-model mean of that quantity. Note that we compute these values for only a single forcing (the calibration dataset of each glacier model) and therefore arrive immediately at a single value per glacier model over the historical period. These metrics quantify how the magnitude of one model's runoff projection or input precipitation compares to the multi-glacier-model mean.

### 3 Results

#### 185 3.1 Annual Glacier Runoff by Basin


Across the majority of basins, the three glacier models project similar inter-annual runoff changes with a noticeable difference in magnitude such that each model's projected runoff appears to be translated up or down relative to the others (Figures 2 and A1). The glacier model that predicts the most runoff varies by river basin (e.g., Figure 2) with some regional consistency. OGGM projects the largest amount of glacier runoff across the entire century for nearly all basins located within Alaska,  
 190 Iceland, the European Alps, central Asia and the low latitudes. GloGEM projects the largest annual runoff totals for all basins located within North Asia, the Southern Andes, and New Zealand. PyGEM projects the most runoff in Scandinavia. [Analyses](#)

of climatological and geographical factors (e.g. historical aridity index, mean precipitation, central latitude, mean glacier altitude) did not reveal regionally consistent drivers of inter-glacier-model differences in runoff; we elaborate on two drivers of differences at global scale in the Discussion. While the evolution of runoff over the century differs between SSPs, the dependence of annual glacier runoff on SSP appears consistent across glacier models. Generally, maritime regions exhibit a more extreme dependence on emission scenarios. In most coastal North American basins (e.g. Yukon, Alsek, and Copper), more severe emission scenarios result in an increase in precipitation and thus an increase in annual runoff across the century (this manuscript Figure 2, and Wimberly, 2024b).



**Figure 2.** Projected multi-GCM median annual glacier runoff for each glacier model and SSP for selected basins: Yukon basin (1.2% initial glacier cover, North America), the Glomaa basin (0.7%, Europe), the Rhone basin (1.0% Europe), the Aral Sea basin (0.9%, Asia), and the Serrano basin (1.1%, South America). The shaded area shows the interquartile range of the 12 single-GCM projections. Annotations on each figure panel state the multi-glacier-model range (n=3, bold text) and multi-GCM range (n=12) of projections, both in  $\text{km}^3\text{yr}^{-1}$ , computed following Sect. 2.2. These basins were selected to sample the range of regions in which glacier models project more or less absolute runoff.

During the historical period (2000–2019), GloGEM outputs forced by only the ERA5 reanalysis are highly variable, while OGGM and PyGEM forced by GCM output produce smoother median projections with an inter-GCM range (e.g. Serrano basin in Fig. 2). The modeler choice to simulate the historical period with climate reanalysis (as in GloGEM) or bias-corrected GCM output (as in OGGM or PyGEM) affects the range in the baseline for future runoff projections.



**Figure 3.** Smoothed (20-year-11-year rolling mean) annual multi-GCM median glacier runoff projections as a percent change relative to the historical (2000–2019) average under SSP 2-4.5. The multi-glacier-model and multi-GCM range are given as in Fig. 2, however, in units of %.

Inter-glacier-model range in annual runoff is comparable to inter-GCM range in most basins (‘Glacier Model Range’ and ‘GCM Range’ annotations, Figure 2). For the Yukon, Glomaa, and Rhone basin, the inter-GCM ranges are slightly larger than the inter-glacier-model ranges. By contrast, the glacier model ranges are larger than the inter-GCM ranges in the Aral Sea and Serrano river basins, where the range in historical baseline runoff is also large.

Normalizing runoff as a percent change relative to the model’s historical mean dramatically reduces the range in glacier model projections (Figure 3). While the Yukon basin’s glacier model range was comparable to its GCM range, the normalized glacier model range was one-seventh of the normalized GCM range. Even more drastically, the Serrano basin’s projections have a glacier model range three times the GCM range, but normalization decreases inter-glacier-model range to one third the GCM range. Beyond significantly reducing inter-glacier-model uncertainty, normalization allows for a more direct visual comparison of differences in glacier model tendencies. For example, Figure 3 reveals that PyGEM runoff decreases much more rapidly from its projected year of peak water than the other two glacier models in the Aral Sea, Serrano, and Rhone basins. We

present both absolute and normalized examples here to illustrate their differences; we encourage those interested in a particular  
215 region to carefully study runoff series of both types (see Sect. 4 as well).

### 3.2 Timing of Peak Water

In most basins, the projected year of peak water is similar across glacier models (Figure 4 and A1). Generally, the inter-  
glacier-model range is minimal compared to the inter-GCM range (Figure 4(b)). For SSP-2.45, the median (across all basins)  
inter-GCM range is 40 years while the median inter-glacier-model range is 6 years. However, the timing of peak water does vary  
220 considerably between glacier models for a few basins (e.g., Clutha basin, see Fig. A1). These differences primarily appear in  
basins with minimal runoff variability over time, i.e., projected runoff is more-or-less constant over the century. In such basins,  
the year of peak water is predominantly determined by any shorter timescale fluctuations that persist through the smoothing  
process of the rolling mean analysis. The SSP range of the ~~Thus, these peak water estimates are determined primarily by GCM~~  
variability and have a high degree of uncertainty.


225 Basins that have already passed observed peak water may nevertheless show an early-century year of peak water is generally  
larger than the glacier model range, but smaller than the GCM range, in the projections. This discrepancy arises because glacier  
models are initialized under the assumption of equilibrium, whereas many glaciers are far from equilibrium. As a result, runoff  
may exhibit an artificial peak at the beginning of the simulation that does not align with real-world observations. The Skagit  
River basin, which is well past peak water based on observations, highlights this limitation in our simulated peak water years  
230 (Riedel and Larrabee, 2016; Pelto et al., 2022).

### 3.3 Seasonality


In most basins, all three glacier models show the seasonal glacier runoff peak shifting earlier by the end of the century (Figure  
5 and A3). GloGEM tends to maintain wider, less defined peaks, i.e., more months with a high fraction of maximum monthly  
runoff, while OGGM and PyGEM have narrower seasonal peaks at the end of the century compared with 2000–2019 (Figure  
235 A2). The shift in the timing of glacier runoff is highly affected by the variability in GCMs as the spread associated with  
individual GCMs is considerably wider at the end of the century relative to the start of the century (Figure 5).

Future changes in seasonal cycle show some differences across different climate regions. Many basins in the midlatitudes of  
North America, South America, and Europe tend to show some months of increasing runoff and some decreasing, consistent  
with a change in shape of the seasonal peak (e.g. Rhone, Figure 5). By contrast, basins in tropical South America have a weaker  
240 seasonal cycle overall, consistent with a tropical climate, and show relatively small changes in monthly runoff (e.g. Magdalena,  
Figure 5). A few basins in arid central Asia (e.g. Indus, Tarim He, Aral Sea) and maritime North America (Yukon, Copper,  
Skagit) show a net increase in magnitude of the seasonal glacier runoff peak in most simulations (Figure 5 and A2). Annual  
runoff projections for these basins peak in the mid- to late century and do not diminish much afterward (Figure A1).

In several heavily populated Asian basins (Yellow River, Yangtze, Brahmaputra, Ganges, Indigirka, Salween, Mekong) all  
245 three glacier models project the magnitude of the seasonal runoff peak to substantially decrease by the end of century (Figure  
5 and A2). Annual peak water in these basins occurs at the start of the century (Figure A1). As such, the 2000–2019 reference



**Figure 4.** Subfigure (a) shows the projected peak water year for all 75 basins (rows) for each GCM (columns) and glacier model (panels) for SSP 2-4.5. The GCMs (i.e., pixels) are ordered by ascending year of peak water, which varies across basins and glacier models (i.e., the first pixel in the first row is not necessarily the same GCM as the first pixel in the second row or the first pixel in the first row of the second panel). The black text laid over the pixels gives the multi-GCM median year of peak water. Subfigure (b) depicts the inter-GCM ( $n=12$ , at SSP-2.45), inter-glacier-model ( $n=3$ , at SSP-2.45), and inter-SSP ( $n=4$ ) range of year of peak water.



**Figure 5.** Variability in relative monthly glacier runoff for each glacier model for five example basins over 2000–2019 (top row), 2040–2059 (second row), 2080–2099 (middle-third row), and the change between these two the beginning of and end of the century periods (bottom row). Monthly glacier runoff for 2000–2019 and 2080–2099 is normalized by the maximum monthly runoff over the 2000–2019 period, computed for each glacier model and GCM individually. The percent change over the two time periods is expressed as a percentage of the 2000–2019 maximum monthly runoff, shown only for the multi-GCM median. Heavier lines show multi-GCM median for each glacier model; light lines show individual GCMs. Results for all basins are shown in Figures A2- A3.

period is likely on the ascending branch of the “peak water” curve, while the end of century is far past the peak, when melt-season runoff has tapered off (Huss and Hock, 2018).


There are several basins where one of the three glacier models projects a net increase in the seasonal runoff peak (e.g., 250 Susitna, Colville, Nushagak, Dramselva, Irrawaddy, Santa, Majes, Ocona) while the others project a decrease; across those basins, it is not consistent which glacier model projects an increase and which project a decrease. The glacier model that projects an increase in seasonal runoff is also not necessarily the glacier model that projects the highest absolute runoff. For example, GloGEM is the only glacier model that projects an increase in monthly runoff for the Susitna basin (Figure 5), but its end-of-century annual runoff is similar to OGGM. Likewise, PyGEM projects the greatest net increase in monthly runoff 255 across multiple months for the Jokulsa a Fjollum, but projects the lowest absolute runoff at the end of century for that basin (Figures A1-A2). Projections of seasonal change in such basins may benefit from a more detailed regional study using a multi-glacier-model ensemble and validating against historical observations where available.

## 4 Discussion

We have found that different metrics of future glacier runoff – absolute annual runoff, runoff change from a historical baseline, year of peak water, and change in seasonal runoff distribution – have different dominant sources of uncertainty in 21st century projections. While the multi-GCM and multi-glacier-model ranges have statistical limitations, comparing them to each other indicates whether glacier models produce broadly similar projections (Sect. 2.2). Absolute annual runoff projections are at least as affected by the choice of glacier model as by the choice of forcing GCM. For 29 of the 75 examined basins, the inter-glacier-model range in annual runoff is larger than the inter-GCM range (Figures 2 and A1). For the remaining 46 basins the inter-glacier-model range is relatively comparable to the inter-GCM range.

Glacier models have generally been calibrated ~~to match per glacier using single glacier~~ mass change observations (Hugonnet et al., 2021) from global datasets (Hugonnet et al., 2021), rather than local or ~~regional~~ regional-scale glacier runoff metrics, ~~since the runoff data required does not exist~~. Runoff has been a variable of secondary interest to model development focused on glacier contributions to sea level (e.g. Hock et al., 2019; Marzeion et al., 2020) ; further, local and regional-scale runoff data is too limited to permit its use in global model calibration (van Tiel et al., 2020b). The scarcity of historical observations of glacier change, and the different approaches to initialization in each glacier model, lead to wide spread in model initial conditions that persist in the future projections (Eis et al., 2019; Schuster et al., 2023a; ?)(Eis et al., 2019; Schuster et al., 2023a; Aguayo et al., 2023). Expressing projections as a percent difference from a historical baseline – where the baseline may be different for each model – corrects this spread in initial conditions and makes the future changes more consistent across glacier models (Figure 3). Studies using glacier runoff projections should account for the wide spread in glacier model initial conditions, for example by ~~analysing~~ analyzing percent change rather than absolute runoff, or by calibrating new regional model runs with additional observations (such as winter accumulation) that are not globally available.

Variability in glacier runoff projections between glacier models and GCMs is driven by differences in temperature (i.e., melt) and solid and liquid precipitation. While the future climate data was consistent across the glacier models, the models varied in the reference climate datasets used to bias correct the future climate data, as well as the calibration framework used to estimate the degree-day factors, temperature biases, and precipitation factors (Sect. 2.1). We thus evaluate the relation between projected glacier runoff and precipitation over the historical period, since the latter captures both the reference climate data and adjustments caused by the calibrated precipitation factors. We find that the extent to which one glacier model projects more or less runoff than the others for a given basin is directly related to its precipitation adjustment (Figure 6) and the area over which precipitation is summed for runoff. Further, the linear correlation explicitly shows that the inter-model offsets in the magnitudes of projected runoff (Figure 2) are driven by modeler decisions related to the reference climate data and calibration frameworks used. However, though the relation is clearly linear, the slope of the regression lines are all less than 1. This indicates that while increasing precipitation will increase runoff, runoff is also affected by the amount of melt occurring. The melt in turn depends on additional calibrated model parameters (degree day factors and temperature biases) and the implementation of the ~~melt parametrization (Schuster et al., 2023a)~~ surface-type dependent melt parametrization (Schuster et al. 2023a; and see Methods above) as well as the air temperature forcing.



**Figure 6.** Relative average annual runoff (2000–2099) for SSP 2-4.5 vs. relative average (annual) historical (2000–2019) precipitation. The above values are made "relative" by dividing the multi-GCM median annual runoff (or average historical precipitation) by the mean of all 3 glacier models and averaging across the period such that, if one model projects 20 % more runoff than the mean of all three models, its relative runoff value is 1.2 (see Sect. 2.2) The slopes of our calculated regression lines and the corresponding correlation coefficients are provided for each glacier model. [The RGI region, within which the given basin resides, is indicated by the shape of the marker.](#)

[For example, Schuster et al. \(2023a\) found slightly larger projected runoff when surface-type distinction was not used and smaller runoff when monthly temperature data, instead of superimposed daily data, were applied \(OGGM vs PyGEM GloGEM, which use both surface-type distinction and superimposed daily data\).](#) Differences in model adjusted precipitation are further amplified or reduced by secondary variables. **For example** [Notably](#), differences in a basin's initially-glaciated area (apparent from Supplementary Data A1) would result in differing input precipitation totals even prior to glacier-model calibration. Such differences may arise due to slight discrepancies in the number of glaciers simulated by each model in a given basin (Supplementary Data A1) or as a result of OGGM's dynamic spinup routine for initial area (as contrasted with GloGEM and PyGEM's use of the RGI outline, Sect. 2.1).

300 Projections of peak water year vary across single GCMs – in some cases spanning the entire 21st century – but are consistent across glacier models for each basin (Figure 4). Indeed, the century-wide range in peak water projections between GCMs suggests that single-GCM simulations cannot be trusted to accurately project peak water years (Figure 4). Further, the influence

of GCM on runoff varies by basin, i.e., there is no single GCM with consistently earlier or later peak water years than the others, which makes it difficult to interpret single-GCM results in isolation. Users interested in the timing of peak water could use any one of the three glacier models for projections, but should be sure to use the full ensemble of simulations forced by different GCMs.

The sensitivity of peak water year to future warming scenario is consistent across GCMs and glacier models, but not consistent across basins. Where peak water occurs early in the century, runoff declines after the peak in every scenario. In basins where peak water occurs later, the timing of peak water depends more clearly on SSP scenario (see Fig. 4 (b) and Fig. A4). Basins with later peak water tend to be more heavily glaciated (not shown) and/or in regions that receive more precipitation (e.g. Copper, Yukon, Indus). In those basins, increasing future warming may melt more ice later in the century as well as alter precipitation trends. Both phenomena would serve to push the peak water year later in the century with increasing radiative forcing, which produces the apparent sensitivity of peak water year to SSP scenario (Figure A4). Although our study did not analyze runoff components due to data limitations, the partitioning of runoff into glacier wastage or “excess melt”, equilibrium ice melt, seasonal snow melt, and precipitation could support additional insights into peak water year sensitivity to SSP scenario. For example, Rounce et al. (2020a) found that basins with large excess meltwater signals (e.g. Amu Darya, Indus, and Tarim) had a later-century peak water whose timing was sensitive to SSP scenario. Moore et al. (2020) found that for several smaller Canadian basins, excess melt had already passed its peak in 2019 and that observed streamflow was already below what would be expected if glaciers were in equilibrium. The latter study emphasises the need for glacio-hydrological modelling studies with evolving glacier cover.

Runoff seasonality changes in a globally consistent way across all three glacier models, with seasonal runoff peaks shifting earlier and decreasing in magnitude in most basins (Figures 5 and A2-A3). Seasonality of simulated runoff is controlled by the temperature downscaled to the glacier sites as well as by the partitioning between solid and liquid precipitation. Each glacier model uses different methods for those tasks, which we expect to produce some slight differences among their projections. In the basins with apparent differences in multi-GCM median seasonal runoff, we note that GloGEM tends to maintain wider seasonal peaks toward the end of the century (e.g. Columbia through Yukon panels of Figure A2). OGGM tends to maintain the strongest seasonal peaks – i.e. monthly runoff in the peak month that is closest to the historical maximum monthly runoff (e.g. Colville, Indigirka, Balkhash, Mekong, Amazon, Ocona, Titicaca basins in Figure A2). ~~These slight differences in seasonality may reflect differences among the models in the underlying components of runoff (ice melt, snow melt, liquid precipitation) all of which have distinct seasonal curves; a detailed analysis of hydrological components could guide basin-level seasonal runoff projections.~~ PyGEM tends to project the strongest percent change in runoff distribution (Figure A3), which is consistent with PyGEM’s historical baseline values being the lowest in many basins (Figure A1): the same absolute change in monthly runoff will be a larger percentage of a small baseline value. ~~Despite~~ These slight differences in seasonality may reflect differences among the models in the underlying components of runoff (ice melt, snow melt, liquid precipitation) all of which have distinct seasonal curves; a detailed analysis of hydrological components could guide basin-level seasonal runoff projections. However, despite these small differences, we emphasise the broad similarity across glacier models in projected seasonal runoff decreasing and shifting earlier in the year.

In both arid central Asia and maritime North America, there are several basins where all three models project increases in the magnitude of the seasonal runoff peak at the end of the century (Sect. 3.3 and Figure A2). Both regions are projected to see increased winter (December–February) precipitation (Lee et al., 2021); if the magnitude of the seasonal runoff peak depends most strongly on winter snow accumulation, increased snowfall could drive a larger seasonal runoff peak in both regions. Another possible driver would be stronger seasonal melting in both regions due to summer warming in glaciated areas that were previously close to the freezing point; the seasonal warming signals in the CMIP6 projections are not robust in these regions (Lee et al., 2021), so further regional analysis would be needed to confirm that possibility. An analysis of projected solid versus liquid precipitation in each basin — not pursued here due to data limitations — would also clarify the regional dynamics contributing to projected increases in seasonal runoff.

Although the runoff projections from different glacier models have large offsets in absolute annual runoff, the same projections expressed in relative terms – percent change from a historical baseline – ~~are much more similar~~ exhibit strongly reduced offsets (Figures 2-3). The temporal pattern of runoff increase, peak water, and subsequent runoff decrease in each basin is similar for all three glacier models; as a result, the year of peak water and its response to increasing climate warming has little glacier model uncertainty (Figures 4 and A1). Future changes in runoff seasonality also show similarities in basins with similar peak water years (Sect. 3.3). Thus, although “peak water” is limited to a very general description of glacier runoff, which does not account for broader basin hydrology or within-basin differences among glaciers, we believe that the year of peak water, with its consistency across glacier models, can serve as a helpful heuristic for other runoff metrics.

We emphasize that the results we present are a sample of 12 GCM realizations per scenario, selected from the more than 250 plausible realizations from the CMIP6 archive. Our results thus likely under-sample the true uncertainty arising from CMIP6 projections. ~~In addition, all~~ All GCMs struggle to represent precipitation processes, particularly in areas of high relief typical of glaciated basins ~~–~~ (Douville et al., 2021). In addition, the spread in surface air temperature among GCM realizations, which we have not quantified here, will produce differences in melt and in how much precipitation falls as solid or liquid phase, and that will affect both magnitude and timing of glacier runoff. As such, there is a high likelihood of systematic biases in the GCM forcing that cannot be quantified and that are not easily addressed with bias correction (e.g., if bias impacts longer timescale climate variability and change). Those interested in water availability for a specific basin should conduct a regional analysis to select an ensemble of GCMs that best represents precipitation and temperature trends and variability for that basin.

Similarly, the glacier runoff projections we ~~analyse~~ analyze come from the three globally-capable glacier evolution models. No other global “glacier-centric” runoff projections are publicly available, and we do not quantify glacier model parameter uncertainty in any of our results. As such, our results do not map a full probability distribution of future glacier runoff (?). (Aguayo et al., 2023). Glacier models’ differing approaches to handling precipitation input have a strong effect on their runoff output (Figure 6). In a given basin, a glacier model forced with more precipitation during the historical period (either due to the choice of reanalysis data or model calibration) usually also projects more runoff during the 21st century. A multiplicative “precipitation factor” that scales up GCM precipitation will have a corresponding effect on the precipitation component of glacier runoff (Sect. 2.1). The three glacier models also used different climate reanalysis datasets for their bias correction and calibration period (Sect. 2.1.1-2.1.3), but all were calibrated to match the same global glacier mass change dataset (Hugonnet

et al., 2021). A glacier model that is calibrated with higher precipitation may therefore also produce higher melt to match the observed mass change (Schuster et al., 2023a), which means that the influence of the precipitation factor is not limited to the first order precipitation response, but on the partitioning and seasonality of ice melt as well.


Ultimately, the importance of glacier runoff in a given basin depends on the local hydrology. Glacier runoff may infiltrate to groundwater reservoirs (Somers et al., 2016; Mackay et al., 2020), sustain high-altitude wetlands (López-Moreno et al., 2022), flow through proglacial lakes or streams, evaporate while transiting arid downstream regions (Wang et al., 2013), and encounter human management interventions [such as diverting runoff towards agricultural lands, hydroelectric plants, or storage reservoirs.](#) Simulating those dynamics will require coupled glacio-hydrological models (Hanus et al., 2024). Ideally, projections of regional water availability will calibrate and apply a coupled glacio-hydrological model; however, such models are strongly under-constrained, perhaps even more so than the glacier models we have presented (Somers and McKenzie, 2020; Drenkhan et al., 2023). Without sufficient calibration data, one of the coupled model components may inadvertently be calibrated to over-compensate for the shortcomings of the other. Although we have presented an analysis of glacier runoff in isolation, our work highlights the urgent need for additional observations of mountain hydrology (Somers and McKenzie, 2020) and meteorology (Shahgedanova et al., 2021) to support glacio-hydrological simulations.

## 5 Conclusions


Aggregation of annual runoff series among the three glacier models reveals regionally-dependent offsets. The offsets arise from differences in historical baseline conditions for the three glacier models; their effect can be removed by normalizing relative to those conditions to produce percent change in runoff. Remaining differences are the result of different approaches to correcting GCM precipitation across the glacier models, which can only be refined with more complete observations of mountain hydrology. Despite differences in annual runoff, changes in runoff seasonality are qualitatively consistent across glacier models, with most basins' seasonal runoff peaks moving earlier and having lower magnitude than their historical runoff peaks. [These robust seasonal shifts could be expected to impact drought risk in mountain areas, though changes in glacial drought buffering were not apparent in an analysis of the Standardized Precipitation-Evapotranspiration Index \(Ultee et al., 2022\).](#) The inter-GCM range in basin peak water year, which is otherwise consistent across glacier models, highlights the need for judicious selection of climate forcings to simulate hydrologic change in glaciated river basins.

*Code and data availability.* The OGGM standard projections are a variant of Schuster et al. (2023b). The PyGEM standard projections are archived as Rounce et al. (2022). Aggregated runoff for each basin for all three glacier models is available in CSV format at DataDryad (Wimberly, 2024c). Analysis code is archived on Zenodo (Wimberly, 2024a; Ultee, 2024). Code for the interactive map is archived on Zenodo (Wimberly, 2024b). Data processing relied on the xarray Python package (Hoyer and Hamman, 2017).

## Appendix A: Global results - Total Annual Runoff Projections



**Figure A1.** Total annual runoff projections for all 75 glaciated major river basins given SSP 2-4.5. The multi-GCM range is highlighted in the same way as Fig. 2. In this plot we have also indicated the calculated year of peak water with a vertical dashed line. The color of each line corresponds to the glacier evolution model color indicated by the legend.




**Figure A2.** Seasonal cycle of runoff in 2080–2099, expressed as a fraction of maximum monthly runoff during the 2000–2019 baseline period, as in the middle row of Figure 5, here for all basins. Heavier lines show multi-GCM median for each glacier model; light lines show individual GCMs. Lines that cross the dashed line at 1.0 show a seasonal peak with a higher magnitude than the historical.


Table A1: Initial (Year 2000) Areas and Number of Glaciers for each Glacier Model

Basin	GloGEM		OGGM		PyGEM	
	Initial Area [km <sup>2</sup> ]	Number of Glaciers	Initial Area [km <sup>2</sup> ]	Number of Glaciers	Initial Area [km <sup>2</sup> ]	Number of Glaciers
Alsek	5498.55	1390	5614.80	1391	5499.49	1391
Amazon	1402.74	1552	1490.85	1557	1400.20	1555
Aral Sea	12522.45	16103	13309.70	16211	12557.75	16210
Azopardo	30.16	87	31.18	88	30.21	88
Baker	2200.19	1955	2229.16	1957	2206.68	1957
Biobio	74.64	137	78.99	138	74.66	138
Brahmaputra	10529.95	11505	10751.82	11508	10528.21	11507
Chico	33.80	109	36.20	109	33.84	109
Chuy	380.86	719	388.55	719	381.09	719
Cisnes	71.79	180	72.39	180	71.70	180
Clutha	156.82	660	164.13	660	132.25	557
Colorado	1300.15	1743	1413.58	1746	1300.96	1746
Columbia	1952.34	3433	2200.04	3437	1922.12	3436
Colville	33.42	103	38.74	103	33.46	103
Copiapo	35.18	71	39.24	71	35.20	71
Copper	16166.33	3305	16274.66	3305	16174.21	3305
Danube	405.32	822	419.54	843	411.26	841
Dramselva	31.55	50	35.49	50	31.62	50
Fraser	2538.10	2632	2686.05	2632	2536.99	2631
Ganges	7902.78	6552	7952.92	6552	7906.05	6552
Glomaa	273.36	284	289.56	284	273.40	284
Har Us Nuur	319.54	566	340.08	567	319.67	566
Huasco	33.80	89	38.52	89	33.83	89
Indigirka	171.41	329	189.96	331	171.94	331
Indus	27211.12	23419	28143.11	23424	27201.24	23422
Irrawaddy	47.59	144	49.76	144	47.46	143
Jokulsa A Fjollum	1097.90	10	1098.33	10	1098.65	10
Kalixalven	38.49	59	39.35	59	38.49	59
Kamchatka	196.40	135	208.12	135	196.53	135
Kuban	173.34	285	165.83	297	162.29	258
Kuskokwim	1055.76	874	1099.66	874	1056.66	874
Lagarfljot	154.29	10	154.41	10	154.39	10
Lake Balkhash	2271.51	3067	2370.86	3142	2328.32	3142
Lulealven	256.33	244	267.30	244	256.48	244
Mackenzie	1428.24	2130	1481.82	2130	1457.07	2129
Magdalena	46.73	30	49.34	30	46.76	30
Majes	30.66	64	32.87	64	30.63	64
Mekong	171.30	405	179.27	405	171.33	405
Nass	1230.51	819	1279.38	819	1231.58	819
Negro	47.69	194	53.16	194	47.72	194
Nelson	370.85	586	387.60	586	373.42	586
Nushagak	89.33	116	96.92	116	89.37	116
Ob	762.81	1608	832.35	1609	763.40	1608
Ocona	34.86	34	37.03	34	34.87	34
Olfusa	683.60	41	686.59	41	684.12	40
Palena	349.26	651	354.57	651	349.10	650

Basin	GloGEM		OGGM		PyGEM	
	Initial Area [km <sup>2</sup> ]	Number of Glaciers	Initial Area [km <sup>2</sup> ]	Number of Glaciers	Initial Area [km <sup>2</sup> ]	Number of Glaciers
Pascua	2437.00	1098	2452.38	1098	2437.52	1098
Po	313.30	765	323.83	765	313.41	765
Puelo	217.21	416	220.76	416	217.08	416
Rapel	234.13	245	235.69	246	234.27	246
Rhine	336.58	796	351.85	796	336.92	796
Rhone	915.64	1168	931.44	1169	917.29	1169
Salween	1265.22	2134	1309.88	2134	1265.38	2134
Santa	354.78	409	353.41	409	354.43	409
Santa Cruz	3010.24	461	3024.34	462	3011.12	461
Serrano	940.73	166	950.48	167	941.32	167
Skagit	200.46	801	244.58	802	181.03	802
Skeena	764.20	1168	816.39	1168	764.77	1168
Stikine	3578.02	2264	3618.35	2275	3581.22	2275
Susitna	4292.82	1256	4363.96	1256	4295.75	1256
Svarta	277.38	19	278.86	19	277.50	19
Taku	1337.94	639	1367.25	639	1338.67	639
Talas	68.85	174	72.29	174	68.85	174
Tarim He	26116.42	20736	26528.58	20791	26116.95	20783
Thjorsa	972.28	22	973.43	22	972.32	21
Titicaca	279.87	308	294.22	310	280.02	310
Tornealven	34.25	64	37.31	64	34.26	64
Uvs Nuur	48.24	54	51.52	54	48.26	54
Valdivia	45.80	33	45.87	33	45.76	33
Yangtze	1647.63	1538	1741.54	1633	1710.23	1633
Yelcho	238.94	616	247.73	618	238.96	618
Yellow River	177.42	287	183.77	287	177.38	287
Ysyk-Kol	507.88	807	514.07	807	507.86	807
Yukon	9996.19	2948	10109.20	2994	10027.76	2994



**Figure A3.** Seasonal shift in runoff from early century (2000–2019) to late (2080–2099), as in the bottom row of Figure 5, here for all basins. Shifts are expressed in differences per month as a fraction of historical (2000–2019) maximum monthly runoff for each basin. Heavier lines show multi-GCM median for each glacier model; light lines show individual GCMs. Positive y-axis values indicate a net increase in that month’s runoff at the end of the century compared with the baseline period (2000–2019); negative y-axis values indicate a decrease.



**Figure A4.** Range of predicted year of peak water across emission scenarios (SSPs) compared to the multi-glacier-model median year of peak water for SSP 2-4.5. Points of the same color represent different basins within one RGI region.

*Author contributions.* LU designed the study, with input from SC, JM, MH, DR, and FM. MH, DR, and LS conducted global glacier model simulations and shared output with LU. FW aggregated all model output data, produced all visualizations, interpreted the results, and drafted the manuscript. EH conducted preliminary analysis on CMIP5 output, and supported data processing and visualization. All authors  
405 \*\*\*contributed to revising the manuscript and approved its final form\*\*\*.

*Competing interests.* The authors declare that no competing interests are present.

*Acknowledgements.* FW was supported by Middlebury College Undergraduate Research funding. LS is recipient of a DOC Fellowship of the Austrian Academy of Sciences at the Department of Atmospheric and Cryospheric Sciences, Universität Innsbruck (No. 25928). LS  
410 and FM have received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 101003687. This text reflects only the author's view and that the Agency is not responsible for any use that may be made of the information it contains. EH was supported by Chalmers University of Technology graduate funding. MH was supported by the European Union's Horizon 2020 research and innovation programme (PROTECT; grant agreement number 869304). DR was supported by National Aeronautics and Space Administration grants 80NSSC20K1296 and 80NSSC20K1595. JM's contribution was funded by the Natural Environment Research  
415 Council (NERC) MCNC Grant TerraFirma NE/W004895/1.

The authors thank Brandon Tober for producing global ERA5 runs of PyGEM that supported the analysis of precipitation factor among the glacier models.

## References

- Aguayo, R., Maussion, F., Schuster, L., Schaefer, M., Caro, A., Schmitt, P., Mackay, J., Ultee, L., Leon-Muñoz, J., and Aguayo, M.: Assessing the glacier projection uncertainties in the Patagonian Andes (40–56° S) from a catchment perspective, *EGUosphere*, 2023, 1–41, <https://doi.org/10.5194/egusphere-2023-2325>, 2023.
- Ahmed, R., Wani, G. F., Ahmad, S. T., Sahana, M., Singh, H., and Ahmed, P.: A review of glacial lake expansion and associated glacial lake outburst floods in the Himalayan region, *Earth Systems and Environment*, 5, 695–708, <https://doi.org/https://doi.org/10.1007/s41748-021-00230-9>, 2021.
- Bosson, J.-B., Huss, M., and Osipova, E.: Disappearing world heritage glaciers as a keystone of nature conservation in a changing climate, *Earth's Future*, 7, 469–479, 2019.
- Bosson, J.-B., Huss, M., Cauvy-Fraunié, S., Clément, J.-C., Costes, G., Fischer, M., Poulénard, J., and Arthaud, F.: Future emergence of new ecosystems caused by glacial retreat, *Nature*, 620, 562–569, 2023.
- Christian, J. E., Whorton, E., Carnahan, E., Koutnik, M., and Roe, G.: Differences in the transient responses of individual glaciers: a case study of the Cascade Mountains of Washington State, USA, *Journal of Glaciology*, 68, 751–763, <https://doi.org/DOI: 10.1017/jog.2021.133>, 2022.
- Compagno, L., Huss, M., Miles, E. S., McCarthy, M. J., Zekollari, H., Dehecq, A., Pellicciotti, F., and Farinotti, D.: Modelling supraglacial debris-cover evolution from the single-glacier to the regional scale: an application to High Mountain Asia, *The Cryosphere*, 16, 1697–1718, <https://doi.org/10.5194/tc-16-1697-2022>, 2022.
- Douville, H., Raghavan, K., Renwick, J., Allan, R., Arias, P., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T., Gergis, J., Jiang, D., Khan, A., Pokam Mba, W., Rosenfeld, D., Tierney, J., and Zolina, O.: Water Cycle Changes, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 1055–1210, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.
- Drenkhan, F., Buytaert, W., Mackay, J. D., Barrand, N. E., Hannah, D. M., and Huggel, C.: Looking beyond glaciers to understand mountain water security, *Nature Sustainability*, 6, 130–138, <https://doi.org/10.1038/s41893-022-00996-4>, 2023.
- Eis, J., Maussion, F., and Marzeion, B.: Initialization of a global glacier model based on present-day glacier geometry and past climate information: an ensemble approach, *The Cryosphere*, 13, 3317–3335, <https://doi.org/10.5194/tc-13-3317-2019>, 2019.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., van den Broeke, M. R., and Paul, F.: A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*, 340, 852–857, <https://doi.org/10.1126/science.1234532>, 2013.
- Global Runoff Data Centre: GRDC Major River Basins, <http://grdc.bafg.de>, [https://grdc.bafg.de/GRDC/EN/02\\_srvcs/22\\_gslrs/221\\_MRB/riverbasins\\_node.html](https://grdc.bafg.de/GRDC/EN/02_srvcs/22_gslrs/221_MRB/riverbasins_node.html), 2020.
- Hanus, S., Schuster, L., Burek, P., Maussion, F., Wada, Y., and Viviroli, D.: Coupling a large-scale glacier and hydrological model (OGGM v1.5.3 and CWatM V1.08) – Towards an improved representation of mountain water resources in global assessments, *EGUosphere*, 2024, 1–31, <https://doi.org/10.5194/egusphere-2023-2562>, 2024.

- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.
- Hock, R., Bliss, A., Marzeion, B., Giesen, R. H., Hirabayashi, Y., Huss, M., Radić, V., and Slangen, A. B.: GlacierMIP –A model intercomparison of global-scale glacier mass-balance models and projections, *Journal of Glaciology*, 65, 453–467, <https://doi.org/DOI:10.1017/jog.2019.22>, 2019.
- 465 Hoyer, S. and Hamman, J.: xarray: N-D labeled Arrays and Datasets in Python, *Journal of Open Research Software*, 5, <https://doi.org/https://doi.org/10.5334/jors.148>, 2017.
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- 470 Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Frontiers in Earth Science*, 3, <https://doi.org/10.3389/feart.2015.00054>, 2015.
- Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 135–140, <https://doi.org/10.1038/s41558-017-0049-x>, 2018.
- Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., and Baillie, J. E. M.: Importance and vulnerability of the world’s water towers, *Nature*, 577, 364–369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- Jacobsen, D., Milner, A. M., Brown, L. E., and Dangles, O.: Biodiversity under threat in glacier-fed river systems, *Nature Climate Change*, 2, 361–364, <https://doi.org/10.1038/nclimate1435>, 2012.
- 480 Kaser, G., Großhauser, M., and Marzeion, B.: Contribution potential of glaciers to water availability in different climate regimes, *Proceedings of the National Academy of Sciences*, 107, 20 223–20 227, <https://doi.org/10.1073/pnas.1008162107>, 2010.
- Kinouchi, T., Nakajima, T., Mendoza, J., Fuchs, P., and Asaoka, Y.: Water security in high mountain cities of the Andes under a growing population and climate change: A case study of La Paz and El Alto, Bolivia, *Water Security*, 6, 100, 2019.
- 485 Lange, S., Menz, C., Gleixner, S., Cucchi, M., Weedon, G. P., Amici, A., Bellouin, N., Müller Schmied, H., Hersbach, H., Buontempo, C., and Cagnazzo, C.: WFDE5 over land merged with ERA5 over the ocean (W5E5 v2.0), ISIMIP Repository, <https://doi.org/https://doi.org/10.48364/ISIMIP.342217>, 2021.
- Lee, J.-Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J. P., Engelbrecht, F., Fischer, E., Fyfe, J. C., Jones, C., Maycock, A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Future global climate: Scenario-based projections and near-term information, in: *Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T.,

- Yelekçi, O., Yu, R., and Zhou, B., pp. 553–672, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.006>, 2021.
- 495 López-Moreno, J. I., Rojas-Heredia, F., Ceballos, J. L., Morán-Tejeda, E., Alonso-Gonzalez, E., Vidaller, I., Deschamps-Berger, C., and Revuelto, J.: Recent evolution of glaciers in the Cocuy-Güicán Mountains (Colombian Andes) and the hydrological implications, *Land Degradation & Development*, 33, 2606–2618, <https://doi.org/https://doi.org/10.1002/ldr.4336>, 2022.
- Mackay, J. D., Barrand, N. E., Hannah, D. M., Krause, S., Jackson, C. R., Everest, J., MacDonald, A. M., and ÓDochartaigh, B. É.: Proglacial groundwater storage dynamics under climate change and glacier retreat, *Hydrological Processes*, 34, 5456–5473, <https://doi.org/https://doi.org/10.1002/hyp.13961>, 2020.
- 500 Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, *The Cryosphere*, 6, 1295–1322, <https://doi.org/10.5194/tc-6-1295-2012>, 2012.
- Marzeion, B., Hock, R., Anderson, B., Bliss, A., Champollion, N., Fujita, K., Huss, M., Immerzeel, W., Kraaijenbrink, P., Malles, J.-H., Maussion, F., Radić, V., Rounce, D. R., Sakai, A., Shannon, S., van de Wal, R., and Zekollari, H.: Partitioning the Uncertainty of Ensemble  
505 Projections of Global Glacier Mass Change, *Earth’s Future*, p. e2019EF001470, <https://doi.org/10.1029/2019EF001470>, 2020.
- Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A. H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C. T., and Marzeion, B.: The Open Global Glacier Model (OGGM) v1.1, *Geoscientific Model Development*, 12, 909–931, <https://doi.org/10.5194/gmd-12-909-2019>, 2019.
- Maussion, F., Rothenpieler, T., Dusch, M., Schmitt, P., Vlug, A., Schuster, L., Champollion, N., Li, F., Marzeion, B., Oberrauch, M., Eis,  
510 J., Fischer, A., Landmann, J., Jarosch, A., luzpaz, Hanus, S., Rounce, D., Castellani, M., Bartholomew, S. L., Minallah, S., bowen-belongstonature, Merrill, C., Otto, D., Loibl, D., Rosa, G., Ultee, L., Thompson, S., anton ub, and Gregor, P.: OGGM/oggm: v1.6.1, <https://doi.org/10.5281/zenodo.8287580>, 2023.
- McCarthy, M., Meier, F., Fatichi, S., Stocker, B. D., Shaw, T. E., Miles, E., Dussaillant, I., and Pellicciotti, F.: Glacier contributions to river discharge during the current Chilean megadrought, *Earth’s Future*, 10, 2022.
- 515 Moore, R., Pelto, B., Menounos, B., and Hutchinson, D.: Detecting the effects of sustained glacier wastage on streamflow in variably glacierized catchments, *Frontiers in Earth Science*, 8, 136, 2020.
- Oerlemans, J., ed.: *On the Response of Valley Glaciers to Climatic Change*, Springer Netherlands, Dordrecht, 1989.
- Pelto, M. S., Dryak, M., Pelto, J., Matthews, T., and Perry, L. B.: Contribution of Glacier Runoff during Heat Waves in the Nooksack River Basin USA, *Water*, 14, <https://doi.org/10.3390/w14071145>, 2022.
- 520 RGI Consortium: *Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0*, Technical Report, Global Land Ice Measurements from Space, 2017.
- Riedel, J. L. and Larrabee, M. A.: Impact of Recent Glacial Recession on Summer Streamflow in the Skagit River, *Northwest Science*, 90, 5–22, <https://doi.org/10.3955/046.090.0103>, 2016.
- Rounce, D. R., Hock, R., and Shean, D. E.: Glacier Mass Change in High Mountain Asia Through 2100 Using the Open-Source Python  
525 Glacier Evolution Model (PyGEM), *Frontiers in Earth Science*, 7, 331, <https://doi.org/10.3389/feart.2019.00331>, 2020a.
- Rounce, D. R., Khurana, T., Short, M. B., Hock, R., Shean, D. E., and Brinkerhoff, D. J.: Quantifying parameter uncertainty in a large-scale glacier evolution model using Bayesian inference: application to High Mountain Asia, *Journal of Glaciology*, 66, 175–187, <https://doi.org/10.1017/jog.2019.91>, 2020b.

- Rounce, D. R., Hock, R., McNabb, R. W., Millan, R., Sommer, C., Braun, M. H., Malz, P., Maussion, F., Mouginot, J., Seehaus, T. C., and Shean, D. E.: Distributed Global Debris Thickness Estimates Reveal Debris Significantly Impacts Glacier Mass Balance, *Geophysical Research Letters*, 48, e2020GL091311, <https://doi.org/https://doi.org/10.1029/2020GL091311>, 2021.
- Rounce, D. R., Hock, R., and Maussion, F.: Global PyGEM-OGGM Glacier Projections with RCP and SSP Scenarios, Version 1, <https://doi.org/10.5067/P8BN9VO9N5C7>, 2022.
- Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78–83, <https://doi.org/10.1126/science.abo1324>, 2023.
- Schuster, L., Rounce, D. R., and Maussion, F.: Glacier projections sensitivity to temperature-index model choices and calibration strategies, *Annals of Glaciology*, pp. 1–16, <https://doi.org/DOI:10.1017/aog.2023.57>, 2023a.
- Schuster, L., Schmitt, P., Vlug, A., and Maussion, F.: OGGM/oggm-standard-projections-csv-files: v1.0 (v1.0), Zenodo, <https://doi.org/https://doi.org/10.5281/zenodo.8286065>, 2023b.
- Shahgedanova, M., Adler, C., Gebrekirstos, A., Grau, H. R., Huggel, C., Marchant, R., Pepin, N., Vanacker, V., Viviroli, D., and Vuille, M.: Mountain Observatories: Status and Prospects for Enhancing and Connecting a Global Community, *Mountain Research and Development*, 41, A1, <https://doi.org/10.1659/MRD-JOURNAL-D-20-00054.1>, 2021.
- Somers, L. D. and McKenzie, J. M.: A review of groundwater in high mountain environments, *WIREs Water*, 7, e1475, <https://doi.org/https://doi.org/10.1002/wat2.1475>, 2020.
- Somers, L. D., Gordon, R. P., McKenzie, J. M., Lautz, L. K., Wigmore, O., Glose, A., Glas, R., Aubry-Wake, C., Mark, B., Baraer, M., and Condom, T.: Quantifying groundwater–surface water interactions in a proglacial valley, Cordillera Blanca, Peru, *Hydrological Processes*, 30, 2915–2929, <https://doi.org/10.1002/hyp.10912>, 2016.
- Taylor, C., Robinson, T. R., Dunning, S., Rachel Carr, J., and Westoby, M.: Glacial lake outburst floods threaten millions globally, *Nature Communications*, 14, 487, 2023.
- Taylor, K. E., Durack, P. J., Elkington, M., Guilyardi, E., Hassell, D., Lautenschlager, M., and Stockhause, M.: CMIP6 Participation Guidance for Modelers, <https://pcmdi.llnl.gov/CMIP6/Guide/modelers.html#7-archivingpublishing-output>, 2022.
- Ultee, L.: ehultee/glacio-rf-comp: Discussion release, <https://doi.org/10.5281/zenodo.11406333>, 2024.
- Ultee, L., Coats, S., and Mackay, J.: Glacial runoff buffers droughts through the 21st century, *Earth System Dynamics*, 13, 935–959, <https://doi.org/10.5194/esd-13-935-2022>, 2022.
- van Tiel, M., Kohn, I., Van Loon, A. F., and Stahl, K.: The compensating effect of glaciers: Characterizing the relation between interannual streamflow variability and glacier cover, *Hydrological Processes*, 34, 553–568, <https://doi.org/10.1002/hyp.13603>, 2020a.
- van Tiel, M., Stahl, K., Freudiger, D., and Seibert, J.: Glacio-hydrological model calibration and evaluation, *Wiley Interdisciplinary Reviews: Water*, 7, e1483, 2020b.
- Wang, A., Miao, Y., Kong, X., and Wu, H.: Future changes in global runoff and runoff coefficient from CMIP6 multi-model simulation under SSP1-2.6 and SSP5-8.5 scenarios, *Earth's Future*, 10, e2022EF002910, <https://doi.org/10.1029/2022EF002910>, 2022.
- Wang, H., Chen, Y., Li, W., and Deng, H.: Runoff responses to climate change in arid region of northwestern China during 1960–2010, *Chinese Geographical Science*, 23, 286–300, <https://doi.org/10.1007/s11769-013-0605-x>, 2013.
- WGMS: Fluctuations of Glaciers Database. World Glacier Monitoring Service, Zurich, Switzerland, <https://doi.org/10.5904/wgms-fog-2020-08>, 2020.

- WGMS: Fluctuations of Glaciers Database, World Glacier Monitoring Service (WGMS), Zurich, Switzerland, <https://doi.org/10.5904/wgms-fog-2024-01>, 2024.
- Wiersma, P., Aerts, J., Zekollari, H., Hrachowitz, M., Drost, N., Huss, M., Sutanudjaja, E. H., and Hut, R.: Coupling a global glacier model to a global hydrological model prevents underestimation of glacier runoff, *Hydrology and Earth System Sciences*, 26, 5971–5986, <https://doi.org/10.5194/hess-26-5971-2022>, 2022.
- 570 Wimberly, F.: `finnwimberly/glacial_runoff_intercomparison`: Latest Release, <https://doi.org/10.5281/zenodo.11398017>, 2024a.
- Wimberly, F.: `finnwimberly/glacial_runoff_mapping`: Initial Release, <https://doi.org/10.5281/zenodo.11397990>, 2024b.
- Wimberly, F.: Processed Runoff Files, DataDryad, <https://doi.org/doi:10.5061/dryad.pk0p2ngxf>, 2024c.
- Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, *The Cryosphere*, 13, 1125–1146, <https://doi.org/10.5194/tc-13-1125-2019>, 2019.
- 575 Zekollari, H., Huss, M., Schuster, L., Maussion, F., Rounce, D. R., Aguayo, R., Champollion, N., Compagno, L., Hugonnet, R., Marzeion, B., Mojtabavi, S., and Farinotti, D.: 21<sup>st</sup> century global glacier evolution under CMIP6 scenarios and the role of glacier-specific observations, *EGU sphere*, 2024, 1–33, <https://doi.org/10.5194/egusphere-2024-1013>, 2024.
- Zemp, M., Gärtner-Roer, I., Nussbaumer, S. U., Welty, E. Z., Dussaillant, I., and Bannwart, J., eds.: Global glacier change bulletin No. 5 (2020–2021), ISC (WDS) / IUGG (IACS) / UNEP / UNESCO / WMO, World Glacier Monitoring Service, Zurich, Switzerland, based on database version <https://doi.org/10.5904/wgms-fog-2023-09>, 2023.
- 580