

Surface Observation Constrained High Frequency Coal Mine Methane Emissions in Shanxi China Reveal More Emissions than Inventories, Consistency with Satellite Inversion

Fan Lu¹, Qin Kai^{1*}, Jason Blake Cohen^{1*}, Qin He¹, Pravash Tiwari¹, Wei Hu¹, Chang Ye¹, Yanan Shan¹, Qing Xu¹, Shuo Wang¹, Qiansi Tu²

¹Jiangsu Key Laboratory of Coal-Based Greenhouse Gas Control and Utilization, School of Environment and Spatial Informatics, China University of Mining and Technology, Xuzhou, China

²School of Mechanical Engineering, Tongji University, Shanghai, China

*Correspondence to Kai Qin (qinkai@cumt.edu.cn) and Jason Blake Cohen (jasonbc@alum.mit.edu)

Abstract This work focuses on Changzhi, Shanxi China, a city and surrounding rural region with one of the highest atmospheric concentrations of methane (CH₄) worldwide (campaign-wide minimum/mean/standard deviation/max observations: 2.0, 2.9, 1.3, and 16 ppm) due to a rapid increase in the mining, production, and use of coal over the past decade. An intensive 15-day surface observation campaign of CH₄ concentration is used to drive a new analytical, mass-conserving method to compute and attribute CH₄ emissions. Observations made in concentric circles at 1km, 3km, and 5km around a high production high gas coal mine yielded emissions of 0.73, 0.28, and 0.15 ppm min⁻¹ respectively. Attribution used a 2-box mass conserving model to identify the known mine's emissions from 0.042-5.3 ppm min⁻¹, and a previously unidentified mine's emission from 0.22-7.9 ppm min⁻¹. These results demonstrate the importance of simultaneously quantifying both the spatial and temporal distribution of CH₄ emissions to better control regional-scale CH₄ emissions. Results of the attribution are used in tandem with observations of boundary layer height to quantify policy-relevant emissions from the two coal mines as 6860±3520 kg h⁻¹ and 1010±347 kg h⁻¹ respectively. Both mines display a fat tail distribution, with respective 25th, median, and 75th percentile values of [1600, 3070, 10500] kg h⁻¹ and [755, 1090, 1420] kg h⁻¹. These findings are demonstrated to be higher than CH₄ emissions from equivalent oil and gas

Formatted: Right: 0.63 cm

operations in the USA, with one about double and the other similar to day-to-day emissions inverted over 5-years using TROPOMI over the same region.

Key Words: *CH₄; Top-down emissions; Mass-conserving Model; Attribution; Mining*

1 Introduction

Emissions of methane (CH₄) contribute the second most to direct anthropogenic longwave radiative forcing (Etminan et al., 2016; Li et al., 2022). Since CH₄ has a lifetime from 9.5 to 12.5 years (Li et al., 2022; Prather et al., 2012), controlling CH₄ emissions can provide an opportunity to mitigate peak loading and slow the rate of net global warming. [Emissions from fossil fuel are one of the largest sources of anthropogenic CH₄](#) (Kirschke et al., 2013; Saunio et al., 2020a). Since China is the world's largest producer and consumer of coal (Bournazian, 2016), [Coal mines contribute up to 33%-40% of China's CH₄ emissions](#) (Janssens-Maenhout et al., 2017; Miller et al., 2019; Peng et al., 2016). Although China enacted [coal mines CH₄ \(CMM\)](#) regulations in 2010, CMM continues grow (Kerr and Yang, 2009; Miller et al., 2019). CH₄ emission estimates [remain exist](#) uncertain in both space and time (Brandt et al., 2014; Saunio et al., 2020b). They also generally have a fat tail distribution, wherein a small number of samples have extremely large emissions that overwhelm emissions under average conditions (Duren et al., 2019; Plant et al., 2022). For these reasons, new approaches to quantify, reduce uncertainty, and attribute CH₄ emissions are necessary and can provide support for policies aiming to control and mitigate CMM (Cao, 2017).

Bottom-up (BU) quantification of emissions requires a priori knowledge of source locations and diversity, which tends to not represent real-world conditions. Top-down (TD) approaches analyze concentration data with improving accuracy (Allen, 2014; Rigby et al., 2019; Varon et al., 2018; Vaughn et al., 2018), specifically combining surface (Heerah et al., 2021; Katzenstein et al., 2003; Shi et al., 2023), aircraft (Karion et al., 2013; Shi et al., 2022; Tong et al., 2023; Vinković et al., 2022), and/or satellite (Wecht et al., 2014) CH₄ observations with atmospheric models. Some TD approaches use physically realistic but complex chemical transport models (Bloom et al., 2017), others use plume models (Goldsmith et al., 2012), and others still use data driven

approaches (Buchwitz et al., 2017). [Uncertainties in whether or not to consider the basic formulation of plume models as being reasonable under the observed conditions are rarely addressed holistically or in detail, including but not limited to: stable wind speed and direction \(Varon et al., 2018\), no outside sources intersecting the plume \(Varon et al., 2018\), no significant enhancement in the background concentrations \(Irakulis-Loitxate et al., 2021\), no pooling or other non-linear behavior within the plume \(Bruno et al., 2024\), or where to draw the boundaries of a plume \(He et al., 2024\). Similarly, large-scale chemical transport models tend to be quite stiff, with internal parameters and processes not being variable, even if on the subgrid scale this may introduce considerable uncertainty \(Cohen and Prinn, 2011, Cohen and Wang 2014, Qin et al 2013; Hu et al., 2024; Lu et al., 2024\).](#)

Airborne remote sensing is a highly technical and costly approach to record CH₄ fluxes from landfills, coal basins, and oil and gas production (Krautwurst et al., 2021; Krautwurst et al., 2017; Kuhlmann et al., 2023), which suffers from not being able to monitor CH₄ emissions over long periods of time or in regions where the source is not well constrained (Brandt et al., 2014; Gorchov Negron et al., 2020; Hiller et al., 2014; Mehrotra et al., 2017; Molina et al., 2010). Satellite remote sensing can measure CH₄ under specific orbits where the source is known and identified (Jacob et al., 2016; Jacob et al., 2022; Plant et al., 2022; Varon et al., 2018; Zhang et al., 2020), but only after being calibrated by upward looking remotely sensed measurements (Tu et al., 2022), and only when the atmosphere is rain, cloud and aerosol free (Cohen and Prinn, 2011; Reuter et al., 2019; Sadavarte et al., 2021). TROPOMI and GOSAT have both been shown to be data-rich at times (Butz et al., 2012; Hu et al., 2018; Jacob et al., 2016), but severely limited at other times (Butz et al., 2012; Kuze et al., 2009). Even when these satellites have sufficient data to compute emissions from other species, frequently CH₄ emissions cannot be computed (Li et al., 2023; Qin et al., 2023b) due to insufficient signal strength, and uncertainties which are both non-understood and mis-constrained (Povey and Grainger, 2015).

Ground-based remote sensing provides higher accuracy versus satellite observations (Heerah et al., 2021; Luther et al., 2022; Tu et al., 2022). EM27/SUN measurements have approximated CH₄ emissions in Poland (Luther et al., 2019; Luther et al., 2022). However, these instruments are expensive, require calibration, and have limited data collection due to solar signal strength.

This work employs a high-frequency surface-based observation platform of CH₄ concentration which is portable, economical, and unaffected by most environmental factors. The observations are combined with a new mass-conserving methodology based on temporal transformation of the spatially derived mass-conserving framework successfully applied to NO₂ (Li et al., 2023; Qin et al., 2023b). This work focuses on Shanxi, one of the densest coal mining regions in the world, accounting for approximately 10% total global coal production (Lin and Liu, 2010; Qin et al., 2023a). Continuous observations were made around known coal mines, unknown sources, and of background conditions. [The 2-box model used the high-frequency emissions calculated using these data to drive for attributing](#) emissions to the known mine and a second low production mine previously thought insignificant. The results provide insights into the spatial distribution of CH₄ emissions, demonstrate rapid adoption of practical methods globally, and enable source attribution.

2 Method and Data

2.1 Study Site and Campaign Design

[Changzhi, Shanxi is located in a basin, with coal mines and associated coal use industries densely distributed throughout both flat central regions and around the mountainous edges \(Figure 1\), many of which are classified as high CH₄ emitting mines. Due to this combination, city-wide background CH₄ concentrations are very high and have large variation in time.](#) Due to this combination, province-wide background CH₄ concentrations are very high and have large variation in time. This study mainly focuses on two coal mines: one mine is classified as having high amounts of CH₄ [emissions](#) per unit of production and an annual coal production of 4 million tons (CM-A), and the other is unclassified for CH₄ [emissions](#) per unit of production and having an annual coal production of 3 million tons (CM-B) (Qin et al., 2023a). [Instruments](#) were positioned

Deleted: of

Deleted: High-frequency emissions calculated using these data were used to drive a 2-box model to attribute

Deleted: Observations

Formatted: Right: 0.63 cm

along concentric circles located 1km, 3km, and 5km from CM-A, over an approximation of the four ordinal directions: east, west, south, north (Figure 2). All locations were planned to be far away from known anthropogenic sources, leading to a net total 12 measurement points. As later discovered, CM-B is located approximately 1km southwest from the measurement point located at 5km west.




Figure 1. Topographic map of Changzhi, and its location in Shanxi Province (bottom left). The triangles represent the locations of all individual coal mines (including underground and abandoned mines), where the triangle color represents the emissions amount: high (red), middle (green), low (blue), and very-low (grey). The red stars represent the two coal mines in this work.





Figure 2. Locations of four individual coal mines (Green filled houses), a power plant (Red flag), and the 12 observation locations presented in this work (red double-outlined triangles). Distance from CM-A are given as concentric circles at 1km (blue), 3km (orange), and 5km (green).



Deleted:

Formatted: Right: 0.63 cm

2.2 Measuring CH₄ Concentration

Atmospheric CH₄ concentrations at 5 m above the surface were observed daily at 1 Hz from 8:30 to 17:00 local time in August 2022 using two portable greenhouse gas analyzers (LGR-915-0011, California, USA). Three different locations (1 km, 3 km and 5 km) were selected daily along a single direction from the CM-A, allowing a more consistent and precise calculation of the spatial gradient (Table S1). In order to reduce the time errors, using two portable greenhouse gas analyzers randomly selecting the three observation points during the daily measurements and without fixed sequence. [During field experiments, each day in a background location we confirmed that the standard baseline was found within its reliable range. Further, to ensure measurement consistency when the instrument's location was changed, after relocation, we rechecked the condition against the standard baseline. Another, the instrument was allowed to stabilize and any air from the previous location was flushed out, with the first 10 minutes of measurement data discarded, to ensure that only local air was observed.](#)

The CH₄ [concentrations](#) data was averaged minute-by-minute to match observed wind data, and subsequently used to compute CH₄ emissions. As show in Figure [S1 and S2](#), the CH₄ concentrations data is highly correlated with rapid changes in both the wind speed and direction.

Observations made in clean locations with a wind direction not from the mine are subsequently considered for background sites. The lowest and least variable CH₄ concentrations observations are found on August 23 in the south (2.08 ± 0.08 ppm) (Figure [S2](#)). It is important to note that although this site has the minimum concentrations observed in this work, these values are significantly higher than the global latitude-band background. Three other locations and days were observed with relatively low mean and not significantly large variation: August 19 in the east (2.63 ± 0.35 ppm), August 22 in the east (2.65 ± 0.51 ppm) (Figure [S1](#)), and August 22 in the south (2.60 ± 0.55 ppm) (Figure [S2](#)). [These results highlight the importance of accurately determining background concentrations in mass-balance emissions estimates. Unlike satellite-based emissions assessments, which often rely on separating plumes from global latitude bands or climatological background states \(Buchwitz et al.,](#)


2017; Irakulis-Loitxate et al., 2021; Lauvaux et al., 2022; Sadavarte et al., 2021), in situ measurements typically determine background concentrations by sampling upwind or outside the plume (Brantley et al. 2014). In this study, we have taken the spirit of the latter approach a step further, to ensure that concentrations observed as relatively clean are actually representative of locally background air. This way ensures that our background concentrations were representative of the local conditions not influenced directly by the site of interest (Figure S2). This method provides a more accurate approach when the baseline itself also changes, as in the specific locations sampled in this work.

2.3 Meteorological Data

The wind speed and direction were obtained from local meteorological stations with a temporal frequency of 1min. As show in Figures 3 and 4 the overall wind was dominated by a southerly direction (38.0% of observations between 150° and 210°) and found to be moderately slow (69.9% of observations were between 1 m s⁻¹ and 4 m s⁻¹). The 10th and 90th percentiles of wind direction (54° and 312°) and wind speed (1 m s⁻¹ and 5.1 m s⁻¹) respectively, indicate that high frequency sampling reveals a small number of relatively large changes are observed, which are expected to lead to a “fat-tail” type of distribution of subsequently computed CH₄ emissions (Delkash et al., 2016).

The temperature and pressure data were measured by a handheld meteorological instrument (HWS1000, ZOGLAB, China) with an accuracy of ± 0.5 °C for temperature and ± 0.5 hPa for pressure, ensuring reliable data collection. The meteorological instrument was calibrated according to the manufacturer’s guidelines prior to use. Measurements were taken at 5 s intervals to capture temporal variations in the atmospheric conditions. The temperature and pressure data were averaged minute-by-minute to match observed wind data, and subsequently used to convert CH₄ emissions unit (ppm min⁻¹) into policy-relevant unit (kg h⁻¹). The boundary layer data were obtained from <https://zenodo.org/records/6498004> (Guo et al., 2022) based on a merging of reanalysis data with observations (Guo et al., 2024).

Deleted: These results imply that the practice adopted by the community to separate a plume from the global latitude band or climatological background state is not applicable in the locations sampled in this paper (Buchwitz et al., 2017; Irakulis-Loitxate et al., 2021; Lauvaux et al., 2022; Sadavarte et al., 2021). For this reason, a new quantitative approach is presented to understand and quantify what is actually a source and what is not. This approach is applicable under conditions both encountered globally as well as those under the uniquely high and variable conditions observed herein



Formatted: Right: 0.63 cm




Figure 3. The wind rose of all observed wind speeds from August 10, 2022 to August 25, 2022.




Figure 4. Probability density function (PDF) for all observed wind direction (yellow) and wind speed (blue) from August 10, 2022 to August 25, 2022.

2.4 Quantitative Estimation of CH₄ Emissions

A mass conserving approach was used to estimate the CH₄ emissions using high frequency observations of CH₄ concentrations and meteorological data, hereafter called the Mass Conserving Model of Measured CMM (MCM²). This approach is based on previous dynamic emissions estimates of tropospheric atmospheric column observations of short-lived NO₂ (Li et al., 2023; Qin et al., 2023b), and CH₄ (Hu et al., 2024) but has never been applied to surface observations or at minute-frequency scale. Adopting this approach to solve for CH₄ emissions is done starting with the continuity equation for the conservation of mass (Equation 1), reorganizing the individual terms and converting coordinates from space to time (Equation 2) and finally combining the terms (Equation 3) as follows:

$$\frac{\partial[CH_4]}{\partial t} = E_{CH_4} - \nabla(\mathbf{U} \times [CH_4]) \quad (1)$$

Deleted: 5

Deleted: in connection with the

Deleted: in general, or CH₄ in specific

Deleted: E_{CH_4}

Formatted: Right: 0.63 cm

$$\nabla(U \times [CH_4]) = [CH_4] \times \nabla U + U \times \nabla[CH_4]$$

$$= \alpha \times \left([CH_4] \times \frac{\partial U}{\partial t} + U \times \frac{\partial [CH_4]}{\partial t} \right) \quad (2)$$

$$\frac{\partial [CH_4]}{\partial t} = ECH_4 - \alpha \times \left([CH_4] \times \frac{\partial U}{\partial t} + U \times \frac{\partial [CH_4]}{\partial t} \right) \quad (3)$$

where $[CH_4]$ is the CH_4 concentration (ppm), $\frac{\partial}{\partial t}$ is the temporal derivative operator, ECH_4 is the CH_4 emissions flux ($ppm \text{ min}^{-1}$), U is the wind speed ($m \text{ s}^{-1}$), and the ∇ is a mathematical operator acting on spatially distributed variables. However, when considering motion along one-dimension, the relationship between distance, speed, and time can be used to rewrite the spatial derivatives of $\nabla([CH_4])$ and $\nabla(U)$ as temporal derivatives (Brasseur and Jacob, 2017), where α is a conversion coefficient between distance and wind speed.

The gradient term used in these equations take into account the local topography of Shanxi, which is known for its significant features and surrounding mountains. These geographical features can impact the transport and dispersion of CH_4 , and their effects are incorporated into the wind field in the continuity equation. Notably, when dealing with a non-divergent wind field, the gradient term simplifies to the term $(U \times \nabla[CH_4])$ (Sun, 2022). Uncertainty analysis was conducted before calculating the CH_4 emissions to ensure only reliable data was used, since observed variation of CH_4 concentrations over time is influenced not only by CH_4 emissions, but also changes in wind speed and pressure. Specifically, $[CH_4] \times \frac{\partial U}{\partial t}$ represents the change in CH_4 concentrations influenced by pressure, while $U \times \frac{\partial [CH_4]}{\partial t}$ represents the change in CH_4 concentrations influenced by advection. Furthermore, since there is uncertainty in the observations, this work takes a conservative approach, and only considers data when the threshold given by equation (4) is observed to be considered influenced by emissions (a lower threshold can be selected like 25% or 15% et al., but uncertainty will increase).

$$U \times \frac{\partial [CH_4]}{\partial t} / \nabla(U \times [CH_4]) > 30\% \quad (4)$$

The remaining data (approximately 22%, presented as red circle indicators in Figure 5) is not processed in the emissions calculation as the signal is most likely due to a combination of observational uncertainty and white noise (Prinn et al., 1987; Conrad et al., 2023).

Deleted: E_{CH_4}

Deleted: t

Deleted: s the time (min)

Deleted: E_{CH_4}

Deleted: and

Deleted: .

Deleted: The

Deleted: operator that takes the gradient on spatially distributed variables

Deleted: u

Deleted: 6

Formatted: Right: 0.63 cm




Figure 5. Time series of CH₄ concentrations (top, blue), background concentrations (top, red), wind direction (bottom, blue) and wind speed (bottom, red) measured 5km west of CM-A on August 18, 2022. MCM² computed CH₄ emissions (top, numbers) (ppm min⁻¹) are computed for all regions where the observations are enhanced compared with the background for at least 3 consecutive observations, and which further pass the noise threshold (Equation 4).

2.5 Uncertainty Analysis

In order to reduce the uncertainty of the CH₄ emissions estimation, only data above the threshold given by equation (4) is considered. Prior to this, uncertainty analysis was also conducted on the relevant variables in actual experiments. As shown in Figure 6, a 5% uncertainty was assigned to both the CH₄ concentrations and wind speed data, and the CH₄ emissions was calculated. The uncertainty analysis results indicate that probability distribution of all possible calculated emissions are consistent, and the errors are smaller than 5% in each case, consistent with Equation 3 leading to a dampening of the uncertainty, as also observed in Qin et al. (2023a) study. Therefore, we believe that the results of CH₄ emissions in this study can be trusted.

Deleted: 6

Deleted: 7

Deleted: flux

Formatted: Right: 0.63 cm




Figure 6. The PDFs of uncertain analysis results (C represents CH_4 concentrations, W represents wind speed).

2.6 Attribution Analysis

A 2-box mass conserving model (based on equation 5) was used to attribute CH_4 emissions from the more than one suspected source of CH_4 emissions in the 5k west. An overview of the MCM² and 2-box mass conserving model used in this work is provided in Figure 7. The changes in CH_4 concentrations ($C_{\text{coal mine}}$ (ppm)) over time t (min) at the observation point, is driven by CH_4 emissions ($E_{\text{coal mine}}$ (ppm min⁻¹)) from the upwind coal mine and the CH_4 concentration blown away from the coal mine by the wind U (m s⁻¹), and the CH_4 background concentration ($C_{\text{background}}$ (ppm)) as demonstrated in Figure 7.

$$\frac{\partial C_{\text{coal mine}}}{\partial t} = E_{\text{coal mine}} + U \times C_{\text{background}} - U \times C_{\text{coal mine}} \quad (5)$$

All observed CH_4 concentrations data and estimated CH_4 emissions data are used when wind direction is capable of transporting the CH_4 from either CM-A or CM-B towards the observation site (Figure 10), while the remaining data is not used. A discretized version of Equation 5 is given in Equation 6 and solved using a first order finite difference approach:

$$C_{\text{Coal mine}_{\tau_{i+1}}} - C_{\text{coal mine}_{\tau_i}} = E_{\text{coal mine}_{\tau_i}} + U_{\tau_i} \times C_{\text{background}_{\tau_i}} - U_{\tau_i} \times C_{\text{coal mine}_{\tau_i}} \quad (6)$$

where τ_i and τ_{i+1} are the current and next time step, and the other terms are defined as in equation 5.

Deleted: 7

Moved (insertion) [1]

Moved up [1]: $C_{\text{coal mine}}$ (ppm)

Moved (insertion) [2]

Moved up [2]: $E_{\text{coal mine}}$ (ppm min⁻¹)

Deleted: gradient computed using

Deleted: 8

Deleted: 8

Formatted: Right: 0.63 cm

All possible sets of steady-state CH_4 concentrations are computed using all possible combinations of CH_4 emissions and observed CH_4 concentrations as boundary and initial conditions and running the equation forward to equilibrium.

The analyze probabilistically by comparing the modeled CH_4 concentrations probability density function (PDF) with the observed CH_4 concentrations PDF. Differences between the PDFs are clearly associated with the different wind directions and hence geophysical locations of the CH_4 emissions sources can be distinguished.

Deleted: computed concentrations are

Deleted: d




Figure 7. Overview of the MCM2 and 2-box mass conserving model used in this work.




Figure 8. Schematic diagram of the 2-Box model.

Deleted: 8

Formatted: Right: 0.63 cm

2.7 Converting Emissions into Policy-Relevant Units

In order to compare the emissions with some other studies, the units (ppm min^{-1}) were converted into policy-relevant units (kg h^{-1}), although as outlined below, this conversion leads to a larger uncertainty range. According to the attribution analysis in Section 2.6, when the wind direction is located within a 60° arc of coal mine A or coal mine B (Figure 8), the respective CH_4 emissions which successfully passed attribution were assigned to the respective coal mine. Therefore, based on the wind direction, the CH_4 emissions of coal mine A reported in this study use all of the data from the north 1 km CH_4 station, the CH_4 emissions of coal mine B considered all of the data from the west 5 km CH_4 station, and the background data came from all of the non-emissions data, adopting the following equation (7) to convert the units from ppm min^{-1} to kg h^{-1} :

$$E'_{\text{CH}_4} = E_{\text{CH}_4} \times \rho_{\text{air}} \times H \times A \times 60 \quad (7)$$

$$\rho_{\text{air}} = \frac{P \times M_{\text{air}}}{R \times T} \quad (8)$$

Where E'_{CH_4} is the CH_4 emissions with unit of kg h^{-1} , E_{CH_4} is the CH_4 emissions with unit of ppm min^{-1} , ρ_{air} is the dry gas density (kg m^{-3}) (based on equation 8), H is the height of the vertical rise that the emissions undergo within their first minute (m), A is the area (m^2) swept over an arc (Figure 9), which ranges linearly from 60° under slow wind conditions to 30° over very fast wind conditions, based on the wind speed when the direction is found to lead to successful attribution. P is the atmosphere pressure (Pa) over the sampling duration, M_{air} is the molecular weight of dry air, which is a fixed constant ($28.97 \times 10^{-3} \text{ kg mol}^{-1}$), R is the universal gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$), T is the air temperature (K) over the sampling duration.

Deleted: .

Deleted: 8

Deleted: CH_4 emissions of coal mine A were screened from the CH_4 emissions captured at North 1km, and CH_4 emissions of coal mine B were screened from the CH_4 emissions captured at West 5km

Deleted: E'_{CH_4} (E_{CH_4})

Deleted: E_{CH_4}

Deleted: E'_{CH_4}

Deleted: (

Deleted:)

Deleted: E_{CH_4}

Deleted: (

Deleted:)

Formatted: Right: 0.63 cm




Figure 9. The control volume for CH₄ emissions unit conversion from ppm min⁻¹ to kg h⁻¹.

Two different assumptions are made for the vertical extent of the plume rise, since the emissions are computed minute-by-minute which is shorter than the adjustment time throughout the entire boundary layer (Vaughn et al., 2018; Zinchenko et al., 2002). The first is to assume it has mixed within the bottom one fourth of the boundary layer, and the second is that it has mixed based on a steady vertical rise equal to one tenth of the horizontal wind. In this work, results using both assumptions will be presented.

3 Results and Discussion

3.1 Spatial Distribution Characteristics of CH₄ Concentration Around Coal Mine


Time series of CH₄ concentration, wind speed, and direction at 1 km, 3 km and 5 km north of CM-A are given in Figure 10. The wind direction predominantly blew from CM-A towards the observation point (wind direction is between 150° and 210°), for about 60% of the daily observation time. Only one day (August 15) observed at 1km north with a significant amount of wind from the west (wind direction is between 240°

Deleted: 9a


Deleted: and Figure 9b

Formatted: Right: 0.63 cm

and 300°), accounting for approximately 92.8% of the observation time on that day. Consistent with CM-A being the major source at 1km, when the wind blew from the south, the CH₄ concentration (3.45±0.79 ppm) was both higher and had a larger variation than when the wind blew from the west (2.40±0.17 ppm) which was similar to background conditions. This is consistent with there being no known significant sources to the west from this observation location, as shown in Figure 2. Similarly, under faster than average wind conditions from the direction of CM-A (on August 21 the mean wind was 5.70 m s⁻¹ with 14.9% of observations faster than 7 m s⁻¹), the observed CH₄ concentrations were slightly lower, yet similarly variable (3.17±0.82 ppm). All of these findings are consistent with transport dominating the CH₄ concentrations at 1 km north, and that high frequency wind and CH₄ concentration observations are required in tandem to compute the required spatial gradients in the CH₄ emissions, otherwise there is no basis to objectively separate the effects of the emitting region (CM-A) from the background.



Deleted: The wind direction blew from CM-A towards the observations (between 150° and 210°) 59.2% of the time, with only one day observed at 1km north (August 15) with a significant amount of wind from the west (between 240° and 300°) 92.8% of the time...



Deleted:

Formatted: Right: 0.63 cm

Figure 10. Time series of CH₄ concentrations (ppm), wind speed (m s⁻¹) and wind direction (°) measured at 1km (blue line), 3km (red line) and 5km (yellow line) located north of CM-A on two different days.




Figure 11. Time series of CH₄ concentrations (ppm), wind speed (m s⁻¹) and wind direction (°) measured at 1km (blue line), 3km (red line) and 5km (yellow line) located north of CM-A on two different days.

A similar set of findings were observed at 3 km north, while 5 km north is generally similar to the CH₄ background concentrations. At 3 km north, when the wind was from the south (59.3% of data), the CH₄ concentration was lower and more variable (3.16 ± 1.48 ppm, with 78.7% of observations below 3.0 ppm) than at 1 km north, consistent with advection from CM-A and a relatively stable atmosphere with a small contribution from diffusion between the plume and the background. When the wind blew from other directions, the distribution of CH₄ concentrations broadened considerably, with a range from background (2.25 ppm) through extremely polluted (16.2 ppm). One subset of this was observed on August 15 (observed over a total of 61 mins of observations, 6.68% of the total observations at 3 km north) when the wind was

Deleted: Figure 9. Time series of CH₄ concentration (ppm) (blue), wind speed (m s⁻¹) (yellow) and wind direction (°) (orange lines) measured at 1 km (solid), 3km (dashed) and 5km (dash-dot) located north (top) and west (bottom) of CM-A on four different days.

Formatted: Right: 0.63 cm

430 from the west and slow, where the CH₄ concentration was (5.44±2.82 ppm), as depicted
431 in Figure 10. The data on this day aligned with the presence of a major highway west
432 of the observation site, which was observed in-person to have heavy traffic consisting
433 of vehicles carrying coal (which could still be outgassing) as well as others powered by
434 compressed natural gas (CNG) (Figure 2). At 5 km north the overall CH₄ concentrations
435 (2.40±0.28 ppm) was generally lower than at 3km and had much lower variability,
436 consistent with background CH₄.

Deleted: 9

437 Time series of CH₄ concentrations measured at 1 km, 3 km and 5 km west of CM-
438 A and corresponding wind direction and speed are given in Figure 11. Overall, the main
439 wind direction is from the south 98.4% of the time at 1km, 74.5% of the time at 3km,
440 and 70.2% of the time at 5 km, and the wind speed was very high when measuring CH₄
441 at 1 km west, with an average value of 4.28±1.13 m s⁻¹ and a maximum of 7.4 m s⁻¹.
442 This set of findings is consistent with clean upwind sources. Accordingly at 1km west,
443 the observed CH₄ concentrations were slightly higher than CH₄ background
444 concentrations and had similar variability to 1km and 3km north (2.71±0.94 ppm and
445 86.5% of the data below 3 ppm). At 3 km west, CH₄ concentrations was observed to be
446 similar to the CH₄ background concentrations (2.32±0.09 ppm). The only exception
447 was found at 1 km west between 9:00 am and 9:30 am on August 17, in which all of the
448 observations of CH₄ concentrations were greater than 4 ppm. Since the areas to the west
449 from 1 km west contains mostly farmland, there was no expected strong sources of CH₄,
450 as shown in Figure 2. This indicates that during this special short time, the observed
451 slow increase and rapid fall-off in CH₄ concentrations must be due an unidentified
452 sources, or a change in the boundary layer or vertical mixing structure.

Deleted: 9c

Deleted: and Figure 9d

Deleted: was

453 Following this, it was anticipated that the 5 km west site would exhibit background
454 types of conditions, however the observed data deviates significantly. Wind speed was
455 low (1.63±0.54 m s⁻¹, maximum 3.0 m s⁻¹), CH₄ concentrations were both very high
456 and exhibited substantial temporal variability (5.83±2.99 ppm, 66.7% of CH₄
457 concentrations data exceeding 4 ppm, and peak of 15.3 ppm), and 70.2% of the CH₄
458 concentrations observations were from the south as demonstrated in Figure 12d,e,f.
459 From Figure 1, it can be seen that there is another coal mine (CM-B) located about 1

Deleted: was

Deleted: 10d

Formatted: Right: 0.63 cm

km away from the 5 km west measurement point, to its southwest, although CM-B has an annual production of about 3 million tons (smaller than CM-A) and not considered to be high gas (like CM-A), and therefore was not previously considered important. The overlap of high CH_4 concentrations with low a priori emissions, suggests that formal attribution is essential to quantitatively confirm whether CM-B is the source responsible for both typical conditions at 5 km west, as well as the long-range transport event at 1 km west.

CH_4 concentrations and wind observations in all directions except to the west, and except for the small number of special events documents above, exhibit PDFs that show there is a decrease in CH_4 concentrations the further the distance from CM-A (Figures 12 and 13), indicating that CM-A is consistent with the major CH_4 emission sources in these regions. These decreases of CH_4 concentrations were observed in terms of the median, mean, distribution width, and percentage over 4.0 ppm all decreasing from 1 km north to 3 km north and again from 3 km north to 5 km north.




Figure 12. Probability density map for CH_4 concentration and wind rose measured at 1 km north (a), 1 km west (d), 3 km north (b), 3 km west (e), 5 km north (c), and 5 km west (f) of CM-A and corresponding wind rose chart.

Deleted:

Deleted: 10

Deleted: 11

Deleted: are

Deleted: 10

Deleted: top left

Deleted: bottom left

Deleted: top center

Deleted: bottom center

Deleted: top right

Deleted: bottom right

Deleted: .

Formatted: Right: 0.63 cm




Figure 13. Probability density map for CH₄ concentrations and wind rose measured at 1 km east (a), 1 km south (d), 3 km east (b), 3 km south (e), 5 km east (c), and 5 km south (f) of CM-A and corresponding wind rose chart.

The observed CH₄ concentrations gradient as one moves westward from CM-A is inconsistent with the other ordinal directions (Figure 12d, e, f). While there was a small decrease in the mean and distribution breadth from 1 km west to 3 km west, there was a large increase in the median, mean, distribution width, percentage over 4.0 ppm from 3 km west to 5 km west. Furthermore, the data at 5 km west was found to be skewed differently than at the other sites, with approximately 70% of the CH₄ concentrations data greater than 4.0 ppm. The data clearly indicates that the 5 km west site behaves more like a CH₄ emission source region than even the 1 km north site.

3.2 Quantification and Emission Characteristics of CMM

The CH₄ emissions have been estimated at each of the observation points, with 25.7% of CH₄ concentrations observations yielding emissions results. The PDFs of the CH₄ emissions (Figures 14 and 15) reveal that the three stations in the north and the 5 km west station all are relatively high and variable, while the remainder are relatively low and non-variable. Among all the CH₄ emissions results, the highest median, mean, maximum, and breadth of the distribution are all observed at 5 km west. Another. The 3 km south location has the lowest CH₄ emissions of all points observed (by median), with a respective median, mean, maximum, and percentage greater than 1.0 ppm min⁻¹ of (0.03 ppm min⁻¹, 0.26 ppm min⁻¹, 0.90 ppm min⁻¹, 0%) (Figure 15), and is

Deleted: 11

Deleted: Probability density map for CH₄ concentration measured at 1 km, 3 km and 5 km east and south of CM-A and corresponding wind rose chart.

Deleted: 10d

Deleted: computed

Deleted: 12

Deleted: 13

Deleted: /

Deleted: 13

Formatted: Right: 0.63 cm

subsequently considered representative of CH_4 background emissions in this work. It is important to note that there is no area within this region that has 0 ppm min⁻¹ emissions and that the minimum CH_4 concentration on average is about 2.23 ppm (Figure 13), both of which are considered very high or polluted compared with most other current studies (Irakulis-Loitxate et al., 2021; Sadavarte et al., 2021).




Figure 14. Probability density functions (PDF) of CH_4 emissions located at 1 km north (a), 3 km north (b), 5 km north (c), 1 km west (d), 3 km west (e), and 5 km west (f) of CM-A, including median, mean, maximum, and minimum statistics.




Figure 15. Probability density functions (PDF) of CH_4 emissions located at 1 km east (a), 3 km east (b), 5 km east (c), 1 km south (d), 3 km south (e), and 5 km south (f) of CM-A, including median, mean, maximum, and

Deleted: 11

Deleted: 12

Deleted: computed

Deleted: 13

Deleted: computed

Formatted: Right: 0.63 cm

545 minimum statistics.

546 The spatial distribution [characteristics](#) of the CH₄ emissions is similar to that of
547 the CH₄ concentrations observations (Figure [14](#)). First, there is a decrease as
548 [observation points](#) moves northward along the axis away from CM-A, with the median,
549 mean, maximum, and percentage of emissions greater than 1.0 ppm min⁻¹ at 1 km north
550 (0.73 ppm min⁻¹, 1.18 ppm min⁻¹, 5.67 ppm min⁻¹, and 42%) all larger than at 3km north
551 (0.28 ppm min⁻¹, 0.72 ppm min⁻¹, 3.41 ppm min⁻¹, and 29%). The [CH₄ emissions](#) values
552 at 3 km north are also larger than those at 5 km north, which respectively are 0.11 ppm
553 min⁻¹, 0.18 ppm min⁻¹, and 0.59 ppm min⁻¹, and 0%. The subset of [CH₄ emissions](#) under
554 low wind speed conditions exhibited a larger decline from 1 km to 3 km and from 3 km
555 to 5 km. The observations are further consistent with transport from a single dominant
556 source located at CM-A being the primary driving factor, and diffusion from other
557 industrial sources in Changzhi city center being a secondary factor.

558 Consistent with there being few to no sources impacting the 1 km west and 3 km
559 west sites, except for considerably less transport from CM-A the [CH₄ emissions](#) PDFs
560 at these sites (Figure [14](#)) demonstrate low [CH₄ emissions](#) and low variability, with the
561 respective median, mean, maximum, and percentage of [CH₄ emissions](#) greater than 1.0
562 ppm min⁻¹ at 1 km west being 0.28 ppm min⁻¹, 0.55 ppm min⁻¹, 3.03 ppm min⁻¹, and
563 16% and at 3km west being even lower (0.08 ppm min⁻¹, 0.10 ppm min⁻¹, 0.27 ppm
564 min⁻¹, and 0%). However, the CH₄ emissions [values](#) at 5 km west were the highest and
565 most variable of all results [in this work](#), with the respective statistics being 1.45 ppm
566 min⁻¹, 1.82 ppm min⁻¹, 7.92 ppm min⁻¹, and 60%. Furthermore, the skewness of the
567 distribution at 5 km west (which has 30% of the CH₄ emissions above 2.0 ppm min⁻¹)
568 is much larger than at 1 km north (which only has 15% of emissions above 2.0 ppm
569 min⁻¹). Combining these pieces of information, at first look it seems that the site at 5
570 km west is not related to the [CH₄ emissions](#) from CM-A, or at best are a mixture of [CH₄](#)
571 emissions from CM-A and those at another site, herein proposed to be CM-B. The
572 remainder of this study focuses on disentangling and attributing contributions from
573 CM-A and CM-B at 5 km west, with the observations at the remaining sites ruled out
574 in terms of having a contribution from CM-B.

Deleted: Characteristics

Deleted: 12

Deleted: one

Deleted: computed

Deleted: 12

Deleted: computed

Deleted: computed

Formatted: Right: 0.63 cm

3.3 Attribution of CH₄ Emissions

This work applied the 2-box model at the 5 km west site and quantified the contribution of both CM-A and CM-B CH₄ emissions to the observed CH₄ concentration distributions as given in Figure 16. First, the results of the 2-box model produce CH₄ concentrations PDFs which overlap with the overall observed CH₄ concentrations PDF, indicating that the results are reasonable. Second, space of the CH₄ emissions from the two different two coal mines do not overlap, and cover two independent portions of the observed CH₄ concentrations PDF. Specifically, the 30%, 50%, and 70% values of CH₄ concentrations observed at 5km west are 3.68 ppm, 5.18 ppm, and 6.86 ppm respectively. The CH₄ emissions from CM-A yield a CH₄ concentration less than 4 ppm most of the time, with a 30%, 50%, 70%, and maximum concentration of 2.96 ppm, 3.15 ppm, 3.31 ppm, and 4.60 ppm, while the CH₄ emissions from CM-B yield a CH₄ concentration more than 5 ppm most of the time, with a minimum, 30%, 50%, 70%, and maximum concentration of 4.76 ppm, 5.20 ppm, 5.68 ppm, and 6.18 ppm.

Overall, the CH₄ emissions from CM-B cover well the observed CH₄ concentration values from the range of 50% to 70%, with a single high value around the 90% value, while the CH₄ emissions from CM-A cover well the observed CH₄ concentration values in the range from 10% to 30%. One weakness is that the length of observation time is not as comprehensive as at the other observation sites, and therefore it is possible that had more observation time been made, the CH₄ contributions from CM-B would have filled more of the space between the 70% and 90% levels, and some combination of sources from CM-A and CM-B would have better filled the space between the 30% and 50% levels. The results indicate to a high degree of certainty that the effect of CH₄ emissions from the two respective coal mines on CH₄ concentrations at west 5 km are distinct, with CM-A the source of emissions in the lower range of the CH₄ concentration distribution and CM-B the source for emissions in the higher CH₄ concentration range, covering values in the middle and upper range. Improvements in modeling, additional observations, considering possible contributions from additional missing sources, and consideration of longer-range transport could add further improvement and better

Deleted: 14

Deleted: computed


Deleted: s

Deleted: s

Deleted: The results indicate to a high degree of certainty that the emissions from the two respective coal mines are distinct

Formatted: Right: 0.63 cm

618 explore the intermediate range of observed concentrations.



619 **Figure 16. Probability density functions (PDF) of CH₄ concentrations (a) observed at the wet 5km site (blue),**
 620 **and concentrations simulated by the 2-box model resulting from emissions attributed to (b) Coal mine A (red),**
 621 **and (c) Coal mine B (orange). The respective 30th, 50th, and 70th percentiles of the distributions are given in**
 622 **text, outlined by the corresponding color box.**

624 3.4 Policy-Relevant Emissions

625 In order to compare the values of CH₄ emissions from the Shanxi coal mines computed
 626 in section 2.7, the units (ppm min⁻¹) are transformed into units of (kg h⁻¹) via a
 627 conversion factor based on equations (7) and (8). This conversion increases the overall
 628 uncertainty, since since it involves approximations of the area swept, the boundary layer
 629 height, and and other uncertainties. In this study, average CH₄ emissions from CM-A
 630 and CM-B are 6860±3520 kg h⁻¹ and 1010±347 kg h⁻¹, and the CH₄ emissions range is
 631 from 102 kg hr⁻¹ to 19000 kg h⁻¹ and 185 kg hr⁻¹ to 1720 kg h⁻¹ repectively (Table2).
 632 Both mines display a fat tail distribution, with respective 25th, median, and 75th
 633 percentile values of [1600, 3070, 10500] kg h⁻¹ and [755, 1086, 1416] kg h⁻¹ respectively.
 634 These findings are demonstrated to be higher than CH₄ emissions from equivalent oil
 635 and gas operations in the USA (Chen et al., 2022), with one site being roughly double
 636 and the other similar to and slightly lower than day-to-day emissions inverted over 5-
 637 years from TROPOMI (Hu et al., 2024) over the same region. This is consistent with
 638 the fact that the results herein target very high frequency and spatially confined
 639 emissions, while satellites provide day-to-day values over a larger pixel area, as well as
 640 associated significant uncertainties involved in converting from ppm to kg. In specific,

Deleted: Figure 14. The PDFs of CH₄ concentration measured at 5 km west (blue) and simulated using the 2-Box model under conditions when the source is CM-A (red), and when the sources is CM-B (orange), including representative 30%, 50%, and 70% bounds are in.

Formatted: Right: 0.63 cm

at CM-A, the minimum value (102 kg h⁻¹) of CH₄ emissions is greater than the minimum value (8 kg h⁻¹) of CH₄ emissions inverted from TROPOMI, the maximum value (19.0 ton h⁻¹) of CH₄ emissions is less than the maximum value (24.9 ton h⁻¹) of CH₄ emissions from TROPOMI. Similarly, at CM-B, the minimum value (185 kg h⁻¹) of CH₄ emissions is greater than the minimum value (20 kg h⁻¹) of CH₄ emissions inverted from TROPOMI, the maximum value (1720 kg h⁻¹) of CH₄ emissions is much smaller than the maximum value (37.3 ton h⁻¹) of CH₄ emissions from TROPOMI. In this study, observations were made within 1 km of the coal mines on a minute-to-minute basis, while the TROPOMI observed the XCH₄ over a space scale (5.5×7 km²) and on a day-to-day average basis, the higher temporal resolution of our in-situ measurements offers an advantage in capturing short-term variations and extreme values within the study period when compared with TROPOMI's results, due to both the fat-tail distribution and the strong temporal variation of the observations. For this reason, it is likely that the sampling time (two days) at CM-B was insufficient to fully capture the fat tail of the CH₄ emissions. The estimation of CH₄ emissions from coal mine B in this study carries multiple sources of uncertainty. First, the site has a smaller amount of data than the other sites, and therefore may be less statistically representative. Second, the site generally has a lower production level and is thought to have a lower emission, however, its method of venting is also not known and there could be other such reasons why it could have very variable emissions over time. Third, the TROPOMI emissions themselves are occurring on different grids, which have different properties in terms of surface albedo, aerosols, and other factors, and therefore will be more uncertain at site B than site A, since site A was constrained using ground-trothed data. Finally, there likely are additional sources in the same upwind regions and direction, but further from the observations than site B, which would require additional measurements in the other three ordinal directions from site B to further disentangle. More accurate CH₄ emission would require additional monitoring data over a longer time, as well as in the other three ordinal directions around site B. Furthermore, coal mine B has a relatively small coal production compared to coal mine A. Given that coal mine B is only about 6 km away from coal mine A and shares the same geological environment, its CH₄ emissions

should theoretically be lower than those from coal mine A, but only assuming if the ventilation technology and work schedules were consistent with each other, for which we have no a priori information.

Table 2. The CH₄ emissions (kg h⁻¹) of CM-A and CM-B using different observation methods and statistical methods. H means the height.

Coal mines	High frequency ground observation CH ₄ emissions (kg h ⁻¹)								TROPOMI inverted CH ₄ emissions (kg h ⁻¹)			
	<u>H: Lowest</u> boundary layer				<u>H: Vertical</u> wind×time							
	Mean±SE	Min	Median	Max	Mean±SE	Min	Median	Max	Mean±SE	Min	Median	Max
CM-A	6860±3520	102	3070	19000	8200±4400	67	3160	22800	5500±700	8	2130	24900
CM-B	1010±347	185	1090	1720	200±65	34	211	337	6200±1000	20	1450	37300

4 Conclusions

This study presents a high frequency ground observation campaign and a new analytical top down emissions estimation approach to quantify the emissions of CH₄ from a high gas coal mine region with multiple mines. The base observations are made using a portable greenhouse gas analyzer in connection with meteorological and other optical measurements. Observations have been made over 15 days at a frequency of 1 Hz, at various locations of known distance from an existing high production coal mine. The high frequency observations are then used in connection with a mass conserving modeling platform to estimate the CH₄ emissions rate. A mass-conserving Two-Box model was used for attribution analysis in this study. The results show that the spatial characteristics of CH₄ concentration/emissions are consistent with the distance from a well characterized of single coal mine within 5km distance, and CH₄ emissions demonstrate clear first order effects of both transport and diffusion, with CH₄ emission rates of 0.73, 0.28 and 0.15 ppm min⁻¹ at 1, 3 and 5 km downwind respectively. At 5 km north the overall CH₄ concentration (2.40±0.28 ppm) was generally lower than at 3km and had much lower variability, consistent with background CH₄, which demonstrate that the CMM emissions mainly affect the surrounding area within 5 km distance. However, the overlap of two coal mines (CM-A and CM-B) have a far more complex distribution of CH₄ emissions intensity, ranging as high as 1.82 ppm min⁻¹, which is much higher than the emissions of single source at any directions. Another, the

Deleted: Given that the emissions are generally larger at the higher production coal mine, they are consistent with the concept that over geologically similar environments, higher coal mine production leads to increased CH₄ emissions, although the increase is not linear as most current models assume.

Formatted: Right: 0.63 cm

background concentration of surface CH₄ in the mining areas is very high compared with other studies, with a value always at and above 2.23 ppm. Finally, in order to compare these results with results from other parts of the world, the subset of emissions which successfully underwent attribution were converted into the unit of kg h⁻¹ using an approximation of the volume swept by the wind and other approximations of the atmosphere. The resulting values were found to be 6860±3520 kg h⁻¹ and 1010±347 kg h⁻¹ respectively, which are higher than CH₄ emissions from equivalent oil and gas operations in the USA, and in one case are higher than but in the other case similar to day-to-day emissions inverted from 5-years of TROPOMI over the same region.

This work demonstrates that high frequency surface observations of CH₄, in combination with high frequency observations of wind can provide deep insights into emissions by accounting for high frequency changes in space and time at the same time, which tend to be missing from models which used more idealized approaches (such as average plume shapes and sizes, levels of coal production, and interpreting gradients from a small number of fixed images). A significant source of CH₄ emissions from a previously unknown or improperly classified mine may pose a vastly different range of observed concentration as well as computed emissions than expected. The importance of observations at both high frequency and regional spatial coverage are demonstrated, and a set of practical methods that are freely open and can be adopted and modified rapidly are provided. The approach to source attribution used herein can provide insights to policymakers to formulate regional emission control policies and provide a check on or a priori assumption for the new generation of advance satellite-based top-down emissions estimates, while demonstrating that spatial attribution is a critical next-step for satellite approximations and CH₄ control policies.

Data Availability. All underlying data herein are available for access by the editors and reviewers at <https://figshare.com/s/1a393772d7b72ae17e62> and will be made available to the community upon publication.

Author contributions. K.Q., J.B.C. and F.L. designed the research; F.L., C.Y., and Y.S. collected the data; J.B.C. and F.L. analyzed the data; Q.T., W. H. and Q.X. provided the support for data analysis and drawing; Q.H., S.W gave suggestion on running the 2-Box model; F.L. wrote the manuscript with

Formatted: Right: 0.63 cm

inputs from J.B.C., Q.H. and P.T.; All authors discussed the results and contributed to the final manuscript.

Competing interests. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgments. We sincerely appreciate all the scientists, engineers, and students who participated in the field campaigns, maintained the measurement instruments, and helped with and collection and processing of the data.

Finacial support. This study was funded by the National Natural Science Foundation of China (42075147, 42375125) and the Fundamental Research Funds for the Central Universities (2023KYJD1003, JP230021).

References

Allen, D. T.: Methane emissions from natural gas production and use: reconciling bottom-up and top-down measurements, *Curr. Opin. Chem.*, 5, 78-83, <https://doi.org/10.1016/j.coche.2014.05.004>, 2014.

Bloom, A. A., Bowman, K. W., Lee, M., Turner, A. J., Schroeder, R., Worden, J. R., Weidner, R., McDonald, K. C., and Jacob, D. J.: A global wetland methane emissions and uncertainty dataset for atmospheric chemical transport models (WetCHARTs version 1.0), *Geosci. Model Dev.*, 10, 2141-2156, <https://gmd.copernicus.org/articles/10/2141/2017/>, 2017.

Bournazian, J.: US. Energy Information Administration, <https://hdl.handle.net/1813/4504>, 2016.

Brandt, A. R., Heath, G. A., Kort, E. A., O'Sullivan, F., Pétron, G., Jordaan, S. M., Tans, P., Wilcox, J., Gopstein, A. M., Arent, D., Wofsy, S., Brown, N. J., Bradley, R., Stucky, G. D., Eardley, D., and Harriss, R.: Methane Leaks from North American Natural Gas Systems, *Science*, 343, 733-735, <https://www.science.org/doi/abs/10.1126/science.1247045>, 2014.

Brantley, H. L., Thoma, E. D., Squier, W. C., Guven, B. B., and Lyon, D.: Assessment of methane emissions from oil and gas production pads using mobile measurements, *Environ. Sci. Technol.*, 48(24), 14508-14515, <https://pubs.acs.org/doi/full/10.1021/es503070q>, 2014.

Brasseur, G. P., and Jacob, D. J.: *Model Equations and Numerical Approaches*, in: *Modeling of atmospheric chemistry*, Cambridge University Press Publishing, 84-91, <https://books.google.com.hk/books>, 2017.

Bruno, J. H., Jervis, D., Varon, D. J., and Jacob, D. J.: U-Plume: automated algorithm for plume detection and source quantification by satellite point-source imagers, *Atmos. Meas. Tech.*, 17, 2625-2636, <https://amt.copernicus.org/articles/17/2625/2024/>, 2024.

Buchwitz, M., Schneising, O., Reuter, M., Heymann, J., Krautwurst, S., Bovensmann, H., Burrows, J. P., Boesch, H., Parker, R. J., Somkuti, P., Detmers, R. G., Hasekamp, O. P., Aben, I., Butz, A., Frankenberg, C., and Turner, A. J.: Satellite-derived methane hotspot emission estimates using a fast data-driven method, *Atmos. Chem. Phys.*, 17, 5751-5774, <https://acp.copernicus.org/articles/17/5751/2017/>, 2017.

Formatted: Right: 0.63 cm

777 Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: TROPOMI aboard Sentinel-5
778 Precursor: Prospective performance of CH₄ retrievals for aerosol and cirrus loaded atmospheres, Remote
779 Sens. Environ., 120, 267-276, <https://doi.org/10.1016/j.rse.2011.05.030>, 2012.

780 Cao, X.: Policy and regulatory responses to coalmine closure and coal resources consolidation for
781 sustainability in Shanxi, China, J. Clean. Prod., 145, 199-208,
782 <https://doi.org/10.1016/j.jclepro.2017.01.050>, 2017.

783 Chen, Y., Sherwin, E. D., Berman, E. S. F., Jones, B. B., Gordon, M. P., Wetherley, E. B., Kort, E. A.,
784 and Brandt, A. R.: Quantifying Regional Methane Emissions in the New Mexico Permian Basin with a
785 Comprehensive Aerial Survey, Environ. Sci. Technol., 56, 4317-4323,
786 <https://doi.org/10.1021/acs.est.1c06458>, 2022.

787 Cohen, J. B., and Prinn, R. G.: Development of a fast, urban chemistry metamodel for inclusion in global
788 models, Atmos. Chem. Phys., 11, 7629-7656, <https://acp.copernicus.org/articles/11/7629/2011/>, 2011.

789 Cohen, J. B., and Wang, C.: Estimating global black carbon emissions using a top-down Kalman Filter
790 approach, J. Geophys Res: Atmos., 119, 307-323, <https://doi.org/10.1002/2013JD019912>, 2014.

791 Conrad, B. M., Tyner, D. R., and Johnson, M. R.: Robust probabilities of detection and quantification
792 uncertainty for aerial methane detection: Examples for three airborne technologies. Remote Sens.
793 Environ., 288, 113499, <https://doi.org/10.1016/j.rse.2023.113499>, 2023.

794 Delkash, M., Zhou, B., Han, B., Chow, F. K., Rella, C. W., and Imhoff, P. T.: Short-term landfill methane
795 emissions dependency on wind, Waste Management, 55, 288-298,
796 <https://doi.org/10.1016/j.wasman.2016.02.009>, 2016.

797 Duren, R. M., Thorpe, A. K., Foster, K. T., Rafiq, T., Hopkins, F. M., Yadav, V., Bue, B. D., Thompson,
798 D. R., Conley, S., Colombi, N. K., Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Falk, M., Herner,
799 J. D., Croes, B. E., Green, R. O., and Miller, C. E.: California's methane super-emitters, Nature, 575,
800 180-184, <https://doi.org/10.1038/s41586-019-1720-3>, 2019.

801 Etminan, M., Myhre, G., Highwood, E. J., and Shine, K. P.: Radiative forcing of carbon dioxide, methane,
802 and nitrous oxide: A significant revision of the methane radiative forcing, Geophys. Res. Lett., 43, 12614-
803 12623, <https://doi.org/10.1002/2016GL071930>, 2016.

804 Goldsmith, C. D., Chanton, J., Abichou, T., Swan, N., Green, R., and Hater, G.: Methane emissions from
805 20 landfills across the United States using vertical radial plume mapping, J. Air Waste Manage., 62, 183-
806 197, <https://doi.org/10.1080/10473289.2011.639480>, 2012.

807 Gorchoy Negron, A. M., Kort, E. A., Conley, S. A., and Smith, M. L.: Airborne Assessment of Methane
808 Emissions from Offshore Platforms in the U.S. Gulf of Mexico, Environ. Sci. Technol., 54, 5112-5120,
809 <https://doi.org/10.1021/acs.est.0c00179>, 2020.

810 Guo, J., Zhang, J., Shao, J., Chen, T., Bai, K., Sun, Y., Li, N., Wu, J., Li, R., Li, J., Guo, Q., Cohen, J. B.,
811 Zhai, P., Xu, X., and Hu, F.: A merged continental planetary boundary layer height dataset based on high-
812 resolution radiosonde measurements, ERA5 reanalysis, and GLDAS: Earth Syst. Sci. Data, v. 16, no. 1,
813 p. 1-14, <https://essd.copernicus.org/articles/16/1/2024/essd-16-1-2024.html>, 2024.

814 Heerah, S., Frausto-Vicencio, I., Jeong, S., Marklein, A. R., Ding, Y., Meyer, A. G., Parker, H. A., Fischer,
815 M. L., Franklin, J. E., Hopkins, F. M., and Dubey, M.: Dairy Methane Emissions in California's San
816 Joaquin Valley Inferred With Ground-Based Remote Sensing Observations in the Summer and Winter,
817 Journal of Geophys Res: Atmo., 126, e2021JD034785, <https://doi.org/10.1029/2021JD034785>, 2021.

818 He, T.-L., Boyd, R. J., Varon, D. J., and Turner, A. J.: Increased methane emissions from oil and gas
819 following the Soviet Union's collapse, PNAS, 121, e2314600121,
820 <https://doi.org/10.1073/pnas.2314600121>, 2024.

821 Hiller, R. V., Neiningner, B., Brunner, D., Gerbig, C., Bretscher, D., Künzle, T., Buchmann, N., and Eugster,
822 W.: Aircraft-based CH₄ flux estimates for validation of emissions from an agriculturally dominated area
823 in Switzerland, Journal of Geophys Res: Atmos., 119, 4874-4887,

Formatted: Right: 0.63 cm

824 <https://doi.org/10.1002/2013JD020918>, 2014.

825 Hu, H., Landgraf, J., Detmers, R., Borsdorff, T., Aan de Brugh, J., Aben, I., Butz, A., and Hasekamp, O.:
 826 Toward Global Mapping of Methane With TROPOMI: First Results and Intersatellite Comparison to
 827 GOSAT, *Geophys. Res. Lett.*, 45, 3682-3689, <https://doi.org/10.1002/2018GL077259>, 2018.

828 Irakulis-Loitxate, I., Guanter, L., Liu, Y.-N., Varon, D. J., Maasakkers, J. D., Zhang, Y., Chulakadabba,
 829 A., Wofsy, S. C., Thorpe, A. K., Duren, R. M., Frankenberg, C., Lyon, D. R., Hmiel, B., Cusworth, D.
 830 H., Zhang, Y., Segl, K., Gorroño, J., Sánchez-García, E., Sulprizio, M. P., Cao, K., Zhu, H., Liang, J., Li,
 831 X., Aben, I., and Jacob, D. J.: Satellite-based survey of extreme methane emissions in the Permian basin,
 832 *Sci. Adv.*, 7, eabf4507, <https://www.science.org/doi/abs/10.1126/sciadv.abf4507>, 2021.

833 Jacob, D. J., Turner, A. J., Maasakkers, J. D., Sheng, J., Sun, K., Liu, X., Chance, K., Aben, I., McKeever,
 834 J., and Frankenberg, C.: Satellite observations of atmospheric methane and their value for quantifying
 835 methane emissions, *Atmos. Chem. Phys.*, 16, 14371-14396,
 836 <https://acp.copernicus.org/articles/16/14371/2016/>, 2016.

837 Jacob, D. J., Varon, D. J., Cusworth, D. H., Dennison, P. E., Frankenberg, C., Gautam, R., Guanter, L.,
 838 Kelley, J., McKeever, J., Ott, L. E., Poulter, B., Qu, Z., Thorpe, A. K., Worden, J. R., and Duren, R. M.:
 839 Quantifying methane emissions from the global scale down to point sources using satellite observations
 840 of atmospheric methane, *Atmos. Chem. Phys.*, 22, 9617-9646,
 841 <https://acp.copernicus.org/articles/22/9617/2022/>, 2022.

842 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., and Petrescu, A. M. R.: EDGAR v4.3.2
 843 Global Atlas of the three major Greenhouse Gas Emissions for the period 1970–2012, *Earth Syst. Sci.*
 844 *Data*, 1-55, <https://doi.org/10.5194/essd-2017-79>, 2017.

845 Karion, A., Sweeney, C., Pétron, G., Frost, G., Michael Hardesty, R., Kofler, J., Miller, B. R., Newberger,
 846 T., Wolter, S., Banta, R., Brewer, A., Dlugokencky, E., Lang, P., Montzka, S. A., Schnell, R., Tans, P.,
 847 Trainer, M., Zamora, R., and Conley, S.: Methane emissions estimate from airborne measurements over
 848 a western United States natural gas field, *Geophys. Res. Lett.*, 40, 4393-4397,
 849 <https://doi.org/10.1002/grl.50811>, 2013.

850 Katzenstein, A. S., Doeze, L. A., Simpson, I. J., Blake, D. R., and Rowland, F. S.: Extensive regional
 851 atmospheric hydrocarbon pollution in the southwestern United States, *PNAS*, 100, 11975-11979,
 852 <https://www.pnas.org/doi/abs/10.1073/pnas.1635258100>, 2003.

853 Kerr, T., and Yang, M.: Coal mine methane in China: A budding asset with the potential to bloom, IEA
 854 Information Paper, 1-35, [https://www.iea.org/reports/coal-mine-methane-in-china-a-budding-asset-](https://www.iea.org/reports/coal-mine-methane-in-china-a-budding-asset-with-the-potential-to-bloom)
 855 [with-the-potential-to-bloom](https://www.iea.org/reports/coal-mine-methane-in-china-a-budding-asset-with-the-potential-to-bloom), 2009.

856 Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P.,
 857 Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L.,
 858 Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque,
 859 J.-F., Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D.,
 860 Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I.
 861 J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van
 862 Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and
 863 sinks, *Nat. Geosci.*, 6, 813-823, <https://doi.org/10.1038/ngeo1955>, 2013.

864 Krautwurst, S., Gerilowski, K., Borchardt, J., Wildmann, N., Gałkowski, M., Swolkień, J., Marshall, J.,
 865 Fiehn, A., Roiger, A., Ruhtz, T., Gerbig, C., Necki, J., Burrows, J. P., Fix, A., and Bovensmann, H.:
 866 Quantification of CH₄ coal mining emissions in Upper Silesia by passive airborne remote sensing
 867 observations with the Methane Airborne MAPper (MAMAP) instrument during the CO₂ and Methane
 868 (CoMet) campaign, *Atmos. Chem. Phys.*, 21, 17345-17371,
 869 <https://acp.copernicus.org/articles/21/17345/2021/>, 2021.

870 Krautwurst, S., Gerilowski, K., Jonsson, H. H., Thompson, D. R., Kolyer, R. W., Iraci, L. T., Thorpe, A.
 871 K., Horstjann, M., Eastwood, M., Leifer, I., Vigil, S. A., Krings, T., Borchardt, J., Buchwitz, M.,
 872 Fladeland, M. M., Burrows, J. P., and Bovensmann, H.: Methane emissions from a Californian landfill,
 873 determined from airborne remote sensing and in situ measurements, *Atmos. Meas. Tech.*, 10, 3429-3452,
 874 <https://amt.copernicus.org/articles/10/3429/2017/>, 2017.

Formatted: Right: 0.63 cm

875 Kuhlmann, G., Brunner, D., Emmenegger, L., Schwietzke, S., Zavala-Araiza, D., Thorpe, A., Hueni, A.,
876 and Röckmann, T.: Quantifying methane super-emitters from oil and gas production in Romania with the
877 AVIRIS-NG imaging spectrometer, EGU General Assembly 2023, Vienna, Austria., 24–28 Apr 2023
878 EGU23-6751.

879 Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon
880 observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse
881 gases monitoring, *Appl. Optics.*, 48, 6716-6733, [https://opg.optica.org/ao/abstract.cfm?URI=ao-48-35-](https://opg.optica.org/ao/abstract.cfm?URI=ao-48-35-6716)
882 [6716](https://opg.optica.org/ao/abstract.cfm?URI=ao-48-35-6716), 2009.

883 Lauvaux, T., Giron, C., Mazzolini, M., d'Aspremont, A., Duren, R., Cusworth, D., Shindell, D., and Ciais,
884 P.: Global assessment of oil and gas methane ultra-emitters, *Science*, 375, 557-561,
885 <https://www.science.org/doi/abs/10.1126/science.abj4351>, 2022.

886 Li, Q., Fernandez, R. P., Hossaini, R., Iglesias-Suarez, F., Cuevas, C. A., Apel, E. C., Kinnison, D. E.,
887 Lamarque, J.-F., and Saiz-Lopez, A.: Reactive halogens increase the global methane lifetime and
888 radiative forcing in the 21st century, *Nat. Commun.*, 13, 2768, [https://doi.org/10.1038/s41467-022-](https://doi.org/10.1038/s41467-022-30456-8)
889 [30456-8](https://doi.org/10.1038/s41467-022-30456-8), 2022.

890 Li, X., Cohen, J. B., Qin, K., Geng, H., Wu, L., Wu, X., Yang, C., Zhang, R., and Zhang, L.: Remotely
891 Sensed and Surface Measurement Derived Mass-Conserving Inversion of Daily High-Resolution NO_x
892 Emissions and Inferred Combustion Technologies in Energy Rich Northern China, *Atmos. Chem. Phys.*,
893 23, 8001-8019, <https://egusphere.copernicus.org/preprints/2023/egusphere-2023-2/>, 2023,.

894 Lin, B.-q., and Liu, J.-h.: Estimating coal production peak and trends of coal imports in China, *Energ.*
895 *Policy*, 38, 512-519, <https://doi.org/10.1016/j.enpol.2009.09.042>, 2010.

896 [Lu, L., Cohen, J. B., Qin, K., Li, X., and He, Q.: Identifying Missing Sources and Reducing NO_x](https://doi.org/10.1016/j.enpol.2009.09.042)
897 [Emissions Uncertainty over China using Daily Satellite Data and a Mass-Conserving Method.](https://doi.org/10.1016/j.enpol.2009.09.042)
898 [EGUsphere \[ACCEPTED\], https://doi.org/10.5194/egusphere-2024-1903, 2024.](https://doi.org/10.1016/j.enpol.2009.09.042)

899 Luther, A., Kleinschek, R., Scheidweiler, L., Defratyka, S., Stanisavljevic, M., Forstmaier, A., Dandocsi,
900 A., Wolff, S., Dubravica, D., Wildmann, N., Kostinek, J., Jöckel, P., Nickl, A. L., Klausner, T., Hase, F.,
901 Frey, M., Chen, J., Dietrich, F., Nećki, J., Swolkień, J., Fix, A., Roiger, A., and Butz, A.: Quantifying
902 CH₄ emissions from hard coal mines using mobile sun-viewing Fourier transform spectrometry, *Atmos.*
903 *Meas. Tech.*, 12, 5217-5230, <https://amt.copernicus.org/articles/12/5217/2019/>, 2019.

904 Luther, A., Kostinek, J., Kleinschek, R., Defratyka, S., Stanisavljević, M., Forstmaier, A., Dandocsi, A.,
905 Scheidweiler, L., Dubravica, D., Wildmann, N., Hase, F., Frey, M. M., Chen, J., Dietrich, F., Nećki, J.,
906 Swolkień, J., Knot, C., Vardag, S. N., Roiger, A., and Butz, A.: Observational constraints on methane
907 emissions from Polish coal mines using a ground-based remote sensing network, *Atmos. Chem. Phys.*,
908 22, 5859-5876, <https://acp.copernicus.org/articles/22/5859/2022/>, 2022.

909 Mehrotra, S., Faloona, I., Suard, M., Conley, S., and Fischer, M. L.: Airborne Methane Emission
910 Measurements for Selected Oil and Gas Facilities Across California, *Environ. Sci. Technol.*, 51, 12981-
911 12987, <https://doi.org/10.1021/acs.est.7b03254>, 2017.

912 Miller, S. M., Michalak, A. M., Detmers, R. G., Hasekamp, O. P., Bruhwiler, L. M. P., and Schwietzke,
913 S.: China's coal mine methane regulations have not curbed growing emissions, *Nat. Commun.*, 10, 303,
914 <https://doi.org/10.1038/s41467-018-07891-7>, 2019.

915 Molina, L. T., Madronich, S., Gaffney, J. S., Apel, E., de Foy, B., Fast, J., Ferrare, R., Herndon, S.,
916 Jimenez, J. L., Lamb, B., Osornio-Vargas, A. R., Russell, P., Schauer, J. J., Stevens, P. S., Volkamer, R.,
917 and Zavala, M.: An overview of the MILAGRO 2006 Campaign: Mexico City emissions and their
918 transport and transformation, *Atmos. Chem. Phys.*, 10, 8697-8760,
919 <https://acp.copernicus.org/articles/10/8697/2010/>, 2010.

920 Peng, S., Piao, S., Bousquet, P., Ciais, P., Li, B., Lin, X., Tao, S., Wang, Z., Zhang, Y., and Zhou, F.:
921 Inventory of anthropogenic methane emissions in mainland China from 1980 to 2010, *Atmos. Chem.*
922 *Phys.*, 16, 14545-14562, <https://acp.copernicus.org/articles/16/14545/2016/>, 2016.

923 Plant, G., Kort, E. A., Murray, L. T., Maasakkers, J. D., and Aben, I.: Evaluating urban methane emissions

Formatted: Right: 0.63 cm

924 from space using TROPOMI methane and carbon monoxide observations, *Remote Sen. Environ.*, 268,
925 112756, <https://doi.org/10.1016/j.rse.2021.112756>, 2022.

926 Povey, A. C., and Grainger, R. G.: Known and unknown unknowns: uncertainty estimation in satellite
927 remote sensing, *Atmos. Meas. Tech.*, 8, 4699-4718, <https://amt.copernicus.org/articles/8/4699/2015/>,
928 2015.

929 Prather, M. J., Holmes, C. D., and Hsu, J.: Reactive greenhouse gas scenarios: Systematic exploration of
930 uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, L09803,
931 <https://doi.org/10.1029/2012GL051440>, 2012.

932 Prinn, R., Cunnold, D., Rasmussen, R., Simmonds, P., Alyea, F., Crawford, A., Fraser, P., and Rosen, R.:
933 Atmospheric Trends in Methylchloroform and the Global Average for the Hydroxyl Radical, *Science*,
934 238, 945-950, <https://www.science.org/doi/abs/10.1126/science.238.4829.945>, 1987.

935 Qin, K., Hu, W., He, Q., Lu, F., and Cohen, J. B.: Individual Coal Mine Methane Emissions Constrained
936 by Eddy-Covariance Measurements: Low Bias and Missing Sources, *EGUsphere*, 1-49,
937 <https://egusphere.copernicus.org/preprints/2023/egusphere-2023-1210/>, 2023a.

938 Qin, K., Lu, L., Liu, J., He, Q., Shi, J., Deng, W., Wang, S., and Cohen, J. B.: Model-free daily inversion
939 of NO_x emissions using TROPOMI (MCMFE-NO_x) and its uncertainty: Declining regulated emissions
940 and growth of new sources, *Remote Sens. Environ.*, 295, 113720,
941 <https://doi.org/10.1016/j.rse.2023.113720>, 2023b.

942 Reuter, M., Buchwitz, M., Schneising, O., Krautwurst, S., O'Dell, C. W., Richter, A., Bovensmann, H.,
943 and Burrows, J. P.: Towards monitoring localized CO₂ emissions from space: co-located regional CO₂
944 and NO₂ enhancements observed by the OCO-2 and S5P satellites, *Atmos. Chem. Phys.*, 19, 9371-9383,
945 <https://acp.copernicus.org/articles/19/9371/2019/>, 2019.

946 Rigby, M., Park, S., Saito, T., Western, L. M., Redington, A. L., Fang, X., Henne, S., Manning, A. J.,
947 Prinn, R. G., Dutton, G. S., Fraser, P. J., Ganesan, A. L., Hall, B. D., Harth, C. M., Kim, J., Kim, K. R.,
948 Krummel, P. B., Lee, T., Li, S., Liang, Q., Lunt, M. F., Montzka, S. A., Mühle, J., O'Doherty, S., Park,
949 M. K., Reimann, S., Salameh, P. K., Simmonds, P., Tunncliffe, R. L., Weiss, R. F., Yokouchi, Y., and
950 Young, D.: Increase in CFC-11 emissions from eastern China based on atmospheric observations, *Nature*,
951 569, 546-550, <https://doi.org/10.1038/s41586-019-1193-4>, 2019.

952 Sadavarte, P., Pandey, S., Maasakkers, J. D., Lorente, A., Borsdorff, T., Denier van der Gon, H.,
953 Houweling, S., and Aben, I.: Methane Emissions from Superemitting Coal Mines in Australia Quantified
954 Using TROPOMI Satellite Observations, *Environ. Sci. Technol.*, 55, 16573-16580,
955 <https://doi.org/10.1021/acs.est.1c03976>, 2021.

956 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A.,
957 Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P.,
958 Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N.,
959 Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin,
960 M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M.,
961 Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov,
962 S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores,
963 F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C.,
964 Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H.,
965 Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N.,
966 Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D.,
967 Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The
968 Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561-1623,
969 <https://essd.copernicus.org/articles/12/1561/2020/>, 2020a.

970 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., and Zhuang, Q.: The Global Methane Budget 2000–
971 2017, *Earth System Science Data*, 12, 1561-1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020b.

972 Shi, T., Han, G., Ma, X., Mao, H., Chen, C., Han, Z., Pei, Z., Zhang, H., Li, S., and Gong, W.: Quantifying
973 factory-scale CO₂/CH₄ emission based on mobile measurements and EMISSION-PARTITION model:
974 cases in China, *Environ. Res. Lett.*, 18, 034028, <https://dx.doi.org/10.1088/1748-9326/acbee7>, 2023.

Formatted: Right: 0.63 cm

975 Shi, T., Han, Z., Han, G., Ma, X., Chen, H., Andersen, T., Mao, H., Chen, C., Zhang, H., and Gong, W.:
 976 Retrieving CH₄-emission rates from coal mine ventilation shafts using UAV-based AirCore observations
 977 and the genetic algorithm–interior point penalty function (GA-IPPF) model, *Atmos. Chem. Phys.*, 2,
 978 13881-13896, <https://acp.copernicus.org/articles/22/13881/2022/>, 2022.

979 Sun, K.: Derivation of Emissions From Satellite-Observed Column Amounts and Its Application to
 980 TROPOMI NO₂ and CO Observations, *Geophys. Res. Lett.*, 49, e2022GL101102,
 981 <https://doi.org/10.1029/2022GL101102>, 2022.

982 Tong, X., van Heuven, S., Scheeren, B., Kers, B., Hutjes, R., and Chen, H.: Aircraft-Based AirCore
 983 Sampling for Estimates of N₂O and CH₄ Emissions, *Environ. Sci. Technol.*, 57, 15571-15579,
 984 <https://doi.org/10.1021/acs.est.3c04932>, 2023.

985 Tu, Q., Hase, F., Schneider, M., García, O., Blumenstock, T., Borsdorff, T., Frey, M., Khosrawi, F.,
 986 Lorente, A., Alberti, C., Bustos, J. J., Butz, A., Carreño, V., Cuevas, E., Curcoll, R., Diekmann, C. J.,
 987 Dubravica, D., Ertl, B., Estruch, C., León-Luis, S. F., Marrero, C., Morgui, J. A., Ramos, R., Scharun, C.,
 988 Schneider, C., Sepúlveda, E., Toledano, C., and Torres, C.: Quantification of CH₄ emissions from waste
 989 disposal sites near the city of Madrid using ground- and space-based observations of COCCON,
 990 TROPOMI and IASI, *Atmos. Chem. Phys.*, 22, 295-317, <https://acp.copernicus.org/articles/22/295/2022/>,
 991 2022.

992 Varon, D. J., Jacob, D. J., McKeever, J., Jervis, D., Durak, B. O. A., Xia, Y., and Huang, Y.: Quantifying
 993 methane point sources from fine-scale satellite observations of atmospheric methane plumes, *Atmos.*
 994 *Meas. Tech.*, 11, 5673-5686, <https://amt.copernicus.org/articles/11/5673/2018/>, 2018.

995 Vaughn, T. L., Bell, C. S., Pickering, C. K., Schwietzke, S., Heath, G. A., Pétron, G., Zimmerle, D. J.,
 996 Schnell, R. C., and Nummedal, D.: Temporal variability largely explains top-down/bottom-up difference
 997 in methane emission estimates from a natural gas production region, *PNAS*, 115, 11712-11717,
 998 <https://www.pnas.org/doi/abs/10.1073/pnas.1805687115>, 2018.

999 Vinković, K., Andersen, T., de Vries, M., Kers, B., van Heuven, S., Peters, W., Hensen, A., van den Bulk,
 1000 P., and Chen, H.: Evaluating the use of an Unmanned Aerial Vehicle (UAV)-based active AirCore system
 1001 to quantify methane emissions from dairy cows, *Sci. Total Environ.*, 831, 154898,
 1002 <https://doi.org/10.1016/j.scitotenv.2022.154898>, 2022.

1003 Wecht, K. J., Jacob, D. J., Sulprizio, M. P., Santoni, G. W., Wofsy, S. C., Parker, R., Bösch, H., and
 1004 Worden, J.: Spatially resolving methane emissions in California: constraints from the CalNex aircraft
 1005 campaign and from present (GOSAT, TES) and future (TROPOMI, geostationary) satellite observations,
 1006 *Atmos. Chem. Phys.*, 14, 8173-8184, <https://acp.copernicus.org/articles/14/8173/2014/>, 2014.

1007 Zhang, Y., Gautam, R., Pandey, S., Omara, M., Maasakkers, J. D., Sadavarte, P., Lyon, D., Nesser, H.,
 1008 Sulprizio, M. P., Varon, D. J., Zhang, R., Houweling, S., Zavala-Araiza, D., Alvarez, R. A., Lorente, A.,
 1009 Hamburg, S. P., Aben, I., and Jacob, D. J.: Quantifying methane emissions from the largest oil-producing
 1010 basin in the United States from space, *Sci. Adv.*, 6, eaaz5120,
 1011 <https://www.science.org/doi/abs/10.1126/sciadv.aaz5120>, 2020.

1012 Zinchenko, A. V., Paramonova, N. N., Privalov, V. I., and Reshetnikov, A. I.: Estimation of methane
 1013 emissions in the St. Petersburg, Russia, region: An atmospheric nocturnal boundary layer budget
 1014 approach, *J. Geophys. Res.-Atmos.*, 107, ACH 2-1-ACH 2-11, <https://doi.org/10.1029/2001JD001369>,
 1015 2002.

1016

1017