



# IMPROVING LARGE-SCALE SNOW ALBEDO MODELLING USING A CLIMATOLOGY OF LIGHT-ABSORBING PARTICLE DEPOSITION

Manon Gaillard<sup>1,2</sup>, Vincent Vionnet<sup>1</sup>, Matthieu Lafaysse<sup>3</sup>, Marie Dumont<sup>3</sup>, and Paul Ginoux<sup>4</sup>

<sup>1</sup>Environment and Climate Change Canada, Meteorological Research Division, Dorval, Canada

<sup>2</sup>Ecole Polytechnique, Palaiseau, FRANCE

<sup>3</sup>Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Etudes de la Neige, 38000 Grenoble, France

<sup>4</sup>NOAA/OAR, Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, USA

**Correspondence:** Vincent Vionnet (vincent.vionnet@ec.gc.ca)

**Abstract.** Light-absorbing particles (LAPs) deposited at the snow surface significantly reduce its albedo and strongly affect the snow melt dynamics. The explicit simulation of these effects with advanced snow radiative transfer models is generally associated with a large computational cost. Consequently, many albedo schemes used in snowpack models still rely on empirical parameterizations that do not account for the spatial variability of LAP deposition. In this study, a new strategy of intermediate complexity that includes the effects of spatially variable LAP deposition on snow albedo is tested with the snowpack model Crocus. It relies on an optimization of the parameter that controls the evolution of snow albedo in the visible range. Optimized values for multi-year snow albedo simulations with Crocus were generated at ten reference experimental sites spanning a large variety of climates across the world. A regression was then established between these optimal values and climatological deposition of LAP on snow at the location of the experimental sites extracted from a global climatology developed in this study. This regression was finally combined with the global climatology to obtain an LAP-informed and spatially variable parameter for the Crocus albedo parameterization. The revised parameter improved snow albedo simulations on average by 10% with the largest improvements found in the Arctic (more than 25%). This methodology can be applied to other land surface models using the global climatology of LAP deposition on snow developed for this study.

## 15 1 Introduction

Snow is a key component of the Earth surface energy balance and water cycle (Armstrong and Brun, 2009; Flanner et al., 2011) and provides critical water resources for ecosystems and industrial applications (irrigation, hydro-power, ...) (Sturm et al., 2017; Immerzeel et al., 2020). Snow albedo, the fraction of incident solar radiation reflected by the snow, strongly impacts the surface radiative balance and influences the mass balance of the snow cover through modified snow melt and sublimation (Qu and Hall,



20 2006; Painter et al., 2017; Skiles et al., 2018; Réveillet et al., 2022). Snow albedo depends on different factors including snow physical properties (e.g. grain size), solar conditions (e.g. solar zenith angle, presence of clouds), and the abundance and optical properties of light absorbing particles (Warren and Wiscombe, 1980; Tuzet et al., 2019; He and Flanner, 2020).

Light-absorbing particles (LAPs) are small impurities often deposited from the atmosphere such as mineral dust (Painter et al., 2010) and black carbon (Flanner et al., 2007), but can also be living organisms such as algae (Cook et al., 2017). Black  
25 carbon (BC) is the optically absorbing portion of soot and originates from the incomplete combustion of fossil and biofuels (Bond et al., 2013). Mineral dust originates from arid and semi-arid landscapes and can be transported by the wind over long distances to other parts of the globe (e.g. Di Mauro et al., 2015). When LAPs are present in snow, they darken the snow surface and decrease its albedo in the visible range (e.g., Warren and Wiscombe, 1980). The broadband albedo of fresh snow can drop  
30 an increase of snow grain size (through accelerated snow metamorphism), which results in a decrease in snow albedo in the near-infrared range and further increases radiative forcing from the LAPs present in the snowpack. There are therefore two main effects of LAP presence in snow: a direct effect through darkening of the surface, and an associated feedback through grain-coarsening (Painter et al., 2007; Skiles et al., 2018).

Snow radiative transfer models of various complexity have been developed to simulate the impact of LAPs on snow albedo  
35 (Warren and Wiscombe, 1980; Flanner and Zender, 2005; Libois et al., 2013; He and Flanner, 2020; He, 2022). They simulate snow albedo in the visible and near-infrared wavelengths for given snowpack optical properties (specific surface area or optical grain size) and LAP contents. Radiative transfer models have been coupled to snowpack models to simulate the temporal and spatial evolution of snow radiative properties and account for the albedo feedbacks (Flanner et al., 2007; Tuzet et al., 2017). For example, Libois et al. (2015) and Tuzet et al. (2017) have coupled the Two-stream Analytical Radiative TransfER in Snow  
40 (TARTES) radiative transfer scheme (Libois et al., 2013) with the detailed snowpack model Crocus (Vionnet et al., 2012; Lafaysse et al., 2017). Among numerous applications, these advanced coupled models have been recently used to (i) quantify the impact of LAPs on snow cover evolution in mountainous terrain (Réveillet et al., 2022), (ii) study the impacts of snow cover on energy fluxes over the Tibetan Plateau (Hao et al., 2023), and (iii) assess the influence of LAPs on snowpack stability (Dick et al., 2023).

45 Several challenges arise when combining a snow radiative transfer model with a snow model. They are generally associated with the computational costs and the spectral resolution of the radiative transfer model (Flanner et al., 2007), although methodologies have been developed to optimize spectral snow albedo calculation (van Dalum et al., 2019; Veillon et al., 2021). The need for additional atmospheric forcings (LAP deposition fluxes) to drive the snow model (Tuzet et al., 2017) represents another challenge. For this reason, snowpack models used in land surface and hydrological models still often rely on empirical  
50 parameterizations to describe the temporal evolution of snow albedo (Pedersen and Winther, 2005; Lee et al., 2023). Simple time/temperature-dependent parameterizations (e.g., Verseghy, 1991; Douville et al., 1995) have been developed to represent the combined effects on snow albedo of multiple snow aging processes (increase in grain size due snow metamorphism, LAP deposition, ...). These parameterizations use fixed time constants that have been optimized using sparse observational data, restricting their extension to untested environments and time periods (e.g. Mölders et al., 2008). More advanced snow models



55 implemented in land surface schemes such as BATS (Dickinson, 1993), JULES (Best et al., 2011) and ISBA (Vionnet et al.,  
2012; Decharme et al., 2016) simulate the snow albedo evolution in different large spectral bands and include explicitly the ef-  
fect of optical grain size on albedo in these different bands. These models represent only indirectly the impact of LAPs on snow  
albedo in the visible range through parameters that require calibration. For example, the default version of the Crocus snow  
60 scheme used in support of operational avalanche forecasting (Vionnet et al., 2012) relies on an aging coefficient approximately  
representing the darkening of the snow with time. This parameter was calibrated in the French Alps (Brun et al., 1992) and  
does not account for the spatial variability of LAP deposition. Applications of the model to other climates such as Antarctica  
where LAP deposition is extremely low require to tune the rate of albedo decrease in the visible range (Brun et al., 2011).  
Such tuning is not possible when the model is applied at large scales, which limits the quality of continental-scale snowpack  
simulations with Crocus (Brun et al., 2013; Mortimer et al., 2020).

65 The objective of this study is to develop a methodology to improve large scale simulations of snow albedo in snowpack  
schemes by taking into account the spatial variability of LAP deposition. This methodology is applied in this paper to the  
default snow albedo parameterization of the detailed snowpack model Crocus to allow a better robustness of the model when  
applied at large spatial scales. Optimized parameters for snow albedo simulations with Crocus were generated at ten reference  
experimental sites spanning a large variety of climates. A regression was then established between these optimal parameters  
70 and climatological deposition of LAPs (BC and dust) on snow at the location of the experimental sites extracted from a global  
climatology developed in this study. This regression was finally combined with the global climatology to obtain LAP-informed  
and spatially-variable parameters for the Crocus albedo parameterization. This methodology could be applied to optimize  
parameters controlling the albedo evolution in the visible range in other snowpack schemes (e.g., Dickinson, 1993; Best et al.,  
2011; Decharme et al., 2016). The paper is organized as follows. Section 2 presents the snow albedo parameterization in  
75 Crocus. Section 3 then details the model configuration, the evaluation data and method as well as the datasets used to build  
the global climatology of LAP deposition on snow. Results are presented in Section 4 followed by a discussion in Section 5.  
Finally, Section 6 summarizes the results and offers concluding remarks.

## 2 Snow albedo parameterization in Crocus

Crocus is a multilayer snowpack model used for avalanche hazard forecasting, climate, and hydrology applications (Brun et al.,  
80 1992; Vionnet et al., 2012; Lafaysse et al., 2017). It simulates the seasonal evolution of the physical properties of the snowpack  
and its vertical layering. For each snow layer, Crocus simulates the evolution of the thickness, density, liquid water content,  
temperature, age, and snow microstructure represented by the snow specific surface area and a semi-empirical variable. This  
study relies on the version of Crocus that has recently been implemented as an additional option for snow simulations in the  
Soil, Vegetation and Snow version 2 (SVS-2) land surface scheme (Garnaud et al., 2019; Vionnet et al., 2022). SVS-2 is the land  
85 surface scheme used at Environment and Climate Change Canada (ECCC) in preparation of the Terrestrial Snow Mass Mission  
(Derksen et al., 2021). Within SVS-2, Crocus is coupled to a multi-layer soil model including soil freezing (Boone et al., 2000).  
In this version of Crocus, the maximum number of simulated snow layers was set at 20 to test a viable configuration for an



**Table 1.** Equations representing snow albedo for the three spectral bands in Crocus. The parameters are as follows:  $d_{opt}$  (m) is the optical grain diameter of the snow,  $P$  (Pa) is the mean pressure at the site,  $P_{CDP}$  (Pa) is the mean pressure at the Col de Porte site,  $A$  (days) is the age of the snow, and  $\gamma$  (days) is the snow aging coefficient. Adapted from Table 4 in Vionnet et al. (2012).


Spectral band	Spectral albedo $\alpha_{spectral\ band}$
0.3—0.8 $\mu\text{m}$	$\alpha_{0.3-0.8\mu\text{m}} = \max(0.6, \alpha_i - \Delta\alpha_{age})$ <p>where : <math>\alpha_i = \min(0.92, 0.96 - 1.58\sqrt{d_{opt}})</math>                      and : <math>\Delta\alpha_{age} = \min(1, \max(\frac{P}{P_{CDP}}, 0.5)) \times 0.2\frac{A}{\gamma}</math></p>
0.8—1.5 $\mu\text{m}$	$\alpha_{0.8-1.5\mu\text{m}} = \max(0.3, 0.9 - 15.4\sqrt{d_{opt}})$
1.5—2.8 $\mu\text{m}$	$\alpha_{1.5-2.8\mu\text{m}} = 346.3d' - 32.31\sqrt{d'} + 0.88$ <p>where : <math>d' = \min(d_{opt}, 0.0023)</math></p>

eventual operational implementation covering the whole Canadian territory at 500-m resolution; such a configuration needs to balance accuracy and computational time. The rest of this section describes the default snow albedo parameterization in Crocus.

Snow albedo in Crocus is split between three spectral bands of incoming solar radiation: one in the visible (0.3 – 0.8  $\mu\text{m}$ ), and two in the near infrared (0.8 – 1.5  $\mu\text{m}$  and 1.5 – 2.8  $\mu\text{m}$ ). The albedo in each spectral band is calculated using a different equation (Table 1). In the visible band, snow albedo depends mostly on the amount of LAPs, which is parameterized by the age of snow, and on the snow microstructure, represented by the optical diameter of snow (Carmagnola et al., 2014). In the near-infrared bands, the spectral albedo depends only on the optical diameter of snow. To obtain the total albedo, the model assumes by default that the incoming shortwave radiation is split into the three bands as follows: 71% in the 0.3—0.8  $\mu\text{m}$  range, 21% in the 0.8—1.5  $\mu\text{m}$  range, and 8% in the 1.5—2.8  $\mu\text{m}$  range. The total albedo is therefore defined as:

$$\alpha = 0.71\alpha_{0.3-0.8\mu\text{m}} + 0.21\alpha_{0.8-1.5\mu\text{m}} + 0.08\alpha_{1.5-2.8\mu\text{m}} \quad (1)$$

The impact of the deposition of light absorbing particles is parameterized by the age of snow through a snow ageing coefficient,  $\gamma$  (Table 1). This parameter controls the impact of the snow age  $A$  on the temporal evolution of the albedo in the visible band, as shown on Fig. 1. If  $\gamma$  is high, the albedo decreases slowly with the snow age, which is characteristic of a snowpack that receives low LAP deposition. In contrast, if  $\gamma$  is low, the effect of snow aging on albedo is larger, so that the snow albedo decreases more quickly, which can be associated with higher LAP concentrations at the snow surface. Figure 1 illustrates the dependency of the snow albedo to the snow age for different values of gamma and a constant value of the optical diameter. The default value of  $\gamma$  used until now in Crocus simulations is 60 days. It has been set during the early developments of Crocus at the Col de Porte experimental site in the French Alps (Brun et al., 1992). When applied in Antarctica, Brun et al. (2011) used a value of 900 days to take into account the exceptionally clean atmosphere over the Antarctic plateau so that the snow albedo decrease was drastically reduced in the visible range.



**Figure 1.** Graphic representation of the dependency of the snow albedo in the visible spectral band (0.3 – 0.8 $\mu$ m) to the snow age for different values of  $\gamma$  and a constant value of the optical diameter,  $d_{opt} = \frac{6}{\rho_{ice}SSA}$ , where  $\rho_{ice} = 917 \text{ kgm}^{-3}$  (at 0°C, Libois et al. (2015)) and  $SSA = 50 \text{ m}^2\text{kg}^{-1}$  (maximum for fresh snow at the Dome C site, Libois et al. (2015))

The narrowband albedo in the visible range also includes an empirical term that was initially designed to reflect the impact of elevation on the concentration of LAPs in the snow when applied in the French Alps for avalanche hazard forecasting (Brun et al., 1992). This element is the pressure term,  $A_P$  in the last line of the 0.3 – 0.8  $\mu$ m albedo equations in Table 1:

$$A_P = \min(1, \max(\frac{P}{P_{CDP}}, 0.5)), \quad (2)$$

where  $P_{CDP}=870 \text{ hPa}$  is the mean pressure at the Col de Porte experimental site in the French Alps (1325 m) (Morin et al., 2012).  $A_P$  decreases with the site's elevation, thus decreasing the impact of the snow age on spectral albedo and mimicking the effect of lower LAP concentrations at high elevations. The idea behind this mechanism is that sites which are at a high elevation in the French Alps are supposedly further away from pollution sources such as roads and cities located in valleys, and consequently have lower concentrations of LAPs. However, the added value of this parameterization has never been objectively evaluated and the relation between LAP impact and elevation could be more complex, as suggested in Réveillet et al. (2022), where a stronger influence of LAPs on the snow melt onset date was found at higher elevations. In the context of this work, considering large scale applications of Crocus, the pressure term was not considered when computing  $\Delta\alpha_{age}$  (Table 1). Instead, the impact of elevation on the deposition of LAPs was included by considering the altitudinal gradient of LAP deposition around each study site as described in Section 4.2.2.

### 3 Data and methods

First, the skills of snow albedo simulations with Crocus were examined for different values of the snow ageing coefficient,  $\gamma$ , at 10 reference sites around the world, to find optimal ranges of  $\gamma$  for each site. Then, a global climatology of mean-annual



LAP deposition rates on snow was computed, from which the values at each of the 10 reference sites were extracted. A cross analysis was then carried out to find a simple relationship between the LAP climatology and the optimal ranges of  $\gamma$  at the reference sites. This relationship was finally applied to the global climatology of LAP depositions, to obtain a global dataset of new and improved  $\gamma$  values that can be used for large scale application of Crocus.

### 130 3.1 Snow albedo simulations

#### 3.1.1 Data

This study used data from 10 sites spanning various snow cover types such as taiga snow, alpine snow and maritime snow (Table 2 and Fig. 2). These sites were selected because of the availability of reference meteorological data to drive snowpack simulations and reference snow measurements (including snow albedo) to evaluate the simulations. Six of these sites were  
135 taken from the ESM-SnowMIP dataset (Ménard et al., 2019), a series of ten sites with standardized and quality-checked observations and meteorological driving data. Among the ten original ESM-SnowMIP sites, the three located in forested areas (Old Aspen, Old Black Spruce, and Old Jack Pine) were not selected due to the additional effect of forest debris on snow albedo (Melloh et al., 2001). Out of the seven remaining sites, snow albedo measurements were not provided for two sites – Reynolds Mountain East (Idaho, USA), and Sodankylä (Finland) – in the original ESM-SnowMIP dataset (Ménard et al., 2019).  
140 However, the Sodankylä site has more recent albedo measurements which were not included in the ESM-SnowMIP dataset but were made available by the Finn. Met. Inst. (2018). A corresponding meteorological forcing file was built extending in time the methodology of Essery et al. (2016) as detailed in Sect. S2.1 in Supplementary Material. Therefore, Sodankylä was included with this alternative observation and meteorological dataset.


To include more diverse locations, several other sites were also considered in the analysis. Three sites in the Canadian Arctic  
145 region (Bylot, Umiujaq, and Trail Valley Creek) were added, as well as one alpine site in Austria (Kühtai). This yielded a total of 10 sites for this study. To guarantee homogeneity between all ten sites, the observed daily-average albedo at the added sites was computed from hourly incoming and outgoing shortwave radiation using the same methods as for the ESM-SnowMIP sites (Morin et al., 2012; Ménard et al., 2019). The only difference was the addition of a filter ensuring that the hourly incoming radiation was greater than the outgoing radiation. Hours that did not meet this requirement were not included in the computation  
150 of the daily-averaged albedo, and days where less than 5 hours were compliant were not included. Hourly incoming longwave radiation from the ERA5 reanalysis (Hersbach et al., 2020) was used at Bylot to drive Crocus due to uncertainty in the observed longwave radiation at this site (Domine et al., 2021). For all the other sites, the observed longwave radiation was used.

On top of observed albedo data, daily observations of snow depth ( $SD$ ) were also extracted at these ten sites. They were used to select days for the evaluation of albedo simulations (Section 3.1.2), and to evaluate the final results of this study (Section  
155 3.3).



**Table 2.** List of the meteorological sites used in this study and some of their characteristics.

Site Name	Code	Source	Time Period	Elevation (m)	Latitude, Longitude (°)	Snow Cover
Bylot	BYL	Domine et al. (2021)	2014-2019	22	73.15, -80.00	Taiga
Col de Porte	CDP	Ménard et al. (2019)	1994-2014	1325	45.30, 5.77	Alpine
Kühtai	KUT	Krajčí et al. (2017), Günther et al. (2019)	1990-2013	1920	47.21, 11.01	Alpine
Sapporo	SAP	Ménard et al. (2019)	2005-2015	15	43.08, 141.34	Maritime
Senator Beck	SNB	Ménard et al. (2019)	2005-2015	3714	37.91, -107.73	Alpine
Sodankylä	SOD	Finn. Met. Inst. (2018), Corcket (2023)	2012-2022	179	67.37, 26.63	Taiga
Swamp Angel	SWA	Ménard et al. (2019)	2005-2015	3371	37.91, -107.71	Alpine
Trail Valley Creek	TVC	Tutton et al. (2024)	2013-2018	91	68.75, -133.50	Taiga
Umiujaq	UMQ	Lackner et al. (2023)	2012-2020	130	56.56, -76.48	Taiga
Weissfluhjoch	WFJ	Ménard et al. (2019)	1996-2016	2536	46.83, 9.81	Alpine



**Figure 2.** Map showing the location of the experimental sites used in this study.

### 3.1.2 Methods

Multi-year snowpack simulations were carried out with the snowpack model Crocus within SVS2 at the ten experimental sites described in Sect. 3.1.1. The model configurations (meteorological forcing heights, soil texture, vegetation type, etc.) were obtained from the reference papers describing each site (Table 2). A spin-up period of four years at each site (the first four





160 years of forcing dataset) was considered to provide initial conditions for the surface and soil column. The physical options used in the multiphysics version of Crocus are detailed in Table A1.

Simulations were first run at each of the ten sites for values of  $\gamma$  between 5 and 900 days, with an eleven-point distribution based on an exponential: [5, 10, 20, 30, 60, 90, 150, 250, 400, 650, 900] days. The maximal value of 900 days corresponds to the value used by Brun et al. (2011) in Antarctica. The minimum values of 5 days corresponds to a decrease in snow albedo of 165 0.2 in one week (Fig. 1), representative of maximal albedo decrease during LAP deposition events (e.g. Dumont et al., 2020). A range of  $\gamma$  containing the values which yield the most skilled simulations was found at each site, using a method described further on in this section. After this first round of simulations with  $\gamma$  in the [5, 900] days interval, a second round of simulations was run at each site for a zoomed interval around the range of best  $\gamma$  values found during the first round. The zoomed intervals were manually chosen for each site as a compromise between granularity of the explored  $\gamma$  values and reasonable computing 170 time. As in the first run, the intervals were based on exponentials. This second round yielded more refined ranges of best  $\gamma$  values at each site. For the evaluation, the simulated snow albedo was averaged over each day to retain an aggregated daily value similar to the observations.

Two criteria were imposed to select the days used to evaluate the simulated snow albedo. These criteria were all compiled for each site and year into a single mask which was then applied to all observation and simulation series for this site and year. 175 The first criterion ensured that the observed albedo value had been collected over a fully snow-covered ground and therefore corresponded to an actual snow albedo value: observed SD had to be higher than 20 cm. For high-latitude sites (above 60°N), this criterion was relaxed to 10 cm because there is less precipitation at these sites, so SD did not reach far over 20 cm certain years. Observed albedo also had to be higher than 0.5, which was a looser version of the criteria used in Lafaysse et al. (2017), where all albedo values below 0.6 were discarded. Finally, days where SD was null in the simulation with the lowest value of 180  $\gamma$  were discarded, keeping only days where there was in fact snow on the ground in all the simulations. This assumption is reasonable because simulated albedo is only computed from the surface layer physical properties so that the simulated albedo is not modified in case of thin simulated snowpacks.

The second criterion was to select periods when the impact of LAPs on snow albedo was significant. These periods physically correspond to days where the snow has aged, accumulating dry depositions of LAPs, and possibly gathering at the surface 185 LAPs from the lower layers if melting occurred (Tuzet et al., 2017). The snow age of the simulation with the highest  $\gamma$  had to be higher than a given value  $A_{lim}$  (in days). Several criteria guided the choice of  $A_{lim}$ :

- $A_{lim}$  had to be high enough for the impact of LAPs to be perceptible.
- $A_{lim}$  had to be high enough to discriminate between different values of  $\gamma$ , as its effect becomes increasingly noticeable with the increase of snow age (Fig. 1).
- 190 –  $A_{lim}$  had to be high enough to reduce the chance of having snow accumulated on the incoming solar radiation sensors and affecting the quality of the data (Lapo et al., 2015).
- $A_{lim}$  had to be low enough to retain a sufficient number of good quality days.





In fine, the value  $A_{lim} = 5$  days was retained as providing a compromise between these criteria.

Two metrics were used to quantify the skills of the snow albedo simulations: the bias and the root mean square error (RMSE).

195 The bias quantifies the tendency of a simulation to systematically overestimate or underestimate the snow albedo. Its definition is as follows to compare the observed (*obs*) and simulated (*sim*) times series for a given site, year, and value of  $\gamma$ :

$$Bias_{site,year,\gamma} = \mathbb{E}[sim_{site,year,\gamma}] - \mathbb{E}[obs_{site,year,\gamma}], \quad (3)$$

where  $\mathbb{E}[x]$  is the mean of  $x$ . The RMSE was also used to complement the bias. Its definition is as follows:

$$RMSE_{site,year,\gamma} = \sqrt{\mathbb{E}[(sim_{site,year,\gamma} - obs_{site,year,\gamma})^2]}. \quad (4)$$

200 For each site, and for each simulation with a given value of  $\gamma$ , the scores were computed for each year. A group of yearly scores for each value of  $\gamma$  was obtained at each site. The median over the years was computed to derive a single value of the score for each  $\gamma$  at the site.

Albedo measurements are subject to instrumental uncertainties in both intrinsic precision and snow accumulation on incoming radiation receptors (e.g. Lejeune et al. (2019); Lapo et al. (2015)). This means that there was a need to account for observa-  
205 tion uncertainty when computing the error metrics, as two slightly different scores for two values of  $\gamma$  could turn out to be statistically equivalent. In such a case, a range of best  $\gamma$  values would need to be considered instead of a single best  $\gamma$  value at each site. This range would contain the value with the best score at the given site, as well as all the other values which are statistically equivalent to it. Taking inspiration from Lafaysse et al. (2017), and focusing on the RMSE score, a statistical method described in Appendix B was used to find the range of best  $\gamma$  values at each site.

210

### 3.2 Climatology of light-absorbing particles deposition on snow

The goal of this section is to obtain a global climatology of daily LAP deposition rates on snow. Due to the important variability of the length of the snow season around the world, global snow data was needed to select only deposition over snow. A climatology of daily LAP deposition data was therefore combined with snow cover data in order to retain only LAP depositions  
215 on snow-covered ground.

#### 3.2.1 Data

The LAP deposition data was taken from Zhao et al. (2018). This dataset has been generated with the NOAA/OAR Geophysical Fluid Dynamics Laboratory Coupled Model version 4 (CM4; Zhao et al. (2018)) running for 37 years (1979-2015) on a C192 grid (192 grid points on each edge of the cube projected on Earth). GFDL-CM4 is driven by IPCC CMIP6 forcings, including  
220 the emissions of BC (fossil fuel burning, aircraft, shipping, and biomass burning). Dust emissions are calculated on-line with a cubic dependency on surface winds using a fixed dust source function, and a constant threshold of wind erosion (Ginoux et al., 2001). The aerosols are removed from the atmosphere by in- and below-cloud deposition, dry deposition at the surface, and gravitational settling for dust particles. The model results are interpolated from C192 to a Cartesian latitude-longitude grid



of 0.5 by 0.5 degrees resolution. The dataset consists of median BC and dust deposition rates (in  $\text{kgm}^{-2}\text{s}^{-1}$ ) over the period  
225 1979-2015, for each day of the year (see an example on Fig. S1 in the Supplementary Material). The deposition rates for each  
type of LAP correspond to the total deposition rates (including both wet and dry depositions). This dataset has been described  
and evaluated in Zhao et al. (2018). The deposition fluxes at the surface are given for an elevation corresponding to the ground  
elevation plus half of the thickness of the first atmospheric layer ( $\sim 15$  m). This elevation is referred to as the elevation of the  
GFDL grid in the rest of the paper.

230 For the daily snow cover data, the Global Multisensor Automated Snow and Ice Mapping System (GMASI) product (Romanov,  
2017) was used. Based on a combination of satellite observations in the visible, infrared, and microwave bands, it provides  
a daily partition of surface cover between water, bare ground, ice-covered water, and snow-covered ground, at a global scale  
from 1988 to the present (see an example on Fig. S2 in the Supplementary Material). This data is discretized along a  $0.04^\circ$  by  
 $0.04^\circ$  longitude-latitude grid, which is equivalent to about 4 kilometers by 4 kilometers at the equator. There are two versions  
235 of GMASI available: version 3 runs from 1988 to 2018, while version 4 is available from 2006 to 2023. In this study, the earlier  
version 3 was used up until 2005, after which it was replaced by the current version 4. The authors of the GMASI product  
made the choice to assume the permanent absence of snow between  $-25^\circ\text{N}$  and  $+25^\circ\text{N}$  longitudes, except in South America  
where this is only true East of  $-60^\circ\text{E}$  to capture the presence of snow in the Andes. This means that the climatology produced  
in this study has the same characteristics.

### 240 3.2.2 Methods

The climatology of dust and BC deposition over snow-covered ground was computed using the GFDL and GMASI datasets.  
Proportions of snow-covered ground for each day of the year were first computed over the GMASI grid by averaging the  
GMASI product over the period 1988-2015. A weighted average was then computed to obtain the mean daily LAP deposition  
over snow-covered ground: for each day the median deposition rates were first multiplied by the corresponding proportion of  
245 snow-covered ground, and these products were summed over the full year. Then this sum was divided by the sum over the year  
of all proportions of snow-covered ground. For a given location and for a given type of LAP (BC or mineral dust), the mean  
daily deposition rate on snow,  $D$  (in  $\text{kg m}^{-2} \text{d}^{-1}$ ), is therefore written as:

$$D = \frac{\sum_{d=1}^{365} \delta_d p_d}{\sum_{d=1}^{365} p_d} \times 60 \times 60 \times 24, \quad (5)$$

where  $\delta_d$  is the median deposition for that type of LAP over 1981-2015 on the day of the year  $d$  (in  $\text{kg m}^{-2} \text{s}^{-1}$ ), and  $p_d$   
250 is the proportion of snow-covered ground over 1988-2015 for the day of the year  $d$ . February 29 was ignored for leap years,  
as if all years over the period had 365 days. Two other limitations can be identified in this calculation: (i) GMASI begins in  
1988, while the GFDL dataset accounts for years between 1981 and 1987 and (ii) the GFDL dataset gives the median of the  
deposition rates and not the mean values.

Because of their different optical properties, dust and BC have different radiative impacts on the snowpack for the same  
255 deposited mass (Clarke et al., 2004). To present a single climatological value of LAP deposition, it was therefore convenient to



express dust deposition in terms of equivalent BC deposition. This corresponds to the BC deposition that would have the same integrated radiative impact as the considered dust deposition, over the studied spectral bands. Tuzet et al. (2019) examined expressions for an equivalent BC concentration in snow  $c_{eq,BC}$  integrated over the 0.35 – 0.9  $\mu\text{m}$  spectral band. An average of the slope in Fig. 1 of Tuzet et al. (2019) yielded the following expression:

$$260 \quad c_{eq,BC} = c_{BC} + 0.0033c_{dust}, \quad (6)$$

where  $c_{BC}$  is the BC concentration in snow and  $c_{dust}$  is the dust concentration, all concentrations being expressed in  $\text{ng g}^{-1}$ . This relationship can also be used on deposition rates, by linearity.

From the global climatology, values of dust, BC, and equivalent BC deposition were extracted for each of the experimental sites described in Section 3.1.1. The effect of elevation on the climatological BC and dust depositions was also investigated in the vicinity of the five mountain sites: Col de Porte, Kühltai, Senator Beck, Swamp Angel, and Weissfluhjoch. The elevation of the GFDL climatology grid cell closest to each site was first compared to the real elevation at the site. The regional gradient of LAP deposition with elevation was then determined around each site. It was computed using a linear regression of the total LAP deposition rate as a function of the elevation using the grid cells surrounding each site in a 25-cell by 25-cell square (which was approximately equivalent to a 5-cell by 5-cell square of the GFDL grid). An elevation-corrected total LAP deposition rate at each site was then computed using the linear regression and the actual elevation of the site and the corresponding elevation of the GFDL dataset. The addition of half of the first atmospheric layer of the GFDL dataset was neglected here, as it was very small compared to the sites' ground elevations.


### 3.3 Cross-analysis

The final step of the analysis consisted in identifying a relationship between the range of statically equivalent best  $\gamma$  values and the LAP deposition rate at each site. A specific linear regression method, described in Appendix C, was proposed to find the optimal relationship between the statically-equivalent range of  $\gamma$  and the individual LAP deposition rate.

To assess the added value of this LAP-dependent parameterization of gamma in large scale simulations compared to the default constant setting, a final evaluation was carried out to quantify the impact of using the new values of  $\gamma$  on simulated snow albedo and snow depth at the ten experimental sites. The RMSE was computed to compare simulated and observed albedo and  $SD$  at each site and for each year, for two values of  $\gamma$ : the default value of 60 days, and the new value resulting from the regression proposed in this study. The median of the yearly scores was then considered, as in Section 3.2.2. For each year, the RMSE was computed when the ground was fully covered by snow as described in Section 3.2.2. The filter on the snow age was not applied. The relative change in RMSE resulting the use of the new value of  $\gamma$  was finally computed at each site as:

$$I_v = \frac{\widehat{RMSE}_{v,60} - \widehat{RMSE}_{v,\gamma_R}}{\widehat{RMSE}_{v,60}} \quad (7)$$

285 where  $v$  is the considered variable (snow albedo or  $SD$ ) and  $\gamma_R$  is the new value obtained from this study.




**Figure 3.** Time series of observed (black dots) and simulated (colored lines, one for each value of  $\gamma$ ) values for  $SD$  (in meters, top), snow age of the surface layer (in days, middle), and albedo (no unit, bottom) at the Col de Porte site over the 1998 – 1999 snow year. The grey vertical lines mark the days which have been selected for the evaluation. Note that observed data are not available for snow age.

## 4 Results

### 4.1 Snow albedo simulations

Figure 3 shows an example of the seasonal evolution of observed and simulated snow depth and snow albedo for the Col de Porte experimental site during winter 1998-1999. The age of surface snow is also shown for the simulations with different values of  $\gamma$ . As expected, simulations with low values of  $\gamma$  are associated with a faster decrease in snow albedo than simulations with high values of  $\gamma$  during all the periods following snowfall that refreshes the snow surface. Snow depth starts diverging between the different simulations in the second half of the snow season once the peak of snow depth has been reached. Snow depth in simulations with low values of  $\gamma$  decreases faster due to larger melt rates associated with lower snow albedo and increased snow compaction due to the presence of liquid water in the snowpack. The snow cover totally vanished 13 days later in the simulation using  $\gamma = 900$  days compared to the simulation with  $\gamma = 5$  days. The vertical grey lines on Fig. 3 show the days selected to compute the albedo evaluation using the criteria listed in Sect. 3.1.2. As designed in Sect. 3.1.2, they cover periods of decreasing albedo, far enough from snowfall events (increases in  $SD$ ). The average number of selected days per snow season was examined at each site and can be found in Supplementary Material (Table S1). On average over all sites and snow seasons, 41 days were retained per season, and all sites but two are grouped around this value. The two outliers are Sapporo with 12 days selected due to its particularly short snow season, and Weissfluhjoch with 73 days.



**Figure 4.** Box plot showing the distribution of the yearly scores at the Col de Porte site from 1994 to 2014, for the first round of evaluation (a) over the [5, 900] days range and (b) for the second round of evaluation zoomed in the [7, 55] days range. The median over the years for each  $\gamma$  is shown in white. The box plots show the interquartile values, and outliers are plotted as circles. The grey rectangle along the  $x$  axis shows the best  $\gamma$  range derived from the evaluation process. Note that the vertical axes are not identical for plots (a) and (b).


Error metrics (bias and RMSE) for snow albedo were then computed for each year at each site. Figure 4 shows an example of the distribution of these error metrics for different values of  $\gamma$  at the Col de Porte experimental site. Each boxplot summarizes the distribution of the score for all the available years for a given value of  $\gamma$ . Results for the two rounds of evaluation considering two ranges of values for  $\gamma$  (Sect. 3.1.2) are shown on Fig. 4. At Col de Porte, the optimal values of  $\gamma$  ranged from 13 to 20 days: these values were associated with the lowest values of RMSE and had snow albedo biases close to zero. The box plots for all other sites for the second round of evaluation can be seen in Supplementary Material (Fig. S3 to S11).

The resulting optimal ranges of  $\gamma$  obtained at each site were compiled in Fig. 5 (logarithmic scale on Fig. 5a and linear scale on Fig. 5b). The optimal values of  $\gamma$  strongly depart from the default value of 60 days used in Crocus. The highest values of  $\gamma$ , in the 400 – 800 days range, were found at the Canadian Arctic sites of Bylot and Trail Valley Creek. Optimal values ranging around 100 – 200 days were found at Sodankylä, a site in northern Scandinavia, Umiujaq, a low Arctic site in Canada and at the high-elevation alpine site of Weissfluhjoch. Values close to the default 60 days values were optimal at Senator Beck and Swamp Angel, the two high elevation Colorado sites. Finally, the lowest values of  $\gamma$  appeared at Col de Porte, Sapporo, and Kühtai.

## 4.2 Climatology of light-absorbing particles deposition on snow

### 4.2.1 Global climatology


From the GMASI product, a probability of snow cover at each location for each day of the year was computed, by dividing the number of years where there was snow cover at the location by the number of years in the dataset. Summing these probabilities



**Figure 5.** Optimal range of  $\gamma$  at each site, in (a) logarithmic and (b) linear scale.

over all days of the year yielded the average number of snow cover days per year at each location over the 1988-2016 period. Figure 6 shows the spatial variability at a global scale with a strong latitudinal trend in the Northern Hemisphere. Mountainous areas (mostly the Himalayas, the European Alps, and the Canadian and US Rockies) appear very clearly as they are more often snow-covered than the lower-elevation surrounding regions. As expected, Antarctica and Greenland are characterized by the presence of snow cover that extends all year long. The no-snow mask applied in the Southern Hemisphere in the GMASI product (see Sect. 3.1.2) can be clearly seen in South America (east of the Andes), in South Africa and in Australia.

Global maps of mean annual dust and BC deposition rates on snow are presented in Fig. 7. Some spatial patterns are visible: BC deposition is higher in South-East Asia and Europe (Fig. 7a) whereas low BC deposition rates are found in the Canadian Arctic and in Eastern Siberia. Mineral dust is mainly deposited in regions close to the main source regions such as the Middle East, the Andes (close to the Altiplano and Patagonian deserts), and to a lesser extent the West of the United States (Fig. 7b). The lowest deposition rates of BC and dust on snow are found in Antarctica. Mineral dust deposition rates are overall higher than BC deposition rates, but the impact of mineral dust on albedo is much lower than the impact of BC (about three orders of magnitude according to Eq. 6; Clarke et al. (2004)). For this reason, the total LAP deposition rates on snow (Fig. 7c) have similar orders of magnitude to BC deposition rates, with a spatial variability that reflects the variability found in the climatology of both dust and BC deposition on snow.




**Figure 6.** Global map of average snow cover days per year, derived from the GMA SI dataset.

#### 4.2.2 Climatology at the reference sites

The local gradients of total LAP deposition with elevation were computed around each mountainous site, as described in Sect. 3.2.2, and are shown on Fig. 8. Senator Beck and Swamp Angel are represented on the same figure (Fig. 8c) since they fell within the same grid cell. The comparison of the actual elevation at each site with the elevation of the corresponding grid cell in the GFDL dataset, as well as the value of the gradient and the regression coefficient of each linear regression are also shown in Fig. 8. These altitudinal gradients revealed contrasted evolutions of LAP deposition rates with elevation around the different sites. Indeed, the two sites located in the Rockies display an increase in LAP deposition as a function of elevation whereas a decrease with elevation is found for the three sites located in the European Alps with similar values of the altitudinal gradient.

The LAP climatologies were then extracted from Fig. 7c at the ten experimental sites considered in the study and were adjusted using the altitudinal gradients for the five mountainous sites. Figure 9 presents the initial and adjusted values of LAP deposition rates for all sites. The sites displayed a wide range of LAP deposition rates, with over two orders of magnitude of difference between the least contaminated site (Bylot in Northern Canada) and the most contaminated (Sapporo in Japan). The Canadian Arctic sites received the least deposition, while the sites in the European Alps had the highest deposition rates after





**Figure 7.** Global maps of annual mean (a) BC, (b) dust, and (c) total LAP (BC + dust in equivalent BC) deposition rates over snow. The logarithmic color scale highlights the orders of magnitude of difference between BC (a) and dust (b) deposition rates.


Sapporo. The two Colorado sites as well as Sodankylä received mid-range deposition rates. The largest altitudinal correction was performed at the Weissfluhjoch site due to the large difference between the actual elevation of the site and the corresponding GFDL elevation. Only minor corrections were applied to the other mountain sites. At Swamp Angel and Senator Beck, the correction was positive whereas it was negative at the other sites, in agreement with the altitudinal gradients shown on Fig. 8.

### 4.3 Cross-analysis

The linear regression method presented in Section 3.3 was applied to find a relationship between the range of statistically equivalent values of  $\gamma$  and the climatological LAP deposition at each site. It is presented in Fig. 10, before and after the correction for elevation was applied to the mountainous sites. The regression with the altitudinal correction had the following characteristics:

$$\log(\gamma) = -0.641 \log(D_{LAP}) - 2.99, \quad (8)$$

$$R^2 = 0.642, \quad (9)$$




**Figure 8.** Local gradient of the total LAP climatological deposition rate with elevation and associated linear regression at (a) Col de Porte, (b) Kühtai, (c) Swamp Angel and Senator Beck, and (d) Weissfluhjoch. Note that the axes are not identical between the plots. The gradients have the units  $\text{kg m}^{-2} \text{d}^{-1}$ .

where  $D_{LAP}$  is the climatological deposition rate of LAPs over snow expressed in  $\text{kg m}^{-2} \text{d}^{-1}$  of equivalent BC. The regression without altitudinal correction was very similar with a slightly lower regression coefficient of  $R^2 = 0.613$ .

360 The two linear regressions are very similar (Fig. 10), but the correction for elevation provided a slight improvement in the regression coefficient, so that this regression is used as the reference regression in the rest of this study. Updated values of  $\gamma$  (referred to as  $\gamma_R$ ) were obtained at each site from Eq. 9.


The benefit of using  $\gamma_R$  compared to the current default in Crocus (60 days) was then assessed in terms of simulated snow albedo and snow depth at the 10 experimental sites. Results are shown on Fig. 11. Improvements in snow albedo simulations  
 365 (i.e., decrease in RMSE) are found at 7 sites (Fig. 11a). The average relative decrease in RMSE over the 10 sites was 10% for snow albedo. The albedo simulations were improved most at the three Canadian Arctic sites (decrease in RMSE larger than 25%), because the default value of 60 days was very small compared to the value of  $\gamma_R$  at these sites (Fig. 10). At Col de Porte, Kühtai and Sapporo, the opposite effect was observed: the default value was too high compared to  $\gamma_R$  and the ideal range of  $\gamma$  at these sites. In Sodankylä,  $\gamma_R$  fell very close to the ideal range of  $\gamma$  at this site and provided a moderate improvement  
 370 in albedo simulations (decrease in RMSE by 6 %). Albedo simulations at the two Colorado sites and at Weissfluhjoch were degraded when using  $\gamma_R$ , which is further away from their optimal ranges of  $\gamma$  than the 60 days default (Fig. 10).



**Figure 9.** Graph representing climatological LAP deposition rates at all sites, before and after the correction of elevation effects. Arrows indicate these corrections, at mountainous sites.

The snow depth (SD) simulations show different improvement tendencies compared to snow albedo (Fig. 11b) with an average relative decrease in RMSE over the 10 sites of 3% (10 % for albedo). SD simulations were improved at 6 out of the 10 experimental sites. Among these 6 sites, 5 of them showed simultaneous improvements in albedo and SD simulations. SD simulations at Col de Porte were degraded when using  $\gamma_R$  (increase in RMSE by 10%) despite strong improvements in albedo simulations (decrease in RMSE by 13%). A similar behavior was observed for the SD simulations at Sodankylä. Using  $\gamma_R$  degraded SD simulations at Weissfluhjoch (increase in RMSE by 12%) in agreement with the degradation found for snow albedo (increase in RMSE by 13%). The largest RMSE for SD simulations (using both configurations of the model) was found at Senator Beck. This site is exposed to wind-induced snow redistribution which cannot be reproduced by the model in point-scale mode. The same issue affected a large number of models used in the ESM SnowMIP exercise (Menard et al., 2021).


Finally, using the relation obtained between the climatology of LAP deposition and  $\gamma$  (Eq. 9), a map of optimal  $\gamma$  values was derived from the global climatology (Fig. 12). The horizontal resolution of this global map is the same as the global climatology of LAP deposition on snow (0.04° by 0.04°). Threshold values for  $\gamma$  were set at 5 and 900 days. The map shows that the lowest values of  $\gamma$  (in the range 5 to 50 days) should be used in the regions that receive the largest deposition of LAP, notably South-East Asia and Europe (Fig. 7c). The highest values of  $\gamma$  (above 500 days) should be used in regions which receive low LAP depositions, namely Antarctica, Greenland and the high Canadian Arctic.



**Figure 10.** Graph representing the optimal ranges of  $\gamma$  as a function of total LAP deposition rates in equivalent BC, in logarithmic scales, and the corresponding linear regressions with and without correction for elevation at the mountain sites (the correction is represented by the arrows). The points which have been represented with circles in each range are the ones closest to the final regression (Eq. 9), which are used to compute the regression coefficient (see method in Appendix C).

## 5 Discussion


Multiyear simulations with the Crocus snowpack scheme have been carried out to determine the ranges of values of the snow aging coefficient  $\gamma$  giving the best snow albedo simulations at ten sites covering different climates (Fig. 2 and Table 2). A large variability of the optimal range of values for  $\gamma$  was found across the ten sites (Fig. 5). These results confirm that a single value of  $\gamma$  cannot be used for all climates around the world, and this parameter needs to be tuned as proposed by Brun et al. (2011). In particular, the values in the 500- to 900-day range found in this study for the two sites located in the Canadian Arctic (Trail Valley Creek and Bylot Island) are consistent with the values used for polar snow by Brun et al. (2011) and Woolley et al. (2024). On the other hand, Fig. 5 reports optimal  $\gamma$  values in the range 13-20 days at the Col de Porte experimental site. These values differ from the default value of 60 days that was optimized by Brun et al. (1992) for this site. However, Brun et al. (1992) did not use albedo measurements to tune the value of  $\gamma$ . Instead, they relied on measurements of surface temperature to show that  $\gamma=60$  days provided good simulations of the snow surface energy balance. Lafaysse et al. (2017) have shown that the energy balance simulated by Crocus is associated with additional sources of uncertainties, in particular due to the formulation



**Figure 11.** For each site, relative improvement of simulation skills for (a) albedo and (b) snow depth, from this study compared to the default 60 days value, and absolute improvements for (c) albedo and (d) snow depth, where the black dots are the default 60 days value and the bars are the results from this study.

of turbulent fluxes (Martin and Lejeune, 1998). The direct optimization of  $\gamma$  on snow albedo proposed in this study reduces the impact of these additional sources of uncertainties on the estimation of the optimal values of this parameter.

The optimal values of  $\gamma$  shown in Fig. 5 have then been combined with climatological values of LAP deposition on snow at the ten experimental sites to determine a regression between these two variables (Fig. 10). This logarithmic linear regression confirms the dependency of  $\gamma$  on the LAP deposition rate which influences snow albedo in the visible range. 64% of the variation in  $\gamma$  is explained by the regression model. The largest discrepancies between the regression and the optimal values of  $\gamma$  are found for the Weissfluhjoch and Kühtai sites, located in the European Alps. These two sites are geographically close to each other (approximately 150 km apart) with optimal values of  $\gamma$  in the range 134-245 days at the Weissfluhjoch site and in the range 7-10 days at Kühtai. According to the GFDL climate model, these two sites present similar climatological values of LAP deposition on snow. The correction for elevation proposed in this study only slightly differentiated these climatological values. The discrepancy between the two sites may result from the inability of the global climatology of LAP deposition on snow to capture fine-scale features of LAP deposition in mountainous terrain and the presence of local LAP sources. The Weissfluhjoch site is located well above an alpine valley whereas the Kühtai snow site is located in a valley. The GFDL climate model at 50-km grid spacing (Zhao et al., 2018) cannot accurately reproduce orographic precipitations, which are essential for wet deposition of LAPs, nor channelling of dust flow through valleys (Baladima et al., 2022).



**Figure 12.** Global map showing the optimal value of  $\gamma$  derived from Eq. 9.

The climatology of LAP deposition on snow would therefore benefit from inputs of chemistry transport models at 10-km  
415 resolution for global applications (e.g. Bessagnet et al., 2017)) and at even higher resolutions for regional applications (e.g.  
Baladima et al., 2022)). Such products would allow to refine the altitudinal adjustments applied to the local climatology to  
correct for the difference of elevation between the climatology and the actual elevation of the sites. The results obtained with  
the current climatology suggest that the local altitudinal gradients of LAP climatological deposition rate are characteristic  
of each mountain range (Fig. 8). The climatological LAP deposition rate decreases with elevation in the European Alps  
420 (around Weissfluhjoch, Col de Porte, and Kuthai) whereas it increases with elevation in the Colorado Rocky Mountains (USA,  
around Senator Beck and Swamp Angel). The default snow albedo parameterization in Crocus is consistent with the altitudinal  
gradient found in the European Alps since it uses a pressure term (Eq. 2) to increase the value of  $\gamma$  with elevation, indirectly  
representing a decrease in LAP deposition with elevation. However, this term is not suitable for the Colorado Rocky Mountains  
nor potentially for other mountain ranges. This result justifies why the pressure term was not considered in this study and  
425 highlights the need to carefully revise this term when Crocus is applied in different mountain ranges across the world.

The regression between  $\gamma$  and the LAP climatological deposition rates relied on a total of 10 points corresponding to the  
ten experimental sites considered in this study (Fig. 10). The size of the sample limits the robustness of the regression. This  
limitation is explained by the fact that the LAP deposition outputs of the GFDL CM4 model only contain the median deposition



430 rates for each day of the year, based on the 1979-2015 period. As a consequence, the ideal values of gamma at each site were determined by aggregating all the years of data available at each site to obtain a single range of gamma for each site, hence the ten points considered in the regression. An alternative would be to consider a dataset such as the MERRA2 atmospheric reanalysis (Gelaro et al., 2017), which provides global daily estimates of wet and dry LAP depositions rates at 50-km resolution since 1980. Combined with GMASI, such a product would allow to derive average LAP deposition rates on snow for each year and each site. These could be compared with the corresponding yearly optimal values of  $\gamma$ . This would increase the size of  
435 the sample considered in the regression ten-fold. Adding new sites in the analysis is another option to increase the size of this sample, but it requires reference sites with quality-controlled meteorology and dedicated snow observations such as those available in the ESM-SnowMIP dataset (Ménard et al., 2019).

The values of  $\gamma$  determined with the logarithmic linear regression improved snow albedo simulations with Crocus at seven out of the ten sites (Fig. 11a and c), with the largest improvements found for the Arctic sites (decrease in RMSE larger  
440 than 25%). These results confirm the robustness of the approach at the sites where the regression was developed. Further evaluations using independent sites are required to confirm the transferability of the approach. Simulations of snow depth with Crocus including the LAP-dependent  $\gamma$  were also slightly improved compared to the default version of Crocus (Fig. 11b and d; average decrease in RMSE of 3% at the ten sites). However, certain sites such as Col de Porte and Sodankylä showed degradations in the simulations of snow depth despite improvements in the simulation of snow albedo. This behavior may  
445 be explained by the fact that the quality of snow depth simulations by the model is influenced by other sources of errors in the different model parameterizations (Lafaysse et al., 2017; Günther et al., 2019). Ensemble snowpack simulations (Lafaysse et al., 2017) could be used to quantify the uncertainties associated with these different parameterizations and identify the role of the adjusted values of  $\gamma$ .

The methodology developed in this study relies on climatological values of LAP deposition on snow, so that it cannot represent the impact of individual LAP deposition events which can nonetheless strongly influence the evolution of snow albedo  
450 (e.g. Di Mauro et al., 2015; Dumont et al., 2020). Their representation can only be achieved by coupling a snowpack model with a more numerically expensive snow radiative transfer model and using LAP deposition fluxes as additional forcing (Flanner et al., 2007; Tuzet et al., 2017; Réveillet et al., 2022). Instead, the methodology proposed in this study aims at capturing the impact of the large spatial variability of LAP deposition on the simulation of snow albedo. It has been applied to the default  
455 albedo parameterization of the Crocus snow scheme and presents a strong potential to improve continental-scale snow simulations (Brun et al., 2013; Mortimer et al., 2020), which serve as a basis for climate trend-analysis (Mudryk et al., 2024). The same methodology can be applied to optimize parameters that indirectly represent the impact of LAPs on snow albedo in the visible range in other snowpack models explicitly including the effect of optical grain size on albedo (Dickinson, 1993; Best et al., 2011; Decharme et al., 2016). The time constants in more simple time-dependent snow albedo parameterizations (e.g.,  
460 Verseghy, 1991; Douville et al., 1995; Pedersen and Winther, 2005) could also be optimized using the global climatology proposed in this study. However, this optimization would require a careful analysis to make the distinction between the combined effects of several snow aging processes contributing to the decrease of snow albedo with time, such as LAP depositions and the increase in optical grain size due to snow metamorphism.





## 6 Conclusions

465 The goal of this study was to develop a methodology of intermediate complexity to improve large scale simulations of snow  
albedo in snowpack schemes by taking into account the spatial variability of LAP deposition (BC and dust). Toward this goal,  
a global climatology of LAP deposition over snow has been built by combining a climatology of daily LAP deposition from  
a global climate model with a global remotely-sensed snow cover dataset to retain only deposition over snow, providing an  
estimation of the average BC and dust deposition rates over snow for the period 1981-2015. The two types of LAPs were then  
470 expressed in terms of equivalent black carbon and summed to obtain the climatology of total LAP deposition. This climatology  
is available for scientific use (Gaillard et al., 2024a).

The default albedo parameterization in the detailed snowpack scheme Crocus has then been improved using this climatology  
by optimizing a parameter that controls the snow albedo evolution in the visible range (known as the aging coefficient,  $\gamma$ ). This  
coefficient implicitly represents the darkening of the snow with time due to LAP deposition with low (high) values indicative  
475 of darkened (clean) snow. Multi-year snowpack simulations were carried out with Crocus at ten reference sites covering a  
large variety of climates. The range of optimal values of  $\gamma$  yielding the most skilled albedo simulations was found at each site,  
accounting for uncertainties in the observation of snow albedo. This analysis revealed a wide variety and a good dispersion of  
the ranges of  $\gamma$  at the ten chosen sites: ideal  $\gamma$  values went from 7 days at Kühtai, in the Austrian Alps, to 898 days at Bylot  
and Trail Valley Creek, in the Canadian Arctic.

480 A logarithmic linear regression was then applied between the optimal ranges of  $\gamma$  and the LAP deposition rates at the ten  
sites extracted from the climatology. These LAP deposition rates ranged over two orders of magnitude between the Canadian  
Arctic sites and Sapporo in Japan. The logarithmic linear relationship was finally combined with the global climatology to  
obtain an LAP-informed and spatially variable  $\gamma$  parameter for the Crocus albedo parameterization. The revised parameter  
improved, on average, snow albedo simulations by 10% with the largest improvements found in the Arctic (more than 25%).  
485 The global dataset with the optimal values of  $\gamma$  is available for the users of Crocus (Gaillard et al., 2024b).

Future work will test the impact of the spatially optimized values of  $\gamma$  in snowpack simulations with Crocus over large  
domains such as Canada. The methodology detailed in this paper can be applied to optimize parameters controlling the albedo  
evolution in the visible range in other snowpack schemes (e.g., Dickinson, 1993; Best et al., 2011; Decharme et al., 2016)  
without requiring coupling with a more expensive snow radiative transfer model.

490 *Code and data availability.* The global climatology of LAP deposition on snow is available at <https://doi.org/10.5281/zenodo.11554783>  
(Gaillard et al., 2024a). The LAP-informed  $\gamma$  values are available at <https://doi.org/10.5281/zenodo.11554926> (Gaillard et al., 2024b). The  
code of Crocus within the land surface scheme SVS2 used in this study is available at [https://github.com/VVionnet/MESH\\_SVS/tree/master](https://github.com/VVionnet/MESH_SVS/tree/master).  
Integration of these developments in the version of Crocus available in the SURFEX modelling platform will be done after the publication  
of this study.



**Table A1.** Physical options of the multiphysics version of Crocus used in this study. The names of the options refer to the Crocus namelist (Lafaysse et al., 2017)

Physical process	Name of the option	Reference
Snowfall density	V21	Vionnet et al. (2012)
Metamorphism	B21	Baron (2023)
Turbulent fluxes	RI2	Lafaysse et al. (2017)
Thermal conductivity	Y81	Yen (1981)
Liquid water holding	B92	Brun et al. (1992)
Compaction	B92	Brun et al. (1992)
Snowdrift	VI13	Vionnet et al. (2013)

## 495 Appendix A: Parameterizations used in Crocus

### Appendix B: Method to select statistically-equivalent intervals of $\gamma$

For a given site, the score estimator for each value of  $\gamma$  is:

$$\widehat{RMSE}_{site,\gamma} = \mathcal{M}_{years}(RMSE_{site,year,\gamma}), \quad (B1)$$

where  $\mathcal{M}_{years}$  is the median over all the years available at this site and  $RMSE_{site,year,\gamma}$  is the yearly RMSE score as expressed in Eq. 4. This score is uncertain because of the observational error. It is modelled as proposed in Lafaysse et al. (2017) using a normal distribution as a first-order approximation:

$$\mathbb{N}_{site,\gamma} = \mathcal{N}(\widehat{RMSE}_{site,\gamma}, \sigma_{RMSE}), \quad (B2)$$

where the variance  $\sigma_{RMSE}$  is the estimated observational uncertainty corresponding to the RMSE score. The a priori value for  $\sigma_{RMSE}$  suggested in Lafaysse et al. (2017) is 0.069. This value was estimated at the Col de Porte site, whose incoming radiation receptors are equipped with automatic wipers to avoid the inexactitudes from snowfall accumulation on the sensors. Most of the other sites in this study are not similarly equipped, though some have frequent manual wiping of the sensors. Therefore, this value of  $\sigma_{RMSE}$  is potentially underestimated for most of the sites. In this study, it was nonetheless retained as a first order estimate of the observational uncertainty corresponding to the RMSE score. Numerically, 1001 values of  $\mathbb{N}_{site,\gamma}$  were randomly generated to represent the distribution.

The statistically-equivalent range of  $\gamma$  was finally computed by considering  $\tilde{\gamma}$  the value of  $\gamma$  which gave the best score estimator  $\widehat{RMSE}_{site,\gamma}$ . An independent samples t-test was applied between  $\mathbb{N}_{site,\tilde{\gamma}}$  and each of the other distributions  $\mathbb{N}_{site,\gamma}$ , to assess if the difference between them was statistically significant or not. For this test, a confidence interval of 90% was used, as suggested in Lafaysse et al. (2017). All values of  $\gamma$  which were not declared statistically different from  $\tilde{\gamma}$  by the t-test were added to the range of best  $\gamma$  values for the considered site. A range of statistically equivalent ideal  $\gamma$  values was obtained for each site and considered in the rest of the study.



### Appendix C: Method for the linear regression

This method slightly differs from classic linear regression methods that do not consider a uniform range of statically equivalent values for the target variable. The cost function used was inspired from the mean squared error (MSE), defined as follows for an independent variable  $x$  and a dependent variable  $y$ :

$$520 \quad MSE = \frac{1}{N} \sum_{i=1}^N (ax_i + b - y_i)^2, \quad (C1)$$

where the scatter plot from which to derive the regression is  $(x_i, y_i)_{i \in [1, N]}$ , and  $a$  and  $b$  are the regression coefficients such that  $y = ax + b$ . This MSE was therefore the sum of squared distances between the predicted value of  $y$  and the actual value of  $y$ . To account for the ranges of  $y$  (which in our study represents  $\gamma$ ), the actual value of  $y$  was replaced with the range of values  $Y$ :

$$525 \quad MSE = \frac{1}{N} \sum_{i=1}^N \Delta_i^2, \quad (C2)$$

$$\text{with } \Delta_i = \begin{cases} 0 & \text{if } y_{i, \min} \leq ax_i + b \leq y_{i, \max} \\ ax_i + b - y_{i, \min} & \text{if } y_{i, \min} \geq ax_i + b \\ ax_i + b - y_{i, \max} & \text{if } y_{i, \max} \leq ax_i + b \end{cases} \quad (C3)$$

where  $y_{i, \min}$  and  $y_{i, \max}$  are the minimal and maximal values within the range  $Y_i$ . A gradient descent method was used to optimize  $a$  and  $b$  following this cost function, using a learning rate of 0.01, 1000 iterations, and initial  $a$  and  $b$  values derived from an automatic standard linear regression for a random value within each range of  $\gamma$ .

530 Once the optimal  $a$  and  $b$  were found, the points closest to the regression within each range were found. These are the points which were taken into account during the final iteration of the gradient descent. If the regression had been carried out over these points instead of the ranges  $Y$ , the resulting  $a$  and  $b$  would have been the same. For this reason, these points were used to compute the regression coefficient  $R^2$ :

$$R^2 = 1 - \frac{\sum_{i=1}^N (ax_i + b - y_i)^2}{\sum_{i=1}^N (ax_i + b - \bar{y}_i)^2} \quad (C4)$$

$$535 \quad \text{with } y_i = \begin{cases} ax_i + b & \text{if } y_{i, \min} \leq ax_i + b \leq y_{i, \max} \\ ax_i + b - y_{i, \min} & \text{if } y_{i, \min} \geq ax_i + b \\ ax_i + b - y_{i, \max} & \text{if } y_{i, \max} \leq ax_i + b \end{cases} \quad (C5)$$

where  $\bar{y}_i$  is the average of  $(y_i)_{i \in [1, N]}$ . This coefficient was useful to evaluate the quality of the linear regression.

*Author contributions.* MG, VV, ML and MD were responsible for designing the methodology of the study. MG carried out the numerical simulations and the analysis. VV supervised the project. MD provided expertise on LAP deposition. ML advised on the statistical method



540 to account for observation uncertainties in the optimal  $\gamma$  values, and provided the meteorological forcing file for the Sodankylä site. PG provided and advised on the GFDL dataset. MG and VV drafted the manuscript and all authors participated in reviewing and editing the paper.

*Competing interests.* Some authors are members of the editorial board of The Cryosphere.

545 *Acknowledgements.* The authors would like to thank Antoine Corcket, Georgina Woolley, and David Günther, who have provided some of the observation and meteorological forcing data for the reference sites of this study. Many thanks also to Gonzalo Leonardini for his help on formatting the observation data at some of the reference meteorological sites. Marie Dumont has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (IVORI; grant no. 949516).



## References

- Armstrong, R. and Brun, E.: Snow and climate: physical processes, surface energy exchange and modelling, *Journal of Glaciology*, 55, 384–384, <https://doi.org/10.3189/002214309788608741>, 2009.
- 550 Baladima, F., Thomas, J. L., Voisin, D., Dumont, M., Junquas, C., Kumar, R., Lavaysse, C., Marelle, L., Parrington, M., and Flemming, J.: Modeling an extreme dust deposition event to the French alpine seasonal snowpack in April 2018: Meteorological context and predictions of dust deposition, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035 745, 2022.
- Baron, M.: Modelling soil thermal regimes in high altitue open environnement of the french Alps : effect of wind-induced snow transport and vegetation (in French), Ph.D. thesis, Universite Grenoble Alpes, Grenoble, France, 2023.
- 555 Bessagnet, B., Menut, L., Colette, A., Couvidat, F., Dan, M., Mailler, S., Létinois, L., Pont, V., and Rouïl, L.: An evaluation of the CHIMERE Chemistry Transport Model to simulate dust outbreaks across the Northern Hemisphere in March 2014, *Atmosphere*, 8, 251, 2017.
- Best, M. J., Pryor, M., Clark, D., Rooney, G. G., Essery, R., Ménard, C., Edwards, J., Hendry, M., Porson, A., Gedney, N., et al.: The Joint UK Land Environment Simulator (JULES), model description–Part 1: energy and water fluxes, *Geoscientific Model Development*, 4, 677–699, 2011.
- 560 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., et al.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of geophysical research: Atmospheres*, 118, 5380–5552, 2013.
- Boone, A., Masson, V., Meyers, T., and Noilhan, J.: The influence of the inclusion of soil freezing on simulations by a soil–vegetation–atmosphere transfer scheme, *Journal of Applied Meteorology and Climatology*, 39, 1544–1569, 2000.
- 565 Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *Journal of Glaciology*, 38, 13–22, <https://doi.org/10.3189/S0022143000009552>, 1992.
- Brun, E., Six, D., Picard, G., Vionnet, V., Arnaud, L., Bazile, E., Boone, A., Bouchard, A., Genthon, C., Guidard, V., et al.: Snow/atmosphere coupled simulation at Dome C, Antarctica, *Journal of Glaciology*, 57, 721–736, 2011.
- Brun, E., Vionnet, V., Boone, A., Decharme, B., Peings, Y., Valette, R., Karbou, F., and Morin, S.: Simulation of Northern Eurasian Local  
570 Snow Depth, Mass, and Density Using a Detailed Snowpack Model and Meteorological Reanalyses, *Journal of Hydrometeorology*, 14, 203 – 219, <https://doi.org/10.1175/JHM-D-12-012.1>, 2013.
- Carmagnola, C., Morin, S., Lafaysse, M., Domine, F., Lesaffre, B., Lejeune, Y., Picard, G., and Arnaud, L.: Implementation and evaluation of prognostic representations of the optical diameter of snow in the SURFEX/ISBA-Crocus detailed snowpack model, *The Cryosphere*, 8, 417–437, 2014.
- 575 Clarke, A., Shinozuka, Y., Kapustin, V., Howell, S., Huebert, B., Doherty, S., Anderson, T., Covert, D., Anderson, J., Hua, X., Ii, K., McNaughton, C., Carmichael, G., and Weber, R.: Size distributions and mixtures of dust and black carbon aerosol in Asian outflow: Physiochemistry and optical properties J, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2003JD004378>, 2004.
- Cook, J. M., Hodson, A. J., Taggart, A. J., Mernild, S. H., and Tranter, M.: A predictive model for the spectral “bioalbedo” of snow, *Journal of Geophysical Research: Earth Surface*, 122, 434–454, <https://doi.org/10.1002/2016JF003932>, 2017.
- 580 Corcket, A.: Local-scale evaluation of subarctic snowpack simulations by Crocus at Sodankylä Finland, with Météo France and CNRS, 2023.
- Decharme, B., Brun, E., Boone, A., Delire, C., Le Moigne, P., and Morin, S.: Impacts of snow and organic soils parameterization on northern Eurasian soil temperature profiles simulated by the ISBA land surface model, *The Cryosphere*, 10, 853–877, 2016.



- Derksen, C., King, J., Belair, S., Garnaud, C., Vionnet, V., Fortin, V., Lemmetyinen, J., Crevier, Y., Plourde, P., Lawrence, B., et al.: Development of the Terrestrial Snow Mass Mission, in: 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, pp. 585 614–617, IEEE, 2021.
- Di Mauro, B., Fava, F., Ferrero, L., Garzonio, R., Baccolo, G., Delmonte, B., and Colombo, R.: Mineral dust impact on snow radiative properties in the European Alps combining ground, UAV, and satellite observations, *Journal of geophysical research: Atmospheres*, 120, 6080–6097, 2015.
- Dick, O., Viallon-Galinier, L., Tuzet, F., Hagenmuller, P., Fructus, M., Reuter, B., Lafaysse, M., and Dumont, M.: Can Saharan dust deposition impact snowpack stability in the French Alps?, *The Cryosphere*, 17, 1755–1773, 2023.
- Dickinson, R., H.-S. A. K.: Biosphere-Atmosphere Transfer Scheme (BATS) Version 1e as Coupled to the NCAR Community Climate Model, Tech. rep., <https://doi.org/10.5065/D67W6959>, 1993.
- Domine, F., Lackner, G., Sarrazin, D., Poirier, M., and Belke-Brea, M.: Meteorological, snow and soil data (2013–2019) from a herb tundra permafrost site at Bylot Island, Canadian high Arctic, for driving and testing snow and land surface models, *Earth System Science Data*, 585 13, 4331–4348, <https://doi.org/10.5194/essd-13-4331-2021>, 2021.
- Douville, H., Royer, J.-F., and Mahfouf, J.-F.: A new snow parameterization for the Météo-France climate model: Part I: validation in stand-alone experiments, *Climate Dynamics*, 12, 21–35, 1995.
- Dumont, M., Tuzet, F., Gascoïn, S., Picard, G., Kutuzov, S., Lafaysse, M., Cluzet, B., Nheili, R., and Painter, T.: Accelerated snow melt in the Russian Caucasus mountains after the Saharan dust outbreak in March 2018, *Journal of Geophysical Research: Earth Surface*, 125, 600 e2020JF005 641, 2020.
- Essery, R., Kontu, A., Lemmetyinen, J., Dumont, M., and Ménard, C. B.: A 7-year dataset for driving and evaluating snow models at an Arctic site (Sodankylä, Finland), *Geoscientific Instrumentation, Methods and Data Systems*, 5, 219–227, 2016.
- Finn. Met. Inst.: Finnish Meteorological Institute - Radiation and snow depth data at Sodankylä Peatland Site, <https://litdb.fmi.fi/peatland.php>, observations from the Arctic Space Centre, 2018.
- 605 Flanner, M., Shell, K., Barlage, M., Perovich, D., and Tschudi, M.: Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008, *Nature geoscience*, 4, 151–155, <https://doi.org/10.1038/NGEO1062>, 2011.
- Flanner, M. G. and Zender, C. S.: Snowpack radiative heating: Influence on Tibetan Plateau climate, *Geophysical research letters*, 32, 2005.
- Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD008003>, 2007.
- 610 Gaillard, M., Vionnet, V., Lafaysse, M., Dumont, M., and Ginoux, P.: Global climatology of light-absorbing particle deposition on snow [Dataset], *Zenodo*, <https://doi.org/10.5281/zenodo.11554783>, 2024a.
- Gaillard, M., Vionnet, V., Lafaysse, M., Dumont, M., and Ginoux, P.: Improved snow ageing parameter for large-scale albedo modelling with Crocus [Dataset], *Zenodo*, <https://doi.org/10.5281/zenodo.11554926>, 2024b.
- Garnaud, C., Bélair, S., Carrera, M. L., Derksen, C., Bilodeau, B., Abrahamowicz, M., Gauthier, N., and Vionnet, V.: Quantifying snow mass mission concept trade-offs using an observing system simulation experiment, *Journal of Hydrometeorology*, 20, 155–173, 2019.
- 615 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419 – 620 5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.



- Ginoux, P., Chin, M., Tegen, I., Prospero, J., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research*, 106, 20 255–20 274, <https://doi.org/10.1029/2000JD000053>, 2001.
- Günther, D., Marke, T., Essery, R., and Strasser, U.: Uncertainties in Snowpack Simulations—Assessing the Impact of Model Structure, Parameter Choice, and Forcing Data Error on Point-Scale Energy Balance Snow Model Performance, *Water Resources Research*, 55, 2779–2800, <https://doi.org/10.1029/2018WR023403>, 2019.
- Hao, D., Bisht, G., Rittger, K., Bair, E., He, C., Huang, H., Dang, C., Stillinger, T., Gu, Y., Wang, H., et al.: Improving snow albedo modeling in the E3SM land model (version 2.0) and assessing its impacts on snow and surface fluxes over the Tibetan Plateau, *Geoscientific Model Development*, 16, 75–94, 2023.
- He, C.: Modelling light-absorbing particle–snow–radiation interactions and impacts on snow albedo: fundamentals, recent advances and future directions, *Environmental Chemistry*, 2022.
- He, C. and Flanner, M.: Snow albedo and radiative transfer: Theory, modeling, and parameterization, *Springer Series in Light Scattering: Volume 5: Radiative Transfer, Remote Sensing, and Light Scattering*, pp. 67–133, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B., Elmore, A., et al.: Importance and vulnerability of the world’s water towers, *Nature*, 577, 364–369, 2020.
- Krajči, P., Kimbauer, R., Parajka, J., Schöber, J., and Blöschl, G.: The Kühltai data set: 25 years of lysimetric, snow pillow, and meteorological measurements, *Water Resources Research*, 53, 5158–5165, <https://doi.org/10.1002/2017WR020445>, 2017.
- Lackner, G., Domine, F., Sarrazin, D., Nadeau, D., and Belke-Brea, M.: Meteorological, snow and soil data, CO<sub>2</sub>, water and energy fluxes, from a low-Arctic valley in the forest-tundra ecotone of Northern Quebec, *Earth System Science Data Discussions*, 2023, 1–28, <https://doi.org/10.5194/essd-2023-7>, 2023.
- Lafaysse, M., Cluzet, B., Dumont, M., Lejeune, Y., Vionnet, V., and Morin, S.: A multiphysical ensemble system of numerical snow modelling, *The Cryosphere*, 11, 1173–1198, <https://doi.org/10.5194/tc-11-1173-2017>, 2017.
- Lapo, K. E., Hinkelman, L. M., Landry, C. C., Massmann, A. K., and Lundquist, J. D.: A simple algorithm for identifying periods of snow accumulation on a radiometer, *Water Resources Research*, 51, 7820–7828, <https://doi.org/10.1002/2015WR017590>, 2015.
- Lee, W. Y., Gim, H.-J., and Park, S. K.: Parameterizations of Snow Cover, Snow Albedo and Snow Density in Land Surface Models: A Comparative Review, *Asia-Pacific Journal of Atmospheric Sciences*, pp. 1–26, 2023.
- Lejeune, Y., Dumont, M., Panel, J.-M., Lafaysse, M., Lapalus, P., Le Gac, E., Lesaffre, B., and Morin, S.: 57 years (1960–2017) of snow and meteorological observations from a mid-altitude mountain site (Col de Porte, France, 1325 m of altitude), *Earth System Science Data*, 11, 71–88, 2019.
- Libois, Q., Picard, G., France, J., Arnaud, L., Dumont, M., Carmagnola, C., and King, M.: Influence of grain shape on light penetration in snow, *The Cryosphere*, 7, 1803–1818, 2013.
- Libois, Q., Picard, G., Arnaud, L., Dumont, M., Lafaysse, M., Morin, S., and Lefebvre, E.: Summertime evolution of snow specific surface area close to the surface on the Antarctic Plateau, *The Cryosphere*, 9, 2383–2398, <https://doi.org/10.5194/tc-9-2383-2015>, 2015.
- Martin, E. and Lejeune, Y.: Turbulent fluxes above the snow surface, *Annals of Glaciology*, 26, 179–183, 1998.
- Melloh, R. A., Hardy, J. P., Davis, R. E., and Robinson, P. B.: Spectral albedo/reflectance of littered forest snow during the melt season, *Hydrological Processes*, 15, 3409–3422, 2001.





- Ménard, C. B., Essery, R., Barr, A., Bartlett, P., Derry, J., Dumont, M., Fierz, C., Kim, H., Kontu, A., Lejeune, Y., Marks, D., Niwano, M., Raleigh, M., Wang, L., and Wever, N.: Meteorological and evaluation datasets for snow modelling at 10 reference sites: description of in situ and bias-corrected reanalysis data, *Earth System Science Data*, 11, 865–880, <https://doi.org/10.5194/essd-11-865-2019>, 2019.
- Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., Brutel-Vuilmet, C., Burke, E., Cuntz, M., Dai, Y., et al.: Scientific and human errors in a snow model intercomparison, *Bulletin of the American Meteorological Society*, 102, E61–E79, 2021.
- Mölders, N., Luijting, H., and Sassen, K.: Use of atmospheric radiation measurement program data from Barrow, Alaska, for evaluation and development of snow-albedo parameterizations, *Meteorology and Atmospheric Physics*, 99, 199–219, 2008.
- 665 Morin, S., Lejeune, Y., Lesaffre, B., Panel, J.-M., Poncet, D., David, P., and Sudul, M.: An 18-yr long (1993–2011) snow and meteorological dataset from a mid-altitude mountain site (Col de Porte, France, 1325 m alt.) for driving and evaluating snowpack models, *Earth System Science Data*, 4, 13–21, <https://doi.org/10.5194/essd-4-13-2012>, 2012.
- Mortimer, C., Mudryk, L., Derksen, C., Luoju, K., Brown, R., Kelly, R., and Tedesco, M.: Evaluation of long-term Northern Hemisphere snow water equivalent products, *The Cryosphere*, 14, 1579–1594, <https://doi.org/10.5194/tc-14-1579-2020>, 2020.
- 670 Mudryk, L., Mortimer, C., Derksen, C., Elias Chereque, A., and Kushner, P.: Benchmarking of SWE products based on outcomes of the SnowPEX+ Intercomparison Project, *EGUsphere*, 2024, 1–28, 2024.
- Painter, T., Barrett, A., Landry, C., Neff, J., Cassidy, M., Lawrence, C., McBride, K., and Farmer, L.: Impact of disturbed desert soil on duration of mountain snow cover, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL030284>, 2007.
- Painter, T., Skiles, M., Deems, J., Brandt, W., and Dozier, J.: Variation in Rising Limb of Colorado River Snowmelt Runoff Hydrograph Controlled by Dust Radiative Forcing in Snow, *Geophysical Research Letters*, 45, <https://doi.org/10.1002/2017GL075826>, 2017.
- 675 Painter, T. H., Deems, J. S., Belnap, J., Hamlet, A. F., Landry, C. C., and Udall, B.: Response of Colorado River runoff to dust radiative forcing in snow, *Proceedings of the National Academy of Sciences*, 107, 17 125–17 130, <https://doi.org/10.1073/pnas.0913139107>, 2010.
- Pedersen, C. A. and Winther, J.-G.: Intercomparison and validation of snow albedo parameterization schemes in climate models, *Climate Dynamics*, 25, 351–362, 2005.
- 680 Qu, X. and Hall, A.: Assessing snow albedo feedback in simulated climate change, *Journal of Climate*, 19, 2617–2630, 2006.
- Romanov, P.: Global Multisensor Automated satellite-based Snow and Ice Mapping System (GMASI) for cryosphere monitoring, *Remote Sensing of Environment*, 196, 42–55, <https://doi.org/10.1016/j.rse.2017.04.023>, 2017.
- Réveillet, M., Dumont, M., Gascoïn, S., Lafaysse, M., Nabat, P., Ribes, A., Nheili, R., Tuzet, F., Ménégoz, M., Morin, S., Picard, G., and Ginoux, P.: Black carbon and dust alter the response of mountain snow cover under climate change, *Nature Communications*, 13, <https://doi.org/10.1038/s41467-022-32501-y>, 2022.
- 685 Skiles, M., Flanner, M., Cook, J., Dumont, M., and Painter, T.: Radiative forcing by light-absorbing particles in snow, *Nature Climate Change*, 8, <https://doi.org/10.1038/s41558-018-0296-5>, 2018.
- Sturm, M., Goldstein, M. A., and Parr, C.: Water and life from snow: A trillion dollar science question, *Water Resources Research*, 53, 3534–3544, 2017.
- 690 Tutton, R., Darkin, B., Essery, R., Griffith, J., Hould Gosselin, G., Marsh, P., Sonnentag, O., Thorne, R., and Walker, B.: A hydro-meteorological dataset from the taiga-tundra ecotone in the western Canadian Arctic: Trail Valley Creek, Northwest Territories (1991-2023) (DRAFT VERSION), <https://doi.org/10.5683/SP3/BXV4D>, 2024.
- Tuzet, F., Dumont, M., Lafaysse, M., Picard, G., Arnaud, L., Voisin, D., Lejeune, Y., Charrois, L., Nabat, P., and Morin, S.: A multilayer physically based snowpack model simulating direct and indirect radiative impacts of light-absorbing impurities in snow, *The Cryosphere*, 11, 2633–2653, <https://doi.org/10.5194/tc-11-2633-2017>, 2017.
- 695



- Tuzet, F., Dumont, M., Arnaud, L., Voisin, D., Lamare, M., Larue, F., Revuelto, J., and Picard, G.: Influence of light-absorbing particles on snow spectral irradiance profiles, *The Cryosphere*, 13, 2169–2187, <https://doi.org/10.5194/tc-13-2169-2019>, 2019.
- van Dalum, C. T., van de Berg, W. J., Libois, Q., Picard, G., and van den Broeke, M. R.: A module to convert spectral to narrowband snow albedo for use in climate models: SNOWBAL v1. 2, *Geoscientific Model Development*, 12, 5157–5175, 2019.
- 700 Veillon, F., Dumont, M., Amory, C., and Fructus, M.: A versatile method for computing optimized snow albedo from spectrally fixed radiative variables: VALHALLA v1. 0, *Geoscientific Model Development*, 14, 7329–7343, 2021.
- Verseghy, D. L.: CLASS—A Canadian land surface scheme for GCMs. I. Soil model, *International Journal of Climatology*, 11, 111–133, 1991.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geoscientific Model Development*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- 705 Vionnet, V., Guyomarc’h, G., Bouvet, F. N., Martin, E., Durand, Y., Bellot, H., Bel, C., and Puglièse, P.: Occurrence of blowing snow events at an alpine site over a 10-year period: Observations and modelling, *Advances in water resources*, 55, 53–63, 2013.
- Vionnet, V., Verville, M., Fortin, V., Brugman, M., Abrahamowicz, M., Lemay, F., Thériault, J. M., Lafaysse, M., and Milbrandt, J. A.: Snow Level From Post-Processing of Atmospheric Model Improves Snowfall Estimate and Snowpack Prediction in Mountains, *Water Resources Research*, 58, e2021WR031778, <https://doi.org/10.1029/2021WR031778>, e2021WR031778 2021WR031778, 2022.
- 710 Warren, S. G. and Wiscombe, W. J.: A Model for the Spectral Albedo of Snow. II: Snow Containing Atmospheric Aerosols, *Journal of Atmospheric Sciences*, 37, 2734 – 2745, [https://doi.org/10.1175/1520-0469\(1980\)037<2734:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2734:AMFTSA>2.0.CO;2), 1980.
- Woolley, G. J., Rutter, N., Wake, L., Vionnet, V., Derksen, C., Essery, R., Marsh, P., Tutton, R., Walker, B., Lafaysse, M., and Pritchard, D.: Multi-physics ensemble modelling of Arctic tundra snowpack properties, *EGUsphere*, 2024, 1–37, 2024.
- 715 Yen, Y.-C.: Review of thermal properties of snow, ice, and sea ice, vol. 81, US Army, Corps of Engineers, Cold Regions Research and Engineering Laboratory, 1981.
- Zhao, M., Golaz, J.-C., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S.-M., Freidenreich, S. M., Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A. R., Liang, Z., Lin, P., Lin, S.-J., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik, V., Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G., Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B., and Xiang, B.: The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 1. Simulation Characteristics With Prescribed SSTs, *Journal of Advances in Modeling Earth Systems*, 10, 691–734, <https://doi.org/10.1002/2017MS001208>, 2018.
- 720