

Dynamics of CO₂ and CH₄ fluxes in Red Sea mangrove soils

Jessica Breavington^{1,2,3}, Alexandra Steckbauer^{1,2,3}, Chuancheng Fu^{1,2,3}, Mongi Ennasri^{1,2,3}, Carlos M. Duarte^{1,2,3}

5 ¹Marine Science Program, Biological and Environmental Science and Engineering Division (BESE), King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Kingdom of Saudi Arabia

²Red Sea Research Center (RSRC), King Abdullah University of Science and Technology, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Kingdom of Saudi Arabia

10 ³Computational Bioscience Research Center (CBRC), King Abdullah University of Science and Technology, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Kingdom of Saudi Arabia

Correspondence to: Jessica Breavington (jessicabreavington@gmail.com)

Abstract. Red Sea mangroves have a lower carbon burial rate than the global average, whereby small greenhouse gas fluxes may offset a large proportion of carbon burial. Monthly soil core sampling was conducted across 2 years at two sites within a central Eastern Red Sea mangrove stand to examine carbon dioxide (CO₂) and methane (CH₄) fluxes under dry and inundated conditions. Fluxes were highly variable, characterized by a prevalence of low emissions punctuated by bursts of high emissions. At the landward site, average \pm SE (median) flux from the soil-air interface was 3111 ± 929 (811) $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and 1.68 ± 0.63 (0.26) $\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ under light conditions, and 8657 ± 2269 (1615) $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and 0.84 ± 0.79 (0.59) $\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ under dark conditions. Average \pm SE (median) sea-air fluxes were -55 ± 165 (-79) $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and 0.12 ± 0.23 (0.08) $\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ under light conditions, and 27 ± 48 (53) $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and 0.16 ± 0.13 (0.09) $\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ in dark conditions. The seaward site recorded higher CH₄ flux, averaging 18.7 ± 8.18 (1.7) and 17.1 ± 4.55 (7.7) $\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ in light and dark conditions. Mean fluxes offset 94.5 % of carbon burial, with a median of 4.9 % skewed by extreme variability. However, reported CO₂ removal by total alkalinity emission from carbonate dissolution greatly exceeded both processes and drives the role of these ecosystems as intense CO₂ sinks.

25

1 Introduction

30 Mangrove forests thrive in estuarine and intertidal zones within latitudes of 0 ° to 40 ° (Rosentreter et al., 2018a), storing a significant amount of organic carbon and providing numerous ecosystem services, including coastal protection and biodiversity enhancement (Curran et al., 2002; Howard et al., 2014). Mangroves offer a promising nature-based solution to

mitigate global warming due to their high sequestration of soil organic carbon (C_{org}), while offering coastal protection to sea level rise (Duarte et al., 2013). Carbon preservation in mangrove soils is facilitated by waterlogged, anoxic conditions that limit the decay of organic matter (OM). However, as mangroves exist at the boundary between terrestrial and marine environments, the capacity for carbon sequestration varies depending on multiple factors such as the tidal range, sediment and nutrient inputs.

Mangroves in the Red Sea are subject to extreme environmental conditions that restrict their growth and productivity. The Red Sea is one of the warmest and most saline seas globally, characterized by oligotrophic and nutrient-limited conditions. Moreover, central Saudi Arabia experiences extreme aridity, with an average annual precipitation of 60 mm (Gabr et al., 2017). Consequently, *Avicennia marina* is the predominant mangrove species in the Red Sea, existing at the thresholds of its physiochemical physiological tolerance. It is one of the most highly adapted mangrove species to the high salinity and aridity, and found predominantly as monospecific mangrove stands (Khalil, 2015). *Rhizophora mucronata* are also found within the Red Sea but predominantly in Southern regions where there is lower salinity (Khalil, 2015). Due to the absence of permanent rivers, mangroves in the Saudi Arabian Red Sea typically form narrow fringing bands along the coastline. In the central Red Sea, the distribution of mangroves is also constrained by the small tidal range, which is typically less than 1.5 m (Blanco-Sacristán et al., 2022). The conditions in the Red Sea result in reduced growth of *A. marina* with trees only reaching 2-3 meters compared to over 16 meters in Australia (Mackey, 1993). The oligotrophic conditions prevalent in the Red Sea result in stunted growth and dwarf forms of mangroves due to nutrient limitation (Almahasheer et al., 2016b). As a result, mangroves in the Red Sea have one of the lowest rates carbon sequestration rates, approximately $15 \pm 1 \text{ g } C_{org} \text{ m}^{-2} \text{ yr}^{-1}$, compared with a global average estimated at $163 \text{ g } C_{org} \text{ m}^{-2} \text{ yr}^{-1}$ (Almahasheer et al., 2017; Breithaupt et al., 2012; Sanderman et al., 2018).

Greenhouse gas (GHG) fluxes, involving the release of carbon dioxide (CO_2) or methane (CH_4), in mangrove soils partially offset their role in removing atmospheric CO_2 , which is at its highest in the past 800,000 years (Tripathi et al., 2009), contributing to radiative heating of the atmosphere and a global temperature increase at a rate of $1.7 \text{ }^\circ\text{C}$ per century since the beginning of the industrial revolution (IPCC, 2014; Marcott et al., 2013). CH_4 is the second most important GHG associated with climate change (Forster et al., 2007), and substantially more potent than CO_2 , with a global warming potential (GWP) approximately 28 times greater (Myhre et al., 2014). The low carbon sequestration rates of Red Sea mangroves may be offset by GHG fluxes. However, a lack of dynamic estimates for GHG fluxes from arid mangrove soils in the Red Sea preclude such assessment. To date, and to the best of our knowledge, only one other known study has provided estimates of GHG fluxes from mangrove soil in the Red Sea over a short period in this region. These fluxes ranged from -3452 to $7500 \text{ } \mu\text{mol } \text{CO}_2 \text{ m}^{-2} \text{ d}^{-1}$ and 0.9 to $13.3 \text{ } \mu\text{mol } \text{CH}_4 \text{ m}^{-2} \text{ d}^{-1}$ across different locations in the Red Sea (Sea et al., 2018). Therefore, it is difficult to reliably quantify the role of GHG emissions in offsetting CO_2 removal by carbon sequestration in Red Sea

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Subscript

Formatted: Superscript

Formatted: Superscript

Formatted: Subscript

mangrove soils, which is important to creating accurate carbon budgets for arid mangroves. Furthermore, GHG flux estimations exhibit wide variation due to factors such as location, environmental conditions, and study design.

Intertidal conditions in mangrove forests allow for flux measurements directly from the soil to the air (soil-air interface) or through the sea-air interface, with different transfer velocity equations introducing variability in the flux estimates (Akhand et al., 2021; Call et al., 2015). Additionally, flux measurements can be measured *in situ* or through controlled *ex situ* laboratory studies, with variations in chamber design, that can be closed, or open with circulating air. Recent advancements in measurement technology, particularly with the growing use of cavity ring-down spectroscopy (CRDS), enable high-accuracy measurements even at low gas concentrations, but accurate comparison with other methods, such as eddy flux covariance can be challenging (Brannon et al., 2016). ~~Considering these methodological variations is essential when comparing results across studies to explain the high variability observed across locations.~~ Furthermore, environmental variables and ~~physiochemical~~physicochemical properties should be considered to comprehensively understand the variability of GHG emissions from mangrove soils. A comprehensive understanding of carbon stores and emissions in mangrove ecosystems is required to accurately determine the net climate benefits from mangrove coverage and restoration efforts (Lovelock et al., 2022). The Red Sea is one of the few regions where mangrove coverage has been steadily increasing over the past four decades, underscoring the importance of accurate carbon budgets for Red Sea mangroves (Almahasheer et al., 2016a).

Here, we quantify the dynamics of CO₂ and CH₄ fluxes from mangrove soils in a mangrove stand in the highly arid Central Red Sea to assess the scale of soil carbon burial offset by GHG flux. We also test the effect of various physical and chemical soil properties on GHG fluxes. This study represents the first effort to simultaneously measure CO₂ and CH₄ fluxes from both the sea-air and soil-air interfaces in Red Sea mangroves over a time series relevant to providing needed insights into the dynamics of carbon cycling in this unique ecosystem.

2 Methods

2.1 Sampling location

Sampling was conducted at two adjacent monospecific *Avicenna marina* mangrove stands in Thuwal, on the eastern coast of the Central Red Sea (22.340787°N, 39.087991°E) (Fig. 1). Soil cores for CO₂ and CH₄ flux were collected over two years, from April 2021 to May 2023, on a monthly basis, except when this was prevented by logistical challenges. The first sampling location was approximately 150 m inland from the coast, referred to as the landward site, with an elevation approximately 0.75 m above sea level. This landward site was characterised by a strong seasonal microtidal influence, with a tidal range of less than 0.5 m, resulting in a scarce tidal inundation during the summer months and a more regular inundation during winter. The second site was located approximately 200 m from the landward site, referred to as the seaward site.

100 Sampling of this second site was conducted over a narrower time window between September and October 2022: weekly sampling for two consecutive weeks, followed by a two-week break to minimise disturbance to the site and allow for a greater range in temperature, and then resumed for another two weeks of weekly sampling, resulting in a total of four sampling events. This seaward site experienced continuous water coverage across the sampling period and resultingly, was subject to fewer environmental extremes than the landward site.



105 **Figure 1: Mangrove sampling sites as indicated by red circles. Inset: Location of sampling in the Eastern Central Red Sea (from ESRI Satellite).**

2.2 Core collection

110 Two sets of cores were collected each month. The first set of cores comprised four large clear PVC cylinders (height: 30 cm, diameter: 9.6 cm) inserted into the soil to a depth of 10 cm and retrieved without disturbing the soil layers. During sampling roots and undecomposed organic matter were avoided as the aim was to estimate GHG emission from the soil rather than the mangrove trees. If water was present during sampling, it was retained within the cylinder up to a maximum depth of 10 cm to ensure a minimum of 10 cm of air for incubation, and without disturbing the soil-water interface. Initially, in April, May, and July (2021) collection of the soil cores was conducted in the early morning hours allowing for sufficient time to transport and process soil cores with a stabilisation period for 1 hour between sealing the core and taking the T0 gas sample at 7am (following the protocol of Sea et al. 2018). Subsequent sampling events, until the study conclusion, were conducted late

115 afternoon on the day before for logistical reasons, with the cores left unsealed in the incubator under darkness to mirror
night-time conditions. Leaving the cores unsealed and undisturbed was to allow for regular gas exchange following
disturbance of the soil caused by collection in the field, and to avoid the creation of anoxic conditions in the sediment and
water overnight before the start of the experiment. If water was present at the time of sample collection, extra water was
120 sampled with the sediment cores and also placed in the incubator to keep the temperature stable. On the morning of the start
of the experiment, water was exchanged, and the air-water interface of the sealed cores was allowed to stabilize for 1-hour
before the collection of the T0 gas measurement at 7am (Sea et al., 2018). The 1-hour stabilization was not required for cores
without the water phase there was no water to exchange, and no water-soil interface to influence gas exchange dynamics.
There were no significant differences in T0 concentrations with or without water.

125 The second set of cores (height: 9 cm, diameter: 2.5 cm) were collected immediately next to the large cores and used
to assess the physical and chemical properties of the soil, including conductivity, total carbon (TC), total organic carbon
(C_{org}), total nitrogen (TN), bulk density (BD), and water content (WC). Both sets of cores were transferred into a cooler and
transported to the lab for processing within an hour after sampling. Temperature and tidal inundation at the site were
continuously recorded via *in situ* temperature and water level loggers (U22-001 v2 and U20L-04, Onset Computer Corp.,
130 Massachusetts, USA).

2.3 Flux measurements

GHG flux was measured from the soil-air interface or the sea-air interface, depending on the presence of water at the time of
sampling. The four replicate cores were immediately transported to the laboratory and placed in an incubator (I-30L,
Percival, Geneva Scientific LLC, Fontana Wisconsin, USA). Temperature was set to the average temperature in the field as
135 determined by readings from *in situ* temperature loggers (U22-001 v2, Onset Computer Corp., Massachusetts, USA) over the
past 72 hours from the time of sampling. After sampling, an airtight lid was fitted to the bottom of the core, and opaque tape
was wrapped around the outside of the core to cover the soil phase, to avoid light exposure to the sides of the soil. The top of
each core was left unsealed and kept in the incubator overnight to equilibrate. Immediately before the onset of sampling, the
top of the cores were sealed with airtight lids. Three gas samples of 25 mL per core were taken starting at 7 am (T0), after 12
140 hours of light (T1), and the final sample (T2) after 12 hours of darkness. For the duration of the light condition, incubator
lights were set to 100 % intensity at 125 $\mu\text{mol m}^{-2} \text{sec}^{-1}$ irradiance (I-30L, Percival, Geneva Scientific LLC, Fontana
Wisconsin, USA). Gas samples were taken using a syringe and valve system. The syringes with gas samples were connected
to a G2201-*i* CRDS analyser (Picarro, Santa Clara, California USA), coupled to a Small Sample Introduction Module
(SSIM2), to measure CO₂, CH₄, $\delta^{13}\text{C-CO}_2$ and $\delta^{13}\text{C-CH}_4$.

145 CO₂ and CH₄ concentrations were converted from dry mole fractions in parts per million (ppm) to $\mu\text{mol m}^{-2} \text{d}^{-1}$ (24 hours)
using the linear change in concentration between the 12-hour sampling periods (Brannon et al., 2016; Tete et al., 2015) (Eq.
1).

Formatted: Superscript

$$F = \frac{dC}{dt} \left(\frac{PV}{RAT} \right) \quad [\text{Eq. 1}]$$

Where F is flux of CO₂ or CH₄ (μmol m⁻²); dC/dt is the linear concentration change of CO₂ or CH₄ over 12 hours from T0 to T1 to measure light fluxes or T1 to T2 to measure fluxes under the dark condition; P is the pressure (Pa) calculated using Boyles law which was used to correct the pressure in the headspace after taking 25 mL air at each sampling point; V is the volume of gas (m³) in the cylinder headspace; R is the ideal gas constant (8.314 J mol⁻¹ K⁻¹); A is the area of soil (m²); and T is temperature (K).

The CO₂ equivalent (g CO₂-eq m⁻² y⁻¹) of the combined CO₂ and CH₄ fluxes was calculated for the flux across the sea-air and soil-air interfaces (Eqs. 2 & 3). Mangrove carbon storage was calculated using estimates from previous studies in the Red Sea. Using 55 g CO₂-eq m⁻² y⁻¹ for the soil carbon burial rate (Almahasheer et al., 2017), and 1266 g CO₂-eq m⁻² yr⁻¹ for CO₂ uptake from total alkalinity (TA) enhancement determined at this site (Saderne et al., 2021).

$$\frac{\mu\text{mol m}^{-2}\text{d}^{-1}}{1,000,000} \times 365 = \text{mol m}^{-2} \text{y}^{-1} \quad [\text{Eq. 2}]$$

$$(\text{mol CO}_2 \text{ m}^{-2} \text{y}^{-1} \times 44) + (\text{mol CH}_4 \text{ m}^{-2} \text{y}^{-1} \times 16 \times 28) = \text{g CO}_2 - \text{eq m}^{-2} \text{y}^{-1}$$

[Eq. 3]

Where CO₂ = 44 g mol⁻¹, CH₄ = 16 g mol⁻¹, and CH₄ global warming potential (GWP) over a 100-year horizon = 28 (IPCC, 2014).

2.4 Soil chemical and physical variables

The soil samples from the small cores were fully dried at 60 °C to a constant weight. Samples were ground using an agate pestle and mortar for analysis of total carbon (TC), total organic carbon (C_{org}), total nitrogen (TN) and soil conductivity (EC_{1:5}). For C_{org}, a 10 ± 1 mg sample was acidified with 5 μl of 3 mol HCL L⁻¹ in silver capsules. Samples were dried for 30 min at 60 °C and acidification was repeated a minimum of 3 times, or until no bubbles were observed during the addition of HCL to remove all carbonates before being fully dried and wrapped in tin capsules for organic elemental analysis (Flash 2000, ThermoFisher Scientific, Massachusetts, USA). Soil organic carbon (C_{org}) and inorganic carbon (C_{inorg}) for 0-3 cm soil depth was calculated using bulk density (Howard et al., 2014), and with the following formulas (Eqs. 4 & 4-65):

$$\text{Bulk Density (g cm}^{-3}\text{)} = \frac{\text{Oven-dry sample mass (g)}}{\text{Sample volume (m}^3\text{)}} \quad \text{---}$$

[Eq. 4]

$$C_{org} (\text{mg C}_{org} \text{ cm}^{-3}) = \text{Bulk Density (g cm}^{-3}\text{)} \times \left(\frac{\text{TOC (\%)}}{100} \right) \times 1000 \quad \text{---}$$

[Eq. 54]

$$C_{inorg} (mg C cm^{-3}) = Bulk Density (g cm^{-3}) \times \left(\frac{TC (\%) - TOC (\%)}{100} \right) \times 1000$$

[Eq. 65]

Conductivity was measured using an electrical conductivity (EC) sensor (Inlab® 738 ISM, Mettler Toledo, Schwerzenbach, Switzerland). Prior to measurement the sensor was calibrated with 12.88 mS/cm potassium chloride as produced by the manufacturer (Mettler Toledo). For the surface soil, 5 ± 0.01 g of soil was used with 25 mL water for a 1:5 ratio of 1 part soil to 5 parts Milli-Q water. The samples were mixed on an orbital shaker (VWR®) following a typical protocol for the EC_{1:5} method for high salinity soils (Hardie and Doyle, 2012; Kargas et al., 2018).

2.5 Data analysis

Differences in mean soil properties and GHG flux between sampling sites, and wet and dry conditions were evaluated for significance by means of Mann-Whitney U test in R Studio (v.4.1.2). A correlation matrix showing significance between GHG fluxes, isotope signatures, soil properties and environmental variables using Spearman rank correlation coefficient was created with the use of ‘Scipy’ package (v1.11.1) in Python (v3.11.5). In addition, a random forest algorithm was used to model the influence of environmental, and temporal variables on CO₂ flux in light and dark conditions through the use of regression trees utilizing bootstrapping techniques (Breiman, 2001). The models were built in Python v.3.9.13 and Jupyter Notebook v.6.4.12 using the RandomForestRegressor from the SciKit-Learn package v.1.0.2. Only data from the landward site was used in the models due to the greater number of observations and longer sampling period. 80% of data was randomly selected and used for training, with the remaining 20% used for validation.

To optimize model accuracy and minimise overfitting, we utilized the R² metric, which is an easy-to-interpret standardised measure of linear association (Fox et al., 2017), and implemented a 5-fold cross-validation to assess how the model generalizes to unseen data and reduces the risk of overfitting. Hyperparameter tuning for the number of trees, maximum depth, minimum sample split, and minimum sample leaf, was utilized to maximise the R² metric. Furthermore, a baseline accuracy threshold was defined for feature selection, where $R^2 \geq 0.6$ and the average 5-fold cross-validation (CV) score ≥ 0.4 . Backward elimination of variables based on these performance metrics was systematically performed to maximise the number of variables included within each model without sacrificing model performance to ensure the maximum predictive power without overfitting (Genuer et al., 2010; Speiser et al., 2019). These models were used to map feature importance of the variables retained from the feature selection stage. All other figures were made with the use of ggplot in R Studio (v.4.1.2).

3 Results

3.1 Soil properties

210 The most pronounced variation in soil characteristics between wet and dry sampling conditions at the landward site was observed in conductivity (EC), averaging 22.6 mS cm⁻¹ under dry conditions compared to 9.25 mS cm⁻¹ under wet conditions (Table 1), although all locations were classified under the ‘extreme’ salinity class (Hardie and Doyle, 2012). EC and WC were the only soil properties to demonstrate significant differences (p<0.001) under wet and dry sampling conditions at the landward site. The largest contrast between the two sampling sites was evident in the C_{inorg} concentration, with the seaward site exhibiting a significantly higher (p<0.001) mean C_{inorg} (94.51 ± 3.37 mg C cm⁻³) compared to the landward site (66.64 ± 1.16 mg C cm⁻³) under wet and dry sampling conditions. Additionally, the seaward site had a lower C_{org} concentration, averaging 5.53 mg C_{org} cm⁻³ (0.34 % ± 0.017 %) compared to an average of 9.52 mg C_{org} cm⁻³ (0.72 % ± 0.021 %) at the landward site throughout the entire sampling period. C_{org} was significantly greater (p<0.001) at the landward site under dry conditions, averaging 2.43 mg C_{org} cm⁻³ more than the seaward site. Under wet conditions, there was a smaller but still significant difference (p<0.05) of 2.29 mg C_{org} cm⁻³ between the landward and seaward sites.

225 **Table 1: Average soil properties (± SE) for the top 3cm of soil in at the landward site in dry and wet sampling conditions, and at the continually inundated seaward site. C:N = C:N (molar ratio), C_{org}= Organic carbon, C_{inorg} = Inorganic carbon, WC = Water content, EC_{1:5} = Electrical conductivity (1:5 soil:water ratio). Mean values for sampling sites and conditions with no common letter show significant differences (Mann-Whitney U test, p<0.05).**

Sample location and condition	C:N	C _{org} (mg C _{org} cm ⁻³)	C _{inorg} (mg C cm ⁻³)	WC (%)	EC _{1:5} (mS cm ⁻¹)
Landward – Dry	12.67 ± 0.43 ^a	7.96 ± 0.24 ^a	65.38 ± 1.51 ^a	40.63 ± 2.30 ^a	22.61 ± 1.71 ^a
Landward – Wet	12.44 ± 0.43 ^a	7.82 ± 0.46 ^a	68.43 ± 1.79 ^a	49.57 ± 1.49 ^b	9.25 ± 0.47 ^b
Seaward – Wet	11.49 ± 2.29 ^b	5.53 ± 0.95 ^b	93.51 ± 3.37 ^b	34.97 ± 1.16 ^a	5.71 ± 0.23 ^c

Formatted: Font: Not Bold


Formatted: Font: Bold

Formatted: Font: Bold

3.2 Highly variable CO₂ and CH₄ fluxes

230 Between April 2021 to May 2023 twenty months were sampled at the landward site. Nine were under inundated conditions measuring flux from the sea-air interface, and eleven months of sampling was under dry conditions, measuring fluxes from the soil-air interface. Five months could not be sampled due to logistical issues. At the landward site the CO₂ flux varied between -3136 μmol CO₂ m⁻² d⁻¹ in the light condition to 37,644 μmol CO₂ m⁻² d⁻¹ in the dark condition (Fig. 2). The average fluxes combined across the soil-air and sea-air interfaces were 1686 ± 546 μmol CO₂ m⁻² d⁻¹ under the light conditions and three times larger, 4774 ± 1337 μmol CO₂ m⁻² d⁻¹, under dark conditions (Table. 2). The net daily flux over the full incubation period combining light and dark fluxes was 3178 ± 806 μmol CO₂ m⁻² d⁻¹ (range: -811 to 28,048 μmol CO₂ m⁻² d⁻¹).


235 ¹). On average, the soil was a net source of CO₂ to the atmosphere in all conditions except the light CO₂ flux from the sea-air interface at the landward site (-55 μmol CO₂ m⁻² d⁻¹).



240 **Figure 2: Median values of CO₂ flux for each month and condition (light and dark) at the landward site. The thick line inside the box represents the median value of the data, and 25th and 75th percentiles denoted by the box ends. The whiskers extend to the minimum and maximum values within 1.5 times the interquartile range, outliers are marked by black points. Blue shading: periods of net flux from the sea-air interface. No shading: periods of net flux from the soil air interface. Note: The axis label for the time scale is non-continuous, as months without sampling are not shown.**

245

The average CH₄ flux at the landward site was 0.98 ± 0.37 μmol CH₄ m⁻² d⁻¹ under the light conditions, 0.54 ± 0.44 μmol CH₄ m⁻² d⁻¹ under dark conditions (Fig. 3). The net daily flux over the 24-hour incubation period was 0.74 ± 0.23 μmol CH₄ m⁻² d⁻¹ (range: -1.47 to 5.71 μmol CH₄ m⁻² d⁻¹).



250 **Figure 3: Median values of CH₄ flux for each month and condition (light and dark) at the landward site. The thick line inside the box represents the median value of the data, and 25th and 75th percentiles denoted by the box ends. The whiskers extend to the minimum and maximum values within 1.5 times the interquartile range, outliers are marked by black points. Blue shading: periods of net flux from the sea-air interface. No shading: periods of net flux from the soil air interface. Note: The axis label for the time scale is non-continuous, as months without sampling are not shown.**

255

At the seaward site, only sea-air flux was measured given the constant inundation, and there was a lower CO₂ flux compared to the overall mean CO₂ flux from landward site (Table 2). However, there was a higher mean and median sea-air CO₂ flux when compared with only the sea-air fluxes from the landward site (Fig. 4). CH₄ flux was also significantly higher than that at the landward site (Table 2). The average flux was 18.67 and 17.12 µmol CH₄ m⁻² d⁻¹ in light and dark conditions, respectively.

260

The isotopic signature of CO₂ averaged -12.02 ± 0.14 ‰ at the landward site and -11.75 ± 0.46 ‰ at the seaward site. Despite the lighter isotope at the landward site there was no significant difference in $\delta^{13}\text{C-CO}_2$ between sites (Mann-Whitney U test, $p = 0.0795$). The isotopic signature of the CH₄ averaged -46.24 ± 0.58 ‰ at the landward site and -48.18 ± 0.67 ‰ at the seaward site, with no significant difference in $\delta^{13}\text{C-CH}_4$ between sites (Mann-Whitney U test, $p = 0.3684$). The

265

$\delta^{13}\text{C}$ signature of CO_2 and CH_4 did not change significantly across seasons. However, significant correlations ($p > 0.05$) were observed between core replicates and inorganic carbon ($\text{C}_{\text{inorg-3cm}}$) with $\delta^{13}\text{C}\text{-CO}_2$, and well as between electrical conductivity ($\text{EC}_{4.5}$) and $\delta^{13}\text{C}\text{-CH}_4$ (Fig. S1).

Formatted: Subscript

Formatted: Subscript

Table 2: Summary of CO_2 and CH_4 fluxes and combined $\text{CO}_2\text{-eq}$ flux offset (using the total flux over the 24-hour incubation) of carbon burial (C burial) and total alkalinity enhancement (TA) for the landward and seaward study sites (Carbon burial data adapted from Almahasheer et al., 2017 and Saderne et al., 2021). Mean values for sampling sites and conditions with no common letter show significant differences (Mann-Whitney U test, $p < 0.05$).

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold


Sampling site and condition	Light CO_2 ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$)	Dark CO_2 ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ d}^{-1}$)	Light CH_4 ($\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$)	Dark CH_4 ($\mu\text{mol CH}_4 \text{ m}^{-2} \text{ d}^{-1}$)	$\text{CO}_2\text{-eq}$ ($\text{g CO}_2\text{-eq m}^{-2} \text{ y}^{-1}$)	C burial offset by flux (%)	C burial and TA offset by flux (%)
Landward							
Sea-air mean	-55.2 ^{bc}	27.5 ^a	0.12 ^a	0.19 ^a	-0.2 ^a	-0.4 ^a	-0.01 ^a
Soil-air mean	3110.8 ^{bc}	8657.4 ^b	1.25 ^{ab}	0.83 ^b	94.7 ^b	172.1 ^b	7.2 ^b
Sea-air mean	-55.2	27.5	0.12	0.19	-0.2	-0.4	-0.01
Combined mean	1686.1 ^c	4774.0 ^{ab}	0.98 ^a	0.54 ^c	52.0 ^c	94.5 ^c	3.9 ^c
Median	216.2	115.3	0.38	0.17	2.7	4.9	0.2
Min	-3135.7	-5799.6	0.03	0.04	-71.7	-130.4	-5.4
Max	18547.2	37644.0	5.71	3.96	452.0	821.8	34.2
Seaward							
Sea-air mean	2832.5 ^{bc}	2244.9 ^b	18.67 ^b	17.12 ^d	43.7 ^b	79.5 ^b	3.3 ^b
Median	2187.7	1669.4	1.75	7.69	31.7	57.7	2.4
Min	-69.4	-2770.4	-0.35	-0.51	-22.9	-41.6	-1.7
Max	10361.4	7785.7	101.9	51.2	158.2	287.7	12.0


Fluxes were generally a net source of $\text{CO}_2\text{-eq}$ to the atmosphere (Table 2). On average, 95% of soil carbon burial was offset by GHG flux at the seaward/landward site. However, the estimates were highly skewed, so that the mean value does not represent the central tendency, which was best represented by the median flux, which was equivalent to 4.9% offset relative to carbon burial rates. Median $\text{CO}_2\text{-eq}$ fluxes only offset 4.9% of the carbon burial rate at the same site.

When incorporating the CO_2 drawdown of TA enhancement, 3.9% (mean) and 0.2% (median) of carbon sequestration potential was offset by the GHG fluxes measured in this study at the landward site (Table 2). At the seaward site, the greater flux and GWP of CH_4 resulted in a greater median offset of carbon burial compared to the landward site but the mean offset at the landward site remained higher due to the very large upper-range CO_2 fluxes. Generally, the $\text{CO}_2\text{-eq}$ offset was

significantly higher when fluxes were measured between the soil-air interface, compared to measurements between the sea-air interface (Fig. 4).

285





290 **Figure 4: Boxplot comparison of mean (red diamond), IQR (boxes), median (black line) and outliers (black points/white circles) of CO₂-eq flux across sites, with landward sites separated by dry and wet sampling conditions. “Total” shows the combined g CO₂-eq m⁻² y⁻¹ for both CO₂ and CH₄. The right panel displays the full range of CO₂-eq flux, while the left panel provides a zoomed view for comparison of smaller fluxes under wet conditions.**

3.3 Drivers of flux variation

295 There were several significant correlations relating to GHG flux with environmental and soil properties. CO₂ flux demonstrated a significant correlation with water volume under both conditions (light condition, p=0.008; dark condition, p=0.032) (Fig. S1). Light CH₄ flux significantly correlated with temperature (p=0.007) and water content (p=0.009) while dark CH₄ flux correlated with water content (p=0.043) and electrical conductivity (p=0.018). (Fig. S1)

Formatted: Subscript

Formatted: Subscript

300 Random forest modelling for CO₂ flux under light conditions yielded the maximum predictive power, with an R² value of 0.62, when using only 8 variables. Inclusion of any additional variables resulted in a model performance below the baseline

threshold ($R^2 \geq 0.6$, CV-score ≥ 0.4). Of the 8 variables, temperature is the most important single variable in correctly predicting CO₂ flux under light conditions with the feature importance of temperature exceeding 0.3, compared to all other variables in the model which have a feature importance below 0.2 (Fig. 5). Temporal variables (year and month) featured among the 8 selected variables, with the year of sampling being the second-most important variable in predicting CO₂ flux under light conditions (0.19 importance).

305




Figure 5: Importance of environmental, temporal, and ~~physiochemical~~ ~~physicochemical~~ -variables in predicting CO₂ flux under light conditions at the landward sampling site ($R^2=0.62$, cross-validation average (n=5) of 0.48 after feature engineering).

CO₂ flux under dark conditions was accurately predicted with the inclusion of 15 out of the 17 possible variables (Fig. 6).

310

The most important single variable in predicting CO₂ flux under dark conditions was $\delta^{13}\text{C-CH}_4$ (0.46 importance) by a large margin. $\delta^{13}\text{C-CH}_4$ averaged -47.5 ± 0.25 ‰ in dry conditions and -44.75 ± 1.2 ‰ under wet conditions, with a large range from -54.84 ‰ to -21.12 ‰. There was a negative correlation between $\delta^{13}\text{C-CH}_4$ and CO₂ flux in both dark (-0.2) and light (-0.22) conditions. As with the model for light CO₂ flux, the year of sampling was also the second most important feature in predicting dark CO₂ flux (0.14 importance).

315

~~The remaining 13 variables all had a feature importance below 0.1, but this combination contributed towards a good R^2 score of 0.63. Although the remaining 13 variables all had a feature importance below 0.1 this combination contributed towards an R^2 score of 0.63.~~

In both models the core replicate was of minor importance (Figs. 5 & 6). The season during sampling was not included in the random forest model due to its nature as a categorical variable and high collinearity with other variables. Instead, Kruskal-Wallis ANOVA showed that the season had significant relationships with temperature ($p < 0.001$), water volume in the cores during incubation ($p < 0.05$), dark CH_4 flux ($p < 0.05$) and light CH_4 flux ($p < 0.05$). However, GHG flux and soil water content (WC) were not significantly influenced by seasonality.




Figure 6: Importance of environmental, temporal, and physiochemical variables in predicting CO₂ flux under dark conditions at the landward sampling site ($R^2=0.63$, cross-validation average (n=5) of 0.43 after feature engineering).

4 Discussion

4.1 Small but highly variable GHG fluxes

The CO₂ and CH₄ fluxes reported in this study are, in general, a small source of GHG emissions, but include episodic events of high flux. The results fall within the lower range of fluxes previously reported, with CO₂ flux between -16,900 to 629,200 μmol CO₂ m⁻² d⁻¹, and CH₄ flux ranging from -2.1 to 25,974 μmol CH₄ m⁻² d⁻¹ (Sea et al., 2018). A review of 140 mangrove

studies reported the global average CO₂ flux of 56,800 ± 890 μmol CO₂ m⁻² d⁻¹ (Rosentreter et al., 2018b), while a CH₄ flux of 4,557.0 ± 1,102.1 μmol CH₄ m⁻² d⁻¹ was found across 54 mangrove studies, with a total of 110 flux observations (Al-Haj and Fulweiler, 2020). Our results for sea-air fluxes in particular are many orders of magnitude smaller than other studies with similar methodologies (Jacotot et al., 2018). Two defining characteristics of the soil in this study is the low C_{org} and high salinity, which may reduce CO₂ and CH₄, respectively (Ouyang et al., 2017; Poffenbarger et al., 2011).

*While comparisons can, and should, be drawn across different studies, the methodology of the study should be considered when interpreting results. For example, in-situ studies have the advantage of natural conditions with minimal disturbance caused by sampling whereas *ex-situ* studies, such as incubation techniques, allow for greater control of variables but typically cannot entirely replicate *in situ* conditions such as diel temperature variation, changes in light intensity and meteorological conditions (Toczydlowski et al., 2020; Sjögersten et al., 2018). For example, one study found mangrove ecosystem flux of CH₄ was the most variable on a daily basis due to meteorological variables and plant activities, both of which were excluded in this study (Liu et al., 2022). However, this study utilized incubations to maintain stringent control of environmental variables during the measurement period. The caveat of this approach is that it limits applicability to field conditions, but is useful in separating the effects of individual drivers of GHG flux variation from mangrove soil and minimising the number of confounding variables (Bond-Lamberty et al., 2016). An additional element of variation comes from different measurement techniques, as results can differ markedly between laser-based spectrometers, chamber-based systems, and eddy covariance measurements (Brannon et al., 2016; Podgrajsek et al., 2014). All studies compared in Table 3 are of *in situ* design, but there are a range of techniques and calculations used. These elements of variability complicate comparison across studies. There is often a large variation in GHG flux across studies and it should be considered whether this variation is due to environmental conditions or different study designs. For example, in the same study site, CH₄ fluxes from eddy covariance measurements have been lower than closed static chamber designs (Gnanamoorthy et al., 2022).*

Table 3: Comparative assessments of average mangrove fluxes under light conditions and standardised to μmol m⁻² d⁻¹ (± SE where data is available), unless otherwise specified. Fluxes for this study calculated using data from both sites. More comprehensive review papers can be found for CO₂ (Rosentreter et al., 2018b), and CH₄ (Al-Haj and Fulweiler, 2020).

Study	Location	CO ₂ flux (μmol CO ₂ m ⁻² d ⁻¹)	CH ₄ flux (μmol CH ₄ m ⁻² d ⁻¹)	Interface
This study	Central Red Sea, Saudi Arabia	5788 ± 1341	1.25 ± 0.34	Soil-air
This study	Central Red Sea, Saudi Arabia	3059 ± 679	3.67 ± 1.15	Sea-air
Das and Mandal,	Sundarbans, India	Range: 17,460 to	Range: 100 to 310	Soil-air

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Subscript

Formatted: Font: Italic

Formatted: Subscript

2022		70,000		
Hien et al., 2018	Northern Viet Nam	95,500 ± 89,280		Soil-air
Leopold et al., 2013	New Caledonia	91,800 ± 78,200		Soil-air
Chen et al., 2010	South China	Range: 560 to 20,560	Range: 10.1 to 5168.6	Soil-air
Kitpakornsanti et al., 2022	Thailand	62,160 ± 22,560	92.64 ± 48.24	Soil-air
Rosentreter et al., 2018a	Queensland estuary, Australia	156,900 ± 94,700		Soil-air & sea-air
Kitpakornsanti et al., 2022	Thailand	39,840 ± 15,840	59.28 ± 35.28	Sea-air
Akhand et al., 2021	Iriomote Island, Japan	2352 ± 2208 to 54,072 ± 50,976		Sea-air
Call et al., 2015	Queensland bay, Australia	Range: 9400 to 629,200	Range: 13.1 to 632.9	Sea-air
Chen et al., 2010	South China	Range: 560 to 20,560	Range: 10.10 to 5168.6	Soil-air
Bouillon et al., 2008	Global average	59,000 ± 52,000		Sea-air
Rosentreter et al., 2018b	Global estimate	56,800 ± 890		N/A
Al-Haj and Fulweiler, 2020	Global estimate		4557.0 ± 1,102.1	N/A

4.2 Drivers of flux variation

Although the landward and seaward study sites were within the same mangrove stand, there were considerably higher fluxes of CO₂ and CH₄ at the sea-air interface of the seaward site (43.7 g CO₂-eq m⁻² y⁻¹), compared to $-0.2 \text{ g CO}_2\text{-eq m}^{-2}\text{-y}^{-1}$ between the sea-air interface at the landward site ($-0.2 \text{ g CO}_2\text{-eq m}^{-2}\text{-y}^{-1}$). The main distinguishing environmental factor between the two sites appears to be the frequency and magnitude of tidal inundation as the landward site was microtidal, with long periods without tidal inundation. There is a strong semi-annual seasonal influence on tides in central Red Sea. Extremely hot summer months coincide with low mean sea-level states (Sultan et al., 1996), and in winter, the normally prevailing northwest winds are met by southeast winds, forming the Red Sea Convergence Zone in the centre of the Red Sea, resulting in higher mean sea levels (Langodan et al., 2017). This is supported by our analysis, showing the significance

($p < 0.05$) of season on the water volume captured in the soil cores during sampling. There was also a statistically significant seasonal influence on light and dark CH_4 flux. This seasonal effect is likely modulated by temperature variation, which proved to be an important element of light CO_2 flux in our random forest model. Additionally, higher temperatures in summer increase sub-surface soil temperature which can increase CH_4 emissions due to the temperature-dependency of microbial methanogens (Liu et al., 2020). The frequent absence of tidal inundation in summer exacerbates this effect as the high latent heat capacity of water could otherwise help regulate soil temperatures. Therefore, seasonality may exert a dual impact on methane emissions, explaining the significance.

A second important factor in CH_4 flux is salinity, measured by electrical conductivity in this study which demonstrated a lower mean in the seaward location. This may explain the higher CH_4 emissions from this site as salinity is reported to have a negative influence on CH_4 flux (Liu et al., 2020). Hypersaline mangrove environments are associated with low methane emissions (Cotovicz et al., 2024; Sea et al., 2018), because high salinity suppresses microbial activity and biogeochemical processes, reducing GHG cycling (Zhu et al., 2021). There is a proposed salinity threshold of 18 ppt, where CH_4 flux may become negligible which is significantly below the salinity found in the Red Sea (Alhassan and Aljhdali, 2021; Poffenbarger et al., 2011). ~~The causes for the large differences in GHG flux between sites within the same mangrove stand are not fully resolved. The causes for the large differences in GHG flux between sites within the same mangrove stand are not fully resolved, although it is likely that there is microscale variation due, in part, to different gas transport processes. The release of CH_4 from the soil via ebullition has particularly high spatial variability within sampling sites (Baulch, 2011; Chuang et al., 2017). Furthermore, the episodic nature of ebullition events may distort the flux calculation which assumes a linear concentration change over time, as is the case with diffusive flux (Jacotot et al., 2018). The possibility of active ebullition in saline, undisturbed mangrove ecosystems require further investigation, as to-date, no study has found ebullition to be a significant pathway of CH_4 release under these conditions (Cotovicz et al., 2024). Considering this small-scale variability, it is important to emphasise the need for comprehensive assessments in individual mangrove ecosystems as GHG flux is highly site-specific. However, it is important to emphasise the need for comprehensive assessments to determine the true magnitude of GHG flux in a given mangrove ecosystem considering this small-scale variability.~~ Commonly, spatial variation in GHG fluxes is inferred from a few plots within the study site (Castillo et al., 2017). However, this ~~is~~ method is likely to result in larger errors in estimates without attempting to determine factors driving this variation.

As evidenced by the monthly and site-specific flux variation, environmental and soil physio-chemical factors are important in regulating mangrove soil GHG fluxes. In the literature, there are a multitude of variables suggested to influence CO_2 and CH_4 flux from mangrove soils. The variables reported to affect CO_2 fluxes include soil organic carbon (SOC), nitrogen, phosphate, iron, ammonium, porosity, and tidal range (Chen et al., 2010; Jacotot et al., 2018; Sugiana et al., 2023; Wang et al., 2016). The variables reported to affect CH_4 flux include SOC, ammonium, porewater salinity, redox potential, soil temperature, air temperature and tidal range (Allen et al., 2007; Chen et al., 2010; Jacotot et al., 2018; Sugiana et al., 2023;

Formatted: Subscript

400 Wang et al., 2016). Furthermore, additional factors have been suggested as general influences on overall mangrove GHG
flux such as temperature fluctuations, soil moisture content, soil grain size, and tidal patterns (Hien et al., 2018; Ouyang
et al., 2017). Many of these factors are inferred by a correlational relationship with GHG flux, with many variables likely to be
colinear making causality difficult to determine. ~~Additionally, diverse methodologies for equipment and measurements make
it challenging to draw generalised conclusions and comparisons.~~

405 An advantage of the random forest algorithm is that it allows many variables to be taken into account, with the ability to
uncover non-linear relationships, its resistance to outliers, and the ability to test the model on other datasets (Smorkalov,
2022). ~~However, there were variables mentioned above that were found to be important in GHG flux in other studies but
were not measured in this study, for example, ammonium, iron, and soil grain size. There are limitations on the number of
410 variables relative to a fairly small number of observations as in this study (Kiers and Smilde, 2007), along with practical
limitations of time and resources. There is substantial scope in future research to comprehensively investigate more variables
than those reported here over a longer sampling period, or with more frequent observations. An analysis of a greater number
of chemical and physical characteristics of the soil beyond carbon and nitrogen would be particularly relevant for GHG flux
(Nóbrega et al., 2014; Chen et al., 2010). This limitation must be acknowledged when interpreting our results as there may
415 have been significantly important factors which were not measured and thus not considered in our analysis of the most
important drivers of GHG flux.~~

The random forest modelling we conducted suggested temperature to be the most important factor in predicting light CO₂
flux and δ¹³C-CH₄ to be the most important factor in accurately predicting dark CO₂ flux. To the best of our knowledge, the
420 isotopic signature of methane is a variable that has not been previously suggested as an important predictor of mangrove CO₂
flux. ~~Notably, there was no statistically significant correlation between δ¹³C-CH₄ and dark CO₂ flux, contrary to the random
forest model, suggesting this finding may be a result of overfitting from the random forest model or there may be more
complex non-linear relationships uncovered by machine learning which are not detected by simple correlation. This may be a
feature of random forest overfitting or that the δ¹³C-CH₄ signature masks other variables which are the true driving influence
425 of CO₂.~~

However, the mean δ¹³C-CH₄ at the landward and seaward sites (-47.2‰ and -48.1‰) were considerably less enriched than
the -80.6‰ δ¹³C-CH₄ found in a similar study on Red Sea mangrove GHG flux (Sea et al., 2018). The difference may be due
to a number of factors including methanogenesis or oxidisation, although both factors are unlikely to directly influence CO₂
430 emissions. A previous study has found mangroves with the lightest δ¹³C-CO₂ and δ¹³C-CH₄ to have the lowest CO₂ flux
further suggesting there may be a link between δ¹³C-CH₄ and CO₂ flux (Sea et al., 2018). Variations in δ¹³C-CH₄ is highly
likely to be driven by microbial processes, for example, methanotrophic bacteria which oxidize a fraction of total CH₄
production resulting in a more positive δ¹³C-CH₄. A range of -65 ‰ to -50 ‰, similar to this study, found that aceticlastic

methanogenesis (produced from acetate) dominates (Ouyang et al., 2024; Teh et al., 2005). Additionally, in a previous study
435 on mangrove forests in Mexico it was found that 30–70% of the total CO₂ measured was produced by methanogenesis
(Sanchez-Carrillo et al., 2021). Anaerobic oxidation can also form CO₂ (Shukla et al., 2013). These are possible explanations
for our results demonstrating high importance of $\delta^{13}\text{C}$ -CH₄ as a predictor of CO₂ flux. ~~however investigation of the soil
microbial community would be necessary to draw solid conclusions on the mechanisms underlying this statistical
relationship. However, to better understand the origin and fate of CH₄ from mangrove soils, methanogenesis should be
440 studied directly through the determination of $\delta^{13}\text{C}$ of the methyl group of acetate (Govert and Conrad, 2009) or an isotope
mass-balance approach (Sánchez-Carrillo et al., 2021), along with an investigation of the soil microbial community.~~

Formatted: Superscript

From our random forest models, the most important soil variables for CO₂ flux were C:N, and TC for light conditions, and
soil water content (WC-3cm), and water volume for dark conditions. All factors have previously been documented to play a
445 role in CO₂ emissions (Chen et al., 2010). Preservation of TC is related to factors such as water level and inundation time,
and where low OC burial efficiency increases soil respiration (Breithaupt et al., 2019). C:N is a good predictor of soil
microbial respiration (Fang and Moncrieff, 2005), and has previously been found to have a significant positive correlation
with mangrove CO₂ flux (Hien et al., 2018). Furthermore, soil respiration exhibits diurnal patterns which may explain the
high importance of carbon and nitrogen concentrations in predicting light CO₂ flux but not dark CO₂ flux (Jin et al., 2013).
450 C:N may also be a good predictor for CO₂ flux variability because of its relationship with the labile carbon pool, influenced
by microbial biomass which will vary by month and season depending on the suitability of conditions for microbial growth
(Padhy et al., 2020). Secondly, soil water content has been found to exert a negative influence on CO₂ flux, but have positive
relationship with C_{org} (Ouyang et al., 2017). However, there is also likely to be covariation among water content and
variables not measured in this study, such as soil porosity, grain size, and density of crab burrows which can increase CO₂
455 flux (Booth et al., 2019; Ouyang et al., 2017). This implies that the interpretation of GHG flux variability should be carefully
considered to ensure that non-linear relationships between multiple interrelated variables are accounted for.

In both models, ~~the year had the second-highest predictive importance. There are a few theories for the importance of this
factor. The growth and flowering cycles of *A. marina* mangroves in the Red Sea are not annual (Almahasheer et al., 2016). In
460 theory, increased growth over a given year may result in increased soil carbon pools for microbial respiration, directly
impacting GHG flux. However, this cannot be tested as mangrove growth was not measured in the present study.
Alternatively, the importance of the year of sampling may be artificially inflated in our models due to the presence of water
during 4 of the 5 months sampled in 2021 while subsequent years were dominated by dry sampling conditions. However,
there were also climatic variables and extreme weather patterns for the region across the 3-year period. Central Saudi Arabia
465 experienced widespread greening in 2023 due to higher-than-average rainfall (Van Dijk et al., 2023), potentially also
facilitating mangrove growth. It is likely that a combination of these 3 factors explain the predictive importance of the
sampling year, and emphasise the importance of long-term flux measurements to capture variations resulting from climatic~~

Formatted: Font: Italic

470 changes, and perennial life-cycles of mangroves. Also, in both models, the core was the of minor importance in predicting
CO₂ flux, which shows good replicability across the four cores sampled each month. This is supported by the correlation
analysis, where the core replicate had no significant relationships with CO₂ or CH₄ flux under any conditions. However, there
were significant relationships between the core and soil physicochemical properties such as C_{org}, TN, and δ¹³C-CO₂ (Fig.
S1). This is likely due to microscale differences in the deposition of organic matter, and microbial communities, which is an
element of natural variation in response to environmental conditions (Padhy et al., 2020). ~~core replicates and inorganic~~
carbon (C_{inorg} 3cm) with δ¹³C-CO₂, and well as between electrical conductivity (EC_{1:5}) and δ¹³C-CH₄ (Fig. S1).

475 The random forest model for the CO₂ dark condition had a ~~good~~ R² score comparable to models previously used to predict
SOC stock (Moreno Muñoz et al., 2024), while maintaining the majority of variables (15 of 17), with a comparable
performance with models published in previous studies to predict SOC stock (Moreno Muñoz et al., 2024). However, based
480 on the R² metric, the model for light CO₂ flux performed poorly on a higher number of variables suggesting that many of
these variables simply added ‘noise’ to the predictions, without adding predictive power (Fox et al., 2017). It is likely the
models’ performance, particularly for light CO₂ could be improved with the addition of other unmeasured factors such as
clay or sulphur content which were found to be important predictors of soil CO₂ flux in sugarcane with random forest
modelling (Tavares et al., 2018). The CH₄ flux was not modelled due to the importance of microbial activities in CH₄
cycling, which would not be accurately captured by the variables measured in this study (Das et al., 2018; Liu et al., 2020;
485 Yu et al., 2020).

4.3 Implications for mangrove carbon budget

Despite the small magnitude of fluxes reported in this study compared to global estimates, they deserve consideration in the
net carbon sequestration of Red Sea mangroves given their low carbon burial rate (Almahasheer et al., 2017). The carbon
sequestration offset by the CO₂-eq of the combined CO₂ and CH₄ fluxes measured in this study ranged between -130% and
490 822%. A negative CO₂-eq implies net GHG removal from the atmosphere. There was an important difference between the
mean and median offset of carbon sequestration by the combined CO₂ and CH₄ fluxes, which were 94.5% and 4.9%,
respectively. The median estimate is less affected by extreme values, and is, therefore, more representative of the central
tendency of the offset, while the mean estimate fully captures the large variability in the long-term dataset of this study.
Previous studies have also highlighted extreme variability where global mean emissions of CH₄ flux were ~16 times higher
495 than the median estimate (Al-Haj and Fulweiler, 2020). Highly skewed data is appropriate to use only if it accurately
reflecting the true distribution of fluxes and not sampling bias (Rosentreter and Williamson, 2020). In this study, averages
are likely to be an accurate statistic, given the controls on sampling location and consistent samples times each month over
the full study period. This means that whereas the combined CO₂ and CH₄ fluxes were relatively small compared to reported
mean organic carbon sequestration by the Red Sea mangrove stands studied, these are subject to occasional large emissions
500 that offset much of the carbon removed.

Prior studies have found GHG emissions to offset between 9.3 to 32.7% of the organic carbon sequestration of mangrove forests (Chen et al., 2016). A large component of this variability is dependent whether fluxes are measured between the sea-air or soil-air interface (Table 3). CO₂ emissions, which are the biggest contributor to CO₂-eq emissions are greatly affected when being measured between the sea-air or soil-air interface. When CO₂ is released from the soil into the water column it enters the carbonate system and can be converted to bicarbonate or carbonate ions (Zeebe and Wolf-Gladrow, 2001). As a result, the majority of CO₂ emitted from the soil undergoes dissolution in the water column before it is released to the air. This explains the lower CO₂-eq from the sea-air interface of -0.4 g CO₂-eq m⁻² y⁻¹ compared to 172.1 g CO₂-eq m⁻² y⁻¹ for the soil-air interface when soils are directly exposed to air. Typically, when fluxes are measured from the sea-air interface, equilibration equations are used to account for the changes in carbonate chemistry in the seawater (Akhand et al., 2021; Call et al., 2015). However, the aim of this study was to compare GHG flux to the air between interfaces, so the calculations used here only consider linear changes in concentration across timepoints emphasising diffusive fluxes to the atmosphere over other methods of gas transfer such as bubble ebullition (Jacotot et al., 2018). Overall, the CO₂-eq released to the atmosphere is a significant offset to carbon burial given the carbon burial rate of Red Sea mangroves is just 55 g CO₂-eq m⁻² y⁻¹ (15 g C_{org} m⁻² y⁻¹), over 10-fold lower than the global average of 598 g CO₂-eq m⁻² y⁻¹ (163 g C_{org} m⁻² y⁻¹) (Almahasheer et al., 2017; Breithaupt et al., 2012).

However, in the Red Sea region mangrove soils have a high carbonate content, our estimates of C_{inorg} fall within the upper range of previously reported figures for Red Sea mangroves, which are higher than global average estimates (Garcias-Bonet et al., 2019; Saderne et al., 2019). Furthermore, mangrove soil in the same location as the present study has 76% ± 2% (dry weight) CaCO₃, which is attributed to their growth on underlying carbonate platforms formed by Pleistocene coral reefs (Saderne et al., 2018). As a result, there is an additional factor to consider; the role of total alkalinity (TA) enhancement from carbonate dissolution in the mangrove soils, which increases the capacity for seawater to absorb CO₂ from the atmosphere (Alongi, 2022; Saderne et al., 2019). Mangroves in the Red Sea are characterized as important TA sources (Saderne et al., 2019), which are driven by high metabolic activity in their soil and multi-stage biogeochemical processes such as carbonate dissolution, denitrification, sulfate reduction, and ammonification (Baldry et al., 2020; Saderne et al., 2021; Sippo et al., 2016).

CaCO₃ dissolution is particularly relevant to the central Red Sea, as one mole of dissolved CaCO₃ results in the uptake of 0.6 mol of atmospheric CO₂ (Frankignoulle et al., 1995). The dissolution of the large CaCO₃ pool in the soils of Red Sea mangroves present a substantial additional carbon sink, exceeding by 23-fold the C_{org} burial rate for the central Red Sea (Almahasheer et al., 2017; Saderne et al., 2021). Although the lower carbon burial in soil means GHG fluxes are a large offset to the soil carbon burial, TA enhancement brings the carbon sink value of the mangrove stand in this study to 360 g C m⁻² yr⁻¹, which is 2.2-fold above global mean mangrove C_{org} (Saderne et al., 2021). In the present study the CO₂-eq of GHG

535 fluxes represent a small offset (3.9% on average) to the combined carbon sequestration of this mangrove stand when
accounting for carbon burial and TA enhancement combined.

5 Conclusion

The long-term flux variability captured in this study provides valuable insights into the role of GHG flux in offsetting carbon
burial in Red Sea mangrove soils. Our study involved an improved temporal resolution, in terms of the overall duration and
540 frequency of assessments, beyond most previous assessments. This is important because our results show that CO₂ and CH₄
fluxes are typically a small carbon offset compared to carbon burial in soils but is punctuated with episodic GHG emission
bursts that suffice to offset a large fraction of carbon burial. This aspect of GHG flux dynamics may be missed by studies
with poorer temporal coverage.

When considering the carbon budget of the Central Red Sea mangrove stand considered in this study, our results show the
545 overriding importance of TA enhancement from carbonate dissolution, which is emerging as a major component of
mangrove CO₂ removal, not yet captured in blue carbon projects. Our results also showed the direct exposure of mangrove
soils to the atmosphere drastically enhances GHG emissions compared to emissions during tidal flooding. Environmental
conditions helped explain variability in CO₂ emissions, whereas those in CH₄ emissions seem to be dominated by the
dynamics of the microbial community responsible for methanogenesis and methane oxidation.

550

Code and data availability

555 All data to support the findings of this study are available in FigShare. Raw data for the landward site available at:
10.6084/m9.figshare.26085898. Combined site data across sea-air and soil-air interfaces available at:
10.6084/m9.figshare.26085928. Code and associated data for Random Forest algorithms available at:
10.6084/m9.figshare.26085940.

560 *Competing interests.*

The authors declare that they have no conflict of interest.

Author contribution

565

C.M.D conceived the research. A.S, C.F, J.B, and C.M.D designed the study. A.S, C.F, and J.B performed the field sampling. A.S, C.F, J.B, and M.E performed laboratory work. M.E performed gas measurements. A.S, C.F, and J.B conducted data analysis. J.B produced display figures and wrote the manuscript. All authors contributed and approved the manuscript.

570

Acknowledgements

This research was funded by King Abdullah University of Science and Technology through baseline (BAS/1/1071- 01-01) 495 funding for CMD. We extend thanks to Asalata Kotikalapudi for help with sample processing, Reny P. Devassy,

575

Jennifer Thompson, Naira Pluma, Elisa Laiolo and Anastasiia Martynova for their assistance with fieldwork. We thank Larissa Frühe for her assistance with R plots.

References

- 580 Akhand, A., Watanabe, K., Chanda, A., Tokoro, T., Chakraborty, K., Moki, H., Tanaya, T., Ghosh, J., and Kuwae, T.: Lateral carbon fluxes and CO₂ evasion from a subtropical mangrove-seagrass-coral continuum, *Sci. Total Environ.*, 752, 142190, <https://doi.org/10.1016/j.scitotenv.2020.142190>, 2021.
- Al-Haj, A. N. and Fulweiler, R. W.: A synthesis of methane emissions from shallow vegetated coastal ecosystems, *Global Change Biology*, 26, 2988–3005, <https://doi.org/10.1111/gcb.15046>, 2020.
- 585 Alhassan, A. B. and Aljahdali, M. O.: Nutrient and physicochemical properties as potential causes of stress in mangroves of the central Red Sea, *PLOS ONE*, 16, e0261620, <https://doi.org/10.1371/journal.pone.0261620>, 2021.
- Allen, D. E., Dalal, R. C., Rennenberg, H., Meyer, R. L., Reeves, S., and Schmidt, S.: Spatial and temporal variation of nitrous oxide and methane flux between subtropical mangrove sediments and the atmosphere, *Soil Biology and Biochemistry*, 39, 622–631, <https://doi.org/10.1016/j.soilbio.2006.09.013>, 2007.
- 590 Almahasheer, H., Aljowair, A., Duarte, C. M., and Irigoien, X.: Decadal stability of Red Sea mangroves, *Estuarine, Coastal and Shelf Science*, 169, 164–172, <https://doi.org/10.1016/j.ecss.2015.11.027>, 2016a.
- Almahasheer, H., Duarte, C. M., and Irigoien, X.: Nutrient Limitation in Central Red Sea Mangroves, *Frontiers in Marine Science*, 3, 2016b.
- Almahasheer, H., Serrano, O., Duarte, C. M., Arias-Ortiz, A., Masque, P., and Irigoien, X.: Low Carbon sink capacity of Red Sea mangroves, *Sci Rep*, 7, 9700, <https://doi.org/10.1038/s41598-017-10424-9>, 2017.
- Alongi, D.: Impacts of Climate Change on Blue Carbon Stocks and Fluxes in Mangrove Forests, *FORESTS*, 13, <https://doi.org/10.3390/f13020149>, 2022.
- Baldry, K., Saderne, V., McCorkle, D., Churchill, J., Agust, S., and Duarte, C.: Anomalies in the carbonate system of Red Sea coastal habitats, *BIOGEOSCIENCES*, 17, 423–439, <https://doi.org/10.5194/bg-17-423-2020>, 2020.
- 600 [Baulch, H. M., Dillon, P. J., Maranger, R., and Schiff, S. L.: Diffusive and ebullitive transport of methane and nitrous oxide from streams: Are bubble-mediated fluxes important?. *Journal of Geophysical Research: Biogeosciences*, 116\(G4\), <https://doi.org/10.1029/2011JG001656>, 2011.](https://doi.org/10.1029/2011JG001656)
- Blanco-Sacristán, J., Johansen, K., Duarte, C. M., Daffonchio, D., Hoteit, I., and McCabe, M. F.: Mangrove distribution and afforestation potential in the Red Sea, *Science of The Total Environment*, 843, 157098, <https://doi.org/10.1016/j.scitotenv.2022.157098>, 2022.
- [Bond-Lamberty, B., Smith, A. P., and Bailey, V.: Temperature and moisture effects on greenhouse gas emissions from deep active-layer boreal soils. *Biogeosciences*, 13\(24\), 6669-6681, <https://doi.org/10.5194/bg-13-6669-2016>, 2016.](https://doi.org/10.5194/bg-13-6669-2016)
- 610 Booth, J. M., Fusi, M., Marasco, R., Mbobo, T., and Daffonchio, D.: Fiddler crab bioturbation determines consistent changes in bacterial communities across contrasting environmental conditions, *Sci Rep*, 9, 3749, <https://doi.org/10.1038/s41598-019-40315-0>, 2019.

Formatted: Left

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Default Paragraph Font, Font: (Default) +Body (Times New Roman), 10 pt, Not Bold

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 10 pt, Not Bold

- 615 Bouillon, S., Borges, A. V., Castañeda-Moya, E., Diele, K., Dittmar, T., Duke, N. C., Kristensen, E., Lee, S. Y., Marchand, C., Middelburg, J. J., Rivera-Monroy, V. H., Smith III, T. J., and Twilley, R. R.: Mangrove production and carbon sinks: A revision of global budget estimates, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2007GB003052>, 2008.
- Brannon, E. Q., Moseman-Valtierra, S. M., Rella, C. W., Martin, R. M., Chen, X., and Tang, J.: Evaluation of laser-based spectrometers for greenhouse gas flux measurements in coastal marshes, *Limnology and Oceanography: Methods*, 14, 466–476, <https://doi.org/10.1002/lom3.10105>, 2016.
- 620 ~~Bre~~Breavington, J., Steckbauer, A., Fu, C., Ennasri, M., Duarte, C. M.: Landward site raw data, Figshare, available at: [10.6084/m9.figshare.26085898](https://doi.org/10.6084/m9.figshare.26085898), last access: 23 June 2024.
- Breavington, J., Steckbauer, A., Fu, C., Ennasri, M., Duarte, C. M.: Landward and seaward fluxes and soil properties, Figshare, available at: [10.6084/m9.figshare.26085928](https://doi.org/10.6084/m9.figshare.26085928), last access: 23 June
- Breavington, J., Steckbauer, A., Fu, C., Ennasri, M., Duarte, C. M.: Random forest variable importance, Figshare, available at: [10.6084/m9.figshare.26085940](https://doi.org/10.6084/m9.figshare.26085940), last access: 23 June 2024..
- 625 Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Breithaupt, J. L., Smoak, J. M., Smith III, T. J., Sanders, C. J., and Hoare, A.: Organic carbon burial rates in mangrove sediments: Strengthening the global budget, *Global Biogeochemical Cycles*, 26, <https://doi.org/10.1029/2012GB004375>, 2012.
- 630 Breithaupt, J. L., Smoak, J. M., Sanders, C. J., and Troxler, T. G.: Spatial Variability of Organic Carbon, CaCO₃ and Nutrient Burial Rates Spanning a Mangrove Productivity Gradient in the Coastal Everglades, *Ecosystems*, 22, 844–858, <https://doi.org/10.1007/s10021-018-0306-5>, 2019.
- Call, M., Maher, D. T., Santos, I. R., Ruiz-Halpern, S., Mangion, P., Sanders, C. J., Erler, D. V., Oakes, J. M., Rosentreter, J., Murray, R., and Eyre, B. D.: Spatial and temporal variability of carbon dioxide and methane fluxes over semi-diurnal and spring–neap–spring timescales in a mangrove creek, *Geochimica et Cosmochimica Acta*, 150, 211–225, <https://doi.org/10.1016/j.gca.2014.11.023>, 2015.
- 635 Castillo, J., Apan, A., Maraseni, T., and Salmo, S.: Soil greenhouse gas fluxes in tropical mangrove forests and in land uses on deforested mangrove lands, *CATENA*, 159, 60–69, <https://doi.org/10.1016/j.catena.2017.08.005>, 2017.
- Chen, G., Chen, B., Yu, D., Tam, N. F. Y., Ye, Y., and Chen, S.: Soil greenhouse gas emissions reduce the contribution of mangrove plants to the atmospheric cooling effect, *Environmental Research Letters*, 11, 124019, <https://doi.org/10.1088/1748-9326/11/12/124019>, 2016.
- 640 Chen, G. C., Tam, N. F. Y., and Ye, Y.: Summer fluxes of atmospheric greenhouse gases N₂O, CH₄ and CO₂ from mangrove soil in South China, *Science of The Total Environment*, 408, 2761–2767, <https://doi.org/10.1016/j.scitotenv.2010.03.007>, 2010.
- 645 [Chuang, P. C., Young, M. B., Dale, A. W., Miller, L. G., Herrera-Silveira, J. A., and Paytan, A.: Methane fluxes from tropical coastal lagoons surrounded by mangroves, Yucatán, Mexico. *Journal of Geophysical Research: Biogeosciences*, 122\(5\), 1156-1174 <https://doi.org/10.1002/2017JG003761>, 2017.](https://doi.org/10.1002/2017JG003761)

- Cotovicz, L. C., Abril, G., Sanders, C. J., Tait, D. R., Maher, D. T., Sippo, J. Z., Holloway, C., Yau, Y. Y. Y., and Santos, I. R.: Methane oxidation minimizes emissions and offsets to carbon burial in mangroves, *Nat. Clim. Chang.*, 14, 275–281, <https://doi.org/10.1038/s41558-024-01927-1>, 2024.
- 650 Curran, S., Kumar, A., Lutz, W., and Williams, M.: Interactions between Coastal and Marine Ecosystems and Human Population Systems: Perspectives on How Consumption Mediates This Interaction, *Ambio*, 31, 264–268, 2002.
- Das, N. and Mandal, S.: Microbial populations regulate greenhouse gas emissions in Sundarban mangrove ecosystem, India, *Acta Ecologica Sinica*, 42, 641–652, <https://doi.org/10.1016/j.chnaes.2021.07.011>, 2022.
- 655 Das, S., Ganguly, D., Chakraborty, S., Mukherjee, A., and De, T.: Methane flux dynamics in relation to methanogenic and methanotrophic populations in the soil of Indian Sundarban mangroves, *MARINE ECOLOGY-AN EVOLUTIONARY PERSPECTIVE*, 39, <https://doi.org/10.1111/maec.12493>, 2018.
- Duarte, C., Losada, I., Hendriks, I., Mazarrasa, I., and Marba, N.: The role of coastal plant communities for climate change mitigation and adaptation, *NATURE CLIMATE CHANGE*, 3, 961–968, <https://doi.org/10.1038/NCLIMATE1970>, 2013.
- 660 Fang, C. and Moncrieff, J. B.: The variation of soil microbial respiration with depth in relation to soil carbon composition, *Plant Soil*, 268, 243–253, <https://doi.org/10.1007/s11104-004-0278-4>, 2005.
- Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., van Dorland, R., Bodeker, G., Boucher, O., Collins, W. D., Conway, T. J., Dlugokencky, E., Elkins, J. W., Etheridge, D., Foukal, P., Fraser, P., Geller, M., Joos, F., Keeling, C. D., Kinne, S., Lassey, K., Lohmann, U., Manning, A. C., Montzka, S., Oram, D., O’Shaughnessy, K., Piper, S., Plattner, G.-K., Ponater, M., Ramankutty, N., Reid, G., Rind, D., Rosenlof, K., Sausen, R., Schwarzkopf, D., Solanki, S. K., Stenchikov, G., Stuber, N., Takemura, T., Textor, C., Wang, R., Weiss, R., and Whorf, T.: Changes in Atmospheric Constituents and in Radiative Forcing, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the 4th Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, USA, 2007.
- 670 Fox, E. W., Hill, R. A., Leibowitz, S. G., Olsen, A. R., Thornbrugh, D. J., and Weber, M. H.: Assessing the accuracy and stability of variable selection methods for random forest modeling in ecology, *Environ Monit Assess*, 189, 316, <https://doi.org/10.1007/s10661-017-6025-0>, 2017.
- 675 Frankignoulle, M., Pichon, M., and Gattuso, J.-P.: Aquatic Calcification as a Source of Carbon Dioxide, in: *Carbon Sequestration in the Biosphere*, Berlin, Heidelberg, 265–271, https://doi.org/10.1007/978-3-642-79943-3_18, 1995.
- Gabr, S. S., Morsy, E. A., El Bastawesy, M. A., Habeebullah, T. M., and Shaaban, F. F.: Exploration of potential groundwater resources at Thuwal area, north of Jeddah, Saudi Arabia, using remote sensing data analysis and geophysical survey, *Arab J Geosci*, 10, 509, <https://doi.org/10.1007/s12517-017-3295-3>, 2017.
- 680 Gnanamoorthy, P., Chakraborty, S., Nagarajan, R., Ramasubramanian, R., Selvam, V., Burman, P.K.D., Sarathy, P.P., Zeeshan, M., Song, Q. and Zhang, Y.: Seasonal variation of methane fluxes in a mangrove ecosystem in south India: An eddy covariance-based approach. *Estuaries and Coasts*, 1-16, <https://doi.org/10.1007/s12237-021-00988-1>, 2022.
- 685 Garcias-Bonet, N., Delgado-Huertas, A., Carrillo-de-Albornoz, P., Anton, A., Almahasheer, H., Marbà, N., Hendriks, I. E., Krause-Jensen, D., and Duarte, C. M.: Carbon and Nitrogen Concentrations, Stocks, and Isotopic Compositions in Red Sea Seagrass and Mangrove Sediments, *Frontiers in Marine Science*, 6, 2019.

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Genuer, R., Poggi, J.-M., and Tuleau-Malot, C.: Variable selection using random forests, *Pattern Recognition Letters*, 31, 2225–2236, <https://doi.org/10.1016/j.patrec.2010.03.014>, 2010.

690 [Goevert, D., and Conrad, R.: Effect of substrate concentration on carbon isotope fractionation during acetoclastic methanogenesis by *Methanosarcina barkeri* and *M. acetivorans* and in rice field soil. *Applied and Environmental Microbiology*, 75\(9\), 2605-2612, <https://doi.org/10.1128/AEM.02680-08>, 2009.](#)

Hardie, M. and Doyle, R.: Measuring Soil Salinity, in: *Plant Salt Tolerance: Methods and Protocols*, edited by: Shabala, S. and Cuin, T. A., Humana Press, Totowa, NJ, 415–425, https://doi.org/10.1007/978-1-61779-986-0_28, 2012.

695 Hien, H. T., Marchand, C., Aimé, J., and Cuc, N. T. K.: Seasonal variability of CO₂ emissions from sediments in planted mangroves (Northern Viet Nam), *Estuarine, Coastal and Shelf Science*, 213, 28–39, <https://doi.org/10.1016/j.ecss.2018.08.006>, 2018.

Howard, J., Hoyt, S., Isensee, K., Telszewski, M., and Pidgeon, E.: *Coastal blue carbon: methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrasses*, 2014.

700 IPCC, Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., Church, J. A., Clarke, L., Dahe, Q., Dasgupta, P., Dubash, N. K., Edenhofer, O., Elgizouli, I., Field, C. B., Forster, P., Friedlingstein, P., Fuglestvedt, J., Gomez-Echeverri, L., Hallegatte, S., Hegerl, G., Howden, M., Jiang, K., Jimenez Cisneroz, B., Kattsov, V., Lee, H., Mach, K. J., Marotzke, J., Mastrandrea, M. D., Meyer, L., Minx, J., Mulugetta, Y., O'Brien, K., Oppenheimer, M., Pereira, J. J., Pichs-Madruga, R., Plattner, G.-K., Pörtner, H.-O., Power, S. B., Preston, B., Ravindranath, N. H., Reisinger, A., Riahi, K., Rusticucci, M., Scholes, R., Seyboth, K., Sokona, Y., Stavins, R., Stocker, T. F., Tschakert, P., van Vuuren, D., and van Ypersele, J.-P.: IPCC, 2014: *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Pachauri, R. K. and Meyer, L., IPCC, Geneva, Switzerland, 151 pp., 2014.

710 Jacotot, A., Marchand, C., and Allenbach, M.: Tidal variability of CO₂ and CH₄ emissions from the water column within a *Rhizophora* mangrove forest (New Caledonia), *Science of The Total Environment*, 631–632, 334–340, <https://doi.org/10.1016/j.scitotenv.2018.03.006>, 2018.

Jin, L., Lu, C.-Y., Ye, Y., and Ye, G.-F.: Soil Respiration in a Subtropical Mangrove Wetland in the Jiulong River Estuary, China, *Pedosphere*, 23, 678–685, [https://doi.org/10.1016/S1002-0160\(13\)60060-0](https://doi.org/10.1016/S1002-0160(13)60060-0), 2013.

715 Kargas, G., Chatzigiakoumis, I., Kollias, A., Spiliotis, D., and Kerkides, P.: An Investigation of the Relationship between the Electrical Conductivity of the Soil Saturated Paste Extract ECE with the Respective Values of the Mass Soil/Water Ratios 1:1 and 1:5 (EC1:1 and EC1:5), *Proceedings*, 2, 661, <https://doi.org/10.3390/proceedings2110661>, 2018.

[Khalil, A. S.: Mangroves of the red sea. *The Red Sea: The formation, morphology, oceanography and environment of a young ocean basin*, 585-597, \[https://doi.org/10.1007/978-3-662-45201-1_33\]\(https://doi.org/10.1007/978-3-662-45201-1_33\), 2015.](#)

720 [Kiers, H. A., and Smilde, A. K.: A comparison of various methods for multivariate regression with highly collinear variables. *Statistical Methods and Applications*, 16, 193- 228, <https://doi.org/10.0017/s10260-006-0025-5>, 2007.](#)

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Formatted: Line spacing: single

725 Kitpakornsanti, K., Pengthamkeerati, P., Limsakul, A., Worachananant, P., and Diloksumpun, S.: Greenhouse gas emissions from soil and water surface in different mangrove establishments and management in Ranong Biosphere Reserve, Thailand, *Regional Studies in Marine Science*, 56, 102690, <https://doi.org/10.1016/j.rsma.2022.102690>, 2022.

Langodan, S., Cavaleri, L., Vishwanadhapalli, Y., Pomaro, A., Bertotti, L., and Hoteit, I.: The climatology of the Red Sea - part 1: the wind: THE WIND CLIMATOLOGY OF THE RED SEA, *Int. J. Climatol*, 37, 4509–4517, <https://doi.org/10.1002/joc.5103>, 2017.

730 Leopold, A., Marchand, C., Deborde, J., Chaduteau, C., and Allenbach, M.: Influence of mangrove zonation on CO₂ fluxes at the sediment–air interface (New Caledonia), *Geoderma*, 202–203, 62–70, <https://doi.org/10.1016/j.geoderma.2013.03.008>, 2013.

735 Liu, J., Zhou, Y., Valach, A., Shortt, R., Kasak, K., Rey-Sanchez, C., Hemes, K. S., Baldocchi, D., and Lai, D. Y. F.: Methane emissions reduce the radiative cooling effect of a subtropical estuarine mangrove wetland by half, *Glob Change Biol*, 26, 4998–5016, <https://doi.org/10.1111/gcb.15247>, 2020.

[Liu, J., Valach, A., Baldocchi, D., & Lai, D. Y.: Biophysical controls of ecosystem-scale methane fluxes from a subtropical estuarine mangrove: Multiscale, nonlinearity, asynchrony and causality. *Global Biogeochemical Cycles*, 36\(6\), e2021GB007179, <https://doi.org/10.1029/2021GB007179>, 2022.](https://doi.org/10.1029/2021GB007179)

740 Lovelock, C. E., Barbier, E., and Duarte, C. M.: Tackling the mangrove restoration challenge, *PLOS Biology*, 20, e3001836, <https://doi.org/10.1371/journal.pbio.3001836>, 2022.

[Mackey, A.: Biomass of the mangrove *Avicennia marina* \(Forsk.\) Vierh. near Brisbane, south-eastern Queensland. *Marine and Freshwater Research* 44, 721–725, <https://doi.org/10.1071/MF9930721>, 1993.](https://doi.org/10.1071/MF9930721)

745 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A Reconstruction of Regional and Global Temperature for the Past 11,300 Years, *Science*, 339, 1198–1201, <https://doi.org/10.1126/science.1228026>, 2013.

Moreno Muñoz, A. S., Guzmán Alvis, Á. I., and Benavides Martínez, I. F.: A random forest model to predict soil organic carbon storage in mangroves from Southern Colombian Pacific coast, *Estuarine, Coastal and Shelf Science*, 299, 108674, <https://doi.org/10.1016/j.ecss.2024.108674>, 2024.

750 Myhre et al (Ed.): Anthropogenic and Natural Radiative Forcing, in: Anthropogenic and Natural Radiative Forcing. In: Stocker, Thomas (ed.): Climate change 2013: the physical science basis; Working Group I contribution to the fifth assessment report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 659–740, <https://doi.org/10.1017/CBO9781107415324.018>, 2014.

755 [Nóbrega, G.N., Ferreira, T.O., Neto, M.S., Queiroz, H.M., Artur, A.G., Mendonça, E.D.S., Silva, E.D.O., and Otero, X.L.: Edaphic factors controlling summer \(rainy season\) greenhouse gas emissions \(CO₂ and CH₄\) from semiarid mangrove soils \(NE-Brazil\). *Science of the Total Environment*, 542, 685–693, <https://doi.org/10.1016/j.scitotenv.2015.10.108>, 2014.](https://doi.org/10.1016/j.scitotenv.2015.10.108)

Ouyang, X., Lee, S. Y., and Connolly, R. M.: Structural equation modelling reveals factors regulating surface sediment organic carbon content and CO₂ efflux in a subtropical mangrove, *Science of The Total Environment*, 578, 513–522, <https://doi.org/10.1016/j.scitotenv.2016.10.218>, 2017.

760 Ouyang, X., Guo, F., and Lee, S. Y.: Multiple drivers for carbon stocks and fluxes in different types of mangroves, *Science of The Total Environment*, 906, 167511, <https://doi.org/10.1016/j.scitotenv.2023.167511>, 2024.

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto

Formatted: Bibliography

Padhy, S., Bhattacharyya, P., Dash, P., Reddy, C., Chakraborty, A., and Pathak, H.: Seasonal fluctuation in three mode of greenhouse gases emission in relation to soil labile carbon pools in degraded mangrove, Sundarban, India, *SCIENCE OF THE TOTAL ENVIRONMENT*, 705, <https://doi.org/10.1016/j.scitotenv.2019.135909>, 2020.

765 [Podgrajsek, E., Sahlée, E., Bastviken, D., Ho Ist, J., Lindroth, A., Tranvik, L., et al.: Comparison of floating chamber and eddy covariance measurements of lake greenhouse gas fluxes. *Biogeosciences* 11, 4225–4233, <https://doi.org/10.5194/bg-11-4225-2014>, 2014.](#)

770 Poffenbarger, H. J., Needelman, B. A., and Megonigal, J. P.: Salinity Influence on Methane Emissions from Tidal Marshes, *Wetlands*, 31, 831–842, <https://doi.org/10.1007/s13157-011-0197-0>, 2011.

Rosentreter, J. A. and Williamson, P.: Concerns and uncertainties relating to methane emissions synthesis for vegetated coastal ecosystems, *Global Change Biology*, 26, 5351–5352, <https://doi.org/10.1111/gcb.15201>, 2020.

775 Rosentreter, J. A., Maher, D. T., Erler, D. V., Murray, R., and Eyre, B. D.: Factors controlling seasonal CO₂ and CH₄ emissions in three tropical mangrove-dominated estuaries in Australia, *Estuarine, Coastal and Shelf Science*, 215, 69–82, <https://doi.org/10.1016/j.ecss.2018.10.003>, 2018a.

Rosentreter, J. A., Maher, D. T., Erler, D. V., Murray, R., and Eyre, B. D.: Seasonal and temporal CO₂ dynamics in three tropical mangrove creeks – A revision of global mangrove CO₂ emissions, *Geochimica et Cosmochimica Acta*, 222, 729–745, <https://doi.org/10.1016/j.gca.2017.11.026>, 2018b.

780 Saderne, V., Cusack, M., Almahasheer, H., Serrano, O., Masqué, P., Arias-Ortiz, A., Krishnakumar, P. K., Rabaoui, L., Qurban, M. A., and Duarte, C. M.: Accumulation of Carbonates Contributes to Coastal Vegetated Ecosystems Keeping Pace With Sea Level Rise in an Arid Region (Arabian Peninsula), *Journal of Geophysical Research: Biogeosciences*, 123, 1498–1510, <https://doi.org/10.1029/2017JG004288>, 2018.

785 Saderne, V., Gerdali, N. R., Macreadie, P. I., Maher, D. T., Middelburg, J. J., Serrano, O., Almahasheer, H., Arias-Ortiz, A., Cusack, M., Eyre, B. D., Fourqurean, J. W., Kennedy, H., Krause-Jensen, D., Kuwae, T., Lavery, P. S., Lovelock, C. E., Marba, N., Masqué, P., Mateo, M. A., Mazarrasa, I., McGlathery, K. J., Oreska, M. P. J., Sanders, C. J., Santos, I. R., Smoak, J. M., Tanaya, T., Watanabe, K., and Duarte, C. M.: Role of carbonate burial in Blue Carbon budgets, *Nat Commun*, 10, 1106, <https://doi.org/10.1038/s41467-019-08842-6>, 2019.

790 Saderne, V., Fusi, M., Thomson, T., Dunne, A., Mahmud, F., Roth, F., Carvalho, S., and Duarte, C. M.: Total alkalinity production in a mangrove ecosystem reveals an overlooked Blue Carbon component, *Limnology and Oceanography Letters*, 6, 61–67, <https://doi.org/10.1002/lol2.10170>, 2021.

Sanchez-Carrillo, S., Garatuza-Payan, J., Sanchez-Andres, R., Cervantes, F., Bartolome, M., Merino-Ibarra, M., and Thalasso, F.: Methane Production and Oxidation in Mangrove Soils Assessed by Stable Isotope Mass Balances, *WATER*, 13, <https://doi.org/10.3390/w13131867>, 2021.

795 Sanderman, J., Hengl, T., Fiske, G., Solvik, K., Adame, M. F., Benson, L., Bukoski, J. J., Carnell, P., Cifuentes-Jara, M., Donato, D., Duncan, C., Eid, E. M., Ermgassen, P. zu, Lewis, C. J. E., Macreadie, P. I., Glass, L., Gress, S., Jardine, S. L., Jones, T. G., Nsombo, E. N., Rahman, M. M., Sanders, C. J., Spalding, M., and Landis, E.: A global map of mangrove forest soil carbon at 30 m spatial resolution, *Environ. Res. Lett.*, 13, 055002, <https://doi.org/10.1088/1748-9326/aabe1c>, 2018.

Sea, M. A., Garcias-Bonet, N., Saderne, V., and Duarte, C. M.: Carbon dioxide and methane fluxes at the air–sea interface of Red Sea mangroves, *Biogeosciences*, 15, 5365–5375, <https://doi.org/10.5194/bg-15-5365-2018>, 2018.

- 800 Shukla, P. N., Pandey, K. D., and Mishra, V. K.: Environmental Determinants of Soil Methane Oxidation and Methanotrophs, *Critical Reviews in Environmental Science and Technology*, 43, 1945–2011, <https://doi.org/10.1080/10643389.2012.672053>, 2013.
- Sippo, J. Z., Maher, D. T., Tait, D. R., Holloway, C., and Santos, I. R.: Are mangroves drivers or buffers of coastal acidification? Insights from alkalinity and dissolved inorganic carbon export estimates across a latitudinal transect, *Global Biogeochemical Cycles*, 30, 753–766, <https://doi.org/10.1002/2015GB005324>, 2016.
- 805 [Sjögersten, S., Aplin, P., Gauci, V., Peacock, M., Siegenthaler, A., and Turner, B. L.: Temperature response of ex-situ greenhouse gas emissions from tropical peatlands: Interactions between forest type and peat moisture conditions. *Geoderma* 324, 47–55. <https://doi.org/10.1016/j.geoderma.2018.02.029>, 2018.](https://doi.org/10.1016/j.geoderma.2018.02.029)
- 810 Smorkalov, I. A.: Soil Respiration Variability: Contributions of Space and Time Estimated Using the Random Forest Algorithm, *Russ J Ecol*, 53, 295–307, <https://doi.org/10.1134/S1067413622040051>, 2022.
- Speiser, J. L., Miller, M. E., Tooze, J., and Ip, E.: A comparison of random forest variable selection methods for classification prediction modeling, *Expert Systems with Applications*, 134, 93–101, <https://doi.org/10.1016/j.eswa.2019.05.028>, 2019.
- 815 Sugiana, I. P., Faiqoh, E., Adame, M. F., Indrawan, G. S., Andiani, A. A. E., Dewi, I. G. A. I. P., and Dharmawan, I. W. E.: Soil greenhouse gas fluxes to the atmosphere during the wet season across mangrove zones in Benoa Bay, Indonesia, *Asian J. Atmos. Environ*, 17, 13, <https://doi.org/10.1007/s44273-023-00014-9>, 2023.
- Sultan, S., Ahmad, F., and Elghribi, N.: Sea level variability in the central Red Sea, *Oceanologica Acta*, 19, 1996.
- Tavares, R. L. M., Oliveira, S. R. de M., Barros, F. M. M. de, Farhate, C. V. V., Souza, Z. M. de, and Scala Junior, N. L.: Prediction of soil CO₂ flux in sugarcane management systems using the Random Forest approach, *Sci. agric. (Piracicaba, Braz.)*, 75, 281–287, <https://doi.org/10.1590/1678-992X-2017-0095>, 2018.
- 820 Teh, Y. A., Silver, W. L., and Conrad, M. E.: Oxygen effects on methane production and oxidation in humid tropical forest soils, *Global Change Biology*, 11, 1283–1297, <https://doi.org/10.1111/j.1365-2486.2005.00983.x>, 2005.
- 825 Tete, E., Viaud, V., and Walter, C.: Organic carbon and nitrogen mineralization in a poorly-drained mineral soil under transient waterlogged conditions: an incubation experiment, *European Journal of Soil Science*, 66, 427–437, <https://doi.org/10.1111/ejss.12234>, 2015.
- [Toczylowski, A. J. Z., Slesak, R. A., Kolka, R. K., and Venterea, R. T.: Temperature and water-level effects on greenhouse gas fluxes from black ash \(*Fraxinus nigra*\) wet land soils in the Upper Great Lakes region, USA. *Appl. Soil Ecol.* 153103565. <https://doi.org/10.1016/j.apsoil.2020.103565>, 2020.](https://doi.org/10.1016/j.apsoil.2020.103565)
- 830 Tripathi, A. K., Roberts, C. D., and Eagle, R. A.: Coupling of CO₂ and Ice Sheet Stability Over Major Climate Transitions of the Last 20 Million Years, *Science*, 326, 1394–1397, <https://doi.org/10.1126/science.1178296>, 2009.
- [Van Dijk, A.I.J.M., H.E. Beck, E. Boergens, R.A.M. de Jeu, W.A. Dorigo, T. Frederikse, A. Güntner, J. Haas, J. Hou, W. Preimesberger, J. Rahman, P.R. Rozas Larraondo, and R. van der Schalie: Global Water Monitor 2023, Summary Report. Global Water Monitor \(www.globalwater.online\), 2023.](https://www.globalwater.online)
- 835

Wang, H., Liao, G., D'Souza, M., Yu, X., Yang, J., Yang, X., and Zheng, T.: Temporal and spatial variations of greenhouse gas fluxes from a tidal mangrove wetland in Southeast China, *Environ Sci Pollut Res*, 23, 1873–1885, <https://doi.org/10.1007/s11356-015-5440-4>, 2016.

840 Yu, X., Yang, X., Wu, Y., Peng, Y., Yang, T., Xiao, F., Zhong, Q., Xu, K., Shu, L., He, Q., Tian, Y., Yan, Q., Wang, C.,
Wu, B., and He, Z.: *Sonneratia apetala* introduction alters methane cycling microbial communities and increases methane
845 emissions in mangrove ecosystems, *Soil Biology and Biochemistry*, 144, 107775, <https://doi.org/10.1016/j.soilbio.2020.107775>, 2020.

Zeebe, R. E. and Wolf-Gladrow, D.: *CO₂ in Seawater: Equilibrium, Kinetics, Isotopes*, Gulf Professional Publishing, 382
845 pp., 2001.

Zhu, X., Sun, C., and Qin, Z.: Drought-Induced Salinity Enhancement Weakens Mangrove Greenhouse Gas Cycling, *Journal
of Geophysical Research: Biogeosciences*, 126, e2021JG006416, <https://doi.org/10.1029/2021JG006416>, 2021.