

Earth observation reveals reduced winter wheat growth and the importance of plant available water during drought

Hanna Sjulgård¹, Lukas Valentin Graf^{2,3}, Tino Colombi^{1,4}, Juliane Hirte³, Thomas Keller^{1,3,§}, Helge Aasen^{3,§}

¹ Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

² Crop Science, Institute for Agricultural Science, ETH Zurich, Zürich, Switzerland

³ Department of Agroecology and Environment, Agroscope, Zürich, Switzerland

⁴ School of Biosciences, University of Nottingham, Sutton Bonington, United Kingdom

§ These authors contributed equally

Correspondence to: Hanna Sjulgård (hanna.sjulgard@slu.se)

Abstract. Drought poses increasing challenges to global food production. Knowledge about the influence of drought on crop development and ~~about~~ the role of soil properties for crop drought severity is important in drought risk analysis and for mitigating drought impacts at the landscape level ~~is important to guide climate change adaptation. Satellite earth observations can provide area wide insights into crop growth processes that may help identify risk factors and quantify vulnerability to drought.~~ Here, we evaluate the potential of tested if satellite images from Sentinel-2 to reveal interactions could be used to assess the impacts of drought on crop development and the influence of plant growth soil properties on crop drought responses at the landscape scale, and soil parameters during variable weather conditions what the responses were. As a case study, we ~~assess~~assessed winter wheat growth on 13 fields belonging to commercial farmers in southern Sweden in a dry year (2018) and a year with normal weather conditions (2021). To track crop growth, green leaf area index (GLAI) was estimated from satellite imagery using a radiative transfer model. Proxies for winter wheat growth rate, peak GLAI, and the timing of peak GLAI were derived from the GLAI development at the single field level.

We then compared the crop growth proxies between the two years and across the fields, and related them the year-to-year differences between fields to measured soil properties. We found a lower estimated growth rates, lower peak GLAI and earlier peak GLAI in the dry year compared to the year with normal weather conditions. An increase in A higher peak GLAI in the dry year was also shown to be related to a higher growth rate, and this was not shown in the year with normal precipitation. Differences in crop development between years were large for some fields but small for other fields, suggesting that soil properties play a role in crop response to drought. We found that fields with a higher amount of plant available water capacity had better crop performance a higher growth rate in the dry year and smaller relative differences in growth rate between the two years. The observed lower growth rate, lower peak GLAI, and earlier peak in This shows the dry year compared importance of soils to the year with normal weather mitigate drought conditions, demonstrate which will likely become more relevant in an increasingly drier climate. Our case study demonstrates that satellite derived crop growth proxies

~~can identify crop responses to drought events, and that satellite imagery can be used to quantify plant-discover impacts of soil-weather interactions properties on crop development at scales relevant to commercial farming. Our investigation serves as a first step towards supporting drought risk management, drought adaptation and communication activities on this important topic.~~

1 Introduction

Extreme weather events such as droughts have become more frequent and severe in recent years due to climate change, posing challenges to global food and feed production (IPCC 2022). Drought is one of the main climatic constraints limiting crop growth and crop productivity (Fahad et al. 2017; Matiu et al. 2017; Ru et al. 2023). Water is crucial for plant growth, and plants can respond to water limitation through different mechanisms, such as reducing water losses through transpiration by closing their stomata (Huang et al. 2020) or by reducing leaf area (Wasaya et al. 2023). In turn, the photosynthesis rate and thus carbon acquisition decrease. Plants may also accelerate their development to complete the plant life cycle before the occurrence of a severe water deficit (Abid et al. 2018; Seleiman et al. 2021). The impact of drought on crops is complex and depends on several factors including the plant species and variety, the developmental stage of plants, the timing, duration and severity of the drought (Gray and Brady 2016), as well as the properties of the soil (Bodner et al. 2015).

The capacity of soil to sustain plant growth and crop productivity is affected by biological, chemical and physical soil properties, which collectively determine the soil conditions for plant growth (Stockdale et al. 2002). Soils ~~with higher resilience to drought that~~ allow water to infiltrate and can store ~~moisture~~ sufficient amounts of water to sustain plant growth can mitigate drought conditions (Rockström 2003; Bodner et al. 2015). Higher ~~moisture in the~~ soil moisture may also benefit nutrient uptake during drought. ~~A, while a~~ water deficit could lead to a lack of nutrients in crops as nutrients are mainly transported into plants through water uptake (He and Dijkstra 2014). Plant roots must also be able to penetrate the soil to access water and nutrient resources, where a high penetration resistance, which increases ~~with~~ under dry conditions, could impede root growth and resource accessibility (Bengough et al. 2011; Colombi et al. 2018). Recent research also ~~shows~~ provides evidence that certain rhizosphere microbiomes might enhance plant growth during dry conditions (Rolli et al. 2015; Rubin et al. 2017; de Vries et al. 2020). Therefore, ~~the~~ soil properties are of high importance to sustain crop growth during drought.

~~Plant growth dynamics can be quantified with ecophysiological properties such as the green leaf area index (GLAI), which is the ratio of photosynthetically active leaf area to ground area (Watson 1947). Previous studies demonstrated that the influence of soil properties and soil-borne stress on plant growth can be detected using GLAI. For example, positive relationships between GLAI and soil water content have been found (Chen et al. 2021), and GLAI at the heading stage of winter wheat has been shown to decrease with a high degree of soil compaction (Lipiec et al. 1991) in field experiments. The growth rate estimated from GLAI has also been shown to be related to soil carbon and nitrogen contents (Hirooka et al. 2017). However, there is still limited information about how soil properties affect crop development under various weather conditions, especially~~

at scales relevant to commercial agriculture (i.e., at the landscape scale). Research at larger scales than field experiments is particularly important since the heterogeneity of environmental factors in the landscape is more complex than what can be investigated in typical field plot experiments:

~~Monitoring crop growth at the landscape scale can be done with satellite remote sensing, for example using the twin constellation of Sentinel-2A and 2B. The Sentinel satellites provide a high spatial resolution optical imagery of up to 10 metres, and a high revisit time of one to four days depending on the latitude (Drusch et al. 2012). Plant growth dynamics can be quantified with ecophysiological properties such as the green leaf area index (GLAI), which is the ratio of photosynthetically active leaf area to ground area (Watson 1947). Previous studies using field experiments demonstrated that the influence of soil properties and soil-borne stress on plant growth can be detected using GLAI. For example, positive relationships between GLAI and soil water content have been found (Chen et al. 2021), and GLAI at the heading stage of spring barley has been shown to decrease with a high degree of soil compaction (Lipiec et al. 1991). The growth rate estimated from GLAI has also been shown to be related to soil organic carbon and nitrogen contents (Hirooka et al. 2017). In addition, the GLAI may vary by crop species, scales and environmental factors (Kang et al. 2002; Kang et al. 2016; Lawal et al. 2022), and there is still limited information about how soil properties affect crop GLAI development under extreme weather conditions, at scales relevant to commercial agriculture (i.e., at the landscape scale). Pot and field plot experiments are needed to understand single factors, but conducting research at larger scales is important to capture the heterogeneity of environmental factors in the landscape.~~

~~Monitoring crop growth at the landscape scale can be done with satellite remote sensing, for example using the twin constellation of Sentinel-2A and 2B.~~ The Sentinel-2 multispectral sensors have been shown to be suitable for estimating GLAI for different crop species (Clevers et al. 2017; Revill et al. 2019; Dong et al. 2020; Ali et al. 2021). One promising way to interpret satellite data for ecophysiological traits is the use of radiative transfer models. ~~Radiative transfer models that~~ describe the relationship between leaf and canopy traits and spectral properties of plants using physical principles (Jacquemoud et al. 1996; Myneni et al. 1997; Verhoef 1998). Thus, in contrast to the widely used vegetation indices, there is no need to establish empirical relationships between vegetation indices and crop traits (Atzberger et al. 2011). Those empirical relationships are usually not transferable in space and time, and hence not suitable for studies at the landscape scale. In addition, vegetation indices such as the widely used Normalized Difference Vegetation Index (NDVI) saturate at low biomass levels (Myneni and Williams 1994; Prabhakara et al. 2015), which is undesirable for a reliable and robust quantification of plant growth. The combination of satellite images and radiative transfer models allows estimating GLAI on a large scale.

The use of satellite-derived GLAI for crop growth characterization and productivity has become common in recent years (Punalekar et al. 2018; Peng et al. 2019; Dong et al. 2020; He et al. 2021; Graf et al. 2023), ~~but knowledge about how extreme weather, such as drought, affects GLAI development remains scarce. The year 2018 was characterised by an unusually dry and hot spring and summer in northern Europe (Wileke et al. 2020). These extreme conditions had a large impact on agriculture, and around 40% of the cropping areas with winter wheat in Northern Europe had yields below the 10th percentile (Beillouin et al. 2023), and many remote sensing studies motivate their work by the potential of remote sensing to detect crop stress. Still,~~

~~studies that a) demonstrate how extreme weather, such as drought, affects GLAI development, and b) provide the link to environmental variables, such as soil properties, to explain the observed differences remain scarce. Investigating if satellite images can be used to identify crop stress responses at the landscape scale, and if the importance of soil properties can be identified under drought stress at agricultural fields, could motivate the use of satellite images in crop monitoring at farm fields. 2020). The probability of droughts is increasing due to climate change (Wileke et al. 2020), posing increasing challenges to crop production. Studying the impact of drought on plant growth and development, and the role of soil properties in aggravating or mitigating drought impacts, can provide crucial knowledge for future climate change adaptations.~~ In the present study, the aims were to:

- i) analyse winter wheat development in farm fields within a region in southern Sweden by quantifying GLAI based on Sentinel-2 data,
- ii) investigate ~~theif~~ impact of drought on winter wheat growth can be identified using satellite images at the farm fields by comparing the GLAI development ~~by comparing between~~ a dry year (year 2018) and a year with normal weather conditions (year 2021), and
- iii) examine ~~the influence of if differences in~~ soil properties ~~on relate to~~ differences between GLAI development across fields and between the two years.

2 Materials and Methods

2.1 Study area and meteorological data

The study area was located in the south of Sweden at a latitude of approximately 58.5°, spanning 160 km from ~~the~~ west to ~~the~~ east (Fig. 1), and is characterized by a humid continental climate (Peel et al. 2007). Winter wheat is the major crop cultivated in Sweden in general and in the study area (Sjulgård et al. 2022). We included 13 fields in this study, belonging to commercial farmers ~~that~~. The fields were cultivated with winter wheat (*Triticum aestivum* L.) in both 2018 and 2021, and ~~for which~~ detailed soil data were available for all fields. All fields were managed conventionally and they were not irrigated. The farmers manage their fields according to best practices, but detailed information about crop and soil management practises was only available from some of the farmers. We therefore minimised the variation in management practises between years by selecting fields that were managed by the same farmer in 2018 and 2021, and with the same crop cultivated in both years.






Fig. 1. A map of Sweden with county borders showing the location of the study area (left), and a map displaying the locations of the 13 fields (right). A small blue circle indicates the location of one field, and a larger blue circle indicates two fields close to each other.

The centroid coordinates of the fields were used to obtain the daily temperature and precipitation data for each field. The meteorological data were obtained from the “PTHBV database,” available from the Swedish Meteorological and Hydrological Institute (SMHI). The data include gridded and interpolated daily mean temperature and precipitation at a resolution of 4 km by 4 km (SMHI 2023). The interpolation is based on 700 meteorological stations across Sweden and considers orographic effects (Berg et al. 2015). Differences in weather conditions between fields and years were assessed by the De Martonne Aridity Index (DMI; De Martonne 1926)), defined as:

$$DMI = \frac{P_m}{T_m + 10} \quad (1)$$

where P_m is the monthly total precipitation (mm) and T_m is the monthly average temperature ($^{\circ}\text{C}$). A higher DMI indicates wetter conditions, while a lower DMI value indicates drier conditions. Weather conditions during the main winter wheat growing period (May to July) in 2018 and 2021 were contrasting: 2018 was unusually dry, while 2021 was “normal” (Fig. 2). The lack of precipitation has been referred to as the main reason for the large yield losses observed in 2018 (Bakke et al. 2020, Beillouin et al. 2020). In the summer 2018, May was already unusually warm and dry in Sweden (SMHI 2018). Between May and July, the DMI was on average $1.2 \text{ mm}/^{\circ}\text{C}^{-1}$ (SD = 0.06) per month for the 13 fields in 2018, which was

drier than the long-term average between 1991 and 2020 of $2.9 \text{ mm/}^\circ\text{C}^{-1}$ (SD = 0.19) for the same period. In 2021, the DMI was closer to the long-term average with a monthly mean during the summer of $3.2 \text{ mm/}^\circ\text{C}^{-1}$ (SD = 0.16), and with in May to July. In both years, DMI was similar values between the across fields.






Fig.

Fig. 1. A map of Sweden with county borders showing the location of the study area (left), and the map displaying the locations of the fields (right). 2. Weather conditions in 2018 and 2021 at the locations of the 13 fields. Locally estimated scatterplot smoothing curves plotting the average a) daily temperature and b) precipitation against calendar date for the 13 fields in 2018 and 2021. c) The average DMI (De Martonne Aridity Index) during the months May to July in 2018 and 2021, and the long-term mean May to July between 1991 and 2020 with error bars indicating the standard deviation.

~~A small blue circle indicates the location of one field, and a larger blue circle indicates two fields close to each other.~~

2.2 GLAI derived from satellite data

The twin constellations of the Sentinel-2A and B satellites have a revisit time of two days in the study area. Downloading and processing of Sentinel-2 data were performed using the open-source Python Earth Observation Data Analysis Library (EOdal, Graf et al. 2022). The Sentinel-2 scenes were obtained for the years 2018 and 2021 from Microsoft Planetary Computer. 20 m and 10 m bands were obtained, and the Sentinel-2 scenes and 20 m bands were resampled to 10 m using nearest-neighbour interpolation to generate equal spatial resolution. The Sentinel-2 scenes were cropped to only retain pixels within the 13 fields based on a shapefile containing the field boundaries. From the resampled scene classification layer, only pixels from the scene classification layer class 4 (vegetation) and class 5 (bare soil) were kept to filter out pixels containing clouds, snow, shadow, and dark areas. Further filtering was performed to remove dates with a cloud cover of $\geq 10\%$ on a field-per-field basis.

~~The GLAI was derived from the radiative transfer model PROSAIL, following the approach described in Graf et al. (2023). A lookup table consisting of 50,000 spectra was generated by running the PROSAIL model in forward mode for each Sentinel-2 scene. We randomly generated combinations of leaf and canopy parameters according to a uniform or Gaussian distribution (Table S1). View and illumination geometry were set to scene specific values extracted from the Sentinel 2 scene~~

~~metadata~~, Tab. S2; Graf et al. 2023; Woher et al. 2020; Danner et al. 2021). ~~View and illumination geometry were set to scene-specific values extracted from Sentinel-2 scene metadata.~~ Building on the workflow of Graf et al. (2023), known empirical relationships between GLAI and chlorophyll a and b, and GLAI and the carotenoid content of leaves were used to increase the physiological plausibility of the input parameter combinations. For GLAI retrieval, we compared the Sentinel-2 pixel spectra with the PROSAIL simulated spectra using the mean absolute error as a cost function. We then used the median of the 5000 (10%) best matching simulated spectra in terms of the smallest mean absolute error to derive a GLAI value per Sentinel-2 pixel.

For each Sentinel-2 scene, an average value of GLAI was calculated per field. A smoothed curve was fitted to the GLAI time series by the locally estimated scatterplot smoothing (~~LOESS~~) method with a span of 0.3 (Fig. 23). The smoothed curve was also used to identify and remove outliers that were missed by the scene classification layer and the cloud filtering (Fig. S2S1).

2.3 Crop growth curve parametrisation

The air temperature sum (~~TSUM~~ T_{sum}) at ~~the location of~~ each field was assessed by adding up the daily mean temperatures exceeding ~~the~~ threshold value of 0 °C, where growth for winter wheat starts (Porter and Gawith 1999), from the 1st of January following:

$$T_{sum}UM = \sum_{i=1}^j T_i \times \sigma_i$$

$$\sigma_i = \begin{cases} 0 & \text{if } T_i \leq 0 \text{ °C} \\ 1 & \text{if } T_i > 0 \text{ °C} \end{cases} \quad (2)$$

~~Where~~ T_i is the daily mean temperature and j is the number of days.

From the GLAI development curve, characteristic properties were calculated to estimate ~~the~~ crop growth rate, green biomass, and timing of ~~development~~ heading stage for each field and year (Fig. 23). GLAI increases early in the season due to leaf production in the vegetative growth phase (Bhattacharya 2019). Growth rate during the vegetative growth phase was estimated from the slope of a linear plateau curve with an endpoint at the start of the upper plateau. The linear plateau model was fitted to the GLAI values with a start at a temperature sum of 200 °C (corresponding to the end of April) when ~~the~~ GLAI started to increase around the beginning of ~~the~~ stem elongation (Chen et al. 2009). The GLAI development curve is typically bell-shaped, with the peak GLAI observed around the heading stage for winter wheat (Feng et al. 2019). The timing of the peak GLAI was assessed from the corresponding temperature sum- (Fig. 3). The ~~peak~~ GLAI ~~at the peak~~ indicates the maximum green biomass (Lambert et al. 2018; Skakun et al. 2019), and was assessed from the smoothed GLAI curve.




Fig. 23. Example from one of the fields showing the green leaf area index (GLAI) temporal development curve. We obtained proxies for the growth rate from the slope between a temperature sum of 200 °C until the start of the plateau, (dashed red line), the peak GLAI from the maximum GLAI, and the timing of peak GLAI from the temperature sum at the peak GLAI. The raw GLAI values are shown by black dots and the smoothed GLAI is shown by the black curve.

2.4 Soil sampling and analyses

Soil sampling was conducted in June and in the beginning of July in 2021. Loose soil samples and undisturbed soil cores were collected from the topsoil at five locations in each field. Sampling locations within each field were arranged in a quincunx, with one point in the middle of the field and the others at least a few metres from the field borders. Loose soil samples were taken with a shovel from 0-20 cm depth. The five samples taken in each field were pooled into a plastic bag and the resulting composite sample was air-dried. Five undisturbed soil cores (5 cm in height, 7.2 cm inner diameter) were collected at a depth of 10 cm in each field. The soil core samples were wrapped airtight and stored at 4 °C until further processing.

Soil organic matter content was determined by loss of ignition from the loose soil samples. Cation exchange capacity was analysed using an inductively coupled plasma–optical emission spectrometer (ICP-OES) to obtain the base cations in the soil samples. The base cations and acidity titration were used to calculate the cation exchange capacity at pH 7. Soil water content at the permanent wilting point (-1500 kPa) was determined with pressure plate extractors. Soil water content at field capacity was assessed by equilibrating the soil cores to -10 kPa ~~on ceramic plates (ecoTec, Bonn). Plant available water capacity was obtained by calculating the difference in soil water content between field capacity and the permanent wilting point. (i.e., field capacity; Krueger and Ochsner (2024)) on ceramic plates (ecoTec, Bonn). Plant available water capacity was obtained by calculating the difference in gravimetric soil water content between field capacity and the permanent wilting point.~~ Dry soil bulk density was determined on the undisturbed soil core samples by drying the samples at 105 °C for 48 h. Soil texture including clay (< 0.002 mm) content was determined from the loose soil samples by sedimentation ('pipette' method).

2.5 Statistical analyses

GLAI development responses to drought were analysed by comparing ~~the~~ differences in crop growth proxies (i.e., growth rate, peak GLAI, and the timing of peak) between the dry year (2018) and the year with normal weather conditions (2021). A two-tailed t-test was applied to determine whether there was a significant difference in growth rate, peak GLAI, and the timing of the peak GLAI between the two years. Spearman correlation was used to assess ~~the~~ relationships between soil properties. Multiple linear regression was used to assess relationships between the crop growth proxies, and between soil properties, while accounting for the average monthly DMI (May – July) of the corresponding year. To relate soil properties to differences in ~~the~~ growth rate, peak GLAI, and the timing of the peak GLAI between ~~the~~ years, the relative difference of ~~the~~ crop growth proxies (ΔGP) between the years 2018 and 2021 was calculated as:

$$\Delta GP = \frac{GP_{2021} - GP_{2018}}{GP_{2018}} \times 100\% \quad (3)$$

~~Where~~ where GP is ~~the~~ the crop growth ~~proxies~~ proxy (i.e. growth rate, peak GLAI or the timing of the peak GLAI ~~in~~ for year 2018 and 2021).


A variance decomposition method proposed by Zuber and Strimmer (2011), called Correlation-Adjusted coRelation (CAR) scores, was used to determine the relative importance of the soil properties for the growth rate, peak GLAI, and the timing of the peak GLAI in each year (i.e., for 2021 and 2018), and for the relative difference of the crop growth proxies between the years. CAR scores provide a criterion for variable ranking in linear regression based on the Mahalanobis-decorrelation of covariates (Zuber and Strimmer 2011). The direction of the relationships and p-values were obtained from univariate linear regressions between the crop growth proxies and the soil properties for each year, and for the relative difference of crop growth proxies ~~in~~ between 2021 ~~to~~ and 2018, respectively. ~~A significance level of $p < 0.05$ was used. The statistical~~ Statistical analyses were carried out in R version 4.2.1 (R Core Team 2022), ~~and~~ the CAR scores were calculated from the R package “relaimpo” (Groemping and Lehrkamp 2023) ~~and the linear mixed models using the “lme4” package (Bates et al. 2015).~~

3 Results

3.1 Growth patterns across years

~~The differences~~ Differences in crop development between years varied ~~between~~ across fields, where certain fields showed a large difference in growth rate, peak GLAI and the timing of the peak GLAI between years, while others ~~showed~~ had only small differences (Fig. 34, Fig. S2). For growth rate and ~~the~~ peak GLAI, four fields ~~in total~~ had an increase from 2018 to 2021 of less than 10%, while some fields had a difference of 50-59%. The difference between ~~the~~ years in the timing of the peak

GLAI was lower in comparison, with four fields having an increase <10% and three fields a decrease <10%, while the maximum difference was 30-39% ~~between the years~~ (Fig. 34, Fig. S2).






Fig. 3. Temporal evolution of GLAI during 2018 (dry year) and 2021 (year with normal weather conditions) for two different fields with a) a large difference between the years and b) a small difference between the years. c) Number of fields by the percentage difference in the crop growth proxies (i.e., peak GLAI, growth rate, and temperature sum at peak GLAI) between the years 2018 (dry year) and 2021 (year with normal weather conditions).

Growth rate was lower during the dry year (2018) compared to the year with normal weather conditions (2021; Fig. 45), indicating reduced plant growth in response to drought. The growth rate during the reproductive period was on average 19% lower in the dry year (2018) than in the year 2021 with closer to normal weather conditions, and we found a significant effect of the year on growth rate ($p < 0.001$; Fig. 4a5a). The peak GLAI was in general lower during the dry year compared to the year with normal weather conditions ($p < 0.001$; Fig. 6e5c), with an average difference of 28% between the two years. The timing of peak GLAI occurred significantly earlier, i.e. at a lower temperature sum, during the dry year, with the peak GLAI around a temperature sum of 775 °C in the dry year and 881 °C in the year with normal weather conditions ($p = 0.015$; Fig. 4d)-5d).





Fig. 45. Crop growth proxies obtained from the temporal evolution of green leaf area index (GLAI) in the dry year (2018) and the year with normal weather conditions (2021); a) growth rate, b) peak GLAI, and c) temperature sum at peak GLAI. Data show yearly average (black dots), median, upper and lower quartiles (box), and minimum and maximum values (whiskers). P-values from the t-test are displayed for the differences between the years (number of fields, n=13).

~~Relationships Fig. 4. Crop growth proxies obtained from the temporal evolution of green leaf area index (GLAI) in the dry year 2018 and the year with normal weather conditions 2021; a) growth rate, b) peak GLAI and c) temperature sum at peak~~

GLAI, during the dry year (2018) and the year with normal weather conditions (2021). Data show yearly average (black dots), median, upper and lower quartiles, and minimum and maximum values. P-values from the t-test are displayed for the differences between the years (number of fields, n=13).

The relationships among the different crop growth proxies showed a positive correlation between growth rate and the peak GLAI in the dry year (year 2018), while the relationship was not significant during the year with normal weather conditions. The timing of the peak GLAI had no significant relationship to the growth rate or to the peak GLAI for either years



(Fig. 5).

Tab. S3).

Fig. 5. Spearman correlation coefficients among the crop growth proxies that were derived from green leaf area index (GLAI) dynamics; growth rate, peak GLAI and temperature sum at peak GLAI, during a) dry year (year 2018) and b) year with normal weather conditions (year 2021). Blue colour indicates a positive correlation coefficient (number of fields n = 13).

3.2 Relationships between soil properties and crop development

On average across the thirteen 13 fields, plant available water capacity was 0.23%, m³ m⁻³ bulk density was 1.5 g cm⁻³, cation exchange capacity was 16 cmol kg⁻¹, soil organic matter content was 3.6%, and clay content was 31% (Tab. S2S1). Some soil properties were related to each other, with positive correlations between soil organic matter content and cation exchange capacity, and between clay content and cation exchange capacity- (p < 0.05; Fig. S4). Negative relationships were found between clay content and bulk density, and between clay content and plant available water capacity (p < 0.05; Fig. S4).

~~The soil~~Soil properties explained together 15%, 54% and 27% of the variations across fields in growth rate, peak GLAI, and ~~the~~ timing of peak GLAI, respectively, in the year with normal weather conditions (year 2021). However, none of the soil properties was significantly related to growth rate, peak GLAI or ~~the~~ timing of the peak GLAI in 2021 (Fig. 6a). In the dry year (year 2018), ~~the~~ soil properties explained together 44%, 40% and 55% of the variation in growth rate, peak GLAI and ~~the~~ timing of peak GLAI, respectively. Plant available water capacity was significantly related to crop growth rate in 2018, with a positive association of increased crop growth with higher plant available water capacity ($p < 0.05$). In addition, plant available water capacity explained 21% of the variation in growth rate across fields in the dry year. There were no significant relationships between the other soil properties ~~to~~and growth rate, peak GLAI or ~~the~~ timing of ~~the~~ peak GLAI in 2018 (Fig. 6b). ~~For~~Plant available water capacity was the most important soil property in explaining the relative difference between the year with normal weather conditions (2021) ~~to~~and the dry year (2018), ~~plant available water capacity was the most important soil property.~~ The relative difference in growth rate between the years was negatively related to plant available water capacity ($p < 0.05$), and plant-available water capacity explained 30% of the variation in the difference in growth rates between the years.




Fig. 6. Explained variation in growth rate, peak GLAI and temperature sum at peak GLAI, calculated from correlation-adjusted correlation (CAR) scores, by the soil properties clay content, soil organic matter content (SOM), bulk density, cation exchange capacity (CEC) and plant available water capacity (PAWC) on growth rate, peak GLAI and temperature sum at peak GLAI in a) a year with normal weather conditions (year 2021), b) a dry year (year 2018), and c) the relative difference between the years 2021 and 2018. The p-values and the positive or negative relationships between each soil property and the crop growth proxies were obtained from univariate linear regressions. The rings are 10%, 20%, 30% and 40% explained variation, starting from the smallest, and the last ring is the outside border of the plot area.

4 Discussion

4.1 The impact of drought on crop development

In the present study, we used satellite ~~imagery~~ images to assess winter wheat development in farm fields by quantifying GLAI based on Sentinel-2 data. We investigated whether the impact of drought on winter wheat growth-GLAI development and development at the landscape scale. In combination with measurements of relationships between soil properties, we could identify specific soil properties that were related to the potential to mitigate the impacts of and GLAI development during drought- could be identified by using satellite images. The early growing season in 2018 was exceptionally dry and warm, (Fig. 2), resulting in generally-reduced winter wheat ~~crop~~ development compared to 2021 (Fig. 5) that had close to long-term ~~summer~~-average weather conditions (Fig. 4). Previous research has shown negative effects of drought on crop yield at the landscape and country scale (Zipper et al. 2016; Ray et al. 2018; Sjulgård et al. 2023), and lower growth rate and lower peak GLAI during water-limited conditions have been found in field trials in which GLAI was measured at the canopy (Meinke et al. 1997; Boedhram et al. 2001). The lower crop growth rate and development~~the earlier GLAI peak~~ during drought that we observed in our study demonstrate that Sentinel-2 derived estimates of crop growth proxies can be used to detect drought responses in crop development at the landscape scale.

The dry conditions early in the growing season in year 2018 resulted in lower peak GLAI compared to the year with normal weather conditions, with an average difference of 40% between years (Fig. 4). In earlier studies, 5). Peak GLAI is a proxy of the peak GLAI has been used to provide yield estimations for different crops- maximum green biomass (Lambert et al. 2018; WaldnerSkakun et al. 2019; Yamamoto et al. 2023). Large reductions in winter wheat yields were reported in Sweden during 2018 due to the exceptional drought, with 44% lower total crop production compared to the previous five year average (SCB 2018). Reduced), and reduced wheat biomass during drought has been shown in earlier studies (Villegas et al. 2001; Zhang et al. 2018), and according). According to Villegas et al. (2001), the decrease in biomass und droughtis during drought was mainly due to a lower growth rate. Similarly, we found lower a positive relationship between crop growth rate during the reproductive period and peak GLAI in the dry year compared to, but not in the year with normal weather conditions, with an average difference between the years of 28% (Fig. 4a). Our results showed no relationships between growth rate and the peak GLAI during the normal weather year, but a positive relationship between growth rate and the peak GLAI in the dry year (Fig. 5). (Tab. S3). The positive relationship in the dry year shows the importance of a suggests that a faster growth is important to obtain higher growth rate for maximum biomass and in turn higher total biomass production yield during dry conditions, highlighting the importance of faster growth to mitigate drought impacts. Theand the non-significant relationship during the year with normal weather conditions suggests suggests that growth rate is not as critical for biomass accumulation during normal weather conditions.

Using farm fields, earlier research has shown that leaf area (He et al. 2020; Sun et al. 2024) and peak GLAI (Lambert et al. 2018, Yamamoto et al. 2023) can be related to crop yield. For the fields with yield data available in this study (six fields) together with additional 23 farm fields in the same region, there was a strong correlation between higher peak GLAI and higher

winter wheat yield in year 2021 shown in Sjulgård (2024). The peak GLAI was reached earlier ~~(i.e., at a lower temperature sum)~~, during the dry year, ~~and since~~. Since the peak GLAI has been associated with heading growth stage (Feng et al. 2019), this might indicate a shift in phenology during dry conditions. Some studies have shown that plants ~~might~~ develop faster during drought to reach flowering earlier and complete the life cycle before severe water shortage occurs (Abid et al. 2018; Seleiman et al. 2021). However, we did not find a significant ~~correlation~~ relationship between the timing of ~~the~~ peak GLAI and growth rate or peak GLAI in our data, which would imply that the timing of the heading growth stage did not influence the overall crop performance (Fig. 5). ~~Earlier research studying the impact of the timing of the peak GLAI on wheat yield is not unambiguous: some studies found no relationships between timing of peak GLAI and crop yield (Irfan-Ullah et al. 2021; Mroz et al. 2023), while an earlier heading have been related to higher wheat yield due to a longer grain filling period in other studies (Fewolde et al. 2006; Mohan et al. 2022). Tab. S3).~~

4.2. The influence of soil properties on crop development

We found that differences in crop development between the two years ~~but also between~~ varied across fields. ~~There was~~ When comparing 2018 and 2021, we identified a large difference of up to 50-59% ~~increase~~ in growth rate and peak GLAI for certain fields and up to 30-30% in the ~~crop growth proxies between 2018 and 2021, timing of peak GLAI~~, while there was a smaller difference for other fields (Fig. 3). ~~The timing of peak GLAI increased in some fields and decrease in other fields between the years, however, the differences were small (<9%). Due to similar~~ 4, Fig. S2). As weather conditions ~~(i.e., similar DMI)~~ across all fields within a specific year, ~~were similar (Fig. 2c)~~, the varying crop responses to drought stress ~~between~~ among fields imply that additional factors than the weather must have had an impact on crop development. Here, we show that soil properties ~~had an influence on~~ influenced the crop growth proxies. In 2018, a positive relationship between plant available water capacity and growth rate ~~shows~~ demonstrates the importance of sufficient soil water retention to sustain crop growth during drought (Fig. 6). ~~In addition, fields~~ Fields with lower plant available water capacity had a larger relative difference in growth rate between the ~~years: dry and normal year~~. Earlier studies have ~~also shown the importance of the~~ that soil water ~~status to retain a sufficient amount of moisture~~ retention is crucial for crop performance during drought (Wang et al. 2009; Huang et al. 2020). Accordingly, the performance of crops grown on soils with high plant available water capacity has been found less affected by changes in rainfall compared to crops grown on soils with low plant available water capacity (Wang et al. 2017). The ~~importance~~ relevance of plant available water capacity ~~for on crop growth during~~ drought ~~resistance found~~ identified in our study ~~suggests~~ demonstrates that ~~the influence of soil water retention is a crucial soil property that explains crop responses to~~ properties on crop development can be detected during drought ~~stress at the landscape level and a key target property to mitigate drought impacts~~ scale by ~~soil management using~~ Sentinel-2 derived GLAI.

~~The other~~ Other soil properties assessed in this study were not correlated with estimates of growth rate, peak GLAI or timing of ~~the~~ peak GLAI in 2018, and none of the soil properties was significantly related to the crop growth proxies in 2021 (Fig. 6). Clay content only explained a small part of the variation in crop growth proxies, but influenced other soil properties such as cation exchange capacity, bulk density, and plant available water capacity (Fig. S4). Cation exchange capacity only explained

a low part of the variation ~~of the~~ⁱⁿ crop growth proxies. All ~~the~~ fields were above the recommended cation exchange capacity for crop production of 10 cmol kg⁻¹ (Tab. ~~S2S1~~) (Chowdhury et al. 2021), implying that cation exchange capacity was not a limiting factor for crop development. Our findings that bulk density had no direct relationship with the crop growth proxies may seem to contradict the study of Lipiec et al. (1991) ~~that~~, who found decreasing GLAI at the heading stage of spring barley with ~~a higher~~^{increasing} degree of soil compaction. However, in our fields, bulk density was not critically high, with an average bulk density of 1.5 g cm⁻³ (Tab. ~~S2S1~~). We found no relationships between ~~the estimates of~~^{crop} growth rate, peak GLAI or timing of ~~the~~ peak GLAI and soil organic matter content. Earlier studies have shown positive effects of soil organic matter content on soil fertility (Lal 2009; Fageria 2012; Oldfield et al. 2019) and on crop productivity during drought (Kane et al. 2021; Mahmood et al. 2023), however, negative effects of soil organic matter content on crop yields have also been found in Sweden (Kirchmann et al. 2020).

~~In our study, soil sampling was conducted in 2021 only. With soil properties changing over time, this may introduce uncertainty in the relationships between the soil properties and the crop development that we established for year 2018. Nevertheless, a number of studies has shown only small year-to-year changes in soil organic carbon content (Krauss et al. 2020), water content at field capacity (Alam et al. 2014) and bulk density (Alam et al. 2014; Alnaimy et al. 2020) within given soil management systems. In addition to soil properties and weather conditions, crop development is influenced by soil and crop management practices such as fertilization (Agenbag and Maree 1991; Shankar et al. 2021), tillage (Agenbag and Maree 1991; Abagandura et al. 2017), sowing date and crop variety selection (Ihsan et al. 2016; Minoli et al. 2022). We minimised variation in these factors by selecting fields that were managed by the same farmer in 2018 and in 2021. Future research could examine the response of different soil and crop management strategies or the impact of crop variety on drought response to assist farmers in extreme weather mitigation.~~

4.3 Limitations and motivations

~~In our study, soil sampling was conducted in 2021 only. With soil properties changing over time, this may introduce uncertainty in the relationships between soil properties and crop development that we established for year 2018. However, a number of studies has shown only small year-to-year changes in soil organic carbon content (Krauss et al. 2020), water content at field capacity (Alam et al. 2014) and bulk density (Alam et al. 2014; Alnaimy et al. 2020) within given soil management systems. The small differences between years suggests that the soil properties probably are rather similar between the years 2021 and 2018 within the same fields included in this study. In addition to soil properties and weather conditions, crop development is influenced by soil and crop management practices such as fertilization (Agenbag and Maree 1991; Shankar et al. 2021), tillage (Agenbag and Maree 1991; Abagandura et al. 2017), sowing date and crop variety selection (Ihsan et al. 2016; Minoli et al. 2022). Earlier studies have shown differences in leaf area index between farming systems, with higher leaf area index in conventional in comparison to organic systems (Petcu et al. 2011; Pużyńska et al. 2021). In this study, all fields selected were conventionally managed to reduce these differences. Additional information about for example the winter wheat varieties, sowing date and fertilization levels were not available from all farmers. However, winter wheat is sown within a short time~~

window around the middle of September in the study region (Andersson 1983; SCB 1993). A change of a few days in sowing date of winter wheat has been shown to have limited influence on crop yield (Ding et al. 2016), and would therefore not substantially influence our findings. According to Stenberg et al. (2005), the average inorganic fertilizer used for winter wheat cultivation is 160 N kg ha⁻¹ (SD = 19) in Östergötland and 170 N kg ha⁻¹ (SD = 27) in Västra Götaland, respectively, based on extensive data collection between 2000 and 2003. This shows that the fertilization levels between the two counties that covered our study region are similar and that the variation between years is in general low.

The varying soil and crop management practices among fields, and the different availability of baseline data (e.g. soil management and input history) is one of the challenges with on-farm research, but such studies are essential to evaluate the use of satellite data in the context of commercial farms (Doole et al. 2023). Our results show that satellite derived GLAI can be used to identify environmental stress response on plants, and this could help farmers to monitor crops and to identify when stresses occur. The influence of soil properties on crop response during drought demonstrates the importance of accounting for soil properties when evaluating the impact of drought on crops.

5 Conclusion

The impact of drought on winter wheat development was shown by comparing Sentinel-2 derived GLAI development ~~of~~during a dry year (2018) and a year with normal weather ~~condition~~conditions (2021) across ~~thirteen~~13 fields belonging to commercial ~~farmers~~farm fields in southern Sweden. We observed lower crop growth rate, lower peak GLAI and earlier peak GLAI during the dry year (2018) compared to the year with normal weather conditions ~~(2021)~~. Our data ~~showed~~revealed the importance of a faster crop growth to obtain more biomass during dry conditions, while the growth rate was less crucial for crop performance during the year with normal weather conditions. ~~The differences~~Differences in crop development between the years demonstrate that stress related crop response to changing environmental conditions can be detected by monitoring crops using satellite images at the landscape level. ~~The variation in crop growth proxies between fields suggest that differences in soil properties across fields play a critical role for the differences in drought response of winter wheat. We,~~ and this could be useful for farmers to monitor their crops and identify when the plants are stressed. In addition, we found that plant available water capacity was important for crop growth rate during the dry year, ~~and that the relative difference in growth rate between the years was lower with higher plant available water capacity.~~ This suggests that ~~water retention is a target soil property to mitigate effects of drought~~satellite imagery can be used to discover soil impacts on crop development at scales relevant to commercial farming. The inclusion of soil properties in satellite images analyses could further improve the accuracy of the prediction of drought stress on crops.

Data availability

The Sentinel-2 scenes were obtained from Microsoft Planetary Computer, and the downloading and processing were performed using the open-source Python Earth Observation Data Analysis Library (EOdal, <https://github.com/EOA-team/eodal>). Data of the growth proxies are available from the corresponding author upon request. The precipitation and temperature data are available from the Swedish Meteorological and Hydrological Institute website <https://www.smhi.se/data/ladda-ner-data/griddade-nederbord-och-temperaturdata-ptbvy>.

Author contributions

Funding was acquired by TC, TK and HA. HS, TK and HA contributed to project conceptualization. HS performed the investigations with advise from LVG, JH, TC, TK and HA. LVG implemented the programming code in EOdal. HS wrote the paper, where the LVG, JH, TC, TK and HA contributed to the review and editing.

Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

This work was partly funded by the Swedish Farmers' Foundation for Agricultural Research (Stiftelsen Lantbruksforskning, SLF, grant number: O-19-23-309) and by Formas (The Swedish Research Council for Sustainable Development; grant no: 2022-00218), which is greatly acknowledged. Hanna Sjulgård acknowledges financial support from the Lennart Hjelm's foundation, Lukas Valentin Graf and Helge Aasen acknowledge funding by the Swiss National Science Foundation for the project “PhenomEn” (grant number IZCOZ0_198091), and Tino Colombi acknowledges funding from the University of Nottingham (Nottingham Research Fellowship). The authors want to thank all the farmers involved in this study for allowing soil sampling on their fields. We also thank Laura Martínez, Leah Eitelberg and Loraine ten Damme for help with the soil sampling, and Ana María Mingot Soriano, Anna Oskar and Jan Fiedler for laboratory measurements.

References

- Abagandura GO, Nasr GE-DM, Moumen NM. 2017. Influence of Tillage Practices on Soil Physical Properties and Growth and Yield of Maize in Jabal al Akhdar, Libya. *Open Journal of Soil Science*. 7(7):118–132. <https://doi.org/10.4236/ojss.2017.77010><https://doi.org/10.4236/ojss.2017.77010>
- Abid M, Ali S, Qi LK, Zahoor R, Tian Z, Jiang D, Snider JL, Dai T. 2018. Physiological and biochemical changes during drought and recovery periods at tillering and jointing stages in wheat (*Triticum aestivum* L.). *Sci Rep*. 8:4615. <https://doi.org/10.1038/s41598-018-21441-7>
- Agenbag GA, Maree PCJ. 1991. Effect of tillage on some soil properties, plant development and yield of spring wheat (*Triticum aestivum* L.) in stony soil. *Soil and Tillage Research*. 21(1):97–112. [https://doi.org/10.1016/0167-1987\(91\)90008-L](https://doi.org/10.1016/0167-1987(91)90008-L)
- Alam MK, Islam MM, Salahin N, Hasanuzzaman M. 2014. Effect of Tillage Practices on Soil Properties and Crop Productivity in Wheat-Mungbean-Rice Cropping System under Subtropical Climatic Conditions. *The Scientific World Journal*. 2014:e437283. <https://doi.org/10.1155/2014/437283>
- Ali AM, Savin I, Poddubskiy A, Abouelghar M, Saleh N, Abutaleb K, El-Shirbeny M, Dokukin P. 2021. Integrated method for rice cultivation monitoring using Sentinel-2 data and Leaf Area Index. *The Egyptian Journal of Remote Sensing and Space Science*. 24(3, Part 1):431–441. <https://doi.org/10.1016/j.ejrs.2020.06.007>
- Alnaimy M, Zelenakova M, Vranayova Z, Abu-Hashim M. 2020. Effects of Temporal Variation in Long-Term Cultivation on Organic Carbon Sequestration in Calcareous Soils: Nile Delta, Egypt. *Sustainability*. 12(11):4514. <https://doi.org/10.3390/su12114514>
- [Andersson B. 1983. Odlingstekniska försök med höstvet. Institutionen för Växtodling, Sveriges lantbruksuniversitet, Uppsala.\(121\).](#)
- Atzberger C, Richter K, Vuolo F, Darvishzadeh R, Schlerf M. 2011. Why confining to vegetation indices? Exploiting the potential of improved spectral observations using radiative transfer models. In: *Remote Sensing for Agriculture, Ecosystems, and Hydrology XIII*. Vol. 8174. ~~[place unknown]~~: SPIE; p. 263–278. <https://doi.org/10.1117/12.898479>
- [Bakke SJ, Ionita M, Tallaksen LM. 2020. The 2018 northern European hydrological drought and its drivers in a historical perspective. *Hydrol Earth Syst Sci*. 24\(11\):5621–5653. https://doi.org/10.5194/hess-24-5621-2020](#)
- [Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*. 67:1–48. https://doi.org/10.18637/jss.v067.i01](#)
- Beillouin D, Schauburger B, Bastos A, Ciais P, Makowski D. 2020. Impact of extreme weather conditions on European crop production in 2018. *Philos Trans R Soc Lond B Biol Sci*. 375(1810):20190510. <https://doi.org/10.1098/rstb.2019.0510>
- Bengough AG, McKenzie BM, Hallett PD, Valentine TA. 2011. Root elongation, water stress, and mechanical impedance: a review of limiting stresses and beneficial root tip traits. *Journal of Experimental Botany*. 62(1):59–68. <https://doi.org/10.1093/jxb/erq350>

- Bhattacharya A. 2019. Chapter 3 - Water-Use Efficiency Under Changing Climatic Conditions. In: Bhattacharya A, editor. Changing Climate and Resource Use Efficiency in Plants. ~~[place unknown]~~ Academic Press; p. 111–180. <https://doi.org/10.1016/B978-0-12-816209-5.00003-9>
- Bodner G, Nakhforoosh A, Kaul H-P. 2015. Management of crop water under drought: a review. *Agron Sustain Dev.* 35(2):401–442. <https://doi.org/10.1007/s13593-015-0283-4>
- Boedhram N, Arkebauer TJ, Batchelor WD. 2001. Season-Long Characterization of Vertical Distribution of Leaf Area in Corn. *Agronomy Journal.* 93(6):1235–1242. <https://doi.org/10.2134/agronj2001.1235>

Chen S, Du T, Wang S, Parsons D, Wu D, Guo X, Li D. 2021. Quantifying the effects of spatial-temporal variability of soil properties on crop growth in management zones within an irrigated maize field in Northwest China. *Agricultural Water Management*. 244:106535. <https://doi.org/10.1016/j.agwat.2020.106535>

Chen Y, Carver BF, Wang S, Zhang F, Yan L. 2009. Genetic loci associated with stem elongation and winter dormancy release in wheat. *Theor Appl Genet*. 118(5):881–889. <https://doi.org/10.1007/s00122-008-0946-5>

Chowdhury S, Bolan N, Farrell M, Sarkar B, Sarker JR, Kirkham MB, Hossain MZ, Kim G-H. 2021. Chapter Two - Role of cultural and nutrient management practices in carbon sequestration in agricultural soil. In: Sparks DL, editor. *Advances in Agronomy*. Vol. 166. [place-unknown]: Academic Press; p. 131–196. <https://doi.org/10.1016/bs.agron.2020.10.001>

Clevers JGPW, Kooistra L, Van den Brande MMM. 2017. Using Sentinel-2 Data for Retrieving LAI and Leaf and Canopy Chlorophyll Content of a Potato Crop. *Remote Sensing*. 9(5):405. <https://doi.org/10.3390/rs9050405>

Colombi T, Torres LC, Walter A, Keller T. 2018. Feedbacks between soil penetration resistance, root architecture and water uptake limit water accessibility and crop growth – A vicious circle. *Science of The Total Environment*. 626:1026–1035. <https://doi.org/10.1016/j.scitotenv.2018.01.129>
<https://doi.org/10.1016/j.scitotenv.2018.01.129>

[Danner M, Berger K, Woche M, Mauser W, Hank T. 2021. Efficient RTM-based training of machine learning regression algorithms to quantify biophysical & biochemical traits of agricultural crops. ISPRS Journal of Photogrammetry and Remote Sensing. 173:278–296. https://doi.org/10.1016/j.isprsjsprs.2021.01.017](https://doi.org/10.1016/j.isprsjsprs.2021.01.017)

[Ding DY, Feng H, Zhao Y, He JQ, Zou YF, Jin JM. 2016. Modifying Winter Wheat Sowing Date as an Adaptation to Climate Change on the Loess Plateau. Agronomy Journal. 108\(1\):53–63. https://doi.org/10.2134/agronj15.0262](https://doi.org/10.2134/agronj15.0262)

Dong T, Liu Jiangui, Qian B, He L, Liu Jane, Wang R, Jing Q, Champagne C, McNairn H, Powers J, et al. 2020. Estimating crop biomass using leaf area index derived from Landsat 8 and Sentinel-2 data. *ISPRS Journal of Photogrammetry and Remote Sensing*. 168:236–250. <https://doi.org/10.1016/j.isprsjsprs.2020.08.003>

~~[Drusch M, Del Bello U, Carlier S, Colin O, Fernandez V, Gascon F, Hoersch B, Isola C, Laberinti P, Martimort P, et al. 2012. Sentinel 2: ESA's Optical High Resolution Mission for GMES Operational Services. Remote Sensing of Environment. 120:25–36. https://doi.org/10.1016/j.rse.2011.11.026](https://doi.org/10.1016/j.rse.2011.11.026)~~

[Doole G, Tozer K, Sauermaann C, Stevens D, Ward J. 2023. Challenges and opportunities for conducting on-farm research. Journal of New Zealand Grasslands.:85–93. https://doi.org/10.33584/jnzg.2023.85.3659](https://doi.org/10.33584/jnzg.2023.85.3659)

Fageria NK. 2012. Role of Soil Organic Matter in Maintaining Sustainability of Cropping Systems. *Communications in Soil Science and Plant Analysis*. 43(16):2063–2113. <https://doi.org/10.1080/00103624.2012.697234>

Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins S, Saud S, et al. 2017. Crop Production under Drought and Heat Stress: Plant Responses and Management Options. *Frontiers in Plant Science*. 8.

Feng W, Wu Y, He L, Ren X, Wang Yangyang, Hou G, Wang Yonghua, Liu W, Guo T. 2019. An optimized non-linear vegetation index for estimating leaf area index in winter wheat. *Precision Agric*. 20(6):1157–1176. <https://doi.org/10.1007/s11119-019-09648-8>

- Graf LV, Merz QN, Walter A, Aasen H. 2023. Insights from field phenotyping improve satellite remote sensing based in-season estimation of winter wheat growth and phenology. *Remote Sensing of Environment*. 299:113860. <https://doi.org/10.1016/j.rse.2023.113860>
- Graf LV, Perich G, Aasen H. 2022. EOA-team/eodal_notebooks: EOdal Notebooks (Graf et al., 2022) [Internet]. [accessed 2023 Jul 20]. <https://doi.org/10.5281/ZENODO.7278252>
- Gray SB, Brady SM. 2016. Plant developmental responses to climate change. *Developmental Biology*. 419(1):64–77. <https://doi.org/10.1016/j.ydbio.2016.07.023>
- Groemping U, Lehrkamp M. 2023. relaimpo: Relative Importance of Regressors in Linear Models [Internet]. [accessed 2023 Dec 22]. <https://cran.r-project.org/web/packages/relaimpo/index.html>
- He L, Wang R, Mostovoy G, Liu Jane, Chen JM, Shang J, Liu Jianguai, McNairn H, Powers J. 2021. Crop Biomass Mapping Based on Ecosystem Modeling at Regional Scale Using High Resolution Sentinel-2 Data. *Remote Sensing*. 13(4):806. <https://doi.org/10.3390/rs13040806>
- He M, Dijkstra FA. 2014. Drought effect on plant nitrogen and phosphorus: a meta-analysis. *New Phytologist*. 204(4):924–931. <https://doi.org/10.1111/nph.12952><https://doi.org/10.1111/nph.12952>
- [He J, Shi Y, Zhao J, Yu Z, He J, Shi Y, Zhao J, Yu Z. 2020. Strip rotary tillage with subsoiling increases winter wheat yield by alleviating leaf senescence and increasing grain filling. *Crop Journal*. 8\(2\):327–340.](#)
- Hirooka Y, Homma K, Maki M, Sekiguchi K, Shiraiwa T, Yoshida K. 2017. Evaluation of the dynamics of the leaf area index (LAI) of rice in farmer's fields in Vientiane Province, Lao PDR. *J Agric Meteorol*. 73(1):16–21. <https://doi.org/10.2480/agrmet.D-14-00021><https://doi.org/10.2480/agrmet.D-14-00021>
- Huang Z, Liu Y, Tian F-P, Wu G-L. 2020. Soil water availability threshold indicator was determined by using plant physiological responses under drought conditions. *Ecological Indicators*. 118:106740. <https://doi.org/10.1016/j.ecolind.2020.106740>
- Ihsan MZ, El-Nakhlawy FS, Ismail SM, Fahad S, daur I. 2016. Wheat Phenological Development and Growth Studies As Affected by Drought and Late Season High Temperature Stress under Arid Environment. *Frontiers in Plant Science*. 7.
- IPCC. 2022. Climate Change 2022: Impacts, Adaptation and Vulnerability Working Group II Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. ~~[place—unknown]~~. <https://doi.org/10.1017/9781009325844>
- ~~Irfan Ullah M, Mahpara S, Bibi R, Ullah Shah R, Ullah R, Abbas S, Ihsan Ullah M, Hassan AM, El Shehawi AM, Brestic M, et al. 2021. Grain yield and correlated traits of bread wheat lines: Implications for yield improvement. *Saudi J Biol Sci*. 28(10):5714–5719. <https://doi.org/10.1016/j.sjbs.2021.06.006>~~
- Jacquemoud S, Ustin SL, Verdebout J, Schmuck G, Andreoli G, Hosgood B. 1996. Estimating leaf biochemistry using the PROSPECT leaf optical properties model. *Remote Sensing of Environment*. 56(3):194–202. [https://doi.org/10.1016/0034-4257\(95\)00238-3](https://doi.org/10.1016/0034-4257(95)00238-3)

- Kane DA, Bradford MA, Fuller E, Oldfield EE, Wood SA. 2021. Soil organic matter protects US maize yields and lowers crop insurance payouts under drought. *Environ Res Lett.* 16(4):044018. <https://doi.org/10.1088/1748-9326/abe492>
- [Kang S, Zhang F, Hu X, Zhang J. 2002. Benefits of CO2 enrichment on crop plants are modified by soil water status. *Plant and Soil.* 238\(1\):69–77. <https://doi.org/10.1023/A:1014244413067>](https://doi.org/10.1023/A:1014244413067)
- [Kang Y, Özdoğan M, Zipper SC, Román MO, Walker J, Hong SY, Marshall M, Magliulo V, Moreno J, Alonso L, et al. 2016. How Universal Is the Relationship between Remotely Sensed Vegetation Indices and Crop Leaf Area Index? A Global Assessment. *Remote Sensing.* 8\(7\):597. <https://doi.org/10.3390/rs8070597>](https://doi.org/10.3390/rs8070597)
- Kirchmann H, Börjesson G, Bolinder MA, Kätterer T, Djodjic F. 2020. Soil properties currently limiting crop yields in Swedish agriculture – An analysis of 90 yield survey districts and 10 long-term field experiments. *European Journal of Agronomy.* 120:126132. <https://doi.org/10.1016/j.eja.2020.126132>
- Krauss M, Berner A, Perrochet F, Frei R, Niggli U, Mäder P. 2020. Enhanced soil quality with reduced tillage and solid manures in organic farming – a synthesis of 15 years. *Sci Rep.* 10(1):4403. <https://doi.org/10.1038/s41598-020-61320-8>
- [Krueger ES, Ochsner TE. 2024. Traditional matric potential thresholds underestimate soil moisture at field capacity across Oklahoma. *Soil Science Society of America Journal.* 88\(5\):1678–1690. <https://doi.org/10.1002/saj2.20733>](https://doi.org/10.1002/saj2.20733)
- Lal R. 2009. Challenges and opportunities in soil organic matter research. *European Journal of Soil Science.* 60(2):158–169. <https://doi.org/10.1111/j.1365-2389.2008.01114.x>
- Lambert M-J, Traoré PCS, Blaes X, Baret P, Defourny P. 2018. Estimating smallholder crops production at village level from Sentinel-2 time series in Mali's cotton belt. *Remote Sensing of Environment.* 216:647–657. <https://doi.org/10.1016/j.rse.2018.06.036>
- [Lawal S, Sitch S, Lombardozzi D, Nabel JEMS, Wey H-W, Friedlingstein P, Tian H, Hewitson B. 2022. Investigating the response of leaf area index to droughts in southern African vegetation using observations and model simulations. *Hydrology and Earth System Sciences.* 26\(8\):2045–2071. <https://doi.org/10.5194/hess-26-2045-2022>](https://doi.org/10.5194/hess-26-2045-2022)
- Lipiec J, Håkansson I, Tarkiewicz S, Kossowski J. 1991. Soil physical properties and growth of spring barley as related to the degree of compactness of two soils. *Soil and Tillage Research.* 19(2):307–317. [https://doi.org/10.1016/0167-1987\(91\)90098-I](https://doi.org/10.1016/0167-1987(91)90098-I)
- Mahmood S, Nunes MR, Kane DA, Lin Y. 2023. Soil health explains the yield-stabilizing effects of soil organic matter under drought. *Soil & Environmental Health.* 1(4):100048. <https://doi.org/10.1016/j.seh.2023.100048>
- Matiu M, Ankerst DP, Menzel A. 2017. Interactions between temperature and drought in global and regional crop yield variability during 1961-2014. *PLOS ONE.* 12(5):e0178339. <https://doi.org/10.1371/journal.pone.0178339>
- Meinke H, Hammer GL, van Keulen H, Rabbinge R, Keating BA. 1997. Improving wheat simulation capabilities in Australia from a cropping systems perspective: water and nitrogen effects on spring wheat in a semi-arid environment. In: van Ittersum MK, van de Geijn SC, editors. *Developments in Crop Science.* Vol. 25. ~~{place unknown}~~: Elsevier; p. 99–112. [https://doi.org/10.1016/S0378-519X\(97\)80012-8](https://doi.org/10.1016/S0378-519X(97)80012-8)

- Minoli S, Jägermeyr J, Asseng S, Urfels A, Müller C. 2022. Global crop yields can be lifted by timely adaptation of growing periods to climate change. *Nat Commun.* 13:7079. <https://doi.org/10.1038/s41467-022-34411-5>
- ~~Mohan D, Mamrutha HM, Khobra R, Singh G, Singh GP. 2022. Relevance of height, heading and maturity in productivity enhancement of wheat. INDIAN JOURNAL OF GENETICS AND PLANT BREEDING. 82(01):31–37. <https://doi.org/10.31742/IJGPB.82.1.5>~~
- ~~Mroz T, Dieseth JA, Lillemo M. 2023. Grain yield and adaptation of spring wheat to Norwegian growing conditions is driven by allele frequency changes at key adaptive loci discovered by genome wide association mapping. Theor Appl Genet. 136(9):191. <https://doi.org/10.1007/s00122-023-04424-9>~~
- Myneni RB, Ramakrishna R, Nemani R, Running SW. 1997. Estimation of global leaf area index and absorbed par using radiative transfer models. *IEEE Transactions on Geoscience and Remote Sensing.* 35(6):1380–1393. <https://doi.org/10.1109/36.649788>
- Myneni RB, Williams DL. 1994. On the relationship between FAPAR and NDVI. *Remote Sensing of Environment.* 49(3):200–211. [https://doi.org/10.1016/0034-4257\(94\)90016-7](https://doi.org/10.1016/0034-4257(94)90016-7)
- Oldfield EE, Bradford MA, Wood SA. 2019. Global meta-analysis of the relationship between soil organic matter and crop yields. *SOIL.* 5(1):15–32. <https://doi.org/10.5194/soil-5-15-2019>
- Peng Y, Zhu T, Li Y, Dai C, Fang S, Gong Y, Wu X, Zhu R, Liu K. 2019. Remote prediction of yield based on LAI estimation in oilseed rape under different planting methods and nitrogen fertilizer applications. *Agricultural and Forest Meteorology.* 271:116–125. <https://doi.org/10.1016/j.agrformet.2019.02.032>
- ~~Petcu E, Ion T, Pompiliu M, Petcu V. 2011. Effect of organic and conventional farming systems on some physiological indicators of winter wheat. *Romanian Agricultural Research.*(28).~~
- Porter JR, Gawith M. 1999. Temperatures and the growth and development of wheat: a review. *European Journal of Agronomy.* 10(1):23–36. [https://doi.org/10.1016/S1161-0301\(98\)00047-1](https://doi.org/10.1016/S1161-0301(98)00047-1)
- Prabhakara K, Hively WD, McCarty GW. 2015. Evaluating the relationship between biomass, percent groundcover and remote sensing indices across six winter cover crop fields in Maryland, United States. *International Journal of Applied Earth Observation and Geoinformation.* 39:88–102. <https://doi.org/10.1016/j.jag.2015.03.002>
- Punalekar SM, Verhoef A, Quaife TL, Humphries D, Birmingham L, Reynolds CK. 2018. Application of Sentinel-2A data for pasture biomass monitoring using a physically based radiative transfer model. *Remote Sensing of Environment.* 218:207–220. <https://doi.org/10.1016/j.rse.2018.09.028>
- ~~Pużyńska K, Synowiec A, Pużyński S, Bocianowski J, Klima K, Lepiarczyk A. 2021. The Performance of Oat-Vetch Mixtures in Organic and Conventional Farming Systems. *Agriculture.* 11(4):332. <https://doi.org/10.3390/agriculture11040332>~~
- R Core Team. 2022. R: The R Project for Statistical Computing [Internet]. [accessed 2023 Dec 22]. <https://www.r-project.org/>
- Ray RL, Fares A, Risch E. 2018. Effects of Drought on Crop Production and Cropping Areas in Texas. *Agricultural & Environmental Letters.* 3(1):170037. <https://doi.org/10.2134/ael2017.11.0037>

- Revell A, Florence A, MacArthur A, Hoad SP, Rees RM, Williams M. 2019. The Value of Sentinel-2 Spectral Bands for the Assessment of Winter Wheat Growth and Development. *Remote Sensing*. 11(17):2050. <https://doi.org/10.3390/rs11172050>
- Rockström J. 2003. Resilience building and water demand management for drought mitigation. *Physics and Chemistry of the Earth, Parts A/B/C*. 28(20):869–877. <https://doi.org/10.1016/j.pce.2003.08.009>
- Rolli E, Marasco R, Vigani G, Ettoumi B, Mapelli F, Deangelis ML, Gandolfi C, Casati E, Previtali F, Gerbino R, et al. 2015. Improved plant resistance to drought is promoted by the root-associated microbiome as a water stress-dependent trait. *Environmental Microbiology*. 17(2):316–331. <https://doi.org/10.1111/1462-2920.12439>
- Ru C, Hu X, Chen D, Wang W, Zhen J, Song T. 2023. Individual and combined effects of heat and drought and subsequent recovery on winter wheat (*Triticum aestivum* L.) photosynthesis, nitrogen metabolism, cell osmoregulation, and yield formation. *Plant Physiology and Biochemistry*. 196:222–235. <https://doi.org/10.1016/j.plaphy.2023.01.038>
- Rubin RL, van Groenigen KJ, Hungate BA. 2017. Plant growth promoting rhizobacteria are more effective under drought: a meta-analysis. *Plant Soil*. 416(1):309–323. <https://doi.org/10.1007/s11104-017-3199-8>
- SCB. [1993. Jordbruksstatistisk årsbok 1993. Bulls Tryckeriaktiebolag.](#)
- SCB. 2018. Production of cereal crops, dried pulses and oilseed crops in 2018. *Statistiska meddelanden*. JO 19 SM 1801.
- Seleiman MF, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid HH, Battaglia ML. 2021. Drought Stress Impacts on Plants and Different Approaches to Alleviate Its Adverse Effects. *Plants*. 10(2). <https://doi.org/10.3390/plants10020259>
- Shankar T, Banerjee M, Malik GC, Dutta S, Maiti D, Maitra S, Alharby H, Bamagoos A, Hossain A, Ismail IA, EL Sabagh A. 2021. The Productivity and Nutrient Use Efficiency of Rice–Rice–Black Gram Cropping Sequence Are Influenced by Location Specific Nutrient Management. *Sustainability*. 13(6):3222. <https://doi.org/10.3390/su13063222>
- Sjulgård H, Colombi T, Keller T. 2022. Spatiotemporal patterns of crop diversity reveal potential for diversification in Swedish agriculture. *Agriculture, Ecosystems & Environment*. 336:108046. <https://doi.org/10.1016/j.agee.2022.108046>
- Sjulgård H, Keller T, Garland G, Colombi T. 2023. Relationships between weather and yield anomalies vary with crop type and latitude in Sweden. *Agricultural Systems*. 211:103757. <https://doi.org/10.1016/j.agsy.2023.103757>
- [Sjulgård H. 2024. The potential of agricultural management to alleviate extreme weather impacts on Swedish crop production. Acta Universitatis Agriculturae Sueciae \(2024:84\). https://doi.org/10.54612/a.107ri2j3pt](#)
- Skakun S, Vermote E, Franch B, Roger J-C, Kussul N, Ju J, Masek J. 2019. Winter Wheat Yield Assessment from Landsat 8 and Sentinel-2 Data: Incorporating Surface Reflectance, Through Phenological Fitting, into Regression Yield Models. *Remote Sensing*. 11(15):1768. <https://doi.org/10.3390/rs11151768>~~https://doi.org/10.3390/rs11151768~~<https://doi.org/10.3390/rs11151768>
- SMHI. 2018. [Maj 2018 - Sommarväder med rekordvärme | SMHI \[Internet\]. \[accessed 2024 Oct 3\]. https://www.smhi.se/klimat/klimatet-da-och-nu/manadens-vader-och-vatten-sverige/manadens-vader-i-sverige/maj-2018-sommarvader-med-rekordvarme-1.134781](#)
- SMHI. 2023. [Nedladdning av griddad nederbörd- och temperaturdata \(PTHBV\) \[Internet\]. \[accessed 2023 Mar 8\]. https://www.smhi.se/data/ladda-ner-data/griddade-nederbord-och-temperaturdata-ptbvy](#)

- Stockdale E a., Shepherd M a., Fortune S, Cuttle S p. 2002. Soil fertility in organic farming systems – fundamentally different? *Soil Use and Management*. 18(s1):301–308. <https://doi.org/10.1111/j.1475-2743.2002.tb00272.x>
- ~~Tewelde H, Fernandez CJ, Erickson CA. 2006. Wheat Cultivars Adapted to Post-Heading High Temperature Stress. *Journal of Agronomy and Crop Science*. 192(2):111–120. <https://doi.org/10.1111/j.1439-037X.2006.00189.x>~~
- ~~Stenberg M, Bjurling E, Gruvaeus I, Gustafsson K. 2005. *Gödslingsrekommendationer och optimala kvävegivor för lönsamhet och kväveeffektivitet i praktisk spannmålsodling*. Skara: SLU.~~
- ~~Sun Q, Gilgen AK, Wittwer R, von Arx G, van der Heijden MGA, Klaus VH, Buchmann N. 2024. Drought effects on trait space of winter wheat are independent of land management. *New Phytol*. 243(2):591–606. <https://doi.org/10.1111/nph.19851>~~
- Verhoef W. 1998. Theory of radiative transfer models applied in optical remote sensing of vegetation canopies [PhD]. S.l.
- Villegas D, Aparicio N, Blanco R, Royo C. 2001. Biomass Accumulation and Main Stem Elongation of Durum Wheat Grown under Mediterranean Conditions. *Annals of Botany*. 88(4):617–627. <https://doi.org/10.1006/anbo.2001.1512>
- de Vries FT, Griffiths RI, Knight CG, Nicolitch O, Williams A. 2020. Harnessing rhizosphere microbiomes for drought-resilient crop production. *Science*. 368(6488):270–274. <https://doi.org/10.1126/science.aaz5192>
- Waldner F, Horan H, Chen Y, Hochman Z. 2019. High temporal resolution of leaf area data improves empirical estimation of grain yield. *Sci Rep*. 9(1):15714. <https://doi.org/10.1038/s41598-019-51715-7>
- Wang B, Liu DL, Asseng S, Macadam I, Yu Q. 2017. Modelling wheat yield change under CO₂ increase, heat and water stress in relation to plant available water capacity in eastern Australia. *European Journal of Agronomy*. 90:152–161. <https://doi.org/10.1016/j.eja.2017.08.005>
- Wang E, Cresswell H, Xu J, Jiang Q. 2009. Capacity of soils to buffer impact of climate variability and value of seasonal forecasts. *Agricultural and Forest Meteorology*. 149(1):38–50. <https://doi.org/10.1016/j.agrformet.2008.07.001>
- Wasaya A, Rehman I, Mohi Ud Din A, Hayder Bin Khalid M, Ahmad Yasir T, Mansoor Javaid M, El-Hefnawy M, Brestic M, Rahman MA, El Sabagh A. 2023. Foliar application of putrescine alleviates terminal drought stress by modulating water status, membrane stability, and yield-related traits in wheat (*Triticum aestivum* L.). *Frontiers in Plant Science*. 13.
- Watson DJ. 1947. Comparative Physiological Studies on the Growth of Field Crops: I. Variation in Net Assimilation Rate and Leaf Area between Species and Varieties, and within and between Years. *Annals of Botany*. 11(1):41–76. <https://doi.org/10.1093/oxfordjournals.aob.a083148>
- ~~Wileke RAI, Kjellström E, Lin C, Matei D, Moberg A, Tyrlis E, Wocher M, Berger K, Danner M, Mauser W, Hank T. 2020. The extremely warm summer RTM-based dynamic absorption integrals for the retrieval of 2018 in Sweden—set in a historical context. *biochemical vegetation traits. International Journal of Applied Earth System Dynamics*. 11(4):1107–1121. *Observation and Geoinformation*. 93:102219. <https://doi.org/10.5194/esd-11-1107-1016/j.jag.2020.102219>~~
- Yamamoto S, Hashimoto N, Homma K. 2023. Evaluation of LAI Dynamics by Using Plant Canopy Analyzer and Its Relationship to Yield Variation of Soybean in Farmer Field. *Agriculture*. 13(3):609. <https://doi.org/10.3390/agriculture13030609>

- Zhang J, Zhang S, Cheng M, Jiang H, Zhang X, Peng C, Lu X, Zhang M, Jin J. 2018. Effect of Drought on Agronomic Traits of Rice and Wheat: A Meta-Analysis. *International Journal of Environmental Research and Public Health*. 15(5):839. <https://doi.org/10.3390/ijerph15050839>
- Zipper SC, Qiu J, Kucharik CJ. 2016. Drought effects on US maize and soybean production: spatiotemporal patterns and historical changes. *Environ Res Lett*. 11(9):094021. <https://doi.org/10.1088/1748-9326/11/9/094021>
- Zuber V, Strimmer K. 2011. High-Dimensional Regression and Variable Selection Using CAR Scores. *Statistical Applications in Genetics and Molecular Biology*. 10(1). <https://doi.org/10.2202/1544-6115.1730>