

DiuSST: A conceptual model of diurnal warm layers for idealized atmospheric simulations with interactive SST

Reyk Börner^{1,2}, Jan O. Haerter^{1,3,4,5}, and Romain Fiévet^{1,6}

¹Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

²Department of Mathematics and Statistics, University of Reading, Reading, UK

³Integrated Modeling, Leibniz Center for Tropical Marine Research, Bremen, Germany

⁴Physics and Earth Sciences, Constructor University Bremen, Bremen, Germany

⁵Department of Physics and Astronomy, University of Potsdam, Potsdam, Germany

⁶Scientific Computing Lab, Max Planck Institute for Meteorology, Hamburg, Germany

Correspondence: Reyk Börner (reyk.boerner@reading.ac.uk)

Abstract. The diurnal variability of sea surface temperature (SST) may play an important role for cloud organization above the tropical ocean, with implications for precipitation extremes, storminess, and climate sensitivity. Recent cloud-resolving simulations demonstrate how imposed diurnal SST oscillations can strongly, and delicately, impact mesoscale convective organization. In spite of this nuanced interaction, many idealized modeling studies of tropical convection either assume a constant, homogeneous SST or, in case of a responsive sea surface, represent the upper ocean by a slab with fixed thickness. Here we show that slab ocean models with constant heat capacity fail to capture the wind-dependence of observed diurnal sea surface warming. To alleviate this shortcoming, we present a simple, yet explicitly depth-resolved model of upper-ocean temperature dynamics under atmospheric forcing. Our modular scheme describes turbulent mixing as diffusion with a wind-dependent diffusivity, in addition to a bulk mixing term and heat fluxes entering as sources and sinks. Using observational data, we apply Bayesian inference to calibrate the model. In contrast with a slab model, our model captures the exponential reduction of the diurnal warming amplitude with increasing wind speed. Further, our model performs comparably to a more elaborately parameterized diurnal warm layer model. Formulated as a single partial differential equation with three key tuning parameters, the model is suitable as an interactive numerical boundary condition for idealized atmospheric simulations.

1 Introduction

The role of clouds in a changing climate remains an open question that challenges predictions of climate sensitivity, regional precipitation patterns and extreme weather events (Bony et al., 2015; Siebesma et al., 2020). Fundamental processes in cloud dynamics remain insufficiently understood, particularly how convective clouds cluster and interact with their environment. This knowledge gap calls for idealized modeling approaches simple enough to distill mechanisms, yet relatable to the real world.

In the tropics, clouds organize across a wide range of spatial and temporal scales (Moncrieff, 2010). Diurnally, thunderstorms often cluster in mesoscale convective systems (MCSs), which are associated with extreme precipitation and the genesis of tropical cyclones (Tan et al., 2015; Schumacher and Rasmussen, 2020). Intraseasonally, variability is dominated by the

Madden-Julian Oscillation (MJO), an eastward-propagating zone of strong deep convective activity (Madden and Julian, 1972; Zhang, 2005). Although the MJO is known to couple to the large-scale circulation and impact weather around the globe, it remains difficult to model (Jiang et al., 2020; DeMott et al., 2015).

25 Due to the multiscale interaction of tropical convection, small-scale processes may be key to understanding large-scale patterns (Slingo et al., 2003). Indeed, it is increasingly acknowledged that the diurnal variability of sea surface temperature (SST) can play an important role for atmospheric dynamics (Li et al., 2001; Clayson and Chen, 2002; Bernie et al., 2008; Bellenger et al., 2010; Haerter et al., 2020). Observations draw a clear link between strong diurnal SST oscillations and a diurnal cycle of marine cumulus convection (Johnson et al., 1999). Furthermore, the diurnal cycle of SST, by enhancing air-sea
30 heat transfer, could help trigger the active phase of the MJO (Seo et al., 2014; Woolnough et al., 2007; Zhang, 2005; Zhao and Nasuno, 2020; Karlowska et al., 2023). Neglecting diurnal SST variations may lead to a bias on the order of 10 W m^{-2} in monthly-averaged surface heat fluxes (Weihs and Bourassa, 2014), and even stronger biases on shorter timescales. These findings emphasize the relevance of resolving diurnal air-sea interactions when modeling convective organization.

However, many idealized modeling studies of tropical convection prescribe a constant SST in space and time. A popular modeling framework is *radiative-convective equilibrium* (RCE), where constant solar forcing is balanced by outgoing longwave radiation above a constant, homogeneous SST (Tompkins and Craig, 1998). Under RCE, the moisture field is known to spontaneously self-organize into convective clusters separated by extended dry regions (Bretherton et al., 2005). This mechanism termed *convective self-aggregation* has been associated with real-world features such as MCS formation. Yet, self-aggregation is hampered in the realistic limit of fine horizontal model resolution, calling the realism of the mechanism into question (Yanase
40 et al., 2020). Recent studies, imposing a diurnal oscillation of SST, demonstrate the emergence of *diurnal self-aggregation* even at fine spatial resolution (Haerter et al., 2020; Jensen et al., 2022). Likewise, spatial variations in SST have been shown to imprint themselves on the moisture field (Müller and Hohenegger, 2020; Shamekh et al., 2020b; Skyllingstad et al., 2019). Since SST variability can substantially alter the spatio-temporal patterns of marine tropical convection, there is a need to incorporate higher-fidelity SST representations when investigating convective processes.

45 Over the past years, the modeling community studying convective organization has largely addressed the issue of an interactive SST by coupling the atmosphere to a single-layer slab with fixed heat capacity, which absorbs and re-emits heat according to parameterized surface fluxes (Hohenegger and Stevens, 2016; Shamekh et al., 2020a; Coppin and Bony, 2017; Tompkins and Semie, 2021; Wing et al., 2017). These works report overall that a responsive SST slows down or even prevents the onset of convective aggregation. However, in this paper we show that slab models are inadequate for producing realistic diurnal SST
50 warming: they fail to capture its wind-dependence by neglecting upper ocean mixing. Yet, by no means can wind be neglected even in strongly idealized studies, given that cold pools, bringing gusty winds, are increasingly appreciated as an integral mechanism in convective self-organization (Tompkins, 2001; Böing, 2016; Haerter, 2019; Nissen and Haerter, 2021). To address this inconsistency, we present a simplified 1D model that significantly improves the representation of diurnal SST variability compared to a slab ocean while being adaptable and affordable for coupling to cloud-resolving atmospheric models.

55 SST plays a key role in governing the heat and moisture exchange between the atmosphere and ocean (Seo et al., 2023). Whereas observed diurnal amplitudes of surface temperature are typically largest over land, a diurnal cycle of SST is also

common throughout the tropical ocean (Kawai and Wada, 2007). Under strong insolation and calm conditions, a diurnal warm layer forms during the day, raising skin SST (measured directly at the surface) by up to 3-4 K. Field studies have observed extreme diurnal warming events of 5 K or more, though these often lie in the extra-tropics or coastal regions (Gentemann et al., 2008; Minnett, 2003; Ward, 2006). Importantly, the amplitude of diurnal warming is highly sensitive to wind speed, with observations suggesting an approximately exponential decay of diurnal warming with increasing wind speed (Gentemann et al., 2003; Börner, 2021).

To first order, the diurnal warm layer results from a competition between insolation-driven thermal stratification and wind-driven mixing. On a calm, clear day after sunrise, the incident solar radiation quickly heats up the upper ocean, creating a stably stratified density profile. This stratification leads to suppressed vertical heat exchange, hence trapping further heat near the surface in a positive feedback which can produce strong surface warming. Later, in the early afternoon when the net surface heat flux changes sign, surface cooling causes unstable density stratification and initiates vertical mixing, resulting in a deepening of the diurnal warm layer. At night, convective mixing typically acts to “reset” the temperature profile. Furthermore, wind stress induces turbulent mixing in the water column. Known as the cool skin effect, the molecular skin layer at the sea surface is typically a few tenths of a degree colder than the water a millimeter deeper (Donlon et al., 2002; Wong and Minnett, 2018).

Upper ocean heat transfer has been modeled on various levels of complexity over the last fifty years, ranging from fully turbulence-resolving models to simplified bulk models and empirical models (Kawai and Wada, 2007). Turbulence-resolving models make up the most realistic but also computationally expensive category (Kondo et al., 1979; Mellor and Yamada, 1982; Large et al., 1994; Kantha and Clayson, 1994; Noh and Jin Kim, 1999; Stull and Kraus, 1987). Though such ocean models have been coupled to a weather forecasting model (Noh et al., 2011), they are arguably too costly and complex for the purpose of idealized atmospheric simulations at high horizontal resolution. The simplest models, derived from empirical relations, specialize on estimating the daily SST amplitude (Webster et al., 1996; Price et al., 1987; Kawai and Kawamura, 2002) or its hourly evolution (Li et al., 2001; Zeng et al., 1999; Gentemann et al., 2003) based on averaged atmospheric data. These models are not designed for the high temporal resolution of atmospheric large-eddy simulations (on the order of seconds), and sometimes not physics-informed.

Alternatively, simplified models based on boundary layer physics have been developed in the spirit of subdividing the water column into layers described by bulk dynamical equations. A popular candidate, which we compare against here, is the prognostic ZB05 scheme by Zeng and Beljaars (2005) with improvements by Takaya et al. (2010). It computes the sea skin temperature from an integrated mixed layer equation combined with a cool skin scheme, assuming a power-law temperature profile. Other bulk models include the PWP model (Price et al., 1986) and its developments (Fairall et al., 1996, 2003; Gentemann et al., 2009; Schiller and Godfrey, 2005). Widely applied in weather and climate modeling (Brunke et al., 2008), such models could offer a suitable balance between physical rigor and computational cost. Nonetheless, they do not necessarily cater to the needs of idealized atmospheric modeling. First, some bulk models require unknown input about the oceanic background state. Second, the integral descriptions of heat and momentum transfer are typically based on elaborate parameterizations, making the models less intuitive and difficult to tune in sensitivity experiments. Third, these models do not resolve the vertical

temperature profile (apart from the parameterized profile in Gentemann et al. (2009)), limiting their use as an interface to sub-surface processes that might be sensitive to diurnal warming, such as microbial ecosystems (Wurl et al., 2017).

In this paper, we present an idealized model of diurnal sea surface warming that explicitly resolves temperature in the upper few meters of the ocean while being conceptually simple and efficient. Derived from first principles as a modified heat equation, the model consists of a single partial differential equation controlled by three key parameters, taking insolation and atmospheric conditions as forcing input. After describing the model (section 2), we use a Bayesian approach to calibrate and evaluate the model based on cruise observations from the tropical Eastern Pacific (section 3). We compare the performance of our model to both a typical slab model and the ZB05 scheme (section 4). Whereas slab models fail to describe diurnal SST variability, our model accurately reproduces key features of diurnal warming, such as its wind dependence, near-surface heat trapping, and skin cooling. Our model performs comparably to the ZB05 scheme, which is more rigorously derived but does not resolve temperature vertically. In section 4.4, we discuss the implications of our results for the atmospheric heat and moisture budget. We believe that our modular scheme can be useful both as a wind-responsive ocean surface for cloud-resolving modeling and as an interface to study further air-sea-biosphere interactions in an idealized setting.

2 Model description

Our approach relies on the following basic assumptions. First, we assume that the oceanic, atmospherically-driven diurnal temperature variability is constrained to within a few meters from the sea surface. We thus define the *foundation depth* z_f at which diurnal temperature changes become negligible. Conceptually, this partitions the water column into the *diurnal layer* above z_f and the *foundation layer* below. Second, we assume that we may separate the scales of diurnal warming from the ocean's slow internal temperature variability. The foundation layer may then be considered as an infinite heat reservoir with constant *foundation temperature* T_f . Third, we neglect any horizontal inhomogeneities and flows, allowing us to treat the problem as one-dimensional. The goal is then to determine the evolution of the vertical temperature profile within the diurnal layer, in response to a given time sequence of atmospheric forcing. A schematic sketch of the setup is depicted in Fig. 1.

2.1 Main equation

We consider the sea temperature $T(z, t)$ as a function of time $t \geq 0$ and vertical coordinate $z \in (z_f, 0]$, where $z = 0$ defines the air-sea interface and $z < 0$ is below the surface. Given an initial profile $T(z, 0)$, foundation temperature T_f , and atmospheric forcing $\mathcal{F}(t)$, we propose a single prognostic partial differential equation (PDE) for the time evolution of $T \equiv T(z, t)$:

$$\frac{\partial T}{\partial t} = \underbrace{\frac{\partial}{\partial z} \left(\kappa(z, t) \frac{\partial T}{\partial z} \right)}_{\text{diffusion}} - \underbrace{\mu \frac{T - T_f}{z - z_f}}_{\text{mixing}} + \underbrace{\frac{1}{\rho_w c_p} \frac{\partial Q(z, t)}{\partial z}}_{\text{source/sink}}. \quad (1)$$

Here $\kappa > 0$ represents the (time- and depth-dependent) diffusivity, $\mu > 0$ is a constant which we term mixing coefficient, and the constants ρ_w and c_p denote the density and specific heat capacity at constant pressure of sea water, respectively. Finally, Q




Figure 1. Schematic of the simplified upper ocean model, illustrating the processes considered.

120 is the net vertical heat flux at depth z , defined positive downwards (into the ocean). Explicit expressions of these quantities in relation to the forcing \mathcal{F} follow below.

The three terms on the right-hand side of equation (1) provide strongly idealized representations of different physical processes. Motivated by the heat equation, the first term describes the vertical *diffusion* of heat due to turbulent eddies within the diurnal layer. In addition, internal processes such as convection may cause additional vertical *mixing* of water masses between the diurnal layer and the foundation layer. This is crudely incorporated by the second term, which relaxes T to the foundation temperature with relaxation time scale $t_\mu = (z - z_f)/\mu$. Here the depth dependence reflects the intuition that water near the foundation depth will mix faster with foundation-layer water compared to water near the surface. Lastly, the third term comprises all *sources and sinks* of heat, both at the air-sea interface and within the diurnal layer.

130 The foundation temperature $T(z_f, t) = T_f \forall t$ acts as a Dirichlet boundary condition at z_f . At the air-sea interface ($z = 0$), the sea temperature evolves according to the net surface heat flux $Q_0(t) \equiv Q(0, t)$ (see section 2.3), which closes the energy budget.

2.2 Wind-driven mixing and stratification

Wind stress at the sea surface induces shear instability, causing vertical turbulent heat transport in the upper ocean. This implies that the diffusivity $\kappa(z, t)$ in our model should depend on the wind speed $u(t)$. Since the magnitude of wind stress on a water surface is approximately proportional to the square of the wind speed (Smith, 1988; Edson et al., 2013), we propose to model the diffusivity κ as

$$\kappa(z, t) := \kappa_{\text{mol}} + \kappa_0 \varphi(z) \left(\frac{u(t)}{u_0} \right)^2, \quad (2)$$

where κ_{mol} and κ_0 are coefficients of molecular and (vertical) eddy diffusion, respectively. Typically, molecular heat conduction is negligible ($\kappa_{\text{mol}} \ll \kappa_0$). Note the scaling of the eddy diffusion term with u^2 , which we non-dimensionalize by the reference wind speed $u_0 = 1 \text{ m s}^{-1}$ simply to ensure that κ_0 has units of diffusivity.

140 The vertical diffusivity profile $\varphi(z)$ in Eq. (2) approximates how turbulent mixing varies with depth. Physically, thermal density stratification inhibits turbulent heat transfer, since vertical mixing of a stratified fluid is energetically unfavorable. Additionally, the characteristic size of eddies decreases when approaching the air-sea boundary (Pope, 2000). Both effects motivate suppressing the diffusivity near the surface, where stratification becomes largest during diurnal heating. For simplicity, we limit our study to a time-independent linear profile,

$$\varphi(z) = 1 + \sigma \left(\frac{z}{z_f} - 1 \right), \quad (3)$$

145 parameterized by the surface suppressivity $\sigma \in [0, 1]$. If $\sigma > 0$, then φ increases linearly with depth from $\varphi(0) = 1 - \sigma$ at the surface to $\varphi(z_f) = 1$ at the foundation depth.

It may seem counter-intuitive to decrease the diffusivity towards the surface when it is driven by surface winds. In this idealized setup, however, the diffusion term simultaneously accounts for density stratification (in a time-averaged sense). Furthermore, surface wind stress can accelerate the warm layer, leading to turbulence due to shear instability at its base (Hughes
150 et al., 2020). Alternative choices for the diffusivity profile are discussed in section 5 and the appendix.

2.3 Air-sea interaction

In our model, the state of the atmosphere enters as the forcing $\mathcal{F}(t) \equiv (R_{sw,\downarrow}(t), u(t), T_a(t), q_v(t))$, comprising the downward solar, or “shortwave”, irradiance $R_{sw,\downarrow}$, horizontal wind speed u , air temperature T_a , and specific humidity q_v . These quantities refer to some reference height above the sea surface (usually 10 m, where they are typically measured (Friehe and
155 Schmitt, 1976). The diurnal layer interacts with the atmosphere through the absorption and reflection of shortwave radiation, the absorption and emission of thermal (“longwave”) radiation, as well as sensible and latent heat exchange. While shortwave radiation penetrates the sea surface and is absorbed at a range of depths, the other fluxes act within micrometers of the air-sea interface (Wong and Minnett, 2018).

At the air-sea interface, the net surface heat flux $Q_0(t)$ entering the water body is given by

$$Q_0(t) = R_{sw}(t) + R_{lw}(t) + Q_s(t) + Q_l(t), \quad (4)$$

160 where $R_{sw} = (1 - \mathcal{R})R_{sw,\downarrow}$ is the penetrating shortwave irradiance after subtracting from $R_{sw,\downarrow}$ the fraction \mathcal{R} that is reflected at the sea surface (see appendix). The net longwave radiative flux R_{lw} and sensible heat flux Q_s can point into or out of the ocean, whereas the latent heat flux Q_l is always negative. We use standard bulk formulae to describe the surface heat fluxes

(Friehe and Schmitt, 1976; DeCosmo et al., 1996; Wells and King-Hele, 1990; Large and Yeager, 2009),

$$R_{lw}(t) = \sigma_{SB} (T_a(t)^4 - T(0,t)^4), \quad (5a)$$

$$Q_s(t) = \rho_a c_{p,a} C_s u(t) (T_a(t) - T(0,t)), \quad (5b)$$

$$Q_l(t) = \rho_a C_l L_v u(t) (q_v(t) - q_{sat}(T(0,t))), \quad (5c)$$

given in terms of the Stefan-Boltzmann constant σ_{SB} , the density and specific heat capacity of air, ρ_a and $c_{p,a}$, respectively, as well as the Stanton number C_s , Dalton number C_l , and latent heat of vaporization L_v . We approximate these coefficients by constants based on literature values (see table 1), which are roughly valid for wind speeds on the order of 1 to 10 m s^{-1} and air-sea temperature differences around 1 K, measured 10 m above sea level (Wells and King-Hele, 1990). Note that Eq. (5a) simply applies the Stefan-Boltzmann law for black body radiation. When coupled to an atmospheric model, the downward longwave radiative flux can be taken from the atmospheric model output. Furthermore, the latent heat flux, Eq. (5c), involves the saturation specific humidity $q_{sat}(T)$. The temperature dependence of q_{sat} obeys the Clausius-Clapeyron relation, which is approximated by the empirical formula

$$q_{sat}(T) \approx \frac{611.2}{\rho_a r_w T} \exp\left(\frac{17.67(T - 273.15)}{T - 29.65}\right), \quad (6)$$

where r_w denotes the gas constant of water vapor and T enters in units of kelvins (Alduchov and Eskridge, 1996).

We assume that the penetrating shortwave radiation R_{sw} is attenuated exponentially as it propagates downward through the water column,

$$Q(z,t) = R_{sw}(t) \exp\left(\frac{\alpha z}{\cos \phi'(t)}\right) \quad \text{for } z < 0, \quad (7)$$

where $\alpha > 0$ is the *attenuation coefficient* and ϕ' denotes the refracted solar angle. It follows from the sun's angle relative to the surface normal, $\phi \in [0, \pi/2)$, by Snell's refraction law,

$$\phi'(t) = \arcsin\left(\frac{n_a}{n_w} \sin \phi(t)\right), \quad (8)$$

where n_w and n_a denote the refractive indices of sea water and air, respectively.

To summarize, our idealized model is controlled by three key parameters: (i) the eddy diffusivity κ_0 governs the magnitude of wind-driven turbulent heat diffusion; (ii) the mixing coefficient μ sets the time scale of relaxation to the foundation temperature; and (iii) the attenuation coefficient α regulates how deep shortwave radiation penetrates. Realistic values of these parameters are estimated from observational evidence (section 3). We list all model constants and coefficients in Table 1.

2.4 Numerical implementation

To solve Eq. (1), we discretize the spatial coordinate z and numerically integrate the resulting system of ordinary differential equations in time (see Appendix A and Börner (2024) for a software implementation in Python). Spatial derivatives are approximated by second-order accurate finite differences on a non-uniform vertical grid. We set the foundation depth to $z_f = -10\text{m}$,

Quantity	Symbol	Unit	Value
Eddy diffusivity	κ_0	$\text{m}^2 \text{s}^{-1}$	1.34×10^{-4} *
Mixing coefficient	μ	m s^{-1}	2.85×10^{-3} *
Attenuation coefficient	α	m^{-1}	3.52 *
Surface suppressivity	σ	–	0.8
Foundation temperature	T_f	K	298.19 **
Foundation depth	z_f	m	–10
Molecular diffusivity (water)	κ_{mol}	$\text{m}^2 \text{s}^{-1}$	1×10^{-7}
Specific heat (water)	c_p	$\text{J K}^{-1} \text{kg}^{-1}$	3850
Specific heat (air)	$c_{p,a}$	$\text{J K}^{-1} \text{kg}^{-1}$	1005
Density (water)	ρ_w	kg m^{-3}	1027
Density (air)	ρ_a	kg m^{-3}	1.1
Refractive index (water)	n_w	–	1.34
Refractive index (air)	n_a	–	1.00
Stanton number	C_s	–	1.3×10^{-3}
Dalton number	C_i	–	1.5×10^{-3}
Latent heat of vaporization	L	J kg^{-1}	2.5×10^6
Stefan-Boltzmann const.	σ_{SB}	$\text{W m}^{-2} \text{K}^{-4}$	5.67×10^{-8}
Gas constant (water vapor)	r_w	$\text{J K}^{-1} \text{kg}^{-1}$	461.51
Grid spacing at surface	Δz_0	m	0.1
Number of vertical grid points	N	–	40

* determined from data via Bayesian inference, see table E1.

** corresponds to mean 3 m water temperature of MOCE-5 data set

Table 1. Model constants and their default values used.

where diurnal temperature variability is mostly negligible (Kawai and Wada, 2007). The grid spacing is set to $\Delta z_0 = 0.1$ m at the sea surface and increases by a stretch factor $\epsilon \approx 1.04$ with each consecutive grid point below, such that a total of $N = 40$ grid points cover the diurnal layer $z \in (z_f, 0]$ (excluding the boundary point at z_f). Specifically, the depth z_n of the n -th grid point is given by

$$z_n = -\Delta z_0 \left(\frac{1 - \epsilon^n}{1 - \epsilon} \right), \quad n \in \{0, \dots, N\}, \quad (9)$$

190 with ϵ set such that $z_N = z_f$.

We implement the time integration as an explicit Euler scheme, which is numerically stable if the chosen time step Δt satisfies the Courant-Friedrichs-Lewy (CFL) condition,

$$C_i := \max_n \left[2 \frac{\kappa(z_n, t_i) \Delta t_i}{(\Delta z_n)^2} \right] \leq 1, \quad n = 0, 1, 2, \dots, N-1, \quad (10)$$

where n and i index the discrete space and time coordinates, respectively. Via the time-dependent diffusivity κ , the CFL condition depends on time, specifically on the current wind speed. To minimize computational cost, we use an adaptive time step Δt_i that maintains a maximal CFL number of $\mathcal{C}_i = 0.95$ at each instant of time t_i . Consequently, the time step Δt_i scales inversely with the square of the wind speed. We impose a cut-off wind speed $u_{\max} = 10 \text{ m s}^{-1}$, causing any wind speed $u > u_{\max}$ to be replaced by u_{\max} when computing the diffusivity (eq. (2)). In the model simulations of this study, Δt_i ranged from 0.004 to 10 s with a median time step of 5-6 s.

Comparing the explicit Euler scheme with analogous implementations of the fourth-order Runge-Kutta and implicit Euler methods, we find that the explicit Euler scheme is fastest unless the implicit method is used at very large time step, in which case the solution lacks accuracy.

3 Model calibration and evaluation

Our model parameterizes diurnal warming in terms of three unknown constants: the diffusivity coefficient κ_0 , mixing coefficient μ , and attenuation coefficient α . In principle, their values depend on the ocean properties at a given time and location. In this section, we first analyze how the parameters affect diurnal warming, and then use observational data to estimate realistic values via Bayesian inference. Finally, we evaluate the performance of the calibrated model against observations.

3.1 Parameter sensitivity under idealized forcing

As an initial step, we perform a sensitivity study where we force the model with idealized atmospheric pseudo-data representing a calm and clear tropical day. Setting the foundation temperature to $T_f = 300 \text{ K}$, we let the air temperature $T_a(t)$ oscillate harmonically around T_f with a diurnal amplitude¹ of $\Delta T = 2 \text{ K}$. Similarly, we impose a harmonically oscillating horizontal wind speed with a mean of $\bar{u} = 2 \text{ m s}^{-1}$ and amplitude $\Delta u = 2 \text{ m s}^{-1}$, peaking at midnight (this choice generates favorable conditions for diurnal warming and nighttime mixing). Such profiles read:

$$T_a(t) = T_f - \frac{\Delta T}{2} \cos(2\pi t/t_0); \quad u(t) = \bar{u} + \frac{\Delta u}{2} \cos(2\pi t/t_0), \quad (11)$$

where t is local sun time (with respect to midnight) and $t_0 = 1 \text{ d}$. The solar irradiance is given by

$$R_{\text{sw},\downarrow}(t) = \begin{cases} -R_{\text{max}} \cos(2\pi t/t_0) & \text{if } \cos(2\pi t/t_0) < 0 \\ 0 & \text{otherwise,} \end{cases} \quad (12)$$

where we set the peak insolation $R_{\text{max}} = 1000 \text{ W m}^{-2}$. Lastly, we fix the specific humidity at a constant value of $q_v = 15 \text{ g kg}^{-1}$. Under these conditions, we may expect peak diurnal surface warming of around 2-3 K (Minnett, 2003).

Using the atmospheric pseudo-data described above, we now run two-day-long model simulations, consecutively varying one model parameter of the set $\{\kappa_0, \mu, \alpha\}$ while fixing the other two at select default values (see Fig. 2). Increasing the eddy

¹This amplitude is about twice the diurnal amplitude recorded in the observational dataset of this study, where the air temperature oscillates by around 1 K with the maximum in the afternoon.




Figure 2. Model sensitivity under variation of the parameters κ_0 , μ , and α . In each sensitivity experiment, we force the model with two days of idealized atmospheric data (see main text), successively varying one parameter while fixing the other two. The top panels depict simulated surface warming ΔSST as a function of time for different values of the eddy diffusivity (left), mixing coefficient (center), and attenuation coefficient (right). The bottom panels show corresponding vertical temperature profiles at 2 a.m. (dashed lines, shifted by -1 K) and 2 p.m. (solid lines) on the second day of simulation. Values of the varied parameter are given in the respective legend; fixed parameter values are $\kappa_0 = 1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$, $\mu = 1 \times 10^{-3} \text{ m s}^{-1}$, and $\alpha = 3 \text{ m}^{-1}$. Further model settings are detailed in table 1 (here $T_f = 300 \text{ K}$).

diffusivity κ_0 enhances the heat transport within the diurnal layer, flattening the vertical temperature gradient, diminishing surface warming and advancing the time of peak surface warming. A decrease in the mixing coefficient μ corresponds to a slower removal of heat from the diurnal layer into the deeper ocean. This causes increased heat trapping in the upper ocean, hence stronger surface warming and a deeper warm layer. Particularly, for $\mu < 10^{-3} \text{ m s}^{-1}$, excess heat remains in the diurnal layer throughout the night, accumulating heat on the next day. Setting μ to a sufficiently large value ensures that the temperature profile can “reset” at night, as is often observed (Kawai and Wada, 2007). Finally, the attenuation coefficient α determines the depth range in which solar radiation is absorbed. For $\alpha > 1 \text{ m}^{-1}$, more than 60% of radiation is absorbed within 1 m of the surface, leading to strong surface warming. Reducing α causes radiation to reach deeper, where it is more quickly transported into the foundation ocean via the mixing term.

In Fig. 2, the similarity in model response between the parameters κ_0 and α suggests that they might play a similar dynamical role. However, this is not the case. The two parameters control different aspects of the air-sea coupling. The eddy diffusivity κ_0 determines the coupling to wind, whereas the attenuation coefficient α modulates the oceanic heat uptake due to solar radiation. Correlations between model parameters are discussed below (section 3.3).

3.2 Observational data

To infer realistic values for the parameter set $\{\kappa_0, \mu, \alpha\}$, we now conduct a case study where we force the model with real observational data and compare the modeled diurnal warming with the observed signal. Here we use cruise data (available from Börner (2024)) from the Fifth Marine Optical Characterization Experiment (MOCE-5), conducted during October 1999 off the Mexican west coast (Minnett, 2003; Ward, 2006). The route led along the coast of the Baja California peninsula, both in the open Pacific Ocean and within the Gulf of California, thus including offshore as well as more coastal conditions (see Fig. 3a).

The research vessel *Melville* was equipped with an infrared radiometer of type M-AERI (Marine-Atmospheric Emitted Radiance Interferometer, Minnett et al. 2001). This instrument provides precise measurements of sea skin temperature by detecting infrared radiation emitted from within micrometers of the ocean surface. Using skin SST for calibration, rather than bulk SST measured at up to 1 m depth, is crucial when modeling air-sea interactions because the atmosphere senses only the sea skin. Additionally, solar irradiance $R_{\text{sw},\downarrow}$, wind speed u , air temperature T_a , and water temperature at 3 m depth were recorded at time intervals of approximately 10 to 12 minutes (see Fig. 3). Unfortunately, the data set does not contain air humidity. We therefore assume a constant specific humidity of $q_v = 15 \text{ g kg}^{-1}$ throughout the time series, which may be considered typical for the tropical ocean.

Throughout the following analysis, following Minnett (2003), we define *diurnal warming* ΔSST as the temperature difference between the sea skin (that is, the water directly at the surface) and a reference depth d ,

$$\Delta\text{SST}(t) := T(0, t) - T(d, t),$$

where $d = -3 \text{ m}$ for the present data set and $T(0, t)$ is given by the radiometric SST measurements. Diurnal warming events exceeding 1°C are observed on several days, with ΔSST reaching up to 5°C on Oct 13 (Fig. 3c; see also Ward (2006)). The




Figure 3. Observational data set used in this study. a) Travel route of the MOCE-5 cruise in the Pacific Ocean and Gulf of California, colored by the day since departure near San Diego, USA. b) Diurnal warming ΔSST by location, as recorded during the cruise (data points with higher ΔSST are enlarged). c) Time series of observed diurnal warming, ΔSST , showing the individual data points. Gray shaded intervals indicate the training data used for Bayesian inference. Panels d) and e) display the time series of radiometric skin SST (blue), air temperature (red), horizontal wind speed (green), and downward shortwave irradiance (orange). Note that we omit data recorded after Oct 16 due to extended temporal gaps in the data set.

data set also includes days without any substantial diurnal warming, such as on Oct 2 and Oct 9. These days correlate with relatively high wind speeds, while the strong warming events in the second week, such as on Oct 10, 13, and 14, coincide with low winds especially during midday. Note that the time series includes diurnal warming events at different locations (Fig. 3b) and covers a wide range of SST values from 290 to 305 K (Fig. 3d). Horizontal wind speeds (corrected for the ship’s motion) did not exceed 10 m s^{-1} ; downward solar irradiance peaked at around $800\text{ to }900\text{ W m}^{-2}$ each day.

While the MOCE-5 cruise data has a temporal resolution of around 10 minutes, our model requires a time step on the order of seconds to meet the CFL condition (eq. (10)). This necessitates interpolating the atmospheric data between data points; we apply linear interpolation with a wind-adaptive time step as described in section 2.4.

3.3 Bayesian parameter estimation

260 The MOCE-5 observations allow us to estimate the free parameters of our model using Bayesian inference. Given the data \mathcal{D} , the conditional probability $P(\Theta|\mathcal{D})$ that a certain value of the parameter set $\Theta = \{\kappa_0, \mu, \alpha\}$ represents the “true” model follows from Bayes’ rule,

$$P(\Theta|\mathcal{D}) \propto P(\mathcal{D}|\Theta)P(\Theta), \quad (13)$$

that is, the *posterior* distribution $P(\Theta|\mathcal{D})$ is proportional to the product of the *likelihood* $P(\mathcal{D}|\Theta)$ and the *prior* distribution $P(\Theta)$. The likelihood quantifies how well the model with given parameter settings Θ describes the observed data, while the
265 prior distribution encapsulates previous knowledge of suitable parameter ranges (Gelman et al., 2013).

To estimate the posterior distribution, we first partition the data time series into a six-day *training* set and an about eight-day *validation* set (see gray shading in Fig. 3c). The training data are used to evaluate the likelihood function $\mathcal{L}(\Theta) \equiv P(\mathcal{D}|\Theta)$, which is computed in the following way. For given Θ , the model is run using the time series of the variables
270 $\mathcal{D} = \{R_{sw,\downarrow}, u, T_a, q_v\}$ as atmospheric forcing. We set the foundation temperature T_f to the mean observed sea temperature at 3 m depth, $T_f = 298.19\text{K}$, and shift the air temperature T_a according to the current deviation of the 3 m-temperature from its mean T_f . This maintains the observed air-sea temperature contrast at all times. Then, we compare the modeled time series of diurnal warming, $\Delta\text{SST}^{\text{model}}$, to the observed diurnal warming signal $\Delta\text{SST}^{\text{obs}}$ in order to compute the likelihood,

$$\mathcal{L}(\Theta) \propto \exp\left(-\sum_j \frac{(\Delta\text{SST}_j^{\text{model}}(\Theta) - \Delta\text{SST}_j^{\text{obs}})^2}{\Sigma_j^2}\right). \quad (14)$$

275 Here the index j runs through all data points (over time) in the training set, and Σ_j denotes a weight (see appendix).

Approximating the posterior distribution in parameter space is achieved via Markov chain Monte Carlo (MCMC) sampling using the `emcee` package in Python (Foreman-Mackey et al., 2013), based on an affine-invariant algorithm proposed by Goodman and Weare (2010). Further information on the Bayesian inference procedure and choice of prior is provided in the
appendix.

280 After convergence, the sampled posterior distribution shows a single peak in each parameter direction, indicating a well-defined optimal value for each parameter (Fig. 4). From the projected distributions, we compute the maximum, median, and mean value as parameter estimates (tab. E1). The maximum value, or *maximum a posteriori* (MAP) estimate, corresponds to the most likely parameter value and will be used for the subsequent analysis.

The two-dimensional projections of the posterior distribution (Fig. 4) indicate possible correlations between the three pa-
285 rameters. In particular, a positive correlation between κ_0 and α is visible. However, part of the correlation likely reflects the fact that the solar irradiance (modulated by α) and wind speed (modulated by κ_0) are themselves negatively correlated in the training data (not shown).




Figure 4. Posterior distribution of the parameter set $\{\kappa_0, \mu, \alpha\}$, as obtained from Bayesian MCMC sampling. Plots on the upper diagonal are 1D projections of the parameter space, showing for each parameter the prior (orange) and sampled posterior (black) distribution (vertical axis not to scale). Blue lines indicate the MAP values (table E1). The 2D projections of parameter space depict scatter points and smoothed iso-contours of the posterior. Plotted using `corner.py` (Foreman-Mackey, 2016).

3.4 Model performance against observations

After estimating the parameters κ_0 , μ , and α via Bayesian inference, we now evaluate the performance of the calibrated model
290 against the full observational data set, using the settings given in table 1. We force the model with the observed 16-day time
series of solar irradiance, wind speed, and air temperature, interpolated in time to match the CFL condition (Eq. (10)). As in
the previous section, we set the foundation temperature to the observed mean $T_f = 298.19\text{K}$ and adjust the air temperature
time series accordingly.

Fig. 5 illustrates the results of this simulation in comparison with the observed diurnal warming. Similar to the observations,
295 the model produces a wide range of diurnal warming amplitudes from 0.2 K on day 9 to almost 4 K on day 13, which coincides
with the maximal observed diurnal warming of 4.9 K (Fig. 5a). Modeled diurnal warming peaks align in time with the obser-
vations. On days 6 and 7, the model overestimates ΔSST by more than 1 K (see section 5). The variation in diurnal warming
amplitudes links closely to wind speed (Fig. 7b). To some extent, the model captures skin cooling at night, where ΔSST drops
below 0K in agreement with the observations (e.g. on the nights of days 3-4, 6-7, 9-10, and 12-13, see Fig. 5a). Overall, the
300 reduction of ΔSST due to skin cooling is less by about 0.07 K in the model than in observations.

For the whole time series, the modeled and observed ΔSST are correlated with a Pearson correlation coefficient of 0.74 (Fig.
6b). Considering only the training data, the correlation coefficient is 0.82, whereas the value for all data points outside of the
training data is 0.67. This evidences that our model has predictive skill in situations which the model has not been calibrated
to. In particular, the model predicts both the absence of diurnal warming on day 9 and the strong consecutive warming event
305 on day 10 (Fig. 5a). The model performance is further discussed in comparison with other models (sections 4 and 5).

3.5 Depth-resolved temperature profile

Rather than simulating diurnal warming only at the surface, our model provides the vertical temperature profile within the
upper 10 m of the ocean (Figs. 5b, c). In agreement with observations, heat trapping under calm and clear conditions occurs
in the uppermost meter (Soloviev and Lukas, 1997; Ward, 2006; Gentemann et al., 2009). Following peak warming, the model
310 exhibits a deepening of the warm layer between noon and sunset, as seen in Price et al. (1986) and Soloviev and Lukas (1997).

Analyzing the temporal evolution of the temperature profile on days 13 and 14 reveals the close connection to the wind
(Fig. 5b), with increased winds causing a fast diffusion of the warm layer on day 13 after noon and overall weaker temperature
gradients on day 14. Night-time skin cooling is visible as a slightly negative near-surface temperature gradient between around
22.00h and 06.00h (Fig. 5c). The fact that the model produces qualitatively realistic temperature profiles, even though its
315 parameters were merely calibrated with respect to the temperature difference between the surface and a reference depth,
supports the physical basis of the conceptual model.




Figure 5. Simulation results for the calibrated model, forced with the observational data set. a) Modeled (red) and observed (blue) time series of diurnal warming. Grey shaded intervals indicate the training data used for calibration. The subpanel below depicts wind speed (green, smoothed) for reference. b) Modeled sea temperature $\Delta T = T - T_f$ as a function of depth and time, shown for day 13, 00.00h to day 15, 00.00h (local sun time). The green line indicates wind speed. c) Modeled vertical temperature profiles on day 13 (black) and 14 (orange), plotted at intervals of two hours (shifted by 0.5 K per hour). The scale bar indicates a temperature difference of 1 K along the x-axis.

4 Comparison with other models

Our model (hereafter referred to as *DiuSST*) is intended for use as an interactive SST boundary in idealized atmospheric simulations. This calls for a comparison with existing models that could be selected for this purpose. First, we consider a slab
 320 ocean model of the type previously used in atmospheric convection studies (Hohenegger and Stevens, 2016; Shamekh et al., 2020a; Coppin and Bony, 2017; Tompkins and Semie, 2021). Second, we test the prognostic scheme by Zeng and Beljaars (2005), which has been widely applied in weather and climate modeling but, to the best of our knowledge, not been employed by the idealized convection community. Finally, we compare all models against the observations of the MOCE-5 data set and discuss their effect on air-sea heat exchange.

We use a single-layer slab with temperature dynamics described by

$$\dot{T}(t) = \frac{Q_0(t) - S}{\rho_w c_p h} - \xi_1 T(t) - \xi_2 \int_0^t T(t') dt', \quad (15)$$

where T denotes the slab temperature relative to T_f , h is the slab thickness, S represents a constant heat sink, and the net surface heat flux Q_0 is given by eq. (4). The constants ρ_w , c_p , and T_f are listed in table 1. In addition to the heat sink S , we include two correction terms that are sometimes added to control the slab temperature: a linear relaxation and an integral
330 correction to prevent temperature drift. Their strengths are tuned via the parameters ξ_1 and ξ_2 , respectively. Here $t > 0$ and $t = 0$ is the time of the initial condition.

To compare this slab model with our model, we calibrate the parameter set $\Theta_{\text{slab}} = \{h, S, \xi_1, \xi_2\}$ using Bayesian inference, taking the same data and settings as when calibrating our model (see section 3.3). In the case of the slab, diurnal warming ΔSST directly corresponds to the slab temperature anomaly T . The resulting parameter estimates are given in table E1. The
335 Bayesian MCMC sampling converges to $S = 92 \text{ W m}^{-2}$, a realistic approximation of net oceanic heat uptake in the tropics. Simulation results are discussed together with the other models (section 4.3).

4.2 ZB05 model

The ZB05 model by Zeng and Beljaars (2005) consists of two components: a bulk equation describing the temperature evolution in the diurnal warm layer of depth d , and a skin layer equation representing the cool skin effect, that is, cooling due to surface heat fluxes within the upper millimeter of the ocean (Wong and Minnett, 2018; Fairall et al., 1996). The sea skin temperature relative to the foundation temperature T_d is thus given by the sum of warm layer heating ΔT and skin cooling δT ,

$$\Delta\text{SST}(t) = \Delta T(t) + \delta T(t).$$

In order to integrate over the warm layer, the model assumes a power-law temperature profile, parameterized by an empirical shape parameter $\nu = 0.3$. The integrated equation reads (Eq. (11) in Zeng and Beljaars (2005)):

$$340 \quad \frac{d(\Delta T)}{dt} = \frac{Q_0 - R(-d)}{d\rho_w c_p \nu / (\nu + 1)} - \frac{(\nu + 1)k u_* \Delta T}{d\phi_t(d/L)}, \quad (16)$$

where $\Delta T = T_{-\delta} - T_d$ denotes the temperature difference between the skin layer depth δ and the reference depth $d = 3 \text{ m}$, Q_0 is the net surface heat flux (Eq. (4)), $R(-d)$ is the downward-penetrating shortwave radiation at depth d , and $k = 0.4$ is the von Karman constant. Furthermore, u_* represents the friction velocity which we compute in terms of the wind speed u and drag coefficient $C_D = 1.3 \times 10^{-3}$ according to $u_* = \sqrt{\rho_a C_D u^2 / \rho_w}$ (Trenberth et al., 1989; Kara et al., 2007). The stability
345 function ϕ_t is derived from Monin-Obukhov similarity theory and depends on the Monin-Obukhov length L . Here we use the refined formulation of Takaya et al. (2010) for ϕ_t .

In describing the cool skin layer we follow the scheme by Fairall et al. (1996, 2003), as implemented by the NOAA Physical Sciences Laboratory (2023). The evolution of the skin layer thickness δ and skin layer temperature difference T_d are computed

based on each other in an iterative fashion. The reason why we do not directly use Eq. (6) in Zeng and Beljaars (2005) to
350 determine δ is that this equation differs from the referenced version in Fairall et al. (1996) and leads to numerical instabilities.

4.3 Comparison of model performance

Overall, we find that both our DiuSST model and the ZB05 model give a similar, adequate performance when tested on the MOCE-5 dataset (Figs. 6, 7, 8). On the contrary, the slab model fails to describe the observed diurnal SST variability, even though it has been optimized on the training set of this data.

355 Specifically, the slab model is unable to capture the wind-dependent variation in diurnal warming amplitudes (Fig. 6a). The slab simulation exhibits a rather invariant diurnal amplitude of just above $\Delta\text{SST} = 1\text{ K}$, largely independent of wind speed (Fig. 7b). The poor agreement between the slab and the observations manifests itself in the relatively low Pearson correlation coefficient of 0.50 and a standard mean error of 0.36 K (over the whole data set). In comparison, the DiuSST and ZB05 models yield a Pearson correlation of 0.74 and 0.72, respectively, each deviating from the observations by around 0.29 K on average.
360 Even more than DiuSST, the ZB05 model overpredicts diurnal warming on days 6 and 7 (Fig. B1). ZB05 exhibits more high-frequency variability during diurnal warming events compared to DiuSST, which we attribute to the more sensitive skin layer model explicitly included in ZB05. Night-time skin cooling is overall stronger in ZB05 relative to DiuSST, agreeing better with observations during the second half of the time series (days 9-16) but exaggerating skin cooling during the first, windier half (days 1-7, see Fig. B1). While capturing some of the observed night-time dynamics of ΔSST , the Slab tends to produce excess
365 skin cooling especially during strong winds.

Fig. 7b) highlights the improved wind dependence of ΔSST in DiuSST compared to the Slab model. Similar to observations and ZB05, our model exhibits a roughly exponential decay of peak diurnal warming with increasing wind speed. To provide a more quantitative analysis, we bin each diurnal warming time series by wind speed, selecting a bin size of 0.5 ms^{-1} . For each bin, we calculate hourly averages of ΔSST as a function of local sun time and take the maximum of these hourly averages as
370 an estimate of the mean diurnal warming amplitude for the given wind speed (Fig. 8). Based on exponential fits, ΔSST as a function of wind speed decays with a scaling constant of roughly 2 ms^{-1} for the observations, DiuSST and ZB05 models (Tab. 2). In the case of the Slab, the exponent is almost 30 ms^{-1} , confirming the weak wind sensitivity of the diurnal amplitude. The expected diurnal warming amplitude under calm conditions, approximated by the intercept of the fit at $u = 0$, is underestimated (overestimated) by DiuSST (ZB05) by about 0.5 K each. The Slab diurnal warming amplitude is significantly too low.

375 4.4 Air-sea heat fluxes

The diurnal evolution of SST impacts the atmosphere by regulating the heat, moisture and momentum transfer at the air-sea interface. Over the course of the observational data interval (October 1-16), we compute the net surface heat flux $Q_{\text{net}}(t) = R_{\text{lw}}(t) + Q_1(t) + Q_s(t)$ (excluding shortwave radiation) for each model simulation (Fig. 6b, $Q_{\text{net}} < 0$ corresponds to net heat export from ocean to atmosphere). In addition, we compute $Q_{\text{net}}(t)$ for the given atmospheric forcing under the assumption
380 that SST is fixed at $T(0, t) = T_f$ for all t .




Figure 6. Comparison of the calibrated DiuSST, ZB05, and Slab models. a) Time series of diurnal warming ΔSST as modeled by DiuSST (red), ZB05 (dark blue) and the slab (orange) for a select time interval. Points (light blue) indicate the observations. b) Net surface heat loss to the atmosphere, $Q_{net} = R_{lw} + Q_1 + Q_s$ (excluding shortwave radiation), for the model (red) and slab (orange). c) Difference in Q_{net} between our model and the slab (black) as well as our model and a fixed SST at T_f (gray). d) Correlations between the modeled and observed ΔSST for our model (top) as well as ZB05 and the slab (bottom). The Pearson correlation coefficients (Corr.) are given. In the top panel, circles (crosses) mark training (validation) data points.

Data	intercept y_0 (K)	exponent a (ms^{-1})
Observations	4.59	1.98
DiuSST	4.12	2.08
ZB05	5.04	1.74
Slab	1.04	29.63

$$\text{Fit function: } \Delta SST(u) = y_0 \exp(-u/a)$$

Table 2. Exponential fits of the wind dependence of diurnal warming (see Fig. 8).




Figure 7. Diurnal warming in the observations and simulations with the DiuSST, ZB05, and Slab models (from left to right), showing all data points of the 16-day time series. a) ΔSST as a function of local sun time, colored by wind speed. b) ΔSST as a function of wind speed, colored by local sun time.

The oceanic heat loss to the atmosphere can differ by up to around 50 W m^{-2} between DiuSST and the Slab (Fig. 6c). Q_{net} is more negative for DiuSST when it produces a larger sea skin temperature compared to the Slab, and vice versa. During strong diurnal warming events, the difference ΔQ_{net} is even larger when comparing DiuSST to a fixed SST, exceeding 50 W m^{-2} on Oct 13. We conclude that using a fixed SST as an oceanic boundary condition leads to biases in surface heat fluxes particularly during strong diurnal warming events, whereas using a slab model causes biases under strong insolation both in windy and calm conditions. Between DiuSST and ZB05, differences in the net surface heat flux are relatively small and mainly due to differences in the magnitude of skin cooling.

5 Discussion

This study aims at contributing a diurnal warm layer model that offers a simple improvement over slab ocean models previously used in idealized atmosphere-ocean studies. Upper ocean heat transfer involves complex processes from wave breaking and Langmuir circulations to biological productivity (Edson et al., 2007; Noh et al., 2004). To first order, however, the diurnal variability of SST is governed by the competing effects of solar absorption and wind-driven turbulent mixing. This permits our reductionist modeling approach, which approximates the temperature dynamics in the diurnal warm layer mainly as a function




Figure 8. Wind dependence of the amplitude of diurnal warming, comparing the observations against the predictions by the DiuSST, ZB05, and Slab models (see figure legend). The data points represent the mean diurnal warming amplitude (see main text). Solid lines indicate least-squares exponential fits, with parameters given in tab. 2. Wind speeds above 6.5 m s^{-1} are not shown due to a small number of raw data points.

of wind speed and insolation. For detailed realism of diurnal warm layer dynamics, more comprehensive models (as listed in
 395 the introduction) will be appropriate.

5.1 Model limitations and extensions

Due to its one-dimensional setup, DiuSST neglects horizontal flows and heat exchange. Furthermore, turbulence is dissipated
 instantaneously, i.e. there is no memory of momentum due to the wind stress history. These restrictions could explain the poor
 performance of DiuSST (and ZB05, likewise) on days 6 and 7, when the amplitude of diurnal warming is overestimated by
 400 more than 1 K (Fig. 5a). Despite strong insolation and low winds before noon on these days, observed diurnal warming does
 not exceed 1 K. This suggests the presence of enhanced turbulent vertical mixing not captured by the models, possibly due
 to non-local effects such as horizontal currents or rough seas, either advected from a windy region or remnant from a windy
 episode preceding the observations.

For simplicity, DiuSST represents stratification in the diurnal warm layer only in a time-averaged sense via the diffusivity
 405 profile $\varphi(z)$ which suppresses turbulent diffusion near the surface. In reality, upper-ocean water column stability depends on the
 interplay between time- and depth-dependent density gradients and turbulence dissipation (Hughes et al., 2020). Instead of the
 highly simplified linear profile given in eq. (3), a nonlinear, dynamic diffusivity profile could incorporate the nature of upper
 ocean turbulence more realistically. For instance, a state-dependent profile, $\varphi = \varphi(z, T(z, t))$, could reflect the temperature
 dependence of stratification. We discuss alternative diffusivity profiles in D.

410 The mixing term controlled by μ ensures that the diurnal layer temperature relaxes back to the foundation temperature, resetting the temperature profile at night. Combined with the diffusion term, it can only serve as a crude account of the complex mixing processes in the upper ocean (Hughes et al., 2020), such as convective overturning or internal waves. However, the contribution from this term is typically small compared to the other terms, particularly near the surface. In some cases, it may be realistic to further decrease μ to allow for multi-day warming events (Jia et al., 2023).

415 With a vertical resolution set to 10 cm at the surface, our numerical implementation is too coarse to explicitly resolve the cool skin layer, which forms within millimeters from the air-sea interface. Nonetheless, the model still captures skin cooling at night, indicated by slightly negative values of ΔSST , in agreement with the observations (Fig. 5). In fact, the model considers the cool skin effect in a coarse-grained sense, averaging heat fluxes over the upper 10 cm of the water column. Of course, the grid resolution could be increased, albeit at the cost of a smaller integration time step to maintain numerical stability.

420 For future refinement, further dynamical processes could be built into DiuSST via the three key modular terms. A time-dependent diffusivity profile would allow to parameterize the effect of precipitation, which we presently neglect in the model. Rain freshwater pools can either enhance heat trapping or mixing, depending on the competing effects of salinity and temperature (Webster et al., 1996; Bellenger et al., 2017; Witte et al., 2023). As another example, a dynamic attenuation coefficient $\alpha(t)$ would allow to account for changes in the seawater optical properties, e.g. due to microbial activity (Wurl et al., 2017).

425 5.2 Data-driven parameter estimation

While the parameters $\{\kappa_0, \alpha, \mu\}$ are fixed in our model, they generally depend on oceanic conditions that may vary in space and time. A major benefit of Bayesian inference is that the model parameters can readily be re-calibrated to additional data of interest.

The MOCE-5 cruise data used here cover measurements across approximately ten degrees latitude and longitude, both in
430 the open Pacific Ocean and in coastal waters of the Gulf of California, where dynamical and optical properties of sea water are likely to have differed. The observed foundation temperature varied by several Kelvin during the cruise, whereas in the model we fix T_f to the observed mean. Thus, spatial heterogeneity of water properties probably constitutes a main error source between model and observations. For example, on day 13, the modeled maximum of diurnal warming is about 1 K below the observed value of $\Delta\text{SST} \approx 4.9\text{K}$. This difference may be attributed to spatial variations of optical water properties affecting
435 the attenuation coefficient α . The cruise vessel’s location on day 13 near the Midriff Islands is known for high phytoplankton concentrations, which enhances the absorption of shortwave radiation (Álvarez Borrego, 2012). Indeed, we can accurately model the maximum of ΔSST on day 13 by increasing α , but this reduces the model performance across the time series overall.

At the same time, the heterogeneity of the training data presents a benefit, as it reduces the risk of overfitting and promises
440 a more generally applicable calibration. Even with constant parameters obtained from Bayesian inference, our model captures the observed variability of ΔSST despite the heterogeneity of the MOCE-5 dataset. For idealized atmospheric modeling, we argue that one is usually interested in average sea properties for conditions of interest, or a parameter sensitivity experiment. While the present calibration could be specific to the background conditions of the study region, the model formulation itself

is not. The parameter posterior distribution reported here can be used directly as a prior distribution for a re-calibration based
445 on relevant additional data.

To validate the vertical profiles generated by DiuSST, depth-resolved observations are needed. Incorporating information at
depth in the calibration can further constrain the parameter estimates, help resolve apparent parameter correlations, and inform
the choice of diffusivity profile φ . However, the vertical temperature profiles measured during the MOCE-5 cruise (Ward,
2006) are subject to various other influences besides diurnal warming (Gentemann et al., 2009), and currently not publicly
450 available. Validating the profiles is thus left for future work.

5.3 Inaptitude of slab ocean

In contrast to the DiuSST and ZB05 models, the Slab model fails to capture the observed diurnal warming dynamics, including
the wind dependence of ΔSST . This is because the evolution of slab temperature depends on wind speed only via the latent
and sensible surface heat flux, whereas wind strongly influences vertical heat transport via the diffusion term in DiuSST or
455 via the stability function in ZB05. In other words, the wind dependence of upper-ocean turbulence controls the effective heat
capacity of the upper ocean, but the slab's heat capacity is fixed by the slab thickness h . This issue also pertains to ocean
models with a surface layer of a meter thickness or more, which may reproduce a mean diurnal cycle but not the full range of
diurnal variability (Voltaire et al., 2022).

One might argue that the slab exhibits too little variability because the parameter ξ_1 , the inverse timescale of temperature
460 relaxation, is set too large. Indeed, decreasing ξ_1 allows for stronger diurnal warming but also leads to excessive nighttime
cooling and a slow temperature decline in the afternoon, thus worsening the agreement with the observations overall (see Fig.
F1). In fact, the shape of the diurnal warming curve produced by a slab with small corrector coefficients ξ_1 and ξ_2 resembles
the diurnal temperature evolution observed at around 1 m depth, e.g., by moored buoys (Börner, 2021). This highlights that
slab oceans mimic bulk SST rather than skin SST. Also other models of the diurnal SST cycle are sometimes calibrated with
465 respect to bulk SST. However, skin SST is the relevant quantity for atmospheric studies, since the atmosphere only senses the
temperature of the sea skin.

5.4 Interactive diurnal SST in atmospheric simulations

As a more realistic alternative to a slab model, the DiuSST and ZB05 models can be coupled to high-resolution cloud-resolving
atmospheric models, e.g. to study how air-sea interactions impact atmospheric convection in idealized setups.

470 The ZB05 model appeals with its rigorous derivation from Monin-Obukhov similarity theory, condensed into one integrated
bulk equation for the warm layer plus the cool skin layer scheme. This makes it computationally fast, though calculating
the skin layer depth δ involves an implicit equation that requires iterative solving (see Eqs. (5) and (6) in Zeng and Beljaars
(2005)). The empirical shape parameter ν of the vertical temperature profile could serve as a tuning parameter in sensitivity
experiments. While ZB05 matches fairly well with the observations used in this study, its performance has varied in other
475 comparative studies (Bellenger and Duvel, 2009; Jia et al., 2023). Running DiuSST requires integrating a discretized PDE,
making it slightly slower than ZB05 in the Python scripts implemented here (Börner, 2024) but not significantly – especially

relative to the cost of a coupled atmospheric component. Compared to ZB05 or similar existing diurnal warm layer schemes, modelers may find DiuSST easier to tune, interpret and adapt to their purposes due to the conceptually simple modular structure. The fact that DiuSST resolves the vertical temperature profile opens up opportunities to study, for example, how biological upper-ocean processes interact with weather and climate across the air-sea interface.

480 Recently, we implemented our upper ocean model as an interactive sea surface boundary condition in the System for Atmospheric Modeling (SAM) model (Khairoutdinov and Randall, 2003). At each horizontal grid point of the atmospheric model, the surface boundary condition is independently updated at each time step by numerically integrating eq. (A1) to the next time step, based on the local atmospheric forcing \mathcal{F} . This produces a responsive, spatially heterogeneous sea surface whose temperature feeds back into the atmospheric boundary layer. According to first tests, the difference in computation time between this
485 setup and a coupled slab ocean is unnoticeable.

The influence of our model on the spatio-temporal patterns of convection is subject of future research. As a global kilometer-scale simulation study by Shevchenko et al. (2023) suggests, diurnal warm layers can have a significant impact on the atmosphere locally where diurnal warming is strong; yet their role for the large-scale tropical cloud field and convective organization
490 remains unclear.

6 Conclusions

This paper presents a simple, one-dimensional prognostic model of diurnal sea surface temperature variability in the tropical ocean, described by Eq. (1) and formulated as a numerical scheme in Eq. (A1). Written as a single partial differential equation, the model describes upper ocean heat transfer through three idealized terms, controlled by three tuning parameters: an eddy
495 diffusivity κ_0 , a bulk mixing rate μ and an attenuation coefficient α . κ_0 controls the strength of wind-driven turbulent heat transport, μ determines the relaxation rate towards the foundation temperature and α specifies how deeply solar radiation penetrates into the ocean.

First, we used Bayesian inference to estimate the values of these parameters based on an observational data set recorded on the MOCE-5 cruise in the Eastern Pacific. Then, we compared the performance of our model with two other models of diurnal
500 SST dynamics: a slab ocean model, as previously used to mimic a responsive sea surface in atmospheric simulations, and the diurnal warm layer scheme ZB05 (Zeng and Beljaars, 2005).

Our results show that slab models with fixed heat capacity cannot capture diurnal SST variability realistically. Instead, our model reproduces an exponential dependence of the diurnal warming amplitude on wind speed, in accordance with observations and similar to ZB05. By introducing a diffusion term that scales with surface wind stress, we propose a simple solution that
505 offers significantly improved results compared to a slab model and parameterizes the basic features of upper ocean turbulence in a physically interpretable way. Enhancing the model in the future, e.g. by refining the diffusivity profile to include effects of precipitation, is facilitated by the concise modular code. To confirm the model's validity for diverse conditions, the Bayesian approach offers a natural way to update the parameter estimates based on additional data.

Due to its numerical and conceptual simplicity, we envision the model presented in this study to serve as a generic interactive
 510 boundary condition for oceans in idealized cloud-resolving simulations of the atmosphere. This will enable a wind-responsive
 SST field that evolves under and feeds back into the atmospheric fluid dynamics. Given the multiscale interaction between
 the diurnal cycle and large-scale patterns of tropical convection, it becomes increasingly clear that we must consider diurnal
 warm layer dynamics to better understand the mechanisms of marine cloud organization. Our work thus hopes to contribute to
 bridging the gap between idealized studies of convective aggregation and real-world process understanding.

515 *Code and data availability.* A documented Python implementation of the models (DiuSST, ZB05 and Slab), the MOCE-5 observational
 dataset, as well as a tutorial for running the code are available under the DOI 10.5281/zenodo.13363481 at [https://github.com/
 reykboerner/diusst](https://github.com/reykboerner/diusst) (Börner, 2024).

Video supplement. A video presentation introducing the model is available at https://www.youtube.com/watch?v=KdOWF_fzRLE.

Appendix A: DiuSST model – Discretized model equation

520 The discretized form of eq. (1), using an explicit Euler scheme in time and finite differences on a non-uniform grid in space,
 reads

$$\begin{aligned}
 \frac{T_n^{i+1} - T_n^i}{\Delta t_i} = & \kappa(z_n, t_i) \left[(T_{n+1}^i - 2T_n^i + T_{n-1}^i) \cdot \left(\frac{dn}{dz} \Big|_{z_n} \right)^2 + \frac{T_{n+1}^i - T_{n-1}^i}{2} \cdot \frac{d^2n}{dz^2} \Big|_{z_n} \right] \\
 & + \frac{\partial \kappa(z, t_i)}{\partial z} \Big|_{z_n} \left[\frac{T_{n+1}^i - T_{n-1}^i}{2} \cdot \frac{dn}{dz} \Big|_{z_n} \right] - \mu \frac{T_n^i - T_f}{|z_n - z_f|} \\
 & + \frac{1}{\rho_w c_p} \left[(Q_{n+1}^i - Q_n^i) \cdot \frac{dn}{dz} \Big|_{z_n} \right], \quad (\text{for } n = 0, \dots, N-1), \tag{A1}
 \end{aligned}$$

where n is the depth index and i the time index (nomenclature as in the main text, see also table 1). Here $\kappa(z, t)$ is given by eq.
 (2); its derivative by z is computed analytically. The derivatives dn/dz and d^2n/dz^2 ,

$$\frac{dn}{dz} = \left[\ln \epsilon \cdot \left(\frac{\Delta z_0}{1 - \epsilon} + z \right) \right]^{-1}; \quad \frac{d^2n}{dz^2} = \left[-\ln \epsilon \cdot \left(\frac{\Delta z_0}{1 - \epsilon} + z \right)^2 \right]^{-1}, \tag{A2}$$

map the finite differences onto the non-uniform grid spacing, where the stretch factor $\epsilon > 1$ solves the equation $\epsilon(\Delta z_0, N) =$
 525 $1 + (1 - \epsilon^N) \Delta z_0 / z_f$ (see also eq. (9)). At the foundation depth, the Dirichlet boundary condition is $T_N^i = T_f$ for all i . At
 the surface ($n = 0$), we require a choice on dealing with the free boundary. We introduce a dummy grid point ($n = -1$) at
 $z_{-1} = \Delta z_0$, at which we set the temperature to $T_{-1}^i = T_0^i$ for all i (essentially assuming that the temperature gradient vanishes




Figure B1. Comparison of observed diurnal warming ΔSST (light blue) against the predictions of the calibrated DiuSST (red dashed), ZB05 (dark blue), and calibrated Slab (orange dash-dotted) models, showing the full 16-day time series recorded on the MOCE-5 cruise.

at the air-sea interface). The surface heat flux (at $n = 0$) is given by

$$Q_0^i := R_{\text{sw}}(t_i) + R_{\text{lw}}(t_i) + Q_s(t_i) + Q_1(t_i). \quad (\text{A3})$$

For $n > 0$, the heat flux $Q_n^i \equiv Q(z_n, t_i)$ is given by eq. (7). Using a forward difference for Q_n^i ensures that the integrated heat flux over the domain corresponds to the total heat uptake within the diurnal layer. Note that the diffusion part of eq. (1) involves two terms in eq. (A1) owing to the chain rule of differentiation.

Appendix B: Model comparison over full MOCE5 time series

Simulation results over the whole time interval of the MOCE-5 observations are shown in Fig. B1 for all models. DiuSST and ZB05 produce similar results, with ZB05 occasionally overshooting the observations by more than 1 K around diurnal warming peaks, e.g. on days 6, 10 and 14. DiuSST produces a more accurate nighttime skin temperature throughout the windy first half of the timeseries (days 2-8), whereas ZB05 yields better nighttime skin temperatures during the second half of the timeseries. The slab produces a constant diurnal warming amplitude throughout, overestimating ΔSST on windy days and underestimating it on calm days.




Figure C1. Difference in simulated Δ SST of the DiuSST model without reflection and refraction vs. the original DiuSST model, for the parameters given in Tab. 1 and forcing with MOCE-5 data.

Appendix C: Surface reflection of solar radiation in DiuSST

540 Based on Fresnel's equations, assuming unpolarized light, the reflected fraction \mathcal{R} of irradiance incident on the (flat and smooth) air-sea interface is given by $\mathcal{R} = (\mathcal{R}_\perp + \mathcal{R}_\parallel)/2$, where the contributions from the two polarization directions read

$$\mathcal{R}_\perp = \left(\frac{n_a \cos \phi - n_w \sqrt{1 - \left(\frac{n_a}{n_w} \sin \phi\right)^2}}{n_a \cos \phi + n_w \sqrt{1 - \left(\frac{n_a}{n_w} \sin \phi\right)^2}} \right)^2 \quad (\text{C1})$$

$$\mathcal{R}_\parallel = \left(\frac{n_a \sqrt{1 - \left(\frac{n_a}{n_w} \sin \phi\right)^2} - n_w \cos \phi}{n_a \sqrt{1 - \left(\frac{n_a}{n_w} \sin \phi\right)^2} + n_w \cos \phi} \right)^2. \quad (\text{C2})$$

Here ϕ is the solar angle with respect to the surface normal; n_a and n_w denote the refractive index of air and water, respectively (see table 1). Thus, we approximate the transmitted solar irradiance R_{sw} entering the water body as

$$R_{\text{sw}}(t) = (1 - \mathcal{R}(\phi(t))) R_{\text{sw},\downarrow}(t), \quad (\text{C3})$$

where $R_{\text{sw},\downarrow}$ is the downward shortwave irradiance above the sea surface.

545 When neglecting surface reflection and the effect of surface refraction on the absorption of downward shortwave radiation, the simulated Δ SST (under the same parameter settings) can be larger by up to 0.2 K for the MOCE-5 case study (Fig. C1). However, note that if we would have calibrated the model without reflection and refraction, this difference would likely be compensated by adjusting the parameters.

Appendix D: Alternative diffusivity profiles

550 In the main text, we introduce a highly idealized linear diffusivity profile $\varphi(z)$ (eq. (3), called LIN hereafter). Here we briefly discuss two alternatives that have been investigated in the course of this study. Consider the profile

$$\varphi(z, t) = \frac{1 - S(t)\sigma \exp(z/\lambda)}{1 - S(t)\sigma \exp(z_f/\lambda)}, \quad (\text{D1})$$

given in terms of the suppressivity $\sigma \in [0, 1]$, the trapping depth $\lambda > 0$, and the (time-dependent) stability function S .

First, set $S(t) = 1$ for all t , giving a time-independent exponential profile (EXP). If $\sigma > 0$, then φ decreases exponentially towards the sea surface, where $\varphi(0) = 1 - \sigma$. The parameter λ , setting the curvature of the profile, determines the depth scale
555 at which the diffusivity is suppressed. In the limit $\lambda \rightarrow \infty$, we obtain the linear profile in eq. (3).

Second, we investigate a state-dependent stability function (STAB) that depends on the vertical temperature gradient $\Delta T_\lambda(t) := T(0, t) - T(\lambda, t)$, given by

$$S(t) = \begin{cases} \min(1, \Delta T_\lambda(t)/T_{\text{strat}}) & \text{if } \Delta T_\lambda(t) > 0 \\ 0 & \text{otherwise,} \end{cases} \quad (\text{D2})$$

where T_{strat} is a reference temperature at which thermal stratification dominates over turbulent mixing. The physical rationale behind this is that density stratification due to near-surface heat trapping locally enhances water column stability, inhibiting
560 vertical diffusion. As soon as the sea skin temperature cools again with respect to the temperature at depth λ , diffusivity increases, mimicking enhanced afternoon mixing due to an unstable temperature profile.

Comparing simulations with the LIN and EXP profiles, we find that EXP performs slightly better than LIN for the observational data set considered in this study, particularly with regards to night-time skin cooling (Fig. D1). Though arguably more realistic than LIN and EXP, the STAB profile does not necessarily perform better because it kills the coarse-grained skin
565 cooling effect in our model. Comparing the effect of EXP and STAB on the vertical temperature profile (Fig. D2), we see that STAB counteracts near-surface temperature inversions, creating a mixed warm layer of near-constant temperature that better matches observations Soloviev and Lukas (1997). However, STAB would have to be combined with a skin layer scheme as in ZB05 in order to capture the sea skin temperature in the presence of skin cooling. Due to the overall similarity of modeled SST warming between the LIN, EXP, and STAB versions, we choose to present the simplest option, LIN, in the main text.

570 Appendix E: Bayesian inference

Consider a model $\mathcal{M}(\Theta)$ controlled by the parameter set Θ . Given the data \mathcal{D} , the probability that a certain value of Θ represents the truth follows from Bayes' rule,

$$P(\Theta|\mathcal{D}) = \frac{P(\mathcal{D}|\Theta)P(\Theta)}{P(\mathcal{D})}. \quad (\text{E1})$$

Here $P(A|B)$ denotes the conditional probability of A given B . On the right-hand side, $P(\mathcal{D}|\Theta)$ is the *likelihood* of observing the data under the assumption that Θ represents the true model. The *prior* probability $P(\Theta)$ quantifies our knowledge of the




Figure D1. Comparison of modeled diurnal warming ΔSST between different diffusivity profiles φ discussed in D and the main text. LIN (solid red line) corresponds to the linear profile (eq. (3)) used throughout the main text. EXP (dash-dotted blue) and STAB (dashed orange) show results for the corresponding versions of φ described in D. For each case, the model parameters κ_0 , μ , and α are calibrated specifically via Bayesian inference. All other model settings are kept constant, as given in table 1.

575 parameter set before observing the data. Lastly, the denominator states the probability of the data being true, which is independent of the choice of model and merely adds a normalization factor. Hence the *posterior* probability $P(\Theta|\mathcal{D})$ is proportional to the product of the likelihood times the prior probability.

In practice, inferring the posterior distribution $P(\Theta|\mathcal{D})$ of the parameters Θ from the data \mathcal{D} involves three steps: 1) construct the prior distribution $P(\Theta)$ in parameter space based on previous knowledge or belief; 2) define a suitable likelihood function
 580 $\mathcal{L}(\Theta) \equiv P(\mathcal{D}|\Theta)$; and 3) compute the (un-normalized) posterior distribution by evaluating the product $\mathcal{L}(\Theta)P(\Theta)$.

E1 Choice of prior

Our sensitivity study under idealized forcing (section 3.1) provides orientation on physical ranges of the parameters $\{\kappa_0, \mu, \alpha\}$ of the DiuSST model. In the literature, common values of oceanic turbulent vertical diffusivity are on the order of $\kappa \sim 10^{-4} \text{ m}^2 \text{ s}^{-1}$, varying with location and depth Denman and Gargett (1983). Stating a physical value for the attenuation coefficient α is difficult since, in reality, attenuation of shortwave radiation depends strongly on wavelength and the biochemical
 585 composition of the seawater. Empirical values for the diffusive attenuation coefficient of photosynthetically active radiation range from $\sim 10^{-2} \text{ m}^{-1}$ to 10 m^{-1} Son and Wang (2015). However, we note that the model parameters are conceptual, particularly the mixing coefficient μ , such that they do not necessarily represent directly observable physical quantities.




Figure D2. Comparison of vertical temperature profiles, produced by the model, with diffusivity profile EXP (dark blue) and STAB (orange). The top panel shows the vertical profile every two hours, shifted by 1 K per hour along the horizontal axis. The bottom panels show the temperature difference ΔT with respect to the foundation temperature as a function of depth and time for the EXP model (left) and STAB model (right).

Reflecting our limited knowledge, we impose uniform prior distributions for the parameters κ_0 and α but constrain their range to $\kappa \in [0, 5 \times 10^{-4}] \text{ m}^2 \text{ s}^{-1}$ and $\alpha \in [0.05, 10] \text{ m}^{-1}$. The parameter μ requires more subtle treatment because it acts mainly near the foundation depth, whereas the likelihood function evaluates temperature differences near the surface (see below). Fig. 2 indicates that excess heat remains in the interior of the diurnal layer if μ falls below $\sim 1 \times 10^{-4} \text{ m s}^{-1}$. Based on this insight, we define a normally distributed prior for μ with mean $6 \times 10^{-3} \text{ m s}^{-1}$ and standard deviation $1.5 \times 10^{-3} \text{ m s}^{-1}$. Thus the three-dimensional prior distribution $P(\Theta)$ is uniform and bounded along the κ_0 and α directions but Gaussian with respect to μ .

Note that the posterior distribution with respect to μ differs clearly from the prior distribution, confirming that the data added information on the parameter μ (Fig. 4).

E2 Defining the likelihood function

Our *training data* \mathcal{D} consists of a six-day subset of the diurnal warming time series from the MOCE-5 cruise, as indicated by the gray shading in Fig. 3. Specifically, we select days 2-4 and 13-15 of October 1999 as training data, while all other days from 1 to 16 October make up the *validation data*. By this choice, the training data contains warming events of different amplitudes, covering the observed range from a few tenths of a Kelvin up to 5 K. Moreover, the two sub-intervals composing \mathcal{D} correspond to different geographical regions (open Pacific vs. Gulf of California), presumably featuring differing environmental conditions. This ensures that the resulting parameter estimates will be valid for a broader range of conditions.

We choose a likelihood function that decays exponentially with the weighted square error between model and data,

$$\mathcal{L} \propto \exp \left(- \sum_j \frac{(\Delta\text{SST}_{\mathcal{M}}(t_j) - \Delta\text{SST}_{\mathcal{D}}(t_j))^2}{\Sigma_j^2} \right), \quad (\text{E2})$$

where the index j runs through all data points in \mathcal{D} and $\Delta\text{SST}_{\mathcal{M}}$ ($\Delta\text{SST}_{\mathcal{D}}$) denotes the modeled (observed) diurnal warming at the corresponding time of observation. Specifically, $\Delta\text{SST}_{\mathcal{M}}(t_j) = T(0, t_j) - T(z_d, t_j)$, with z_d being the depth of the vertical grid point closest to the observational reference depth $d = -3$ m. Furthermore, the standard error Σ_j determining the relative weight of the j -th data point is defined as

$$\Sigma_j = 2\epsilon_j(1 + v_j/v_{\max}), \quad (\text{E3})$$

where ϵ_j denotes the uncertainty of the 3 m temperature measurement, as stated in the data set; v_j is the current boat speed and v_{\max} is the maximum boat speed throughout the data set. This choice reflects a reduced confidence in measurements taken while the ship was moving rapidly. Indeed, the MOCE-5 time series features several short, isolated warming spikes of around 1-2°C during nighttime hours, which cannot be explained by atmospheric forcing but could have been caused by the boat's movement. Since the ship mainly moved at night while often remaining at fixed locations during the day, Σ_j emphasizes daylight data.

E3 Computing the posterior distribution

To approximate the posterior distribution, we perform Markov chain Monte Carlo (MCMC) sampling using the `emcee` package in Python Foreman-Mackey et al. (2013). It is based on the affine-invariant GW10 algorithm proposed by Goodman and Weare 2010. The algorithm explores the parameter space with multiple interdependent walkers whose moves efficiently deal with correlations, yielding fast convergence.

At each step of the Markov chain, the likelihood function, Eq. (E2), is evaluated for each walker at its respective position Θ in parameter space. Each evaluation requires simulating the model for the duration of the training data, which makes the sampling computationally expensive. For each of 24 walkers we generate 4000 steps, initialized at the value $\{\kappa_0, \mu, \alpha\} = \{10^{-4} \text{ m}^2 \text{ s}^{-1}, 6 \times 10^{-3} \text{ m s}^{-1}, 4 \text{ m}^{-1}\}$. Finding an auto-correlation length of around 50 steps (along the Markov chain), this gives around 80 independent samples.

We summarize the resulting parameter estimates in Tab. E1, listing common statistical estimators as obtained from the posterior distribution of the DiuSST (Fig. 4) and Slab (Fig. E1) models, respectively. Bayesian inference for the Slab model was per-

formed in analogy to the DiuSST model but in a four-dimensional parameter space $\{h, S, \xi_1, \xi_2\}$ and with the slab temperature anomaly representing $\Delta\text{SST}_{\mathcal{M}}$ in the likelihood function. The run was initialized at values $\{h, S, \xi_1, \xi_2\} = \{1, 100, 10^{-4}, 10^{-9}\}$ with uniform priors for h, ξ_1, ξ_2 and a Gaussian prior for S with mean 100 W m^{-2} and standard deviation 10 W m^{-2} .

Model	Parameter	MAP	Mean	Median
DiuSST	$\kappa_0 \text{ (m}^2\text{s}^{-1}\text{)}$	1.34×10^{-4}	1.41×10^{-4}	1.40×10^{-4}
	$\mu \text{ (ms}^{-1}\text{)}$	2.85×10^{-3}	3.00×10^{-3}	2.96×10^{-3}
	$\alpha \text{ (m}^{-1}\text{)}$	3.52	3.83	3.70
Slab	$h \text{ (m)}$	1.20	1.20	1.20
	$S \text{ (W m}^{-2}\text{)}$	92.67	93.86	93.94
	ξ_1	1.19×10^{-4}	1.20×10^{-4}	1.18×10^{-4}
	ξ_2	3.1×10^{-11}	5.3×10^{-11}	3.7×10^{-11}

Table E1. Parameter estimates obtained from the sampled posterior distribution.

630 Appendix F: Slab model parameter sensitivity

As mentioned in the main text, the Bayesian optimization procedure applied to the Slab model acts mainly by ramping up the relaxation parameter ξ_1 , which constrains diurnal warming to 1 K. Larger diurnal amplitudes are achieved by reducing ξ_1 (see Fig. F1). However, in that case the slab also cools excessively at night, and the diurnal warming peak shifts into the late afternoon, with a slow temperature decrease after peak warming.

635 *Author contributions.* JOH and RF conceived and supervised the study. All authors developed the model and methodology. RB produced the code, acquired and curated the data, conducted the investigation and formal analysis, performed the model evaluation and validation, and prepared the figures. RB prepared the manuscript with input from all co-authors.

Competing interests. The authors declare no competing interests.

640 *Acknowledgements.* We thank Gorm Gruner Jensen, Peter Ditlevsen, and Chong Jia for useful discussions, as well as Peter Minnett for providing the MOCE-5 dataset and helpful feedback on the manuscript. We further thank the two anonymous reviewers for their constructive feedback which helped strengthen the manuscript. The development and deployment of the instruments used during the MOCE-5 cruise was funded by NASA. R. F. and J. O. H. gratefully acknowledge funding from the Villum Foundation (grant no. 13168). J. O. H. acknowledges funding from the European Research Council under the European Union’s Horizon 2020 Research and Innovation programme (grant no.




Figure E1. Posterior distribution for the parameters of the slab model, obtained from Bayesian MCMC sampling. Compare with Fig. 4, which shows the results for the DiuSST model.




Figure F1. Performance of the slab model as a function of its control parameter ξ_1 . The solid red (Model) and blue (Slab A) lines show simulations of our calibrated model and the slab model, respectively, with parameter values given in table E1. The orange curve (Slab B) shows a slab simulation with $h = 1.1$ m, $S = 70$ W m⁻², $\xi_1 = 1 \times 10^{-7}$ and $\xi_2 = 0$. Note that for Slab B, ξ_1 is three orders of magnitude smaller compared to Slab A.

771859) and the Novo Nordisk Foundation Interdisciplinary Synergy Program (grant no. NNF19OC0057374). R.B. acknowledges funding
 645 from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 956170 (CriticalEarth).

References

- Alduchov, O. A. and Eskridge, R. E.: Improved Magnus form approximation of saturation vapor pressure, *Journal of Applied Meteorology and Climatology*, 35, 601–609, 1996.
- 650 Bellenger, H. and Duvel, J.-P.: An Analysis of Tropical Ocean Diurnal Warm Layers, *Journal of Climate*, 22, 3629–3646, <https://doi.org/10.1175/2008JCLI2598.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2009.
- Bellenger, H., Takayabu, Y. N., Ushiyama, T., and Yoneyama, K.: Role of Diurnal Warm Layers in the Diurnal Cycle of Convection over the Tropical Indian Ocean during MISMO, *Monthly Weather Review*, 138, 2426–2433, <https://doi.org/10.1175/2010MWR3249.1>, publisher: American Meteorological Society Section: *Monthly Weather Review*, 2010.
- 655 Bellenger, H., Drushka, K., Asher, W., Reverdin, G., Katsumata, M., and Watanabe, M.: Extension of the prognostic model of sea surface temperature to rain-induced cool and fresh lenses, *Journal of Geophysical Research: Oceans*, 122, 484–507, <https://doi.org/10.1002/2016JC012429>, 2017.
- Bernie, D. J., Guilyardi, E., Madec, G., Slingo, J. M., Woolnough, S. J., and Cole, J.: Impact of resolving the diurnal cycle in an ocean–atmosphere GCM. Part 2: A diurnally coupled CGCM, *Climate dynamics*, 31, 909–925, 2008.
- 660 Böing, S.: An object-based model for convective cold pool dynamics, *Mathematics of Climate and Weather Forecasting*, 2, 2016.
- Bony, S., Stevens, B., Frierson, D. M. W., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., Watanabe, M., and Webb, M. J.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261–268, <https://doi.org/10.1038/ngeo2398>, number: 4 Publisher: Nature Publishing Group, 2015.
- Bretherton, C. S., Blossey, P. N., and Khairoutdinov, M.: An energy-balance analysis of deep convective self-aggregation above uniform SST, *Journal of the atmospheric sciences*, 62, 4273–4292, 2005.
- 665 Brunke, M. A., Zeng, X., Misra, V., and Beljaars, A.: Integration of a prognostic sea surface skin temperature scheme into weather and climate models, *Journal of Geophysical Research*, 113, D21 117, <https://doi.org/10.1029/2008JD010607>, 2008.
- Börner, R.: Modeling diurnal sea surface warming in the tropical ocean, Master’s thesis, Niels Bohr Institute, University of Copenhagen, https://nbi.ku.dk/english/theses/masters-theses/reyk-borner/boerner_MSc_thesis.pdf, 2021.
- 670 Börner, R.: DiuSST - Model code and data, <https://doi.org/https://doi.org/10.5281/zenodo.13363481>, 2024.
- Clayson, C. A. and Chen, A.: Sensitivity of a coupled single-column model in the tropics to treatment of the interfacial parameterizations, *Journal of climate*, 15, 1805–1831, https://journals.ametsoc.org/view/journals/clim/15/14/1520-0442_2002_015_1805_soacsc_2.0.co_2.xml, 2002.
- Coppin, D. and Bony, S.: Internal variability in a coupled general circulation model in radiative-convective equilibrium, *Geophysical Research Letters*, 44, 5142–5149, <https://doi.org/10.1002/2017GL073658>, 2017.
- 675 DeCosmo, J., Katsaros, K. B., Smith, S. D., Anderson, R. J., Oost, W. A., Bumke, K., and Chadwick, H.: Air-sea exchange of water vapor and sensible heat: The Humidity Exchange Over the Sea (HEXOS) results, *Journal of Geophysical Research: Oceans*, 101, 12 001–12 016, <https://doi.org/https://doi.org/10.1029/95JC03796>, 1996.
- DeMott, C. A., Klingaman, N. P., and Woolnough, S. J.: Atmosphere-ocean coupled processes in the Madden-Julian oscillation, *Reviews of Geophysics*, 53, 1099–1154, <https://doi.org/https://doi.org/10.1002/2014RG000478>, 2015.
- 680 Denman, K. L. and Gargett, A. E.: Time and space scales of vertical mixing and advection of phytoplankton in the upper ocean, *Limnology and Oceanography*, 28, 801–815, <https://doi.org/10.4319/lo.1983.28.5.0801>, 1983.

- Donlon, C. J., Minnett, P. J., Gentemann, C., Nightingale, T. J., Barton, I. J., Ward, B., and Murray, M. J.: Toward Improved Validation of Satellite Sea Surface Skin Temperature Measurements for Climate Research, *Journal of Climate*, 15, 353–369, [https://doi.org/10.1175/1520-0442\(2002\)015<0353:TIVOSS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0353:TIVOSS>2.0.CO;2), publisher: American Meteorological Society Section: Journal of Climate, 2002.
- Edson, J., Crawford, T., Crescenti, J., Farrar, T., Frew, N., Gerbi, G., Helmis, C., Hristov, T., Khelif, D., and Jessup, A.: The coupled boundary layers and air–sea transfer experiment in low winds, *Bulletin of the American Meteorological Society*, 88, 341–356, 2007.
- Edson, J. B., Jampana, V., Weller, R. A., Bigorre, S. P., Plueddemann, A. J., Fairall, C. W., Miller, S. D., Mahrt, L., Vickers, D., and Hersbach, H.: On the Exchange of Momentum over the Open Ocean, *Journal of Physical Oceanography*, 43, 1589–1610, <https://doi.org/10.1175/JPO-D-12-0173.1>, publisher: American Meteorological Society Section: Journal of Physical Oceanography, 2013.
- Fairall, C. W., Bradley, E. F., Godfrey, J. S., Wick, G. A., Edson, J. B., and Young, G. S.: Cool-skin and warm-layer effects on sea surface temperature, *Journal of Geophysical Research: Oceans*, 101, 1295–1308, <https://doi.org/10.1029/95JC03190>, 1996.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk Parameterization of Air–Sea Fluxes: Updates and Verification for the COARE Algorithm, *Journal of Climate*, 16, 571–591, [https://doi.org/10.1175/1520-0442\(2003\)016<0571:BPOASF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0571:BPOASF>2.0.CO;2), publisher: American Meteorological Society Section: Journal of Climate, 2003.
- Foreman-Mackey, D.: corner.py: Scatterplot matrices in Python, *Journal of Open Source Software*, 1, 24, <https://doi.org/10.21105/joss.00024>, 2016.
- Foreman-Mackey, D., Hogg, D. W., Lang, D., and Goodman, J.: emcee : The MCMC Hammer, *Publications of the Astronomical Society of the Pacific*, 125, 306–312, <https://doi.org/10.1086/670067>, 2013.
- Friehe, C. A. and Schmitt, K. F.: Parameterization of air–sea interface fluxes of sensible heat and moisture by the bulk aerodynamic formulas, *Journal of Physical Oceanography*, 6, 801–809, 1976.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B.: *Bayesian Data Analysis*, CRC Press, 2013.
- Gentemann, C. L., Donlon, C. J., Stuart-Menteth, A., and Wentz, F. J.: Diurnal signals in satellite sea surface temperature measurements, *Geophysical Research Letters*, 30, <https://doi.org/https://doi.org/10.1029/2002GL016291>, 2003.
- Gentemann, C. L., Minnett, P. J., Borgne, P. L., and Merchant, C. J.: Multi-satellite measurements of large diurnal warming events, *Geophysical Research Letters*, 35, <https://doi.org/https://doi.org/10.1029/2008GL035730>, 2008.
- Gentemann, C. L., Minnett, P. J., and Ward, B.: Profiles of ocean surface heating (POSH): A new model of upper ocean diurnal warming, *J. Geophys. Res.*, 114, C07017, <https://doi.org/10.1029/2008JC004825>, 2009.
- Goodman, J. and Weare, J.: Ensemble samplers with affine invariance, *Communications in Applied Mathematics and Computational Science*, 5, 65–80, <https://doi.org/10.2140/camcos.2010.5.65>, 2010.
- Haerter, J. O.: Convective Self-Aggregation As a Cold Pool-Driven Critical Phenomenon, *Geophys. Res. Lett.*, 46, 4017–4028, <https://doi.org/10.1029/2018GL081817>, 2019.
- Haerter, J. O., Meyer, B., and Nissen, S. B.: Diurnal self-aggregation, *npj Clim. Atmos. Sci.*, 3, 1–11, <https://doi.org/10.1038/s41612-020-00132-z>, publisher: Springer US _eprint: 2001.04740, 2020.
- Hohenegger, C. and Stevens, B.: Coupled radiative convective equilibrium simulations with explicit and parameterized convection, *Journal of Advances in Modeling Earth Systems*, 8, 1468–1482, <https://doi.org/https://doi.org/10.1002/2016MS000666>, 2016.
- Hughes, K. G., Moum, J. N., and Shroyer, E. L.: Heat Transport through Diurnal Warm Layers, *Journal of Physical Oceanography*, 50, 2885–2905, <https://doi.org/10.1175/JPO-D-20-0079.1>, publisher: American Meteorological Society Section: Journal of Physical Oceanography, 2020.

- Jensen, G. G., Fiévet, R., and Haerter, J. O.: The Diurnal Path to Persistent Convective Self-Aggregation, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002 923, <https://doi.org/https://doi.org/10.1029/2021MS002923>, 2022.
- Jia, C., Minnett, P. J., and Luo, B.: Significant Diurnal Warming Events Observed by Sairdron at High Latitudes, *Journal of Geophysical Research: Oceans*, 128, <https://doi.org/https://doi.org/10.1029/2022JC019368>, 2023.
- 725 Jiang, X., Adames, \. F., Kim, D., Maloney, E. D., Lin, H., Kim, H., Zhang, C., DeMott, C. A., and Klingaman, N. P.: Fifty years of research on the Madden-Julian Oscillation: Recent progress, challenges, and perspectives, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030 911, 2020.
- Johnson, R. H., Rickenbach, T. M., Rutledge, S. A., Ciesielski, P. E., and Schubert, W. H.: Trimodal Characteristics of Tropical Convection, *Journal of Climate*, 12, 2397–2418, [https://doi.org/10.1175/1520-0442\(1999\)012<2397:TCOTC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2397:TCOTC>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Climate*, 1999.
- 730 Kantha, L. H. and Clayson, C. A.: An improved mixed layer model for geophysical applications, *Journal of Geophysical Research: Oceans*, 99, 25 235–25 266, <https://doi.org/https://doi.org/10.1029/94JC02257>, 1994.
- Kara, A. B., Wallcraft, A. J., Metzger, E. J., Hurlburt, H. E., and Fairall, C. W.: Wind Stress Drag Coefficient over the Global Ocean, *Journal of Climate*, 20, 5856–5864, <https://doi.org/10.1175/2007JCLI1825.1>, publisher: American Meteorological Society Section: *Journal of*
- 735 *Climate*, 2007.
- Karlowska, E., Matthews, A. J., Webber, B. G. M., Graham, T., and Xavier, P.: The effect of diurnal warming of sea-surface temperatures on the propagation speed of the Madden–Julian oscillation, *Quarterly Journal of the Royal Meteorological Society*, n/a, <https://doi.org/https://doi.org/10.1002/qj.4599>, 2023.
- Kawai, Y. and Kawamura, H.: Evaluation of the Diurnal Warming of Sea Surface Temperature Using Satellite-Derived Marine Meteorological
- 740 Data, *Journal of Oceanography*, p. 10, 2002.
- Kawai, Y. and Wada, A.: Diurnal sea surface temperature variation and its impact on the atmosphere and ocean: A review, *Journal of Oceanography*, 63, 721–744, <https://doi.org/10.1007/s10872-007-0063-0>, 2007.
- Khairoutdinov, M. F. and Randall, D. A.: Cloud Resolving Modeling of the ARM Summer 1997 IOP: Model Formulation, Results, Uncertainties, and Sensitivities, *Journal of the Atmospheric Sciences*, 60, 607–625, [https://doi.org/10.1175/1520-0469\(2003\)060<0607:CRMOTA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<0607:CRMOTA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2003.
- 745 Kondo, J., Sasano, Y., and Ishii, T.: On Wind-Driven Current and Temperature Profiles with Diurnal Period in the Oceanic Planetary Boundary Layer, *Journal of Physical Oceanography*, 9, 360–372, [https://doi.org/10.1175/1520-0485\(1979\)009<0360:OWDCAT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1979)009<0360:OWDCAT>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Physical Oceanography*, 1979.
- 750 Laboratory, N. P. S.: COARE-algorithm v3.6, <https://github.com/NOAA-PSL/COARE-algorithm>, 2023.
- Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air–sea flux data set, *Clim Dyn*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.
- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Reviews of Geophysics*, 32, 363, <https://doi.org/10.1029/94RG01872>, 1994.
- 755 Li, W., Yu, R., Liu, H., and Yu, Y.: Impacts of diurnal cycle of SST on the intraseasonal variation of surface heat flux over the western Pacific warm pool, *Advances in Atmospheric Sciences*, 18, 793–806, 2001.
- Madden, R. A. and Julian, P. R.: Description of global-scale circulation cells in the tropics with a 40–50 day period, *Journal of Atmospheric Sciences*, 29, 1109–1123, 1972.

- Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Reviews of Geophysics*, 20, 760 851–875, <https://doi.org/https://doi.org/10.1029/RG020i004p00851>, 1982.
- Minnett, P. J.: Radiometric measurements of the sea-surface skin temperature: The competing roles of the diurnal thermocline and the cool skin, *Int. J. Remote Sens.*, 24, 5033–5047, <https://doi.org/10.1080/0143116031000095880>, 2003.
- Minnett, P. J., Knuteson, R. O., Best, F. A., Osborne, B. J., Hanafin, J. A., and Brown, O. B.: The Marine-Atmospheric Emitted Radiance Interferometer: A High-Accuracy, Seagoing Infrared Spectroradiometer, *Journal of Atmospheric and Oceanic Technology*, 18, 994–1013, 765 [https://doi.org/10.1175/1520-0426\(2001\)018<0994:TMAERI>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<0994:TMAERI>2.0.CO;2), publisher: American Meteorological Society Section: Journal of Atmospheric and Oceanic Technology, 2001.
- Moncrieff, M. W.: The Multiscale Organization of Moist Convection and the Intersection of Weather and Climate, in: *Climate Dynamics: Why Does Climate Vary?*, pp. 3–26, American Geophysical Union (AGU), ISBN 978-1-118-67039-2, <https://doi.org/10.1029/2008GM000838>, 2010.
- 770 Müller, S. K. and Hohenegger, C.: Self-Aggregation of Convection in Spatially Varying Sea Surface Temperatures, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001 698, <https://doi.org/https://doi.org/10.1029/2019MS001698>, 2020.
- Nissen, S. B. and Haerter, J. O.: Circling in on convective self-aggregation, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035 331, 2021.
- Noh, Y. and Jin Kim, H.: Simulations of temperature and turbulence structure of the oceanic boundary layer with the improved near-surface 775 process, *Journal of Geophysical Research: Oceans*, 104, 15 621–15 634, <https://doi.org/10.1029/1999JC900068>, 1999.
- Noh, Y., Min, H. S., and Raasch, S.: Large Eddy Simulation of the Ocean Mixed Layer: The Effects of Wave Breaking and Langmuir Circulation, *Journal of Physical Oceanography*, 34, 720–735, [https://doi.org/10.1175/1520-0485\(2004\)034<0720:LESOTO>2.0.CO;2](https://doi.org/10.1175/1520-0485(2004)034<0720:LESOTO>2.0.CO;2), publisher: American Meteorological Society Section: Journal of Physical Oceanography, 2004.
- Noh, Y., Lee, E., Kim, D.-H., Hong, S.-Y., Kim, M.-J., and Ou, M.-L.: Prediction of the diurnal warming of sea surface temperature using an 780 atmosphere-ocean mixed layer coupled model, *Journal of Geophysical Research*, 116, C11 023, <https://doi.org/10.1029/2011JC006970>, 2011.
- Pope, S. B.: *Turbulent flows*, Cambridge University Press, 2000.
- Price, J. F., Weller, R. A., and Pinkel, R.: Diurnal cycling: Observations and models of the upper ocean response to diurnal heating, cooling, and wind mixing, *Journal of Geophysical Research*, 91, 8411, <https://doi.org/10.1029/JC091iC07p08411>, 1986.
- 785 Price, J. F., Weller, R. A., Bowers, C. M., and Briscoe, M. G.: Diurnal response of sea surface temperature observed at the long-term upper ocean study (34°N, 70°W) in the Sargasso Sea, *Journal of Geophysical Research: Oceans*, 92, 14 480–14 490, <https://doi.org/https://doi.org/10.1029/JC092iC13p14480>, 1987.
- Schiller, A. and Godfrey, J. S.: A diagnostic model of the diurnal cycle of sea surface temperature for use in coupled ocean-atmosphere models, *Journal of Geophysical Research: Oceans*, 110, <https://doi.org/10.1029/2005JC002975>, 2005.
- 790 Schumacher, R. S. and Rasmussen, K. L.: The formation, character and changing nature of mesoscale convective systems, *Nature Reviews Earth & Environment*, 1, 300–314, <https://doi.org/10.1038/s43017-020-0057-7>, number: 6 Publisher: Nature Publishing Group, 2020.
- Seo, H., Subramanian, A. C., Miller, A. J., and Cavanaugh, N. R.: Coupled Impacts of the Diurnal Cycle of Sea Surface Temperature on the Madden–Julian Oscillation, *Journal of Climate*, 27, 8422–8443, <https://doi.org/10.1175/JCLI-D-14-00141.1>, publisher: American Meteorological Society Section: Journal of Climate, 2014.
- 795 Seo, H., O’Neill, L. W., Bourassa, M. A., Czaja, A., Drushka, K., Edson, J. B., Fox-Kemper, B., Frenger, I., Gille, S. T., Kirtman, B. P., Minobe, S., Pendergrass, A. G., Renault, L., Roberts, M. J., Schneider, N., Small, R. J., Stoffelen, A., and Wang, Q.: Ocean Mesoscale

- and Frontal-Scale Ocean–Atmosphere Interactions and Influence on Large-Scale Climate: A Review, *Journal of Climate*, 36, 1981–2013, <https://doi.org/10.1175/JCLI-D-21-0982.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2023.
- 800 Shamekh, S., Muller, C., Duvel, J.-P., and d’Andrea, F.: Self-Aggregation of Convective Clouds With Interactive Sea Surface Temperature, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002164, 2020a.
- Shamekh, S., Muller, C., Duvel, J.-P., and d’Andrea, F.: How do ocean warm anomalies favor the aggregation of deep convective clouds?, *Journal of the Atmospheric Sciences*, 77, 3733–3745, 2020b.
- Shevchenko, R., Hohenegger, C., and Schmitt, M.: Impact of Diurnal Warm Layers on Atmospheric Convection, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD038473, <https://doi.org/10.1029/2022JD038473>, 2023.
- 805 Siebesma, A. P., Bony, S., Jakob, C., and Stevens, B.: *Clouds and Climate: Climate Science’s Greatest Challenge*, Cambridge University Press, 2020.
- Skyllingstad, E. D., Szoek, S. P. d., and O’Neill, L. W.: Modeling the Transient Response of Tropical Convection to Mesoscale SST Variations, *Journal of the Atmospheric Sciences*, 76, 1227–1244, <https://doi.org/10.1175/JAS-D-18-0079.1>, publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2019.
- 810 Slingo, J., Inness, P., Neale, R., Woolnough, S., and Yang, G.-Y.: Scale interactions on diurnal to seasonal timescales and their relevance to model systematic errors, *Annals of Geophysics*, 46, <https://doi.org/10.4401/ag-3383>, 2003.
- Smith, S. D.: Coefficients for sea surface wind stress, heat flux, and wind profiles as a function of wind speed and temperature, *Journal of Geophysical Research*, 93, 15467, <https://doi.org/10.1029/JC093iC12p15467>, 1988.
- Soloviev, A. and Lukas, R.: Observation of large diurnal warming events in the near-surface layer of the western equatorial Pacific warm pool, *Deep Sea Research Part I: Oceanographic Research Papers*, 44, 1055–1076, [https://doi.org/10.1016/S0967-0637\(96\)00124-0](https://doi.org/10.1016/S0967-0637(96)00124-0), 1997.
- 815 Son, S. and Wang, M.: Diffuse attenuation coefficient of the photosynthetically available radiation $K_d(\text{PAR})$ for global open ocean and coastal waters, *Remote Sensing of Environment*, 159, 250–258, <https://doi.org/10.1016/j.rse.2014.12.011>, 2015.
- Stull, R. B. and Kraus, E. B.: The transient model of the upper ocean, *Journal of Geophysical Research: Oceans*, 92, 10745–10755, 1987.
- Takaya, Y., Bidlot, J.-R., Beljaars, A. C. M., and Janssen, P. A. E. M.: Refinements to a prognostic scheme of skin sea surface temperature, *Journal of Geophysical Research*, 115, C06009, <https://doi.org/10.1029/2009JC005985>, 2010.
- 820 Tan, J., Jakob, C., Rossow, W. B., and Tselioudis, G.: Increases in tropical rainfall driven by changes in frequency of organized deep convection, *Nature*, 519, 451–454, <https://doi.org/10.1038/nature14339>, 2015.
- Tompkins, A. M.: Organization of tropical convection in low vertical wind shears: The role of cold pools, *Journal of the atmospheric sciences*, 58, 1650–1672, 2001.
- 825 Tompkins, A. M. and Craig, G. C.: Radiative–convective equilibrium in a three-dimensional cloud-ensemble model, *Quarterly Journal of the Royal Meteorological Society*, 124, 2073–2097, <https://doi.org/10.1002/qj.49712455013>, 1998.
- Tompkins, A. M. and Semie, A. G.: Impact of a mixed ocean layer and the diurnal cycle on convective aggregation, *Journal of Advances in Modeling Earth Systems*, <https://doi.org/10.1029/2020MS002186>, 2021.
- Trenberth, K. E., Large, W. G., and Olson, J. G.: The Effective Drag Coefficient for Evaluating Wind Stress over the Oceans, *Journal of Climate*, 2, 1507–1516, [https://doi.org/10.1175/1520-0442\(1989\)002<1507:TEDCFE>2.0.CO;2](https://doi.org/10.1175/1520-0442(1989)002<1507:TEDCFE>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Climate*, 1989.
- 830 Voldoire, A., Roehrig, R., Giordani, H., Waldman, R., Zhang, Y., Xie, S., and Bouin, M.-N.: Assessment of the sea surface temperature diurnal cycle in CNRM-CM6-1 based on its 1D coupled configuration, *Geoscientific Model Development*, 15, 3347–3370, <https://doi.org/10.5194/gmd-15-3347-2022>, 2022.

- 835 Ward, B.: Near-surface ocean temperature, *Journal of Geophysical Research: Oceans*, 111, <https://doi.org/10.1029/2004JC002689>, 2006.
- Webster, P. J., Clayson, C. A., and Curry, J. A.: Clouds, radiation, and the diurnal cycle of sea surface temperature in the tropical western Pacific, *J. Clim.*, 9, [https://doi.org/10.1175/1520-0442\(1996\)009<1712:CRATDC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<1712:CRATDC>2.0.CO;2), 1996.
- Weih, R. R. and Bourassa, M. A.: Modeled diurnally varying sea surface temperatures and their influence on surface heat fluxes, *Journal of Geophysical Research: Oceans*, 119, 4101–4123, <https://doi.org/https://doi.org/10.1002/2013JC009489>, 2014.
- 840 Wells, N. C. and King-Hele, S.: Parametrization of tropical ocean heat flux, *Quarterly Journal of the Royal Meteorological Society*, 116, 1213–1224, <https://doi.org/10.1002/qj.49711649511>, 1990.
- Wing, A. A., Emanuel, K., Holloway, C. E., and Muller, C.: *Convective Self-Aggregation in Numerical Simulations: A Review*, *Surv. Geophys.*, 38, 1173–1197, <https://doi.org/10.1007/s10712-017-9408-4>, ISBN: 1071201794 Publisher: Springer Netherlands, 2017.
- Witte, C. R., Zappa, C. J., and Edson, J. B.: The Response of Ocean Skin Temperature to Rain: Observations and Implications for Parameter-
845 ization of Rain-Induced Fluxes, *Journal of Geophysical Research: Oceans*, 128, e2022JC019146, <https://doi.org/10.1029/2022JC019146>, 2023.
- Wong, E. W. and Minnett, P. J.: The Response of the Ocean Thermal Skin Layer to Variations in Incident Infrared Radiation, *Journal of Geophysical Research: Oceans*, 123, 2475–2493, <https://doi.org/https://doi.org/10.1002/2017JC013351>, 2018.
- Woolnough, S. J., Vitart, F., and Balmaseda, M. A.: The role of the ocean in the Madden–Julian Oscillation: Implications for MJO prediction,
850 *Quarterly Journal of the Royal Meteorological Society*, 133, 117–128, <https://doi.org/10.1002/qj.4>, 2007.
- Wurl, O., Ekau, W., Landing, W. M., and Zappa, C. J.: Sea surface microlayer in a changing ocean—A perspective, *Elementa: Science of the Anthropocene*, 5, 2017.
- Yanase, T., Nishizawa, S., Miura, H., Takemi, T., and Tomita, H.: New Critical Length for the Onset of Self-Aggregation of Moist Convection, *Geophysical Research Letters*, 47, e2020GL088763, <https://doi.org/10.1029/2020GL088763>, 2020.
- 855 Zeng, X. and Beljaars, A.: A prognostic scheme of sea surface skin temperature for modeling and data assimilation, *Geophysical Research Letters*, 32, <https://doi.org/https://doi.org/10.1029/2005GL023030>, 2005.
- Zeng, X., Zhao, M., Dickinson, R. E., and He, Y.: A multiyear hourly sea surface skin temperature data set derived from the TOGA TAO bulk temperature and wind speed over the tropical Pacific, *Journal of Geophysical Research: Oceans*, 104, 1525–1536, <https://doi.org/10.1029/1998JC900060>, 1999.
- 860 Zhang, C.: Madden-Julian Oscillation, *Reviews of Geophysics*, 43, 2005.
- Zhao, N. and Nasuno, T.: How Does the Air-Sea Coupling Frequency Affect Convection During the MJO Passage?, *Journal of Advances in Modeling Earth Systems*, 12, <https://doi.org/https://doi.org/10.1029/2020MS002058>, 2020.
- Álvarez Borrego, S.: Phytoplankton biomass and production in the Gulf of California: a review, *Botanica Marina*, 55, <https://doi.org/10.1515/bot.2011.105>, 2012.