

1 **Seasonal Air Concentration Variability, Gas/Particle Partitioning, Precipitation**
2 **Scavenging, and Air-Water Equilibrium of Organophosphate Esters in Southern**
3 **Canada**

4 Yuening Li,¹ Faqiang Zhan,¹ Chubashini Shunthirasingham,² Ying Duan Lei,¹ Jenny Oh,^{1,3}
5 Amina Ben Chaaben,⁴ Zhe Lu,⁴ Kelsey Lee,⁵ Frank A. P. C. Gobas,⁵ Hayley Hung,² Frank
6 Wania^{1,3*}

7 ¹ *Department of Physical and Environmental Sciences, University of Toronto Scarborough,*
8 *1265 Military Trail, Toronto, Ontario, Canada M1C 1A4*

9 ² *Environment and Climate Change Canada, Downsview, 4905 Dufferin St, North York,*
10 *Ontario, Canada M3H 5T4*

11 ³ *Department of Chemistry, University of Toronto Scarborough, 1265 Military Trail, Toronto,*
12 *Ontario, Canada M1C 1A4*

13 ⁴ *Institut des Sciences de la Mer de Rimouski, Université du Québec à Rimouski, 300 allée*
14 *des Ursulines, Rimouski, Québec, Canada G5L 3A1*

15 ⁵ *School of Resource and Environmental Management, Simon Fraser University, 8888*
16 *University Dr, Burnaby, British Columbia, Canada V5A 1S6*

17 *Corresponding author: frank.wania@utoronto.ca

18 **Abstract**

19 In response to increasing production and application volumes, organophosphate esters
20 (OPEs) have emerged as pervasively detected contaminants in various environmental media,
21 with concentrations often exceeding those of traditional organic contaminants. Despite the
22 recognition of the atmosphere's important role in dispersing OPEs and a substantial number
23 of studies quantifying OPEs in air, investigations into atmospheric phase distribution
24 processes are rare. Using measurements of OPEs in the atmospheric gas and particle phase,
25 in precipitation and in surface water collected in Southern Canada, we explored the seasonal
26 concentration variability, gas/particle partitioning behaviour, precipitation scavenging, and
27 air-water equilibrium status of OPEs. Whereas consistent seasonal trends were not observed
28 for OPEs concentrations in precipitation or atmospheric particles, gas phase concentrations
29 of several OPEs were elevated during the summer in suburban Toronto and at two remote
30 sites on Canada's east and west coast. Apparent enthalpies of air-surface exchange fell
31 mainly within or slightly above the range of air/water and air/octanol enthalpies of exchange,
32 indicating the influence of local air-surface exchange processes and/or seasonally variable
33 source strength. While many OPEs were present in notable fraction in both gas and particle
34 phase, no clear relationship with compound volatility was apparent, although there was a
35 tendency for higher particle-bound fractions at lower temperature. High precipitation
36 scavenging ratios for OPEs measured at the two coastal sites are consistent with low air-
37 water partitioning ratios and the association with particles. Although beset by large
38 uncertainties, air-water equilibrium calculations suggest net deposition of gaseous OPEs
39 from the atmosphere to the Salish Sea and the St. Lawrence River and Estuary. The
40 measured seasonal concentration variability is likely less a reflection of temperature driven
41 air-surface exchange and instead indicates that more OPE enter, or are formed in, the
42 atmosphere in summer. More research is needed to better understand the atmospheric gas-
43 particle partitioning behaviour of the OPEs and how it may be influenced by transformation
44 reactions.

45 **Key words:**

46 OPEs, air, precipitation, water, partitioning, air-water exchange, relative abundance

47 **1. INTRODUCTION**

48 Organophosphate esters (OPEs) are synthetic organophosphorus compounds consisting of a
49 central phosphate molecule substituted with non-halogenated, halogenated alkyl, or aryl
50 groups. Widely used as flame retardants, plasticizers, stabilizers, and defoaming agents in
51 various industries and consumer products (Environment and Climate Change Canada,
52 2023abcd; Salamova et al., 2016; van der Veen and de Boer, 2012), OPEs are typically
53 physically incorporated into materials rather than chemically bonded (Wang et al., 2020b;
54 Wong et al., 2018), facilitating their release into the environment. Following restrictions on
55 many brominated flame retardants, e.g. through listing in the Stockholm Convention, OPEs
56 use has increased, reaching 620 kilotons globally in 2013, accounting for 30% of total flame
57 retardant usage (Sühring et al., 2016; Xie et al., 2022). The extensive application of OPEs,
58 coupled with their potential for long-range atmospheric transport (Na et al., 2020; Sühring
59 et al., 2016) and persistence (Möller et al., 2012; Salamova et al., 2014), has resulted in their
60 ubiquitous presence in the environment (Han et al., 2020; Li et al., 2019a, b; Lu et al., 2017;
61 Mi et al., 2023; Regnery and Püttmann, 2009; Stackelberg et al., 2007), often at
62 concentrations exceeding those of traditional flame retardants and plasticizers (Salamova et
63 al., 2014; Shoeib et al., 2014; Zhao et al., 2021b). Given their potential toxicity (Gu et al.,
64 2019; Li et al., 2020; Rosenmai et al., 2021; Wang et al., 2022; Yan and Hales, 2019, 2020),
65 understanding the fate, occurrence, and distribution of OPEs in the environment is critical
66 for assessing their ecological and human health impacts.

67 The atmosphere plays a key role in the dispersion and transport of OPEs, with concentrations
68 and spatial and temporal variability in air being influenced by emission sources, atmospheric
69 transport, chemical transformation (Liu et al., 2023; Liu and Mabury, 2019) and deposition
70 processes. The distribution of OPEs between different atmospheric phases (gas phase,
71 particles, precipitation) affects these processes and is influenced by their partition properties.
72 Most studies on OPEs in the atmosphere report concentrations in the particle phase, whereas
73 studies on the presence in the gas phase are far more limited, which may be related to the
74 relatively short half-lives of gas phase OPEs (Shi et al., 2024; Zhang et al., 2016). However,
75 gaseous OPEs can constitute 15% to 65% of atmospheric OPEs (Möller et al., 2011), and
76 diffusive air-water gas exchange of OPEs can be 2-3 orders of magnitude higher than dry
77 particle deposition (Castro-Jiménez et al., 2016; Ma et al., 2021), highlighting the need for
78 more research on OPE vapours.

79 Precipitation acts as a major pathway for the removal and redistribution of OPEs from the

80 atmosphere to aquatic and terrestrial environments (Shi et al., 2024). It can scavenge and
81 deposit both gas-phase and particle-bound OPEs. Depending on regional emissions,
82 temperature, precipitation type, and the physicochemical properties of the OPEs (Lei and
83 Wania, 2004), the wet deposition flux of OPEs can be significantly larger than the dry
84 deposition flux (Kim and Kannan, 2018). Despite its importance, fewer than ten studies have
85 reported OPE concentrations in precipitation (Bacaloni et al., 2008; Casas et al., 2021; Fries
86 and Püttmann, 2003; Kim and Kannan, 2018; Marklund et al., 2005b; Mihajlović and Fries,
87 2012; Regnery and Püttmann, 2009; Zhang et al., 2020), and only one study has reported
88 precipitation scavenging ratios for atmospheric OPEs (Casas et al., 2021).

89 OPEs can enter water bodies through air-water gas exchange (Castro-Jiménez et al., 2016;
90 Ma et al., 2021), wet and dry deposition (Castro-Jiménez et al., 2016; Kim and Kannan,
91 2018; Ma et al., 2021), wastewater effluent (Marklund et al., 2005a), industrial and
92 municipal discharges (Bacaloni et al., 2008; Fries and Püttmann, 2003), and surface runoff
93 (Awonaike et al., 2021; Regnery and Püttmann, 2010). Some OPEs, including tris(1-chloro-
94 2-propyl) phosphate (TCPP) and tris (phenyl) phosphate (TPhP), have been detected in fish
95 (Ma et al., 2013; Sundkvist et al., 2010). A comprehensive understanding of the
96 environmental fate and occurrence of OPEs, and in particular a better understanding of the
97 contribution that the atmosphere makes for the delivery of OPEs to aquatic ecosystems,
98 would benefit from investigations that quantify OPE concentrations in multiple
99 environmental media sampled in the same area and at the same time. Despite the substantial
100 number of studies on OPEs in the environment, those examining OPEs across three or more
101 phases are very rare (He et al., 2019; Li et al., 2019b; Mi et al., 2023) and most studies focus
102 on just one or two media, usually gas and/or particle phases (Li et al., 2018; Ma et al., 2022;
103 Sühling et al., 2016; Zhao et al., 2021b) or the water phase (Choo and Oh, 2020; Ding et al.,
104 2015; McDonough et al., 2018; Shi et al., 2020). No previous study has investigated OPEs
105 in atmospheric gas and particle phases, precipitation, and surface water simultaneously.

106 To address this research gap and gather information on the contribution that the atmosphere
107 makes to OPEs in coastal waters of Southern Canada, we aimed to characterize the
108 occurrence, behavior, and fate of OPEs in different atmospheric phases. We measured OPEs
109 in precipitation and atmospheric gas and particle phase for one year at two remote sites on
110 Canada's East and West coast, respectively, and complemented this dataset with the results
111 of a year-long measurement campaign of OPEs in the gas and particle phase in Toronto. We
112 further used passive samplers to gather data on the spatial variability of OPE concentrations

113 in the atmospheric gas phase and in water in the two coastal regions. The passive air
114 sampling data have been presented previously (Li et al., submitted). This unique data set
115 allowed us to estimate the gas-particle distribution in the atmosphere, precipitation
116 scavenging ratios, and the state of air-surface water equilibrium, often in their seasonal
117 dependence or their variability between urban, rural and remote locales. Finally, we used
118 this dataset to explore the relative abundance of OPEs in the different types of samples.

119 **2. MATERIALS AND METHODS**

120 **2.1 Active Air Sampling and Precipitation Collection.** 24-hour air samples were collected
121 monthly for one year using a high-volume active air sampler (AAS); twelve at a location on
122 Saturna Island, British Columbia (BC) (48.7753N, -123.1283W), and twelve in the vicinity
123 of Tadoussac, Quebec (QC) (48.1415N, -69.6991W). Forty-eight consecutive week-long
124 AASs were taken with a mid-volume pump in the Eastern suburbs of Toronto (43.78371 N,
125 -79.19027 W) (Li et al., 2023a, b, 2024). At all three sites, polyurethane foam
126 (PUF)/XAD/PUF sandwiches and glass-fiber filters (GFFs) were used to collect OPEs in
127 the gas and particle phase, respectively. Precipitation samples (PCPNs) were collected at the
128 AAS sampling locations in BC and QC during the same months as the air samples and the
129 sampling length was ~ 30 days (Oh et al., 2023; Zhan et al., 2023).

130 **2.2 Passive Air and Water Sampling.** In QC, 86 passive air samplers (PASs) were
131 deployed at 71 unique sampling sites on either shore of the St. Lawrence River and Estuary,
132 including in Montreal and Quebec City between 2019 and 2022. In BC, 83 PASs were
133 deployed at 47 sites in the lower mainland around Vancouver and on the Canadian shore of
134 the Salish Sea during different time periods between 2020 and 2022. More details are given
135 in Table S5 in the Supporting Information (SI) of Li et al. (submitted) and in previous
136 publications (Oh et al., 2023; Zhan et al., 2023).

137 Forty-eight low-density polyethylene (LDPE) based passive water samplers (PWSs) were
138 spiked with performance reference compounds (PRCs), deployed at 10 sites in BC and 10
139 sites in QC, and collected after deployment lasting 20-35 days in BC and 27-70 days in QC.
140 Detailed information on the PWS sampling is provided in the SI (Table S13) and previous
141 publications (Oh et al., 2023; Zhan et al., 2023).

142 **2.3 Sample Analysis.** Prior to extraction all samples were spiked with seven isotopically
143 labeled OPEs (Table S1) as surrogates. XAD from the PASs, the PUF/XAD sandwiches and
144 GFFs from the AASs were extracted using a Dionex Accelerated Solvent Extractor 350. The

145 PCPN and PWS samples were extracted using liquid-liquid extraction with dichloromethane
146 and soaking in hexane, respectively. Extracts were concentrated to 0.5 mL using a rotary
147 evaporator and nitrogen blow-down. Triamyl phosphate was added into the concentrated
148 extracts as an injection standard. Gas chromatography-tandem mass spectrometry (GC-
149 MS/MS) was used to detect and quantify 16 OPEs (Tables S1 and S2).

150 **2.4 Quality Assurance and Quality Control.** All extraction and concentration procedures
151 were carried out in a trace analytical laboratory. The glassware was cleaned using a machine
152 with detergents, then rinsed with deionized water, and finally baked with GFFs at 450 °C in
153 a muffle furnace for 24 hours. Experimental materials that came into contact with samples
154 or extracts were thoroughly cleaned and rinsed three times with solvents (acetone and
155 hexane, or dichloromethane) before use. Field blanks, procedure blanks, and solvent blanks
156 were prepared with each batch of extractions and analyses (Oh et al., 2023; Zhan et al., 2023).
157 OPEs were not found in procedure or solvent blanks. Only a few analytes were present in
158 the field blanks, and for these, the average detected amount was subtracted from the amounts
159 of target chemicals in the field samples. Method detection limits (MDLs) were calculated as
160 three times the standard deviations of levels in field blanks when analytes were detected
161 (signal-to-noise ratio (S/N) > 3); otherwise, MDLs were based on concentrations at which
162 S/N is 3 (Desimoni and Brunetti, 2015). MDLs are provided in the Supplementary
163 Information (Tables S5, S8, S10, S11, and S13). The average recoveries of five surrogates
164 in AASs, PCPNs, and PWSs ranged from 78% to 232% (Table S3). The concentrations
165 reported have been corrected for recovery.

166 **2.5 Data Analysis.** Water concentration of OPEs were calculated from the amounts
167 quantified in PWS extracts following the method by Booij and Smedes (Booij et al., 2003;
168 Booij and Smedes, 2010), with details provided by Oh et al. (2023).

169 The fraction of an OPE in the particle phase (Φ , %) was obtained by dividing the particle-
170 phase concentration by the sum of concentrations in the gas and particle phase. Gas-particle
171 partition ratios K_{PA} ($\text{m}^3 \text{air g}^{-1} \text{aerosol}$) were derived by dividing the measured concentrations
172 of an OPE in the particle phase (pg m^{-3}) by the product of the concentrations of particles less
173 than 2.5 μm in diameter ($\text{PM}_{2.5}$, g m^{-3}) obtained from nearby national air pollution
174 surveillance program (NAPS) stations (Table S8) and the measured concentrations of this
175 OPE in the gas phase (pg m^{-3}). More detail is given in previous publications (Li et al., 2023a;
176 Oh et al., 2023; Zhan et al., 2023).

177 Measured scavenging ratios (SRs) were calculated as the ratios between the concentrations
178 of an OPE in precipitation and air (sum of gas and particle phase). We also estimated SRs
179 by assuming equilibrium of OPE between the atmospheric gas phase and water droplets (Oh
180 et al., 2023), and that all OPEs are sorbed to the same particles, which are scavenged with a
181 scavenging ratio W_P of 200,000 (Kim et al., 2006). An estimated SR thus is $(1-\Phi)K_{WA} +$
182 ΦW_P , where K_{WA} is the temperature-adjusted partition ratio between water and air ($K_{WA} =$
183 K_{AW}^{-1} , Table S4).

184 The fugacities of OPEs in water f_w , at average sea surface temperature T_w in K, were
185 calculated using $C_w \cdot K_{AW}(T_w) \cdot R \cdot T_w$, and those in air (f_A), at average air temperature T_A in
186 K, were derived with $C_A \cdot R \cdot T_A$, where C_w and C_A are the OPE concentrations (mol m^{-3}) in
187 water and air, respectively, and R is the gas constant.

188 3. RESULTS

189 **3.1 OPEs in the Atmospheric Gas Phase.** The gas phase concentrations obtained during
190 the three one-year AAS campaigns in Tadoussac, on Saturna Island, and in Toronto are given
191 in Table S5. The gas phase concentrations obtained by passive air sampling in QC and BC
192 have been previously reported (Li et al., submitted) with tri-n-butyl phosphate (TBP),
193 tris(2-chloroethyl) phosphate (TCEP), tris(1-chloro-2-propyl) phosphate (TCPP), and tris
194 (phenyl) phosphate (TPhP) being reliably and ubiquitously detected. Due to the higher
195 sampling volumes of the AAS ($\sim 520 \text{ m}^3$) compared to the PAS (less than 200 m^3), more
196 OPEs could be detected above the MDL in the AAS. At all three locations, TBP, TCEP,
197 TCPP, TPhP, and 2-ethylhexyl-diphenyl phosphate (EHDPP) were present above the MDL.
198 Additionally, triethyl phosphate (TEP) was detected on Saturna Island, TEP,
199 tris(1,3-dichloro-2-propyl) phosphate (TDCPP), and tris (2-butoxyethyl) phosphate (TBEP)
200 were detected in Tadoussac, and tri-propyl phosphate (TPrP) and TDCPP were detected in
201 Toronto. We are **not** comparing here the gas phase concentrations recorded in our study with
202 those reported previously, because that had already been done in Li et al. (submitted).

203 For the four most frequently detected OPEs, it is possible to compare the levels obtained
204 with the AASs on Saturna Island and in Tadoussac and by PASs at the nearby sites L43 and
205 S57. On Saturna Island, the PAS deployment at site L43 overlapped with the timeframe of
206 the AASs (between May and October 2020) (Table S8). In Tadoussac, the deployment
207 period of the PAS at S57 (November 2019 – August 2020) preceded the AASs sampling by
208 about one year (December 2020 – September 2021). Except for TBP and TPhP on Saturna

209 Island, PAS levels generally trended lower than AAS levels at both locations, albeit within
210 a factor of 5. One contributing factor to this difference could be the episodic 24-hour active
211 air sampling's inability to represent long-term concentration levels compared to PAS. For
212 instance, AAS-measured concentrations of TBP in Tadoussac ranged from below detection
213 to approximately 200 pg m^{-3} . Another factor could be the spatial distribution variability of
214 atmospheric OPEs. Despite our efforts to use PAS data from sites closest to AAS locations
215 for comparison, the PAS and AAS sampling sites were not identical. To support the
216 hypothesis that spatial and temporal variability in OPE concentrations contributes to the
217 discrepancy, we also compared AAS and PAS results for hexachlorobutadiene (HCB) and
218 hexachlorobenzene (HCB), which exhibit uniform spatial distribution and consistent
219 concentrations over time, using the same samples as for the OPEs. PAS levels for these two
220 compounds closely aligned with AAS levels within a factor of 1.5. Similarly,
221 halomethoxybenzene levels from PASs and AASs were within a factor of 3 (Zhan et al.,
222 2023).

223 **3.2 OPEs in Atmospheric Particle Phase.** The concentrations of five OPEs (TBP, TCEP,
224 TCPP, TPhP, EHDPP) in the atmospheric particles from the three AAS sampling locations
225 are compiled in Tables S9. Except for TPhP and EHDPP which were not detected in particles
226 from Saturna Island, all five OPEs most frequently detected in the gas phase could also be
227 quantified in particle samples. Again, TCPP is the most abundant OPE at all three sites.


228 Concentration levels on Saturna Island and in suburban Toronto are similar and almost one
229 order of magnitude higher than those in Tadoussac. The averaged TBP levels of 7 pg m^{-3} , 3
230 pg m^{-3} , and 9 pg m^{-3} on Saturna Island, in Tadoussac, and in Toronto are lower than those
231 in Antarctica (23 pg m^{-3}) (Wang et al., 2020a) and two order of magnitude lower than those
232 detected in cities in the Great Lakes area (130 pg m^{-3}) in 2012 (Salamova et al., 2013).
233 Except in Tadoussac (1 pg m^{-3}), the TCEP levels of 50 pg m^{-3} and 17 pg m^{-3} on Saturna
234 Island and in Toronto are higher than those in Antarctica (5 pg m^{-3}) (Wang et al., 2020a) and
235 ca. 2~4 times lower than those detected in cities in the Great Lakes region (89 pg m^{-3})
236 (Salamova et al., 2013), and two orders of magnitude lower than the reported median
237 concentration in Quebec City and near the St Lawrence River (1903 pg m^{-3}) (Sühring et al.,
238 2016). TCPP in Tadoussac, 3 pg m^{-3} , is comparable to its level in Antarctica (6 pg m^{-3})
239 (Wang et al., 2020a), and TCPP on Saturna Island (122 pg m^{-3}) and in Toronto (90 pg m^{-3})
240 are three and four times lower than those detected in cities in Great Lakes area (321 pg m^{-3})
241 (Salamova et al., 2013), and one order of magnitude lower than the detected level in Quebec

242 City and near the St Lawrence River (1557 pg m^{-3}) (Sühring et al., 2016). TPhP and EHDPP
243 levels in Antarctica (1 pg m^{-3}) (Wang et al., 2020a) are close to those in Tadoussac (2 pg m^{-3}),
244 and one order of magnitude lower than levels in Toronto (12 pg m^{-3}) and those in Quebec
245 City and near the St Lawrence River (51 pg m^{-3}) (Sühring et al., 2016). The relatively higher
246 concentration levels of certain OPEs in Antarctica, such as TBP, may be due to preferential
247 partitioning of TBP to particles at low temperatures. Compared to the sites in the Great Lakes
248 region (Salamova et al., 2013) as well as Quebec City and near the St Lawrence River region
249 (Sühring et al., 2016), our sampling sites were more rural, which could explain lower OPE
250 concentrations.

251 **3.3 OPEs in Precipitation.** Eight OPEs, i.e., TEP, TBP, TCEP, TCPP, TDCPP, TPhP,
252 TBEP, and EHDPP, were reliably detected in the precipitation samples from Saturna Island
253 and Tadoussac (Table 1 & Table S11). Concentrations are generally higher on Saturna Island
254 than in Tadoussac. The OPE levels in Tadoussac were comparable to those in Antarctica
255 (Casas et al., 2021). TDCPP detected on Saturna Island and in Tadoussac are two times to
256 one order of magnitude higher than the levels in Antarctica (Casas et al., 2021), Nanjing
257 (Zhang et al., 2020), and New York (Kim and Kannan, 2018), and our measured EHDPP
258 levels were higher than those detected in Antarctica. Overall, except for TDCPP and EHDPP,
259 the average OPE concentrations detected in our study were comparable or one order of
260 magnitude lower than literature data (Table 1). Except for TBP in Tadoussac, OPE
261 concentrations varied greatly between months, whereby no distinct and consistent seasonal
262 trends were discernible (Table S11), which is consistent with previous observations
263 (Regnery and Püttmann, 2009).

264 **3.4 OPEs in Water.** The OPEs concentrations in water, obtained with PWSs deployed in
265 the summer 2021, are reported in Table S13. Their spatial patterns are displayed in Figures
266 1, S1, and S2. In BC, OPEs had elevated levels in the interior of Burrard Inlet close to Port
267 Moody (V1 and V2), at the southern mouth of the Fraser River (V5), and at some sites
268 around populated areas in Victoria, BC (V6-V8). TBP, TCPP, and TDCPP had higher
269 concentrations close to an industrial area near Esquimalt (V10). In QC, highest OPE water
270 concentrations were usually detected at site W5, in the Saint Lawrence River close to an
271 industrial area in Sorel-Tracy, rather than at sites in Montreal (W1 and W2) or Québec City
272 (W8 and W9). W4 also had elevated concentrations for some OPEs such as TBP. Water
273 concentrations at the one sampling site in the Saint Lawrence Estuary were much lower than
274 in the river. Overall, the spatial patterns suggest that the water concentrations of OPEs were

275 related to both industrial activities and human populations in BC, whereas industrial
 276 activities might have relatively higher impact on water concentrations of OPEs in QC.



277
 278 **Figure 1** Spatial patterns of OPEs in the water in British Columbia (left panel) and Quebec (right
 279 panel). The inserted maps at the bottom right of each panel show the sampling sites
 280 located within Victoria (left panel) and Quebec City (right panel). Concentrations in
 281 duplicate samples were averaged. The stacked bars indicate the total concentrations
 282 levels of all detected OPEs and individual OPE. Various colors are used for different
 283 OPEs. The dispersion plume of the Montreal waste water treatment plant enters the
 284 river at 45 40' N, 73 28' W and stays on the north side of the river (Marcogliese et al.,
 285 2015), therefore, the OPEs in the dispersion plume might not be sampled at W3 and
 286 W4. The concentration scale is shown to the right of the maps, which were created
 287 using the basemap of MATLAB, copyrighted to Esri, TomTom, Garmin, SafeGraph,
 288 GeoTechnologies, Inc, METI/NASA, USGS, Bureau of Land Management, EPA, NPS,
 289 US Census Bureau, USDA, USFWS, NRCan, and Parks Canada.

290 4. DISCUSSION

291 **4.1 Seasonality and Temperature Dependence.** Clear and consistent seasonal trends were
 292 not observed for particle-bound OPEs at any location during the one year of sampling
 293 (Tables S9). However, OPE gas phase concentrations at all three AAS sampling sites clearly
 294 varied seasonally, allowing us to investigate the influence of temperature on those
 295 concentrations (Figure 2). Except for EHDPP and TDCPP, concentrations of frequently
 296 detected OPEs increased with increasing ambient temperature. The logarithm of the partial
 297 pressures of OPEs ($\ln p$) were linearly regressed against the reciprocal of absolute
 298 temperatures ($1/T$) (Clausius-Clapeyron (CC) relationship), with the slopes, R^2 values, and
 299 p values summarized in Table S7. Regressions for TBP were significant at the three AAS
 300 sites ($p < 0.05$), whereas EHDPP did not show significant relationships at any site. The CC

301 relationships for other OPEs were only significant ($p < 0.05$) at some locations, i.e., TCEP
302 and TCPP on Saturna Island, TEP and TPhP in Tadoussac, and TPrP, TCPP, TCEP, and
303 TDCPP in Toronto. In cases with $R^2 > 0.10$, the trends indicate higher partial pressures at
304 higher temperatures.

305 Apparent enthalpies of air-surface exchange (ΔH_{AS-app}) were obtained from the slopes of the
306 CC relationships with $R^2 > 0.30$ and $p < 0.05$, and compared with enthalpies of exchange
307 between air and water (ΔH_{AW}) and between air and octanol (ΔH_{AO}), estimated using poly-
308 parameter linear free energy relationships (UFZ-LSER database v 3.2.1 [Internet], 2024)
309 (Table S7). Values of ΔH_{AS-app} that are similar to ΔH_{AW} and ΔH_{AO} have been interpreted as
310 being indicative of a dominant contribution of temperature-driven local air-surface
311 exchanges on the air concentration at a site (Bidleman et al., 2023; Wania et al., 1998; Zhan
312 et al., 2023). If ΔH_{AS-app} is much lower than ΔH_{AW} and ΔH_{AO} , advection from elsewhere is
313 presumed to play a more important role. ΔH_{AS-app} values of OPEs at the three sampling sites
314 were mostly within the uncertainty range of ΔH_{AW} and ΔH_{AO} . In several instances the
315 temperature dependence of air concentrations was even larger than might be expected from
316 air-surface equilibrium, i.e. ΔH_{AS-app} was larger than ΔH_{AW} and ΔH_{AO} . Examples are the
317 ΔH_{AS-app} values of TCPP on Saturna Island and in Toronto, as well as those of TBP and
318 TPhP in Tadoussac, and, to a smaller extent also TCEP on Saturna Island and in Toronto,
319 and TEP in Tadoussac.

320 This may simply be a result of high uncertainty, considering the relatively small number of
321 samples available for deriving the CC relationships for Saturna Island and Tadoussac. It
322 could also suggest that temperature influences not only the exchange between air and surface
323 but also the OPE source strength to the atmosphere. This source strength could be correlated
324 with temperature, e.g., because of enhanced release of OPEs from materials at higher
325 temperatures or higher indoor-outdoor exchange rates in summer. Furthermore, the
326 formation of TCPP, TCEP, and TPhP from precursor compounds (i.e., tris(2-
327 chloroisopropyl) phosphite (TCPPi), tris(2-chloroethyl) phosphite (TCEPi), and triphenyl
328 phosphite (TPhPi) by reaction with ozone could be higher in summer (Liu et al., 2023; Liu
329 and Mabury, 2019; Turygin et al., 2018; Zhang et al., 2021), when photooxidant
330 concentrations tend to be higher. Even though TCPP is widely used in large quantities, the
331 spatial distribution and usage of its precursor TCPPi has not been reported. The high ΔH_{AS-}
332 $_{app}$ of TEP and TBP in Tadoussac may also be related to the conversion of their
333 corresponding phosphite esters. The value of the measured ΔH_{AS-app} may potentially contain

334 information on the contribution of the transformation of OPAs to OPEs in the atmosphere,
335 i.e., the extent to which ΔH_{AS-app} exceeds ΔH_{AW} and ΔH_{AO} may indicate the extent of such
336 transformation. However, this would be beset by high uncertainties considering the complex
337 set of factors influencing the ΔH_{AS-app} .

338 Incidentally, at 11 sites in BC, where PASs were deployed at least three times during
339 different seasons with different average temperatures, higher OPEs concentrations were
340 generally also observed during warmer deployments (Tables S6 and Table S5 in the SI of Li
341 et al. submitted). 33 out of 55 CC relationships using these PAS data had $R^2 > 0.5$, and 27 of
342 these 33 were negative (Table S6). Considering the limited number of data points (3~4) for
343 PAS sites with multiple deployments in different seasons, ΔH_{AS-app} values may have high
344 uncertainties (Table S6) and were therefore not compared with theoretical values.


345

346 **Table 1** Summary of levels OPEs in precipitation reported in the literature and our study.

Region & locations	Years	Concentration, ng/L								References	Note	
		TEP	TBP	TCEP	TCPP	TDCPP	TPhP	TBEP	EHDPP			
Literature data												
Livingston Island, Antarctic	2018	2.1	1.0	3.1	26.0	1.9				0.11	(Casas et al., 2021)	
Nanning, China	N/A		4.0	15	38	2.1	1.0				(Zhang et al., 2020)	Mean
Osnabrueck, Germany	2011			187	372	46					(Mihajlović and Fries, 2012)	Median
Bahnbrücke, Germany	2001		911	121					394		(Fries and Püttmann, 2003)	
Rome, Italy	2007	46	46	155	686	404			112		(Bacaloni et al., 2008)	Mean
Martignano, Italy	2007	12	11	19	28	108			38		(Bacaloni et al., 2008)	
New York, USA	2017	17.7	3.9	5.7	61.8	11.7	11.0				(Kim and Kannan, 2018)	Mean
Our study												
Saturna Island	2020	3.0	4.0	15.6	25.7	20.9	1.0	13.7	1.2		Our study	Mean
Tadoussac	2021	1.2	0.6	2.6	5.1	37.0	0.6	2.8	0.5		Our study	Mean

347 The concentrations of OPEs in snow and rain water samples from five locations in Germany during 2007-2008 were reported (Regnery and Püttmann, 2009). However, as we could not calculate
 348 the average OPE concentrations in precipitation, we did not include these data in this table.

349 Marklund et al.(2005b) reported the concentrations of OPEs in combined dry and wet deposition samples, as there are no data for precipitation samples, therefore, these data were not included in
 350 this table either.



351

352 **Figure 2** Seasonal variability in ambient temperature (right y axis) and gas phase concentrations
 353 of OPEs (left y axis) in the atmosphere of Saturna Island, BC (2019-2020, top),
 354 Tadoussac, QC (2020-2021, middle), and Toronto, Ontario (2020-2021, bottom). Only
 355 data for OPEs with detection frequency higher than 50% are shown.

356 **4.2 Gas-Particle Partitioning.** The fraction of the OPEs in the particle phase (Φ , %) are
 357 given in Table S9. As more than 50% of Φ values for TPhP and EHDPP in Tadoussac were
 358 calculated using values $<$ MDL, these data are not discussed further. Overall, Φ ranged
 359 between 32% and 68 % and varied between OPEs and location. Among the five OPEs, the
 360 Φ of TCEP is the smallest at almost all three AAS sites. The Φ values for TBP and TCPP
 361 on Saturna Island (both ca. 66%) are 12 % higher than that for TCEP (54%). In Tadoussac,

362 the Φ for TCPP (38%) is comparable to that for TCEP (34%), whereas Φ for TBP (52%) is
363 the highest among three OPEs. In Toronto, the Φ values of the five major OPEs were in the
364 sequence of TCEP (50%) = TBP (50%) < TPhP (54%) < TCPP (56%) < EHDPP (68%).
365 This sequence is opposite to that found above the North Atlantic Ocean and in the Arctic
366 (Wu et al., 2020).

367 Theoretically, TCEP, TCPP, and TBP have very similar volatility with logarithmic
368 equilibrium partition ratios between octanol and air ($\log K_{OA}$) around 9 and $\log (K_{PA} / \text{m}^3 \text{g}^{-1})$
369 of ~ 1 at 15 °C estimated using the UFZ-LSER website (UFZ-LSER database v 3.2.1, 2024)
370 (Table S9). These three chemicals are expected to be largely in the gas phase at ambient
371 temperatures. TPhP and EHDPP have estimated $\log K_{OA}$ values > 12 and $\log (K_{PA} / \text{m}^3 \text{g}^{-1})$
372 of ~ 5 at 15 °C which would indicate strong particle sorption in the atmosphere. However,
373 the unexpectedly low fraction observed in the particle phase may suggest that TPhP and
374 EHDPP are emitted at higher temperatures and are not in a state of equilibrium between gas
375 and particle phase (Zhao et al., 2021a). Alternatively, the fraction of TPhP and EHDPP in
376 the gas phase may have been overestimated if very fine and ultrafine particles containing
377 these OPEs passed through the glass fiber filters (Zhao et al., 2021b). While it has been
378 suggested that the composition of the particles (Li et al., 2017b), relative humidity (Li et al.,
379 2017b; Wu et al., 2020), and degradation of OPEs in gas and particle phases may also
380 influence the gas-particle partitioning of OPEs, we do not have the empirical data to explore
381 the influence of these factors on our measurements.

382 The calculated Φ at the three AAS sites increases with decreasing ambient temperatures.
383 This is consistent with lower temperatures favoring partitioning to particles (Table S9). This
384 is also reflected in the positive linear relationships between the $\ln K_{PA}$ and reciprocal
385 temperature (in K) in Tadoussac, Saturna Island, and Toronto (Table S10).


386 **4.3 Scavenging Ratios.** Measured SRs could be calculated for eight OPEs and ranged
387 mainly from 10^4 to 10^7 (Table S12). These SRs are highly uncertain because of the
388 uncertainty in the measured concentrations and because we combine a monthly precipitation
389 sample with a 24-hour air sample taken during the same month. The estimated SRs are also
390 uncertain due to the possibly high uncertainty in the estimated K_{WA} and the assumptions
391 regarding equilibrium partitioning of OPE vapors between air and water droplets and the
392 value and constancy of W_P . Despite these uncertainties, estimated SRs for TBP and EHDPP
393 are generally around 10^5 and therefore comparable to the measured ones, which indicates
394 that equilibrium between precipitation and these chemicals in the atmosphere was achieved.

395 The estimated SRs for other OPEs are mostly within the range of 2×10^6 to 10^9 and therefore
396 orders of magnitude higher than the measured SRs. At very high values, exceeding a
397 threshold of $\sim 10^6$, the SR concept loses its usefulness, because the atmosphere will
398 essentially be cleansed of such compounds at the onset of a precipitation event and
399 subsequent precipitation will simply dilute the concentrations (Lei and Wania, 2004). As
400 such, measured SRs that are smaller than these very high estimated ones are not too
401 surprising.

402 **4.4 Diffusive Air-Water Gas Exchange.** The water-air equilibrium status was evaluated
403 using fugacity ratios (f_w/f_A), whereby f_w/f_A values lower (higher) than 1 indicate a tendency
404 for net deposition (volatilization). The estimated fugacity ratios for five OPEs (TBP, TCEP,
405 TCPP, TPhP, and EHDPP) are given in Table S14. This estimation of f_w/f_A incurs substantial
406 uncertainty because of uncertainty in K_{AW} and the passive sampling rates, and because it
407 involves combining air and water data obtained during different time periods (Oh et al., 2023;
408 Zhan et al., 2023). Nevertheless, the f_w/f_A values in BC and QC were so far below unity, that
409 one can confidently assert that all five OPEs were net deposited from atmosphere to water.
410 Ma et al. (2021) also reported that almost all five OPEs, except TBP, underwent net gas
411 phase deposition in the Lower Great Lakes Region.

412 **4.5 Relative Abundance of OPEs in Different Environmental Media.** The frequent
413 detection of TCPP, TCEP, TBP, and TPhP in PASs, the gas and particle phase of the AAS, ,
414 PWSs, and PCPNs in QC and BC allows us to investigate the relative abundance of these
415 OPEs in different environmental media (Figure 3). Chlorinated compounds (TCPP and
416 TCEP) were dominant in all environmental media regardless of sampling locations, which
417 is consistent with observations in gaseous and aqueous phases in the Great Lakes region (Ma
418 et al., 2021). Specifically, TCPP was the most abundant of the four OPEs in all types of
419 samples, except for the gas phase in Tadoussac and the PWS. By reporting the relative
420 abundance of the OPEs in PASs separately for industrial, urban, and rural sites, we find a
421 consistent pattern in both QC and BC, namely that the relative abundance of halogenated
422 OPEs (TCPP and TCEP) decreased from industrial (65% in QC and 79% in BC) to urban
423 (59% in QC and 63% in BC) to rural sites (52% in QC and 53% in BC) with a concomitant
424 increase of two nonhalogenated OPEs (TBP and TPhP). This is consistent with previous
425 studies (Kurt-Karakus et al., 2018; Zhang et al., 2019), but contrasts with the predominance
426 of TCPP and TCEP reported for Antarctic air (Wang et al., 2020a). The higher abundance
427 of TPhP at rural sites would be consistent with a relatively higher long-range transport

428 potential (LRTP) estimated with the improved OECD Pov and LRTP Screening Tool
 429 (OECD Tool) (Breivik et al., 2022) (Table S15). Even though the observed higher
 430 abundance of TBP in remote areas is inconsistent with its relatively low estimated LRTP,
 431 Sühning et al. (2020) indicated that non-chlorinated OPEs could be subject to LRTP.



432
 433 **Figure 3** The relative abundance of four frequently detected OPEs in passive air samples (PAS),
 434 gas phase active air samples (AAS-G), glass fiber filter samples (GFF), precipitation
 435 samples (PCPN), and passive water samples (PWS) in QC and BC. Due to the large
 436 concentration ranges, geomean was used for calculating the relative abundance for
 437 PASs

438 The relative abundance of OPEs in the gas and particle phase at the three AAS sites was
 439 similar. Mirroring the observation in the PASs, the relative contribution of chlorinated OPEs
 440 was higher in urban and industrial Toronto than in rural Saturnia and remote Tadoussac.
 441 Chen et al. (2019) and Salamova et al. (2013) reported a similar trend for OPEs in dust and
 442 atmospheric particles, respectively. Interestingly, in the atmospheric gas and particle phases,
 443 the higher abundance for TPhP in Tadoussac compared to Toronto suggests that long-range
 444 transport was dominant, despite Toronto being a highly populated city. This observation
 445 aligns with the above findings of a low ΔH_{AS-app} , the low median concentration level in
 446 Toronto compared to that in Tadoussac (Table S5), as well as the spatial patterns, cluster

447 analyses, and a weak linear relationship with population previously reported in Li et al.
448 (submitted). Collectively, these pieces of evidence indicate that TPhP is more closely related
449 to industrial activities and subject to long-range transport.

450 The abundance of chlorinated OPEs in PCPN and PWS was much greater than those of the
451 nonhalogenated OPEs, which is consistent with the dominance of TCPP and TCEP in the
452 dissolved phase of water sampled from urban and rural watersheds in Toronto (Awonaike et
453 al., 2021). Based on section 4.4, net diffusion from the atmosphere to water occurred in BC
454 and QC, therefore, the high abundance of chlorinated OPEs can be explained by their
455 relatively high K_{WA} (Tables S4 & S15).

456 **5. IMPLICATIONS**

457 Some observations made here are conforming with general expectations regarding the
458 environmental behaviour of semi-volatile organic chemicals, such as higher gas phase
459 concentrations and a decrease in the particle bound fraction at higher ambient temperatures.
460 Also, the measured precipitation scavenging ratios, while high, can be reconciled with
461 equilibrium partitioning ratios of gaseous OPEs that favour aqueous phases over the gas
462 phase. Other observations are more puzzling, such as the general lack of a clear relationship
463 between OPE volatility and the observed gas-particle partitioning behaviour. Furthermore,
464 the strong temperature dependence of OPE gas phase concentrations that indicates the
465 importance of temperature-driven local air-surface exchange processes is not entirely
466 consistent with the low air/water fugacity ratios which suggest that gaseous air-water
467 exchange is strongly depositional. One possible explanation is that the measured seasonal
468 concentration variability is less a reflection of temperature driven air-surface exchange and
469 instead indicates that more OPE enter, or are formed in, the atmosphere in summer. Potential
470 mechanisms are (i) an increased release of OPEs at higher temperatures from outdoor
471 materials to which they have been added (Kemmlein et al. 2003), (ii) a faster ventilation of
472 OPE emitted indoors (Stamp et al., 2022, Han et al., 2024), (iii) more active industrial
473 activities, such as construction, using products containing OPEs in the summer months, and
474 (iv) the atmospheric oxidation of organophosphite precursors (Liu et al., 2023).

475 Our data also highlight that the understanding of the atmospheric dispersion potential of
476 OPEs is still incomplete. While a relatively high long range transport potential for aryl-OPE
477 (TPhP) is consistent with the results from the OECD Tool (Kung et al., 2022; Sühling et al.,
478 2020), the higher or constant relative abundance for TBP at remote sites does not align with

479 predictions, which indicate a limited LRTP for TBP. This, too, may be related to the
480 unpredictable gas-particle partitioning behaviour of the OPEs and the role of gas and multi-
481 phase transformation processes, e.g. the possibility that TBP originates in part from the
482 transformation of precursors. More research is needed to better understand the atmospheric
483 gas-particle partitioning behaviour of the OPEs and to elucidate the role that transformation
484 reactions may play in this regard.

485 **Code and data availability**

486 All data generated for this project are contained in the Supplement.

487 **Supplement**

488 The supplement related to this article is available online at: xxxxxx.

489 **Author contributions**

490 YL, FZ, and JO prepared and extracted the PASs and the Toronto AASs. YL and FZ also
491 took the Toronto AAS. YDL prepared standards. CS prepared, obtained, and extracted
492 samples from Saturna Island and Tadoussac as well as the PWSs and analyzed the particle
493 samples. KL and FAPCG deployed and retrieved PASs and PWSs in British Columbia. ABC,
494 ZL, HH, FZ, and FW deployed/retrieved PASs and PWSs in Quebec. YL compiled and
495 interpreted data. YL wrote the manuscript under the guidance of FW with input by the other
496 co-authors. HH coordinated the project. All authors reviewed the manuscript.

497 **Competing interests**

498 The contact author has declared that none of the authors has any competing interests.

499 **Acknowledgements**

500 We thank Geri Crooks, Alexandre Costa, Yannick Lapointe, Louis-Georges Esquilat,
501 Jocelyn Praud, Sandrine Vigneron, François Gagnon, Jonathan Pritchard, Alessia Colussi,
502 Nicolas Alexandrou, Abigaëlle Dalpé Castilloux, Christian Boutot, Bruno Cayouette, Fella
503 Moualek, Frédéric Bélanger, Claude Lapierre, Félix Ledoux, Samuel Turgeon, Sarah
504 Duquette and the CAPMON team for their assistance in deploying samplers and providing
505 facilities/permissions to the sampling locations.

506 **Financial support**

507 This research has been supported by Environment and Climate Change Canada under the
508 Whale Initiative 1.0 (grants no. GCXE20S008, GCXE20S010, GCXE20S011), and a

509 Connaught scholarship to Yuening Li.

510 **References**

- 511 Awonaike, B., Lei, Y. D., and Wania, F.: Precipitation-induced transport and phase
512 partitioning of organophosphate esters (OPEs) in urban and rural watersheds, *Environ. Sci.*
513 *Water Res. Technol.*, 7, 2106–2115, <https://doi.org/10.1039/D1EW00329A>, 2021.
- 514 Bacaloni, A., Cucci, F., Guarino, C., Nazzari, M., Samperi, R., and Laganà, A.: Occurrence
515 of organophosphorus flame retardant and plasticizers in three volcanic lakes of Central Italy,
516 *Environ. Sci. Technol.*, 42, 1898–1903, <https://doi.org/10.1021/es702549g>, 2008.
- 517 Bidleman, T., Andersson, A., Brorström-Lundén, E., Brugel, S., Ericson, L., Hansson, K.,
518 and Tysklind, M.: Halomethoxybenzenes in air of the Nordic region, *Environ. Sci.*
519 *Ecotechnology*, 13, 1–7, <https://doi.org/10.1016/j.ese.2022.100209>, 2023.
- 520 Booiij, K. and Smedes, F.: An Improved Method for Estimating in Situ Sampling Rates of
521 Nonpolar Passive Samplers, *Environ. Sci. Technol.*, 44, 6789–6794,
522 <https://doi.org/10.1021/es101321v>, 2010.
- 523 Booiij, K., Hofmans, H. E., Fischer, C. V., and Van Weerlee, E. M.: Temperature-Dependent
524 Uptake Rates of Nonpolar Organic Compounds by Semipermeable Membrane Devices and
525 Low-Density Polyethylene Membranes, *Environ. Sci. Technol.*, 37, 361–366,
526 <https://doi.org/10.1021/es025739i>, 2003.
- 527 Breivik, K., McLachlan, M. S., and Wania, F.: The Emissions Fractions Approach to
528 Assessing the Long-Range Transport Potential of Organic Chemicals, *Environ. Sci.*
529 *Technol.*, 56, 11983–11990, <https://doi.org/10.1021/acs.est.2c03047>, 2022.
- 530 Casas, G., Martinez-Varela, A., Vila-Costa, M., Jiménez, B., and Dachs, J.: Rain
531 Amplification of Persistent Organic Pollutants, *Environ. Sci. Technol.*, 55, 12961–12972,
532 <https://doi.org/10.1021/acs.est.1c03295>, 2021.
- 533 Castro-Jiménez, J., González-Gaya, B., Pizarro, M., Casal, P., Pizarro-Álvarez, C., and
534 Dachs, J.: Organophosphate ester flame retardants and plasticizers in the global oceanic
535 atmosphere, *Environ. Sci. Technol.*, 50, 12831–12839,
536 <https://doi.org/10.1021/acs.est.6b04344>, 2016.
- 537 Chen, Y., Zhang, Q., Luo, T., Xing, L., and Xu, H.: Occurrence, distribution and health risk
538 assessment of organophosphate esters in outdoor dust in Nanjing, China: Urban vs. rural
539 areas, *Chemosphere*, 231, 41–50, <https://doi.org/10.1016/j.chemosphere.2019.05.135>, 2019.
- 540 Choo, G. and Oh, J.-E.: Seasonal occurrence and removal of organophosphate esters in
541 conventional and advanced drinking water treatment plants, *Water Res.*, 186, 116359,
542 <https://doi.org/10.1016/j.watres.2020.116359>, 2020.
- 543 Desimoni, E. and Brunetti, B.: About Estimating the Limit of Detection by the Signal to
544 Noise Approach, *Pharm. Anal. Acta*, 6, 1–4, <https://doi.org/10.4172/2153-2435.1000355>,
545 2015.
- 546 Ding, J., Shen, X., Liu, W., Covaci, A., and Yang, F.: Occurrence and risk assessment of
547 organophosphate esters in drinking water from Eastern China, *Sci. Total Environ.*, 538, 959–
548 965, 2015.
- 549 Draft screening assessment flame retardants group: [https://www.canada.ca/en/environment-](https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-substances/draft-screening-assessment-flame-retardants-group.html#toc7)
550 [climate-change/services/evaluating-existing-substances/draft-screening-assessment-flame-](https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-substances/draft-screening-assessment-flame-retardants-group.html#toc7)
551 [retardants-group.html#toc7](https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-substances/draft-screening-assessment-flame-retardants-group.html#toc7), last access: 17 November 2023.

552 Risk management scope for TPHP, BDPD, BDMEPPP, IDDP, IPPP and TEP:
553 [https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-
555 substances/risk-management-scope-tphp-bdpd-bdmeppp-iddp-ipp-ipp-tep.html](https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-
554 substances/risk-management-scope-tphp-bdpd-bdmeppp-iddp-ipp-ipp-tep.html), last access: 17
November 2023.

556 Screening assessment alkyl aryl phosphites: [https://www.canada.ca/en/environment-
climate-change/services/evaluating-existing-substances/screening-assessment-alkyl-aryl-
phosphites.html](https://www.canada.ca/en/environment-
557 climate-change/services/evaluating-existing-substances/screening-assessment-alkyl-aryl-
558 phosphites.html), last access: 17 November 2023.

559 Updated draft screening assessment - Certain organic flame retardants substance grouping -
560 TCPP and TDCPP: [https://www.canada.ca/en/environment-climate-
563 change/services/evaluating-existing-substances/updated-draft-screening-assessment-
organic-flame-retardants-substance-grouping-tcpp-tdcpp.html](https://www.canada.ca/en/environment-climate-
561 change/services/evaluating-existing-substances/updated-draft-screening-assessment-
562 organic-flame-retardants-substance-grouping-tcpp-tdcpp.html), last access: 17 November
2023.

564 Fries, E. and Püttmann, W.: Monitoring of the three organophosphate esters TBP, TCEP and
565 TBEP in river water and ground water (Oder, Germany), *J. Environ. Monit.*, 5, 346–352,
566 <https://doi.org/10.1039/b210342g>, 2003.

567 Gu, J., Su, F., Hong, P., Zhang, Q., and Zhao, M.: 1H NMR-based metabolomic analysis of
568 nine organophosphate flame retardants metabolic disturbance in Hep G2 cell line, *Sci. Total
569 Environ.*, 665, 162–170, <https://doi.org/10.1016/j.scitotenv.2019.02.055>, 2019.

570 Han, X., Hao, Y., Li, Y., Yang, R., Wang, P., Zhang, G., Zhang, Q., and Jiang, G.:
571 Occurrence and distribution of organophosphate esters in the air and soils of Ny-Ålesund
572 and London Island, Svalbard, Arctic, *Environ. Pollut.*, 263, 114495,
573 <https://doi.org/10.1016/j.envpol.2020.114495>, 2020.

574 He, M. J., Lu, J. F., and Wei, S. Q.: Organophosphate esters in biota, water, and air from an
575 agricultural area of Chongqing, western China: Concentrations, composition profiles,
576 partition and human exposure, *Environ. Pollut.*, 244, 388–397,
577 <https://doi.org/10.1016/j.envpol.2018.10.085>, 2019.

578 Kim, H.-K., Shin, Y.-S., Lee, D.-S., Song, B.-J., and Kim, J.-G.: Estimation of Rain
579 Scavenging Ratio for Particle Bound Polycyclic Aromatic Hydrocarbons and
580 Polychlorinated Biphenyls, <https://doi.org/10.4491/eer.2006.11.1.033>, 2006.

581 Kim, U. J. and Kannan, K.: Occurrence and Distribution of Organophosphate Flame
582 Retardants/Plasticizers in Surface Waters, Tap Water, and Rainwater: Implications for
583 Human Exposure, *Environ. Sci. Technol.*, 52, 5625–5633,
584 <https://doi.org/10.1021/acs.est.8b00727>, 2018.

585 Kung, H. C., Hsieh, Y. K., Huang, B. W., Cheruiyot, N. K., and Chang-Chien, G. P.: An
586 Overview: Organophosphate Flame Retardants in the Atmosphere, *Aerosol Air Qual. Res.*,
587 22, <https://doi.org/10.4209/aaqr.220148>, 2022.

588 Kurt-Karakus, P., Alegria, H., Birgul, A., Gungormus, E., and Jantunen, L.:
589 Organophosphate ester (OPEs) flame retardants and plasticizers in air and soil from a highly
590 industrialized city in Turkey, *Sci. Total Environ.*, 625, 555–565,
591 <https://doi.org/10.1016/j.scitotenv.2017.12.307>, 2018.

592 Lei, Y. D. and Wania, F.: Is rain or snow a more efficient scavenger of organic chemicals?,
593 *Atmos. Environ.*, 38, 3557–3571, <https://doi.org/10.1016/j.atmosenv.2004.03.039>, 2004.

594 Li, C., Chen, J., Xie, H. Bin, Zhao, Y., Xia, D., Xu, T., Li, X., and Qiao, X.: Effects of
595 Atmospheric Water on ·OH-initiated Oxidation of Organophosphate Flame Retardants: A
596 DFT Investigation on TCPP, *Environ. Sci. Technol.*, 51, 5043–5051,

597 <https://doi.org/10.1021/acs.est.7b00347>, 2017a.

598 Li, J., Xie, Z., Mi, W., Lai, S., Tian, C., Emeis, K. C., and Ebinghaus, R.: Organophosphate
599 Esters in Air, Snow, and Seawater in the North Atlantic and the Arctic, *Environ. Sci.*
600 *Technol.*, 51, 6887–6896, <https://doi.org/10.1021/acs.est.7b01289>, 2017b.

601 Li, J., Tang, J., Mi, W., Tian, C., Emeis, K. C., Ebinghaus, R., and Xie, Z.: Spatial
602 Distribution and Seasonal Variation of Organophosphate Esters in Air above the Bohai and
603 Yellow Seas, China, *Environ. Sci. Technol.*, 52, 89–97,
604 <https://doi.org/10.1021/acs.est.7b03807>, 2018.

605 Li, J., Zhao, L., Letcher, R. J., Zhang, Y., Jian, K., Zhang, J., and Su, G.: A review on
606 organophosphate Ester (OPE) flame retardants and plasticizers in foodstuffs: Levels,
607 distribution, human dietary exposure, and future directions, *Environ. Int.*, 127, 35–51,
608 <https://doi.org/10.1016/j.envint.2019.03.009>, 2019a.

609 Li, J., Cao, H., Mu, Y., Qu, G., Zhang, A., Fu, J., and Jiang, G.: Structure-Oriented Research
610 on the Antiestrogenic Effect of Organophosphate Esters and the Potential Mechanism,
611 *Environ. Sci. Technol.*, 54, 14525–14534, <https://doi.org/10.1021/acs.est.0c04376>, 2020.

612 Li, W., Wang, Y., and Kannan, K.: Occurrence, distribution and human exposure to 20
613 organophosphate esters in air, soil, pine needles, river water, and dust samples collected
614 around an airport in New York state, United States, *Environ. Int.*, 131, 105054,
615 <https://doi.org/10.1016/j.envint.2019.105054>, 2019b.

616 Li, Y., Zhan, F., Lei, Y. D., Shunthirasingham, C., Hung, H., and Wania, F.: Field calibration
617 and PAS-SIM model evaluation of the XAD-based passive air samplers for semi-volatile
618 organic compounds, *Environ. Sci. Technol.*, 57, 9224–9233,
619 <https://doi.org/10.1021/acs.est.3c00809>, 2023a.

620 Li, Y., Zhan, F., Shunthirasingham, C., Lei, Y. D., Hung, H., and Wania, F.: Unbiased
621 Passive Sampling of All Polychlorinated Biphenyls Congeners from Air, *Environ. Sci.*
622 *Technol. Lett.*, 10, 565–572, <https://doi.org/10.1021/acs.estlett.3c00271>, 2023b.

623 Li, Y., Zhan, F., Su, Y., Lei, Y. D., Shunthirasingham, C., Zhou, Z., Abbatt, J. P. D., Hung,
624 H., and Wania, F.: Uptake behavior of polycyclic aromatic compounds during field
625 calibrations of the XAD-based passive air sampler across seasons and locations, *Atmos.*
626 *Meas. Tech.*, 17, 715–729, <https://doi.org/10.5194/amt-17-715-2024>, 2024.

627 Li, Y., Zhan, F., Shunthirasingham, C., Lei, Y. D., Oh, J., Weng, C., Chaaben, A. Ben, Lu,
628 Z., Lee, K., Gobas, F. A. P. C., Hung, H., and Wania, F.: Inferring Atmospheric Sources of
629 Gaseous Organophosphate Esters from Spatial Patterns, submitted, n.d.

630 Liu, Q., Liu, R., Zhang, X., Li, W., Harner, T., Saini, A., Liu, H., Yue, F., Zeng, L., Zhu, Y.,
631 Xing, C., Li, L., Lee, P., Tong, S., Wang, W., Ge, M., Wang, J., Wu, X., Johannessen, C.,
632 Liggio, J., Li, S. M., Hung, H., Xie, Z., Mabury, S. A., and Abbatt, J. P. D.: Oxidation of
633 commercial antioxidants is driving increasing atmospheric abundance of organophosphate
634 esters: Implication for global regulation, *One Earth*, 6, 1202–1212,
635 <https://doi.org/10.1016/j.oneear.2023.08.004>, 2023.

636 Liu, R. and Mabury, S. A.: Organophosphite Antioxidants in Indoor Dust Represent an
637 Indirect Source of Organophosphate Esters, *Environ. Sci. Technol.*, 53, 1805–1811,
638 <https://doi.org/10.1021/acs.est.8b05545>, 2019.

639 Lu, Z., Martin, P. A., Burgess, N. M., Champoux, L., Elliott, J. E., Baressi, E., De Silva, A.
640 O., de Solla, S. R., and Letcher, R. J.: Volatile Methylsiloxanes and Organophosphate Esters
641 in the Eggs of European Starlings (*Sturnus vulgaris*) and Congeneric Gull Species from

642 Locations across Canada, *Environ. Sci. Technol.*, 51, 9836–9845,
643 <https://doi.org/10.1021/acs.est.7b03192>, 2017.

644 Ma, Y., Cui, K., Zeng, F., Wen, J., Liu, H., Zhu, F., Ouyang, G., Luan, T., and Zeng, Z.:
645 Microwave-assisted extraction combined with gel permeation chromatography and silica gel
646 cleanup followed by gas chromatography-mass spectrometry for the determination of
647 organophosphorus flame retardants and plasticizers in biological samples, *Anal. Chim. Acta*,
648 786, 47–53, <https://doi.org/10.1016/j.aca.2013.04.062>, 2013.

649 Ma, Y., Vojta, S., Becanova, J., Habtemichael, A. Z., Adelman, D. A., Muir, D., and
650 Lohmann, R.: Spatial distribution and air–water exchange of organophosphate esters in the
651 lower Great Lakes, *Environ. Pollut.*, 286, 117349,
652 <https://doi.org/10.1016/j.envpol.2021.117349>, 2021.

653 Ma, Y., Luo, Y., Zhu, J., Zhang, J., Gao, G., Mi, W., Xie, Z., and Lohmann, R.: Seasonal
654 variation and deposition of atmospheric organophosphate esters in the coastal region of
655 Shanghai, China, *Environ. Pollut.*, 300, 118930,
656 <https://doi.org/10.1016/j.envpol.2022.118930>, 2022.

657 Marcogliese, D. J., Blaise, C., Cyr, D., de Lafontaine, Y., Fournier, M., Gagné, F., Gagnon,
658 C., and Hudon, C.: Effects of a major municipal effluent on the St. Lawrence River: A case
659 study, *Ambio*, 44, 257–274, <https://doi.org/10.1007/s13280-014-0577-9>, 2015.

660 Marklund, A., Andersson, B., and Haglund, P.: Organophosphorus flame retardants and
661 plasticizers in Swedish sewage treatment plants, *Environ. Sci. Technol.*, 39, 7423–7429,
662 <https://doi.org/10.1021/es051013l>, 2005a.

663 Marklund, A., Andersson, B., and Haglund, P.: Traffic as a source of organophosphorus
664 flame retardants and plasticizers in snow, *Environ. Sci. Technol.*, 39, 3555–3562,
665 <https://doi.org/10.1021/es0482177>, 2005b.

666 McDonough, C. A., De Silva, A. O., Sun, C., Cabrerizo, A., Adelman, D., Soltwedel, T.,
667 Bauerfeind, E., Muir, D. C. G., and Lohmann, R.: Dissolved organophosphate esters and
668 polybrominated diphenyl ethers in remote marine environments: Arctic surface water
669 distributions and net transport through Fram Strait, *Environ. Sci. Technol.*, 52, 6208–6216,
670 <https://doi.org/10.1021/acs.est.8b01127>, 2018.

671 Mi, L., Xie, Z., Zhang, L., Waniek, J. J., Pohlmann, T., Mi, W., and Xu, W.:
672 Organophosphate Esters in Air and Seawater of the South China Sea: Spatial Distribution,
673 Transport, and Air–Sea Exchange, *Environ. Heal.*, 1, 191–202,
674 <https://doi.org/10.1021/envhealth.3c00059>, 2023.

675 Mihajlović, I. and Fries, E.: Atmospheric deposition of chlorinated organophosphate flame
676 retardants (OFR) onto soils, *Atmos. Environ.*, 56, 177–183,
677 <https://doi.org/10.1016/j.atmosenv.2012.03.054>, 2012.

678 Möller, A., Xie, Z., Caba, A., Sturm, R., and Ebinghaus, R.: Organophosphorus flame
679 retardants and plasticizers in the atmosphere of the North Sea, *Environ. Pollut.*, 159, 3660–
680 3665, <https://doi.org/10.1016/j.envpol.2011.07.022>, 2011.

681 Möller, A., Sturm, R., Xie, Z., Cai, M., He, J., and Ebinghaus, R.: Organophosphorus flame
682 retardants and plasticizers in airborne particles over the Northern Pacific and Indian Ocean
683 toward the polar regions: Evidence for global occurrence, *Environ. Sci. Technol.*, 46, 3127–
684 3134, <https://doi.org/10.1021/es204272v>, 2012.

685 Na, G., Hou, C., Li, R., Shi, Y., Gao, H., Jin, S., Gao, Y., Jiao, L., and Cai, Y.: Occurrence,
686 distribution, air-seawater exchange and atmospheric deposition of organophosphate esters

687 (OPEs) from the Northwestern Pacific to the Arctic Ocean, *Mar. Pollut. Bull.*, 157, 111243,
688 <https://doi.org/10.1016/j.marpolbul.2020.111243>, 2020.

689 Oh, J., Shunthirasingham, C., Lei, Y. D., Zhan, F., Li, Y., Dalpé Castilloux, A., Ben Chaaben,
690 A., Lu, Z., Lee, K., Gobas, F. A. P. C., Eckhardt, S., Alexandrou, N., Hung, H., and Wania,
691 F.: The atmospheric fate of 1,2-dibromo-4-(1,2-dibromoethyl)cyclohexane (TBECH):
692 spatial patterns, seasonal variability, and deposition to Canadian coastal regions, *Atmos.*
693 *Chem. Phys.*, 23, 10191–10205, <https://doi.org/10.5194/acp-23-10191-2023>, 2023.

694 Regnery, J. and Püttmann, W.: Organophosphorus Flame Retardants and Plasticizers in Rain
695 and Snow from Middle Germany, *CLEAN – Soil, Air, Water*, 37, 334–342,
696 <https://doi.org/10.1002/clen.200900050>, 2009.

697 Regnery, J. and Püttmann, W.: Occurrence and fate of organophosphorus flame retardants
698 and plasticizers in urban and remote surface waters in Germany, *Water Res.*, 44, 4097–4104,
699 <https://doi.org/10.1016/j.watres.2010.05.024>, 2010.

700 Rosenmai, A. K., Winge, S. B., Möller, M., Lundqvist, J., Wedebye, E. B., Nikolov, N. G.,
701 Lilith Johansson, H. K., and Vinggaard, A. M.: Organophosphate ester flame retardants have
702 antiandrogenic potential and affect other endocrine related endpoints in vitro and in silico,
703 *Chemosphere*, 263, 127703, <https://doi.org/10.1016/j.chemosphere.2020.127703>, 2021.

704 Salamova, A., Ma, Y., Venier, M., and Hites, R. A.: High Levels of Organophosphate Flame
705 Retardants in the Great Lakes Atmosphere, *Environ. Sci. Technol. Lett.*, 1, 8–14,
706 <https://doi.org/10.1021/ez400034n>, 2013.

707 Salamova, A., Hermanson, M. H., and Hites, R. A.: Organophosphate and Halogenated
708 Flame Retardants in Atmospheric Particles from a European Arctic Site, *Environ. Sci.*
709 *Technol.*, 48, 6133–6140, <https://doi.org/10.1021/es500911d>, 2014.

710 Salamova, A., Peverly, A. A., Venier, M., and Hites, R. A.: Spatial and temporal trends of
711 particle phase organophosphate ester concentrations in the atmosphere of the great lakes,
712 *Environ. Sci. Technol.*, 50, 13249–13255, <https://doi.org/10.1021/acs.est.6b04789>, 2016.

713 Shi, T., Li, R., Fu, J., Hou, C., Gao, H., Cheng, G., Zhang, H., Jin, S., Kong, L., and Na, G.:
714 Fate of organophosphate esters from the Northwestern Pacific to the Southern Ocean:
715 Occurrence, distribution, and fugacity model simulation, *J. Environ. Sci.*, 137, 347–357,
716 <https://doi.org/10.1016/j.jes.2023.03.001>, 2024.

717 Shi, Y., Zhang, Y., Du, Y., Kong, D., Wu, Q., Hong, Y., Wang, Y., Tam, N. F. Y., and
718 Leung, J. Y. S.: Occurrence, composition and biological risk of organophosphate esters
719 (OPEs) in water of the Pearl River Estuary, South China, *Environ. Sci. Pollut. Res.*, 27,
720 14852–14862, <https://doi.org/10.1007/s11356-020-08001-1>, 2020.

721 Shoeib, M., Ahrens, L., Jantunen, L., and Harner, T.: Concentrations in air of organobromine,
722 organochlorine and organophosphate flame retardants in Toronto, Canada, *Atmos. Environ.*,
723 99, 140–147, <https://doi.org/10.1016/j.atmosenv.2014.09.040>, 2014.

724 Stackelberg, P. E., Gibs, J., Furlong, E. T., Meyer, M. T., Zaugg, S. D., and Lippincott, R.
725 L.: Efficiency of conventional drinking-water-treatment processes in removal of
726 pharmaceuticals and other organic compounds, *Sci. Total Environ.*, 377, 255–272,
727 <https://doi.org/10.1016/j.scitotenv.2007.01.095>, 2007.

728 Storey, J. M. E., Luo, W., Isabelle, L. M., and Pankow, J. F.: Gas/Solid Partitioning of
729 Semivolatile Organic Compounds to Model Atmospheric Solid Surfaces as a Function of
730 Relative Humidity. 1. Clean Quartz, *Environ. Sci. Technol.*, 29, 2420–2428,
731 <https://doi.org/10.1021/es00009a039>, 1995.

732 Sührling, R., Diamond, M. L., Scheringer, M., Wong, F., Pućko, M., Stern, G., Burt, A.,
733 Hung, H., Fellin, P., Li, H., and Jantunen, L. M.: Organophosphate esters in Canadian Arctic
734 air: Occurrence, levels and trends, *Environ. Sci. Technol.*, 50, 7409–7415,
735 <https://doi.org/10.1021/acs.est.6b00365>, 2016.

736 Sührling, R., Scheringer, M., Rodgers, T. F. M. M., Jantunen, L. M., and Diamond, M. L.:
737 Evaluation of the OECD P OV and LRTP screening tool for estimating the long-range
738 transport of organophosphate esters, *Environ. Sci. Process. Impacts*, 22, 207–216,
739 <https://doi.org/10.1039/c9em00410f>, 2020.

740 Sundkvist, A. M., Olofsson, U., and Haglund, P.: Organophosphorus flame retardants and
741 plasticizers in marine and fresh water biota and in human milk, *J. Environ. Monit.*, 12, 943–
742 951, <https://doi.org/10.1039/B921910B>, 2010.

743 Turygin, V. V., Sokhadze, L. A., Golubeva, Y. Y., Platonova, L. V., Afanas'eva, A. A.,
744 Nazarenko, D. I., and Shvetsova-Shilovskaya, T. N.: New Approach to Obtain Neutral Ester
745 of Phosphoric Acid: Tris(2-chloroisopropyl) Phosphate, *Theor. Found. Chem. Eng.*, 52,
746 643–647, <https://doi.org/10.1134/S0040579518040280>, 2018.

747 UFZ-LSER database v 3.2.1 [Internet]: <http://www.ufz.de/lserd>, last access: 13 January
748 2024.

749 van der Veen, I. and de Boer, J.: Phosphorus flame retardants: Properties, production,
750 environmental occurrence, toxicity and analysis, *Chemosphere*, 88, 1119–1153,
751 <https://doi.org/10.1016/j.chemosphere.2012.03.067>, 2012.

752 Wang, C., Wang, P., Zhao, J., Fu, M., Zhang, L., Li, Y., Yang, R., Zhu, Y., Fu, J., Zhang,
753 Q., and Jiang, G.: Atmospheric organophosphate esters in the Western Antarctic Peninsula
754 over 2014–2018: Occurrence, temporal trend and source implication, *Environ. Pollut.*, 267,
755 115428, <https://doi.org/10.1016/j.envpol.2020.115428>, 2020a.

756 Wang, X., Zhu, Q., Yan, X., Wang, Y., Liao, C., and Jiang, G.: A review of organophosphate
757 flame retardants and plasticizers in the environment: Analysis, occurrence and risk
758 assessment, *Sci. Total Environ.*, 731, 139071,
759 <https://doi.org/10.1016/j.scitotenv.2020.139071>, 2020b.

760 Wang, X., Luu, T., Beal, M. A., Barton-Maclaren, T. S., Robaire, B., and Hales, B. F.: The
761 Effects of Organophosphate Esters Used as Flame Retardants and Plasticizers on Granulosa,
762 Leydig, and Spermatogonial Cells Analyzed Using High-Content Imaging, *Toxicol. Sci.*,
763 186, 269–287, <https://doi.org/10.1093/toxsci/kfac012>, 2022.

764 Wania, F., Haugen, J. E., Lei, Y. D., and Mackay, D.: Temperature dependence of
765 atmospheric concentrations of semivolatile organic compounds, *Environ. Sci. Technol.*, 32,
766 1013–1021, <https://doi.org/10.1021/es970856c>, 1998.

767 Wong, F., de Wit, C. A., and Newton, S. R.: Concentrations and variability of
768 organophosphate esters, halogenated flame retardants, and polybrominated diphenyl ethers
769 in indoor and outdoor air in Stockholm, Sweden, *Environ. Pollut.*, 240, 514–522,
770 <https://doi.org/10.1016/j.envpol.2018.04.086>, 2018.

771 Wu, Y., Venier, M., and Salamova, A.: Spatioseasonal Variations and Partitioning Behavior
772 of Organophosphate Esters in the Great Lakes Atmosphere, *Environ. Sci. Technol.*, 54,
773 5400–5408, <https://doi.org/10.1021/acs.est.9b07755>, 2020.

774 Xie, Z., Wang, P., Wang, X., Castro-Jiménez, J., Kallenborn, R., Liao, C., Mi, W., Lohmann,
775 R., Vila-Costa, M., and Dachs, J.: Organophosphate ester pollution in the oceans, *Nat. Rev.*
776 *Earth Environ.*, 3, 309–322, <https://doi.org/10.1038/s43017-022-00277-w>, 2022.

777 Yan, H. and Hales, B. F.: Effects of Organophosphate Ester Flame Retardants on
778 Endochondral Ossification in Ex Vivo Murine Limb Bud Cultures, *Toxicol. Sci.*, 168, 420–
779 429, <https://doi.org/10.1093/toxsci/kfy301>, 2019.

780 Yan, H. and Hales, B. F.: Exposure to tert-Butylphenyl Diphenyl Phosphate, an
781 Organophosphate Ester Flame Retardant and Plasticizer, Alters Hedgehog Signaling in
782 Murine Limb Bud Cultures, *Toxicol. Sci.*, 178, 251–263,
783 <https://doi.org/10.1093/toxsci/kfaa145>, 2020.

784 Zhan, F., Shunthirasingham, C., Li, Y., Oh, J., Lei, Y. D., Ben Chaaben, A., Dalpé Castilloux,
785 A., Lu, Z., Lee, K., Gobas, F. A. P. C., Alexandrou, N., Hung, H., and Wania, F.: Sources
786 and environmental fate of halomethoxybenzenes, *Sci. Adv.*, 9, eadi8082,
787 <https://doi.org/10.1126/sciadv.adi8082>, 2023.

788 Zhang, Q., Li, X., Wang, Y., Zhang, C., Cheng, Z., Zhao, L., Li, X., Sun, Z., Zhang, J., Yao,
789 Y., Wang, L., Li, W., and Sun, H.: Occurrence of novel organophosphate esters derived from
790 organophosphite antioxidants in an e-waste dismantling area: Associations between hand
791 wipes and dust, *Environ. Int.*, 157, 106860, <https://doi.org/10.1016/j.envint.2021.106860>,
792 2021.

793 Zhang, W., Wang, P., Li, Y., Wang, D., Matsiko, J., Yang, R., Sun, H., Hao, Y., Zhang, Q.,
794 and Jiang, G.: Spatial and temporal distribution of organophosphate esters in the atmosphere
795 of the Beijing-Tianjin-Hebei region, China, *Environ. Pollut.*, 244, 182–189,
796 <https://doi.org/10.1016/j.envpol.2018.09.131>, 2019.

797 Zhang, X., Sühring, R., Serodio, D., Bonnell, M., Sundin, N., and Diamond, M. L.: Novel
798 flame retardants: Estimating the physical-chemical properties and environmental fate of 94
799 halogenated and organophosphate PBDE replacements, *Chemosphere*, 144, 2401–2407,
800 <https://doi.org/10.1016/j.chemosphere.2015.11.017>, 2016.

801 Zhang, Z., Lin, G., Lin, T., Zhang, R., Jin, L., and Di, Y.: Occurrence, behavior, and fate of
802 organophosphate esters (OPEs) in subtropical paddy field environment: A case study in
803 Nanning City of South China, *Environ. Pollut.*, 267, 115675,
804 <https://doi.org/10.1016/j.envpol.2020.115675>, 2020.

805 Zhao, F., Riipinen, I., and MacLeod, M.: Steady-State Mass Balance Model for Predicting
806 Particle–Gas Concentration Ratios of PBDEs, *Environ. Sci. Technol.*, 55, 9425–9433,
807 <https://doi.org/10.1021/acs.est.0c04368>, 2021a.

808 Zhao, S., Tian, L., Zou, Z., Liu, X., Zhong, G., Mo, Y., Wang, Y., Tian, Y., Li, J., Guo, H.,
809 and Zhang, G.: Probing Legacy and Alternative Flame Retardants in the Air of Chinese
810 Cities, *Environ. Sci. Technol.*, 55, 9450–9459, <https://doi.org/10.1021/acs.est.0c07367>,
811 2021b.

812