

Anthropogenic emission controls reduce summertime ozone-temperature sensitivity in the United States

Shuai Li^{1,2,3}, ~~Xiao Lu^{1,2,3,*}~~, Haolin Wang^{1,2,3}, ~~Xiao Lu^{1,2,3,*}~~

¹ School of Atmospheric Sciences, Sun Yat-sen University, and Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, Guangdong 519082, China

² Guangdong Provincial Observation and Research Station for Climate Environment and Air Quality Change in the Pearl River Estuary, Zhuhai, Guangdong 519082, China

³ Key Laboratory of Tropical Atmosphere-Ocean System, Ministry of Education, Zhuhai, China, Zhuhai, Guangdong 519082, China

Correspondence to Xiao Lu (luxiao25@mail.sysu.edu.cn)

Abstract

The ozone-temperature sensitivity is widely used to ~~predict~~infer the impact of future climate warming on ozone. However, trends in ozone-temperature sensitivity and possible drivers remained unclear. Here, we show that the observed summertime surface ozone-temperature sensitivity, defined as the slope of the best-fit line of daily anomaly in ozone versus maximum temperature ($m_{\Delta O_3-\Delta T_{max}}$), has decreased by 50% during 1990-2021 in the continental United States (CONUS), with a mean decreasing rate of -0.57 ppbv/K/decade ($p < 0.01$) across 608 monitoring sites. We conduct high-resolution GEOS-Chem simulations in 1995-2017 to interpret the $m_{\Delta O_3-\Delta T_{max}}$ trends and underlying mechanisms in the CONUS. The simulations identify the dominant role of anthropogenic nitrogen oxides (NO_x) emission reduction in the observed $m_{\Delta O_3-\Delta T_{max}}$ decrease. We find that approximately 76% of the simulated decline in $m_{\Delta O_3-\Delta T_{max}}$ can be attributed to the temperature-indirect effects arising from the shared collinearity of other meteorological effects (such as humidity, ventilation, and transport) on ozone. The remaining portion (24%) is mostly due to the temperature-direct effects, in particular four explicit temperature-dependent processes, including the biogenic volatile organic compounds (BVOCs) emissions, soil NO_x emissions, dry deposition, and the thermal decomposition of peroxyacetyl nitrate (PAN). With reduced anthropogenic NO_x emissions, the expected ozone enhancement from temperature-driven BVOCs emissions, dry deposition, and PAN decomposition decreases, contributing to the decline in $m_{\Delta O_3-\Delta T_{max}}$. However, soil NO_x emissions increase $m_{\Delta O_3-\Delta T_{max}}$ with anthropogenic NO_x emission reduction, indicating an increasing role of soil NO_x emissions in shaping the ozone-temperature sensitivity. As indicated by the decreased $m_{\Delta O_3-\Delta T_{max}}$, model simulations estimate that reduced anthropogenic NO_x emissions from 1995 to 2017 have lowered ozone enhancement from low to high temperatures by 6.8 ppbv averaged over the CONUS, significantly reducing the risk of extreme ozone pollution events under high temperatures. Our study illustrates the dependency of ozone-temperature sensitivity on anthropogenic emission levels that should be considered in the future ozone mitigation in a warmer climate.

1. Introduction

Surface ozone harms human health and causes loss of crop yields (Feng et al., 2022; Mills et al., 2018; Monks et al., 2015; Turner et al., 2016); [Wang et al., 2024](#)). It is chemically generated from its precursors including nitrogen oxides (NO_x), volatile organic compounds (VOCs), and carbon monoxide (CO) in the presence of sunlight. The natural sources, chemical kinetics, deposition, and transport of ozone and its precursors are significantly influenced by meteorology and climate (Fiore et al., 2012; Fu and Tian, 2019; Jacob and Winner, 2009; Lu et al., 2019b), shaping the strong sensitivity of surface ozone concentration to meteorological parameters such as temperature. Quantification of ozone-meteorology sensitivity provides a useful tool for predicting daily variation of ozone and for understanding climate-chemistry interactions, yet how anthropogenic emission levels may affect the sensitivity remains unclear. Here, we examine whether long-term anthropogenic control of ozone precursors has changed the response of summertime ozone to daily variations in temperature in the United States (US) and the underlying mechanisms.

High temperature is expected to increase ozone concentrations in polluted environment, through boosting biogenic VOCs (BVOCs) and soil NO_x emissions, accelerating photochemical kinetics of ozone formation, and suppressing ozone dry deposition (Hudman et al., 2012; Lin et al., 2020; Porter and Heald, 2019; Pusede et al., 2015; Romer et al., 2018; Varotsos et al., 2019). In addition, temperature-dependent meteorological parameters, such as solar radiation and humidity, and temperature-related meteorological effects, such as air stagnancy, enhanced ventilation, and regional transport, can also influence surface ozone level (Kerr et al., 2019; Lu et al., 2019b; Porter and Heald, 2019; Zhang et al., 2022a). Such effects can be reflected in but at the same time complicate the ozone-temperature relationship. Still, temperature is often used as a proxy to synthesize the effects of meteorology and climate on ozone. Previous studies have documented a robust positive ozone-temperature sensitivity in NO_x-rich environment, typically defined as the slope of the best-fit line for ozone and temperature ($d[O_3]/dT$), of 2-8 ppbv/K across the US, Europe, and China (Bloomer et al., 2009; Gu et al., 2020; Ning et al., 2022; Pusede et al., 2014; Sillman and Samson, 1995; Varotsos et al., 2019). The positive $d[O_3]/dT$ also indicates an ozone climate change penalty, *i.e.* future warming may deteriorate ozone air quality in the absence of changes in anthropogenic emission activities (Zhang et al., 2022b). The climate penalty requires additional anthropogenic emission reductions to offset the ozone increase in a warmer climate (Wu et al., 2008); [Zanis et al., 2022](#)).


While the overall positive ozone-temperature relationship is well recognized, how ozone-temperature sensitivity has changed remains much less explored. Some studies have reported the weakening of regional ozone-temperature sensitivity in California, the Midwestern US, and the eastern US based on observations, and supposed reduction in local anthropogenic emissions as a possible driver (Bloomer et al., 2009; [Hembeck et al., 2022](#); Jing et al., 2017; Rasmussen et al., 2013). In contrast, Fu et al. (2015) reports large interannual variations in ozone-temperature sensitivity in the southeast US that may be tied to climate variability. Model simulations project a decrease in ozone-temperature sensitivity in future scenarios with lower anthropogenic emissions in the US (Nolte et al., 2021). These studies indicate that the surface ozone-temperature sensitivity has been shifting with significant regional variations in the US, yet an up-to-date view on the long-term and continental-scale

65 trends is currently missing. In particular, the quantitative assessment of underlying mechanisms driving long-term changes in surface ozone-temperature sensitivity remains rather unclear, limiting the application of this important metric in predicting future ozone evolution.

70 In this study, we analyze the present-day (2017-2021) and long-term trends (1990-2021) in the summertime surface ozone-temperature sensitivity relationship in the continental US (CONUS), combining observational monitoring network and state-of-art chemical modeling. We utilize the GEOS-Chem chemical transport model to quantify the role of anthropogenic emission reduction in the long-term trends in the ozone-temperature sensitivity, and investigate the underlying mechanisms. We also examine the benefit of reduced ozone-temperature sensitivity in ozone mitigation during high temperatures that frequently cause severe ozone pollution extremes.

2. Materials and Methods

2.1 Surface ozone measurement in the US



75

Figure 1. Locations of the 319 urban sites (crosses) and 289 rural sites (dots) across the continental US used in this study. Sites are categorized in seven regions, including Northwestern US (NWUS), Southwestern US (SWUS), Northeastern US (NEUS), Southeastern US (SEUS), Midwestern US (Midwest), the mountainous western US (Intermountain West), and Central Plains of the US (Plains). The underlying figure shows terrain elevation.

80

We obtain hourly measurements of surface ozone concentration from the US Environmental Pollution Agency (EPA) Air Quality System (AQS) data program. Our study period covers 1990-2021, in total of 32 years, with a focus on boreal

summertime (June, July, August). We derive the daily maximum 8-hour average (MDA8) ozone concentrations from the hourly data, and select sites with valid summertime ozone measurements for at least 24 years (i.e. $\geq 75\%$) in the 1990-2021 period and for at least 3 years in 2017-2021 (Text S1). A total of 608 sites are selected, including 319 urban sites and 289 rural sites (based on EPA categorization). We follow previous studies to categorize the sites into seven geographic areas (Nolte et al., 2021; Rasmussen et al., 2012), including the Northwestern US (NWUS), Southwestern US (SWUS), Northeastern US (NEUS), Southeastern US (SEUS), Midwestern US (Midwest), the mountainous western US (Intermountain West), and Central Plains of the US (Plains) (Figure 1).

2.2 Temperature data

The AQS dataset also provides measurements of temperature that could be ideally used in quantifying the ozone-temperature relationship at individual sites. However, the temperature measurement is largely missing, with only 170 sites (<30% of the total 608 sites selected for analysis) providing long-term (at least 24 years) records, which is insufficient to support our analysis. Here we derive the gridded ($0.5^\circ \times 0.625^\circ$) data of temperature at 2 meters above the ground from the Modern-Era Retrospective analysis for Research and Applications (Version 2, MERRA-2) dataset (Gelaro et al., 2017), which consistently serves as input for the GEOS-Chem chemical transport model (Section 2.4). We align the gridded temperature data with in-situ ozone measurement based on the coordinates of individual sites. Evaluation of the MERRA-2 gridded data with in-situ measurements of temperature at available sites shows an excellent agreement between the two with a mean bias (MB) of 0.3-1.0 K and the correlation coefficient (r) of 0.96-0.98 for years after 2000, however, the two datasets have slightly larger disparities in the earliest part of our study period (e.g. MB=0.4 K, $r=0.87$ for year 1990) (Figure S1). We also compare temperature trends from MERRA-2 with observations over the period 1990-2021 (Table S1). While the overall trends are consistent, there are notable overestimation (e.g. NEUS, Plains) and underestimation (e.g. SEUS and SWUS) in different regions, which may lead to biases in interpreting the observed ozone-temperature sensitivity (as observed ozone variation responds to “true” air temperature).

2.3 Definition of ozone-temperature sensitivity

Our goal is to examine the response of summertime MDA8 ozone concentration to the variation in daily maximum temperature (T_{\max}) across the US, and the trends in such response from 1990 to 2021. We use T_{\max} instead of daytime temperature or mean temperature as strong correlation coefficients between MDA8 ozone and T_{\max} have been revealed in previous studies (e.g. Steiner et al., 2010; Fu et al., 2015). Ozone levels in the US have experienced significant decreasing trends since 1980s due to anthropogenic emission control measures (Gaudel et al., 2018; Kim et al., 2006; Lin et al., 2017; Simon et al., 2015). The higher ozone concentration in earlier years may obfuscate the long-term trends in ozone-temperature sensitivity, if the ozone-temperature sensitivity were derived by the raw measurements. Therefore, we first subtract the monthly-mean MDA8 ozone concentration and T_{\max} from each daily record to derive their daily anomaly (ΔO_3 and ΔT_{\max}) at

individual sites for each year. This process allows us to remove the seasonal (monthly) influences and also the 1990-2021 trends in ozone concentration and temperature. We then define the summertime ozone-temperature sensitivity ($m_{\Delta O_3-\Delta T_{max}}$) at individual sites as the slope of the best-fit line of daily ΔO_3 versus ΔT_{max} . Fu et al. (2015) also applied similar process to quantify ozone-temperature sensitivity across the southeast US. We calculate the mean values of $m_{\Delta O_3-\Delta T_{max}}$ over the sites across the contiguous US (CONUS) or individual regions to represent the regional-mean ozone-temperature relationship. Trends in $m_{\Delta O_3-\Delta T_{max}}$ over each site are estimated using the linear regression method, with a 5-year smoothing average applied to the yearly $m_{\Delta O_3-\Delta T_{max}}$ to filter the interannual variability. The trends of the mean $m_{\Delta O_3-\Delta T_{max}}$ values across the sites are used to represent regional mean trends in the ozone-temperature sensitivity.

2.4 GEOS-Chem model simulation

We use the GEOS-Chem version 11-02-rc chemical transport model (Bey et al., 2001) to interpret summertime ozone-temperature sensitivity and its trend in the United States. The GEOS-Chem model is driven by MERRA-2 assimilated meteorological data. We conduct simulations over the North America nested-grid domain (140°-40° W, 10°-70° N) at the horizontal resolution of 0.5°(latitude) × 0.625°(longitude), ~~with boundary conditions archived from consistent). The global simulations at 2° × 2.5° resolution-~~ providing the boundary conditions were configured consistently with the nested simulations (simulation time, chemical schemes, emission inventory, etc., see discussions below). GEOS-Chem model describes a state-of-art ozone-NO_x-VOCs-aerosol-halogen tropospheric chemistry scheme, and also includes online calculation of emissions, dry and wet depositions of gases and aerosols. Anthropogenic emissions in this study are from the Community Emissions Data System (CEDS v-2021-04-21), in which the interannual variability in the US emissions are scaled based on the US National Emissions Inventory (US NEI) (McDuffie et al., 2020). The CEDS inventory indicates a significant decrease in anthropogenic NO_x, NMVOCs, CO emissions over the CONUS of 62.5, 70.8, 48.0% respectively from 1995-2017.

GEOS-Chem is capable ~~for~~ of simulating the temperature's influences on ozone through chemical kinetics, natural emissions, transport, and dry deposition. Chemical kinetics in GEOS-Chem are modularized based on the Jet Propulsion Laboratory (JPL) and International Union of Pure and Applied Chemistry (IUPAC) scheme (IUPAC, 2011; Sander et al, 2013), with temperature input from the hourly MERRA-2 reanalysis data. GEOS-Chem also includes online calculation of temperature-dependent natural emissions. Biogenic emissions are parameterized following The Model of Emissions of Gases and Aerosols from Nature (MEGAN version v2.1) algorithm (Guenther et al., 2012), in which biogenic emissions are calculated based on temperature, solar radiation, leaf area index (LAI), and other parameters. Biogenic emissions increase exponentially with temperature, but emissions of some BVOCs are inhibited at higher temperatures. Soil NO_x emissions are calculated based on nitrogen availability in soil, edaphic conditions such as soil temperature and moisture, and other gridded parameters such as vegetation type using the Berkeley-Dalhousie Soil NO_x Parameterization ~~as described in Hudman et al. (2012). According to the BDSNP scheme, soil emission is an exponential function of temperature when the soil temperature is between 0 and 30 °C and is constant when the soil temperature exceeds 30 °C. Some studies have shown continuous increases~~

of soil NO_x emissions at temperature higher than 30 °C over (BDSNP) as described in Hudman et al. (2012). ~~California (Oikawa et al., 2015; Wang et al., 2021), but here we do not implement this scheme in our simulation due to the lack of supporting emissions in other regions of the CONUS.~~ Lightning NO_x emissions are parameterized based on cloud-top heights with the spatial distribution of flash rates constrained by satellite observations (Murray et al., 2012). Biomass burning emissions are from the BB4CMIP (biomass burning emissions for CMIP6) inventory (van Marle et al., 2017), in which emissions after 1997 are consistent with the Global Fire Emissions Database version 4 (GFED4) inventory (van der Werf et al., 2017). However, temperature's impacts on anthropogenic NO_x and VOCs emissions (Liu et al., 2024; Wu et al., 2024) are not considered in our simulation.

Dry deposition of both gas and aerosols is calculated online based on the resistance-in-series algorithm (Wesely, 1989). Surface temperature influences deposition velocity through a stomatal resistance term, which remains low within normal temperatures (e.g. 10-30 °C) but rises at two extremes (below 0 °C and above 40 °C) (Porter and Heald, 2019), contributing to local ozone increases at high temperatures. Wet deposition for water-soluble aerosols and ~~gasgases~~ in GEOS-Chem is described by Liu et al. (2001) and Amos et al. (2012), but it has small direct effects on ozone due to the low solubility of both NO_x and ozone.

Model simulations are summarized in Table 1. We ~~conducted~~conduct a BASE simulation for July for every two years from 1995 to 2017, with one-month simulation (June) as model spin-up: for both the global and regional simulations. The initial chemical fields are closed to conditions for July 2005. The one-month spin-up time can be considered sufficient in this case as the ozone in the urban boundary layer typically has a lifetime ranging from hours to days. However, it may be short for ozone in the free troposphere where ozone has a lifetime of orders of weeks (Monks et al., 2015). To demonstrate this, we conducted an additional set of experiments, starting with a global simulation at 2°×2.5° resolution from 1st January 2017 to 1st August 2017. The global simulation on 1st June 2017 was then interpolated into the high-resolution nested grid to drive the high-resolution simulation from 1st June 2017 to 1st August 2017. A comparison of surface MDA8 ozone concentrations and ozone-temperature sensitivity between the two sets of simulations is shown in Figure S2. We find that the differences between the simulations with 1-month and 6-month spin-up times had only minor impacts on ozone concentrations and $m_{\Delta O_3-\Delta T_{max}}$. The average differences between the two simulations were only 2.3% for ozone concentrations and 0.3% for $m_{\Delta O_3-\Delta T_{max}}$, with high spatial consistency ($r > 0.99$). This confirms that using a 1-month spin-up time for the simulation should not affect the analysis and conclusions. However, a longer spin-up time is favorable for generating global chemical fields when sufficient computational resources are available. The BASE simulation applies the yearly-varied anthropogenic emissions and includes all the abovementioned temperature-dependent mechanisms. We then conduct two simulations for the same 1995-2017 period, but in which the domestic anthropogenic NO_x (1995E) or VOCs (1995EAVOCs) emissions in US are fixed to 1995 level.

We conducted 14 additional sets of sensitivity experiments to explore the role of different mechanisms in the ozone-temperature sensitivity and its trend. First, we separate the effect of temperature on ozone through direct and indirect effects. Here, the temperature-direct effect is defined as the effect directly parameterized with temperature in GEOS-Chem, including natural emissions of BVOCs and soil NO_x, the chemical kinetics, dry deposition, and others. In comparison, temperature-

180 indirect effect is defined as the effect not directly parameterized with temperature but is also influenced or reflected by
temperature, for example humidity, radiation, and transport. The simulation strategy is to remove the daily variation of
temperature (while keeping the diurnal cycle) and its influence on ozone daily variations. For this purpose, we generate the
mean diurnal cycle of temperature averages over all 31 days in July 2017 at each grid cell. We then feed this normalized
185 effect of temperature on ozone. Comparison of the FTEMP simulation and BASE simulation yields a quantitative assessment
of the direct effect.

For the temperature-direct effect, we further follow Porter and Heald (2019) to explore the role of four temperature-
dependent mechanisms on the ozone-temperature sensitivity. These four mechanisms are BVOCs emissions, soil NO_x
emissions, thermal decomposition of peroxyacetyl nitrate (PAN, whose decomposition is strongly correlated to temperature),
190 and dry deposition. We feed the normalized temperature data (remove daily variation but keep diurnal cycle) into the
calculations of all or each of the four temperature-dependent mechanisms in the GEOS-Chem. For the temperature-indirect
effect, we additionally examine the role of transport in the ozone-temperature sensitivity. This is done by generating a
meteorological field that retains only the daily variation of three-dimensional wind field and boundary layer height (PBLH)
and removes the daily variation of all other meteorological elements, which is used into the GEOS-Chem (~~FTRANS~~-TRANS).
195 Interpretation of these sensitivity simulations are summarized in Table S2.

We conduct the above simulations ~~withat~~ both ~~inthe~~ 2017 and 1995 emission level, allowing us to explore the role of
these mechanisms in the changes in ozone-temperature sensitivity with anthropogenic NO_x emissions reduction, which has not
been addressed in previous modeling studies. Except for the BASE, 1995E, and 1995E-AVOCs simulation, other simulations
are only conducted for year 2017 (the latest year with available anthropogenic emission inventory when the simulations were
200 conducted) as sensitivity tests.

Table 1 Configurations of Model Simulations

Cases	Simulation time	Description
BASE	July,1995-2017 (biennially)	Default simulation with yearly-varied anthropogenic emissions and all temperature-dependent mechanisms
1995E	Same as BASE	Same as BASE, but anthropogenic NO _x emissions fixed in 1995
1995EAVOCs	Same as BASE	Same as BASE, but anthropogenic VOCs emissions fixed in 1995
BASE-FTEMP	July, 2017	Same as BASE, but with normalized temperature field in the model
1995E-FTEMP	July, 2017	Same as 1995E, but with normalized temperature field in the model

BASE- <u>FTRANSTRANS</u>	July, 2017	Same as BASE, but with normalized all meteorological elements except three-dimensional wind field and PBLH
1995E- <u>FTRANSTRANS</u>	July, 2017	Same as 1995E, but with normalized all meteorological elements except three-dimensional wind field and PBLH
BASE- <u>FALLF4PATHS</u>	July, 2017	Same as BASE, but remove four mechanisms temperature dependence by normalized temperature for BVOCs emissions, Soil NO _x emissions, PAN decomposition, and dry deposition
1995E- <u>FALLF4PATHS</u>	July, 2017	Same as 1995E, but remove four mechanisms temperature dependence by normalized temperature for BVOCs emissions, Soil NO _x emissions, PAN decomposition, and dry deposition
BASE-FBVOC	July, 2017	Same as BASE, but remove BVOCs temperature dependence by normalized temperature for biogenic VOC emissions
1995E -FBVOC	July, 2017	Same as 1995E, but remove BVOCs temperature dependence by normalized temperature for biogenic VOC emissions
BASE-FSNO_x	July, 2017	Same as BASE, but remove Soil NO _x temperature dependence by normalized temperature for soil NO _x emissions
1995E-FSNO_x	July, 2017	Same as 1995E, but remove Soil NO _x temperature dependence by normalized temperature for soil NO _x emissions
BASE-FPAN	July, 2017	Same as BASE, but remove PAN temperature dependence by normalized temperature for PAN decomposition
1995E-FPAN	July, 2017	Same as 1995E, but remove PAN temperature dependence by normalized temperature for PAN decomposition
BASE-FDEP	July, 2017	Same as BASE, but remove dry deposition temperature dependence by normalized temperature for dry deposition
1995E-FDEP	July, 2017	Same as 1995E, but remove dry deposition temperature dependence by normalized temperature for dry deposition

3. Results

3.1 Present-day level and trends of summertime ozone-temperature sensitivity in the continental US





Figure 2. Present-day summertime ozone concentrations and ozone-temperature sensitivity in the continental US. (a,b) Distribution of summertime (June, July, August) $r_{\Delta O_3-\Delta T_{max}}$ and $m_{\Delta O_3-\Delta T_{max}}$ at individual sites averaged in 2017-2021. Black borders indicate sites with a p-value<0.01 for $r_{\Delta O_3-\Delta T_{max}}$ or $m_{\Delta O_3-\Delta T_{max}}$. (c) Summer mean MDA8 ozone concentrations at individual sites. Urban sites are represented by circle and rural sites by square. Mean values and standard deviations over the sites are shown in the inset.

Figure 2(a) presents the widespread positive correlation coefficients between summertime daily MDA8 ozone and T_{max} ($r_{\Delta O_3-\Delta T_{max}}$) across the CONUS sites. 604 out of the total 608 sites are showing positive $r_{\Delta O_3-\Delta T_{max}}$ (570/608 sites with p-value<0.01) in the present-day (2017-2021), with a mean $r_{\Delta O_3-\Delta T_{max}}$ value of 0.41±0.17 (mean ± standard deviation across the sites) averaged over all sites. Urban sites show slightly higher $r_{\Delta O_3-\Delta T_{max}}$ values than rural sites. Figure 2(b) shows that the present-day mean $m_{\Delta O_3-\Delta T_{max}}$ (see Section 2.3 for the definition) values averaged for the 608 sites are 1.52±0.76 ppbv/K, with the $m_{\Delta O_3-\Delta T_{max}}$ values at urban sites higher by 18% than those averaged for the rural sites (1.64±0.78 versus 1.39±0.72 ppbv/K). These results reflect the expected ozone increases with temperature in NO_x -rich environment, which are more commonly found in urban than rural areas.

We find distinct variability in the spatial distributions of both $r_{\Delta O_3-\Delta T_{max}}$ and $m_{\Delta O_3-\Delta T_{max}}$ (Figure 2, Table S13). The Midwest and NEUS regions show the highest mean $m_{\Delta O_3-\Delta T_{max}}$ values reaching 2.07±0.64 ($r_{\Delta O_3-\Delta T_{max}}$ =0.50±0.12) and

1.9699±0.65 ppbv/K ($r_{\Delta O_3-\Delta T_{max}}=0.52\pm 0.09$), followed by NWUS with mean $m_{\Delta O_3-\Delta T_{max}}$ of 1.54±0.3938 ppbv/K ($r_{\Delta O_3-\Delta T_{max}}=0.63\pm 0.0807$). The Intermountain West and Plains region show the lowest mean $m_{\Delta O_3-\Delta T_{max}}$ of less than 1.1 ppbv/K in both urban and rural sites with mean $r_{\Delta O_3-\Delta T_{max}}$ lower than 0.2526, indicating daily ozone variation in this region is not strongly affected by temperature. We also find that the spatial distribution of ozone-temperature sensitivity does not follow that of the MDA8 ozone level (Figure 2c), as the highest summertime MDA8 ozone concentrations are over the SWUS and Intermountain West Regions. The higher $m_{\Delta O_3-\Delta T_{max}}$ in the NEUS and Midwest regions than in other regions may reflect the stronger daily variation of ozone due to rapid shift of synoptic patterns (e.g. mid-latitude cyclones) in this region during summer (Leibensperger et al., 2008). Additionally, changes in other mid-latitude dynamic systems, such as meridional movement by the mid-latitude jet, also play a significant role in shaping the regional ozone-temperature sensitivity (Barnes and Fiore, 2013; Kerr et al., 2020; Zhang et al., 2022c). We observe a decreasing gradient in both $r_{\Delta O_3-\Delta T_{max}}$ and $m_{\Delta O_3-\Delta T_{max}}$ from north to south in the eastern United States, which aligns with previous findings (Camalier et al., 2007; Tawfik and Steiner, 2013). This observed north-to-south shift may be related to the transition in land-atmosphere coupling mechanisms due to soil moisture limitations in the southern regions (Tawfik and Steiner, 2013). The low $m_{\Delta O_3-\Delta T_{max}}$ in the Intermountain region largely reflects the strong background ozone influences (including stratospheric intrusion, long-range transport of wildfire or anthropogenic plumes) instead of local photochemical production (Jaffe et al., 2018; Zhang et al., 2014). These background sources may contribute to high ozone there but are not directly modulated by local temperature.

Previous studies report a decrease in ozone concentration at extreme high temperature over the US (Shen et al., 2016; Steiner et al., 2010). Here we investigate how the suppression of ozone concentration influences the overall ozone-temperature sensitivity. We identify occurrences of ozone suppression and the critical temperature (i.e. beyond which ozone increases are suppressed) at individual sites every year following the criteria described in Ning et al (2022). We find that while ozone suppression at extreme high temperature can be detected at 477 out of 608 sites in 2017-2021, excluding data above the critical temperature only changes the present-day mean $m_{\Delta O_3-\Delta T_{max}}$ by 2.6%. It indicates that such phenomenon does not change the overall positive ozone-temperature sensitivity.







Figure 3. Observed decrease in summertime ozone-temperature sensitivity in 1990-2021. (a) Time series of the summertime $m_{\Delta O_3-\Delta T_{max}}$ averaged over the CONUS sites, with a 5-year smoothing average applied to the yearly $m_{\Delta O_3-\Delta T_{max}}$ to filter the interannual variability. $m_{\Delta O_3-\Delta T_{max}}$ for urban and rural sites are shown in red and blue lines, respectively. Shaded areas represent the range of mean values $\pm 30\%$ of standard deviation across the sites. The CONUS $m_{\Delta O_3-\Delta T_{max}}$ trends are shown inset. Anthropogenic NO_x emissions in the CONUS are shown in grey line. (b) Spatial distributions of long-term trends in $m_{\Delta O_3-\Delta T_{max}}$ in 1990-2021 across the US. Only sites with $r_{\Delta O_3-\Delta T_{max}}$ p-values < 0.01 with $r_{\Delta O_3-\Delta T_{max}}$ are shown. Both directions and colors of the vectors indicate the $m_{\Delta O_3-\Delta T_{max}}$ trends.

The present-day (2017-2021) ozone-temperature sensitivity is lower than the reported values for earlier years (i.e. 2-7 ppbv/K reported in 2000 (Bloomer et al., 2009; Fu et al., 2015; Rasmussen et al., 2013), though different definitions of ozone-sensitivity sensitivity were applied), suggesting that the ozone-temperature sensitivity may have experienced significant reduction in recent decades. Figure 3 illustrates this feature from long-term observations in 1990-2021. We find in Figure 3a that mean $m_{\Delta O_3-\Delta T_{max}}$ for the CONUS decreased by 50% from 3.0 ppbv/K in 1990 to 1.5 ppbv/K in 2021 with a mean decreasing rate of -0.57 ppbv/K/decade ($p < 0.01$). $m_{\Delta O_3-\Delta T_{max}}$ over the CONUS urban sites was higher than rural sites by 0.50 ppbv/K in the early 1990s. However, urban sites exhibit a faster decline rate of $m_{\Delta O_3-\Delta T_{max}}$ (-0.61 ppbv/K/decade, $p < 0.01$) compared to rural (-0.53 ppbv/K/decade, $p < 0.01$), narrowing the disparity in $m_{\Delta O_3-\Delta T_{max}}$ between the two. At the same time, the mean $r_{\Delta O_3-\Delta T_{max}}$ decreased from 0.51 in 1990 to 0.40 in 2021 (Figure S2S3). The significant decrease in both $m_{\Delta O_3-\Delta T_{max}}$ and $r_{\Delta O_3-\Delta T_{max}}$ all imply a much weaker response of ozone to temperature in present-day compared to that in three decades ago. While some studies have shown observed decreases in some regions (e.g. California as described in Steiner et al. (2010)), such significant decreases of ozone-temperature sensitivity over the CONUS have not been presented in previous studies to the best of our knowledge.

The decreasing trends in $m_{\Delta O_3-\Delta T_{max}}$ are widespread across the CONUS sites (Figure 3b), but spatial and temporal variabilities exist. 411419 sites (6869%) out of the total 608 sites are showing negative trends with $p < 0.01$ (481492 sites with $p < 0.05$). The largest decreases are in the NEUS region, where $m_{\Delta O_3-\Delta T_{max}}$ values exceeded 4.3 ppbv/K in the 1990s but have

steadily decreased by -0.8381 ppbv/K/decade, reaching 1.78 ppbv/K in 2021. The SWUS region also shows a large decrease in $m_{\Delta O_3-\Delta T_{max}}$ by -0.60 ppbv/K/decade ($p < 0.01$). A distinct feature in the SWUS is the notably high urban-rural disparity in $m_{\Delta O_3-\Delta T_{max}}$ (4.7 versus 1.9 ppbv/K) in the early 1990s (Figure S3S4), but this disparity has been significantly reduced as urban sites exhibit a much larger $m_{\Delta O_3-\Delta T_{max}}$ trend (-0.88 ppbv/K/decade, $p < 0.01$) than rural sites (-0.34 ppbv/K/decade, $p < 0.01$), particularly in early 1990s. The SEUS and Midwest regions also show decreases in $m_{\Delta O_3-\Delta T_{max}}$ with a mean rate of -0.6462 and -0.52 ppbv/K. However, we notice an increase of $m_{\Delta O_3-\Delta T_{max}}$ in 1990-2000 for the SEUS region and in ~~1990~~1999-2005 for the Plains region (Figure S3-S4). The increase in ozone-temperature sensitivity in these two regions explains the $m_{\Delta O_3-\Delta T_{max}}$ plateau in CONUS during the 1996-2004 period. Fu et al. (2015) attributes the increase ozone-temperature sensitivity in 1990-2000 in the SEUS to variations in regional ozone advection tied to climate variability. This further underscores the significant influence of climate variability on $m_{\Delta O_3-\Delta T_{max}}$ trends. The region with the ~~least~~smallest $m_{\Delta O_3-\Delta T_{max}}$ trends is the Intermountain region (-0.08 ppbv/K/decade).

3.2 Simulated long-term trends in ozone-temperature sensitivity and attribution to anthropogenic emission reduction



Comparison of observed and simulated $m_{\Delta O_3-\Delta T_{max}}$, 1995-2017






Figure 4. Evaluation of GEOS-Chem simulated $m_{\Delta O_3-\Delta T_{max}}$ in July, 1995-2017. (a) Spatial distributions of the observed (circles) and simulated (from the BASE simulation, shaded) $m_{\Delta O_3-\Delta T_{max}}$ during July averaged over 1995 to 2017. (b) Scatterplots of the observed and simulated $m_{\Delta O_3-\Delta T_{max}}$ for July in each simulated year from 1995 to 2017. Mean values, standard deviations for the CONUS sites from the observation and GEOS-Chem model, and their correlation coefficients (r) in different regions are shown in the inset.

We now apply the GEOS-Chem chemical transport model to interpret the trends in ozone-temperature sensitivity over the CONUS. Figure S5S6 compares the spatial distribution of observed and simulated mean surface MDA8 ozone concentrations in July at the 608 sites averaged for 12 years (1995-2017 biennially). Our GEOS-Chem simulation captures the spatial distributions of surface MDA8 ozone across the CONUS, although showing some high bias of MDA8 ozone of 11 ppbv, as also reported in other surface ozone air quality studies using the GEOS-Chem model (Lu et al., 2019a; Travis and Jacob, 2019). Most importantly, the model largely reproduces the spatial pattern of observed $m_{\Delta O_3-\Delta T_{max}}$, with a high correlation coefficient of 0.6867 and a small positive mean bias of 0.4911 ppbv/K (4.7%) at the 608 sites for the monthly $m_{\Delta O_3-\Delta T_{max}}$ values at all sites (Figure 4). Table S2S4 further shows the simulated and observed $m_{\Delta O_3-\Delta T_{max}}$ and their correlation coefficients (r) across different periods and regions. The model demonstrates relatively better performance of $m_{\Delta O_3-\Delta T_{max}}$ across the CONUS in 2001-2011 compared to other periods, with small mean absolute bias (0.0301-0.18 ppbv/K, 1%-8%) and high correlation coefficients (0.67-0.7470). The simulated $m_{\Delta O_3-\Delta T_{max}}$ in the Eastern United States (NEUS, SEUS, Midwest, and Plains) is in better agreement with the observed values than in the Western United States, with r ranging from 0.4950 to 0.7976. The above analyses support that the GEOS-Chem model well captures the overall ozone-temperature sensitivity in the period of 1995-2017.


(a) Observed vs simulated $m_{\Delta O_3-\Delta T_{max}}$ trends




(b) $m_{\Delta O_3-\Delta T_{max}}$ trends in different emission scenarios




(c) Anthropogenic NO_x emission trends



(d) Contribution of NO_x emission reduction to $m_{\Delta O_3-\Delta T_{max}}$ trends





315

Figure 5. GEOS-Chem simulated decrease in summertime ozone-temperature sensitivity and the attribution to reduction in anthropogenic NO_x emission. (a) Spatial distributions of the observed (circles) and simulated (from the BASE simulation, shaded) $m_{\Delta O_3-\Delta T_{max}}$ trends during July from 1995 to 2017. Mean trends $\pm 95\%$ confidence level for the CONUS sites from the observation and GEOS-Chem model, and the correlation coefficients (r) of $m_{\Delta O_3-\Delta T_{max}}$ trends between the two are shown inset. (b) Time series of the observed and simulated $m_{\Delta O_3-\Delta T_{max}}$ in July during 1995-2017 (biennially) at CONUS sites. Results from the BASE simulation and a sensitivity simulation with anthropogenic NO_x emissions fixed at 1995 level (1995E) are compared. Colored circles denote the July anthropogenic NO_x emissions in the CONUS. (c) Spatial distribution of anthropogenic NO_x emission trends during July from 1995 to 2017. Trends are calculated for each model grid. Emissions trends aggregated over the CONUS are insets. (d) Contribution of anthropogenic NO_x emissions to $m_{\Delta O_3-\Delta T_{max}}$ trends, estimated as the difference in the $m_{\Delta O_3-\Delta T_{max}}$ trend between BASE and 1995E simulations. Mean trends $\pm 95\%$ confidence level is shown inset.

320

325

Figure 5(a) further compares the observed and GEOS-Chem simulated 1995-2017 trends in $m_{\Delta O_3-\Delta T_{max}}$ in July across the CONUS. The following analysis applies biennial data from 1995 to 2017 to align with the GEOS-Chem simulations. The overall observed $m_{\Delta O_3-\Delta T_{max}}$ trends in July, as depicted in Figure 5a, are similar to those in June-July-August period as presented

330 in Figure 3b, but there are slight differences at individual sites reflecting the difference in the time frame. We find that driven by yearly-varied meteorological fields and anthropogenic emissions, GEOS-Chem model successfully reproduces the decline of $m_{\Delta O_3-\Delta T_{max}}$ across the CONUS, showing a spatial correlation coefficient of 0.4746 with observed $m_{\Delta O_3-\Delta T_{max}}$ trends ($p < 0.01$). In particular, the model reproduces the much larger $m_{\Delta O_3-\Delta T_{max}}$ decreases in the eastern CONUS (the NEUS, Midwest, and SEUS) compared to other regions, consistent with the observations. However, the model has difficulty in capturing the magnitude of observed $m_{\Delta O_3-\Delta T_{max}}$ trends. The model shows a mean $m_{\Delta O_3-\Delta T_{max}}$ trend of -0.28 ppbv/K/decade over the CONUS that accounts for 42% of the observed trends of -0.67 ppbv/K/decade. ~~This largely reflects the model's bias in capturing the $m_{\Delta O_3-\Delta T_{max}}$ decreases in the SWUS region.~~ Figure 5b also shows that the model's underestimation of $m_{\Delta O_3-\Delta T_{max}}$ trends is primarily attributed to an overestimation of $m_{\Delta O_3-\Delta T_{max}}$ from 2013 to 2017 and an underestimation from 1995 to 1999. The consistency between the observed and simulated $m_{\Delta O_3-\Delta T_{max}}$ trends also shows regional differences. As shown in Figure S8, the model reproduces the interannual variation of $m_{\Delta O_3-\Delta T_{max}}$ well in the Plains and Intermountain West regions, and also captures 65% of the observed trend in the NWUS. However, in other regions, the model only captures less than 50% of the observed $m_{\Delta O_3-\Delta T_{max}}$ trends, with either an overestimation in 2013-2017 or underestimation in 1995-1999.

340 Our GEOS-Chem simulation has successfully reproduced the observed long-term ozone trend averaged over the CONUS (-6.1 ppbv/decade in GEOS-Chem vs -6.5 ppbv/decade in observations) (Table S5). However, capturing the long-term trends in $m_{\Delta O_3-\Delta T_{max}}$ can be more challenging than that of ozone concentrations, as it involves the combined uncertainty in temperature data, simulated ozone concentrations, and the parameterization of ozone-temperature response. The underestimation of $m_{\Delta O_3-\Delta T_{max}}$ from 1995 to 1999 may be partly attributed to the larger bias in MERRA-2 temperature dataset such that the observed $m_{\Delta O_3-\Delta T_{max}}$ may be biased compared to other periods (Figure S1), and such bias can propagate to the derivation of observed $m_{\Delta O_3-\Delta T_{max}}$ based on MERRA-2 dataset. Excluding the 1995, 1997, and 1999 records improve the model's ability in capturing observed $m_{\Delta O_3-\Delta T_{max}}$ trends (~~-0.46 ppbv/K/decade in GEOS-Chem vs -0.82 ppbv/K/decade, 56%~~), especially for the SWUS region (~~Figure 5b, Figure S7~~) in the CONUS (-0.46 ppbv/K/decade in GEOS-Chem vs -0.80 ppbv/K/decade, 58%). In particular, for the NEUS, Midwest, and SWUS, the model's ability to capture observed $m_{\Delta O_3-\Delta T_{max}}$ trends improves from 44%, 49%, and 23% to 83%, 66%, and 54%, respectively. The simulated ozone-temperature sensitivity for 2013-2017 shows an overestimation, particularly in the SEUS and Midwest regions (Figure S8). Christiansen et al. (2024) suggested that the CEDS inventory overestimates post-2010 anthropogenic NO_x emissions, especially in the eastern United States, which may lead to overestimation of ozone-temperature sensitivity in these regions. The GEOS-Chem model also misses several pathways in describing the responses of ozone to temperature, such as the responses in anthropogenic emission and land-atmosphere interaction through soil and vegetation. This will be discussed in detail in Section 4. We do not further differentiate the simulated $m_{\Delta O_3-\Delta T_{max}}$ trends at urban and rural sites because the model resolution at about 50 km may be too coarse for such separation.

360 Previous studies have implied reductions of anthropogenic emissions would result in a decrease in the ozone-temperature sensitivity (Bloomer et al., 2009). Here we explicitly test this ~~propose by theory using~~ our sensitivity experiments with anthropogenic NO_x emissions in US fixed at the 1995 level (1995E). Figure 5b shows that once the anthropogenic NO_x

emissions were fixed in 1995, the GEOS-Chem ~~would~~ simulate no decrease in $m_{\Delta O_3-\Delta T_{max}}$ (instead a positive trend by 0.13 ppbv/K/decade averaged over all sites, $p=0.12$). This implies that the change in anthropogenic NO_x emissions alone decreases $m_{\Delta O_3-\Delta T_{max}}$ by -0.41 ppbv/K/decade for all 608 sites, compared to the observed $m_{\Delta O_3-\Delta T_{max}}$ trend of -0.67 ppbv/K/decade, and is apparently the dominant driver of the observed decrease in $m_{\Delta O_3-\Delta T_{max}}$. In comparison, the simulation with only anthropogenic VOCs emissions fixed at the 1995 level shows ~~neglectable~~negligible difference in $m_{\Delta O_3-\Delta T_{max}}$ compared to the BASE simulation. We note that the difference in $m_{\Delta O_3-\Delta T_{max}}$ trend between BASE and 1995E simulations is highly consistent with the spatial distribution of anthropogenic NO_x emission trends ($r=0.40$, $p<0.01$) (Figure 5c), further confirming that NO_x emission reduction is an important driver of the decline in $m_{\Delta O_3-\Delta T_{max}}$. Figure ~~S7S8~~ illustrates that the regions with $m_{\Delta O_3-\Delta T_{max}}$ being mostly affected by anthropogenic NO_x emission reductions are located in the eastern CONUS (the NEUS, Midwest, and SEUS), while other regions are less affected.

3.3 The underlying mechanisms for the decrease in ozone-temperature sensitivity with reduced NO_x emissions




Figure 6. Mechanisms for the decrease in $m_{\Delta O_3-\Delta T_{max}}$ with anthropogenic NO_x emission reduction. (a) Changes in $m_{\Delta O_3-\Delta T_{max}}$ due to the difference in anthropogenic NO_x emissions in 2017 and 1995, estimated as the difference in $m_{\Delta O_3-\Delta T_{max}}$ between the BASE and 1995E simulation for July 2017. (b) The contribution of the temperature-indirect effect to $m_{\Delta O_3-\Delta T_{max}}$ with changes in anthropogenic NO_x emissions, estimated as the difference of $m_{\Delta O_3-\Delta T_{max}}$ between BASE-FTEMP and 1995E-FTEMP (Section 2.4). (c) The contribution of the temperature-direct effect, estimated as the difference of $m_{\Delta O_3-\Delta T_{max}}$ between BASE and 1995E minus the difference between BASE-FTEMP and 1995E-FTEMP. Mean, maximum, and minimum values of the contributions among all CONUS sites are shown inset.

Our analyses above prove that the reduction in anthropogenic NO_x emissions is the dominant driver of the observed long-term decrease in $m_{\Delta O_3-\Delta T_{max}}$ in the CONUS. We next examine how the changes in anthropogenic NO_x emissions have altered processes controlling ozone's response to temperature. Previous studies have shown that temperature's impacts on surface ozone concentrations involve acceleration of chemical reaction rates, in particular the thermal decomposition of PAN,

increased natural emissions of BVOCs and soil reactive nitrogen, and inhabitation of ozone dry deposition (Lu et al., 2019b; Porter and Heald, 2019; Steiner et al., 2010). Some studies also argued that the temperature-related covariance with other meteorological phenomena such as drought (low humidity), stagnancy, and transport may be more important in determining $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ (Kerr et al., 2019; Porter and Heald, 2019; Zhang et al., 2022a, c). Based on these previous studies, we focus on the changes of these impacts on $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ with anthropogenic emission reduction in the US.

We illustrate in Figure 6 the simulated changes in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ in July 2017 through temperature-direct effects and temperature-indirect effects associated with the anthropogenic reduction of NO_x . Figure 6a shows that the reduction of anthropogenic NO_x emissions from 1995 to 2017 alone decreased $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ by 0.86 ppbv/K in July 2017 (estimated as the difference between the BASE simulation and 1995E simulation). The decreases are larger in the eastern US (including NEUS, Midwest, and SEUS), reaching 1.37, 1.28, and 1.00 ppbv/K, respectively. When the temperature-direct effect is all removed from the GEOS-Chem simulation (Section 2.4), the reduction of anthropogenic NO_x emissions from 1995 to 2017 would decrease $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ by 0.65 ppbv/K in July 2017. It indicates that only a relatively small portion of the decrease in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ (24%, 0.21 ppbv/K compared with 0.86 ppbv/K) with anthropogenic NO_x reduction can be attributed to the temperature-direct effect (Figure 6c), yet the remaining is explained by temperature-indirect effect. Our results agree with Porter and Heald (2019), which shows that the collinearity between temperature and other meteorological variables played a significant role in determining the overall ozone-temperature relationship. Here, we further demonstrate that the temperature-indirect effect also dominates the decline in ozone-temperature sensitivity with anthropogenic NO_x emission reduction.




Figure 7. The different temperature-indirect effects for the decrease in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ with anthropogenic NO_x emission reduction. (a) the contribution of the transport to $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ decrease with changes in anthropogenic NO_x emissions, estimated as the difference of $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ between BASE-TRANS and 1995E-TRANS (Section 2.4). (b) the contribution of the other temperature-indirect effect, estimated as the difference of $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ between BASE-FTEMP and 1995E-FTEMP minus the difference between BASE-TRANS and 1995E-TRANS. Mean, maximum, and minimum values of the contributions among all CONUS sites are shown inset.

The temperature-indirect effect on ozone mainly includes the influence of temperature-relevant meteorological parameters such as humidity (as an indicator of content of water vapor) and shortwave radiation on ozone photochemistry, and


also the effect of transport (including stagnancy and regional transport). We further distinguish the impact of transport (by normalizing all meteorological elements except three-dimensional wind field and PBLH as input in the GEOS-Chem model) and the other indirect effects on the decrease in $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}}$ with emission reduction. As shown in Figure 7, transport (-0.28 ppbv/K) and other indirect effects (-0.37 ppbv/K) such as humidity and radiation show comparable contribution to the decline in $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}}$, but the spatial patterns show large disparity. The temperature-indirect effect excluding transport (Figure 7b) on $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}}$ shows a more uniform decline with reduced emissions in most regions across the CONUS, with a larger decrease in Southeast US. This may reflect the impact of solar radiation on BVOCs emission in Southeast US, and The radiation received by vegetation in the southeastern United States is highly collinear with temperature and also plays an important role in BVOC emissions (Guenther et al., 2012), which may reflect its potential for ozone formation reduces with the decline in anthropogenic NO_x emissions. In comparison, the transport effect has larger impacts on the $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}\text{T}_{\text{max}}}$ trend (Figure 7a) with reduced NO_x emissions in the northeastern US, where transport has the largest contribution to the mean $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}\text{T}_{\text{max}}}$ values (Figure ~~S9S10~~) as also reported in Kerr et al. (2019), ~~but~~. Some studies have demonstrated that changes in mid-latitude weather systems can significantly influence the ozone-temperature sensitivity by affecting pollutant transport (Barnes and Fiore, 2013; Kerr et al., 2020), which could be the underlying mechanism explaining the role of transport in contributing to the decrease of ozone-temperature sensitivity with emission reductions. But we find that ~~it causes~~these effects cause an increase in $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}\text{T}_{\text{max}}}$ in the southern US- in July 2007. Nevertheless, the impact of ~~transportation~~transport on $m_{\Delta\text{O}_3-\Delta\text{T}_{\text{max}}\text{T}_{\text{max}}}$ largely depends on the ~~transportation~~transport pattern itself, and it would be more ideally investigated through long-term simulations rather than the one-month study we conducted.

420

425

430

435



440 **Figure 8. The different temperature-~~indirect~~direct effects for the decrease in $m_{\text{AO3-ATmax}}$ with anthropogenic NO_x emission reduction. (a) Combined contribution of the four temperature-dependent mechanisms (BVOCs emissions, dry deposition, PAN decomposition, and soil NO_x emissions) to $m_{\text{AO3-ATmax}}$ with changes in anthropogenic NO_x emissions, estimated as the difference of $m_{\text{AO3-ATmax}}$ between BASE and 1995E minus the difference between BASE-~~FALLF4PATHS~~ and 1995E-~~FALLF4PATHS~~ (Section 2.4). (b)-(e) Individual contribution of BVOCs emissions, dry deposition, PAN decomposition, and soil NO_x emissions) to $m_{\text{AO3-ATmax}}$ with changes in anthropogenic NO_x emissions, respectively. Mean, maximum, and minimum values of the contributions among all CONUS sites are shown inset. Note that the data range of each figure is different.**

445

The temperature-direct effects on ozone-temperature sensitivity include the explicit impacts of temperature on BVOCs and soil NO_x emissions, chemical kinetics, and dry deposition. Figure 8 shows the additive and individual impacts from the

four temperature-dependent mechanisms (BVOCs, dry deposition, PAN decomposition, and soil NO_x) on the decrease in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$. Comparison of Figure 8a with Figure 6c shows that the contribution of the four temperature-dependent mechanisms contributes to almost all of the $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ decreases attributable to temperature-direct effect (-0.19 ppbv/K versus -0.21 ppbv/K).

We find that the ozone-temperature sensitivity contributed by BVOCs emissions has significantly reduced with anthropogenic emission control (Figure 8b). In July 2017, BVOCs emissions alone would have contributed to ozone-temperature sensitivity by 0.2 ppbv/K if anthropogenic emissions had remained at 1995 levels, with a particularly large contribution of 0.5 ppbv/K over the parts of eastern US where anthropogenic NO_x emissions are high and ozone formation is sensitivity to VOCs emissions (Figure [S10dS11d](#)). However, with anthropogenic NO_x emission decreased to the 2017 level, the contribution of BVOCs emissions decreases to 0.03 ppbv/K averaged over the CONUS sites and -0.01 ppbv/K averaged over the SEUS region (Figure [S10eS11c](#)). This suggests that the reduction in anthropogenic NO_x emission has shifted the ozone formation regime to a less VOCs-sensitive state, in which ozone concentrations are much less sensitive to increased BVOCs at high temperatures. Ozone-temperature sensitivity contributed by dry deposition also reduced by -0.03 ppbv/K averaged over the CONUS sites with anthropogenic emission reduction (Figure 8c).

The thermal decomposition of PAN contributes to 0.43 ppbv/K of the overall $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ over the CONUS (Figure [S10gS11g](#)), with larger contribution of 0.7 ppbv/K over the eastern US states. This is also consistent with Porter and Heald (2019), which shows the PAN decomposition explains a large fraction of the ozone-temperature sensitivity compared to other mechanisms such as BVOCs emissions and dry deposition. The PAN concentrations averaged over the CONUS decrease by 27% with the reduction in anthropogenic NO_x emissions (Figure [S11S12](#)). Nevertheless, $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ contributed by PAN decomposition only shows minor change with the reduction in anthropogenic NO_x emission of -0.02 ppbv/K averaged over the CONUS (Figure 8d), reflecting the offset between $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ increase in the central and western US and decrease in the eastern US. A possible reason is that, with the reduction of anthropogenic NO_x emissions, ozone formation in the central and western US becomes more NO_x-sensitive, as such the decomposition of PAN increases ozone-temperature sensitivity. The decrease in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ contributed by PAN decomposition in the eastern US may mainly reflect the reduction of PAN concentration with anthropogenic NO_x emission reduction (Figure [S11S12](#)).

Unlike the other mechanisms, $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ contributed by the temperature-dependent soil NO_x emissions increases by 0.03 ppbv/K averaged over the CONUS with anthropogenic NO_x emission reduction. The increase in $m_{\Delta\text{O}_3-\Delta\text{Tmax}}$ reflects the competitive effect between natural soil (from both natural pool and agricultural fertilizer, but are conventionally categorized as natural sources) and anthropogenic (from fossil fuel) NO_x emissions on ozone formation (Lu et al., 2021; Tan et al., 2023). Soil emissions become an increasingly important source of nitrogen for ozone formation with decreases anthropogenic NO_x emission levels- ([Guo et al., 2018](#); [Geddes et al., 2022](#)). As soil emissions are larger at higher temperatures, they contribute to an increasing ozone-temperature sensitivity. The above analysis reveals an increasing importance of soil NO_x emissions in determining ozone-temperature sensitivity in a future with low anthropogenic NO_x emissions.

3.4 Ozone mitigation benefit through the declined ozone-temperature sensitivity




Figure 9. Decreased $m_{\Delta O_3-\Delta T_{max}}$ offers ozone mitigation benefit at high temperatures. (a) Simulated ozone concentration in different T_{max} bins in the State of New York in July 2013, 2015, and 2017. Data are binned to 2K intervals. Results from the BASE simulation (black) and 1995E simulation (grey) are shown. The blue bars represent the ozone enhancement for each temperature bin compared to 291K from the BASE simulation. The bar marked by the grey boarder denotes ozone enhancement for each temperature bin compared to 291K from the 1995E simulation. Thus, the red bar (difference between the grey and blue bars) estimated the decrease in ozone enhancement due to the reduction of anthropogenic emissions from 1995 to 2017. (b) Distributions of ozone mitigation benefit in July due to the decreased $m_{\Delta O_3-\Delta T_{max}}$, estimated as the ozone enhancement from the lowest 0-10% to 90-100% temperatures bins in the 1995E minus that in the BASE at each grid in July (2013,2015 and 2017). Mean, max, and min values for the 608 sites are shown inset.

The significant decrease in $m_{\Delta O_3-\Delta T_{max}}$ over the CONUS indicates that controlling anthropogenic emission not only reduces the mean ozone levels, but also reduces the response of ozone to temperature. Consequently, this reduction lowers the risk of extreme ozone pollution and associated health damage at high temperatures, presenting an appealing benefit for ozone mitigation. We illustrate the benefit by reducing $m_{\Delta O_3-\Delta T_{max}}$ in ozone migration mitigation taking the State of New York as an example, as it has high $m_{\Delta O_3-\Delta T_{max}}$ and large population exposed to ozone pollution. Figure 9a shows the GEOS-Chem simulated ozone in July for three years (2013, 2015, 2017) at different T_{max} bins. As expected, MDA8 ozone increases with temperature rise. Ozone difference between the highest temperature bin (307K) and the lowest temperature bin (291K) is 31 ppbv in the BASE simulation, comparable to observations (26 ppbv between 307 and 291K). If anthropogenic NO_x emissions were fixed at the 1995 level, however, the predicted ozone difference between the 307K and 291K would be enlarged to 41 ppbv. This means that NO_x emission reductions cause an “additional” ozone concentration reduction of 10 ppbv from 291 to 307 K, as reflected in the significant decline in $m_{\Delta O_3-\Delta T_{max}}$. Such benefit of reducing $m_{\Delta O_3-\Delta T_{max}}$ is typically larger at higher temperature. Similar phenomenon can be found in other regions with significant decrease in $m_{\Delta O_3-\Delta T_{max}}$ (Figure S12S13).

Figure 9b further quantifies the beneficial effect of anthropogenic emission reduction on ozone mitigation through reducing $m_{\Delta O_3-\Delta T_{max}}$ over the CONUS. This can be estimated as the suppression of ozone increase between high (90-100th percentile of T_{max} in July 2013, 2015, 2017) and low temperature range (the lowest 10th percentile of T_{max}) due to anthropogenic

NO_x emission reduction from 1995 to 2017. We find that the additional ozone mitigation benefit by reducing $m_{\Delta O_3-\Delta T_{max}}$ is 6.8 ppbv averaged across the CONUS. The benefit is more pronounced in the eastern US, where emission reductions are more prominent, reaching a maximum of 19.4 ppbv. This benefit significantly reduces the probability of ozone exceedance (MDA8 ozone > 70 ppbv) during high-temperature conditions (above the 90th percentile of T_{max}), from 70% (estimated from the 1995E simulation) to 28% (from the BASE simulation). The results show that emission controlled on ozone precursors in the US have effectively reduced the ozone surge at high temperatures across the CONUS, and alleviated the combined health damage in the joint occurrences of heat and ozone extremes, highlighting the importance of continuous anthropogenic emission control on ozone mitigation in a warming future.

4. Summary and Discussion

We have estimated in this study the present-day (2017-2021) distributions and long-term (1995-2021) trends in summertime surface ozone-temperature sensitivity in the CONUS, combining observational monitoring network and GEOS-Chem simulations at a resolution of about 50km. We find a clear pattern that the observed $m_{\Delta O_3-\Delta T_{max}}$ for the CONUS decreased by 50% from 3.0 ppbv/K in 1990 to 1.5 ppbv/K in 2021 with a mean decreasing rate of -0.57 ppbv/K/decade ($p < 0.01$), with urban sites showing faster trends than rural sites (-0.61 vs -0.53 ppbv/K/decade), indicating a much weaker response of ozone to temperature in present-day compared to that in three decades ago. During the period from 1990 to 2021, anthropogenic NO_x emissions in the United States decreased by approximately 69%, and the eastern United States, where stricter anthropogenic emission controls were implemented, is the core region where ozone-temperature sensitivity has declined the most. The GEOS-Chem simulations driven by year-specific anthropogenic emission inventory and MERRA-2 reanalysis meteorological fields well reproduce the distribution and magnitude of multi-year mean $m_{\Delta O_3-\Delta T_{max}}$, and capture 42% of the observed trends in $m_{\Delta O_3-\Delta T_{max}}$ in 1995-2017. The model simulation shows that the decline in anthropogenic NO_x emission over the CONUS is the dominant driver of the $m_{\Delta O_3-\Delta T_{max}}$ decrease. Mechanically, approximately 76% of the simulated decline in $m_{\Delta O_3-\Delta T_{max}}$ can be attributed to the temperature-indirect effects arising from the shared collinearity of other meteorological effects (such as humidity, ventilation, and transport) on ozone. The remaining portion explaining the decrease in $m_{\Delta O_3-\Delta T_{max}}$ with anthropogenic NO_x emission reduction is mostly attributed to four direct temperature-dependent processes, in which $m_{\Delta O_3-\Delta T_{max}}$ decrease through the pathways of BVOCs emissions, dry deposition, and PAN decomposition (mostly in the eastern US), while soil NO_x emissions increase $m_{\Delta O_3-\Delta T_{max}}$ with anthropogenic NO_x emission reduction.

Our study illustrates that anthropogenic controls on NO_x emissions have significantly reduced the response of surface ozone concentration to the variation of temperature, offering a compelling advantage for ozone mitigation at high temperatures. The model simulation estimates that the reduction of anthropogenic NO_x emissions from 1995 to 2017 decreases the ozone enhancement from low to high temperatures by 6.8 ppbv on average across the CONUS (reaching 19 ppbv in the part of eastern US). The ozone-temperature sensitivity remains a crucial factor in quantifying the impact of climate on ozone. Our research demonstrates that anthropogenic emission changes not only alleviate current ozone pollution but also help mitigate potential

future increases in ozone concentrations due to climate change. It also indicates the dependency of ozone-temperature sensitivity on anthropogenic emission levels that should be considered in projecting future ozone concentration in a warmer climate.

545 Nevertheless, there is significant room for improving the ability in capturing the ozone-temperature relationship in the chemical transport model. The GEOS-Chem simulations do not account for the response of anthropogenic NO_x and VOCs emissions to temperature. Recent studies have shown that these emissions can increase simulated regional ozone-temperature sensitivity by up to 7% and 14% (Kerr et al., 2019; Wu et al., 2024). The parameterization of several temperature-dependent processes is limited or even missing in the model. For example, the dry deposition scheme used in this study lacks the temperature response of non-stomatal pathways (Clifton et al., 2020), which could introduce uncertainty in simulated $m_{\Delta O_3-\Delta T_{max}}$ particularly in vegetation-rich regions such as the southeastern United States. Additionally, according to the BDSNP scheme used in this study, soil NO_x emissions are modeled as an exponential function of temperature between 0 and 30 °C, remaining constant at temperatures above 30 °C. However, some studies have reported continuous increases in soil NO_x emissions at temperatures higher than 30 °C in regions such as California (Oikawa et al., 2015; Wang et al., 2021). The absence of other temperature-dependent natural emissions, such as soil Nitrous acid (HONO) (Tan et al., 2023), may also lead to an underestimation of ozone responses to extreme temperatures in the GEOS-Chem simulations. Uncertainties in the biomass burning emission inventory (Fasullo et al., 2022) limit the accuracy of ozone-temperature sensitivity simulations in fire-impacted regions, such as the mountainous western United States. The 50 km resolution of the model may not fully capture sub-grid meteorological variations, which can play an important role in reproducing extreme conditions at site-level scales. Our study demonstrates that ozone-temperature sensitivity is highly responsive to changes in emissions, emphasizing the importance of more accurate anthropogenic emissions inventory for interpreting the ozone-temperature relationship. Further efforts are needed to enhance the model's ability to capture long-term trends in the ozone response to temperature (including underlying weather conditions and transport patterns), and to better unravel the mechanisms driving the observed ozone-temperature relationship, in particular the role of transport and ventilation.

Data availability

565 The observational data used in this study is open-access as described in the study. Data from GEOS-Chem modeling that support the findings of this study can be accessed by contacting the corresponding authors (Xiao Lu, luxiao25@mail.sysu.edu.cn).

Acknowledgement

This study is supported by the National Key Research and Development Program of China (2023YFC3706104), **Guangdong**

570 Basic and Applied Basic Research project (No. 2020B0301030004), and the Young Elite Scientists Sponsorship Program by
CAST (2023QNRC001), the National Natural Science Foundation of China (42105103), and the Fundamental Research Funds
for the Central Universities (Sun Yat-sen University, 241gqb004).

Author contribution

575 X.L. designed the study. S.L. performed the model simulations and data analyses with significant input from H.L.W.. S.L. and
X.L. wrote the manuscript.

Competing interests.

The contact author has declared that none of the authors has any competing interests

References

580 Amos, H. M., Jacob, D. J., Holmes, C. D., Fisher, J. A., Wang, Q., Yantosca, R. M., Corbitt, E. S., Galarneau, E., Rutter, A. P.,
Gustin, M. S., Steffen, A., Schauer, J. J., Graydon, J. A., Louis, V. L. S., Talbot, R. W., Edgerton, E. S., Zhang, Y., and
Sunderland, E. M.: Gas-particle partitioning of atmospheric Hg(II) and its effect on global mercury deposition,
Atmospheric ~~Chem. Phys.~~ Chemistry and Physics, 12, 591–603, <https://doi.org/10.5194/acp-12-591-2012>, 2012.

585 Barnes, E. A. and Polvani, L.: Response of the Midlatitude Jets, and of Their Variability, to Increased Greenhouse Gases in the
CMIP5 Models. <https://doi.org/10.1175/JCLI-D-12-00536.1>, 2013.

Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz,
M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, ~~J.~~
Geophys. Res. Journal of Geophysical Research: Atmospheres, 106, 23073–23095,
<https://doi.org/10.1029/2001JD000807>, 2001.

590 Bloomer, B. J., Stehr, J. W., Piety, C. A., Salawitch, R. J., and Dickerson, R. R.: Observed relationships of ozone air pollution
with temperature and emissions, *Geophys. Res. Lett.*, 36, L09803, <https://doi.org/10.1029/2009GL037308>, 2009.

Camalier, L., Cox, W., and Dolwick, P.: The effects of meteorology on ozone in urban areas and their use in assessing ozone
trends, *Atmospheric Environment*, 41, 7127–7137, <https://doi.org/10.1016/j.atmosenv.2007.04.061>, 2007.

595 [Christiansen, A., Mickley, L. J., and Hu, L.: Constraining long-term NO_x emissions over the United States and Europe using nitrate wet deposition monitoring networks, Atmospheric Chemistry and Physics, 24, 4569–4589, <https://doi.org/10.5194/acp-24-4569-2024>, 2024.](https://doi.org/10.5194/acp-24-4569-2024)

600 [Clifton, O. E., Fiore, A. M., Massman, W. J., Baublitz, C. B., Coyle, M., Emberson, L., Fares, S., Farmer, D. K., Gentine, P., Gerosa, G., Guenther, A. B., Helmig, D., Lombardozi, D. L., Munger, J. W., Patton, E. G., Pusede, S. E., Schwede, D. B., Silva, S. J., Sörgel, M., Steiner, A. L., and Tai, A. P. K.: Dry Deposition of Ozone Over Land: Processes, Measurement, and Modeling, Reviews of Geophysics, 58, e2019RG000670, <https://doi.org/10.1029/2019RG000670>, 2020.](https://doi.org/10.1029/2019RG000670)

[Fasullo, J. T., Lamarque, J.-F., Hannay, C., Rosenbloom, N., Tilmes, S., DeRepentigny, P., Jahn, A., and Deser, C.: Spurious Late Historical-Era Warming in CESM2 Driven by Prescribed Biomass Burning Emissions, Geophysical Research Letters, 49, e2021GL097420, <https://doi.org/10.1029/2021GL097420>, 2022.](https://doi.org/10.1029/2021GL097420)

605 [Feng, Z., Xu, Y., Kobayashi, K., Dai, L., Zhang, T., Agathokleous, E., Calatayud, V., Paoletti, E., Mukherjee, A., Agrawal, M., Park, R. J., Oak, Y. J., and Yue, X.: Ozone pollution threatens the production of major staple crops in East Asia, Nat- Food, 3, 47–56, <https://doi.org/10.1038/s43016-021-00422-6>, 2022.](https://doi.org/10.1038/s43016-021-00422-6)

610 [Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S., Eyring, V., Folberth, G. A., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. A., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and climate, Chem. Soc. Rev., 41, 6663–6683, <https://doi.org/10.1039/C2CS35095E>, 2012.](https://doi.org/10.1039/C2CS35095E)

[Fu, T.-M. and Tian, H.: Climate Change Penalty to Ozone Air Quality: Review of Current Understandings and Knowledge Gaps, Curr-~~Pollut.~~ \[Pollution\]\(https://doi.org/10.1007/s40726-019-00115-6\) Rep-~~s~~, 5, 159–171, <https://doi.org/10.1007/s40726-019-00115-6>, 2019.](https://doi.org/10.1007/s40726-019-00115-6)

615 [Fu, T.-M., Zheng, Y., Paulot, F., Mao, J., and Yantosca, R. M.: Positive but variable sensitivity of August surface ozone to large-scale warming in the southeast United States, \[Nat-Nature\]\(https://doi.org/10.1038/nclimate2567\) Clim- Change, 5, 454–458, <https://doi.org/10.1038/nclimate2567>, 2015.](https://doi.org/10.1038/nclimate2567)

620 [Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P.-F., Cuesta, J., Cuevas, E., Doniki, S., Dufour, G., Ebojje, F., Foret, G., Garcia, O., Granados-Muñoz, M. J., Hannigan, J. W., Hase, F., Hassler, B., Huang, G., Hurtmans, D., Jaffe, D., Jones, N., Kalabokas, P., Kerridge, B., Kulawik, S., Latter, B., Leblanc, T., Le Flochmoën, E., Lin, W., Liu, J., Liu, X., Mahieu, E., McClure-Begley, A., Neu, J. L., Osman, M., Palm, M., Petetin, H., Petropavlovskikh, I., Querel, R., Rahpoe, N., Rozanov, A., Schultz, M. G., Schwab, J., Siddans, R., Smale, D., Steinbacher, M., Tanimoto, H., Tarasick, D. W., Thouret, V., Thompson, A. M., Trickl, T., Weatherhead, E., Wespes, C., Worden, H.](https://doi.org/10.1002/2019GL084011)

625 M., Vigouroux, C., Xu, X., Zeng, G., and Ziemke, J.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, ~~Elem. Sci. Anthr.~~Elementa: Science of the Anthropocene, 6, 39, <https://doi.org/10.1525/elementa.291>, 2018.

Geddes, J. A., Pusede, S. E., and Wong, A. Y. H.: Changes in the Relative Importance of Biogenic Isoprene and Soil NO_x Emissions on Ozone Concentrations in Nonattainment Areas of the United States, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036361, <https://doi.org/10.1029/2021JD036361>, 2022.

630 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Silva, A. M. da, Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), ~~J. Clim.~~Journal of Climate, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.

635 Gu, Y., Li, K., Xu, J., Liao, H., and Zhou, G.: Observed dependence of surface ozone on increasing temperature in Shanghai, China, ~~Atmos. Environ.~~Atmospheric Environment, 221, 117108, <https://doi.org/10.1016/j.atmosenv.2019.117108>, 2020.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, ~~Geosci-Geoscientific~~ Model ~~Dev.~~Development, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.

640 Guo, J. J., Fiore, A. M., Murray, L. T., Jaffe, D. A., Schnell, J. L., Moore, C. T., and Milly, G. P.: Average versus high surface ozone levels over the continental USA: model bias, background influences, and interannual variability, *Atmospheric Chemistry and Physics*, 18, 12123–12140, <https://doi.org/10.5194/acp-18-12123-2018>, 2018.

645 Hembeck, L., Dickerson, R. R., Canty, T. P., Allen, D. J., and Salawitch, R. J.: Investigation of the Community Multiscale air quality (CMAQ) model representation of the Climate Penalty Factor (CPF), *Atmospheric Environment*, 283, 119157, <https://doi.org/10.1016/j.atmosenv.2022.119157>, 2022.

Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C., and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints, ~~Atmospheric Chem. Phys.~~Chemistry and Physics, 12, 7779–7795, <https://doi.org/10.5194/acp-12-7779-2012>, 2012.

650 IUPAC: Task group on atmospheric chemical kinetic data evaluation by International Union of Pure and Applied Chemistry (IUPAC), available at: <http://iupac.pole-ether.fr/> (last access: 22 June 2023), 2013.

- Jacob, D. J. and Winner, D. A.: Effect of climate change on air quality, ~~Atmos. Environ.~~Atmospheric Environment, 43, 51–63, <https://doi.org/10.1016/j.atmosenv.2008.09.051>, 2009.
- Jaffe, D. A., Cooper, O. R., Fiore, A. M., Henderson, B. H., Tonnesen, G. S., Russell, A. G., Henze, D. K., Langford, A. O., Lin, M., and Moore, T.: Scientific assessment of background ozone over the U.S.: Implications for air quality management, ~~Elem. Sci. Anthr.~~Elementa: Science of the Anthropocene, 6, 56, <https://doi.org/10.1525/elementa.309>, 2018.
- Jing, P., Lu, Z., and Steiner, A. L.: The ozone-climate penalty in the Midwestern U.S., ~~Atmos. Environ.~~Atmospheric Environment, 170, 130–142, <https://doi.org/10.1016/j.atmosenv.2017.09.038>, 2017.
- Kerr, G. H., Waugh, D. W., Strode, S. A., Steenrod, S. D., Oman, L. D., and Strahan, S. E.: Disentangling the Drivers of the Summertime Ozone-Temperature Relationship Over the United States, *J. Geophys. Res.* ~~Atmospheres~~Atmos., 124, 10503–10524, <https://doi.org/10.1029/2019JD030572>, 2019.
- Kerr, G. H., Waugh, D. W., Steenrod, S. D., Strode, S. A., and Strahan, S. E.: Surface Ozone-Meteorology Relationships: Spatial Variations and the Role of the Jet Stream. *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032735, <https://doi.org/10.1029/2020JD032735>, 2020.
- Kim, S.-W., Heckel, A., McKeen, S. A., Frost, G. J., Hsie, E.-Y., Trainer, M. K., Richter, A., Burrows, J. P., Peckham, S. E., and Grell, G. A.: Satellite-observed U.S. power plant ~~NO_x~~NO_x emission reductions and their impact on air quality, *Geophys. Res. Lett.*, 33, L22812, <https://doi.org/10.1029/2006GL027749>, 2006.
- Leibensperger, E. M., Mickley, L. J., and Jacob, D. J.: Sensitivity of US air quality to mid-latitude cyclone frequency and implications of 1980–2006 climate change, *Atmospheric* ~~Chem. Phys.~~Chemistry and Physics, 8, 7075–7086, <https://doi.org/10.5194/acp-8-7075-2008>, 2008.
- Lin, M., Horowitz, L. W., Payton, R., Fiore, A. M., and Tonnesen, G.: US surface ozone trends and extremes from 1980 to 2014: quantifying the roles of rising Asian emissions, domestic controls, wildfires, and climate, ~~Atmospheric~~Atmos. Chem. Phys., 17, 2943–2970, <https://doi.org/10.5194/acp-17-2943-2017>, 2017.
- Lin, M., Horowitz, L. W., Xie, Y., Paulot, F., Malyshev, S., Shevliakova, E., Finco, A., Gerosa, G., Kubistin, D., and Pilegaard, K.: Vegetation feedbacks during drought exacerbate ozone air pollution extremes in Europe, *Nat. Clim. Change*, ~~Change~~Chang., 10, 444–451, <https://doi.org/10.1038/s41558-020-0743-y>, 2020.
- Liu, H., Jacob, D. J., Bey, I., and Yantosca, R. M.: Constraints from 210Pb and 7Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, ~~J. Geophys. Res.~~Journal of

[Geophysical Research](https://doi.org/10.1029/2000JD900839): Atmospheres, 106, 12109–12128, <https://doi.org/10.1029/2000JD900839>, 2001.

680 Liu, S., Shu, L., Zhu, L., Song, Y., Sun, W., Chen, Y., Wang, D., Pu, D., Li, X., Sun, S., Li, J., Zuo, X., Fu, W., Yang, X., and Fu, T.-M.: Underappreciated Emission Spikes From Power Plants During Heatwaves Observed From Space: Case Studies in India and China, [Earth's Future](https://doi.org/10.1029/2023EF003937), 12, e2023EF003937, <https://doi.org/10.1029/2023EF003937>, 2024.

Lu, X., Zhang, L., Chen, Y., Zhou, M., Zheng, B., Li, K., Liu, Y., Lin, J., Fu, T.-M., and Zhang, Q.: Exploring 2016–2017 surface ozone pollution over China: source contributions and meteorological influences, [Atmospheric Chem. Phys.](https://doi.org/10.5194/acp-19-8339-2019), 19, 8339–8361, <https://doi.org/10.5194/acp-19-8339-2019>, 2019a.

685 Lu, X., Zhang, L., and Shen, L.: Meteorology and Climate Influences on Tropospheric Ozone: a Review of Natural Sources, Chemistry, and Transport Patterns, [Curr. Pollut. Pollution Rep.](https://doi.org/10.1007/s40726-019-00118-3), 5, 238–260, <https://doi.org/10.1007/s40726-019-00118-3>, 2019b.

690 Lu, X., Ye, X., Zhou, M., Zhao, Y., Weng, H., Kong, H., Li, K., Gao, M., Zheng, B., Lin, J., Zhou, F., Zhang, Q., Wu, D., Zhang, L., and Zhang, Y.: The underappreciated role of agricultural soil nitrogen oxide emissions in ozone pollution regulation in North China, [Nat. Commun.](https://doi.org/10.1038/s41467-021-25147-9), 12, 5021, <https://doi.org/10.1038/s41467-021-25147-9>, 2021.

695 van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), [Geosci. Model Dev.](https://doi.org/10.5194/gmd-10-3329-2017), 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.

McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), [Earth Syst. Sci. Data](https://doi.org/10.5194/essd-12-3413-2020), 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.

700 Mills, G., Pleijel, H., Malley, C. S., Sinha, B., Cooper, O. R., Schultz, M. G., Neufeld, H. S., Simpson, D., Sharps, K., Feng, Z., Gerosa, G., Harmens, H., Kobayashi, K., Saxena, P., Paoletti, E., Sinha, V., and Xu, X.: Tropospheric Ozone Assessment Report: Present-day tropospheric ozone distribution and trends relevant to vegetation, [Elem. Sci. Anthr. Elementa: Science of the Anthropocene](https://doi.org/10.1525/elementa.302), 6, 47, <https://doi.org/10.1525/elementa.302>, 2018.

705 Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von Schneidmesser, E., Sommariva, R., Wild, O., and Williams, M. L.:

Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, [Atmospheric Atmos. Chem. Phys.](https://doi.org/10.5194/acp-15-8889-2015), 15, 8889–8973, <https://doi.org/10.5194/acp-15-8889-2015>, 2015.

710 Murray, L. T., Jacob, D. J., Logan, J. A., Hudman, R. C., and Koshak, W. J.: Optimized regional and interannual variability of lightning in a global chemical transport model constrained by LIS/OTD satellite data, [J. Geophys. Res. Journal of Geophysical Research: Atmospheres](https://doi.org/10.1029/2012JD017934), 117, <https://doi.org/10.1029/2012JD017934>, 2012.

Ning, G., Wardle, D. A., and Yim, S. H. L.: Suppression of Ozone Formation at High Temperature in China: From Historical Observations to Future Projections, [Geophys. Geophysical Research Letters](https://doi.org/10.1029/2021GL097090), ~~Res. Lett.~~ 49, <https://doi.org/10.1029/2021GL097090>, 2022.

715 Nolte, C. G., Spero, T. L., Bowden, J. H., Sarofim, M. C., Martinich, J., and Mallard, M. S.: Regional temperature-ozone relationships across the U.S. under multiple climate and emissions scenarios, [J. Journal of the Air & Waste Manag. Assoc., Management Association](https://doi.org/10.1080/10962247.2021.1970048), 71, 1251–1264, <https://doi.org/10.1080/10962247.2021.1970048>, 2021.

Oikawa, P. Y., Ge, C., Wang, J., Eberwein, J. R., Liang, L. L., Allsman, L. A., Grantz, D. A., and Jenerette, G. D.: Unusually high soil nitrogen oxide emissions influence air quality in a high-temperature agricultural region, [Nat. Commun.](https://doi.org/10.1038/ncomms9753), 6, 8753, <https://doi.org/10.1038/ncomms9753>, 2015.

720 Porter, W. C. and Heald, C. L.: The mechanisms and meteorological drivers of the summertime ozone–temperature relationship, [Atmospheric Atmos. Chem. Phys.](https://doi.org/10.5194/acp-19-13367-2019), 19, 13367–13381, <https://doi.org/10.5194/acp-19-13367-2019>, 2019.

725 Pusede, S. E., Gentner, D. R., Wooldridge, P. J., Browne, E. C., Rollins, A. W., Min, K.-E., Russell, A. R., Thomas, J., Zhang, L., Brune, W. H., Henry, S. B., DiGangi, J. P., Keutsch, F. N., Harrold, S. A., Thornton, J. A., Beaver, M. R., St. Clair, J. M., Wennberg, P. O., Sanders, J., Ren, X., VandenBoer, T. C., Markovic, M. Z., Guha, A., Weber, R., Goldstein, A. H., and Cohen, R. C.: On the temperature dependence of organic reactivity, nitrogen oxides, ozone production, and the impact of emission controls in San Joaquin Valley, California, [Atmospheric Atmos. Chem. Phys.](https://doi.org/10.5194/acp-14-3373-2014), 14, 3373–3395, <https://doi.org/10.5194/acp-14-3373-2014>, 2014.

Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and Recent Trends in the Chemistry of Continental Surface Ozone, [Chem. Rev.](https://doi.org/10.1021/cr5006815), 115, 3898–3918, <https://doi.org/10.1021/cr5006815>, 2015.

730 Rasmussen, D. J., Fiore, A. M., Naik, V., Horowitz, L. W., McGinnis, S. J., and Schultz, M. G.: Surface ozone-temperature relationships in the eastern US: A monthly climatology for evaluating chemistry-climate models, [Atmos. Environ., Atmospheric Environment](https://doi.org/10.1016/j.atmosenv.2011.11.021), 47, 142–153, <https://doi.org/10.1016/j.atmosenv.2011.11.021>, 2012.

Rasmussen, D. J., Hu, J., Mahmud, A., and Kleeman, M. J.: The Ozone–Climate Penalty: Past, Present, and Future, *Environ. Sci. Technol.*, 47, 14258–14266, <https://doi.org/10.1021/es403446m>, 2013.

735 Romer, P. S., Duffey, K. C., Wooldridge, P. J., Edgerton, E., Baumann, K., Feiner, P. A., Miller, D. O., Brune, W. H., Koss, A. R., de Gouw, J. A., Misztal, P. K., Goldstein, A. H., and Cohen, R. C.: Effects of temperature-dependent NO_x emissions on continental ozone production, *Atmospheric ~~Chem. Phys.~~ Chemistry and Physics*, 18, 2601–2614, <https://doi.org/10.5194/acp-18-2601-2018>, 2018.

740 Sander, S. P., Golden, D., Kurylo, M., Moortgat, G., Wine, P., Ravishankara, A., Kolb, C., Molina, M., Finlayson-Pitts, B., and Huie, R.: Chemical kinetics and photochemical data for use in atmospheric studies, JPL Publ., 06-2, 684 pp., 2011.

Shen, L., Mickley, L. J., and Gilleland, E.: Impact of increasing heat waves on U.S. ozone episodes in the 2050s: Results from a multimodel analysis using extreme value theory, *~~Geophys. Res. Lett.~~ Geophysical Research Letters*, 43, 4017–4025, <https://doi.org/10.1002/2016GL068432>, 2016.

745 Sillman, S. and Samson, P. J.: Impact of temperature on oxidant photochemistry in urban, polluted rural and remote environments, *~~J. Geophys. Res.~~ Journal of Geophysical Research: Atmospheres*, 100, 11497–11508, <https://doi.org/10.1029/94JD02146>, 1995.

Simon, H., Reff, A., Wells, B., Xing, J., and Frank, N.: Ozone Trends Across the United States over a Period of Decreasing NO_x and VOC Emissions, *Environ. Sci. Technol.*, 49, 186–195, <https://doi.org/10.1021/es504514z>, 2015.

750 Steiner, A. L., Davis, A. J., Sillman, S., Owen, R. C., Michalak, A. M., and Fiore, A. M.: Observed suppression of ozone formation at extremely high temperatures due to chemical and biophysical feedbacks, *Proc. Natl. Acad. Sci. U.S.A.*, 107, 19685–19690, <https://doi.org/10.1073/pnas.1008336107>, 2010.

Tan, W., Wang, H., Su, J., Sun, R., He, C., Lu, X., Lin, J., Xue, C., Wang, H., Liu, Y., Liu, L., Zhang, L., Wu, D., Mu, Y., and Fan, S.: Soil Emissions of Reactive Nitrogen Accelerate Summertime Surface Ozone Increases in the North China Plain, *Environ. Sci. Technol.*, 57, 12782–12793, <https://doi.org/10.1021/acs.est.3c01823>, 2023.

755 [Tawfik, A. B. and Steiner, A. L.: A proposed physical mechanism for ozone-meteorology correlations using land–atmosphere coupling regimes, *Atmospheric Environment*, 72, 50–59, <https://doi.org/10.1016/j.atmosenv.2013.03.002>, 2013.](https://doi.org/10.1016/j.atmosenv.2013.03.002)

Travis, K. R. and Jacob, D. J.: Systematic bias in evaluating chemical transport models with maximum daily 8 h average (MDA8) surface ozone for air quality applications: a case study with GEOS-Chem v9.02, *~~Geosci. Geoscientific~~ Model Dev. Development*, 12, 3641–3648, <https://doi.org/10.5194/gmd-12-3641-2019>, 2019.

760 Turner, M. C., Jerrett, M., Pope, C. A., Krewski, D., Gapstur, S. M., Diver, W. R., Beckerman, B. S., Marshall, J. D., Su, J.,
Crouse, D. L., and Burnett, R. T.: Long-Term Ozone Exposure and Mortality in a Large Prospective Study, *Am. J. Respir.
Crit. Care Med.*, 193, 1134–1142, <https://doi.org/10.1164/rccm.201508-1633OC>, 2016.

Varotsos, K. V., Giannakopoulos, C., and Tombrou, M.: Ozone-temperature relationship during the 2003 and 2014 heatwaves
in Europe, *Reg. Environ. Change*, 19, 1653–1665, <https://doi.org/10.1007/s10113-019-01498-4>, 2019.

765 [Wang, H., Lu, X., Palmer, P., Zhang, L., Lu, K., Li, K., Nagashima, T., Koo, J., Tanimoto, H., Wang, H., Gao, M., He, C., Wu,
K., Fan, S., and Zhang, Y.: Deciphering decadal urban ozone trends from historical records since 1980, *National Science
Review*, In press, 2024](#)

Wang, Y., Ge, C., Garcia, L. C., Jenerette, G. D., Oikawa, P. Y., and Wang, J.: Improved modelling of soil NO_x emissions in a
high temperature agricultural region: role of background emissions on NO₂ trend over the US, *Environ. Res. Lett.*, 16,
770 084061, <https://doi.org/10.1088/1748-9326/ac16a3>, 2021.

van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E.,
Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016,
Earth Syst. Sci. System Science Data, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.

Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models,
775 *Atmospheric Environ.* (1967), 23, 1293–1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.

Wu, S., Mickley, L. J., Leibensperger, E. M., Jacob, D. J., Rind, D., and Streets, D. G.: Effects of 2000–2050 global change on
ozone air quality in the United States, *J. Geophys. Res.*, 113, D06302, <https://doi.org/10.1029/2007JD008917>, 2008.

Wu, W., Fu, T.-M., Arnold, S. R., Spracklen, D. V., Zhang, A., Tao, W., Wang, X., Hou, Y., Mo, J., Chen, J., Li, Y., Feng, X.,
Lin, H., Huang, Z., Zheng, J., Shen, H., Zhu, L., Wang, C., Ye, J., and Yang, X.: Temperature-Dependent Evaporative
780 Anthropogenic VOC Emissions Significantly Exacerbate Regional Ozone Pollution, *Environ. Sci. Technol.*,
<https://doi.org/10.1021/acs.est.3c09122>, 2024.

[Zanis, P., Akritidis, D., Turnock, S., Naik, V., Szopa, S., Georgoulas, A. K., Bauer, S. E., Deushi, M., Horowitz, L. W., Keeble,
J., Sager, P. L., O'Connor, F. M., Oshima, N., Tsigaridis, K., and Noije, T. van: Climate change penalty and benefit on
surface ozone: a global perspective based on CMIP6 earth system models, *Environ. Res. Lett.*, 17, 024014,
785 <https://doi.org/10.1088/1748-9326/ac4a34>, 2022.](#)

Zhang, J., Gao, Y., Leung, L. R., Luo, K., Wang, M., Zhang, Y., Bell, M. L., and Fan, J.: Disentangling the mechanism of

temperature and water vapor modulation on ozone under a warming climate, *Environ. Res. Lett.*, 17, 124032, <https://doi.org/10.1088/1748-9326/aca3bc>, 2022a.

790

Zhang, J., Gao, Y., Leung, L. R., Luo, K., Wang, M., Zhang, Y., Bell, M. L., and Fan, J.: Isolating the modulation of mean warming and higher-order temperature changes on ozone in a changing climate over the contiguous United States, *Environ. Res. Lett.*, 17, 094005, <https://doi.org/10.1088/1748-9326/ac8695>, 2022b.

Zhang, L., Jacob, D. J., Yue, X., Downey, N. V., Wood, D. A., and Blewitt, D.: Sources contributing to background surface ozone in the US Intermountain West, *Atmospheric ~~Chem. Phys.~~ Chemistry and Physics*, 14, 5295–5309, <https://doi.org/10.5194/acp-14-5295-2014>, 2014.

795

Zhang, X., Waugh, D. W., Kerr, G. H., and Miller, S. M.: Surface Ozone-Temperature Relationship: The Meridional Gradient Ratio Approximation, *~~Geophys. Res. Lett.~~ Geophysical Research Letters*, 49, e2022GL098680, <https://doi.org/10.1029/2022GL098680>, 2022c.

800