

Multispecies expression of coccolithophore vital effects with changing CO₂ concentrations and pH in the laboratory with insights for reconstructing CO₂ levels in geological history

Goulwen Le Guevel^{1,2}, Fabrice Minoletti¹, Carla Geisen², Gwendoline Duong³, Virginia Rojas¹ and Michaël Hermoso²

¹Institut des Sciences de la Terre de Paris (UMR 7193 ISTeP), CNRS, Sorbonne Université, 75005 Paris, France

²Laboratoire d'Océanologie et de Géosciences (UMR 8187 LOG), Université du Littoral Côte d'Opale, CNRS, Université de Lille, 62930 Wimereux, France

³Laboratoire d'Océanologie et de Géosciences (UMR 8187 LOG), Université de Lille, CNRS, Université du Littoral Côte d'Opale, 62930 Wimereux, France

Correspondence to: Michael Hermoso (michael.hermoso@univ-littoral.fr)

Abstract. The coccolith sedimentary and micropalaeontological archive has fostered great interest for palaeoclimate applications. Indeed, the geochemistry of coccolith calcite has the potential to reconstruct both palaeo-CO₂ concentrations and palaeo-temperature of seawater. Studying coccolith geochemistry aims at better understanding the changes in the vital effect of coccoliths with changes in environmental parameters, especially the carbonate chemistry of seawater. To this aim, we need to deconvolve the biological imprint from the environmental signals recorded in the composition of coccolith-biominerals. We have undertaken large-scale culture experiments of four strains of coccolithophores strains with of various sizes and growth rates, grown under eight CO₂/pH conditions typifying the long-term CO₂ evolution of the Cenozoic Era. We propose an assessment of the expression of the vital effects for *Emiliania huxleyi*, *Gephyrocapsa oceanica*, *Helicosphaera carteri* and *Coccolithus braarudii* with simultaneous changes in Dissolved Inorganic Carbon (DIC) and pH in the medium resulting in variations in dissolved CO_{2-aq} (CO_{2-aq}) availability to the cells. We have identified a distinct isotopic response of *C. braarudii* to pCO₂ levels on either both sides of the 600 ppmv (pH 7.89) condition. We propose that this discrepancy is the result of a modification of the proton efflux across the plasma membrane through (voltage-dependent proton channels). We further show that as the CO₂ level rises and pH decreases (from 200 to 500 ppmv and from 8.29 to 7.96 pH units, respectively), a significant increase in $\delta^{13}\text{C}_{\text{coccolith}}$ of *C. braarudii* is expressed, along with a coeval decrease in $\delta^{13}\text{C}_{\text{org}}$. The constant physiological parameters of *C. braarudii* (growth rate, PIC, POC) across the 200 to 500 ppmv interval support the idea that the change in $\delta^{13}\text{C}_{\text{coccolith}}$ is only a the consequence of a lower fractionation between dissolved CO₂ and organic matter. Meanwhile, the small (less carbon-limited) cells of *E. huxleyi* and *G. oceanica* are less carbon-limited and do not exhibit any change in their carbon vital effects with changes in carbonate chemistry of the environment across the whole CO₂ spectrum. Using this new biogeochemical framework, we have established a calibration between CO_{2-aq} concentration and the differential vital effect ($\Delta\delta^{13}\text{C}$) between isotopically-invariant small *G. oceanica* and large coccoliths *C. braarudii*, whose vital effect is CO₂-

dependent at low CO₂. The CO₂-Δδ¹³C transfer equation allows palaeo-pCO₂ reconstructions based on isotope changes explained by physiological processes, especially at low and medium CO₂ levels.

35 1 Introduction

The atmospheric concentration of carbon dioxide (pCO₂) is a key parameter controlling global climate through its radiative forcing on Earth's surface temperatures. The reconstruction of past pCO₂ is challenging to handle as many caveats exist in the palaeo-CO₂ barometry methods. The magnitude of the carbon isotope fractionation between membrane lipids called alkenones produced by the coccolithophores – found in sediments – and ambient aqueous CO₂ constitutes the basis of the foremost
40 palaeo-pCO₂ proxy, referred to as ε_{p-alk} (Pagani et al., 2010; Zhang et al., 2019, 2020). This approach has led to significant advances in the understanding of the past CO₂ concentration, but its application still has some limitations. Conversely to the coccolith archive, alkenone molecules are not ubiquitously preserved in marine sediments and the assessment of pCO₂ involves a cascade of calculations requiring assumptions made on the δ¹³C value of CO₂aq. Furthermore, this method has recently been shown to overestimate pCO₂ below 270 ppmv (Badger et al., 2019), potentially leading to underestimated pCO₂ above 270
45 ppmv. Alongside this proxy, the boron isotope fractionation in foraminifera tests has increasingly been used to reconstruct palaeo-pH (Rae et al., 2021). Combined with another parameter of the carbonate chemistry system as DIC or alkalinity, and with temperature, palaeo-pH can be used for palaeo-pCO₂ reconstructions (Foster, 2008; Sanyal et al., 1995). This proxy also suffers from uncertainties due to the evaluation of δ¹¹B of past seawater among others unknown factors (Klochko et al., 2006; Tripathi et al., 2011).

50 In this study, we investigated the fractionation of carbon and oxygen isotopes in phytoplanktonic organic matter and coccolith calcite driven by changes in the carbonate chemistry of coccolithophores culture media. We have conducted *in vivo* culture experiments, that provide a means to constrain the cellular and isotopic responses of coccolithophores to environmental changes. Our culture experiments were conducted to quantify the isotopic departure of coccoliths from inorganic calcite, called
(vital effect). The latter only depends on physico-chemical parameters of the environment such as δ¹⁸O_{sw}, δ¹³C_{CO2}, temperature,
55 salinity, and pH (Zeebe and Wolf-Gladrow, 2001). However, when calcite is biomineralised intracellularly, biological parameters such as growth rate, cell size, and the PIC/POC ratio - which refers to the distribution of carbon between particulate organic carbon (POC) and particulate inorganic carbon (PIC) produced by calcifying organisms - also influence this fractionation (Dudley et al., 1986; McClelland et al., 2017; Rickaby et al., 2010). The causes behind the changes in the isotopic fractionation between the different cellular compartments are to be investigated at various stages of carbon fixation. Aqueous
60 CO₂, the main form of DIC entering coccolithophorid cells, diffuses through the plasma membrane via passive diffusion (Gutknecht et al., 1977). Aqueous CO₂ diffusion is governed primarily by Fick's first law, but also by the specific conditions of the cell boundary layer on the external side of the plasma membrane (Reinfelder, 2011; Wolf-Gladrow and Riebesell, 1997). In addition to passive CO₂ diffusion, certain taxa of coccolithophores such as *Emiliania* possess CCMs (Carbon Concentration Mechanisms), allowing the intake of HCO₃⁻ through active transmembrane transport (Bach et al., 2013; Parker and Boron,

65 2013; Romero et al., 2004). As HCO_3^- is enriched in ^{13}C compared to CO_2 , the greater acquisition of HCO_3^- , the higher $\delta^{13}\text{C}$ values in the intracellular DIC pool and ultimately the higher $\delta^{13}\text{C}$ of coccolith calcite.


Once inside the cell, carbon is utilised by two main pathways: the fixation into organic matter via photosynthesis in chloroplasts and precipitation of calcite forming coccoliths within the coccolith vesicle. Carbon fixation into organic matter by the enzyme RuBisCO (ribulose 1,5-bisphosphate carboxylase/oxygenase) (Ellis, 1979) leads to fractionation as it preferentially fixes the
70 lighter isotopes (^{12}C) (Laws et al., 2001; Popp et al., 1998). Consequently, organic matter is relatively depleted in ^{13}C (very negative $\delta^{13}\text{C}_{\text{org}}$) compared to the carbon source and compared to the intracellular DIC pool (Bidigare et al., 1997; Guy et al., 1993; Rau et al., 1996). The fractionation induced by RuBisCO is one component of a more general fractionation between the CO_2_{aq} from the external environment and the organic matter synthesised within the cell, which leads to an enrichment, noted ϵ_p (Jasper and Hayes, 1990; Pagani et al., 1999). In addition to being dependent on the fractionation associated with RuBisCO,
75 ϵ_p is influenced by physiological parameters such as growth rate, cell size, and the presence of CCMs (Laws et al., 1995; Popp et al., 1998).

In this work, we provide both $\delta^{13}\text{C}_{\text{coccolith}}$ and $\delta^{13}\text{C}_{\text{org}}$ values with changes in $p\text{CO}_2$ and pH to explain the mechanisms responsible for the carbon isotope fractionation within the cell. The present study stems from several lines of evidence that the isotopic offset between coccoliths of different sizes conveys a specific $p\text{CO}_2$ signal as shown in sedimentary records by the
80 work of Bolton and Stoll (2013). This empirical observation has been repeatedly reported in culture data (Hermoso, 2015; Hermoso et al., 2016b; McClelland et al., 2017; Rickaby et al., 2010). This approach used to derive palaeo- CO_2 concentrations still needs to be fully constrained ~~by the implementing of new cultures campaigns to with a view to -consolidateing the calibration~~, although a proof of concept exists (Bolton and Stoll, 2013; Godbillot et al., 2022; Hermoso et al., 2016b, 2020; Tremblin et al., 2016). Modelling studies fed by culture data have identified and quantified the main forcing parameters behind
85 the magnitude of carbon isotope vital effect in coccolith calcite: growth rate, cell size, the partitioning of CO_2 in particulate inorganic matter and particulate organic matter (PIC/POC ratio), among other ancillary parameters (McClelland et al., 2017). Four strains of geological-relevant coccolithophores, representing a wide diversity of growth rate and coccolith and cell sizes, were cultured to study how the composition of the culture medium (DIC, pH) influences the magnitude of coccoliths vital effects and to establish transfer functions between these vital effects and the aqueous CO_2 concentrations and pH. In contrast
90 with previous culture studies and to accurately mimicking the carbonate chemistry of the ocean through the Cenozoic Era, we cultured these calcifying microalgae with coupled $p\text{CO}_2$ and pH perturbations of the medium. The culture conditions varied from 200 ppmv/8.29 pH units (pre-industrial) to 1400 ppmv/7.55 pH units (thought to represent the Mid-Eocene levels) keeping all other parameters constant. Particular attention has been paid to conditions similar to those of the last 12 Myrs (narrow step of 100 ppmv for $p\text{CO}_2$ between 200 and 500 ppmv) because of the scarcity of available data.

95 **2 Materials and methods**

2.1 Strains

We cultured four coccolithophore strains kindly provided by the biological station of Roscoff: RCC1200 *Coccolithus braarudii*, RCC1323 *Helicosphaera carteri*, RCC1314 *Gephyrocapsa oceanica*, and RCC1256 *Emiliana huxleyi* (Figure 1). These species represent a wide range of coccolith taxa with various coccosphere and cell sizes. The rationale of this choice is that they belong to the most abundant groups found in Neogene pelagic sediments (Bolton et al., 2012; Claxton et al., 2022). If comparable with our own study (mode of culture, temperature, pH etc...), the results of previously-published biogeochemical work on the same species will be presented along with our own data in the figures.



105 **Figure 1: Scanning electron microscope (SEM) images of the four studied strains. Top images show the coccoliths-coccospheres and bottom images show the coccospheres-coccoliths. The scale bars represent 1 μm.**

The cultures were undertaken at the Maison de la Recherche en Environnement Naturel (ULCO-LOG) in Wimereux (Northern France) in 2023–2024.

110 **2.2 Medium preparation**

Our culture experiments were designed to represent pH and $p\text{CO}_2$ from the last greenhouse period (Eocene) to the preindustrial Holocene oceanic conditions (Rae et al., 2021; Sosdian et al., 2018). Variable $p\text{CO}_2$ between 200 and 1400 ppmv at constant alkalinity of 2300 $\mu\text{mol/kg}$ have been used to calculate the target pH (total scale) with CO2sys program (Lewis and Wallace, 1998)(Bakker et al., 2016) See table 24 for details on the chemical parameters of the culture medium.

115 Artificial seawater with a salinity of 34 psu was prepared following the ESAW recipe of Berges ~~et al., Franklin and Harrison,~~
(2001). The use of artificial seawater was preferred over natural seawater in order to facilitate the production of medium with
CO₂ level under 400 ppmv. Phosphate, nitrate, silica, Fe-EDTA, trace metals and vitamins were added to produce K/2 media
according to Keller et al. (1987). The DIC concentration was reached adding different amounts of NaHCO₃ (Sigma – batch
CAS 144-55-8) (Table 1). ~~The equivalence between the quantity of NaHCO₃ added and the resulting DIC concentration of~~
120 ~~DIC induced by this was checked beforehand with an Isotope and Gas Concentration Analyzer Picarro G2131-i coupled with~~
~~an Apollo SciTech DIC-δ¹³C analyser AS-D1 at LOCEAN laboratory (Sorbonne University) and by potentiometric dosing at~~
~~the SNAPO-CO2 service facility at LOCEAN laboratory (Sorbonne University).~~ The pH was adjusted with HCl and NaOH
addition until the target pH was reached. This treatment led to a change in the relative abundance of the DIC species (see
Bjerrum plot of carbonate speciation versus pH, Zeebe and Wolf-Gladrow, 2001). Lastly, the medium was sterilised with a
125 0.22 µm ~~filtration-filtering~~ step and stored without headspace in amber-coloured bottles at 15 °C in the dark.

Condition	pCO ₂ (ppmv)	pH	NaHCO ₃ added (mg/Lsw)
1	200	8.29	164.68
2	300	8.15	171.94
3	400	8.04	176.59
4	500	7.96	179.9
5	600	7.89	182.41
6	900	7.73	187.47
7	1200	7.61	190.74
8	1400	7.55	192.44

Table 1: Amount of NaHCO₃ added to prepare the different media used in this study (conditions 1 to 8). The amount of NaHCO₃ corresponds to the production of one litre of artificial seawater.

2.3 Culture growth

130 The cultures were undertaken at the Maison de la Recherche en Environment Naturel (ULCO - LOG) in Wimereux (Northern
France) in 2023 – 2024. The coccolithophore strains were first acclimatised to the medium during about 10 generations in 25
cm² polystyrene flasks. The cells were then inoculated in triplicate experiments in culture bottles of increasing size (75 cm²
polystyrene flasks, 600 mL and 2300 mL polycarbonate bottles) until the target cell number of the culture was reached. ~~We~~
~~grew the cultures until reaching~~~~The target cell number corresponds to~~ a sufficient biomass for our analyses (between 2 and 10
mg for the small cells *E. huxleyi* and *G. oceanica* and between 50 and 60 mg for the large cells *H. carteri* and *C. braarudii*),
135 while ensuring that the cultures remained adequately diluted (~~less than 8 000 cell/ml for *C. braarudii* and for *H. carteri* and~~
~~less than 20 000 cell/ml for *E. huxleyi* and for *G. oceanica*, in accordance with Riebesell et al. (2011).~~
The cell cultures were maintained at constant temperature (15 °C) with a 14/10 day-length-night cyclicity and an irradiance of
150 µmol.photons.m⁻².s⁻¹. The cells were regularly resuspended to avoid cell clustering and kept at relatively low
concentrations to maintain homogeneous carbon bioavailability and light access during all the experiment. A daily control of

Mis en forme : Indice

Mis en forme : Exposant

Mis en forme : Centré

Tableau mis en forme

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Centré

Mis en forme : Légende

Mis en forme : Indice

Mis en forme : Indice

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Non Italique

the culture health was made with a reverse optical microscope under x400 magnification. A final control of the coccoliths was made after the harvest with a Zeiss Supra 55 VP SEM at Sorbonne University.

Cell numeration and coccosphere sizes were measured using a Beckman Coulter Counter Multisizer 4e calibrated with 10.16 μm latex beads. The measurements were always performed at the same time of the day (8:30 to 10:00 am) to avoid biases related to the growth of the cell during the day phase.

The growth rate μ in day^{-1} of a microorganism culture corresponds to the increase of the cell number by time units commonly calculated with the formula:

$$\mu (\text{day}^{-1}) = \frac{\ln(c_{\text{final}}) - \ln(c_{\text{initial}})}{t_{\text{final}} - t_{\text{initial}}}, \quad (1)$$

where c is the cell concentration and $t_{\text{final}} - t_{\text{initial}}$ is the number of days between the initial and the final cell concentration measurements. To take into account all the concentration measurements, the μ was derived from the slope of the linear

regression of the function $f(t_x - t_{\text{initial}}) = \ln(c_x)$ where x is the day of the measurement of c_x .

2.4 Isotopic reference of the culture medium

The $\delta^{13}\text{C}$ of the Dissolved Inorganic Carbon ($\delta^{13}\text{C}_{\text{DIC}}$) and the DIC concentrations were measured in 11 samples of various CO_2/pH with an Isotope and Gas Concentration Analyzer Picarro G2131-i coupled with an Apollo SciTech DIC- $\delta^{13}\text{C}$ analyser AS-D1 at LOCEAN laboratory (Sorbonne University). The medium samples were preserved in 500 mL glass bottle with ground neck with 0,3 mL saturated HgCl_2 solution. The measurements are calibrated with in-house standards. Since we used the same NaHCO_3 powder batch for the different experiments, all the culture media had the same $\delta^{13}\text{C}_{\text{DIC}}$ values (-12.17‰ V-PDB (± 0.07)).

The $\delta^{18}\text{O}$ of seawater was measured by cavity ring down spectroscopy (CRDS) using a Picarro instrument (model L2130-i Isotopic H_2O) at LOCEAN, Sorbonne University. The in-house standards (freshwater) are calibrated using IAEA references V-SMOW (Vienna Standard Mean Ocean Water) and GISP (Greenland Ice Sheet Precipitation). The $\delta^{18}\text{O}$ of the inorganic reference ($\delta^{18}\text{O}_i$) in ‰ V-PDB is calculated from the $\delta^{18}\text{O}_{\text{sw}}$ and temperature (15°C) according to the equation of Kim and O'Neil 1997 modified by Tremblin, et al. (2016). The $\delta^{18}\text{O}_{\text{sw}}$ slightly changed between successive culture campaigns: -6.44‰ V-SMOW (± 0.03) for *C. braarudii* and *G. oceanica*, -6.67‰ V-SMOW (± 0.03) for *H. carteri* and -6.36‰ V-SMOW (± 0.15) for *E. huxleyi*.

The oxygen vital effects are calculated as the difference between the $\delta^{18}\text{O}$ of the coccoliths (VE^{18}O) in ‰ V-PDB and the $\delta^{18}\text{O}$ of a theoretical inorganic calcite ($\delta^{18}\text{O}_i$) in ‰ V-PDB:

$$\text{VE}^{18}\text{O} = \delta^{18}\text{O}_{\text{coccoliths}} - \delta^{18}\text{O}_i, \quad (2)$$

2.5 Isotopic analysis on calcite

Two methods were used to collect the culture residues for further isotopic measurements of coccolith calcite:

170 - RCC1200 *C. braarudii* and RCC1323 *H. carteri* culture residues were centrifuged during 15 minutes at 4500 rpm in 500 mL centrifuge bottles. After removing the supernatant, samples were rinsed with three cycles of centrifugation with replacement of the supernatant with neutralised demineralised water and homogenisation between each cycle. The rinsing is employed to remove the salt from the culture water. The culture residues were finally dried at 40 °C after the removal of the last supernatant.

- RCC1256 *E. huxleyi* and RCC1314 *G. oceanica* samples were collected and rinsed three times with neutralised demineralised water onto polycarbonate filtering membranes, and then dried in a desiccator. We applied a different harvesting method for those strains because of their small size, that induced too much material loss during the centrifugation process. All culture residues were stored at 5 °C after harvest.

Isotopic measurements on calcite were performed at the ISTE-P laboratory (Sorbonne University). Between 30 and 60 µg of the samples were digested with 100% phosphoric acid at 70°C in pre-evacuated vials using a Kiel IV carbonate device. The evolved CO₂ was purified in a cryogenic trapping system and carbon and oxygen isotope compositions were measured in an isotope-ratio mass spectrometer DELTA V advantage (Thermo Scientific) with a dual inlet introduction system. The carbon and oxygen isotope composition were expressed in the delta notation as a value relative to the Vienna Pee Dee Belemnite (V-PDB) and reported in ‰. $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values were calibrated using NBS-19 and NBS-18 international standards. Internal reproducibility and accuracy were monitored by replicate analysis of in-house calcite standard Marceau ($\delta^{13}\text{C} = 2.12\text{‰}$ and $\delta^{18}\text{O} = -1.87\text{‰}$), being the measured values for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ $2.10 \pm 0.08\text{‰}$ (1 σ) and $-1.89 \pm 0.10\text{‰}$ (1 σ), respectively. The external reproducibility, obtained from replicate analysis of the samples, is better than 0.05‰ (1 σ) for $\delta^{13}\text{C}_{\text{coccolith}}$ and 0.10‰ (1 σ) for $\delta^{18}\text{O}_{\text{coccolith}}$. Transfer equations between coccolith isotope ratios and CO₂ levels are proposed in this study. The residual errors are evaluated through Monte Carlo analysis code with 1,000,000 iterations and an uncertainty of 0.17‰ for the differential vital effect between small and large coccolithes $\Delta\delta^{13}\text{C}_{\text{small-large}}$ (1 σ).

190 2.6 Carbon isotope analysis of the organic matter

The culture residues of *C. braarudii*, *G. oceanica* and *E. huxleyi* were gathered onto glass microfiber filters (GFF), previously burnt at 400 °C for 3 hours to remove all trace of residual organic matter. After gathering of the culture residues, the GFF filters were rinsed three times with neutralized demineralized (Millipore®) water (adjusted to pH 7 by adding sodium hydroxide) neutralised demineralised water and then stored at -18 °C. We then gently scratched the superficial part of the GFF to collect all the culture residue. The culture residues were decarbonated with hydrochloric acid 2N overnight. The samples were then rinsed to eliminate the excess of hydrochloric acid and released Ca²⁺ and alkalinity, and then dried at 35 °C. Between 10 and 50 µg of decarbonated culture residues (only organic matter) were weighted and loaded into tin capsules.

The carbon isotope composition of the organic matter was measured at the Stable Isotope Geochemistry laboratory at IPGP (Institut de Physique du Globe de Paris). A Flash EA 1112 elemental analyser coupled in continuous helium flow to an isotope ratio mass spectrometer Thermo Fisher Scientific DELTA V Plus was used for the *C. braarudii* samples cultured between 200 and 500 ppmv (8.29 and 7.96 pH units). For *G. oceanica* and *E. huxleyi* for all culture conditions, and for *C. braarudii* between 500 and 1400 ppmv (7.96 and 7.55), we used an Elementar ~~vario~~-Vario PYRO cube analyser coupled with the same

Mis en forme : Exposit

spectrometer DELTA V Plus at IPGP. Three organic-rich internal standards, calibrated against international standards, were used to calculate the $\delta^{13}\text{C}$ values of the samples, reported in ‰ with respect to V-PDB. Replicate analysis of standards yielded an internal reproducibility better than 0.10‰ (1 σ). Accuracy was assessed by measuring an in-house Tyrosine standard ($\delta^{13}\text{C}$ = -23.23‰), for which the estimated error was better than 0.20‰. The external reproducibility for $\delta^{13}\text{C}_{\text{org}}$, obtained from replicate analysis of the samples, is better than 0.07‰.

2.7 Carbon content analysis

The PIC:POC ratios of the four strains studied under various $p\text{CO}_2$ /pH conditions were measured at the Wimereux marine station using a Thermo Fisher Flash 2000 elemental analyzer. The culture residues were collected and stored in the same way for analyses of the $\delta^{13}\text{C}_{\text{org}}$. The calibration used for reconstructing carbon content was done with various amounts of acetanilide with known amount of carbon and nitrogen. Samples for POC analysis were previously acidified. The filters were encapsulated in a tin disk before analysis.

Culture conditions	Temperature (°C)	Salinity (psu)	Light period length (h)	Dark period length (h)	pH (total scale)	TA (μmol/kg)	DIC (μmol/kg)	CO ₂ (μmol/kg)	HCO ₃ ⁻ (μmol/kg)	CO ₃ ²⁻ (μmol/kg)	pCO ₂ (ppmv)
	controlled (incubator)	measured	controlled (incubator)	controlled (incubator)	Adjusted / measured	calculated (CO2sys)	calculated (CO2sys) <i>calculated (CO2sys) Fixed with NaHCO₃</i>	calculated (CO2sys)	calculated (CO2sys)	calculated (CO2sys)	initial target
1	15	33.86	14	10	8.29	2256.0	1913.6	7.51	1669.3	236.9	200
2	15	33.86	14	10	8.15	2275.0	2011.6	11.27	1813.9	186.5	300
3	15	33.86	14	10	8.04	2248.7	2042.2	15.03	1877.4	149.8	400
4	15	33.86	14	10	7.96	2272.4	2100.3	18.78	1951.9	129.6	500
5	15	33.86	14	10	7.89	2272.2	2128.8	22.54	1993.6	112.6	600
6	15	33.86	14	10	7.73	2269.0	2183.6	33.81	2068.9	80.9	900
7	15	33.86	14	10	7.61	2246.3	2199.7	45.08	2092.6	62.0	1200
8	15	33.86	14	10	7.55	2262.2	2233.8	52.59	2126.3	54.9	1400

Table 1: Experimental parameters of the mediums. The salinity has been measured in the initial medium. The temperature and the light and dark period length are controlled by incubator set up. The total alkalinity (TA) and the DIC, CO₂, HCO₃⁻ and CO₃²⁻ concentrations are calculated by CO2sys program. $p\text{CO}_2$ and pH correspond to the culture conditions tested.

3 Results

3.1 Physiological parameters and carbon content

3.1.1 Growth rates

The growth rates of the four strains do not show any statistical trend with changes in the carbonate chemistry of the medium (*C. braarudii*: $r^2 = 0.04$ and $p > 0.05$, *H. carteri*: $r^2 = 0.08$ and $p > 0.05$, *G. oceanica*: $r^2 = 0.11$ and $p > 0.05$, *E. huxleyi*: $r^2 = 0.38$ and $p < 0.01$) (Figure 2). Especially, the linear regression between replicated values of growth rates and CO₂ concentration below 600 ppmv gives a p-value of 0.052 and a r^2 of 0.41. More than 85% of the data are comprised within a $\pm 15\%$ relative range of the mean growth rate for each strain (coloured band in Fig. 2). *E. huxleyi* and *G. oceanica* have the higher mean

Tableau mis en forme

Mis en forme : Indice

Mis en forme : Indice

Mis en forme : Exposant

growth rate (0.87 d^{-1} and 0.67 d^{-1} respectively), while the larger *C. braarudii* and *H. carteri* have a mean growth rate of 0.58 d^{-1} and 0.24 d^{-1} , respectively (Figure 2). These absolute values are similar to those obtained in previous studies for *E. huxleyi*, *G. oceanica* and *C. braarudii* (Phelps et al., 2021; Rickaby et al., 2010). *C. braarudii* has been reported with decreasing growth rates with rising CO_2 levels (Hermoso et al., 2016b; Rickaby et al., 2010).




Figure 2: Specific growth rate (μ) of the four cultured strains with respect to CO_2 level and pH of the culture media. The empty dots show all the replicate data and the filled dots denote the mean growth rates. Growth rates are given in d^{-1} (left axis) and as a relative deviation of the mean growth rate of each strain (right axis). The coloured bands correspond to the range of $\pm 15\%$ of variation from the mean growth rate. More than 85% of the data are comprised within this range.

3.1.2 Coccosphere sizes

The sizes of the coccospheres – i.e., the coccolithophore exoskeletons surrounding the cells, composed of coccoliths – approximate the evolution of cell size. The evolution of this parameter with CO₂ and pH varies among the different strains ~~varies with the CO₂/pH treatments~~. *C. braarudii* displays the largest coccospheres amongst the cultured strains. The mean coccosphere diameter of *C. braarudii* increases from 16.7 to 18.4 µm with increasing CO₂ levels from 200 to 600 ppmv and decreasing pH from 8.29 to 7.89 units of pH. The +1.70 µm increase in the coccosphere size represent a relative variation of 10.2% ($r^2 = 0.86$, $p < 0.01$). For higher $p\text{CO}_2$ and lower pH, the distribution of the coccosphere sizes is not linked with the carbonate chemistry (Figure 3). The size of *H. carteri* presents a hyperbolic trend ($r^2 = 0.72$, $p < 0.01$) of the coccosphere size with increasing CO₂ level and decreasing pH (+0.69 µm from 200 ppmv/8.29 pH units to 600 ppmv/7.89 pH units, and -1.05 µm from 600 ppmv/7.89 pH units to 1400 ppmv/7.55 pH units) (Figure 3). The size of the coccospheres produced by *G. oceanica* decreases from 8.0 to 7.5 µm with increasing CO₂ levels from 200 to 1200 ppmv and decreasing pH from 8.29 to 7.55 (-0.51 µm, a relative variation of 6.4%, $r^2 = 0.67$, $p < 0.01$) (Figure 3). The coccosphere size of the small species *E. huxleyi* exhibit a +0.22 µm increase with increasing $p\text{CO}_2$ and decreasing pH until 600 ppmv/7.89 pH units (a relative variation of 4.4%, $r^2 = 0.77$, $p < 0.001$). At higher $p\text{CO}_2$ and lower pH conditions, the coccosphere size is constant ($r^2 = 0.04$, $p > 0.05$) (Figure 3). These absolute values and trends in coccosphere size with changes in CO₂ level and pH are in line with previously reported coccosphere and cell sizes from previous culture studies (Hermoso, 2015; Hermoso et al., 2014, 2016b; Phelps et al., 2021; Rickaby et al., 2010).

Mis en forme : Indice





Figure 3: Evolution of the coccosphere sizes of the four cultured strains as a function of the CO_2 and pH culture conditions. The coccosphere sizes are given as the mean diameter of the coccospheres. They are given in μm (left axis) and as a percent relative to the mean coccosphere size of each strain (right axis). The empty points show the replicate data and the filled points show the mean coccosphere sizes.

3.1.3 PIC:POC ratios

The PIC:POC ratio allows quantifying the respective allocation of carbon in its two fixation pathways (photosynthesis and calcification), which has isotopic implications on organic matter and coccoliths. The PIC:POC ratios of the four species do not show any statistically significant trend with changes in carbonate chemistry. Among the species cultured, *C. braarudii* has the

highest PIC:POC ratio (mean value of 2.23 ± 0.49), followed by *H. carteri* (mean value of 1.44 ± 0.35), *G. oceanica* (0.98 ± 0.20) and *E. huxleyi* (0.50 ± 0.09) (Figure 4). Those data are consistent with previously published PIC:POC data (Krug et al., 2011; Langer et al., 2006; Müller et al., 2010; Rickaby et al., 2010; Riebesell et al., 2000).



270 **Figure 4: Particulate inorganic (square) and organic (triangle) carbon in ng per cell for each of the four species through the $p\text{CO}_2$ interval of the study (lower x-axis) and the aqueous CO_2 (upper x-axis). The PIC:POC ratios are also shown. The empty dots are all the replicate data and the filled points are the mean values for each species/condition.**

3.2 Carbon isotope ratios of coccolith calcite

275 The species *E. huxleyi* does not show any significant trend in $\delta^{13}\text{C}_{\text{coccolith}}$ with changes in $p\text{CO}_2$ and pH ($r^2 = 0.16$, $p > 0.05$). The taxon *H. carteri* exhibits a minor increase of less than 0.5‰ in $\delta^{13}\text{C}_{\text{coccolith}}$ with increasing CO_2 level et decreasing pH ($r^2 = 0.49$, $p < 0.001$) (Figure 5). A key feature of the dataset pertains to the biogeochemical response of *C. braarudii*. The distribution of the isotope data with changes in carbonate chemistry can be divided into two distinct trends for *C. braarudii* and *G. oceanica*. The first trend corresponds to low CO_2 level from 200 to 500 ppmv and high pH from 8.29 to 7.96. The second stage corresponds to high CO_2 level from 600 to 1400 ppmv and low pH from 7.89 to 7.55 pH units. At low CO_2 levels and high pH, the $\delta^{13}\text{C}_{\text{coccolith}}$ of *C. braarudii* increases with increasing $p\text{CO}_2$ and decreasing pH (+2.41‰ V-PDB, a relative variation of 14%, $r^2 = 0.83$, $p < 0.001$). Regarding *G. oceanica*, it exhibits a small $\delta^{13}\text{C}_{\text{coccolith}}$ increase with increasing $p\text{CO}_2$ and decreasing pH (+0.41‰ V-PDB, a relative variation of 3%, $r^2 = 0.61$, $p < 0.01$). Compared to low $p\text{CO}_2$ levels and high pH, the $\delta^{13}\text{C}_{\text{coccolith}}$ of *C. braarudii* increases with a less steep slope at high CO_2 levels and low pH (+0.92‰ V-PDB, a relative variation of 7%, $r^2 = 0.90$, $p < 0.001$). The $\delta^{13}\text{C}_{\text{coccolith}}$ of *G. oceanica* is steady at high $p\text{CO}_2$ and low pH ($r^2 = 0.10$, $p > 0.05$). These $\delta^{13}\text{C}$ data do not exhibit any consistent trend with changes in growth rate, nor with PIC and POC. *G. oceanica* and *E. huxleyi* show a carbon vital effect similar to that obtained in the work of Rickaby et al. (2010) and Hermoso et al. (2016). It is also similar to the dataset published in McClelland et al. (2017). The vital effects of *C. braarudii* in this study have a similar evolution with previously published results (Hermoso et al., 2016b; Rickaby et al., 2010) with a 1 ‰ shift towards more negative values (Figure 5). The $\delta^{13}\text{C}$ ratios of *C. braarudii* are relatively similar to the results of Chauchan and Rickaby (2024), albeit with slightly lower absolute values in Chauchan and Rickaby's this latter work. The $\delta^{13}\text{C}$ values observed by Chauchan and Rickaby (-2024) for *G. oceanica* and for *E. huxleyi* are very negative for CO_2 concentrations greater than 20 $\mu\text{mol/kg}$ ($\delta^{13}\text{C}_{\text{coccolith}}$ - $\delta^{13}\text{C}_{\text{DIC}}$ values between -2.5 and -8 ‰ V-PDB), which is very surprising contrasting with in comparison with the previous studies published in literature (Rickaby et al., 2010, Hermoso et al., 2016, McClelland et al., 2017).

295 One of the results of thisour study is that growth rates remain unchanged despite the imposed changes in CO_2 concentration and pH for all strains, and that cell size remains unchanged despite imposed changes in CO_2 concentration and pH for the small strains (namely *G. oceanica* and *E. huxleyi*). It thus turns out that the CO_2 concentration is the only parameter that dictates the carbon limitation here (Figures A3 and Figure A4). This is why we initiallyoriginally decided not to represent the isotopic data as a function of the carbon limitation (Bidigare et al., 1997), but only as a function of CO_2 level (Figures A3 and A4).

300

- Mis en forme : Exposant
- Mis en forme : Police :Italique
- Mis en forme : Exposant
- Mis en forme : Police :Italique
- Mis en forme : Police :Italique
- Mis en forme : Indice
- Mis en forme : Exposant
- Mis en forme : Indice
- Mis en forme : Exposant
- Mis en forme : Indice
- Mis en forme : Indice
- Mis en forme : Indice
- Mis en forme : Police :Italique
- Mis en forme : Police :Italique
- Mis en forme : Indice
- Mis en forme : Indice




Figure 5: Evolution of the carbon isotopic ratios of the four studied strains with increasing CO₂ concentration and decreasing pH. The data are represented as the $\delta^{13}\text{C}$ difference-fractionation between coccoliths and DIC. The empty points are all the replicate data and the filled points are the means (circles: *Coccolithus braarudii*, RCC 1200; diamonds: *Gephyrocapsa oceanica*, RCC 1314; squares: *Helicosphaera carteri*; pentagons: *Emiliania huxleyi*). The grey band represents the carbonate chemistry condition where a shift in the carbon and oxygen isotope ratio occurs (between 500 and 600 ppmv). The pH indicated below only refers to the pH of the cultures of this study. The sources of previously reported data (smaller symbols) are inset.

3.3 Oxygen isotope ratios of coccolith calcite

The $\delta^{18}\text{O}_{\text{coccolith}}$ of *G. oceanica* is constant regardless ambient conditions ($r^2 = 0.02$, $p > 0.05$), as for $\delta^{18}\text{O}_{\text{coccolith}}$ values of *E. huxleyi* ($r^2 = 0.15$, $p > 0.05$). The mean magnitude of the oxygen vital effects of *G. oceanica* and *E. huxleyi* are positive (+1.05‰ and +1.92‰, respectively), in good agreement with previously published data (Dudley et al., 1986; Hermoso et al., 2016b, a; Rickaby et al., 2010; Stevenson et al., 2014; Ziveri et al., 2003).

As already shown in the literature, *H. carteri* have a $\delta^{18}\text{O}_{\text{coccolith}}$ close to the inorganic reference (mean $\delta^{18}\text{O}_{\text{coccolith}}$ of +0.42‰ (Ziveri et al., 2003). Indeed, at low CO₂ concentrations and high pH, the $\delta^{18}\text{O}_{\text{coccolith}}$ of *H. carteri* is slowly decreasing (-0.22‰,

Mis en forme : Français (France)

a relative variation of 4%, $r^2 = 0.57$, $p < 0.01$). From 600 ppmv and 7.89 pH units, the $\delta^{18}\text{O}_{\text{coccolith}}$ of *H. carteri* is stable ($r^2 \approx 0$, $p > 0.05$) (Figure 6).

The $\delta^{18}\text{O}_{\text{coccolith}}$ of *C. braarudii* is constant at low $p\text{CO}_2$ and high pH with a mean value of $-0.81 \pm 0.13\text{‰}$ V-PDB and is also constant at high $p\text{CO}_2$ and low pH with a mean value of $-0.37 \pm 0.12\text{‰}$ V-PDB. A significant difference in the *C. braarudii* VE^{18}O is registered between high and low CO_2 ambient conditions in the culture medium. This $+0.5\text{‰}$ shift occurs between 500 and 600 ppmv. While we present evidence of a negative vital effect of coccoliths for *C. braarudii*, previous studies have shown a $\delta^{18}\text{O}_{\text{coccolith}}$ close to that of the inorganic for this species. (Hermoso, 2015; Hermoso et al., 2016a; Rickaby et al., 2010; Stevenson et al., 2014).




Figure 6: Evolution of the oxygen isotopic ratios of the four studied strains with increasing CO_2 concentration and decreasing pH. The data are represented as the vital effect (left axis) and as the isotopic difference-fractionation between the seawater and the coccoliths $\delta^{18}\text{O}$ (right axis). The empty points are all the replicate data and the filled points are the means (circles: *Coccolithus braarudii*, RCC 1200; diamonds: *Gephyrocapsa oceanica*, RCC 1314; squares: *Helicosphaera carteri*; pentagons: *Emiliania huxleyi*). The vertical gray band represents the carbonate chemistry condition where a shift in the carbon and oxygen isotope ratio occurs. The black line shows the calculated isotope ratio of an inorganic calcite (see Materials and methods). The pH scales indicated below only refers to the data published in this study. The sources of previously reported data (smaller symbols) are inset.

Mis en forme : Français (France)


Mis en forme : Français (France)

Mis en forme : Français (France)

3.4 Carbon isotope ratios of organic matter

335 The carbon isotopic ratio of organic matter of *C. braarudii* is decreasing with increasing CO₂ level (from 200 to 500 ppmv) and decreasing pH (from 8.29 to 7.89 pH units) ($r^2 = 0.93$, $p < 0.001$). This result is mirrored by a large $\delta^{13}\text{C}_{\text{cocolith}}$ increase (+2.4‰ V-PDB) (Figure 5). Above 600 ppmv (below 7.89 pH units), $\delta^{13}\text{C}_{\text{org}}$ of *C. braarudii* are stable and close to the values observed for the 200 ppmv/8.29 pH units condition ($-30.77 \pm 0.36\text{‰}$ V-PDB) (Figure 7). $\delta^{13}\text{C}_{\text{org}}$ of *G. oceanica* do not change with changes in $p\text{CO}_2$ and pH (mean of $-29.12 \pm 0.41\text{‰}$ V-PDB), similar to what was described for the changes in $\delta^{13}\text{C}_{\text{cocolith}}$.

340



345 Figure 7: Carbon isotopic ratios of organic matter compared to $\delta^{13}\text{C}_{\text{CO}_2(\text{aq})}$ of the strains studied with changes in CO₂ level and pH. The empty points are all the replicate data and the filled points are the means. The grey band represents the carbonate chemistry condition where a shift in the carbon and oxygen isotope ratio occurs. The sources of previously reported data (smaller symbols) are inset.

4 Discussion

The aim of our study is to quantify the environmental forcing exerted by CO₂ availability and ambient pH on the expression of vital effects of the coccolithophores with a view to develop and further palaeoenvironmental proxies. Our experimental results shed light onto the link between the environment, cellular growth and the efficiency of carbon fixation, as these parameters collectively control the expression of biologically-induced fractionation in organic and inorganic calcite. Published literature has revealed the role of carbon acquisition throughout the cell membrane and intracellular utilisation of carbon on the magnitude of the vital effect through laboratory and modelling studies (Laws et al., 1995; McClelland et al., 2017; Popp et al., 1998; Rau et al., 1996) Bolton and Stoll (2013) coined the concept of the demand-to-supply ratio to characterise the isotopic implication of carbon trafficking within the cells with large implication of the carbon isotope vital effect at different timescales. ~~From~~ Based on these studies, it ~~has emerged~~ appears that the carbon isotopic composition of coccolith calcite was primarily controlled by the interplay ~~between~~ of the inorganic carbon fixation in the coccolith vesicles and ~~that of~~ organic carbon fixation in the chloroplast. At constant source of external carbon, the main driver dictating the $\delta^{13}\text{C}_{\text{coccolith}}$ is the amount and isotopic composition of organic matter produced through photosynthetic carbon fixation (POC/cell and $\delta^{13}\text{C}_{\text{org}}$ respectively) (McClelland et al., 2017).

As the foremost finding of our study, we show that *C. braarudii* exhibits a large $\delta^{13}\text{C}_{\text{coccolith}}$ increase (+2.4‰ V-PDB) mirrored by a $\delta^{13}\text{C}_{\text{org}}$ decrease (-2.6‰ V-PDB) from 200 ppmv and 8.29 pH units to 500 ppmv and 7.96 pH units (low $p\text{CO}_2$ and high pH conditions) (Figure 8). The correlation between $\delta^{13}\text{C}_{\text{coccolith}}$ and $\delta^{13}\text{C}_{\text{org}}$ of *C. braarudii* is not seen at high $p\text{CO}_2$ and low pH (600 to 1400 ppmv/7.89 to 7.55 pH units) (Figures 5 and 7). Based on this observation, we will treat separately the data obtained at low $p\text{CO}_2$ /high pH from those obtained at high $p\text{CO}_2$ /low pH in the following sections. It is also noteworthy that the small species *G. oceanica* and *E. huxleyi* do not show statistically-significant changes in ~~the magnitude of~~ $\delta^{13}\text{C}_{\text{coccolith}}$ ~~fractionation neither in coccolithophores calcite nor in the organic matter or~~ $\delta^{13}\text{C}_{\text{org}}$ ~~values coeval~~ with changes in CO₂ level and pH (neither at low $p\text{CO}_2$ and high pH, nor over the whole interval). ~~The invariant nature of inorganic and bulk organic isotopic composition with our CO₂ and pH treatment is an important result for research dealing with es-about-alkenone carbon isotope fractionations and would require closer examination by molecular geochemists.~~

4.1 The biogeochemical causes for the changes in $\delta^{13}\text{C}_{\text{coccolith}}$ of *C. braarudii* at low $p\text{CO}_2$ and high pH

Given that organic compounds are significantly ¹²C-enriched relative to CO₂ with typical $\delta^{13}\text{C}_{\text{org}}$ values around -25‰, the more organic matter produced, the isotopically heavier the residual intracellular carbon pool. As calcification derives from this latter pool, coccoliths produced by highly photosynthetic coccolithophore cells exhibit relatively higher $\delta^{13}\text{C}$ values. This phenomenon is particularly expressed in small cells such as *E. huxleyi* and *G. oceanica* that are characterised by high POC contents and the highest $\delta^{13}\text{C}$ values (Ziveri et al., 2003) (Figures 4 and 5). We can hypothesise that a change in the efficiency of POC production with changes in carbon availability can induce a change in $\delta^{13}\text{C}_{\text{coccolith}}$ values, as it is apparent for *C. braarudii* with +2.4‰ V-PDB shift between low and high $p\text{CO}_2$ conditions (Figure 5). There is no apparent change in POC,

Mis en forme : Indice

380 PIC/POC or growth rates with the increase in $\delta^{13}\text{C}$ at low $p\text{CO}_2$ and high pH (Figures 2 and 4), despite an increase in chlorophyll
a concentration within the cells between 200 and 500 ppmv (Figure A2). These observations exclude a control of the amount
of organic matter produced on the isotopic signature of the internal pool. Furthermore, a change in the efficiency of POC
production could not explain the coeval -2.6‰ and progressive decrease in $\delta^{13}\text{C}_{\text{org}}$ values within this low $p\text{CO}_2$ interval. Thus,
another biogeochemical process has to be sought.

385 A second means to explain coeval changes in both $\delta^{13}\text{C}_{\text{org}}$ and $\delta^{13}\text{C}_{\text{coccolith}}$ can rely on a change in the isotopic composition of
the carbon acquired by the cell (CO_2 vs. HCO_3^-). Carbon concentrating mechanisms (Giordano et al., 2005) can induce a shift
in internal carbon pool to which a contribution by HCO_3^- ions by active transport becomes significant (Bolton and Stoll, 2013).
Under this circumstance, the internal carbon may exhibit higher $\delta^{13}\text{C}$ values, as there is a typical 9‰ equilibrium fractionation
between CO_2 and HCO_3^- (Zeebe and Wolf-Gladrow, 2001). However, early biological work aiming at characterising inducible
390 CCMs in phytoplankton has revealed the lack of such active strategies of carbon acquisition in the large and ancestral
Coccolithus taxon (Nimer et al., 1999). Furthermore, the build-up of the carbon pool with a proportion of HCO_3^- would have
led to higher $\delta^{13}\text{C}$ values in both organic and inorganic pool, which is not what is observed at lowest $p\text{CO}_2$ (conversely, the
 $\delta^{13}\text{C}_{\text{coccolith}}$ values are lower) (Figures 5 and 7). These two lines of evidence rule out the CCM hypothesis.

Culture and wild coccolithophore data have revealed that the carbon isotope composition of the organic matter decreases with
395 increasing CO_2 concentration (Bidigare et al., 1997). The magnitude of carbon isotope fractionation between the CO_2 substrate
and the organic matter is thus not constant and can change with the when the amount concentration/availability of ambient CO_2
changes, forming the ground of the $\varepsilon_{\text{p-alk}}$ proxy (Pagani, 2002; Popp et al., 1998). This modulation of the carbon kinetic
fractionation by RuBisCO not only has consequences on the isotopic signature of the organic matter of which the compound-
specific as alkenones, but also on the residual carbon pool, and ultimately on coccolith calcite.

400 Culture and wild coccolithophore data have revealed that the carbon isotope composition of the organic matter decreases with
increasing CO_2 concentration (Bidigare et al., 1997). This biogeochemical control is compatible with the 2.6‰ decrease
observed in $\delta^{13}\text{C}_{\text{org}}$ of *C. braarudii* between 200 and 500 ppmv ($r^2 = 0.93$, $p < 0.001$) (Figure 7). As the inorganic and organic
carbon pool are linked at least from an isotopic perspective, we can interpret the +2.4‰ change in $\delta^{13}\text{C}_{\text{coccolith}}$ as the result of
a modulation of kinetic fractionation of the organic matter by the preferential enhanced fixation of ^{12}C in organic matter by the
405 enzyme RuBisCO —, itself caused by via the increasing alleviated CO_2 availability limitation (Figure 8).

As we do not observe any change in the growth rate nor in the POC per cell between low and high $p\text{CO}_2$ conditions, we suggest
that the cells compensate for the decrease in carbon bioavailability in a different way. Indeed, the lower surface-to-volume
ratio at 200 ppmv compared to 500 ppmv (cell diameter of $16.7 \pm 0.2 \mu\text{m}$ at 200 ppmv and $17.6 \pm 0.4 \mu\text{m}$ at 500 ppmv, in Fig.
3) may compensate for the lower supply with a lower demand to sustain growth rate (Bolton and Stoll, 2013; Rau et al., 1996).

410 This subject point will be clarified further discussed in Section 4.2 for the high CO_2 level and low pH conditions.

Mis en forme : Exposant

Mis en forme : Indice




Figure 8: Scatter plot showing $\delta^{13}\text{C}_{\text{org}}$ versus $\delta^{13}\text{C}_{\text{coccolith}}$ values with changes in $p\text{CO}_2$ and pH (circles: *Coccolithus braarudii*, RCC 1200; diamonds: *Gephyrocapsa oceanica*, RCC 1314). Grey levels correspond to growth conditions (white: 200 ppmv/8.29 pH units, light grey: 300 ppmv/8.15 pH units, dark grey: 400 ppmv/8.04 pH units, black: 500 ppmv/7.96 pH units).

4.2 Abrupt change in the isotope biogeochemistry of *Coccolithophore* cells *Coccolithus braarudii* in response to alleviated carbon limitation and enhanced proton concentration in the cell (*Coccolithus braarudii* versus smaller species)

The data for *C. braarudii*, and in particular the link between CO_2 availability and the isotopic composition of coccoliths and organic matter, cannot be explained by a uniform biogeochemical framework, as a gap is seen for *C. braarudii* between 500 and 600 ppmv (Figures 5 and 7). The data show that above 600 ppmv, there is no correlation between ambient CO_2 levels and $\delta^{13}\text{C}_{\text{org}}$. Furthermore, there is no covariation between $\delta^{13}\text{C}_{\text{org}}$ and $\delta^{13}\text{C}_{\text{coccolith}}$ values of *C. braarudii* at high $p\text{CO}_2$ and low pH, in contrast to the covariation observed at low $p\text{CO}_2$ and high pH. The latter point suggests that the organic-to-inorganic forcing that occurs at low $p\text{CO}_2$ levels no longer operates at high $p\text{CO}_2$ levels. The fact that there is no organic-to-inorganic forcing at high $p\text{CO}_2$ is presumably due to the alleviation of carbon limitation, or at least due to a lower carbon demand-to-supply ratio (Bolton et al., 2012). Likewise, coccosphere sizes exhibit no discernible trend with CO_2 concentrations at high CO_2 levels (Figure 3), in contrast to the findings at low CO_2 levels, which were tentatively attributed to a way to enhance CO_2 influx to

Mis en forme : Police :Italique

the cells. This biogeochemical feature would indicate that the forcing of CO₂ availability on the apparent ¹³C/¹²C fractionation
430 between the organic matter and calcite occurs only below 600 ppmv. Meanwhile, small species (*E. huxleyi* and *G. oceanica*)
show unchanged isotopic values with the pCO₂/pH treatments. As explained in Rickaby et al. (2010) and Hermoso et al. (2016),
the high surface to volume ratio of the small cells induces no carbon limitation at low CO₂ level, hence no impact of the change
of pCO₂ and pH on the carbon isotopic system.

Elevated pCO₂ conditions are accompanied by greater proton concentrations in the environment (more acidic conditions).
435 Calcifying phytoplankton such as the Coccolithophores has to efflux the excess protons generated by calcification by an active
process operated by transmembrane Hv channels (Kottmeier et al., 2022; Taylor et al., 2011). Previous biological studies have
demonstrated that the opening of Hv channels are affected by intracellular pH (Taylor et al., 2011). Hv channels are closed at
low pH to counter the influx of protons into the cell from the ambient environment. The pH threshold for the closure of the Hv
channels has been assessed to be within the range of 8.1 – 7.5 pH units (Kottmeier et al., 2022). Therefore, assuming that the
440 threshold stands between 7.96 and 7.89 pH units, the closure of the Hv channels is a good candidate for the 500-600 ppmv gap
apparent in our dataset (Figures 5, 6 and 7). The closure of the Hv channels can thus result in an accumulation of H⁺ ions
within the cytosol, leading to an intense decline in intracellular pH (Kottmeier et al., 2022; Taylor et al., 2011). The release of
protons by coccolithogenesis is even more influential on intracellular pH for species such as *C. braarudii*, which produces a
large amount of PIC compared to smaller cells (Figure 4). The latter cells (*G. oceanica* and *E. huxleyi*) indeed do not exhibit
445 such a gap in δ¹³C_{org} between 500 and 600 ppmv (small mean PIC values of 0.025 ±0.004 ng/cell and 0.006 ±0.001 ng/cell in
the 600-1400 ppmv interval, respectively). Interestingly, the gap in *C. braarudii* δ¹³C_{org} between low and high CO₂ levels is
not seen in δ¹³C_{coccolith}. This observation could be the consequence of a decrease in the organic/inorganic carbon pool coupling
when carbon supply is high, unlike below 500 ppmv when the small carbon pool imposes a high isotopic dependence between
organic matter and calcite through Rayleigh distillation governed by the organic carbon fixation.

450 Around the same pCO₂/pH limit as presented for δ¹³C_{coccolith} of *C. braarudii*, we observe a change in δ¹⁸O_{coccolith} values. The
mean VE¹⁸O value is -0.81 ±0.13‰ V-PDB at low pCO₂/high pH conditions, while the mean VE¹⁸O value is -0.37 ±0.12‰
V-PDB at high pCO₂/low pH conditions (Figure 6). The gap in δ¹⁸O_{coccolith} can be explained by the change in pH. When the
pH in the environment is low, it has been demonstrated that the ¹⁸O/¹⁶O equilibration time between DIC species and H₂O is
shorter compared to high pH conditions (Uzdowski et al., 1991). The isotopic equilibration between δ¹⁸O_{DIC} and δ¹⁸O_{sw} due to
455 the assimilation of isotopically heavy CO₂ into the cell is consequently more complete at low pH than at high pH. As a
consequence, δ¹⁸O_{DIC} (and consequently δ¹⁸O_{coccolith}) values are closer to δ¹⁸O_{sw} at high pCO₂ and low pH (Figure 6). The
δ¹⁸O_{coccolith} difference between high and low pH levels is consistent with a change in pH homeostasis strategy between 7.96
and 7.89 pH units that can be responsible for the δ¹³C before and after 500/600 ppmv. It has to be noted that coccolithophores
possess internal Carbonic Anhydrase (CA) (Paneth and O'Leary, 1985). CA could also accelerate the sluggish H₂O-DIC
460 equilibration isotopic time for the oxygen system, but kinetic of isotopic re-equilibration of the internal carbon pool within the
cell compartments is not unknown here.

Mis en forme : Indice

4.3 Palaeoclimatic implications of the CO₂ proxy based on $\delta^{13}\text{C}_{\text{coccolith}}$

4.3.1 $\Delta\delta^{13}\text{C}_{\text{small-large}}$ evolution with changes in CO₂ levels and pH

465 The increase in $\delta^{13}\text{C}_{\text{coccolith}}$ of *C. braarudii* with rising $p\text{CO}_2$ and decreasing pH shows potential as a proxy for reconstructing
past carbonate chemistry, and consequently atmospheric $p\text{CO}_2$. However, utilizing absolute carbon vital effects of coccoliths
for palaeo- $p\text{CO}_2$ reconstructions requires accurate knowledge of the $\delta^{13}\text{C}_{\text{inorganic}}$ (DIC) of past oceans, which remains
challenging (Hermoso et al., 2020). To overcome this issue, previous studies (Bolton et al., 2012; Bolton and Stoll, 2013;
Godbillot et al., 2022; Hermoso et al., 2016b; McClelland et al., 2017; Tremblin et al., 2016) have proposed the use of the
470 offset between the $\delta^{13}\text{C}_{\text{coccolith}}$ of small and large species, known as the differential vital effect (noted $\Delta\delta^{13}\text{C}$, Eq. (3)). Indeed,
the $\delta^{13}\text{C}_{\text{coccolith}}$ of the small cells (Noelaerhabdaceae) is steady regardless of the $p\text{CO}_2$ levels due to their relatively large carbon
pool, akin to *C. braarudii* under carbon replete conditions (at high $p\text{CO}_2$ and low pH). The coccolith-based palaeo- $p\text{CO}_2$ proxy
is independent of the isotope ratio of ambient DIC, as both small and large coccoliths are produced by cells growing in the
same shallow water. In practical terms, $\Delta\delta^{13}\text{C}_{\text{small-large}}$ can be expressed as follows:

475
$$\Delta\delta^{13}\text{C}_{\text{small-large}} = \delta^{13}\text{C}_{\text{small coccolith}} - \delta^{13}\text{C}_{\text{large coccolith}} , \quad (3)$$

where $\delta^{13}\text{C}$ is expressed in ‰ V-PDB. The small coccoliths used in this work are those of *G. oceanica* and the large coccoliths
are those of *C. braarudii*.

$\Delta\delta^{13}\text{C}$ and $p\text{CO}_2$ data highlight a linear relationship across the range of 200 to 500 ppmv (i.e., pH values between 8.29 and
7.96), that we previously explained by a modulation in the carbon isotope fractionation between $\text{CO}_{2\text{aq}}$ and the organic matter
480 (Figure 9, Equation (4)). The uncertainties associated with the constants in Eq. (4) represent the standard deviation obtained
with a linear regression model.

$$\text{CO}_{2\text{aq}} (\pm 3.17) = -4.65(\pm 0.72) \times \Delta\delta^{13}\text{C}_{G. oceanica - C. braarudii} + 26.90(\pm 2.21) , \quad (4)$$

where $\text{CO}_{2\text{aq}}$ is expressed in $\mu\text{mol.kg}^{-1}$ and $\Delta\delta^{13}\text{C}$ is expressed in ‰ V-PDB.




Figure 9: Calibration between CO_2 and $\Delta\delta^{13}\text{C}_{\text{small-large}}$ obtained from culture experiments. The results of our study are shown in red (replicate data as dots, the linear regression made between 200 and 500 ppmv corresponds to the solid line and the confidence interval is shown with dotted lines). The brown curve is the equation recalculated from Rickaby et al. (2010) dataset. The relation encompassing the 200-1400 ppmv spectrum is shown in Figure A1.

4.3.2 Palaeoclimate application of carbon isotope culture data

In this study, we **refine-further develop** a geological CO_2 -sensitive probe by the use of the $\Delta\delta^{13}\text{C}_{\text{small-large}}$ isotopic offset. This probe forms the basis for a palaeo- CO_2 proxy transferable to sedimentary records from geological periods with low CO_2 levels, such as the Neogene and the Quaternary (Bolton and Stoll, 2013; Godbillot et al., 2022; Hermoso et al., 2020). Providing a new palaeo- $p\text{CO}_2$ proxy for those periods is of key interest, as it has been shown that alkenones are less sensitive to low and medium CO_2 level changes (Badger et al., 2019). Another strength of the $p\text{CO}_2$ proxy presented here is that it is based on coccolith calcite, which can be separated from other sedimentary components according to their size ranges by microfiltering and centrifuging protocols (Minoletti et al., 2008; Stoll and Ziveri, 2002; Zhang et al., 2018, 2021).

In addition, our study provides a biogeochemical explanation for $\Delta\delta^{13}\text{C}_{\text{small-large}}$ changes with CO_2 level and pH, thus supporting the reliability of these equations for palaeoclimate applications. We also demonstrate that the $\Delta\delta^{13}\text{C}_{\text{small-large}}\text{-CO}_2$ calibration can be extended to the whole interval studied, i.e., above 600 ppmv (Figure A1). Despite the fact that the link between $\Delta\delta^{13}\text{C}_{\text{small-large}}$ and CO_2 levels is less constrained and understood at high CO_2 level than at low CO_2 levels, the relationship is statistically supported on the whole interval ($r^2 = 0.80$, $p < 0.001$) (Figure A1). Thus, this calibration can potentially be used to reconstruct palaeo- CO_2 levels throughout the entire Cenozoic era.

505 ~~It has to be mentioned that using *C. braarudii* as a culture-model for large cells for with a palaeoclimatic applications perspective~~
~~is supported by the fact that *C. pelagicus* coccoliths were present in all sediments from the equator to the mid latitudes (50°N)~~
~~in the Atlantic and Pacific oceans before the Plio-Pleistocene boundary (Sato et al., 2004). Nowadays, even if *C. braarudii* is~~
~~not (anymore) a species thriving at low latitudes.~~
~~One promising research avenue would be the study of the co-variation of CO₂/pH levels crossed with temperature changes~~
510 ~~in new culture campaigns. This approach will could have the potential to revel the synergistic effect of discrete various controls~~
~~create a data set that on cell growth rates and refine can be used with three palaeoclimate data i.e., CO₂/pH/temperature the~~
~~biogeochemical understanding of the vital effects.~~

- Mis en forme : Police :Italique
- Mis en forme : Police :Italique
- Mis en forme : Police :Italique
- Mis en forme : Indice
- Mis en forme : Indice

5 Conclusion

The combined study of the impact of CO₂ concentration and pH on the fractionation of carbon and oxygen isotopes in
515 coccolithophores (in their organic matter and calcite biominerals) provides an explanation of the cause of variations in
 $\Delta\delta^{13}\text{C}_{\text{small-large}}$. One of the major findings of this study is the coeval variation of $\delta^{13}\text{C}_{\text{coccolith}}$ and $\delta^{13}\text{C}_{\text{org}}$ in *Coccolithus braarudii*
with changes in $p\text{CO}_2$ between 200 and 500 ppmv and pH between 8.29 and 7.96. Combined with the fact that physiological
parameters (growth rates, PIC, and POC) of *C. braarudii* remain unchanged despite changes in the availability of carbon, these
results indicate that the cause of variations in $\delta^{13}\text{C}_{\text{coccolith}}$ is an environmental-driven change in the magnitude of the
520 fractionation between ambient CO₂ aq and organic matter produced by *C. braarudii*. Above 500 ppmv, and for pH values below
7.96, greater carbon availability induces isotopic decoupling between $\delta^{13}\text{C}_{\text{org}}$ and $\delta^{13}\text{C}_{\text{coccolith}}$. On the other hand, small species
exhibit no change in $\delta^{13}\text{C}_{\text{coccolith}}$ or $\delta^{13}\text{C}_{\text{org}}$ in response to changes in CO₂ levels and pH. By comparing $\delta^{13}\text{C}_{\text{coccolith}}$ of small (*G.*
oceanica or *E. huxleyi*) with those of large cells (*C. braarudii*), we have established an insightful $\Delta\delta^{13}\text{C-CO}_2$ transfer equation
relevant for the Neogene and Quaternary timeslices.




Figure A1: Calibration between CO_2 concentrations and $\Delta\delta^{13}C_{\text{small-large}}$ obtained from culture experiments at low and high CO_2 levels. The results of our study are shown in red. Replicate data are illustrated with red dots. The linear regression made between 200 and 500 ppmv corresponds to the solid line and the confidence interval is shown with dotted lines. The dotted line shows the logarithmic regression made for the data on the whole interval (200 to 500 ppmv). The brown curve is the equation recalculated from Rickaby et al. (2010) dataset.




Figure A2: Chlorophyll a concentration (pg/cell) as a function of CO₂ and pH. The empty points are all the replicate data and the filled points are the means (circles: *Coccolithus braarudii*, RCC 1200; diamonds: *Gephyrocapsa oceanica*, RCC 1314; squares: *Helicosphaera carteri*; pentagons: *Emiliania huxleyi*). Measurements of chlorophyll a concentration were conducted using fluorometry according to the SOMLIT national protocol (Yentsch and Menzel, 1963). A Turner Design Trilogy fluorometer was employed to measure the fluorescence of our samples before and after acidification. The samples were acidified with 10 μ L of hydrochloric acid per ml of acetone extract and left in the dark for 2 minutes between the two measurements. The equation used to calculate the chlorophyll concentration is as follows (Lorenzen, 1967).




Figure A3: Carbon (A) and oxygen (B) isotopic effect with changes in the carbon limitation. Carbon limitation is calculated as the product of growth rate (μ , in day^{-1}) and coccosphere volume (V , in μm^3) divided by the product of CO₂ concentration (CO₂, in $\mu\text{mol.kg}^{-1}$) and coccosphere surface (S , in μm^2), according to Bidigare et al., 1997.

Mis en forme : Légende

Mis en forme : Exposant

Mis en forme : Exposant

Mis en forme : Indice

Mis en forme : Indice

Mis en forme : Exposant

Mis en forme : Exposant

545




Figure A4: Carbon (A) and oxygen (B) isotopic effect with changes in the carbon limitation calculated as for Figure A3 with the difference that the CO_2 concentration is a constant ($25.82 \mu\text{mol.kg}^{-1}$, mean CO_2 concentration of the medium conditions in this study).

Data availability:

550

All numerical data generated in this study are included in the Supplement. The Available culture residues can be obtained upon demand (Michael.Hermoso@univ-littoral.fr).

Supplement:

The data related to this article will be available online upon publication at: <https://doi.org/10.5281/zenodo.12187457>

Author contributions:

555

This study was conceived by MH and FM. Experiments were undertaken by GLG, CG and MH. Measurements were done by GLG, CG, GD and VR. Data were analysed by GLG, MH and FM. The paper was written by GLG, FM and MH with inputs from the other authors.

Competing interests:

The authors declare that they have no conflict of interest.

Mis en forme : Légende

Mis en forme : Indice

Mis en forme : Exposant

Mis en forme : Indice

Mis en forme : Anglais (Royaume-Uni)

560 **Acknowledgements:**

We thank L. Emmanuel and A. Guittet for help with the calcite isotopic analyses and O. Boudouma for help on the SEM at the laboratory IStEP, Sorbonne University. We are also grateful to G. Reverdin, J. Demange, C. Waelbroeck and J. Fin for isotopic and DIC concentration analyses in the culture medium at LOCEAN, Sorbonne University. [Part of the AT and CT data used in this study were analyzed at the SNAPO-CO2 service facility at LOCEAN laboratory and supported by CNRS-INSU and OSU Ecce-Terra.](#) For help with isotopic analyses on organic matter, we thank M. Ader, [Virginia Rojas](#) and G. Bardoux from IPGP. Our thanks also go to I. Probert at the biological station of Roscoff for providing the strains. We acknowledge with thanks the financial support from the French *Agence Nationale de la Recherche* (ANR) – Project CARCLIM under reference ANR-17-CE01-0004-01 and from the Graduate school IFSEA under reference ANR-21-EXES-0011 (France 2030 programme). We also benefited financial support from the French *Centre National de la Recherche Scientifique* (CNRS-INSU) Project TOPCAPI within the TelluS-SYSTER programme.

References

Bach, L. T., MacKinder, L. C. M., Schulz, K. G., Wheeler, G., Schroeder, D. C., Brownlee, C., and Riebesell, U.: Dissecting the impact of CO₂ and pH on the mechanisms of photosynthesis and calcification in the coccolithophore *Emiliania huxleyi*, *New Phytologist*, 199, 121–134, <https://doi.org/10.1111/nph.12225>, 2013.

575 Badger, M. P. S., Chalk, T. B., Foster, G. L., Bown, P. R., Gibbs, S. J., Sexton, P. F., Schmidt, D. N., Pälke, H., Mackensen, A., and Pancost, R. D.: Insensitivity of alkenone carbon isotopes to atmospheric CO₂ at low to moderate CO₂ levels, *Climate of the Past*, 15, 539–554, <https://doi.org/10.5194/cp-15-539-2019>, 2019.

[Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosea, C., Harasawa, S., Jones, S. D., Nakaoka, S. I., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W. J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., Van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high quality fCO₂ data in version 3 of the Surface Ocean CO₂ Atlas \(SOCAT\), <https://doi.org/10.5194/essd-8-383-2016>, 15 September 2016.](#)

- 590 Berges, J. A., Franklin, D. J., and Harrison, P. J.: Evolution of an artificial seawater medium: Improvements in enriched seawater, artificial water over the last two decades, *J Phycol*, 37, 1138–1145, <https://doi.org/10.1046/j.1529-8817.2001.01052.x>, 2001.
- Bidigare, R., Freeman, H., Hanson, L., Hayes, M., Jasper, P., King, L., Millero, J., Popp, N., Steinberg, A., and Wakeham, G.: Consistent fractionation of ^{13}C in nature and in the laboratory: Growth-rate effects in some haptophyte algae, *Global Biogeochem Cycles*, 11, 279–292, 1997.
- 595 Bolton, C. T. and Stoll, H. M.: Late Miocene threshold response of marine algae to carbon dioxide limitation, *Nature*, 500, 558–562, <https://doi.org/10.1038/nature12448>, 2013.
- Bolton, C. T., Stoll, H. M., and Mendez-Vicente, A.: Vital effects in coccolith calcite: Cenozoic climate- pCO_2 drove the diversity of carbon acquisition strategies in coccolithophores?, *Paleoceanography*, 27, <https://doi.org/10.1029/2012PA002339>, 2012.
- 600 [Chauhan, N., Rickaby, R.E.M., Size-dependent dynamics of the internal carbon pool drive isotopic vital effects in calcifying phytoplankton, *Geochimica et Cosmochimica Acta*, Volume 373, Pages 35-51, 2024, https://doi.org/10.1016/j.gca.2024.03.033](https://doi.org/10.1016/j.gca.2024.03.033)
- Claxton, L. M., McClelland, H. L. O., Hermoso, M., and Rickaby, R. E. M.: Eocene emergence of highly calcifying coccolithophores despite declining atmospheric CO_2 , *Nat Geosci*, 2, <https://doi.org/10.1038/s41561-022-01006-0>, 2022.
- 605 Dudley, W. C., Blackwelder, P., Brand, L., and Duplessy, J. C.: Stable isotopic composition of coccoliths, *Mar Micropaleontol*, 10, 1–8, [https://doi.org/10.1016/0377-8398\(86\)90021-6](https://doi.org/10.1016/0377-8398(86)90021-6), 1986.
- Ellis, R. J.: The most abundant protein in the world, *Trends Biochem Sci*, 4, 241–244, [https://doi.org/10.1016/0968-0004\(79\)90212-3](https://doi.org/10.1016/0968-0004(79)90212-3), 1979.
- 610 Foster, G. L.: Seawater pH, pCO_2 and $[\text{CO}_2\text{-3}]$ variations in the Caribbean Sea over the last 130 kyr: A boron isotope and B/Ca study of planktic foraminifera, *Earth Planet Sci Lett*, 271, 254–266, <https://doi.org/10.1016/j.epsl.2008.04.015>, 2008.
- Giordano, M., Beardall, J., and Raven, J. A.: CO_2 concentrating mechanisms in algae: Mechanisms, environmental modulation, and evolution, *Annu Rev Plant Biol*, 56, 99–131, <https://doi.org/10.1146/annurev.arplant.56.032604.144052>, 2005.
- 615 Godbillot, C., Minoletti, F., Bassinot, F., and Hermoso, M.: Parallel between the isotopic composition of coccolith calcite and carbon levels across Termination II: developing a new paleo- CO_2 probe, *Climate of the Past*, 18, 449–464, <https://doi.org/10.5194/cp-18-449-2022>, 2022.
- Gutknecht, J., Bisson, M. A., and Tosteson, F. C.: Diffusion of carbon dioxide through lipid bilayer membranes: effects of carbonic anhydrase, bicarbonate, and unstirred layers., *J Gen Physiol*, 69, 779–794, <https://doi.org/10.1085/jgp.69.6.779>, 1977.
- 620 Guy, R. D., Fogel, M. L., and Berry, J. A.: Photosynthetic fractionation of the stable isotopes of oxygen and carbon, *Plant Physiol*, 101, 37–47, <https://doi.org/10.1104/pp.101.1.37>, 1993.
- Hermoso, M.: Control of ambient pH on growth and stable isotopes in phytoplanktonic calcifying algae, *Paleoceanography*, 30, 1100–1112, <https://doi.org/10.1002/2015PA002844>, 2015.

Hermoso, M., Horner, T. J., Minoletti, F., and Rickaby, R. E. M.: Constraints on the vital effect in coccolithophore and dinoflagellate calcite by oxygen isotopic modification of seawater, *Geochim Cosmochim Acta*, 141, 612–627, <https://doi.org/10.1016/j.gca.2014.05.002>, 2014.

Hermoso, M., Minoletti, F., Aloisi, G., Bonifacie, M., McClelland, H. L. O., Labourdette, N., Renforth, P., Chaduteau, C., and Rickaby, R. E. M.: An explanation for the ^{18}O excess in Noelaerhabdaceae coccolith calcite, *Geochim Cosmochim Acta*, 189, 132–142, <https://doi.org/10.1016/j.gca.2016.06.016>, 2016a.

Hermoso, M., Chan, I. Z. X., McClelland, H. L. O., Heux, A. M. C., and Rickaby, R. E. M.: Vanishing coccolith vital effects with alleviated carbon limitation, *Biogeosciences*, 13, 301–312, <https://doi.org/10.5194/bg-13-301-2016>, 2016b.

Hermoso, M., McClelland, H. L. O., Hirst, J. S., Minoletti, F., Bonifacie, M., and Rickaby, R. E. M.: Towards the use of the coccolith vital effects in palaeoceanography: A field investigation during the middle Miocene in the SW Pacific Ocean, *Deep Sea Res 1 Oceanogr Res Pap*, 160, <https://doi.org/10.1016/j.dsr.2020.103262>, 2020.

Jasper, J. P. and Hayes, J. M.: A carbon isotope record of CO_2 levels during the late Quaternary, *Nature*, 347, 462–464, <https://doi.org/10.1038/347462a0>, 1990.

Keller, M. D., Selvin, R. C., Claus, W., and Guillard, R. R. L.: Media for the culture of oceanic ultraphytoplankton, *J Phycol*, 23, 633–638, <https://doi.org/10.1111/j.1529-8817.1987.tb04217.x>, 1987.

Klochko, K., Kaufman, A. J., Yao, W., Byrne, R. H., and Tossell, J. A.: Experimental measurement of boron isotope fractionation in seawater, *Earth Planet Sci Lett*, 248, 276–285, <https://doi.org/10.1016/j.epsl.2006.05.034>, 2006.

Kottmeier, D. M., Chrachri, A., Langer, G., Helliwell, K. E., Wheeler, G. L., and Brownlee, C.: Reduced H^+ channel activity disrupts pH homeostasis and calcification in coccolithophores at low ocean pH, <https://doi.org/10.1073/pnas.2022>.

Krug, S. A., Schulz, K. G., and Riebesell, U.: Effects of changes in carbonate chemistry speciation on *Coccolithus braarudii*: A discussion of coccolithophorid sensitivities, *Biogeosciences*, 8, 771–777, <https://doi.org/10.5194/bg-8-771-2011>, 2011.

Langer, G., Geisen, M., Baumann, K. H., Kläs, J., Riebesell, U., Thoms, S., and Young, J. R.: Species-specific responses of calcifying algae to changing seawater carbonate chemistry, *Geochemistry, Geophysics, Geosystems*, 7, <https://doi.org/10.1029/2005GC001227>, 2006.

Laws, E. A., Popp, B. N., Bidigare, R. R., Kennicutt, M. C., and Macko, S. A.: Dependence of phytoplankton carbon isotopic composition on growth rate and $[\text{CO}_2]_{\text{aq}}$: Theoretical considerations and experimental results, *Geochim Cosmochim Acta*, 59, 1131–1138, 1995.

Laws, E. A., Popp, B. N., Bidigare, R. R., Riebesell, U., Burkhardt, S., and Wakeham, S. G.: Controls on the molecular distribution and carbon isotopic composition of alkenones in certain haptophyte algae, *Geochemistry, Geophysics, Geosystems*, 2, <https://doi.org/10.1029/2000gc000057>, 2001.

[Lewis, E. and Wallace, D.: Program Developed for \$\text{CO}_2\$ System Calculations, 1998, doi:10.2172/639712.](#)

Lorenzen, C. J.: Determination of chlorophyll and pheo-pigments: spectrophotometric equations, *Limnol Oceanogr*, 12, 343–346, <https://doi.org/10.4319/lo.1967.12.2.0343>, 1967.

McClelland, H. L. O., Bruggeman, J., Hermoso, M., and Rickaby, R. E. M.: The origin of carbon isotope vital effects in coccolith calcite, *Nat Commun*, 8, <https://doi.org/10.1038/ncomms14511>, 2017.

Minoletti, F., Hermoso, M., and Gressier, V.: Separation of sedimentary micron-sized particles for palaeoceanography and calcareous nannoplankton biogeochemistry, *Nat Protoc*, 4, 14–24, <https://doi.org/10.1038/nprot.2008.200>, 2008.

Müller, M. N., Schulz, K. G., and Riebesell, U.: Effects of long-term high CO₂ exposure on two species of coccolithophores, *Biogeosciences*, 7, 1109–1116, <https://doi.org/10.5194/bg-7-1109-2010>, 2010.

[Nimer, N. A., Ling, M. X., Brownlee, C. and Merrett, M. J.: Inorganic carbon limitation, exofacial carbonic anhydrase activity, and plasma membrane redox activity in marine phytoplankton species, *J. Phycol.* 1205, 1200–1205, 1999, <https://onlinelibrary.wiley.com/doi/10.1046/j.1529-8817.1999.3561200.x>](https://onlinelibrary.wiley.com/doi/10.1046/j.1529-8817.1999.3561200.x)

Pagani, M.: The alkenone - CO₂ proxy and ancient atmospheric carbon dioxide, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 360, 609–632, <https://doi.org/10.1098/rsta.2001.0959>, 2002.

Pagani, M., Arthur, M. A., and Freeman, K. H.: Miocene evolution of atmospheric carbon dioxide, *Paleoceanography*, 14, 273–292, <https://doi.org/10.1029/1999PA900006>, 1999.

Pagani, M., Liu, Z., LaRiviere, J., and Ravelo, A. C.: High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations, *Nat Geosci*, 3, 27–30, <https://doi.org/10.1038/ngeo724>, 2010.

[Paneth P. and O'Leary M. H., Carbon isotope effect on dehydration of bicarbonate ion catalyzed by carbonic anhydrase. *Biochemistry* 24, 5143–5147, 1985.](https://doi.org/10.1016/0021-9258(85)90001-1)

Parker, M. D. and Boron, W. F.: The Divergence, Actions, Roles, and Relatives of Sodium-Coupled Bicarbonate Transporters, *Physiol Rev*, 93, 803–959, <https://doi.org/10.1152/physrev.00023.2012>, 2013.

Phelps, S. R., Hennon, G. M. M., Dyhrman, S. T., Hernández Limón, M. D., Williamson, O. M., and Polissar, P. J.: Carbon Isotope Fractionation in Noelaerhabdaceae Algae in Culture and a Critical Evaluation of the Alkenone Paleobarometer, *Geochemistry, Geophysics, Geosystems*, 22, 1–20, <https://doi.org/10.1029/2021GC009657>, 2021.

Popp, B. N., Laws, E. A., Bidigare, R. R., Dore, J. E., Hanson, K. L., and Wakeham, S. G.: Effect of Phytoplankton Cell Geometry on Carbon Isotopic Fractionation, *Geochim Cosmochim Acta*, 62, 69–77, [https://doi.org/10.1016/S0016-7037\(97\)00333-5](https://doi.org/10.1016/S0016-7037(97)00333-5), 1998.

Rae, J. W. B., Zhang, Y. G., Liu, X., Foster, G. L., Stoll, H. M., and Whiteford, R. D. M.: Atmospheric CO₂ over the Past 66 Million Years from Marine Archives, *Annu Rev Earth Planet Sci*, 49, 609–641, <https://doi.org/10.1146/annurev-earth-082420-063026>, 2021.

Rau, G. H., Riebesell, U., and Wolf-Gladrow, D.: A model of photosynthetic ¹³C fractionation by marine phytoplankton based on diffusive molecular CO₂ uptake, *Mar Ecol Prog Ser*, 133, 275–285, <https://doi.org/10.3354/meps133275>, 1996.

Reinfelder, J. R.: Carbon concentrating mechanisms in eukaryotic marine phytoplankton, *Ann Rev Mar Sci*, 3, 291–315, <https://doi.org/10.1146/annurev-marine-120709-142720>, 2011.

Rickaby, R. E. M., Henderiks, J., and Young, J. N.: Perturbing phytoplankton: Response and isotopic fractionation with
 690 changing carbonate chemistry in two coccolithophore species, *Climate of the Past*, 6, 771–785, <https://doi.org/10.5194/cp-6-771-2010>, 2010.

Riebesell, U., Zondervan, I., Rost, B., Tortell, P. D., Zeebe, R. E., and Morel, F. M. M.: Reduced calcification of marine
 plankton in response to increased atmospheric CO₂, *Nature*, 407, 364–367, <https://doi.org/10.1038/35030078>, 2000.

695 [Riebesell, U., Fabry, V. J., Hansson, L. and Gattuso, J.-P.: Guide to best practices for ocean acidification research and data reporting, \[reprinted edition including erratum\], Publications Office of the European Union, 258p, 2011, DOI 10.2777/66906](#)

Romero, M. F., Fulton, C. M., and Boron, W. F.: The SLC4 family of HCO₃⁻ transporters, *Pflügers Arch*, 447, 495–509,
<https://doi.org/10.1007/s00424-003-1180-2>, 2004.

Sanyal, A., Hemming, N. G., Hanson, G. N., and Broecker, W. S.: Evidence for a higher pH in the glacial ocean from boron
 isotopes in foraminifera, *Nature*, 373, 234–236, <https://doi.org/10.1038/373234a0>, 1995.

700 [Sato, T., Yuguichi, S., Takayama, T., Kameo, K., Drastic change in the geographical distribution of the cold-water nannofossil *Coccolithus pelagicus* \(Wallich\) Schiller at 2.74 Ma in the late Pliocene, with special reference to glaciation in the Arctic Ocean, *Marine Micropaleontology* 52, 1–4, pp 181-19, 2004, <https://doi.org/10.1016/j.marmicro.2004.05.003>](#)

Sosdian, S. M., Greenop, R., Hain, M. P., Foster, G. L., Pearson, P. N., and Lear, C. H.: Constraining the evolution of Neogene
 ocean carbonate chemistry using the boron isotope pH proxy, *Earth Planet Sci Lett*, 498, 362–376,
 705 <https://doi.org/10.1016/j.epsl.2018.06.017>, 2018.

Stevenson, E. I., Hermoso, M., Rickaby, R. E. M., Tyler, J. J., Minoletti, F., Parkinson, I. J., Mokadem, F., and Burton, K. W.:
 Controls on stable strontium isotope fractionation in coccolithophores with implications for the marine Sr cycle, *Geochim
 Cosmochim Acta*, 128, 225–235, <https://doi.org/10.1016/j.gca.2013.11.043>, 2014.

Stoll, H. M. and Ziveri, P.: Separation of monospecific and restricted coccolith assemblages from sediments using differential
 710 settling velocity, *Mar Micropaleontol*, 46, 209–221, [https://doi.org/10.1016/S0377-8398\(02\)00040-3](https://doi.org/10.1016/S0377-8398(02)00040-3), 2002.

Taylor, A. R., Chrachri, A., Wheeler, G., Goddard, H., and Brownlee, C.: A voltage-gated H⁺ channel underlying pH
 homeostasis in calcifying Coccolithophores, *PLoS Biol*, 9, <https://doi.org/10.1371/journal.pbio.1001085>, 2011.

Tremblin, M., Hermoso, M., and Minoletti, F.: Equatorial heat accumulation as a long-term trigger of permanent Antarctic ice
 sheets during the Cenozoic, *Proceedings of the National Academy of Sciences*, 113, 11782–11787,
 715 <https://doi.org/10.1073/pnas.1608100113>, 2016.

Tripathi, A. K., Roberts, C. D., Eagle, R. A., and Li, G.: A 20 million year record of planktic foraminiferal B/Ca ratios:
 Systematics and uncertainties in pCO₂ reconstructions, *Geochim Cosmochim Acta*, 75, 2582–2610,
<https://doi.org/10.1016/j.gca.2011.01.018>, 2011.

720 Usdowski, E., Michaelis, J., Botcher, M. E., and Hoef, J.: Factors for the oxygen isotope equilibrium fractionation between
 aqueous and gaseous CO₂, carbonic acid, bicarbonate, carbonate, and water (19°C), *Zeitschrift für Physikalische Chemie*, 170,
 237–249, 1991.

Wolf-Gladrow, D. and Riebesell, U.: Diffusion and reactions in the vicinity of plankton: A refined model for inorganic carbon transport, *Mar Chem*, 59, 17–34, [https://doi.org/10.1016/S0304-4203\(97\)00069-8](https://doi.org/10.1016/S0304-4203(97)00069-8), 1997.

Yentsch, C. S. and Menzel, D. W.: A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence, *Deep-Sea Research and Oceanographic Abstracts*, 10, 221–231, [https://doi.org/10.1016/0011-7471\(63\)90358-9](https://doi.org/10.1016/0011-7471(63)90358-9), 1963.

Zeebe, R. and Wolf-Gladrow, D.: CO₂ in Seawater: Equilibrium, Kinetics, Isotopes, in: *Oceanography Book Series*, vol. 65, Amsterdam, 346, 2001.

Zhang, H., Stoll, H., Bolton, C., Jin, X., and Liu, C.: Technical note: A refinement of coccolith separation methods: Measuring the sinking characteristics of coccoliths, *Biogeosciences*, 15, 4759–4775, <https://doi.org/10.5194/bg-15-4759-2018>, 2018.

Zhang, H., Liu, C., Mejía, L. M., and Stoll, H.: Technical note: Accelerate coccolith size separation via repeated centrifugation, *Biogeosciences*, 18, 1909–1916, <https://doi.org/10.5194/bg-18-1909-2021>, 2021.

Zhang, Y. G., Pearson, A., Benthien, A., Dong, L., Huybers, P., Liu, X., and Pagani, M.: Refining the alkenone-pCO₂ method I: Lessons from the Quaternary glacial cycles, *Geochim Cosmochim Acta*, 260, 177–191, <https://doi.org/10.1016/j.gca.2019.06.032>, 2019.

Zhang, Y. G., Henderiks, J., and Liu, X.: Refining the alkenone-pCO₂ method II: Towards resolving the physiological parameter ‘b,’ *Geochim Cosmochim Acta*, 281, 118–134, <https://doi.org/10.1016/j.gca.2020.05.002>, 2020.

Ziveri, P., Stoll, H., Probert, I., Klaas, C., Geisen, M., Ganssen, G., and Young, J.: Stable isotope “vital effects” in coccolith calcite, *Earth Planet Sci Lett*, 210, 137–149, [https://doi.org/10.1016/S0012-821X\(03\)00101-8](https://doi.org/10.1016/S0012-821X(03)00101-8), 2003.